

THE STRUCTURE OF QUASI-TRANSITIVE GRAPHS AVOIDING A MINOR WITH APPLICATIONS TO THE DOMINO PROBLEM

LOUIS ESPERET, UGO GIOCANTI, AND CLÉMENT LEGRAND-DUCHESNE

ABSTRACT. An infinite graph is quasi-transitive if its vertex set has finitely many orbits under the action of its automorphism group. In this paper we obtain a structure theorem for locally finite quasi-transitive graphs avoiding a minor, which is reminiscent of the Robertson-Seymour Graph Minor Structure Theorem. We prove that every locally finite quasi-transitive graph G avoiding a minor has a tree-decomposition whose torsos are finite or planar; moreover the tree-decomposition is canonical, i.e. invariant under the action of the automorphism group of G . As applications of this result, we prove the following.

- Every locally finite quasi-transitive graph attains its Hadwiger number, that is, if such a graph contains arbitrarily large clique minors, then it contains an infinite clique minor. This extends a result of Thomassen (1992) who proved it in the (quasi-)4-connected case and suggested that this assumption could be omitted. In particular, this shows that a Cayley graph excludes a finite minor if and only if it avoids the countable clique as a minor.
- Locally finite quasi-transitive graphs avoiding a minor are accessible (in the sense of Thomassen and Woess), which extends known results on planar graphs to any proper minor-closed family.
- Minor-excluded finitely generated groups are accessible (in the group-theoretic sense) and finitely presented, which extends classical results on planar groups.
- The domino problem is decidable in a minor-excluded finitely generated group if and only if the group is virtually free, which proves the minor-excluded case of a conjecture of Ballier and Stein (2018).

1. INTRODUCTION

1.1. A structure theorem. A central result in modern graph theory is the Graph Minor Structure Theorem of Robertson and Seymour [RS03], later extended to infinite graphs by Diestel and Thomas [DT99]. This theorem states that any graph G avoiding a fixed minor has a *tree-decomposition*, such that each piece of the decomposition, called a *torso*, is close to being embeddable on a surface of bounded genus (the notions of tree-decomposition and torso will be defined in the next section). A natural question is the following: if the graph G has non trivial symmetries, can we make these symmetries apparent in the tree-decomposition? In other words, do graphs avoiding a fixed minor have a tree-decomposition as above, but with the additional constraint that the decomposition is *canonical*, i.e., invariant under the action of the automorphism group of G ? In this paper we answer this question positively for infinite, locally finite graphs G that are *quasi-transitive*, i.e., the vertex set of G has finitely many orbits under the action of the automorphism group of G . This additional restriction, which is a way of saying that the graph G is highly

All authors are partially supported by the French ANR Project GrR (ANR-18-CE40-0032). L. Esperet and U. Giocanti are partially supported by the French ANR Project TWIN-WIDTH (ANR-21-CE48-0014-01), and by LabEx PERSYVAL-lab (ANR-11-LABX-0025).

symmetric, has the advantage of making the structure theorem much cleaner: instead of being almost embeddable on a surface of bounded genus, each torso of the tree-decomposition is now simply finite or planar.

Theorem 1.1 (see Theorem 4.1). *Every locally finite quasi-transitive graph avoiding the countable clique as a minor has a canonical tree-decomposition whose torsos are finite or planar.*

The tree-decomposition in Theorem 1.1 will be obtained by refining the tree-decomposition obtained in the following more detailed version of the result, which might be useful for applications.

Theorem 1.2 (see Theorem 4.3). *Every locally finite quasi-transitive graph G avoiding the countable clique as a minor has a canonical tree-decomposition with adhesion at most 3 in which each torso is a minor of G , and is planar or has bounded treewidth.*

Interestingly, the proof does not use the original structure theorem of Robertson and Seymour [RS03] or its extension to infinite graphs by Diestel and Thomas [DT99]. Instead, we rely mainly on a series of results and tools introduced by Grohe [Gro16a] to study decompositions of finite 3-connected graphs into quasi-4-connected components, together with a result of Thomassen [Tho92] on locally finite quasi-4-connected graphs. The main technical contribution of our work consists in extending the results of Grohe to infinite, locally finite graphs and in addition, making sure that the decompositions we obtain are canonical (in a certain weak sense). Our proof crucially relies on a recent result of Carmesin, Hamann, and Miraftab [CHM22], which shows that there exists a canonical tree-decomposition that distinguishes all tangles of a given order (in our case, of order 4).

Thomassen proved that if a locally finite quasi-transitive graph has only one end, then this end must be thick [Tho92, Proposition 5.6] (see below for a definition of thick end). At some point of our proof, we also need to show the stronger result (see Proposition 4.7), of independent interest, that for any $k \geq 1$, a locally finite quasi-transitive graph cannot have only one end of degree k .

We now discuss some applications of Theorem 1.1.

1.2. Hadwiger number. As a consequence of Theorem 1.1, we obtain a result on the Hadwiger number of locally finite quasi-transitive graphs. The *Hadwiger number* of a graph G is the supremum of the sizes of all finite complete minors in G . We say that a graph G *attains its Hadwiger number* if the supremum above is attained, that is if it is either finite, or G contains an infinite clique minor. Thomassen [Tho92] proved that every locally finite quasi-transitive 4-connected graph attains its Hadwiger number, and suggested that the 4-connectedness assumption might be unnecessary. We prove that this is indeed the case.

Theorem 1.3 (see Theorem 5.1). *Every locally finite quasi-transitive graph attains its Hadwiger number.*

We will indeed prove a stronger statement, namely that every locally finite quasi-transitive graph avoiding the countable clique as a minor also avoids a finite graph with crossing number 1 as a minor.

1.3. Accessibility in graphs. We now introduce the notion of accessibility in graphs considered by Thomassen and Woess [TW93]. To distinguish it from the related notion in groups (see below), we will call it vertex-accessibility in the remainder of the paper. A *ray* in an infinite graph G is an infinite one-way path in G . Two rays of G are *equivalent* if there are infinitely many disjoint paths between them in G (note that this is indeed an equivalence relation). An *end* of G is an equivalence class of rays in G . When there is a finite set X of vertices of G , two distinct components C_1, C_2 of $G - X$, and two distinct ends ω_1, ω_2 of G such that for each $i = 1, 2$, all but finitely many vertices of all (equivalently any) rays of ω_i are in C_i , we say that X *separates* ω_1 and ω_2 . A graph G is *vertex-accessible* if there is an integer k such that for any two distinct ends ω_1, ω_2 in G , there is a set of at most k vertices that separates ω_1 and ω_2 .

It was proved by Dunwoody [Dun07] (see also [Ham18b, Ham18a] for a more combinatorial approach) that locally finite quasi-transitive planar graphs are vertex-accessible. Here we extend the result to locally finite quasi-transitive graphs excluding the countable clique K_∞ (and not necessarily K_5 and $K_{3,3}$) as a minor, and in particular to locally finite quasi-transitive graphs from any proper minor-closed family.

Theorem 1.4 (see Theorem 5.2). *Every locally finite quasi-transitive K_∞ -minor-free graph is vertex-accessible.*

1.4. Accessibility in groups. The notion of vertex-accessibility introduced above is related to the notion of accessibility in groups. Given a finitely generated group Γ , and a finite set of generators S , the *Cayley graph* of Γ with respect to the set of generators S is the edge-labeled graph $\text{Cay}(\Gamma, S)$ whose vertex set is the set of elements of Γ and where for every two elements $g, h \in \Gamma$ we put an arc (g, h) labeled with $a \in S$ when $h = a \cdot g$. Cayley graphs have to be seen as highly symmetric graphs; in particular they are transitive: the right action of the group Γ onto itself can be easily seen to induce a transitive group action on $\text{Cay}(\Gamma, S)$. It is known that the number of ends of a Cayley graph of a finitely generated group does not depend of the choice of generators, so we can talk about the number of ends of a finitely generated group. A classical theorem of Stallings [Sta72] states that if a finitely generated group Γ has more than one end, it can be split as a non-trivial free product with finite amalgamation, or as an HNN-extension over a finite subgroup. If any group produced by the splitting still has more than one end we can keep splitting it using Stallings theorem. If the process eventually stops (with Γ being obtained from finitely many 0-ended or 1-ended groups using free products with amalgamation and HNN-extensions), then Γ is said to be *accessible*. Thomassen and Woess [TW93] proved that a finitely generated group is accessible if and only if at least one of its locally finite Cayley graphs is vertex-accessible, if and only if all of its locally finite Cayley graphs are vertex-accessible.

A finitely generated group is *minor-excluded* if at least one of its Cayley graphs avoids a finite minor. Similarly a finitely generated group is *K_∞ -minor-free* if one of its Cayley graphs avoids the countable clique as a minor, and *planar* if one of its Cayley graphs is planar. Note that planar groups are minor-excluded and Theorem 1.3 immediately implies that a finitely generated group is minor-excluded if and only if it is K_∞ -minor-free.

Droms [Dro06] proved that finitely generated planar groups are finitely presented, while Dunwoody [Dun85] proved that finitely presented groups are accessible, which implies that finitely generated planar groups are accessible. Theorem 1.4 immediately implies the following, which extends this result to all minor-excluded finitely generated groups, and equivalently to all finitely generated K_∞ -minor-free groups.

Corollary 1.5. *Every finitely generated K_∞ -minor-free group is accessible.*

In fact, combining Theorem 1.1 with techniques introduced by Hamann [Ham18b, Ham18a] in the planar case, we prove the following stronger result which also implies Corollary 1.5 using the result of Dunwoody [Dun85] that all finitely presented groups are accessible.

Theorem 1.6 (see Corollary 5.7). *Every finitely generated K_∞ -minor-free group is finitely presented.*

1.5. The domino problem. We refer to [ABJ18] for a detailed introduction to the domino problem. A *coloring* of a graph G with colors from a set Σ is simply a map $V(G) \rightarrow \Sigma$. The *domino problem* for a finitely generated group Γ together with a finite generating set S is defined as follows. The input is a finite alphabet Σ and a finite set $\mathcal{F} = \{F_1, \dots, F_p\}$ of forbidden *patterns*, which are colorings with colors from Σ of the closed neighborhood of the neutral element 1_Γ in the Cayley graph $\text{Cay}(\Gamma, S)$, viewed as an edge-labeled subgraph of $\text{Cay}(\Gamma, S)$ (recall that each element of S corresponds to a different label). The problem then asks if there is a coloring of $\text{Cay}(\Gamma, S)$ with colors from Σ , such that for each $v \in \Gamma$, the coloring of the closed neighborhood of v in $\text{Cay}(\Gamma, S)$ (viewed as an edge-labeled subgraph of $\text{Cay}(\Gamma, S)$), is not isomorphic to any of the colorings F_1, \dots, F_p , where we consider isomorphisms preserving the edge-labels (equivalently, isomorphisms corresponding to the right multiplication by elements of Γ).

It turns out that the decidability of the domino problem for (Γ, S) is independent of the choice of the finite generating set S , hence we can talk of the decidability of the domino problem for a finitely generated group Γ . If we consider $\Gamma = (\mathbb{Z}^2, +)$, then the domino problem corresponds exactly to the well-known Wang tiling problem, which was shown to be undecidable by Berger in [Ber66]. On the other hand, there is a simple greedy procedure to solve the domino problem in free groups, which admit trees as Cayley graphs. More generally, the domino problem is decidable in virtually free groups, which can equivalently be defined as finitely generated groups having a locally finite Cayley graph of bounded treewidth [ABJ18, Ant11]. A remarkable conjecture of Ballier and Stein [BS18] asserts that these groups are the only ones for which the domino problem is decidable.

Conjecture 1.7 (Domino problem conjecture [BS18]). *A finitely generated group has a decidable domino problem if and only if it is virtually free.*

Recall that virtually free groups are precisely the groups having a locally finite Cayley graph of bounded treewidth. Since having bounded treewidth is a property that is closed under taking minor, it is natural to ask whether Conjecture 1.7 holds for minor-excluded groups (or equivalently, using Theorem 1.3, to K_∞ -minor-free groups). Using Corollary 1.5, together with classical results on planar groups and recent results on fundamental groups of surfaces [ABM19], we prove that this is indeed the case.

Theorem 1.8 (see Theorem 5.8). *A finitely generated K_∞ -minor-free group has a decidable domino problem if and only if it is virtually free.*

1.6. Overview of the proof of Theorems 1.1 and 1.2. Consider a locally finite quasi-transitive graph G that excludes the countable clique K_∞ as a minor. The graph G is said to be *quasi-4-connected* if it is 3-connected and for every set $S \subseteq V(G)$ of size 3 such that $G - S$ is not connected, $G - S$ has exactly two connected components and one of them consists of a single vertex. Thomassen proved that if G is quasi-4-connected, then G is planar or has finite treewidth [Tho92], which implies Theorem 1.2 in this case (with a trivial tree-decomposition consisting of a single node).

To deal with the more general case, the first step is to obtain a canonical tree-decomposition of G of adhesion at most 2 in which all torsos are minors of G that are 3-connected graphs, cycles, or complete graphs on at most 2 vertices. The existence of such a decomposition in the finite case is a well-known result of Tutte [Tut84] and was proved in the locally finite case in [DSS98]. For our proof we need to go one step further. Grohe [Gro16a] proved that every finite graph G has a tree-decomposition of adhesion at most 3 whose torsos are minors of G and are complete graphs on at most 4 vertices or quasi-4-connected graphs. A crucial step for us would be to prove a version of this result in which the tree-decomposition would be canonical, and which would hold for locally finite graphs.

However, as observed by Grohe, even in the finite case the decomposition he obtains is not canonical in general. Our main technical contribution is to extend the result of Grohe [Gro16a] mentioned in the previous paragraph to locally finite graphs, while making sure that most of the construction (except the very end) is canonical. For this, we proceed in two steps. First, we use a result of [CHM22] to find a canonical tree-decomposition of any 3-connected graph G that distinguishes all its tangles of order 4. Using this result, we show that we can assume that the graph under consideration admits a unique tangle \mathcal{T} of order 4. We then follow the main arguments from [Gro16a] and show that G has a canonical tree-decomposition of adhesion 3 which is a star and whose torsos are all minors of G and finite, except for the torso H associated to the center of the star, which has the following property: there exists a matching $M \subseteq E(H)$ which is invariant under the action of the automorphism group of G and such that the graph $H' := H/M$ obtained after the contraction of the edges of M is quasi-transitive, locally finite, and quasi-4-connected. In particular, a result of Thomassen [Tho92] then implies that H' is planar or has bounded treewidth. We then prove that even if H itself is not necessarily quasi-4-connected, it is still planar or has bounded treewidth, which is enough to conclude the proof of Theorem 1.2. The final step to prove Theorem 1.1 consists in refining the tree-decomposition to make sure that torsos of bounded treewidth are replaced by torsos of finite size (moreover, this refinement has to be done in a canonical way).

1.7. Related work. Independently of our work, Carmesin and Kurkofka [CK23] recently worked on decompositions of 3-connected graphs with an approach that differs from Grohe's approach. They obtained a canonical decomposition into basic pieces consisting in quasi-4-connected graphs, wheels or thickenings of $K_{3,m}$ for $m \geq 0$. It is possible that their approach could also imply some of the applications we describe. However it is not clear to us whether this work directly implies the

existence of canonical tree-decompositions with the properties described in Theorems 1.1 and 1.2, as they consider *mixed separations*, i.e. separations containing both vertices and edges, while we focus on vertex-separations. Vertex-separations yield tree-decompositions, while mixed separations do not yield tree-decompositions in a traditional sense.

1.8. Organization of the paper. We start with some preliminary definitions and results about graphs and groups in Section 2. Section 3 is dedicated to the study of canonical tree-decompositions. It contains the main technical contribution of the paper, a partial extension of results of Grohe [Gro16a] to infinite graphs. Section 4 contains the proof of Theorems 1.1 and 1.2. Section 5 is dedicated to the main applications of Theorems 1.1 and 1.2. We conclude with a number of open problems in Section 6.

2. PRELIMINARIES

For every $n \in \mathbb{N}$, we let $[n] := \{1, 2, \dots, n\}$.

2.1. Graphs. In what follows, we will consider undirected graphs $G = (V(G), E(G))$ which are *simple* (i.e. without loops and multi-edges), unless specifically stated otherwise. The set of vertices $V(G)$ will always be finite or infinite countable, and we will always assume our graphs to be connected. Most of the time G will be *locally finite*, meaning that every vertex has finite degree (the only graphs that will not satisfy this additional requirement will be trees of the tree-decompositions). We equip a graph G with its shortest-path metric d_G . For any set of vertices $X \subseteq V(G)$, the *neighborhood* of X in G is denoted by

$$N_G(X) := \{u \in V(G) \setminus X : \exists v \in X, uv \in E(G)\}.$$

When the graph G is clear from the context we will drop the subscript and write $N(X)$ instead of $N_G(X)$. For each $v \in V(G)$, we set $N_G(v) := N_G(\{v\})$. For every graph G and every subset of vertices $X \subseteq V(G)$, we denote by $G[X]$ the *subgraph of G induced by X* , which is the graph with vertex set X whose edge set consists of all the pairs uv such that $uv \in E(G)$. We let $G - X := G[V(G) \setminus X]$. We denote by $G\llbracket X \rrbracket$ the graph with vertex set X whose edge set consists of all the pairs uv such that $uv \in E(G)$ or there exists a connected component C of $G - X$ such that $\{u, v\} \subseteq N(C)$. The graphs $G\llbracket X \rrbracket$ are called the *torsos* of G .

For each $n \in \mathbb{N}$, we let K_n denote the complete graph with vertex set $[n]$. We let K_∞ be the *countable clique*, that is the infinite complete graph with vertex set \mathbb{N} . This graph is sometimes also denoted by K_{\aleph_0} , which is less ambiguous, but we prefer to keep the notation K_∞ as used by Thomassen in [Tho92], since we reuse a number of results proved in his paper.

Minors and models. Given two graphs G, H , we say that H is a *minor* of G if it can be obtained from G after removing some vertices and edges, and contracting edges. A *model* of H in G is a family $(V_v)_{v \in V(H)}$ of pairwise disjoint vertex subsets of G such that each V_v induces a connected subgraph of G , and for each $uv \in E(H)$, there exists $u' \in V_u, v' \in V_v$ such that $u'v' \in E(G)$. Note that H is a minor of G if and only if there is a model of H in G . When $V(H) \subseteq V(G)$, a model $(V_v)_{v \in V(H)}$ of H in G is said to be *faithful* if for each $v \in V(H), v \in V_v$. H is a *faithful minor* of G if it admits a faithful model in G . Consider for instance a graph G with a subset

of vertices $X \subseteq V(G)$ such that $G - X$ is connected and only two vertices of X (call them x and y) have a neighbor in $G - X$. Then the torso $G[[X]]$ (as defined above), is a faithful minor of G and it consists of the graph $G[X]$ with the addition of the edge xy (if it is not already present in G).

Connectedness. For every $k \geq 0$, a graph G is k -connected if it has at least $k + 1$ vertices and for every subset $S \subseteq V(G)$ of at most $k - 1$ vertices, the graph $G - S$ is connected. We recall that a graph is said to be *quasi-4-connected* if it is 3-connected and for every set $S \subseteq V(G)$ of size 3 such that $G - S$ is not connected, $G - S$ has exactly two connected components and one of them consists of a single vertex.

Rays and ends. A *ray* in a graph G is an infinite simple one-way path $P = (v_1, v_2, \dots)$. A *subray* P' of P is a ray of the form $P' = (v_i, v_{i+1}, \dots)$ for some $i \geq 1$. We say that a ray *lives* in a set $X \subseteq V(G)$ if one of its subrays is included in X . We define an equivalence relation \sim over the set of rays $\mathcal{R}(G)$ by letting $P \sim P'$ if and only if for every finite set of vertices $S \subseteq V(G)$, there is a component of $G - S$ that contains infinitely many vertices from both P and P' . When G is infinite, this is equivalent to saying that for any finite set $S \subseteq V(G)$, P and P' are living in the same component of $G - S$. The *ends* of G are the elements of $\mathcal{R}(G)/\sim$, the equivalence classes of rays under \sim . For every $X \subseteq V(G)$, we say that an end ω *lives in* X if one of its rays lives in X .

When there is a set X of vertices of G , two distinct components C_1, C_2 of $G - X$, and two distinct ends ω_1, ω_2 of G such that for each $i = 1, 2$, ω_i lives in C_i , we say that X *separates* ω_1 and ω_2 . A graph G is *vertex-accessible* if there is an integer k such that for any two distinct ends ω_1, ω_2 in G , there is a set of at most k vertices that separates ω_1 and ω_2 . The *degree* of an end ω is the supremum number $k \in \mathbb{N} \cup \{\infty\}$ of pairwise disjoint rays that belong to ω . By a result of Halin [Hal65], this supremum is a maximum i.e. if an end ω has infinite degree, then there exists an infinite countable family of pairwise disjoint rays belonging to ω . An end is *thin* if it has finite degree, and *thick* otherwise. It is an easy exercise to check that for every end ω of finite degree k and every end $\omega' \neq \omega$, there is a set of size at most k that separates ω from ω' .

The interested reader is referred to Chapter 8 in [Die17] for more background and important results in infinite graph theory.

2.2. Groups and Cayley graphs. An *automorphism* of a graph G is a graph isomorphism from G to itself (i.e., a bijection from $V(G)$ to $V(G)$ that maps edges to edges and non-edges to non-edges). The set of automorphisms of G has a natural group structure (as a subgroup of the symmetric group over $V(G)$); the group of automorphisms of G is denoted by $\text{Aut}(G)$.

For a graph G and a group Γ , we will say that Γ *acts by automorphisms on* G (or simply that Γ *acts on* G when the context is clear) if every element of Γ induces an automorphism g of G , such that the induced application $\Gamma \rightarrow \text{Aut}(G)$ is a group morphism. We will usually use the right multiplicative notation $x \cdot g$ instead of $g(x)$ for $g \in \Gamma$, $x \in V(G)$. For every $X \subseteq V(G)$, $\Gamma' \subseteq \Gamma$ and $g \in \Gamma$, we let $X \cdot g := g(X) = \{x \cdot g : x \in X\}$ and $X \cdot \Gamma' := \bigcup_{g \in \Gamma'} X \cdot g$. We denote the set of orbits of $V(G)$ under the action of Γ by G/Γ (Γ naturally induces an equivalence relation on $V(G)$, relating elements in the same orbit of Γ). For every subset $X \subseteq V(G)$ we

let $\text{Stab}_\Gamma(X) := \{g \in \Gamma : X \cdot g = X\}$ denote the *stabilizer* of X , which is always a subgroup of Γ . For each $x \in X$, we let $\Gamma_x := \text{Stab}_\Gamma(\{x\})$.

Quasi-transitive graphs. The action of a group Γ on a graph G is said to be *vertex-transitive* (or simply *transitive*) when there is only one orbit in G/Γ , i.e. when for every two vertices $u, v \in V(G)$ there exists an element $g \in \Gamma$ such that $u \cdot g = v$. The action of Γ on G is said to be *quasi-transitive* if there is only a finite number of orbits in G/Γ . We say that G is *transitive* (resp. *quasi-transitive*) if it admits a transitive (resp. quasi-transitive) group action.

It was proved, first for finitely generated groups and then in the more general graph-theoretic context, that the number of ends of a quasi-transitive graph is either $0, 1, 2$ or ∞ [Fre44, Hop44, DJM93]. A graph with a single end is said to be *one-ended*.

Finitely presented groups. A *group presentation* is a pair $\langle S|R \rangle$ where S is a set of letters called *generators* and R a set of finite words over the alphabet S called *relators*. We will always assume that S is finite and closed under taking inverse, i.e. that every generator $a \in S$ comes with an associated *inverse* $a^{-1} \in S$ (which may be equal to a) such that $(a^{-1})^{-1} = a$, and we assume that $aa^{-1} \in R$. We say that $\langle S|R \rangle$ is *finite* when both S and R are finite. The group associated to $\langle S|R \rangle$ is the group $F(S)/N(R)$, where $F(S)$ is the free group over S and $N(R)$ is the normal closure of R in $F(S)$, i.e. the set of elements of $F(S)$ of the form $(w_1 \cdot r_1 \cdot w_1^{-1}) \cdots (w_\ell \cdot r_\ell \cdot w_\ell^{-1})$ for any $\ell \in \mathbb{N}$, $r_1, \dots, r_\ell \in R$ and $w_1, \dots, w_\ell \in F(S)$. For simplicity, we will also denote this group with $\langle S|R \rangle$. Note that by definition of $\langle S|R \rangle$, since $aa^{-1} \in R$ for every element $a \in S$, the (formal) inverse a^{-1} of a in S is indeed the inverse of a in the group $\langle S|R \rangle$.

Cayley graphs. The *Cayley graph* of a finitely generated group Γ with respect to the finite set of generators S is the edge-labeled graph $\text{Cay}(\Gamma, S)$ whose vertex set is the set of elements of Γ and where for every two $g, h \in \Gamma$ we add an arc (g, h) labeled with $a \in S$ when $h = a \cdot g$. Note that $\text{Cay}(\Gamma, S)$ is always locally finite (as we assumed S to be finite), connected and that the group Γ acts transitively by right multiplication on $\text{Cay}(\Gamma, S)$. As mentioned above we will always assume that the set S of generators is symmetric, so whenever there is an arc (u, v) labeled with $a \in S$, the graph also contained the arc (v, u) labeled a^{-1} . In this case we can consider the non-labeled version of $\text{Cay}(\Gamma, S)$ as an undirected simple graph, with a single edge uv instead of each pair of arcs (u, v) and (v, u) .

We say that a finitely generated group Γ is *planar* (resp. *minor-excluded*) if it admits a finite generating set S such that $\text{Cay}(\Gamma, S)$ is planar (resp. does not contain every finite graph as a minor). Similarly, we say that Γ is K_∞ -minor-free if it admits a finite generating set S such that $\text{Cay}(\Gamma, S)$ is K_∞ -minor-free.

Accessibility. In order to give a precise definition of accessibility in groups (without using Stallings theorem and the notions of free product with amalgamation and HNN-extension explicitly as we did in the introduction), we first need to define the notion of a graph of groups. The reader is referred to [Ser80, Chapter 1] for more details on graphs of groups and Bass-Serre theory.

A *graph of groups* consists of a pair (G, \mathcal{G}) such that $G = (V, E)$ is a graph (possibly having loops and multi-edges), and \mathcal{G} is a family of *vertex-groups* Γ_v for each $v \in V$,

edge-groups Γ_{uv}, Γ_{vu} for each edge $uv \in E$, and of group isomorphisms $\phi_{u,v} : \Gamma_{uv} \rightarrow \Gamma_{vu}$ for each $uv \in E(G)$, such that Γ_{uv} and Γ_{vu} are respectively subgroups of Γ_u and Γ_v for each $uv \in E$.

Let T be a spanning tree of G and $\langle S_v | R_v \rangle$ be a presentation of Γ_v for each $v \in V$. The *fundamental group* $\Gamma := \pi_1(G, \mathcal{G})$ of the graph of groups (G, \mathcal{G}) is defined as the group having as generators the set:

$$S := (\sqcup_{v \in V} S_v) \sqcup (\sqcup_{e \in E \setminus E(T)} \{t_e\})$$

and as relations:

- the relations of each R_v ;
- for every edge $uv \in E(T)$ and every $g \in \Gamma_{uv} \subseteq \Gamma_u$, the relation $\phi_{uv}(g) = g$;
- for every edge $e = uv \in E \setminus E(T)$ and every $g \in \Gamma_{uv}$, the relation $t_e \phi_{uv}(g) t_e^{-1} = g$.

It can be shown that for a given graph of groups (G, \mathcal{G}) , the definition of its fundamental group does not depend of the choice of the spanning tree T (see for example [Ser80, Section I.5.1]).

Remark 2.1. At the beginning of this section we have defined Γ_v as the stabilizer of $\{v\}$ by the action of Γ on a graph G and the reader might be worried about a possible confusion with the notation Γ_v of vertex-groups above. On the one hand the vertex-group Γ_v has a close connection with the stabilizer of $\{v\}$ in this context, so the objects are not completely unrelated, and on the other hand vertex-groups will only be used at the very end (in Subsection 5.4, where stabilizers will not be used at all), so hopefully there should not be any risk of confusion.

A group Γ is said to be *accessible* if it is the fundamental group of a finite graph of groups G with finite vertex set $V(G)$ such that:

- the vertex-groups have at most one end, and
- the edge-groups are finite.

By Bass-Serre theory [Ser80, DD89], accessible groups are exactly those groups acting on trees without edge-inversion, such that the vertex-stabilizers have at most one end and the edge-stabilizers of the action are finite.

As mentioned in the introduction, Thomassen and Woess [TW93] obtained the following connection between accessibility in groups and vertex-accessibility in graphs.

Theorem 2.2 ([TW93]). *A finitely generated group is accessible if and only if it admits a Cayley graph which is vertex-accessible.*

It was open for a long time whether there exist finitely generated groups which are not accessible, and Dunwoody answered this question negatively in [Dun93]. On the other hand he also proved the following result.

Theorem 2.3 ([Dun85]). *Every finitely presented group is accessible.*

In particular, planar groups form a proper subclass of accessible groups.

Theorem 2.4 ([Dro06]). *Every finitely generated planar group is finitely presented, and thus accessible.*

Virtually free groups. A group is *virtually free* if it contains a finitely generated free group as a subgroup of finite index. For graphs of groups, this property can be related to the structure of the vertex groups as follows.

Theorem 2.5 ([KPS73]). *A finitely generated group is virtually free if and only if it is the fundamental group of a finite graph of groups in which all vertex-groups are finite.*

3. TREE-DECOMPOSITIONS AND TANGLES

3.1. Separations and canonical tree-decompositions. As we will be heavily relying on results of Grohe [Gro16a], we use his notation for all objects related to separations and tangles. A *separation* in a graph $G = (V, E)$ is a triple (Y, S, Z) such that Y, S, Z are pairwise disjoint, $V = Y \cup S \cup Z$ and there is no edge between vertices of Y and Z . A separation (Y, S, Z) is *proper* if Y and Z are nonempty. In this case, S is a *separator* of G .

The separation (Y, S, Z) is said to be *tight* if there are some components C_Y, C_Z respectively of $G[Y], G[Z]$ such that $N_G(C_Y) = N_G(C_Z) = S$. The *order* of a separation (Y, S, Z) is $|S|$ and the *order* of a family \mathcal{N} of separations is the supremum of the orders of its separations. In what follows, we will always consider sets of separations of finite order. We will denote $\text{Sep}_k(G)$ (respectively $\text{Sep}_{<k}(G)$) the set of all separations of G of order k (respectively less than k).

The following lemma was originally stated in [TW93] for transitive graphs, but the same proof immediately implies that the result also holds for quasi-transitive graphs.

Lemma 3.1 (Corollary 4.3 in [TW93]). *Let G be a locally finite graph. Then for every $v \in V(G)$ and $k \geq 1$, there is only a finite number of tight separations (Y, S, Z) of order k in G such that $v \in S$. Moreover, for any group Γ acting quasi-transitively on G and any $k \geq 1$, there is only a finite number of Γ -orbits of tight separations of order at most k in G .*

Canonical tree-decompositions. A *tree-decomposition* of a graph G is a pair (T, \mathcal{V}) where T is a tree and $\mathcal{V} = (V_t)_{t \in V(T)}$ is a family of subsets V_t of $V(G)$ such that:

- $V(G) = \bigcup_{t \in V(T)} V_t$;
- for every nodes t, t', t'' such that t' is on the unique path of T from t to t'' , $V_t \cap V_{t''} \subseteq V_{t'}$;
- every edge $e \in E(G)$ is contained in an induced subgraph $G[V_t]$ for some $t \in V(T)$.

Note that in our definition of tree-decomposition, we allow T to have vertices of infinite degree. The sets V_t for every $t \in V(T)$ are called the *bags* of (T, \mathcal{V}) , and the induced subgraphs $G[V_t]$ the *parts* of (T, \mathcal{V}) . The *width* of (T, \mathcal{V}) is the supremum of $|V_t| - 1$, for $t \in V(T)$. Note that the width of a tree-decomposition can be infinite. The sets $V_t \cap V_{t'}$ for every $tt' \in E(T)$ are called the *adhesion sets* of (T, \mathcal{V}) and the *adhesion* of (T, \mathcal{V}) is the supremum of the sizes of its adhesion sets (possibly infinite). We also let $V_\infty(T) \subseteq V(T)$ denote the set of nodes $t \in V(T)$ such that V_t is infinite.

For a group Γ acting (by automorphisms) on a graph G , we say that a tree-decomposition (T, \mathcal{V}) of G is *canonical with respect to Γ* , or simply Γ -*canonical*, if Γ

induces a group action on T such that for every $\gamma \in \Gamma$ and $t \in V(T)$, $V_t \cdot \gamma = V_{t \cdot \gamma}$. By definition of a group action on a graph, $t \mapsto t \cdot \gamma$ is an automorphism of T for any $\gamma \in \Gamma$. In particular, for every $\gamma \in \Gamma$, note that γ sends bags of (T, \mathcal{V}) to bags, and adhesion sets to adhesion sets. When (T, \mathcal{V}) is $\text{Aut}(G)$ -canonical, we simply say that it is *canonical*.

Remark 3.2. If (T, \mathcal{V}) is a Γ -canonical tree-decomposition of a graph G , then Γ acts both on G and T , so there are two different notions of a stabilizer of a node $t \in V(T)$: $\Gamma_t = \text{Stab}_\Gamma(t)$ (where we consider the action of Γ on T), and $\text{Stab}_\Gamma(V_t)$ (where we consider the action of Γ on G). Observe that for any $t \in V(T)$ we have $\Gamma_t \subseteq \text{Stab}_\Gamma(V_t)$. The reverse inclusion does not hold in general (when there are adjacent nodes $s, t \in V(T)$ with $V_s = V_t$, automorphisms of T exchanging s and t stabilize $V_s = V_t$ without stabilizing s or t). However, if $t \in V(T)$ is such that $\{t' \in V(T) : V_{t'} = V_t\} = \{t\}$, then $\Gamma_t = \text{Stab}_\Gamma(V_t)$. In particular, if (T, \mathcal{V}) has finite adhesion, then every bag V_t with $t \in V_\infty(T)$ appears only once in the decomposition, and thus for each such node $t \in V_\infty(T)$ we have $\Gamma_t = \text{Stab}_\Gamma(V_t)$. The property that the two notions of stabilizers coincide for infinite bags when the canonical tree-decomposition has finite adhesion will be used repeatedly in the remainder of the paper.

Edge-separations and torsos. Consider a tree-decomposition (T, \mathcal{V}) of a graph G , with $\mathcal{V} = (V_t)_{t \in V(T)}$. Let A be an orientation of the edges of $E(T)$, i.e. a choice of either (t_1, t_2) or (t_2, t_1) for every edge $t_1 t_2$ of T . For an arbitrary pair $(t_1, t_2) \in A$, and for each $i \in \{1, 2\}$, let T_i denote the component of $T - \{t_1 t_2\}$ containing t_i . Then the *edge-separation* of G associated to (t_1, t_2) is (Y_1, S, Y_2) with $S := V_{t_1} \cap V_{t_2}$ and $Y_i := \bigcup_{s \in V(T_i)} V_s \setminus S$ for $i \in \{1, 2\}$.

Given a separation (Y, S, Z) and an automorphism γ of a graph G , let $(Y, S, Z) \cdot \gamma := (Y \cdot \gamma, S \cdot \gamma, Z \cdot \gamma)$. If $\Gamma \subseteq \text{Aut}(G)$ and \mathcal{N} is a family of separations of G , we say that \mathcal{N} is Γ -invariant if for every $(Y, S, Z) \in \mathcal{N}$ and $\gamma \in \Gamma$, we have $(Y, S, Z) \cdot \gamma \in \mathcal{N}$. Note that if (T, \mathcal{V}) is Γ -canonical, then the associated set of edge-separations is Γ -invariant.

The *torsos* of (T, \mathcal{V}) are the graphs with vertex set V_t and edge set $E(G[V_t])$ together with the edges xy such that x and y belong to a common adhesion set of (T, \mathcal{V}) . Note that this definition coincides with the general definition of torso $G[V_t]$ we gave in Subsection 2.1 when the edge-separations of (T, \mathcal{V}) are tight. To prevent any ambiguity between these definitions, we will always ensure that the tree-decompositions we work with have this property (i.e., all the associated edge-separations are tight).

Remark 3.3. If (T, \mathcal{V}) is a Γ -canonical tree-decomposition of a locally finite graph G whose edge-separations are tight, then by Lemma 3.1 the action of Γ on $E(T)$ must induce a finite number of orbits. In particular, Γ must also act quasi-transitively on $V(T)$.

The *treewidth* of a graph G is the infimum of the width of (T, \mathcal{V}) , among all tree-decompositions (T, \mathcal{V}) of G . Note that adding to a tree-decomposition of bounded width the restriction that it must be canonical can be very costly in the finite case: while it is well known that every cycle graph C_n on n vertices has treewidth 2,

the example below shows that in any canonical tree-decomposition of C_n , some bag contains all the nodes of C_n .

Example 3.4. Let C_n be the cycle graph on n elements. Note that the additive group \mathbb{Z}_n acts transitively by rotation on C_n . We let a be a generator of \mathbb{Z}_n of order n . Let $(T, (V_t)_{t \in V(T)})$ be a \mathbb{Z}_n -canonical tree-decomposition of C_n . Without loss of generality we may assume that T is finite, by contracting every edge $tt' \in E(T)$ such that $V_t = V_{t'}$. We may also assume that no edge tt' of T is *inverted* by a , i.e. such that $(t, t') \cdot a = (t', t)$, as if it was the case we could subdivide the edge tt' (i.e. add a new vertex t^* between t and t') and let $V_{t^*} := V_t \cap V_{t'}$. If we let T' be the tree of the tree-decomposition obtained after performing such a subdivision, note that the obtained tree-decomposition is still \mathbb{Z}_n -canonical as a induces an automorphism of T' that stabilizes the vertex t^* and acts on $V(T)$ the same way that it did before the subdivision. After this operation none of the edges tt^* , t^*t' is inverted by a . It is an easy exercise to prove that if no edge of T is inverted by a , there exists a vertex $t \in V(T)$ stabilized by a , and hence by all the elements of \mathbb{Z}_n . Then as \mathbb{Z}_n acts transitively on G , we must have $V_t = V(C_n)$ for such a $t \in V(T)$.

It turns out that in the quasi-transitive locally-finite case, if a graph has bounded treewidth then adding the restriction that the tree-decomposition must be canonical is fairly inexpensive if we only care about having finite bags.

Theorem 3.5 (Theorem 7.5 in [HLMR22], [MS83], [Woe89], [TW93]). *Let G be a connected quasi-transitive locally finite graph. Then the following are equivalent:*

- G has finite treewidth;
- all the ends of G are thin;
- there exists $k \geq 1$ such that every end of G has degree at most k ;
- there exists a canonical tree-decomposition of G with tight edge-separations and finite width.

Note that the final item above is not stated explicitly in Theorem 7.5 in [HLMR22], but it can be easily deduced from this result (see also Theorem 2.4 in [Ham25], which uses the result of [HLMR22] together with our Lemmas 3.1 and 3.13).

Separations of order at most 3. If G is not connected, then the tree-decomposition (T, \mathcal{V}) where T is a star whose central bag is empty and where we put a bag for each connected component of G can easily be seen to be a canonical tree-decomposition with adhesion 0, as every automorphism of G acts on T by permuting some branches. If we start from a connected graph G , it is well-known that the block cut-tree of G is a canonical tree-decomposition $(T, (V_t)_{t \in V(T)})$ of G whose adhesion sets have size 1 and such that for each $t \in V(T)$, $G[V_t] = G[V_t]$ has either size at most 2 or is 2-connected. A similar result holds for separations of order 2 (this was proved by Tutte [Tut84] in the finite case, and generalized to infinite graphs in [DSS98]).

Theorem 3.6 ([DSS98]). *Every locally finite graph G has a canonical tree-decomposition of adhesion at most 2, whose torsos are minors of G and are complete graphs of order at most 2, cycles, or 3-connected graphs.*

For separations of order 3, a similar result was obtained by Grohe for finite graphs [Gro16a].

Theorem 3.7 ([Gro16a]). *Every finite graph G has a tree-decomposition of adhesion at most 3 whose torsos are minors of G and are complete graphs on at most 4 vertices or quasi-4-connected graphs.*

Our main technical contribution will be to extend Theorem 3.7 to locally finite graphs, while making sure that most of the construction (except the very end) is canonical. More precisely, we reproduce in Subsections 3.5 and 3.6 the main steps of the work of [Gro16a] and give the additional arguments to extend them to locally finite graphs. A consequence is that Theorem 3.7 extends to locally finite graphs. However in our case, the main difficulty we face is that the decomposition of Grohe is not canonical (although some parts of the construction are canonical, which will be crucial for our purposes). To give a rough idea of the bulk of the problem it is helpful to consider Example 3.8 below, which was introduced in [Gro16a] in the finite case.




FIGURE 1. Left: a finite section of the 3-connected infinite graph obtained by replacing in the infinite hexagonal planar grid each vertex by a triangle with three vertices of degree 3. Right: a finite section of the quasi-4-connected torso $G[V_{z_0}]$ of (T, \mathcal{V}) . Note that it does not depend of the choice of V_{z_0} .

Example 3.8. Consider the 3-connected infinite planar graph H obtained from the infinite hexagonal planar grid by replacing each vertex by a triangle with three vertices of degree 3 (see Figure 1 (left) for a finite part of this graph). We let M be the set of edges connecting pairs of triangles, or equivalently the set of edges that do not belong to any triangle (note that M is a perfect matching). The tree-decomposition (T, \mathcal{V}) of H obtained by extending the ideas in [Gro16a] to the infinite case has an infinite bag V_{z_0} obtained by selecting one endpoint of each edge of M (which is equivalent to fixing an orientation of each of these edges). The tree T is a subdivision of a star with center z_0 , and its other bags are finite. While there are many different choices for V_{z_0} , none of them gives a canonical tree-decomposition. Indeed one can check more generally that no tree-decomposition of H satisfying the properties of Theorem 3.7 can be canonical. To see this, assume for the sake of contradiction that such a decomposition (T, \mathcal{V}) exists. Then one of its edge-separations should be proper of order 3. Note that the only such separations separate a subgraph of a triangle from the rest of the graph. Let (Y, S, Z) be such a separation, such that

Z is finite. Then there exists an edge e from M with one endpoint in Z and the other in S . Note that there exists an automorphism $\gamma \in \text{Aut}(H)$ exchanging the two endpoints of e . In particular, as (T, \mathcal{V}) is canonical, both $(Y, S, Z) \cdot \gamma$ and (Y, S, Z) must be edge-separations of (T, \mathcal{V}) , which can be seen to be impossible.

This example illustrates the fact that in general it is impossible to obtain a *canonical* tree-decomposition having exactly the properties described in Theorem 3.7. However note that here, if we want a canonical tree-decomposition whose torsos are either planar or finite, it is sufficient to take the trivial tree-decomposition with a single bag containing all the vertices. This is exactly what our proof will do when applied to this graph. More precisely, we note that on this example, the set M of edges is invariant under the action of $\text{Aut}(H)$. Based on this observation, our solution to obtain a canonical tree-decomposition will be to start with the same decomposition as that of [Gro16a], but to keep the two endpoints of each edge of M instead of choosing only one of its endpoints as above.

Combining canonical tree-decompositions. Let (T, \mathcal{V}) and (T', \mathcal{V}') be tree-decompositions of two graphs G, G' , respectively. We say that (T, \mathcal{V}) and (T', \mathcal{V}') are *isomorphic* if there exists an isomorphism φ from G to G' , and an isomorphism ψ from T to T' such that for each $t \in V(T)$, we have: $V'_{\psi(t)} = \varphi(V_t)$.

Let G be a graph and let Γ be a group acting on G . Let (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, be a Γ -canonical tree-decomposition of G , and recall that for any $t \in V(T)$, $\Gamma_t = \text{Stab}_\Gamma(t)$ denotes the stabilizer of the node t in the action of Γ on the tree T . For each $t \in V(T)$, let (T_t, \mathcal{V}_t) be a Γ_t -canonical tree-decomposition of $G[V_t]$. Our goal will be to refine (T, \mathcal{V}) by combining it with the tree-decompositions $(T_t, \mathcal{V}_t)_{t \in V(T)}$. If we want the resulting refined tree-decomposition of G to be Γ -canonical, we need to impose a condition on the tree-decompositions $(T_t, \mathcal{V}_t)_{t \in V(T)}$ (namely that they are consistent with the action of Γ on G). This is captured by the following definition. For $g \in \Gamma$ and $t \in V(T)$, we define $(T_t, \mathcal{V}_t) \cdot g := (T_t, \mathcal{V}_t \cdot g)$ as the tree-decomposition of $G[V_t \cdot g] = G[V_{t \cdot g}]$ with underlying tree T_t and bags $\mathcal{V}_t \cdot g := (V_s \cdot g)_{s \in V(T_t)}$. Observe that $(T_t, \mathcal{V}_t) \cdot g$ is $\Gamma_{t \cdot g}$ -canonical. We say that the construction $t \mapsto (T_t, \mathcal{V}_t)$ is Γ -*canonical* if for each $g \in \Gamma$ and $t \in V(T)$, the tree-decompositions $(T_t, \mathcal{V}_t) \cdot g$ and $(T_{t \cdot g}, \mathcal{V}_{t \cdot g})$ are isomorphic. We emphasize here that the first tree-decomposition is indexed by T_t , while the second is indexed by $T_{t \cdot g}$.

The *trivial tree-decomposition* of a graph G consists of a tree T with a single node, whose bag is $V(G)$. Note that the trivial tree-decomposition is canonical.

Lemma 3.9. *Assume that G is locally finite, Γ is a group acting on G , and (T, \mathcal{V}) is a Γ -canonical tree-decomposition of G with finitely bounded adhesion. Let $\{t_i : i \in I_\infty\}$ be a set of representatives of the Γ -orbits of $V_\infty(T)$, indexed by some set I_∞ . Assume that for every $i \in I_\infty$ there exists a Γ_{t_i} -canonical tree-decomposition $(T_{t_i}, \mathcal{V}_{t_i})$ of the torso $G[V_{t_i}]$ of finitely bounded adhesion. Then we can find some family $(T_t, \mathcal{V}_t)_{t \in V(T)}$ extending the family $(T_{t_i}, \mathcal{V}_{t_i})_{i \in I_\infty}$ such that each (T_t, \mathcal{V}_t) is a Γ_t -canonical tree-decomposition of $G[V_t]$, the construction $t \mapsto (T_t, \mathcal{V}_t)$ is Γ -canonical, and for each $t \in V(T) \setminus V_\infty(T)$, (T_t, \mathcal{V}_t) is the trivial tree-decomposition of $G[V_t]$.*

Proof. First we check that for each $g \in \Gamma$ and every $t \in V_\infty(T)$, $g^{-1} \cdot \Gamma_t \cdot g = \Gamma_{t \cdot g}$. For this we claim that we only need to prove the inclusion $g^{-1} \cdot \Gamma_t \cdot g \subseteq \Gamma_{t \cdot g}$, as the converse then follows from replacing (g, t) by $(g^{-1}, t \cdot g)$. Let $y \in V_{t \cdot g} = V_t \cdot g$ (where the equality follows from the assumption that (T, \mathcal{V}) is Γ -canonical) and let $h \in \Gamma_t$.

Then $y = x \cdot g$ for some $x \in V_t$ and we have:

$$y \cdot (g^{-1} \cdot h \cdot g) = x \cdot g \cdot (g^{-1} \cdot h \cdot g) = (x \cdot h) \cdot g.$$

Since $h \in \Gamma_t$, we have $x \cdot h \in V_t$, we thus get that $y \cdot (g^{-1} \cdot h \cdot g) \in V_t \cdot g = V_{t \cdot g}$, so we just proved that every element of $g^{-1} \cdot \Gamma_t \cdot g$ stabilizes $V_{t \cdot g}$. Since $t \in V_\infty(T)$, V_t and $V_{t \cdot g}$ are infinite and it follows from Remark 3.2 that $\text{Stab}_\Gamma(V_{t \cdot g}) = \Gamma_{t \cdot g}$. This implies that every element of $g^{-1} \cdot \Gamma_t \cdot g$ lies in $\Gamma_{t \cdot g}$, and thus $g^{-1} \cdot \Gamma_t \cdot g \subseteq \Gamma_{t \cdot g}$, as desired.

We complete I_∞ into a set I of representatives of the Γ -orbits of $V(T)$, and for each $i \in I \setminus I_\infty$ we let $(T_{t_i}, \mathcal{V}_{t_i})$ denote the trivial tree-decomposition of $G[V_{t_i}]$. For each $t \in V(T)$, we let $g \in \Gamma$ and $i \in I$ be such that $t = t_i \cdot g$ and let $(T_t, \mathcal{V}_t) := (T_{t_i}, \mathcal{V}_{t_i}) \cdot g$. We check that (T_t, \mathcal{V}_t) is well-defined: for any two $g, g' \in \Gamma$ such that $t_i \cdot g = t_i \cdot g' = t$, we have $g' \cdot g^{-1} \in \Gamma_{t_i}$. As $(T_{t_i}, \mathcal{V}_{t_i})$ is Γ_{t_i} -canonical, $(T_{t_i}, \mathcal{V}_{t_i}) \cdot g' \cdot g^{-1} = (T_{t_i}, \mathcal{V}_{t_i})$ so we have $(T_{t_i}, \mathcal{V}_{t_i}) \cdot g = (T_{t_i}, \mathcal{V}_{t_i}) \cdot g'$ and (T_t, \mathcal{V}_t) is well-defined for each $t \in V(T)$. The fact that the construction $t \mapsto (T_t, \mathcal{V}_t)$ is Γ -canonical immediately follows from the definition. Finally, the tree-decomposition (T_t, \mathcal{V}_t) is Γ_t -canonical because if it is not trivial, and $i \in I_\infty$ and $g \in \Gamma$ are such that $t_i \cdot g = t$, then $\Gamma_t = g^{-1} \cdot \Gamma_{t_i} \cdot g$ and thus Γ_t induces a group action on $(T_t, \mathcal{V}_t) = (T_{t_i}, \mathcal{V}_{t_i}) \cdot g$. \square

Given two tree-decompositions $(T, \mathcal{V}), (T', \mathcal{V}')$ of a graph G , with $\mathcal{V} = (V_t)_{t \in V(T)}$ and $\mathcal{V}' = (V'_t)_{t \in V(T')}$, we say that (T', \mathcal{V}') *refines* (T, \mathcal{V}) with respect to some family $(T_t, \mathcal{V}_t)_{t \in V(T)}$ of tree-decompositions if for every $t \in V(T)$, T_t is a subtree of T' such that $V_t = \bigcup_{s \in V(T_t)} V'_s$ and the trees $(T_t)_{t \in V(T)}$ are pairwise vertex-disjoint, cover $V(T')$ and for every edge $uv \in E(T)$, there exist $u' \in V(T_u), v' \in V(T_v)$ such that $u'v' \in E(T')$.

We say that (T', \mathcal{V}') is a *subdivision* of (T, \mathcal{V}) if T' is obtained from T after considering a subset $E' \subseteq E(T)$ and doing the following for every edge $tt' \in E'$: we subdivide the edge tt' (by adding a new vertex t^* between t and t'), and we add a corresponding bag $V_{t^*} := V_t \cap V_{t'}$ in the tree-decomposition. Note that if (T', \mathcal{V}') is a subdivision of (T, \mathcal{V}) , the two tree-decompositions have the same edge-separations.

The following result from [CHM22] will allow us to construct canonical tree-decompositions inductively:

Proposition 3.10 (Proposition 7.2 in [CHM22]). *Assume that G is locally finite, Γ is a group acting on G and (T, \mathcal{V}) is a Γ -canonical tree-decomposition of G with finitely bounded adhesion, with $\mathcal{V} = (V_t)_{t \in V(T)}$. Assume that for every $t \in V(T)$, there exists a Γ_t -canonical tree-decomposition (T_t, \mathcal{V}_t) of the torso $G[V_t]$ of finitely bounded adhesion such that the edge-separations induced by (T_t, \mathcal{V}_t) in $G[V_t]$ are tight and pairwise distinct, and the construction $t \mapsto (T_t, \mathcal{V}_t)$ is Γ -canonical. Then there exists a Γ -canonical tree-decomposition (T', \mathcal{V}') of G that refines (T, \mathcal{V}) with respect to a family $(T'_t, \mathcal{V}'_t)_{t \in V(T)}$ such that for each $t \in V(T)$, (T'_t, \mathcal{V}'_t) is a Γ_t -canonical tree-decomposition of $G[V_t]$ which is a subdivision of (T_t, \mathcal{V}_t) , and such that every adhesion set of (T', \mathcal{V}') is either an adhesion set of (T, \mathcal{V}) or an adhesion set of some (T_t, \mathcal{V}_t) for some $t \in V(T)$. Moreover, the construction $t \mapsto (T'_t, \mathcal{V}'_t)$ is Γ -canonical.*

Remark 3.11. In the original statement of [CHM22, Proposition 7.2], the fact that each tree-decomposition (T'_t, \mathcal{V}'_t) is Γ_t -canonical and that the construction $t \mapsto (T'_t, \mathcal{V}'_t)$ is also Γ -canonical is not stated explicitly, however the authors show it explicitly in the proof.

Hence putting Lemma 3.9 together with Proposition 3.10, we immediately get:

Corollary 3.12. *Assume that G is locally finite, Γ is a group acting on G , and (T, \mathcal{V}) a Γ -canonical tree-decomposition of G of finitely bounded adhesion, with $\mathcal{V} = (V_t)_{t \in V(T)}$. Let $\{t_i : i \in I_\infty\}$ denote a set of representatives of the orbits $V_\infty(T)/\Gamma$ such that for each $i \in I_\infty$, there exists a Γ_{t_i} -canonical tree-decomposition $(T_{t_i}, \mathcal{V}_{t_i})$ of $G[V_{t_i}]$ with finitely bounded adhesion, such that the edge-separations induced by each $(T_{t_i}, \mathcal{V}_{t_i})$ in $G[V_{t_i}]$ are tight and pairwise distinct. Then there exists a Γ -canonical tree-decomposition of G that refines (T, \mathcal{V}) with respect to some family $(T'_t, \mathcal{V}'_t)_{t \in V(T)}$ of Γ_t -canonical tree-decompositions of $G[V_t]$ such that for each $i \in I_\infty$, $(T'_{t_i}, \mathcal{V}'_{t_i})$ is a subdivision of $(T_{t_i}, \mathcal{V}_{t_i})$, and for every $t \in V(T) \setminus V_\infty(T)$, (T'_t, \mathcal{V}'_t) is the trivial tree-decomposition of $G[V_t]$. Moreover, the construction $t \mapsto (T'_t, \mathcal{V}'_t)$ is Γ -canonical.*

The main objects of study of this paper are canonical tree-decompositions of quasi-transitive graphs. A crucial property is that the torsos or parts of the tree-decomposition are themselves quasi-transitive. This is proved in [HLMR22, Proposition 4.5] in the special case where Γ acts transitively on $E(T)$. We give here a more general proof, which is self-contained.

Lemma 3.13. *Let $k \in \mathbb{N}$, let G be a locally finite graph, and let Γ be a group acting quasi-transitively on G . Let (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, be a Γ -canonical tree-decomposition of G whose edge-separations are tight and have order at most k . Then, for any $t \in V(T)$, the group $\Gamma_t := \text{Stab}_\Gamma(t)$ induces a quasi-transitive action on $G[V_t]$, and thus also on $G[V_t]$.*

Proof. Let $G, \Gamma, (T, \mathcal{V})$ be as described above. By Lemma 3.1, as the edge-separations are tight and of bounded size, there are only finitely many Γ -orbits of $E(T)$. We fix an orientation A of $E(T)$. Let $e_1, \dots, e_m \in A$ be representatives of each of the Γ -orbits of A , and let $\bar{e}_1, \dots, \bar{e}_m$ denote their inverse pairs. Fix any node $t \in V(T)$.

We consider an arbitrary vertex $z \in V(G)$, and let $t_0 \in V(T)$ be such that $z \in V_{t_0}$. Let $\Omega_z := z \cdot \Gamma$ denote the Γ -orbit of z . We define the following subset of $V_t \cap \Omega_z$.

$$\Theta_z := \{y \in V_t \cap \Omega_z : y = z \cdot \gamma \text{ for some } \gamma \text{ such that } t_0 \cdot \gamma = t\}.$$

We first show that Γ_t acts transitively on Θ_z . Let $y, y' \in \Theta_z$ and $\gamma, \gamma' \in \Gamma$ be such that $y = z \cdot \gamma$, $y' = z \cdot \gamma'$ and $t_0 \cdot \gamma = t_0 \cdot \gamma' = t$. Then if we set $\alpha := \gamma'^{-1} \cdot \gamma$, we have

$$t \cdot \alpha = t \cdot \gamma'^{-1} \cdot \gamma = t_0 \cdot \gamma = t,$$

and thus $\alpha \in \Gamma_t$. As $y' \cdot \alpha = y$, this shows that Γ_t acts transitively on Θ_z .

For $i \in [m]$, we define:

$$\Psi_i := \{y \in V_t \cap V_{t'} : \text{there exists } \gamma \in \Gamma \text{ such that } (t, t') = e_i \cdot \gamma\}$$

$$\Psi_{i+m} := \{y \in V_t \cap V_{t'} : \text{there exists } \gamma \in \Gamma \text{ such that } (t, t') = \bar{e}_i \cdot \gamma\}.$$

We observe that if a vertex of $V_t \cap \Omega_z$ does not lie in Θ_z , it has to lie in one of sets Ψ_i for $i \in [2m]$. To see this, let $y \in V_t \cap \Omega_z$, and $\gamma \in \Gamma$ be such that $y = z \cdot \gamma$. If $y \notin \Theta_z$, then $t_0 \cdot \gamma \neq t$. In this case the unique path in T from t to $t_0 \cdot \gamma$ contains at least one edge. Let t' be the neighbor of t on this path. As $V_{t_0 \cdot \gamma} = V_{t_0} \cdot \gamma$ and $z \in V_{t_0}$, we have $y \in V_t \cap V_{t_0 \cdot \gamma}$. Hence as (T, \mathcal{V}) is a tree-decomposition, $y \in V_t \cap V_{t'}$. Thus if we let i be such that $(t, t') = e_i \cdot \beta$ or $(t, t') = \bar{e}_i \cdot \beta$ for some $\beta \in \Gamma$, we obtain that $y \in \Psi_i \cup \Psi_{i+m}$. This shows that V_t is covered by the union of the sets $\Theta_z, z \in V(G)$ (there are at most $|V(G)/\Gamma|$ such sets), and the sets $\Psi_i, i \in [2m]$.

We now show that Γ_t acts quasi-transitively on each Ψ_i , $i \in [2m]$. Let $i \in [m]$, $y_1, y_2 \in \Psi_i$, $t_1, t_2 \in V(T)$ and $\beta_1, \beta_2 \in \Gamma$ such that $(t, t_1), (t, t_2) \in E(T)$, $(t, t_1) = e_i \cdot \beta_1$ and $(t, t_2) = e_i \cdot \beta_2$. We set $\alpha := \beta_1^{-1} \cdot \beta_2$ and note that α sends the directed edge $e_i \cdot \beta_1$ to $e_i \cdot \beta_2$. Let S_i be the separator of G associated to the edge-separation induced by the edge e_i in (T, \mathcal{V}) . The previous remark implies that α sends $S_i \cdot \beta_1$ to $S_i \cdot \beta_2$ and that $\alpha \in \Gamma_t$. As for every $i \in [m]$, S_i has size at most k , we just proved that the action of Γ_t on Ψ_i induces at most k orbits. The case $i \in \{m+1, \dots, 2m\}$ is exactly the same.

As V_t is covered by the union of the sets Θ_z , $z \in V(G)$ (there are at most $|V(G)/\Gamma|$ such sets, and Γ_t acts transitively on each of these sets), and the sets Ψ_i , $i \in [2m]$ (and the action of Γ_t on each of these sets induces at most k orbits), we have $|V_t/\Gamma_t| \leq 2km + |V(G)/\Gamma|$, which implies that Γ_t acts quasi-transitively on $G[V_t]$. As (T, \mathcal{V}) is Γ -canonical, for each $g \in \Gamma$ and each edge e lying inside some adhesion set of the tree-decomposition, g sends e to a pair of vertices in another adhesion set of the tree-decomposition (and this pair of vertices must thus be joined by an edge in the corresponding torso). It follows that any automorphism $g \in \Gamma_t$ of $G[V_t]$ is also an automorphism of the torso $G[V_t]$. Hence, Γ_t also acts quasi-transitively on $G[V_t]$. \square

Note that Lemma 3.13 still holds if we only require $E(T)/\Gamma$ to be finite (instead of requiring the edge-separations of (T, \mathcal{V}) to be tight).

3.2. Tangles. Tangles were introduced by Robertson and Seymour [RS91] and play a fundamental role in their proof of the Graph Minor Structure Theorem. We will consider here the equivalent definition used by Grohe [Gro16a]. A *tangle of order k* in G is a subset \mathcal{T} of $\text{Sep}_{<k}(G)$ such that

- (T1) For all separations $(Y, S, Z) \in \text{Sep}_{<k}(G)$, either $(Y, S, Z) \in \mathcal{T}$ or $(Z, S, Y) \in \mathcal{T}$;
- (T2) For all separations $(Y_1, S_1, Z_1), (Y_2, S_2, Z_2), (Y_3, S_3, Z_3) \in \mathcal{T}$, either $Z_1 \cap Z_2 \cap Z_3 \neq \emptyset$ or there exists an edge with an endpoint in each Z_i .

Note that (T2) with $(Y_i, S_i, Z_i) = (Y, S, Z)$ for each $i \in [3]$ implies in particular that for every separation $(Y, S, Z) \in \mathcal{T}$, $Z \neq \emptyset$. A tangle in G will be called a G -tangle, for brevity. Intuitively, a G -tangle is a consistent orientation of the separations of G , pointing towards a highly connected region of G . We refer the reader to [RS91, Gro16a] for background on tangles. In general when G is finite there is a one-to-one correspondence between the G -tangles of order 1 and the connected components of G , between the G -tangles of order 2 and the *biconnected* components of G , and between the G -tangles of order 3 and the *triconnected* components of G (we omit the definitions as these notions will not be needed in the remainder of the paper, and instead refer the interested reader to [RS91] and [Gro16b]).

In infinite graphs, tangles can also be seen as a notion generalizing the notion of ends: for each end ω of a graph G and every $k \geq 2$, we define the G -tangle \mathcal{T}_ω^k of order k induced by ω by:

$$\mathcal{T}_\omega^k := \{(Y, S, Z) \in \text{Sep}_{<k}(G) : \omega \text{ lives in a component of } Z\}.$$

The fact that \mathcal{T}_ω^k is indeed a tangle is a folklore result. One of the basic properties of tangles is that for any fixed model of a graph H in a graph G , any H -tangle of order k induces a G -tangle of order k . More precisely, if $\mathcal{M} = (M_v)_{v \in V(H)}$ is a model of H in G and (Y, S, Z) is a separation of order less than k in G , then its *projection*

with respect to \mathcal{M} is the separation $\pi_{\mathcal{M}}(Y, S, Z) = (Y', S', Z')$ of H of order less than k defined by: $Y' := \{v \in V(H) : M_v \subseteq Y\}$, $S' := \{v \in V(H) : M_v \cap S \neq \emptyset\}$ and $Z' := \{v \in V(H) : M_v \subseteq Z\}$.

A proof of the following result can be found in [RS91, (6.1)] or in a more similar version in [Gro16a, Lemma 3.11]. Its proof extends to the locally finite case.

Lemma 3.14 (Lemma 3.11 in [Gro16a]). *Let G be a locally finite graph. Let $\mathcal{M} = (M_v)_{v \in V(H)}$ be a model of a graph H in G and \mathcal{T}' be an H -tangle of order $k \geq 2$. Then the set*

$$\mathcal{T} := \{(Y, S, Z) \in \text{Sep}_{<k}(G) : \pi_{\mathcal{M}}(Y, S, Z) \in \mathcal{T}'\}$$

is a G -tangle of order k , called the lifting of \mathcal{T}' in G with respect to \mathcal{M} .

Remark 3.15. Assume that \mathcal{M} is a faithful model of H in G with the property that for each $(Y', S', Z') \in \text{Sep}_{<k}(H)$, there exists some $(Y, S, Z) \in \text{Sep}_{<k}(G)$ such that $\pi_{\mathcal{M}}(Y, S, Z) = (Y', S', Z')$ and $S' = S$. Then the function that maps every tangle of order k in H to its lifting in G with respect to \mathcal{M} is injective. To see this, consider two distinct tangles $\mathcal{T}'_1 \neq \mathcal{T}'_2$ of order k in H . Then there exists some $(Y', S', Z') \in \mathcal{T}'_1$ such that $(Z', S', Y') \in \mathcal{T}'_2$. If we consider $(Y, S, Z) \in \text{Sep}_{<k}(G)$ such that $\pi_{\mathcal{M}}(Y, S, Z) = (Y', S', Z')$, we have $(Y, S, Z) \in \mathcal{T}_1$ and $(Z, S, Y) \in \mathcal{T}_2$, where for each $i \in \{1, 2\}$, \mathcal{T}_i denotes the lifting of \mathcal{T}'_i with respect to \mathcal{M} . It then follows that $\mathcal{T}_1 \neq \mathcal{T}_2$, as desired. Note that if (T, \mathcal{V}) is a tree-decomposition with finitely bounded adhesion and $t \in V(T)$ is such that $G[[V_t]]$ is a faithful minor of G , then any faithful model \mathcal{M} of $G[[V_t]]$ has the property we just described.

If $\mathcal{M} = (M_v)_{v \in V(H)}$ is a model of H in G , and \mathcal{T} is a tangle of G , then $\mathcal{T}' := \{\pi_{\mathcal{M}}(Y, S, Z) : (Y, S, Z) \in \mathcal{T}\}$ is called the *projection* of \mathcal{T} . Note that \mathcal{T}' is not a tangle in general. Projecting is the converse operation of lifting in the sense that if \mathcal{M} is faithful, and \mathcal{T} and \mathcal{T}' are tangles of G and H , then \mathcal{T} is the lifting of \mathcal{T}' if and only if \mathcal{T}' is the projection of \mathcal{T} .

We define a partial order \prec over the set of separations of a graph G by letting for every two separations $(Y, S, Z), (Y', S', Z')$, $(Y, S, Z) \preceq (Y', S', Z')$ if and only if $S \cup Z \subsetneq S' \cup Z'$ or $(S \cup Z = S' \cup Z'$ and $S \subsetneq S')$. Intuitively, $(Y, S, Z) \preceq (Y', S', Z')$ means that (Y, S, Z) points towards a direction in a more accurate way than (Y', S', Z') does. Note that our definition of \preceq is the same as in [Gro16a, Subsection 3.2] and slightly differs from the more conventional one of [RS91, CHM22].

A partially ordered set $(X, <)$ is said to be *well-founded* if every strictly decreasing sequence of elements of X is finite. In particular, if $(X, <)$ is well-founded then for every $x \in X$, there exists $y \in X$ which is minimal with respect to $<$ and such that $y \leq x$. In the remainder of the paper, whenever we consider a minimal separation or a well-founded family of separations, we always implicitly refer to the partial order \prec defined in the paragraph above.

We will distinguish two types of tangles in infinite graphs:

- the *region tangles*, defined as those which are well-founded (with respect to the order \prec), and
- the *evasive tangles*, which contain some infinite decreasing sequence of separations (with respect to the order \prec).

The tangles we consider in this work will always have order at most 4. Note that if G is 3-connected, an evasive tangle \mathcal{T} of order 4 is exactly a tangle \mathcal{T}_ω^4 induced

by an end ω of degree 3. On the other hand, a region tangle is either a tangle of order 4 induced by some end ω of degree at least 4, or a tangle which is not induced by an end. For example, one can check that both graphs in Figure 1 have a unique tangle of order 4 which is the tangle induced by their unique end (which is thick), and this tangle is a region tangle in both cases.

We say that a separation (Y, S, Z) *distinguishes* two tangles $\mathcal{T}, \mathcal{T}'$ if $(Y, S, Z) \in \mathcal{T}$ and $(Z, S, Y) \in \mathcal{T}'$, or vice versa. We say that (Y, S, Z) distinguishes \mathcal{T} and \mathcal{T}' *efficiently* if there is no separation of smaller order distinguishing \mathcal{T} and \mathcal{T}' . A tree-decomposition (T, \mathcal{V}) *distinguishes* a set of tangles \mathcal{A} if for every two distinct tangles $\mathcal{T}, \mathcal{T}' \in \mathcal{A}$ there exists an edge-separation of (T, \mathcal{V}) distinguishing \mathcal{T} and \mathcal{T}' . A separation is called *relevant* with respect to \mathcal{A} if it distinguishes at least two tangles of \mathcal{A} . A tree-decomposition is *nice* (with respect to \mathcal{A}) if all its edge-separations are relevant (with respect to \mathcal{A}).

We will need the following result, which extends earlier results of [RS91, DHL18], and which is a canonical version of one of the main results of the grid-minor series in the locally finite case. We will only use it with $k = 4$, but we nevertheless state the result in its most general form.

Theorem 3.16 (Theorem 7.3 in [CHM22]). *Let $k \geq 1$ and let G be a locally finite graph. Then there exists a canonical tree-decomposition (T, \mathcal{V}) of G that efficiently distinguishes the set \mathcal{A}_k of tangles of order at most k and that is nice with respect to \mathcal{A}_k .*

Remark 3.17. The fact that (T, \mathcal{V}) is nice in Theorem 3.16 is not explicit in the original statement, however it directly follows from the proof. Moreover, the proof also ensures that the edge-separations of (T, \mathcal{V}) are pairwise distinct.

3.3. An example. We give here an example of a one-ended graph that excludes some minor and has infinitely many region tangles of order 4. We show how to distinguish them on this example with a canonical tree-decomposition. As the application of Theorem 3.16 allowing to distinguish all tangles of order 4 is the very first step of our proof of Theorems 4.1 and 4.3, this example may also be useful to have some intuition on it.

We consider the infinite graph G (a finite section of which is illustrated in Figure 2), which is obtained from the infinite triangular grid by adding in each triangular face $f = v_1v_2v_3$ three vertices $\{w_1, w_2, w_3\}$ inducing a $K_{3,3}$ with the vertices of the triangle, and another vertex z connected to each of the w_i 's. G has two types of tangles of order 4: one is the tangle \mathcal{T}_ω^4 induced by the unique end ω of G , and all the others are the tangles \mathcal{T}_f^4 pointing towards each face $f = v_1v_2v_3$ of the triangular grid; more precisely, \mathcal{T}_f^4 has the same set of separations as \mathcal{T}_ω^4 except for $(G - A, \{v_1, v_2, v_3\}, \{w_1, w_2, w_3, z\}) \in \mathcal{T}_f^4$, where $A := \{v_1, v_2, v_3, w_1, w_2, w_3, z\}$. Note that with respect to our definition, all the tangles of order 4 of G are region tangles. We represented with red arrows the two separations of \mathcal{T}_f^4 that are minimal with respect to the order \prec but which are not minimal separations of \mathcal{T}_ω^k (for one fixed face f). The three red arrows crossing the red triangle correspond to the minimal separation $(G - A, \{v_1, v_2, v_3\}, \{w_1, w_2, w_3, z\})$ of \mathcal{T}_f^4 that points towards the triangular face $v_1v_2v_3$, while the three arrows directed away from z correspond to the minimal separation $(\{z\}, \{w_1, w_2, w_3\}, \{v_1, v_2, v_3\} \cup (G - A))$ of \mathcal{T}_f^4 . The tree-decomposition (T, \mathcal{V}) of Theorem 3.16 distinguishing all the tangles of order 4 is such that T is a




FIGURE 2. A useful example.

star with center $t_0 \in V(T)$ such that $G[V_{t_0}]$ is the infinite planar triangular grid. Then T has one vertex t_f for each face $f = v_1v_2v_3$ of $G[V_{t_0}]$ and the bag V_{t_f} is finite and contains the 7 vertices $\{v_1, v_2, v_3, w_1, w_2, w_3, z\}$ associated to f . Note that such a tree-decomposition enjoys the properties of Theorems 4.1 and 4.3. However, this is not always the case and we need in general to decompose further some torsos of the tree-decomposition given by Theorem 3.16 in order to obtain such a decomposition.

3.4. Tangles of order 4: orthogonality and crossing-lemma. In this section we introduce some notions from [Gro16a] and briefly explain how to extend them in the locally finite case. Unless specified otherwise, we assume in the whole section that the graphs we consider are locally finite and 3-connected.

A separation $(Y, S, Z) \in \text{Sep}_{<4}(G)$ is said to be *degenerate* if

- (Y, S, Z) has order 3,
- $G[S]$ is an independent set, and
- $|Y| = 1$.

The following result from [Gro16a] immediately generalizes to locally finite graphs:

Lemma 3.18 (Lemma 4.13 and Remark 4.14 in [Gro16a]). *Let G be a locally finite 3-connected graph, and (Y, S, Z) be a proper separation of order 3. Then $G[Z \cup S]$ is a faithful minor of G if and only if (Y, S, Z) is non-degenerate.*

We say that the edge-separations of a tree-decomposition (T, \mathcal{V}) are *non-degenerate* if for every $e \in E(T)$, none of the two edge-separations associated to e are degenerate.

Lemma 3.19. *Let G be a locally finite 3-connected graph and let (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, be a tree-decomposition of G whose edge-separations have order 3 and are non-degenerate. Then $G[V_t]$ is a faithful minor of G for each $t \in V(T)$.*

Proof. Let $t \in V(T)$, and t' be a neighbor of t in T . Let $(Y_{t'}, S_{t'}, Z_{t'})$ be the edge-separation of G associated to the (oriented) edge $(t', t) \in E(T)$, that is $S_{t'} = V_t \cap V_{t'}$, $V_{t'} \subseteq Y_{t'} \cup S_{t'}$, and $V_t \subseteq Z_{t'} \cup S_{t'}$. By Lemma 3.18, there is a faithful model $(M_v^{t'})_{v \in (Z_{t'} \cup S_{t'})}$ of $G[[Z_{t'} \cup S_{t'}]]$ in G . As the only edges of $G[[Z_{t'} \cup S_{t'}]]$ that are not edges of G must be between pairs of vertices of $S_{t'}$, we may assume that every $M_v^{t'}$ has size 1, except possibly when $v \in S_{t'}$, in which case the only vertices distinct from v that $M_v^{t'}$ can have must lie in $Y_{t'}$. For every $v \in V_t$, we let:

$$M_v := \bigcup_{\substack{t' \in V(T), \\ tt' \in E(T)}} M_v^{t'}.$$

We show that $(M_v)_{v \in V_t}$ is a faithful model of $G[[V_t]]$ in G . As (T, \mathcal{V}) is a tree-decomposition, for every two distinct neighbors t', t'' of t in T , $Y_{t'} \cap Y_{t''} = \emptyset$ so we must have $M_v^{t'} \cap M_v^{t''} = \{v\}$ and $M_u^{t'} \cap M_u^{t''} = \emptyset$ for each distinct vertices $u, v \in V_t$. As $(M_v^{t'})_{v \in (Z_{t'} \cup S_{t'})}$ is a model, we have $M_u^{t'} \cap M_v^{t'} = \emptyset$ for each $u \neq v \in V_t$. It follows that $M_u \cap M_v = \emptyset$ for each distinct $u, v \in V_t$. Now if $uv \in E(G[[V_t]])$ and $uv \notin E(G)$, there must exist some edge-separation $(Y_{t'}, S_{t'}, Z_{t'})$ such that $u, v \in S_{t'}$ and there exists a path from u to v in $G[S_{t'} \cup Y_{t'}]$. In particular, there must exist $u' \in M_u^{t'}$ and $v' \in M_v^{t'}$ such that $u'v' \in E(G)$. As $u' \in M_u$ and $v' \in M_v$, we proved that $(M_v)_{v \in V_t}$ is a faithful model of $G[[V_t]]$ in G . \square

For every tangle \mathcal{T} of a graph G , we denote by \mathcal{T}_{\min} its set of minimal separations (here and in the remainder, minimality of separations is always with respect to the partial order \preceq defined above). If \mathcal{T} has order 4, then we let \mathcal{T}_{nd} be its set of non-degenerate minimal separations.

Remark 3.20. Let G be locally finite, let \mathcal{T} be a G -tangle of order 4, and let (Y, S, Z) be a degenerate separation of G . Then $(Y, S, Z) \in \mathcal{T}$. This is a direct consequence of [Gro16a, Lemma 3.3], which states that if \mathcal{T} is a tangle of order k then for every separation (Y, S, Z) of order $k-1$ such that $|Y \cup S| \leq \frac{3}{2}(k-1)$ we have $(Y, S, Z) \in \mathcal{T}$.

For every tangle \mathcal{T} of order 4, we let:

$$X_{\mathcal{T}} := \bigcap_{\substack{(Y, S, Z) \in \mathcal{T}, \\ (Y, S, Z) \text{ is non-degenerate}}} (Z \cup S).$$

Note that if \mathcal{T} is an evasive tangle, then $X_{\mathcal{T}}$ is empty. In this case, and because G is 3-connected, there exists a unique end ω of degree 3 such that for any finite subset \mathcal{S} of \mathcal{T} , the end ω lies in

$$\bigcap_{\substack{(Y, S, Z) \in \mathcal{S}, \\ (Y, S, Z) \text{ is non-degenerate}}} (Z \cup S).$$

Remark 3.21. If $(Y, S, Z), (Y', S', Z') \in \mathcal{T}$ are such that $(Y', S', Z') \preceq (Y, S, Z)$ and (Y, S, Z) is non-degenerate, then it is easy to see that (Y', S', Z') is also non-degenerate (recall that G is 3-connected). Also if \mathcal{T} is a region tangle, for every $(Y, S, Z) \in \mathcal{T}$, there exists a separation $(Y', S', Z') \in \mathcal{T}_{\min}$ such that $(Y', S', Z') \preceq (Y, S, Z)$. These observations imply that if \mathcal{T} is a region tangle of order 4 and G is 3-connected and locally finite, then:

$$X_{\mathcal{T}} = \bigcap_{(Y,S,Z) \in \mathcal{T}_{\text{nd}}} (Z \cup S).$$

Two separations $(Y_1, S_1, Z_1), (Y_2, S_2, Z_2)$ are *orthogonal* if $(Y_1 \cup S_1) \cap (Y_2 \cup S_2) \subseteq S_1 \cap S_2$ (see Figure 3a). A set \mathcal{N} of separations is said to be *orthogonal* if its separations are pairwise orthogonal. One can easily show that the set of minimal separations of a (region) tangle of order at most 3 is orthogonal. This does not hold for tangles of order 4, but Grohe [Gro16a] proved that for tangles of order 4, minimal separations can only cross in a restricted way. Two separations (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) are *crossing* if $Y_1 \cap Y_2 = S_1 \cap S_2 = \emptyset$ and there is an edge $s_1 s_2 \in E(G)$, with $S_1 \cap Y_2 = \{s_1\}$ and $S_2 \cap Y_1 = \{s_2\}$ (see Figure 3b). In this case, we call $s_1 s_2$ the *crossedge* of (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) . We denote by $E_{\text{nd}}^{\times}(\mathcal{T})$ the set of crossedges of \mathcal{T}_{nd} . Lemma 4.16 from [Gro16a] generalizes to region tangles of order 4 of locally finite graphs:

Lemma 3.22 (Lemma 4.16 and Corollary 4.20 in [Gro16a]). *Let G be a locally finite 3-connected graph. Let \mathcal{T} be a region G -tangle of order 4. Then every two distinct minimal separations of \mathcal{T} are either crossing or orthogonal. Moreover, $E_{\text{nd}}^{\times}(\mathcal{T})$ forms a matching in G .*




FIGURE 3. Interaction between minimal separations. The white zones represent empty sets while the grey represent potentially non-empty sets.

In [Gro16a], orthogonal sets of separations are presented as the nice case, as they allow to efficiently find quasi-4-connected regions. We show that, up to some additional assumptions, this observation still holds in the locally finite case. We recall that for a tangle \mathcal{T} of order 4, \mathcal{T}_{nd} denote its set of minimal non-degenerate separations.

Lemma 3.23. *Let G be a locally finite 3-connected graph. Let \mathcal{T} be a region G -tangle of order 4. Assume that \mathcal{T}_{nd} is orthogonal. Then $X_{\mathcal{T}} \neq \emptyset$ and the torso $G[[X_{\mathcal{T}}]]$ has size 3 or is a quasi-4-connected minor of G .*

Proof. If every separation of order 3 in G is degenerate, then G is quasi-4-connected and all the separations of \mathcal{T}_{nd} are non-proper. It follows that $G = G[[X_{\mathcal{T}}]]$ and the desired properties hold.

Assume now that G has a proper non-degenerate separation (Y, S, Z) of order 3. As \mathcal{T} is a region tangle, there is a separation $(Y', S', Z') \in \mathcal{T}_{\text{min}}$ such that

$(Y', S', Z') \preceq (Y, S, Z)$. As observed in Remark 3.21, $(Y', S', Z') \in \mathcal{T}_{\text{nd}}$. We claim that $S' \subseteq X_{\mathcal{T}}$ so $X_{\mathcal{T}} \neq \emptyset$: let $(Y_0, S_0, Z_0) \in \mathcal{T}_{\text{nd}} \setminus \{(Y', S', Z')\}$. As (Y_0, S_0, Z_0) and (Y', S', Z') are orthogonal, we must have: $S' \cap Y_0 = \emptyset$ so $S' \subseteq Z_0 \cup S_0$. As \mathcal{T} is a region tangle, the equality $X_{\mathcal{T}} = \bigcap_{(Y,S,Z) \in \mathcal{T}_{\text{nd}}} (Z \cup S)$ holds, and thus we proved that $S' \subseteq X_{\mathcal{T}}$, and so $X_{\mathcal{T}} \neq \emptyset$. Moreover, as G is 3-connected, the separations of \mathcal{T}_{min} have order 3 so $|X_{\mathcal{T}}| \geq |S'| \geq 3$.

We now assume that $|X_{\mathcal{T}}| \geq 4$ and show that $G[[X_{\mathcal{T}}]]$ is quasi-4-connected. Since G is 3-connected and $|X_{\mathcal{T}}| \geq 4$, $G[[X_{\mathcal{T}}]]$ is 3-connected (any proper separation of order at most 2 in $G[[X_{\mathcal{T}}]]$ would induce a proper separation of order at most 2 in G). If $|X_{\mathcal{T}}| = 4$, then $G[[X_{\mathcal{T}}]]$ is clearly also quasi-4-connected so we can assume that $|X_{\mathcal{T}}| \geq 5$. Suppose that $G[[X_{\mathcal{T}}]]$ is not 4-connected and let (Y_0, S_0, Z_0) be a proper separation of $G[[X_{\mathcal{T}}]]$ of order at most 3. We will prove that $|Y_0| = 1$ or $|Z_0| = 1$, which will immediately imply that $G[[X_{\mathcal{T}}]]$ is quasi-4-connected. We let Y_1 be the union of all connected components of $G - S_0$ that intersect Y_0 or have a neighbor in Y_0 . Let $S_1 := S_0$ and $Z_1 := V(G) \setminus (S_0 \cup Y_1)$. By definition of the torso $G[[X_{\mathcal{T}}]]$, (Y_1, S_1, Z_1) is a proper separation of order at most 3 in G , hence we must have $|S_0| = 3$ as G is 3-connected. Assume first that $(Y_1, S_1, Z_1) \in \mathcal{T}$. If (Y_1, S_1, Z_1) is non-degenerate, then $X_{\mathcal{T}} \cap Y_0 \subseteq X_{\mathcal{T}} \cap Y_1 = \emptyset$ by definition of $X_{\mathcal{T}}$. It follows that $Y_0 = \emptyset$, which contradicts the assumption that (Y_0, S_0, Z_0) is proper. If (Y_1, S_1, Z_1) is degenerate, then $|Y_0| \leq |Y_1| = 1$ so as (Y_0, S_0, Z_0) is proper we must have $|Y_0| = 1$. The case $(Z_1, S_1, Z_1) \in \mathcal{T}$ is symmetric. Hence we proved that every separation (Y, S, Z) of $G[[X_{\mathcal{T}}]]$ of order at most 3 satisfies $|Y| \leq 1$ or $|Z| \leq 1$ so we are done.

Finally the fact that $G[[X_{\mathcal{T}}]]$ is a minor of G easily follows from Lemma 3.19: we consider the tree-decomposition (T, \mathcal{V}) where T is a star with a central vertex z_0 and one edge $z_0 z_i$ for each $(Y_i, S_i, Z_i) \in \mathcal{T}_{\text{nd}}$. We let $V_{z_0} := X_{\mathcal{T}}$ and $V_{z_i} := Y_i \cup S_i$ for each $(Y_i, S_i, Z_i) \in \mathcal{T}_{\text{nd}}$. The fact that (T, \mathcal{V}) is a tree-decomposition follows from the orthogonality of \mathcal{T}_{nd} . Hence by Lemma 3.19, $G[[X_{\mathcal{T}}]]$ is a minor of G . \square

Whenever \mathcal{T}_{nd} is not orthogonal, Lemma 3.23 does not hold anymore and if we want to obtain a canonical tree-decomposition, we will need to consider a larger set, whose torso is not necessarily quasi-4-connected, but can be defined uniquely from the structural properties of \mathcal{T} , which will ensure that the resulting decomposition is canonical. For every region tangle \mathcal{T} of order 4 we let:

$$R_{\mathcal{T}} := \left(\bigcup_{(Y,S,Z) \in \mathcal{T}_{\text{nd}}} S \right) \cup \left(\bigcap_{(Y,S,Z) \in \mathcal{T}_{\text{nd}}} Z \right).$$

The set $R_{\mathcal{T}}$ corresponds to the set called $R^{(0)}$ in [Gro16a, Section 4.5]. Note that we always have $X_{\mathcal{T}} \subseteq R_{\mathcal{T}}$ and that equality holds when \mathcal{T}_{nd} is orthogonal. To illustrate the definition of $R_{\mathcal{T}}$, it is helpful to go back to the graph G on Figure 1 (left). Then G has a unique tangle \mathcal{T} of order 4, the set \mathcal{T}_{nd} is the set of separations (Y, S, Z) of order 3 where Y is a triangular face of G , $S = N(Y)$ and $Z = V(G) \setminus (Y \cup S)$. Hence on this example, the set of crossededges $E_{\text{nd}}^{\times}(G)$ is the set of edges joining two triangular faces, and $R_{\mathcal{T}} = V(G)$.

While the proof of the following result was originally written for finite graphs, it immediately generalizes to locally finite graphs.

Lemma 3.24 (Lemma 4.32 in [Gro16a]). *If G is a locally finite 3-connected graph and if \mathcal{T} is a region tangle of order 4 in G , then $G[[R_{\mathcal{T}}]]$ is a faithful minor of G .*

For each $(Y, S, Z) \in \mathcal{T}_{\text{nd}}$, the *fence* $\text{fc}(S)$ of S in G is the union of

- the subset of vertices of S that are not the endpoint of some crossed edge of \mathcal{T} , and
- the subset of vertices s' such that ss' is a crossed edge of \mathcal{T} and $s \in S$.

In particular, as the crossed edges form a matching in G (Lemma 3.22), $|\text{fc}(S)| = |S| = 3$ for each $(Y, S, Z) \in \mathcal{T}_{\text{nd}}$. A consequence of Lemma 3.22 is the following:

Lemma 3.25. *Let G be a locally finite 3-connected graph and \mathcal{T} be a region G -tangle of order 4. Then G has a tree-decomposition (T, \mathcal{V}) of adhesion 3 where $\mathcal{V} = (V_t)_{t \in V(T)}$ and T is a star with central vertex z_0 such that $V_{z_0} = R_{\mathcal{T}}$. If moreover, \mathcal{T} is the unique G -tangle of order 4, then (T, \mathcal{V}) is canonical and every bag except possibly V_{z_0} is finite.*

Proof. We let:

$$V(T) := \{z_0\} \cup \{z_C : C \text{ connected component of } G - R_{\mathcal{T}}\}$$

where we choose the z_C 's to be pairwise distinct nodes. We let T be the star with vertex set $V(T)$ and central vertex z_0 , and we define $\mathcal{V} = (V_t)_{t \in V(T)}$ by setting $V_{z_0} := R_{\mathcal{T}}$, and for each connected component C of $G - R_{\mathcal{T}}$: $V_{z_C} := C \cup N(C)$. It is not hard to check that (T, \mathcal{V}) is a tree-decomposition of G . By [Gro16a, Lemma 4.31] (whose proof extends to the locally finite case), for each component C of $G - R_{\mathcal{T}}$ there exists a unique separator S such that $(Y, S, Z) \in \mathcal{T}_{\text{nd}}$ for some separation (Y, S, Z) , $N(C) = \text{fc}(S)$ and $C \subseteq Y$. This implies that (T, \mathcal{V}) has adhesion 3, so in particular its edge-separations are tight.

We now prove the second part of Lemma 3.25 and assume that \mathcal{T} is the unique tangle of order 4 of G . Then \mathcal{T} is $\text{Aut}(G)$ -invariant, and (T, \mathcal{V}) is clearly canonical. If some V_{z_C} is infinite for some $z_C \neq z_0$, then as G is locally finite, $G[V_{z_C}]$ has at least one infinite connected component, and hence there exists some end ω living in $G[V_{z_C}]$. In particular, ω induces some G -tangle \mathcal{T}_ω of order 4. We let $(Y, S, Z) \in \mathcal{T}_{\text{nd}}$ be the separation given by [Gro16a, Lemma 4.31] such that $N(C) = \text{fc}(S)$ and $C \subseteq Y$. Then (Y, S, Z) distinguishes \mathcal{T}_ω from \mathcal{T} , which contradicts the uniqueness of \mathcal{T} in G . \square

Note that in the non-orthogonal case, the tree-decomposition from Lemma 3.25 is not the same as the one from [Gro16a], as the torso $G[R_{\mathcal{T}}]$ associated to the center of the star might not be quasi-4-connected. However, we will prove in Subsection 3.7 that $G[R_{\mathcal{T}}]$ still enjoys the same useful properties as a quasi-transitive quasi-4-connected graph, namely it is either planar or has bounded treewidth. We note that the crucial ingredient that allows us to obtain a canonical tree-decomposition in the second part of Lemma 3.25 (contrary to Grohe's decomposition) is the assumption that \mathcal{T} is the unique G -tangle of order 4. So one of the most important steps in the proof of our main results will be a reduction to the case where graphs have a single tangle of order 4.

3.5. Contracting a single crossed edge. In what follows, we let G be a locally finite 3-connected graph, and \mathcal{T} be a region tangle of order 4 in G . Recall that by Lemma 3.22 the set $E_{\text{nd}}^\times(\mathcal{T})$ of the crossed edges forms a matching. We will see that contracting a crossed edge results in a 3-connected graph G' that has a tangle \mathcal{T}' of order 4 induced by \mathcal{T} [Gro16a, Subsection 4.5]. More precisely, \mathcal{T}' contains as a subset the projection of \mathcal{T} with respect to the minor G' of G . We give here an

overview of the lemmas stated in [Gro16a, Subsection 4.5] which all hold when G is locally finite instead of finite, by using the exact same proofs. The only additional property that we need in the locally finite case is that \mathcal{T}' is still a region tangle, which is proved in Lemma 3.29 below.

In the remainder of this subsection, we let (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) be two crossing separations of \mathcal{T}_{nd} with crossed edge $s_1 s_2$. *Contracting* $s_1 s_2$ consists in deleting s_1 and s_2 and adding a new vertex s' whose neighborhood is equal to $N_G(s_1) \cup N_G(s_2) \setminus \{s_1, s_2\}$. We denote by G' the graph obtained after contracting $s_1 s_2$. The *projection* (referred to as *contraction* in [Gro16a]) of a set X of vertices of G is defined as

$$X^\vee := \begin{cases} X & \text{if } X \cap \{s_1, s_2\} = \emptyset \\ X \setminus \{s_1, s_2\} \cup \{s'\} & \text{if } X \cap \{s_1, s_2\} \neq \emptyset. \end{cases}$$

Given a set X' of vertices of G' , the *expansion* X'_\wedge of X' is defined as

$$X'_\wedge := \begin{cases} X' & \text{if } s' \notin X' \\ X' \setminus \{s'\} \cup \{s_1, s_2\} & \text{if } s' \in X'. \end{cases}$$

Observe that for all $X' \subseteq V(G')$, we have $(X'_\wedge)^\vee = X'$ and for all $X \subseteq V(G)$, we have $X \subseteq (X^\vee)_\wedge$ (where the inclusion might be strict). We also define for every $(Y, S, Z) \in \text{Sep}_{<4}(G)$:

$$(Y, S, Z)^\vee := \begin{cases} (Y^\vee \setminus \{s'\}, S^\vee, Z^\vee \setminus \{s'\}) & \text{if } S \cap \{s_1, s_2\} \neq \emptyset \\ (Y^\vee, S^\vee, Z^\vee) & \text{if } S \cap \{s_1, s_2\} = \emptyset. \end{cases}$$

Note that $(Y, S, Z)^\vee$ is exactly the projection $\pi_{\mathcal{M}}(Y, S, Z)$ with respect to the model $\mathcal{M} = (\{v\}_\wedge)_{v \in V(G')}$ of G' in G .

In the context of finite graphs, [Gro16a] proves the following lemmas that extend directly to the locally finite case:

Lemma 3.26 (Corollary 4.24 in [Gro16a]). *The graph G' resulting from the contraction of $s_1 s_2$ is 3-connected.*

Lemma 3.27 (Lemmas 4.26 and 4.27 in [Gro16a]). *There exists a tangle \mathcal{T}' of order 4 in G' containing the projection of \mathcal{T} with respect to the model $\mathcal{M} = (\{v\}_\wedge)_{v \in V(G')}$.*

Note that the projection of \mathcal{T} with respect to \mathcal{M} is exactly the set $\{(Y, S, Z)^\vee : (Y, S, Z) \in \mathcal{T}\}$. In the remainder of this subsection, we let \mathcal{T}' be the tangle given by Lemma 3.27. In [Gro16a], the author gives an explicit definition of \mathcal{T}' , but for the sake of clarity we only summarize here the properties of \mathcal{T}' that will be of interest for our purposes.

Note that the inclusion $\{(Y, S, Z)^\vee : (Y, S, Z) \in \mathcal{T}\} \subseteq \mathcal{T}'$ is strict in general, as some separations from $\text{Sep}_{<4}(G')$ might not be projections of separations from $\text{Sep}_{<4}(G)$. The next lemma intuitively states that every separation of \mathcal{T}' is close to an element from $\{(Y, S, Z)^\vee : (Y, S, Z) \in \mathcal{T}\} \subseteq \mathcal{T}'$.

Lemma 3.28 (Definition of \mathcal{T}' and Lemmas 4.23 and 4.25 in [Gro16a]). *For every separation $(Y', S', Z') \in \mathcal{T}'$ such that $s' \in S'$ and $G'[Z']$ is connected, there exists a separation $(Y, S, Z) \in \mathcal{T}$ such that $S^\vee = S'$ and $Z \setminus S_\vee = Z' \setminus \{s_1, s_2\} = Z'$.*

As Lemma 3.28 is not exactly stated this way in [Gro16a], we briefly sketch how to obtain it. If $(Y', S', Z') \in \mathcal{T}'$ is such that $s' \in S'$, then by [Gro16a, Lemma 4.25] there exists a (unique) connected component C of $G \setminus S'_\vee$ such that the separations (Y'', S'', Z'') of \mathcal{T}' such that $S'' = S'$ are exactly the ones such that $C \subseteq Z''$, and for

which every separation $(Y, S, Z) \in \mathcal{T}$ such that $S^\vee = S'$ satisfies $C \subseteq Z$. [Gro16a, Lemmas 4.23, 4.25] and the fact that \mathcal{T} is a tangle ensure the existence of a separation $(Y, S, Z) \in \mathcal{T}$ such that $S^\vee = S'$ and $C = Z \setminus \{s_1, s_2\} = Z \setminus S$. In particular by Lemma 3.27 the projection $(Y, S, Z)^\vee = (Y \setminus S', S', C)$ is in \mathcal{T}' so if we assume that $G'[Z']$ is connected, the choice of C imposes $C \subseteq Z'$, thus $Z' = C$. It implies that (Y, S, Z) satisfies the property described in Lemma 3.28.

Lemma 3.29. *\mathcal{T}' is a region tangle.*

Proof. Assume for the sake of contradiction that \mathcal{T}' contains an infinite strictly decreasing sequence of separations $(Y'_n, S'_n, Z'_n)_{n \in \mathbb{N}}$. By Lemma 3.26, G' is 3-connected, so the only possible non-proper separation (Y_n, S_n, Z_n) is for $n = 0$, thus we may assume that all the separations (Y_n, S_n, Z_n) are tight. By Lemma 3.1, there are finitely many integers n for which $s' \in S'_n$. Up to extracting an infinite subsequence, one can assume that for all n , either $s' \in Y'_n$ or $s' \in Z'_n$. If there exists N such that $s' \in Y'_N$ then for all $n \geq N$ we must have $s' \in Y_n$ by definition of \preceq . Up to extracting another infinite subsequence, we can assume that either $s' \in Y'_n$ for all n or $s' \in Z'_n$ for all n . As a result, and because \mathcal{T}' contains the projection of \mathcal{T} with respect to \mathcal{M} , $((Y'_n)^\wedge, S'_n, (Z'_n)^\wedge)_{n \in \mathbb{N}}$ is an infinite decreasing sequence of separations of order 3 in \mathcal{T} , contradicting the fact that \mathcal{T} is well-founded. \square

We conclude this subsection with the following result relating the degeneracy of minimal separations in G and G' . Its proof is the same as the proof of [Gro16a, Lemma 4.28], which directly translates to the locally finite case. To be more precise, we also need the additional assumption that \mathcal{T}' is a region tangle to make the proof work, which is given by Lemma 3.29.

Lemma 3.30 (Lemma 4.28 and Corollary 4.29 in [Gro16a]). *Either G' is 4-connected and $\mathcal{T}'_{\min} = \{(\emptyset, \emptyset, V(G'))\}$, or*

$$\mathcal{T}'_{\min} = \{(Y, S, Z)^\vee : (Y, S, Z) \in \mathcal{T}_{\min} \text{ and } S^\vee \text{ is a separator of } G'\}.$$

In the latter case, for all $(Y, S, Z) \in \mathcal{T}_{\min}$, (Y, S, Z) is non-degenerate if and only if $(Y, S, Z)^\vee$ is non-degenerate. Moreover, $E_{\text{nd}}^\times(\mathcal{T}') = E_{\text{nd}}^\times(\mathcal{T}) \setminus \{s_1 s_2\}$.

3.6. Contracting all the crossedges. In the previous subsection we studied the consequences of contracting a single crossed edge in G . However, in our application we will need to contract *all* crossed edges of $E_{\text{nd}}^\times(\mathcal{T})$ (which form a matching in G). We now study how this affects G .

Before going further, we will need to introduce some notation, extending the notation from [Gro16a] to the infinite case. For convenience we write $M := E_{\text{nd}}^\times(\mathcal{T})$ (and recall that M is a matching in G). For every subset $L \subseteq M$ of crossed edges, we let $G^{\setminus L/}$ be the graph obtained from G after contracting each edge $uv \in L$ into a new vertex $s_{u,v}$. Note that the order in which the edges are contracted is irrelevant in the definition of $G^{\setminus L/}$.

We denote $\bar{L} = M \setminus L$. In this section we will also often use the notation $L - L'$ instead of $L \setminus L'$, to avoid any possible confusion when reading superscripts (for instance we will write $G^{\setminus L - L'/}$ instead of $G^{\setminus L \setminus L'/}$).

For every $L \subseteq M$, for every vertex $x \in V(G)$, we let

$$x^{\setminus L/} := \begin{cases} x & \text{if } x \in X \setminus V(L) \\ s_{u,v} & \text{if } x \text{ is the endpoint of a crossed edge } uv \in L. \end{cases}$$

For every subset $X \subseteq V(G)$ of vertices, we let $X^{\setminus L/} := \{x^{\setminus L/} : x \in X\}$ be the projection of X to $G^{\setminus L/}$.

Remark 3.31. Note that for every disjoint subsets $K, L \subseteq M$ and for all $X \subseteq V(G)$, $X^{\setminus K \cup L/} = (X^{\setminus L/})^{\setminus K/} = (X^{\setminus K/})^{\setminus L/}$.

For every $X' \subseteq V(G^{\setminus M/})$ and $L \subseteq M$, let $X'_{/L\setminus}$ denote the maximal set $X \subseteq V(G^{\setminus L/})$ such that $X^{\setminus L/} = X'$. In other words $X'_{/L\setminus}$ is the set of vertices obtained after “uncontracting” the edges of L in X . Note that with the notation introduced above we have $G = G^{\setminus \emptyset/}$. Given a separation (Y, S, Z) of G , we define

$$(Y, S, Z)^{\setminus L/} := (Y^{\setminus L/} \setminus S^{\setminus L/}, S^{\setminus L/}, Z^{\setminus L/} \setminus S^{\setminus L/}).$$

Note that when $L = \{s_1 s_2\}$ consists of a single edge, we recover the notions of the previous subsection; with our previous notation this gives: $x^{\setminus L/} = x^\vee$, $X^{\setminus L/} = X^\vee$ and $(Y, S, Z)^{\setminus L/} = (Y, S, Z)^\vee$.

For each finite subset of crossed edges $L \subseteq M$, and every enumeration (e_1, \dots, e_ℓ) of the edges of L , we let $\mathcal{T}^{\setminus (e_1, \dots, e_\ell)/}$ denote the tangle of $G^{\setminus L/}$ obtained after iteratively applying Lemma 3.27 to the graphs $G_0 := G, G_1, \dots, G_\ell$ with $G_i := G^{\setminus \{e_1, \dots, e_i\}/}$ for each $i \in [\ell]$.

Lemma 3.32 (Lemma 4.30 (5) in [Gro16a]). *For every enumeration (e_1, \dots, e_ℓ) of a finite set $L \subseteq M$ of crossed edges and every permutation σ of $[\ell]$, $\mathcal{T}^{\setminus (e_1, \dots, e_\ell)/} = \mathcal{T}^{\setminus (e_{\sigma(1)}, \dots, e_{\sigma(\ell)})/}$.*

In the remainder of the subsection, for every finite subset $L \subseteq M$, we will denote with $\mathcal{T}^{\setminus L/}$ the unique tangle associated to any enumeration of L given by Lemma 3.32.

Intuitively when G is finite, one of the main properties of $\mathcal{T}^{\setminus L/}$ is that separations of $\mathcal{T}_{\text{nd}}^{\setminus L/}$ are in correspondence with separations of \mathcal{T}_{nd} , and that the only crossing pairs between elements of $\mathcal{T}^{\setminus L/}$ correspond to pairs which were already crossing in \mathcal{T} . Thus after each contraction, we reduce the number of crossed edges, hence when $L = M$, the family $\mathcal{T}_{\text{nd}}^{\setminus L/}$ must be orthogonal and we can apply results from previous subsections to the graph $G^{\setminus L/}$. We now show formally how to extend the relevant proofs of [Gro16a] to the locally finite case.

In [Gro16a], the author proved that if G is finite and 3-connected, for every $L \subseteq M$, the graph $G^{\setminus L/}$ is 3-connected and that $\mathcal{T}^{\setminus L/}$ is a tangle of order 4 induced by \mathcal{T} in $G^{\setminus L/}$. Using the results from the previous subsection, this immediately extends to $G^{\setminus L/}$ and $\mathcal{T}^{\setminus L/}$ when G is locally finite and L is finite, by induction on the size of L .

Theorem 3.33 (Generalization of Lemma 4.30 in [Gro16a]). *Let $L \subseteq M$ be a finite set of crossed edges. Then we have*

- (1) $G^{\setminus L/}$ is 3-connected
- (2) $\mathcal{T}^{\setminus L/}$ is a region tangle of order 4 of $G^{\setminus L/}$ such that

$$\mathcal{T}_{\text{min}}^{\setminus L/} = \{(Y, S, Z)^{\setminus L/} : (Y, S, Z) \in \mathcal{T}_{\text{min}} \text{ such that } S^{\setminus L/} \text{ is a separator of } G^{\setminus L/}\}$$

or $\mathcal{T}_{\text{min}}^{\setminus L/} = \{(\emptyset, \emptyset, V(G'))\}$ if $L = M$ is finite and $G^{\setminus L/}$ is 4-connected.

- (3) $E_{\text{nd}}^\times(\mathcal{T}^{\setminus L/}) = E_{\text{nd}}^\times(\mathcal{T}) \setminus L$
- (4) $\mathcal{T}^{\setminus L/}$ contains the projection $\{(Y, S, Z)^{\setminus L/} : (Y, S, Z) \in \mathcal{T}\}$ of \mathcal{T} with respect to the model $\mathcal{M} = (\{v\}_{/L\setminus})_{v \in V(G^{\setminus L/})}$.

We will now extend Theorem 3.33 to the case where $L \subseteq M$ is infinite. Given a set $X \subseteq V(G^{\setminus M/})$, we denote by $M(X) \subseteq M$ the subset of edges of G contracted to a vertex in X .

Lemma 3.34. *The graph $G^{\setminus M/}$ is 3-connected.*

Proof. Assume for the sake of contradiction that $G^{\setminus M/}$ has a separator S of order at most 2. Then the set $L := M(S)$ has size at most 2 and S is a separator of order 2 of $G^{\setminus L/}$. This contradicts Theorem 3.33. \square

We now let $L \subseteq M$ be any (not necessarily finite) subset of crossed edges and give a general definition of $\mathcal{T}^{\setminus L/}$ extending the previous one. For every $(Y', S', Z') \in \text{Sep}_{<4}(G^{\setminus L/})$, we let $L' := M(S')$. Note that L' is finite, and that $(Y'_{/\overline{L\setminus}}, S', Z'_{/\overline{L\setminus}})$ is a separation of order at most 3 in $G^{\setminus L'/}$. We define $\mathcal{T}^{\setminus L/}$ as the family of separations (Y', S', Z') of $G^{\setminus L/}$ such that $(Y'_{/\overline{L\setminus}}, S', Z'_{/\overline{L\setminus}}) \in \mathcal{T}^{\setminus L'/}$. Note that when L is finite, $(Y', S', Z') = (Y'_{/\overline{L\setminus}}, S', Z'_{/\overline{L\setminus}})^{\setminus L-L'}$ and thus iterative applications of Lemma 3.27 together with Lemma 3.32 imply that our definition of $\mathcal{T}^{\setminus L/}$ coincides with the one we gave above for finite subsets $L \subseteq M$.

Thanks to Remark 3.31, for all $L \subseteq M$, $\mathcal{T}^{\setminus M/} = (\mathcal{T}^{\setminus L/})^{\setminus \overline{L/}}$ and $G^{\setminus M/} = (G^{\setminus L/})^{\setminus \overline{L/}}$. We say that a set $X \subseteq V(G)$ hits the edges of L once if for all $e \in L$, $|X \cap e| = 1$.

Lemma 3.35. *$\mathcal{T}^{\setminus M/}$ is a region tangle of order 4 in $G^{\setminus M/}$ such that $\{(Y, S, Z)\}^{\setminus M/} : (Y, S, Z) \in \mathcal{T}\} \subseteq \mathcal{T}^{\setminus M/}$.*

Proof. We first prove that $\mathcal{T}^{\setminus M/}$ is a tangle of order 4. To prove (T1), let $(Y', S', Z') \in \text{Sep}_{<4}(G^{\setminus M/})$ and $L := M(S')$. Then L has size at most 3, so by Theorem 3.33, $\mathcal{T}^{\setminus L/}$ is a region tangle of order 4 of $G^{\setminus L/}$. As $(Y, S, Z) := (Y'_{/\overline{L\setminus}}, S', Z'_{/\overline{L\setminus}})$ is a separation of order 3 of $G^{\setminus L/}$ and $\mathcal{T}^{\setminus L/}$ is a tangle of order 4, either $(Y, S, Z) \in \mathcal{T}^{\setminus L/}$ or $(Z, S, Y) \in \mathcal{T}^{\setminus L/}$. By definition of $\mathcal{T}^{\setminus M/}$ we then have either $(Z', S', Y') \in \mathcal{T}^{\setminus M/}$ or $(Y', S', Z') \in \mathcal{T}^{\setminus M/}$, implying that $\mathcal{T}^{\setminus M/}$ satisfies (T1).

To prove (T2), let $(Y'_1, S'_1, Z'_1), (Y'_2, S'_2, Z'_2), (Y'_3, S'_3, Z'_3) \in \mathcal{T}^{\setminus M/}$. Let $L := M(S'_1 \cup S'_2 \cup S'_3)$. Once again L is finite with size at most 9 and for all $i \in \{1, 2, 3\}$, $(Y_i, S_i, Z_i) := ((Y'_i)_{/\overline{L\setminus}}, S'_i, (Z'_i)_{/\overline{L\setminus}})$ is a separation of order 3 of $G^{\setminus L/}$.

Claim 3.36. *For every $i \in \{1, 2, 3\}$, $(Y_i, S_i, Z_i) \in \mathcal{T}^{\setminus L/}$.*

Proof of the Claim: Assume that $i = 1$, the other cases being symmetric. We let $L_1 := M(S'_1)$. Then by definition of $\mathcal{T}^{\setminus M/}$, $(Y''_1, S''_1, Z''_1) := ((Y'_1)_{/\overline{L_1\setminus}}, S'_1, (Z'_1)_{/\overline{L_1\setminus}}) \in \mathcal{T}^{\setminus L_1/}$. Our goal is to show that $(Y_1, S_1, Z_1) = (Y''_1, S''_1, Z''_1)^{\setminus L-L_1/}$. As both L and L_1 are finite and $L_1 \subseteq L$, iterative applications of Lemmas 3.27 and Lemma 3.32 imply $\mathcal{T}^{\setminus L/}$ must contain the projection of $\mathcal{T}^{\setminus L_1/}$ with respect to the model $\mathcal{M} = (\{v\}_{/(L \setminus L_1)\setminus})_{v \in V(G) \setminus L/}$. Thus if we succeed to prove that

$$(1) \quad (Y_1, S_1, Z_1) = (Y''_1, S''_1, Z''_1)^{\setminus L-L_1/},$$

we immediately obtain that $(Y_1, S_1, Z_1) \in \mathcal{T}^{\setminus L/}$, which concludes the claim.

To prove that (1) holds, note first that every edge of M is contracted in $G^{\setminus M/}$ so in particular it has its endpoints in exactly one of the three sets $(Y'_1)_{/M\setminus}$, $(S'_1)_{/M\setminus}$ and $(Z'_1)_{/M\setminus}$. In particular by definition of L_1 , the edges of L_1 are all disjoint from $(Y'_1)_{/M\setminus}$ and thus $(Y'_1)_{/M\setminus} = (Y'_1)_{/\overline{L_1\setminus}}$. This implies that $Y_1 = (Y'_1)_{/\overline{L\setminus}} = ((Y'_1)_{/\overline{L_1\setminus}})^{\setminus L-L_1/} =$

$(Y_1'')^{\setminus L-L_1/}$. As S_1' is disjoint from Y_1' in $G^{\setminus M/}$, it is also disjoint from Y_1' in $G^{\setminus L/}$ so we have $Y_1 = (Y_1'')^{\setminus (L \setminus L_1)/} \setminus S_1''$. Symmetric arguments give $Z_1 = (Z_1'')^{\setminus L-L_1/} \setminus S_1''$, and as $S_1'' = S_1' = S_1$, we get the desired equality. \diamond

By Theorem 3.33, $\mathcal{T}^{\setminus L/}$ is a region tangle of order 4 so Claim 3.36 implies that either $Z_1 \cap Z_2 \cap Z_3 \neq \emptyset$ or there exists an edge of $G^{\setminus L/}$ with both endpoints in $Z_1 \cup Z_2 \cup Z_3$. If $Z_1 \cap Z_2 \cap Z_3 \neq \emptyset$, then $Z_1' \cap Z_2' \cap Z_3' = (Z_1 \cap Z_2 \cap Z_3)^{\setminus \bar{L}/} \neq \emptyset$. Otherwise, there is an edge $e \in E(G^{\setminus L/})$ that has an endpoint in each Z_i , in which case, either $Z_1' \cap Z_2' \cap Z_3' \neq \emptyset$ if $e \in \bar{L}$, or $e^{\setminus \bar{L}/}$ is an edge of $G^{\setminus M/}$ which has an endpoint in each Z_i' . This proves (T2) and shows that $\mathcal{T}^{\setminus M/}$ is a tangle of order 4.

We now prove the inclusion $\{(Y, S, Z)^{\setminus M/} : (Y, S, Z) \in \mathcal{T}\} \subseteq \mathcal{T}^{\setminus M/}$. Let $(Y, S, Z) \in \mathcal{T}$ and $L := M(S)$. By Theorem 3.33 (4), $(Y, S, Z)^{\setminus L/} \in \mathcal{T}^{\setminus L/}$. Write $(Y', S', Z') = (Y, S, Z)^{\setminus M/}$ and note that $(Y'_{/\bar{L}}, S', Z'_{/\bar{L}}) = (Y, S, Z)^{\setminus L/}$, thus by definition of $\mathcal{T}^{\setminus M/}$, $(Y, S, Z)^{\setminus M/} \in \mathcal{T}^{\setminus M/}$.

We now prove that $\mathcal{T}^{\setminus M/}$ is a well-founded set. For the sake of contradiction, let $((Y'_n, S'_n, Z'_n))_{n \in \mathbb{N}}$ be an infinite decreasing sequence of separations of $\mathcal{T}^{\setminus M/}$. The contradiction will follow from the next claim

Claim 3.37. *There exists an infinite decreasing sequence $((Y''_n, S''_n, Z''_n))_{n \in \mathbb{N}}$ in $\mathcal{T}^{\setminus M/}$ such that for each $n \geq 0$, $(Y''_n, S''_n, Z''_n) = (Y_n, S_n, Z_n)^{\setminus M/}$ for some $(Y_n, S_n, Z_n) \in \mathcal{T}$.*

Proof of the Claim: By Lemma 3.1 and because $G^{\setminus M/}$ is 3-connected, for any $n \in \mathbb{N}$, there are finitely many separations (Y'_m, S'_m, Z'_m) such that $S'_n \cap S'_m \neq \emptyset$. Therefore, up to considering a subsequence, we can assume that for all n , $S'_n \subseteq Y'_{n+1}$ and $S'_{n+1} \subseteq Z'_n$. In particular, as by Lemma 3.34 $G^{\setminus M/}$ is 3-connected, S'_{n+1} is included in some connected component C_n of $G^{\setminus M/}[Z'_n]$. Then we have

$$(Y'_{n+1}, S'_{n+1}, Z'_{n+1}) \preceq (V(G^{\setminus M/}) \setminus (S'_n \cup C_n), S'_n, C_n) \preceq (Y'_n, S'_n, Z'_n),$$

implying that $(V(G^{\setminus M/}) \setminus (S'_n \cup C_n), S'_n, C_n) \in \mathcal{T}^{\setminus M/}$. Hence we may also assume up to replacing (Y'_n, S'_n, Z'_n) with $(V(G^{\setminus M/}) \setminus (S'_n \cup C_n), S'_n, C_n)$ that for each $n \geq 0$, $G^{\setminus M/}[Z'_n]$ is connected.

For each $n \geq 0$, we let $L_n := M(S'_n)$. Connectedness of $G^{\setminus M/}[Z'_n]$ then implies that $G^{\setminus L_n/}[(Z'_n)_{/\bar{L}_n}]$ is connected. Observe that $|L_n| \leq 3$ successive applications of Lemma 3.28 imply that there exists some separation $(Y_n, S_n, Z_n) \in \mathcal{T}$ such that $S_n^{\setminus L_n/} = S'_n$ and $Z_n \setminus ((S'_n)_{/L_n \setminus}) = (Z'_n)_{/\bar{L}_n \setminus}$. We let $(Y''_n, S''_n, Z''_n) := (Y_n, S_n, Z_n)^{\setminus M/}$. By Theorem 3.33 (4), $(Y_n, S_n, Z_n)^{\setminus L_n/} \in \mathcal{T}^{\setminus L_n/}$. Moreover,

$$(Y_n, S_n, Z_n)^{\setminus L_n/} = ((Y''_n)_{/\bar{L}_n \setminus}, S''_n, (Z''_n)_{/\bar{L}_n \setminus}),$$

thus by definition of $\mathcal{T}^{\setminus M/}$, we have $(Y''_n, S''_n, Z''_n) = ((Y_n, S_n, Z_n)^{\setminus L_n/})^{\setminus \bar{L}_n/} \in \mathcal{T}^{\setminus M/}$.

Note that $S''_n = S_n^{\setminus M/} = S_n^{\setminus L_n/} = S'_n$ and

$$Z''_n = Z_n^{\setminus M/} \setminus S'_n = (Z_n \setminus ((S'_n)_{/L_n \setminus}))^{\setminus \bar{L}_n/} = ((Z'_n)_{/\bar{L}_n \setminus})^{\setminus \bar{L}_n/} = Z'_n.$$

We thus deduce that $(Y'_n, S'_n, Z'_n) = (Y''_n, S''_n, Z''_n)$, so $((Y'_n, S'_n, Z'_n))_{n \in \mathbb{N}}$ is an infinite decreasing sequence in $\mathcal{T}^{\setminus M/}$ satisfying the desired properties. \diamond

It remains to show how to derive a contradiction from Claim 3.37. For this let $((Y''_n, S''_n, Z''_n))_{n \in \mathbb{N}}$ and $((Y_n, S_n, Z_n))_{n \in \mathbb{N}}$ be as in Claim 3.37. Again, up to considering a subsequence, we can assume that for all n , $S''_n \subseteq Y''_{n+1}$ and $S''_{n+1} \subseteq Z''_n$. As $S''_n \cap S''_{n+1} = \emptyset$, the separators S_n and S_{n+1} cannot contain two vertices of a common crossed edge

of M . Thus, $(S''_n)_{/M\setminus} \subseteq Y_{n+1}$ and $(S''_{n+1})_{/M\setminus} \subseteq Z_n$ and hence $(Y_{n+1}, S_{n+1}, Z_{n+1}) \prec (Y_n, S_n, Z_n)$. This proves that $((Y_n, S_n, Z_n))_{n \in \mathbb{N}}$ is an infinite decreasing sequence of separations of G with respect to \mathcal{T} , contradicting the fact that \mathcal{T} is a region tangle. \square

Lemma 3.38. *Either $G^{\setminus M/}$ is 4-connected and $\mathcal{T}_{\min}^{\setminus M/} = \{(\emptyset, \emptyset, V(G)^{\setminus M/})\}$ or*

$$\mathcal{T}_{\min}^{\setminus M/} = \{(Y, S, Z)^{\setminus M/} : (Y, S, Z) \in \mathcal{T}_{\min} \text{ such that } S^{\setminus M/} \text{ is a separator of } G^{\setminus M/}\}.$$

Finally, we have $E_{\text{nd}}^{\times}(G^{\setminus M/}) = \emptyset$.

Proof. Assume that $G^{\setminus M/}$ is not 4-connected. We first prove the direct inclusion. Let $(Y', S', Z') \in \mathcal{T}_{\min}^{\setminus M/}$, $L := M(S')$ and $(Y_0, S_0, Z_0) := (Y'_{/\overline{L}}, S', Z'_{/\overline{L}})$. Then by definition of $\mathcal{T}^{\setminus M/}$, $(Y_0, S_0, Z_0) \in \mathcal{T}^{\setminus L/}$. We prove that (Y_0, S_0, Z_0) is a minimal element of $\mathcal{T}^{\setminus L/}$. Assume for a contradiction that there exists $(Y_1, S_1, Z_1) \prec (Y_0, S_0, Z_0)$ in $\mathcal{T}^{\setminus L/}$. Then $(Y_1, S_1, Z_1)^{\overline{L/}} \preceq (Y_0, S_0, Z_0)^{\overline{L/}} = (Y', S', Z')$ and $(Y_1, S_1, Z_1)^{\overline{L/}} \neq (Y', S', Z')$ because $\overline{L} \cap S' = \emptyset$. Moreover, by Theorem 3.33 (2), we may assume that (Y'_1, S'_1, Z'_1) is minimal and that $(Y_1, S_1, Z_1) = (Y'_1, S'_1, Z'_1)^{\setminus L/}$ for some $(Y'_1, S'_1, Z'_1) \in \mathcal{T}$. Thus as $(Y_1, S_1, Z_1)^{\overline{L/}} = ((Y'_1, S'_1, Z'_1)^{\setminus L/})^{\overline{L/}} = (Y'_1, S'_1, Z'_1)^{\setminus M/}$, we must have $(Y_1, S_1, Z_1) \in \mathcal{T}^{\setminus M/}$, contradicting the minimality of (Y_0, S_0, Z_0) in $\mathcal{T}^{\setminus L/}$. Hence $(Y', S', Z') = (Y_0, S_0, Z_0)^{\overline{L/}}$ for some $(Y_0, S_0, Z_0) \in \mathcal{T}_{\min}^{\setminus L/}$. Again we can apply Theorem 3.33 and write $(Y_0, S_0, Z_0) = (Y, S, Z)^{\setminus L/}$ for some $(Y, S, Z) \in \mathcal{T}_{\min}$. Thus we have:

$$(Y', S', Z') = ((Y, S, Z)^{\setminus L/})^{\overline{L/}} = (Y, S, Z)^{\setminus M/},$$

so we are done with the direct inclusion.

Conversely, let $(Y_1, S_1, Z_1) \in \mathcal{T}_{\min}$ such that $S_1^{\setminus M/}$ is a separator of $G^{\setminus M/}$. Note that by previous inclusion and because $\mathcal{T}^{\setminus M/}$ is a region tangle, it is enough to prove that for any $(Y_2, S_2, Z_2) \in \mathcal{T}$ such that $(Y_2, S_2, Z_2)^{\setminus M/} \preceq (Y_1, S_1, Z_1)^{\setminus M/}$, we have $(Y_2, S_2, Z_2)^{\setminus M/} = (Y_1, S_1, Z_1)^{\setminus M/}$. Let $(Y_2, S_2, Z_2) \in \mathcal{T}$ such that $(Y_2, S_2, Z_2)^{\setminus M/} \preceq (Y_1, S_1, Z_1)^{\setminus M/}$, $L := M(S_1 \cup S_2)$ and $\overline{L} := M \setminus L$. For $i \in \{1, 2\}$, the edges in \overline{L} are either contained in Y_i or in Z_i . Note that for each i and $L \subseteq M$:

$$S_i^{\setminus L/} \cup (Z_i^{\setminus L/} \setminus S_i^{\setminus L/}) = (Z_i \cup S_i)^{\setminus L/}$$

and

$$S_i^{\setminus L/} \cup (Y_i^{\setminus L/} \setminus S_i^{\setminus L/}) = (Y_i \cup S_i)^{\setminus L/}.$$

Thus as $(Y_1, S_1, Z_1)^{\setminus M/} \preceq (Y_2, S_2, Z_2)^{\setminus M/}$, we have $(S_1 \cup Z_1)^{\setminus M/} \subseteq (S_2 \cup Z_2)^{\setminus M/}$. By previous remark that no edge of \overline{L} is contained in Z_i , this implies that $(S_1 \cup Z_1)^{\setminus L/} \subseteq (S_2 \cup Z_2)^{\setminus L/}$. Likewise $(S_1 \cup Y_1)^{\setminus L/} \subseteq (S_2 \cup Y_2)^{\setminus L/}$. As a result, $(Y_1, S_1, Z_1)^{\setminus L/} \preceq (Y_2, S_2, Z_2)^{\setminus L/}$. By Theorem 3.33 (2), $(Y_1, S_1, Z_1)^{\setminus L/} \in \mathcal{T}_{\min}^{\setminus L/}$, thus we have $(Y_1, S_1, Z_1)^{\setminus L/} = (Y_2, S_2, Z_2)^{\setminus L/}$ and $(Y_1, S_1, Z_1)^{\setminus M/} = (Y_2, S_2, Z_2)^{\setminus M/}$, showing that $(Y_1, S_1, Z_1)^{\setminus M/} \in \mathcal{T}_{\min}^{\setminus M/}$.

We now prove that $E_{\text{nd}}^{\times}(G^{\setminus M/}) = \emptyset$. Assume that there are two crossing non-degenerate minimal 3-separations (Y''_1, S''_1, Z''_1) and (Y''_2, S''_2, Z''_2) in $\mathcal{T}^{\setminus M/}$, let $L = M(S''_1 \cup S''_2)$. For $i \in \{1, 2\}$, all crossed edges of $G^{\setminus L/}$ lie in Y''_i or in Z''_i , hence $(Y''_i, S''_i, Z''_i) = ((Y''_i)_{/\overline{L}}, S''_i, (Z''_i)_{/\overline{L}})$ is the only 3-separation of $\text{Sep}_{<4}(G^{\setminus L/})$ such that $(Y''_i, S''_i, Z''_i)^{\overline{L/}} = (Y''_i, S''_i, Z''_i)$. Since $(Y''_i, S''_i, Z''_i) \in \mathcal{T}_{\min}^{\setminus M/}$, from what we just proved, we must have $(Y''_i, S''_i, Z''_i) \in \mathcal{T}_{\min}^{\setminus M/}$. Note that the separations (Y''_i, S''_i, Z''_i) are non-degenerate in $G^{\setminus L/}$. Furthermore, as $L = M(S''_1 \cup S''_2)$, (Y''_1, S''_1, Z''_1) and (Y''_2, S''_2, Z''_2)

must be also crossing in $G^{\setminus L/}$, but this contradicts $E_{\text{nd}}^{\times}(T^{\setminus L/}) = E_{\text{nd}}^{\times}(\mathcal{T}) \setminus L$ (third item of Theorem 3.33). \square

For each $L \subseteq M$, we let $R^{\setminus L/} := R_{\mathcal{T}}^{\setminus L/}$. Note that for each $L \subseteq M$, $(R_{\mathcal{T} \setminus L/})_{/L \setminus} = R_{\mathcal{T}}$. Thus together with Lemma 3.23, this immediately gives the following, which is the locally finite extension of one of the main results from [Gro16a]:

Theorem 3.39. *Let G be a locally finite 3-connected graph, and let \mathcal{T} be a region tangle of order 4 in G . Let $M := E_{\text{nd}}^{\times}(\mathcal{T})$ be the set of crossed edges between non-degenerate minimal separations of \mathcal{T} . Then the graph $G^{\setminus M/} \llbracket R^{\setminus M/} \rrbracket$ is a quasi-4-connected minor of G .*

In order to obtain a proof of Theorem 3.7 in the locally finite case, one can either reuse the arguments from [Gro16a, Section 5], or equivalently adapt our proof from Subsection 4.3.

3.7. Planarity after uncontracting crossed edges. The difficulty is that in general, nice properties of $G^{\setminus M/} \llbracket R^{\setminus M/} \rrbracket$ are not satisfied anymore by $G \llbracket R_{\mathcal{T}} \rrbracket$. To circumvent this and find a quasi-4-connected region in G , it is proved in [Gro16a] that for every subset X' of $R_{\mathcal{T}}$ obtained by deleting one endpoint of each edge of M , the graph $G \llbracket X' \rrbracket$ is isomorphic to $G^{\setminus M/} \llbracket R^{\setminus M/} \rrbracket$. However we cannot choose such a subset X' canonically in general, as illustrated in Example 3.8. Despite the fact that uncontracting the edges of M does not preserve the quasi-4-connectivity of torsos, we now prove that at least planarity is preserved by this operation.

Proposition 3.40. *If $G^{\setminus M/} \llbracket R^{\setminus M/} \rrbracket$ is planar, then so is $G \llbracket R_{\mathcal{T}} \rrbracket$.*

This is obtained by combining the following two lemmas:

Lemma 3.41. *For every subset $L \subseteq M$, we denote $\bar{L} := M \setminus L$. Assume that for every finite subset $L \subseteq M$, $G^{\setminus \bar{L}/} \llbracket R^{\setminus \bar{L}/} \rrbracket$ is planar. Then $G \llbracket R_{\mathcal{T}} \rrbracket$ is also planar.*

Proof. Assume for the sake of contradiction that G enjoys the properties described above but that $G \llbracket R_{\mathcal{T}} \rrbracket$ is not planar. Then by Wagner's theorem [Wag37], $G \llbracket R_{\mathcal{T}} \rrbracket$ admits F as a minor, for some $F \in \{K_5, K_{3,3}\}$. Note that we can find a model $(V_v)_{v \in V(F)}$ of F such that each set V_v is finite. Then $X := \bigcup_{v \in V(F)} V_v$ is a finite subset of $V(G)$ and as G is locally finite, there are only finitely many edges in $M(X^{\setminus M/})$ (recall that for each subset $X' \subseteq V(G^{\setminus M/})$, $M(X')$ is the set of crossed edges of M that contract to a vertex in X'). We let $L := M(X^{\setminus M/})$ denote this finite set of edges and note that the sets V_v are also subsets of $V(G^{\setminus L/})$. It follows that $(V_v)_{v \in V(F)}$ is also a model of F in $G^{\setminus \bar{L}/} \llbracket R^{\setminus \bar{L}/} \rrbracket$, a contradiction. \square

Lemma 3.42 (Planar contraction of a single crossed edge). *Let G be locally finite and 3-connected, and \mathcal{T} be a region tangle of order 4 in G . Let (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) be two minimal non-degenerate crossing separations of \mathcal{T} . Let $s_1 s_2$ be the corresponding crossed edge and G' be the graph obtained from G after contracting $s_1 s_2$. Let $(Y'_i, S'_i, Z'_i) := (Y_i, S_i, Z_i)^{\vee}$ be the projection of (Y_i, S_i, Z_i) to G' for each $i \in \{1, 2\}$. Let $R := R_{\mathcal{T}} \subseteq V(G)$ and $R' := R^{\vee}$. If $G' \llbracket R' \rrbracket$ is planar, then so is $G \llbracket R \rrbracket$.*

Proof. We let $H := G \llbracket R \rrbracket$ and $H' := G' \llbracket R' \rrbracket$ and for $i \in \{1, 2\}$, we write $S_i = \{s_i, t_i, r_i\}$ such that $s_1 s_2$ is the crossed edge between (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) . Since (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) are crossing, the edge $s_1 s_2$ belongs to $E(G \llbracket R \rrbracket) \subseteq E(H)$. Note that in particular we have $6 = |S_1 \cup S_2| \leq |R|$.

Claim 3.43. *The neighborhood of s_1 in H is:*

$$N_H(s_1) = \{s_2\} \cup (\text{fc}(S_2) \setminus \{s_1\}),$$

and $\text{fc}(S_2)$ is a triangle in H .

Proof of the Claim: Note that by definition of the torso, the projection $(Y_i \cap R, S_i \cap R, Z_i \cap R)$ of (Y_i, S_i, Z_i) to H is a separation of H . Hence the only possible neighbors of s_1 in H must lie in $(R \cap Z_1 \cap Y_2) \cup \{t_2, r_2\}$ (see Figure 3b).

Note that as G is 3-connected, H must also be 3-connected: this comes from the fact that $|V(H)| \geq 6$ and from the observation that any separator $S \subseteq R = V(H)$ of H is also a separator of G . Thus in particular every vertex of H has degree at least 3.

Then, as $|\{s_2\} \cup (\text{fc}(S_2) \setminus \{s_1\})| = 3$, it is enough to prove that $N_H(s_1) \subseteq \{s_2\} \cup (\text{fc}(S_2) \setminus \{s_1\})$ as equality will be immediately implied as $d_H(s_1) \geq 3$. For this we let $t \in N_H(s_1) \setminus \{s_2\}$. We distinguish two cases:

- If $t \in S_2$, then without loss of generality let $t = t_2$. First note that if t is not an endpoint of some crossed edge then $t \in S_2 \cap \text{fc}(S_2)$ and there is nothing to prove (see Figure 4). Thus we assume that there exists a crossed edge $t_2 s_3$ incident to t_2 for some s_3 and we prove that this case implies a contradiction, which will imply the desired inclusion. As $t_2 \neq s_2$ and $E_{\text{nd}}^\times(\mathcal{T})$ is a matching, we have $s_3 \neq s_1$ and there exists $(Y_3, S_3, Z_3) \in \mathcal{T}_{\text{nd}}$ that crosses (Y_2, S_2, Z_2) via the crossed edge $t_2 s_3$. As (Y_1, S_1, Z_1) and (Y_2, S_2, Z_2) cross, we have $s_1 \in Y_2$. As (Y_2, S_2, Z_2) and (Y_3, S_3, Z_3) cross, we have $t_2 \in Y_3$ and $S_3 \setminus \{s_3\} \subseteq Z_2$. As we assumed that $s_1 t_2 \in E(H)$, $s_1 \neq s_3$ and as $t_2 \in Y_3$, we must have $s_1 \in Y_3 \cup (S_3 \setminus \{s_3\})$. This implies a contradiction as $Y_2 \cap Y_3 = \emptyset$ and $Y_2 \cap (S_3 \setminus \{s_3\}) = \emptyset$.
- If $t \notin S_2$, then we must have: $t \in R \cap Y_2 \cap Z_1$ and by definition of R , as $t \notin \bigcap_{(Y,S,Z) \in \mathcal{T}_{\text{nd}}} Z$, this means that $t \in S_3$ for some $(Y_3, S_3, Z_3) \in \mathcal{T}_{\text{nd}} \setminus \{(Y_1, S_1, Z_1), (Y_2, S_2, Z_2)\}$. By Lemma 3.22, (Y_2, S_2, Z_2) and (Y_3, S_3, Z_3) are either orthogonal or crossing. If we were in the former case, then we should have $S_3 \cap Y_2 = \emptyset$, which is impossible as $t \in S_3 \cap Y_2$. Hence (Y_2, S_2, Z_2) and (Y_3, S_3, Z_3) are crossing, and if s_3 denotes the endpoint of the crossed edge between S_2 and S_3 , as $S_3 \cap Y_2 = \{s_3\}$ we must have $t = s_3$ so we are done as $s_3 \in \text{fc}(S_2) \setminus \{s_1\}$.

The fact that $\text{fc}(S_2)$ forms a clique follows from [Gro16a, Lemma 4.33]. \diamond

Note that by symmetry, Claim 3.43 also implies that we have $N_H(s_2) = \{s_1\} \cup (\text{fc}(S_1) \setminus \{s_2\})$ and that $\text{fc}(S_1)$ is a clique.

Recall that by Wagner's theorem [Wag37], a graph is planar if and only if it is K_5 and $K_{3,3}$ -minor free. Hence, it is enough to prove that if H contains K_5 or $K_{3,3}$ as a minor, then so does H' . We write $\text{fc}(S_i) = \{s_{3-i}, u_i, v_i\}$ for $i \in \{1, 2\}$, and we recall that the vertex of H' resulting from the contraction of s_1 and s_2 is denoted by s' .

Claim 3.44. *If H contains a K_5 -minor, then so does H' .*

Proof of the Claim: Let (V_1, \dots, V_5) be a model of K_5 in H . Let V'_1, \dots, V'_5 be the projection of the sets V_i to H' . If s_1 and s_2 are in the same set V_i , then (V'_1, \dots, V'_5) is also a model of K_5 in H' , so we can assume that the vertices s_1 and s_2 belong to distinct sets V_i , say $s_1 \in V_1$ and $s_2 \in V_2$. As by Claim 3.43, s_1 has degree 3 in H , we have $V_1 \neq \{s_1\}$, so V_1 contains one neighbor of s_1 distinct of s_2 , say u_2 . Since u_2 and v_2 are adjacent in H , the edge $u_2 s'$ in H' has an endpoint in $V'_1 \setminus \{s'\}$ and




FIGURE 4. Left: The graph G when S_2 is incident to exactly 2 crossedges. Here t_2 is part of no crossedge and S_2 and S_3 are crossing via the crossedge r_2s_3 . Hence, $\text{fc}(S_2) = \{s_1, t_2, s_3\}$.

Right: The graph G' obtained after contracting the crossedge s_1s_2 . The dashed edges are edges that appear in H and H' respectively. The situation is identical when S_2 is incident to 3 crossedges, but harder to illustrate in 2 dimensions.

an endpoint in V'_2 . Moreover as $u_2v_2 \in E(H')$, the set $V'_1 \setminus \{s'\}$ is connected in H' . Thus $(V'_1 \setminus \{s'\}, V'_2, V'_3, V'_4, V'_5)$ is a model of K_5 in H' , as desired. \diamond

Claim 3.45. *If H contains a $K_{3,3}$ -minor, then H' contains a $K_{3,3}$ -minor or a K_5 -minor.*

Proof of the Claim: Let (V_1, \dots, V_6) be a model of $K_{3,3}$ in H , such that V_i is adjacent to V_j if i and j have different parities. Let V'_1, \dots, V'_6 be their projection to H' . If s_1 and s_2 are in the same set V_i , then (V'_1, \dots, V'_6) is also a model of $K_{3,3}$ in H' , so we can assume that the vertices s_1 and s_2 belong to distinct sets V_i , say $s_1 \in V_1$ and $s_2 \notin V_1$.

If $u_2 \in V_1$, then the edges $s'u_2$ and u_2v_2 in H' ensure that $V'_1 \setminus \{s'\}$ remains connected and that $(V'_1 \setminus \{s'\}, V'_2, \dots, V'_6)$ is a model of $K_{3,3}$ in H' . Thus we can assume that $u_2 \notin V_1$ and similarly $v_2 \notin V_1$. Since V_1 is connected, we must have $V_1 = \{s_1\}$.

As s_1 has degree three and V_1 is adjacent to V_2, V_4 and V_6 , this implies that s_2, u_2 and v_2 must belong to different sets V_{2i} , say $s_2 \in V_2, u_2 \in V_4$ and $v_2 \in V_6$. By applying the same reasoning as for s_2 , we obtain $V_2 = \{s_2\}, u_1 \in V_3$ and $v_1 \in V_5$. But then $(\{s'\}, V'_3, V'_4, V'_5, V'_6)$ is a model of K_5 in H' . \diamond

This concludes the proof of Lemma 3.42. \square

Proof of Proposition 3.40. Assume that $G^{\setminus M}/\llbracket R^{\setminus M} \rrbracket$ is planar and for every $L \subseteq M$, set $\bar{L} := M \setminus L$. Then, using Lemma 3.42, we can easily prove by induction on $|L| \in \mathbb{N}$ that for any finite set $L \subseteq M$, $G^{\setminus \bar{L}}/\llbracket R_{\mathcal{T}}^{\setminus \bar{L}} \rrbracket$ is planar. In order to be able to use induction, we also need to observe that for the contraction of a single crossedge, the equality $R_{\mathcal{T}'} = R_{\mathcal{T}}^{\vee}$ holds. This is proved in [Gro16a, Section 4.5] and can be deduced from item (2) of Theorem 3.33. We thus conclude by Lemma 3.41 that $G\llbracket R_{\mathcal{T}} \rrbracket$ is also planar. \square

4. THE STRUCTURE OF QUASI-TRANSITIVE GRAPHS AVOIDING A MINOR

4.1. Main results. Our main result in this section is the following more precise version of Theorem 1.1.

Theorem 4.1. *Let G be a locally finite graph excluding K_∞ as a minor and let Γ be a group with a quasi-transitive action on G . Then there is an integer k such that G admits a Γ -canonical tree-decomposition (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, whose torsos $G[V_t]$ either have size at most k or are Γ_t -quasi-transitive 3-connected planar minors of G . Moreover, the edge-separations of (T, \mathcal{V}) are tight.*

Remark 4.2. A natural question is whether we can bound the maximum size k of a finite bag in Theorem 4.1 by a function of the forbidden minor, when G excludes some finite minor instead of the countable clique K_∞ . By taking the free product of the cyclic groups \mathbb{Z}_k and \mathbb{Z} , with their natural sets of generators, we obtain a 4-regular Cayley graph consisting of cycles of length k arranged in a tree-like way. This graph has no K_4 minor, but in any canonical tree-decomposition, each cycle C_k has to be entirely contained in a bag, and thus there is no bound on the size of a bag as a function of the forbidden minor in Theorem 4.1. We can replace \mathbb{Z}_k in this construction by the toroidal grid $\mathbb{Z}_k \times \mathbb{Z}_k$, and obtain a Cayley graph with no K_8 -minor, such that the bags in any (non-necessarily canonical) tree-decomposition of finitely bounded adhesion are arbitrarily large.

We will also prove the following version of Theorem 1.2 at the same time.

Theorem 4.3. *Let G be a locally finite graph excluding K_∞ as a minor and let Γ be a group with a quasi-transitive action on G . Then there is an integer k such that G admits a Γ -canonical tree-decomposition (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, of adhesion at most 3, and whose torsos $G[V_t]$ are Γ_t -quasi-transitive minors of G which are either planar or have treewidth at most k . The edge-separations of (T, \mathcal{V}) are all non-degenerate.*

Remark 4.4. If we carefully consider the proof of Theorem 4.3, we can check that if G has only one end (as in the example of Figure 2), then the tree-decomposition we obtain has adhesion 3 and consists of a star with one infinite bag associated to its central vertex z_0 , and finite bags on its branches. In particular, $G[V_{z_0}]$ cannot have bounded treewidth, as otherwise it would have more than one end, hence it must be planar. Thus, Theorem 4.3 implies that every one-ended locally finite quasi-transitive graph that excludes a minor can be obtained from a one-ended quasi-transitive planar graph by attaching some finite graphs on it along separators of order at most 3.

4.2. Tools. Our proof of Theorems 4.1 and 4.3 mainly consists in an application of Theorem 3.39 together with the following result of Thomassen:

Theorem 4.5 (Theorem 4.1 in [Tho92]). *Let G be a locally finite, quasi-transitive, quasi-4-connected graph G . If G has a thick end, then G is either planar or admits the countable clique K_∞ as a minor.*

A direct consequence of Theorem 4.5 is the following, which will be our base case in what follows.

Corollary 4.6. *Let G be a quasi-transitive, quasi-4-connected, locally finite graph which excludes the countable clique K_∞ as a minor. Then G is planar or has finite treewidth.*

Proof. Assume that G is non-planar. As G is K_∞ -minor free, by Theorem 4.5, all its ends have finite degree. Then by Theorem 3.5, G has finite treewidth. \square

Thomassen proved that if a quasi-transitive graph has only one end, then this end must be thick [Tho92, Proposition 5.6]. We prove the following generalization, which might be of independent interest.

Proposition 4.7. *Let $k \geq 1$ be an integer, and let G be a locally finite quasi-transitive graph. Then G cannot have exactly one end of degree exactly k .*

Proof. Assume without loss of generality that G is connected, since otherwise each component of G is also quasi-transitive locally finite, and we can restrict ourselves to a single component containing an end of degree exactly k . Let Γ be a group acting quasi-transitively on G . Assume that G has an end ω of degree exactly k for some integer $k \geq 1$. As explained in [TW93, Section 4], there exists an infinite sequence of sets S_0, S_1, \dots of size k such that for each $i \geq 0$, S_{i+1} belongs to the component G_i of $G_{i-1} - S_i$ where ω lives (where we set $G_0 := G$), and such that there exist k vertex-disjoint paths $P_{1,i}, \dots, P_{k,i}$ from the k vertices of S_i to the k vertices of S_{i+1} . By concatenating these paths, we obtain k vertex-disjoint rays in G living in ω . As G is connected and locally finite, note that up to extracting a subsequence of $(S_i)_{i \geq 0}$, we may assume that the k paths $P_{1,i}, \dots, P_{k,i}$ are in the same component of $G - (S_i \cup S_{i+1})$. Hence if we set $(Y_i, S_i, Z_i) := (G - (G_i \cup S_i), S_i, G_i)$ for each $i \geq 1$, (Y_i, S_i, Z_i) is a tight separation such that for each $i \geq 1$, $(Y_{i+1}, S_{i+1}, Z_{i+1}) \prec (Y_i, S_i, Z_i)$. Hence by Lemma 3.1, as there are only finitely many Γ -orbits of tight separations of size k , there exist $i < j$ and $g \in \Gamma$ such that $(Y_i, S_i, Z_i) \cdot g = (Y_j, S_j, Z_j)$. Assume without loss of generality that $(i, j) = (0, 1)$. Note that by definition of \prec , the action of g preserves the order \prec , i.e. for each $(Y, S, Z) \prec (Y', S', Z')$, we must have $(Y, S, Z) \cdot g \prec (Y', S', Z') \cdot g$. We now consider the sequence of separations $(Y'_i, S'_i, Z'_i)_{i \geq 0}$ defined for each $i \geq 0$ by: $(Y'_i, S'_i, Z'_i) := (Y_0, S_0, Z_0) \cdot g^i$. Then the sequence $(Y'_i, S'_i, Z'_i)_{i \geq 0}$ is strictly decreasing according to \prec . Recall that there exist k vertex-disjoint paths from S_0 to S_1 that extend to k disjoint rays belonging to ω . Then for each $i \geq 0$, there exist k vertex-disjoint paths from S'_i to S'_{i+1} such that their concatenations consists in k vertex-disjoint rays that belong to some end ω' of degree exactly k (the fact that the end has degree at most k follows from the fact that all the sets S'_i are separators of size k in G). If $\omega' \neq \omega$ then we are done, so we assume that $\omega' = \omega$. Now, observe that the sequence $(Y''_i, S''_i, Z''_i)_{i \geq 0}$ defined for each $i \geq 0$ by $(Y''_i, S''_i, Z''_i) := (Y_0, S_0, Z_0) \cdot g^{-i}$ also satisfies that for each $i \geq 0$, there exists k vertex-disjoint paths $P''_{j,i} := P_{j,0} \cdot g^{-i}$ for $j \in [k]$ from S''_{i+1} to S''_i . If we consider the k vertex-disjoint rays obtained from the concatenation of the paths $P''_{j,k}$, these rays must belong to the same end ω'' as for each i , the paths $P''_{j,k}$ are in the same component of $G - (S''_i \cup S''_{i+1})$. The end ω'' must have degree exactly k as each (Y''_i, S''_i, Z''_i) is a separation of order k . Moreover the sequence $(Y''_i, S''_i, Z''_i)_{i \geq 0}$ is strictly increasing according to \prec , hence ω and ω'' cannot live in the same component of $G - S_0$. Thus we found an end ω'' distinct from ω of degree k . \square

Proposition 4.7 and its proof are reminiscent of Halin's classification of the different types of action an automorphism of a quasi-transitive locally finite graph G can have on the ends of G [Hal73, Theorem 9]. However it is not clear for us whether Proposition 4.7 can be seen as an immediate corollary of Halin's work.

4.3. Proof of Theorems 4.1 and 4.3. Let G be a locally-finite quasi-transitive graph excluding K_∞ as a minor and let Γ be a group inducing a quasi-transitive action on G . Let (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, be a Γ -canonical tree-decomposition of adhesion at most 2 obtained by applying Theorem 3.6 to G . By Lemma 3.13, for each $t \in V_t$, Γ_t acts quasi-transitively on $G_t := G[V_t]$. Moreover, as G_t is a minor of G , it must also exclude K_∞ as a minor.

We let t_1, \dots, t_m be representatives of the orbits of $V(T)/\Gamma$. For each finite torso G_{t_i} of (T, \mathcal{V}) , we define $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ as the trivial tree-decomposition of G_{t_i} (in which the tree \tilde{T}_{t_i} contains a single node). For each infinite, 3-connected torso G_{t_i} of (T, \mathcal{V}) , we let $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ be a Γ_{t_i} -canonical tree-decomposition of G_{t_i} obtained by applying Theorem 3.16 to G_{t_i} , i.e. $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ distinguishes efficiently all the tangles of G_{t_i} of order 4. By Remark 3.17, the edge-separations of $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ in G_{t_i} are all distinct. By Remark 3.20, the edge-separations of $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ in G_{t_i} are non-degenerate. Hence by Lemma 3.19, the torsos of $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$ are minors of G_{t_i} . We now use Corollary 3.12 and find a refinement (T_1, \mathcal{V}_1) of (T, \mathcal{V}) with respect to some family $(T_t, \mathcal{V}_t)_{t \in V(T)}$ of Γ_t -canonical tree-decompositions of G_t such that the construction $t \mapsto (T_t, \mathcal{V}_t)_{t \in V(T)}$ is Γ -canonical and such that for each $i \in I$, $(T_{t_i}, \mathcal{V}_{t_i})$ is a subdivision of $(\tilde{T}_{t_i}, \tilde{\mathcal{V}}_{t_i})$. Since the construction $t \mapsto (T_t, \mathcal{V}_t)_{t \in V(T)}$ is Γ -canonical, for each $t \in V(T_1)$ the decomposition (T_t, \mathcal{V}_t) is Γ_t -canonical and efficiently distinguishes the tangles of order 4 of G_t (by a slight abuse of notation, we keep denoting by G_t the torso of the tree-decomposition (T_1, \mathcal{V}_1) associated to the node $t \in V(T_1)$). Note that by construction, the adhesion sets of (T_1, \mathcal{V}_1) have size at most 3 and all the edge-separations are tight. Moreover, the torsos of each tree-decomposition (T_t, \mathcal{V}_t) are minors of G_t for each $t \in V(T)$, and as the torsos of (T, \mathcal{V}) are minors of G , we also have that the torsos of (T_1, \mathcal{V}_1) are minors of G . In particular, they also exclude K_∞ as a minor. Moreover, by Lemma 3.13, for each $t \in V(T_1)$, Γ_t acts quasi-transitively on G_t . By Lemma 3.1, since all edge-separations of (T_1, \mathcal{V}_1) are tight and have order at most 3, the graph G_t is locally finite for each $t \in V(T_1)$.

Claim 4.8. *For each $t \in V(T_1)$ such that G_t is infinite, G_t is 3-connected and has a unique tangle \mathcal{T}_t of order 4. Moreover \mathcal{T}_t is a Γ_t -invariant region tangle and every end of G_t has degree at least 4.*

Proof of the Claim: Consider a node $t \in V(T_1)$ such that G_t is infinite. As all torsos are cycles, subgraphs of complete graphs of size at most 3, or 3-connected, G_t itself is 3-connected. Since G_t is connected and infinite, it contains some end ω . Let $\mathcal{T}_t := \{(Y, S, Z), |S| \leq 3 \text{ and } \omega \text{ lives in } Z\}$ be defined in G_t . Note that \mathcal{T}_t is a tangle of order 4 in G_t . As G_t is a minor of G , by Lemma 3.14 every tangle \mathcal{T}' of order 4 in G_t induces a tangle \mathcal{T} of order 4 in G , and by Remark 3.15 this mapping is injective. Moreover, note that if (Y, S, Z) is an edge-separation of (T_1, \mathcal{V}_1) such that $V_t \subseteq Z \cup S$, then if \mathcal{M} is any faithful model of G_t in G , the projection $(Y', S', Z') := \pi_{\mathcal{M}}(Y, S, Z)$ is such that $Y' = \emptyset$. Thus $(Y', S', Z') \in \mathcal{T}'$, hence $(Y, S, Z) \in \mathcal{T}$. This means that every edge-separation of (T_1, \mathcal{V}_1) is oriented toward t by \mathcal{T} . Hence if G_t admits two distinct tangles $\mathcal{T}'_1, \mathcal{T}'_2$ of order 4, the two associated tangles $\mathcal{T}_1, \mathcal{T}_2$ given by Lemma 3.14 must be distinct and not distinguished by (T_1, \mathcal{V}_1) , a contradiction. This proves the existence and uniqueness of a tangle \mathcal{T}_t of order 4 in G_t .

Note that as Γ_t acts on G_t and \mathcal{T}_t is the unique tangle of order 4 in G_t , the tangle \mathcal{T}_t is Γ_t -invariant (as a family of separations).

We can also observe that if the end ω in G_t has degree at most 3, then by Proposition 4.7, G_t has another end ω' of degree at most 3 and the construction of \mathcal{T}_t using the end ω' instead of ω yields a different tangle of order 4, which contradicts the uniqueness of \mathcal{T}_t . So every end of G_t has degree at least 4.

It remains to prove that \mathcal{T}_t is a region tangle. If not we can find an infinite decreasing sequence of separations of order 3 in G_t , and this sequence defines an end of degree 3 in G_t , which contradicts the fact that every end of G_t has degree at least 4. \diamond

We will need to decompose further the infinite torsos of the tree-decomposition (T_1, \mathcal{V}_1) . Let $t \in V(T_1)$ be such that G_t is infinite, and let \mathcal{T}_t be the region tangle of order 4 in G_t given by Claim 4.8. We let $M_t := E_{\text{nd}}^\times(\mathcal{T}_t)$ denote the set of crossed edges of \mathcal{T}_t , (T'_t, \mathcal{V}'_t) be the Γ_t -canonical tree-decomposition of G_t given by Lemma 3.25, and $z_0 \in V(T'_t)$ be the center of the star T'_t . By Lemmas 3.13 and 3.24, the graph $H := G_t \llbracket V'_{z_0} \rrbracket$ is a Γ_t -quasi-transitive faithful minor of G_t , thus it must also exclude K_∞ as a minor.

Now we observe that Γ_t induces a quasi-transitive group action on $H^{\setminus M_t/}$: for each $w \in V(H^{\setminus M_t/})$ and every $\gamma \in \Gamma_t$, we set:

$$w \cdot \gamma := \begin{cases} s_{u \cdot \gamma, v \cdot \gamma} & \text{if } w = s_{u, v}, \text{ for some } \{u, v\} \in M_t, \text{ and} \\ w \cdot \gamma & \text{otherwise,} \end{cases}$$

where we recall that the notation $s_{u, v}$, for $\{u, v\} \in M_t$, is introduced at the beginning of Subsection 3.6. As M_t is Γ_t -invariant, we easily see that the mapping γ defines a bijection over $V(H^{\setminus M_t/})$. We let the reader check that it gives a graph isomorphism of $H^{\setminus M_t/}$. Note that the number of Γ_t -orbits of $V(H^{\setminus M_t/})$ is at most the number of Γ_t -orbits of $V(H)$, hence it must be finite.

As $H^{\setminus M_t/}$ is a minor of H , it also excludes the countable clique K_∞ as a minor. It follows from Theorem 3.39 that $H^{\setminus M_t/}$ is quasi-4-connected. Hence, by Corollary 4.6, $H^{\setminus M_t/}$ either has finite treewidth or it is planar. It is not hard to observe that the treewidth of H is at most twice the treewidth of $H^{\setminus M_t/}$ so in particular if we are in the first case, H has also bounded treewidth. In the second case, Proposition 3.40 implies that H is also planar. In both cases, we obtain that (T'_t, \mathcal{V}'_t) is a Γ_t -canonical tree-decomposition of G_t with non-degenerate edge-separations, adhesion 3 and where each torso is a minor of G_t and has either bounded treewidth or is planar. Eventually we can use Proposition 3.10 together with Lemma 3.9 as we did before to find a tree-decomposition (T^*, \mathcal{V}^*) of G with the properties of Theorem 4.3.

We now explain how to derive Theorem 4.1: every torso $G \llbracket V_t \rrbracket$ of (T^*, \mathcal{V}^*) which is neither finite nor planar must have bounded treewidth, hence by Theorem 3.5 it must admit a Γ_t -canonical tree-decomposition where each torso has bounded width. Exactly as before we can apply Corollary 3.12 to find a refinement of (T^*, \mathcal{V}^*) with the properties of Theorem 4.1. \square

5. APPLICATIONS

5.1. The Hadwiger number of quasi-transitive graphs. We say that a graph H is *singly-crossing* if H can be embedded in the plane with a single edge-crossing. It was observed by Paul Seymour that Theorem 4.3 bears striking similarities with a structure theorem of Robertson and Seymour [RS93] related to the exclusion of a singly-crossing graph as a minor. Their theorem states that if H is singly-crossing,

then there is a constant k_H such that any graph excluding H as a minor has a tree-decomposition with adhesion at most 3 in which all torsos are planar or have treewidth at most k_H . On the other hand, for any integer k there is a finite singly-crossing graph H_k such that any graph with a tree-decomposition with adhesion at most 3 in which all torsos are planar or have treewidth at most k must exclude H_k as a minor (this can be seen by taking H_k to be a 4-connected triangulation of a sufficiently large grid, and adding an edge between two non-adjacent vertices lying on incident faces). Using this observation, the following strengthening of Theorem 1.3 is now an immediate consequence of Theorem 4.3.

Theorem 5.1. *For every locally finite quasi-transitive graph G avoiding the countable clique K_∞ as a minor, there is a finite singly-crossing graph H such that G is H -minor-free. In particular there is an integer k such that G is K_k -minor-free.*

Note that in this application we have not used explicitly the property that the underlying tree-decomposition was canonical, but it is used implicitly in the sense that this is what guarantees that the treewidth of the torsos is uniformly bounded in Theorem 4.3.

5.2. Accessibility in quasi-transitive graphs avoiding a minor. We now prove Theorem 1.4, which we restate here for convenience.

Theorem 5.2. *Every locally finite quasi-transitive graph avoiding the countable clique K_∞ as a minor is vertex-accessible.*

Proof. Let G be a locally finite graph avoiding the countable clique K_∞ as a minor, with a group Γ acting quasi-transitively on G . Let (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, be a Γ -canonical tree-decomposition of G of adhesion at most 3 obtained by applying Theorem 4.3 to G . In particular all torsos are quasi-transitive minors of G , and all non-planar torsos have bounded treewidth. Observe that any two ends living in different parts of the tree-decomposition are separated by the separator of size at most 3 of an edge-separation of the tree-decomposition. Consider any node $t \in V(T)$. If $G[V_t]$ is planar then since it is quasi-transitive (by Lemma 3.13) and locally finite, $G[V_t]$ is vertex-accessible [Dun07, Theorem 3.8] and thus there is an integer k_t such that all pairs of ends lying in $G[V_t]$ can be separated by a set of at most k_t vertices. If $G[V_t]$ has bounded treewidth then by Theorem 3.5 there is a integer k_t such that all ends of $G[V_t]$ have degree at most k_t , and thus all pairs of ends lying in $G[V_t]$ can be separated by a set of at most k_t vertices. As $V(T)/\Gamma$ is finite; there is only a finite number of possible values for the integers k_t , $t \in V(T)$, and thus their maximum k is well-defined. We have proved that every pair of ends in G can be separated by a set of at most $\max\{k, 3\}$ vertices, which concludes the proof. \square

5.3. Finite presentability of minor-excluded groups. A *walk* in a graph G is a finite sequence of vertices $W = (v_1, \dots, v_k)$ where for each $i \in [k-1]$, $v_i = v_{i+1}$ or $v_i v_{i+1} \in E(G)$. We call W a *closed walk* when $v_1 = v_k$ or $v_1 v_k \in E(G)$. We let $\mathcal{W}(G)$ denote the set of closed walks of G . If W is a closed walk that contains a *spur*, i.e. if there is some i such that $v_{i-1} = v_{i+1}$, then we say that $W' = (v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k)$ is obtained from W by *deleting the spur*. The inverse operation of *adding a spur* consists in adding a neighbor of v between v_i and v_{i+1} in the walk W if $v_i = v_{i+1} = v$. Similarly, by *deleting a repetition* we mean replacing W by $(v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k)$ if $v_i = v_{i+1}$, and the inverse operation of *adding a repetition* consists in replacing

W by $(v_1, \dots, v_{i-1}, v_i, v_i, v_{i+1}, \dots, v_k)$, for some $1 \leq i \leq k$. The *rotation* of W is the walk $(v_2, v_3, \dots, v_k, v_1)$, and the *reflection* of W is the walk $(v_k, v_{k-1}, \dots, v_2, v_1)$.

If $W = (v_1, \dots, v_k)$ and $W' = (v'_1, \dots, v'_\ell)$ are two walks such that $v_k = v'_1$, then their *sum* is the walk $W \cdot W' := (v_1, \dots, v_k = v'_1, \dots, v'_\ell)$. We will say that a set of closed walks \mathcal{W} *generates* another set of closed walks \mathcal{W}' if every element of \mathcal{W}' can be obtained from elements of \mathcal{W} by adding and deleting spurs and repetitions, and performing sums, reflections and rotations.

The following result was proved in [Ham18b, Theorem 5.12] when G is a quasi-transitive locally finite planar graph. We reuse some of the arguments of the proof of [Ham18b, Proposition 5.9] and combine them with our structure theorem to extend the result to graphs excluding the countable clique K_∞ as a minor.

Theorem 5.3. *Let G be a locally finite graph excluding the countable clique K_∞ as a minor and let Γ be a group acting quasi-transitively on G . Then the set of closed walks of G admits a Γ -invariant generating set with finitely many Γ -orbits.*

Proof. We consider a Γ -canonical tree-decomposition (T, \mathcal{V}) , with $\mathcal{V} = (V_t)_{t \in V(T)}$, given by Theorem 4.1. We let A denote the set of pairs $\{x, y\}$ of vertices of G for which there exists an edge-separation (Y, S, Z) of (T, \mathcal{V}) such that $x, y \in S$ and $xy \notin E(G)$. By Remark 3.3, as the edge-separations associated to (T, \mathcal{V}) are tight, $E(T)/\Gamma$ is finite. As (T, \mathcal{V}) has finitely bounded adhesion, this implies that there is a finite number of Γ -orbits of A . We let $\{x_1, y_1\}, \dots, \{x_\ell, y_\ell\}$ be representatives of these orbits. For each $j \in [\ell]$ we let P_j be a path from x_j to y_j (which always exists, since the edge-separations are tight). For each $\{x, y\} \in A$, we consider the representative $\{x_j, y_j\}$ in the Γ -orbit of $\{x, y\}$, and we define $f(x, y)$ as the image of the path P_j under an automorphism that maps $\{x_j, y_j\}$ to $\{x, y\}$. Note that $f(x, y)$ is an (x, y) -path in G .

We let G^+ be the graph obtained from G by adding all possible edges xy such that $xy \in E(G[V_t])$ for some $t \in V(T)$. In other words the edge-set of G^+ is exactly $E(G) \uplus A$. For each walk W in G^+ , we define the walk $f(W)$ in G as the walk obtained from W by replacing every edge (x, y) of W such that $\{x, y\} \in A$ by $f(x, y)$ (this definition extends the definition of f above, which applied to walks (x, y) of length 1 in G^+). For each set of walks $\mathcal{S} \subseteq \mathcal{W}(G^+)$, we let $f(\mathcal{S}) := \{f(W) \in \mathcal{W}(G), W \in \mathcal{S}\}$.

Claim 5.4. *For every $W \in \mathcal{W}(G^+)$, if $W_1, \dots, W_k \in \mathcal{W}(G^+)$ generate W in $\mathcal{W}(G^+)$, then $f(W)$ is generated by $f(W_1), \dots, f(W_k)$ in $\mathcal{W}(G)$.*

Proof of the Claim: Let $W \in \mathcal{W}(G^+)$ be generated by $W_1, \dots, W_k \in \mathcal{W}(G^+)$. We prove by induction on the number of operations needed to generate W from W_1, \dots, W_k that $f(W)$ is generated by the closed walks $f(W_1), \dots, f(W_k)$.

If $W = W_i$ for some $i \in [k]$, then the result is immediate. Assume that W is obtained from some closed walk W' after performing a rotation on W' , and that W' is generated by W_1, \dots, W_k . We write $W = (v_1, \dots, v_r)$. If $v_1 = v_2$ or $v_1 v_2 \in E(G)$, then $f(W)$ is obtained after performing a single rotation on $f(W')$. If $v_1 v_2 \in E(G^+) \setminus E(G) = A$, then $f(W)$ is obtained after performing $|f(v_1, v_2)|$ rotations to $f(W')$. In any case if we assume by the induction hypothesis that $f(W')$ is generated by $f(W_1), \dots, f(W_k)$, we are immediately done. The case where W is obtained after performing a reflection on W' or adding/removing a repetition on a walk W' is even simpler.

Now assume that W is the concatenation of two walks $W', W'' \in \mathcal{W}(G^+)$ for which the induction hypothesis holds. Then we observe by definition of f that $f(W) = f(W') \cdot f(W'')$. Then again we conclude by the induction hypothesis that $f(W)$ is generated by $f(W_1), \dots, f(W_k)$.

Assume now that $W = (v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_r)$ is obtained from $W' = (v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_r)$ after adding the spur v between v_{i-1} and v_{i+1} with $v_{i-1} = v_{i+1}$. We let $x := v_{i-1}$ and $y := v_i$ and distinguish two cases:

- If $xy \in E(G)$, then we observe that by definition of f , $f(W)$ is obtained from $f(W')$ after adding the same spur so we are done using the induction hypothesis on W' .
- If $xy \in A$, then $f(W)$ must be of the form $U_1 \cdot f(x, y) \cdot f(x, y)^{-1} \cdot U_2$, where $U_1 \cdot U_2 = f(W')$. This means that $f(W)$ can be obtained from $f(W') = U_1 \cdot U_2$ by adding $|f(x, y)|$ spurs, hence the induction hypothesis on W' implies that $f(W)$ is generated by $f(W_1), \dots, f(W_k)$.

Finally assume that $W = (v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_r)$ is obtained from $W' = (v_1, \dots, v_{i-1}, v_i, v_{i+1}, \dots, v_r)$ after deleting the spur v_i . Again we let $x := v_{i-1} = v_{i+1}$ and $y := v_i$ and distinguish two cases:

- If $xy \in E(G)$, then as above, $f(W)$ is obtained from $f(W')$ after the removal of a spur and we are immediately done by applying the induction hypothesis on W' .
- If $xy \in A$, then we claim that $f(W)$ is generated by $f(W')$ as $f(W) = U_1 \cdot U_2$, with $U_1 := f((v_1, \dots, x))$ and $U_2 := f((x, \dots, v_r))$, and $f(W') = U_1 \cdot f(x, y) \cdot f(x, y)^{-1} \cdot U_2$. This shows that $f(W)$ is obtained from $f(W')$ after deleting $|f(x, y)|$ spurs and we can conclude by the induction hypothesis applied to W' that $f(W)$ is generated by $f(W_1), \dots, f(W_k)$.

This concludes the proof of Claim 5.4. \diamond

Claim 5.5. $\mathcal{W}(G)$ is generated by $\bigcup_{t \in V(T)} f(\mathcal{W}(G[V_t]))$.

Proof of the Claim: Let $W \in \mathcal{W}(G)$. First, note that W can be generated in G^+ by closed walks of $\bigcup_{t \in V(T)} \mathcal{W}(G[V_t])$. This comes from the following observation: fix any edge $t_1 t_2$ in T , with associated separation (Y, S, Z) in G . Then any closed walk W in G can be written as the sum of closed walks in $G^+[Y \cup S]$ and $G^+[S \cup Z]$, followed by the removal of spurs corresponding to the edges of the adhesion S . Thus we proved that $\mathcal{W}(G)$ is generated by $\bigcup_{t \in V(T)} \mathcal{W}(G[V_t])$ in G^+ .

Now observe that as for each $W \in \mathcal{W}(G)$, $f(W) = W$, Claim 5.4 implies that $\mathcal{W}(G)$ is generated by $\bigcup_{t \in V(T)} f(\mathcal{W}(G[V_t]))$ in G . \diamond

As G is locally finite, note that for every pair $\{x, y\} \in A$, there are only finitely many paths of the form $P_j \cdot \gamma$ for some $(j, \gamma) \in [\ell] \times \Gamma$ having x and y as endpoints. For each $\{x, y\} \in A$, we let $\mathcal{P}_{x,y}$ denote the set of all such paths and $\mathcal{C}_{x,y}$ denote the set of all closed walks of the form $P \cdot P'^{-1}$ with $P, P' \in \mathcal{P}_{x,y}$. Then $\mathcal{C}_{x,y}$ is finite for each $\{x, y\} \in A$ and the set

$$\mathcal{C} := \bigcup_{\{x,y\} \in A} \mathcal{C}_{x,y}$$

is a Γ -invariant subset of $\mathcal{W}(G)$ with a finite number of Γ -orbits. We also consider the set of closed walks \mathcal{C}' of $\mathcal{W}(G^+)$ of the form xPy for each $\{x, y\} \in A$ and $P \in \mathcal{P}_{x,y}$. Note that $f(\mathcal{C}') \subseteq \mathcal{C}$.

By Remark 3.3, $V(T)/\Gamma$ is finite. As for every $t \in V(T)$, $G[V_t]$ is either finite or Γ_t -quasi-transitive planar, by [Ham18b, Theorem 25] the set $\mathcal{W}(G[V_t])$ of closed walks of $G[V_t] = G^+[V_t]$ has a generating set of cycles with finitely many Γ_t -orbits. We consider representatives t_1, \dots, t_m of each of the finitely many orbits $V(T)/\Gamma$, and for each $i \in [m]$, we let \mathcal{W}_i be a finite set of closed walks of $G[V_{t_i}] = G^+[V_{t_i}]$ such that $\mathcal{W}_i \cdot \Gamma_{t_i}$ generates $\mathcal{W}(G[V_{t_i}])$.

Claim 5.6. *The set*

$$\left(\bigcup_{i=1}^m f(\mathcal{W}_i) \right) \cdot \Gamma \cup \mathcal{C}$$

generates $\bigcup_{t \in V(T)} f(\mathcal{W}(G[V_t]))$ in $\mathcal{W}(G)$.

Proof of the Claim: First, note that for each $i \in [m]$, Claim 5.4 implies that $f(\mathcal{W}_i \cdot \Gamma_{t_i})$ generates $f(\mathcal{W}(G[V_{t_i}]))$.

We first let $i \in [m]$ and show that $f(\mathcal{W}_i) \cdot \Gamma_{t_i} \cup \mathcal{C}$ generates $f(\mathcal{W}_i \cdot \Gamma_{t_i})$. We let $W_i \in \mathcal{W}_i$ and $\gamma \in \Gamma_{t_i}$. One has to be careful as in general the walks $f(W_i \cdot \gamma)$ and $f(W_i) \cdot \gamma$ are not the same. Nevertheless we show that $f(W_i \cdot \gamma)$ is generated by $f(W_i) \cdot \gamma$ and by the walks of \mathcal{C} , which is enough to conclude. Indeed, it is not hard to see that $f(W_i \cdot \gamma)$ is generated in $\mathcal{W}(G^+)$ by $f(W_i) \cdot \gamma$ and by the walks $xPy \in \mathcal{C}'$ for each pair $\{x, y\} \in A$ of consecutive vertices of W_i , where $P \in \mathcal{P}_{x,y}$. Thus by Claim 5.4, $f(W_i \cdot \gamma) = f(f(W_i \cdot \gamma))$ is generated by $f(W_i) \cdot \gamma = f(f(W_i) \cdot \gamma)$ and by the walks of $f(\mathcal{C}') \subseteq \mathcal{C}$.

To conclude with the proof of the claim we let $t \in V(T)$, and $(i, \gamma) \in [m] \times \Gamma$ be such that $t = t_i \cdot \gamma$. We let $W \in \mathcal{W}(G[V_t])$. Then there exist $\gamma \in \Gamma$ and $W' \in \mathcal{W}(G[V_{t_i}])$ such that $W = W' \cdot \gamma$. The exact same arguments as in previous paragraph also apply to prove that $f(W)$ is generated by $f(W') \cdot \gamma$ together with the walks of \mathcal{C} . As we just proved above that $f(W')$ can be generated by finitely elements from $f(\mathcal{W}_i) \cdot \Gamma_{t_i} \cup \mathcal{C}$, we conclude that $f(W)$ can be generated by finitely elements from

$$(f(\mathcal{W}_i) \cdot \Gamma_{t_i} \cup \mathcal{C}) \cdot \gamma \cup \mathcal{C} \subseteq (f(\mathcal{W}_i) \cdot \Gamma) \cup \mathcal{C},$$

as desired. ◇

Combining Claims 5.5 and 5.6, we obtain that $\mathcal{W}(G)$ is generated by $(\bigcup_{i=1}^m f(\mathcal{W}_i)) \cdot \Gamma \cup \mathcal{C}$ in $\mathcal{W}(G)$. As \mathcal{C} has a finite number of Γ -orbits, this concludes the proof of Theorem 5.3. □

We obtain the following consequence of Theorem 5.3 in the group setting, which is a restatement of Theorem 1.6.

Corollary 5.7. *Let Γ be a finitely generated K_∞ -minor-free group. Then Γ is finitely presented.*

Proof. We let $G := \text{Cay}(\Gamma, S)$ be a locally finite Cayley graph of Γ which is K_∞ -minor-free, and consider the right action of Γ on G . Let \mathcal{W} denote a finite set of representative of the Γ -orbits of the Γ -invariant generating set of closed walks of G obtained by applying Theorem 5.3 to G . Note that for every $W \in \mathcal{W}$, if r_W denotes the sequence of labels of W with respect to S , the set of closed walks of G labeled by r_W is exactly the orbit $W \cdot \Gamma$ of W . Thus it is not hard to check that $\langle S \mid r_W, W \in \mathcal{W} \rangle$ is a finite presentation of Γ . □

Note that Corollary 1.5 can now be deduced either from Corollary 5.7, combined with the result of Dunwoody [Dun85] stating that every finitely presented group is accessible, or from Theorem 1.4, combined with the result of Thomassen and Woess [TW93] stating that groups admitting a vertex-accessible Cayley graphs are accessible.

5.4. The domino problem in minor-excluded groups. In this section we prove the following version of Theorem 1.8.

Theorem 5.8. *Let Γ be a finitely generated group excluding the countable clique K_∞ as a minor. Then the domino problem is undecidable on Γ if and only if*

- Γ is one-ended, or
- Γ has an infinite number of ends and has a one-ended planar subgroup which is finitely generated.

In particular, these situations correspond exactly to the cases where Γ is not virtually free.

We will need the following results on one-ended planar groups. One-ended planar groups are exactly infinite *planar discontinuous groups*, i.e. groups that act properly discontinuously on simplicial complexes homeomorphic to \mathbb{R}^2 where every nontrivial automorphism fixes neither a face nor maps an edge (u, v) to its inverse (v, u) (see [ZVC80, Section 4] for a complete survey on these groups).

Theorem 5.9 ([BN46, Fox52]). *Every planar discontinuous group contains the fundamental group of a closed orientable surface as a subgroup of finite index.*

It follows that every one-ended planar group contains the fundamental group of a closed orientable surface of genus $g \geq 1$ as a subgroup (if $g = 0$ the fundamental group is trivial and cannot be a subgroup of finite index). We will use Theorem 5.9 in combination with the following result of Aubrun et al. [ABM19] on the domino problem in surface groups (note that the case $g = 1$ corresponds to the undecidability of the domino problem in \mathbb{Z}^2).

Theorem 5.10 ([ABM19]). *For any $g \geq 1$, the domino problem in the fundamental group of the closed orientable surface of genus g is undecidable.*

For groups with infinitely many ends we will use the existence of finite graphs of groups (which follows from the accessibility of the groups we consider). The following result of Babai [Bab77] will be crucial.

Theorem 5.11 ([Bab77]). *If Γ' is a finitely generated subgroup of Γ , then there exists some finite generating set S' of Γ' such that $\text{Cay}(\Gamma', S')$ is a minor of $\text{Cay}(\Gamma, S)$.*

Finally we will need the following result of Thomassen, which is tightly related to Theorem 4.5, and which will allow us to reduce the K_∞ -minor-free case to the planar case.

Theorem 5.12 (Theorem 5.7 in [Tho92]). *Let G be a locally finite, transitive, non-planar, one-ended graph. Then G contains the countable clique K_∞ as a minor.*

We are now ready to prove Theorem 5.8.

Proof. Assume that Γ has a Cayley graph $G = \text{Cay}(\Gamma, S)$ excluding K_∞ as a minor. If Γ has 0 end, then it is finite and the domino problem is decidable. If Γ has 2

ends then Γ is virtually \mathbb{Z} and the domino problem is also decidable. If Γ is one-ended, then since G is transitive and excludes the countable clique K_∞ as a minor, it follows from Theorem 5.12 that G is (one-ended) planar. By Theorem 5.9, Γ contains the fundamental group Γ' of a closed surface of genus $g \geq 1$ as a subgroup. By Theorem 5.10, the domino problem is undecidable for Γ' . It was proved that for every finitely generated subgroup Γ' of a finitely generated group Γ , if the domino problem is undecidable for Γ' then it is also undecidable for Γ [ABJ18, Proposition 9.3.30]. This implies that if Γ is one-ended, the domino problem is undecidable for Γ .

Assume now that Γ has an infinite number of ends. By Corollary 1.5, Γ is accessible and by Theorem 2.5, if Γ is not virtually free then one of the (finitely presented) vertex-groups Γ_u (which is a subgroup of Γ) in its associated finite graph of groups is one-ended. By Theorem 5.11, Γ_u must also have a locally finite Cayley graph that excludes K_∞ as a minor. By the paragraph above, the domino problem is undecidable for Γ_u , and since Γ_u is a subgroup of Γ , the domino problem is also undecidable for Γ . \square

6. OPEN PROBLEMS

The first problem we consider is related to Conjecture 1.7, which states that a group has a decidable domino problem if and only if it is virtually free (or equivalently, it has bounded treewidth). Using the planar case of Theorem 5.8, this conjecture would be a direct consequence of a positive answer to the following problem.

Problem 6.1. *Is it true that every finitely generated group which is not virtually free has a finitely generated one-ended planar subgroup?*

It was pointed out to us by Agelos Georgakopoulos (personal communication) that the lamplighter group $L = \mathbb{Z}_2 \wr \mathbb{Z}$ provides a negative answer to Problem 6.1. This follows from a characterization of all subgroups of L by Grigorchuk and Kravchenko [GK14], which implies that every finitely generated subgroup of L is either finite or isomorphic to another lamplighter group (and the latter cannot be planar).

A related problem on the graph theory side is the following.

Problem 6.2. *Is it true that every locally finite quasi-transitive graph of unbounded treewidth contains a quasi-transitive planar graph of unbounded treewidth as a subgraph?*

It could be the case that Cayley graphs of lamplighter groups also provide a negative answer to Problem 6.2, but this is not clear (the assumption of being quasi-transitive and subgraph containment are both much weaker than being a Cayley graph and subgroup containment).

A crucial component of our result is the strategy of Grohe [Gro16a] to obtain a canonical tree-decomposition up to the contraction of a matching (see also Theorem 3.39). At one point we tried to figure out whether the following was true (in the end it turned out that we did not need to answer these questions, but we believe they might be of independent interest).

Problem 6.3. *Let G be a locally finite quasi-transitive graph. Is there a proper coloring of G with a finite number of colors such that the colored graph G itself is quasi-transitive (where automorphisms have to preserve the colors of the vertices)?*

Problem 6.4. *Let G be a locally finite quasi-transitive graph. Is there an orientation of the edges of G such that the oriented graph G itself is quasi-transitive (where automorphisms have to preserve the orientation of the edges)?*

Note that Problem 6.4 has a positive answer for Cayley graphs whose generating set does not contain any element of order 2. An example showing that Problem 6.3 has a negative answer was recently constructed by Hamann and Möller. It was then observed by Abrishami, Esperet and Giocanti, and independently by Norin and Przytycki, that a variant of this example could be used to provide a negative answer to Problem 6.4 as well.

Finally, we started the introduction of the paper by mentioning the following natural question: do graphs avoiding a fixed minor have a tree-decomposition in the spirit of the Graph Minor Structure Theorem of Robertson and Seymour [RS03], but with the additional constraint that the decomposition is canonical? We gave a positive answer to this question for quasi-transitive graphs but the general case is still open (although Example 3.4 and Remark 4.2 give natural limitations to the properties of such a canonical tree-decomposition).

Acknowledgement. The authors would like to thank Marthe Bonamy and Vincent Delecroix for their involvement in the early stages of the project and all the subsequent discussions. The second author would like to thank Reinhard Diestel, Raphael Jacobs, Paul Knappe and Matthias Hamann for the helpful discussions and for pointing out a simplification of our initial proof of Theorem 5.3. We also thank Johannes Carmesin and Jan Kurkofka for sharing their recent results on canonical decompositions of 3-connected graphs. Finally, we would like to express our gratitude to the reviewers of the journal version of the paper for their careful reading and the large number of corrections and suggestions. The quality of the paper benefited greatly from their feedback.

REFERENCES

- [ABJ18] Nathalie Aubrun, Sebastián Barbieri, and Emmanuel Jeandel. About the Domino Problem for Subshifts on Groups. In V. Berthé and M. Rigo, editors, *Sequences, Groups, and Number Theory*, Trends in Mathematics, pages 331–389. Birkhäuser, Cham, 2018.
- [ABM19] Nathalie Aubrun, Sebastián Barbieri, and Etienne Moutot. The domino problem is undecidable on surface groups. In Peter Rossmanith, Pinar Heggernes, and Joost-Pieter Katoen, editors, *44th International Symposium on Mathematical Foundations of Computer Science, MFCS 2019, August 26-30, 2019, Aachen, Germany*, volume 138 of *LIPICs*, pages 46:1–46:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
- [Ant11] Yago Antolín. On Cayley graphs of virtually free groups. *Groups Complexity Cryptology*, 3(2):301–327, 2011.
- [Bab77] László Babai. Some applications of graph contractions. *Journal of Graph Theory*, 1(2):125–130, 1977.
- [Ber66] Robert L. Berger. The undecidability of the domino problem. *Memoirs of the American Mathematical Society*, 1966.
- [BN46] Svend Bundgaard and Jakob Nielsen. Forenklede beviser for nogle sætninger i fladetopologien. *Matematisk Tidsskrift. B*, pages 1–16, 1946.

- [BS18] Alexis Ballier and Maya Stein. The domino problem on groups of polynomial growth. *Groups, Geometry, and Dynamics*, 12(1):93–105, 2018.
- [CHM22] Johannes Carmesin, Matthias Hamann, and Babak Miraftab. Canonical trees of tree-decompositions. *Journal of Combinatorial Theory, Series B*, 152:1–26, 2022.
- [CK23] Johannes Carmesin and Jan Kurkofka. Canonical decompositions of 3-connected graphs. In *64th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2023, Santa Cruz, CA, USA, November 6-9, 2023*, pages 1887–1920. IEEE, 2023.
- [DD89] Warren Dicks and Martin J. Dunwoody. *Groups Acting on Graphs*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1989.
- [DHL18] Reinhard Diestel, Fabian Hundertmark, and Sahar Lemanczyk. Profiles of separations: in graphs, matroids, and beyond. *Combinatorica*, 39, 11 2018.
- [Die17] Reinhard Diestel. *Graph Theory: 5th edition*. Springer Graduate Texts in Mathematics. Springer-Verlag, 2017.
- [DJM93] Reinhard Diestel, H. A. Jung, and Rögnvaldur G. Möller. On vertex transitive graphs of infinite degree. *Archiv der Mathematik*, 60:591–600, 1993.
- [Dro06] Carl Droms. Infinite-ended groups with planar Cayley graphs. *Journal of Group Theory*, 9(4):487–496, 2006.
- [DSS98] Carl Droms, Brigitte Servatius, and Herman Servatius. The structure of locally finite two-connected graphs. *Electronic Journal of Combinatorics*, 2, 01 1998.
- [DT99] Reinhard Diestel and Robin Thomas. Excluding a countable clique. *Journal of Combinatorial Theory, Series B*, 76(1):41–67, 1999.
- [Dun85] Martin J. Dunwoody. The accessibility of finitely presented groups. *Inventiones Mathematicae*, 81:449–458, 1985.
- [Dun93] Martin J. Dunwoody. *An Inaccessible Group*, volume 1 of *London Mathematical Society Lecture Note Series*, page 75–78. Cambridge University Press, 1993.
- [Dun07] Martin J. Dunwoody. Planar graphs and covers. *arXiv preprint arXiv:0708.0920*, 2007.
- [Fox52] Ralph H. Fox. On Fenchel’s Conjecture about F-Groups. *Matematisk Tidsskrift. B*, pages 61–65, 1952.
- [Fre44] Hans Freudenthal. Über die Enden diskreter Räume und Gruppen. *Commentarii Mathematici Helvetici*, 17:1–38, 1944.
- [GK14] Rostislav Grigorchuk and Rostyslav Kravchenko. On the lattice of subgroups of the lamplighter group. *International Journal of Algebra and Computation*, 24(6):837–877, 2014.
- [Gro16a] Martin Grohe. Quasi-4-connected components. In Ioannis Chatzigiannakis, Michael Mitzenmacher, Yuval Rabani, and Davide Sangiorgi, editors, *43rd International Colloquium on Automata, Languages, and Programming, ICALP 2016, July 11-15, 2016, Rome, Italy*, volume 55 of *LIPICs*, pages 8:1–8:13. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016.
- [Gro16b] Martin Grohe. Tangles and connectivity in graphs. In Adrian-Horia Dediu, Jan Janousek, Carlos Martín-Vide, and Bianca Truthe, editors, *Language and Automata Theory and Applications - 10th International Conference, LATA 2016, Prague, Czech Republic, March 14-18, 2016, Proceedings*, volume 9618 of *Lecture Notes in Computer Science*, pages 24–41. Springer, 2016.
- [Hal65] R. Halin. Über die maximalzahl fremder unendlicher wege in graphen. *Mathematische Nachrichten*, 30(1-2):63–85, 1965.
- [Hal73] Rudolf Halin. Automorphisms and endomorphisms of infinite locally finite graphs. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 39:251–283, 1973.
- [Ham18a] Matthias Hamann. Accessibility in transitive graphs. *Combinatorica*, 38(4):847–859, 2018.
- [Ham18b] Matthias Hamann. Planar transitive graphs. *Electron. J. Comb.*, 25(4):research paper p4.8, 18, 2018.
- [Ham25] Matthias Hamann. Quasi-transitive K_∞ -minor free graphs. *European Journal of Combinatorics*, 124:104056, 2025.

- [HLMR22] Matthias Hamann, Florian Lehner, Babak Miraftab, and Tim Rühmann. A Stallings type theorem for quasi-transitive graphs. *Journal of Combinatorial Theory, Series B*, 157:40–69, 2022.
- [Hop44] Heinz Hopf. Enden offener Räume und unendliche diskontinuierliche Gruppen. *Commentarii Mathematici Helvetici*, 16:81–100, 1943/44.
- [KPS73] Abraham Karrass, Alfred Pietrowski, and Donald Solitar. Finite and infinite cyclic extensions of free groups. *Journal of the Australian Mathematical Society*, 16(4):458–466, 1973.
- [MS83] David E. Muller and Paul E. Schupp. Groups, the theory of ends, and context-free languages. *Journal of Computer and System Sciences*, 26(3):295–310, 1983.
- [RS91] Neil Robertson and Paul Seymour. Graph minors. X. obstructions to tree-decomposition. *Journal of Combinatorial Theory, Series B*, 52(2):153–190, 1991.
- [RS93] Neil Robertson and Paul Seymour. Excluding a graph with one crossing. In *Graph structure theory*, pages 669–675. American Mathematical Society, 1993.
- [RS03] Neil Robertson and Paul Seymour. Graph minors. XVI. Excluding a non-planar graph. *Journal of Combinatorial Theory, Series B*, 89(1):43–76, 2003.
- [Ser80] Jean-Pierre Serre. *Trees*. Springer-Verlag, Berlin, 1980. Translated from the French by John Stillwell.
- [Sta72] John Stallings. *Group theory and three-dimensional manifolds*. Yale University Press, 1972.
- [Tho92] Carsten Thomassen. The Hadwiger number of infinite vertex-transitive graphs. *Combinatorica*, 12:481–491, 1992.
- [Tut84] William T. Tutte. *Graph Theory*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1984.
- [TW93] Carsten Thomassen and Wolfgang Woess. Vertex-transitive graphs and accessibility. *Journal of Combinatorial Theory, Series B*, 58(2):248–268, 1993.
- [Wag37] Klaus Wagner. Über eine eigenschaft der ebenen komplexe. *Mathematische Annalen*, 114(1):570–590, 1937.
- [Woe89] Wolfgang Woess. Graphs and groups with tree-like properties. *Journal of Combinatorial Theory, Series B*, 47(3):361–371, 1989.
- [ZVC80] Heiner Zieschang, Elmar Vogt, and Hans-Dieter Coldewey. *Surfaces and planar discontinuous groups. Revised and expanded transl. from the German by J. Stillwell*, volume 835 of *Lect. Notes Math*. Springer, Cham, 1980.

(L. Esperet) UNIV. GRENOBLE ALPES, CNRS, LABORATOIRE G-SCOP, GRENOBLE, FRANCE
Email address: louis.esperet@grenoble-inp.fr

(U. Giocanti) UNIV. GRENOBLE ALPES, CNRS, LABORATOIRE G-SCOP, GRENOBLE, FRANCE
Email address: ugo.giocanti@grenoble-inp.fr

(C. Legrand-Duchesne) CNRS, LABRI, UNIVERSITÉ DE BORDEAUX, BORDEAUX, FRANCE
Email address: clement.legrand@u-bordeaux.fr