# Construction of determinants for the six-vertex model with domain wall boundary conditions

Mikhail D. Minin, Andrei G. Pronko, and Vitaly O. Tarasov

Abstract. We consider the problem of construction of determinant formulas for the partition function of the six-vertex model with domain wall boundary conditions. In pioneering works of Korepin and Izergin a determinant formula was proposed and proved using a recursion relation. In later works, another determinant formulas were given by Kostov for the rational case and by Foda and Wheeler for the trigonometric case. Here, we develop an approach in which the recursion relation is replaced by a system of algebraic equations with respect to one set of spectral parameters. We prove that this system has a unique solution. The result can be easily given as a determinant parametrized by an arbitrary basis of polynomials. In particular, the choice of the basis of Lagrange polynomials with respect to the second set of spectral parameters leads to the Izergin–Korepin representation, and the choice of the monomial basis leads to the Kostov and Foda–Wheeler representations.

## CONTENTS



Key words and phrases. Bethe ansatz, vertex models, domain wall boundary conditions, determinant representations.

## 1. Introduction

<span id="page-1-0"></span>The six-vertex model with domain wall boundary conditions (DWBC) was introduced by Korepin in [\[1\]](#page-16-2) for calculation of norms of Bethe states in the framework of the quantum inverse scattering method (QISM) [\[2,](#page-16-3) [3\]](#page-16-4). Korepin showed that the partition function can be uniquely determined by a list of conditions, and among those a key role is played by a certain recursion relation. A milestone result was later obtained by Izergin in [\[4\]](#page-17-0) where he obtained a determinant formula satisfying those conditions. A detailed exposition can be found in [\[5\]](#page-17-1).

In more recent studies, it was found that the six-vertex model with DWBC and their generalizations, the so-called partial DWBC (pDWBC), appear in the route of calculation of structure constants (three-point functions) in  $\mathcal{N}=4$  supersymmetric Yang–Mill theory [\[6–](#page-17-2)[10\]](#page-17-3). In the context of these studies, and on the grounds of Slavnov's determinant for the scalar product of Bethe states [\[11\]](#page-17-4), Kostov in [\[12,](#page-17-5) [13\]](#page-17-6) obtained a determinant formula for the partition function of the six-vertex model with pDWBC for the case of rational Boltzmann weights. The Kostov's result appeared to be intriguing, since when pDWBC are specified to DWBC it remains different from the Izergin–Korepin representation.

Further progress was made by Foda and Wheeler in [\[14\]](#page-17-7), who argued that the six-vertex model with pDWBC in the case of the rational weights can be treated starting from the one with DWBC, by sending several lattice parameters to infinity. In this procedure, Izergin–Korepin representation leads to a determinant formula which is different from the Kostov's one, but it turns out possible to prove that these two representations are equivalent to each other. Foda and Wheeler have also provided a trigonometric generalization of the Kostov's formula, albeit for a severe restriction on the Boltzmann weights, which must be chosen with broken arrow-reversal symmetry and obeying the stochasticity condition. As we explain in the sequel, for DWBC these restrictions can be lifted and the formula can be used for generic weights.

In the present paper, we propose a modified version of the original approach of Korepin and Izergin, where we replace the Korepin's recursion relation by a system of equations with respect to one (out of two) sets of spectral parameters. We prove that this system of equations, supplemented by the requirements of symmetry and polynomiality, has a unique solution that equals the partition function of the six-vertex model with DWBC, see Propositions  $2-5$ . On the other hand, the solution can be easily written down as a determinant parametrized by an arbitrary basis of polynomials. For clarity, we treat the rational and trigonometric cases separately; our main results are stated in Theorem [1](#page-11-1) (see Sect. [4.2\)](#page-11-0) for the rational case, and in Theorem [2](#page-15-1) (see Sect. [5.2\)](#page-15-0), for the trigonometric case. The proofs of the uniqueness statements, Propositions [3](#page-10-0) and [5,](#page-14-0) follow the idea used in [\[15\]](#page-17-8).

Choosing particular bases of polynomials, we recover previously known representations for the partition functions. For instance, we obtain the Izergin–Korepin representation by taking the basis of Lagrange interpolating polynomials with respect to the second set of spectral parameters. On the other hand, picking up the basis of monomials leads to the Kostov and Foda–Wheeler representations.

Although we restrict ourselves here only to treating the six-vertex model with DWBC, we believe that it can be extended to pDWBC as well. Indeed, at least for the rational case this can be done straightforwardly, while the trigonometric case looks subtle and requires more thorough analysis. It is useful to mention that for



<span id="page-2-1"></span>FIGURE 1. Allowed vertex configurations of the six-vertex model and their Boltzmann weights

the trigonometric case of the model with pDWBC in the limit where the lattice domain become a semi-infinite strip shape, the partition function can be given in terms of a Pfaffian [\[16\]](#page-17-9). Another directions to be explored concern more general fixed boundary conditions and calculation of correlation functions in these models [\[17](#page-17-10)[–21\]](#page-17-11).

We organize our paper as follows. In the next section, we recall definition of the six-vertex model with DWBC and list known determinant representations for the partition function. In Sect. [3,](#page-4-0) we formulate the model within the QISM formalism and provide various important ingredients which fix defining properties of the partition function. In Sect. [4](#page-9-0) we study the partition function in the rational case; our main result is the determinant representation which depends on an arbitrary basis of polynomials. In Sect. [5](#page-13-0) we give a generalization of this result to the trigonometric case.

## 2. The six-vertex model with DWBC

<span id="page-2-0"></span>The six-vertex model or the ice-type model, originally is defined in terms of arrows placed on edges of a square lattice. On the vertical edges the arrows point up or down and on the horizontal edges the arrows point left or right. Vertex configurations must contain equal numbers of incoming and outgoing arrows around each lattice vertex. The six allowed vertex configurations, in the standard order (following the conventions adopted in [\[22\]](#page-17-12)), are given in Fig. [1.](#page-2-1)

Domain wall boundary conditions (DWBC) can be defined for the six-vertex model on a finite domain obtained by intersection of N horizontal and N vertical lines, the so-called  $N \times N$  lattice. The DWBC mean that all arrows on the top and bottom boundaries are incoming, on the left and right boundaries all arrows are outgoing, see Fig. [2.](#page-3-0)

To use the quantum inverse scattering method (QISM) in our study (for a detailed exposition of the QISM, see, e.g., [\[3\]](#page-16-4)), we consider the inhomogeneous version of the model in which the Boltzmann weights are taken to be site-dependent in the following way. We label vertical lines from right to left and horizontal lines from top to bottom. We assign variables  $\lambda_j$ ,  $j = 1, \ldots, N$ , to the vertical lines and variables  $\nu_k, k = 1, \ldots, N$ , to the horizontal lines. The Boltzmann weight of the vertex of the ith type lying at the intersection of the jth vertical and kth horizontal lines depends only on the variables corresponding to these lines:  $w_i = w_i(\lambda_i, \nu_k)$ .

The partition function, usually denoted as  $Z_N$ , is defined as follows:

$$
Z_N = \sum_{\mathcal{C} \in \Omega} \prod_{j,k=1}^N w_{i(\mathcal{C},j,k)}(\lambda_j, \nu_k).
$$



<span id="page-3-0"></span>FIGURE 2. An  $N \times N$  lattice with DWBC: all arrows on the top and bottom boundaries are incoming and those on the left and right boundaries are outgoing

Here,  $\Omega$  denotes the set of all allowed configurations obeying the DWBC, and  $i(\mathcal{C}, j, k)$  is the type of the vertex in the configuration C at the intersection of the jth vertical and kth horizontal lines.

In this paper, we consider the Boltzmann weights invariant under the reversal of all arrows,

$$
w_1 = w_2 =: a
$$
,  $w_3 = w_4 =: b$ ,  $w_5 = w_6 =: c$ .

In the rational case, the Boltzmann weights are

<span id="page-3-3"></span><span id="page-3-2"></span>
$$
a(\lambda, \nu) = \lambda - \nu + 1, \qquad b(\lambda, \nu) = \lambda - \nu, \qquad c = 1 \tag{2.1}
$$

and in the trigonometric case are

$$
a(\lambda, \nu) = \sin \gamma(\lambda - \nu + 1), \qquad b(\lambda, \nu) = \sin \gamma(\lambda - \nu), \qquad c = \sin \gamma. \tag{2.2}
$$

An explicit formula for the partition function via the determinant of an  $N \times N$ matrix was derived in [\[5\]](#page-17-1) for both the rational and trigonometric cases,

$$
Z_N = (-1)^{\frac{N(N-1)}{2}} \frac{\prod_{j,k=1}^N a(\lambda_j, \nu_k) b(\lambda_j, \nu_k)}{\nu_N(\{\lambda\}) \nu_N(\{\nu\})} \times \det \left[ \frac{c}{a(\lambda_j, \nu_k) b(\lambda_j, \nu_k)} \right]_{j,k=1,\dots,N} . \quad (2.3)
$$

Here,  $\{\lambda\}$  and  $\{\nu\}$  stand for ordered sets  $\{\lambda\} \equiv \lambda_1, \ldots, \lambda_N$  and  $\{\nu\} \equiv \nu_1, \ldots, \nu_N$ , and the factors  $v_N(\{\lambda\})$  and  $v_N(\{\nu\})$  denote the "Vandermonde determinants" over these sets, namely, in the rational case

<span id="page-3-5"></span><span id="page-3-4"></span><span id="page-3-1"></span>
$$
v_N(\{\lambda\}) = \prod_{1 \le j < k \le N} (\lambda_k - \lambda_j),\tag{2.4}
$$

and in the trigonometric case

$$
v_N(\{\lambda\}) = \prod_{1 \le j < k \le N} \sin \gamma (\lambda_k - \lambda_j). \tag{2.5}
$$

Note that the factor  $(-1)^{\frac{N(N-1)}{2}}$  in [\(2.3\)](#page-3-1) can be canceled out by reversing the order of the double product in one of the two Vandermonde factors in the denominator.

In papers [\[12,](#page-17-5) [13\]](#page-17-6), Kostov gave another determinant formula for the partition function in the rational case (see also [\[14\]](#page-17-7)):

<span id="page-4-2"></span>
$$
Z_N = \frac{\prod_{j,k=1}^N a(\lambda_j, \nu_k)}{v_N(\{\lambda\})} \det \left[ \lambda_j^{k-1} - (\lambda_j + 1)^{k-1} \prod_{l=1}^N \frac{b(\lambda_j, \nu_l)}{a(\lambda_j, \nu_l)} \right]_{j,k=1,...,N} .
$$
 (2.6)

A trigonometric version of formula [\(2.6\)](#page-4-2) can be extracted from the result of Foda and Wheeler [\[14\]](#page-17-7) who proposed a determinant formula for the six-vertex model with pDWBC with asymmetric weights restricted by the stochastisity condition (see formula (4.19) therein and also discussion in [\[23\]](#page-17-13)). For DWBC however, one can ignore these restrictions on weights because the numbers  $n_i$  of vertices of types  $i = 1, \ldots, 6$  in each configuration are related by  $n_1 = n_2, n_3 = n_4$ , and  $n_5 + N = n_6$ , hence the asymmetry can be removed. The Foda–Wheeler representation specialized to DWBC and recast in our notation yields the following formula for the partition function:

<span id="page-4-3"></span>
$$
Z_N = \frac{\prod_{j,k=1}^N a(\lambda_j, \nu_k)}{(2\mathbf{i})^{N(N-1)/2} v_N(\{\lambda\})} \prod_{j=1}^N e^{-(N-2)\mathbf{i}\gamma \lambda_j - \mathbf{i}\gamma \nu_j}
$$
  
 
$$
\times \det \left[ e^{2\mathbf{i}\gamma \lambda_j(k-1)} \left( 1 - e^{\mathbf{i}\gamma(2k-N)} \prod_{l=1}^N \frac{b(\lambda_j, \nu_l)}{a(\lambda_j, \nu_l)} \right) \right]_{j,k=1,\dots,N} . \quad (2.7)
$$

It seems that this formula is new in the context of the DWBC, and we stress again that it is valid for arbitrary weights. Furthermore, we also establish an analogous formula

$$
Z_N = \frac{\prod_{j,k=1}^N a(\lambda_j, \nu_k)}{(2\mathbf{i})^{N(N-1)/2} v_N(\{\lambda\})} \prod_{j=1}^N e^{-N\mathbf{i}\gamma\lambda_j + \mathbf{i}\gamma\nu_j}
$$
  
 
$$
\times \det \left[ e^{2\mathbf{i}\gamma\lambda_j(k-1)} \left( 1 - e^{\mathbf{i}\gamma(2k-2-N)} \prod_{l=1}^N \frac{b(\lambda_j, \nu_l)}{a(\lambda_j, \nu_l)} \right) \right]_{j,k=1,\dots,N}.
$$
 (2.8)

Below we derive general determinant representations of the partition function in both rational and trigonometric cases. The expressions above will arise under specializations of the polynomial bases entering our representations.

# <span id="page-4-4"></span>3. QISM and a basis of the Bethe vectors

<span id="page-4-0"></span>In this section, we introduce standard objects of QISM and formulate a proposition concerning some useful basis of the (off-shell) Bethe vectors. We also show that there exists a family of null Bethe vectors, which provide a system of equations for the partition function.

<span id="page-4-1"></span>3.1. QISM formulation. We start with the consideration of the vector space  $\mathbb{C}^2$  with a standard basis

$$
|\!\!\uparrow\rangle=\begin{pmatrix}1\\0\end{pmatrix},\qquad |\!\!\downarrow\rangle=\begin{pmatrix}0\\1\end{pmatrix}
$$

that we denote as spin-up and spin-down states, respectively. The states on the edges correspond to the spin states as follows:

$$
\uparrow, \rightarrow \Leftrightarrow |\uparrow\rangle \text{ or } \langle \uparrow|,
$$
  

$$
\downarrow, \leftarrow \Leftrightarrow |\downarrow\rangle \text{ or } \langle \downarrow|,
$$

where the bra-vectors are corresponding row vectors

<span id="page-5-0"></span>
$$
\langle \uparrow | = \begin{pmatrix} 1 & 0 \end{pmatrix}, \qquad \langle \downarrow | = \begin{pmatrix} 0 & 1 \end{pmatrix}.
$$

We associate vector space  $\mathbb{C}^2$  to each vertical and horizontal line of the lattice.

We introduce a quantum L-operator which can be defined as matrix of Boltzmann weights. We associate operator  $L_{ik}(\lambda_i, \nu_k)$  with the vertex being at the intersection of the jth vertical and the kth horizontal lines. This operator acts non-trivially in the direct product of two vector spaces. The first one is the "vertical" space  $V_j = \mathbb{C}^2$ , associated with the *j*th vertical line, and the second one is the "horizontal" space  $\mathcal{H}_k = \mathbb{C}^2$ , associated with the kth horizontal line. We regard the "vertical" space  $V_j$  as auxiliary and the "horizontal" space  $\mathcal{H}_k$  as quantum. Graphically, the L-operator acts from top to bottom and from right to left. In other words, arrow states on the top and right edges of the vertex correspond to the "in" indices of the matrix of the L-operator, and arrow states on the bottom and left edges correspond to the "out" indices.

The L-operator is given by

$$
L_{jk}(\lambda,\nu) = \begin{pmatrix} a(\lambda,\nu)\pi_k^+ + b(\lambda,\nu)\pi_k^- & c\sigma_k^- \\ c\sigma_k^+ & b(\lambda,\nu)\pi_k^+ + a(\lambda,\nu)\pi_k^- \end{pmatrix}_{[\mathcal{V}_j]}.
$$
(3.1)

Here,  $\sigma_k^{\pm}$  and  $\pi_k^{\pm}$  denote basis operators acting in the space  $\mathcal{H}_k$ ,

$$
\sigma^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \sigma^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \pi^+ = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \pi^- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix},
$$

and the superscript  $[\mathcal{V}_i]$  in [\(3.1\)](#page-5-0) indicates that this is a matrix in the space  $\mathcal{V}_i$ .

Now we construct the monodromy matrix  $T_i(\lambda_i)$ , multiplying L-operators along the jth vertical line,

$$
T_j(\lambda_j) = L_{jN}(\lambda_j, \nu_N) \cdots L_{j2}(\lambda_j, \nu_2) L_{j1}(\lambda_j, \nu_1) = \begin{pmatrix} A(\lambda_j) & B(\lambda_j) \\ C(\lambda_j) & D(\lambda_j) \end{pmatrix}_{[\mathcal{V}_j]}.
$$
 (3.2)

Operators  $A(\lambda)$ ,  $B(\lambda)$ ,  $C(\lambda)$  and  $D(\lambda)$  act in the total quantum space  $\mathcal{H} = \otimes_{k=1}^{N} \mathcal{H}_k$ . Note that these operators depend also on the parameters  $\nu_1, \ldots, \nu_N$ , that is  $A(\lambda) =$  $A(\lambda; \nu_1, \ldots, \nu_N)$ , etc. Although this dependence is important, we omit it in the notation.

To express the partition function in the operator form, we note that the boundary conditions on the right boundary correspond to the vacuum-state

$$
|\Uparrow\rangle:=|\Uparrow_N\ldots\Uparrow_1\rangle:=|\Uparrow_N\rangle\otimes\cdots\otimes|\Uparrow_1\rangle,
$$

where  $|\uparrow_k\rangle$  is the basis vector in the space  $\mathcal{H}_k$ ,  $k = 1, \ldots, N$ . The boundary conditions on the left boundary are described by the bra-vector

<span id="page-5-2"></span><span id="page-5-1"></span>
$$
\langle \Downarrow| := \langle \downarrow_N \ldots \downarrow_1|.
$$

The partition function reads:

$$
Z_N = \langle \Downarrow | B(\lambda_N) \cdots B(\lambda_2) B(\lambda_1) | \Uparrow \rangle.
$$
 (3.3)

Notice that the "all spins up" state  $|\Uparrow\rangle$  is annihilated by the operator  $C(\lambda)$ ,

<span id="page-6-3"></span><span id="page-6-2"></span>
$$
C(\lambda)|\Uparrow\rangle = 0.\tag{3.4}
$$

<span id="page-6-5"></span><span id="page-6-1"></span>.

It is an eigenstate of the operators  $A(\lambda)$  and  $D(\lambda)$ ,

$$
A(\lambda)|\Uparrow\rangle = a(\lambda)|\Uparrow\rangle, \qquad D(\lambda)|\Uparrow\rangle = d(\lambda)|\Uparrow\rangle,\tag{3.5}
$$

with the eigenvalues

$$
a(\lambda) = \prod_{j=1}^{N} a(\lambda, \nu_j), \qquad d(\lambda) = \prod_{j=1}^{N} b(\lambda, \nu_j).
$$
 (3.6)

We will use these properties later.

<span id="page-6-0"></span>3.2. The Yang-Baxter algebra. The key relation of QISM is the RLLrelation, which reads

$$
R_{jk}(\lambda,\mu) [L_{jl}(\lambda,\nu) \otimes L_{kl}(\mu,\nu)] = [L_{kl}(\mu,\nu) \otimes L_{jl}(\lambda,\nu)] R_{jk}(\lambda,\mu).
$$

Here  $R_{jk}(\lambda,\mu)$  is the R-matrix that acts non-trivially in the tensor product of two auxiliary spaces  $\mathcal{V}_j \otimes \mathcal{V}_k$  and has the following form as a 4  $\times$  4 matrix,

$$
R_{jk}(\lambda,\mu) = \begin{pmatrix} f(\lambda,\mu) & 0 & 0 & 0 \\ 0 & 1 & g(\lambda,\mu) & 0 \\ 0 & g(\lambda,\mu) & 1 & 0 \\ 0 & 0 & 0 & f(\lambda,\mu) \end{pmatrix}_{[\mathcal{V}_j \otimes \mathcal{V}_k]}
$$

The functions  $f(\lambda, \mu)$  and  $g(\lambda, \mu)$  are given by

$$
f(\lambda, \mu) = \frac{a(\lambda, \mu)}{b(\lambda, \mu)}, \qquad g(\lambda, \mu) = \frac{c}{b(\lambda, \mu)}.
$$

The RLL-relation leads to the same relation for the monodromy matrix, the RTT-relation

$$
R_{jk}(\lambda,\mu) [T_j(\lambda) \otimes T_k(\mu)] = [T_k(\mu) \otimes T_j(\lambda)] R_{jk}(\lambda,\mu). \tag{3.7}
$$

This relation encodes commutation relations between the operators  $A(\lambda)$ ,  $B(\lambda)$ ,  $C(\lambda)$  and  $D(\lambda)$ , called the Yang-Baxter algebra. For our purpose, we will use only some of these commutation relations, namely, the commutation relation of the B-operator with itself,

<span id="page-6-4"></span>
$$
[B(\lambda), B(\mu)] = 0,
$$

and the following commutation relation between the A- and B-operators:

$$
A(\mu)B(\lambda) = f(\lambda, \mu)B(\lambda)A(\mu) + g(\mu, \lambda)B(\mu)A(\lambda).
$$
\n(3.8)

In addition to these relations, we also use below an explicit expression for the eigenvalue  $a(\lambda)$  under the action of the operator  $A(\lambda)$  on the vacuum vector.

<span id="page-7-0"></span>3.3. The quantum determinant. The Yang–Baxter algebra [\(3.7\)](#page-6-1) has a central element, called the quantum determinant [\[24,](#page-17-14) [25\]](#page-17-15). It is defined as follows:

$$
Det_q T(\lambda) := D(\lambda)A(\lambda - 1) - C(\lambda)B(\lambda - 1).
$$
 (3.9)

Another formula for  $Det_q T(\lambda)$  is

$$
Det_q T(\lambda) = A(\lambda)D(\lambda - 1) - B(\lambda)C(\lambda - 1).
$$
\n(3.10)

The fact that the quantum determinant is a central element of the algebra means that it commutes with all the entries of the monodromy matrix,

$$
[\text{Det}_{\mathbf{q}} T(\lambda), X(\mu)] = 0, \qquad X \in \{A, B, C, D\}.
$$

The action of the quantum determinant on the quantum space is proportional to the identity operator. The proportionality coefficient can be computed by acting with the expression  $(3.10)$  on the state  $|\Uparrow\rangle$  using  $(3.4)$  and  $(3.5)$ , that yields

<span id="page-7-4"></span><span id="page-7-3"></span><span id="page-7-2"></span>
$$
Det_q T(\lambda) = a(\lambda)d(\lambda - 1).
$$
 (3.11)

The properties of the quantum determinant follow from the commutation relations encoded in [\(3.7\)](#page-6-1), for a detailed exposition see, e.g., [\[3\]](#page-16-4), Ch. VII.

<span id="page-7-1"></span>3.4. Eigenvectors of the A-operator. Here we consider a construction of the eigenvectors of the operator  $A(\lambda)$ .

<span id="page-7-7"></span>PROPOSITION 1. All the eigenvectors of the operator  $A(\mu)$  are given by the following non-zero Bethe vectors

<span id="page-7-6"></span>
$$
B(\nu_{j_1}-1)B(\nu_{j_2}-1)\cdots B(\nu_{j_k}-1)|\!\!\Uparrow\rangle,
$$

with the eigenvalues

$$
\Lambda(\mu; \nu_{j_1}, \dots, \nu_{j_k}) = a(\mu) \prod_{l=1}^k f(\nu_{j_l} - 1, \mu), \tag{3.12}
$$

where  $k \leq N$ , and  $\nu_{j_l} \neq \nu_{j_m}$  for any  $l, m = 1, \ldots, k, l \neq m$ .

PROOF. Let us consider the Bethe vector  $B(\nu_{j_1} - 1)|\Uparrow\rangle$ . We act on the Bethe vector with  $A(\mu)$  and use commutation relation [\(3.8\)](#page-6-4) to get

$$
A(\mu)B(\nu_{j_1} - 1)|\Uparrow\rangle = a(\mu)f(\nu_{j_1} - 1, \mu)B(\nu_{j_1} - 1)|\Uparrow\rangle
$$
  
+ a(\nu\_{j\_1} - 1)g(\mu, \nu\_{j\_1} - 1)B(\mu)|\Uparrow\rangle.

The second term equals zero because the eigenvalue of the operator  $A(\mu)$ , the function  $a(\mu)$ , has zeros at the points  $\mu = \nu_j - 1$ ,  $j = 1, ..., n$ , see [\(3.6\)](#page-6-5), [\(2.1\)](#page-3-2), [\(2.2\)](#page-3-3). Hence, the Bethe vector  $B(\nu_{j_1} - 1)|\uparrow\rangle$  is an eigenvector of the operator  $A(\mu)$  with the eigenvalue

<span id="page-7-5"></span>
$$
\Lambda(\mu; \nu_{j_1}) = a(\mu) f(\nu_{j_1} - 1, \mu),
$$

unless it is the null vector.

To show that the Bethe vector is non-zero, we consider the quantum determinant [\(3.9\)](#page-7-3). We act with  $\text{Det}_{q} T(\nu_{j_1})$  on the vacuum-state

$$
\begin{aligned} \text{Det}_{\mathbf{q}} \, T(\nu_{j_1})|\Uparrow\rangle &= D(\nu_j) a(\nu_{j_1} - 1)|\Uparrow\rangle - C(\nu_{j_1}) B(\nu_{j_1} - 1)|\Uparrow\rangle \\ &= -C(\nu_{j_1}) B(\nu_{j_1} - 1)|\Uparrow\rangle \end{aligned} \tag{3.13}
$$

and use the fact that  $\mathrm{Det}_{q} T(\nu_{j_1})$  is proportional to the identity operator multiplied by the  $a(\nu_{j_1})d(\nu_{j_1}-1)$ , see [\(3.11\)](#page-7-4). Therefore, the right-hand side of [\(3.13\)](#page-7-5) is not equal to zero, hence the Bethe vector  $B(\nu_{j_1} - 1)|\Uparrow\rangle$  does not vanish as well.

The rest of the proof is by induction on the number of B-operators. It is fairly straightforward and we omit the details.

Since  $\nu_{j_l} \neq \nu_{j_m}$  for  $l, m = 1, ..., N$ , and  $l \neq m$ , we get all  $2^N$  non-zero eigenvectors of the operator  $A(\lambda)$  with different eigenvalues given by [\(3.12\)](#page-7-6).  $\square$ 

<span id="page-8-0"></span>3.5. Null Bethe vectors. An important application of the explicit form for the eigenstates of the operator  $A(\mu)$  is the following fact, playing a conceptual role in what follows.

<span id="page-8-3"></span>LEMMA 1. The vector  $B(\nu_j)B(\nu_j-1)|\Uparrow\rangle$  equals zero for each  $j=1,\ldots,N$ .

PROOF. We proof the assertion by contradiction. We act with the operator  $A(\mu)$  on the vector  $B(\nu_j)B(\nu_j-1)|\Uparrow\rangle$  and use commutation relation [\(3.8\)](#page-6-4) to get

$$
A(\mu)B(\nu_j)B(\nu_j-1)|\Uparrow\rangle = f(\nu_j,\mu)B(\nu_j)A(\mu)B(\nu_j-1)|\Uparrow\rangle
$$
  
- 
$$
g(\nu_j,\mu)B(\mu)A(\nu_j)B(\nu_j-1)|\Uparrow\rangle.
$$

The second term vanishes because the eigenvalue of operator  $A(\nu_i)$  is equal to zero, see [\(3.12\)](#page-7-6). Hence, the vector  $B(\nu_j)B(\nu_j-1)|\Uparrow\rangle$  is either an eigenvector of the operator  $A(\mu)$  with the eigenvalue

$$
\Lambda(\mu; \nu_j, \nu_j - 1) = a(\mu)f(\nu_j, \mu)f(\nu_j - 1, \mu)
$$

or the null vector. By Proposition [1,](#page-7-7) we have already constructed the complete set of eigenvectors of the operator  $A(\mu)$ , and none of eigenvalues [\(3.12\)](#page-7-6) is equal to  $\Lambda(\mu; \nu_j, \nu_j - 1)$ . Therefore, the vector  $B(\nu_j)B(\nu_j - 1)|\Uparrow\rangle$  is equal to zero.

<span id="page-8-1"></span>**3.6. Spin flip operators.** The operators  $B(\nu_j)$ ,  $j = 1, \ldots, N$ , play the role of operators flipping spins for certain vectors. More precisely, we have the following simple but useful property.

<span id="page-8-4"></span>LEMMA 2. The operator  $B(\nu_j)$  flips the jth spin in the vector  $|\uparrow_N \dots \uparrow_{j+1} \uparrow_j \downarrow_{j-1}$  $\ldots \downarrow_1$ , namely

$$
B(\nu_j)|\uparrow_N \ldots \uparrow_{j+1} \uparrow_j \downarrow_{j-1} \ldots \downarrow_1\rangle = a(\nu_j)|\uparrow_N \ldots \uparrow_{j+1} \downarrow_j \downarrow_{j-1} \ldots \downarrow_1\rangle, \tag{3.14}
$$

where  $a(\lambda)$  is the vacuum eigenvalue of the operator  $A(\lambda)$ , given by [\(3.6\)](#page-6-5).

PROOF. We represent the B-operator as the  $(1, 2)$ -entry of the monodromy matrix,  $B(\lambda) = [T(\lambda)]_{1,2}$ . Acting with monodromy matrix  $T(\nu_j)$  using its definition [\(3.2\)](#page-5-1) on the vector  $|\uparrow_N \dots \uparrow_{j+1} \uparrow_j \downarrow_{j-1} \dots \downarrow_1 \rangle$  and picking up the relevant entry, yields

$$
B(\nu_j)|\uparrow_N \dots \uparrow_{j+1} \uparrow_j \downarrow_{j-1} \dots \downarrow_1 \rangle
$$
\n
$$
= \left[ \begin{pmatrix} a(\nu_j, \nu_N)|\uparrow_N \rangle & c|\downarrow_N \rangle \\ 0 & b(\nu_j, \nu_N)|\uparrow_N \rangle \end{pmatrix} \dots \begin{pmatrix} a(\nu_j, \nu_{j+1})|\uparrow_{j+1} \rangle & c|\downarrow_{j+1} \rangle \\ 0 & b(\nu_j, \nu_{j+1})|\uparrow_{j+1} \rangle \end{pmatrix} \times \begin{pmatrix} a(\nu_j, \nu_j)|\uparrow_j \rangle & c|\downarrow_j \rangle \\ 0 & 0 \end{pmatrix} \begin{pmatrix} b(\nu_j, \nu_{j-1})|\downarrow_{j-1} \rangle & 0 \\ c|\uparrow_{j-1} \rangle & a(\nu_j, \nu_{j-1})|\downarrow_{j-1} \rangle \end{pmatrix} \dots
$$
\n
$$
\times \dots \begin{pmatrix} b(\nu_j, \nu_1)|\downarrow_1 \rangle & 0 \\ c|\uparrow_1 \rangle & a(\nu_j, \nu_1)|\downarrow_1 \rangle \end{pmatrix} \right]_{1,2}
$$
\n
$$
= c \prod_{\substack{k=1 \ k \neq j}}^N a(\nu_j, \nu_k)|\uparrow_N \dots \uparrow_{j+1} \downarrow_j \downarrow_{j-1} \dots \downarrow_1 \rangle.
$$

Since  $a(\nu_i, \nu_i) = c$  and taking into account [\(3.6\)](#page-6-5), we get [\(3.14\)](#page-8-2).

<span id="page-8-2"></span>

As a corollary, the operator  $B(\nu_i) \cdots B(\nu_1)$  applied to the vector  $|\!\!\Uparrow\rangle$  flips j spins,

$$
B(\nu_j)\cdots B(\nu_1)|\Uparrow\rangle=\prod_{k=1}^j a(\nu_k)|\Uparrow_N\ldots\Uparrow_{j+1}\downarrow_j\ldots\downarrow_1\rangle.
$$

In the important case  $j = N$ ,

<span id="page-9-3"></span>
$$
B(\nu_N) \cdots B(\nu_1) |\Uparrow\rangle = \prod_{k=1}^N a(\nu_k) |\Downarrow\rangle, \tag{3.15}
$$

that is closely related to the partition function  $Z_N$ , see [\(3.3\)](#page-5-2).

#### 4. Determinant representations in the rational case

<span id="page-9-0"></span>In this section, we state the properties of the partition function  $Z_N(\{\lambda\})$  in the rational case. We prove that the partition function can be written as a determinant involving an arbitrary basis of polynomials. We also show that representations [\(2.3\)](#page-3-1) and [\(2.6\)](#page-4-2) arise under specializations to the Lagrange and monomial bases, respectively.

<span id="page-9-1"></span>4.1. Properties of the partition function. We consider the partition function  $Z_N$  defined by [\(3.3\)](#page-5-2). We will study it as a function of the variables  $\lambda_1, \ldots, \lambda_N$ , considering the variables  $\nu_1, \ldots, \nu_N$  as parameters. As before, we use the notation  $\{\lambda\} \equiv \lambda_1, \ldots, \lambda_N$ . We begin with describing several properties of the function  $Z_N(\{\lambda\})$  and, then we will show that a function with these properties is unique.

<span id="page-9-2"></span>PROPOSITION 2. The partition function  $Z_N(\{\lambda\})$  of the six-vertex model with DWBC in the rational parametrization, as a function of the variables  $\lambda_1, \ldots, \lambda_N$ has the following properties:

- (1) It is a symmetric polynomial in  $\lambda_1, \ldots, \lambda_N$ ;
- (2) The degree in each of the variables  $\lambda_1, \ldots, \lambda_N$  equals  $N-1$ ;
- (3) For each pair of variables, say  $\lambda_1, \lambda_2$ , one has

$$
Z_N(\nu_j, \nu_j - 1, \lambda_3, \dots, \lambda_N) = 0, \qquad j = 1, \dots, N; \tag{4.1}
$$

(4) As  $\{\lambda\} = \{\nu\}$ , the following holds:

<span id="page-9-5"></span><span id="page-9-4"></span>
$$
Z_N(\{\nu\}) = \prod_{j,k=1}^N (\nu_j - \nu_k + 1).
$$
 (4.2)

PROOF. The first and second properties follow from the explicit form of the L-operator, see [\(3.1\)](#page-5-0), and the commutativity of the B-operators. The third property is due to Lemma [1.](#page-8-3) The fourth property follows from relations  $(3.3)$  and  $(3.15)$ .  $\Box$ 

As a remark, notice that one can prove property [\(4.2\)](#page-9-4) without Lemma [2](#page-8-4) using Korepin's reduction relation  $[1]$ . The starting point is the fact that L-operator [\(3.1\)](#page-5-0) with coinciding arguments is just the permutation operator,  $L_{jk}(\nu, \nu) = P_{jk}$ . Consider the vertex at the top right corner of the lattice, see Fig. [2.](#page-3-0) The DWBC mean that at  $\lambda_1 = \nu_1$  only the arrow configuration of type 6 can contribute, see

Fig. [1.](#page-2-1) This observation immediately yields the relation

$$
Z_N(\nu_1, \lambda_2, ..., \lambda_N) = \prod_{k=2}^N (\nu_1 - \nu_k + 1) \prod_{j=2}^N (\lambda_j - \nu_1 + 1) \times Z_{N-1}(\lambda_2, ..., \lambda_N; {\{\nu\}} \setminus \nu_1), \quad (4.3)
$$

where it is indicated in the notation that the partition function  $Z_{N-1}$  depends on the set of parameters  $\{\nu\} \setminus \nu_1 \equiv \nu_2, \ldots, \nu_N$ . Repeating relation [\(4.3\)](#page-10-1) iteratively for the remaining variables by setting subsequently  $\lambda_2 = \nu_2, \lambda_3 = \nu_3, \ldots, \lambda_N = \nu_N$ , one obtains [\(4.2\)](#page-9-4).

Let us now study the system of equations [\(4.1\)](#page-9-5), [\(4.2\)](#page-9-4) and show that it has a unique solution.

<span id="page-10-0"></span>PROPOSITION 3. Let  $P_N(\{\lambda\})$  be a polynomial in the variables  $\lambda_1, \ldots, \lambda_N$ , depending on the parameters  $\{\nu\} = \nu_1, \ldots, \nu_N$ , that has the following properties:

- (1) It is symmetric in  $\lambda_1, \ldots, \lambda_N$ ;
- (2) The degree in each of the variables  $\lambda_1, \ldots, \lambda_N$  equals  $N-1$ ;
- (3) For each pair of variables, say  $\lambda_1, \lambda_2$ , one has

$$
P_N(\nu_j, \nu_j - 1, \lambda_3, \dots, \lambda_N) = 0, \qquad j = 1, \dots, N; \tag{4.4}
$$

(4) As  $\{\lambda\} = \{\nu\}$ , the following holds:

<span id="page-10-4"></span><span id="page-10-2"></span><span id="page-10-1"></span>
$$
P_N(\{\nu\}) = \prod_{j,k=1}^N (\nu_j - \nu_k + 1).
$$

Then,  $P_N(\{\lambda\}) = Z_N(\{\lambda\}).$ 

PROOF. The proof is by induction on N. The base of induction at  $N = 1$  is clear.

For the induction step, expand the polynomial  $P_N(\{\lambda\})$  in the variable  $\lambda_1$  using the Lagrange interpolating polynomials of degree  $N-1$  for the points  $\nu_1, \ldots, \nu_N$ ,

<span id="page-10-3"></span>
$$
P_N(\{\lambda\}) = \sum_{i=1}^N P_N(\nu_i, \lambda_2, \dots, \lambda_N) \prod_{\substack{j=1 \ j \neq i}}^N \frac{\lambda_1 - \nu_j}{\nu_i - \nu_j}.
$$
\n(4.5)

It follows from [\(4.4\)](#page-10-2) and the symmetry of the variables  $\lambda_1, \ldots, \lambda_N$  that

$$
P_N(\nu_i, \lambda_2, \dots, \lambda_N) = S_{N-1,i}(\lambda_2, \dots, \lambda_N) \prod_{k=2}^N (\lambda_k - \nu_i + 1) \prod_{j=1}^N (\nu_i - \nu_j + 1), \tag{4.6}
$$

 $i = 1, \ldots, N$ , for some polynomials  $S_{N-1,i}(\lambda_2, \ldots, \lambda_N)$  in  $\lambda_2, \ldots, \lambda_N$ . The second product in [\(4.6\)](#page-10-3) provides a convenient normalization of the polynomials  $S_{N-1,i}$ .

It is straightforward to verify from the properties  $(1) - (4)$  of  $P_N({\lambda})$  that for each  $i = 1, \ldots, N$ , the polynomial  $S_{N-1,i}(\lambda_2, \ldots, \lambda_N)$  has the following properties:

- (1) It is symmetric in  $\lambda_2, \ldots, \lambda_N$ ;
- (2) The degree in each of the variables  $\lambda_2, \ldots, \lambda_N$  equals  $N-2$ ;
- (3) For each pair of the variables, say  $\lambda_2, \lambda_3$ , one has

$$
S_{N-1,i}(\nu_j, \nu_j-1, \lambda_4, \ldots, \lambda_N) = 0, \qquad j = 1, \ldots, i-1, i+1, \ldots, N;
$$

(4) The following holds:

$$
S_{N-1,i}(\{\nu\} \setminus \nu_i) = \prod_{\substack{j,k=1 \ j,k \neq i}}^N (\nu_j - \nu_k + 1),
$$

where we use the notation  $\{\nu\} \setminus \nu_i := \nu_1, \ldots, \nu_{i-1}, \nu_{i+1}, \ldots, \nu_N$ . Hence by the induction assumption, for each  $i = 1, \ldots, N$ , one has

$$
S_{N-1,i}(\lambda_2,\ldots,\lambda_N)=Z_{N-1}(\lambda_2,\ldots,\lambda_N;\{\nu\}\setminus\nu_i),\qquad(4.7)
$$

indicating in the right-hand side the dependence on the parameters  $\{\nu\}$  explicitly.

Formulas [\(4.5\)](#page-10-4), [\(4.6\)](#page-10-3), [\(4.7\)](#page-11-2) show that  $P_N({\{\lambda\}})$  is uniquely determined by its properties (1)-(4). Therefore,  $P_N(\{\lambda\}) = Z_N(\{\lambda\})$  by Proposition [2.](#page-9-2)

<span id="page-11-0"></span>**4.2. Determinant representation.** Consider polynomials  $p_1(\lambda), \ldots, p_N(\lambda)$ in one variable of degree at most  $N-1$ ,

<span id="page-11-2"></span>N

$$
p_i(\lambda) = \sum_{j=1}^N p_{ij} \lambda^{j-1}, \qquad i = 1, \dots, N.
$$

Set

$$
Q_N(\{p\}) = \det[p_{ij}]_{i,j=1,...,N},
$$

where  $\{p\} = (p_1, \ldots, p_N)$ . The polynomials  $p_1(\lambda), \ldots, p_N(\lambda)$  form a basis of polynomials in  $\lambda$  of degree  $N - 1$  if and only if  $Q_N(\{p\}) \neq 0$ . Notice the equality

$$
\det\left[p_k(\lambda_j)\right]_{j,k=1,\ldots,N} = Q_N(\{p\}) \prod_{1 \leq j < k \leq N} (\lambda_k - \lambda_j),\tag{4.8}
$$

that follows from reading the formula

<span id="page-11-4"></span><span id="page-11-3"></span>
$$
p_k(\lambda_j) = \sum_{i=1}^N p_{ki} \lambda_j^{i-1}
$$

as a product of matrices

$$
[p_k(\lambda_j)]_{j,k=1,\ldots,N} = [p_{ki}]_{i,k=1,\ldots,N} [ \lambda_j^{i-1}]_{i,j=1,\ldots,N},
$$

and the Vandermonde determinant formula.

Our main result in the rational case can be formulated as follows.

<span id="page-11-1"></span>THEOREM 1. The partition function of the six-vertex model with DWBC with the rational weights  $(2.1)$  can be written in the form

$$
Z_N(\{\lambda\}) = \frac{\det\left[p_k(\lambda_j)a(\lambda_j) - p_k(\lambda_j+1)d(\lambda_j)\right]_{j,k=1,\ldots,N}}{\det\left[p_k(\lambda_j)\right]_{j,k=1,\ldots,N}}\tag{4.9}
$$

where  $p_1(\lambda), \ldots, p_N(\lambda)$  form a basis of polynomials in  $\lambda$  of degree  $N-1$ , and  $a(\lambda)$ and  $d(\lambda)$  are the eigenvalues of the operators  $A(\lambda)$  and  $D(\lambda)$ :

<span id="page-11-5"></span>
$$
a(\lambda) = \prod_{k=1}^{N} (\lambda - \nu_k + 1), \qquad d(\lambda) = \prod_{k=1}^{N} (\lambda - \nu_k).
$$
 (4.10)

Notice formula [\(4.8\)](#page-11-3) for the denominator in [\(4.9\)](#page-11-4).

**PROOF.** First observe that the denominator in  $(4.9)$  is a nonzero polynomial since  $p_1(\lambda), \ldots, p_N(\lambda)$  form a basis of polynomials in  $\lambda$  of degree  $N-1$ , see formula  $(4.8).$  $(4.8).$ 

Denote by  $P_N(\{\lambda\})$  the right-hand side of formula [\(4.9\)](#page-11-4). The numerator in  $(4.9)$  is an antisymmetric polynomial in  $\lambda_1, \ldots, \lambda_N$ , and hence, is divisible by the denominator. Therefore,  $P_N(\{\lambda\})$  is a symmetric polynomial in  $\lambda_1, \ldots, \lambda_N$ . To prove the theorem, we will show that  $P_N(\{\lambda\})$  obeys also properties  $(2) - (4)$  described in Proposition [3.](#page-10-0) Then  $P_N({\{\lambda\}}) = Z_N({\{\lambda\}})$  by Proposition 3.

For the second property, observe that both  $a(\lambda)$  and  $d(\lambda)$  are monic polynomials of degree N. Thus for each  $k = 1, ..., N$ , the polynomial  $p_k(\lambda)a(\lambda) - p_k(\lambda+1)d(\lambda)$ has degree at most  $2N-2$ . Therefore, the degree of  $P_N({\lambda})$  in each of the variables  $\lambda_1, \ldots, \lambda_N$  does not exceed  $N-1$ .

The third property follows from the equalities

<span id="page-12-0"></span>
$$
a(\nu_j - 1) = d(\nu_j) = 0, \qquad j = 1, \dots, N,
$$
\n(4.11)

so that after the substitution  $\lambda_1 = \nu_j$ ,  $\lambda_2 = \nu_j - 1$ , the first two rows of the matrix  $[p_k(\lambda_j)a(\lambda_j) - p_k(\lambda_j+1)d(\lambda_j)]_{j,k=1,\ldots,N}$  become proportional,



and its determinant vanishes. Hence,  $P_N(\nu_j, \nu_j - 1, \lambda_3, \dots, \lambda_N) = 0$ .

To verify the fourth property, we take  $\{\lambda\} = \{\nu\}$ . Taking into account the second equality in [\(4.11\)](#page-12-0), we get

$$
\det [p_k(\nu_j)a(\nu_j) - p_k(\nu_j + 1)d(\nu_j)]_{j,k=1,...,N} \n= \det [p_k(\nu_j)a(\nu_j)]_{j,k=1,...,N} = \left(\prod_{j=1}^N a(\nu_j)\right) \det [p_k(\nu_j)]_{j,k=1,...,N}.
$$

Therefore by the first of formulas [\(4.10\)](#page-11-5),

$$
P_N(\{\nu\}) = \prod_{j,k=1}^N (\nu_j - \nu_k + 1),
$$

that completes the proof of Theorem [1.](#page-11-1)

Let us now turn to examples. The first example to consider is the celebrated de-terminant formula [\(2.3\)](#page-3-1). To get it, we choose  $p_1(\lambda), \ldots, p_N(\lambda)$  to be the interpolating polynomials of degree  $N-1$  for the points  $\nu_1, \ldots, \nu_N$ ,

$$
p_k(\lambda) = \prod_{\substack{l=1\\l \neq k}}^N (\lambda - \nu_l), \qquad k = 1, \dots, N.
$$

Since  $p_k(\lambda) = d(\lambda)/(\lambda - \nu_k)$  and  $p_k(\lambda + 1) = a(\lambda)/(\lambda - \nu_k + 1)$ , we have

$$
p_k(\lambda_j)a(\lambda_j) - p_k(\lambda_j+1)d(\lambda_j)
$$
  
= 
$$
\frac{d(\lambda_j)a(\lambda_j)}{\lambda_j - \nu_k} - \frac{a(\lambda_j)d(\lambda_j)}{\lambda_j - \nu_k+1} = \frac{a(\lambda_j)d(\lambda_j)}{(\lambda_j - \nu_k)(\lambda_j - \nu_k+1)},
$$

and for the numerator in formula [\(4.9\)](#page-11-4), we obtain

$$
\det [p_k(\nu_j)a(\nu_j) - p_k(\nu_j+1)d(\nu_j)]_{j,k=1,\dots,N}
$$
  
= 
$$
\left(\prod_{j=1}^N a(\lambda_j)d(\lambda_j)\right) \det \left[\frac{1}{(\lambda_j-\nu_k)(\lambda_j-\nu_k+1)}\right]_{j,k=1,\dots,N}.
$$
 (4.12)

For the denominator in formula [\(4.9\)](#page-11-4), we use first formula [\(4.8\)](#page-11-3) and evaluate the factor  $Q_N({p})$  employing the same formula with  $\{\lambda\} = {\nu}$ ,

$$
Q_N(\{p\}) = \frac{\det [p_k(\nu_j)]_{j,k=1,...,N}}{\prod_{1 \le j < k \le N} (\nu_k - \nu_j)} = \frac{\det \left[ \delta_{kj} \prod_{i=1, i \ne k}^N (\nu_k - \nu_i) \right]_{j,k=1,...,N}}{\prod_{1 \le j < k \le N} (\nu_k - \nu_j)} = \prod_{1 \le j < k \le N} (\nu_j - \nu_k). \tag{4.13}
$$

Taking together formulas  $(2.4)$ ,  $(4.9)$ ,  $(4.12)$ , and  $(4.13)$ , we get formula  $(2.3)$  in the rational case.

The second example we would like to mention is the choice of  $p_1(\lambda), \ldots, p_N(\lambda)$ as monomials,

<span id="page-13-3"></span><span id="page-13-2"></span>
$$
p_k(\lambda) = \lambda^{k-1}, \qquad k = 1, \dots, N,
$$

In this case, representation [\(4.9\)](#page-11-4) clearly reproduces Kostov's determinant formula  $(2.6).$  $(2.6).$ 

### 5. Determinant representation in the trigonometric case

<span id="page-13-0"></span>In this section, we generalize the results of the previous section to the trigonometric case. As before, our main result is the determinant representation for the partition function that depends on an arbitrary basis of polynomials. Particular examples are given by representations [\(2.3\)](#page-3-1) and [\(2.7\)](#page-4-3).

<span id="page-13-1"></span>5.1. Properties of the partition function. As we have observed studying the rational case, an important role is played by a polynomial dependence of the partition function on the inhomogeneity parameters. The polynomial dependence in the trigonometric case can be introduced by the "change of variables"

<span id="page-13-4"></span>
$$
x_j = q^{2\lambda_j}, \qquad y_k = q^{2\nu_k}, \qquad q = e^{i\gamma},
$$

so the Boltzmann weights [\(2.2\)](#page-3-3) in the trigonometric case are

$$
a(\lambda_j, \nu_k) = \frac{qx_j - y_kq^{-1}}{2i(x_jy_k)^{1/2}}, \quad b(\lambda_j, \nu_k) = \frac{x_j - y_k}{2i(x_jy_k)^{1/2}}, \quad c(\lambda_j, \nu_k) = \frac{q - q^{-1}}{2i}.
$$
 (5.1)

Then the partition function of the six-vertex model with DWBC in the trigonometric case has the form

<span id="page-13-5"></span>
$$
Z_N(\{\lambda\};\{\nu\}) = \frac{\tilde{Z}_N(\{x\};\{y\})}{(2i)^{N^2} \prod_{j=1}^N (x_j y_j)^{(N-1)/2}},\tag{5.2}
$$

where  $\widetilde{Z}_N({x}; {y})$  is a polynomial in  ${x} \equiv x_1, \ldots, x_N$  and  ${y} \equiv y_1, \ldots, y_N$ .

Indeed, for every allowed configuration of the six-vertex with DWBC, each horizontal and each vertical line contains an odd number of arrow reversing vertices (the 5th and 6th types, see Fig. [1\)](#page-2-1), that is, of the c-weight vertices. Therefore, the weight of every configuration is a polynomial of the variables  $x_1, \ldots, x_N, y_1, \ldots, y_N$ , divided by the product  $\prod_{j=1}^{N} (x_j y_j)^{(N-1)/2}$ .

We apply the method developed in the rational case to the function  $Z_N$ . We begin with the counterpart of Proposition [2.](#page-9-2)

<span id="page-14-6"></span>PROPOSITION 4. The partition function  $\widetilde{Z}_N({x})$  as a function of the variables  $x_1, \ldots, x_N$  has the following properties:

- (1) It is a symmetric polynomial in  $x_1, \ldots, x_N$ ;
- (2) The degree in each of the variables  $x_1, \ldots, x_N$  equals  $N-1$ ;
- (3) For each pair of variables, say  $x_1, x_2$ , one has

$$
\widetilde{Z}_N(y_j, q^{-2}y_j, x_3, \dots, x_N) = 0, \qquad j = 1, \dots, N;
$$
\n(5.3)

(4) As  $\{x\} = \{y\}$ , the following holds:

<span id="page-14-2"></span><span id="page-14-1"></span>
$$
\widetilde{Z}_N({y}) = (q - q^{-1})^N \prod_{\substack{j,k=1 \ j \neq k}}^N (qy_j - q^{-1}y_k).
$$
\n(5.4)

PROOF. The proof is completely similar to the rational case, see Proposition [2.](#page-9-2)

Now we show that the system of equations [\(5.3\)](#page-14-1), [\(5.4\)](#page-14-2) has a unique solution.

<span id="page-14-0"></span>PROPOSITION 5. Let  $P_N({x})$  be a polynomial in the variables  $x_1, \ldots, x_N$ , depending on the parameters  $\{y\} = y_1, \ldots, y_N$ , that has the following properties:

- (1) It is a symmetric polynomial in  $x_1, \ldots, x_N$ ;
- (2) The degree in each of the variables  $x_1, \ldots, x_N$  equals  $N-1$ ;
- (3) For each pair of variables, say  $x_1, x_2$ , one has

$$
P_N(y_j, q^{-2}y_j, x_3, \dots, x_N) = 0, \qquad j = 1, \dots, N; \tag{5.5}
$$

(4) As  $\{x\} = \{y\}$ , the following holds:

<span id="page-14-3"></span>
$$
P_N({y}) = (q - q^{-1})^N \prod_{\substack{j,k=1 \ j \neq k}}^N (qy_j - q^{-1}y_k).
$$

Then,  $P_N(\lbrace x \rbrace) = \widetilde{Z}_N(\lbrace x \rbrace).$ 

Proof. The proof is the same as that of Proposition [3.](#page-10-0) We reproduce it in the condensed form for clarity.

The proof is by induction on N. The base of induction at  $N = 1$  is clear. For the induction step, we expand the polynomial  $P_N(\lbrace x \rbrace)$  in the variable  $x_1$ 

<span id="page-14-5"></span><span id="page-14-4"></span>
$$
P_N(\{x\}) = \sum_{i=1}^N P_N(y_i, x_2, \dots, x_N) \prod_{\substack{j=1 \ j \neq i}}^N \frac{x_1 - y_j}{y_i - y_j},\tag{5.6}
$$

and use [\(5.5\)](#page-14-3) to get

$$
P_N(y_i, x_2, \dots, x_N) = S_{N-1,i}(x_2, \dots, x_N)
$$
  
 
$$
\times (q - q^{-1}) \prod_{k=2}^N (qx_k - q^{-1}y_i) \prod_{\substack{j=1 \ j \neq i}}^N (qy_i - q^{-1}y_j), \quad (5.7)
$$

 $i = 1, \ldots, N$ , for some polynomials  $S_{N-1,i}(x_2, \ldots, x_N)$  in  $x_2, \ldots, x_N$ .

For each  $i = 1, ..., N$ , the polynomial  $S_{N-1,i}(x_2,...,x_N)$  has the properties:

- (1) It is symmetric in  $x_2, \ldots, x_N$ ;
- (2) The degree in each of the variables  $x_2, \ldots, x_N$  equals  $N-2$ ;
- (3) For each pair of the variables, say  $x_2, x_3$ , one has

$$
S_{N-1,i}(y_j, q^{-2}y_j, x_4, \dots, x_N) = 0, \qquad j = 1, \dots, i-1, i+1, \dots, N;
$$

(4) The following holds:

$$
S_{N-1,i}(\{y\} \setminus y_i) = (q - q^{-1})^{N-1} \prod_{\substack{j,k=1 \ j,k \neq i, \ j \neq k}}^N (qy_j - q^{-1}y_k),
$$

where we use the notation  $\{y\} \setminus y_i := y_1, \ldots, y_{i-1}, y_{i+1}, \ldots, y_N$ . Hence by the induction assumption, for each  $i = 1, \ldots, N$ , one has

<span id="page-15-2"></span>
$$
S_{N-1,i}(x_2,...,x_N) = \widetilde{Z}_{N-1}(x_2,...,x_N;\{y\} \setminus y_i),
$$
\n(5.8)

where we indicate in the right-hand side the dependence on the parameters  $\{y\}$ explicitly.

Formulas [\(5.6\)](#page-14-4), [\(5.7\)](#page-14-5), [\(5.8\)](#page-15-2) show that  $P_N({x})$  is uniquely determined by its properties (1) – (4). Therefore,  $P_N(\lbrace x \rbrace) = \widetilde{Z}_N(\lbrace x \rbrace)$  by Proposition [4.](#page-14-6)

<span id="page-15-0"></span>5.2. Determinant representation. Our main result in the trigonometric case can be formulated as follows.

<span id="page-15-1"></span>THEOREM 2. The partition function of the six-vertex model with DWBC with the trigonometric weights  $(2.2)$  can be written in the form

<span id="page-15-3"></span>
$$
\widetilde{Z}_N(\{x\}) = \frac{\det \left[ p_k(x_j) \widetilde{a}(x_j) - q^{-N+2} p_k(q^2 x_j) \widetilde{d}(x_j) \right]_{j,k=1,...,N}}{\left( \prod_{k=1}^N y_k \right) \det [p_k(x_j)]_{j,k=1,...,N}},\tag{5.9}
$$

and in the form

$$
\widetilde{Z}_N(\{x\}) = \frac{\det \left[ p_k(x_j) \widetilde{a}(x_j) - q^{-N} p_k(q^2 x_j) \widetilde{d}(x_j) \right]_{j,k=1,...,N}}{\left( \prod_{j=1}^N x_j \right) \det [p_k(x_j)]_{j,k=1,...,N}}.
$$
(5.10)

Here  $p_1(x), \ldots, p_N(x)$  form a basis of polynomials in x of degree  $N-1$ , and  $\tilde{a}(x)$ ,  $\tilde{d}(x)$  are

<span id="page-15-4"></span>
$$
\tilde{a}(x) = \prod_{k=1}^{N} (qx - q^{-1}y_k), \qquad \tilde{d}(x) = \prod_{k=1}^{N} (x - y_k).
$$

Notice formula [\(4.8\)](#page-11-3),

$$
\det[p_k(x_j)]_{j,k=1,\dots,N} = Q_N(\{p\}) \prod_{1 \le j < k \le N} (x_k - x_j).
$$

PROOF. The proof is similar to the rational case. We verify that the right-hand sides of formulas [\(5.9\)](#page-15-3) and [\(5.10\)](#page-15-4) obeys properties  $(1) - (4)$  stated in Proposition [4.](#page-14-6) This is done along the same lines as in the proof of Theorem [1.](#page-11-1)  $\Box$ 

Note that the functions  $\tilde{a}(x)$  and  $\tilde{d}(x)$  are related to vacuum eigenvalues of the A- and D-operators,

$$
a(\lambda) = \frac{\tilde{a}(x)}{(2\mathbf{i})^N x^{N/2} \prod_{j=1}^N y_j^{1/2}}, \qquad d(\lambda) = \frac{\tilde{d}(x)}{(2\mathbf{i})^N x^{N/2} \prod_{j=1}^N y_j^{1/2}}, \qquad x = q^{2\lambda}.
$$

Let us now turn to examples. The first example to consider is determinant formula [\(2.3\)](#page-3-1) in the trigonometric case. To get it, we choose  $p_1(x), \ldots, p_N(x)$  to be the interpolating polynomials of degree  $N-1$  for the points  $y_1, \ldots, y_N$ ,

$$
p_k(x) = \prod_{\substack{l=1 \\ l \neq k}}^N (x - y_l), \qquad k = 1, ..., N.
$$

Since  $p_k(x) = \tilde{d}(x)/(x - y_k)$ ,  $p_k(q^2x) = q^{N-1}\tilde{a}(x)/(qx - q^{-1}y_k)$ , we have

$$
p_k(x_j)\tilde{a}(x_j) - q^{-N+2}p_k(q^2x_j)\tilde{d}(x_j)
$$
  
= 
$$
\frac{\tilde{d}(x_j)\tilde{a}(x_j)}{x_j - y_k} - \frac{q\tilde{a}(x_j)\tilde{d}(x_j)}{qx_j - q^{-1}y_k} = \frac{(q - q^{-1})y_k\tilde{a}(x_j)\tilde{d}(x_j)}{(x_j - y_k)(qx_j - q^{-1}y_k)}
$$

and

$$
p_k(x_j)\tilde{a}(x_j) - q^{-N}p_k(q^2x_j)\tilde{d}(x_j)
$$
  
= 
$$
\frac{\tilde{d}(x_j)\tilde{a}(x_j)}{x_j - y_k} - \frac{q^{-1}\tilde{a}(x_j)\tilde{d}(x_j)}{qx_j - q^{-1}y_k} = \frac{(q - q^{-1})x_j\tilde{a}(x_j)\tilde{d}(x_j)}{(x_j - y_k)(qx_j - q^{-1}y_k)}.
$$

Using these formulas to transform the numerators in formulas  $(5.9)$ ,  $(5.10)$ , and formulas [\(4.8\)](#page-11-3), [\(4.13\)](#page-13-3) to deal with the denominators, we get in both cases

$$
\widetilde{Z}_N(\{x\}) = \frac{\prod_{j=1}^N \widetilde{a}(x_j) \widetilde{d}(x_j)}{\prod_{1 \le j < k \le n} (x_k - x_j)(y_j - y_k)} \det \left[ \frac{q - q^{-1}}{(x_j - y_k)(qx_j - q^{-1}y_k)} \right]_{j,k=1,\ldots,N}.
$$

Taking into account formulas  $(2.5)$ ,  $(3.6)$ ,  $(5.1)$ , and  $(5.2)$ , the last equality yields formula [\(2.3\)](#page-3-1) in the trigonometric case.

The second example is the choice of  $p_1(x), \ldots, p_N(x)$  as monomials,

$$
p_k(x) = x^{k-1}, \qquad k = 1, \ldots, N,
$$

In this case, representation [\(5.9\)](#page-15-3) gives formula [\(2.7\)](#page-4-3) and representation [\(5.10\)](#page-15-4) gives formula  $(2.8)$ .

# 6. Acknowledgments

<span id="page-16-0"></span>The first and second authors (M.D.M. and A.G.P.) are supported in part by the BASIS Foundation, grant  $\#21$ -7-1-32-3. The first author (M.D.M.) also acknowledges partial support from the Leonhard Euler International Mathematical Institute, agreement #075–15–2019–1620.

#### <span id="page-16-1"></span>References

- <span id="page-16-2"></span>[1] V. E. Korepin, Calculations of norms of Bethe wave functions, Commun. Math. Phys. 86 (1982), 391–418.
- <span id="page-16-3"></span>[2] L. A. Takhtadjan and L. D. Faddeev, The quantum method of the inverse problem and the Heisenberg XYZ model, Russ. Math. Surveys 34 (1979), no. 5, 11-68.
- <span id="page-16-4"></span>[3] V. E. Korepin, N. M. Bogoliubov, and A. G. Izergin, Quantum inverse scattering method and correlation functions, Cambridge University Press, Cambridge, 1993.
- <span id="page-17-0"></span>[4] A. G. Izergin, Partition function of the six-vertex model in the finite volume, Sov. Phys. Dokl. 32 (1987), 878–879.
- <span id="page-17-1"></span>[5] A. G. Izergin, D. A. Coker, and V. E. Korepin, Determinant formula for the six-vertex model, J. Phys. A 25 (1992), 4315–4334.
- <span id="page-17-2"></span>[6] J. Escobedo, N. Gromov, A. Sever, and P. Viera, Tailoring three-point functions and integrability, JHEP 09 (2011), 028, arXiv:[1012.2475](https://arxiv.org/abs/1012.2475).
- [7] J. Escobedo, N. Gromov, A. Sever, and P. Viera, Tailoring three-point functions and integra-bility II. Weak/strong coupling match, JHEP 09 (2011), 029, arXiv:[1104.5501](https://arxiv.org/abs/1104.5501).
- [8] N. Gromov, A. Sever, and P. Viera, Tailoring three-point functions and integrability III. Classical tunneling, JHEP 07 (2012), 044, arXiv:[1111.2349](https://arxiv.org/abs/1111.2349).
- [9] O. Foda,  $N = 4$  SYM structure constants as determinants, JHEP 03 (2012), 096, arXiv: [1111.4663](https://arxiv.org/abs/1111.4663).
- <span id="page-17-3"></span>[10] Y. Jiang, S. Komatsu, I. Kostov, and D. Serban, The hexagon in the mirror: the three-point function in the SoV representation, J. Phys. A 49 (2016), 174007, arXiv:[1506.09088](https://arxiv.org/abs/1506.09088).
- <span id="page-17-4"></span>[11] N. A. Slavnov, Calculation of scalar products of wave functions and form factors in the framework of the algebraic Bethe ansatz, Theor. Math. Phys. 79 (1989), no. 2, 502–508.
- <span id="page-17-5"></span>[12] I. Kostov, Classical limit of the three-point function of  $N = 4$  supersymmetric Yang-Mills Theory from integrability, Phys. Rev. Lett. 108 (2012), 261604, arXiv:[1203.6180](https://arxiv.org/abs/1203.6180).
- <span id="page-17-6"></span>[13] I. Kostov, Three-point function of semiclassical states at weak coupling, J. Phys. A 45 (2012), 494018, arXiv:[1205.4412](https://arxiv.org/abs/1205.4412).
- <span id="page-17-7"></span>[14] O. Foda and M. Wheeler, *Partial domain wall partition functions*, JHEP 213 (2012), 186, arXiv:[1205.4400](https://arxiv.org/abs/1205.4400).
- <span id="page-17-8"></span>[15] A. Nakayashiki, S. Pakuliak, and V. Tarasov, On solutions of the KZ and qKZ equations at level zero, Ann. Inst. Henri Poincaré  $71$  (1999), 459–496, arXiv:[q-alg/9712002](https://arxiv.org/abs/q-alg/9712002).
- <span id="page-17-9"></span>[16] A. G. Pronko and G. P. Pronko, Off-shell Bethe states and the six-vertex model, J. Math. Sci. (N.Y.) 242 (2019), 742–752.
- <span id="page-17-10"></span>[17] G. A. P. Ribeiro, General scalar products in the arbitrary six-vertex model, J. Stat. Mech. Theor. Exp. (2011), P11015, arXiv:[1109.3686](https://arxiv.org/abs/1109.3686).
- [18] I. R. Passos and G. A. P. Ribeiro, Boundary correlations for the six-vertex model with reflecting end boundary condition, J. Stat. Mech. Theor. Exp. (2019), 083101, arXiv:[1904.06383](https://arxiv.org/abs/1904.06383).
- [19] M. D. Minin and A. G. Pronko, Boundary polarization of the rational six-vertex model on a semi-infinite lattice, J. Math. Sci. (N.Y.)  $257$  (2021), 537-550.
- [20] M. D. Minin and A. G. Pronko, Boundary one-point function of the rational six-vertex model with partial domain wall boundary conditions: explicit formulas and scaling properties, SIGMA (2021), 111 (29 pp), arXiv:[2108.06190](https://arxiv.org/abs/2108.06190).
- <span id="page-17-11"></span>[21] F. Colomo, G. Di Guilio, and A. G. Pronko, Six-vertex model on a finite lattice: integral representations for nonlocal correlation functions, Nucl. Phys. B 972 (2021), 115535 (42 pp), arXiv:[2107.13358](https://arxiv.org/abs/2107.13358).
- <span id="page-17-12"></span>[22] R. J. Baxter, Exactly solved models in statistical mechanics, Academic Press, San Diego, CA, 1982.
- <span id="page-17-13"></span>[23] P. Bleher and K. Liechty, Six-vertex model with partial domain wall boundary conditions: Ferroelectric phase, J. Math. Phys. 56 (2015), 023302, arXiv:[1407.8483](https://arxiv.org/abs/1407.8483).
- <span id="page-17-14"></span>[24] A. G. Izergin and V. E. Korepin, Lattice model connected with nonlinear Schrödinger equation, Sov. Phys. Dokl. 26 (1981), 653–654.
- <span id="page-17-15"></span>[25] V. E. Korepin, Analysis of bilinear relation for the six-vertex model, Sov. Phys. Dokl. 27 (1982), 612–613.

Email address: mmd@pdmi.ras.ru

Email address: agp@pdmi.ras.ru

Email address: vt@pdmi.ras.ru

Steklov Mathematical Institute, Fontanka 27, St. Petersburg, 191023, Russia