

Determinantal Sieving

Eduard Eiben*

Tomohiro Koana†

Magnus Wahlström‡

April 6, 2023

Abstract

We introduce a new, remarkably powerful tool to the toolbox of algebraic FPT algorithms, *determinantal sieving*. Given a polynomial $P(x_1, \dots, x_n)$ over a field \mathbb{F} of characteristic 2, on a set of variables $X = \{x_1, \dots, x_n\}$, and a linear matroid $M = (X, \mathcal{I})$ over \mathbb{F} of rank k , in 2^k evaluations of P we can sieve for those terms in the monomial expansion of P which are multilinear and whose support is a basis for M . The known tools of *multilinear detection* and *constrained multilinear detection* then correspond to the case where M is a uniform matroid, respectively the truncation of a disjoint union of uniform matroids. More generally, let the *odd support* of a monomial m be the set of variables which have odd degree in m . Using 2^k evaluations of P , we can sieve for those terms m whose odd support spans M . Applying this framework to well-known efficiently computable polynomial families allows us to simplify, generalize and improve on a range of algebraic FPT algorithms, such as:

- Solving q -MATROID INTERSECTION in time $O^*(2^{(q-2)k})$ and q -MATROID PARITY in time $O^*(2^{qk})$, improving on $O^*(4^{qk})$ (Brand and Pratt, ICALP 2021)
- T -CYCLE, COLOURFUL (s, t) -PATH, COLOURFUL (S, T) -LINKAGE in undirected graphs, and the more general RANK k (S, T) -LINKAGE problem over so-called *frameworks* (see Fomin et al., SODA 2023), all in $O^*(2^k)$ time, improving on $O^*(2^{k+|S|})$ respectively $O^*(2^{|S|+O(k^2 \log(k+|\mathbb{F}|))})$
- Many instances of the DIVERSE X paradigm, finding a collection of r solutions to a problem with a minimum mutual distance of d in time $O^*(2^{r^2 d/2})$, improving solutions for k -DISTINCT BRANCHINGS from time $2^{O(k \log k)}$ to $O^*(2^k)$ (Bang-Jensen et al., ESA 2021), and for DIVERSE PERFECT MATCHINGS from $O^*(2^{2^{O(rd)}})$ to $O^*(2^{r^2 d/2})$ (Fomin et al., STACS 2021)

For several other problems, such as SET COVER, STEINER TREE, GRAPH MOTIF and SUBGRAPH ISOMORPHISM, where the current algorithms are either believed to be optimal or are proving exceedingly difficult to improve, we show matroid-based generalisations at no increased cost to the running time. All matroids are assumed to be represented over a field of characteristic 2, and all algorithms use polynomial space. Over general fields, we achieve similar results at the cost of using exponential space by working over the *exterior algebra*. For a class of arithmetic circuits we call *strongly monotone*, this is even achieved without any loss of running time. However, the *odd support* sieving result appears to be specific to working over characteristic 2.

*Dept. of Computer Science, Royal Holloway, University of London, Eduard.Eiben@rhul.ac.uk

†Algorithmics and Computational Complexity, Technische Universität Berlin, tomohiro.koana@tu-berlin.de

‡Dept. of Computer Science, Royal Holloway, University of London, Magnus.Wahlstrom@rhul.ac.uk

Contents

1	Introduction	3
1.1	Determinantal sieving	5
1.1.1	Over general fields	6
1.1.2	Linear matroids	6
1.2	Applications	7
1.2.1	Matroid Covering, Packing and Intersection Problems	7
1.2.2	Fair and Diverse Solutions	9
1.2.3	Undirected paths and linkages	9
1.2.4	Subgraph problems	11
1.2.5	Speeding up dynamic programming	11
2	Preliminaries	13
2.1	Linear matroids	14
2.2	Enumerating polynomials	15
3	Determinantal sieving	16
3.1	Over a field of characteristic 2	16
3.1.1	Sieving for bases	17
3.1.2	Sieving for spanning sets	18
3.2	Over general fields	19
4	Matroid Covering, Packing and Intersection Problems	22
4.1	Rank k Set Cover and Rank k Set Packing.	23
4.2	q -Matroid Parity and q -Matroid Intersection	24
4.3	Odd Coverage	25
5	Balanced Solution and Diverse Collection	26
5.1	Balance-fair X paradigm	26
5.2	Diverse X paradigm	27
6	Path, cycle and linkage problems	29
6.1	The linkage-generating determinant	30
6.1.1	Path enumeration	31
6.1.2	Linkage enumeration	32
6.2	Rank k (S, T) -linkage	33
6.3	Faster Long st -Path and Long Cycle	35
7	Subgraph problems	39
7.1	Finding high-rank connected subgraphs	41
7.2	Independent subgraph isomorphism	44
8	Speeding-up Dynamic Programming	45
8.1	Minimum Equivalent Graph	46
8.2	Eulerian Deletion	47
8.3	Conflict-free Solution	48
9	Conclusions	49
9.1	Open questions	51

1 Introduction

Algebraic algorithms is a non-obvious but remarkably powerful algorithmic paradigm, especially for exact (exponential-time) and parameterized algorithms. To narrow the scope a bit, let us consider more specifically what may be called the *enumerating polynomial* method. Consider a problem of looking for a particular substructure in an object; for example, given a graph G , we may ask if G has a perfect matching, or a path on at least k vertices, etc. (We focus on the decision problem. Given the ability to solve the decision problem, an explicit solution can be found with limited overhead; see Björklund et al. [22, 24] for deeper investigations into this.) For surprisingly many applications, this problem can be reduced to polynomial identity testing: construct a multivariate polynomial $P(X) = P(x_1, \dots, x_n)$ (occasionally referred to as *multivariate generating polynomial* [26]) such that the monomials of P enumerate all instances of the substructure you are looking for, and then test whether $P(X)$ contains at least one monomial or not. The latter is the *polynomial identity testing* (PIT) problem, which can be solved efficiently in randomized polynomial time via the Schwartz-Zippel (a.k.a. DeMillo-Lipton-Schwartz-Zippel) Lemma: given the ability to evaluate $P(X)$, possibly over an extension field of the original field, we can test if it is non-zero [89, 96]. Therefore the challenge lies in constructing an enumerating polynomial that can be sufficiently efficiently evaluated. In particular, it is a priori non-obvious why it would be easier to construct an enumerating polynomial for a problem than to simply solve the problem directly.

In our experience, this ability has two sources. First, there are well-known families of polynomials that can be efficiently evaluated (despite having exponentially many monomial terms) and which can be usefully interpreted combinatorially as enumerating polynomials for certain objects. For example, if G is a bipartite graph with vertex partition $U \cup V$, the *Edmonds matrix* of G over some field \mathbb{F} is a matrix $A \in \mathbb{F}^{U \times V}$ constructed by replacing the non-zero entries of the bipartite adjacency matrix of G by distinct new variables – i.e., for every edge $e \in E(G)$ we define a variable x_e , and we let $A(u, v) = x_{uv}$ if $uv \in E(G)$ and $A(u, v) = 0$ otherwise. If $|U| = |V|$, then $P(X) = \det A$ is a polynomial over the variables $X = \{x_e \mid e \in E(G)\}$, and can easily be seen to be an enumerating polynomial for perfect matchings in G . (Note that we pay no attention to the coefficients of the monomials, which are here either 1 or -1 depending on the matching; in particular, we are not concerned with *counting* the objects.) For our second example, let G be a digraph, and let A be its standard adjacency matrix, modified as above so that non-zero entries $A(u, v)$ are replaced by $x_{(u,v)}$ for distinct new variables $x_{(u,v)}$, $(u, v) \in E(G)$. Then $A^k(u, v)$ enumerates k -edge walks from u to v . Further examples include the Tutte matrix, which provides a way to enumerate perfect matchings in non-bipartite graphs [91]; *branching walks* (due to Nederlof [82]), which are a relaxation of subtrees of a graph similar to how walks are a relaxation of paths; and any number of applications of basic linear algebra, which become especially relevant in the context of *linear matroids* (see below).

Second, there is a rich toolbox of *transformations* of polynomials, by which a given enumerating polynomial can be modified into a more relevant form. We are particularly concerned with what can be called *sieving* operations: transformations applied to a given polynomial $P(X)$ such that every monomial m in the monomial expansion of P either survives (possibly multiplied by some new factor) or is cancelled, depending on the properties of m . For example, consider a graph G with edges partitioned as $E(G) = E_R \cup E_B$ into *red* and *blue* edges. Does G have a perfect matching where precisely half the edges are red (or more generally, with precisely w red edges)? This is known as the EXACT MATCHING problem, and is not known to have a deterministic polynomial-time algorithm. However, there is a simple randomized polynomial-time algorithm using the enumerating polynomial approach (cf. Mulmuley et al. in 1987 [80]). Assume for simplicity that G is bipartite, and let A be the Edmonds matrix of G as above (if G is not bipartite, instead let A be the

Tutte matrix, and replace all mentions of the determinant of A by the *Pfaffian* of A). Introduce a new variable z , and for every edge $uv \in E_R$ multiply $A(u,v)$ by z . Now, a perfect matching M of G with w red edges will correspond to a monomial where the degree of z is w , and we are left asking for monomials in $P(X, z) = \det A$ where the z -component is z^w . Via the standard method of *interpolation*, we can define a second polynomial $P_2(X)$ which enumerates precisely these monomials, and $P_2(X)$ can be evaluated using $O(n)$ evaluations of $P(X)$ (i.e., P_2 *sieves* for monomials in $P(X, z)$ where z has degree w). Thus, applying polynomial identity testing to P_2 gives a randomized polynomial-time algorithm for EXACT MATCHING.

For applications to exact and parameterized algorithms, more powerful transformations are available. The most well known is *multilinear detection*: Given a polynomial $P(X)$, does the monomial expansion of P contain a monomial of degree k which is *multilinear*, i.e., where every variable has degree at most one? Slightly more generally, to avoid undesired cancellations, we consider the following. Let $P(X, Y)$ be a polynomial in two sets of variables X and Y . Say that a monomial m is *k-multilinear in Y* if the total degree of m in Y is k (not counting any contributions from X) and every variable in Y has degree at most one in m . Then the following is known.

Lemma 1.1 (Multilinear detection [16, 19]). *Let $P(X, Y)$ be a polynomial over a field of characteristic 2. There is a polynomial $Q(X, Y)$, that can be computed using $O^*(2^k)$ evaluations of P , such that Q is not identically zero if and only if P contains a monomial that is k -multilinear in Y .*

For example, consider again the case where A is the modified adjacency matrix of a graph G , and scale every entry $A(u, v)$ by a new variable y_v . Then the terms of $A^k(u, v)$ that are multilinear in $Y = \{y_v \mid v \in V\}$ enumerate k -edge *paths* from u to v , i.e., multilinear detection and a PIT algorithm solve the k -PATH problem. This idea was pioneered by Koutis [65] and improved by Williams and Koutis [66, 93], using a different approach based on group algebra; the above polynomial sieving result is by Björklund et al. [16, 19]. Brand et al. [27] proposed yet another approach using *exterior algebra* to develop deterministic algorithms. Multilinear detection and other algebraic sieving has had many applications, including Björklund’s celebrated algorithm for finding undirected Hamiltonian cycles in time $O^*(1.66^n)$ [16] and an algorithm solving k -PATH in time $O^*(1.66^k)$ [19]. See Koutis and Williams [67] for an overview.

Some variations are also known. One arguably simpler variant is when $|Y| = k$ and we wish to sieve for monomials in $P(X, Y)$ where every variable of Y occurs (regardless of their degree). This can be handled over any field in 2^k evaluations of P using *inclusion-exclusion* (and this is a “clean” sieve, that does not change the coefficient of any monomial). This has been used, e.g., in parameterized algorithms for LIST COLOURING [56] and RURAL POSTMAN [58]. Another variant, which is a generalisation of multilinear detection, is *constrained multilinear detection*. Let $P(X, Y)$ be a polynomial. Let C be a set of colours, and for every $q \in C$ let $d_q \in \mathbb{N}$ be the *capacity* of colour q . Let a colouring $c: Y \rightarrow C$ be given. A monomial m is *properly coloured* if, for every $q \in C$, the total degree of all variables in m with colour q is at most d_q . Björklund et al. [23] show the following (again, we allow an additional set of variables X to avoid undesired cancellations).

Lemma 1.2 (Constrained multilinear detection [23]). *Let $P(X, Y)$ be a polynomial over a field of characteristic 2. Let a colouring $c: Y \rightarrow C$ and a list of colour capacities $(d_q)_{q \in C}$ be given. There is a polynomial $Q(X, Y)$, that can be computed using $O^*(2^k)$ evaluations of P , such that Q is not identically zero if and only if P contains a monomial that is k -multilinear in Y and properly coloured.*

Using this method, Björklund et al. [23] solve GRAPH MOTIF and associated optimization variants in time $O^*(2^k)$, which is optimal under the Set Cover Conjecture (SeCoCo) [23, 35].

Although many other variations of algebraically styled FPT algorithms are known [18, 39, 50, 27, 26, 28], the above methods (degree-extraction and multilinear detection) are remarkable in the power and simplicity of their applications. In this paper, we show an extension of this toolbox.

1.1 Determinantal sieving

We introduce *determinantal sieving*, a powerful new sieving operation that drastically extends the power of the tools of multilinear detection and constrained multilinear detection. Let $P(x_1, \dots, x_n)$ be a polynomial over a field \mathbb{F} of characteristic 2, and let $M \in \mathbb{F}^{k \times n}$ be a matrix (e.g., a linear matroid on the ground set $X = \{x_1, \dots, x_n\}$). For a monomial m in P , let $\text{supp}(m)$ be the set of variables x_i of non-zero degree in m . We show a sieving method that, using $O^*(2^k)$ evaluations of P , sieves for those monomials m in P that are multilinear of degree k and such that the matrix $M[\cdot, \text{supp}(m)]$ indexed by the support is non-singular. More precisely, we show the following.

Theorem 1.3 (Basis sieving). *Let $P(X)$ be a polynomial of degree d over a field \mathbb{F} of characteristic 2, and let $M = (X, \mathcal{I})$ be a matroid on X of rank k , represented by a matrix $A \in \mathbb{F}^{k \times X}$. There is a randomized algorithm with running time $O^*(d2^k)$ that tests if there is a multilinear term m in the monomial expansion of $P(X)$ such that the matrix $A[\cdot, \text{supp}(m)]$ is non-singular. The algorithm uses polynomial space, needs only evaluation access to P , has no false positives and produces false negatives with probability at most $2k/|\mathbb{F}|$.*

The proof is remarkably simple, consisting of merely inspecting the result of an application of inclusion-exclusion sieving; see Section 3. We also note that in all our applications, the failure rate can be made arbitrarily small with negligible overhead by moving to an extension field of \mathbb{F} .

We note a useful variant. In many applications, instead of insisting that the term m as a whole forms a basis for A (and thus, has degree precisely k), we may wish to sieve for monomials m whose support *spans* M . For example, if we think of the ground set X as being *coloured* by some colouring $c: X \rightarrow L$ to a label set L , then sieving for a basis corresponds to looking for a solution $S \subseteq X$ that is *rainbow* – i.e., all labels $c(x)$ for $x \in S$ are distinct – whereas we may wish to sieve for solutions $S \subseteq X$ that are merely *colourful* – i.e., at least k distinct colours $c(x)$ occur among elements $x \in S$. If $P(X)$ is multilinear in X , then this can be achieved by evaluating $P(X)$ at a point where each variable $x_i \in X$ is assigned a value $(1 + x'_i)x''_i$ for new variables x'_i, x''_i . The resulting expansion over a monomial m will generate all monomials whose support in X'_i is a subset of the support of m in X , and the additional variables x''_i prevent cancellation against contributions from other monomials m' in $P(X)$. We can thus use Theorem 1.3 to sieve for terms of $P(X)$ whose support spans M .

However, if $P(X)$ is not multilinear, then a useful phenomenon occurs. Since $P(X)$ is evaluated over a field of characteristic 2, some linear factors disappear; e.g., $(1 + x_i)^2 = 1 + x_i^2$, with no linear contribution in x_i . More generally, over characteristic 2, $(1 + x_i)^d$ has a non-zero linear term if and only if d is odd. Hence, we end up sieving for a basis among the variables of odd degree in m only – the *odd support* of m . We get the following.

Theorem 1.4 (Odd sieving). *Let $P(X)$ be a polynomial of degree d over a field \mathbb{F} of characteristic 2, and let $M = (X, \mathcal{I})$ be a matroid on X of rank k , represented by a matrix $A \in \mathbb{F}^{k \times X}$. There is a randomized algorithm with running time $O^*(d2^k)$ that tests if there is a term m in the monomial expansion of $P(X)$ such that the odd support of m spans M . The algorithm uses polynomial space, needs only evaluation access to P , has no false positives and produces false negatives with probability at most $(k + d)/|\mathbb{F}|$.*

For further examples, such as decorating every variable of X by multiple vectors from the matroid, see Section 3. Furthermore, the odd sieving method has applications on its own; see the *Diverse X* and *paths and linkages* examples below.

1.1.1 Over general fields

The aforementioned sieving algorithms only work over fields of characteristic 2. By utilizing *exterior algebra*, we can effectively sieve over arbitrary fields. We will follow the work of Brand et al. [27], who exhibited the power of exterior algebra in parameterized algorithms. Assume that a polynomial $P(X)$ over \mathbb{F} is represented by a skew arithmetic circuit C , a circuit where at least one input of every product gate is an input gate. (If C is not skew, then similar results apply with slightly larger running times.) Following the idea of Brand et al. [27], we attempt to evaluate the circuit C over the exterior algebra $\Lambda(\mathbb{F}^k)$. The exterior algebra is essentially a vector space of dimension 2^k , where the addition is commutative but the multiplication (called *wedge product*) is not (see Section 3.2 for the definition). Thus, naively evaluating over C will not preserve the coefficients of $P(X)$. We present two ways to circumvent this issue.

The first one concerns the restriction on the circuit. We consider *strongly monotone circuits*, which are basically circuit without any “cancellation” whatsoever. We show that the result of evaluating a strongly monotone circuit C over $\Lambda(\mathbb{F}^k)$ turns out nonzero only if $P(X)$ contains a monomial m such that $A[\cdot, \text{supp}(m)]$ is non-singular.

Theorem 1.5. *Let $P(X)$ be a polynomial of degree d over a field \mathbb{F} , represented by a skew strongly monotone arithmetic circuit C , and let $M = (X, \mathcal{I})$ be a matroid on X of rank k , represented by a matrix $A \in \mathbb{F}^{k \times X}$. There is a randomized algorithm with running time $O^*(2^k)$ that tests if there is a multilinear term m in the monomial expansion of $P(X)$ such that the matrix $A[\cdot, \text{supp}(m)]$ is non-singular. The algorithm uses $O^*(2^k)$ space, has no false positives and produces false negatives with probability at most $2k/|\mathbb{F}|$.*

We also provide a way to sieve over arbitrary arithmetic circuits by using the *lift mapping*, which maps every extensor in $\Lambda(\mathbb{F}^k)$ to $\Lambda(\mathbb{F}^{2k})$, a vector space of dimension 4^k . Although the lift mapping costs extra time and space usage, it brings commutativity to the algebra. This allows us to evaluate the circuit over the exterior algebra.

Theorem 1.6. *Let $P(X)$ be a polynomial of degree d over a field \mathbb{F} , represented by a skew arithmetic circuit C , and let $M = (X, \mathcal{I})$ be a matroid on X of rank k , represented by a matrix $A \in \mathbb{F}^{k \times X}$. There is a randomized algorithm with running time $O^*(4^k)$ that tests if there is a multilinear term m in the monomial expansion of $P(X)$ such that the matrix $A[\cdot, \text{supp}(m)]$ is non-singular. The algorithm uses $O^*(4^k)$ space, has no false positives and produces false negatives with probability at most $2k/|\mathbb{F}|$.*

1.1.2 Linear matroids

The most useful applications of determinantal sieving come when the labelling matrix M represents a *linear matroid* over the variable set. A *matroid* is a pair $M = (V, \mathcal{I})$ where V is the ground set and $\mathcal{I} \subseteq 2^V$ a set of *independent sets* in M , subject to the following axioms: (1) $\emptyset \in \mathcal{I}$; (2) If $B \in \mathcal{I}$ and $A \subset B$ then $A \in \mathcal{I}$; and (3) For any $A, B \in \mathcal{I}$ such that $|A| < |B|$ there exists an element $x \in B \setminus A$ such that $(A + x) \in \mathcal{I}$. A *linear matroid* is a matroid M represented by a matrix A with column set V , such that a set $S \subseteq V$ is independent in M if and only if $A[\cdot, S]$ is non-singular.

A more complete overview of matroid theory concepts is given in Section 2, but let us review two particularly relevant matroid constructions. A *uniform matroid* $U_{n,k}$ is the matroid $M = (V, \mathcal{I})$

where $\mathcal{I} = \binom{V}{\leq k}$ (for $|V| = n$), i.e., a set is independent if and only if it has cardinality at most k . Letting M be a uniform matroid in determinantal sieving corresponds to traditional multilinear detection. More generally, a *partition matroid* $M = (V, \mathcal{I})$ is defined by a partition $V = V_1 \cup \dots \cup V_d$ of the ground set and a list of capacities $(c_i)_{i=1}^d$; note that we allow $c_i > 1$ [84]. A set $S \subseteq V$ is independent if and only if $|S \cap V_i| \leq c_i$ for every $i \in [d]$. Constrained multilinear detection corresponds roughly to the case of M being a partition matroid (or more precisely, the truncation of a partition matroid to rank k). Both of these classes can be represented over fields of characteristic 2.

There also exists a range of transformations that can be applied to matroids, with preserved representation; see Section 2.1. Here, we only note the operation of *truncation*: Given a matroid $M = (V, \mathcal{I})$, represented over a field \mathbb{F} , and an integer k , we can in polynomial time *truncate* M to have rank k and otherwise preserve the representation [70, 75]. Thereby, whenever we are looking for a solution of rank k , we may assume that every matroid $M = (V, \mathcal{I})$ in our input is represented by a full-rank matrix of dimension $k \times |V|$.

In the above, we find it particularly interesting that the fastest known method for multilinear detection, which sieves over a random bijective labelling [16, 19], can be seen as a direct analogue of Theorem 1.3 applied to a representation of a uniform matroid. In this sense, the results of this paper come *for free* – they represent the same sieving steps that existing algorithms already perform, only computed on a more carefully chosen set of evaluation points.

1.2 Applications

Given Theorems 1.3–1.6, a large collection of applications can be achieved by combining a suitable enumerating polynomial for a problem with a suitable matroid labelling. Before we undertake a survey, let us more carefully define our terms. Let V be a ground set and $\mathcal{F} \subseteq 2^V$ a set system over V . An *enumerating polynomial* for \mathcal{F} is a polynomial $P(X, Y)$ over a set of variables $X \cup Y$, where $X = \{x_v \mid v \in V\}$, such that the following holds.

1. $P(X, Y)$ is multilinear in X
2. For $S \subseteq V$, there is a monomial m in $P(X, Y)$ whose support in X is S if and only if $S \in \mathcal{F}$.

Similarly, to capture applications of Theorem 1.4 (odd sieving), define a *parity-enumerating polynomial* for \mathcal{F} as a polynomial $P(X, Y)$ where there exists a monomial m whose odd support in X is S , $S \subseteq V$, if and only if $S \in \mathcal{F}$. The definition can be generalized further – for example, if we want to refer to an “enumerating polynomial for walks” we could treat walks as *multisets* of vertices or edges, and adjust the definition accordingly. However, the above suffices for almost all applications.

We next survey results covered by our approach. Our results cover multiple areas, and include both significant speedups of previous results (see Table 1) and generalisations where a previous running time for a problem can be reproduced in a much broader setting. Furthermore, in general, both the proofs and the algorithms are short and simple, given existing families of enumerating polynomials and linear matroids.

1.2.1 Matroid Covering, Packing and Intersection Problems

We begin with a straight-forward application to the SET COVER and SET PACKING problems. Let V be a ground set and $\mathcal{E} \subseteq 2^V$ a collection of sets. Let $M = (V, \mathcal{I})$ be a matroid of rank k , and let t be an integer. In RANK k SET COVER we ask, is there a subcollection $S \subseteq \mathcal{E}$ with $|S| \leq t$ such that $\bigcup S$ spans M ? In RANK k SET PACKING we ask if there is a collection $S \subseteq \mathcal{E}$ of pairwise disjoint

sets with $|S| = t$ such that $\bigcup S$ is a basis of M . (The variant of RANK k SET PACKING where $\bigcup S$ is only required to be independent in M , not a basis, reduces to the above via truncation of M .)

Theorem 1.7. RANK k SET COVER and RANK k SET PACKING for matroids represented over a field of characteristic 2 can be solved in randomized time $O^*(2^k)$ and polynomial space, and in time $O^*(2^{\omega k/2})$ and $O^*(2^k)$ space over general fields.

To achieve this result, we use a simple subset-enumerating polynomial. Assume an input $(V, \mathcal{E}, M, w, k)$ is given, and define a set of variables $X_{v,E}$, $v \in V$, $E \in \mathcal{E}$, as well as a set of fingerprinting variables $Y = \{y_{i,E} \mid i \in [t], E \in \mathcal{E}\}$ to prevent cancellations. Define

$$P(X, Y) = \prod_{i=1}^t \sum_{E \in \mathcal{E}} y_{i,E} \prod_{v \in E} x_{v,E}.$$

Note that $P(X)$ is multilinear. Hence to solve RANK k SET PACKING we associate each variable $x_{v,E}$ with the vector representing v in M , and invoke Theorem 1.3 respectively 1.5 depending on the representation of M . For RANK k SET COVER we simply evaluate P at a point $x_{v,E} \leftarrow 1 + x_{v,E}$ for every $x_{v,E} \in X$ for the same result.

Note that Theorem 1.7 is tight under SeCoCo, since SET COVER corresponds to the simple case where each element $v_i \in V$ is associated with the n -dimensional unit vector e_i .

Theorem 1.7 improves on state of the art even for very simple settings. In MATROID q -PARITY, the input is a matroid $M = (V, \mathcal{I})$, a partition of V into set of size q , and an integer k , and the question is whether there is a packing of k sets that is independent in M . This problem can be solved in polynomial time if $q = 2$ and M is linear, but is hard even for linear matroids if $q \geq 3$. The fastest known algorithm for MATROID q -PARITY by Brand and Pratt runs in time $O^*(4^{qk})$ with exponential space [28], improving on a previous result of Fomin et al. [50] with running time $O^*(2^{\omega qk})$, where $\omega < 2.37$ is the matrix multiplication exponent. We get the following.

Corollary 1.8. MATROID q -PARITY for a linear matroid over a field of characteristic 2 can be solved in randomized time $O^*(2^{qk})$ and polynomial space.

For a related problem, we get a greater speedup. In q -MATROID INTERSECTION, the input is q matroids M_1, \dots, M_q of rank k , and the question is if they have a common basis. Again, this is tractable if $q = 2$, but NP-hard if $q \geq 3$ even for linear matroids. Assume that the matroids are represented by matrices A_1, \dots, A_q over a common field \mathbb{F} and a common ground set V , where w.l.o.g. every matrix A_i has k rows and has rank k over \mathbb{F} . We can use the Cauchy-Binet formula to sieve for solutions more efficiently. Let $X = \{x_v \mid v \in V\}$ be a set of variables and let A_1' be the result of scaling every column v of A_1 by x_v . By the Cauchy-Binet formula,

$$P(X) := \det(A_1' A_2^T) = \sum_{B \in \binom{V}{k}} \det A_1[\cdot, B] \det A_2[\cdot, B] \prod_{v \in B} x_v.$$

Thus $P(X)$ enumerates monomials $\prod_{v \in B} x_v$ for common bases B of A_1 and A_2 , and we only have to sieve for terms that in addition are bases of the remaining $q - 2$ matroids. We get the following.

Theorem 1.9. q -MATROID INTERSECTION for linear matroids represented over a common field \mathbb{F} of characteristic 2 can be solved in randomized time $O^*(2^{(q-2)k})$ and polynomial space.

The previous best result is Brand and Pratt [28], with running time $O^*(4^{qk})$. In particular, for $q = 3$ this improves on the state of the art from $O^*(4^{3k})$ to $O^*(2^k)$. Theorem 1.9 matches the fastest algorithm by Björklund et al. [19] for the much simpler q -DIMENSIONAL MATCHING problem.

As a particular special case, Theorem 1.9 with $q = 3$ implies a polynomial-space, $O^*(2^n)$ -time algorithm for DIRECTED HAMILTONIAN PATH, which despite intense efforts at improvement remains the state of the art for the general case [13, 21, 37].

1.2.2 Fair and Diverse Solutions

Fairness and diversity are important concepts in many areas of research, including artificial intelligence and optimization, and have also seen increased focus in theoretical computer science. We discuss two related problems: finding a balanced-fair solution and a diverse collection of solutions.

The problem of finding a balanced-fair solution arises in many contexts [5, 14, 32, 33], including MATROID INTERSECTION, k -MATCHING, and k -PATH. We define a general problem BALANCED SOLUTION: Given a set E with coloured elements, a collection \mathcal{F} of subsets of E , the goal is to find a set $S \in \mathcal{F}$ of size k such that the number of elements of S with each colour is within certain bounds. We show that this problem can be solved in $O^*(2^k)$ time using basis sieving:

Theorem 1.10. BALANCED SOLUTION *can be solved in $O^*(2^k)$ time if there is an enumerating polynomial for \mathcal{F} that can be efficiently evaluated.*

The problem of finding a diverse collection of solutions is another important optimization problem. Here, the goal is to find not just a single optimal solution, but a collection of solutions that are diverse in some sense. We measure diversity in terms of Hamming distance, i.e., diverse solutions should have a large Hamming distance between them. This problem has received significant attention in the parameterized complexity literature [11, 12, 46, 48, 60]. We discuss a general method based on the odd sieving technique that can be used to find a diverse collection of solutions for a wide range of optimization problems. We define the DIVERSE COLLECTION problem defined as follows. The input is a set E , collections of subsets $\mathcal{F}_1, \dots, \mathcal{F}_k$, and $d_{i,j} \in \mathbb{N}$ for $1 \leq i < j \leq k$, and the goal is to find subsets $S_i \in \mathcal{F}_i$ for each $i \in [k]$ such that $|S_i \Delta S_j| = |(S_i \setminus S_j) \cup (S_j \setminus S_i)| \geq d_{i,j}$ for every i, j . Let $D = \sum_{i < j \in [k]} d_{i,j}$. We use the odd sieving algorithm to obtain an $O^*(2^D)$ -time algorithm. The key here is to use a distinct set of variables for every pair i, j . Thereby, those elements in the intersection of two solutions, having contribution two, can be excluded in the odd sieving.

Theorem 1.11. DIVERSE COLLECTION *can be solved in $O^*(2^D)$ time if all collections \mathcal{F}_i admit enumerating polynomials that can be efficiently evaluated.*

This leads to significant speed-ups compared to existing algorithms, one for DIVERSE MATCHINGS and another for d -DISTINCT BRANCHINGS. The DIVERSE MATCHINGS problem ask whether a given graph contains k perfect matchings M_1, \dots, M_k whose pairwise Hamming distances are all at least d . Fomin et al. [46] give an algorithm with running time $2^{2^{O(kd)}}$. We obtain a faster algorithm running in time $O^*(2^{d \binom{k}{2}})$. In d -DISTINCT BRANCHINGS, we are given a directed graph G , two vertices s, t , and an integer d , and we search for an in-branching rooted at s and out-branching rooted t whose Hamming distance is at least d . This problem can be solved in $O^*(2^d)$ time. In particular, this answers the question of Bang-Jensen et al. [8] whether there exists an $O^*(2^{O(d)})$ -time algorithm. Previously known algorithms run in time $O^*(2^{d^2 \log^2 d})$ [57] and $O^*(d^{O(d)})$ [8].

1.2.3 Undirected paths and linkages

As noted above, among the earliest and most powerful applications of algebraic FPT algorithms is for path and cycle problems. In fact, all the current fastest FPT algorithms for k -PATH – randomized time $O^*(1.66^k)$ for undirected graphs [19] and $O^*(2^k)$ for digraphs [93]; deterministic $O^*(2.55^k)$ time for both variants [90] – ultimately have algebraic underpinnings.

Another highly surprising result was for the T -CYCLE problem (we use the name from Fomin et al. [47] to distinguish more clearly from k -CYCLE). Here, the input is an undirected graph G and a set terminals $T \subseteq V(G)$, and the question is whether G contains a simple cycle C that

passes through all vertices in T . This problem was known to be FPT parameterized by $k = |T|$, using graph structural methods, but the running time was impractical [64]. Björklund, Husfeldt and Taslamán [20] showed an $O^*(2^k)$ -time algorithm, based on cancellations in the evaluation of a large polynomial. Wahlström [92] showed that the problem even has a *polynomial compression* in k , based on a reinterpretation of the previous algorithm in terms of the determinant of a modified Tutte matrix (similar to Björklund’s celebrated $O^*(1.66^n)$ -time algorithm for HAMILTONICITY [16]). It is this latter determinant approach that we build upon in the algorithms for path and linkage problems in this paper.

Let G be an undirected graph and $S, T \subseteq V(G)$ be disjoint vertex sets. An (S, T) -linkage in G is a collection of $|S| = |T|$ pairwise vertex-disjoint paths from S to T – i.e., a vertex-disjoint (S, T) -flow assuming that vertices of $S \cup T$ have capacity 1. Let \mathcal{P} be an (S, T) -linkage for some (G, S, T) . A *padding* of \mathcal{P} is a collection of oriented cycles that covers $G - V(\mathcal{P})$, where every cycle has length at most 2 (i.e., every cycle is either a 2-cycle uvw over some edge $uv \in E(G)$ or a loop v on some vertex $v \in V(G)$). We show that there is an enumerating polynomial for padded (S, T) -linkages.

Lemma 1.12. *Let G be an undirected graph, possibly with loops, and let $S, T \subseteq V(G)$ be disjoint. In polynomial time, we can construct a matrix A with entries from the variable set $X = \{x_e \mid e \in E(G)\}$ such that the polynomial $P(X) = \det A$ evaluated over a field of characteristic 2 enumerates padded (S, T) -linkages of G ; i.e., $P(X)$ is a parity-enumerating polynomial for (S, T) -linkages.*

This result is interesting even when $|S| = |T| = 1$, in which case $P(X)$ enumerates padded (s, t) -paths. We find this remarkable, as normally, a polynomial that is efficiently computable would only be expected to enumerate walks, as opposed to paths or cycles. It is not *too* powerful, since the padding terms from 2-cycles prevent us from using it to solve, e.g., HAMILTONIAN PATH in polynomial time. But it is highly useful for FPT purposes, since Theorem 1.4 allows us to sieve for terms that span a linear matroid M while ignoring the padding-part of each padded linkage. Thus we get the following.

Theorem 1.13. *Let $G = (V, E)$ be an undirected graph and let $M = (V, \mathcal{I})$ be a matroid represented over a field of characteristic 2. Let $S, T \subseteq V(G)$ be disjoint vertex sets and $k \in \mathbb{N}$. In randomized time $O^*(2^k)$ and polynomial space we can find a minimum-length (S, T) -linkage in G that has rank at least k in M (or determine that none exist).*

This result improves and generalizes a number of results. Fomin et al. [47] gave randomized algorithms in time $O^*(2^{k+p})$ for finding a minimum-length colourful (S, T) -linkage, and in time $O^*(2^{p+O(k^2 \log(q+k))})$ for finding a minimum-length (S, T) -linkage of rank at least k in M , where M is represented by a matrix over a finite field of order q and $p = |S| = |T|$.¹ Theorem 1.13 directly generalizes the first result, removing the dependency on p , and improves the running time for the second in the case that M can be represented over a field of characteristic 2. It also significantly simplifies the correctness proof, which in [47] runs to over 20 pages.

As they observe, even the problem COLOURFUL (s, t) -PATH captures a number of problems, including T -CYCLE, LONG (s, t) -PATH and LONG CYCLE (i.e., finding an (s, t) -path, respectively cycle, of length at least k). Finding an (s, t) -path of rank at least k also generalizes the variant LIST T -CYCLE, previously shown to be FPT by Panolan, Saurabh and Zehavi [85].

We also show an improvement to LONG (s, t) -PATH and LONG CYCLE. Fomin et al. [47] ask as an open problem whether these can be solved in time $O^*((2 - \epsilon)^n)$ for some $\epsilon > 0$, given that k -PATH and k -CYCLE have $O^*(1.66^k)$ -time algorithms due to Björklund et al. [19]. We confirm this.

¹The formulation of Fomin et al. [47] is slightly different, but equivalent under simple transformations.

Theorem 1.14. LONG (s, t) -PATH and LONG CYCLE can be solved in randomized time $O^*(1.66^k)$ and polynomial space.

1.2.4 Subgraph problems

Another class of problems where algebraic methods have been important is for the general question of finding subgraphs of a graph G with a given property. We give two results in this domain.

First, let $G = (V, E)$ be a graph and M a matroid over V . Let RANK k CONNECTED SUBGRAPH be the following general problem: Given integers k and t , is there a connected subgraph H of G on at most t vertices such that $V(H)$ has rank at least k in M ?

Theorem 1.15. RANK k CONNECTED SUBGRAPH for a linear matroid M can be solved in randomized time $O^*(2^k)$ and polynomial space if M is represented over a field of characteristic 2, and in randomized time $O^*(2^{\omega k})$ and space $O^*(4^k)$ otherwise.

This result is an application of the powerful notion of *branching walks*, introduced by Nederlof [82], which underlie several FPT algorithms. We rely on Björklund et al. [23] who gave an explicit algorithm for evaluating the branching walk polynomial. As special cases of Theorem 1.15 with various matroids M we recover the $O^*(2^k)$ -time algorithms for STEINER TREE [82] and GROUP STEINER TREE [78] on k terminals, and for GRAPH MOTIF and CLOSEST GRAPH MOTIF [23].

More generally, consider SUBGRAPH ISOMORPHISM, the problem of finding a subgraph of G isomorphic to a given graph H . This is W[1]-hard in general (cf. k -clique), but is FPT parameterized by $|V(H)|$ if H has bounded treewidth. In fact, up to plausible conjectures, the dependency on the treewidth w for known algorithms is optimal for every $w \geq 3$ [29]. Like previous algorithms, we employ the *homomorphism polynomial* (see, e.g., Brand [26]), and show the following.

Theorem 1.16. Let G and H be undirected graphs, $k = |V(H)|$ and $n = |V(G)|$, and let M be a linear matroid over $V(G)$. Let a tree decomposition of H of width w be given. We can find a subgraph H' of G isomorphic to H such that $V(H')$ is independent in M in randomized time $O(k^{O(1)} \cdot 2^k \cdot n^{w+1})$ and polynomial space if M is represented over a field of characteristic 2, and in time $O(k^{O(1)} \cdot 2^{\omega k} \cdot n^{w+1})$ and space $O^*(4^k)$ otherwise.

1.2.5 Speeding up dynamic programming

The *representative sets lemma* [72, 75] is a statement from matroid theory that has seen a multitude of applications in parameterized complexity, both in kernelization [68] and in FPT algorithms [75, 50]. The latter class of application typically consists of a *sped-up dynamic programming* algorithm; e.g., a dynamic programming algorithm over a state space that could potentially contain $n^{O(k)}$ different partial solutions, but where the representative sets lemma is used to prove that it suffices to maintain a set of $2^{O(k)}$ *representative* solutions. This includes algorithms for paths and cycles [50] as well as many more complex questions. We refer to this as a *rep-set DP*.

For many of these applications, faster algorithms are known, even in polynomial space, if randomness is allowed, and the main contribution of the representative sets lemma becomes to enable an almost competitive deterministic FPT algorithm [50, 90]. However, for other applications this is not so clear, and there are many applications of the representative sets lemma where no faster method is known. With the more powerful algebraic sieving methods of this paper, we can revisit some of these applications and show a speed-up of the algorithm, while at the same time reducing the space usage to polynomial space. We give three examples.

In MINIMUM EQUIVALENT GRAPH (MEG), the input is a digraph G , and the task is to find a subgraph G' of G with a minimum number of edges such that G and G' have the same reachability

Table 1: A list of speed-ups over previous results. Results marked with \dagger use exponential space, and those with \S only work over a field of characteristic 2. For the linkage problems, p is the order of the linkage.

Problem	Existing	New
q -MATROID INTERSECTION	$O^*(4^{qk})^\dagger$ [28]	$O^*(2^{(q-2)k})^\S$ $O^*(2^{\min(q, 2q-4) \cdot k})^\dagger$
q -MATROID PARITY	$O^*(4^{qk})^\dagger$ [28]	$O^*(2^{qk})^\dagger, O^*(2^{qk})^\S$
LONG (s, t) -PATH	$O^*(2^k)$ [47]	$O^*(1.66^k)$
COLOURFUL (S, T) -LINKAGE	$O^*(2^{k+p})$ [47]	$O^*(2^k)$
RANK k (S, T) -LINKAGE	$O^*(2^{p+O(k^2 \log(k+ \mathbb{F}))})$ [47]	$O^*(2^k)^\S$
DIVERSE PERFECT MATCHINGS	$O^*(2^{2^{O(D)}})^\dagger$ [46]	$O^*(2^D)$
k -DISTINCT BRANCHINGS	$O^*(k^{O(k)})^\dagger$ [8]	$O^*(2^k)$
MINIMUM EQUIVALENT GRAPH	$O^*(2^{4\omega n})^\dagger$ [50]	$O^*(2^{2n})$
(UN)DIRECTED EULERIAN DELETION	$O^*(2^{(2+\omega)k})^\dagger$ [55]	$O^*(2^k)$
CHORDAL-CONFLICT-FREE MATCHING	$O^*(2^{(2+2\omega)k})^\dagger$ [2]	$O^*(2^{2k})$

relation. Fomin et al. [50] give the first single-exponential algorithm for MEG. They show that MEG ultimately reduces down to finding an in-branching B_1 and an out-branching B_2 with a common root sharing at least ℓ edges, which they solve via rep-set DP in time $O^*(2^{4\omega n})$. We reduce this question to an application of 4-MATROID INTERSECTION and get the following.

Theorem 1.17. *MINIMUM EQUIVALENT GRAPH can be solved in polynomial space and randomized time $O^*(2^{2n})$.*

In (UNDIRECTED/DIRECTED) EULERIAN EDGE DELETION, the input is a graph G (undirected respectively directed), and the question is whether we can remove at most k edges from G so that the resulting graph is Eulerian (i.e., has a closed walk that visits every edge precisely once). Cai and Yang [30] surveyed related problems, but left the above questions open. Cygan et al [38] gave the first FPT algorithms, with running times of $O^*(2^{O(k \log k)})$, and Goyal et al. [55] improved this to $O^*(2^{(2+\omega)k})$ using a rep-set DP approach over the co-graphic matroid. We combine the co-graphic matroid approach with suitable enumerating polynomials to get the following.

Theorem 1.18. *UNDIRECTED EULERIAN EDGE DELETION and DIRECTED EULERIAN EDGE DELETION can be solved in $O^*(2^k)$ randomized time and polynomial space.*

Finally, we consider a more unusual application. Consider a generic problem where we are searching for a subset S with property Π of a ground set V . In the *conflict-free* version, the input additionally contains a graph $H = (V, E)$ and S is required to be an independent set in H . Naturally, this is hard in general (even disregarding the property Π), but multiple authors have considered restricted variants. In particular, if H is chordal then Agrawal et al. [2] show that CONFLICT-FREE MATCHING can be solved in $O^*(2^{(2\omega+2)k})$ time, and Jacob et al. [62] show that CONFLICT-FREE SET COVER can be solved in $O^*(3^n)$ time. We note that the *independent set polynomial* (in our terminology, an enumerating polynomial for independent sets in a graph) can be efficiently evaluated if H is chordal [1], allowing us to speed up both results. See Section 8.3 for details.

Structure of the paper. In Section 2 we cover preliminaries, and in Section 3 we prove the determinantal sieving statements of Theorem 1.3–1.6. In Section 4–8 we cover the applications mentioned in Section 1.2.1–1.2.5, respectively. We conclude in Section 9 with discussion and open problems.

2 Preliminaries

We use standard terminology from parameterized complexity, see, e.g., the book of Cygan et al. [36]. For background on graph theory, see Diestel [44] and Bang-Jensen and Gutin [7].

Let $P(X)$ be a polynomial over a set of variables $X = \{x_1, \dots, x_n\}$. A monomial is a product $m = x_1^{m_1} \cdots x_n^{m_n}$ for nonnegative integers m_1, \dots, m_n . A monomial m is called multilinear if $m_i \leq 1$ for each $i \in [n]$. We say that its *support* is $\{i \in [n] \mid m_i > 0\}$ and that its *odd support* is $\{i \in [n] \mid m_i \equiv 1 \pmod{2}\}$ denoted by $\text{supp}(m)$ and $\text{osupp}(m)$, respectively. We sometimes use the notation X^m for the monomial $m = x_1^{m_1} x_2^{m_2} \cdots x_n^{m_n}$, to clarify that the monomial m does not include a coefficient. For a set of variables $X' = \{x'_1, \dots, x'_n\}$ we will also write $(X')^m = \prod_{i=1}^n (x'_i)^{m_i}$. For a monomial m in the monomial expansion of $P(X)$, we let $P(m)$ denote the coefficient of m in P , i.e., $P(X) = \sum_m P(m)X^m$ where m ranges over all monomials in $P(X)$. The total degree of $P(X)$ is $\max_m \sum_{i \in [n]} m_i$. The Schwartz-Zippel lemma [89, 96] states that a polynomial $P(X)$ of total degree at most d over a field \mathbb{F} becomes nonzero with probability at least $1 - d/|\mathbb{F}|$ when evaluated at uniformly chosen elements from \mathbb{F} , unless $P(X)$ is identically zero.

The following two lemmas are the foundation of our sieving algorithms.

Lemma 2.1 (Interpolation). *Let $P(z)$ be a polynomial over a field \mathbb{F} of degree $n - 1$. Suppose that $P(z_i) = p_i$ for distinct $z_1, \dots, z_n \in \mathbb{F}$. By the Lagrange interpolation,*

$$P(z) = \sum_{i \in [n]} p_i \prod_{j \in [n] \setminus \{i\}} \frac{z - z_j}{z_i - z_j}.$$

Thus, given n evaluations p_1, \dots, p_n of $P(z)$, the coefficient of z^t in $P(z)$ for every $t \in [n]$ can be computed in polynomial time.

Lemma 2.2 (Inclusion-exclusion [92]). *Let $P(Y)$ be a polynomial over a set of variables $Y = \{y_1, \dots, y_n\}$ and a field of characteristic two. For $T \subseteq [n]$, Q be a polynomial identical to P except that the coefficients of monomials not divisible by $\prod_{i \in T} y_i$ is zero. Then, $Q = \sum_{I \subseteq T} P_{-I}$, where $P_{-I}(y_1, \dots, y_n) = P(y'_1, \dots, y'_n)$ for $y'_i = y_i$ if $i \notin I$ and $y'_i = 0$ otherwise.*

Let A be a matrix over a field \mathbb{F} . For a set of rows I and columns J , we denote by $A[I, J]$ the submatrix containing rows I and columns J . If I contains all rows (J contains all columns), then we use the shorthand $A[\cdot, J]$ ($A[I, \cdot]$, respectively).

For a $k \times n$ -matrix A_1 and an $n \times k$ -matrix A_2 , the Cauchy-Binet formula states that

$$\det(A_1 A_2) = \sum_{S \in \binom{[n]}{k}} \det(A_1[\cdot, S]) \det(A_2[S, \cdot]).$$

A square matrix A is called *skew-symmetric* if $A = -A^T$. Suppose that the rows and columns of A are indexed by V . The *Pfaffian* of A is defined by

$$\text{Pf } A = \sum_M \sigma_M \prod_{uv \in M} M[u, v],$$

where M ranges over all perfect matchings of the complete graph $(V, \binom{V}{2})$, and $\sigma_M = \pm 1$ is the sign of M whose definition is not relevant in this work (see e.g., [81]). It is well-known that $\det A = (\text{Pf } A)^2$.

2.1 Linear matroids

We review the essentials of matroid theory, with a focus on linear matroids. For more background, see Oxley [84] and Marx [75]. A *matroid* is a pair $M = (V, \mathcal{I})$ where V is the ground set and $\mathcal{I} \subseteq 2^V$ a set of *independent sets* in M , subject to the following axioms:

1. $\emptyset \in \mathcal{I}$
2. If $B \in \mathcal{I}$ and $A \subset B$ then $A \in \mathcal{I}$
3. For any $A, B \in \mathcal{I}$ such that $|A| < |B|$ there exists an element $x \in B \setminus A$ such that $(A + x) \in \mathcal{I}$.

A *basis* of a matroid M is a maximal independent set. The *rank* $r(M)$ of M is the cardinality of a basis of M . A *linear matroid* is a matroid M represented by a matrix A with column set V , such that a set $S \subseteq V$ is independent in M if and only if the set of columns of A indexed by S is linearly independent. We review some useful matroid constructions, expanded from the introduction. All of the matroids below can be represented over fields of characteristic 2, although in some cases the only known methods for efficiently constructing a representation are randomized.

- A *uniform matroid* $U_{n,k}$ is the matroid $M = (V, \mathcal{I})$ where $\mathcal{I} = \binom{V}{\leq k}$ (for $|V| = n$), i.e., a set is independent if and only if it has cardinality at most k .
- A *partition matroid* $M = (V, \mathcal{I})$ is defined by a partition $V = V_1 \cup \dots \cup V_d$ of the ground set and a list of capacities $(c_i)_{i=1}^d$. A set $S \subseteq V$ is independent if and only if $|S \cap V_i| \leq c_i$ for every $i \in [d]$.
- Given an undirected graph $G = (V, E)$, the *graphic matroid* of G is a matroid $M = (E, \mathcal{I})$ where a set $F \subseteq E$ is independent if and only if it is acyclic. The *cographic matroid* of G is a matroid $M = (E, \mathcal{I})$ where a set $F \subseteq E$ is independent if and only if it preserves connectivity (i.e., G and $G - F$ have the same connected components).
- Let $G = (U \cup V, E)$ be a bipartite graph. The *transversal matroid* of G is the matroid $M = (V, \mathcal{I})$ where a set $S \subseteq V$ is independent if and only if it is matchable in G .
- Let $G = (V, E)$ be a digraph and $T \subseteq V$ a set of terminals. A set $S \subseteq V$ is *linked* to T if there is a collection of $|S|$ pairwise vertex-disjoint paths from S to T . The set of all sets $S \subseteq V$ that are linked to T form a matroid called a *gammoid*.

If $M = (V, \mathcal{I})$ is a matroid, the *dual matroid* of M is the matroid $M^* = (V, \mathcal{I}')$ where a set $S \subseteq V$ is independent in M^* if and only if $V \setminus S$ contains a basis of M . Given a representation for M , a representation for M^* can be constructed in deterministic polynomial time. Given two matroids $M_1(V_1, \mathcal{I}_1)$ and $M_2(V_2, \mathcal{I}_2)$ on disjoint sets V_1 and V_2 , the *disjoint union* of M_1 and M_2 is the matroid $M = M_1 \vee M_2 = (V, \mathcal{I})$ where $V = V_1 \cup V_2$ and a set $S \subseteq V$ is independent if and only if $S \cap V_1 \in \mathcal{I}_1$ and $S \cap V_2 \in \mathcal{I}_2$. More generally, for any two matroids $M_1 = (V_1, \mathcal{I}_1)$ and $M_2 = (V_2, \mathcal{I}_2)$ the *matroid union* $M = M_1 \vee M_2$ is the matroid $M = (V, \mathcal{I})$ where $V = V_1 \cup V_2$ and $\mathcal{I} = \{I_1 \cup I_2 \mid I_1 \in \mathcal{I}_1, I_2 \in \mathcal{I}_2\}$. Given representations of M_1 and M_2 , a representation for $M_1 \vee M_2$ can be constructed in randomized polynomial time. The *extension* of a matroid $M = (V, \mathcal{I})$ by rank d is the matroid $M \vee M'$ where M' is the uniform matroid of rank d over V .

For a matroid $M = (V, \mathcal{I})$, the *k-truncation* of M is the matroid $M' = (V, \mathcal{I}')$ where for $S \subseteq V$, $S \in \mathcal{I}'$ if and only if $S \in \mathcal{I}$ and $|S| \leq k$. Given a representation of M over a field \mathbb{F} , a representation of the *k-truncation* of M over an extension field of \mathbb{F} can be constructed in polynomial time [70, 75].

Given two matroids $M_1 = (V, \mathcal{I})$ and $M_2 = (V, \mathcal{I})$, the *matroid intersection* problem is to find a common basis B of M_1 and M_2 . Matroid intersection can be solved in polynomial time, with a variety of methods [88]. In this paper, with a focus on linear matroids, we note that the Cauchy-Binet formula implies an enumerating polynomial for matroid intersection, and thereby a randomized efficient algorithm. More generally, given a matroid $M = (V, \mathcal{I})$ and a partition E of V into pairs, the *matroid matching* (or *matroid parity*) problem is to find a basis B of M which is the union of $|B|/2$ pairs. Matroid matching is infeasible in general, but efficiently solvable over linear matroids [73, 88].

2.2 Enumerating polynomials

Let V be a ground set and $\mathcal{F} \subseteq 2^V$ be a set family over V . An *enumerating polynomial* over a set of variables $\{x_v \mid v \in V\}$ and auxiliary variables Y over a field \mathbb{F} is

$$P(X, Y) = \sum_{S \in \mathcal{F}} Q_S(Y) \prod_{v \in S} x_v,$$

where $Q_S(Y)$ for $S \in \mathcal{F}$ is a polynomial over \mathbb{F} that is not identically zero. We give useful examples of enumerating polynomials that can be efficiently evaluated below.

***k*-walks.** For a directed graph $G = (V, E)$, two vertices $s, t \in V$, and an integer k , an enumerating polynomial for k -walks from s to t is defined as follows. Let $X = \{x_{s,0}\} \cup \{x_{v,i} \mid v \in V, i \in [k]\} \cup \{x_{e,i} \mid e \in E, i \in [k]\}$ be variables. For every A_i define a $V \times V$ -matrix A_i with

$$A_i[u, v] = \begin{cases} x_{v,i} x_{uv,i} & \text{if } uv \in E \\ 0 & \text{otherwise.} \end{cases}$$

Then, the polynomial

$$P(X) = x_{s,0} \cdot e_s^T A_1 A_2 \cdots A_k e_t,$$

where e_s and e_t are the unit vectors with $e_s[s] = 1$ and $e_t[t] = 1$, enumerates all (labelled) k -walks from s to t . The polynomial can be defined for undirected graphs analogously.

Matroid Intersections. For linear matroids $M_1 = (V, \mathcal{I}_1), M_2 = (V, \mathcal{I}_2)$ with the ground set V represented by $A_1, A_2 \in \mathbb{F}^{k \times V}$, let $X = \{x_v \mid v \in V\}$ be variables for V . Then, by the Cauchy-Binet formula, the polynomial

$$P(X) = \det A_1 A_X A_2^T = \prod_{B \in \binom{V}{k}} \det A_1[\cdot, B] \det A_2[\cdot, B] \prod_{v \in B} x_v,$$

where A_X is a diagonal matrix of dimension $V \times V$ with $A_X[v, v] = x_v$ for every $v \in V$, enumerates all matroid intersection terms. Particularly, we obtain an effective evaluation of an enumerating polynomial for branchings in directed graphs. Recall that an out-branching (in-branching) is a rooted tree with every arc oriented away from (towards) the root. This can be expressed as the intersection of a graphic matroid and a partition matroid, where the partition matroid ensures that every vertex has in-degree (out-degree) at most one. Hence this is a special case of matroid intersection and an enumerating polynomial for out-branchings and in-branchings can be efficiently evaluated. Alternatively, one can use the directed matrix-tree theorem (see [21, 54]).

Perfect matchings. For an undirected graph $G = (V, E)$ (with a fixed ordering $<$ on V), the *Tutte matrix* is defined by

$$A[u, v] = \begin{cases} x_{uv} & \text{if } uv \in E \text{ and } u < v \\ -x_{uv} & \text{if } uv \in E \text{ and } v < u \\ 0 & \text{otherwise.} \end{cases}$$

Then, the Pfaffian $\text{Pf } A$ enumerates all perfect matchings, which can be efficiently evaluated using an elimination procedure. For an integer k , all k -matchings (matchings with k edges) also can be enumerated: Introduce $n - 2k$ vertices that are adjacent to all vertices in G ; the Pfaffian of the resulting graph enumerates all k -matchings.

3 Determinantal sieving

3.1 Over a field of characteristic 2

We show that, with only evaluation access to a polynomial (over a field of characteristic 2), we can sieve for those terms in its monomial expansion spanning a linear matroid. We will give two sieving algorithms, one that sieves for terms that are also independent (basis sieving) and the other that sieves for terms whose odd support sets are spanning (odd sieving). One could derive basis sieving from odd sieving using polynomial interpolation (Lemma 2.1). We will, however, give a direct proof for basis sieving as well because basis sieving itself has applications. Typically, basis sieving is useful when we are search for a solution of size exactly k (regardless of whether the objective is maximisation or minimisation). Odd sieving is particularly powerful when we want to exclude variables in the support set with even (typically 2) contributions. See Sections 5.2 and 6 for such applications.

We begin with a support statement. This is the central observation for our sieving algorithms.

Lemma 3.1. *Let $A \in \mathbb{F}^{k \times k}$ be a matrix over a field \mathbb{F} of characteristic 2 and define the polynomial*

$$P(y_1, \dots, y_k) = \prod_{i=1}^k \sum_{j=1}^k y_j A[j, i].$$

Then the coefficient of $\prod_{i=1}^k y_i$ in P is $\det A$.

Proof. Expanding the product into monomials, we get precisely

$$\sum_{f: [k] \rightarrow [k]} \prod_{i=1}^k y_{f(i)} A[f(i), i]$$

where f ranges over all mappings $[k] \rightarrow [k]$. Considering only those terms of the sum which contain all variables y_i , $i \in [k]$ we find that the coefficient of $\prod_{i=1}^k y_i$ is precisely a sum over all transversals of A , i.e., $\det A$, in particular, since \mathbb{F} is of characteristic 2 the sign term of the determinant disappears. \square

If performed over fields of characteristic other than 2, then instead of $\det A$ the coefficient is the permanent of A (while over fields of characteristic 2, the permanent and the determinant agree). To cover applications for fields of other characteristics, we instead use the *exterior algebra*; see Section 3.2. For the below, we focus on applications over fields of characteristic 2.

3.1.1 Sieving for bases

The following is the most immediate application of Lemma 3.1 (proving Theorem 1.3 from the introduction). We also add an observation about tighter running time when the polynomial is already homogeneous.

Theorem 3.2 (Basis sieving). *Let $X = \{x_1, \dots, x_n\}$ be a set of variables and let $P(X)$ be a polynomial of degree d over a field \mathbb{F} of characteristic 2. Let $A \in \mathbb{F}^{k \times n}$ be a matrix representing a matroid $M = (X, \mathcal{I})$. In time $O^*(d2^k)$ and polynomial space, using evaluation access to P , we can test if the monomial expansion of P contains a multilinear monomial m whose support is a basis for M . Our algorithm is randomized with no false positives and failure probability at most $2k/|\mathbb{F}|$. If P is homogeneous in X , then the polynomial overhead disappears and the running time is $O(2^k(nk \cdot f + T))$ where f is the time for a field operation and T is the time to evaluate P .*

Proof. Let $P_k(X)$ denote the homogeneous degree k part of $P(X)$, i.e., for every monomial m of total degree k the coefficient of m in P_k is $P_k(m) = P(m)$, and for every other monomial m we have $P_k(m) = 0$. We can evaluate $P_k(X)$ deterministically using $O(d)$ evaluations of $P(X)$, by evaluating $f_X(z) := P(zx_1, \dots, zx_n)$ for a new variable z and computing the coefficient of z^k in the resulting polynomial $f_X(z)$ using interpolation (Lemma 2.1). Note that by assumption, any monomial m that is multilinear and is a basis for M will be of total degree k , thus we proceed with working over $P_k(X)$.

Introduce a set of variables $Y = \{y_1, \dots, y_k\}$ and define a new polynomial

$$P'(X, Y) = P_k \left(x_1 \sum_{i=1}^k y_i A[i, 1], \dots, x_n \sum_{i=1}^k y_i A[i, n] \right).$$

Let $Q(X, Y)$ be the result of sieving for terms in $P'(X, Y)$ which are of degree at least one in every variable $y_i, i \in [k]$. $Q(X, Y)$ can be computed from 2^k evaluations of P' (hence of P), using the method of inclusion-exclusion in Lemma 2.2. Since the definition of Q is linear over P' , it suffices to consider its effect on a single monomial at a time. Let $m = x_1^{m_1} \dots x_n^{m_n}$ be a monomial in the expansion of P . Let (i_1, \dots, i_k) be the sequence of non-zero indices of m repeated with multiplicity according to degree, in non-decreasing order; e.g., a monomial $x_1^3 x_4^2$ corresponds to sequence $(1, 1, 1, 4, 4)$. In the evaluation of P' , m can be written as a product

$$P(m) \cdot X^m \cdot \prod_{p=1}^k \sum_{j=1}^k y_j A[j, i_p]$$

where $P(m)$ is the coefficient of m in P . Using Lemma 3.1, the contribution of the monomial m to $Q(X, Y)$ is precisely

$$P(m) \cdot X^m \cdot Y \cdot \det A[\cdot, (i_1, \dots, i_k)]$$

where $Y = \prod_{i=1}^k y_i$ and $A' = A[\cdot, (i_1, \dots, i_k)]$ denotes the matrix consisting of columns i_j of A included with multiplicity. Now, if m is not multilinear, then the resulting matrix A' has a repeated column and is clearly singular, so m does not contribute to Q . If m is multilinear, then m contributes a non-zero value to Q if and only if the support of m spans M . Furthermore, since the first part $P(m)X^m$ of this expression is precisely the value of the original monomial m in P , no further algebraic cancellation occurs in Q . Hence Q enumerates monomials corresponding to multilinear monomials in P whose support spans M . The result now follows from a random evaluation of P using Schwartz-Zippel. In particular, Q has degree $2k$ since the sieving started from $P_k(X)$.

For the case that P is homogeneous, we can bypass the phase of extracting $P_k(X)$ and use $P(X)$ directly. The polynomial $Q(X, Y)$ is defined as a sum over 2^k evaluations of $P(X)$ with arguments $x_j \sum_{i=1}^k y_i A[i, j]$, $j \in X$. The precise running time follows with no additional tricks. \square

We note a variant of this. Instead of every variable x_v being associated with only one column v of A , we may wish for each variable to be associated with multiple columns.

Corollary 3.3. *Let $X = \{x_1, \dots, x_n\}$ be a set of variables and let $P(X)$ be a polynomial of degree d over a field \mathbb{F} of characteristic 2. Suppose that M is a linear matroid of rank k over a set V and that each variable x_i is associated with a pairwise disjoint subset $\Gamma_i \subseteq V$ of size γ_i . In time $O^*(d2^k)$ and polynomial space, using evaluation access to P , we can test if the monomial expansion of P contains a multilinear monomial m such that $\bigcup_{i \in \text{supp}(m)} \Gamma_i$ is a basis for M . Our algorithm is randomized with no false positives and failure probability at most $2k/|\mathbb{F}|$.*

Proof. Define a new set of variables $X' = \{x'_{i,v} \mid i \in [n], v \in \Gamma_i\}$, and apply Theorem 3.2 to the polynomial $P'(X')$ resulting from an evaluation where

$$x_i = \prod_{v \in \Gamma_i} x'_{i,v}$$

for every $x_i \in X$. A monomial m in $P(X)$ effectively vanishes if at least one of the following holds: (i) m is not multilinear, (ii) the sets Γ_i for $i \in \text{supp}(m)$ are not pairwise disjoint, or (iii) $\sum_{i \in \text{supp}(m)} \gamma_i \neq k$. All surviving terms are thus multilinear monomials m such that $\bigcup_{i \in \text{supp}(m)} \Gamma_i$ is a basis for M . \square

3.1.2 Sieving for spanning sets

We give the odd sieving algorithm, proving Theorem 1.4. The proof is similar to that of basis sieving. Let us illustrate why only the odd support sets pass through the sieve. To sieve for spanning sets, we basically need to replace each variable x_i with $1 + x_i$. Then, $(1 + x_i)^{m_i} = 1 + m_i x_i + \binom{m_i}{2} x_i^2 + \dots$ (for a monomial m) becomes $1 + m_i x_i$ because only multilinear terms survive, and further reduces to 1 if m_i is even. So a variable with even contributions effectively diminishes. We give the formal proof:

Theorem 3.4 (odd sieving). *Let $P(X)$ be a polynomial over a variable set $X = \{x_1, \dots, x_n\}$ over a field \mathbb{F} of characteristic 2 with degree d . Suppose that M is a linear matroid of rank k over a set V and that each variable x_i is associated with a pairwise disjoint subset $\Gamma_i \subseteq V$ of size γ_i . Given black-box (evaluation) access to a polynomial $P(X)$, we can test in randomized $O^*(2^k)$ time with failure probability at most $\delta = (d + k)/|\mathbb{F}|$ and in polynomial space, whether P contains a term in the monomial expansion of $P(X)$ such that $\Gamma_S = \bigcup_{i \in S} \Gamma_i$ is a basis of M , where $S \subseteq X$ is a subset of its odd support set with $\sum_{i \in S} \gamma_i = k$.*

Proof. We will define a polynomial Q such that it evaluates to non-zero with probability at least $1 - \delta$ if it contains a monomial as stated in the lemma with and to zero otherwise. Let $A \in \mathbb{F}^{k \times m}$ be the matrix representing the linear matroid $M = (X, \mathcal{I})$. For every $i \in [n]$, we define

$$x_i^* = x_i'' (1 + z^{\gamma_i} x_i' \prod_{q \in \Gamma_i} \sum_{p \in [k]} y_p A[p, q]),$$

where x_i' , x_i'' for $i \in [n]$, y_p for $p \in [k]$, and z are new variables. Let $X' = \{x_1', \dots, x_n'\}$, $X'' = \{x_1'', \dots, x_n''\}$, and define a polynomial $Q(X', X'')$ that sieves for those terms in the monomial

expansion of $P^* = P(x_1^*, \dots, x_n^*)$ that contain precisely k contributions of z and which contain y_p for each $p \in [k]$. By Lemmas 2.1 and 2.2, $Q(X', X'')$ can be evaluated using $O^*(2^k)$ evaluations of P .

The expansion in P^* corresponding to m is

$$\begin{aligned} P_m^* &= P(m) \cdot (X'')^m \cdot \prod_{i \in \text{supp}(m)} (1 + z^{\gamma_i} x'_i \prod_{q \in \Gamma_i} \sum_{p \in [k]} y_p A[p, q])^{m_i} \\ &= P(m) \cdot (X'')^m \cdot \sum_{m^*} \prod_{i \in \text{supp}(m^*)} \binom{m_i}{m_i^*} (z^{\gamma_i} x'_i \prod_{q \in \Gamma_i} \sum_{p \in [k]} y_p A[p, q])^{m_i^*}, \end{aligned}$$

where m^* ranges over all monomials that divide m . The last equality is due to the binomial theorem. By Lemma 3.1, the coefficient of $z^k \prod_{i \in [qk]} y_i$ in $P^*(m)$ is thus

$$Q_m = P(m) \cdot (X'')^m \cdot \sum_{m'} \det A_{m'} \prod_{i \in \text{supp}(m')} \binom{m_i}{m'_i} (x'_i)^{m'_i},$$

where m' ranges over all monomials of degree k that divide m and $A_{m'}$ is the $k \times k$ -matrix that contain m'_i copies of $A[\cdot, \Gamma_i]$ for each $i \in \text{supp}(m')$. If $A_{m'_i}$ contains duplicate columns (i.e., $m'_i \geq 2$ for some i), then $\det A_{m'_i} = 0$, and thus we may assume that m' is multilinear. Hence, we obtain

$$Q_m = P(m) \cdot (X'')^m \cdot \sum_{m'} \det A_{m'} \prod_{i \in \text{supp}(m')} m_i x'_i,$$

where m' ranges over all multilinear monomials of degree k that divides m . Since \mathbb{F} has characteristic 2, the summand correspond to m' is non-zero only if $\text{supp}(m')$ is contained in the odd support of m .

On the other hand, for every monomials m and m' such that m' divides m and $A_{m'}$ is nonsingular, there is a term

$$P(m) \cdot \left(\prod_{i \in \text{supp}(m')} m_i \right) (X'')^m \cdot (X')^{m'} \det A_{m'}.$$

Since the variables x'_i and x''_i are newly added variables, this term does not cancel against any other term from the expansion of $Q(m)$. More specifically, these variables uniquely indicate the combination of the monomials m and m' . We evaluate Q for variables x'_i, x''_i randomly chosen from \mathbb{F} . Since Q has degree most $d + k$, by the Schwartz-Zippel lemma, the probability that Q evaluates to zero at most $(d + k)/|\mathbb{F}|$. \square

3.2 Over general fields

We give two sieving algorithms for general fields. First, we present an algorithm for what we call *strongly monotone circuits*—circuits without any cancellation, informally speaking. Our second algorithm works for arbitrary arithmetic circuits albeit with a worse running time.

To sieve over general fields, we use the exterior algebra. For a field \mathbb{F} , $\Lambda(\mathbb{F}^k)$ is a 2^k -dimensional vector space where each basis e_I is defined by a subset $I \subseteq [k]$. Each element $a = \sum_{I \subseteq [k]} a_I e_I$ is called an *extensor*. For $i \in \{0, \dots, k\}$, we denote by $\Lambda^i(\mathbb{F}^k)$ the vector subspace spanned by bases e_I with $|I| = i$. For instance, $\Lambda^0(\mathbb{F}^k)$ is isomorphic to \mathbb{F} and $\Lambda^1(\mathbb{F}^k)$ is isomorphic to the vector space \mathbb{F}^k , so we will use them interchangeably. The addition in $\Lambda(\mathbb{F}^k)$ is defined in the element-wise manner. The multiplication in $\Lambda(\mathbb{F}^k)$ is called *wedge product*, and it is defined as follows: If

$I \cap J \neq \emptyset$, then $e_I \wedge e_J = 0$. If I and J are disjoint, then $e_I \wedge e_J = (-1)^{\sigma(I,J)} e_{I \cup J}$, where $\sigma(I, J) = \pm 1$ is the sign of the permutation that maps the concatenation of I and J each in increasing order into the increasing sequence of $I \cup J$. Over vectors $v, v' \in \mathbb{F}^k$, we have anti-commutativity, i.e., $v \wedge v' = -v' \wedge v$, and in particular, $v \wedge v = 0$. The key property of exterior algebra is that for a matrix $A \in \mathbb{F}^{k \times k}$ with $a_i = A[\cdot, i]$, we have $a_1 \wedge \cdots \wedge a_k = \det A \cdot e_{[k]}$, where $e_{[k]}$ is the basis extensor $e_1 \wedge \cdots \wedge e_k$. For instance, when $k = 2$,

$$\begin{aligned} & (a_{11}e_1 + a_{21}e_2) \wedge (a_{12}e_1 + a_{22}e_2) \\ &= a_{11}a_{12} \cdot e_1 \wedge e_1 + a_{11}a_{22} \cdot e_1 \wedge e_2 + a_{21}a_{12} \cdot e_2 \wedge e_1 + a_{21}a_{22} \cdot e_2 \wedge e_2 \\ &= 0 + a_{11}a_{22} \cdot e_1 \wedge e_2 - a_{21}a_{12} \cdot e_1 \wedge e_2 + 0 = (a_{11}a_{22} - a_{12}a_{21}) \cdot e_1 \wedge e_2. \end{aligned}$$

So a matrix is nonsingular if and only if the wedge product of its columns are nonzero.

An extensor $a \in \Lambda(\mathbb{F}^k)$ is *decomposable* if there are vectors v_1, \dots, v_ℓ such that $a = v_1 \wedge \cdots \wedge v_\ell$. A decomposable extensor a is zero if the vectors v_1, \dots, v_ℓ are linearly dependent. For two decomposable extensors a, a' , it holds that $a \wedge a' = \pm a' \wedge a$ (this is generally not the case, e.g., $e_1 \wedge (e_2 \wedge e_3 + e_4) = e_1 \wedge e_2 \wedge e_3 + e_1 \wedge e_4$ and $(e_2 \wedge e_3 + e_4) \wedge e_1 = e_1 \wedge e_2 \wedge e_3 - e_1 \wedge e_4$).

The sum of two extensors can be computed with 2^k field operations. The wedge product $a \wedge a'$ of two extensors $a \in \Lambda(\mathbb{F}^k)$ and $b \in \Lambda^i(\mathbb{F}^k)$ can be computed with $2^k \binom{k}{i}$ field operations according to the definition (hence $O^*(2^k)$ time for $i \in O(1)$). In general, there is an $O(2^{\omega k/2})$ -time algorithm to compute the wedge product, given implicitly by Włodarczyk [94] (see the thesis of Brand [26] for a more explicit exposition).

Suppose that a polynomial $P(X)$ is represented by an arithmetic circuit C . An *arithmetic circuit* is a directed acyclic graph with a single sink (called output gate) in which every source is labelled by a variable x_i or an element of \mathbb{F} (called input gate) and every other node is labelled by addition (called sum gate) or multiplication (called product gate). We will assume that every sum and product gate has in-degree 2. An arithmetic circuit is called *skew* (δ -skew) if at least one input of every product gate is an input gate (has polynomial degree at most δ , respectively). An arithmetic circuit over the field of rationals \mathbb{Q} is called *monotone* if every constant is nonnegative. We say that an arithmetic circuit (over any field) is *strongly monotone* if the following hold:

- All evaluations are multilinear, which implies that each input to the sum or product gate can be represented as a set family \mathcal{F} over X and coefficients $c: \mathcal{F} \rightarrow \mathbb{F} \setminus \{0\}$.
- For every sum gate with two inputs (\mathcal{F}, c) and (\mathcal{F}', c') , $\mathcal{F} \cap \mathcal{F}' = \emptyset$.
- For every product gate with two inputs (\mathcal{F}, c) and (\mathcal{F}', c') , $S \cup S'$ is distinct for every $S \in \mathcal{F}$ and $S' \in \mathcal{F}'$.

At first glance, the condition for strong monotonicity may seem very restrictive. However, any “cancellation-free” circuit without can be turned into an equivalent strongly monotone circuit, often without blowing up its size, simply by making d copies of sub-circuits for each gate with out-degree $d > 1$. Note that, for every input gate g for the variable x with out-degree d , we will have d input gates each labelled by a new variable say x_i . By associating the variables x_i with one vector, the resulting circuit is essentially equivalent to the original. See e.g., $O^*(2^{qk})$ -time algorithm for q -MATROID PARITY (Theorem 4.3) in Section 4.

Theorem 3.5. *Let C be a strongly monotone arithmetic circuit computing a multilinear polynomial $P(X)$ of degree d over a variable set $X = \{x_1, \dots, x_n\}$ and a field \mathbb{F} . Suppose that M is a linear matroid of rank k over a set V and that each variable x_i is associated with a subset $\Gamma_i \subseteq V$ of size γ_i , and that the subsets Γ_i are pairwise disjoint. We can test in randomized $O^*(2^{\omega k/2})$ time with*

failure probability $d/|\mathbb{F}|$ and in $O^*(2^k)$ space, whether there is a term m in the monomial expansion of $P(X)$ such that $\bigcup_{i \in \text{supp}(m)} \Gamma_i$ is a basis of M . The running time can be improved to $O^*(2^k)$ if C is δ -skew for $\delta \in O(1)$ and $\gamma_i \in O(1)$ for all i .

Proof. Fixing an arbitrary ordering of the input of each product gate (this is necessary because the wedge product is not commutative), we evaluate the circuit C over $\Lambda(\mathbb{F}^k)$ by plugging in the extensor $x_i = x'_i a_i$ for every $i \in [n]$, where x'_i is a new variable and $a_i = \bigwedge_{q \in \Gamma_i} A[\cdot, q]$ (the order of wedge products is not important here). Let $r \in \Lambda(\mathbb{F}^k)$ denote the result. Note that with each variable x'_i substituted by a random element from \mathbb{F} , the extensor r can be computed in time $O^*(2^{\omega k/2})$ (and $O^*(2^k)$ if C is skew and $\max_{i \in [n]} \gamma_i \in O(1)$). We will show that the coefficient of $e_{[k]}$ is nonzero with high probability given that there is a monomial constituting a basis of M .

We show by induction on the number of gates that

$$r = \sum_m \pm P(m) \cdot (X')^m \bigwedge_{i \in \text{supp}(m)} a_i,$$

where m ranges over all monomials m of $P(X)$. For every monomial m , the sign \pm depends on the ordering on product gates. There are two cases depending on whether the last gate g in C is a sum gate or product gate. Let $Q(X) = \sum_{m_Q} Q(m_Q) m_Q$ and $Q'(X) = \sum_{m'_Q} Q'(m'_Q) m'_Q$ denote its input. By the induction hypothesis, suppose that the result of evaluating over the exterior algebra $\Lambda(\mathbb{F}^k)$ is

$$q = \sum_{m_Q} \pm Q(m_Q) \left(\prod_{i \in \text{supp}(m_Q)} x'_i \right) \bigwedge_{i \in \text{supp}(m_Q)} a_i \text{ and } q' = \sum_{m'_Q} \pm Q'(m'_Q) \left(\prod_{i \in \text{supp}(m'_Q)} x'_i \right) \bigwedge_{i \in \text{supp}(m'_Q)} a_i.$$

First, suppose that g is a sum gate. By the strong monotonicity of C , each term m in P corresponding to Q (and Q') has coefficient $Q(m)$ (and $Q'(m)$, respectively). Thus, for every monomial m in P , there is a term $\prod_{i \in \text{supp}(m)} x'_i \bigwedge_{i \in \text{supp}(m)} a_i$ (with the coefficient $\pm Q(m)$ or $\pm Q'(m)$) in r . We stress that the strong monotonicity is crucial here; suppose that Q and Q' share a term with opposite signs. This term should cancel out in P , but it does not necessarily when evaluated over $\Lambda(\mathbb{F}^k)$ as the sign may be flipped.

Next, suppose that g is a product gate. By the strong monotonicity of C , each term m in P corresponding to a pair of monomials, one from $Q(m)$ and the other from $Q'(m)$. We need to verify that for every monomial m of $P(X)$ with $\bigwedge_{i \in \mathcal{S}(m)} a_i \neq 0$, the corresponding terms in $q \wedge q'$ and $q' \wedge q$ have nonzero coefficients. Note that

$$q \wedge q' = \sum_{m_Q, m'_Q} \pm Q(m_Q) Q'(m'_Q) \left(\prod_{i \in \text{supp}(m_Q) \cup \text{supp}(m'_Q)} x'_i \right) \bigwedge_{i \in \text{supp}(m_Q)} a_i \wedge \bigwedge_{i \in \text{supp}(m'_Q)} a_i.$$

Since $\bigwedge_{i \in \text{supp}(m_Q)} a_i$ and $\bigwedge_{i \in \text{supp}(m'_Q)} a_i$ are both decomposable, $\bigwedge_{i \in \text{supp}(m'_Q)} a_i \wedge \bigwedge_{i \in \text{supp}(m_Q)} a_i = \pm \bigwedge_{i \in \text{supp}(m_Q)} a_i \wedge \bigwedge_{i \in \text{supp}(m'_Q)} a_i$, and consequently, $q' \wedge q$ have the same form possibly with opposite signs. In particular, this shows that the induction is correct regardless of how two inputs of product gates are ordered.

Thus, the term corresponding to $e_{[k]}$ in r is

$$\sum_m \pm P(m) \left(\prod_{i \in \text{supp}(m)} x'_i \right) \bigwedge_{i \in \text{supp}(m)} a_i = \sum_m \pm P(m) \cdot (X')^m \cdot \det A[\cdot, \text{supp}(m)] \cdot e_{[k]},$$

where m ranges over all monomials in P with $\sum_{i \in \text{supp}(m)} \gamma_i = k$. Observe that there is no further cancellation between any two terms. We evaluate the coefficient of $e_{[k]}$ in r at random coordinates. By the Schwartz-Zippel lemma, the result follows. \square

We also provide a sieving algorithm for general arithmetic circuits. The idea is again, to evaluate the circuit over the exterior algebra. In order to deal with the issue of non-commutativity, we use the *lift mapping* $\bar{\phi}: \Lambda(\mathbb{F}^k) \rightarrow \Lambda(\mathbb{F}^{2k})$, where $\bar{\phi}(v) = v_1 \wedge v_2$ for $v_1 = (v \ 0)^T, v_2 = (0 \ v)^T \in \mathbb{F}^{2k}$. It has proven useful in similar settings in previous work [25, 27]. The subalgebra generated by the image of $\bar{\phi}$ is commutative, because

$$\bar{\phi}(v) \wedge \bar{\phi}(v') = v_1 \wedge v_2 \wedge v'_1 \wedge v'_2 = v'_1 \wedge v_1 \wedge v_2 \wedge v'_2 = v'_1 \wedge v'_2 \wedge v_1 \wedge v_2 = \bar{\phi}(v') \wedge \bar{\phi}(v).$$

Note that the sign is preserved when the transposition occurs twice. Moreover, if $a_1 \wedge \dots \wedge a_k = \det A \cdot e_{[k]}$, then we have $\bar{\phi}(a_1) \wedge \dots \wedge \bar{\phi}(a_k) = (\det A)^2 \cdot e_{[2k]}$.

Theorem 3.6. *Let C be an arithmetic circuit computing a polynomial $P(X)$ over a variable set $X = \{x_1, \dots, x_n\}$ and a field \mathbb{F} . Suppose that M is a linear matroid of rank k over a set V and that each variable x_i is associated with a subset $\Gamma_i \subseteq V$ of size γ_i , and that the subsets Γ_i are pairwise disjoint. We can test in randomized $O^*(2^{\omega k})$ time with failure probability $k/|\mathbb{F}|$ and in $O^*(4^k)$ space, whether C contains a term m in the monomial expansion of $P(X)$ such that $\bigcup_{i \in \text{supp}(m)} \Gamma_i$ is a basis of M . The running time can be improved to $O^*(4^k)$ if C is δ -skew for $\delta \in O(1)$ and $\gamma_i \in O(1)$ for all i .*

Proof. We evaluate the circuit over the subalgebra of $\Lambda(\mathbb{F}^{2qk})$ by plugging the extensor $x_i = x'_i a_i$, where x'_i is a new variable and $a_i = \bigwedge_{i \in \Gamma_i} \phi(A[\cdot, q])$. Let $r \in \Lambda(\mathbb{F}^{2k})$ denote the result. Note that with each variable x_i substituted with a random element from \mathbb{F} , the extensor r can be computed in time $O^*(2^{\omega k})$ (and $O^*(2^k)$ if C is skew and $\max_{i \in [n]} \gamma_i \in O(1)$). As in the proof of Theorem 3.5, we will show that the coefficient of $e_{[2k]}$ is nonzero with high probability given that there is a monomial constituting a basis of M .

Since the evaluation is over a commutative algebra, for every monomial m in P , there is a “term” in the expansion of r :

$$P(m) \cdot \prod_{i \in \text{supp}(m)} (x'_i)^{m_i} \cdot \bigwedge_{i \in \text{supp}(m)} a_i.$$

It is straightforward to prove by induction as in the proof of Theorem 3.5 that its coefficient is $P(m)$ (without any sign flip). The crucial difference (i.e., no sign flip) arises from the commutativity of the underlying algebra. The term corresponding to $e_{[2k]}$ in r is

$$\sum_m P(m) \left(\prod_{i \in \text{supp}(m)} x'_i \right) \bigwedge_{i \in \text{supp}(m)} a_i = \sum_m P(m) \cdot (X')^m \cdot (\det A[\cdot, \text{supp}(m)])^2 \cdot e_{[2k]},$$

where m ranges over all monomials in P with $\sum_{i \in \text{supp}(m)} \gamma_i = k$. Note that the determinant is squared due to the lift mapping. Also, observe that there is no cancellation between any two terms. We evaluate the coefficient of $e_{[2k]}$ in r at random coordinates. By the Schwartz-Zippel lemma, the result follows. \square

4 Matroid Covering, Packing and Intersection Problems

For our first application section, we review some fairly straightforward results regarding matroid variants of the SET COVER and SET PACKING problems defined in Section 1 and related problems. We start off by recalling the definitions.

SET COVER and SET PACKING are classical NP-hard problems. For both problems, the input is a ground set V , a set system $\mathcal{E} \subseteq 2^V$ over V , and an integer t . SET COVER asks whether there is a subcollection $S \subseteq \mathcal{E}$ such that $|S| \leq t$ and $\bigcup S = V$, i.e., S covers V , and SET PACKING asks whether there is a subcollection $S \subseteq \mathcal{E}$ of t pairwise disjoint sets. Their matroid variants RANK k SET COVER and RANK k SET PACKING are defined as follows. We are given as input a set V , a set family $\mathcal{E} \subseteq 2^V$, a matroid $M = (V, \mathcal{I})$ of rank k , and an integer t . In RANK k SET COVER, the question is whether there is a subcollection $S \subseteq \mathcal{E}$ with $|S| \leq t$ such that $\bigcup S$ has rank k . RANK k SET PACKING asks for a subcollection S of pairwise disjoint t sets such that $\bigcup S$ has rank k (i.e., it is a basis of M). Note that SET COVER is the special case of RANK k SET COVER where M is the free matroid, i.e., all subsets of V are independent, and $k = |V|$. Similarly, SET PACKING is the special case of RANK k SET PACKING where M is the uniform matroid of rank $k = |\bigcup S|$ for a solution S . One may also consider the apparently more general variant of RANK k SET PACKING where one does not require that $\bigcup S$ is a basis for M , but only that it is independent. However, this reduces to RANK k SET PACKING by iterating over the acceptable cardinalities $|\bigcup S|$ and applying matroid truncation.

For $q \in O(1)$, the q -SET PACKING problem is SET PACKING in which every set has cardinality q . The q -DIMENSIONAL MATCHING problem is a well-studied special case of q -SET PACKING, where V is partitioned into q sets V_1, \dots, V_q , and every set in \mathcal{E} is from $V_1 \times \dots \times V_q$. The matroid analogs to q -DIMENSIONAL MATCHING and q -SET PACKING are q -MATROID INTERSECTION and q -MATROID PARITY, respectively. In q -MATROID INTERSECTION, we are given as input q matroids M_1, \dots, M_q over the same ground set V and an integer k , and the question is whether there is a subset $S \subseteq V$ of size k that is independent in M_i (equivalently, a basis for M_i by applying matroid truncation) for every $i \in [q]$. In q -MATROID PARITY, we are given as input a matroid $M = (V, \mathcal{I})$ with V partitioned into disjoint sets $\mathcal{E} = \{E_1, \dots, E_m\}$ each of size q , and an integer k , the question is whether a collection S of k sets from V_1, \dots, V_m , such that $\bigcup S$ has rank qk in M . The problems q -MATROID PARITY and q -MATROID INTERSECTION generalize q -SET PACKING (when M is the uniform matroid) and q -DIMENSIONAL MATCHING (when M_i is the partition matroid over V_i), respectively.

We survey the known results for these problems. The fastest known algorithm for SET COVER in terms of $n = |V|$ is $O^*(2^n)$, which can be achieved either via classical dynamic programming or by inclusion-exclusion. It is a major open problem whether this can be improved; Cygan et al. [35] propose the *Set Cover Conjecture* (SeCoCo) that effectively conjectures that this is not possible, analogous to the more commonly used strong exponential-time hypothesis (SETH). More precisely, SeCoCo states that for every $\varepsilon > 0$ there exist $d \in \mathbb{N}$ such that SET COVER on n elements where all sets of size at most d cannot be solved in $O^*(2^{(1-\varepsilon)n})$ time [35]. The currently known fastest algorithms for q -DIMENSIONAL MATCHING and q -SET PACKING are by Björklund et al. [19]. (The reader is referred to [19] for a series of previous improvements on these problems e.g., [31, 65, 66].) Their running time bounds are $O^*(2^{(q-2)k})$ and $O^*(2^{(q-\varepsilon_q)k})$, respectively, where $\varepsilon_q < 2$ is a constant depending on q , tending to zero as $q \rightarrow \infty$. Finally, the fastest known algorithms for q -MATROID INTERSECTION and q -MATROID PARITY run in time $O^*(4^{qk})$ [28].

4.1 Rank k Set Cover and Rank k Set Packing.

We start with RANK k SET COVER, reiterating Theorem 1.7. We will assume that the solution size is exactly t . We will use the polynomial-space sieving algorithm (Theorem 3.4) if the underlying field has characteristic 2, and the sieving algorithm for strongly monotone circuits (Theorem 3.5) otherwise. To that end, we construct a polynomial as follows. Let $X = \{x_{v,E} \mid v \in V, E \in \mathcal{E}\}$ and

$Y = \{y_{i,E} \mid i \in [t], E \in \mathcal{E}\}$ be a set of variables. For RANK k SET COVER, we define

$$\begin{aligned} P(X, Y) &= \prod_{i \in [t]} \sum_{E \in \mathcal{E}} y_{i,E} \prod_{v \in E} (1 + x_{v,E}) \\ &= \sum_{f: [t] \rightarrow \mathcal{E}} \prod_{i \in [t]} y_{i,f(i)} \prod_{v \in f(i)} (1 + x_{v,f(i)}) = \sum_{f: [t] \rightarrow \mathcal{E}} \sum_{\substack{E_1, \dots, E_t \\ E_i \subseteq f(i), i \in [t]}} \left(\prod_{i \in [t]} y_{i,f(i)} \right) \left(\prod_{i \in [t]} \prod_{v \in E_i} x_{v,f(i)} \right) \end{aligned}$$

For RANK k SET PACKING, we tweak the polynomial slightly:

$$P(X, Y) = \prod_{i \in [t]} \sum_{E \in \mathcal{E}} y_{i,E} \prod_{v \in E} x_{v,E} = \sum_{f: [t] \rightarrow \mathcal{E}} \prod_{i \in [t]} y_{i,f(i)} \prod_{v \in f(i)} x_{v,f(i)}.$$

Note that the function $f: [t] \rightarrow \mathcal{E}$ plays the role of choosing t sets from \mathcal{E} . For every $f: [t] \rightarrow \mathcal{E}$, there is a distinct monomial and thus no further algebraic cancellation occurs. Let $A \in \mathbb{F}^{k \times V}$ be the linear representation of M . We may assume that A has exactly k rows by truncating M . Let $P'(X)$ be the result of substituting every variable $y_{i,E}$ with a uniformly chosen random element from \mathbb{F} . Mapping every variable $x_{v,E}$ to the column vector $A[:, v]$, we use the sieving algorithm. If \mathbb{F} has characteristic 2, then Theorem 3.4 gives an $O^*(2^k)$ -time algorithm. Otherwise, we use Theorem 3.5. Note that $P'(X)$ can be realized by a strongly monotone circuit. Thus, we have:

Theorem 4.1 (Restatement of Theorem 1.7). *RANK k SET COVER for matroids represented over a field \mathbb{F} can be solved in $O^*(2^k)$ time and polynomial space if \mathbb{F} has characteristic 2 and in $O^*(2^{\omega k/2})$ time and $O^*(2^k)$ space in general.*

Theorem 4.2. *RANK k SET PACKING for matroids represented over a field \mathbb{F} can be solved in $O^*(2^k)$ time and polynomial space if \mathbb{F} has characteristic 2 and in $O^*(2^{\omega k/2})$ time and $O^*(2^k)$ space in general.*

4.2 q -Matroid Parity and q -Matroid Intersection

Next, we discuss q -MATROID PARITY. Let $X = \{x_E \mid E \in \mathcal{E}\}$ be a set of variables. We define a polynomial:

$$P(X) = \prod_{E \in \mathcal{E}} (1 + x_E) = \sum_{j \in [|\mathcal{E}|]} \sum_{S \in \binom{\mathcal{E}}{j}} \prod_{E \in S} x_E.$$

We apply the sieving algorithm by associating every x_E with q columns $A[:, E]$. To sieve over general fields, observe that the polynomial $P(X)$ can be computed using a 1-skew strongly monotone circuit. Using the basis sieving algorithm (Cor. 3.3 and Theorem 3.5), we obtain:

Theorem 4.3. *q -MATROID PARITY for matroids represented over a field \mathbb{F} can be solved in $O^*(2^{qk})$ time (and polynomial space if \mathbb{F} has characteristic 2).*

Since q -MATROID INTERSECTION is a special case of q -MATROID PARITY, we also obtain:

Corollary 4.4. *q -MATROID INTERSECTION for matroids represented over a field \mathbb{F} can be solved in $O^*(2^{qk})$ time (and polynomial space if \mathbb{F} has characteristic 2).*

We obtain a greater speedup for q -MATROID INTERSECTION by using the Cauchy-Binet formula. Suppose that $A_i \in \mathbb{F}^{k \times V}$ represents the matroid M_i . Let $X = \{x_v \mid v \in V\}$ be a set of variables and let A'_1 be the result of scaling every column v of A_1 by x_v . By the Cauchy-Binet formula,

$$P(X) := \det(A'_1 A_2^T) = \sum_{B \in \binom{V}{k}} \det A_1[\cdot, B] \det A_2[\cdot, B] \prod_{v \in B} x_v.$$

Thus $P(X)$ enumerates monomials $\prod_{v \in B} x_v$ for common bases B of A_1 and A_2 , and we only have to sieve for terms that in addition are bases of the remaining $q - 2$ matroids. We get the following.

Theorem 4.5 (Restatement of Theorem 1.9). *q -MATROID INTERSECTION for linear matroids represented over a common field \mathbb{F} of characteristic 2 can be solved in randomized time $O^*(2^{(q-2)k})$ and polynomial space.*

For fields of characteristic other than 2, this does not represent a speedup over Cor. 4.4 since the circuit computing $\det A'_1 A_2^T$ is not strongly monotone. However, we do obtain a speedup for general \mathbb{F} for the special case $q = 3$. Observe that every entry in $A'_1 A_2^T$ has polynomial degree at most 1. It is known that the determinant of a symbolic matrix can be computed with a skew circuit [74]. Thus, there is a 1-skew circuit computing $\det(A'_1 A_2^T)$. Using the sieving algorithm of Theorem 3.6 for general arithmetic circuits, we obtain:

Theorem 4.6. *q -MATROID INTERSECTION for linear matroids can be solved in $O^*(4^{(q-2)k})$ time. In particular, the bound is $O^*(4^k)$ for $q = 3$.*

It is an interesting open question whether q -MATROID PARITY can be solved in $O^*(2^{(q-\varepsilon)k})$ for $\varepsilon > 0$ when $q \geq 3$ is constant. Note that an enumerating polynomial for 2-matroid parity (let us call it *matroid matching* for clarity) can be efficiently evaluated using the linear representation of Lovász [73]: Suppose that A represents a matroid $M = (V, \mathcal{I})$ with V partitioned into pairs $P_i = \{v_i, v'_i\}$. If x_i is a variable representing the pair P_i , then the Pfaffian $\text{Pf } B$, where

$$B = \sum_i x_i (A[\cdot, v_i] A^T[v'_i, \cdot] - A[\cdot, v'_i] A^T[v_i, \cdot]),$$

enumerates all matroid matching terms. Lovász [73] only showed that the rank of B equals twice the maximum matroid matching size, but $\text{Pf } B$ indeed enumerates all matroid matching terms. We refer to the textbook of Murota [81, Section 7.3.4] for this fact (the exposition concerns an alternative equivalent formulation of matroid matching proposed by Geelen and Iwata [53]). The trick employed by Björklund et al. [19] to speed up q -SET PACKING of “reducing” (via colour-coding type arguments) to q -DIMENSIONAL MATCHING, however, seemingly does not work for the matroid analogs. The simple idea of having the variable x_i encode $q - 2$ columns in the matroid matching enumerating polynomial fails because the space spanned by vectors in the matroid matching is not necessarily orthogonal to the other of $q - 2$ columns.

4.3 Odd Coverage

Finally, let us discuss another corollary of Theorem 3.4 on a variant of SET COVER, called ODD COVERAGE. The input is a set family \mathcal{E} over V and integers t, p . The question is whether there is a subcollection $S \subseteq \mathcal{E}$ with $|S| = t$ such that there are at least p elements $v \in V$ with $|\{E \in S \mid v \in E\}| \bmod 2 = 1$ (i.e., v is covered an odd number of times). Over a set of variables $X = \{x_v \mid v \in V\}$

and $Y = \{y_E \mid E \in \mathcal{E}\}$, define

$$P(X, Y, z) = \prod_{E \in \mathcal{E}} \left(1 + zy_E \prod_{v \in E} x_v \right).$$

The coefficient of z^t then enumerates the subcollections of size t . Note that there is a solution if and only if there is a monomial (over X) whose odd support set is size at least p . Thus, the odd sieving algorithm implies:

Theorem 4.7. ODD COVERAGE can be solved in $O^*(2^p)$ time and polynomial space.

An $O^*(2^p)$ -time (and exponential-space) algorithm for a special case is known, given by Saurabh and Zehavi [87]. They studied the following problem: given a graph $G = (V, E)$ and integers t, p , is there a set S of exactly t vertices such that there are at least p edges with one endpoint in S and the other in $V \setminus S$? Note that this is a special case of ODD COVERAGE in which every element occurs in two sets.

5 Balanced Solution and Diverse Collection

As noted, given an efficient enumerating polynomial $P(X)$ for a category of objects, and given a representable matroid M over X , we can use our methods out-of-the-box to sieve for objects in the collection that are independent or spanning in M . In this section, we survey two applications. The first concerns the problem of finding a *balance-fair* solution. A balanced-fairness is, in a way, a stronger notion of colourfulness; every colour should appear not only once, but also almost equally frequently. We note that with an efficient enumerating polynomial at hand, our sieving algorithm can find a balanced-fair solution. The second addresses another problem category, of finding a *diverse* collection of objects, with prescribed pairwise minimum distances. Utilizing the odd sieving method (Theorem 3.4), we show a general way to find a diverse collection.

5.1 Balance-fair X paradigm

There is a recent trend in pursuing fairness especially in artificial intelligence applications (see e.g., the work of Chierichetti et al. [32]). There are many notions of fairness known in the literature. Here, we consider the problem of finding a *balanced* solution. We assume that every object is assigned a colour from a set C . For $\alpha \leq \beta \in \mathbb{N}$, a set S of objects is said to be (α, β) -balanced if $\alpha \leq |S_c| \leq \beta$ for every colour $c \in C$, where $S_c \subseteq S$ denotes the objects in S with colour c . The problem of finding a balanced solution has been studied in the context of MATROID INTERSECTION [33], k -MATCHING [5], and k -PATH [14]. We define a general problem called BALANCED SOLUTION as follows. The input is a set E , a collection of (possibly exponentially many) subsets $\mathcal{F} \in 2^E$ of E , a set of colours C , a colouring $\chi: E \rightarrow C$, and integers k, α, β . The question is whether there is a set $S \in \mathcal{F}$ of size k such that $\alpha \leq |S \cap \chi^{-1}(c)| \leq \beta$. We observe that the basis sieving (Theorem 3.2) solves this problem in time $O^*(2^k)$, if an enumerating polynomial for \mathcal{F} can be evaluated in polynomial time. To set up the matroid constraint, we can use the observation of Bentert et al. [14] that there is a linear matroid M of rank k with coloured objects as its ground set such that a set of k objects is (α, β) -balanced if and only if it is a basis for M . We thus obtain from the definitions:

Theorem 5.1. BALANCED SOLUTION can be solved in $O^*(2^k)$ time if there is an enumerating polynomial for \mathcal{F} that can be efficiently evaluated.

In particular, this implies $O^*(2^k)$ -time algorithms for balanced-fair variants of MATROID INTERSECTION, k -MATCHING, and k -PATH (see Section 2.2 for the enumerating polynomials). In particular, for k -PATH we use the enumerating polynomial for k -walks, and give all copies $x_{v,i}$ for a vertex v the same label in the matroid M , thereby ensuring that any surviving monomial represents a path. This is an improvement over the existing algorithms, all of which run in $O^*(2^{ck})$ time for some $c > 1$.

5.2 Diverse X paradigm

In the so-called “diverse X paradigm” (X being the placeholder for an optimization problem), we seek—rather than a single solution—a diverse collection of solutions, where the diversity is measured in terms of the Hamming distance, i.e., the size of the symmetric difference. Recently, there is an increasing number of publications studying the problem of finding diverse solutions from the parameterized complexity perspective [11, 12, 46, 48, 60].

The DIVERSE COLLECTION problem is defined as follows. For a set E , let \mathcal{F}_i be a collection of (potentially exponentially many) subsets of E for each $i \in [k]$. Given $d_{i,j} \in \mathbb{N}$ for $i < j \in [k]$, the problem asks to determine the existence of subsets $S_i \in \mathcal{F}_i$ for $i \in [k]$ such that $|S_i \Delta S_j| \geq d_{i,j}$ for each $i < j \in [k]$. Here, $S_i \Delta S_j$ denotes the symmetric difference $(S_i \setminus S_j) \cup (S_j \setminus S_i)$. We show that if all collections \mathcal{F}_i admit enumerating polynomials $P_i(X)$ that can be efficiently evaluated, then DIVERSE COLLECTION can be solved in $O^*(2^D)$ time, where $D = \sum_{i < j \in [k]} d_{i,j}$.

Let $X' = \{x_e^{\{i,j\}} \mid i, j \in [k], e \in E\}$ and $Y = \{y_{i,e} \mid i \in [k], e \in E\}$ be variables. We define

$$P(X', Y) = \prod_{i \in [k]} P'_i(X', Y)$$

where $P'_i(X', Y)$ is the result of plugging $x_e = y_{i,e} \prod_{j \in [k] \setminus \{i\}} x_e^{\{i,j\}}$ in the enumerating polynomial $P_i(X) = \sum_{S_i \in \mathcal{F}_i} c(i, S_i) \prod_{e \in S_i} x_e$ for coefficients $c(i, S_i) \in \mathbb{F}$. The variables $x_e^{\{i,j\}}$ will play a key role in ensuring that $|S_i \Delta S_j| \geq d_{i,j}$. Let us expand $P(X', Y)$ into a sum of monomials:

$$P(X', Y) = \sum_{i \in [k], S_i \in \mathcal{F}_i} \prod_{i \in [k]} c(i, S_i) \cdot \prod_{i \in [k], e \in S_i} y_{i,e} \cdot \prod_{i < j \in [k]} \prod_{e \in S_i \Delta S_j} x_e^{\{i,j\}}$$

With $S_i \in \mathcal{F}_i$ fixed for each $i \in [k]$, we have

$$\prod_{i \in [k], e \in S_i} \prod_{j \in [k] \setminus \{i\}} x_e^{\{i,j\}} = \prod_{i < j \in [k]} \prod_{e \in S_i} x_e^{\{i,j\}} \prod_{e \in S_j} x_e^{\{i,j\}} = \prod_{i < j \in [k]} \prod_{e \in S_i \Delta S_j} x_e^{\{i,j\}} \prod_{e \in S_i \cap S_j} (x_e^{\{i,j\}})^2.$$

We therefore have

$$P(X', Y) = \sum_{i \in [k], S_i \in \mathcal{F}_i} \prod_{i \in [k]} c(i, S_i) \cdot \prod_{i \in [k], e \in S_i} y_{i,e} \cdot \prod_{i < j \in [k]} \prod_{e \in S_i \Delta S_j} x_e^{\{i,j\}} \prod_{e \in S_i \cap S_j} (x_e^{\{i,j\}})^2.$$

For every collection of k -tuples (S_1, \dots, S_k) with $S_i \in \mathcal{F}_i$, there is a distinct monomial in $P(X', Y)$. We use the odd sieving algorithm of Theorem 3.4. More precisely, we add constraints such that for every $\{i, j\} \subseteq [k]$, there are at least $d_{i,j}$ variables $x_e^{\{i,j\}}$ in the odd support set. This ensures that each pairwise Hamming distance is at least $d_{i,j}$. Note that these constraints can be realized in a matroid of rank D by taking the direct sum of $\binom{k}{2}$ partition matroids each of rank $d_{i,j}$. Thus, we obtain:

Theorem 5.2. DIVERSE COLLECTION can be solved in $O^*(2^D)$ time if all collections \mathcal{F}_i admit enumerating polynomials that can be efficiently evaluated.

Remark 5.3. Our approach can be adapted to solve the weighted variant considered by Fomin et al. [48]. For the weighted variant, every element e has a positive weight $w_e \in \mathbb{N}$, and we require S_i and S_j to have $\sum_{e \in S_i \Delta S_j} w_e \geq d_{i,j}$, rather than $|S_i \Delta S_j| \geq d_{i,j}$. To deal with weights, simply replace each variable $x_e^{\{i,j\}}$ with the product of w_e variables $x_{e,1}^{\{i,j\}} x_{e,2}^{\{i,j\}} \cdots x_{e,w_e}^{\{i,j\}}$.

Remark 5.4. A variant of DIVERSE COLLECTION where we wish to maximise the sum of all pairwise Hamming distances (that is, $\sum_{i < j \in [k]} |S_i \Delta S_j| \geq D_+$) are also studied in the literature [11, 12, 59, 60]. A similar approach yields an FPT algorithm with running time $O^*(2^{D_+})$. Using the same polynomial $P(X', Y)$, we require that there should be at least D_+ variables $x_e^{\{i,j\}}$ in the odd support set. Obviously, this can be done using a uniform matroid of rank D_+ . Thus, the sieving algorithm of Theorem 3.4 gives an $O^*(2^{D_+})$ -time algorithm.

We discuss several corollaries of Theorem 5.2. First, we consider DIVERSE PERFECT MATCHINGS: we are given an undirected graph G , an integer k , and $\binom{k}{2}$ integers $d_{i,j}$ for $i < j \in [k]$, and we want to find k perfect matchings M_1, \dots, M_k with $|M_i \Delta M_j| \geq d_{i,j}$ for every $i < j \in [k]$. Let $d = d_{1,2}$ and $D = \sum_{i < j \in [k]} d_{i,j}$. This problem is NP-hard even for $k = 2$ [61]. Fomin et al. [46] gave an $O^*(4^d)$ -time algorithm for the special case $k = 2$. Later, Fomin et al. [48] proved that DIVERSE PERFECT MATCHINGS is FPT for the case $d_{i,j} = d$ for all $i < j \in [k]$, giving an algorithm running in time $O^*(2^{2^{O(dk)}})$. Since the pfaffian is an enumerating polynomial for perfect matchings, we obtain:

Corollary 5.5. DIVERSE PERFECT MATCHINGS can be solved in $O^*(2^D)$ time.

Our approach also works for diverse matroid problems DIVERSE BASES and DIVERSE COMMON INDEPENDENT SETS, which were introduced by [48]. In DIVERSE BASES, we are given a matroid M and $k, d_{i,j} \in \mathbb{N}$ for $i < j \in [k]$, and the question is whether M has bases B_1, \dots, B_k such that $|B_i \Delta B_j| \geq d_{i,j}$ for all $i < j \in [k]$. In DIVERSE COMMON INDEPENDENT SETS, we are given two matroids and $k, d_{i,j} \in \mathbb{N}$ for $i < j \in [k]$, and the question is whether a collection of sets I_1, \dots, I_k that are independent in both matroids such that $|I_i \Delta I_j| \geq d_{i,j}$ for all $i < j \in [k]$. The previous known algorithms of Fomin et al. [48] solve DIVERSE BASES and DIVERSE COMMON INDEPENDENT SETS in time $O^*(2^{O(k^2 d \log kd)})$ and $O^*(2^{O(k^3 d^2 \log kd)})$, respectively when $d_{i,j} = d$.

Corollary 5.6. DIVERSE BASES on linear matroids can be solved in $O^*(2^D)$ time.

Corollary 5.7. DIVERSE COMMON INDEPENDENT SETS on linear matroids can be solved in $O^*(2^D)$ time.

Theorem 5.2 also has an implication for the k -DISTINCT BRANCHING problem. Its input is a directed graph G , two vertices s and t , and an integer k . The problem asks whether G admits an out-branching (V, B_s^+) rooted at s and in-branching (V, B_t^-) rooted at t such that $|B_s^+ \Delta B_t^-| \geq k$. The NP-hardness is even for $s = t$ and $k = 2n - 2$ [6]. Since Bang-Jensen and Yeo [10] asked whether k -DISTINCT BRANCHING is FPT for $s = t$, this problem has been studied in parameterized complexity. We briefly survey the history here. Bang-Jensen et al. [9] gave an FPT algorithm for strongly connected graphs. Later, Gutin et al. [57] showed that k -DISTINCT BRANCHING on arbitrary directed graphs can be solved in $O^*(2^{O(k^2 \log^2 k)})$ time for $s = t$. Very recently, Bang-Jensen et al. [8] designed an $O^*(2^{O(k \log k)})$ -time algorithm. They asked whether k -DISTINCT BRANCHINGS can be solved in $O^*(2^{O(k)})$ time. As a corollary of Theorem 5.2, we answer this question in the affirmative. Recall that the determinant of the symbolic Laplacian matrix yields an enumerating polynomial for out-branchings and for in-branchings by reversing arcs (see [21, 54]). Thus, Theorem 5.2 implies:

Corollary 5.8. k -DISTINCT BRANCHINGS can be solved in $O^*(2^k)$ time.

6 Path, cycle and linkage problems

One of the main application areas of algebraic algorithms in parameterized complexity is for path and cycle problems. Indeed, one of the earliest examples of an algebraic FPT algorithm was for k -PATH, finding a path on k vertices in a possibly directed graph, ultimately improved to time $O^*(2^k)$ [65, 93, 66]. Another breakthrough result in the area is Björklund’s algorithm for HAMILTONICITY, finding a Hamiltonian path in an undirected graph, in time $O^*(1.66^n)$ [16], and more generally solving k -PATH in undirected graphs in time $O^*(1.66^k)$ [19]. In fact, even the apparently simple question of k -PATH, k -CYCLE and HAMILTONICITY problems remains a highly active area of research. This is particularly true in *directed* graphs; however, in this section we restrict ourselves to undirected graphs. We also restrict ourselves solely to matroids represented over fields of characteristic 2, since we need the power of the odd support sieving method (Theorem 3.4).

Another, subtly different problem is to find a cycle of length *at least* k , which we refer to as the LONG CYCLE problem. Unlike the corresponding “LONG PATH” problem, being able to find a k -cycle in time $O^*(c^k)$ does not guarantee being able to solve LONG CYCLE in the same time. On directed graphs, the first algorithm for LONG CYCLE with running time $O^*(2^{O(k)})$ was given by Fomin et al. [50] using the *representative families* approach to algorithm design (cf. Sections 1.2.5 and 8); the current record is $O^*(4^k)$ by Zehavi [95]. For undirected graphs, the currently fastest algorithm for LONG CYCLE is by reduction to the more general LONG (s, t) -PATH problem. Note that, again unlike unrooted LONG PATH, asking for an st -path of length at least k is a sensible question that does not trivially reduce to k -PATH. In turn, the fastest algorithm for LONG st -PATH is by Fomin et al. [47] in time $O^*(2^k)$; see below.

In a different direction, in the problem T -CYCLE (a.k.a. K -CYCLE), the input is an undirected graph G and a set of vertices $T \subseteq V(G)$, and the question is whether there is a simple cycle in G that visits every vertex in T . As mentioned in the introduction, this problem was known to be FPT using an algorithm working over heavy graph structural methods [64], and it was a major surprise when Björklund, Husfeldt and Taslamán [20] showed an $O^*(2^{|T|})$ -time algorithm based on polynomial cancellations. Specifically, they defined a polynomial, roughly corresponding to walks without U-turns, and showed in an intricate argument that an $O^*(2^{|T|})$ -time sieving step over this polynomial tests for T -cycles in G . Wahlström [92] adapted Björklund’s *determinant sums* method [16] to the T -CYCLE problem and thereby showed that it even allows for a *polynomial compression*, i.e., a reduction in polynomial time to an object of size $|T|^{O(1)}$ from which the existence of a T -cycle can be decided.

Recently, Fomin et al. [47] considered problems pushing the envelope on the method of Björklund, Husfeldt and Taslamán [20], showing more involved cancellation-based algorithms for more general path and cycle problems, and also extending the scope to *linkages*. Let $G = (V, E)$ be a graph, and $S, T \subseteq V$ be vertex sets. An (S, T) -*linkage* in G is a set \mathcal{P} of pairwise vertex-disjoint (S, T) -paths. The *order* of the linkage is $p = |\mathcal{P}|$. We say the linkage is *perfect* if $|\mathcal{P}| = |S| = |T|$. Let the COLOURFUL (S, T) -LINKAGE problem refer to the following question. Let $G = (V, E)$, $S, T \subseteq V$ and an integer k be given. Furthermore, let $c: V \rightarrow [n]$ be a not necessarily proper vertex colouring, also given as input. Then the question is: Does G contain a perfect (S, T) -linkage using vertices of at least k colours? (More generally, one may ask of an (S, T) -linkage of order p , but this is essentially equivalent as we can create new sets S' and T' of p vertices each, and connect them to S and T .) Fomin et al. showed, using complex polynomial cancellation arguments, that COLOURFUL (S, T) -LINKAGE can be solved in time $O^*(2^{k+p})$ where $p = |S| = |T|$ [47]. We show the following

improvement.

Theorem 6.1. COLOURFUL (S, T) -LINKAGE for undirected graphs can be solved in randomized time $O^*(2^k)$ and polynomial space.

As Fomin et al. note, even the problem COLOURFUL (s, t) -PATH (being the case where $|S| = |T| = 1$) has a multitude of applications. Among others, this implies solving LONG (s, t) -PATH and LONG CYCLE in time $O^*(2^k)$ – i.e., in an undirected graph, find an (s, t) -path, respectively a cycle, of length *at least* k in time $O^*(2^k)$, and T -CYCLE in time $O^*(2^{|T|})$, among other results. All of these improve on or match the previous state of the art.

Fomin et al. also consider the more general setting of *frameworks*. Let $G = (V, E)$ be an undirected graph and $M = (V, \mathcal{I})$ a matroid over the vertex set of G . Let $S, T \subseteq V$ and let k be an integer. Fomin et al. show that if M is represented over a finite field of order q , then an (S, T) -linkage of rank at least k in M can be found in time $O^*(2^{p+O(k^2 \log(k+q))})$ [47]. We note that if M is represented over a field of characteristic 2, then we get a significant speedup over their algorithm.

Theorem 6.2. Given an undirected graph $G = (V, E)$, a matroid M over V represented over a field of characteristic 2, sets $S, T \subseteq V$ and an integer k , in randomized time $O^*(2^k)$ and polynomial space we can find a perfect (S, T) -linkage in G which has rank at least k in M .

Theorem 6.1 follows from Theorem 6.2 by an appropriate matroid construction. Let $M = (V, \mathcal{I})$ be the linear matroid with a representation where each vertex $v \in V$ is associated with the $c(v)$ -th n -dimensional unit vector $e_{c(v)}$. Then a linkage has rank at least k if and only if it visits vertices of at least k different colours.

We note that *directed* variants of the above results are excluded, as it is NP-hard to find a directed (s, t) -path with even two distinct colours (see Fomin et al. [47]).

Finally, Fomin et al. [47] ask as an open question whether LONG (s, t) -PATH and LONG CYCLE can be solved in $O((2 - \varepsilon)^k)$ for any $\varepsilon > 0$. We show this in the affirmative, giving an algorithm for both problems that matches the running time for UNDIRECTED HAMILTONICITY. Our algorithm is a mild reinterpretation of the *narrow sieves* algorithm for k -PATH [19], rephrased in terms of an external matroid labelling the vertices of G .

Theorem 6.3. LONG CYCLE and LONG (s, t) -PATH can be solved in randomized time $O^*(1.66^k)$ and polynomial space.

All of the above theorems follow from the same underlying enumerating polynomial result. At the heart of the HAMILTONICITY algorithm of Björklund [16], and the polynomial compression for T -CYCLE of Wahlström [92], is the result that given a graph $G = (V, E)$ and $s, t \in V$, there is a particular *almost* symmetric matrix A_{st} such that $\det A_{st}$ effectively enumerates (s, t) -paths, with some additional “padding” terms (see below). We note that this statement can be generalized to linkages: Given $G = (V, E)$ and $S, T \subseteq V$ there is a matrix A_{ST} such that $\det A_{ST}$ enumerates padded perfect (S, T) -linkages. Furthermore, the “padding” is compatible with the odd sieving approach of Theorem 3.4. We review this construction next.

6.1 The linkage-generating determinant

We now present the algebraic statements that underpin the algorithms in this section. Like the rest of the paper, these algorithms are based on algebraic sieving over a suitable enumerating polynomial. Here, we present this polynomial, in the form of a *linkage-enumerating determinant*.

6.1.1 Path enumeration

We begin with the simpler case of enumerating st -paths. This result is from Wahlström [92], repeated for completeness, but is also implicitly present in Björklund [16]. We note st -path enumeration is still far from a trivial conclusion, since we want to enumerate only *paths* without also enumerating st -walks. Indeed, there is a catch, since otherwise we could solve HAMILTONICITY in polynomial time by searching for an st -path term of degree n . Specifically, we generate *padded* st -paths, which is a union of st -paths and 2-cycles; details follow.

Let $G = (V, E)$ be an undirected graph and $s, t \in V$ be vertices. We show that a modified Tutte matrix of G can be used to produce a polynomial that effectively enumerates st -paths in G . This was previously used in the polynomial compression for the T -cycle problem [92].

Let $X = \{x_e \mid e \in E\}$ be a set of edge variables. Let P be an st -path in G and define

$$X(P) = \prod_{e \in E(P)} x_e.$$

A *2-cycle term* over (G, X) is a term x_e^2 for some $e \in E$; note that as a polynomial, if $e = uv$ then this term corresponds to the closed walk uvu in G . A *padded st -path term* for an st -path P is a term

$$X(P) \cdot \prod_{e \in M} x_e^2,$$

where M is a (not necessarily perfect) matching of $G - V(P)$. We assume by edge subdivision that $st \notin E$.

Lemma 6.4. *There is a matrix A_{st} over a field of characteristic 2 such that $\det A_{s,t}$ enumerates padded st -path terms.*

Proof. Let A be the Tutte matrix of G over a sufficiently large field of characteristic 2. Define A_{st} by letting $A[v, v] = 1$ for every $v \in V \setminus \{s, t\}$, $A[s, t] = 0$, $A[t, s] = 1$ and $A[t, v] = 0$ for every $v \in V - s$. We claim that $\det A_{st}$ enumerates padded st -path terms as described. This follows from arguments in Wahlström [92]. Viewing the rows and columns of A_{st} as vertices of G , each term of $\det A_{st}$ can be viewed as an *oriented cycle cover* of G , i.e., a partition of V into oriented cycles (which may include cycles of length 1 where a diagonal entry of A_{st} is used). Due to the modifications made to A_{st} above, t has s as its unique out-neighbour in every oriented cycle cover, and for every vertex $v \in V \setminus \{s, t\}$ the loop term on v can be used in the cycle cover. Furthermore, every other edge of the graph is bidirected (i.e., symmetric). Hence, if a cycle cover \mathcal{C} contains any cycle C of at least three edges which does not use the arc ts , then the orientation of C can be reversed to produce a distinct oriented cycle cover \mathcal{C}' , corresponding to a distinct term of the determinant. Let a *reversible cycle* in an oriented cycle cover \mathcal{C} be a cycle C in \mathcal{C} which contains at least three edges and does not use the arc ts . To argue that all oriented cycle covers with reversible cycles cancel over a field of characteristic 2, we define the following pairing. Fix an arbitrary ordering $<$ on V . For each oriented cycle cover \mathcal{C} with at least one reversible cycle, select such a cycle $C \in \mathcal{C}$ by the earliest incidence of a vertex of C according to $<$, and let \mathcal{C}' be the result of reversing C in \mathcal{C} . Since the selection of C is independent of orientation, this map defines a pairing between \mathcal{C} and \mathcal{C}' . By the symmetry of A_{st} , \mathcal{C} and \mathcal{C}' contribute precisely the same term to $\det A_{st}$. Generalising the argument, every oriented cycle cover \mathcal{C} with at least one reversible cycle cancels in $\det A_{st}$ in characteristic 2.

For any oriented cycle cover \mathcal{C} that is not cancelled by this argument, we note that the monomial contributed by \mathcal{C} to $\det A_{st}$ is unique (recall that there is one distinct variable x_e for every edge

e of G). Furthermore, let $e = uv$ be an edge such that x_e occurs in a monomial m of $\det A_{st}$ corresponding to an oriented cycle cover \mathcal{C} . If e occurs in a 2-cycle in \mathcal{C} , then m contains x_e^2 ; otherwise e occurs in the st -cycle C with a passage such as uvw , and x_e has degree 1 in m . \square

Note the slightly subtle interaction between 2-cycle-terms in $\det A_{st}$ and applications of Theorem 3.4. Since variables in 2-cycle-terms have even degree, they are not relevant for the matroid basis sieving of the algorithm, which will therefore effectively sieve directly over st -paths in G . However, the 2-cycle terms prevent us from (for example) finding a Hamiltonian st -path in polynomial time by sieving for terms of degree n . (However, using a weight-tracing variable it is possible to find a *shortest* solution, and to check for the existence of an *odd* or *even* solution.)

6.1.2 Linkage enumeration

Through the same principle, we can construct a matrix whose determinant enumerates perfect (S, T) -linkages. Let $G = (V, E)$ be an undirected graph and let $S, T \subseteq V$ where $|S| = |T|$. As above, define a set of edge variables $X = \{x_e \mid e \in E\}$. For an (S, T) -linkage \mathcal{P} , define

$$X(\mathcal{P}) = \prod_{e \in E(\mathcal{P})} x_e.$$

A *padded* (S, T) -linkage term for an (S, T) -linkage \mathcal{P} is defined as a term

$$X(\mathcal{P}) \cdot \prod_{e \in M} x_e^2,$$

where again M is a (not necessarily perfect) matching in $G - V(\mathcal{P})$. Since we are interested in perfect (S, T) -linkages we make some simplifications. If there is a vertex $v \in S \cap T$, simply delete v from G , S and T since the only possible path on v in a perfect (S, T) -linkage is the length-0 path v . Hence we assume $S \cap T = \emptyset$. We also assume by edge subdivision that $S \cup T$ is an independent set: Note that no (S, T) -linkage needs to use an edge of $G[S]$ or $G[T]$, and a perfect (S, T) -linkage cannot use such an edge. Furthermore, any edge between S and T can be safely subdivided without altering the structure of linkages (and, e.g., give the subdividing vertex the zero vector in the matroid representation).

Lemma 6.5. *There is a matrix A_{ST} over a field of characteristic 2 such that $\det A_{ST}$ enumerates padded perfect (S, T) -linkage terms.*

Proof. We follow the proof of Lemma 6.4, suitably modified. Let A be the Tutte matrix of G over a sufficiently large field of characteristic 2. Let $S = \{s_1, \dots, s_p\}$ and $T = \{t_1, \dots, t_p\}$ with arbitrary ordering. We obtain A_{ST} from A by letting $A[v, v] = 1$, $A[v, s] = 0$, $A[t, v] = 0$ for every $s \in S$, $t \in T$ and $v \in V \setminus (S \cup T)$. Furthermore, we let $A[t_i, s_i] = 1$ for $i \in [p]$ and $A[t_i, s_j] = 0$ otherwise. Essentially, one can think of A_{ST} as the Tutte-like matrix on the directed graph G' where every $v \in V \setminus (S \cup T)$ has a self-loop and the incoming arcs of S and outgoing arcs of T are replaced by the induced matching $\{t_i s_i \mid i \in [p]\}$.

We claim that $\det A_{ST}$ enumerates padded perfect (S, T) -linkage terms as described. This mimics the argument of Lemma 6.4: the terms of $\det A_{ST}$ have a one-to-one correspondence with oriented cycle covers of G' . Note that all other edges of G' not incident with $S \cup T$ are bidirected (i.e., symmetric). Hence, if a cycle cover \mathcal{C} contains any cycle C of length at least 3 disjoint from $S \cup T$, then the orientation of C can be reversed to produce a distinct oriented cycle cover \mathcal{C}' . We call such a cycle *reversible*. To argue that all oriented cycle covers with reversible cycles cancel

over a field of characteristic 2, we define the following pairing. Fix an arbitrary ordering $<$ on V . For each oriented cycle cover \mathcal{C} with at least one reversible cycle, select such a cycle $C \in \mathcal{C}$ by the earliest incidence of a vertex of C according to $<$, and let \mathcal{C}' be the result of reversing C in \mathcal{C} . Since the selection of C is independent of orientation, this map defines a pairing between \mathcal{C} and \mathcal{C}' . By the symmetry of A_{ST} , \mathcal{C} and \mathcal{C}' contribute precisely the same term to $\det A_{ST}$. Generalising the argument, every oriented cycle cover \mathcal{C} with at least one reversible cycle cancels in $\det A_{ST}$ in characteristic 2.

It remains to show that any oriented cycle cover where every cycle either intersects $S \cup T$ or has length at most 2 corresponds to a monomial that does not cancel in $\det A_{ST}$, and that such terms are precisely padded perfect (S, T) -linkages. Let \mathcal{C} be such an oriented cycle cover. We note that the monomial contributed by \mathcal{C} is a unique “fingerprint” of \mathcal{C} as an *undirected* cycle cover, since all edges correspond to distinct variables. Hence if \mathcal{C} is cancelled, it has to be against a distinct oriented cycle cover \mathcal{C}' over the same underlying set of undirected edges. However, reversing a cycle C of length at most 2 yields precisely the same oriented cycle C again, and any cycle C intersecting $S \cup T$ is non-reversible. The latter follows since the only edge leaving T in G' is a directed edge to S , so reversing C leads to attempting to use a non-existing edge from S to T . Hence any oriented cycle cover \mathcal{C} that consists of cycles intersecting $S \cup T$, 2-cycles and 1-cycles survives cancellation.

We next show that the surviving oriented cycle cover terms correspond directly to padded perfect (S, T) -linkages. For any perfect (S, T) -linkage \mathcal{P} padded with a matching M , we can construct a non-cancelled oriented cycle cover: connect the paths of \mathcal{P} up using the edges $t_i s_i$, $i \in [p]$. This defines a vertex-disjoint cycle packing on $V(\mathcal{P})$ which covers all of $S \cup T$. The number of cycles in the cycle cover depends on how the paths in \mathcal{P} connect their endpoints, but the number of cycles is immaterial to the correctness; it is enough that \mathcal{P} produces a unique non-padded term. Together with M and 1-cycles we get an oriented cycle cover with no reversible cycles.

Finally, let \mathcal{C} be a surviving oriented cycle cover, let $\mathcal{C}' \subseteq \mathcal{C}$ be the set of cycles of length at least 3 (which includes every cycle on $S \cup T$ by construction), and let \mathcal{P} be the set of paths produced by deleting any arcs ts , $t \in T$, $s \in S$ from the cycles of \mathcal{C}' . We claim that \mathcal{P} is a perfect (S, T) -linkage. Indeed, since \mathcal{C} is a cycle cover every vertex of $S \cup T$ occurs in a cycle, and there are no cycles on $S \cup T$ of length 1 or 2. Hence $S \cup T$ occur in \mathcal{P} . Furthermore, they clearly occur as endpoints, and oriented such that every path in \mathcal{P} leads from S to T . The cycles of $\mathcal{C} \setminus \mathcal{C}'$ correspond to the padding of the term produced. Thus, $\det A_{ST}$ enumerates all padded (S, T) -linkage terms. \square

6.2 Rank k (S, T) -linkage

We now formally note Theorem 6.2 (from which Theorem 6.1 follows). We begin by bridging the gap between edge variables (from Lemma 6.5) and vertex variables (from the labels of matroid M).

Lemma 6.6. *Let $G = (V, E)$ be an undirected graph and $S, T \subseteq V$ disjoint vertex sets so that $G[S \cup T]$ is edgeless. Let $X_V = \{x_v \mid v \in V\}$ and $X_E = \{x_e \mid e \in E\}$. There is an efficiently computable polynomial $P(X_V, X_E)$ such that the following hold:*

1. *For every perfect (S, T) -linkage \mathcal{P} there is a monomial in $P(X_V, X_E)$ whose odd support corresponds to $V(\mathcal{P}) \cup E(\mathcal{P})$*
2. *For every monomial m in $P(X_V, X_E)$ with odd support $U \subseteq X_V$ and $F \subseteq X_E$, F is the edge set of a perfect (S, T) -linkage \mathcal{P} where $U \subseteq V(\mathcal{P})$*

Proof. Let A_{ST} be the matrix constructed in Lemma 6.5. Thus, $\det A_{ST}$ enumerates padded perfect (S, T) -linkages over the variable set X_E . We evaluate $\det A_{ST}$ with an assignment where

$$x_{uv} \leftarrow x_{uv}(x_u + x_v),$$

and define

$$P(X_V, X_E) = \det A_{ST} \cdot \prod_{s \in S} x_s.$$

We claim that this produces monomials precisely as described.

First, let \mathcal{P} be a perfect (S, T) -linkage, and let $m = X(\mathcal{P}) = \prod_{e \in E(\mathcal{P})} x_e$, with no padding (i.e., with 1-cycles on all other vertices). Then m is a monomial produced by $\det A_{ST}$. We consider the expansion of m into monomials over $X_V \cup X_E$ resulting from the evaluation. We claim that the term

$$\prod_{v \in V(\mathcal{P})} x_v \cdot \prod_{e \in E(\mathcal{P})} x_e$$

is contributed multilinearly precisely once in $P(X_V, X_E)$. Indeed, for every edge $e = uv \in E(\mathcal{P})$, effectively the expansion has to select either the contribution x_u or x_v . Since $P(X_V, X_E)$ is “pre-padded” by $\prod_{s \in S} x_s$, every edge sv leaving $s \in S$ in \mathcal{P} must be oriented to produce x_v instead, or otherwise x_s gets even degree. It follows that the only production that covers every variable x_v for $v \in V(\mathcal{P})$ is when every edge uv , oriented in \mathcal{P} from S to T as (u, v) , contributes its head variable x_v .

Conversely, assume that $P(X_V, X_E)$ has a monomial m where the odd support consists of $U \subseteq X_V$ and $F \subseteq X_E$. Then there is a perfect (S, T) -linkage \mathcal{P} such that $F = \{x_e \mid e \in E(\mathcal{P})\}$. We claim that every variable $x_v \in U$ comes from an edge variable $x_e \in F$. Indeed, the only other production of x_v would be from a padding 2-cycle uvu , which contributes

$$(x_{uv}(x_u + x_v))^2 = x_{uv}^2(x_u^2 + x_v^2)$$

since we are working over a field of characteristic 2. Since no padding cycles intersect $S \cup T$ and since padding 2-cycles evidently do not contribute to the odd support of m , the conclusion follows. \square

We can now finish the result. We recall the statement of the theorem.

Theorem 6.2. *Given an undirected graph $G = (V, E)$, a matroid M over V represented over a field of characteristic 2, sets $S, T \subseteq V$ and an integer k , in randomized time $O^*(2^k)$ and polynomial space we can find a perfect (S, T) -linkage in G which has rank at least k in M .*

Proof. Let $I = (G, M, S, T, k)$ be the input. As noted before Lemma 6.5 we can safely modify I so that $S \cap T = \emptyset$ and $G[S \cup T]$ is edgeless. Let $X_V = \{x_v \mid v \in V\}$ and $X_E = \{x_e \mid e \in E\}$ and let $P(X_V, X_E)$ be the polynomial of Lemma 6.6. Let A_M be the representation of M truncated to rank k and dimension $k \times |V|$. Recall that this can be constructed efficiently, possibly by moving to an extension field \mathbb{F} (see Section 2.1). Furthermore, we assume $|\mathbb{F}| = \Omega(n)$ for the sake of vanishing error probability; again, this can be arranged by moving to an extension field. Now, we use Theorem 3.4 to sieve over $P(X_V, X_E)$ for a monomial whose odd support in X_V spans A_M . By Lemma 6.6, if there is such a monomial m then the vertex set of the monomial is contained in a perfect (S, T) -linkage \mathcal{P} , hence there is a perfect (S, T) -linkage \mathcal{P} such that $V(\mathcal{P})$ spans A_M . Conversely, if there is a perfect (S, T) -linkage \mathcal{P} such that $V(\mathcal{P})$ spans A_M , then there is also a monomial m of $P(X_V, X_E)$ such that the odd support of m spans A_M . The running time and failure probability comes from Theorem 3.4 and $|\mathbb{F}|$. \square

We note a handful of consequences.

Corollary 6.7. *The following problems can be solved in randomized time $O^*(2^k)$ and polynomial space.*

1. Finding a perfect (S, T) -linkage of total length at least k
2. In a vertex-coloured graph, finding a perfect (S, T) -linkage which uses at least k different colours
3. Given a set of terminals $K \subseteq V(G)$ with $|K| = k$, finding a perfect (S, T) -linkage that visits every vertex of K
4. Given a matroid M over $V \cup E$ of total rank k , represented over a field of characteristic 2, finding a perfect (S, T) -linkage \mathcal{P} such that $E(\mathcal{P}) \cup V(\mathcal{P})$ is independent in M

Furthermore, for each of these settings we can find a shortest solution, or a shortest solution of odd, respectively even total length.

Proof. The first three applications follow as in the discussion at the start of this section, by assigning appropriate vectors to the vertices of G . To additionally find a shortest, respectively shortest odd/even solution, attach a weight-tracing variable z to every edge variable x_e and look for a non-zero term in the sieving whose degree in z is minimum (respectively minimum subject to having odd/even degree). For every perfect (S, T) -linkage \mathcal{P} , there is a multitude of padded productions, but there is a unique monomial m where every vertex $v \notin V(\mathcal{P})$ is padded using a 1-cycle (such that only edge variables corresponding to $E(\mathcal{P})$ occur in m). Finding this minimum degree therefore corresponds to finding the shortest length $|\mathcal{P}|$ for a solution \mathcal{P} . Finally, note that padding terms always come in pairs, hence padding m does not change its parity in z .

For the final case, first let $R = S \cap T$. We delete R from S, T and G , contract R in M , and set $k \leftarrow k - |R|$. We then proceed as follows. Attach a weight-tracing variable z to every edge variable x_e . Guess the value of $k_e = |E(\mathcal{P})|$ and note that $|V(\mathcal{P})| = |E(\mathcal{P})| + |S|$ for every perfect (S, T) -linkage; indeed, since $S \cap T = \emptyset$, every path contains an edge, hence every path $P \in \mathcal{P}$ is a tree with $|V(P)| = |E(P)| + 1$. Thus set $k' = 2k_e + |S|$, restricting the guess for k_e to values such that $k' \leq k$, and truncate M to rank k' . Use interpolation to extract terms of $P(X_V, X_E)$ of degree k_e in X_E and use Theorem 3.2 to check for multilinear monomials that span the truncation of M . As above, if there is a solution \mathcal{P} , with the parameter $k_e = |E(\mathcal{P})|$ as guessed, then there is also a monomial m in $P(X_V, X_E)$ of degree precisely k_e in X_E such that m is multilinear and its support corresponds precisely to $E(\mathcal{P}) \cup V(\mathcal{P})$. Furthermore, m is of degree precisely k' , hence m precisely spans the truncation of M . Any term in $P(X_V, X_E)$ of total degree k_e in X_E that is not of this form will either contribute fewer than $k_e + |S|$ variables from $V(\mathcal{P})$ or will fail to be multilinear, and hence will fail to pass Theorem 3.2. \square

6.3 Faster Long st -Path and Long Cycle

Fomin et al. [47] ask whether LONG st -PATH or LONG CYCLE – i.e., the problem of finding, respectively, an st -path or a cycle of length at least k – can be solved in time $O^*((2 - \varepsilon)^k)$ for any $\varepsilon > 0$, given that there is an algorithm solving k -CYCLE in time $O^*(1.66^k)$ by Björklund et al. [19]. We answer in the affirmative, showing that the algorithm of Björklund et al. can be modified to solve LONG st -PATH in time $O^*(1.66^k)$ by working over the cycle-enumerating determinant of Lemma 6.4. A corresponding algorithm for LONG CYCLE follows, by iterating over all choices of st as an edge of the cycle. We prove the following.

Theorem 6.8. *Let $G = (V, E)$ be an undirected graph and $s, t \in V$. There is a randomized algorithm that finds an st -path in G of length at least k in time $O^*((4(\sqrt{2} - 1))^k) = O^*(1.66^k)$ and polynomial space.*

The result takes the rest of the subsection. Like Björklund et al. [19], the algorithm is based around randomly partitioning the vertex set of G as $V = V_1 \cup V_2$, then use algebraic sieving to look for an st -path P that splits “agreeably” between V_1 and V_2 , in time better than $O^*(2^{|P|})$. More specifically, we pick integers k_x and k_2 and define a matroid $M = M(k_2, k_x)$ of rank $r = \eta k$ for some $\eta < 3/4$, and prove that any st -path P that (1) intersects V_2 in precisely k_2 vertices, (2) contains precisely k_x edges that cross between V_1 and V_2 , and (3) has an edge set that spans M necessarily has $|V(P)| \geq k$. We can then look for such a path by working over Lemma 6.4. The details follow.

For a partition $V = V_1 \cup V_2$, let $E_1 = E(G[V_1])$, $E_2 = E(G[V_2])$ and $E_X = E \setminus (E_1 \cup E_2)$ so that $E = E_1 \cup E_X \cup E_2$ partitions E . Given a partition $V = (V_1, V_2)$ and integers r_1 and r_2 , define a matroid $M(r_1, r_2)$ as follows. Let M_1 be a uniform matroid over E_1 of rank r_1 . Let M_2 be the transversal matroid over $E_X \cup E_2$ with hidden set V_2 , where each $e \in E_X \cup E_2$ is linked to every $v \in e \cap V_2$ in V_2 . Furthermore, truncate M_2 to have rank r_2 . That is, a set $F \subseteq E_X \cup E_2$ is independent in M_2 if and only if $|F| \leq r_2$ and F has a set of distinct representatives in V_2 . Let $M(r_1, r_2)$ over ground set E be the disjoint union of M_1 and M_2 .

Lemma 6.9. *Let a partition $V = V_1 \cup V_2$ be given with corresponding edge partition $E = E_1 \cup E_X \cup E_2$. Furthermore let $s, t \in V$, and $M = M(r_1, r_2)$ for some r_1, r_2 . Let P be an st -path and let $C = P + st$ be the corresponding cycle. Assume there is a set $F \subseteq E(C)$ such that $E_2 \cap E(C) \subseteq F$ and F is independent in M , and let $k_x = |F \cap E_X|$. Then $|V(C)| \geq |F| + k_x$.*

Proof. Decompose C cyclically into edges in E_1 and V_1 -paths, i.e., paths whose endpoints lie in V_1 and whose internal vertices lie in V_2 , where we require each V_1 -path to have at least one internal vertex. Let P' be a V_1 -path. We claim that the initial and final edges of P' cannot both be in F . Indeed, all internal edges of P' (except the initial and final edges) lie in F , and the full set of edges $E(P')$ intersect only $|E(P')| - 1$ distinct vertices in V_2 , i.e., $E(P')$ is dependent in M . Since this argument applies to every V_1 -path in C separately, and since the V_1 -paths partition the edges of $E(C) \cap (E_X \cup E_2)$, we conclude

$$|(E(C) \cap E_X) \setminus F| \geq k_x.$$

Hence $|V(C)| = |E(C)| \geq |F| + k_x$. □

We show that this implies an algorithm for detecting long st -paths.

Lemma 6.10. *Let $V = V_1 \cup V_2$ be a partition and k, k_2 and k_x be integers. Let $\mu = \max(k_2, k - k_x/2)$. There is a randomized, polynomial-space algorithm with running time $O^*(2^\mu)$ that detects the existence of an st -path P such that $|V(P)| \geq k$, $|V(P) \cap V_2| = k_2$ and $|E(P + st) \cap E_X| = k_x$.*

Proof. Let $C = P + st$. Let $\ell_1 = \max(0, k - k_x/2 - k_2)$ and note $\mu = k_2 + \ell_1$. Construct the matroid $M = M(\ell_1, k_2)$. By Lemma 6.9, it suffices to prove that there exists a set $F \subseteq E(C)$ independent in M such that $E(C) \cap E_2 \subseteq F$, $|F| \geq k - k_x/2$ and F contains at least $k_x/2$ crossing edges. We use odd support sieving over the construction of Lemma 6.4 to detect such terms.

Let A_{st} be the matrix of Lemma 6.4 such that $\det A_{st}$ enumerates padded st -paths, over a variable set $X = \{x_e \mid e \in E(G)\}$. Create additional variables z_x and z_2 . For every edge $uv \in E$ we associate the vector $M(uv)$ of M with the variable x_{uv} . We modify A_{st} as follows. For every edge $uv \in E_1$, set

$$A[u, v] = A[v, u] = x_{uv}.$$

For every edge $uv \in E_X$, set

$$A[u, v] = A[v, u] = z_x x_{uv}.$$

For every edge $uv \in E_2$, set

$$A[u, v] = A[v, u] = z_2 x_{uv}.$$

Finally, let $P(X)$ be the coefficient of $z_x^{k_x} z_2^{k_2 - k_x/2}$ in $\det A_{st}$. (Clearly, we can reject if k_x is odd.) We claim that there is a path P with $|V(P)| \geq k$, $|V(P) \cap V_2| = k_2$ and $|E(P + st) \cap E_X| = k_x$ if and only if $P(X)$ contains a term whose odd support spans M .

First, let $C = P + st$ be a simple cycle meeting the conditions. Orient C arbitrarily cyclically and let $F \subseteq E(C)$ consist of every edge oriented towards a vertex of V_2 together with ℓ_1 further edges of $E(C) \cap E_1$; note $|E(C) \cap E_1| \geq \ell_1$. Also, F contains precisely $k_x/2$ crossing edges. Furthermore, clearly F is a basis for M . Let $m = \prod_{uv \in E(C)} A[u, v]$; m occurs in $\det A_{st}$ (using only loops for padding). Furthermore, the degree of z_x in m is precisely k_x and the degree of z_2 is precisely $k_2 - k_x/2$. Hence m also occurs in $P(X)$. This proves one direction of the equivalence.

Conversely, let m be a term in $P(X)$ whose odd support spans M , and let F be a subset of the odd support of m such that F is a basis for M . Let C be a cycle such that m corresponds to a padding of C . By design, m contains precisely k_x crossing edges and $k_2 - k_x/2$ edges of E_2 , counting both C and any 2-cycles in the padding. Now, every vertex of $V_2 \cap V(C)$ contributes two endpoints in $E(C)$, every edge of $E_2 \cap E(C)$ represents two such endpoints, and every edge of $E_X \cap E(C)$ represents one such endpoint. Hence $|V_2 \cap V(C)| = |E_2 \cap E(C)| + |E_X \cap E(C)|/2$. Since not all edges counted in the degrees of z_x and z_2 must come from C itself, this is upper bounded by $(k_2 - k_x/2) + k_x/2 = k_2$, with equality only if no padding 2-cycle uses an edge of $E_X \cup E_2$. Furthermore, since F spans M , F represents precisely k_2 edges incident with distinct vertices of V_2 , and since we sieve in the odd support F cannot use edges from any padding 2-cycles. We conclude that C contains precisely k_x crossing edges and is incident with exactly k_2 vertices of V_2 . Finally, $|V(C)| \geq r(M) + k_x/2 \geq k$ by Lemma 6.9. The running time and failure probability follow from Theorem 3.4. \square

It now only remains to combine Lemma 6.10 with a carefully chosen random partition strategy for $V = V_1 \cup V_2$.

Proof of Theorem 6.8. Let P be an st -path and $C = P + st$ a cycle, and let $|V(C)| = ck$, $c \geq 1$. Sample a partition $V = V_1 \cup V_2$ by placing every vertex v into V_2 independently at random with some probability p . For fix choices of p , k_2 and ℓ_1 we estimate the probability that C contains precisely k_2 vertices of V_2 and precisely ℓ_1 edges of E_1 . First, the probability that $|V(C) \cap V_2| = k_2$ is

$$p_1(p, k_2) := \binom{ck}{k_2} p^{k_2} (1-p)^{ck-k_2}.$$

In particular, all $\binom{ck}{k_2}$ colourings of $V(C)$ with k_2 members of V_2 are equally likely. Now, let us count the number among those colourings where there are precisely k_x transitions between V_1 and V_2 . To eliminate edge cases, assume $k_2 < |V(P)|$, $k_x < 2k_2$ and $k_x < 2(ck - k_2)$ and that k_x is even, so that k_x is achievable. To describe the outcomes, consider an initial shorter cycle of k_2 elements, all of which are coloured V_2 , and consider the different ways to place $ck - k_2$ vertices coloured V_1 between these so that there are precisely k_x transitions between V_1 and V_2 . Counting cyclically, this implies that there are precisely $k_x/2$ blocks of vertices coloured V_1 . There are precisely

$$\binom{ck - k_2 - 1}{k_x/2 - 1}$$

ordered sequences of $k_x/2$ positive numbers that sum to $ck - k_2$. Indeed, these can be thought of as placing all $ck - k_2$ elements in a sequence (coding the number $ck - k_2$ in unary) and selecting $k_x/2 - 1$ out of the $ck - k_2 - 1$ gaps between elements to insert a break between blocks. For every

such ordered sequence, we similarly select

$$\binom{k_2}{k_x/2}$$

positions in the cycle of V_2 -vertices into which to insert the blocks. Hence, given an outcome with $|V(C) \cap V_2| = k_2$ the probability of precisely k_x crossing edges is

$$p_2(k_2, k_x) := \binom{ck}{k_2}^{-1} \binom{ck - k_2 - 1}{k_x/2 - 1} \binom{k_2}{k_x/2}.$$

Thus the total probability of meeting both conditions is

$$p_3(p, k_2, k_x) = p_1(p, k_2)p_2(k_2, k_x) = p^{k_2}(1-p)^{ck-k_2} \binom{ck - k_2 - 1}{k_x/2 - 1} \binom{k_2}{k_x/2}.$$

Given such an outcome, we can then detect a cycle by Lemma 6.10 in time $O^*(2^\mu)$ where $\mu = \max(k_2, k - k_x/2)$. By repeating the algorithm $\Theta^*(p_3(p, k_2, k_x))$ times, we get a high probability of success, with a total running time of

$$O^*(2^\mu/p_3(p, k_2, k_x)) = n^{O(1)} \frac{2^\mu}{p^{k_2}(1-p)^{ck-k_2} \binom{ck-k_2-1}{k_x/2-1} \binom{k_2}{k_x/2}}.$$

The choice of p , k_2 and k_x will depend on c , but since there are only $n - k + 1$ possible values of $|V(C)| \geq k$ we may repeat the algorithm for every such value. We follow approximately the analysis used by Björklund et al. [19]. Due to μ , the algorithm has two modes, depending on the value of c . The expected value of $k_x/2$ depends on c and p , and is maximised at $p = 1/2$ and $ck/4$. When c is close to 1 setting $p = 1/2$ yields $E[k_2] = ck/2 < k - E[k_x/2] = (1 - c/4)k$, hence the algorithm is dominated by $\mu = k - k_x/2$, and the best strategy is to maximise k_x . Here, the analysis of Björklund et al. applies. At some crossover point (e.g., $c = 4/3$ if we use the naïve values $p = 1/2$, $k_2 = ck/2$, $k_x/2 = ck/4$) the algorithm at $p = 1/2$ becomes dominated by $\mu = k_2$, and the best strategy is to pick p so that $E[k_2] = E[k - k_x/2]$. Let us consider the second case first. We refrain from optimizing the running time for these values (since this regime does not represent the limiting behaviour of the algorithm) and use $k_2 = cpk$ and $k_x/2 = cp(1-p)k$. Then we set p so that

$$cpk = k - cp(1-p)k \Rightarrow p = 1 - \sqrt{1 - \frac{1}{c}}.$$

It can easily be checked that with $k_2 = pck$ and $k_x = 2p(1-p)ck$, we get $1/p_3(p, k_2, k_x) = O^*(1)$, hence the running time of the algorithm in this regime is $O^*(2^{c(1-\sqrt{1-1/c})k})$, where the exponent decreases with increasing c (approaching $k/2$) and at $c = 4/3$ it becomes $O^*(2^{2k/3}) = O^*(1.59^k)$. Now we focus on the regime $c < 4/3$, in which case we set $p = 1/2$ and $k_2 = ck/2$ (to maximise the expected number of crossing edges). We set $k_x = 2cp(1-p)k + 2\beta ck = (1/2 + 2\beta)ck$ where $\beta > 0$ is a parameter to optimize. We are in the case $\mu = k - k_x/2$. Consider the effect of increasing k_x by 2. Noting that

$$\binom{n}{k+1} = \binom{n}{k} \cdot \frac{n-k}{k+1},$$

the total running time is multiplied by a factor

$$\frac{1/2}{\frac{(1/4-\beta)ck}{(1/4+\beta)ck} \cdot \frac{(1/4-\beta)ck}{(1/4+\beta)ck}}.$$

For the best possible value of β , will equal $1 \pm o(1)$, since otherwise we can improve the running time by raising or lowering k_x . Thus

$$1/4 + \beta = (1/4 - \beta)\sqrt{2} \Rightarrow \beta = \frac{\sqrt{2} - 1}{4(1 + \sqrt{2})} = \frac{(\sqrt{2} - 1)^2}{4} = \frac{3}{4} - \frac{1}{\sqrt{2}}$$

by multiplying with $\sqrt{2} - 1$ in the next-to-last step. We now revisit the total running time. By Stirling's approximation,

$$\binom{n}{\alpha n} = n^{O(1)} \cdot \left(\frac{1}{\alpha^\alpha (1 - \alpha)^{1 - \alpha}} \right)^n.$$

(see [19] for a derivation). Plugging the values ($p = 1/2$, $k_2 = ck/2$, $k_x/2 = (1/4 + \beta)ck = (1 - 1/\sqrt{2})ck$) into the running time and simplifying we get (up to a polynomial factor)

$$\frac{2^{ck} 2^{k - (1 - 1/\sqrt{2})ck}}{\binom{ck/2}{(1 - 1/\sqrt{2})ck}^2} = 2^k 2^{(1/\sqrt{2})ck} (2 - \sqrt{2})^{(2 - \sqrt{2})ck} (\sqrt{2} - 1)^{(\sqrt{2} - 1)ck} = 2^k 2^{ck} (\sqrt{2} - 1)^{ck},$$

which equals $O^*((4(\sqrt{2} - 1))^k) = O^*(1.66^k)$ for the basic case $ck = k$, and is a decreasing function in c . Therefore, this analysis applies for values of c up to the crossover point where $\mu = k_2$. Switching from the running time $2^\mu = 2^{k - k_x/2}$ to $2^\mu = 2^{k_2}$ for the above values of k_2 and k_x represents multiplying the running time by

$$2^{k_2 - k + k_x/2} = 2^{ck/2 - k + (1 - 1/\sqrt{2})ck}$$

hence the running time after the crossover point, with the above parameters, is some function $O^*(\xi^{ck})$, $\xi > 1$. We evaluate the formula at $c = 4/3$ and find it reaches $O^*(1.62^k)$ at this point. Hence no further case distinctions are needed. In summary, our algorithm has the following steps:

1. Repeat the below with every target value $ck \in \{k, \dots, n\}$.
2. If $c \leq 4/3$, set $p = 1/2$, $k_2 = ck/2$ and $k_x = 2(1 - 1/\sqrt{2})ck$.
3. If $c > 4/3$, set $p = 1 - \sqrt{1 - 1/c}$, $k_2 = pck$ and $k_x = 2p(1 - p)ck$.
4. Repeat $\Omega(n/p_3(p, k_2, k_x))$ times: Compute a partition $V = V_1 \cup V_2$ by placing every vertex $v \in V$ into V_2 independently at random with probability p . Use Lemma 6.10 with partition (V_1, V_2) and arguments k, k_2, k_x to detect a cycle in G with the given parameters. If successful, return YES.
5. If every attempt fails, return NO.

□

7 Subgraph problems

Another major area of applications of algebraic methods in FPT algorithms is problems of detecting a particular kind of subgraph. This is of course very broad (and in the form just described would arguably cover all of Section 6 as well), so we focus on two topics. Essentially, these topics correspond to two families of enumerating polynomials: *branching walks* and *homomorphic images*.

The first topic, which is already very broad, is detecting whether a given graph G contains a connected subgraph meeting a particular condition. The parameter here may either be the size

of the subgraph or a parameter related to the condition itself. We review some examples. The first application of branching walks that we are aware of was for the well-known STEINER TREE problem. Here the input is a graph $G = (V, E)$ and a set of terminals $T \subseteq V$, $|T| = k$, and the task is to find a smallest connected subgraph of G that spans T (i.e., a subtree of G whose vertex set contains T). Nederlof [82] showed a polynomial-space, $O^*(2^k)$ -time algorithm for this problem, using an inclusion-exclusion sieving algorithm. We may also consider generalisations of the problem. In GROUP STEINER TREE, the terminals come in k groups and the task is to find the smallest subtree that contains a terminal of each group. In DIRECTED STEINER OUT-TREE, the graph is directed, and the task is to find the smallest out-tree (i.e., a directed subtree of G where every arc leads away from the root) which spans the terminals. Misra et al. [78] showed that both of these variants can be solved in the same running time $O^*(2^k)$.

Another problem, perhaps of an apparently very different nature, is GRAPH MOTIF. The precise definition is slightly involved, but in its base variant the input contains a graph $G = (V, E)$, a vertex colouring $c: V \rightarrow [n]$, an integer k , and a capacity d_q for every colour $q \in c(V)$. The task is to find a subtree T of G on k vertices such that every colour q occurs in at most d_q vertices of T . As surveyed in the introduction, this problem led to the development of the *constrained multilinear detection* method by Björklund et al., leading to a solution in time $O^*(2^k)$ for GRAPH MOTIF as well as several weighted and approximate variants [23].

In our first application in this section, we note a generalisation of the above results.

Theorem 7.1. *Let $G = (V, E)$ be an undirected graph and M be a matroid over V or over E . Let $k, w \in \mathbb{N}$. If M is represented over a field of characteristic 2, then in randomized time $O^*(2^k)$ and polynomial space we can detect the existence of a connected subgraph H of G such that $V(H)$ (respectively $E(H)$) has rank at least k in M and $|V(H)| \leq w$ (respectively $|E(H)| \leq w$). If M is represented over any other field, then the algorithm needs $O^*(2^{\omega k})$ time and $O^*(4^k)$ space.*

The second result regards subgraph isomorphism. Given graphs G and H , the problem of checking whether H is a subgraph of G parameterized by $|V(H)|$ can be either FPT, as when H is a path, or W[1]-hard, as when H is a clique. More generally, SUBGRAPH ISOMORPHISM is FPT by $|V(H)|$ if H comes from a family of graphs with bounded treewidth (originally shown using the colour-coding technique of Alon et al. [4]), and there is good evidence that no more general such class exists [29, 76]. In general, the parameterized complexity of subgraph isomorphism problems has been extensively and meticulously investigated [63, 77].

In fact, one of the fastest methods for SUBGRAPH ISOMORPHISM works via an arithmetic circuit for evaluating the *homomorphism polynomial* [52], allowing for the randomized detection of a subgraph H with $|V(H)| = k$ and of treewidth w in time $O(2^k n^{w+1})$. Furthermore, the exponent $w + 1$ here is optimal, up to plausible conjectures [29]. We observe that this running time is compatible with an additional constraint that the copy of H found in G should be independent in a given linear matroid.

Theorem 7.2. *Let G and H be undirected graphs, $k = |V(H)|$ and $n = |V(G)|$. Let a tree decomposition of H of width w be given. Also let M be a matroid over $V(G)$. If M is represented over a field of characteristic 2, then in randomized time $O(2^k \cdot k^{O(1)} \cdot n^{w+1})$ and polynomial space we can detect whether there is a subgraph of G isomorphic to H whose vertex set is independent in M . Similarly, given M over $V(G) \cup E(G)$ in time $O(2^{k+E(H)} \cdot k^{O(1)} \cdot n^{w+1})$ we can detect a subgraph of G isomorphic to H whose edge and vertex set, are independent in M . Here, we assume that field operations over \mathbb{F} take at most $k^{O(1)}$ time. Over a general field, the algorithm needs $O^*(4^r)$ space and the running time becomes $O(2^{\omega r} k^{O(1)} n^{w+1})$, or $O(4^r k^{O(1)} n^{w+1})$ if a path decomposition of H of width w is provided instead of a tree decomposition. Here, $r = k$ if M is over $V(G)$ and $r = k + |E(G)|$ if M is over $V(G) \cup E(G)$.*

7.1 Finding high-rank connected subgraphs

For our first application, generalizing STEINER TREE and GRAPH MOTIF, we apply the concept of branching walks. Informally, branching walks in a graph G are a relaxation of subtrees of G , similar to how walks are a generalisation of paths. More formally, a branching walk in G can be described as a tree T and a homomorphism mapping T into G . Let us recall the definitions.

Definition 7.3. Let G and H be undirected graphs. A *homomorphism* from G to H is a mapping $\varphi: V(G) \rightarrow V(H)$ such that for every edge $uv \in E(G)$, $\varphi(u)\varphi(v) \in E(H)$.

Branching walks were defined by Nederlof [82]. We use the more careful definition of Björklund et al. [23].

Definition 7.4. Let G be a graph. A *branching walk* $W = (T, \varphi)$ is an ordered, rooted tree T and a homomorphism φ from T to G . We assume w.l.o.g. that $V(G) = \{1, \dots, n\}$ and $V(T) = \{1, \dots, |V(T)|\}$, where $V(T)$ is ordered according to the preorder traversal of T . We say that W *starts from* the vertex $\varphi(1)$ in G . The *size* of W is $|V(T)|$ and its *span* is $\varphi(V(T))$. W *visits* a vertex $v \in V(G)$ if $v \in \varphi(V(T))$. W is *simple* if φ is injective. Finally, W is *properly ordered* if for any two sibling nodes $a, b \in V(T)$ with $a < b$ we have $\varphi(a) < \varphi(b)$.

The ordering here is a technical device to make the map non-ambiguous. Björklund et al. define a generating polynomial (or in our terms, an enumerating polynomial) for properly ordered branching walks. We recall their construction next. Fix a host graph $G = (V, E)$ and a size k for the branching walk. Introduce two sets of variables $X = \{x_v \mid v \in V(G)\}$ and $Y = \{y_{(u,v)}, y_{(v,u)} \mid uv \in E(G)\}$. For a properly ordered branching walk $W = (T, \varphi)$ in G , define the corresponding monomial

$$m(W, X, Y) = x_{\varphi(1)} \prod_{ab \in E(T): a < b} y_{(\varphi(a), \varphi(b))} x_{\varphi(b)}.$$

As Björklund et al. show, $m(W, X, Y)$ is multilinear if and only if W is simple, and W can be reconstructed from the factors of $m(W, X, Y)$. Given a target size k for W and a starting vertex $s \in V(G)$, define

$$P_{k,s}(X, Y) = \sum_W m(W, X, Y) \quad \text{and} \quad P_k(X, Y) = \sum_{s \in V(G)} P_{k,s}(X, Y),$$

where the sum goes over all properly ordered branching walks of size k in G that start from s . Björklund et al. show that $P_{k,s}(X, Y)$ can be evaluated in time polynomial in $n + k$ (in fact, in $O(k^2 m)$ field operations, where $m = |E(G)|$) [23].

We now visit our target result, Theorem 7.1. We observe the key property of branching walks that make them algorithmically useful: A *minimal* branching walk spanning a given vertex set is always a subtree of G .

Lemma 7.5. *Let $G = (V, E)$ be a graph and let $U \subseteq V$ be a set of vertices such that $G[U]$ is connected. Then there is a properly ordered branching walk W in G with span U and size $|U|$. Furthermore, any branching walk with span U and size $|U|$ is simple.*

Proof. Clearly, every branching walk W with span U needs size at least $|U|$. For existence, let T be an arbitrary spanning tree of $G[U]$ where the nodes of T are ordered in preorder traversal, such that at every vertex the lowest-index unvisited child is visited first. Let $W = (T, \varphi)$ where φ is the inverse of the resulting vertex ordering of T . Then W is a properly ordered branching walk with size $|U|$ and span U . \square

Say that a branching walk W in G is *semisimple* if $m(W, X, Y)$ is multilinear in Y (only) and for every edge $uv \in E(G)$ at most one of the variables $y_{(u,v)}$ and $y_{(v,u)}$ occurs in $m(W, X, Y)$. The *edge span* of a branching walk $W = (T, \varphi)$ is the set of edges $uv \in E(G)$ that are the image of $E(T)$ under φ . We note a similar result for branching walks visiting the edge set of a connected subgraph.

Lemma 7.6. *Let H be a connected subgraph in G with $U = V(H)$ and $F = E(H)$. There is a properly ordered branching walk W in G with span $V(H)$, edge span F , and size $|F|+1$. Furthermore, every such W is semisimple, and there is no such W of size less than $|F| + 1$.*

Proof. Let $W = (T, \varphi)$ be a branching walk with edge span F . Since T is a tree, T has size at least $|F|+1$. Furthermore, if there is a branching walk W with size $|F|+1$ and edge span F , then it must be semisimple. We construct such a walk W . Let $W = (T, \varphi)$ be the branching walk constructed by starting at an arbitrary node of U and doing a DFS-like traversal, except that instead of following an edge uv from a vertex u only when v is unvisited, follow the edge uv from u if the edge uv has not been traversed yet. Furthermore, at every vertex u perform these traversals by following the unvisited edge uv with lowest target vertex v . This defines a properly ordered branching walk $W = (T, \varphi)$ with precisely $|F|$ edges, hence with size $|F| + 1$. \square

Given this, and given the ability to evaluate $P_k(X, Y)$, Theorem 7.1 follows easily.

Proof of Theorem 7.1. Let (G, M, k, w) be the input, where M is represented by a matrix A over a field of characteristic 2. We assume by truncation that A has dimension $k \times V(G)$ and rank k , and let $w = \min(w, |E(G)|)$. Furthermore, ensure that A is over a field \mathbb{F} of size $\omega(n^2)$, e.g., $\mathbb{F} = GF(2^{c \log n})$ for $c > 2$. For each $\ell = k, \dots, w$ let $P_\ell(X, Y)$ be the branching walk polynomial for branching walks of size ℓ over variable sets X and Y defined above. Use Theorem 3.4 with the vectors of A associated with X or with Y , correspondingly (depending on whether M is a matroid over $V(G)$ or over $E(G)$) and assume that for some ℓ , Theorem 3.4 reports that $P_\ell(X, Y)$ contains a monomial m whose odd support spans A . Then $m = m(W, X, Y)$ for a branching walk W . Let $W = (T, \varphi)$ and let $S \subseteq \varphi(V(T)) \cup \varphi(E(T))$ correspond to the subset of the odd support of m that spans A , $|S| = k$. Since W is a branching walk, S is the vertex set or edge set of a connected subgraph of G on at most ℓ vertices. Hence (G, M, k, w) is a YES-instance.

On the other hand, assume that (G, M, k, w) is a YES-instance and let H be a subgraph of minimum cardinality that spans A . Let $\ell = |V(H)|$ respectively $\ell = |E(H)| + 1$ for the vertex respectively edge version of the problem. By Lemma 7.5 respectively Lemma 7.6, there is a branching walk W with span $V(H)$, respectively edge span $E(H)$, which is simple, respectively semisimple. Thus $P_\ell(X, Y)$ contains a monomial $m = m(W, X, Y)$ which is multilinear in X respectively in Y , and which spans A . Note that $P_\ell(X, Y)$ is a homogeneous polynomial of degree $2\ell - 1 < n^2$, so the probability of a false negative for $P_\ell(X, Y)$ is $o(1)$. Hence with probability $1 - o(1)$, the algorithm reports that the input is a YES-instance. The running time follows from Theorem 3.4.

In the case that M is represented over some other field \mathbb{F} , the same analysis applies (including assuming $|\mathbb{F}| = \omega(n^2)$), but the running time and space complexity follow from Theorem 3.6 instead. \square

We note some applications of this result. First, consider the basic STEINER TREE problem, and let $G = (V, E)$ and $T \subseteq V$ be an input, $T = \{t_1, \dots, t_k\}$. Define a k -dimensional matroid M over V by letting vertex t_i be associated with vector e_i , and every other vertex associated with the k -dimensional zero vector. Then a connected subgraph H of G spans M if and only if $T \subseteq V(H)$. We can cover GROUP STEINER TREE with a similar construction. Let the input be $(G = (V, E), \mathcal{T})$,

with terminal grouping $\mathcal{T} = \{T_1, \dots, T_k\}$, $T_i \subseteq V$ for each i . We assume the terminal sets are pairwise disjoint by adding pendants: for every $T_i \in \mathcal{T}$ and every vertex $t \in T_i$, add a pendant t^i to t and replace T_i by the set $\{t^i \mid t \in T_i\}$. This raises the size of a minimum solution by precisely k vertices. We can now apply label e_i to every vertex in T_i and the zero vector as label to every other vertex and proceed as above.

Next, let us review how to use matroid constructions to solve the various optimization variants of GRAPH MOTIF surveyed by Björklund et al. [23]. Let $(G = (V, E), c, k, (d_q)_{q \in c(V)})$ be a GRAPH MOTIF instance. Additionally, we consider the following operations, mimicking the EDIT DISTANCE problem. Let H be a connected subgraph of G with k vertices. Let $C = C(H)$ be the multiset of vertex colours in H , i.e., $C(H) = \{c(v) \mid v \in V(H)\}$ with element multiplicities preserved. To *substitute* a colour $q \in C$ for another colour $q' \in c(V)$, we remove one copy of q from C and add a copy of q' . To *insert* a colour q , we add a copy of q to C . To *delete* a colour q , we remove a copy of q from C . Furthermore, let a multiset Q of colours be given. We wish to decide whether there is a connected subgraph H of G with $|V(H)| = k$ such that $C(H)$ can be transformed into Q by making at most k_s substitutions, at most k_i insertions, and at most k_d deletions. Individually, these operations correspond well to standard matroid transformations. Let M be the partition matroid over $V(D)$ where every colour class $c^{-1}(q)$ has capacity d_q . Then using M in Theorem 7.1 directly solves GRAPH MOTIF. Further allowing substitutions, insertions and/or deletions can be handled by combinations of extensions and truncations over M . We consider the following general case.

Lemma 7.7. *Let $G = (V, E)$ be a graph and c a vertex colouring of G . Let $k_s, k_d, k_i \in \mathbb{N}$ and a multiset Q be given. There is a matroid M over V , representable over a field of characteristic 2, such that any set of k vertices from V forms a basis of M if and only if the multiset $C(H)$ can be transformed into Q by making at most k_s substitutions, k_d deletions and k_i insertions.*

Proof. We note that since $C(H)$ and Q are multisets, without element order, finding the minimum cost for a transformation is much simpler than in EDIT DISTANCE. Let $C = C(H)$ and let $C_0 = C \cap Q$ be the multiset intersection. Let $a = \min(|C| - |C_0|, |Q| - |C_0|, k_s)$.

Claim 7.1. *$C(H)$ can be transformed into Q with the operation limits prescribed if and only if $|C_0| + a \geq \max(|Q| - k_i, k - k_d)$.*

Proof of claim: The transformation can use a substitutions, and thereafter it has either exhausted C , Q or the budget k_s . In either case, it thereafter needs to use $|C| - |C_0| - a$ deletions and $|Q| - |C_0| - a$ insertions, i.e., $k_d \geq |C| - (|C_0| + a)$ and $k_i \geq |Q| - (|C_0| + a)$. The result follows. \diamond

Hence, let $r = |C_0| + a$ be the number of vertices from $V(H)$ that can be matched against Q by using at most k_s substitutions. We wish to accept a vertex set $V(H)$ if and only if $r \geq k_0 := \max(|Q| - k_i, k - k_d)$. We construct a matroid for this purpose. If $k_0 > k$, reject the parameters. Otherwise, execute the following construction sequence.

1. Let M_1 be the partition matroid over $V(G)$ where for every colour q , the set $c^{-1}(q)$ has capacity in M_1 corresponding to its count in Q .
2. Let M_2 be M_1 with the rank extended by k_s .
3. Let M_3 be M_2 truncated to rank k_0 .
4. Let M be M_3 extended by additional rank $k - k_0$.

Indeed, let S be a basis of M . Then there is a set $S' \subseteq S$ with $|S'| = k_0$ that is a basis of M_3 , implying that using at most k_s substitutions, S' can be matched into Q , and $|S'| = k_0$. Conversely,

if S is a set of k vertices, let $C' = C(S) \cap Q$, and assume that $|C'| + \min(k - |C'|, |Q| - |C'|, k_s) \geq k_0$. Then there is a set $S' \subseteq S$ consisting of k_0 vertices that can be matched into Q using at most k_s substitutions, i.e., S' is independent in M_3 , and S is a basis of M . \square

Since there are only $O(k^3)$ valid options for the integers k_s , k_d and k_i , by repeating this construction we can clearly sieve for a connected subgraph H of size k with a minimum *cost* of transformation, using costs as given in Björklund et al. [23]. Another option, of attaching weight-tracing variables tracing the number of substitutions and deletions, similarly to the algorithm in [23], would of course also be possible, but our purpose here was to illustrate the matroid construction.

Further variations are clearly also possible, e.g., as in the notion of *balanced solutions* considered in Section 5.1 one may look for subgraphs with both upper and lower bounds $d_q \leq |V(H) \cap c^{-1}(q)| \leq e_q$ for every colour class q .

Interestingly, both STEINER TREE and GRAPH MOTIF are SeCoCo-hard, i.e., under the set cover conjecture they cannot be solved in time $O^*(2^{(1-\varepsilon)k})$ for any $\varepsilon > 0$ [35]. Hence improving the algorithm of Theorem 7.1 is certainly SeCoCo-hard as well.

7.2 Independent subgraph isomorphism

Next, we review the SUBGRAPH ISOMORPHISM problem. Let G and H be undirected graphs, and introduce a variable set $X = \{x_v \mid v \in V(G)\}$. Let $k = |V(H)|$ and $n = |V(G)|$. The *homomorphism polynomial* is the polynomial

$$\sum_{\varphi: V(H) \rightarrow V(G)} \prod_{v \in V(H)} x_{\varphi(v)}$$

where the sum goes over homomorphisms φ . If a treewidth decomposition of width w is given for H , then the homomorphism polynomial can be evaluated in time $f(k) \cdot n^{w+1}$ for a modest function $f(k)$ [52]. In particular, we follow the exposition of Brand [26] who shows that in time $O(c^k + n^{w+1})$ for $c < 2$ we can both compute a tree decomposition of width w for H and construct an algebraic circuit of total size $O(k \cdot n^{w+1})$ which evaluates the homomorphism polynomial.

Since the homomorphism polynomial has no negative terms working over, e.g., the reals there is no concern for cancellations. However, since we want to work over fields of characteristic 2, we introduce additional terms in the way of *algebraic fingerprinting* (cf. [67]), to prevent cancellations. In fact, we introduce two sets of additional variables for algorithmic convenience. Let $X' = \{x'_{i,v} \mid i \in V(H), v \in V(G)\}$ and $Y = \{y_e \mid e \in V(G)\}$. Then we define the *decorated homomorphism polynomial*

$$P_{H \rightarrow G}(X, X', Y) = \sum_{\varphi: V(H) \rightarrow V(G)} \prod_{i \in V(H)} x_{\varphi(i)} x'_{i, \varphi(i)} \prod_{ij \in E(H)} y_{\varphi(i)\varphi(j)},$$

where the variables $y_{\varphi(i)\varphi(j)} = y_{\varphi(j)\varphi(i)}$ are taken without order on its subscript terms, and where φ ranges over all homomorphisms from H to G . Then clearly, every homomorphism φ has a unique algebraic “fingerprint” monomial $m(\varphi)$, since it is encoded in the X' -factors of $m(\varphi)$.

It is easy to modify the construction of Brand [26, Section 4.6.1] to construct a circuit for (or directly compute) $P_{H \rightarrow G}(X, X', Y)$ at a slightly larger polynomial cost in k .

Proposition 7.8. *Let a tree decomposition of with w for H be given. Then in time $k^{O(1)}n^{w+1}$ we can construct an algebraic circuit of size $k^{O(1)}n^{w+1}$ that computes $P_{H \rightarrow G}$. Furthermore, if the decomposition is a path decomposition, then the circuit can be made skew.*

We can now show Theorem 7.2. This follows the obvious path, with some extra care taken to ensure that our polynomial term remains n^{w+1} .

Proof of Theorem 7.2. Let G and H be given, as well as a tree decomposition of H of width w . Note that a homomorphism $\varphi: H \rightarrow G$ represents a subgraph of G isomorphic to H if and only if φ is injective on $V(H)$, which holds if and only if $\prod_{i \in V(H)} x_{\varphi(i)}$ is multilinear.

Let M be a linear matroid, and let $r = k$ if M is over $V(G)$ or $r = k + |E(H)|$ if M is over $V(G) \cup E(G)$. We assume that M is represented by a matrix A with r rows and rank r . We also assume A is over a sufficiently large field \mathbb{F} (where in fact some $|\mathbb{F}| = \Omega(k^2)$ suffices since only the degree of $P_{H \rightarrow G}$ is important for correctness). We now employ the sieving of Theorem 3.2. Note that $P_{H \rightarrow G}$ is homogeneous of degree $2k + |E(H)|$ (and homogeneous in X and Y separately). Using the precise running time bound from Theorem 3.2, we note that field operations over \mathbb{F} can be performed in $\log^{O(1)} |\mathbb{F}| = \tilde{O}(1)$ time, independent of n . A larger field \mathbb{F} could be given in the input, but in this case operations over \mathbb{F} take only $k^{O(1)}$ time by assumption. If there is a subgraph H' of G isomorphic to H and independent in M , then $P_{H \rightarrow G}$ will contain a term m corresponding to that map φ , thus m is multilinear in $X \cup Y$ and Theorem 3.2 applies and will detect H' with high probability. Conversely, assume that Theorem 3.2 detects a monomial m such that m (in X , respectively in $X \cup Y$) is multilinear and contains a basis for M . Then m must be multilinear in X , since m spans M . Since the monomials of $P_{H \rightarrow G}$ are in 1-to-1 correspondence with homomorphisms $\varphi: H \rightarrow G$, m must represent a homomorphism φ which is injective over $V(H)$. Thus H' is isomorphic to H as noted above, and is independent in M . If M is over $V(G) \cup E(G)$, then the same argument and algorithm applies except that we are sieving over both the variable sets X and Y .

The running time over a general field follows from Theorem 3.6. In particular, Brand [26] notes that the homomorphism polynomial circuit can be made skew if constructed over a path decomposition, and not otherwise. \square

The result with a matroid over $V(G) \cup E(G)$ could also be achieved by simply subdividing every edge of H and G , but this would blow up $|V(G)|$ and hence the polynomial factor of the algorithm.

As with our other matroid applications, there is a range of consequences, including (e.g.) finding a colourful copy of H in a vertex-coloured graph; finding a copy of H in G subject to capacity constraints on vertex classes; finding a copy H' of H in G such that $G - E(H')$ is connected; and all the other applications of matroid constraints covered in this paper.

8 Speeding-up Dynamic Programming

The notion of *representative sets* for linear matroids plays an essential role in the design of FPT algorithms [50, 51], as well as kernelization [68]. For a matroid $M = (V, \mathcal{I})$, a set $X \subseteq V$ is said to *extend* a set Y , if X and Y are disjoint and $X \cup Y$ is independent in M . The *representative set lemma*, due to Lovász [72] and Marx [75], states the following: Let $M = (V, \mathcal{I})$ be a linear matroid of rank k , and $\mathcal{Y} \subseteq 2^V$ be a collection of subsets of V . Then, there is a subcollection $\mathcal{Y}' \subseteq \mathcal{Y}$ (which can be computed “efficiently”) of size at most 2^k that *represents* \mathcal{Y} , i.e., for every $X \subseteq V$, there is a set in \mathcal{Y} extending X if and only if such a set exists in \mathcal{Y}' . There are plethora of dynamic programming FPT algorithms in the literature, where the table size is reduced from n^k to 2^k , using the representative set lemma. In this section, we exemplify how to use determinantal sieving in place of such dynamic programming approaches in three applications, MINIMUM EQUIVALENT SUBGRAPH, EULERIAN DELETION, and CONFLICT-FREE SOLUTION. We improve the running time over existing algorithms, while saving space usage to polynomial.

8.1 Minimum Equivalent Graph

MINIMUM EQUIVALENT GRAPH is defined as follows. We are given a directed graph $G = (V, E)$ and an integer k , and the question is whether there is a subgraph $G' = (V, E')$ with at most k edges with the same reachability pattern, i.e., for every $u, v \in V$, there is a uv -path in G if and only if there is in G' . Fomin et al. [50] show that MINIMUM EQUIVALENT GRAPH reduces to the following question: Are there a pair B_1, B_2 where B_1 is an in-branching and B_2 is an out-branching in G , with a common root v , such that they have at least ℓ edges in common? We phrase this as a matroid-theoretical problem.

Let MATROID INTERSECTION INTERSECTION refer to the following problem. The input is four matroids M_1, \dots, M_4 over the same ground set U , each of rank k , and an integer $\ell > 0$. The question is whether there are bases $B_A \in M_1 \cap M_2$ and $B_B \in M_3 \cap M_4$ such that $|B_A \cap B_B| \geq \ell$. This captures the above question, since rooted in- and out-branchings can be constructed as the intersection of a graphic matroid and a suitable partition matroid.

Lemma 8.1. *For matroids represented over a common field \mathbb{F} of characteristic 2, MATROID INTERSECTION INTERSECTION can be solved in $O^*(2^{2k})$ time.*

Proof. We define a new ground set $U^* = U_1 \cup U_2$, where U_1 is a copy of U and $U_2 = U \times U$. We also define new matroids M'_1, \dots, M'_4 as follows. Let $A_i, i = 1, \dots, 4$ be the representation of M_i . Then M'_i is represented by a matrix A'_i where for $x \in U_1$ we have $A'_i[\cdot, x] = A_i[\cdot, x]$, and for $(x, y) \in U_2$ we have $A'_i[\cdot, (x, y)] = A_i(x)$ for $i = 1, 2$ and $A'_i[\cdot, (x, y)] = A_i(y)$ for $i = 3, 4$.

Claim 8.1. *The rank of M'_i for each $i = 1, \dots, 4$ is k .*

Proof of claim: Since each column of A'_i is a copy of a column from A_i , clearly the rank of A'_i is at most the rank of A_i . Conversely, since every column of A_i occurs as a column of A'_i , the rank of A'_i is at least the rank of A_i . \diamond

We show that this reduces MATROID INTERSECTION INTERSECTION to a kind of “weighted” instance of 4-MATROID INTERSECTION over matroids of rank k .

Claim 8.2. *The input instance is positive if and only if there is a common basis of M'_1 through M'_4 that contains at least ℓ elements from U_1 .*

Proof of claim: The idea is the following. Let B_A and B_B be solutions to the problem. Split (B_A, B_B) as $B_0 = B_A \cap B_B$, $B_1 = B_A \setminus B_B$ and $B_2 = B_B \setminus B_A$, $|B_0| = r$ for some $r \geq \ell$. Write $B_0 = \{u_1, \dots, u_r\}$, $B_1 = \{x_1, \dots, x_{k-r}\}$ and $B_2 = \{y_1, \dots, y_{k-r}\}$. Define the new set

$$S = B_0 \cup \{(x_i, y_i) \mid i \in [k-r]\}$$

where $B_0 \subseteq U_1$ and $S \setminus B_0 \subseteq U_2$. Then S is a common basis of M'_1 through M'_4 . Indeed, for M'_1 and M'_2 the matrix $A'_i[\cdot, S]$ induced by S is a copy of $A_i[\cdot, B_A]$ and for M'_3 and M'_4 the matrix $A'_i[\cdot, S]$ is a copy of $A_i[\cdot, B_B]$. These are bases by assumption. Furthermore S contains $r \geq \ell$ elements from U_1 .

Conversely, assume that S is a common basis of M'_1 through M'_4 with $|S \cap U_1| = r$ for some $r \geq \ell$. Extract the sets

$$B_A = (S \cap U_1) \cup \{x \mid (x, y) \in S \cap U_2\} \quad \text{and} \quad B_B = (S \cap U_1) \cup \{y \mid (x, y) \in S \cap U_2\}.$$

We claim that B_A is a basis for M_1 and M_2 , and B_B a basis for M_3 and M_4 . Indeed, we have $|B_A| = |B_B| = |S|$ since otherwise one matrix $A'_i[\cdot, S]$ would contain duplicated columns. Thus the matrices $A'_i[\cdot, S]$ and $A_i[\cdot, B_A]$ (respectively $A_i[\cdot, B_B]$) are identical up to column ordering. \diamond

Hence we are left to solve the question whether there exists a common basis for M'_1 through M'_4 with at least ℓ elements from U_1 . For this, we proceed as in the algorithm for 4-MATROID INTERSECTION. By the Cauchy-Binet formula, there is a polynomial $P(X)$ over $X = \{x_u \mid u \in U\}$ which enumerates common bases of M'_1 and M'_2 . Introduce a new variable z and evaluate $P(X)$ at a value where x_u is multiplied by z for $u \in U_1$. Let $P'_r(X)$ be the coefficient of z^r in $P(X)$ under this evaluation. The question now reduces to asking if there is a monomial m in $P'_r(X)$ for any $r \geq \ell$ such that m is independent in M'_3 and M'_4 , which can be solved using Cor. 3.3 in time $O^*(2^{2k})$. \square

Since rooted in-branchings and out-branchings can be represented via matroid intersection over matroids of rank $n - 1$, an $O^*(2^{2n})$ -time algorithm for MINIMUM EQUIVALENT GRAPH follows.

8.2 Eulerian Deletion

An undirected (or directed) graph is said to be *Eulerian* if it admits a closed walk that visits every edge (or arc, respectively.) exactly once. It is known that an undirected graph is Eulerian if and only if it is connected and *even*, i.e., every vertex has an even degree and that a directed graph is Eulerian if and only if weakly connected and *balanced*, i.e., every vertex has the same number of in-neighbors and out-neighbors [7]. UNDIRECTED EULERIAN EDGE DELETION (DIRECTED EULERIAN EDGE DELETION) is the problem of determining whether the input undirected (directed) graph has an edge (arc) set S of size at most k such that $G \setminus S$ is Eulerian. Cai and Yang [30] initiated the parameterized analysis of these problems among other related problems. The parameterized complexity of UNDIRECTED EULERIAN EDGE DELETION and DIRECTED EULERIAN EDGE DELETION was left open by Cai and Yang [30]. Cygan et al. [38] designed the first FPT algorithms with running time $O^*(2^{O(k \log k)})$ based on the colour-coding technique. Later, Goyal et al. [55] gave improved algorithms running in time $O^*(2^{(2+\omega)k})$ using a representative set approach.

We briefly describe their approach on UNDIRECTED EULER DELETION (the directed version is similar). Let T denote the set of vertices of odd degree in G . (Note that T must have even cardinality.) An edge set S is called a *T-join* if T is exactly the set of vertices of odd degree in the graph (V, S) . In other words, a *T-join* is the collection of T -paths \mathcal{P} between disjoint pairs of vertices in T and cycles that are edge-disjoint (see e.g., [88]). Thus, every (inclusion-wise) minimal *T-join* consists of $|T|/2$ paths connecting disjoint pairs of vertices in T . We will say that a *T-join* is *semi-minimal* if it is the union of $|T|/2$ edge-disjoint walks between disjoint pairs of T . Cygan et al. [38] observed that an edge set S with $|S| \leq k$ is a solution for UNDIRECTED EULER DELETION if and only if S is a *T-join* and $G \setminus S$ is connected. The algorithm of Goyal et al. [55] employs a dynamic programming approach; there is an entry for every subset of $T' \subseteq T$ (which may have size up to $2k$), which stores *T*-walks between disjoint pairs of T' . They may be n^k such walks, so the number of walks to store can be reduced using representative sets of co-graphic matroids.

We give an $O^*(2^k)$ -time (and polynomial-space) algorithm. The improvements are twofold. First, we avoid computing the representative families. Second, we avoid dynamic programming over the subsets of T . Let $X = \{x_e \mid e \in E\}$ be a set of edge variables.

Lemma 8.2. *There is a polynomial $P(X)$ that can be efficiently evaluated with its multilinear terms of degree k enumerating all minimal *T*-joins S of size k and (not necessarily all) semi-minimal *T*-joins of size k .*

Proof. Let A be a symmetric matrix indexed by V , where $A[u, v] = x_{uv}$ if $uv \in E$ and $A[u, v] = 0$ otherwise. For every $v \in V$, let e_v be the $|V|$ -dimensional vector where $e_v[v] = 1$ and $e_v[v'] = 0$ for

each $v' \in V \setminus \{v\}$. We define a skew-symmetric matrix A' indexed by T , where for every $u, v \in T$,

$$A'[u, v] = \sum_{\ell \in [k]} e_u^T A^\ell e_v,$$

which enumerates all (u, v) -walks of length up to k . Note that this is the *unlabelled* walk polynomial, as opposed to the labelled walk polynomial defined in Section 2. We claim that the degree- k terms of $\text{Pf } A'$ yield the desired polynomial. Recall that the Pfaffian enumerates all perfect matching on the complete graph on T . Thus, every multilinear term in the monomial expansion corresponds to a set of T -walks that connect disjoint pairs of T with no edge occurring twice or more. Each multilinear term thus corresponds to a semi-minimal T -join in G . In particular, there is an algebraically independent term correspond to every minimal T -join. (This may not be the case for semi-minimal T -joins.) \square

We solve **UNDIRECTED EULERIAN DELETION** as follows. Assume that there is no solution of size at most $k - 1$. To find a solution of size exactly k , let $P_k(X)$ be the degree- k part of the polynomial $P(X)$ defined in Lemma 8.2. Note that $P_k(X)$ can be evaluated via polynomial interpolation. We use the basis sieving algorithm (Theorem 3.2) over the cographic matroid M_k of rank k . Note that a multilinear term corresponding to a semi-minimal T -join vanishes during the sieving step, since we assumed that there is no solution of size $k - 1$ or smaller. Thus, we sieve for minimal T -join S such that S is a basis for M_k , i.e., $G \setminus S$ is connected. By iterating over all values of k , we can find a minimum solution (or decide that no solution of size k or smaller exists) in time $O^*(2^k)$.

Theorem 8.3. **UNDIRECTED EULERIAN DELETION** can be solved in $O^*(2^k)$ time.

Next, we briefly discuss the **DIRECTED EULERIAN DELETION**. Goyal et al. [55] showed that an arc set S with $|S| \leq k$ is a minimal solution for **DIRECTED EULERIAN DELETION** if and only if S is the union of ℓ arc-disjoint paths $\mathcal{P} = \{P_1, \dots, P_\ell\}$ such that (i) $G \setminus S$ is weakly connected and (ii) there are exactly $\deg^+(v) - \deg^-(v)$ paths starting at every v with $\deg^+(v) > \deg^-(v)$ and $\deg^-(v) - \deg^+(v)$ paths ending at every v with $\deg^-(v) > \deg^+(v)$, where $\ell = \frac{1}{2} \sum_{v \in V} |\deg^+(v) - \deg^-(v)|$.

We define a $V^+ \times V^-$ matrix A' where V^+ (and V^-) is the multiset that contains $|\deg^+(v) - \deg^-(v)|$ copies of v with $\deg^+(v) > \deg^-(v)$ (and $\deg^-(v) > \deg^+(v)$, respectively). Let each entry $A'[v^+, v^-]$ be a polynomial enumerating all walks as in Lemma 8.2. Let $P(X) = \det A$. Then, each term in $P(X)$ corresponds to a set of ℓ walks from V^+ to V^- , where the start and end are distinct elements from V^+ and V^- , respectively. As for **UNDIRECTED EULERIAN DELETION**, using the sieving algorithm of Theorem 3.2 over the cographic matroid of the underlying undirected graph, we obtain:

Theorem 8.4. **DIRECTED EULERIAN DELETION** can be solved in $O^*(2^k)$ time.

8.3 Conflict-free Solution

There has been a line of research studying “conflict-free” variants of classical problems [2, 40, 41, 62]. Consider a problem in which we search for a solution S , which is a subset of the ground set E . In the conflict-free version, the solution should form an independent (i.e., pairwise non-adjacent) set in H , where H is an additionally given graph whose vertices are E and whose edges are “conflicts”. Formally, let us define the problem **CONFLICT-FREE SOLUTION** as follows. The input is a collection \mathcal{F} of (possibly exponentially many) subsets of E , a conflict graph H on E , and an integer t . The problem asks there is a set $S \in \mathcal{F}$ of size t that forms an independent set in H . We give

another immediate consequence of Theorem 3.4 on CONFLICT-FREE SOLUTION. As a by-product, we improve on existing algorithms. Since CONFLICT-FREE SOLUTION is W[1]-hard in general, we restrict the input as follows.

Let $P_H(X)$ over $\{x_v \mid v \in V\}$ be an enumerating polynomial for independent sets in H , i.e., $P_H(X) = \sum_I c_I \prod_{v \in I} x_v$, where I ranges over all independent sets of H and $c_I \in \mathbb{F}$ is a constant. Since it is NP-hard to determine the existence of an independent set of size k , the polynomial $P_H(X)$ cannot be efficiently evaluated, unless $P = NP$. However, when H is from a restricted graph, it can be. Let \mathcal{G}_{IS} be a class of such graphs, i.e., \mathcal{G}_{IS} contains all graphs H such that $P_H(X)$ can be evaluated in polynomial time. Then, \mathcal{G}_{IS} includes, for instance, chordal graphs (graphs that do not contain any cycle of length four or greater as an induced subgraph) as shown by Achlioptas and Zampetakis [1].² Moreover, we will say that a set family \mathcal{F} over E is k -representable if there is a matrix $A \in \mathbb{F}^{k \times \ell}$ over a field \mathbb{F} of characteristic 2 such that every $e \in E$ is associated with a pairwise disjoint subset $\Gamma_e \subseteq [\ell]$, and for any $S \subseteq E$, $S \in \mathcal{F}$ if and only if $A[\cdot, \bigcup_{e \in S} \Gamma_e]$ has full row rank (i.e., contains a non-singular submatrix). By Theorem 3.4, we have

Theorem 8.5. *If \mathcal{F} is k -representable and $H \in \mathcal{G}_{IS}$, then CONFLICT-FREE SOLUTION can be solved in $O^*(2^k)$ time.*

Theorem 8.5 gives the following improvements over existing dynamic programming algorithms:

- **CONFLICT-FREE MATCHING:** Given a graph $G = (V, E)$, a conflict graph $H = (E, E')$, and an integer k , it asks whether G has a conflict-free matching of size k . Agrawal et al. [2] showed that CONFLICT-FREE MATCHING can be solved in $O^*(2^{(2\omega+2)k})$ time when H is chordal. We can solve this problem in $O^*(4^k)$ time because the collection of matchings of size k is $2k$ -representable: Let A be the representation of the uniform matroid over V (with every vertex v copied $\deg(v)$ times) of rank $2k$. Note that a set of k edges spans A if and only if it is pairwise disjoint, i.e., forms a matching.
- **CONFLICT-FREE SET COVER:** Given a collection \mathcal{E} of sets over V (with $|V| = n$), a conflict graph $H = (\mathcal{E}, E')$, and an integer t , and it asks whether G has a conflict-free set cover $S \subseteq \mathcal{E}$ (i.e., $\bigcup S = V$) of size at most t . Jacob et al. [62] gave an $O^*(3^n)$ -time algorithm for CONFLICT-FREE SET COVER when the conflict graph H is chordal. Since set covers are n -representable (mapping every set E to a list of elements (v, E) , $v \in E$ as in Section 4), by Theorem 8.5, it can be solved in $O^*(2^n)$ time.

Incidentally, Jacob et al. [62] showed that CONFLICT-FREE SET COVER is W[1]-hard parameterized by n , even if the conflict graph is bipartite. This implies that the class of bipartite graphs is not contained in \mathcal{G}_{IS} , although a maximum independent set can be found in polynomial time on bipartite graphs. It is perhaps no coincidence that the problem of counting independent sets is #P-hard for bipartite graphs [86].

9 Conclusions

We have presented *determinantal sieving*, a new powerful method for algebraic exact and FPT algorithms that extends the power of multilinear sieving with the ability to sieve for terms in a polynomial that in addition to being multilinear are also independent in an auxiliary linear matroid. This yields significantly improved and generalized results for a range of FPT problems, including

²We remark in passing their algorithm uses divisions; so it does not give an arithmetic circuit.

q -MATROID INTERSECTION in time $O^*(2^{(q-2)k})$ over a field of characteristic 2, improving on a previous result, of $O^*(4^{qk})$ [28], as well as algorithms solving problems over *frameworks* in the same running time as was previously known for the basic existence problem (e.g., SUBGRAPH ISOMORPHISM). Additionally, we showed that over fields of characteristic 2, we can exploit cancellations in monomial expansion to sieve for terms in a polynomial whose *odd support* contains a basis for the auxiliary matroid. This has further applications for a multitude of problems, such as finding diverse solution collections and for parameterized path, cycle and linkage problems. All our algorithms are randomized, and use polynomial space.

Let us mention a few issues that we have not focused on in this paper.

Weighted problems. As in most algebraic algorithms, we can handle solution weights with a pseudopolynomial running time. That is, given an algebraic sieving algorithm for some problem over a ground set V with a running time of $O^*(f(k))$, and given a set of item weights $\omega: V \rightarrow \mathbb{N}$ and a weight target W , we can usually find solutions of weight W in time $O^*(f(k) \cdot W)$ by multiplying every variable $x_v, v \in V$ by $z^{\omega(v)}$ for a new variable z and using interpolation. However, the gold standard would be to reduce the weight dependency to $O^*(f(k) \cdot \log W)$ for the task of finding a min-weight solution, and this appears incompatible with algebraic algorithms. Even for the most classical problem TSP, whose unweighted variant HAMILTONICITY is solvable in $O^*(1.66^n)$ time [16], the $O^*(2^n)$ -barrier has been broken only partially, conditionally, and recently [83].

Counting. Since our most efficient algorithms work over fields of characteristic 2, they do not intrinsically allow us to count the number of solutions. Indeed, for many settings relevant to us, such as k -PATH and bipartite or general matchings, the corresponding counting problems are known to be hard ($\#W[1]$ -hard respectively $\#P$ -hard; see Curticapean [34] for a survey). On the other hand, being able to detect the existence of a witness does have some applications for *approximate* counting. In particular, having access to a decision oracle for *colourful* witnesses, given a colouring of the ground set, implies approximate counting algorithms [43, 42, 15]. Improved, algebraically based FPT approximate counting algorithms are also known for particular problems, such as $\#k$ -PATH [69].

Derandomization. The task of derandomizing our results ranges from doable with known methods to completely infeasible, given the details of the application.

In a typical application of our method, combining a polynomial $P(X)$ and a linear matroid M over X , we have two sources of randomness: Finding a representation of M and the Schwartz-Zippel step of checking whether $P(X)$ is non-zero. For many matroids and matroid constructions, a representation can be found efficiently, including uniform matroids, partition matroids, and graphic and co-graphic matroids, as well as the operations of dualization, disjoint union and truncation [84, 75, 70]. Beyond this, there appears to be a barrier. A deterministic representation of transversal matroids would presumably also lead to a deterministic solution to EXACT MATCHING, which is long open. Since gammoids generalize transversal matroids, and transversal matroids can be constructed via a sequence of matroid union steps over very simple matroids [84], gammoids and non-disjoint union also appear difficult. However, some progress has been made on constructing representations in superpolynomial time that depends on the rank [71, 79].

For the Schwartz-Zippel PIT step, the obstacles are oddly similar. A polynomial $P(X)$ (over \mathbb{Q} or \mathbb{R}) is *combinatorial* if all coefficients are non-negative. In such a case, PIT can be derandomized using the exterior algebra; see Brand [26]. This is compatible with the extensor-based determinantal sieving used in this paper, hence some of our results can be derandomized. However, for results

that depend on odd sieving, or where $P(X)$ corresponds to a determinant or Pfaffian computation, derandomization once again appears infeasible.

9.1 Open questions

Let us highlight some open questions.

One very interesting question is DIRECTED HAMILTONICITY. We note two very different methods for checking Hamiltonicity in bipartite digraphs in time $O^*(c^n)$ for $c < 2$. The first is by Cygan, Kratsch and Nederlof [37], who established a rank bound of $2^{n/2-1}$ on the *perfect matching connectivity matrix*. This leads to a SETH-optimal algorithm for HAMILTONICITY parameterized by pathwidth, and an algorithm for HAMILTONICITY in bipartite digraphs in time $O^*(1.888^n)$. However, the fastest algorithm for HAMILTONICITY in bipartite digraphs follows a polynomial sieving approach by Björklund, Kaski and Koutis [21]. In our terminology, we would describe their algorithm as, given a bipartite digraph $G = (U \cup V, E)$, enumerating subgraphs of G that have in- and out-degree 1 in V and whose underlying undirected graph is a spanning tree of G plus one edge. It then remains to sieve via inclusion-exclusion for those graphs which have non-zero in- and out-degree for every vertex in U as well, which they show can be done in time $O^*(3^{|U|})$ rather than $4^{|U|}$ due to the structure of the problem space. Still, it remains unknown whether HAMILTONICITY in general digraphs can be solved in $O^*(c^n)$ for any $c < 2$.

We would be very interested in a derivation of the $2^{n/2}$ rank bound for the perfect matching connectivity matrix in a less problem-specific manner. We also note, to the best of our knowledge, that the optimal running time for HAMILTONICITY parameterized by treewidth remains open. Finally, can k -PATH be solved in time $O^*(c^k)$ for some $c < 2$ on bipartite digraphs?

Another major problem concerns k DISJOINT PATHS. Given the success of algebraic algorithms for related problems, it would be very interesting to find an algebraic algorithm for this for general k . Björklund and Husfeldt [17] show an algebraic algorithm for $k = 2$, by showing a way to compute the permanent over $\mathbb{Z}_4[X]/(X^m)$, the ring of bounded-degree polynomials over \mathbb{Z}_4 . For the nearly ten years since this result's original publication, we do not know of any developments even for $k = 3$.

As a more down-to-earth problem, what is the best running time for q -MATROID PARITY and (possibly) q -SET PACKING? Recall that Björklund et al. [19] showed that q -DIMENSIONAL MATCHING can be solved in $O^*(2^{(q-2)k})$ time and that q -SET PACKING can be solved in time $O^*(2^{(q-\varepsilon_q)k})$ for some $\varepsilon_q > 0$, essentially by randomized reduction to q -DIMENSIONAL MATCHING. Can q -MATROID PARITY, the generalisation of q -SET PACKING, be solved in $O^*(2^{(q-\varepsilon_q)k})$ time for some $\varepsilon_q > 0$? The difference is most stark for $q = 3$, where 3-MATROID INTERSECTION is solvable in time $O^*(2^k)$, 3-SET PACKING in time $O^*(3.328^k)$ and 3-MATROID PARITY only in time $O^*(8^k)$.

Among other individual problems of interest are to find improvements and the best possible running times for problems such as LONG DIRECTED CYCLE [50, 95] CONNECTED f -FACTOR [49], and more generally parameterized connectivity problems, cf. [3, 45].

Acknowledgments. TK is supported by the DFG project DiPa (NI 369/21).

References

- [1] D. Achlioptas and K. Zampetakis. Local approximations of the independent set polynomial. In *ICALP*, volume 198 of *LIPICs*, pages 8:1–8:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.

- [2] A. Agrawal, P. Jain, L. Kanesh, and S. Saurabh. Parameterized complexity of conflict-free matchings and paths. *Algorithmica*, 82(7):1939–1965, 2020.
- [3] A. Agrawal, P. Misra, F. Panolan, and S. Saurabh. Fast exact algorithms for survivable network design with uniform requirements. *Algorithmica*, 84(9):2622–2641, 2022.
- [4] N. Alon, R. Yuster, and U. Zwick. Color-coding. *J. ACM*, 42(4):844–856, 1995.
- [5] S. Bandyapadhyay, F. V. Fomin, T. Inamdar, and K. Simonov. Proportionally fair matching with multiple groups. *CoRR*, abs/2301.03862, 2023.
- [6] J. Bang-Jensen. Edge-disjoint in- and out-branchings in tournaments and related path problems. *J. Comb. Theory, Ser. B*, 51(1):1–23, 1991.
- [7] J. Bang-Jensen and G. Gutin. *Digraphs - theory, algorithms and applications*. Springer, 2002.
- [8] J. Bang-Jensen, K. V. Klinkby, and S. Saurabh. k -Distinct branchings admits a polynomial kernel. In *ESA*, volume 204 of *LIPICs*, pages 11:1–11:15. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [9] J. Bang-Jensen, S. Saurabh, and S. Simonsen. Parameterized algorithms for non-separating trees and branchings in digraphs. *Algorithmica*, 76(1):279–296, 2016.
- [10] J. Bang-Jensen and A. Yeo. The minimum spanning strong subdigraph problem is fixed parameter tractable. *Discret. Appl. Math.*, 156(15):2924–2929, 2008.
- [11] J. Baste, M. R. Fellows, L. Jaffke, T. Masarík, M. de Oliveira Oliveira, G. Philip, and F. A. Rosamond. Diversity of solutions: An exploration through the lens of fixed-parameter tractability theory. *Artif. Intell.*, 303:103644, 2022.
- [12] J. Baste, L. Jaffke, T. Masarík, G. Philip, and G. Rote. FPT algorithms for diverse collections of hitting sets. *Algorithms*, 12(12):254, 2019.
- [13] R. Bellman. Dynamic programming treatment of the travelling salesman problem. *J. ACM*, 9(1):61–63, 1962.
- [14] M. Bentert, L. Kellerhals, and R. Niedermeier. Finding balance-fair short paths in graphs. *CoRR*, abs/2203.17132, 2022. Accepted to AAI 2023.
- [15] A. Bhattacharya, A. Bishnu, A. Ghosh, and G. Mishra. Faster counting and sampling algorithms using colorful decision oracle. In *STACS*, volume 219 of *LIPICs*, pages 10:1–10:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2022.
- [16] A. Björklund. Determinant sums for undirected Hamiltonicity. *SIAM J. Comput.*, 43(1):280–299, 2014.
- [17] A. Björklund and T. Husfeldt. Shortest two disjoint paths in polynomial time. *SIAM J. Comput.*, 48(6):1698–1710, 2019.
- [18] A. Björklund, T. Husfeldt, P. Kaski, and M. Koivisto. Fourier meets möbius: fast subset convolution. In *STOC*, pages 67–74. ACM, 2007.
- [19] A. Björklund, T. Husfeldt, P. Kaski, and M. Koivisto. Narrow sieves for parameterized paths and packings. *J. Comput. Syst. Sci.*, 87:119–139, 2017.

- [20] A. Björklund, T. Husfeldt, and N. Taslaman. Shortest cycle through specified elements. In *SODA*, pages 1747–1753. SIAM, 2012.
- [21] A. Björklund, P. Kaski, and I. Koutis. Directed Hamiltonicity and out-branchings via generalized Laplacians. In *ICALP*, volume 80 of *LIPICs*, pages 91:1–91:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2017.
- [22] A. Björklund, P. Kaski, and L. Kowalik. Fast witness extraction using a decision oracle. In *ESA*, volume 8737 of *Lecture Notes in Computer Science*, pages 149–160. Springer, 2014.
- [23] A. Björklund, P. Kaski, and L. Kowalik. Constrained multilinear detection and generalized graph motifs. *Algorithmica*, 74(2):947–967, 2016.
- [24] A. Björklund, P. Kaski, L. Kowalik, and J. Lauri. Engineering motif search for large graphs. In *ALLENEX*, pages 104–118. SIAM, 2015.
- [25] C. Brand. Patching colors with tensors. In *ESA*, volume 144 of *LIPICs*, pages 25:1–25:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019.
- [26] C. Brand. *Paths and Walks, Forests and Planes*. PhD thesis, 2019.
- [27] C. Brand, H. Dell, and T. Husfeldt. Extensor-coding. In *STOC*, pages 151–164. ACM, 2018.
- [28] C. Brand and K. Pratt. Parameterized applications of symbolic differentiation of (totally) multilinear polynomials. In *ICALP*, volume 198 of *LIPICs*, pages 38:1–38:19. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [29] K. Bringmann and J. Slusallek. Current algorithms for detecting subgraphs of bounded treewidth are probably optimal. In *ICALP*, volume 198 of *LIPICs*, pages 40:1–40:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [30] L. Cai and B. Yang. Parameterized complexity of even/odd subgraph problems. *J. Discrete Algorithms*, 9(3):231–240, 2011.
- [31] J. Chen, J. Kneis, S. Lu, D. Mölle, S. Richter, P. Rossmanith, S.-H. Sze, and F. Zhang. Randomized divide-and-conquer: Improved path, matching, and packing algorithms. *SIAM Journal on Computing*, 38(6):2526–2547, 2009.
- [32] F. Chierichetti, R. Kumar, S. Lattanzi, and S. Vassilvitskii. Fair clustering through fairlets. In *NIPS*, pages 5029–5037, 2017.
- [33] F. Chierichetti, R. Kumar, S. Lattanzi, and S. Vassilvitskii. Matroids, matchings, and fairness. In *AISTATS*, volume 89 of *Proceedings of Machine Learning Research*, pages 2212–2220. PMLR, 2019.
- [34] R. Curticapean. Counting problems in parameterized complexity. In *IPEC*, volume 115 of *LIPICs*, pages 1:1–1:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018.
- [35] M. Cygan, H. Dell, D. Lokshtanov, D. Marx, J. Nederlof, Y. Okamoto, R. Paturi, S. Saurabh, and M. Wahlström. On problems as hard as CNF-SAT. *ACM Trans. Algorithms*, 12(3):41:1–41:24, 2016.
- [36] M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015.

- [37] M. Cygan, S. Kratsch, and J. Nederlof. Fast Hamiltonicity checking via bases of perfect matchings. *J. ACM*, 65(3):12:1–12:46, 2018.
- [38] M. Cygan, D. Marx, M. Pilipczuk, M. Pilipczuk, and I. Schlotter. Parameterized complexity of Eulerian deletion problems. *Algorithmica*, 68(1):41–61, 2014.
- [39] M. Cygan, J. Nederlof, M. Pilipczuk, M. Pilipczuk, J. M. M. van Rooij, and J. O. Wojtaszczyk. Solving connectivity problems parameterized by treewidth in single exponential time. *ACM Trans. Algorithms*, 18(2):17:1–17:31, 2022.
- [40] A. Darmann, U. Pferschy, and J. Schauer. Determining a minimum spanning tree with disjunctive constraints. In *ADT*, volume 5783 of *Lecture Notes in Computer Science*, pages 414–423. Springer, 2009.
- [41] A. Darmann, U. Pferschy, J. Schauer, and G. J. Woeginger. Paths, trees and matchings under disjunctive constraints. *Discret. Appl. Math.*, 159(16):1726–1735, 2011.
- [42] H. Dell and J. Lapinskas. Fine-grained reductions from approximate counting to decision. *ACM Trans. Comput. Theory*, 13(2):8:1–8:24, 2021.
- [43] H. Dell, J. Lapinskas, and K. Meeks. Approximately counting and sampling small witnesses using a colorful decision oracle. *SIAM J. Comput.*, 51(4):849–899, 2022.
- [44] R. Diestel. *Graph Theory, 4th Edition*, volume 173 of *Graduate texts in mathematics*. Springer, 2012.
- [45] A. E. Feldmann, A. Mukherjee, and E. J. van Leeuwen. The parameterized complexity of the survivable network design problem. In *SOSA*, pages 37–56. SIAM, 2022.
- [46] F. V. Fomin, P. A. Golovach, L. Jaffke, G. Philip, and D. Sagunov. Diverse pairs of matchings. In *ISAAC*, volume 181 of *LIPICs*, pages 26:1–26:12. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [47] F. V. Fomin, P. A. Golovach, T. Korhonen, K. Simonov, and G. Stamoulis. Fixed-parameter tractability of maximum colored path and beyond. In *SODA*, 2023.
- [48] F. V. Fomin, P. A. Golovach, F. Panolan, G. Philip, and S. Saurabh. Diverse collections in matroids and graphs. In *STACS*, volume 187 of *LIPICs*, pages 31:1–31:14. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021.
- [49] F. V. Fomin, P. A. Golovach, F. Panolan, and S. Saurabh. Editing to connected f -degree graph. *SIAM J. Discret. Math.*, 33(2):795–836, 2019.
- [50] F. V. Fomin, D. Lokshtanov, F. Panolan, and S. Saurabh. Efficient computation of representative families with applications in parameterized and exact algorithms. *J. ACM*, 63(4):29:1–29:60, 2016.
- [51] F. V. Fomin, D. Lokshtanov, F. Panolan, and S. Saurabh. Representative families of product families. *ACM Trans. Algorithms*, 13(3):36:1–36:29, 2017.
- [52] F. V. Fomin, D. Lokshtanov, V. Raman, S. Saurabh, and B. V. R. Rao. Faster algorithms for finding and counting subgraphs. *J. Comput. Syst. Sci.*, 78(3):698–706, 2012.

- [53] J. F. Geelen and S. Iwata. Matroid matching via mixed skew-symmetric matrices. *Comb.*, 25(2):187–215, 2005.
- [54] I. M. Gessel and R. P. Stanley. Algebraic enumeration. *Handbook of combinatorics*, 2:1021–1061, 1995.
- [55] P. Goyal, P. Misra, F. Panolan, G. Philip, and S. Saurabh. Finding even subgraphs even faster. *J. Comput. Syst. Sci.*, 97:1–13, 2018.
- [56] G. Z. Gutin, D. Majumdar, S. Ordyniak, and M. Wahlström. Parameterized pre-coloring extension and list coloring problems. *SIAM J. Discret. Math.*, 35(1):575–596, 2021.
- [57] G. Z. Gutin, F. Reidl, and M. Wahlström. k -Distinct in- and out-branchings in digraphs. *J. Comput. Syst. Sci.*, 95:86–97, 2018.
- [58] G. Z. Gutin, M. Wahlström, and A. Yeo. Rural postman parameterized by the number of components of required edges. *J. Comput. Syst. Sci.*, 83(1):121–131, 2017.
- [59] T. Hanaka, Y. Kobayashi, K. Kurita, S. W. Lee, and Y. Otachi. Computing diverse shortest paths efficiently: A theoretical and experimental study. In *AAAI*, pages 3758–3766. AAAI Press, 2022.
- [60] T. Hanaka, Y. Kobayashi, K. Kurita, and Y. Otachi. Finding diverse trees, paths, and more. In *AAAI*, pages 3778–3786. AAAI Press, 2021.
- [61] I. Holyer. The NP-completeness of edge-coloring. *SIAM J. Comput.*, 10(4):718–720, 1981.
- [62] A. Jacob, D. Majumdar, and V. Raman. Parameterized complexity of conflict-free set cover. *Theory Comput. Syst.*, 65(3):515–540, 2021.
- [63] B. M. P. Jansen and D. Marx. Characterizing the easy-to-find subgraphs from the viewpoint of polynomial-time algorithms, kernels, and Turing kernels. In *SODA*, pages 616–629. SIAM, 2015.
- [64] K. Kawarabayashi. An improved algorithm for finding cycles through elements. In *IPCO*, volume 5035 of *Lecture Notes in Computer Science*, pages 374–384. Springer, 2008.
- [65] I. Koutis. Faster algebraic algorithms for path and packing problems. In *ICALP (1)*, volume 5125 of *Lecture Notes in Computer Science*, pages 575–586. Springer, 2008.
- [66] I. Koutis and R. Williams. Limits and applications of group algebras for parameterized problems. In *ICALP (1)*, volume 5555 of *Lecture Notes in Computer Science*, pages 653–664. Springer, 2009.
- [67] I. Koutis and R. Williams. Algebraic fingerprints for faster algorithms. *Commun. ACM*, 59(1):98–105, 2016.
- [68] S. Kratsch and M. Wahlström. Representative sets and irrelevant vertices: New tools for kernelization. *J. ACM*, 67(3):16:1–16:50, 2020.
- [69] D. Lokshtanov, A. Björklund, S. Saurabh, and M. Zehavi. Approximate counting of k -paths: Simpler, deterministic, and in polynomial space. *ACM Trans. Algorithms*, 17(3):26:1–26:44, 2021.

- [70] D. Lokshtanov, P. Misra, F. Panolan, and S. Saurabh. Deterministic truncation of linear matroids. *ACM Trans. Algorithms*, 14(2):14:1–14:20, 2018.
- [71] D. Lokshtanov, P. Misra, F. Panolan, S. Saurabh, and M. Zehavi. Quasipolynomial representation of transversal matroids with applications in parameterized complexity. In *ITCS*, volume 94 of *LIPICs*, pages 32:1–32:13. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018.
- [72] L. Lovász. Flats in matroids and geometric graphs. In *Proc. Sixth British Combinatorial Conf.*, Combinatorial Surveys, pages 45–86, 1977.
- [73] L. Lovász. On determinants, matchings, and random algorithms. In *FCT*, pages 565–574. Akademie-Verlag, Berlin, 1979.
- [74] M. Mahajan and V. Vinay. A combinatorial algorithm for the determinant. In *SODA*, pages 730–738. ACM/SIAM, 1997.
- [75] D. Marx. A parameterized view on matroid optimization problems. *Theor. Comput. Sci.*, 410(44):4471–4479, 2009.
- [76] D. Marx. Can you beat treewidth? *Theory Comput.*, 6(1):85–112, 2010.
- [77] D. Marx and M. Pilipczuk. Everything you always wanted to know about the parameterized complexity of subgraph isomorphism (but were afraid to ask). In *STACS*, volume 25 of *LIPICs*, pages 542–553. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2014.
- [78] N. Misra, G. Philip, V. Raman, S. Saurabh, and S. Sikdar. FPT algorithms for connected feedback vertex set. *J. Comb. Optim.*, 24(2):131–146, 2012.
- [79] P. Misra, F. Panolan, M. S. Ramanujan, and S. Saurabh. Linear representation of transversal matroids and gammoids parameterized by rank. *Theor. Comput. Sci.*, 818:51–59, 2020.
- [80] K. Mulmuley, U. V. Vazirani, and V. V. Vazirani. Matching is as easy as matrix inversion. *Comb.*, 7(1):105–113, 1987.
- [81] K. Murota. *Matrices and matroids for systems analysis*, volume 20. Springer Science & Business Media, 1999.
- [82] J. Nederlof. Fast polynomial-space algorithms using inclusion-exclusion. *Algorithmica*, 65(4):868–884, 2013.
- [83] J. Nederlof. Bipartite TSP in $O(1.9999^n)$ time, assuming quadratic time matrix multiplication. In *STOC*, pages 40–53. ACM, 2020.
- [84] J. Oxley. *Matroid Theory*. Oxford University Press, 2011.
- [85] F. Panolan, S. Saurabh, and M. Zehavi. Parameterized algorithms for list K -cycle. *Algorithmica*, 81(3):1267–1287, 2019.
- [86] J. S. Provan and M. O. Ball. The complexity of counting cuts and of computing the probability that a graph is connected. *SIAM J. Comput.*, 12(4):777–788, 1983.
- [87] S. Saurabh and M. Zehavi. $(k, n - k)$ -Max-Cut: An $O^*(2^k)$ -time algorithm and a polynomial kernel. *Algorithmica*, 80(12):3844–3860, 2018.

- [88] A. Schrijver. *Combinatorial Optimization: Polyhedra and Efficiency*. Springer, 2003.
- [89] J. T. Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *J. ACM*, 27(4):701–717, 1980.
- [90] D. Tsur. Faster deterministic parameterized algorithm for k -path. *Theor. Comput. Sci.*, 790:96–104, 2019.
- [91] W. T. Tutte. The factorization of linear graphs. *Journal of the London Math. Soc.*, 1(2):107–111, 1947.
- [92] M. Wahlström. Abusing the Tutte matrix: An algebraic instance compression for the K-set-cycle problem. In *STACS*, volume 20 of *LIPICs*, pages 341–352. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2013.
- [93] R. Williams. Finding paths of length k in $O^*(2^k)$ time. *Inf. Process. Lett.*, 109(6):315–318, 2009.
- [94] M. Włodarczyk. Clifford algebras meet tree decompositions. *Algorithmica*, 81(2):497–518, 2019.
- [95] M. Zehavi. A randomized algorithm for long directed cycle. *Inf. Process. Lett.*, 116(6):419–422, 2016.
- [96] R. Zippel. Probabilistic algorithms for sparse polynomials. In *EUROSAM*, pages 216–226, 1979.