A new class of copulas having dependence range larger than FGM-type copulas

Swaroop Georgy Zachariah^a, Mohd. Arshad^{a1} and Ashok Kumar Pathak^b

^a Department of Mathematics, Indian Institute of Technology Indore, India ^b Department of Mathematics and Statistics, Central University of Punjab, Bathinda, India

Abstract

We propose a new bivariate symmetric copula with positive and negative dependence properties. The main features of the proposed copula are its simple mathematical structure, wider dependence range compared to FGM copula and its generalizations, and no lower and upper tail dependence. The maximum range of Spearman's Rho of the proposed copula is [-0.5866, 0.5866], which improves the dependence range of the FGM copula and its various generalizations. A new bivariate Rayleigh distribution is developed using the proposed copula, and some statistical properties have been studied. A real data set is analyzed to illustrate the proposed bivariate distribution's relevance in practical contexts.

Keywords: Copula, Kendall's Tau, Spearman's Rho, Rayleigh Distribution, Bivariate Distribution.

Mathematics Subject Classification (2020): 62H05, 62H10.

1 Introduction

Copula plays a significant role in the field of statistics, finance, engineering and medical sciences for modelling dependent data sets. If we have a family of copulas, we naturally have a collection of multivariate distributions with whatever marginal distributions we desire. This feature of the copula is useful in every branch of study where dependence modelling and simulation are an integral part. Sklar (1959) proved that if X and Y are two random variables having marginal cumulative distributions G(x) and H(y) respectively, then there exists a function C such that $F(x, y) = P(X \leq x, Y \leq y) = C(G(x), H(y)), \ \forall (x, y) \in \mathbb{R}^2$. If $G(\cdot)$ and $H(\cdot)$ are continuous, then C is unique: otherwise C is uniquely determined on $\operatorname{Range}(F) \times \operatorname{Range}(G)$. Thus, every joint cumulative distribution function can be expressed as a function $C : \mathcal{I}^2 \to \mathcal{I}$ is said to be a bivariate copula if it satisfies the following conditions:

$$C(u,0) = C(0,v) = 0$$
 and $C(u,1) = u, C(1,v) = v, \forall u, v \in \mathcal{I}$ (1.1)

$$C(u_2, v_2) - C(u_1, v_2) - C(u_2, v_1) + C(u_1, v_1) \ge 0, \ \forall u_1 < u_2, v_1 < v_2 \in \mathcal{I}$$

$$(1.2)$$

¹Corresponding author. E-mail addresses: arshad@iiti.ac.in(M. Arshad), ashokiitb09@gmail.com (Ashok Kumar Pathak), swaroopgeorgy@gmail.com (Swaroop Georgy Zachariah).

where $\mathcal{I} = [0, 1]$ and $\mathcal{I}^2 = [0, 1] \times [0, 1]$. Note that Eq. (1.2) is called the 2-increasing property. For more details about copulas, see Nelsen (2006), Durante and Sempi (2016), Mai and Scherer (2017), Hofert et al. (2018), and references therein.

In recent years, many researchers have used copula functions to construct multivariate distributions. For instances, Achcar et al. (2015) studied bivariate generalized exponential distribution via Farlie-Gumbel-Morgenstern (FGM) copula. Kundu and Gupta (2017) discussed bivariate Birnbaum-Saunders using Gaussian copula. Abd Elaal and Jarwan (2017) studied bivariate generalized exponential distribution from FGM and Plackett copula functions and discussed various estimation procedures. El-Sherpieny et al. (2018) used FGM copula for constructing a bivariate Weibull distribution. Recently, Almetwally and Muhammed (2020) developed bivariate Fréchet distribution using the FGM copula.

In literature, a wide variety of copulas are available; of them, FGM copula received much attention due to its simple mathematical structure and exhibited positive and negative dependence (see Farlie (1960)). Spearman's Rho, denoted by ρ_c , is one of the dependence measures used for measuring the dependence structure captured by a copula. The range of values of ρ_c lies between -1 and 1. Positive values of ρ_c indicate positive dependence and negative values for negative dependence. For the FGM copula, the range of values of Spearman's Rho is very low, i.e., $\rho_c \in [-0.33, 0.33]$ (see Farlie (1960)). So, the FGM copula is unsuitable for modelling data with a high dependence structure. Many researchers attempted to propose FGM-type copula for improving the correlation coefficient. Huang and Kotz (1999) proposed two extended FGM copulas, having $\rho_c \in [-0.33, 0.375]$ and $\rho_c \in [-0.33, 0.391]$ respectively. Bairamov and Kotz (2002) and Bekrizadeh et al. (2015) also extended FGM copula with $\rho_c \in [-0.48, 0.502]$ and $\rho_c \in [-0.50, 0.43]$ respectively. Recently, Chesneau (2022) proposed a polynomial-sine copula exhibiting positive as well as negative dependence with $\rho_c \in [-0.4927, 0.4927]$. In most of these works, we can see that parameters are added to improve Spearman's correlation coefficient range, resulting in a mathematically complex structure and computationally more expensive for estimating unknown parameters. To overcome these drawbacks, we propose a simple bivariate copula without adding any parameter to existing copulas. The proposed copula improves the dependence range of Spearman's correlation of various FGM-type copulas reported in the literature.

The present paper is organized as follows: In Sections 2 and 3, we will introduce a new bivariate copula and study some dependency measures of the proposed copula. In Section 4 a new bivariate Rayleigh distribution is derived from the proposed copula. We also derive expressions for conditional distribution and product moments. A real data is also analyzed using the proposed bivariate Rayleigh distribution as an application.

2 New Bivariate Copula

Consider the following bivariate function

$$C(u,v;\delta,\alpha) = uv + \delta\left(1 - e^{\alpha(u-u^2)}\right)\left(1 - e^{\alpha(v-v^2)}\right), \quad (u,v) \in \mathcal{I}^2,$$
(2.1)

where α and δ are real valued parameters. Assume that the parameter δ depends on α , and α takes arbitrary value in \mathbb{R} . The bivariate function, defined in Eq. (2.1), satisfied the boundary conditions of a bivariate copula given in Eq. (1.1). But we need to find the range of parameter δ for which this function is a valid bivariate copula, i.e., the function in Eq. (2.1) satisfy the 2-increasing property given in Eq. (1.2). Kim et al. (2011) proved that 2-increasing property in

an absolutely continuous copula is equivalent to the condition that copula density $c(u, v; \delta, \alpha)$ is non-negative, i.e.,

$$c(u,v;\delta,\alpha) = \frac{\partial^2 C(u,v)}{\partial u \partial v} = 1 + \delta \alpha^2 g(u) g(v) \ge 0, \qquad (2.2)$$

where $g(t) = (1-2t) e^{\alpha(t-t^2)}, t \in \mathcal{I}$. Clearly, non-negativity of copula density in Eq. (2.2) depends on the behaviour of function $g(\cdot)$ and the value of the parameter δ . For finding the feasible range of δ , we will divide the domain \mathcal{I}^2 of (u, v) into four quadrants as:



Since the sign of g(t) is positive if $t \in (0, \frac{1}{2})$, and negative if $t \in (\frac{1}{2}, 1)$, it follows that the product g(u)g(v) is positive on $R_1 \cup R_3$ and negative on $R_2 \cup R_4$. Thus, the copula density $c(u, v; \delta, \alpha)$ is non-negative if

$$\delta \ge \frac{-1}{\alpha^2 g(u)g(v)}, \quad (u,v) \in R_1 \cup R_3,$$

and

$$\delta \le \frac{-1}{\alpha^2 g(u)g(v)}, \quad (u,v) \in R_2 \cup R_4.$$

Therefore, the copula density $c(u, v; \delta, \alpha)$ is non-negative if

$$\frac{-1}{\alpha^2 \sup_{(u,v) \in R_1 \cup R_3} \{g(u)g(v)\}} \le \delta \le \frac{-1}{\alpha^2 \inf_{(u,v) \in R_2 \cup R_4} \{g(u)g(v)\}}.$$
(2.3)

Since the behaviour of function $g(\cdot)$ depends on $\alpha \in \mathbb{R}$, we will consider the following three cases: **Case I:** When $\alpha = 0$.

In this case, the copula density $c(u, v; \delta, 0) = 1$, which is non-negative, and hence the 2-increasing property holds for arbitrary value of δ . Moreover, the proposed copula (2.1) reduces to the product copula. Note that the product copula is a well known copula, which corresponds to the independence of two random variables.

Case II: When $\alpha \leq 2, \alpha \neq 0$.

In this case, g(t) is a decreasing function on \mathcal{I} with $g(0) = 1, g\left(\frac{1}{2}\right) = 0$ and g(1) = -1. It follows that g(t) takes values in [0, 1] for $t \in \left[0, \frac{1}{2}\right]$, and takes values in [-1, 0) for $t \in \left(\frac{1}{2}, 1\right]$. Thus, the product function g(u)g(v) is bounded above by 1 on $R_1 \cup R_3$, and the upper bound 1 is attended at $(u, v) \in \{(0, 0), (1, 1)\}$. Therefore, $\sup_{(u,v)\in R_1\cup R_3} \{g(u)g(v)\} = 1$. Further, for $(u, v) \in R_2$, g(u) takes values in [-1, 0) and g(v) takes values in [0, 1]. This implies that the product function g(u)g(v)is bounded below by -1 on R_2 , and the lower bound -1 is attended at u = 1, v = 0. Similarly, for $(u, v) \in R_4$, the product function g(u)g(v) is bounded below by -1, which is attended at u = 0, v = 1. Therefore, $\inf_{(u,v)\in R_2\cup R_4} \{g(u)g(v)\} = -1$. Now, using these values in inequality (2.3), we get the feasible range of the parameter δ as

$$\frac{-1}{\alpha^2} \le \delta \le \frac{1}{\alpha^2}.\tag{2.4}$$

Case III: When $\alpha > 2$.

Let $r_1 = \frac{1}{2} - \frac{1}{\sqrt{2\alpha}}$ and let $r_2 = \frac{1}{2} + \frac{1}{\sqrt{2\alpha}}$. Clearly, $0 < r_1 < \frac{1}{2} < r_2 < 1$. It can be observed that g(t) increases on $t \in [0, r_1]$, decreases on $t \in (r_1, r_2)$, and increases on $t \in [r_2, 1]$. Also, g(t) takes positive values on $t \in [0, \frac{1}{2})$ and negative values on $t \in (\frac{1}{2}, 1]$, with $g(\frac{1}{2}) = 0$. Moreover, g(t) is maximum at $t = r_1$ with maximum value $g(r_1) = \sqrt{\frac{2}{\alpha}} \exp\left\{\frac{\alpha}{4} - \frac{1}{2}\right\}$, and g(t) is minimum at $t = r_2$ with minimum value $g(r_2) = -\sqrt{\frac{2}{\alpha}} \exp\left\{\frac{\alpha}{4} - \frac{1}{2}\right\}$. Since the functions g(u) and g(v) are positive on the quadrant R_1 and takes maximum at $u = r_1, v = r_1$, it follows that the product function g(u)g(v) has maximum value $[g(r_1)]^2 = \frac{2}{\alpha} \exp\left\{\frac{\alpha}{2} - 1\right\}$, on $(u, v) \in R_1$. Since the functions g(u) and g(v) is positive and has maximum value $[g(r_2)]^2 = \frac{2}{\alpha} \exp\left\{\frac{\alpha}{2} - 1\right\}$, on $(u, v) \in R_3$. Therefore, $\sup_{(u,v)\in R_1\cup R_3} \{g(u)g(v)\} = \frac{2}{\alpha} \exp\left\{\frac{\alpha}{2} - 1\right\}$. Similarly, we have found that the infimum of

the product function g(u)g(v) on the quadrant $R_2 \cup R_4$ is equal to $g(r_1)g(r_2) = -\frac{2}{\alpha} \exp\left\{\frac{\alpha}{2} - 1\right\}$. Now, using these values in Eq. (2.3), we get the feasible range of δ as

$$-\frac{1}{2\alpha}\exp\left\{1-\frac{\alpha}{2}\right\} \le \delta \le \frac{1}{2\alpha}\exp\left\{1-\frac{\alpha}{2}\right\}.$$
(2.5)

Thus, using the feasible range of the parameter δ given in Eq. (2.4) and Eq. (2.5), we propose the following bivariate copula

$$C(u,v;\delta,\alpha) = uv + \delta\left(1 - e^{\alpha(u-u^2)}\right)\left(1 - e^{\alpha(v-v^2)}\right), \quad (u,v) \in \mathcal{I}^2,$$
(2.6)

where α is a real valued parameter, and $|\delta| \leq \delta^*(\alpha)$. Here,

$$\delta^{\star}(\alpha) = \begin{cases} \frac{1}{\alpha^2}, & \text{if } \alpha \in (-\infty, 2] \setminus \{0\} \\\\ \frac{1}{2\alpha} \exp\left\{1 - \frac{\alpha}{2}\right\}, & \text{if } \alpha > 2. \end{cases}$$

Recall that our proposed copula reduced to the product copula when $\alpha = 0$. Also, the copula density $c(u, v) = \frac{\partial^2 C(u, v)}{\partial u \partial v}$ of the proposed copula is given by

$$c(u,v) = 1 + \alpha^2 \delta (1 - 2u) (1 - 2v) \exp\left\{\alpha \left(u - u^2 + v - v^2\right)\right\}, \quad (u,v) \in \mathcal{I}^2,$$
(2.7)

where $\alpha \in \mathbb{R}$ and $|\delta| \leq \delta^*(\alpha)$. The contour plots of the copula density are shown in Figure 1 for different choices of the parameters α and δ .

3 Measures of Dependence

Copula functions have been widely used for modelling dependence between random variables since they allow the separation of the dependence effect from the effects of the marginal distributions. The dependence between random variables can be measured by several well-known measures of dependence such as Spearman's Rho, Gini's Gamma Coefficient, Kendall's tau, Spearman's footrule coefficient (see Nelsen (2006)) and Blest's measure of rank correlation (see Genest and



Figure 1: Contour plots of copula density c(u, v) for various values of α and δ .

Plante, 2003). These measures are usually based on the ranks of observations rather than their actual values. The range of these dependence measures lies between -1 and 1. A negative value of dependence measure corresponds to negative dependence, zero for independence and a positive value for positive dependence. Most of these measures remain unchanged under strictly increasing transformations of random variables. Since the copula functions of a pair of random variables X and Y are invariant under strictly increasing transformations of X and Y, these dependence measures are expressible in terms of the copulas. Let C(u, v) be the copula function associated with the random variables X and Y. The following are important measures of dependence in terms of copula function:

• Spearman's Rho

$$\rho_c = 12 \int_0^1 \int_0^1 C(u, v) du dv - 3.$$

Gini's Gamma Coefficient

$$\gamma_c = 4 \left\{ \int_0^1 C(u, 1-u) du - \int_0^1 (u - C(u, u)) du \right\}.$$

• Kendall's Tau

$$\tau_c = 4 \int_0^1 \int_0^1 C(u, v) dC(u, v) - 1.$$

• Blest's Measure of Rank Correlation

$$\eta_c = 24 \int_0^1 \int_0^1 (1-u) C(u,v) du dv - 2.$$

• Spearman's footrule

$$\phi_c = 6 \int_0^1 C(u, u) du - 2.$$

Now, we will provide the expressions of the various measures of dependence for the proposed copula function (2.6).

Proposition 3.1. For the copula defined in Eq. (2.6), the Spearman's Rho ρ_C , and Gini's Gamma Coefficient γ_C are given by

$$\begin{split} \rho_{C} = \begin{cases} & 12\delta \left[1 - \sqrt{\frac{\pi}{|\alpha|}} e^{\alpha/4} erfi\left\{\frac{\sqrt{|\alpha|}}{2}\right\} \right]^{2}, \quad if \, \alpha < 0 \\ & 12\delta \left[1 - \sqrt{\frac{\pi}{\alpha}} e^{\alpha/4} erf\left\{\frac{\sqrt{\alpha}}{2}\right\} \right]^{2}, \quad if \, \alpha \geq 0, \end{cases} \\ \gamma_{C} = \begin{cases} & 8\delta \left[1 - 2\sqrt{\frac{\pi}{|\alpha|}} e^{\alpha/4} erfi\left\{\frac{\sqrt{|\alpha|}}{2}\right\} + \sqrt{\frac{\pi}{2|\alpha|}} e^{\alpha/2} erfi\left\{\sqrt{\frac{|\alpha|}{2}}\right\} \right], \quad if \, \alpha < 0 \\ & 8\delta \left[1 - 2\sqrt{\frac{\pi}{\alpha}} e^{\alpha/4} erf\left\{\frac{\sqrt{\alpha}}{2}\right\} + \sqrt{\frac{\pi}{2\alpha}} e^{\alpha/2} erf\left\{\sqrt{\frac{\alpha}{2}}\right\} \right], \quad if \, \alpha \geq 0, \end{cases} \end{split}$$

where $erf(t) = \frac{2}{\sqrt{\pi}} \int_0^t e^{-z^2} dz$ denotes the error function (see Abramowitz and Stegun (1972)) and $erfi(t) = \frac{2}{\sqrt{\pi}} \int_0^t e^{z^2} dz$ denotes the imaginary error function (see Marcinowski and Sadowski (2020)).

Remark 3.1. It can be verified that the expressions of other measures of dependence satisfied the following relation under the copula given in Eq. (2.6).

$$\eta_c = \rho_c = \frac{3}{2}\tau_c, \quad and \quad \phi_c = \frac{3}{4}\gamma_c.$$

Table 1 presents the numerical values of Spearman's rho and Gini's gamma coefficient of the new copula for different values of the copula parameter α . Since the new copula is symmetric, we have shown only the upper boundary values of the dependence measures in Table 1. The lower boundary value is the negative of upper boundary value. It is observed from the Table 1 that Spearman's rho $\rho_c \in [-0.5866, 0.5866]$ when $\alpha = 3.8$, thereby extending the range of the dependence measure Spearman's rho over the popular FGM copula and its various generalizations.

α	δ_{upper}	$ ho_{upper}$	γ_{upper}	α	δ_{upper}	ρ_{upper}	γ_{upper}
-3	0.1111	0.1899	0.1463	1.2	0.6944	0.4264	0.3473
-2.7	0.1372	0.2003	0.1547	1.5	0.4444	0.4544	0.3718
-2.4	0.1736	0.2113	0.1638	1.8	0.3086	0.4845	0.3984
-2.1	0.2268	0.2231	0.1736	2	0.25	0.506	0.4174
-1.8	0.3086	0.2358	0.1841	2.3	0.1871	0.5348	0.4434
-1.5	0.4444	0.2493	0.1954	2.6	0.1425	0.5561	0.4634
-1.2	0.6944	0.2638	0.2076	2.9	0.1099	0.571	0.4783
-0.9	1.2346	0.2794	0.2207	3.2	0.0858	0.5805	0.4888
-0.6	2.7778	0.2961	0.2349	3.5	0.0675	0.5855	0.4956
-0.3	11.1111	0.314	0.2502	3.8	0.0535	0.5866	0.4992
0	0	0	0	4.1	0.0427	0.5845	0.5002
0.3	11.1111	0.3541	0.2845	4.4	0.0342	0.5798	0.499
0.6	2.7778	0.3764	0.3038	4.7	0.0276	0.5729	0.4959
0.9	1.2346	0.4005	0.3247	5	0.0223	0.5643	0.4912

Table 1: Sperman's Rho and Gini's Gamma Coefficient for various values of α

In order to continue our discussion on dependence between random variables, there are some more dependence properties available in the literature. For example, quadrant dependence, totally positive of order 2 (TP_2) , and tail dependence coefficient. For more detailed discussion on these properties, one can see Lehmann (1966), Barlow and Proschan (1975), Drouet Mari and Kotz (2001), Nelsen (2006), Lai and Balakrishnan (2009) and Bhuyan et al. (2020). Now, we will discuss these dependence properties under the copula given in Eq. (2.6).

3.1 Quadrant Dependence

The random vector (X, Y) is said to be positively (negatively) quadrant dependence if

$$P(X \le x, Y \le y) \ge (\le) \ P(X \le x) P(Y \le y), \quad \forall \ (x, y) \in \mathbb{R}^2$$

Similarly, we call copula C(u, v) is positively (negatively) quadrant dependent if

$$C(u,v) \ge (\le) uv, \quad \forall (u,v) \in \mathcal{I}^2.$$

It can be verified that the quadrant dependence of the copula (2.6) depends only on the copula parameter δ . The proof is straightforward, so omitted. Thus, we have the following result.

Proposition 3.2. The copula (2.6) has positive (negative) quadrant dependence if $\delta \ge 0$ ($\delta \le 0$).

3.2 Totally Positive of Order 2 (TP_2)

A bivariate function g(x, y) is said to be totally positive of order 2 (TP_2) if

$$g(x_1, y_1)g(x_2, y_2) - g(x_2, y_1)g(x_2, y_1) \ge 0, \quad \forall x_1 < x_2, y_1 < y_2$$

The TP_2 property is a stronger concept of dependence. If a copula density c(u, v) possesses TP_2 property, then the associated copula C(u, v) has stochastic increasing (SI), right tail increasing (RTI) and positive quadrant dependence (PQD) properties. For more details, see Nelsen (2006), Karlin (1968), and Joe (1997).

Holland and Wang (1987) proved that the function g(x, y) has TP_2 property if

$$\zeta_g(x,y) = \frac{\partial^2 \ln g(x,y)}{\partial x \partial y} \ge 0, \quad \forall (x,y) \in \mathbb{R}^2.$$

Now, using the result of Holland and Wang (1987), we will prove the following result.

Proposition 3.3. The copula density (2.7) has TP_2 property if $\delta \ge 0$ and $\alpha \le 2$.

Proof. We have,

$$\zeta_{c}(u,v) = \frac{\partial^{2} \ln c(u,v)}{\partial x \partial y} \\ = \frac{\delta \alpha^{2} \left[2 - \alpha(1-2u)\right] \left[2 - \alpha(1-2v)\right] \exp\left\{\alpha \left((u-u^{2}) + (v-v^{2})\right)\right\}}{\left(1 + \delta \alpha^{2} \left(1 - 2u\right) \left(1 - 2v\right) \exp\left\{\alpha \left((u-u^{2}) + (v-v^{2})\right)\right\}\right)^{2}} \ge 0,$$

for every $\delta \geq 0$ and $\alpha \leq 2$.

3.3 Tail Dependence Coefficients

The tail dependence coefficients measure the level of dependency among the random variables in the upper-right quadrant and in the lower-left quadrant of \mathcal{I}^2 . In terms of copula, the upper and lower tail dependence coefficients, denoted by λ_U and λ_L respectively, are given by

$$\lambda_L = \lim_{u \to 0^+} \frac{C(u, u)}{u}, \quad \lambda_U = \lim_{u \to 1^-} \frac{1 - 2u + C(u, u)}{1 - u}.$$
(3.1)

It is known that $0 \leq \lambda_L \leq 1$ and $0 \leq \lambda_U \leq 1$. If $\lambda_L \in (0, 1]$, we say the copula C(u, v) has lower tail dependence, and if $\lambda_L = 0$, we say C(u, v) has no lower tail dependence. Similar interpretation can be made for λ_U (see Nelsen (2006), p. 214). Now, we will prove the following result.

Proposition 3.4. The copula (2.6) has no tail dependence.

Proof. Using Eq. (3.1), we have

$$\lambda_L = \lim_{u \to 0^+} \frac{u^2 + \delta \left(1 - e^{\alpha(u - u^2)}\right)^2}{u}$$

= $\delta \lim_{u \to 0^+} \frac{1}{u} \left(1 - \sum_{n=0}^{\infty} \frac{\alpha^n u^n (1 - u)^n}{n!}\right)^2$
= $\delta \lim_{u \to 0^+} \alpha^2 u (1 - u)^2 \left(\sum_{n=2}^{\infty} \frac{\alpha^{n-1} u^{n-1} (1 - u)^{n-1}}{n!}\right)^2$
= 0.

$$\lambda_U = \lim_{u \to 1^-} \frac{1 - 2u + u^2 + \delta \left(1 - e^{\alpha(u - u^2)}\right)^2}{1 - u}$$

=
$$\lim_{u \to 1^-} \frac{(1 - u)^2 + \delta \left(1 - \sum_{n=0}^{\infty} \frac{\alpha^n u^n (1 - u)^n}{n!}\right)^2}{(1 - u)}$$

=
$$\delta \lim_{u \to 1^-} \alpha^2 u^2 (1 - u) \left(\sum_{n=2}^{\infty} \frac{\alpha^{n-1} u^{n-1} (1 - u)^{n-1}}{n!}\right)^2$$

=
$$0.$$

In the next section, we will develop a bivariate Rayleigh distribution as an application of the proposed copula (2.6). We will study some statistical properties of new bivariate Rayleigh distribution, and a real data analysis involving the new distribution is also presented.

4 A New Bivariate Rayleigh distribution

Rayleigh distribution is one of the most popular models in medical sciences, engineering, particle physics and economics. A random variable X follows Rayleigh distribution with parameter λ , denoted by $Rayleigh(\lambda)$, if its cumulative distribution function (CDF) is given by $F(x; \lambda) =$ $1 - e^{-x^2/2\lambda^2}$, $x > 0, \lambda > 0$, and corresponding probability density function (PDF) is given by $f(x;\lambda) = \frac{x}{\lambda^2}e^{-x^2/2\lambda^2}$, $x > 0, \lambda > 0$. Let X and Y be two random variables having $Rayleigh(\lambda_1)$ and $Rayleigh(\lambda_2)$ distributions respectively, and the dependence between X and Y is modelled by the copula (2.6). Then, the join distribution function of X and Y is given by

$$F(x,y;\Theta) = \left(1 - e^{-x^2/2\lambda_1^2} - e^{-y^2/2\lambda_2^2} + e^{-\left(x^2/2\lambda_1^2 + y^2/2\lambda_2^2\right)}\right) + \delta\left(1 - e^{\alpha\left(e^{-x^2/2\lambda_1^2} - e^{-x^2/\lambda_1^2}\right)}\right)$$
$$\left(1 - e^{\alpha\left(e^{-y^2/2\lambda_2^2} - e^{-y^2/\lambda_2^2}\right)}\right),$$
(4.1)

where $x > 0, y > 0, \lambda_1 > 0, \lambda_2 > 0, \alpha \in \mathbb{R}, |\delta| \leq \delta^*(\alpha)$ and $\Theta = (\lambda_1, \lambda_2, \alpha, \delta)$. A non-negative random vector (X, Y) is said to follow bivariate Rayleigh distribution with parameters $\lambda_1, \lambda_2, \alpha$ and δ , if its joint CDF is given by Eq. (4.1) and is denoted by BRD $(\lambda_1, \lambda_2, \alpha, \delta)$. The corresponding joint density function is given by

$$f(x,y;\Theta) = \left(\frac{xy}{\lambda_1^2\lambda_2^2}e^{-(x^2/2\lambda_1^2+y^2/2\lambda_2^2)}\right) \left[1 + \delta\alpha^2 \left(2e^{-x^2/2\lambda_1^2} - 1\right) \left(2e^{-y^2/2\lambda_2^2} - 1\right) \\ \left(\exp\left\{\alpha \left(e^{-x^2/2\lambda_1^2} - e^{-x^2/\lambda_1^2} + e^{-y^2/2\lambda_2^2} - e^{-y^2/\lambda_2^2}\right)\right\}\right)\right],$$
(4.2)

where $x > 0, y > 0, \lambda_1 > 0, \lambda_2 > 0, \alpha \in \mathbb{R}, |\delta| \leq \delta^*(\alpha)$ and $\Theta = (\lambda_1, \lambda_2, \alpha, \delta)$. Surface plots of joint CDF (4.1) and joint density function (4.2) of the BRD family are shown in Figure 2. These figures are constructed using Matlab R2021b. Now, we will provide expressions for conditional



Figure 2: Surface plots of F(x, y) and f(x, y) of the BRD distribution for $\lambda_1 = 3$, $\lambda_2 = 2$, $\delta = 0.5$, $\alpha = 3.8$.

distribution and product moments.

Proposition 4.1. Let $(X, Y) \sim BRD(\lambda_1, \lambda_2, \alpha, \delta)$. Then

- (i) $X \sim Rayleigh(\lambda_1)$ and $Y \sim Rayleigh(\lambda_2)$,
- (ii) the conditional density function of X given Y = y is

$$f(x|y) = \left(\frac{x}{\lambda_1^2} e^{-x^2/2\lambda_1^2}\right) \left[1 + \delta\alpha^2 \left(2e^{-x^2/2\lambda_1^2} - 1\right) \left(2e^{-y^2/2\lambda_2^2} - 1\right) \\ \left(e^{\alpha \left(e^{-x^2/2\lambda_1^2} - e^{-x^2/\lambda_1^2} + e^{-y^2/2\lambda_2^2} - e^{-y^2/\lambda_2^2}\right)}\right)\right],$$

(iii) the conditional distribution function of X given Y = y is

$$F(x|y) = \left(1 - e^{-x^2/2\lambda_1^2}\right) + \delta\alpha \left(e^{\alpha \left(e^{-x^2/2\lambda_1^2} - e^{-x^2/\lambda_1^2}\right)} - 1\right) \left(2e^{-y^2/2\lambda_2^2} - 1\right) \left(e^{\alpha \left(e^{-y^2/2\lambda_2^2} - e^{-y^2/\lambda_2^2}\right)}\right)$$

where $x > 0, y > 0, \lambda_1 > 0, \lambda_2 > 0, \alpha \neq 0, |\delta| \leq \delta^*(\alpha)$ and $\Theta = (\lambda_1, \lambda_2, \alpha, \delta)$.

Proposition 4.2. Let $(X, Y) \sim BRD(\lambda_1, \lambda_2, \alpha, \delta)$. Then (r, s)-th order product moments can be expressed as

$$\begin{split} E(X^r Y^s) = &\lambda_1^r \lambda_2^s 2^{(r+s)/2} \Gamma\left(1+r/2\right) \Gamma\left(1+s/2\right) \\ & \left[1 + \delta \alpha^2 \left(\sum_{k=0}^\infty \frac{\alpha^k}{k!} \sum_{t=0}^k \frac{(-1)^t}{\lambda_1^2} \binom{k}{t} \left(2\left(k+t+2\right)^{-\frac{r+2}{2}} - \left(k+t+1\right)^{-\frac{r+2}{2}}\right)\right) \\ & \left(\sum_{k=0}^\infty \frac{\alpha^k}{k!} \sum_{t=0}^k \frac{(-1)^t}{\lambda_2^2} \binom{k}{t} \left(2\left(k+t+2\right)^{-\frac{s+2}{2}} - \left(k+t+1\right)^{-\frac{s+2}{2}}\right)\right)\right], \end{split}$$

where $\Gamma(t)$ denotes the usual gamma function.

4.1 Real Data Application

We consider the UEFA Champions League data set from 2004 to 2006, reported in Meintanis (2007). In this data set, X and Y represent the time (in minutes) of the first goal scored by Team-A and Team-B, respectively. To check whether the marginal distributions of X and Y support the Rayleigh distribution, we perform Kolmogorov-Smirnov (KS) one-sample test. The KS-test's results suggest that X supports Rayleigh distribution with parameter $\hat{\lambda}_1 = 32.14599$ (p-value=0.934 and KS statistic value=0.088515). Similarly, Y also supports Rayleigh distribution with parameter $\hat{\lambda}_2 = 28.172255$ (p-value=0.07727 and KS statistic value=0.20968). We fit the proposed bivariate Rayleigh distribution, and the results are shown in Table 2. We compare the new BRD model with Marshall Olkin's bivariate exponential distribution (BMOED) by Meintanis (2007), bivariate generalized exponential distribution (BGED) by Mirhosseini et al. (2015), and bivariate generalized Rayleigh distribution (BGRD) proposed by Pathak and Vellaisamy (2022). We use the log-likelihood (LL) function, Akaike Information Criteria (AIC) and Bayesian Information Criteria (BIC) as the comparison criteria. The formulas for AIC and BIC are given by

AIC =
$$2k - 2\ln L$$
, and BIC = $k\ln n - 2\ln L$,

where k is the number parameters in the model, n is the sample size and L is the maximum value of the likelihood function. From Table 2, it is clear that bivariate Rayleigh distribution provides a better fit over BGED, BMOED and BGRD for the UEFA champions league data set.

Table 2:	ML	estimates,	LL,	AIC,	and	BIC	values	for	the	bivarite	distributions	using	UEFA
Champio	n's L	eague data	set.										

Bivariate Distribution	ML Estimates	LL	AIC	BIC
BGED	$\hat{\alpha}_1 = 0.0244, \hat{\alpha}_2 = 0.0304, \hat{\theta} = 0.999$	-340.5234	687.0468	691.8795
BMOED	$\hat{\lambda}_1 = 0.012, \hat{\lambda}_2 = 0.014, \hat{\lambda}_3 = 0.022$	-339.006	684.012	688.8448
BGRD	$\hat{b}_1 = 0.000530, \hat{b}_2 = 0.000836, \hat{\theta} = 0.40331$	-331.879	664.589	672.6436
BRD	$\hat{\lambda}_1 = 33.39429, \hat{\lambda}_2 = 28.08949, \hat{\delta} = 10.39829, \hat{\alpha} = 0.2871858$	-327.256	664.512	668.9557

5 Conclusion and Future Direction

This paper proposes a new bivariate symmetric copula exhibiting positive and negative dependence. The main features of the copula are: (i) it has a simple mathematical structure, (ii) it has a wider dependence range when compared to FGM copula and its generalizations, and (iii) there is no lower and upper tail dependence. Using the proposed copula, we developed a new bivariate Rayleigh distribution (BRD) and discussed some statistical properties. The proposed bivariate model provides a better fit for a real data set. Since we considered only the symmetric version of the bivariate copula, the asymmetric version is still an open problem for new researchers.

Acknowledgments

The first author is thankful to the Indian Institute of Technology Indore for providing financial support for pursuing research.

Conflict of interest

The authors declare no potential conflict of interests.

References

- Abd Elaal, M. K. and R. S. Jarwan (2017). Inference of bivariate generalized exponential distribution based on copula functions. *Applied Mathematical Sciences* 11(24), 1155–1186.
- Abramowitz, M. and I. A. Stegun (1972). Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. National Bureau of Standards Applied Mathematics Series 55. Tenth Printing. ERIC.
- Achcar, J. A., F. A. Moala, M. H. Tarumoto, and L. F. Coladello (2015). A bivariate generalized exponential distribution derived from copula functions in the presence of censored data and covariates. *Pesquisa Operacional* 35(1), 165–186.
- Almetwally, E. M. and H. Z. Muhammed (2020). On a bivariate fréchet distribution. J Stat Appl Probab 9(1), 1–21.
- Bairamov, I. and S. Kotz (2002). Dependence structure and symmetry of huang-kotz fgm distributions and their extensions. *Metrika* 56(1), 55–72.
- Barlow, R. E. and F. Proschan (1975). Statistical theory of reliability and life testing: probability models. Technical report, Florida State Univ Tallahassee.
- Bekrizadeh, H., G. Parham, and M. Zadkarami (2015). Pak. j. statist. 2015 vol. 31 (1), 95-106 an asymmetric generalized fgm copula and its properties. *Pak. J. Statist* 31(1), 95-106.
- Bhuyan, P., S. Ghosh, P. Majumder, and M. Mitra (2020). A bivariate life distribution and notions of negative dependence. *Stat* 9(1), e276.
- Chesneau, C. (2022). A note on a simple polynomial-sine copula. Asian Journal of Mathematics and Applications 2, 1–14.

Drouet Mari, D. and S. Kotz (2001). Correlation and dependence. Imperial College Press.

- Durante, F. and C. Sempi (2016). *Principles of copula theory*, Volume 474. CRC press Boca Raton.
- El-Sherpieny, E., H. Z. Muhammed, and E. M. Almetwally (2018). Fgm bivariate weibull distribution. In Proceedings of the Annual Conference in Statistics (53rd), Computer Science, and Operations Research, Institute of Statistical Studies and Research, Cairo University, pp. 55–77.
- Farlie, D. J. (1960). The performance of some correlation coefficients for a general bivariate distribution. *Biometrika* 47(3/4), 307–323.
- Genest, C. and J.-F. Plante (2003). On blest's measure of rank correlation. *Canadian Journal* of *Statistics 31*(1), 35–52.
- Hofert, M., I. Kojadinovic, M. Mächler, and J. Yan (2018). *Elements of copula modeling with R.* Springer.
- Holland, P. W. and Y. J. Wang (1987). Dependence function for continuous bivariate densities. Communications in Statistics-Theory and Methods 16(3), 863–876.
- Huang, J. S. and S. Kotz (1999). Modifications of the farlie-gumbel-morgenstern distributions. a tough hill to climb. *Metrika* 49(2), 135–145.
- Joe, H. (1997). Multivariate models and multivariate dependence concepts. CRC press.
- Karlin, S. (1968). Total positivity, Volume 1. Stanford University Press.
- Kim, J.-M., E. A. Sungur, T. Choi, and T.-Y. Heo (2011). Generalized bivariate copulas and their properties. *Model Assisted Statistics and Applications* 6(2), 127–136.
- Kundu, D. and R. C. Gupta (2017). On bivariate birnbaum-saunders distribution. American Journal of Mathematical and Management Sciences 36(1), 21–33.
- Lai, C. D. and N. Balakrishnan (2009). *Continuous bivariate distributions*. Springer.
- Lehmann, E. L. (1966). Some concepts of dependence. The Annals of Mathematical Statistics 37(5), 1137–1153.
- Mai, J.-F. and M. Scherer (2017). Simulating copulas: stochastic models, sampling algorithms, and applications, Volume 6. World Scientific.
- Marcinowski, J. and M. Sadowski (2020). Using the erfi function in the problem of the shape optimization of the compressed rod. International Journal of Applied Mechanics and Engineering 25(2), 75–87.
- Meintanis, S. G. (2007). Test of fit for marshall-olkin distributions with applications. *Journal* of Statistical Planning and inference 137(12), 3954–3963.
- Mirhosseini, S. M., M. Amini, D. Kundu, and A. Dolati (2015). On a new absolutely continuous bivariate generalized exponential distribution. *Statistical Methods & Applications* 24(1), 61–83.
- Nelsen, R. B. (2006). An introduction to copulas. Springer Science & Business Media.

- Pathak, A. and P. Vellaisamy (2022). A bivariate generalized linear exponential distribution: properties and estimation. *Communications in Statistics-Simulation and Computation* 51(9), 5426–5446.
- Sklar, M. (1959). Fonctions de repartition an dimensions et leurs marges. Publ. inst. statist. univ. Paris 8, 229–231.