

# The Log-Sobolev inequality for a submanifold in manifolds with asymptotic non-negative intermediate Ricci curvature

Jihye Lee, Fabio Ricci

## Abstract

We prove a sharp Log-Sobolev inequality for submanifolds of a complete non-compact Riemannian manifold with asymptotic non-negative intermediate Ricci curvature and Euclidean volume growth. Our work extends a result of Dong-Lin-Lu [12] which already generalizes Yi-Zheng's [34] and Brendle's [5].

## 1 Introduction

Log-Sobolev inequalities have been already studied in mathematical physics and information theory in the 50's ([29], [14]) then it has been extensively studied in analysis and geometry (see e.g. [17], [16], [24], [22], [13], [10], [25], [26]), with recent developments including their applications to heat kernel and Hamilton-Jacobi equations, inequalities in convex geometry, and their role in optimal transport. Brendle [5] obtained a sharp log-Sobolev inequality for submanifolds in Euclidean space, improving [13], using the Alexandroff-Bakelman-Pucci (ABP) method and optimal transport. Later Yi and Zheng [34] extended this result to Riemannian manifolds with non-negative sectional curvature and Euclidean volume growth conditions. Subsequently Dong-Lin-Lu [12] further extended this to non-negative asymptotic sectional curvature. In this paper, we prove that the logarithmic Sobolev inequality holds under non-negative intermediate Ricci curvature. Furthermore, we extend our result to the asymptotic non-negative intermediate Ricci curvature by introducing error terms that depends only on the curvature's decay at infinity, as in [12].

Let  $(M^n, g)$  be a complete non-compact Riemannian manifold. For  $1 \leq k \leq n - 1$  we consider a  $k$ -dimensional subspace  $P$  of a tangent space  $T_x M$  at  $x \in M$ . Given any tangent vector  $v \in T_x M$  with  $v \perp P$ , the  $k$ -intermediate Ricci curvature ( $k$ th-Ricci curvature) with respect to  $P$  in the direction of  $v$  is defined as

$$\text{Ric}_k(P, v) = \sum_{i=1}^k \text{Sec}(v, e_i) |v|^2,$$

---

Jihye Lee: UC Santa Barbara, Department of Mathematics, email: jihye@ucsb.edu.

Fabio Ricci: UC Santa Barbara, Department of Mathematics, email: FabioRicci@ucsb.edu.

where  $\{e_1, \dots, e_k\}$  is an orthonormal basis of  $P$ . The  $k$ -intermediate Ricci curvature interpolates between Ricci and sectional curvature. A Riemannian manifold has non-negative  $k$ -Ricci curvature if  $\text{Ric}_k(P, v) \geq 0$  for any  $x \in M$ ,  $k$ -dimensional subspace  $P \subset T_x M$ , and a unit tangent vector  $v \in T_x M$  perpendicular to  $P$  (note that some papers [30, 21] require the stronger condition that this holds for all  $v$ , not necessary just the perpendicular ones). This condition is denoted by  $\text{Ric}_k \geq 0$ . In particular, it exhibits a monotonicity property: if  $n \leq m$ , then  $\text{Ric}_n \geq 0$  implies  $\text{Ric}_m \geq 0$ . This curvature condition has been well studied to explore the gap of the global results with sectional curvature bounds and Ricci curvature bounds, see for example: [28, 32, 9, 19, 18, 27, 21, 23, 30]. Our work follows this spirit and generalizes Yi-Zheng's result with sectional curvature lower bound (Theorem 1.1, [34]) and the corresponding asymptotic extension by Dong-Lin-Lu (Theorem 1.1, [12]). Ketterer-Mondino [21] noted that it is possible to characterize lower bounds of  $k$ -intermediate Ricci curvature via optimal transport. We will adopt this approach, developed in [23] and [30], to prove our main result.

Let  $M$  be an  $n$ -dimensional non-compact Riemannian manifold with non-negative Ricci curvature. The asymptotic volume ratio of  $M$  is defined as

$$\theta := \lim_{r \rightarrow \infty} \frac{|B_r(p)|}{\omega_n r^n},$$

where  $p$  is some fixed point in  $M$ ,  $\omega_n$  is the volume of the unit ball in Euclidean space  $\mathbb{R}^n$ , and  $|B_r(p)|$  is the volume of a ball of radius  $r$  centered at  $p$  in  $M$ . The Bishop-Gromov volume comparison assures that the limit exists, it does not depend on the choice of  $p$  and that  $\theta \leq 1$ . We say that  $M$  has Euclidean volume growth if  $\theta > 0$ .

In this paper we are using the definition of asymptotically non negative sectional (resp. Ricci) curvature given in Abresch [1] (see also [2], [20], [35]). Zhu (see theorem 2.1 in [36]) proved the equivalent of the classical Bishop-Gromov volume comparison Theorem with  $\text{Ric} \geq 0$  by replacing  $\mathbb{R}^n$  with a different model space. We will need to define the equivalent of the usual asymptotic volume ratio.

**Definition 1.1** (Abresch [1]). An  $n$ -dimensional complete non-compact Riemannian manifold  $(M, g)$  with base point  $o$  has asymptotically non-negative Sectional curvature (Ricci curvature, respectively) if and only if there exists any non-negative, non-increasing function  $\lambda : [0, \infty) \rightarrow [0, \infty)$  such that the following holds:

- (1)  $b_0(\lambda) = \int_0^\infty t\lambda(t)dt < \infty$
- (2)  $\text{Sec} \geq -\lambda(d(o, p))$  at each point  $p \in M$ . ( $\text{Ric} \geq -(n-1)\lambda(d(o, p))$ , respectively)

The first condition also implies that  $b_1(\lambda) = \int_0^\infty \lambda(t)dt < \infty$ . To extend this notion of asymptotically non-negative sectional curvature to intermediate  $k$ -Ricci curvature, one can simply replace the second condition with:

$$\text{Ric}_k \geq -k\lambda(d(o, p)) \text{ at each point } p \in M.$$

More precisely,  $\text{Ric}_k(P, v) \geq -k\lambda(d(o, p))$  for any  $k$  dimensional subspace  $P \subset T_p M$  and unit tangent vector  $v$  perpendicular to  $P$ .

The usual non-negative sectional curvature (Ricci or  $\text{Ric}_k$  respectively) condition is equivalent to requiring  $\lambda \equiv 0$ . It's immediately evident from the definition to see

that this class of manifolds includes those with non-negative sectional curvature (Ricci or  $\text{Ric}_k$  respectively) outside a compact set and asymptotically flat manifolds as well. Furthermore, the monotonicity of the intermediate curvature still holds: if  $k_1 \leq k_2$ , then  $\text{Ric}_{k_1} \geq -k_1\lambda(d(o, p))$  implies  $\text{Ric}_{k_2} \geq -k_2\lambda(d(o, p))$ .

In this setting, we also replace  $\theta$ , the asymptotic volume ratio of  $M$ , with another similar quantity that will keep track of the geometry at infinity. To achieve this, we define  $h(t)$  as the unique solution of the following ODE:

$$\begin{cases} h''(t) = \lambda(t)h(t), \\ h(0) = 0, h'(0) = 1. \end{cases}$$

We now define the asymptotic volume ratio of  $M$  with respect to  $h$  by

$$\theta_h = \lim_{r \rightarrow \infty} \frac{|B_r(o)|}{n\omega_n \int_0^r h^{n-1}(t)dt},$$

In the above definition,  $|B_r(o)|$  represents the volume of the ball of radius  $r$  centered at  $o$  in  $M$  and  $\omega_n$  represents the volume of the unit ball in  $\mathbb{R}^n$ . In [36], to prove the corresponding version of the Bishop-Gromov volume comparison in this setting, Zhu noted that the following function of  $r$  is non-increasing:

$$\frac{|B_r(o)|}{n\omega_n \int_0^r h^{n-1}(t)dt}.$$

This ensures that the limit exists and  $\theta_h$  is well-defined. Similarly, we say that  $M$  has Euclidean volume growth if  $\theta_h > 0$ . In particular, when a manifold has non-negative intermediate Ricci curvature,  $\lambda(t) \equiv 0$ ,  $h(t) = t$ , and  $\theta_h$  is the usual asymptotic volume ratio  $\theta$ . In the second section of this paper we will show a connection between  $\theta_h$  and an integral of a certain Gaussian function. To state our theorem, we first define, for any non-negative  $t$ , a decreasing function  $P(t)$ :

$$P(t) = (4\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^N} e^{-\frac{(|x|+t)^2}{4}} dx$$

Notice that  $P(0) = 1$ . Below is our main result.

**Theorem 1.2.** *Let  $M^{n+m}$  be a complete, non-compact manifold of dimension  $n + m$  with asymptotically non-negative  $\text{Ric}_k$ , where  $k = \min\{n - 1, m - 1\}$  and Euclidean volume growth. Assume that  $\Sigma^n$  is an  $n$ -dimensional compact submanifold without a boundary. Then, for any positive smooth function  $f$ , we have*

$$\begin{aligned} \int_{\Sigma} f \left( \log f + \log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n + 4b_1^2n^2 + (n + m - 1) \log \frac{1 + b_0}{e^{2r_0b_1 + b_0}} \right) dV(x) \\ - \int_{\Sigma} \frac{|\nabla_{\Sigma} f|^2}{f} - \int_{\Sigma} f |H|^2 \leq \int_{\Sigma} f dV(x) \left( \log \int_{\Sigma} f(x) dV(x) \right), \end{aligned} \quad (1)$$

where  $\theta_h$  is the asymptotic volume ratio of  $M$  with respect to  $h$  and  $H$  is the mean curvature vector of  $\Sigma$  and  $\alpha = (n + m - 1)(2r_0b_1 + b_0)$ , where  $r_0 = \max_{x \in \Sigma} d(o, x)$ .

Similar to the Michael-Simon-Sobolev inequality, the inequality (1) contains a term that involves the mean curvature and depends on the asymptotic volume ratio of  $M$ . If  $\lambda \equiv 0$ , the usual non-negative  $\text{Ric}_k$  condition, in other words  $b_0 = b_1 = 0$ , then our Theorem 1.2 becomes:

**Corollary 1.3.** *Let  $M^{n+m}$  be a complete, non-compact manifold of dimension  $n + m$  with Euclidean volume growth and  $\text{Ric}_k \geq 0$ , where  $k = \min\{n - 1, m - 1\}$ . Let  $\Sigma^n$  be a  $n$ -dimensional compact submanifold without a boundary. Then, for any positive smooth function  $f$ , we have*

$$\begin{aligned} & \int_{\Sigma} f \left( \log f + n + \frac{n}{2} \log(4\pi) + \log \theta \right) dV - \int_{\Sigma} \frac{|\nabla^{\Sigma} f|^2}{f} dV - \int_{\Sigma} f |H|^2 dV \\ & \leq \left( \int_{\Sigma} f dV \right) \log \left( \int_{\Sigma} f dV \right), \end{aligned} \quad (2)$$

where  $\theta$  is the asymptotic volume ratio of  $M$  and  $H$  is the Mean curvature vector of  $\Sigma$ .

This gives the following two consequences by choosing  $f \equiv 1$  in (2) and apply the argument in Section 4 of [34].

**Corollary 1.4.** *Let  $M^{n+m}$  be a complete, non-compact manifold of dimension  $n + m$  with non-negative intermediate  $k$ -Ricci curvature, where  $k = \min\{n - 1, m - 1\}$ . Suppose  $M$  has a Euclidean volume growth. Then, there is no closed minimal submanifold of dimension  $n$  in  $M$ .*

**Corollary 1.5.** *Let  $M^{n+m}$  be a complete, non-compact manifold of dimension  $n + m$  with non-negative intermediate  $k$ -Ricci curvature, where  $k = \min\{n - 1, m - 1\}$ . Suppose there is a closed minimal submanifold of dimension  $n$  in  $M$ . Then, the asymptotic volume ratio of  $M$  is zero.*

In the case of any hypersurface ( $m = 1$ ) under non-negative Ricci curvature condition, these two corollaries have been proved by Agostiniani-Fogagnolo-Mazzieri (see Theorem 1.6 in [3]). We do not recover this case in our result since  $k = \min\{n - 1, m - 1\}$  would be zero. On the other hand, for higher dimension and codimension, these corollaries are also new in the usual non-negative  $k$ -intermediate Ricci curvature setting. When  $k = 1$ , the standard  $\text{Ric}_1 \geq 0$  is equivalent to the non-negative sectional curvature condition. Thus, we recover the corollaries in Yi-Zheng (Corollaries 1.1 and 1.2 in [34]).

The same argument (Section 4, [34]) cannot be applied in the case of asymptotic non-negative  $k$ -intermediate Ricci curvature setting because the error term we have introduced is not scale-invariant and will blow up if we blow down the metric as in [34].

Note that the previous two corollaries do not hold for  $\text{Ric} \geq 0$  total dimension 4; a counterexample is the Eguchi-Hanson metric on  $TS^2$ , which is Ricci flat with Euclidean volume growth and has a totally geodesic submanifold  $S^2$ . See, for example, [4, Page 270].

The proof of Theorem 1.2 relies on the Alexandroff-Bakelman-Pucci (ABP) method, a standard technique in Euclidean space [15]. Cabrè [7] extended this method to manifolds. The classical estimate in  $\mathbb{R}^n$  uses affine functions, but on manifolds, hyperplanes

are replaced with paraboloids interpreted as graphs of the distance squared relative to a point. If a lower bound condition on the curvature is added, this method has been successfully used to prove geometric inequalities (see for example, [33, 31]). In particular, Brendle [6] used this method to prove a monotonicity result in terms of the Jacobian of a certain transport map, while Wang [30] and Ma-Wu [23] independently generalized this method to  $k$ -intermediate Ricci curvature. It will be a key ingredient in our proof.

**Acknowledgements:** The authors gratefully acknowledge the valuable insights and guidance provided by their advisor Guofang Wei through helpful discussions. We would like to thank Frank Morgan for the references [29] and [14]. We would also like to acknowledge Lingen Lu for bringing the manuscript [12] to our attention. Their contribution allowed us to simplify Section 2 in the first version of our work. And thanks Jing Wu for reaching us with the updated version of their paper [23] and useful comments.

## 2 Asymptotic Estimate

In this section we recall an estimate for  $\theta_h$ , which will be used in the final steps of the proof in the next section. We first recall:

**Lemma 2.1** (Lemma 2.1, [34]). *Let  $M$  be a complete non-compact Riemannian manifold of dimension  $N$  with non-negative Ricci curvature. Then,*

$$\theta = \lim_{r \rightarrow \infty} \left( \frac{(4\pi)^{-\frac{N}{2}}}{r^N} \int_M e^{-\frac{d(x,p)^2}{4r^2}} dV(x) \right)$$

for any  $p \in M$ .

The proof is based on the Bishop-Gromov volume comparison. However, in our asymptotic case, we utilize the inequality  $r \leq h(r)$  instead. The upper bound for  $h$ , as provided in [36], is  $h(r) \leq e^{b_0 r}$ . Generally, the presence of  $b_0 \neq 0$  precludes us from replacing  $r^N$  with  $h^N$  in the above lemma. Following Dong-Lin-Lu [12], it is preferable to replace  $r^N$  with  $rh^{N-1}$ , doing so, we first recall the definition of  $P(t)$ :

$$P(t) = (4\pi)^{-\frac{N}{2}} \int_{\mathbb{R}^N} e^{-\frac{(|x|+t)^2}{4}} dx.$$

**Lemma 2.2** (Lemma 2.2, [12]). *Let  $M$  be a complete non-compact Riemannian manifold of dimension  $N$  with asymptotic non-negative Ricci curvature. Then*

$$P(t)\theta_h = \lim_{r \rightarrow \infty} \frac{(4\pi)^{-\frac{N}{2}}}{rh(r)^{N-1}} \int_M e^{-\frac{(d(x,o)+t)^2}{4}} dV(x), \quad (3)$$

where  $o$  is the base point of  $M$ .

This was proved in [12], we are giving the same proof of this statement also here for completeness:

*Proof.* First notice that we can apply De L'Hopital's rule to the limit of  $\theta_h$  and obtain

$$\theta_h = \lim_{r \rightarrow \infty} \frac{|B_r(o)|}{N\omega_N \int_0^r h^{N-1}(t)dt} = \lim_{r \rightarrow \infty} \frac{|\partial B_r(o)|}{N\omega_N h^{N-1}(r)}$$

Now we can compute

$$\begin{aligned} \lim_{r \rightarrow \infty} \frac{(4\pi)^{-\frac{N}{2}}}{rh(r)^{N-1}} \int_M e^{-\frac{(d(x,o)+t)^2}{4}} dV(x) &= \lim_{r \rightarrow \infty} (4\pi)^{-\frac{N}{2}} \int_0^\infty \frac{|\partial B_{rs}(o)|}{h^{N-1}(r)} e^{-\frac{(s+t)^2}{4}} ds \\ &= \lim_{r \rightarrow \infty} (4\pi)^{-\frac{N}{2}} \int_0^\infty \frac{|\partial B_{rs}(o)|}{h^{N-1}(rs)} \frac{h^{N-1}(rs)}{h^{N-1}(r)} e^{-\frac{(s+t)^2}{4}} ds \\ &= (4\pi)^{-\frac{N}{2}} N\omega_N \theta \int_0^\infty s^{N-1} e^{-\frac{(s+t)^2}{4}} ds \\ &= \theta (4\pi)^{-\frac{N}{2}} \int_{\mathbb{R}^n} e^{-\frac{(|x|+t)^2}{4}} dx \\ &= P(t)\theta, \end{aligned}$$

where we used the monotone convergence theorem when we pass the limit under the integral sign.  $\square$

For the next lemma, we aim to substitute the base point  $o$  on the right-hand side of inequality (3) with  $\psi(x)$ , where  $\psi$  is a Borel map whose image is compact.

**Lemma 2.3.** *Let  $M$  be a complete non-compact Riemannian manifold of dimension  $n$  with asymptotic non-negative intermediate Ricci curvature. Then we have*

$$\lim_{r \rightarrow \infty} \frac{(4\pi)^{-\frac{N}{2}}}{rh(r)^{N-1}} \int_M e^{-\frac{(d(x,\psi(x))+t)^2}{4}} dV(x) = \lim_{r \rightarrow \infty} \frac{(4\pi)^{-\frac{N}{2}}}{rh(r)^{N-1}} \int_M e^{-\frac{(d(x,o)+t)^2}{4}} dV(x),$$

where  $\psi : M \rightarrow K$  is any Borel map where  $K \subset M$  a compact subset.

The above lemma is proven, similar to the proof of Lemma 2.2 in [34], by utilizing the triangle inequality.

### 3 Proof of Theorem 1.2

In this section, we prove log Sobolev inequality with non-negative asymptotic intermediate Ricci curvature by following the papers [11, 12, 34].

Let  $M^{n+m}$  be a complete non-compact manifold of dimension  $n+m$  with asymptotically non-negative  $\text{Ric}_k$  and Euclidean volume growth, where  $k = \min\{n-1, m-1\}$ . Assume that  $\Sigma^n$  is an  $n$ -dimensional compact submanifold without a boundary and let  $f$  be any positive smooth function.

We first assume that  $\Sigma$  is connected, which is needed for the existence of a solution to a differential equation. The inequality (1) is invariant when we scale a function  $f$ . Thus, by scaling, we may assume that

$$\int_\Sigma f \log f dV - \int_\Sigma \frac{|\nabla^\Sigma f|^2}{f} dV - \int_\Sigma f |H|^2 dV = 0. \quad (4)$$

Indeed, we can scale the function  $f$  to satisfy the equation (4) because  $\log(cf)$  can be any real number by choosing a suitable constant  $c$ . The left-hand-side of equation (4) comes from the inequality (1). Thus, it suffices to show that

$$\begin{aligned} & \int_{\Sigma} f \left( \log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n + 4b_1^2n^2 + (n+m-1) \log \frac{1+b_0}{e^{2r_0b_1+b_0}} \right) dV \\ & \leq \left( \int_{\Sigma} f dV \right) \log \left( \int_{\Sigma} f dV \right). \end{aligned}$$

In order to prove Isoperimetric or Sobolev inequalities using the ABP method (see [8] and related literature), we need to consider a suitable PDE on a submanifold  $\Sigma$ . To apply our assumption (4), we consider a differential equation as follows:

$$\operatorname{div}_{\Sigma}(f\nabla^{\Sigma}u) = f \log f - \frac{|\nabla^{\Sigma}f|^2}{f} - f|H|^2 \text{ on } \Sigma, \quad (5)$$

where  $\nabla^{\Sigma}$  is the induced Levi-Civita connection on  $\Sigma$ . We do not need to specify a boundary condition since  $\partial\Sigma = \emptyset$ . Using standard PDE theory (see [15] Chapter 6), since  $f$  is a positive function, the operator  $u \mapsto \operatorname{div}_{\Sigma}(f\nabla^{\Sigma}u)$  has Fredholm index 0. Hence, we can find a smooth solution  $u : \Sigma \rightarrow \mathbb{R}$ .

We define a contact set  $A_r$  as the set of all points  $(\bar{x}, \bar{y}) \in T^{\perp}\Sigma$  satisfying

$$ru(x) + \frac{1}{2}d(x, \exp_{\bar{x}}(r\nabla^{\Sigma}u(\bar{x}) + r\bar{y}))^2 \geq ru(\bar{x}) + \frac{1}{2}r^2(|\nabla^{\Sigma}u(\bar{x})|^2 + |\bar{y}|^2) \quad (6)$$

for all  $x \in \Sigma$  as introduced in [6]. This is a natural extension of the Cabrè's idea [7] to the submanifold case.

Let  $\Phi_t : T^{\perp}\Sigma \rightarrow M$  be a map defined by

$$\Phi_t(x, y) = \exp_x(t\nabla^{\Sigma}u(x) + ty).$$

Then we have the following lemma, for which we will recall the proof for the reader's convenience.

**Lemma 3.1** (Lemma 3.2, [34]). *For each  $r \in (0, \infty)$ , we have  $\Phi_r(A_r) = M$ .*

*Proof.* Take any  $p \in M$ . Since  $\Sigma$  is compact, there is  $\bar{x} \in \Sigma$  where the function  $x \mapsto ru(x) + \frac{1}{2}d(x, p)^2$  attains its minimum. Let  $\bar{\gamma}$  be a minimizing geodesic such that  $\bar{\gamma}(0) = \bar{x}$  and  $\bar{\gamma}(r) = p$ . Then the geodesic  $\bar{\gamma}$  minimizes the functional  $u(\gamma(0)) + \frac{1}{2} \int_0^r |\gamma'(t)|^2 dt$  among all smooth curves  $\gamma$  with  $\gamma(0) \in \Sigma$  and  $\gamma(r) = p$ . By the first variational formula, we have

$$\nabla^{\Sigma}u(\bar{x}) - \bar{\gamma}'(0) \in T_{\bar{x}}^{\perp}\Sigma.$$

That is, there is  $\bar{y} \in T_{\bar{x}}^{\perp}\Sigma$  satisfying

$$\nabla^{\Sigma}u(\bar{x}) + \bar{y} = \bar{\gamma}'(0).$$

We can easily check that  $\Phi_r(\bar{x}, \bar{y}) = p$  and  $(\bar{x}, \bar{y}) \in A_r$  using the geodesic  $\bar{\gamma}$ , which implies that  $p \in \Phi_r(A_r)$ .  $\square$

As we see in the above proof, we can define a map  $\psi : M \rightarrow \Sigma$  as  $\psi(p)$  is the point where the function  $ru(x) + \frac{1}{2}d(x, p)^2$  attains its minimum. Then by Lemma 3.1, we have

$$\int_M e^{-\left(\frac{d(\psi(p), p)}{2r} + 2b_1 n\right)^2} dV(p) \leq \int_M \left( \int_{\{(x, y) \in A_r \mid \Phi_r(x, y) = p\}} e^{-\left(\frac{d(x, \Phi_r(x, y))}{2r} + 2b_1 n\right)^2} d\mathcal{H}^0 \right) dV(p), \quad (7)$$

where  $\mathcal{H}^0$  is the counting measure. The left hand side of the inequality (7) is related to the asymptotic volume ratio  $\theta_h$  as we showed in Lemma 2.2. Also, we apply Area formula with the map  $\Phi_r : A_r \rightarrow M$  to the right hand side of (7). That is,

$$\begin{aligned} & \int_M \left( \int_{\{(x, y) \in A_r \mid \Phi_r(x, y) = p\}} e^{-\left(\frac{d(x, \Phi_r(x, y))}{2r} + 2b_1 n\right)^2} d\mathcal{H}^0 \right) dV(p) \\ &= \int_{A_r} e^{-\left(\frac{d(x, \Phi_r(x, y))}{2r} + 2b_1 n\right)^2} |\det D\Phi_r(x, y)| dV(x, y) \\ &= \int_{\Sigma} \left( \int_{T_{\bar{x}}^{\perp} \Sigma} e^{-\left(\frac{d(x, \Phi_r(x, y))}{2r} + 2b_1 n\right)^2} |\det D\Phi_r(x, y)| \chi_{A_r}(x, y) dy \right) dV(x). \end{aligned} \quad (8)$$

We now want to estimate the integrand in the right hand side of (8). Take any  $r > 0$  and  $(\bar{x}, \bar{y})$  in  $A_r$ . Define a geodesic  $\bar{\gamma}(t) = \exp_{\bar{x}}(t\nabla^{\Sigma}u(\bar{x}) + t\bar{y})$  on  $[0, r]$ . Let  $\{e_1, \dots, e_n\}$  be an orthonormal basis of  $T_{\bar{x}}\Sigma$  such that  $\text{Hess}_{\Sigma}u(e_i, e_j) - \langle II(e_i, e_j), \bar{y} \rangle$  is diangonal. Let us denote  $E_i$  by a parallel transport vector field of  $e_i$  along  $\bar{\gamma}$  for each  $1 \leq i \leq n$ . Take any orthonormal frame  $\{e_{n+1}, \dots, e_{n+m}\}$  of  $T^{\perp}\Sigma$  near  $\bar{x}$  such that  $\langle \nabla_{e_i} e_{\alpha}, e_{\beta} \rangle = 0$  at  $\bar{x}$  for all  $1 \leq i \leq n$  and  $n+1 \leq \alpha, \beta \leq n+m$ .

For each  $1 \leq i \leq n$ , let us consider a Jacobi field  $X_i(t)$  along  $\bar{\gamma}(t)$  with the initial conditions

$$\begin{cases} X_i(0) = e_i \\ \langle X_i'(0), e_j \rangle = \text{Hess}_{\Sigma}u(e_i, e_j) - \langle II(e_i, e_j), \bar{y} \rangle & \text{for all } 1 \leq j \leq n \\ \langle X_i'(0), e_{\alpha} \rangle = \langle II(e_i, \nabla^{\Sigma}u(\bar{x})), e_{\alpha} \rangle & \text{for all } n+1 \leq \alpha \leq n+m. \end{cases}$$

For each  $n+1 \leq \alpha \leq n+m$ , consider a Jacobi field  $X_{\alpha}(t)$  along  $\bar{\gamma}(t)$  with the initial conditions

$$X_{\alpha}(0) = 0 \quad \text{and} \quad X'_{\alpha}(0) = e_{\alpha}.$$

Define  $(n+m) \times (n+m)$  matrices  $P(t)$  and  $S(t)$  satisfying  $P_{ij}(t) = \langle X_i(t), E_j(t) \rangle$  and  $S_{ij}(t) = \bar{R}(\bar{\gamma}'(t), E_i(t), \bar{\gamma}'(t), E_j(t))$  for  $1 \leq i, j \leq n+m$ . Since  $X_i(t)$  is a Jacobi field along  $\bar{\gamma}(t)$ , we have  $P''(t) = -P(t)S(t)$ . Define  $Q(t) = P^{-1}(t)P'(t)$ . Then

$$Q'(t) = -(Q(t))^2 - S(t).$$

Let  $tr_n Q(t) = \sum_{i=1}^n Q_{ii}(t)$  and  $tr_m Q(t) = \sum_{\alpha=n+1}^{n+m} Q_{\alpha\alpha}(t)$ . Then

$$(tr_n Q(t))' + \frac{1}{n}(tr_n Q(t))^2 \leq -tr_n S(t)$$

and

$$(tr_m Q(t))' + \frac{1}{m}(tr_m Q(t))^2 \leq -tr_m S(t),$$



where  $tr_n S(t)$  and  $tr_m S(t)$  are defined as a partial trace of  $S(t)$ . Let us note that

$$tr_n S(t) = \sum_{i=1}^n \bar{R}(\bar{\gamma}'(t), E_i(t), \bar{\gamma}'(t), E_i(t)).$$

However, we cannot apply curvature assumption directly because  $\bar{\gamma}'(t)$  may not be perpendicular to the plane spanned by  $E_1(t), \dots, E_n(t)$ . Since  $E_i(t)$  is a parallel vector field along  $\bar{\gamma}(t)$ , it is enough to consider the angle at  $t = 0$ , denoted by  $a$ , between the vector  $\bar{\gamma}'(0)$  and the tangent plane  $T_{\bar{x}}\Sigma$ . Since  $\bar{\gamma}'(0) = \nabla^\Sigma u(\bar{x}) + \bar{y}$ , depending on the vectors  $\nabla^\Sigma u(\bar{x})$  and  $\bar{y}$  the angle is determined.

Let us consider the case  $\nabla^\Sigma u(\bar{x}) \neq 0$  and  $\bar{y} \neq 0$ . In this case,  $\bar{\gamma}'(0)$  is not perpendicular to the tangent plane. After we change the basis as in [23], we get

$$\begin{aligned} tr_n S(t) &= \sin^2(a) |\bar{\gamma}'(0)|^2 Ric_n \left( \frac{\bar{\gamma}'(t)}{|\bar{\gamma}'(t)|}, P_1(t) \right) + \cos^2(a) |\bar{\gamma}'(0)|^2 Ric_{n-1} \left( \frac{\bar{\gamma}'(t)}{|\bar{\gamma}'(t)|}, P_2(t) \right) \\ &\geq -n \sin^2(a) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) - (n-1) \cos^2(a) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) \\ &= (\cos^2(a) - n) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))), \end{aligned} \quad (9)$$

where  $P_1(t)$  is an  $n$ -dimensional subspace spanned by  $E_2(t), \dots, E_{n+1}(t)$  and  $P_2(t)$  is an  $(n-1)$ -dimensional subspace spanned by  $E_2(t), \dots, E_n(t)$ . Similarly, we have

$$tr_m S(t) \geq (\sin^2(a) - m) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))). \quad (10)$$

Indeed, the inequalities (9) and (10) hold for the case  $\nabla^\Sigma u(\bar{x}) = 0$  or  $\bar{y} = 0$ . More specifically, for the case  $\nabla^\Sigma u(\bar{x}) = 0$ , since  $\bar{\gamma}'(0) = \bar{y} \in T_{\bar{x}}^\perp \Sigma$ , the angle  $a = \frac{\pi}{2}$  and we have

$$\begin{aligned} tr_n S(t) &= |\bar{\gamma}'(0)|^2 Ric_n(P_3(t), \bar{\gamma}'(t)) \\ &\geq -n |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) \\ &= (\cos^2(a) - n) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) \end{aligned}$$

and

$$\begin{aligned} tr_m S(t) &= \sum_{\alpha=n+1}^{n+m} \bar{R}(\bar{\gamma}'(t), E_\alpha(t), \bar{\gamma}'(t), E_\alpha(t)) \\ &= \sum_{\alpha=n+2}^{n+m} \bar{R}(\bar{\gamma}'(t), E_\alpha(t), \bar{\gamma}'(t), E_\alpha(t)) \\ &= |\bar{\gamma}'(0)|^2 Ric_{m-1}(P_4(t), \bar{\gamma}'(t)) \\ &\geq -(m-1) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}'(t))) \\ &\geq (\sin^2(a) - m) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))), \end{aligned}$$

where  $P_3(t) = Span(E_1(t), \dots, E_n(t))$  and  $P_4(t) = Span(E_{n+2}(t), \dots, E_{n+m}(t))$  with the assumption  $E_{n+1} = \frac{\bar{y}}{|\bar{y}|}$ . In the same manner, for the case  $\bar{y} = 0$ , since  $\bar{\gamma}'(0) = \nabla^\Sigma u(\bar{x}) \in T_{\bar{x}}\Sigma$ , by assuming  $E_1(t) = \frac{\bar{\gamma}'(t)}{|\bar{\gamma}'(t)|}$ , we can verify that the same inequalities (9) and (10) hold. In all, for  $(\bar{x}, \bar{y}) \in A_r \setminus \{(0, 0)\}$ , we obtain

$$\begin{cases} (tr_n Q(t))' + \frac{1}{n} (tr_n Q(t))^2 \leq (n - \cos^2(a)) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) \\ (tr_m Q(t))' + \frac{1}{m} (tr_m Q(t))^2 \leq (m - \sin^2(a)) |\bar{\gamma}'(0)|^2 \lambda(d(o, \bar{\gamma}(t))) \end{cases}$$

by combining inequalities (9) and (10) with Riccati equation. It follows from the triangle inequality  $d(o, \bar{\gamma}(t)) \geq |d(o, \bar{x}) - d(\bar{x}, \bar{\gamma}(t))|$  and the definition of  $A_r$  that

$$\begin{cases} (tr_n Q(t))' + \frac{1}{n}(tr_n Q(t))^2 \leq (n - \cos^2(a))|\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \\ (tr_m Q(t))' + \frac{1}{m}(tr_m Q(t))^2 \leq (m - \sin^2(a))|\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right). \end{cases}$$

Let  $\phi(t) = e^{\frac{1}{n} \int_0^t \sum_{i=1}^n Q_{ii}(\tau) d\tau}$ . Then

$$\begin{cases} \phi''(t) \leq \frac{n - \cos^2(a)}{n} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \phi(t) \\ \phi'(0) = \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \\ \phi(0) = 1. \end{cases}$$

To use comparison theorems with  $\phi(t)$ , let us define  $\psi_1(t)$  and  $\psi_2(t)$  as a solution to each PDE similarly to [11]:

$$\begin{cases} \psi_1''(t) = \frac{n - \cos^2(a)}{n} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \psi_1(t) \\ \psi_1(0) = 0, \psi_1'(0) = 1 \end{cases}$$

and

$$\begin{cases} \psi_2''(t) = \frac{n - \cos^2(a)}{n} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \psi_2(t) \\ \psi_2(0) = 1, \psi_2'(0) = 0. \end{cases}$$

Then we see that a function  $\psi(t) = \psi_2(t) + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \psi_1(t)$  satisfies the following PDE.

$$\begin{cases} \psi''(t) = \frac{n - \cos^2(a)}{n} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \psi(t) \\ \psi'(0) = \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \\ \psi(0) = 1. \end{cases}$$

By comparing  $\frac{\phi'(t)}{\phi(t)}$  with  $\frac{\psi'(t)}{\psi(t)}$ , we have

$$tr_n Q(t) = n \cdot \frac{\phi'(t)}{\phi(t)} \leq n \cdot \frac{\psi'(t)}{\psi(t)} = \frac{\psi_2' + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \psi_1'}{\psi_2 + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \psi_1}.$$

Define

$$\bar{\phi}(t) = t e^{\frac{1}{m} \int_0^t \sum_{\alpha=n+1}^{n+m} (Q_{\alpha\alpha}(\tau) - \frac{1}{t}) d\tau}.$$

Then

$$\begin{cases} \bar{\phi}'' \leq \frac{m - \sin^2(a)}{m} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \bar{\phi} \\ \bar{\phi}(0) = 0, \quad \bar{\phi}'(0) = 1. \end{cases}$$

To estimate  $\frac{\bar{\phi}'}{\bar{\phi}}$ , let us introduce a  $C^2$  solution  $\bar{\psi}$  to the PDE

$$\begin{cases} \bar{\psi}'' = \frac{m - \sin^2(a)}{m} |\bar{\gamma}'(0)|^2 \lambda \left( \left| d(o, \bar{x}) - t\sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} \right| \right) \bar{\psi} \\ \bar{\psi}(0) = 0, \quad \bar{\psi}'(0) = 1. \end{cases}$$

Thus,

$$\operatorname{tr}_m Q(t) = m \cdot \frac{\bar{\phi}'}{\bar{\phi}} \leq m \cdot \frac{\bar{\psi}'}{\bar{\psi}}.$$

Since  $\frac{d}{dt}(\det P(t)) = \det P(t) \operatorname{tr} Q(t)$ , we have

$$\frac{d}{dt} \log(\det P(t)) = \operatorname{tr} Q(t) \leq n \cdot \frac{\psi'}{\psi} + m \cdot \frac{\bar{\psi}'}{\bar{\psi}} = \frac{d}{dt} (n \log(\psi) + m \log(\bar{\psi})) = \frac{d}{dt} \log(\psi^n \bar{\psi}^m).$$

By integrating the above inequality from  $\epsilon$  to  $t$  and  $\epsilon \rightarrow 0$ , we get

$$\begin{aligned} \det P(t) &\leq \psi^n(t) \bar{\psi}^m(t) = \left( \psi_2(t) + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \psi_1(t) \right)^n \bar{\psi}^m(t) \\ &= \left( \frac{\psi_2(t)}{\psi_1(t)} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n \psi_1^n(t) \bar{\psi}^m(t) \end{aligned}$$

Moreover, comparison theorems imply those inequalities

$$\begin{cases} \frac{\psi_2}{\psi_1} \leq \frac{2b_1(n - \cos^2(a))}{n} \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} \\ \psi_1(t) \leq t e^{\frac{n - \cos^2(a)}{n} (2b_1 d(o, \bar{x}) + b_0)} \\ \bar{\psi}(t) \leq t e^{\frac{m - \sin^2(a)}{m} (2b_1 d(o, \bar{x}) + b_0)}. \end{cases}$$

Thus,

$$\begin{aligned} &\det P(t) \\ &\leq \left( \frac{\psi_2(t)}{\psi_1(t)} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n \psi_1^n(t) \bar{\psi}^m(t) \\ &\leq \left( \frac{2b_1(n - \cos^2(a))}{n} \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n \\ &\quad \cdot t^n e^{(n - \cos^2(a))(2b_1 d(o, \bar{x}) + b_0)} \cdot t^m e^{(m - \sin^2(a))(2b_1 d(o, \bar{x}) + b_0)} \\ &\leq \left( 2b_1 \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n t^{n+m} e^{(n+m-1)(2b_1 d(o, \bar{x}) + b_0)} \\ &\leq \left( 2b_1 \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n t^{n+m} e^{(n+m-1)(2b_1 r_0 + b_0)}, \end{aligned} \tag{11}$$

The following lemma can be also found in the proof of Lemma 3.7 in [34].

**Lemma 3.2.** *For  $(x, y) \in A_r$ , we have:*

$$\Delta_\Sigma u(x) - \langle H(x), y \rangle \leq \log f(x) + \frac{|\nabla^\Sigma u|^2 + |y|^2}{4} - \frac{|2H(x) + y|^2}{4}.$$

*Proof.* Since  $\operatorname{div}_\Sigma(f \nabla^\Sigma u) = f \log f - \frac{|\nabla^\Sigma f|^2}{f} - f|H|^2$ , we get

$$f \Delta_\Sigma u + \langle \nabla^\Sigma f, \nabla^\Sigma u \rangle = f \log f - \frac{|\nabla^\Sigma f|^2}{f} - f|H|^2.$$

Since  $f(x) \neq 0$  for all  $x \in \Sigma$ , we have

$$\Delta_\Sigma u - \langle H(x), y \rangle = \log f - \frac{|\nabla^\Sigma f|^2}{f^2} - |H(x)|^2 - \frac{\langle \nabla^\Sigma f, \nabla^\Sigma u \rangle}{f} - \langle H(x), y \rangle. \quad (12)$$

Note that

$$\begin{aligned} \frac{|\nabla^\Sigma f|^2}{f^2} + \frac{\langle \nabla^\Sigma f, \nabla^\Sigma u \rangle}{f} &= \frac{4|\nabla^\Sigma f|^2 + 4\langle \nabla^\Sigma f, f\nabla^\Sigma u \rangle}{4f^2} \\ &= \frac{4|\nabla^\Sigma f|^2 + 4\langle \nabla^\Sigma f, f\nabla^\Sigma u \rangle + |f\nabla^\Sigma u|^2}{4f^2} - \frac{|f\nabla^\Sigma u|^2}{4f^2} \\ &= \frac{|2\nabla^\Sigma f + f\nabla^\Sigma u|^2}{4f^2} - \frac{|\nabla^\Sigma u|^2}{4} \end{aligned} \quad (13)$$

and

$$|H(x)|^2 + \langle H(x), y \rangle = \frac{|2H(x)|^2 + \langle 4H(x), y \rangle + |y|^2}{4} - \frac{|y|^2}{4} = \frac{|2H(x) + y|^2}{4} - \frac{|y|^2}{4}. \quad (14)$$

Combining identities (12), (13), and (14), we obtain

$$\begin{aligned} \Delta_\Sigma u(x) - \langle H(x), y \rangle &= \log f - \frac{|\nabla^\Sigma f|^2}{f^2} - |H|^2 - \frac{\langle \nabla^\Sigma f, \nabla^\Sigma u \rangle}{f} - \langle H(x), y \rangle \\ &= \log f + \frac{|\nabla^\Sigma u(x)|^2 + |y|^2}{4} - \frac{|2\nabla^\Sigma f + f\nabla^\Sigma u|^2}{4f^2} - \frac{|2H(x) + y|^2}{4} \\ &\leq \log f + \frac{|\nabla^\Sigma u(x)|^2 + |y|^2}{4} - \frac{|2H(x) + y|^2}{4}. \end{aligned}$$

□

**Lemma 3.3.** *For  $(\bar{x}, \bar{y}) \in A_r \setminus \{(0, 0)\}$ , we have:*

$$|\det D\Phi_t(\bar{x}, \bar{y})| \leq e^{\alpha + \frac{n}{t} - n - 4b_1^2 n^2} t^{m+n} f \exp \left( \left( \frac{d(\bar{x}, \Phi_t(\bar{x}, \bar{y}))}{2t} + 2b_1 n \right)^2 \right) \exp \left( -\frac{|2H + \bar{y}|^2}{4} \right),$$

where  $\alpha = (n + m - 1)(2r_0 b_1 + b_0)$ .

*Proof.* Combining inequality (11) and Lemma 3.2, we obtain

$$\begin{aligned} \det P(t) &\leq \left( 2b_1 \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} + \frac{1}{n} (\Delta_\Sigma u(\bar{x}) - \langle H(\bar{x}), \bar{y} \rangle) \right)^n t^{n+m} e^\alpha \\ &\leq \left( 2b_1 \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{1}{t} + \frac{1}{n} \left( \log f + \frac{|\nabla^\Sigma u|^2 + |\bar{y}|^2}{4} - \frac{|2H(\bar{x}) + \bar{y}|^2}{4} \right) \right)^n t^{n+m} e^\alpha \end{aligned}$$

By the inequality  $\lambda \leq e^{\lambda-1}$ , we have

$$\begin{aligned}
& |\det D\Phi_t(\bar{x}, \bar{y})| \\
& \leq \exp \left( 2b_1 n \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{n}{t} + \log f + \frac{|\nabla^\Sigma u|^2 + |\bar{y}|^2}{4} - \frac{|2H + \bar{y}|^2}{4} - n \right) t^{n+m} e^\alpha \\
& = e^{\alpha + \frac{n}{t} - n} t^{m+n} f \exp \left( 2b_1 n \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{|\nabla^\Sigma u|^2 + |\bar{y}|^2}{4} - \frac{|2H + \bar{y}|^2}{4} \right) \\
& \leq e^{\alpha + \frac{n}{t} - n} t^{m+n} f(\bar{x}) \exp \left( 2b_1 n \sqrt{|\nabla^\Sigma u(\bar{x})|^2 + |\bar{y}|^2} + \frac{|\nabla^\Sigma u|^2 + |\bar{y}|^2}{4} - \frac{|2H(\bar{x}) + \bar{y}|^2}{4} \right) \\
& = e^{\alpha + \frac{n}{t} - n} t^{m+n} f(\bar{x}) \exp \left( \left( \frac{\sqrt{|\nabla^\Sigma u|^2 + |\bar{y}|^2}}{2} + 2b_1 n \right)^2 \right) \exp \left( -\frac{|2H + \bar{y}|^2}{4} - 4b_1^2 n^2 \right) \\
& = e^{\alpha + \frac{n}{t} - n - 4b_1^2 n^2} t^{m+n} f(\bar{x}) \exp \left( \left( \frac{d(\bar{x}, \Phi_t(\bar{x}, \bar{y}))}{2t} + 2b_1 n \right)^2 \right) \exp \left( -\frac{|2H(\bar{x}) + \bar{y}|^2}{4} \right).
\end{aligned}$$

□

By combining inequalities (7) and (8) and Lemma 3.3, we obtain

$$\begin{aligned}
& \int_M e^{-\left(\frac{d(\psi(p), p)}{2t} + 2b_1 n\right)^2} dV(p) \\
& \leq \int_\Sigma \left( \int_{T_x^\perp \Sigma} e^{-\left(\frac{d(x, \Phi_t(x, y))}{2t} + 2b_1 n\right)^2} |\det D\Phi_t(x, y)| \chi_{A_t}(x, y) dy \right) dV(x) \\
& = \int_\Sigma \left( \int_{T_x^\perp \Sigma} e^{\alpha + \frac{n}{t} - n - 4b_1^2 n^2} t^{m+n} f(x) \exp \left( -\frac{|2H(x) + y|^2}{4} \right) dy \right) dV(x) \\
& = t^{n+m} e^{\frac{n}{t} - n + \alpha - 4b_1^2 n^2} (4\pi)^{\frac{m}{2}} \int_\Sigma f(x) dV(x).
\end{aligned}$$

Then

$$\frac{(4\pi)^{-\frac{n+m}{2}}}{t(h(t))^{n+m-1}} \int_M e^{-\left(\frac{d(\psi(p), p)}{2t} + 2b_1 n\right)^2} dV(p) \leq \frac{(4\pi)^{-\frac{n}{2}} t^{n+m-1}}{(h(t))^{n+m-1}} e^{\frac{n}{t} - n + \alpha - 4b_1^2 n^2} \int_\Sigma f(x) dV(x) \quad (15)$$

Let us take limit  $t \rightarrow \infty$  to the both sides of the above inequality. Then by Lemma 2.2 and 2.3, we get

$$P(4b_1 n)\theta_h \leq \lim_{t \rightarrow \infty} \frac{t^{n+m-1}}{(h(t))^{n+m-1}} (4\pi)^{-\frac{n}{2}} e^{\frac{n}{t} - n + \alpha - 4b_1^2 n^2} \int_\Sigma f(x) dV(x).$$

Since

$$\lim_{t \rightarrow \infty} \frac{t^{n+m-1}}{(h(t))^{n+m-1}} = \left( \lim_{t \rightarrow \infty} \frac{t}{h(t)} \right)^{n+m-1} = \left( \lim_{t \rightarrow \infty} \frac{1}{h'(t)} \right)^{n+m-1} \leq \frac{1}{(1+b_0)^{n+m-1}},$$

we have

$$P(4b_1 n)\theta_h \leq (4\pi)^{-\frac{n}{2}} e^{-n + \alpha - 4b_1^2 n^2} \frac{1}{(1+b_0)^{n+m-1}} \int_\Sigma f(x) dV(x).$$

By taking a logarithmic function, we obtain

$$\log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n - \alpha + 4b_1^2n^2 + (n+m-1) \log(1+b_0) \leq \log \int_{\Sigma} f(x) dV(x),$$

or equivalently,

$$\log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n + 4b_1^2n^2 + (n+m-1) \log \frac{1+b_0}{e^{2r_0b_1+b_0}} \leq \log \int_{\Sigma} f(x) dV(x).$$

Multiplying the inequality by  $f$  and integrate it over  $\Sigma$ , then

$$\begin{aligned} & \int_{\Sigma} f \left( \log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n + 4b_1^2n^2 + (n+m-1) \log \frac{1+b_0}{e^{2r_0b_1+b_0}} \right) dV(x) \\ & \leq \int_{\Sigma} f dV(x) \left( \log \int_{\Sigma} f(x) dV(x) \right). \end{aligned}$$

By adding our assumption

$$\int_{\Sigma} f \log f - \int_{\Sigma} \frac{|\nabla^{\Sigma} f|^2}{f} - \int_{\Sigma} f |H|^2 = 0,$$

we conclude that

$$\begin{aligned} & \int_{\Sigma} f \left( \log f + \log P(4b_1n)\theta_h + \frac{n}{2} \log(4\pi) + n + 4b_1^2n^2 + (n+m-1) \log \frac{1+b_0}{e^{2r_0b_1+b_0}} \right) dV(x) \\ & - \int_{\Sigma} \frac{|\nabla^{\Sigma} f|^2}{f} - \int_{\Sigma} f |H|^2 \leq \int_{\Sigma} f dV(x) \left( \log \int_{\Sigma} f(x) dV(x) \right). \end{aligned}$$

It now remains to show the case when  $\Sigma$  is disconnected. We follow Brendle [5] and notice that

$$a \log(a) + b \log(b) < a \log(a+b) + b \log(a+b) = (a+b) \log(a+b).$$

The above simple inequality will take care of the right hand side when we apply inequality (1) on each connected component individually. And this completes our proof of Theorem 1.2.

## References

- [1] Uwe Abresch. Lower curvature bounds, Toponogov's theorem, and bounded topology. *Ann. Sci. École Norm. Sup. (4)*, 18(4):651–670, 1985.
- [2] Uwe Abresch and Detlef Gromoll. On complete manifolds with nonnegative Ricci curvature. *J. Amer. Math. Soc.*, 3(2):355–374, 1990.
- [3] Virginia Agostiniani, Mattia Fogagnolo, and Lorenzo Mazzieri. Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature. *Invent. Math.*, 222(3):1033–1101, 2020.

- [4] Michael T. Anderson. Short geodesics and gravitational instantons. *J. Differential Geom.*, 31(1):265–275, 1990.
- [5] Simon Brendle. The logarithmic Sobolev inequality for a submanifold in Euclidean space. *Comm. Pure Appl. Math.*, 75(3):449–454, 2022.
- [6] Simon Brendle. Sobolev inequalities in manifolds with nonnegative curvature. *Comm. Pure Appl. Math.*, 2022. <https://doi.org/10.1002/cpa.22070>.
- [7] Xavier Cabré. Nondivergent elliptic equations on manifolds with nonnegative curvature. *Comm. Pure Appl. Math.*, 50(7):623–665, 1997.
- [8] Xavier Cabré. Isoperimetric, Sobolev, and eigenvalue inequalities via the Alexandroff-Bakelman-Pucci method: a survey. *Chinese Ann. Math. Ser. B*, 38(1):201–214, 2017.
- [9] Yousef K. Chahine. Volume estimates for tubes around submanifolds using integral curvature bounds. *J. Geom. Anal.*, 30(4):4071–4091, 2020.
- [10] Dario Cordero-Erausquin, Bruno Nazaret, and Cédric Villani. A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities. *Adv. Math.*, 182(2):307–332, 2004.
- [11] Yuxin Dong, Hezi Lin, and Lingen Lu. Sobolev inequalities in manifolds with asymptotically nonnegative curvature, 2022. arXiv:2203.14624.
- [12] Yuxing Dong, Hezi Lin, and Lingen Lu. The logarithmic sobolev inequality for a submanifold in manifolds with asymptotically nonnegative sectional curvature. *To appear in Acta Mathematica Scientia, English Series*.
- [13] Klaus Ecker. Logarithmic Sobolev inequalities on submanifolds of Euclidean space. *J. Reine Angew. Math.*, 522:105–118, 2000.
- [14] Paul G. Federbush. Partially alternate derivation of a result of nelson. *Journal of Mathematical Physics*, 10:50–52, 1969.
- [15] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [16] Leonard Gross. Logarithmic Sobolev inequalities. *Amer. J. Math.*, 97(4):1061–1083, 1975.
- [17] Leonard Gross. Logarithmic Sobolev inequalities and contractivity properties of semigroups. In *Dirichlet forms (Varenna, 1992)*, volume 1563 of *Lecture Notes in Math.*, pages 54–88. Springer, Berlin, 1993.
- [18] Luis Guijarro and Frederick Wilhelm. A softer connectivity principle. *Communications in Analysis and Geometry*, 2018.
- [19] Luis Guijarro and Frederick Wilhelm. Restrictions on submanifolds via focal radius bounds. *Mathematical Research Letters*, 27(1):115–139, 2020.

- [20] Atsushi Kasue. Harmonic functions with growth conditions on a manifold of asymptotically nonnegative curvature. I. In *Geometry and analysis on manifolds (Katata/Kyoto, 1987)*, volume 1339 of *Lecture Notes in Math.*, pages 158–181. Springer, Berlin, 1988.
- [21] Christian Ketterer and Andrea Mondino. Sectional and intermediate Ricci curvature lower bounds via optimal transport. *Adv. Math.*, 329:781–818, 2018.
- [22] Michel Ledoux. Concentration of measure and logarithmic Sobolev inequalities. In *Séminaire de Probabilités, XXXIII*, volume 1709 of *Lecture Notes in Math.*, pages 120–216. Springer, Berlin, 1999.
- [23] Hui Ma and Jing Wu. Sobolev inequalities in manifolds with nonnegative intermediate Ricci curvature, 2023. arXiv:2303.09285.
- [24] J. H. Michael and L. M. Simon. Sobolev and mean-value inequalities on generalized submanifolds of  $R^n$ . *Comm. Pure Appl. Math.*, 26:361–379, 1973.
- [25] Shin-ichi Ohta and Asuka Takatsu. Equality in the logarithmic Sobolev inequality. *Manuscripta Math.*, 162(1-2):271–282, 2020.
- [26] Grisha Perelman. The entropy formula for the ricci flow and its geometric applications, 2002. arXiv:math/0211159.
- [27] Philipp Reiser and David J. Wraith. Intermediate Ricci curvatures and Gromov’s Betti number bound, 2022. arXiv:2208.13438.
- [28] Zhongmin Shen. On complete manifolds of nonnegative kth-Ricci curvature. *Trans. Amer. Math. Soc.*, 338(1):289–310, 1993.
- [29] A. J. Stam. Some inequalities satisfied by the quantities of information of fisher and shannon. *Inf. Control.*, 2:101–112 Eqn 2.3, 1959.
- [30] Kai-Hsiang Wang. Optimal transport approach to Michael-Simon-Sobolev inequalities in manifolds with intermediate Ricci curvature lower bounds, 2022. arXiv:2209.06796.
- [31] Yu Wang and Xiangwen Zhang. An Alexandroff-Bakelman-Pucci estimate on Riemannian manifolds. *Adv. Math.*, 232:499–512, 2013.
- [32] Hung-Hsi Wu. Manifolds of partially positive curvature. *Indiana Univ. Math. J.*, 36(3):525–548, 1987.
- [33] Chao Xia and Xiangwen Zhang. ABP estimate and geometric inequalities. *Comm. Anal. Geom.*, 25(3):685–708, 2017.
- [34] Chengyang Yi and Yu Zheng. The logarithmic Sobolev inequality for a submanifold in manifold with nonnegative sectional curvature. *To appear in Chin. Ann. Math.*, 2023. arXiv:2104.05045.
- [35] Yuntao Zhang. Open manifolds with asymptotically nonnegative Ricci curvature and large volume growth. *Proc. Amer. Math. Soc.*, 143(11):4913–4923, 2015.



- [36] Shun-Hui Zhu. A volume comparison theorem for manifolds with asymptotically nonnegative curvature and its applications. *Amer. J. Math.*, 116(3):669–682, 1994.