

Solutions of Spinorial Yamabe-type Problems on S^m : Perturbations and Applications

Takeshi Isobe Tian Xu

Abstract

This paper is part of a program to establish the existence theory for the conformally invariant Dirac equation

$$D_g\psi = f(x)|\psi|_g^{\frac{2}{m-1}}\psi$$

on a closed spin manifold (M, g) of dimension $m \geq 2$ with a fixed spin structure, where $f : M \rightarrow \mathbb{R}$ is a given function. The study on such nonlinear equation is motivated by its important applications in Spin Geometry: when $m = 2$, a solution corresponds to an isometric immersion of the universal covering \widetilde{M} into \mathbb{R}^3 with prescribed mean curvature f ; meanwhile, for general dimensions and $f \equiv \text{constant}$, a solution provides an upper bound estimate for the Bär-Hijazi-Lott invariant.

Comparing with the existing issues, the aim of this paper is to establish multiple existence results in a new geometric context, which have not been considered in the previous literature. Precisely, in order to examine the dependence of solutions of the aforementioned nonlinear Dirac equations on geometrical data, concrete analysis are made for two specific models on the manifold (S^m, g) : the geometric potential f is a perturbation from constant with $g = g_{S^m}$ being the canonical round metric; and $f \equiv 1$ with the metric g being a perturbation of g_{S^m} , that is not conformally flat somewhere on S^m . The proof is variational: solutions of these problems are found as critical points of their corresponding energy functionals. The emphasis is that the solutions are always degenerate: they appear as critical manifolds of positive dimension. This is very different from most situations in elliptic PDEs and classical critical point theory. As corollaries of the existence results, multiple distinct embedded spheres in \mathbb{R}^3 with a common mean curvature are constructed, and furthermore, a strict inequality estimate for the Bär-Hijazi-Lott invariant on S^m , $m \geq 4$, is derived, which is the first result of this kind in the non-locally-conformally-flat setting.

Keywords. Dirac operator; Spinorial Yamabe equation; Perturbation method; conformal class.

T. Isobe: Graduate School of Economics, Hitotsubashi University, 2-1 Naka, Kunitachi, Tokyo 186-8601, Japan; e-mail: t.isobe@r.hit-u.ac.jp

T. Xu: Center for Applied Mathematics, Tianjin University, Nankai District, Tianjin 300072, China; e-mail: xutian@amss.ac.cn

T. Isobe is supported by JSPS KAKENHI Grant Number 20K03674 and T. Xu is supported by the National Science Foundation of China (NSFC 11601370) and the Alexander von Humboldt Foundation of Germany

Mathematics Subject Classification (2010): Primary 53C27; Secondary 35R01

Contents

1	Introduction	2
1.1	Existence results for the PDE problems	5
1.2	Geometric applications	8
1.3	Overview of the method	10
1.4	Organization of the paper	11
2	The precise framework	11
2.1	Reduction to problems on Euclidean spaces	11
2.2	Configuration spaces	12
2.3	The perturbation method with degenerate conditions	13
3	Perturbations of the geometric potential	18
3.1	The topological degree	18
3.2	Proof of the main result	24
4	Perturbations of the background metric	29
4.1	Bourguignon-Gauduchon identification	30
4.2	Basic calculations	31
4.3	Proof the main result	37
A	Appendix	40
A.1	An explanation of the (k, p) -elementary matrix	40
A.2	Some basic estimates	41
A.3	Proofs of Lemma 4.6 and Proposition 4.7	43

1 Introduction

Let (M, g, σ) be an m -dimensional closed spin manifold, $m \geq 2$, with a fixed Riemannian metric g and a fixed spin structure $\sigma : P_{Spin}(M) \rightarrow P_{SO}(M)$. For the metric g , we obtain a spinor bundle $\mathbb{S}(M) \rightarrow M$, which is endowed with a natural hermitian metric $|\cdot|_g$. Sections of $\mathbb{S}(M)$ are usually called spinors. And the compatible covariant derivative on $\mathbb{S}(M)$, denoted by $\nabla^{\mathbb{S}}$, can be locally expressed as

$$\nabla^{\mathbb{S}}\varphi_\alpha = \frac{1}{4} \sum_{i,j=1}^m g(\nabla e_i, e_j) e_i \cdot_g e_j \cdot_g \varphi_\alpha$$

where $\{e_i\}$ is a local orthonormal basis of TM and $\{\varphi_\alpha\}$ is a local spinorial frame in $\mathbb{S}(M)$, the symbol ∇ on the right hand side is the Levi-Civita connection on (M, g) and “ \cdot_g ” stands for the Clifford multiplication. With all these notations, let us introduce the Dirac operator $D_g : C^\infty(M, \mathbb{S}(M)) \rightarrow C^\infty(M, \mathbb{S}(M))$ which is locally formulated as

$$D_g\psi = \sum_{i=1}^m e_i \cdot_g \nabla_{e_i}^{\mathbb{S}}\psi.$$

for a spinor $\psi \in C^\infty(M, \mathbb{S}(M))$.

Under conformal transformations, the Dirac operator behaves analogously to the conformal Laplacian, which arises in the study of the classical Yamabe problem, see for instance [23, 24]. For this reason, a spinorial counterpart of the Yamabe invariant, which is known as the *Bär-Hijazi-Lott invariant* of the Dirac operator, was investigated in [8, 9].

From the variational view point, the Bär-Hijazi-Lott invariant for (M, g, σ) is a positive real number $\lambda_{min}^+(M, g, \sigma)$ given by

$$\lambda_{min}^+(M, g, \sigma) := \inf \left\{ \frac{\left(\int_M |D_g \psi|_g^{\frac{2m}{m+1}} d \text{vol}_g \right)^{\frac{m+1}{m}}}{\int_M (D_g \psi, \psi)_g d \text{vol}_g} \mid \int_M (D_g \psi, \psi)_g d \text{vol}_g > 0 \right\}. \quad (1.1)$$

For $\tilde{g} = f^2 g$ (with $f \in C^\infty(M)$ and $f > 0$) and $\tilde{\psi} = f^{-\frac{m-1}{2}} \psi$ (via the identification of spinor bundles with respect to conformal metrics [23, 24]), we have that $\int_M (D_g \psi, \psi)_g d \text{vol}_g = \int_M (D_{\tilde{g}} \tilde{\psi}, \tilde{\psi})_{\tilde{g}} d \text{vol}_{\tilde{g}}$ and $\int_M |D_g \psi|_g^{\frac{2m}{m+1}} d \text{vol}_g = \int_M |D_{\tilde{g}} \tilde{\psi}|_{\tilde{g}}^{\frac{2m}{m+1}} d \text{vol}_{\tilde{g}}$. Hence, $\lambda_{min}^+(M, g, \sigma)$ is actually a conformal invariant. Furthermore, the Bär-Hijazi-Lott invariant is tightly connected to and behaves much like the classical Yamabe invariant. In fact, it follows from Hijazi's inequality [23] that, for $m \geq 3$, if the Yamabe invariant

$$Y(M, [g]) = \inf_{\tilde{g} \in [g]} \frac{\int_M \text{Scal}_{\tilde{g}} d \text{vol}_{\tilde{g}}}{\text{Vol}(M, \tilde{g})^{\frac{m-2}{m}}}$$

is non-negative, then

$$\lambda_{min}^+(M, g, \sigma) \geq \sqrt{\frac{m}{4(m-1)}} Y(M, [g]). \quad (1.2)$$

And the equality in (1.2) holds for the round spheres.

In a similar way as for the Yamabe invariant, the value of the Bär-Hijazi-Lott invariant for an arbitrary closed spin manifold can not be larger than that for the round sphere (with the same dimension). This is known as the spinorial analogue of Aubin's inequality (see [11]):

$$\lambda_{min}^+(M, g, \sigma) \leq \lambda_{min}^+(S^m, g_{S^m}, \sigma_{S^m}) = \frac{m}{2} \omega_m^{\frac{1}{m}} \quad (1.3)$$

where $(S^m, g_{S^m}, \sigma_{S^m})$ is the m -dimensional sphere equipped with its canonical metric g_{S^m} and its unique spin structure σ_{S^m} , and ω_m stands for the volume of (S^m, g_{S^m}) . In the spirit of this, the next stage would consist in showing that (1.3) is a strict inequality if (M, g) is not conformally equivalent to (S^m, g_{S^m}) . However, the strict inequality in (1.3) is only verified for some special cases (for instance, if M is locally conformally flat, if D_g is invertible and if the so-called *Mass endomorphism* is not identically zero [12], and all rectangular tori [40]), and a general result is still lacking (cf. [10, 20, 22]). The methods that can be used are sometimes similar to the ones of the Yamabe problem, but since we work with Dirac operator and spinors, the reasoning is more involved as the eigenvalues of the Dirac operator tend to both $-\infty$ and $+\infty$ and there is no maximum principle.

By looking at the variational problem (1.1), we can see the corresponding Euler-Lagrange equation is

$$D_g \psi = \lambda_{min}^+(M, g, \sigma) |\psi|_g^{\frac{2}{m-1}} \psi \quad \text{with} \quad \int_M |\psi|_g^{\frac{2m}{m-1}} d \text{vol}_g = 1$$

or equivalently

$$D_g\psi = |\psi|_g^{\frac{2}{m-1}}\psi \quad \text{with} \quad \int_M |\psi|_g^{\frac{2m}{m-1}} d\text{vol}_g = \lambda_{\min}^+(M, g, \sigma)^m \quad (1.4)$$

which is a first order semilinear PDE. On closed spin manifolds, the existence of a solution to Eq. (1.4) was obtained in [9, Theorem 1.1] for $\lambda_{\min}^+(M, g, \sigma) < \lambda_{\min}^+(S^m, g_{S^m}, \sigma_{S^m})$ by using the compactness of the subcritical Sobolev embeddings.

To go a bit further, we can slightly generalize Eq. (1.4) to consider the following one

$$D_g\psi = f(x)|\psi|_g^{\frac{2}{m-1}}\psi \quad \text{on } M, \quad (1.5)$$

where $f : M \rightarrow \mathbb{R}$ is a given function. Geometrically speaking, Eq. (1.5) is of particular interest in dimension $m = 2$ because its solution provides a strong tool for showing the existence of prescribed mean curvature surfaces (here the function f plays the role of the mean curvature). Special cases of such surfaces are constant mean curvature (CMC) surfaces (that is $f \equiv \text{constant}$) which have been studied before by completely different techniques, see for instance [21]. The correspondence between a solution of Eq. (1.5) on a Riemannian surface M and a periodic conformal immersion (possibly with branching points) of the universal covering \widetilde{M} into \mathbb{R}^3 with mean curvature f is known as the spinorial Weierstraß representation. For details in this direction, we refer to [8, 9, 18, 29–31, 36, 42–44] and references therein.

Besides the very special case $f \equiv \text{constant}$, which reduces Eq. (1.5) to Eq. (1.4), there are only a few results about the existence of a solution to Eq. (1.5). For $f \neq \text{constant}$, Raulot [39] obtained an existence criterion for this problem which is similar to the strict inequality in (1.3). Indeed, one of his results shows that if the Dirac operator D_g is invertible, f is positive and satisfies

$$\lambda_{\min, f} \cdot \left(\max_{x \in M} f \right)^{\frac{m-1}{m}} < \lambda_{\min}^+(S^m, g_{S^m}, \sigma_{S^m}), \quad (1.6)$$

where

$$\lambda_{\min, f} := \inf \left\{ \frac{\left(\int_M f^{-\frac{m-1}{m+1}} |D_g\phi|^{\frac{2m}{m+1}} d\text{vol}_g \right)^{\frac{m+1}{m}}}{\int_M (D_g\phi, \phi) d\text{vol}_g} \mid \int_M (D_g\psi, \psi) d\text{vol}_g > 0 \right\},$$

then there exists a solution to Eq. (1.5). However, the validity of condition (1.6) is not very clear. Particularly, the existence results in [39] does not apply for $(M, g) = (S^m, g_{S^m})$ since the strict inequality of (1.6) is not valid on the round spheres in any circumstance. For later developments, we mention that in [27, 28] the author obtained some sufficient conditions about f under which there exists a non-trivial solution to Eq. (1.5) on (S^m, g_{S^m}) , $m \geq 2$. Precisely, up to a simple scaling, [27, 28] studied the case $f(x) = 1 + \varepsilon H(x)$ for some $H \in C^2(S^m)$ and small $\varepsilon > 0$.

Much less is known about the existence of multiple solutions for Eq. (1.5). Examples can only be given on some product manifolds, see [40], or on manifolds that are invariant under some group actions, see [34]. Apart from these cases, we do not know any further multiplicity result. Hence, the purpose of this paper is to discuss new existence and multiplicity results for the problem (1.5) and, when $m = 2$, to give multiplicity results for the immersed spheres in \mathbb{R}^3

with prescribed mean curvature. Note that Eq. (1.5) is invariant under the canonical action of $S^1 = \{z \in \mathbb{C} : |z| = 1\}$ on spinors. Moreover, for the cases $m \equiv 2, 3, 4 \pmod{8}$, the spinor bundle has a quaternionic structure which commutes with Clifford multiplication [19, Section 1.7] or [32, Page 33, Table III]. Thus in these cases, Eq. (1.5) is invariant under the canonical action of the unit quaternions $S^3 = \{q \in \mathbb{H} : |q| = 1\}$. Thus we are interested in multiple existence of S^1 or S^3 -inequivalent solutions.

We are concerned with two concrete problems. First of all, we study the case that the geometric potential in Eq. (1.5) (i.e. the mean curvature function) is a perturbation from constant, namely, we are concerned with the equation

$$D_{g_{S^m}}\psi = (1 + \varepsilon H(x))|\psi|_{g_{S^m}}^{\frac{2}{m-1}}\psi \quad \text{on } S^m, \quad (1.7)$$

for $|\varepsilon| > 0$ small, and we intend to improve the existence results in [27,28]. Secondly, motivated by the (strict) inequality in (1.3), we introduce a new family of metrics g on S^m (close to g_{S^m} but not conformally equivalent) such that the equation

$$D_g\psi = |\psi|_g^{\frac{2}{m-1}}\psi \quad \text{on } S^m, \quad (1.8)$$

has multiple solutions. An important feature of Eq. (1.8) is that (as was indicated in (1.4)) the integral $\int_{S^m} |\psi|_g^{\frac{2m}{m-1}} d\text{vol}_g$ of a solution gives a description of the Bär-Hijazi-Lott invariant. And it would be of particular interest if one can derive a conformal spectral estimate for the Dirac operator D_g so that (1.3) is a strict inequality. As we mentioned before, as far as geometric aspects are concerned, the solutions to both of the equations (1.7) and (1.8) correspond to conformal (branched) immersions of S^2 into \mathbb{R}^3 . However, Eq. (1.8) for $m = 2$ does not reflect the geometric importance since, in view of Riemann mapping theorem, there is only one conformal structure on S^2 .

1.1 Existence results for the PDE problems

Let us start with Eq. (1.7). For a function $H \in C^1(S^m)$, let us denote $\text{Crit}[H]$ the critical set of H . To state our results, we assume the following two standing conditions on H :

(H-1) $H \in C^2(S^m)$ is a Morse function such that $\Delta_{g_{S^m}}H(p) \neq 0$ for $p \in \text{Crit}[H]$.

(H-2) H satisfies that

$$\sum_{p \in \text{Crit}[H], \Delta_{g_{S^m}}H(p) < 0} (-1)^{\mathfrak{M}(H,p)} \neq (-1)^m$$

where $\mathfrak{M}(H, p)$ is the Morse index of H at $p \in \text{Crit}[H]$.

Here we mention that condition (H-2) is the well-known index counting condition which was first introduced in the scalar curvature problem in [13, 17].

In order to describe our general multiplicity result, it is necessary to put some additional requirements on H . Here we are interested in the case that the function H can be decomposed into several components such that each component generates solutions to the Eq. (1.7). For this purpose, for a point $p \in S^m$, let us denote $\pi_p : S^m \setminus \{p\} \rightarrow \mathbb{R}^m$ the standard stereographic projection. Then we introduce the following condition

(H-3) for some $l \in \mathbb{N}$, $l \geq 2$, H has of the form

$$H(x) = \begin{cases} \sum_{i=1}^l H_i(\pi_{p_0}^{-1}(\pi_{p_0}(x) - z_i)) & x \in S^m \setminus \{p_0\} \\ \sum_{i=1}^l H_i(p_0) & x = p_0 \end{cases}$$

where $H_1, \dots, H_l \in C^2(S^m)$ have a common minimum point $p_0 \in S^m$ and $z_1, \dots, z_l \in \mathbb{R}^m$ are fixed.

It is clear that $H_i \circ \pi_{p_0}^{-1}$ defines a C^2 -function on \mathbb{R}^m and $\lim_{|y| \rightarrow \infty} H_i \circ \pi_{p_0}^{-1}(y) = H_i(p_0)$ for each i . The function $H_i(\pi_{p_0}^{-1}(\pi_{p_0}(\cdot) - z_i)) : S^m \setminus \{p_0\} \rightarrow \mathbb{R}$ can be viewed as a translation of H_i on S^m with $H_i(p_0)$ being fixed. Hence the condition (H-3) describes a function that is (approximately) concentrated on the points $\pi_{p_0}^{-1}(z_i) \in S^m \setminus \{p_0\}$, $i = 1, \dots, l$, and is well-defined on S^m . Here, the only reason we keep p_0 being fixed is that we want to present the function H in a simple manner. And, of course, (H-3) can be replaced by other forms. Now we prove the following existence result.

Theorem 1.1. For $m \geq 2$, let \mathcal{H} denote a family of functions in $C^2(S^m)$ as

$$\mathcal{H} := \{H \in C^2(S^m) : H \text{ satisfies the conditions (H-1) and (H-2)}\}.$$

Then

- (1) for a generic function $H \in \mathcal{H}$, there exists $\varepsilon_0 > 0$ such that Eq. (1.7) has at least $2^{\lfloor \frac{m}{2} \rfloor}$ distinct non-trivial S^1 -orbits of solutions for $0 < |\varepsilon| < \varepsilon_0$. Furthermore, when $m \equiv 2, 3, 4 \pmod{8}$, there are at least $2^{\lfloor \frac{m}{2} \rfloor - 1}$ distinct non-trivial S^3 -orbits of solutions for $0 < |\varepsilon| < \varepsilon_0$;
- (2) if $H \in C^2(S^m)$ satisfies condition (H-3) with generic $H_1, \dots, H_l \in \mathcal{H}$ and let $z_1, \dots, z_l \in \mathbb{R}^m$ be located far enough apart, then there exists $\varepsilon_0 > 0$ such that Eq. (1.7) has at least $2^{\lfloor \frac{m}{2} \rfloor} l$ distinct non-trivial S^1 -orbits of solutions for $0 < |\varepsilon| < \varepsilon_0$. Furthermore, when $m \equiv 2, 3, 4 \pmod{8}$, there are at least $2^{\lfloor \frac{m}{2} \rfloor - 1} l$ distinct non-trivial S^3 -orbits of solutions for $0 < |\varepsilon| < \varepsilon_0$;
- (3) if ψ_ε denotes any of the solutions obtained in (1) and (2), then

$$\int_{S^m} (1 + \varepsilon H(x)) |\psi_\varepsilon|_{g_{S^m}}^{\frac{2m}{m-1}} d \text{vol}_{g_{S^m}} = \left(\frac{m}{2}\right)^m \omega_m + O(\varepsilon)$$

as $\varepsilon \rightarrow 0$, moreover, $|\psi_\varepsilon|_{g_{S^m}} > 0$ on S^m provided that $|\varepsilon|$ is small.

Remark 1.2. Here, by ‘‘generic’’ we mean that the existence results holds for H belonging to a residual subset of \mathcal{H} .

Next, we turn to consider the existence results for Eq. (1.8) equipped with a new family of metrics on the spheres. To characterize the new metric g on S^m , we use again the stereographic

projection $\pi_p : S^m \setminus \{p\} \rightarrow \mathbb{R}^m$ (for an arbitrarily fixed $p \in S^m$) to have the following one-to-one correspondence between g on $S^m \setminus \{p\}$ and a metric \tilde{g} on \mathbb{R}^m :

$$\tilde{g} = f^{-2} \cdot (\pi_p^{-1})^* g, \quad f(x) = \frac{2}{1 + |x|^2}, \quad x \in \mathbb{R}^m. \quad (1.9)$$

Clearly, if $\tilde{g} = g_{\mathbb{R}^m}$ is the canonical Euclidean metric, then the metric g on $S^m \setminus \{p\}$ can be extended globally to the standard round metric. In what follows, we assume that \tilde{g} takes the form $\tilde{g} = g_{\mathbb{R}^m} + \varepsilon h$ where h is a smooth symmetric bilinear form on \mathbb{R}^m . In particular, let us consider a specific case

$$\tilde{g}(x) = \text{diag}(\tilde{g}_{11}(x), \dots, \tilde{g}_{mm}(x)) \quad \text{with} \quad \tilde{g}_{ii}(x) = 1 + \varepsilon h_{ii}(x), \quad (1.10)$$

where $h_{ii} : \mathbb{R}^m \rightarrow \mathbb{R}$, $i = 1, \dots, m$, are smooth functions. Let us point out here that, for a general choice of h , the metric g on $S^m \setminus \{p\}$ may be discontinuous at the point p . Hence, in order to extend g globally on S^m , it is natural to require the entries h_{ii} , $i = 1, \dots, m$, and their derivatives behave “nicely” at infinity. To give an idea of our multiplicity results, let us consider h to be compactly supported (that is to say $\cup_i \text{supp } h_{ii}$ is a compact subset of \mathbb{R}^m). And then, we are interested in the situation where \tilde{g} is not conformal to the Euclidean metric.

For the sake of clarity, we construct a simple and effective formulation for perturbations h in (1.10). Let us make this precise.

Definition 1.3. *Given a smooth $m \times m$ diagonal matrix function*

$$h(x) = \text{diag}(h_{11}(x), \dots, h_{mm}(x)) \quad \text{for } x \in \mathbb{R}^m, \quad m \geq 2,$$

and a point $\xi = (\xi_1, \dots, \xi_m) \in \mathbb{R}^m$. For $k \in \{1, \dots, m\}$ and $p \in [1, \infty)$, we say that h is (k, p) -elementary at ξ , if $\xi \notin \text{supp } h_{kk}$ and, for $x = (x_1, \dots, x_m) \in \mathbb{R}^m$ close to ξ and $i \neq k$,

$$h_{ii}(x) = h_{ii}(\xi) + c_i(x_i - \xi_i) + c_k(x_k - \xi_k) + o(|x - \xi|^p)$$

where $c_i \in \mathbb{R}$, $i = 1, \dots, m$, are constants with particularly $c_k \neq 0$. Moreover, if the $o(|x - \xi|^p)$ term vanishes identically in the above local expansion of h_{ii} 's, then we say h is (k, ∞) -elementary at ξ . In this way, we call $p \in [1, \infty) \cup \{\infty\}$ the remainder exponent of h at ξ .

Without breaking the reading, examples and a brief explanation of such (k, p) -elementary matrices are given in the Appendix A.1. It can be seen from Definition 1.3 that “elementary” matrix is a local concept. In the sequel, if it is clear from the context to which dimension we refer, we will simply use the name “elementary matrix” to designate a member h (without mentioning its tag numbers k, p or the location point ξ).

Theorem 1.4. *For $m \geq 4$, let g be the metric defined via (1.9) and (1.10) with*

$$h(x) = h^{(1)}(x) + h^{(2)}(x - x_0), \quad (1.11)$$

where $x_0 \in \mathbb{R}^m$ is fixed, $h^{(1)}$ and $h^{(2)}$ have compact supports and are elementary matrices. Assume additionally that, for $i = 1, 2$,

$$h^{(i)} \text{ has remainder exponent } \begin{cases} p^{(i)} = \infty & m = 4, \\ p^{(i)} > 2 & m = 5, \\ p^{(i)} \geq 2 & m \geq 6. \end{cases}$$

If $|x_0|$ is large enough, then

- (1) Eq. (1.8) has at least two non-trivial solutions ψ_1 and ψ_2 provided that $|\varepsilon| > 0$ is small;
 (2) for each $i = 1, 2$, there holds

$$\int_{S^m} |\psi_i|_{g^{\frac{2m}{m-1}}} d \text{vol}_g = \left(\frac{m}{2}\right)^m \omega_m + C_{i,m} \varepsilon^2 + o(\varepsilon^2),$$

where $C_{i,m} < 0$ is a negative constant depending only on $h^{(i)}$ and the dimension m , $i = 1, 2$.

Remark 1.5. (1) Although we proved the case for h splits into two elementary matrices, Theorem 1.4 extends to the situation where h splits into an arbitrary finite number of elementary matrices. Here the results do not reflect the canonical action of S^1 (or S^3 when $m \equiv 2, 3, 4 \pmod{8}$) as mentioned in Theorem 1.1 since each one of the solutions is generated by the components of h in (1.11) individually.

- (2) For $p \geq 2$, let $[p] \in \mathbb{Z}$ denotes the largest integer that does not exceed p . Definition 1.3 indicates that, when h is (k, p) -elementary at some point ξ , all the j -th partial derivative of each h_{ii} vanish at this point for $j = 2, \dots, [p]$. One possible extension of this condition is to consider the case where the local expansion of h_{ii} has quadratic terms. The proofs can be carried out by using similar arguments.

1.2 Geometric applications

As was mentioned before, the solutions of Eq. (1.7) and Eq. (1.8) has two major applications. The first application is that, when $m = 2$, we obtain existence results for immersed spheres in \mathbb{R}^3 with prescribed mean curvature. And the second application gives us an upper bound estimate for the Bär-Hijazi-Lott invariant.

Remark 1.6. (1) For $m = 2$, Theorem 1.1 implies multiple existence of conformal immersions $S^2 \rightarrow \mathbb{R}^3$ with mean curvature equals to $1 + \varepsilon H$. Precisely, if ψ_ε denotes any one of the solutions obtained in Theorem 1.1, since $|\psi_\varepsilon|_{g_{S^2}} > 0$ we can introduce a conformal metric $g_\varepsilon = |\psi_\varepsilon|_{g_{S^2}}^4 g_{S^2}$ on S^2 . Then, due to the conformal covariance of the Dirac operator, we see that there is a spinor field φ_ε on (S^2, g_ε) such that

$$D_{g_\varepsilon} \varphi_\varepsilon = (1 + \varepsilon H(x)) \varphi_\varepsilon \quad \text{and} \quad |\varphi_\varepsilon|_{g_\varepsilon} \equiv 1.$$

Hence, by the spinorial Weierstraß representation, there is an isometric immersion $\Pi : (S^2, g_\varepsilon) \rightarrow (\mathbb{R}^3, g_{\mathbb{R}^3})$ with mean curvature equals to $1 + \varepsilon H$. Furthermore, since the pull-back of the Euclidean volume form under this immersion is $\Pi^*(d \text{vol}_{g_{\mathbb{R}^3}}) = |\psi_\varepsilon|_{g_{S^2}}^4 d \text{vol}_{g_{S^2}}$, the associated Willmore energy $W(\Pi)$ for this immersion satisfies

$$W(\Pi) = \int_{S^2} (1 + \varepsilon H(x))^2 |\psi_\varepsilon|_{g_{S^2}}^4 d \text{vol}_{g_{S^2}} \leq 4\pi + O(\varepsilon) < 8\pi$$

as $\varepsilon \rightarrow 0$. Due to Li-Yau's inequality [33, Theorem 6], the immersion Π covers points in \mathbb{R}^3 at most once. Hence Π is an embedding.

- (2) The existence result in Theorem 1.4 directly implies a conformal spectral estimate for the Dirac operator D_g on (S^m, g) . In fact, by the definition of Bär-Hijazi-Lott invariant (1.1), we have

$$\lambda_{min}^+(S^m, g, \sigma_{S^m}) \leq \frac{\left(\int_{S^m} |D_g \psi_i|_g^{\frac{2m}{m+1}} d \text{vol}_g \right)^{\frac{m+1}{m}}}{\int_{S^m} (D_g \psi_i, \psi_i)_g d \text{vol}_g} = \left(\int_{S^m} |\psi_i|_g^{\frac{2m}{m-1}} d \text{vol}_g \right)^{\frac{1}{m}} < \frac{m}{2} \omega_m^{\frac{1}{m}}$$

for each $\psi_i, i = 1, 2$, obtained in Theorem 1.4. Hence, the strict inequality in (1.3) holds true for the metric g . It would be very interesting whether analogous results hold for dimension 3.

Now, by summarizing Remark 1.6, we have the following two geometric applications of Theorem 1.1 and 1.4.

Theorem 1.7. *Let H satisfy the conditions of Theorem 1.1 (2), there are at least l -distinct conformal isometric embeddings of (S^2, g_{S^2}) into $(\mathbb{R}^3, g_{\mathbb{R}^3})$ whose images have mean curvature $1 + \varepsilon H$, for $|\varepsilon| > 0$ small.*

Theorem 1.8. *For $m \geq 4$, let g be the metric as in Theorem 1.4. Then the strict inequality*

$$\lambda_{min}^+(S^m, g, \sigma_{S^m}) < \lambda_{min}^+(S^m, g_{S^m}, \sigma_{S^m}) \quad (1.12)$$

holds provided that $|\varepsilon| > 0$ is small.

Remark 1.9. (1) Theorem 1.7 can be viewed as part of a program to understand to what extent one can use mean curvature data to determine the shape of a surface in \mathbb{R}^3 . Usually, a generic immersion is uniquely determined up to a rigid motion by its first fundamental form and its mean curvature function, but there are some exceptions, for instance most constant mean curvature immersions. A classical result by Bonnet states that if there exists a diffeomorphism $\Psi : \Sigma_1 \rightarrow \Sigma_2$ between two closed immersed surfaces Σ_1, Σ_2 of genus zero in \mathbb{R}^3 such that Ψ preserves both the metric and the mean curvature function of the surfaces, then Σ_1 and Σ_2 are congruent in \mathbb{R}^3 (i.e., they differ by a rigid motion). In view of this, one may address the following natural questions:

Question 1. *Given a function $f : S^2 \rightarrow \mathbb{R}$, what conditions on f guarantee the existence of an isometric immersion $\Pi : S^2 \rightarrow \mathbb{R}^3$ realizing f as its mean curvature?*

Question 2. *Assuming that $f : S^2 \rightarrow \mathbb{R}$ is given such that it is the mean curvature function of an isometric immersion $\Pi : S^2 \rightarrow \mathbb{R}^3$. Is it possible that there exists another isometric immersion $\hat{\Pi} : S^2 \rightarrow \mathbb{R}^3$, not congruent with Π , but also realizing f as its mean curvature? If so, what can be said about the pull-back metrics $\Pi^* g_{\mathbb{R}^3}$ and $\hat{\Pi}^* g_{\mathbb{R}^3}$?*

Note that in Question 2 we are not assuming that the immersed spheres $\Pi(S^2)$ and $\hat{\Pi}(S^2)$ are isometric, which is a critical hypothesis of Bonnet's result. Theorem 1.7 provides answers to both Questions 1 and 2: we find sufficient conditions on f such that it can be realized as the mean curvature of distinct isometric embeddings, moreover, the pull-back metrics are conformally related.

(2) Theorem 1.8 provides an explicit family of examples such that the strict inequality in (1.3) holds. We are interested in obtaining this strict inequality for following reasons:

- The strict inequality in (1.3) implies that the invariant $\lambda_{min}^+(M, g, \sigma)$ is attained by a generalized metric, i.e., a metric of the form f^2g where f may have some zeros.
- Using Hijazi’s inequality [23], one obtains a solution of the standard Yamabe problem which consists of finding a metric with constant scalar curvature in the conformal class of g in the case of $m \geq 3$.

With the aim of obtaining this inequality, Ammann et al. considered in [12] the case of a locally conformally flat spin manifold (M, g, σ) with $\ker D_g = \{0\}$ and introduced for a fixed point $p \in M$, a further datum, the *mass endomorphism*. The main result of [12] is that the strict inequality in (1.3) holds, if g is conformally flat around p and the mass endomorphism at p has a positive eigenvalue. Meanwhile, it is also proven in [12] that the mass endomorphism always vanishes in dimension 2 and on flat tori \mathbb{T}^m . To describe the dependence of the mass endomorphism on the Riemannian metric, let $\mathcal{M}_{p,flat}(M)$ be the set of all Riemannian metrics g on M , which are flat on a neighborhood of $p \in M$ and satisfy $\ker D_g = \{0\}$. It is proven in [10], see also [22], that for dimension $m \geq 3$, the subset of all Riemannian metrics with non-vanishing mass endomorphism at p is dense in $\mathcal{M}_{p,flat}(M)$ with respect to the C^∞ -topology. However, for an arbitrarily given metric g , examining the non-vanishingness of its associated mass endomorphism is still a difficult issue.

It can be seen from Theorem 1.8 that, even if the metric g is flat somewhere on M , the appearance of the non-locally-conformally-flat part is the key reason we obtain the strict inequality. It is well known that, for $m \geq 4$, the conformal flatness of (M, g) is characterized by the nullity of the Weyl tensor. However, what is the exact role of the Weyl tensor in the general inequality $\lambda_{min}^+(M, g, \sigma) < \lambda_{min}^+(S^m, g_{S^m}, \sigma_{S^m})$ when (M, g) is non-locally conformally flat remains unexplored. To the best of our knowledge, Theorem 1.8 is the first result in the non-locally-conformally-flat setting. We consider it hence as remarkable that Theorem 1.8 might give us a good chance to uncover the very nature of the Bär-Hijazi-Lott invariant in general cases.

1.3 Overview of the method

The proof of Theorem 1.1 and 1.4 relies on a suitable use of abstract perturbation methods in critical point theory. This idea is very common to many nonlinear problems, such as the classical Yamabe problem and related topics (see for instance [5–7, 16, 35]). It was carried out in detail in the papers [27, 28] that such an idea can be employed to spinorial Yamabe equation (1.7). Roughly speaking, the perturbation methods consist of two main steps: first to use Lyapunov-Schmidt reduction to reduce the original problem to a certain finite dimensional variational problem and then to show that “stable” critical points of the reduced problem produce solutions of the original one.

As was shown in [27, 28], the reduced problem of spinorial Yamabe type equations is very similar to the corresponding problem for elliptic equations at a first glance, see for example [5, 7, 35]. However, unlike the elliptic cases, here the classical abstract framework can not be

used in a straightforward manner. This is because the finite dimensional problem associated to the spinorial Yamabe problem is degenerate, that is, any critical point of the main term of the reduced functional is not isolated, and the collection of these critical points appear as critical manifolds of positive dimension. Thus, it is not clear whether critical points of the reduced functional create true solutions of the original problem. To overcome this difficulty, the author developed, in [27], a variant of Morse-Bott theory and, in [28], a Conley index theory for such variational problem.

In this paper, our approach is different from the ones used in the aforementioned works. We have a two-step reduction procedure before finding a critical point. We start with the Lyapunov-Schmidt reduction to obtain a reduced finite dimensional variational problem, and the new perspective is the observation that the reduced functional is defined on a space endowed with a fiber bundle structure, where the fiber is non-compact and the base space is a close manifold. The key point in our argument is that the degeneracy of the reduced functional essentially occurs on the base manifold. Hence, as a second step, we get a chance to restrict the reduced functional on each fiber rather than on the total space and finally (via implicit function theorem, minimizing/maximizing methods) to show the critical points of the reduced problem give rise to solutions of the original one. An advantage of our approach is that it produces proofs that are rather simpler than the ones of [27, 28]. And this might be meaningful to provide new insight on nonperturbative problems.

1.4 Organization of the paper

This paper is organized as follows: In the next section, we quickly review the transformation of the equations (1.7) and (1.8) into equivalent equations on Euclidean spaces, and then we set up our abstract perturbation framework with degenerate conditions. Section 3 is concerned with our first problem, i.e., Eq. (1.7) where the perturbation is appearing in the geometric potential. And Section 4 deals with the second problem, that is, Eq. (1.8) where the perturbation is involved in the background metric. Both in Section 3 and 4, we first collect properties of the variational problem and then apply the abstract results in Section 2 to obtain the existence results. Finally, in the Appendix, we collect some straight calculations which are needed in the arguments.

2 The precise framework

2.1 Reduction to problems on Euclidean spaces

We first mention that the Eq. (1.7) on S^m is equivalent to an equation on \mathbb{R}^m by conformal equivalence. Precisely, let $H \in C^2(S^m)$ satisfies the hypotheses of Theorem 1.1, we assume H takes its minimum at $p_0 \in S^m$. Denote by $\pi_{p_0} : S^m \setminus \{p_0\} \rightarrow \mathbb{R}^m$ the stereographic projection from p_0 , we have $(\pi_{p_0}^{-1})^* g_{S^m} = f^2 g_{\mathbb{R}^m}$ where g_{S^m} is the round metric on S^m with constant sectional curvature 1 and $f(x) = \frac{2}{1+r^2}$ ($r = |x|$ and $x \in \mathbb{R}^m \cong T_{p_0} S^m$ is the stereographic coordinate with respect to π_{p_0}).

On the other hand, the general equation $D_g \psi = H(x) |\psi|_g^{\frac{2}{m-1}} \psi$ on a manifold (M, g) is invariant under conformal change of the metric. In fact, let $\bar{g} = e^{2u} g$ for some function u on M , there is an isomorphism of vector bundles $F : \mathbb{S}(M, g) \rightarrow \mathbb{S}(M, \bar{g})$ (here $\mathbb{S}(M, g)$ and $\mathbb{S}(M, \bar{g})$

are spinor bundles on M with respect to the metrics g and \bar{g} , respectively) which is a fiberwise isometry such that

$$D_{\bar{g}}(F(\varphi)) = F(e^{-\frac{m+1}{2}u} D_g(e^{\frac{m-1}{2}u} \varphi))$$

for $\varphi \in C^\infty((M, g), \mathbb{S}(M, g))$ (for more detailed definitions and facts about Clifford algebras, spin structures on manifolds and Dirac operators, please consult [19, 32]). Thus, when ψ is a solution to the equation $D_g \psi = H(x) |\psi|_g^{\frac{2}{m-1}} \psi$ on (M, g) , then $\varphi := F(e^{-\frac{m-1}{2}u} \psi)$ also satisfies the same equation on (M, \bar{g}) : $D_{\bar{g}} \varphi = H(x) |\varphi|_{\bar{g}}^{\frac{2}{m-1}} \varphi$ on (M, \bar{g}) .

Applying the above observation to Eq. (1.7) with $M = S^m$, if $\psi \in C^1(S^m, \mathbb{S}(S^m))$ is a solution to the Eq. (1.7), then $\varphi = f^{\frac{m-1}{2}} F(\psi \circ \pi_{p_0}^{-1})$ satisfies the equation

$$D_{g_{\mathbb{R}^m}} \varphi = (1 + \varepsilon a(x)) |\varphi|_{g_{\mathbb{R}^m}}^{\frac{2}{m-1}} \varphi \quad \text{on } \mathbb{R}^m \quad (2.1)$$

where $a(x) = H(\pi_{p_0}^{-1}(x))$.

Conversely, by the regularity theorem for weak solutions for Eq. (1.5) (see [26, Appendix]), for any solution $\varphi \in L^{\frac{2m}{m-1}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ to Eq. (2.1) with Hölder continuous function a , there corresponds to a C^1 -solution $\psi = \varphi \circ \pi_{p_0}$ to Eq. (1.7) on S^m with $H(x) = a(\pi_{p_0}(x))$. Therefore, the study of Eq. (1.7) is equivalent to the study of Eq. (2.1).

For our second problem, i.e., Eq. (1.8), we have a similar equivalent problem on the Euclidean space. Recall that, by (1.9), we have the conformal equivalence $(S^m \setminus \{p\}, g) \cong (\mathbb{R}^m, \tilde{g})$. Thus, by repeating the above arguments, we find that Eq. (1.8) is equivalent to the equation

$$D_{\tilde{g}} \varphi = |\varphi|_{\tilde{g}}^{\frac{2}{m-1}} \varphi \quad \text{on } \mathbb{R}^m \quad (2.2)$$

where $\tilde{g} = g_{\mathbb{R}^m} + \varepsilon h$ as was assumed in (1.9) and (1.10).

For later purpose, we point out that both Eq. (2.1) and (2.2) can be seen as perturbations from the unperturbed equation

$$D_{g_{\mathbb{R}^m}} \psi = |\psi|_{g_{\mathbb{R}^m}}^{\frac{2}{m-1}} \psi \quad \text{on } \mathbb{R}^m. \quad (2.3)$$

Hence, in the sequel, our perturbation framework will be build upon the study of Eq. (2.3) and its associated energy functional

$$\mathcal{J}_0(\psi) = \frac{1}{2} \int_{\mathbb{R}^m} (\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} - \frac{m-1}{2m} \int_{\mathbb{R}^m} |\psi|_{g_{\mathbb{R}^m}}^{\frac{2m}{m-1}} d \text{vol}_{g_{\mathbb{R}^m}} \quad (2.4)$$

where $(\cdot, \cdot)_{g_{\mathbb{R}^m}}$ and $|\cdot|_{g_{\mathbb{R}^m}}$ are the canonical hermitian product and its induced metric on the spinor bundle $\mathbb{S}(\mathbb{R}^m)$.

2.2 Configuration spaces

To treat the Eq. (1.5) and (2.3) from a variational point of view, it is necessary to set up a functional framework. Suitable function spaces are $H^{\frac{1}{2}}(M, \mathbb{S}(M))$ and $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ of spinor fields which are introduced in [26, 27]. For completeness, we give the definitions as follows.

Recall that the Dirac operator D_g on a compact spin manifold (M, g) is self-adjoint on $L^2(M, \mathbb{S}(M))$ and has compact resolvents (see [19, 32]). Particularly, there exists a complete

orthonormal basis ψ_1, ψ_2, \dots of the Hilbert space $L^2(M, \mathbb{S}(M))$ consisting of the eigenspinors of D_g : $D_g\psi_k = \lambda_k\psi_k$. Moreover, $|\lambda_k| \rightarrow \infty$ as $k \rightarrow \infty$.

For $s \geq 0$, we define the operator $|D_g|^s : L^2(M, \mathbb{S}(M)) \rightarrow L^2(M, \mathbb{S}(M))$ by

$$|D_g|^s\psi = \sum_{k=1}^{\infty} |\lambda_k|^s \alpha_k \psi_k,$$

for $\psi = \sum_{k=1}^{\infty} \alpha_k \psi_k \in L^2(M, \mathbb{S}(M))$. Denoted by $H^s(M, \mathbb{S}(M))$ the domain of $|D_g|^s$, then it is clear that $\psi = \sum_{k=1}^{\infty} \alpha_k \psi_k \in H^s(M, \mathbb{S}(M))$ if and only if $\sum_{k=1}^{\infty} |\lambda_k|^{2s} |\alpha_k|^2 < \infty$. And hence $H^s(M, \mathbb{S}(M))$ coincides with the usual Sobolev space of order s , that is $W^{s,2}(M, \mathbb{S}(M))$ (cf. [1, 8]). Furthermore, we can equip $H^s(M, \mathbb{S}(M))$ with the inner product

$$\langle \psi, \varphi \rangle_{s,2} := \operatorname{Re}(|D_g|^s\psi, |D_g|^s\varphi)_2 + \operatorname{Re}(\psi, \varphi)_2$$

where $(\cdot, \cdot)_2$ is the L^2 -inner product on spinors and the induced norm $\|\cdot\|_{s,2}$. In this paper, we are mainly concerned with the space $H^{\frac{1}{2}}(M, \mathbb{S}(M))$ for $M = S^m$. Notice that the spectrum of $D_{g_{S^m}}$ on S^m is bounded away from 0 and one checks easily that $\|\psi\|_{1/2} = \left\| |D_{g_{S^m}}|^{\frac{1}{2}} \psi \right\|_2$ defines an equivalent norm on $H^{\frac{1}{2}}(S^m, \mathbb{S}(S^m))$.

For ease of notation, throughout this paper, we shall denote $L^q := L^q(M, \mathbb{S}(M))$ with the norm $|\psi|_q^q = \int_M |\psi|^q d \operatorname{vol}_g$ for $q \geq 1$ and denote $2^* = \frac{2m}{m-1}$ the critical Sobolev exponent of the embedding $H^{1/2}(M, \mathbb{S}(M)) \hookrightarrow L^q(M, \mathbb{S}(M))$ for $1 \leq q \leq 2^*$.

On \mathbb{R}^m , a similar function space will also be useful in our argument. Let us denote by $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ the set of spinor fields ψ on \mathbb{R}^m such that $\left\| |D_{g_{\mathbb{R}^m}}|^{\frac{1}{2}} \psi \right\|_2^2 < \infty$ and $|\psi|_{2^*} < \infty$ with norm $\|\psi\|_{1/2,2} := \left\| |D_{g_{\mathbb{R}^m}}|^{\frac{1}{2}} \psi \right\|_2 + |\psi|_{2^*}$. Here, $|D_{g_{\mathbb{R}^m}}|^{\frac{1}{2}}$ is defined via the Fourier transformation: $\mathcal{F}(|D_{g_{\mathbb{R}^m}}|^{\frac{1}{2}}\psi)(\xi) = |\xi|^{1/2} \mathcal{F}(\psi)(\xi)$ and $\left\| |D_{g_{\mathbb{R}^m}}|^{\frac{1}{2}} \psi \right\|_2 := \left\| |\cdot|^{1/2} \mathcal{F}(\psi) \right\|_2$. Notice that $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ is isomorphic to $H^{\frac{1}{2}}(S^m, \mathbb{S}(S^m))$ via the stereographic projection. The dual space of $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ will be denoted by $\mathcal{D}^{-\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ with norm $\|\cdot\|_{-1/2,2}$.

2.3 The perturbation method with degenerate conditions

It is clear that, on $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$, the functional \mathcal{J}_0 in (2.4) is well-defined and is of C^2 class. We call \mathcal{J}_0 the unperturbed functional. Its critical points are solutions to the unperturbed spinorial Yamabe equation (2.3). Recall that Eq. (2.1) and (2.2) are perturbations from (2.3) and it is very natural to expect that solutions of Eq. (2.1) and (2.2) can be obtained as perturbations of critical points of \mathcal{J}_0 . In general, a well known Lyapunov-Schmidt reduction technique provides a powerful tool, see for instance [37, Chapter 10] and [15, II, 6] where the variational problem satisfies the Palais-Smale condition and [3, 4, 7] for the case that the Palais-Smale condition fails. For completeness, following [7], we sketch the idea as follows.

Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a Hilbert space with the associated norm $\|\cdot\| := \langle \cdot, \cdot \rangle^{1/2}$. Suppose that $L_0 \in C^2(\mathcal{H}, \mathbb{R})$ and $\Gamma \in C^2(\mathcal{H}, \mathbb{R})$ are given. For $\varepsilon > 0$ small, we consider the perturbed functional

$$L_\varepsilon(z) = L_0(z) + \varepsilon\Gamma(z) + o(\varepsilon). \quad (2.5)$$

Assume that L_0 has a non-degenerate critical manifold $\mathcal{M} \subset \mathcal{H}$, that is,

(A1) \mathcal{M} is a d -dimensional C^2 -submanifold of \mathcal{H} such that $\nabla L_0(z) = 0$ for all $z \in \mathcal{M}$,

(A2) \mathcal{M} is non-degenerate in the sense that for all $z \in \mathcal{M}$, we have $T_z\mathcal{M} = \ker \nabla^2 L_0(z)$,

(A3) $\nabla^2 L_0(z) : \mathcal{H} \rightarrow \mathcal{H}$ is a Fredholm operator with index zero for all $z \in \mathcal{M}$.

Once the conditions (A1)-(A3) are satisfied, the problem of finding critical points of L_ε on \mathcal{H} can be reduced to the same problem on \mathcal{M} for a reduced functional L_ε^{red} (defined below). And critical points of L_ε will be obtained as small perturbations of critical points of L_0 .

Set $\mathcal{W}_z := T_z\mathcal{M}^\perp$, where the orthogonal complement is taking with respect to $\langle \cdot, \cdot \rangle$ in \mathcal{H} . We look for critical points of L_ε in the form $u = z + w$ where $z \in \mathcal{M}$ and $w \in \mathcal{W}_z$. Let $P_z : \mathcal{H} \rightarrow \mathcal{W}_z$ be the orthogonal projection onto \mathcal{W}_z , the Euler-Lagrange equation $\nabla L_\varepsilon(z + w) = 0$ is equivalent to

$$\begin{cases} P_z \nabla L_\varepsilon(z + w) = 0 & (\text{auxiliary equation}) \\ (I - P_z) \nabla L_\varepsilon(z + w) = 0 & (\text{bifurcation equation}). \end{cases} \quad (2.6)$$

Then, under the conditions (A2) and (A3), the auxiliary equation in (2.6) can be solved firstly for w by applying the implicit function theorem: for arbitrary $z \in \mathcal{M}$ there is a unique solution $w = w_\varepsilon(z) \in \mathcal{W}_z$ for small values of ε . Furthermore, on any compact subset $\mathcal{M}_c \subset \mathcal{M}$, one can have a uniform estimate:

$$\mathcal{M}_c \ni z \mapsto w_\varepsilon(z) \in \mathcal{W}_z \text{ is } C^1 \text{ and } \|w_\varepsilon(z)\|, \|w'_\varepsilon(z)\| = O(\varepsilon) \text{ uniformly for } z \in \mathcal{M}_c. \quad (2.7)$$

For the proofs of (2.7) and more detailed explanation about this construction, see [7, Chapter 2].

The next step is to consider the bifurcation equation in (2.6). For this end, we introduce the reduced functional $L_\varepsilon^{red} : \mathcal{M} \rightarrow \mathbb{R}$ by

$$L_\varepsilon^{red}(z) = L_\varepsilon(z + w_\varepsilon(z))$$

Then we have the following theorem

Theorem 2.1 (Theorem 2.12 in [7]). *Suppose (A1)-(A3) are satisfied and assume for a compact subset $\mathcal{M}_c \subset \mathcal{M}$ and $\varepsilon > 0$ small, L_ε^{red} has a critical point $z_\varepsilon \in \mathcal{M}_c$. Then $u_\varepsilon = z_\varepsilon + w_\varepsilon(z_\varepsilon)$ is a critical point of L_ε on \mathcal{H} .*

Thanks to the uniform estimate (2.7), the reduced functional L_ε^{red} is well approximated in the sense that

$$L_\varepsilon^{red}(z) = L_0(z) + \varepsilon \Gamma(z) + o(\varepsilon), \quad \nabla L_\varepsilon^{red}(z) = \varepsilon \nabla \Gamma(z) + o(\varepsilon). \quad (2.8)$$

and $L_0(z)$ is constant on any connected component of \mathcal{M} . Thus if $z \in \mathcal{M}$ is a non-degenerate critical point of Γ in some certain sense (for example, the local degree of $\nabla \Gamma$ at z is non-zero), then z generates a critical point of L_ε on \mathcal{H} (see [7, 15, 37] for details).

For our problem, we intend to apply the above abstract framework to the functionals defined by $L_0 = \mathcal{J}_0$ and some certain perturbation terms on $\mathcal{H} = \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ (see (3.1) and Lemma 4.3 later). As was already shown in [27, Section 5, 6] that \mathcal{J}_0 satisfies (A1)-(A3) for a critical manifold \mathcal{M} defined as

$$\mathcal{M} := \{ \psi_{\lambda, \xi, \gamma} : \lambda > 0, \xi \in \mathbb{R}^m, \gamma \in \mathbb{S}_m, |\gamma| = 1 \}, \quad (2.9)$$

where

$$\psi_{\lambda,\xi,\gamma}(x) = \frac{m^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}}} (\lambda - (x - \xi)) \cdot_{g_{\mathbb{R}^m}} \gamma \quad (2.10)$$

for $\lambda > 0, \xi \in \mathbb{R}^m, \gamma \in \mathbb{S}_m$ with $|\gamma| = 1$ (\mathbb{S}_m is the spinor module, see [19,32]) and $\cdot_{g_{\mathbb{R}^m}}$ denotes the Clifford multiplication with respect to the Euclidean metric. Note that \mathcal{M} is diffeomorphic to $(0, \infty) \times \mathbb{R}^m \times S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$ via the canonical map $(\lambda, \xi, \gamma) \mapsto \psi_{\lambda,\xi,\gamma}$, where $S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$ stands for the $(2^{\lfloor \frac{m}{2} \rfloor + 1} - 1)$ -dimensional unit sphere in \mathbb{S}_m . And hence \mathcal{M} is a non-compact manifold and the dimension of \mathcal{M} is $m + 2^{\lfloor \frac{m}{2} \rfloor + 1}$.

By virtue of Theorem 2.1, we intend to reduce the problem to a finite dimensional variational problem on \mathcal{M} . But in our case, as we will see later, the reduced functional L_ε^{red} is very degenerate on \mathcal{M} : the perturbation term Γ in (2.8) is degenerate on \mathcal{M} (in the sense that the map $\nabla \Gamma(z)|_{T_z \mathcal{M}} : T_z \mathcal{M} \rightarrow \mathbb{R}$ has a non-trivial kernel for any $z \in \mathcal{M}$). Hence, standard methods as in [6, 7, 15, 37] do not apply.

In order to make the perturbation framework applicable, besides the assumptions (A1)-(A3), we assume the following additional conditions for the critical manifold \mathcal{M} :

(A4) \mathcal{M} admits a (globally) trivializable fiber bundle structure over a compact base space \mathcal{N} with projection $\vartheta : \mathcal{M} \rightarrow \mathcal{N}$ and fiber \mathcal{G} . Precisely, there is a fiber preserving diffeomorphism $\iota : \mathcal{G} \times \mathcal{N} \rightarrow \mathcal{M}$ such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{G} \times \mathcal{N} & \xrightarrow{\iota} & \mathcal{M} \\ \text{Proj} \downarrow & & \downarrow \vartheta \\ \mathcal{N} & \xrightarrow{\text{Id}} & \mathcal{N} \end{array}$$

(A5) $T_\gamma \mathcal{N} \subset \ker \nabla(\Gamma \circ \iota)(g, \gamma)$ for any $(g, \gamma) \in \mathcal{G} \times \mathcal{N}$, where we have identified $T_\gamma \mathcal{N}$ as a subspace of the total tangent space $T_{(g,\gamma)}(\mathcal{G} \times \mathcal{N})$.

Remark 2.2. In our application $\mathcal{N} = S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, $\mathcal{G} = (0, +\infty) \times \mathbb{R}^m$ and $\iota(g, \gamma) := \psi_{\lambda,\xi,\gamma}$ for $g = (\lambda, \xi) \in \mathcal{G}$ and $\gamma \in \mathcal{N}$, hence we have a very natural bundle structure on \mathcal{M} . Particularly, we note that there is a continuous action $\mathcal{G} \times \mathcal{M} \rightarrow \mathcal{M}$ such that \mathcal{G} preserves the fibers of \mathcal{M} (i.e. if $(\mu, y) \in \mathcal{G}$ and $\psi_{\lambda,\xi,\gamma} \in \mathcal{M}_\gamma$ then $\psi_{\lambda,\xi,\gamma} * (\mu, y) = \psi_{\lambda\mu,\xi+y,\gamma} \in \mathcal{M}_\gamma$). Hence the critical manifold in (2.9) is essentially a principal \mathcal{G} -bundle. And since it admits a global section, we easily see that \mathcal{M} is trivializable. This is the reason we introduce (A4). Since \mathcal{M} is parameterized via ι , condition (A4) makes the variational problem even clearer: it is equivalent to consider the functional $L_\varepsilon^{red} \circ \iota : \mathcal{G} \times \mathcal{N} \rightarrow \mathbb{R}$. Comparing with the standard theory in [7, 15, 37], the distinct new feature (A5) describes a certain degenerate situation and, particularly, it implies that $\Gamma \circ \iota(g, \gamma)$ depends only on the variables in the fiber space \mathcal{G} . Thus we set $\tilde{\Gamma}(g) = \Gamma \circ \iota(g, \gamma)$. For later use, we distinguish (A5) into the following two cases:

$$\begin{cases} \ker \nabla(\Gamma \circ \iota)(g, \gamma) \neq T_{(g,\gamma)}(\mathcal{G} \times \mathcal{N}) & \text{for some } (g, \gamma) \in \mathcal{G} \times \mathcal{N}, \\ \ker \nabla(\Gamma \circ \iota)(g, \gamma) \equiv T_{(g,\gamma)}(\mathcal{G} \times \mathcal{N}) & \text{for all } (g, \gamma) \in \mathcal{G} \times \mathcal{N}, \end{cases}$$

and we shall establish some abstract results which are useful in the spinorial Yamabe problem.

Case 1: $\ker \nabla(\Gamma \circ \iota)(g, \gamma) \neq T_{(g, \gamma)}(\mathcal{G} \times \mathcal{N})$ for some $(g, \gamma) \in \mathcal{G} \times \mathcal{N}$

Clearly, in this case, $\Gamma \circ \iota(g, \gamma) \neq \text{constant}$ on $\mathcal{G} \times \mathcal{N}$ (equivalently, $\Gamma(z) \neq \text{constant}$ on \mathcal{M}).

Theorem 2.3. *Let $L_0, \Gamma \in C^2(\mathcal{H}, \mathbb{R})$ as in (2.5) and suppose that (A1)-(A5) are satisfied. If there is a $\bar{g} \in \mathcal{G}$ such that \bar{g} is a strict local maximum or minimum of $\tilde{\Gamma}$. Then, for $|\varepsilon|$ small, the functional L_ε has a critical point on \mathcal{M} .*

Proof. It is sufficient to consider $\bar{g} \in \mathcal{G}$ is a strict local minimum and $\varepsilon > 0$, the other cases can be done similarly. Let $\delta > 0$ and $U \subset \mathcal{G}$ be a bounded open neighborhood of \bar{g} such that

$$\tilde{\Gamma}(g) \geq \tilde{\Gamma}(\bar{g}) + \delta, \quad \forall g \in \partial U.$$

By the first expansion in (2.8), we can see

$$L_\varepsilon^{\text{red}} \circ \iota(g, \gamma) - L_\varepsilon^{\text{red}} \circ \iota(\bar{g}, \gamma) = \varepsilon(\tilde{\Gamma}(g) - \tilde{\Gamma}(\bar{g})) + o(\varepsilon) > 0 \quad \forall g \in \partial U, \forall \gamma \in \mathcal{N}$$

for $\varepsilon > 0$ small. Since $\bar{U} \times \mathcal{N}$ is compact, by minimizing the functional $L_\varepsilon^{\text{red}} \circ \iota$ in $\bar{U} \times \mathcal{N}$ and by the above inequality, the minimum is attained in the interior and we obtain a critical point of $L_\varepsilon^{\text{red}} \circ \iota$ and equivalently a critical point of $L_\varepsilon^{\text{red}}$ in $\iota(U \times \mathcal{N}) \subset \mathcal{M}$. Therefore, by Theorem 2.1, it generates a critical point of L_ε . \square

In the hypotheses of Theorem 2.3, the strict local maximum or minimum of $\tilde{\Gamma}$ could possibly be degenerate. If we require certain non-degeneracy on the critical points of $\tilde{\Gamma}$ we can have even more solutions as the following result indicates.

Theorem 2.4. *Let $L_0, \Gamma \in C^2(\mathcal{H}, \mathbb{R})$ as in (2.5) and suppose that (A1)-(A5) are satisfied. If $\tilde{\Gamma}$ is a Morse function on \mathcal{G} and there is an open bounded subset $\Omega \subset \mathcal{G}$ such that the topological degree $\deg(\nabla \tilde{\Gamma}, \Omega, 0) \neq 0$. Then, for $|\varepsilon|$ small, the functional $L_\varepsilon^{\text{red}}$ has at least $\text{Cat}(\mathcal{N})$ critical points on \mathcal{M} .*

Here $\text{Cat}(\mathcal{N})$ denotes the Lusternik-Schnierelman category of \mathcal{N} , namely the smallest integer k such that $\mathcal{N} \subset \cup_{i=1}^k A_k$, where the sets A_k are closed and contractible in \mathcal{N} .

Proof. Fix $\gamma \in \mathcal{N}$ arbitrarily and let us consider the functional $L_\varepsilon^{\text{red}}|_{\partial^{-1}(\gamma)}$. By the assumption $\deg(\nabla \tilde{\Gamma}, \Omega, 0) \neq 0$, there exists $\bar{g} \in \Omega$ such that $\nabla \tilde{\Gamma}(\bar{g}) = 0$ and the local index of $\nabla \tilde{\Gamma}$ at \bar{g} is non-zero. By the second expansion in (2.8) and the continuity property of the topological degree, we have

$$\deg(\nabla L_\varepsilon^{\text{red}}|_{\partial^{-1}(\gamma)}, \iota(\Omega \times \{\gamma\}), 0) = \deg(\nabla \Gamma|_{\partial^{-1}(\gamma)}, \iota(\Omega \times \{\gamma\}), 0) = \deg(\nabla \tilde{\Gamma}, \Omega, 0) \neq 0$$

for small values of ε . Hence, for γ fixed, $L_\varepsilon^{\text{red}} \circ \iota(\cdot, \gamma)$ has a critical point $g(\gamma) \in \Omega$. Moreover, we can choose $g(\gamma)$ such that $g(\gamma) = \bar{g} + o(1)$ as $\varepsilon \rightarrow 0$.

If $\tilde{\Gamma}$ is additionally a Morse function on \mathcal{G} , then by the implicit function theorem, $g(\gamma)$ is smooth as a function of $\gamma \in \mathcal{N}$. Note that we can choose the global branch of critical points $g(\gamma)$ defined for all $\gamma \in \mathcal{N}$ by the condition $g(\gamma) = \bar{g} + o(1)$. Hence, the search for critical points of $L_\varepsilon^{\text{red}}$ on \mathcal{M} is reduced to the one for the following functional defined on \mathcal{N}

$$\gamma \mapsto L_\varepsilon^{\text{red}} \circ \iota(g(\gamma), \gamma).$$

By the well-known Lusternik-Schnierelman theory [41], the above function has at least $\text{Cat}(\mathcal{N})$ critical points and we obtain the assertion. \square

Remark 2.5. In the proof of Theorem 2.4, since $\tilde{\Gamma}$ is a Morse function, any critical point of $\tilde{\Gamma}$ is isolated. Hence, for each critical point \bar{g} of $\tilde{\Gamma}$, there exist at least $\text{Cat}(\mathcal{N})$ critical points $(g(\gamma), \gamma) \in \mathcal{G} \times \mathcal{N}$ of $L_\varepsilon^{\text{red}}$ each of which satisfies $g(\gamma) = \bar{g} + o(1)$ as $\varepsilon \rightarrow 0$.

Case 2: $\ker \nabla(\Gamma \circ \iota)(g, \gamma) \equiv T_{(g, \gamma)}(\mathcal{G} \times \mathcal{N})$ for all $(g, \gamma) \in \mathcal{G} \times \mathcal{N}$

In this case, we have $\Gamma \circ \iota(g, \gamma) \equiv \text{constant}$ on $\mathcal{G} \times \mathcal{N}$. As was pointed out in [7], we need to evaluate further terms in the expansion of $L_\varepsilon^{\text{red}}$. For this purpose, let us develop the expansion (2.5) in powers of ε as

$$L_\varepsilon(z) = L_0(z) + \varepsilon\Gamma(z) + \varepsilon^2\Phi(z) + o(\varepsilon^2). \quad (2.11)$$

Note that $\Gamma \circ \iota(g, \gamma) \equiv \text{constant}$ on $\mathcal{G} \times \mathcal{N}$ is equivalent to $\Gamma(z) \equiv \text{constant}$ on \mathcal{M} . It follows that $\nabla\Gamma(z) \in \mathcal{W}_z := T_z\mathcal{M}^\perp$. Recall that $w_\varepsilon(z)$ is solution to the auxiliary equation $P_z\nabla L_\varepsilon(z + w) = 0$, hence we have $\nabla L_\varepsilon(z + w_\varepsilon(z)) \in T_z\mathcal{M}$. For a fixed $z \in \mathcal{M}$, using Taylor expansion, one sees

$$\begin{aligned} \nabla L_\varepsilon(z + w_\varepsilon(z)) &= \nabla L_0(z + w_\varepsilon(z)) + \varepsilon\nabla\Gamma(z + w_\varepsilon(z)) + o(\varepsilon) \\ &= \nabla^2 L_0(z)[w_\varepsilon(z)] + \varepsilon\nabla\Gamma(z) + \varepsilon\nabla^2\Gamma(z)[w_\varepsilon(z)] + o(\|w_\varepsilon(z)\|) + o(\varepsilon). \end{aligned}$$

Then, from (2.7) and the fact $\nabla L_\varepsilon(z + w_\varepsilon(z)) \in T_z\mathcal{M}$, it follows that

$$\nabla^2 L_0(z)[w_\varepsilon(z)] + \varepsilon\nabla\Gamma(z) + o(\varepsilon) \in T_z\mathcal{M}.$$

And hence we deduce

$$w_\varepsilon(z) = -\varepsilon K_z(\nabla\Gamma(z)) + o(\varepsilon),$$

where K_z stands for the inverse of $\nabla^2 L_0(z)$ restricted to \mathcal{W}_z . Now, we can expand $L_\varepsilon^{\text{red}}(z) := L_\varepsilon(z + w_\varepsilon(z))$ as

$$\begin{aligned} L_\varepsilon^{\text{red}}(z) &= L_0(z) + \frac{1}{2}\nabla^2 L_0(z)[w_\varepsilon(z), w_\varepsilon(z)] + \varepsilon\Gamma(z) + \varepsilon\nabla\Gamma(z)[w_\varepsilon(z)] + \varepsilon^2\Phi(z) + o(\varepsilon^2) \\ &= L_0(z) + \varepsilon\Gamma(z) + \varepsilon^2\left(\Phi(z) - \frac{1}{2}\langle K_z(\nabla\Gamma(z)), \nabla\Gamma(z) \rangle\right) + o(\varepsilon^2). \end{aligned}$$

Here, we emphasize that both $L_0(z)$ and $\Gamma(z)$ are constants on \mathcal{M} . Hence by slightly changing the notation, for each fixed ε , the above expansion can be understood as an analogue of the situation considered in (2.8). Therefore, similar to Theorem 2.3, we have the following result.

Theorem 2.6. Let $L_0, \Gamma, \Phi \in C^2(\mathcal{H}, \mathbb{R})$ as in (2.11) and suppose that (A1)-(A5) are satisfied. If $\Gamma \circ \iota \equiv \text{constant}$ on $\mathcal{G} \times \mathcal{N}$ and there is an open bounded subset $U \subset \mathcal{G}$ such that

$$\inf_{\gamma \in \mathcal{N}} \left(\min_{\partial U} \hat{\Phi}|_{\vartheta^{-1}(\gamma)} - \min_{\bar{U}} \hat{\Phi}|_{\vartheta^{-1}(\gamma)} \right) > 0 \quad \text{or} \quad \sup_{\gamma \in \mathcal{N}} \left(\max_{\partial U} \hat{\Phi}|_{\vartheta^{-1}(\gamma)} - \max_{\bar{U}} \hat{\Phi}|_{\vartheta^{-1}(\gamma)} \right) < 0,$$

where $\hat{\Phi}|_{\vartheta^{-1}(\gamma)} = \hat{\Phi} \circ \iota(\cdot, \gamma)$ and

$$\hat{\Phi}(z) := \Phi(z) - \frac{1}{2}\langle K_z(\nabla\Gamma(z)), \nabla\Gamma(z) \rangle \quad \text{for } z \in \mathcal{M}.$$

Then, for $|\varepsilon|$ small, the functional L_ε has a critical point on \mathcal{M} .

Last but not least, analogous to Theorem 2.4, we have

Theorem 2.7. *Let $L_0, \Gamma, \Phi \in C^2(\mathcal{H}, \mathbb{R})$ as in (2.11) and suppose that (A1)-(A5) are satisfied. Assume further that \mathcal{N} is simply connected. If $\Gamma \circ \iota \equiv \text{constant}$ on $\mathcal{G} \times \mathcal{N}$ and, for each $\gamma \in \mathcal{N}$, if $\hat{\Phi}|_{\vartheta^{-1}(\gamma)} = \hat{\Phi} \circ \iota(\cdot, \gamma)$ is a Morse function on \mathcal{G} and there is an open bounded subset $\Omega(\gamma) \subset \mathcal{G}$ such that the topological degree $\deg(\nabla \hat{\Phi}|_{\vartheta^{-1}(\gamma)}, \iota(\Omega(\gamma) \times \{\gamma\}), 0) \neq 0$. Then, for $|\varepsilon|$ small, the functional $L_\varepsilon^{\text{red}}$ has at least $\text{Cat}(\mathcal{N})$ critical points on \mathcal{M} .*

Proof. By the assumptions, for each $\gamma \in \mathcal{N}$, there exists a critical point $g(\gamma) \in \Omega(\gamma) \subset \mathcal{G}$. For any $\gamma_0 \in \mathcal{N}$, by the Morse condition, we can choose a small neighborhood of γ_0 such that $g(\gamma)$ depends smoothly on γ in this neighborhood. These local branches of critical points are globally continued due to the simply connectivity of \mathcal{N} and we obtain a smooth $g(\gamma)$ defined globally on \mathcal{N} . Then the proof proceed as in Theorem 2.4. \square

3 Perturbations of the geometric potential

In this section we deal with the problem (2.1), namely,

$$D_{g_{\mathbb{R}^m}} \psi = (1 + \varepsilon a(x)) |\psi|_{g_{\mathbb{R}^m}}^{2^*-2} \psi \quad \text{on } \mathbb{R}^m$$

with $2^* = \frac{2m}{m-1}$. The associated functional can be written in the form

$$\mathcal{J}(\psi) = \mathcal{J}_0(\psi) + \varepsilon \Gamma(\psi) \tag{3.1}$$

where

$$\mathcal{J}_0(\psi) = \frac{1}{2} \int_{\mathbb{R}^m} (\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} - \frac{1}{2^*} \int_{\mathbb{R}^m} |\psi|_{g_{\mathbb{R}^m}}^{2^*} d \text{vol}_{g_{\mathbb{R}^m}}$$

and

$$\Gamma(\psi) = -\frac{1}{2^*} \int_{\mathbb{R}^m} a(x) |\psi|_{g_{\mathbb{R}^m}}^{2^*} d \text{vol}_{g_{\mathbb{R}^m}}.$$

We intend to use Theorem 2.4 to obtain Theorem 1.1. In order to do this, we need to have a detailed behavior of the functional Γ on the critical manifold \mathcal{M} mentioned in (2.9)-(2.10).

3.1 The topological degree

Recall that, for a function $H \in C^2(S^m)$, we have assumed H takes its minimum at $p_0 \in S^m$ and denote $\pi_{p_0} : S^m \setminus \{p_0\} \rightarrow \mathbb{R}^m$ the stereographic projection, and then the function a is defined by $a(x) := H(\pi_{p_0}^{-1}(x))$ for $x \in \mathbb{R}^m$.

In what follows, we always assume that $a = H \circ \pi_{p_0}^{-1}$ for some function $H \in C^2(S^m)$ and, moreover, the conditions (H-1) and (H-2) are also satisfied.

Lemma 3.1. *The function a has the following properties:*

- (1) $a \in L^\infty(\mathbb{R}^m) \cap C^2(\mathbb{R}^m)$ and there is a constant $C_0 > 0$ such that

$$|\partial_i a(x)| \leq C_0(1 + |x|^2)^{-1} \quad \text{and} \quad |\partial_i \partial_j a(x)| \leq C_0(1 + |x|^2)^{-\frac{3}{2}}.$$

where $\partial_i = \frac{\partial}{\partial x_i}$ and $\partial_i \partial_j = \frac{\partial^2}{\partial x_i \partial x_j}$.

- (2) a is a Morse function. The number of critical points of a on \mathbb{R}^m is finite and $\Delta_{g_{\mathbb{R}^m}} a(x) \neq 0$ for $x \in \text{Crit}[a]$. Moreover

$$\sum_{x \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(x) < 0} (-1)^{\mathfrak{M}(a,x)} \neq (-1)^m$$

where $\text{Crit}[a]$ stands for the critical set of a on \mathbb{R}^m and $\mathfrak{M}(a, x)$ is the Morse index of a at x .

- (3) there is $R_0 > 0$ such that $\nabla a(x) \cdot x < 0$ for $|x| \geq R_0$ and

$$\int_{B_R} \nabla a(x) \cdot x \, d \text{vol}_{g_{\mathbb{R}^m}} < 0$$

for any $R > R_0$.

Proof. For (1) and (2) we refer to [27, Lemma 4.1] for a detailed proof. Notice that a similar statements of (3) were already proved in [28, Lemma 3.2]. Although, the dimension in [28, Lemma 3.2] was fixed as $m = 3$, the same argument applies and hence we omit the proof. \square

Note that, for $\psi_{\lambda, \xi, \gamma} \in \mathcal{M}$ (see (2.10)),

$$\Gamma(\psi_{\lambda, \xi, \gamma}) = -\frac{m-1}{2m} \int_{\mathbb{R}^m} a(x) |\psi_{\lambda, \xi, \gamma}|_{g_{\mathbb{R}^m}}^{\frac{2m}{m-1}} d \text{vol}_{g_{\mathbb{R}^m}}$$

and

$$|\psi_{\lambda, \xi, \gamma}|_{g_{\mathbb{R}^m}} = \frac{m^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m-1}{2}}}.$$

Then it is clear that $\Gamma(\psi_{\lambda, \xi, \gamma})$ is independent of the factor $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(S_m)$. Hence, in order to study $\Gamma(\psi_{\lambda, \xi, \gamma})$, it sufficient to consider (up to a constant)

$$\Psi(\lambda, \xi) := \int_{\mathbb{R}^m} a(x) |\psi_{\lambda, \xi, \gamma}|_{g_{\mathbb{R}^m}}^{\frac{2m}{m-1}} d \text{vol}_{g_{\mathbb{R}^m}} = m^m \int_{\mathbb{R}^m} \frac{a(\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}$$

where the last equality comes from a change of variable. Now, by Lemma 3.1 (1), one checks easily that, for any $\xi \in \mathbb{R}^m$,

$$\lim_{\lambda \rightarrow 0^+} \Psi(\lambda, \xi) = C_1 a(\xi), \quad \text{where } C_1 = m^m \int_{\mathbb{R}^m} \frac{1}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}, \quad (3.2)$$

$$\partial_\lambda \Psi(\lambda, \xi) = m^m \int_{\mathbb{R}^m} \frac{\nabla a(\lambda x + \xi) \cdot x}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}$$

and

$$\lim_{\lambda \rightarrow 0^+} \partial_\lambda \Psi(\lambda, \xi) = m^m \int_{\mathbb{R}^m} \frac{\nabla a(\xi) \cdot x}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} = 0 \quad (3.3)$$

With the above observation, by employing the ideas of Ambrosetti et al in [5], we intend to extend Ψ on $(-\infty, \infty) \times \mathbb{R}^m$ as

$$\tilde{\Psi}(0, \xi) = C_1 a(\xi), \quad \tilde{\Psi}(\lambda, \xi) = \Psi(-\lambda, \xi) \text{ for } \lambda < 0.$$

By (3.2) and (3.3), $\tilde{\Psi}$ is of C^1 on $(-\infty, \infty) \times \mathbb{R}^m$ and

$$\partial_\lambda \tilde{\Psi}(0, \xi) = 0 \quad \text{for } \xi \in \mathbb{R}^m.$$

Moreover, we have

$$\partial_i \tilde{\Psi}(0, \xi) = \lim_{\lambda \rightarrow 0} m^m \int_{\mathbb{R}^m} \frac{\partial_i a(\lambda x + \xi)}{(1 + |x|^2)^m} d\text{vol}_{g_{\mathbb{R}^m}} = C_1 \partial_i a(\xi) \quad \text{for } \xi \in \mathbb{R}^m.$$

Hence, we find the following one-to-one correspondence:

$$\xi \in \text{Crit}[a] \iff (0, \xi) \in \text{Crit}[\tilde{\Psi}] \quad (3.4)$$

Next, we will characterize the local index of $\tilde{\Psi}$ at critical points of the form $(0, \xi)$ and evaluate the topological degree of $\tilde{\Psi}$ in a bounded subset of $(0, \infty) \times \mathbb{R}^m$. In what follows (for the proofs of Lemma 3.2, Lemma 3.3 and Corollary 3.4), since the calculation is essentially the same as in [28, Section 3], we shall sketch most of the details for general case $m \geq 3$. Here, for the sake of completeness, we mainly specialize to the proofs for the case $m = 2$, where some additional arguments are required.

Lemma 3.2. *There exist at most finitely many critical points of $\tilde{\Phi}$ in the form of $(0, \xi) \in (-\infty, \infty) \times \mathbb{R}^m$. Moreover, these critical points are isolated and local index of $\tilde{\Psi}$ at a critical point $(0, \xi)$ is*

$$\text{ind}(\nabla \tilde{\Phi}, (0, \xi)) = \begin{cases} (-1)^{\mathfrak{M}(a, \xi)} & \text{if } \Delta_{g_{\mathbb{R}^m}} a(\xi) > 0, \\ (-1)^{\mathfrak{M}(a, \xi) + 1} & \text{if } \Delta_{g_{\mathbb{R}^m}} a(\xi) < 0. \end{cases}$$

Proof. For $m \geq 3$, it follows directly from the computation in [27, Section 7] that

$$\lim_{\lambda \rightarrow 0} \partial_\lambda^2 \tilde{\Psi}(\lambda, \xi) = C_2 \Delta_{g_{\mathbb{R}^m}} a(\xi) \quad \text{where } C_2 = m^{m-1} \int_{\mathbb{R}^m} \frac{|x|^2}{(1 + |x|^2)^m} d\text{vol}_{g_{\mathbb{R}^m}}$$

$$\lim_{\lambda \rightarrow 0} \partial_i \partial_j \tilde{\Psi}(\lambda, \xi) = C_1 \partial_i \partial_j a(\xi)$$

and

$$\lim_{\lambda \rightarrow 0} \partial_i \partial_\lambda \tilde{\Psi}(\lambda, \xi) = 0$$

Hence the Hessian of $\tilde{\Phi}$ at $(0, \xi)$ is

$$\nabla^2 \tilde{\Phi}(0, \xi) = \begin{pmatrix} C_2 \Delta_{g_{\mathbb{R}^m}} a(\xi) & O \\ O & C_1 \nabla^2 a(\xi) \end{pmatrix}.$$

If $(0, \xi_0) \in \text{Crit}[\tilde{\Phi}]$, then one sees that

$$\tilde{\Phi}(\lambda, \xi) \sim \tilde{\Phi}(0, \xi_0) + C_2 \Delta_{g_{\mathbb{R}^m}} a(\xi_0) \lambda^2 + C_1 \nabla^2 a(\xi_0) [\xi - \xi_0, \xi - \xi_0]$$

provided (λ, ξ) is close to $(0, \xi_0)$. Hence $(0, \xi_0)$ is isolated and the assertion follows from the multiplicative property of the index.

For $m = 2$, $\partial_\lambda^2 \tilde{\Phi}(0, \xi) = \lim_{\lambda \rightarrow 0} \partial_\lambda^2 \tilde{\Phi}(\lambda, \xi)$ does not exist. Thus, we need to know detailed behavior of $\partial_\lambda \tilde{\Phi}(\lambda, \xi)$ as $\lambda \rightarrow 0$.

To this end, for $\lambda > 0$, let us write

$$\begin{aligned} \partial_\lambda \Phi(\lambda, \xi) &= 4 \sum_{i=1}^2 \int_{\mathbb{R}^2} \frac{\partial_i a(\lambda x + \xi) x_i}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} \\ &= 4 \sum_{i=1}^2 \int_{|x| \leq \frac{1}{\lambda}} \frac{\partial_i a(\lambda x + \xi) x_i}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} + 4 \sum_{i=1}^2 \int_{|x| \geq \frac{1}{\lambda}} \frac{\partial_i a(\lambda x + \xi) x_i}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} \\ &= 4I_1(\lambda, \xi) + 4I_2(\lambda, \xi). \end{aligned}$$

Note that $a \in C^2(\mathbb{R}^2)$, by Taylor's theorem, we have

$$\partial_i a(\lambda x + \xi) = \partial_i a(\xi) + \lambda \nabla \partial_i a(\xi) \cdot x + o(|\lambda x|)$$

where $o(|\lambda x|) \rightarrow 0$ as $|\lambda x| \rightarrow 0$. Hence, we deduce

$$\begin{aligned} I_1(\lambda, \xi) &= \sum_{i=1}^2 \int_{|x| \leq \frac{1}{\lambda}} \frac{\partial_i a(\xi) x_i}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} + \sum_{i=j}^2 \sum_{i=1}^2 \int_{|x| \leq \frac{1}{\lambda}} \frac{\lambda \partial_j \partial_i a(\xi) x_j x_i}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} \\ &\quad + o\left(\int_{|x| \leq \frac{1}{\lambda}} \frac{\lambda |x|^2}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} \right) \\ &= \frac{\lambda}{2} \Delta_{g_{\mathbb{R}^2}} a(\xi) \int_{|x| \leq \frac{1}{\lambda}} \frac{|x|^2}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} + o(\lambda \ln \lambda) \\ &= -\pi \lambda \ln \lambda (\Delta_{g_{\mathbb{R}^2}} a(\xi) + o(1)) + o(\lambda) \end{aligned}$$

and, by Lemma 3.1 (1),

$$I_2(\lambda, \xi) = O\left(\int_{|x| \geq \frac{1}{\lambda}} \frac{|x|}{(1 + |x|^2)^2} d \text{vol}_{g_{\mathbb{R}^2}} \right) = O(\lambda)$$

as $\lambda \rightarrow 0$. Thus, we have

$$\partial_\lambda \Phi(\lambda, \xi) = -4\pi \lambda \ln \lambda (\Delta_{g_{\mathbb{R}^2}} a(\xi) + o(1))$$

as $\lambda \rightarrow 0$. If $(0, \xi_0) \in \text{Crit}[\tilde{\Psi}]$, since $\Delta_{g_{\mathbb{R}^2}} a(\xi_0) \neq 0$, it is clear that the function

$$\lambda \mapsto -4\pi \Delta_{g_{\mathbb{R}^2}} a(\xi_0) \lambda \ln \lambda$$

is homotopic to the function $\lambda \mapsto \Delta_{g_{\mathbb{R}^2}} a(\xi_0) \lambda$ through homotopy which does not take value 0 when λ is close to 0. Hence

$$\text{ind}(\partial_\lambda \tilde{\Psi}(\cdot, \xi_0), 0) = \begin{cases} 1 & \text{if } \Delta_{g_{\mathbb{R}^2}} a(\xi_0) > 0, \\ -1 & \text{if } \Delta_{g_{\mathbb{R}^2}} a(\xi_0) < 0. \end{cases}$$

And therefore, the assertion follows immediately due to the multiplicative property of the index. \square

Lemma 3.3. *For $m \geq 2$, there exists $R > 0$ such that $\text{Crit}[\tilde{\Psi}] \subset B_R^{m+1} := \{(\lambda, \xi) \in (-\infty, \infty) \times \mathbb{R}^m : \lambda^2 + |\xi|^2 \leq R^2\}$ and*

$$\deg(\nabla \tilde{\Psi}, B_R^{m+1}, 0) = (-1)^{m+1}.$$

Proof. Although the arguments in [28] apply to the general case $m \geq 2$ without any difficulty, for completeness, we sketch the proof as follows.

We first show that there exists $R > 0$ such that

$$\nabla \tilde{\Psi}(\lambda, \xi) \cdot (\lambda, \xi) < 0 \quad (3.5)$$

for $(\lambda, \xi) \in (-\infty, \infty) \times \mathbb{R}^m$ with $\lambda^2 + |\xi|^2 \geq R^2$.

To see this, let us compute

$$\begin{aligned} \nabla \tilde{\Psi}(\lambda, \xi) \cdot (\lambda, \xi) &= m^m \int_{\mathbb{R}^m} \frac{\nabla a(\lambda x + \xi) \cdot (\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \\ &= m^m \lambda^{-m} \int_{\mathbb{R}^m} \frac{\lambda^{2m} \nabla a(x) \cdot x}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}. \end{aligned} \quad (3.6)$$

Observe that, for arbitrary $r > 0$, we can decompose the above integral as

$$\begin{aligned} \int_{\mathbb{R}^m} \frac{\lambda^{2m} \nabla a(x) \cdot x}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} &= \int_{|x| \leq r} \frac{\lambda^{2m} \nabla a(x) \cdot x}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad + \int_{|x| \geq r} \frac{\lambda^{2m} \nabla a(x) \cdot x}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \\ &= I_1(r, \lambda, \xi) + I_2(r, \lambda, \xi) \end{aligned}$$

One checks easily that, by the first half of Lemma 3.1 (3), $I_2(r, \lambda, \xi) < 0$ if $r > R_0$. On the other hand, we can decompose $I_1(r, \lambda, \xi)$ as

$$\begin{aligned} I_1(r, \lambda, \xi) &= \int_{|x| \leq r} \frac{\lambda^{2m} (\nabla a(x) \cdot x)_+}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} - \int_{|x| \leq r} \frac{\lambda^{2m} (\nabla a(x) \cdot x)_-}{(\lambda^2 + |x - \xi|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \\ &\leq \max_{|x| \leq r} \frac{\lambda^{2m}}{(\lambda^2 + |x - \xi|^2)^m} \int_{|x| \leq r} (\nabla a(x) \cdot x)_+ d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad - \min_{|x| \leq r} \frac{\lambda^{2m}}{(\lambda^2 + |x - \xi|^2)^m} \int_{|x| \leq r} (\nabla a(x) \cdot x)_- d \text{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

where $(\nabla a(x) \cdot x)_+$ and $(\nabla a(x) \cdot x)_-$ are the positive and the negative parts of $\nabla a(x) \cdot x$, respectively. To proceed, by the second half of Lemma 3.1 (3), we mention that

$$\delta \int_{|x| \leq r} (\nabla a(x) \cdot x)_+ d \text{vol}_{g_{\mathbb{R}^m}} + \int_{|x| \leq r} \nabla a(x) \cdot x d \text{vol}_{g_{\mathbb{R}^m}} < 0 \quad (3.7)$$

provided that $r \geq R_0$ and δ fixed small enough. Set

$$M(r, \lambda, \xi) = \max_{|x| \leq r} \frac{\lambda^{2m}}{(\lambda^2 + |x - \xi|^2)^m} = \begin{cases} \frac{\lambda^{2m}}{(\lambda^2 + (|\xi| - r)^2)^m} & \text{if } |\xi| > r \\ 1 & \text{if } |\xi| \leq r \end{cases}$$

and

$$m(r, \lambda, \xi) = \min_{|x| \leq r} \frac{\lambda^{2m}}{(\lambda^2 + |x - \xi|^2)^m} = \frac{\lambda^{2m}}{(\lambda^2 + (r + |\xi|)^2)^m},$$

we find $\lim_{\lambda^2 + |\xi|^2 \rightarrow \infty} \frac{M(r, \lambda, \xi)}{m(r, \lambda, \xi)} = 1$. Hence, there exists $R_1 > 0$ such that

$$1 \leq \frac{M(r, \lambda, \xi)}{m(r, \lambda, \xi)} \leq 1 + \delta$$

for $\lambda^2 + |\xi|^2 \geq R_1$. Now, by (3.7), we can get the following estimate

$$\begin{aligned} I_1(r, \lambda, \xi) &\leq M(r, \lambda, \xi) \int_{|x| \leq r} (\nabla a(x) \cdot x)_+ d \text{vol}_{g_{\mathbb{R}^m}} - m(r, \lambda, \xi) \int_{|x| \leq r} (\nabla a(x) \cdot x)_- d \text{vol}_{g_{\mathbb{R}^m}} \\ &\leq m(r, \lambda, \xi) \left(\delta \int_{|x| \leq r} (\nabla a(x) \cdot x)_+ d \text{vol}_{g_{\mathbb{R}^m}} + \int_{|x| \leq r} \nabla a(x) \cdot x d \text{vol}_{g_{\mathbb{R}^m}} \right) < 0, \end{aligned}$$

which indicates (3.5).

Since (3.5) suggests $\text{Crit}[\tilde{\Psi}] \subset B_R^{m+1}$ for some $R > 0$ large, it remains to evaluate the topological degree of $\tilde{\Psi}$. Note that (3.5) forces $-t(\lambda, \xi) + (1-t)\nabla\tilde{\Psi}(\lambda, \xi) \neq 0$ for any $(\lambda, \xi) \in \partial B_R^{m+1}$ and $t \in [0, 1]$. It follows that $\nabla\tilde{\Psi}(\cdot, \cdot)$ is homotopic to $-id$ on B_R^{m+1} through homotopy which does not take value 0 on ∂B_R^{m+1} and, particularly,

$$\deg(\nabla\tilde{\Psi}, B_R^{m+1}, 0) = \deg(-id, B_R^{m+1}, 0) = (-1)^{m+1}.$$

This completes the proof. \square

To get the topological degree of Ψ on $(0, \infty) \times \mathbb{R}^m$, let us decompose $\text{Crit}[\tilde{\Psi}]$ as

$$\text{Crit}[\tilde{\Psi}] = \text{Crit}_+[\tilde{\Psi}] \cup \text{Crit}_0[\tilde{\Psi}] \cup \text{Crit}_-[\tilde{\Psi}]$$

where

$$\text{Crit}_\pm[\tilde{\Psi}] = \{(\lambda, \xi) \in \text{Crit}[\tilde{\Psi}] : \pm\lambda > 0\} \quad \text{and} \quad \text{Crit}_0[\tilde{\Psi}] = \{(\lambda, \xi) \in \text{Crit}[\tilde{\Psi}] : \lambda = 0\}.$$

By the symmetry of $\tilde{\Psi}$, we have

$$\text{Crit}_-[\tilde{\Psi}] = \{(\lambda, \xi) : (-\lambda, \xi) \in \text{Crit}_+[\tilde{\Psi}]\}.$$

Thanks to Lemma 3.2 and 3.3, we can find a bounded domain $\Omega_+ \subset (0, \infty) \times \mathbb{R}^m$ such that

$$\Omega_+ \subset B_R^{m+1} \quad \text{and} \quad \text{Crit}_+[\tilde{\Psi}] \subset \Omega_+.$$

As a consequence of Lemma 3.3, we have

Corollary 3.4. $\deg(\nabla\Psi, \Omega_+, 0) \neq 0$.

Proof. Since $\Psi = \tilde{\Psi}|_{(0, \infty) \times \mathbb{R}^m}$, it is sufficient to evaluate $\deg(\nabla\tilde{\Psi}, \Omega_+, 0)$.

Assume to the contrary that $\deg(\nabla\tilde{\Psi}, \Omega_+, 0) = 0$. Let us set

$$\Omega_- = \{(\lambda, \xi) : (-\lambda, \xi) \in \Omega_+\}.$$

Then, it is clear that $\text{Crit}_-[\tilde{\Psi}] \subset \Omega_-$ and, by the assumption, $\deg(\nabla\tilde{\Psi}, \Omega_-, 0) = 0$. Due to the additive property of the degree, we can see from Lemma 3.2 and 3.3

$$\begin{aligned} (-1)^{m+1} &= \deg(\nabla\tilde{\Psi}, B_R^{m+1}, 0) \\ &= \deg(\nabla\tilde{\Psi}, \Omega_+, 0) + \deg(\nabla\tilde{\Psi}, \Omega_-, 0) + \deg(\nabla\tilde{\Psi}, B_R^{m+1} \setminus (\Omega_+ \cup \Omega_-), 0) \\ &= 0 + 0 + \sum_{(0, \xi) \in \text{Crit}_0[\tilde{\Psi}]} \text{ind}(\nabla\tilde{\Psi}, (0, \xi)) \\ &= \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) > 0} \text{ind}(\nabla a, \xi) - \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) < 0} \text{ind}(\nabla a, \xi). \end{aligned} \quad (3.8)$$

On the other hand, by Lemma 3.1 (3), we can proceed analogously to the proof of Lemma 3.3 to get

$$\deg(\nabla a, B_R, 0) = (-1)^m$$

where $B_R = \{x \in \mathbb{R}^m : |x| \leq R\}$. Since

$$\deg(\nabla a, B_R, 0) = \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) > 0} \text{ind}(\nabla a, \xi) + \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) < 0} \text{ind}(\nabla a, \xi),$$

it follows from (3.8) that

$$\sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) > 0} \text{ind}(\nabla a, \xi) = 0 \quad \text{and} \quad \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) < 0} \text{ind}(\nabla a, \xi) = (-1)^m.$$

Therefore, we have

$$\sum_{x \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(x) < 0} (-1)^{\mathfrak{M}(a, x)} = \sum_{\xi \in \text{Crit}[a], \Delta_{g_{\mathbb{R}^m}} a(\xi) < 0} \text{ind}(\nabla a, \xi) = (-1)^m$$

which contradicts to Lemma 3.1 (2). \square

3.2 Proof of the main result

For $l \in \mathbb{N}$, let $a_1, \dots, a_l \in L^\infty(\mathbb{R}^m) \cap C^2(\mathbb{R}^m)$ be functions satisfying the properties mentioned in Lemma 3.1. For $z_1, \dots, z_l \in \mathbb{R}^m$, we consider

$$a(x) = \sum_{i=1}^l a_i(x - z_i).$$

Then for $(\lambda, \xi) \in (0, \infty) \times \mathbb{R}^m$ the function Ψ takes the form

$$\Psi(\lambda, \xi) = \sum_{i=1}^l \Psi_i(\lambda, \xi)$$

where

$$\Psi_i(\lambda, \xi) = m^m \int_{\mathbb{R}^m} \frac{a_i(\lambda x + \xi - z_i)}{(1 + |x|^2)^m} d\text{vol}_{g_{\mathbb{R}^m}}$$

Note that Ψ_i is independent of the variable $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$.

The proofs in the previous section indicates that there exist bounded domains $\Omega_i \subset (0, \infty) \times \mathbb{R}^m$, $i = 1, \dots, l$, such that $\text{Crit}_+[\Psi_i] \subset \Omega_i(z_i)$ and

$$\deg(\nabla \Psi_i, \Omega_i(z_i), 0) \neq 0$$

where $\text{Crit}_+[\Psi_i] = \{(\lambda, \xi) : \nabla \Psi_i(\lambda, \xi) = 0 \text{ and } \lambda > 0\}$ and $\Omega_i(z_i) = \{(\lambda, x) : (\lambda, x - z_i) \in \Omega_i\}$. Then, in order to apply Theorem 2.4, we are left with tasks to show

(T1) $\nabla \Psi_i(\lambda, \xi) \rightarrow 0$ as $\lambda + |\xi - z_i| \rightarrow \infty$;

(T2) for generic choice of a_i ($i = 1, \dots, l$), Ψ_i is a Morse function of (λ, ξ) .

Notice that, if z_1, \dots, z_l are located far apart, $\Omega_i(z_i) \cap \Omega_j(z_j) = \emptyset$ provided $i \neq j$. Thus, if the above two statements are proven, Ψ has at least one critical point in each $\Omega_i(z_i)$ provided that $\min_{1 \leq i \neq j \leq l} |z_i - z_j|$ is large enough. Then, recall that

$$\Gamma(\psi_{\lambda, \xi, \gamma}) = -\frac{m-1}{2m} \Psi(\lambda, \xi) \quad \text{for } \psi_{\lambda, \xi, \gamma} \in \mathcal{M}$$

and

$$\mathcal{J}(\psi) = \mathcal{J}_0(\psi) + \varepsilon \Gamma(\psi) \quad \text{for } \psi \in \mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m)).$$

By applying Theorem 2.4 and Remark 2.5, the functional \mathcal{J} has at least $l \cdot \text{Cat}(S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m))$ critical points. However, these critical points may appear as an S^1 -orbit (or S^3 -orbit when $m = 2, 3, 4 \pmod{8}$) of a single solution. To find non- S^1 -equivalent (or S^3 -equivalent) solutions, we observe that, due to the S^1 -equivariance (or S^3 -equivariance) property of Eq. (2.6), our construction of $w_\varepsilon = w_\varepsilon(z)$ can be done equivariantly with respect to these actions. Namely, $w_\varepsilon(z)$ satisfies $w_\varepsilon(\tau \star z) = \tau \star w_\varepsilon(z)$ for $z \in \mathcal{M}$ and $\tau \in S^1$ (or $\tau \in S^3$), where “ \star ” stands for the corresponding group action. Then the reduced functional $L_\varepsilon^{\text{red}}$ is S^1 (or S^3)-invariant, $L_\varepsilon^{\text{red}} \circ \iota(g, \tau \star \gamma) = L_\varepsilon^{\text{red}} \circ \iota(g, \gamma)$ for all $(g, \gamma) \in \mathcal{G} \times \mathcal{N}$ and $\tau \in S^1$ (or $\tau \in S^3$ when $m = 2, 3, 4 \pmod{8}$). Therefore, $L_\varepsilon^{\text{red}}$ can be reduced to be defined on $\mathcal{G} \times \mathcal{N}/S^1$ (or $\mathcal{G} \times \mathcal{N}/S^3$ when $m = 2, 3, 4 \pmod{8}$). We define $\hat{L}_\varepsilon^{\text{red}}(g, [\gamma]) = L_\varepsilon^{\text{red}}(g, \gamma)$, where $[\gamma] \in \mathcal{N}/S^1$ (or \mathcal{N}/S^3) is the equivalence class represented by $\gamma \in \mathcal{N}$. Since $\mathcal{N} = S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$ is homeomorphic to the unit sphere in $\mathbb{C}^{2^{\lfloor \frac{m}{2} \rfloor}}$ (and in $\mathbb{H}^{2^{\lfloor \frac{m}{2} \rfloor - 1}}$ when $m = 2, 3, 4 \pmod{8}$), $\mathcal{N}/S^1 \cong \mathbb{C}P^{2^{\lfloor \frac{m}{2} \rfloor - 1}}$ ($\mathcal{N}/S^3 \cong \mathbb{H}P^{2^{\lfloor \frac{m}{2} \rfloor - 1} - 1}$ when $m = 2, 3, 4 \pmod{8}$). To estimate the Lusternik-Schnirelmann category of these quotients, recall that the category of a topological space X is estimated from below by the cuplength, $\text{Cat}(X) \geq \text{CL}(X) + 1$. Here the cuplength of X is defined as the greatest integer $k \geq 0$ such that there exist $\omega_1, \dots, \omega_k \in H^*(X; \mathbb{Z})$ so that $\deg \omega_j > 0$ and $\omega_1 \cup \dots \cup \omega_k \neq 0$.

The cohomology rings of $\mathbb{C}P^{2^{\lfloor \frac{m}{2} \rfloor - 1}}$ and $\mathbb{H}P^{2^{\lfloor \frac{m}{2} \rfloor - 1} - 1}$ are respectively isomorphic to truncated polynomial rings $H^*(\mathbb{C}P^{2^{\lfloor \frac{m}{2} \rfloor - 1}); \mathbb{Z}) = \mathbb{Z}[\alpha]/(\alpha^{2^{\lfloor \frac{m}{2} \rfloor}})$ and $H^*(\mathbb{H}P^{2^{\lfloor \frac{m}{2} \rfloor - 1} - 1}; \mathbb{Z}) = \mathbb{Z}[\beta]/(\beta^{2^{\lfloor \frac{m}{2} \rfloor - 1}})$, where $\alpha \in H^2(\mathbb{C}P^{2^{\lfloor \frac{m}{2} \rfloor - 1}); \mathbb{Z})$ and $\beta \in H^4(\mathbb{H}P^{2^{\lfloor \frac{m}{2} \rfloor - 1} - 1}; \mathbb{Z})$. Thus the cuplength of the quotient \mathcal{N}/S^1 and \mathcal{N}/S^3 are $\text{CL}(\mathcal{N}/S^1) = 2^{\lfloor \frac{m}{2} \rfloor} - 1$ and $\text{CL}(\mathcal{N}/S^3) = 2^{\lfloor \frac{m}{2} \rfloor - 1} - 1$ when $m = 2, 3, 4 \pmod{8}$. Therefore, we have $\text{Cat}(\mathcal{N}/S^1) \geq 2^{\lfloor \frac{m}{2} \rfloor}$ and $\text{Cat}(\mathcal{N}/S^3) \geq 2^{\lfloor \frac{m}{2} \rfloor - 1}$ when $m = 2, 3, 4 \pmod{8}$. From this, we conclude that the functional \mathcal{J} has at least $2^{\lfloor \frac{m}{2} \rfloor} l$ critical points whose S^1 -orbits are different in the general case $m \geq 2$ and when $m = 2, 3, 4 \pmod{8}$ there are at least $2^{\lfloor \frac{m}{2} \rfloor - 1} l$ critical points whose S^3 -orbits are different.

To study the zero sets of these solutions, we first observe from Remark 2.5 that: let ψ_ε be one of the solutions obtained above, then ψ_ε converge to some $\psi_{\lambda, \xi, \gamma} \in \mathcal{M}$, as $\varepsilon \rightarrow 0$. Note that

$$\lim_{|x| \rightarrow \infty} f^{-\frac{m-1}{2}}(x) |\psi_{\lambda, \xi, \gamma}(x)|_{g_{\mathbb{R}^m}} = \left(\frac{m}{2}\right)^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}} > 0, \quad \text{where } f(x) = \frac{2}{1 + |x|^2}$$

it follows that $\psi_{\lambda, \xi, \gamma}$ corresponds to a solution φ_0 to the equation

$$D_{g_{S^m}} \varphi = |\varphi|_{g_{S^m}}^{2^* - 2} \varphi \quad \text{on } S^m$$

via stereographic projection (see in Section 2) with $|\varphi_0|_{g_{S^m}} > 0$ on S^m . Hence, by the convergence of $\{\psi_\varepsilon\}$, if we pull-back ψ_ε on S^m we have

$$|(f^{-\frac{m-1}{2}} \psi_\varepsilon) \circ \pi_p|_{g_{S^m}} = |\varphi_0|_{g_{S^m}} + o_\varepsilon(1) > 0$$

as $\varepsilon \rightarrow 0$. And hence, the corresponding solutions to the Eq. (1.7) have no zero on S^m .

In what follows, the proofs of the above statements (T1) and (T2) will be carried out in Lemma 3.5 and Proposition 3.6.

Lemma 3.5. *For each i , $\nabla \Psi_i(\lambda, \xi) \rightarrow 0$ as $\lambda + |\xi - z_i| \rightarrow \infty$.*

Proof. This easily follows from Lemma 3.1 (1) and the dominated convergence theorem. \square

Proposition 3.6. *For $H \in C^2(S^m)$ and $(\lambda, \xi) \in (0, \infty) \times \mathbb{R}^m$, we define*

$$\Psi(\lambda, \xi; H) = m^m \int_{\mathbb{R}^m} \frac{a(\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}$$

where $a = H \circ \pi_p^{-1}$ ($\pi_p : S^m \setminus \{p\} \rightarrow \mathbb{R}^m$ is the stereographic projection with respect to a fixed $p \in S^m$). Then there exists a residual subset $\mathcal{A} \subset C^2(S^m)$ such that, for all $H \in \mathcal{A}$, Ψ is a Morse function of $(0, \infty) \times \mathbb{R}^m$.

Recall that a subset \mathcal{S} in a Banach space \mathcal{B} is residual if \mathcal{S} contains a countable intersection of open dense subset of \mathcal{B} . By the Baire's category theorem, residual set is dense in \mathcal{B} .

To prove the above proposition, we need some preparations. First, we define

$$\mathcal{F}(H, \lambda, \xi) = \nabla \Psi(\lambda, \xi; H) \in \mathbb{R}^{m+1}.$$

Let us consider the following universal critical set

$$\mathcal{C} = \{(H, \lambda, \xi) \in C^2(S^m) \times (0, \infty) \times \mathbb{R}^m : \mathcal{F}(H, \lambda, \xi) = 0\},$$

where $C^2(S^m) \times (0, \infty) \times \mathbb{R}^m$ is a separable Banach manifold modeled by $C^2(S^m) \times \mathbb{R} \times \mathbb{R}^m$ with a norm $\|(K, \mu, \zeta)\| = \|K\|_{C^2(S^m)} + |\mu| + |\zeta|$ for $(K, \mu, \zeta) \in T_{(H, \lambda, \xi)}(C^2(S^m) \times (0, \infty) \times \mathbb{R}^m) = C^2(S^m) \times \mathbb{R} \times \mathbb{R}^m$. Note that $(H, \lambda, \xi) \in \mathcal{C}$ if and only if $(\lambda, \xi) \in (0, \infty) \times \mathbb{R}^m$ is a critical point of $\Psi(\cdot, \cdot; H)$.

Lemma 3.7. $0 \in \mathbb{R}^{m+1}$ is a regular value of \mathcal{F} , that is, the derivative

$$\nabla \mathcal{F}(H, \lambda, \xi) : C^2(S^m) \times \mathbb{R} \times \mathbb{R}^m \rightarrow \mathbb{R}^{m+1}$$

is surjective at any $(H, \lambda, \xi) \in \mathcal{C}$.

Proof. Let $(H, \lambda, \xi) \in \mathcal{C}$, we have

$$\mathcal{F}(H, \lambda, \xi) = \left(m^m \int_{\mathbb{R}^m} \frac{\nabla a(\lambda x + \xi) \cdot x}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}, m^m \int_{\mathbb{R}^m} \frac{\nabla a(\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \right) \in \mathbb{R} \times \mathbb{R}^m.$$

We shall prove that the restriction of $\nabla \mathcal{F}$ to the subspace $C^2(S^m) \times \{0\} \times \{0\}$ is onto. Note that

$$\begin{aligned} \nabla \mathcal{F}(H, \lambda, \xi)[K, 0, 0] &= \left(m^m \int_{\mathbb{R}^m} \frac{\nabla b(\lambda x + \xi) \cdot x}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}, m^m \int_{\mathbb{R}^m} \frac{\nabla b(\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \right) \\ &= \mathcal{F}(K, \lambda, \xi) \end{aligned}$$

where $b = K \circ \pi_p^{-1}$ for $K \in C^2(S^m)$.

Assume to the contrary that the image of $C^2(S^m) \times \{0\} \times \{0\}$ by $\nabla \mathcal{F}(H, \lambda, \xi)$ is a proper subspace of $\mathbb{R}^{m+1} = \mathbb{R} \times \mathbb{R}^m$. Then there exists a non-zero vector $(\lambda_0, \xi_0) \in \mathbb{R} \times \mathbb{R}^m$ with $\xi_0 = (\xi_1, \dots, \xi_m) \in \mathbb{R}^m$ which annihilates the image. That is,

$$0 = \nabla \mathcal{F}(H, \lambda, \xi)[K, 0, 0] \cdot (\lambda_0, \xi_0) = m^m \int_{\mathbb{R}^m} \frac{\nabla b(\lambda x + \xi) \cdot (\lambda_0 x + \xi_0)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} \quad (3.9)$$

for arbitrary $K \in C^2(S^m)$ and $b = K \circ \pi_p^{-1}$.

For $1 \leq i \leq m + 1$, let $\beta_i \in C^\infty(S^m)$ be a smooth function obtained by restricting the i -th coordinate function of \mathbb{R}^{m+1} on S^m , i.e., $\beta_i(x) = x_i|_{S^m}$ where $x = (x_1, \dots, x_{m+1})$. We define $K_i \in C^\infty(S^m)$ by $K_i(x) = \beta_i(\pi_p^{-1}(\frac{\pi_p(x) - \xi}{\lambda}))$. In other words, K_i is the i -th component of the conformal diffeomorphism of S^m which is $\mathbb{R}^m \ni x \mapsto \frac{x - \xi}{\lambda}$ in term of the identification $S^m \cong \mathbb{R}^m \cup \{\infty\}$ by the stereographic projection π_p . Then we have $b_i(x) = K_i \circ \pi_p^{-1}(x) = \beta_i(\pi_p^{-1}(\frac{x - \xi}{\lambda}))$ for $x \in \mathbb{R}^m$.

Now, we substitute K_i and b_i into (3.9). Since $\nabla b_i(x) = \frac{1}{\lambda} \nabla(\beta_i \circ \pi_p^{-1})(\frac{x - \xi}{\lambda})$ and $\nabla b_i(\lambda x + \xi) = \frac{1}{\lambda} \nabla(\beta_i \circ \pi_p^{-1})(x)$, we have

$$\int_{\mathbb{R}^m} \frac{\nabla(\beta_i \circ \pi_p^{-1})(x) \cdot (\lambda_0 x + \xi_0)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}} = 0 \quad (3.10)$$

for $i = 1, \dots, m+1$.

Note that $\beta_i \circ \pi_p^{-1}(x) = \frac{2x_i}{1+|x|^2}$ for $i = 1, \dots, m$ and $\beta_{m+1} \circ \pi_p^{-1}(x) = \frac{|x|^2-1}{1+|x|^2}$ (here, without loss of generality, we assume p is the north pole). We first take $i = m+1$. Since $\nabla(\beta_{m+1} \circ \pi_p^{-1})(x) = \frac{4x}{(1+|x|^2)^2}$, (3.10) gives

$$0 = \int_{\mathbb{R}^m} \frac{4x \cdot (\lambda_0 x + \xi_0)}{(1+|x|^2)^{m+2}} d \text{vol}_{g_{\mathbb{R}^m}} = \lambda_0 \int_{\mathbb{R}^m} \frac{4|x|^2}{(1+|x|^2)^{m+2}} d \text{vol}_{g_{\mathbb{R}^m}},$$

and hence $\lambda_0 = 0$.

For $1 \leq i \leq m$, we have $\partial_j(\beta_i \circ \pi_p^{-1})(x) = \frac{2\delta_{ij}}{1+|x|^2} - \frac{4x_i x_j}{(1+|x|^2)^2}$. Thus, it follows from (3.10) and $\lambda_0 = 0$ that

$$\begin{aligned} 0 &= \int_{\mathbb{R}^m} \frac{1}{(1+|x|^2)^m} \left[\left(\frac{2}{1+|x|^2} - \frac{4x_i^2}{(1+|x|^2)^2} \right) \xi_i - \sum_{j \neq i} \frac{4x_i x_j \xi_j}{(1+|x|^2)^2} \right] d \text{vol}_{g_{\mathbb{R}^m}} \\ &= \xi_i \int_{\mathbb{R}^m} \frac{2(1+|x|^2 - 2x_i^2)}{(1+|x|^2)^{m+2}} d \text{vol}_{g_{\mathbb{R}^m}}. \end{aligned}$$

Since

$$\int_{\mathbb{R}^m} \frac{x_i^2}{(1+|x|^2)^{m+2}} d \text{vol}_{g_{\mathbb{R}^m}} = \frac{1}{m} \int_{\mathbb{R}^m} \frac{|x|^2}{(1+|x|^2)^{m+2}} d \text{vol}_{g_{\mathbb{R}^m}} < \frac{1}{2} \int_{\mathbb{R}^m} \frac{1}{(1+|x|^2)^{m+1}} d \text{vol}_{g_{\mathbb{R}^m}},$$

we have $\xi_i = 0$ for $i = 1, \dots, m$. Therefore we have $(\lambda_0, \xi_0) = 0$ which is a contradiction. Thus the restriction of $\nabla \mathcal{F}$ to the subspace $C^2(S^m) \times \{0\} \times \{0\}$ is onto. \square

Since the kernel of $\nabla \mathcal{F}(H, \lambda, \xi)$ for $(H, \lambda, \xi) \in \mathcal{C}$ has a complemented subspace (since its codimension is finite), by the inverse function theorem, $\mathcal{C} \subset C^2(S^m) \times (0, \infty) \times \mathbb{R}^m$ is a C^1 -submanifold.

Let $\mathcal{P} : \mathcal{C} \rightarrow C^2(S^m)$ be the projection, i.e., $\mathcal{P}(H, \lambda, \xi) = H$. We next prove:

Lemma 3.8. $\mathcal{P} : \mathcal{C} \rightarrow C^2(S^m)$ is a Fredholm map with index 0.

Proof. We first note that the tangent space of \mathcal{C} at $(H, \lambda, \xi) \in \mathcal{C}$ is

$$T_{(H, \lambda, \xi)} \mathcal{C} = \{(K, \mu, \zeta) \in C^2(S^m) \times (0, \infty) \times \mathbb{R}^m : \nabla \mathcal{F}(H, \lambda, \xi)[K, \mu, \zeta] = 0\}.$$

Since $\nabla \mathcal{P}(H, \lambda, \xi)[K, \mu, \zeta] = K$ for $(K, \mu, \zeta) \in T_{(H, \lambda, \xi)} \mathcal{C}$, we see that $(K, \mu, \zeta) \in \ker \nabla \mathcal{P}(H, \lambda, \xi)$ if and only if $K = 0$, and thus

$$\ker \nabla \mathcal{P}(H, \lambda, \xi) \cong \ker \nabla \mathcal{F}(H, \lambda, \xi)[0, \cdot, \cdot] \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m}.$$

On the other hand, let us define a map

$$\begin{aligned} \mathcal{Z} : C^2(S^m) &\longrightarrow \mathbb{R}^{m+1} / \text{im } \nabla \mathcal{F}(H, \lambda, \xi)[0, \cdot, \cdot] \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m} \\ K &\longmapsto [\mathcal{F}(K, \lambda, \xi)] \end{aligned}$$

where $[v]$ stands for the equivalence class of a vector $v \in \mathbb{R}^{m+1}$ in the quotient space. Then, from the fact

$$\nabla \mathcal{F}(H, \lambda, \xi)[K, \mu, \zeta] = \mathcal{F}(K, \lambda, \xi) + \nabla \mathcal{F}(H, \lambda, \xi)[0, \mu, \zeta]$$

for all $(K, \mu, \zeta) \in C^2(S^m) \times \mathbb{R} \times \mathbb{R}^m$, it follows that $\ker \mathcal{Z} = \text{im } \nabla \mathcal{P}(H, \lambda, \xi)$. To proceed, let us claim that \mathcal{Z} is surjective. In fact, since $\nabla \mathcal{F}(H, \lambda, \xi)$ is surjective by Lemma 3.7, we see that for any vector $v \in \mathbb{R}^{m+1}$ there exists $(K, \mu, \zeta) \in C^2(S^m) \times \mathbb{R} \times \mathbb{R}^m$ such that

$$v = \nabla \mathcal{F}(H, \lambda, \xi)[K, \mu, \zeta] = \mathcal{F}(K, \lambda, \xi) + \nabla \mathcal{F}(H, \lambda, \xi)[0, \mu, \zeta].$$

This implies $[v] = [\mathcal{F}(K, \lambda, \xi)]$ in the quotient space $\mathbb{R}^{m+1} / \text{im } \nabla \mathcal{F}(H, \lambda, \xi)[0, \cdot, \cdot] \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m}$. Therefore, the map \mathcal{Z} induces an isomorphism

$$\text{coker } \nabla \mathcal{P}(H, \lambda, \xi) = \frac{C^2(S^m)}{\text{im } \nabla \mathcal{P}(H, \lambda, \xi)} = \frac{C^2(S^m)}{\ker \mathcal{Z}} \cong \frac{\mathbb{R}^{m+1}}{\text{im } \nabla \mathcal{F}(H, \lambda, \xi)[0, \cdot, \cdot] \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m}}.$$

Hence the map \mathcal{P} is Fredholm with $\text{ind } \nabla \mathcal{P}(H, \lambda, \xi) = \text{ind } \nabla \mathcal{F}(H, \lambda, \xi) \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m} = 0$. \square

Proof of Proposition 3.6. By Lemma 3.8, $\mathcal{P} : \mathcal{C} \rightarrow C^2(S^m)$ is a C^1 -Fredholm map between separable C^1 -Banach manifold with index 0. By the Sard-Smale theorem, the set of regular values of \mathcal{P} is residual in $C^2(S^m)$. For any regular value $H \in C^2(S^m)$, the proof of Lemma 3.8 shows that

$$0 = \dim \text{coker } \nabla \mathcal{P}(H, \lambda, \xi) = \dim \frac{\mathbb{R}^{m+1}}{\text{im } \nabla \mathcal{F}(H, \lambda, \xi)[0, \cdot, \cdot] \Big|_{\{0\} \times \mathbb{R} \times \mathbb{R}^m}}.$$

By noting that

$$\nabla \mathcal{F}(H, \lambda, \xi)[0, \mu, \zeta] = \nabla^2 \Psi(\lambda, \xi; H)[(\mu, \zeta)], \quad \text{for } (\mu, \zeta) \in \mathbb{R} \times \mathbb{R}^m = \mathbb{R}^{m+1}$$

we find the Hessian $\nabla^2 \Psi(\lambda, \xi; H)$ is non-degenerate. This means that $\Psi(\cdot, \cdot; H)$ is a Morse function on $(0, \infty) \times \mathbb{R}^m$. This completes the proof. \square

Remark 3.9. Let $H \in C^2(S^m)$ be an arbitrary function satisfying conditions (H-1) and (H-2), then there exists $\delta > 0$ such that $\tilde{H} \in C^2$ with $\|H - \tilde{H}\|_{C^2(S^m)} < \delta$ satisfies the same conditions. Proposition 3.6 implies that for any given $H \in C^2(S^m)$ satisfying conditions (H-1) and (H-2), there exists $\tilde{H} \in C^2(S^m)$, close to H in C^2 -norm, which satisfies the same conditions and the function

$$(\lambda, \xi) \mapsto m^m \int_{\mathbb{R}^m} \frac{\tilde{a}(\lambda x + \xi)}{(1 + |x|^2)^m} d \text{vol}_{g_{\mathbb{R}^m}}$$

is a Morse function, where $\tilde{a}(x) := \tilde{H} \circ \pi_p^{-1}(x)$. Thus our claim “ Ψ is Morse for generic choice of a ” is verified.

4 Perturbations of the background metric

In this section, let us consider the equation

$$D_g \psi = |\psi|_g^{2^* - 2} \psi \quad \text{on } S^m \tag{4.1}$$

involving a perturbed metric, where $2^* = \frac{2m}{m-1}$. Precisely, by using the stereographic projection $\pi_p : S^m \setminus \{p\} \rightarrow \mathbb{R}^m$ (for a fixed $p \in S^m$), the metric g is defined via (1.9) and (1.10). And in this setting, (4.1) is equivalent to the equation

$$D_{\tilde{g}} \psi = |\psi|_{\tilde{g}}^{2^* - 2} \psi \quad \text{on } \mathbb{R}^m. \tag{4.2}$$

4.1 Bourguignon-Gauduchon identification

In this part we will collect some basic results which will enable us to write the energy functional \mathcal{J} for (4.2) in a form suitable for using the perturbation method mentioned before. In order to carry this out, we need to identify Dirac operators and spinor fields on \mathbb{R}^m with respect to the canonical Euclidean metric $g_{\mathbb{R}^m}$ and the new metric \tilde{g} . The construction by Bourguignon and Gauduchon [14] provides us such a necessary identification.

To begin with, let us denote $\odot_+^2(\mathbb{R}^m)$ the space of positive definite symmetric bilinear forms, i.e. metrics on \mathbb{R}^m . Then, for small values of $\varepsilon > 0$, we consider the identity map

$$\begin{aligned} \text{Id} : (\mathbb{R}^m, g_{\mathbb{R}^m}) &\rightarrow (\mathbb{R}^m, \tilde{g}) \\ x &\mapsto x \end{aligned}$$

and the map

$$\begin{aligned} \tilde{G} : \mathbb{R}^m &\rightarrow \odot_+^2(\mathbb{R}^m) \\ x &\mapsto \tilde{G}_x := (\tilde{g}_{ij}(x))_{ij} \end{aligned}$$

which associates to a point $x \in \mathbb{R}^m$ the matrix of the coefficients of the metric \tilde{g} at this point, expressed in the basis $\partial_i = \frac{\partial}{\partial x^i}$, $i = 1, \dots, m$. Notice that $\tilde{G}_x \in \odot_+^2(\mathbb{R}^m)$, hence \tilde{G}_x is invertible and there is a unique matrix $B_x \in \odot_+^2(\mathbb{R}^m)$ such that $B_x^2 = \tilde{G}_x^{-1}$. Let $b_{ij}(x)$, $i, j = 1, \dots, m$, be the entries of B_x , we have

$$\begin{aligned} B_x : (T_x \mathbb{R}^m \cong \mathbb{R}^m, g_{\mathbb{R}^m}) &\rightarrow (T_x \mathbb{R}^m, \tilde{g}_x) \\ v = \sum_k v_k \partial_k &\mapsto B_x(v) := \sum_j \left(\sum_k b_{jk}(x) v_k \right) \partial_j \end{aligned}$$

defines an isometry for each $x \in \mathbb{R}^m$. As the matrix B_x depends smoothly on x , we obtain an isomorphism of $SO(m)$ -principal bundles:

$$\begin{array}{ccc} P_{SO}(\mathbb{R}^m, g_{\mathbb{R}^m}) & \xrightarrow{\eta} & P_{SO}(\mathbb{R}^m, \tilde{g}) \\ \downarrow & & \downarrow \\ \mathbb{R}^m & \xrightarrow{\text{Id}} & \mathbb{R}^m \end{array}$$

where $\eta\{v_1, \dots, v_m\} = \{B(v_1), \dots, B(v_m)\}$ for an oriented frame $\{v_1, \dots, v_m\}$ on $(\mathbb{R}^m, g_{\mathbb{R}^m})$. Note that the map η commutes with the right action of $SO(m)$, it can be lifted to spin structures:

$$\begin{array}{ccc} P_{Spin}(\mathbb{R}^m, g_{\mathbb{R}^m}) & \xrightarrow{\tilde{\eta}} & P_{Spin}(\mathbb{R}^m, \tilde{g}) \\ \downarrow & & \downarrow \\ \mathbb{R}^m & \xrightarrow{\text{Id}} & \mathbb{R}^m \end{array}$$

And hence we obtain an isomorphism between the spinor bundles $\mathbb{S}(\mathbb{R}^m, g_{\mathbb{R}^m})$ and $\mathbb{S}(\mathbb{R}^m, \tilde{g})$:

$$\begin{aligned} \mathbb{S}(\mathbb{R}^m, g_{\mathbb{R}^m}) &:= P_{Spin}(\mathbb{R}^m, g_{\mathbb{R}^m}) \times_{\rho} \mathbb{S}_m \longrightarrow \mathbb{S}(\mathbb{R}^m, \tilde{g}) := P_{Spin}(\mathbb{R}^m, \tilde{g}) \times_{\rho} \mathbb{S}_m \\ \psi = [s, \varphi] &\longmapsto \tilde{\psi} = [\tilde{\eta}(s), \varphi] \end{aligned} \tag{4.3}$$

where ρ is the complex spinor representation and $[s, \varphi]$ stands for the equivalence class of (s, φ) under the action of $Spin(m)$. This identifies the spinor fields.

For the Dirac operators, as was shown by [11, Proposition 3.2], the identification can be expressed in the following formula

$$D_{\tilde{g}}\tilde{\psi} = \widetilde{D_{g_{\mathbb{R}^m}}\psi} + W \cdot_{\tilde{g}} \tilde{\psi} + X \cdot_{\tilde{g}} \tilde{\psi} + \sum_{i,j} (b_{ij} - \delta_{ij}) \tilde{\partial}_i \cdot_{\tilde{g}} \widetilde{\nabla_{\partial_j}\psi} \quad (4.4)$$

where $\cdot_{\tilde{g}}$ denotes the Clifford multiplication with respect to the metric \tilde{g} ,

$$W = \frac{1}{4} \sum_{\substack{i,j,k \\ i \neq j \neq k \neq i}} \sum_{\alpha,\beta} b_{i\alpha} (\partial_\alpha b_{j\beta}) b_{\beta k}^{-1} \tilde{\partial}_i \cdot_{\tilde{g}} \tilde{\partial}_j \cdot_{\tilde{g}} \tilde{\partial}_k,$$

with b_{ij}^{-1} being the entries of the inverse matrix of B , $\tilde{\partial}_i = B(\partial_i)$ and

$$X = \frac{1}{2} \sum_{i,k} \tilde{\Gamma}_{ik}^i \tilde{\partial}_k,$$

with $\tilde{\Gamma}_{ij}^k = \tilde{g}(\tilde{\nabla}_{\tilde{\partial}_i} \tilde{\partial}_j, \tilde{\partial}_k)$ being the Christoffel symbols of the second kind.

Remark 4.1. On spin manifolds, since the tangent bundle is embedded in the bundle of Clifford algebra, vector fields have two different actions on spinors, i.e. the Clifford multiplications and the covariant derivatives. Here, to distinguish the two actions on a spinor ψ , we denote $\partial_i \cdot_{g_{\mathbb{R}^m}} \psi$ the Clifford multiplication of ∂_i and $\nabla_{\partial_i} \psi$ the covariant derivative with respect to the metric $g_{\mathbb{R}^m}$ (respectively, $\tilde{\partial}_i \cdot_{\tilde{g}} \tilde{\psi}$ the Clifford multiplication of $\tilde{\partial}_i$ and $\tilde{\nabla}_{\tilde{\partial}_i} \tilde{\psi}$ the covariant derivative with respect to the metric \tilde{g}). For functions, we shall simply denote $\partial_i u$ for its partial derivative.

4.2 Basic calculations

The following expansions come from elementary calculations.

Lemma 4.2. *Let \tilde{g} be given by (1.10), we have*

$$\sqrt{\det \tilde{G}} = 1 + \frac{\varepsilon}{2} \operatorname{tr} h + \varepsilon^2 \left(\frac{1}{8} (\operatorname{tr} h)^2 - \frac{1}{4} \operatorname{tr}(h^2) \right) + o(\varepsilon^2), \quad (4.5)$$

$$B = I - \frac{\varepsilon}{2} h + \frac{3\varepsilon^2}{8} h^2 + o(\varepsilon^2) \quad (4.6)$$

and

$$B^{-1} = I + \frac{\varepsilon}{2} h - \frac{\varepsilon^2}{8} h^2 + o(\varepsilon^2). \quad (4.7)$$

With the notation of Bourguignon-Gauduchon identification, for a spinor $\tilde{\psi}$ in $\mathbb{S}(\mathbb{R}^m, \tilde{g})$, the energy functional associated to (4.2) is defined as

$$\mathcal{J}(\tilde{\psi}) = \frac{1}{2} \int_{\mathbb{R}^m} (\tilde{\psi}, D_{\tilde{g}} \tilde{\psi})_{\tilde{g}} d \operatorname{vol}_{\tilde{g}} - \frac{1}{2^*} \int_{\mathbb{R}^m} |\tilde{\psi}|_{\tilde{g}}^{2^*} d \operatorname{vol}_{\tilde{g}}. \quad (4.8)$$

It turns out that \mathcal{J} is an ε -involved functional because \tilde{g} is given by (1.10). And now, via the Bourguignon-Gauduchon identification, we have the following expansion:

Lemma 4.3. *Let \tilde{g} be given by (1.10), then*

$$\mathcal{J}(\tilde{\psi}) = \mathcal{J}_0(\psi) + \varepsilon\Gamma(\psi) + \varepsilon^2\Phi(\psi) + o(\varepsilon^2),$$

where

$$\mathcal{J}_0(\psi) = \frac{1}{2} \int_{\mathbb{R}^m} (\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} - \frac{1}{2^*} \int_{\mathbb{R}^m} |\psi|_{g_{\mathbb{R}^m}}^{2^*} d \text{vol}_{g_{\mathbb{R}^m}},$$

$$\Gamma(\psi) = \int_{\mathbb{R}^m} \frac{\text{tr } h}{2} \left[\frac{1}{2} (\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}} - \frac{1}{2^*} |\psi|_{g_{\mathbb{R}^m}}^{2^*} \right] - \frac{1}{4} \sum_i h_{ii} \text{Re} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}}$$

and

$$\begin{aligned} \Phi(\psi) &= \int_{\mathbb{R}^m} \left(\frac{1}{8} (\text{tr } h)^2 - \frac{1}{4} \text{tr}(h^2) \right) \left[\frac{1}{2} (\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}} - \frac{1}{2^*} |\psi|_{g_{\mathbb{R}^m}}^{2^*} \right] \\ &\quad + \frac{1}{16} \sum_i \left(3h_{ii}^2 - 2(\text{tr } h)h_{ii} \right) \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

for $\psi \in \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$.

Proof. We first expand the quadratic part of \mathcal{J} in terms of ε . Using (4.4) and the fact $X \in T\mathbb{R}^m$, we get

$$\text{Re}(\tilde{\psi}, D_{\tilde{g}} \tilde{\psi})_{\tilde{g}} = \text{Re}(\tilde{\psi}, \widetilde{D_{g_{\mathbb{R}^m}} \psi})_{\tilde{g}} + \text{Re}(W \cdot_{\tilde{g}} \tilde{\psi}, \tilde{\psi})_{\tilde{g}} + \sum_{i,j} (b_{ij} - \delta_{ij}) \text{Re}(\tilde{\partial}_i \cdot_{\tilde{g}} \widetilde{\nabla_{\partial_j} \psi}, \tilde{\psi})_{\tilde{g}}.$$

Since the map $\psi \mapsto \tilde{\psi}$ defined in (4.3) is fiberwisely isometric, we obtain

$$\text{Re}(\tilde{\psi}, \widetilde{D_{g_{\mathbb{R}^m}} \psi})_{\tilde{g}} = \text{Re}(\psi, D_{g_{\mathbb{R}^m}} \psi)_{g_{\mathbb{R}^m}}, \quad \text{Re}(\tilde{\partial}_i \cdot_{\tilde{g}} \widetilde{\nabla_{\partial_j} \psi}, \tilde{\psi})_{\tilde{g}} = \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_j} \psi, \psi)_{g_{\mathbb{R}^m}}$$

and

$$\text{Re}(W \cdot_{\tilde{g}} \tilde{\psi}, \tilde{\psi})_{\tilde{g}} = \frac{1}{4} \sum_{\substack{i,j,k \\ i \neq j \neq k \neq i}} \left(\sum_{\alpha,\beta} b_{i\alpha} (\partial_\alpha b_{j\beta}) b_{\beta k}^{-1} \right) \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \partial_j \cdot_{g_{\mathbb{R}^m}} \partial_k \cdot_{g_{\mathbb{R}^m}} \psi, \psi)_{g_{\mathbb{R}^m}}.$$

Plainly, $\text{Re}(W \cdot_{\tilde{g}} \tilde{\psi}, \tilde{\psi})_{\tilde{g}} \equiv 0$ for $m = 2$. For $m \geq 3$, by (4.6) and (4.7), we see that

$$\begin{cases} b_{i\alpha} = \delta_{i\alpha} - \frac{\varepsilon}{2} h_{i\alpha} + \frac{3\varepsilon^2}{8} \sum_l h_{il} h_{l\alpha} + o(\varepsilon^2), \\ \partial_\alpha b_{j\beta} = -\frac{\varepsilon}{2} \partial_\alpha h_{j\beta} + \frac{3\varepsilon^2}{8} \sum_l \left(\partial_\alpha h_{jl} h_{l\beta} + h_{jl} \partial_\alpha h_{l\beta} \right) + o(\varepsilon^2), \\ b_{\beta k}^{-1} = \delta_{\beta k} + \frac{\varepsilon}{2} h_{\beta k} - \frac{\varepsilon^2}{8} \sum_l h_{\beta l} h_{lk} + o(\varepsilon^2). \end{cases}$$

Hence

$$\begin{aligned} b_{i\alpha} (\partial_\alpha b_{j\beta}) b_{\beta k}^{-1} &= -\frac{\varepsilon}{2} \delta_{\beta k} \delta_{i\alpha} \partial_\alpha h_{j\beta} + \frac{\varepsilon^2}{4} \partial_\alpha h_{j\beta} (\delta_{\beta k} h_{i\alpha} - \delta_{i\alpha} h_{\beta k}) \\ &\quad + \frac{3\delta_{\beta k} \delta_{i\alpha} \varepsilon^2}{8} \sum_l \left(\partial_\alpha h_{jl} h_{l\beta} + h_{jl} \partial_\alpha h_{l\beta} \right) + o(\varepsilon^2). \end{aligned}$$

Note that we have assumed $h_{ij} = 0$ for $i \neq j$, we soon get

$$b_{i\alpha}(\partial_\alpha b_{j\beta})b_{\beta k}^{-1} = o(\varepsilon^2).$$

And thus

$$\begin{aligned} \operatorname{Re}(\tilde{\psi}, D_{\tilde{g}}\tilde{\psi})_{\tilde{g}} &= \operatorname{Re}(\psi, D_{g_{\mathbb{R}^m}}\psi)_{g_{\mathbb{R}^m}} - \frac{\varepsilon}{2} \sum_i h_{ii} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} \\ &\quad + \frac{3\varepsilon^2}{8} \sum_i h_{ii}^2 \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} + o(\varepsilon^2) \end{aligned} \quad (4.9)$$

Since $|\tilde{\psi}|_{\tilde{g}} = |\psi|_{g_{\mathbb{R}^m}}$, it follows from (4.5) and (4.9) that

$$\begin{aligned} \mathcal{J}(\tilde{\psi}) &= \mathcal{J}_0(\psi) + \varepsilon \int_{\mathbb{R}^m} \frac{\operatorname{tr} h}{2} \left[\frac{1}{2} (\psi, D_{g_{\mathbb{R}^m}}\psi)_{g_{\mathbb{R}^m}} - \frac{1}{2^*} |\psi|_{g_{\mathbb{R}^m}}^{2^*} \right] \\ &\quad - \frac{1}{4} \sum_i h_{ii} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi) d\operatorname{vol}_{g_{\mathbb{R}^m}} \\ &\quad + \varepsilon^2 \int_{\mathbb{R}^m} \left(\frac{1}{8} (\operatorname{tr} h)^2 - \frac{1}{4} \operatorname{tr}(h^2) \right) \left[\frac{1}{2} (\psi, D_{g_{\mathbb{R}^m}}\psi)_{g_{\mathbb{R}^m}} - \frac{1}{2^*} |\psi|_{g_{\mathbb{R}^m}}^{2^*} \right] \\ &\quad - \frac{\operatorname{tr} h}{8} \sum_i h_{ii} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} \\ &\quad + \frac{3}{16} \sum_i h_{ii}^2 \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi, \psi)_{g_{\mathbb{R}^m}} d\operatorname{vol}_{g_{\mathbb{R}^m}} + o(\varepsilon^2) \end{aligned}$$

which completes the proof. \square

Recall the critical manifold $\mathcal{M} \subset \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ defined in (2.9), an immediate consequence of the above lemma is

Corollary 4.4. For $\psi_{\lambda, \xi, \gamma} \in \mathcal{M}$ with $\lambda > 0$, $\xi \in \mathbb{R}^m$ and $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, we have

$$\Gamma(\psi_{\lambda, \xi, \gamma}) = 0$$

and

$$\Phi(\psi_{\lambda, \xi, \gamma}) = \frac{m^{m-1} \lambda^m}{16} \int_{\mathbb{R}^m} \frac{\operatorname{tr}(h^2) - (\operatorname{tr} h)^2}{(\lambda^2 + |x - \xi|^2)^m} d\operatorname{vol}_{g_{\mathbb{R}^m}}.$$

Proof. For $\psi_{\lambda, \xi, \gamma} \in \mathcal{M}$, one easily checks that it solves the equation

$$D_{g_{\mathbb{R}^m}} \psi_{\lambda, \xi, \gamma} = |\psi_{\lambda, \xi, \gamma}|_{g_{\mathbb{R}^m}}^{2^* - 2} \psi_{\lambda, \xi, \gamma}$$

on \mathbb{R}^m . Furthermore, by the explicit expression of $\psi_{\lambda, \xi, \gamma}$, we have

$$|\psi_{\lambda, \xi, \gamma}|_{g_{\mathbb{R}^m}} = \frac{m^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m-1}{2}}}$$

and

$$\begin{aligned} \nabla_{\partial_i} \psi_{\lambda, \xi, \gamma}(x) &= -\frac{m^{\frac{m+1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}+1}} (x_i - \xi_i) (\lambda - (x - \xi)) \cdot_{g_{\mathbb{R}^m}} \gamma \\ &\quad + \frac{m^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}}} (-\partial_i) \cdot_{g_{\mathbb{R}^m}} \gamma. \end{aligned} \quad (4.10)$$

Hence, by the rules of Clifford multiplication, we get $\partial_i \cdot_{g_{\mathbb{R}^m}} \partial_i = -1$ and

$$\begin{aligned} \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{\lambda, \xi, \gamma}(x) &= -\frac{m^{\frac{m+1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}+1}} (x_i - \xi_i) \partial_i \cdot_{g_{\mathbb{R}^m}} (\lambda - (x - \xi)) \cdot_{g_{\mathbb{R}^m}} \gamma \\ &\quad + \frac{m^{\frac{m-1}{2}} \lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}}} \gamma \end{aligned} \quad (4.11)$$

and

$$\operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{\lambda, \xi, \gamma}, \psi_{\lambda, \xi, \gamma})_{g_{\mathbb{R}^m}} = \frac{m^{m-1} \lambda^m}{(\lambda^2 + |x - \xi|^2)^m} \quad \text{for } i = 1, \dots, m.$$

Therefore,

$$\Gamma(\psi_{\lambda, \xi, \gamma}) = \int_{\mathbb{R}^m} \left(\frac{\operatorname{tr} h}{2} \frac{1}{2m} \frac{m^m \lambda^m}{(\lambda^2 + |x - \xi|^2)^m} - \frac{\operatorname{tr} h}{4} \frac{m^{m-1} \lambda^m}{(\lambda^2 + |x - \xi|^2)^m} \right) d \operatorname{vol}_{g_{\mathbb{R}^m}} \equiv 0$$

and

$$\Phi(\psi_{\lambda, \xi, \gamma}) = \frac{m^{m-1} \lambda^m}{16} \int_{\mathbb{R}^m} \frac{\operatorname{tr}(h^2) - (\operatorname{tr} h)^2}{(\lambda^2 + |x - \xi|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}.$$

□

By Lemma 4.3 and Corollary 4.4, we find Theorem 2.6 is capable of showing the existence issues here. Hence, in what follows, we will investigate the behavior of

$$\hat{\Phi}(z) := \Phi(z) - \frac{1}{2} \langle K_z(\nabla \Gamma(z)), \nabla \Gamma(z) \rangle$$

on a fiber $\vartheta^{-1}(\gamma)$ for a fixed $\gamma \in \mathcal{N} = S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, where K_z stands for the inverse of $\nabla^2 \mathcal{J}_0(z)$ restricted to $\mathcal{W}_z := T_z \mathcal{M}^\perp \subset \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$. Note that $\vartheta^{-1}(\gamma)$ is parameterized by $\mathcal{G} = (0, \infty) \times \mathbb{R}^m$, it is very natural to study the values of $\hat{\Phi}(\psi_{\lambda, \xi, \gamma})$ when $\lambda \rightarrow 0$ and $\lambda + |\xi| \rightarrow \infty$.

Lemma 4.5. *Assume that we are in the hypotheses of Lemma 4.3. There holds*

$$\lim_{\lambda \rightarrow 0} \Phi(\psi_{\lambda, \xi, \gamma}) = C_0 (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi) \quad \text{for any } \xi \in \mathbb{R}^m,$$

where

$$C_0 = \frac{m^{m-1}}{16} \int_{\mathbb{R}^m} \frac{1}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}.$$

Proof. By change of variables, one easily deduce that

$$\Phi(\psi_{\lambda,\xi,\gamma}) = \frac{m^{m-1}}{16} \int_{\mathbb{R}^m} \frac{(\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\lambda x + \xi)}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}.$$

And hence we have

$$\lim_{\lambda \rightarrow 0} \Phi(\psi_{\lambda,\xi,\gamma}) = \frac{m^{m-1}}{16} \int_{\mathbb{R}^m} \frac{1}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi).$$

□

The next two results, i.e. Lemma 4.6 and Proposition 4.7, are the main ingredients needed in our argument. For the sake of clarity, since the proofs of these two results contain the main difficulties of the paper, they are postponed to Appendix A.3.

Lemma 4.6. *Assume that we are in the hypotheses of Lemma 4.3. There holds*

$$\lim_{\lambda \rightarrow 0} \langle K_{\psi_{\lambda,\xi,\gamma}}(\nabla \Gamma(\psi_{\lambda,\xi,\gamma})), \nabla \Gamma(\psi_{\lambda,\xi,\gamma}) \rangle = C_1 (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi) \quad \text{for any } \xi \in \mathbb{R}^m,$$

where

$$C_1 = \frac{m^{m-1}}{4} \int_{\mathbb{R}^m} \frac{|x|^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}}.$$

It follows from Lemma 4.5 and 4.6 that

$$\lim_{\lambda \rightarrow 0} \hat{\Phi}(\psi_{\lambda,\xi,\gamma}) = \left(C_0 - \frac{1}{2} C_1 \right) (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi) = 0.$$

since $C_0 = \frac{1}{2} C_1$. With this observation in mind, to see whether $\hat{\Phi}$ has local maximum or minimum on the fiber $\vartheta^{-1}(\gamma)$, we have to go further to study the asymptotic behavior of $\hat{\Phi}(\psi_{\lambda,\xi,\gamma})$ as $\lambda \rightarrow 0$. By using the concept of elementary matrix in Definition 1.3, the following proposition characterizes the exact expansion of $\hat{\Phi}(\psi_{\lambda,\xi,\gamma})$ with respect to λ small.

Proposition 4.7. *For $m \geq 4$, assume that we are in the hypotheses of Lemma 4.3. Let $k \in \{1, \dots, m\}$, $p \in [2, \infty]$ and $h = \operatorname{diag}(h_{11}, \dots, h_{mm})$ be (k, p) -elementary at a point $\xi \in \mathbb{R}^m$ with $\partial_k h_{ii}(\xi) \equiv c_k \neq 0$, for $i \neq k$. If*

$$\begin{cases} p = \infty & m = 4, \\ p > 2 & m = 5, \\ p \geq 2 & m \geq 6, \end{cases}$$

then

$$\hat{\Phi}(\psi_{\lambda,\xi,\gamma}) = -\frac{3m^{m-2}(m-1)(m-2)c_k^2}{128} \lambda^2 \int_{\mathbb{R}^m} \frac{|x|^2}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} + o(\lambda^2) \quad \text{as } \lambda \rightarrow 0.$$

In particular, $\hat{\Phi}(\psi_{\lambda,\xi,\gamma}) < 0$ for small values of λ .

The next proposition shows the decay of $\hat{\Phi}(\psi_{\lambda,\xi,\gamma})$ as $\lambda + |\xi| \rightarrow \infty$.

Proposition 4.8. *Assume that we are in the hypotheses of Lemma 4.3. Then*

$$\hat{\Phi}(\psi_{\lambda,\xi,\gamma}) \rightarrow 0 \quad \text{as } \lambda + |\xi| \rightarrow \infty.$$

Proof. Let $r : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}^m \setminus \{0\}$ be defined as $r(x) = -\frac{x}{|x|^2}$. Then, for each $(\lambda, \xi, \gamma) \in (0, \infty) \times \mathbb{R}^m \times S^{2i\frac{m}{2}+1-1}(\mathbb{S}_m)$, we can define $\psi_{\lambda,\xi,\gamma}^\sharp(x) = \psi_{\lambda^\sharp,\xi^\sharp,\gamma}(x)$, where $\lambda^\sharp = \frac{\lambda}{\lambda^2+|\xi|^2}$ and $\xi^\sharp = \frac{-\xi}{\lambda^2+|\xi|^2}$. Then $\psi_{\lambda,\xi,\gamma}^\sharp \in \mathcal{M}$ and it follows from a direct computation that that

$$|\psi_{\lambda,\xi,\gamma}^\sharp(x)|_{g_{\mathbb{R}^m}} = |\psi_{\lambda^\sharp,\xi^\sharp,\gamma}(x)|_{g_{\mathbb{R}^m}} = \frac{1}{|x|^{m-1}} |\psi_{\lambda,\xi,\gamma}(r(x))|_{g_{\mathbb{R}^m}}.$$

Roughly speaking, the map $\psi_{\lambda,\xi,\gamma} \mapsto \psi_{\lambda,\xi,\gamma}^\sharp$ can be interpreted as an one-to-one correspondence in \mathcal{M} between the elements at infinity and the elements close to $\psi_{t,0,\gamma}$ with small $t > 0$.

For an arbitrary function $P : \mathbb{R}^m \rightarrow \mathbb{R}$, let us define $P^\sharp(x) = P(r(x))$. By observing that r is conformal with conformal factor $\frac{1}{|x|^4}$, i.e. $r^*g_{\mathbb{R}^m} = \frac{1}{|x|^4}g_{\mathbb{R}^m}$, we find

$$\begin{aligned} \int_{\mathbb{R}^m} P(x) |\psi_{\lambda,\xi,\gamma}(x)|_{g_{\mathbb{R}^m}}^{2^*} d \text{vol}_{g_{\mathbb{R}^m}} &= \int_{\mathbb{R}^m} P(r(x)) |\psi_{\lambda,\xi,\gamma}(r(x))|_{g_{\mathbb{R}^m}}^{2^*} \frac{1}{|x|^{2m}} d \text{vol}_{g_{\mathbb{R}^m}} \\ &= \int_{\mathbb{R}^m} P^\sharp(x) |\psi_{\lambda,\xi,\gamma}^\sharp(x)|_{g_{\mathbb{R}^m}}^{2^*} d \text{vol}_{g_{\mathbb{R}^m}}. \end{aligned} \quad (4.12)$$

Moreover, by notice that

$$\text{Re} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{\lambda,\xi,\gamma}^\sharp(x), \psi_{\lambda,\xi,\gamma}^\sharp(x))_{g_{\mathbb{R}^m}} = \frac{1}{m} |\psi_{\lambda,\xi,\gamma}^\sharp(x)|_{g_{\mathbb{R}^m}}^{2^*}$$

we have

$$\begin{aligned} \text{Re} \int_{\mathbb{R}^m} P(x) (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{\lambda,\xi,\gamma}, \psi_{\lambda,\xi,\gamma})_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \\ = \text{Re} \int_{\mathbb{R}^m} P^\sharp(x) (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{\lambda,\xi,\gamma}^\sharp, \psi_{\lambda,\xi,\gamma}^\sharp)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \end{aligned} \quad (4.13)$$

for $i = 1, \dots, m$.

Now let $\tilde{g}^\sharp(x) := \tilde{g}(r(x))$ and consider the corresponding new perturbed functional \mathcal{J}^\sharp obtained by substituting in (4.8) \tilde{g} with \tilde{g}^\sharp . Analogous to Lemma 4.3, we have an expansion

$$\mathcal{J}^\sharp(\tilde{\psi}) = \mathcal{J}_0(\psi) + \varepsilon \Gamma^\sharp(\psi) + \varepsilon^2 \Phi^\sharp(\psi) + o(\varepsilon^2).$$

And by (4.12) and (4.13), we infer that

$$\Gamma^\sharp(\psi_{\lambda,\xi,\gamma}^\sharp) = \Gamma(\psi_{\lambda,\xi,\gamma}) \quad \text{and} \quad \Phi^\sharp(\psi_{\lambda,\xi,\gamma}^\sharp) = \Phi(\psi_{\lambda,\xi,\gamma}).$$

Furthermore, we also have

$$\hat{\Phi}^\sharp(\psi_{\lambda,\xi,\gamma}^\sharp) = \hat{\Phi}(\psi_{\lambda,\xi,\gamma}).$$

Hence, by applying Lemma 4.5 and 4.6, we have

$$\lim_{\lambda+|\xi| \rightarrow \infty} \hat{\Phi}(\psi_{\lambda,\xi,\gamma}) = \lim_{\lambda+|\xi| \rightarrow \infty} \hat{\Phi}^\sharp(\psi_{\lambda,\xi,\gamma}^\sharp) = \lim_{\substack{\lambda^\sharp \rightarrow 0 \\ \xi^\sharp \rightarrow 0}} \hat{\Phi}^\sharp(\psi_{\lambda^\sharp,\xi^\sharp,\gamma}) = 0,$$

which completes the proof. \square

Remark 4.9. From Proposition 4.7, 4.8 and Theorem 2.6 we can immediately conclude that: there exists a bounded open set $U \subset (0, \infty) \times \mathbb{R}^m$, independent of $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, such that $\hat{\Phi}(\psi_{\lambda, \xi, \gamma})$ achieves a strict minimum in dimension $m \geq 4$ in U , and hence the perturbed spinorial Yamabe equation (4.2) has a solution.

4.3 Proof the main result

The proof will be carried out via several steps, we begin with some basic estimates.

Lemma 4.10. *The operator $K_z = \nabla^2 \mathcal{J}_0(z)^{-1}$ is uniformly bounded for $z \in \mathcal{M}$.*

Proof. For a fixed $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, we denote $z_0 = \psi_{1,0,\gamma}$. For arbitrary $g = (\lambda, \xi) \in (0, \infty) \times \mathbb{R}^m$, we define a right action on elements in $\mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ as

$$(\psi \cdot g)(x) = \lambda^{-\frac{m-1}{2}} \psi\left(\frac{x - \xi}{\lambda}\right) \quad \text{for } \psi \in \mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m)).$$

Then it is easy to check that $z_0 \cdot g = \psi_{\lambda, \xi, \gamma}$ and the inverse action is defined by $g^{-1} = (\lambda^{-1}, -\lambda^{-1}\xi)$. Moreover, by standard Fourier transformation, we can see that the action of g is an isometry, i.e. $\|\psi \cdot g\| = \|\psi\|$ for any $\psi \in \mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ and $g \in (0, \infty) \times \mathbb{R}^m$.

Observe that $\nabla^2 \mathcal{J}_0(z_0)$ is self-adjoint and invertible on $\mathcal{W}_{z_0} = T_{z_0} \mathcal{M}^\perp$, there exists a constant $c_\gamma > 0$ such that for any $w \in \mathcal{W}_{z_0}$ there is $v(w) \in \mathcal{W}_{z_0}$ satisfying $\|v(w)\| = 1$ and

$$\nabla^2 \mathcal{J}_0(z_0)[w, v(w)] = \|\nabla^2 \mathcal{J}_0(z_0)[w]\| \geq c_\gamma \|w\|.$$

Here the subscript in the constant c_γ comes from the fact that z_0 is (implicitly) depending on γ . In what follows, for arbitrary $w \in \mathcal{W}_{z_0}$, we simply denote $w_g = w \cdot g$ for $g = (\lambda, \xi) \in (0, \infty) \times \mathbb{R}^m$ and $z_g := z_0 \cdot g$. We mention that, by change of variables,

$$\begin{aligned} \operatorname{Re} \int_{\mathbb{R}^m} (D_{g_{\mathbb{R}^m}} w_g, v(w)_g)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} &= \operatorname{Re} \int_{\mathbb{R}^m} (D_{g_{\mathbb{R}^m}} w, v(w))_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}}, \\ \operatorname{Re} \int_{\mathbb{R}^m} |z_g|^{\frac{2}{m-1}} (w_g, v(w)_g)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} &= \operatorname{Re} \int_{\mathbb{R}^m} |z_0|^{\frac{2}{m-1}} (w, v(w))_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}^m} |z_g|^{\frac{4-2m}{m-1}} \operatorname{Re}(z_g, w_g)_{g_{\mathbb{R}^m}} \operatorname{Re}(z_g, v(w)_g)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\ = \operatorname{Re} \int_{\mathbb{R}^m} |z_0|^{\frac{4-2m}{m-1}} \operatorname{Re}(z_0, w)_{g_{\mathbb{R}^m}} \operatorname{Re}(z_0, v(w))_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

Hence one deduces

$$\nabla^2 \mathcal{J}_0(z_g)[w, v(w_{g^{-1}})_g] = \nabla^2 \mathcal{J}_0(z_0)[w_{g^{-1}}, v(w_{g^{-1}})] \geq c_\gamma \|w_{g^{-1}}\| = c_\gamma \|w\|$$

Therefore, we see that the operator norm of $\|K_{z_\rho}\|$ is uniformly bounded by $\frac{1}{c_\gamma}$ for all $g \in (0, \infty) \times \mathbb{R}^m$ since $K_{z_g} = \nabla^2 \mathcal{J}_0(z_g)^{-1}$. Note that $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$, where $S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$ is a compact manifold, we have c_γ is bounded away from 0 uniformly in γ . And thanks to the continuity of K_{z_0} with respect to γ , we conclude that K_z is uniformly bounded for all $z \in \mathcal{M}$ and this completes the proof. \square

Lemma 4.11. *Let \tilde{g} be given by (1.10). Assume that the matrix function h is compactly supported. Then*

$$\|\nabla\Gamma(\psi_{\lambda,\xi,\gamma})\| \rightarrow 0 \quad \text{as } \lambda \rightarrow \infty \text{ uniformly in } (\xi, \gamma).$$

Proof. Let $R > 0$ be large enough such that $\text{supp } h \subset B_R$, where B_R is the open ball in \mathbb{R}^m of radius R centered at the origin. For any $v \in \mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$, by (A.12), we have

$$\begin{aligned} |\langle \nabla\Gamma(\psi_{\lambda,\xi,\gamma}), v \rangle| &\leq C \left(\int_{B_R} |\nabla\psi_{\lambda,\xi,\gamma}| \cdot |v| + |\psi_{\lambda,\xi,\gamma}| \cdot |v| d \text{vol}_{g_{\mathbb{R}^m}} \right) \\ &\leq C \left(\int_{B_R} \frac{\lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}}} \cdot |v| + \frac{\lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m-1}{2}}} \cdot |v| d \text{vol}_{g_{\mathbb{R}^m}} \right) \\ &\leq C(\lambda^{-\frac{m+1}{2}} + \lambda^{-\frac{m-1}{2}}) \|v\|. \end{aligned}$$

Thus, $\|\nabla\Gamma(\psi_{\lambda,\xi,\gamma})\| \rightarrow 0$ as $\lambda \rightarrow \infty$ uniformly in (ξ, γ) . \square

Next, let us consider a further situation of the perturbed metric in (1.10). Precisely, we consider the matrix h in the form of (1.11). In this case, the matrix h can be decomposed as $h(x) = h^{(1)}(x) + h^{(2)}(x - x_0)$, and it is clear that $h^{(1)}(\cdot)$ and $h^{(2)}(\cdot - x_0)$ have disjoint supports provided that $|x_0|$ is sufficiently large. In this setting, by substituting in (4.8) with the metrics $\tilde{g}^{(1)} = g_{\mathbb{R}^m} + \varepsilon h^{(1)}$ and $\tilde{g}^{(2)}(\cdot - x_0) = g_{\mathbb{R}^m} + \varepsilon h^{(2)}(\cdot - x_0)$, we obtain $\Gamma^{(1)}$, $\Gamma^{(2), x_0}$, $\Phi^{(1)}$, $\Phi^{(2), x_0}$, etc., the corresponding functionals. It is easy to check that

$$\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma}) = \Gamma^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}) \quad \text{and} \quad \Phi^{(2), x_0}(\psi_{\lambda,\xi,\gamma}) = \Phi^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}).$$

Consequently, we have

$$\hat{\Phi}^{(2), x_0}(\psi_{\lambda,\xi,\gamma}) = \hat{\Phi}^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}).$$

If $|x_0|$ is large enough such that $h^{(1)}$ and $h^{(2)}(\cdot - x_0)$ have disjoint supports, we find

$$\Gamma(\psi_{\lambda,\xi,\gamma}) = \Gamma^{(1)}(\psi_{\lambda,\xi,\gamma}) + \Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma}) = \Gamma^{(1)}(\psi_{\lambda,\xi,\gamma}) + \Gamma^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}),$$

$$\Phi(\psi_{\lambda,\xi,\gamma}) = \Phi^{(1)}(\psi_{\lambda,\xi,\gamma}) + \Phi^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}),$$

and

$$\nabla\Gamma(\psi_{\lambda,\xi,\gamma}) = \nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma}) + \nabla\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma}).$$

In order to get a similar decomposition for $\hat{\Phi}$, we need the following lemma.

Lemma 4.12. *Given $\Lambda > 0$ arbitrarily, then*

$$\|\nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma})\| \cdot \|\nabla\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma})\| \rightarrow 0 \quad \text{as } |x_0| \rightarrow \infty$$

uniformly in $(\lambda, \xi, \gamma) \in (0, \Lambda] \times \mathbb{R}^m \times S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$.

Proof. Similar to Lemma 4.10, we first fix $R > 0$ large such that $\text{supp } h^{(1)} \cup \text{supp } h^{(2)} \subset B_R$. Then, for arbitrary $v \in \mathcal{D}^{1/2}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$,

$$|\langle \nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma}), v \rangle| \leq C^{(1)} \left(\int_{B_R} \frac{\lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m}{2}}} \cdot |v| + \frac{\lambda^{\frac{m-1}{2}}}{(\lambda^2 + |x - \xi|^2)^{\frac{m-1}{2}}} \cdot |v| d \text{vol}_{g_{\mathbb{R}^m}} \right)$$

where $C^{(1)} > 0$ depends on $\|h^{(1)}\|_{L^\infty}$ and $\|\nabla h^{(1)}\|_{L^\infty}$. Hence, if $|\xi| > R$, we see that

$$\|\nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma})\| \leq C^{(1)} \left(\frac{\Lambda^{\frac{m-1}{2}}}{\text{dist}(\xi, B_R)^m} + \frac{\Lambda^{\frac{m-1}{2}}}{\text{dist}(\xi, B_R)^{m-1}} \right)$$

for all $\lambda \in (0, \Lambda]$ and $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$. And similarly, for some $C^{(2)} > 0$, if $|\xi - x_0| > R$ we have

$$\|\nabla\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma})\| = \|\nabla\Gamma^{(2)}(\psi_{\lambda,\xi-x_0,\gamma})\| \leq C^{(2)} \left(\frac{\Lambda^{\frac{m-1}{2}}}{\text{dist}(\xi - x_0, B_R)^m} + \frac{\Lambda^{\frac{m-1}{2}}}{\text{dist}(\xi - x_0, B_R)^{m-1}} \right)$$

for all $\lambda \in (0, \Lambda]$ and $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$.

Observe that

$$\text{dist}(\xi, B_R) + \min\{|\xi|, R\} = |\xi| \quad \text{and} \quad \text{dist}(\xi - x_0, B_R) + \min\{|\xi - x_0|, R\} = |\xi - x_0|.$$

It follows directly from the triangle inequality that, if $|x_0|$ is large (say $|x_0| > 4R$), it is always $\text{dist}(\xi, B_R) \geq \frac{1}{4}|x_0|$ or $\text{dist}(\xi - x_0, B_R) \geq \frac{1}{4}|x_0|$, hence

$$\|\nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma})\| \cdot \|\nabla\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma})\| \leq \frac{C\Lambda^{\frac{m-1}{2}}}{|x_0|^{m-1}} \rightarrow 0$$

as $|x_0| \rightarrow \infty$ uniformly in $(\lambda, \xi, \gamma) \in (0, \Lambda] \times \mathbb{R}^m \times S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$. \square

An immediate consequence of Lemma 4.10-4.12, we have the following decomposition of $\hat{\Phi}(\psi_{\lambda,\xi,\gamma})$.

Corollary 4.13. *There holds*

$$\langle K_{\psi_{\lambda,\xi,\gamma}}(\nabla\Gamma^{(1)}(\psi_{\lambda,\xi,\gamma})), \nabla\Gamma^{(2), x_0}(\psi_{\lambda,\xi,\gamma}) \rangle \rightarrow 0$$

as $|x_0| \rightarrow \infty$, uniformly in $(\lambda, \xi, \gamma) \in (0, \infty) \times \mathbb{R}^m \times S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$. In particular,

$$\hat{\Phi}(\psi_{\lambda,\xi,\gamma}) = \hat{\Phi}^{(1)}(\psi_{\lambda,\xi,\gamma}) + \hat{\Phi}^{(2)}(\psi_{\lambda,\xi-x_0,\gamma}) + o(1),$$

where $o(1) \rightarrow 0$ as $|x_0| \rightarrow \infty$, uniformly in $(\lambda, \xi, \gamma) \in (0, \infty) \times \mathbb{R}^m \times S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$.

Now we are ready to prove our main result.

Proof of Theorem 1.4. For $m \geq 4$, from Proposition 4.7 and 4.8, we can choose

$$U_R^{(1)} := \{(\lambda, \xi) \in (0, \infty) \times \mathbb{R}^2 : \lambda + |\xi| < R\}$$

and

$$U_R^{(2)} := \{(\lambda, \xi) \in (0, \infty) \times \mathbb{R}^2 : \lambda + |\xi - x_0| < R\}$$

with $R > 0$ suitably large such that, for all $\gamma \in S^{2^{\lfloor \frac{m}{2} \rfloor + 1} - 1}(\mathbb{S}_m)$,

$$\min_{(\lambda,\xi) \in \partial U_R^{(1)}} \hat{\Phi}^{(1)}(\psi_{\lambda,\xi,\gamma}) - \min_{(\lambda,\xi) \in U_R^{(1)}} \hat{\Phi}^{(1)}(\psi_{\lambda,\xi,\gamma}) \geq \frac{1}{R}$$

and

$$\min_{(\lambda, \xi) \in \partial U_R^{(2)}} \hat{\Phi}^{(2)}(\psi_{\lambda, \xi - x_0, \gamma}) - \min_{(\lambda, \xi) \in U_R^{(2)}} \hat{\Phi}^{(2)}(\psi_{\lambda, \xi - x_0, \gamma}) \geq \frac{1}{R}$$

Hence, by Corollary 4.13, we find that for $|x_0| \gg R$ there exists $\delta > 0$ such that $\{(\lambda, \xi) : \hat{\Phi}(\psi_{\lambda, \xi, \gamma}) < -\delta\}$ has two disconnected components in $U_R^{(1)}$ and $U_R^{(2)}$ respectively. And therefore, by applying Theorem 2.6, the two distinct local minima of $\hat{\Phi}(\psi_{\lambda, \xi, \gamma})$ give rise to two distinct solutions of Eq. (4.2) (and hence two distinct solutions of Eq. (4.1) via the pull-back of stereographic projection) for small values of ε . \square

A Appendix

A.1 An explanation of the (k, p) -elementary matrix

Here we point out that linear combinations of the (k, p) -elementary matrices, $k = 1, \dots, m$, create a very large family of diagonal matrices. To have a better look at this, let's restrict ourselves in the 3-dimensional space (it will cause no confusion if we are in any other dimensions). First of all, let $h^{(k)}$, $k = 1, 2, 3$, be (k, p) -elementary at a common point $\xi = (\xi_1, \xi_2, \xi_3) \in \mathbb{R}^3$ with correspondingly $c_k \neq 0$ being the non-zero coefficients, i.e., we have the local expressions

$$h^{(1)}(x) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & h_{22}^{(1)}(x) & 0 \\ 0 & 0 & h_{33}^{(1)}(x) \end{pmatrix}, \quad h^{(2)}(x) = \begin{pmatrix} h_{11}^{(2)}(x) & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & h_{33}^{(2)}(x) \end{pmatrix}$$

and

$$h^{(3)}(x) = \begin{pmatrix} h_{11}^{(3)}(x) & 0 & 0 \\ 0 & h_{22}^{(3)}(x) & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

with

$$\left. \begin{aligned} h_{22}^{(1)}(x) &= a_{22}^{(1)} + c_1(x_1 - \xi_1) + c_2^{(1)}(x_2 - \xi_2) + o(|x - \xi|^p) \\ h_{33}^{(1)}(x) &= a_{33}^{(1)} + c_1(x_1 - \xi_1) + c_3^{(1)}(x_3 - \xi_3) + o(|x - \xi|^p) \end{aligned} \right\} \text{for } h^{(1)},$$

$$\left. \begin{aligned} h_{11}^{(2)}(x) &= a_{11}^{(2)} + c_1^{(2)}(x_1 - \xi_1) + c_2(x_2 - \xi_2) + o(|x - \xi|^p) \\ h_{33}^{(2)}(x) &= a_{33}^{(2)} + c_2(x_2 - \xi_2) + c_3^{(2)}(x_3 - \xi_3) + o(|x - \xi|^p) \end{aligned} \right\} \text{for } h^{(2)},$$

$$\left. \begin{aligned} h_{11}^{(3)}(x) &= a_{11}^{(3)} + c_1^{(3)}(x_1 - \xi_1) + c_3(x_3 - \xi_3) + o(|x - \xi|^p) \\ h_{22}^{(3)}(x) &= a_{22}^{(3)} + c_2^{(3)}(x_2 - \xi_2) + c_3(x_3 - \xi_3) + o(|x - \xi|^p) \end{aligned} \right\} \text{for } h^{(3)},$$

where $a_{ii}^{(k)}$ and $c_i^{(k)}$, $i, k = 1, 2, 3$, are some given real constants. Then, a linear combination of $h^{(k)}$, say for instance $\alpha h^{(1)} + \beta h^{(2)} + \gamma h^{(3)}$ for $\alpha, \beta, \gamma \in \mathbb{R}$, gives rise to a matrix of the form (in local expansion around the point ξ)

$$\begin{pmatrix} a_{11} + \mathbf{b}_1 \cdot (x - \xi) & 0 & 0 \\ 0 & a_{22} + \mathbf{b}_2 \cdot (x - \xi) & 0 \\ 0 & 0 & a_{33} + \mathbf{b}_3 \cdot (x - \xi) \end{pmatrix} + o(|x - \xi|^p) \quad (\text{A.1})$$

with $a_{11} = \beta a_{11}^{(2)} + \gamma a_{11}^{(3)}$, $a_{22} = \alpha a_{22}^{(1)} + \gamma a_{22}^{(3)}$, $a_{33} = \alpha a_{33}^{(1)} + \beta a_{33}^{(2)}$ characterizing the zero-order terms and the vector-valued coefficients

$$\mathbf{b}_1 = (\beta c_1^{(2)} + \gamma c_1^{(3)}, \beta c_2, \gamma c_3), \quad \mathbf{b}_2 = (\alpha c_1, \alpha c_2^{(1)} + \gamma c_2^{(3)}, \gamma c_3), \quad \mathbf{b}_3 = (\alpha c_1, \beta c_2, \alpha c_3^{(1)} + \beta c_3^{(2)})$$

characterizing the first-order terms. Such specific local expansion is satisfied by a large class of diagonal matrix functions. Moreover, notice that (A.1) is defined only around one spatial point, we can consider such kind of expressions around an arbitrary finite number of spatial points. And even more, there is no restriction on the global behavior of $h^{(k)}$, $k = 1, 2, 3$. This is why we use the word “*elementary*” in Definition 1.3.

A.2 Some basic estimates

In the sequel we use the notation $f \lesssim g$ for two functions f and g , when there exists a constant $C > 0$ such that $f \leq Cg$. Then, by the explicit expression of $\psi_{1,0,\gamma}$ in (2.10), we have

$$|\psi_{1,0,\gamma}(x)| \lesssim \frac{1}{(1 + |x|^2)^{\frac{m-1}{2}}} \quad \text{and} \quad |\nabla \psi_{1,0,\gamma}(x)| \lesssim \frac{1}{(1 + |x|^2)^{\frac{m}{2}}}.$$

Given $R > \delta > 0$ and let $\lambda \searrow 0$ be a parameter, for arbitrary $\varphi \in \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ with $\|\varphi\| = 1$, we estimate:

$$\begin{aligned} \left| \operatorname{Re} \int_{|x| \geq \frac{\delta}{\lambda}} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \varphi)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\leq \left(\int_{|x| > \frac{\delta}{\lambda}} |\nabla_{\partial_i} \psi_{1,0,\gamma}|^{\frac{2m}{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \right)^{\frac{m+1}{2m}} |\varphi|_{\frac{2m}{m-1}} \\ &\lesssim \left(\int_{\frac{\delta}{\lambda}}^{\infty} \frac{r^{m-1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \lambda^{\frac{m-1}{2}} \quad \text{for } m \geq 2, \end{aligned} \tag{A.2}$$

$$\begin{aligned} \left| \operatorname{Re} \int_{|x| \leq \frac{\delta}{\lambda}} x_j (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \varphi)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{\frac{2m}{m+1} + m - 1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &= \left(\int_0^1 + \int_1^{\frac{\delta}{\lambda}} \frac{r^{\frac{2m}{m+1} + m - 1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \begin{cases} \lambda^{-\frac{1}{2}} & m = 2, \\ |\ln \lambda|^{\frac{2}{3}} & m = 3, \\ 1 & m \geq 4, \end{cases} \end{aligned} \tag{A.3}$$

$$\left| \operatorname{Re} \int_{\frac{\delta}{\lambda} \leq |x| \leq \frac{R}{\lambda}} x_j (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \varphi) d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| \lesssim \left(\int_{\frac{\delta}{\lambda}}^{\frac{R}{\lambda}} \frac{r^{\frac{2m}{m+1} + m - 1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}}$$

$$\lesssim \begin{cases} \lambda^{-\frac{1}{2}} & m = 2, \\ 1 & m = 3, \\ \lambda^{\frac{m-3}{2}} & m \geq 4, \end{cases} \quad (\text{A.4})$$

$$\begin{aligned} \left| \operatorname{Re} \int_{|x| \leq \frac{\delta}{\lambda}} x_j x_k (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \varphi) d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{\frac{4m}{m+1}+m-1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &= \left(\int_0^1 + \int_1^{\frac{\delta}{\lambda}} \frac{r^{\frac{4m}{m+1}+m-1}}{(1+r^2)^{\frac{m^2}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \begin{cases} \lambda^{\frac{m-5}{2}} & 2 \leq m \leq 4, \\ |\ln \lambda|^{\frac{3}{5}} & m = 5, \\ 1 & m \geq 6. \end{cases} \quad (\text{A.5}) \end{aligned}$$

$$\begin{aligned} \left| \operatorname{Re} \int_{|x| \leq \frac{\delta}{\lambda}} (\partial_k \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma}, \varphi) d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{m-1}}{(1+r^2)^{\frac{m(m-1)}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &= \left(\int_0^1 + \int_1^{\frac{\delta}{\lambda}} \frac{r^{m-1}}{(1+r^2)^{\frac{m(m-1)}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \begin{cases} \lambda^{-\frac{1}{2}} & m = 2, \\ |\ln \lambda|^{\frac{2}{3}} & m = 3, \\ 1 & m \geq 4, \end{cases} \quad (\text{A.6}) \end{aligned}$$

$$\begin{aligned} \left| \operatorname{Re} \int_{\frac{\delta}{\lambda} < |x| \leq \frac{R}{\lambda}} (\partial_k \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma}, \varphi) d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\lesssim \left(\int_{\frac{\delta}{\lambda}}^{\frac{R}{\lambda}} \frac{r^{m-1}}{(1+r^2)^{\frac{m(m-1)}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \begin{cases} \lambda^{-\frac{1}{2}} & m = 2, \\ 1 & m = 3, \\ \lambda^{\frac{m-3}{2}} & m \geq 4, \end{cases} \quad (\text{A.7}) \end{aligned}$$

$$\begin{aligned} \left| \operatorname{Re} \int_{|x| \leq \frac{\delta}{\lambda}} x_j (\partial_k \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma}, \varphi) d \operatorname{vol}_{g_{\mathbb{R}^m}} \right| &\lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{\frac{2m}{m+1}+m-1}}{(1+r^2)^{\frac{m(m-1)}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &= \left(\int_0^1 + \int_1^{\frac{\delta}{\lambda}} \frac{r^{\frac{2m}{m+1}+m-1}}{(1+r^2)^{\frac{m(m-1)}{m+1}}} dr \right)^{\frac{m+1}{2m}} \\ &\lesssim \begin{cases} \lambda^{\frac{m-5}{2}} & 2 \leq m \leq 4, \\ |\ln \lambda|^{\frac{3}{5}} & m = 5, \\ 1 & m \geq 6. \end{cases} \quad (\text{A.8}) \end{aligned}$$

In particular, for $m = 5$, we have $\frac{2m}{m+1} = \frac{5}{3}$ and we estimate: for $p > 2$ fixed,

$$\left(\int_{|x| \leq \frac{\delta}{\lambda}} |x|^{\frac{5p}{3}} |\nabla_{\partial_i} \psi_{1,0,\gamma}|^{\frac{5}{3}} d \text{vol}_{g_{\mathbb{R}^m}} \right)^{\frac{3}{5}} \lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{\frac{5p}{3}+4}}{(1+r^2)^{\frac{25}{6}}} dr \right)^{\frac{3}{5}} \lesssim \lambda^{2-p} \quad (\text{A.9})$$

and

$$\left(\int_{|x| \leq \frac{\delta}{\lambda}} |x|^{\frac{5(p-1)}{3}} |\psi_{1,0,\gamma}|^{\frac{5}{3}} d \text{vol}_{g_{\mathbb{R}^m}} \right)^{\frac{3}{5}} \lesssim \left(\int_0^{\frac{\delta}{\lambda}} \frac{r^{\frac{5(p-1)}{3}+4}}{(1+r^2)^{\frac{10}{3}}} dr \right)^{\frac{3}{5}} \lesssim \lambda^{2-p}. \quad (\text{A.10})$$

A.3 Proofs of Lemma 4.6 and Proposition 4.7

Proof of Lemma 4.6. Let $z \in \mathcal{M}$, note that K_z stands for the inverse of $\nabla^2 \mathcal{J}_0(z)$ on $\mathcal{W}_z = T_z \mathcal{M}^\perp$, by setting $\bar{w}_z = -K_z(\nabla \Gamma(z))$, we see that

$$\langle K_z(\nabla \Gamma(z)), \nabla \Gamma(z) \rangle = -\langle \nabla \Gamma(z), \bar{w}_z \rangle = \nabla^2 \mathcal{J}_0(z)[\bar{w}_z, \bar{w}_z]. \quad (\text{A.11})$$

Thanks to the expression of Γ in Lemma 4.3 and the fact $\nabla \Gamma(z) \in \mathcal{W}_z$, we obtain (integrating by parts)

$$\begin{aligned} \langle \nabla \Gamma(z), v \rangle &= \text{Re} \int_{\mathbb{R}^m} \frac{\text{tr } h}{2} \left[\frac{1}{2} [(D_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} + (z, D_{g_{\mathbb{R}^m}} v)_{g_{\mathbb{R}^m}}] - |z|_{g_{\mathbb{R}^m}}^{2^*-2} (z, v)_{g_{\mathbb{R}^m}} \right] \\ &\quad - \frac{1}{4} \sum_i h_{ii} \left((\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} z, v)_{g_{\mathbb{R}^m}} + (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} v, z)_{g_{\mathbb{R}^m}} \right) d \text{vol}_{g_{\mathbb{R}^m}} \\ &= \text{Re} \int_{\mathbb{R}^m} \frac{\text{tr } h}{2} \left[(D_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} - |z|_{g_{\mathbb{R}^m}}^{2^*-2} (z, v)_{g_{\mathbb{R}^m}} \right] d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad + \frac{1}{4} \text{Re} \int_{\mathbb{R}^m} (\nabla(\text{tr } h) \cdot_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad - \frac{1}{2} \sum_i \text{Re} \int_{\mathbb{R}^m} h_{ii} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} z, v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad - \frac{1}{4} \sum_i \text{Re} \int_{\mathbb{R}^m} \partial_i h_{ii} (\partial_i \cdot_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \end{aligned} \quad (\text{A.12})$$

for all $v \in \mathcal{W}_z$, where we have used the facts

$$\begin{aligned} \text{Re} \int_{\mathbb{R}^m} (\text{tr } h)(z, D_{g_{\mathbb{R}^m}} v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} &= \text{Re} \int_{\mathbb{R}^m} (\text{tr } h)(D_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \\ &\quad + \text{Re} \int_{\mathbb{R}^m} (\nabla(\text{tr } h) \cdot_{g_{\mathbb{R}^m}} z, v)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

and

$$\partial_i \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} z, v) = \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} z, v) - \text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} v, z).$$

Note that $z \in \mathcal{M}$ satisfies the equation

$$D_{g_{\mathbb{R}^m}} z = |z|_{g_{\mathbb{R}^m}}^{2^*-2} z \quad \text{on } \mathbb{R}^m,$$

we have

$$\nabla \Gamma(z) = -\frac{1}{2} \sum_i h_{ii} \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} z + \frac{1}{4} \nabla(\text{tr } h) \cdot_{g_{\mathbb{R}^m}} z - \frac{1}{4} \sum_i \partial_i h_{ii} \partial_i \cdot_{g_{\mathbb{R}^m}} z$$

for all $z \in \mathcal{M}$. Moreover, we have \bar{w}_z solves the equation

$$\nabla^2 \mathcal{J}_0(z)[\bar{w}_z] = -\nabla \Gamma(z). \quad (\text{A.13})$$

where

$$\nabla^2 \mathcal{J}_0(z)[\bar{w}_z] = D_{g_{\mathbb{R}^m}} \bar{w}_z - |z|_{g_{\mathbb{R}^m}}^{2^*-2} \bar{w}_z - (2^* - 2) |z|_{g_{\mathbb{R}^m}}^{2^*-4} \text{Re}(z, \bar{w}_z) z.$$

Next, we will explicitly substitute $z = \psi_{\lambda, \xi, \gamma}$ into (A.13). For this purpose, let us set $\bar{w}_{\lambda, \xi, \gamma} = \bar{w}_{\psi_{\lambda, \xi, \gamma}}$ and $w_{\lambda, \xi, \gamma}^*(x) = \lambda^{\frac{m-1}{2}} \bar{w}_{\lambda, \xi, \gamma}(\lambda x + \xi)$. Then, by change of variables, we have $w_{\lambda, \xi, \gamma}^*$ solves

$$\begin{aligned} \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{\lambda, \xi, \gamma}^*] &= \frac{1}{2} \sum_i h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma} - \frac{\lambda}{4} \nabla(\text{tr } h)(\lambda x + \xi) \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma} \\ &\quad + \frac{\lambda}{4} \sum_i \partial_i h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma} \end{aligned} \quad (\text{A.14})$$

Taking to the limit $\lambda \rightarrow 0$, by (A.2), (A.6), (A.7) and the continuity of $K_{\psi_{1,0,\gamma}} = \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})^{-1}$, we see that $w_{\lambda, \xi, \gamma}^* \rightarrow w_{0, \xi, \gamma}^*$ in $\mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$ where $w_{0, \xi, \gamma}^*$ solves the equation

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0, \xi, \gamma}^*] = \frac{1}{2} \sum_i h_{ii}(\xi) \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}. \quad (\text{A.15})$$

The solution of Eq. (A.15) is not unique. It is uniquely determined up to the addition of elements of $\ker \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma}) = T_{\psi_{1,0,\gamma}} \mathcal{M}$. However, by changing of variables again, we find the quantity

$$\lim_{\lambda \rightarrow 0} \langle \nabla \Gamma(\psi_{\lambda, \xi, \gamma}), \bar{w}_{\lambda, \xi, \gamma} \rangle = -\frac{1}{2} \sum_i h_{ii}(\xi) \text{Re} \int_{\mathbb{R}^m} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, w_{0, \xi, \gamma}^*)_{g_{\mathbb{R}^m}} d \text{vol}_{g_{\mathbb{R}^m}}$$

does not depend on the projection of $w_{0, \xi, \gamma}^*$ in $T_{\psi_{1,0,\gamma}} \mathcal{M}$ (since $\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma} \in T_{\psi_{1,0,\gamma}} \mathcal{M}^\perp$). Hence, it is enough for us to construct one solution to (A.15). Observing that Eq. (A.15) is linear in h , one checks easily that

$$w_{0, \xi, \gamma}^*(x) = \frac{1}{2} \sum_i h_{ii}(\xi) x_i \nabla_{\partial_i} \psi_{1,0,\gamma}(x)$$

is indeed a solution. Moreover, following from (4.10), (4.11) and the rules of Clifford multiplication, we see that

$$\text{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, x_i \nabla_{\partial_i} \psi_{1,0,\gamma}) = 0$$

for $i = 1, \dots, m$ and

$$\begin{aligned} & \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, x_j \nabla_{\partial_j} \psi_{1,0,\gamma}) \\ &= \frac{m^m x_i x_j}{(1 + |x|^2)^{m+1}} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} (1 - x) \cdot_{g_{\mathbb{R}^m}} \gamma, \partial_j \cdot_{g_{\mathbb{R}^m}} \gamma) - \frac{m^m x_j^2}{(1 + |x|^2)^{m+1}} \\ &= \frac{m^m x_i x_j}{(1 + |x|^2)^{m+1}} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \gamma, \partial_j \cdot_{g_{\mathbb{R}^m}} \gamma) \\ &\quad - \sum_{\substack{k=1 \\ k \neq i, k \neq j}}^m \frac{m^m x_i x_j x_k}{(1 + |x|^2)^{m+1}} \operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \partial_k \cdot_{g_{\mathbb{R}^m}} \gamma, \partial_j \cdot_{g_{\mathbb{R}^m}} \gamma) - \frac{m^m x_j^2}{(1 + |x|^2)^{m+1}} \end{aligned}$$

for $i \neq j$. Since $\operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \gamma, \partial_j \cdot_{g_{\mathbb{R}^m}} \gamma) \equiv 0$ and $\operatorname{Re}(\partial_i \cdot_{g_{\mathbb{R}^m}} \partial_k \cdot_{g_{\mathbb{R}^m}} \gamma, \partial_j \cdot_{g_{\mathbb{R}^m}} \gamma)$ are constants, we deduce

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \langle \nabla \Gamma(\psi_{\lambda,\xi,\gamma}), \bar{w}_{\lambda,\xi,\gamma} \rangle &= -\frac{1}{2} \sum_i h_{ii}(\xi) \operatorname{Re} \int_{\mathbb{R}^m} (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, w_{0,\xi,\gamma}^*)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\ &= \frac{1}{4} \sum_{\substack{i,j \\ i \neq j}} h_{ii}(\xi) h_{jj}(\xi) \int_{\mathbb{R}^m} \frac{m^m x_j^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\ &= \frac{1}{4} \int_{\mathbb{R}^m} \frac{m^m x_1^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} ((\operatorname{tr} h)^2 - \operatorname{tr}(h^2))(\xi) \end{aligned}$$

Notice that $|x|^2 = x_1^2 + \dots + x_m^2$ and

$$\int_{\mathbb{R}^m} \frac{x_1^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} = \dots = \int_{\mathbb{R}^m} \frac{x_m^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}}$$

we have

$$\lim_{\lambda \rightarrow 0} \langle \nabla \Gamma(\psi_{\lambda,\xi,\gamma}), \bar{w}_{\lambda,\xi,\gamma} \rangle = \frac{m^{m-1}}{4} \int_{\mathbb{R}^m} \frac{|x|^2}{(1 + |x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} ((\operatorname{tr} h)^2 - \operatorname{tr}(h^2))(\xi)$$

as desired. \square

Proof of Proposition 4.7. The procedure is to find the expansion of $\hat{\Phi}(\psi_{\lambda,\xi,\gamma})$ with respect to λ in a neighborhood of 0. In order to do this, we need to find the expansions for both of the functionals $\Phi(\psi_{\lambda,\xi,\gamma})$ and $\langle K_{\psi_{\lambda,\xi,\gamma}}(\nabla \Gamma(\psi_{\lambda,\xi,\gamma})), \nabla \Gamma(\psi_{\lambda,\xi,\gamma}) \rangle$.

For $m \geq 3$, we can assume without loss of generality that $h = \operatorname{diag}(h_{11}, \dots, h_{m-1, m-1}, 0)$ is (m, p) -elementary around the point $\xi \in \mathbb{R}^m$ with $\partial_m h_{ii}(\xi) \equiv c_m \neq 0$, for $1 \leq i \leq m-1$. The other cases can be done in a very similar manner.

By Definition 1.3 and by the change of variables $x \mapsto \lambda x + \xi$, we find

$$h_{ii}(\lambda x + \xi) = h_{ii}(\xi) + \lambda(c_i x_i + c_m x_m) + o(|\lambda x|^p)$$

for $i = 1, \dots, m-1$, and hence

$$\begin{aligned} (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\lambda x + \xi) &= - \sum_{\substack{i,j \\ i \neq j}} (h_{ii} h_{jj})(\lambda x + \xi) \\ &= - \sum_{\substack{i,j \\ i \neq j}} \left[h_{ii}(\xi) h_{jj}(\xi) + \lambda h_{ii}(\xi)(c_j x_j + c_m x_m) + \lambda h_{jj}(\xi)(c_i x_i + c_m x_m) \right. \\ &\quad \left. + \lambda^2 (c_i c_j x_i x_j + c_i c_m x_i x_m + c_j c_m x_j x_m + c_m^2 x_m^2) + o(|\lambda x|^p) \right]. \end{aligned}$$

Terms of order λ in $\Phi(\psi_{\lambda, \xi, \gamma})$ have coefficients like

$$\int_{\mathbb{R}^m} \frac{x_i}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}$$

which are all zero. And the coefficients of λ^2 are of the type

$$\int_{\mathbb{R}^m} \frac{x_i x_j}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}$$

and such integral are non-zero only for $i = j$. Hence, by Lemma 4.5 and $p \geq 2$, we deduce

$$\Phi(\psi_{\lambda, \xi, \gamma}) = C_0 (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi) - A c_m^2 \lambda^2 + o(\lambda^2) \quad (\text{A.16})$$

where

$$A = \frac{m^{m-2}(m-1)(m-2)}{16} \int_{\mathbb{R}^m} \frac{|x|^2}{(1 + |x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}.$$

It remains to find an expansion for $\langle K_{\psi_{\lambda, \xi, \gamma}}(\nabla \Gamma(\psi_{\lambda, \xi, \gamma})), \nabla \Gamma(\psi_{\lambda, \xi, \gamma}) \rangle$. By Eq. (A.14) and the local expansion of h_{ii} , we want to develop $w_{\lambda, \xi, \gamma}^*$ in powers of λ .

We first assume $m \geq 6$. Then, by the computations (A.2)-(A.8) in the Appendix, we have

$$w_{\lambda, \xi, \gamma}^* = w_{0, \xi, \gamma}^* + \lambda \nu_{0, \xi, \gamma}^* + \lambda^2 \mu_{0, \xi, \gamma}^* + o(\lambda^2) \quad \text{in } \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^m, \mathbb{S}(\mathbb{R}^m))$$

where $w_{0, \xi, \gamma}^*$, $\nu_{0, \xi, \gamma}^*$ and $\mu_{0, \xi, \gamma}^*$ satisfy the following equations

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*] = \frac{1}{2} \sum_{i=1}^{m-1} h_{ii}(\xi) \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \quad (\text{A.17})$$

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*] = \frac{1}{2} \sum_{i=1}^{m-1} (c_i x_i + c_m x_m) \partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma} - \frac{m-1}{4} c_m \partial_m \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma} \quad (\text{A.18})$$

and (since the Hessian of h_{ii} at ξ vanishes identically)

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{0,\xi,\gamma}^*] = 0. \quad (\text{A.19})$$

And now we have

$$\begin{aligned} \langle K_{\psi_{\lambda, \xi, \gamma}}(\nabla \Gamma(\psi_{\lambda, \xi, \gamma})), \nabla \Gamma(\psi_{\lambda, \xi, \gamma}) \rangle &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{\lambda, \xi, \gamma}^*, w_{\lambda, \xi, \gamma}^*] \\ &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] + 2 \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*] \lambda \\ &\quad + [2 \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \mu_{0,\xi,\gamma}^*] + \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]] \lambda^2 + o(\lambda^2). \end{aligned}$$

Next, we will calculate the coefficients in front of each power of λ :

1. The zero order term

As was shown by Lemma 4.6, we have

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] = C_1 (\operatorname{tr}(h^2) - (\operatorname{tr} h)^2)(\xi).$$

2. The first order term

By using the equation (A.18), we find

$$\begin{aligned} \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*] &= \frac{1}{2} \sum_{i=1}^{m-1} \operatorname{Re} \int_{\mathbb{R}^m} (c_i x_i + c_m x_m) (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, w_{0,\xi,\gamma}^*)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\ &\quad - \frac{(m-1)c_m}{4} \operatorname{Re} \int_{\mathbb{R}^m} (\partial_m \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma}, w_{0,\xi,\gamma}^*)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}}. \end{aligned}$$

Note that $w_{0,\xi,\gamma}^*$ can be explicitly formulated as

$$w_{0,\xi,\gamma}^* = \frac{1}{2} \sum_{i=1}^{m-1} h_{ii}(\xi) x_i \nabla_{\partial_i} \psi_{1,0,\gamma},$$

we find the integrals in $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]$ are of the type

$$\begin{aligned} \int_{\mathbb{R}^m} \frac{x_i x_j}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}}, \quad \int_{\mathbb{R}^m} \frac{x_i^2 x_j}{(1+|x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}}, \quad \int_{\mathbb{R}^m} \frac{x_i^2 x_k x_j}{(1+|x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}}, \\ \int_{\mathbb{R}^m} \frac{x_i^3}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} \quad \text{and} \quad \int_{\mathbb{R}^m} \frac{x_i x_j x_k x_l}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} \end{aligned}$$

with $i \neq j$, $i \neq k$, $j \neq k$ and $k \neq l$, which are all zero. Hence the first order term vanishes identically.

3. The second order term

By virtue of (A.19), we only need to evaluate $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]$. To this end, it sufficient to solve Eq. (A.18). Analogous to solving Eq. (A.15), we only need to construct one solution to (A.18). Particularly, one checks that

$$\nu_{0,\xi,\gamma}^* = \frac{1}{4} \left(\sum_{i=1}^{m-1} c_i x_i^2 \nabla_{\partial_i} \psi_{1,0,\gamma} - c_m x_m^2 \nabla_{\partial_m} \psi_{1,0,\gamma} \right) - \frac{m-1}{4} c_m x_m \psi_{1,0,\gamma}$$

solves Eq. (A.18). And hence, by the explicit expression of $\psi_{1,0,\gamma}$, we have

$$\begin{aligned}
\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*] &= \frac{1}{2} \sum_{i=1}^{m-1} \operatorname{Re} \int_{\mathbb{R}^m} (c_i x_i + c_m x_m) (\partial_i \cdot_{g_{\mathbb{R}^m}} \nabla_{\partial_i} \psi_{1,0,\gamma}, \nu_{0,\xi,\gamma}^*)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\
&\quad - \frac{(m-1)c_m}{4} \operatorname{Re} \int_{\mathbb{R}^m} (\partial_m \cdot_{g_{\mathbb{R}^m}} \psi_{1,0,\gamma}, \nu_{0,\xi,\gamma}^*)_{g_{\mathbb{R}^m}} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\
&= \frac{1}{2} \sum_{i=1}^{m-1} \int_{\mathbb{R}^m} \left(\frac{m^m c_m^2 x_m^4}{4(1+|x|^2)^{m+1}} - \frac{m^{m-1}(m-1)c_m^2 x_m^2}{4(1+|x|^2)^m} \right) d \operatorname{vol}_{g_{\mathbb{R}^m}} \\
&\quad - \frac{(m-1)c_m^2}{4} \int_{\mathbb{R}^m} \frac{m^{m-1} x_m^2}{4(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} \\
&= -\frac{5m^{m-2}(m-1)(m-2)c_m^2}{64} \int_{\mathbb{R}^m} \frac{|x|^2}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}},
\end{aligned}$$

where the last equality follows from the fact

$$\int_{\mathbb{R}^m} \frac{x_m^4}{(1+|x|^2)^{m+1}} d \operatorname{vol}_{g_{\mathbb{R}^m}} = \frac{3(m+2)}{8m} \int_{\mathbb{R}^m} \frac{x_m^2}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} \quad \text{for } m \geq 3.$$

Now, together with (A.16), we find

$$\hat{\Phi}(\psi_{\lambda,\xi,\gamma}) = -\frac{3m^{m-2}(m-1)(m-2)c_m^2}{128} \lambda^2 \int_{\mathbb{R}^m} \frac{|x|^2}{(1+|x|^2)^m} d \operatorname{vol}_{g_{\mathbb{R}^m}} + o(\lambda^2)$$

as desired.

For $m = 5$, by (A.2)-(A.10), we find

$$w_{\lambda,\xi,\gamma}^* = w_{0,\xi,\gamma}^* + \lambda \nu_{0,\xi,\gamma}^* + \mu_{\lambda,\xi,\gamma}^* + \rho_{\lambda,\xi,\gamma}^* \quad \text{in } \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^5, \mathbb{S}(\mathbb{R}^5)) \quad (\text{A.20})$$

where

$$\begin{aligned}
\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*] &= \frac{1}{2} \sum_{i=1}^4 h_{ii}(\xi) \partial_i \cdot_{g_{\mathbb{R}^5}} \nabla_{\partial_i} \psi_{1,0,\gamma} \\
\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*] &= \frac{1}{2} \sum_{i=1}^4 (c_i x_i + c_5 x_5) \partial_i \cdot_{g_{\mathbb{R}^5}} \nabla_{\partial_i} \psi_{1,0,\gamma} - c_5 \partial_5 \cdot_{g_{\mathbb{R}^5}} \psi_{1,0,\gamma} \\
\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] &= (1-\eta\lambda) \left[\frac{1}{2} \sum_{i=1}^5 h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^5}} \nabla_{\partial_i} \psi_{1,0,\gamma} \right. \\
&\quad \left. - \frac{\lambda}{4} (\nabla \operatorname{tr} h)(\lambda x + \xi) \cdot_{g_{\mathbb{R}^5}} \psi_{1,0,\gamma} + \frac{\lambda}{4} \sum_{i=1}^5 \partial_i h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^5}} \psi_{1,0,\gamma} \right. \\
&\quad \left. - \frac{1}{2} \sum_{i=1}^4 \left[h_{ii}(\xi) + \lambda(c_i x_i + c_5 x_5) \right] \partial_i \cdot_{g_{\mathbb{R}^5}} \nabla_{\partial_i} \psi_{1,0,\gamma} + \lambda c_5 \partial_5 \cdot_{g_{\mathbb{R}^5}} \psi_{1,0,\gamma} \right]
\end{aligned}$$

and

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\rho_{\lambda,\xi,\gamma}^*] = o(\lambda^p) \eta_\lambda (|x|^p |\nabla_{\partial_i} \psi_{1,0,\gamma}| + |x|^{p-1} |\psi_{1,0,\gamma}|).$$

In the above expressions, η_λ is a smooth cut-off function such that $\eta_\lambda(x) \equiv 1$ for $|x| \leq \delta/\lambda$ and $\eta_\lambda(x) \equiv 0$ for $|x| > 2\delta/\lambda$ with $\delta > 0$ being fixed suitably small. We emphasize that, by (A.4), (A.7), (A.9) and (A.10), we have $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] = O(\lambda^2)$ and $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\rho_{\lambda,\xi,\gamma}^*] = o(\lambda^2)$.

By using the expansion (A.20), we have

$$\begin{aligned} \langle K_{\psi_{\lambda,\xi,\gamma}}(\nabla\Gamma(\psi_{\lambda,\xi,\gamma})), \nabla\Gamma(\psi_{\lambda,\xi,\gamma}) \rangle &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{\lambda,\xi,\gamma}^*, w_{\lambda,\xi,\gamma}^*] \\ &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] + 2\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]\lambda \\ &\quad + 2\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \mu_{\lambda,\xi,\gamma}^*] + \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]\lambda^2 + o(\lambda^2). \end{aligned}$$

Since $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] = O(\lambda^2)$, $2^* = \frac{5}{2}$ and $\int_{|x|>\frac{\delta}{\lambda}} |w_{0,\xi,\gamma}^*|^{\frac{5}{2}} d \text{vol}_{g_{\mathbb{R}^5}} \rightarrow 0$ as $\lambda \rightarrow 0$, we find

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \mu_{\lambda,\xi,\gamma}^*] = \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] = o(\lambda^2)$$

Then we see that the coefficient of the first order term vanishes as in the case $m \geq 6$ and the coefficient in front of λ^2 is $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]$. Therefore we can proceed as in the case $m \geq 6$ to obtain our conclusion.

For $m = 4$, by using the stronger asymptotic condition on h , we find

$$w_{\lambda,\xi,\gamma}^* = w_{0,\xi,\gamma}^* + \lambda \nu_{0,\xi,\gamma}^* + \mu_{\lambda,\xi,\gamma}^* \quad \text{in } \mathcal{D}^{\frac{1}{2}}(\mathbb{R}^4, \mathbb{S}(\mathbb{R}^4)) \quad (\text{A.21})$$

where

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*] = \frac{1}{2} \sum_{i=1}^3 h_{ii}(\xi) \partial_i \cdot_{g_{\mathbb{R}^4}} \nabla_{\partial_i} \psi_{1,0,\gamma}$$

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*] = \frac{1}{2} \sum_{i=1}^3 (c_i x_i + c_4 x_4) \partial_i \cdot_{g_{\mathbb{R}^4}} \nabla_{\partial_i} \psi_{1,0,\gamma} - \frac{3}{4} c_4 \partial_4 \cdot_{g_{\mathbb{R}^4}} \psi_{1,0,\gamma}$$

$$\begin{aligned} \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] &= (1 - \eta_\lambda) \left[\frac{1}{2} \sum_{i=1}^4 h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^4}} \nabla_{\partial_i} \psi_{1,0,\gamma} \right. \\ &\quad - \frac{\lambda}{4} \nabla \text{tr}(h)(\lambda x + \xi) \cdot_{g_{\mathbb{R}^4}} \psi_{1,0,\gamma} + \frac{\lambda}{4} \sum_{i=1}^4 \partial_i h_{ii}(\lambda x + \xi) \partial_i \cdot_{g_{\mathbb{R}^4}} \psi_{1,0,\gamma} \\ &\quad \left. - \frac{1}{2} \sum_{i=1}^3 \left[h_{ii}(\xi) + \lambda(c_i x_i + c_4 x_4) \right] \partial_i \cdot_{g_{\mathbb{R}^4}} \nabla_{\partial_i} \psi_{1,0,\gamma} + \frac{3\lambda c_4}{4} \partial_4 \cdot_{g_{\mathbb{R}^4}} \psi_{1,0,\gamma} \right], \end{aligned}$$

and η_λ is a smooth cut-off function as before. Following from (A.2), (A.4) and (A.7), we can see that $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] = O(\lambda^{\frac{3}{2}})$. Then, by using the expansion (A.21), we have

$$\begin{aligned} \langle K_{\psi_{\lambda,\xi,\gamma}}(\nabla\Gamma(\psi_{\lambda,\xi,\gamma})), \nabla\Gamma(\psi_{\lambda,\xi,\gamma}) \rangle &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{\lambda,\xi,\gamma}^*, w_{\lambda,\xi,\gamma}^*] \\ &= \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] + 2\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]\lambda \\ &\quad + 2\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \mu_{\lambda,\xi,\gamma}^*] + \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]\lambda^2 + o(\lambda^2). \end{aligned}$$

Since $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*] = O(\lambda^{\frac{3}{2}})$, $2^* = \frac{8}{3}$ and $(\int_{|x|>\frac{\delta}{\lambda}} |w_{0,\xi,\gamma}^*|^{\frac{8}{3}} d \text{vol}_{g_{\mathbb{R}^4}})^{\frac{3}{8}} = O(\lambda^{\frac{3}{2}})$ as $\lambda \rightarrow 0$, we find

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{0,\xi,\gamma}^*, \mu_{\lambda,\xi,\gamma}^*] = \nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\mu_{\lambda,\xi,\gamma}^*, w_{0,\xi,\gamma}^*] = o(\lambda^2)$$

Hence, apart from the zero order term, the first non-vanishing coefficient is still in front of λ^2 and equals to $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*, \nu_{0,\xi,\gamma}^*]$. Then we can proceed as in the higher dimensional cases. This completes the whole proof. \square

Remark A.1. In 3-dimension, as is computed in (A.3) and (A.6) in the Appendix, the $L^{\frac{3}{2}}$ -norm of the functions $x_j \partial_i \cdot_{g_{\mathbb{R}^3}} \nabla_{\partial_i} \psi_{1,0,\gamma}$ and $\partial_k \cdot_{g_{\mathbb{R}^3}} \psi_{1,0,\gamma}$ become infinite. This causes that $w_{\lambda,\xi,\gamma}^*$ has no more of order λ . Precisely, we see that the equation

$$\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[\nu_{0,\xi,\gamma}^*] = \frac{1}{2} \sum_{i=1}^2 (c_i x_i + c_3 x_3) \partial_i \cdot_{g_{\mathbb{R}^3}} \nabla_{\partial_i} \psi_{1,0,\gamma} - \frac{1}{2} c_3 \partial_3 \cdot_{g_{\mathbb{R}^3}} \psi_{1,0,\gamma}$$

is no longer defined in the dual space of $\mathcal{D}^{1/2}(\mathbb{R}^3, \mathbb{S}(\mathbb{R}^3))$ and we can only expand $w_{\lambda,\xi,\gamma}^*$ in the form

$$w_{\lambda,\xi,\gamma}^* = w_{0,\xi,\gamma}^* + \nu_{\lambda,\xi,\gamma}^*$$

with $\|\nu_{\lambda,\xi,\gamma}^*\| = O(\lambda |\ln \lambda|^{\frac{2}{3}})$. And hence $\nabla^2 \mathcal{J}_0(\psi_{1,0,\gamma})[w_{\lambda,\xi,\gamma}^*, w_{\lambda,\xi,\gamma}^*]$ has no more of order λ^2 . Comparing with the asymptotic expansion of $w_{\lambda,\xi,\gamma}^*$ for $m \geq 4$, it becomes more complicated to determine the exact asymptotic expansion of $w_{\lambda,\xi,\gamma}^*$ in dimension 3. Our approach in this regard up to now have failed.

References

- [1] R. Adams, Sobolev Spaces, Academic Press, New York 1975.
- [2] A. D. Alexandrov, Uniqueness theorems for surfaces in the large, Amer. Math. Soc. Transl. Ser. 2, 21, 441-519, (1962).
- [3] A. Ambrosetti, M. Badiale, Homoclinics: Poincaré-Melnikov type results via a variational approach, Ann. Inst. H. Poincaré Anal. Nonlinéaire 15 (1998), 233-252.
- [4] A. Ambrosetti, M. Badiale, Variational perturbative methods and bifurcation of bound states from the essential spectrum, Proc. Roy. Soc. Edinburgh Sect. A 128 (1998), 1131-1161.
- [5] A. Ambrosetti, J. Garcia Azorero, I. Peral, Perturbation of $\Delta u + u^{\frac{N+2}{N-2}} = 0$, the scalar curvature problem in \mathbb{R}^N , and related topics, J. Funct. Anal. 165 (1999), 117-149.
- [6] A. Ambrosetti, A. Malchiodi, A multiplicity result for the Yamabe problem on S^m , J. Funct. Anal. 168 (1999), 529-561.
- [7] A. Ambrosetti, A. Malchiodi, Perturbation Methods and Semilinear Elliptic Problems on \mathbb{R}^n , Progress in Math. 240, Birkhäuser, Basel-Boston-Berlin (2006).
- [8] B. Ammann, A variational problem in conformal spin geometry, Habilitationsschrift, Universität Hamburg 2003.

- [9] B. Ammann, The smallest Dirac eigenvalue in a spin-conformal class and cmc immersions, *Comm. Anal. Geom.* 17 (2009), no. 3, 429-479.
- [10] B. Ammann, M. Dahl, A. Hermann, E. Humbert, Mass endomorphism, surgery and perturbations, *Ann. Inst. Fourier.* 64 (2014), no. 2, 467-487.
- [11] B. Ammann, J.-F. Grossjean, E. Humbert, B. Morel, A spinorial analogue of Aubin's inequality, *Math. Z.* 260 (2008), 127-151.
- [12] B. Ammann, E. Humbert, B. Morel, Mass endomorphism and spinorial Yamabe type problems on conformally flat manifolds, *Comm. Anal. Geom.* 14 (2006), no. 1, 163-182.
- [13] A. Bahri, J. M. Coron, The scalar curvature problem on the standard three-dimensional sphere, *J. Funct. Anal.* 95 (1991), 106-172.
- [14] J.-P. Bourguignon, P. Gauduchon, Spineurs, opérateurs de Dirac et variations de métriques, *Commun. Math. Phys.* 144 (1992), 581-599.
- [15] K.-C. Chang, *Infinite Dimensional Morse Theory and Multiple Solution Problems*, Birkhäuser Boston (1993).
- [16] S.-Y. A. Chang, P. Yang, A perturbation result in prescribing scalar curvature on S^n , *Duke Math. J.* 64 (1991), 27-69.
- [17] S.-Y. A. Chang, M. Gursky, P. Yang, The scalar curvature equation on 2- and 3-spheres, *Calc. Var.* 1 (1993), 205-229.
- [18] T. Friedrich, On the spinor representation of surfaces in Euclidean 3-space, *J. Geom. Phys.* 28:1-2 (1998), 143-157.
- [19] T. Friedrich, *Dirac Operators in Riemannian Geometry*, Graduate Studies in Mathematics 25 (2000).
- [20] N. Ginoux, *The Dirac Spectrum*, Lecture Notes in Mathematics, vol. 1976. Springer, Berlin 2009.
- [21] K. Grosse-Brauckmann, *Complete embedded constant mean curvature surfaces*, Habilitationsschrift, Universität Bonn, 1998.
- [22] A. Hermann, Generic metrics and the mass endomorphism on spin three-manifolds, *Ann. Global Anal. Geom.* 37 (2010), 163-171.
- [23] O. Hijazi, A conformal lower bound for the smallest eigenvalue of the Dirac operator and Killing spinors, *Comm. Math. Phys.* 104 (1986), 151-162.
- [24] N. Hitchin, Harmonic spinors, *Adv. Math.* 14 (1974), 1-55.

- [25] H. Hopf, Differential geometry in the large. In: Lecture Notes in Mathematics, vol. 1000. Springer, Berlin. Notes taken by Peter Lax and John Gray. With a preface by S.S. Chern (1983).
- [26] T. Isobe, Nonlinear Dirac equations with critical nonlinearities on compact spin manifolds, *J. Funct. Anal.* 260 (2011), 253-307.
- [27] T. Isobe, A perturbation method for spinorial Yamabe type equations on S^m and its applications, *Math. Ann.* 355 (2013), 1255-1299.
- [28] T. Isobe, Spinorial Yamabe type equations on S^3 via Conley index, *Adv. Nonlinear. Stud.* 15(2015), 39-60.
- [29] K. Kenmotsu, Weierstrass formula for surfaces of prescribed mean curvature, *Math. Ann.* 245:2 (1979), 89-99.
- [30] B. G. Konopelchenko, Induced surfaces and their integrable dynamics, *Stud. Appl. Math.* 96:1 (1996), 9-51.
- [31] R. Kusner, N. Schmitt, The spinor representation of surfaces in space, dg-ga/9610005.
- [32] H. B. Lawson, M. L. Michelson, *Spin Geometry*, Princeton University Press (1989).
- [33] P. Li, S.T. Yau, A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue on compact surfaces, *Invent. Math.* 69 (1982), 269-291.
- [34] A. Maalaoui, Infinitely many solutions for the spinorial Yamabe problem on the round sphere, *Nonlinear Differ. Equ. Appl.* 23, 25 (2016).
- [35] A. Malchiodi, The scalar curvature problem on S_n : an approach via Morse theory, *Calc. Var.* 14 (2002), 429-445
- [36] S. Matsutani, Immersion anomaly of Dirac operator on surface in \mathbb{R}^3 , *Rev. Math. Phys.* 11:2 (1999), 171-186.
- [37] J. Mawhin, M. Willem, *Critical Point Theory and Hamiltonian Systems*, Applied Mathematical Sciences 74 (1989), Springer-Verlag.
- [38] A. Mondino, J. Schygulla, Existence of immersed spheres minimizing curvature functionals in non-compact 3-manifolds, *Ann. I.H.Poincaré* 31:4 (2014), 707-724.
- [39] S. Raulot, A Sobolev-like inequality for the Dirac operator, *J. Funct. Anal.* 26 (2009), 1588-1617.
- [40] Y. Sire, T. Xu, A variational analysis of the spinorial Yamabe equation on product manifolds, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* 24:5 (2023), 205-248.

- [41] M. Struwe, Variational Methods, *Ergeb. der Math. u. Grenzgeb.* Vol. 34, Springer 1996.
- [42] I. A. Taimanov, Modified Novikov-Veselov equation and differential geometry of surfaces, *Amer. Math. Soc. Transl. (2)* 179 (1997), 133-151; <http://arxiv.org/dg-ga/9511005>.
- [43] I. A. Taimanov, The Weierstrass representation of closed surfaces in \mathbb{R}^3 , *Funktsional. Anal. i Prilozhen.* 32:4 (1998), 49-62; English transl., *Funct. Anal. Appl.* 32 (1998), 258-267.
- [44] I. A. Taimanov, The Weierstrass representation of spheres in \mathbb{R}^3 , the Willmore numbers, and soliton spheres, *Trudy Mat. Inst. Steklov.* 225 (1999), 339-361; English transl., *Proc. Steklov Inst. Math.* 225 (1999), 225-243.