

---

# A STUDY ON $A_\alpha$ -SPECTRUM AND $A_\alpha$ -ENERGY OF UNITARY ADDITION CAYLEY GRAPHS

---

**Najiya V K, Chithra A V**  
 Department of Mathematics  
 National Institute of Technology, Calicut  
 Kerala, India-673601

najiya\_p190046ma@nitc.ac.in, chithra@nitc.ac.in

**Naveen Palanivel**  
 Department of Mathematics,  
 SASTRA Deemed to be University  
 Thanjavur, India-613401  
 naveenpalanivel.nitc@gmail.com

## ABSTRACT

The unitary addition Cayley graph  $G_n$ ,  $n \in \mathbb{Z}^+$  is the graph whose vertex set is  $\mathbb{Z}_n$ , the ring of integers modulo  $n$  and two vertices  $u$  and  $v$  are adjacent if and only if  $u + v \in \mathbb{U}_n$  where  $\mathbb{U}_n$  is the set of all units of the ring. The  $A_\alpha$ -matrix of a graph  $G$  is defined as  $A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$ ,  $\alpha \in [0, 1]$ , where  $D(G)$  is the diagonal matrix of vertex degrees and  $A(G)$  is the adjacency matrix of  $G$ . In this paper, we investigate the  $A_\alpha$ -eigenvalues for unitary addition Cayley graph and its complement. We determine bounds for  $A_\alpha$ -eigenvalues of unitary addition Cayley graph when its order is odd. Consequently, we compute the  $A_\alpha$ -energy of both  $G_n$  and its complement,  $\overline{G_n}$ , for  $n = p^m$  where  $p$  is a prime number and  $n$  even. Moreover, we obtain some bounds for energies of  $G_n$  and  $\overline{G_n}$  when  $n$  is odd. We also define  $A_\alpha$ -borderenergetic and  $A_\alpha$ -hyperenergetic graphs and observe some classes for each.

**Keywords** Spectrum of a graph, Energy of a graph, Unitary Addition Cayley Graph,  $A_\alpha$ -matrix

## 1 Introduction

A graph  $G = (V, E)$  consists of finite nonempty set  $V$  of  $n$  vertices together with a set  $E$  of  $m$  edges. The degree of a vertex  $v$  is the number of edges incident with it and is denoted as  $deg(v)$ . The complement  $\overline{G}$  of a graph  $G$  has  $V(G)$  as its vertex set, but two vertices are adjacent in  $\overline{G}$  if and only if they are not adjacent in  $G$ . Throughout the paper we consider only simple, undirected graphs.

Let  $\mathcal{G}$  be a group and let  $S$  be a subset of  $\mathcal{G}$  that is closed under taking inverses and does not contain the identity. Then the Cayley graph  $X(\mathcal{G}, S)$  is the graph with vertex set  $\mathcal{G}$  and edge set  $E = \{gh : hg^{-1} \in S\}$ .

In [2] Dejter et al. introduced the concept of unitary Cayley graph. Later in [14] Deepa et al. introduced the unitary addition Cayley graphs(UACG). Naveen et al. in [10, 8] studied the structural properties, adjacency spectrum and energy of UACG.

For a simple graph  $G$  of order  $n$ , the adjacency matrix  $A(G)$  of order  $n$  is a symmetric matrix with

$$a_{ij} = \begin{cases} 1 & \text{if } v_i \sim v_j \\ 0 & \text{otherwise.} \end{cases}$$

In [17] Nikiforov introduced a convex linear combination of the adjacency matrix of  $G$  and degree matrix of  $G$  called the  $A_\alpha$ -matrix. For  $\alpha \in [0, 1]$ ,  $A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$ , where  $D(G)$  is the diagonal matrix with entries as corresponding vertex degrees. Clearly,  $A_0(G) = A(G)$ ,  $A_1(G) = D(G)$  and  $Q(G) = 2A_{\frac{1}{2}}(G) = A(G) + D(G)$ , where  $Q(G)$  is the signless Laplacian matrix. Also  $A_\alpha(G) - A_\beta(G) = (\alpha - \beta)L(G) = (\alpha - \beta)(D(G) - A(G))$ , where  $L(G)$  is the Laplacian matrix of  $G$  and  $\beta \in [0, 1]$ .  $A_\alpha$ -matrix is a generalization of other matrices like  $A(G)$ ,  $Q(G)$ ,  $L(G)$  etc. Thus, it helps in studying spectral properties of those matrices as well as uncountable many convex combinations of  $D(G)$  and  $A(G)$ .

The characteristic polynomial of a real symmetric matrix  $M$  of order  $n$  is defined as  $|\lambda I - M|$  where  $I$  is the  $n$  order identity matrix. The eigenvalues of  $M$ ,  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ , together with their multiplicities gives the spectrum of  $M$ . The spectrum of a graph is the spectrum of the matrix considered. We represent the  $M$ -spectrum of the graph  $G$  as  $\text{spec}_M(G) = \begin{pmatrix} \lambda_1 & \lambda_2 & \dots & \lambda_t \\ m_1 & m_2 & \dots & m_t \end{pmatrix}$  where  $\lambda_i$ 's are the  $t$  distinct eigenvalues of the matrix  $M$  considered with multiplicity  $m_i$ . The roots of the characteristic polynomial of the adjacency matrix of a graph  $G$  are called the adjacency eigenvalues of  $G$ . Similarly, the roots of  $A_\alpha$ -matrix gives the  $A_\alpha$ -spectrum of  $G$ . If  $G$  is an  $r$ -regular graph, then its  $A_\alpha$ -eigenvalues are  $\alpha r + (1 - \alpha) \lambda_i(A(G))$ ,  $i = 1, 2, \dots, n$ . The sum of absolute values of the adjacency eigenvalues of a graph gives the adjacency energy,  $\varepsilon(G)$ , of the graph. The  $A_\alpha$ -energy of a graph  $G$ , for  $\alpha \in [0, 1)$ , is defined as  $\varepsilon_\alpha(G) = \sum_{i=1}^n \left| \lambda_i(A_\alpha(G)) - \frac{2\alpha m}{n} \right|$ ,  $1 \leq i \leq n$ . If  $G$  is regular, then  $\varepsilon_\alpha(G) = (1 - \alpha) \varepsilon(G)$ .

If every vertex of a graph is connected to every other vertices, then it is called a complete graph. Initially researchers believed that complete graphs have the maximum energy among all the graphs on  $n$  vertices. That is, if  $G \neq K_n$  is a graph on  $n$  vertices then  $\varepsilon(G) < 2n - 2$ . But this was disproved by Gutman and gave rise to the question whether there exist graphs whose energy equals or exceeds the energy of complete graph. In [5], Gutman introduced about graphs whose energy exceeds the energy of complete graphs and called it hyperenergetic graphs. Later in 2015, Gong et al. in [4] introduced the notion of borderenergetic graphs, that is the graphs on  $n$  vertices having energy same as that of  $K_n$ . In 2017, Tura in [15] gave the Laplacian analogous of borderenergetic graphs. Several families of borderenergetic graphs can be found in [4, 7, 3, 16].

The tensor product  $A \otimes B$  of two matrices  $A$  and  $B$  of order  $m \times n$  and  $p \times q$ , respectively, is the  $mp \times nq$  block matrix  $[a_{ij}B]$ . All one matrix, identity matrix and anti diagonal matrix, with non zero entries one, of order  $n$  are represented by  $J_n, I_n$  and  $\tilde{D}_n$ , respectively.

Consider a vector  $s = (s_0, s_1, \dots, s_{n-1}) \in \mathbb{R}^n$ . The left circulant matrix  $C_L(s)$  associated with  $s$  is

$$C_L(s) = \begin{bmatrix} s_0 & s_1 & \dots & s_{n-1} \\ s_1 & s_2 & \dots & s_0 \\ \vdots & \vdots & \ddots & \vdots \\ s_{n-1} & s_0 & \dots & s_{n-2} \end{bmatrix}$$

and the right circulant matrix  $C_R(s)$  associated with  $s$  is

$$C_R(s) = \begin{bmatrix} s_0 & s_1 & \dots & s_{n-1} \\ s_{n-1} & s_0 & \dots & s_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ s_1 & s_2 & \dots & s_0 \end{bmatrix}.$$

From [8], the eigenvalues of right circulant matrix are  $\lambda_j = s_0 + s_1\omega^j + s_2\omega^{2j} + \dots + s_{n-1}\omega^{(n-1)j}$ ,  $j = 0, 1, \dots, n-1$ , where  $\omega$  is a complex primitive  $n$ -th root of unity.

The eigenvalues of left circulant matrix are  $\lambda_0, \pm|\lambda_1|, \dots, \pm|\lambda_{(n-1)/2}|$  if  $n$  is odd and  $\lambda_0, \lambda_{n/2}, \pm|\lambda_1|, \dots, \pm|\lambda_{(n-2)/2}|$  if  $n$  is even where  $\lambda_k$  are the eigenvalues of right circulant matrix. For a positive integer  $n$  the Möbius function  $\mu(n)$  is defined as

$$\mu(n) = \begin{cases} 0, & \text{if } n \text{ has a squared prime factor} \\ (-1)^k, & \text{if } n \text{ is a product of } k \geq 0 \text{ distinct primes.} \end{cases}$$

For  $0 \leq k \leq n-1$ , the Ramanujan sum is  $c(k, n) = \mu(t_k) \frac{\phi(n)}{\phi(t_k)}$ , where  $t_k = \frac{n}{\gcd(k, n)}$  and  $\phi(n)$  is the Euler's phi function which is the number of positive integers  $\leq n$  which are relatively prime to  $n$ . Also, the eigenvalues of unitary Cayley graph  $X_n$  are  $\lambda_i(X_n) = \sum_{\substack{1 \leq j < n \\ \gcd(j, n)=1}} \omega^{kj} = c(k, n)$ , where  $\omega$  is a complex primitive  $n$ -th root of unity.

Let  $\mathbb{M}_{m \times n}(\mathbb{R})$  be the set of all  $m \times n$  matrices with real entries. The Frobenius norm for  $M \in \mathbb{M}_{m \times n}(\mathbb{R})$  is defined as

$$\|M\|_F = \sqrt{\sum_{i=1}^m \sum_{j=1}^n |m_{ij}|^2} = \sqrt{\text{trace}(M^T M)},$$

where trace of a square matrix is defined as sum of the elements on the main diagonal. Let  $\lambda_i$ 's be the eigenvalues of the matrix  $M$ . If  $MM^T = M^T M$ , then  $\|M\|_F^2 = \sum_{i=1}^n |\lambda_i(M)|^2$ . The Zagreb index  $\zeta(G)$  of a graph  $G$  is defined as the sum of the squares of the vertices degrees, that is,  $\zeta(G) = \sum_{v \in V(G)} \deg(v)^2$ .

In real world, modelling a problem using graph gives a regular graph or a non-regular graph. In literature, there are more studies happened in regular graphs compared to the non-regular graph. The study of non-regular graph thus has an significant role in vast area of science and technologies. The UACG  $G_n$  is a non-regular graph for odd values of  $n$ . In this paper, the exact  $A_\alpha$ -eigenvalues of UACG and its complement are found for the values  $n = p^m$  and  $n$  even. For the case when  $n$  is odd, it's difficult to find an exact value, but we have found a bound for  $A_\alpha$ -eigenvalues of  $G_n$  and its complement. From the  $A_\alpha$ -eigenvalues, the  $A_\alpha$ -energy for  $G_n$  and  $\overline{G_n}$  are also computed for the values  $n = p^m$  and  $n$  even. We also introduce the concept of  $A_\alpha$ -borderenergetic and  $A_\alpha$ -hyperenergetic graphs and found some graphs which are  $A_\alpha$ -borderenergetic and  $A_\alpha$ -hyperenergetic.

## 2 Preliminaries

**Definition 2.1.** [8] The unitary Cayley graph  $X_n = \text{Cay}(Z_n, \cup_n)$  is the graph with vertex set  $Z_n$ , the integers modulo  $n$  and  $u, v \in Z_n$  are adjacent if and only if  $u - v \in \cup_n$ , where  $\cup_n$  represents set of all units of the ring  $Z_n$ . The graph  $X_n$  is regular with degree  $|\cup_n| = \phi(n)$ .

**Definition 2.2.** [14] For a positive integer  $n > 1$ , the unitary addition Cayley graph(UACG)  $G_n = \text{Cay}^+(Z_n, \cup_n)$  is the graph whose vertex set is  $Z_n = \{0, 1, 2, \dots, n - 1\}$  and the edge set  $E(G_n) = \{ab \mid a, b \in Z_n, a + b \in \cup_n\}$  where  $\cup_n = \{a \in Z_n \mid \gcd(a, n) = 1\}$ . If  $n$  is even, then  $G_n$  is regular and if  $n$  is odd, then  $G_n$  is semi regular .




Figure 1: Unitary addition Cayley graph  $G_9$

**Theorem 2.1.** [14] The UACG  $G_n$  is isomorphic to  $X_n$ , the unitary Cayley graph, if and only if  $n$  is even.

**Theorem 2.2.** [14] Let  $v_i, i = 0, 1, 2, \dots, n - 1$ , be any vertex of the UACG  $G_n$ . Then

$$\deg(v_i) = \begin{cases} \phi(n) & \text{if } n \text{ is even} \\ \phi(n) & \text{if } n \text{ is odd and } \gcd(i, n) \neq 1 \\ \phi(n) - 1 & \text{if } n \text{ is odd and } \gcd(i, n) = 1 \end{cases}$$

**Corollary 2.3.** [14] The total number of edges in the UACG  $G_n$  is

$$|E(G_n)| = \begin{cases} \frac{1}{2}n\phi(n) & \text{if } n \text{ is even} \\ \frac{1}{2}(n-1)\phi(n) & \text{if } n \text{ is odd} \end{cases}$$

**Theorem 2.4.** [6] The energy of unitary Cayley graph  $X_n$  equals  $2^l \phi(n)$ , where  $l$  is the total number of unique prime factors that divides  $n$ .

**Theorem 2.5.** [6] *The complement of unitary Cayley graph  $X_n$  has energy*

$$\varepsilon(\overline{X_n}) = 2(n-1) + (2^k - 2)\phi(n) - \prod_{i=1}^k p_i + \prod_{i=1}^k (2 - p_i),$$

where  $\prod_{i=1}^k p_i$  is the largest divisor of  $n$  that is square-free.

**Theorem 2.6.** [9] *Let  $M, M_1, M_2$  be three real symmetric matrices of order  $n$  such that  $M = M_1 + M_2$ . Then the eigenvalues of these matrices satisfy the following inequalities: for  $1 \leq i \leq n$  and  $0 \leq j \leq \min\{i-1, n-i\}$ ,*

$$\lambda_{i-j}(M_1) + \lambda_{1+j}(M_2) \geq \lambda_i(M) \geq \lambda_{i+j}(M_1) + \lambda_{n-j}(M_2).$$

**Theorem 2.7.** [11] *Let  $G$  be a connected graph with  $n \geq 3$  vertices,  $m$  edges, maximum degree  $\Delta$  and Zagreb index  $\zeta(G)$ . Let  $\alpha \in [0, 1)$ . Then*

$$1. \varepsilon_\alpha(G) \geq \sqrt{2 \left( \alpha^2 \zeta(G) + (1-\alpha)^2 \|A(G)\|_F^2 - \frac{2(\alpha m)^2}{n} \right)}.$$

$$2. \varepsilon_\alpha(G) \geq 4(1-\alpha) \frac{m}{n}.$$

*Equality occurs if and only if  $G$  is a regular graph with one positive and  $n-1$  negative adjacency eigenvalues.*

$$3. \varepsilon_\alpha(G) \geq 2\sqrt{\frac{\zeta(G)}{n} - \frac{4\alpha m}{n}}.$$

*Equality occurs if and only if  $G$  is a regular graph with one positive and  $n-1$  negative adjacency eigenvalues.*

$$4. \varepsilon_\alpha(G) \geq \left( \alpha(\Delta+1) + \sqrt{\alpha^2(\Delta+1)^2 + 4\Delta(1-2\alpha)} \right) - \frac{4\alpha m}{n}$$

*with equality if and only if  $G \cong K_{1,\Delta}$ .*

**Theorem 2.8.** [13] *Let  $G$  be a graph on  $n$  vertices,  $m$  edges and Zagreb index  $\zeta(G)$ . For  $\alpha \in [0, 1)$ , we have*

$$\varepsilon_\alpha(G) \leq \sqrt{n(\alpha^2 \zeta(G) + (1-\alpha)^2 \|A(G)\|_F^2) - 4\alpha^2 m^2}$$

**Lemma 2.9.** [19] *Let  $M_1, M_2, M_3$  and  $M_4$  be matrices with  $M_1$  invertible. Let*

$$\widetilde{M} = \begin{pmatrix} M_1 & M_2 \\ M_3 & M_4 \end{pmatrix}.$$

Then  $\det(\widetilde{M}) = \det(M_1) \det(X - M_3 M_1^{-1} M_2)$ .

If  $M_4$  is invertible, then  $\det(\widetilde{M}) = \det(M_4) \det(M_1 - M_2 M_4^{-1} M_3)$ .

**Definition 2.3.** [1] *The  $M$ -coronal  $\Gamma_M(x)$  of a matrix  $M$  of order  $n$  is defined as the sum of the entries of the matrix  $(xI_n - M)^{-1}$  (if it exists), that is,*

$$\Gamma_M(x) = J_{n \times 1}^T (xI_n - M)^{-1} J_{n \times 1}.$$

If each row sum of an  $n \times n$  matrix  $M$  is constant, say  $t$ , then  $\Gamma_M(x) = \frac{n}{x-t}$ .

**Lemma 2.10.** [18] *Let  $A$  be a real matrix of order  $n$ . Then*

$$\det(yI_n - A - cJ_n) = (1 - c\Gamma_A(y)) \det(yI_n - A).$$

**Lemma 2.11.** [12] *Let  $A$  and  $B$  be symmetric matrices of order  $n$  and  $m$ , respectively. If  $\mu_1, \dots, \mu_n$  and  $\lambda_1, \dots, \lambda_m$  are the eigenvalues of  $A$  and  $B$ , respectively, then the eigenvalues of  $A \otimes B$  are given by  $\mu_j \lambda_i; j = 1, \dots, n; i = 1, \dots, m$ .*

### 3 $A_\alpha$ -spectrum and energy of unitary addition Cayley graphs

This section discusses the  $A_\alpha$ -spectrum and  $A_\alpha$ -energy of UACG for even  $n$  and  $n = p^m$  where  $p$  is a prime number. Also, bound for  $A_\alpha$ -eigenvalues and  $A_\alpha$ -energy are found for odd  $n$ .

**Proposition 3.1.** Let  $n = p^m$ ,  $p \geq 3$ ,  $m \geq 1$ . Then the  $A_\alpha$ -spectrum of UACG  $G_n$  is

$$\left( \begin{array}{ccc} p^{m-1}(p\alpha - 1) - 1 & \frac{x-y}{2} & p^{m-1}(p-1)\alpha - 1 \\ \frac{p-3}{2} & 1 & (p-1)(p^{m-1} - 1) \end{array} \quad \begin{array}{cc} \alpha(p^m - p^{m-1}) & p^{m-1}((p-2)\alpha + 1) - 1 \\ p^{m-1} - 1 & \frac{p-1}{2} \end{array} \quad \begin{array}{c} \frac{x+y}{2} \\ 1 \end{array} \right),$$

where  $x = (1 + \alpha)p^m - 2p^{m-1} - 1$  and  $y = \sqrt{(1 - p^m + \alpha p^m)^2 + (1 - \alpha)4p^{m-1}}$ .

*Proof.* The  $A_\alpha$ -matrix of UACG of order  $p^m$  after proper labelling is  $A_\alpha(G_{p^m}) = J_{p^{m-1}} \otimes C - I_{p^{m-1}} \otimes (C - B)$ , where  $B = \begin{bmatrix} \alpha\phi(p^m) & (1 - \alpha)J_{1 \times p-1} \\ (1 - \alpha)J_{p-1 \times 1} & (1 - \alpha)(J - \tilde{D}) - (1 - \alpha\phi(p^m))I_{p-1} \end{bmatrix}$  and  $C = (1 - \alpha) \begin{bmatrix} 0 & J_{1 \times p-1} \\ J_{p-1 \times 1} & J_{p-1} - \tilde{D} \end{bmatrix}$ .

The matrix  $A_\alpha(G_{p^m})$  is permutationally similar to

$$\hat{A}_\alpha(G_{p^m}) = \begin{bmatrix} \alpha\phi(p^m)I_{p^{m-1}} & (1 - \alpha)J_{p^{m-1} \times \frac{p^m - p^{m-1}}{2}} & (1 - \alpha)J_{p^{m-1} \times \frac{p^m - p^{m-1}}{2}} \\ (1 - \alpha)J_{\frac{p^m - p^{m-1}}{2} \times p^{m-1}} & (1 - \alpha)J + (\alpha\phi(p^m) - 1)I & (1 - \alpha)J - (1 - \alpha)\tilde{D} \otimes J \\ (1 - \alpha)J_{\frac{p^m - p^{m-1}}{2} \times p^{m-1}} & (1 - \alpha)J - (1 - \alpha)\tilde{D} \otimes J & (1 - \alpha)J + (\alpha\phi(p^m) - 1)I \end{bmatrix},$$

where  $\tilde{D}$  is the anti-diagonal matrix with non-zero entries one of order  $\frac{p-1}{2}$ .

Then  $A_\alpha(G_{p^m}) = P\hat{A}_\alpha(G_{p^m})P^{-1}$ ,

where  $P = [A_{11}^T \ A_{12}^T \ \cdots \ A_{1p^{m-2}}^T \ A_{21}^T \ A_{22}^T \ \cdots \ A_{2p^{m-2}}^T \ \cdots \ A_{p1}^T \ A_{p2}^T \ \cdots \ A_{pp^{m-2}}^T]$  is a permutation matrix of order  $p^m$ ,  $m \geq 2$ ,

$$A_{ij} = [a_{st}], a_{st} = \begin{cases} 1 & \text{if } (s, t) \in H_{ij} \\ 0 & \text{otherwise} \end{cases}$$

$s = 1, 2, \dots, p; t = 1, 2, \dots, p^m$  and

$H_{ij} = \{(1, i + (j - 1)p), (2, i + (j - 1)p + p^{m-1}), \dots, (p, i + (j - 1)p + (p - 1)p^{m-1})\}$ ,

$i = 1, 2, \dots, p, j = 1, 2, \dots, p^{m-2}$ .

$P$  becomes the identity matrix when  $m = 1$ .

$$|\lambda I - \hat{A}_\alpha(G_{p^m})| =$$

$$\begin{vmatrix} (\lambda - \alpha\phi(p^m))I_{p^{m-1}} & -(1 - \alpha)J_{p^{m-1} \times \frac{p^m - p^{m-1}}{2}} & -(1 - \alpha)J_{p^{m-1} \times \frac{p^m - p^{m-1}}{2}} \\ -(1 - \alpha)J_{\frac{p^m - p^{m-1}}{2} \times p^{m-1}} & (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J & -(1 - \alpha)J + (1 - \alpha)\tilde{D} \otimes J \\ -(1 - \alpha)J_{\frac{p^m - p^{m-1}}{2} \times p^{m-1}} & -(1 - \alpha)J + (1 - \alpha)\tilde{D} \otimes J & (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J \end{vmatrix}.$$

By Lemma 2.9,

$$|\lambda I - \hat{A}_\alpha(G_{p^m})| = (\lambda - \alpha\phi(p^m))^{p^{m-1}} \times \det S,$$

$$\text{where } S = \begin{bmatrix} (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J & -(1 - \alpha)J + (1 - \alpha)\tilde{D} \otimes J \\ -(1 - \alpha)J + (1 - \alpha)\tilde{D} \otimes J & (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J \end{bmatrix} - \frac{p^{m-1}(1 - \alpha)^2}{\lambda - \alpha\phi(p^m)} J_{p^m - p^{m-1}}$$

$$\det S = \left| (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J - (1 - \alpha)J + (1 - \alpha)\tilde{D} \otimes J - \frac{2p^{m-1}(1 - \alpha)^2}{\lambda - \alpha\phi(p^m)} J \right| \\ \times \left| (\lambda - \alpha\phi(p^m) + 1)I - (1 - \alpha)J + (1 - \alpha)J - (1 - \alpha)\tilde{D} \otimes J \right|.$$

Using Lemma 2.10 and Definition 2.3, we get

$$\begin{aligned} \det S &= \left( 1 - 2 \left( 1 - \alpha + \frac{p^{m-1}(1-\alpha)^2}{\lambda - \alpha\phi(p^m)} \right) \Gamma_{-(1-\alpha)\tilde{D} \otimes J}(\lambda - \alpha\phi(p^m) + 1) \right) \\ &\quad \times \left| (\lambda - \alpha\phi(p^m) + 1)I + (1-\alpha)\tilde{D} \otimes J \right| \left| (\lambda - \alpha\phi(p^m) + 1)I - (1-\alpha)\tilde{D} \otimes J \right| \\ &= \left( 1 - \left( 1 - \alpha + \frac{p^{m-1}(1-\alpha)^2}{\lambda - \alpha\phi(p^m)} \right) \frac{p^m - p^{m-1}}{\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1}} \right) \\ &\quad \left| (\lambda - \alpha\phi(p^m) + 1)^2 I - (1-\alpha)^2 (\tilde{D} \otimes J)^2 \right|. \end{aligned}$$

Therefore,

$$\begin{aligned} |\lambda I - \hat{A}_\alpha(G_{p^m})| &= (\lambda - \alpha\phi(p^m))^{p^{m-1}} \left( 1 - \left( 1 - \alpha + \frac{p^{m-1}(1-\alpha)^2}{\lambda - \alpha\phi(p^m)} \right) \frac{p^m - p^{m-1}}{\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1}} \right) \\ &\quad \left| (\lambda - \alpha\phi(p^m) + 1)^2 I - (1-\alpha)^2 (\tilde{D} \otimes J)^2 \right| \\ &= (\lambda - \alpha\phi(p^m))^{p^{m-1}} \left( 1 - \frac{\left( (1-\alpha)(\lambda - \alpha\phi(p^m)) + p^{m-1}(1-\alpha)^2 \right) (p^m - p^{m-1})}{(\lambda - \alpha\phi(p^m))(\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1})} \right) \\ &\quad \prod_{i=1}^{\frac{p-1}{2}} \left( (\lambda - \alpha\phi(p^m) + 1)^2 - (1-\alpha)^2 (p^{m-1}\mu_i)^2 \right) (\lambda - \alpha\phi(p^m) + 1)^{p^m - p^{m-1} - p + 1}, \end{aligned}$$

where  $\mu_i$ 's are the eigenvalues of  $\tilde{D}$  and they are  $\pm 1$ .

$$\begin{aligned} |\lambda I - \hat{A}_\alpha(G_{p^m})| &= (\lambda - \alpha\phi(p^m))^{p^{m-1}} \left( 1 - \frac{\left( (1-\alpha)(\lambda - \alpha\phi(p^m)) + p^{m-1}(1-\alpha)^2 \right) (p^m - p^{m-1})}{(\lambda - \alpha\phi(p^m))(\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1})} \right) \\ &\quad \left( (\lambda - \alpha\phi(p^m) + 1)^2 - (1-\alpha)^2 (p^{m-1})^2 \right)^{\frac{p-1}{2}} (\lambda - \alpha\phi(p^m) + 1)^{p^m - p^{m-1} - p + 1} \\ &= (\lambda - \alpha\phi(p^m))^{p^{m-1} - 1} (\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1})^{\frac{p-3}{2}} (\lambda - \alpha\phi(p^m) + 1 - (1-\alpha)p^{m-1})^{\frac{p-1}{2}} \\ &\quad \left( (\lambda - \alpha\phi(p^m))(\lambda - \alpha\phi(p^m) + 1 + (1-\alpha)p^{m-1}) - (1-\alpha) \left( (\lambda - \alpha\phi(p^m)) + p^{m-1}(1-\alpha) \right) (p^m - p^{m-1}) \right) \\ &\quad (\lambda - \alpha\phi(p^m) + 1)^{p^m - p^{m-1} - p + 1}. \end{aligned}$$

Thus, the  $A_\alpha$ -spectrum of  $G_{p^m}$  is

$\text{spec}_{A_\alpha}(G_{p^m}) =$

$$\left( \begin{array}{cccccc} p^{m-1}(p\alpha - 1) - 1 & \frac{x-y}{2} & p^{m-1}(p-1)\alpha - 1 & \alpha(p^m - p^{m-1}) & p^{m-1}((p-2)\alpha + 1) - 1 & \frac{x+y}{2} \\ \frac{p-3}{2} & 1 & (p-1)(p^{m-1}-1) & p^{m-1}-1 & \frac{p-1}{2} & 1 \end{array} \right),$$

where  $x = (1 + \alpha)p^m - 2p^{m-1} - 1$  and  $y = \sqrt{(1 - p^m + \alpha p^m)^2 + (1 - \alpha)4p^{m-1}}$ .

□

Next we calculate the  $A_\alpha$ -energy of UACG  $G_n$  for  $n = p^m$ .

**Corollary 3.1.** The  $A_\alpha$ -energy of UACG  $G_{p^m}$ ,  $p \geq 3$  is

$$\varepsilon_\alpha(G_{p^m}) = \frac{3p^m - 5p^{m-1} - p - 1}{2} + \frac{\alpha}{2p} (-3p^{m+1} + 9p^m - 4p^{m-1} + p^2 - 2p + 1) + \frac{p-1}{2p} |(p^m - p)(1 - \alpha) - \alpha| + \left| \frac{x-y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| + \left| \frac{x+y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right|,$$

where  $x = (1 + \alpha)p^m - 2p^{m-1} - 1$  and  $y = \sqrt{(1 - p^m + \alpha p^m)^2 + (1 - \alpha)4p^{m-1}}$ .

*Proof.* The number of vertices and edges of  $G_{p^m}$  is  $p^m$  and  $\frac{(p^m-1)(p^m-p^{m-1})}{2}$  respectively. Thus,

$$\begin{aligned} \varepsilon_\alpha(G_{p^m}) &= \frac{p-3}{2} \left| p^{m-1}(p\alpha - 1) - 1 - \frac{\alpha(p^m - 1)(p-1)}{p} \right| + \left| \frac{x-y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &\quad + (p-1)(p^{m-1} - 1) \left| p^{m-1}(p-1)\alpha - 1 - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &\quad + (p^{m-1} - 1) \left| \alpha(p^m - p^{m-1}) - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &\quad + \frac{p-1}{2} \left| p^{m-1}((p-2)\alpha + 1) - 1 - \frac{\alpha(p^m - 1)(p-1)}{p} \right| + \left| \frac{x+y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &= \frac{p-3}{2} \left( -p^{m-1}(p\alpha - 1) + 1 + \frac{\alpha(p^m - 1)(p-1)}{p} \right) \\ &\quad + (p-1)(p^{m-1} - 1) \left( -p^{m-1}(p-1)\alpha + 1 + \frac{\alpha(p^m - 1)(p-1)}{p} \right) \\ &\quad + (p^{m-1} - 1) \left( \alpha(p^m - p^{m-1}) - \frac{\alpha(p^m - 1)(p-1)}{p} \right) + \left| \frac{x-y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &\quad + \frac{p-1}{2} \left| p^{m-1}((p-2)\alpha + 1) - 1 - \frac{\alpha(p^m - 1)(p-1)}{p} \right| + \left| \frac{x+y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &= \frac{3p^m - 5p^{m-1} - p - 1}{2} + \frac{\alpha}{2p} (-3p^{m+1} + 9p^m - 4p^{m-1} + p^2 - 2p + 1) \\ &\quad + \frac{p-1}{2p} |(p^m - p)(1 - \alpha) - \alpha| + \left| \frac{x-y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right| \\ &\quad + \left| \frac{x+y}{2} - \frac{\alpha(p^m - 1)(p-1)}{p} \right|, \end{aligned}$$

where  $x = (1 + \alpha)p^m - 2p^{m-1} - 1$  and  $y = \sqrt{(1 - p^m + \alpha p^m)^2 + (1 - \alpha)4p^{m-1}}$ . □

**Remark 3.1.** If  $n$  is even, then  $G_n$  is regular and its  $A_\alpha$ -spectrum will be  $\alpha\phi(n) + (1 - \alpha)\lambda_i(A(G_n))$ ,  $i = 1, 2, \dots, n$ . Its  $A_\alpha$ -energy will be  $\varepsilon_\alpha(G_n) = 2^k(1 - \alpha)\phi(n)$ , where  $k$  is the number of distinct prime factors dividing  $n$  and  $\alpha \in [0, 1)$ .

In general, computing exact eigenvalues of UACG for odd values of  $n$  is a tedious work. But, in the following proposition, we determine a bound for  $A_\alpha$ -eigenvalues of UACG  $G_n$  when  $n$  is odd.

**Proposition 3.2.** For an odd integer  $n$ , the  $A_\alpha$ -spectrum of the UACG  $G_n$  satisfy the following inequalities:

$$(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha\phi(n) - 1 \leq \lambda_k(A_\alpha(G_n)) \leq (1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha\phi(n)$$

for  $1 \leq k \leq \frac{(n+1)}{2}$  and

$$-(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha\phi(n) - 1 \leq \lambda_k(A_\alpha(G_n)) \leq -(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha\phi(n)$$

for  $\frac{(n+3)}{2} \leq k \leq n$ .

*Proof.* Let  $A_\alpha = (a_{ij})$ ,  $0 \leq i, j \leq n-1$ , be the  $A_\alpha$  matrix of  $G_n$ , where

$$a_{ij} = \begin{cases} 1 - \alpha & \text{if } \gcd(n, i+j) = 1 \text{ and } i \neq j, \\ \alpha\phi(n) & \text{if } \gcd(n, i+j) \neq 1 \text{ and } i = j, \\ \alpha(\phi(n) - 1) & \text{if } \gcd(n, i+j) = 1 \text{ and } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Then  $A_\alpha = B + C$ , where  $B = [b_{ij}]$ ,  $0 \leq i, j \leq n-1$ ,

$$b_{ij} = \begin{cases} 1 - \alpha & \text{if } \gcd(n, i+j) = 1, \\ 0 & \text{otherwise,} \end{cases}$$

and  $C = [c_{ij}]$ ,  $0 \leq i, j \leq n-1$ ,

$$c_{ij} = \begin{cases} \alpha\phi(n) & \text{if } \gcd(n, i+j) \neq 1 \text{ and } i = j, \\ \alpha\phi(n) - 1 & \text{if } \gcd(n, i+j) = 1 \text{ and } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

According to the definition of  $B$ , it is a left circulant matrix with initial row  $(s_0, s_1, \dots, s_{n-1})$  where

$$s_j = \begin{cases} 1 - \alpha & \text{if } \gcd(j, n) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Thus eigenvalues of  $B$  are  $(1 - \alpha)\mu(t_k)\frac{\phi(n)}{\phi(t_k)}$ , for  $0 \leq k \leq \frac{(n-1)}{2}$  and  $-(1 - \alpha)\mu(t_k)\frac{\phi(n)}{\phi(t_k)}$  for  $\frac{(n+1)}{2} \leq k \leq n-1$ .

Eigenvalues of  $C$  are  $\alpha\phi(n)$  repeated  $n - \phi(n)$  times and  $\alpha\phi(n) - 1$  repeated  $\phi(n)$  times.

The required result is therefore obtained from the eigenvalues of  $B, C$  and Theorem 2.6.  $\square$

In the following proposition, we estimate some bounds for  $A_\alpha$ -energy of UACG  $G_n$  when  $n$  is odd.

**Proposition 3.3.** For an odd integer  $n \geq 3$  and  $\alpha \in [0, 1)$ ,

$$\epsilon_\alpha(G_n) \geq \sqrt{\frac{\phi(n)}{n} [\alpha^2 (\phi(n) (3n^2 - 4n - 1) + 2n) + (1 - 2\alpha)(n - 1)2n\phi(n)].}$$

$$\epsilon_\alpha(G_n) \geq \frac{2(1 - \alpha)(n - 1)\phi(n)}{n}.$$

$$\epsilon_\alpha(G_n) \geq \sqrt{\frac{\phi(n)(\phi(n)(n - 2) + 1)}{n}} - \frac{4\alpha(n - 1)\phi(n)}{n}.$$

$$\epsilon_\alpha(G_n) \geq \alpha(\phi(n) + 1) + \sqrt{\alpha^2(\phi(n) + 1)^2 + 4\phi(n)(1 - \alpha)} - \frac{2\alpha(n - 1)\phi(n)}{n}.$$

$$\epsilon_\alpha(G_n) \leq \sqrt{\phi(n)(n^2(1 - \alpha)^2 - n + 2\alpha n - \alpha^2\phi(n))}.$$

*Proof.* The Zagreb index of UACG is  $\phi(n)^2(n - 2) + \phi(n)$  and  $\|A(G_n)\|_F^2 = (n - 1)\phi(n)$ . Then from Theorems 2.7 and 2.8 the result follows.  $\square$

#### 4 $A_\alpha$ -spectrum and energy of complement of unitary addition Cayley graphs

In this section, we compute the  $A_\alpha$ -spectrum and  $A_\alpha$ -energy of complement of UACG for even  $n$  and  $n = p^m$ . Also we find bounds for  $A_\alpha$ -eigenvalues and  $A_\alpha$ -energy for odd  $n$ .

**Proposition 4.1.** Let  $n = p^m$ ,  $p \geq 3$ ,  $m \geq 1$ . Then the  $A_\alpha$ -spectrum of complement of UACG  $G_n$  is

$$\left( \begin{array}{cccccc} -p^{m-1} & \alpha p^{m-1} - 1 & \alpha p^{m-1} & p^{m-1} - 1 & p^{m-1} \\ \frac{p-1}{2} & p^{m-1} - 1 & (p-1)(p^{m-1} - 1) & 1 & \frac{p-1}{2} \end{array} \right).$$

*Proof.* The  $A_\alpha$ -matrix of complement of UACG  $G_{p^m}$  after proper labelling is  $A_\alpha(\overline{G_{p^m}}) = J_{p^{m-1}} \otimes C - I_{p^{m-1}} \otimes (C - B)$ ,

where  $B = \begin{bmatrix} \alpha(1-p^{m-1}) & 0_{1 \times p-1} \\ 0_{p-1 \times 1} & (1-\alpha)\tilde{D}_{p-1} - \alpha p^{m-1}I \end{bmatrix}$  and  $C = (1-\alpha) \begin{bmatrix} 1 & 0_{1 \times p-1} \\ 0_{p-1 \times 1} & \tilde{D}_{p-1} \end{bmatrix}$ .

The matrix  $A_\alpha(\overline{G_{p^m}})$  is permutationally similar to

$$\hat{A}_\alpha(\overline{G_{p^m}}) = \begin{bmatrix} (1-\alpha)J_{p^{m-1}} - (1-\alpha p^{m-1})I_{p^{m-1}} & 0_{p^{m-1} \times \frac{p^m-p^{m-1}}{2}} & 0_{p^{m-1} \times \frac{p^m-p^{m-1}}{2}} \\ 0_{\frac{p^m-p^{m-1}}{2} \times p^{m-1}} & \alpha p^{m-1}I & (1-\alpha)\tilde{D} \otimes J \\ 0_{\frac{p^m-p^{m-1}}{2} \times p^{m-1}} & (1-\alpha)\tilde{D} \otimes J & \alpha p^{m-1}I \end{bmatrix},$$

where  $\tilde{D}$  is the anti-diagonal matrix of order  $\frac{p-1}{2}$  with all non-zero entries one.

Then  $A_\alpha(\overline{G_{p^m}}) = P\hat{A}_\alpha(\overline{G_{p^m}})P^{-1}$ , where  $P$  is the same matrix as used in proof of Proposition 3.1.

$$|\lambda I - \hat{A}_\alpha(\overline{G_{p^m}})| = \begin{vmatrix} (\lambda + 1 - \alpha p^{m-1})I_{p^{m-1}} - (1-\alpha)J_{p^{m-1}} & 0_{p^{m-1} \times \frac{p^m-p^{m-1}}{2}} & 0_{p^{m-1} \times \frac{p^m-p^{m-1}}{2}} \\ 0_{\frac{p^m-p^{m-1}}{2} \times p^{m-1}} & (\lambda - \alpha p^{m-1})I & -(1-\alpha)\tilde{D} \otimes J \\ 0_{\frac{p^m-p^{m-1}}{2} \times p^{m-1}} & -(1-\alpha)\tilde{D} \otimes J & (\lambda - \alpha p^{m-1})I \end{vmatrix}.$$

By Lemma 2.9,

$$\begin{aligned} |\lambda I - \hat{A}_\alpha(\overline{G_{p^m}})| &= |(\lambda + 1 - \alpha p^{m-1})I - (1-\alpha)J| \begin{vmatrix} (\lambda - \alpha p^{m-1})I & -(1-\alpha)\tilde{D} \otimes J \\ -(1-\alpha)\tilde{D} \otimes J & (\lambda - \alpha p^{m-1})I \end{vmatrix} \\ &= (\lambda + 1 - p^{m-1})(\lambda + 1 - \alpha p^{m-1})^{p^{m-1}-1} (\lambda - \alpha p^{m-1})^{p^m - p^{m-1} - p + 1} \\ &\quad \left( (\lambda - \alpha p^{m-1})^2 - (1-\alpha)^2 p^{2m-2} \right)^{\frac{p-1}{2}}. \end{aligned}$$

Thus, the  $A_\alpha$ -spectrum of  $\overline{G_{p^m}}$  is

$$\text{spec}_{A_\alpha}(\overline{G_{p^m}}) = \left( -\frac{p^{m-1}}{2}, \alpha p^{m-1} - 1, (p-1)(p^{m-1} - 1), p^{m-1} - 1, \frac{p^{m-1}}{2} \right).$$

□

Using Proposition 4.1 we now calculate the  $A_\alpha$ -energy of complement of UACG  $G_n$  for  $n = p^m$ .

**Corollary 4.1.** The  $A_\alpha$ -energy of complement of UACG  $G_{p^m}$ ,  $p \geq 3$  is

$$\varepsilon_\alpha(\overline{G_{p^m}}) = \frac{p^{m+1} - p + \alpha(2 - p - 2p^{m-1} + p^m)}{p} + \left| \frac{p^m - p - \alpha p^m + \alpha}{p} \right|.$$

*Proof.* The number of vertices and edges of  $\overline{G_{p^m}}$  is  $p^m$  and  $\frac{p^{m-1}(p^m-1)}{2}$  respectively. Thus,

$$\begin{aligned} \varepsilon_\alpha(\overline{G_{p^m}}) &= \left( \frac{p-1}{2} \right) \left| -p^{m-1} - \frac{\alpha(p^m-1)}{p} \right| + (p^{m-1}-1) \left| \alpha p^{m-1} - 1 - \frac{\alpha(p^m-1)}{p} \right| \\ &\quad + (p-1)(p^{m-1}-1) \left| \alpha p^{m-1} - \frac{\alpha(p^m-1)}{p} \right| + \left| p^{m-1} - 1 - \frac{\alpha(p^m-1)}{p} \right| \\ &\quad + \left( \frac{p-1}{2} \right) \left| p^{m-1} - \frac{\alpha(p^m-1)}{p} \right| \\ &= \left( \frac{p-1}{2} \right) \left( p^{m-1} + \frac{\alpha(p^m-1)}{p} \right) + (p^{m-1}-1) \left( -\alpha p^{m-1} + 1 + \frac{\alpha(p^m-1)}{p} \right) \\ &\quad + (p-1)(p^{m-1}-1)\alpha + \left| p^{m-1} - 1 - \frac{\alpha(p^m-1)}{p} \right| + \left( \frac{p-1}{2} \right) \left( \frac{p^m - \alpha p^m + \alpha}{p} \right) \\ &= \frac{p^{m+1} - p + \alpha(2 - p - 2p^{m-1} + p^m)}{p} + \left| \frac{p^m - p - \alpha p^m + \alpha}{p} \right| \\ &= \begin{cases} \frac{p^{m+1} + p^m - 2p + \alpha(3 - p - 2p^{m-1})}{p} & \text{if } \alpha \leq \frac{p^m - p}{p^m - 1} \\ \frac{p^{m+1} - p^m + \alpha(1 - p - 2p^{m-1} + 2p^m)}{p} & \text{if } \alpha > \frac{p^m - p}{p^m - 1} \end{cases}. \end{aligned}$$

□

**Remark 4.1.** If  $n$  is even, then  $\overline{G_n}$  is regular and its  $A_\alpha$ -spectrum will be  $n - 1 - \phi(n)$  and  $\alpha(n - \phi(n)) - (1 - \alpha)\lambda_i(A(G_n)) - 1, i = 2, 3, \dots, n$ . Its  $A_\alpha$ -energy will be  $(1 - \alpha) \left( 2n - 2 + (2^k - 2)\phi(n) - \prod_{i=1}^k p_i + \prod_{i=1}^k (2 - p_i) \right)$ , where  $p_1 p_2 \cdots p_k$  is the largest square-free number that divides  $n$ .

In general, computing exact eigenvalues of complement of UACG for odd values of  $n$  is difficult. But, we estimate a bound for  $A_\alpha$ -eigenvalues of complement of UACG  $G_n$  when  $n$  is odd in the following proposition.

**Proposition 4.2.** The  $A_\alpha$ -spectrum of the complement of UACG  $G_n$ , for odd  $n$ , satisfy the ensuing inequalities:

$$(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha n - \alpha\phi(n) - 1 \leq \lambda_k(A(\overline{G_n})) \leq (1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha n - \alpha\phi(n)$$

for  $2 \leq k \leq \frac{(n+1)}{2}$  and

$$-(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha n - \alpha\phi(n) - 1 \leq \lambda_k(A(\overline{G_n})) \leq -(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)} + \alpha n - \alpha\phi(n)$$

for  $\frac{(n+3)}{2} \leq k \leq n$  and

$$n - \phi(n) - 1 \leq \lambda_1(A(\overline{G_n})) \leq n - \phi(n).$$

*Proof.* Let  $\bar{A}_\alpha = (\bar{a}_{ij}), 0 \leq i, j \leq n - 1$ , be the  $A_\alpha$  matrix of  $\overline{G_n}$ , where

$$\bar{a}_{ij} = \begin{cases} 1 - \alpha & \text{if } \gcd(n, i + j) \neq 1 \text{ and } i \neq j, \\ \alpha(n - 1 - \phi(n)) & \text{if } \gcd(i + j, n) \neq 1 \text{ and } i = j, \\ \alpha(n - \phi(n)) & \text{if } \gcd(n, i + j) = 1 \text{ and } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Then  $\bar{A}_\alpha = \tilde{B} + \tilde{C}$ , where  $\tilde{B} = [b_{ij}], 0 \leq i, j \leq n - 1$ ,

$$b_{ij} = \begin{cases} 1 - \alpha & \text{if } \gcd(n, i + j) \neq 1, \\ 0 & \text{otherwise,} \end{cases}$$

and  $\tilde{C} = [c_{ij}], 0 \leq i, j \leq n - 1$ ,

$$c_{ij} = \begin{cases} \alpha n - \alpha\phi(n) - 1 & \text{if } \gcd(n, i + j) \neq 1 \text{ and } i = j, \\ \alpha(n - \phi(n)) & \text{if } \gcd(n, i + j) = 1 \text{ and } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

According to the definition of  $\tilde{B}$ , it is a left circulant matrix with initial row  $(s_0, s_1, \dots, s_{n-1})$  where

$$s_j = \begin{cases} 1 - \alpha & \text{if } \gcd(j, n) = 1, \\ 0 & \text{otherwise.} \end{cases}$$

So eigenvalues of  $\tilde{B}$  are  $(1 - \alpha)(n - \phi(n)), (1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)}$  for  $1 \leq k \leq \frac{(n-1)}{2}$  and  $-(1 - \alpha)\mu(t_k) \frac{\phi(n)}{\phi(t_k)}$  for  $\frac{(n+1)}{2} \leq k \leq n - 1$ .

Eigenvalues of  $\tilde{C}$  are  $\alpha n - \alpha\phi(n) - 1$  repeated  $n - \phi(n)$  times and  $\alpha(n - \phi(n))$  repeated  $\phi(n)$  times.

Then from the eigenvalues of  $\tilde{B}, \tilde{C}$  and Theorem 2.6 we get the desired result. □

In the following proposition, we estimate some bounds for  $A_\alpha$ -energy of complement of UACG  $G_n$  when  $n$  is odd.

**Proposition 4.3.** For an odd integer  $n \geq 3$  and  $\alpha \in [0, 1)$ ,

$$\epsilon_\alpha(\overline{G_n}) \geq$$

$$\sqrt{(n - \phi(n)) (\alpha^2(n - \phi(n))(n^2 + 2n - 2) + 2n\alpha^2(2\phi(n) + n - 1) + 2n(n - 1)(1 - 2\alpha)) + 2n\alpha^2\phi(n)}.$$

$$\epsilon_\alpha(\overline{G}_n) \geq \frac{2(1 - \alpha)(n - 1)(n - \phi(n))}{n}.$$

$$\epsilon_\alpha(\overline{G}_n) \geq 2\sqrt{(n - \phi(n))^2 + \frac{\phi(n)(2(n - \phi(n)) + 1)}{n}} - \frac{2\alpha(n - 1)(n - \phi(n))}{n}.$$

$$\epsilon_\alpha(\overline{G}_n) \geq \alpha(\phi(n) + 4) - n\alpha - \frac{2\alpha\phi(n)}{n} + \sqrt{(n - \phi(n) + 2)(\alpha^2(n - \phi(n) + 2) + 4(1 - 2\alpha))}.$$

$$\epsilon_\alpha(\overline{G}_n) \leq$$

$$\sqrt{\alpha^2 n^4 + n^3(-\alpha - 2\alpha + 2\phi(n) + 3) - n^2(1 + p)(2\alpha^2 + (1 - \alpha)^2) + np(-2\alpha^2 p - 2\alpha + 1) + p^2}.$$

*Proof.* The Zagreb index of complement of UACG is  $(n - \phi(n))^2 n + \phi(n)(2(n - \phi(n)) + 1)$  and  $\|A(\overline{G}_n)\|_F^2 = (n - 1)(n - \phi(n))$ . Then by using Theorems 2.7 and 2.8 we get the desired result.  $\square$

## 5 $A_\alpha$ -borderenergetic and $A_\alpha$ -hyperenergetic graphs

In this section we define  $A_\alpha$  analogous of borderenergetic and hyperenergetic graphs.

On studying about the  $A_\alpha$ -energy we are interested to know whether there is a graph on  $n$  vertices whose  $A_\alpha$ -energy equals the  $A_\alpha$ -energy of complete graph on  $n$  vertices, that is  $2(1 - \alpha)(n - 1)$ . We refer them as  $A_\alpha$ -borderenergetic graphs. Formally, a graph  $G$  is  $A_\alpha$ -borderenergetic if  $\epsilon_\alpha(G) = \epsilon_\alpha(K_n)$ , for some  $\alpha \in [0, 1)$ . In general borderenergetic graphs are not  $A_\alpha$ -borderenergetic, but regular borderenergetic graphs are  $A_\alpha$ -borderenergetic for every value of  $\alpha$ .

The graph whose  $A_\alpha$ -energy exceeding the  $A_\alpha$ -energy of complete graph is called  $A_\alpha$ -hyperenergetic. That is, a graph  $G$  is  $A_\alpha$ -hyperenergetic if  $\epsilon_\alpha(G) \geq \epsilon_\alpha(K_n)$ , for some  $\alpha \in [0, 1)$ .

In the following section we observed some graphs which are  $A_\alpha$ -borderenergetic and  $A_\alpha$ -hyperenergetic.

### Observations

In this section with the help of Matlab software we study about  $A_\alpha$ -energy of UACG and its complement as  $\alpha$  varies.

**Observation 1 :** From Table 1 we observe the following:

1.  $A_\alpha$ -energy of UACG  $G_n$  decreases as  $\alpha$  increases for  $n = p^m$ .
2.  $G_{p^m}$  is  $A_\alpha$ -hyperenergetic for  $p \geq 5, m \geq 2$ .
3. For all  $m, \overline{G}_{p^m}$  is  $A_\alpha$ -hyperenergetic for some  $\alpha$  greater than 0.4.

Eg:  $\overline{G}_9$  is hyperenergetic for  $\alpha > 0.428571428571$ .  
 $\overline{G}_{25}$  is hyperenergetic for  $\alpha > 0.438596491228$ .

	0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	0.9999
$\varepsilon_\alpha(G_9)$	14.717	13.457	12.197	10.940	9.686	8.438	7.2	5.980	4.810	4	3.9996
$\varepsilon_\alpha(\overline{G}_9)$	10	9.8	9.6	9.4	9.2	9	8.8	8.6	8.666	9	9.333
$\varepsilon_\alpha(K_9)$	16	14.4	12.8	11.2	9.6	8	6.4	4.8	3.2	1.6	0.0016
$\varepsilon_\alpha(G_{27})$	50.683	46.618	42.554	38.490	34.427	30.367	26.309	22.257	18.220	14.247	11.998
$\varepsilon_\alpha(\overline{G}_{27})$	34	33.4	32.8	32.2	31.6	31	30.4	29.8	29.2	28.6	29.332
$\varepsilon_\alpha(K_{27})$	52	46.8	41.6	36.4	31.2	26	20.8	15.6	10.4	5.2	0.005
$\varepsilon_\alpha(G_{81})$	158.672	146.206	133.740	121.274	108.809	96.344	83.880	71.418	58.961	46.523	35.996
$\varepsilon_\alpha(\overline{G}_{81})$	106	104.2	102.4	100.6	98.8	97	95.2	93.4	91.6	89.8	89.329
$\varepsilon_\alpha(K_{81})$	160	144	128	112	96	80	64	48	32	16	0.016
$\varepsilon_\alpha(G_5)$	6.472	5.841	5.212	4.588	3.969	3.361	2.772	2.224	1.774	1.546	1.599
$\varepsilon_\alpha(\overline{G}_5)$	4	4.08	4.16	4.24	4.32	4.4	4.48	4.56	4.64	4.72	4.799
$\varepsilon_\alpha(K_5)$	8	7.2	6.4	5.6	4.8	4	3.2	2.4	1.6	0.8	0.001
$\varepsilon_\alpha(G_{25})$	54.413	49.533	44.656	39.778	34.902	30.026	25.153	20.286	15.432	10.641	7.999
$\varepsilon_\alpha(\overline{G}_{25})$	28	27.76	27.52	27.28	27.04	26.8	26.56	26.32	26.08	26.48	27.199
$\varepsilon_\alpha(K_{25})$	48	43.2	38.4	33.6	28.8	24	19.2	14.4	9.6	4.8	0.004
$\varepsilon_\alpha(G_{125})$	294.402	268.722	243.043	217.363	191.684	166.005	140.326	114.648	88.973	63.306	39.996
$\varepsilon_\alpha(\overline{G}_{125})$	148	146.96	145.92	144.88	143.84	142.8	141.76	140.72	139.68	138.64	139.196
$\varepsilon_\alpha(K_{125})$	248	223.2	198.4	173.6	148.8	124	99.2	74.4	49.6	24.8	0.0248
$\varepsilon_\alpha(G_{625})$	1494.401	1364.721	1235.041	1105.361	975.681	846.001	716.321	586.641	456.962	327.285	199.981
$\varepsilon_\alpha(\overline{G}_{625})$	748	742.96	737.92	732.88	727.84	722.8	717.76	712.72	707.68	702.64	699.180
$\varepsilon_\alpha(K_{625})$	1248	1123.2	998.4	873.6	748.8	624	499.2	374.4	249.6	124.8	0.124
$\varepsilon_\alpha(G_{49})$	118.291	107.405	96.520	85.635	74.751	63.867	52.984	42.103	31.226	20.372	11.998
$\varepsilon_\alpha(\overline{G}_{49})$	54	53.742	53.485	53.228	52.971	52.714	52.457	52.2	51.942	52.028	53.141
$\varepsilon_\alpha(K_{49})$	96	86.4	76.8	67.2	57.6	48	38.4	28.8	19.2	9.6	0.009
$\varepsilon_\alpha(G_{121})$	318.183	288.092	258.001	227.911	197.820	167.73	137.639	107.550	77.461	47.378	19.998
$\varepsilon_\alpha(\overline{G}_{121})$	130	129.727	129.454	129.181	128.909	128.636	128.363	128.091	127.818	127.545	129.089
$\varepsilon_\alpha(K_{121})$	240	216	192	168	144	120	96	72	48	24	0.024

Table 1:  $A_\alpha$ -energy of  $G_{p^m}$ ,  $(\overline{G}_{p^m})$  and  $K_{p^m}$  for different values of  $\alpha$ .

**Observation 2 :** From Table 1 we also observe the following:

1. For all  $m$  and some  $\alpha$ ,  $G_{3^m}$  is  $A_\alpha$ -borderenergetic and as  $m$  increases,  $\alpha \rightarrow 0$ .

Eg:  $G_9$  is borderenergetic for  $\alpha = 0.375$ , (see Table 2). However  $G_{p^m}$  is not  $A_\alpha$ -borderenergetic for  $p \geq 5$  and  $m \geq 2$ .

2.  $G_{p^m}$  is not adjacency borderenergetic for any  $p$  and  $m$ .

3. For all  $m$  and  $p \geq 5$ ,  $\overline{G}_{p^m}$  is  $A_\alpha$ -borderenergetic for some  $\alpha \in (0.4, 0.5)$ .

Eg:  $\overline{G}_5$  is borderenergetic for  $\alpha = 0.454545454545$ .

$\overline{G}_{49}$  is borderenergetic for  $\alpha = 0.449541284404$ .

$n$	$\alpha$	$\varepsilon_\alpha(G_n)$	$\varepsilon_\alpha(K_n)$
9	0.375	10	10
27	0.115981467677	45.968963680773	45.968963680773
81	0.037573807884	153.988190738623	153.988190738623
243	0.012405082000	477.995940311853	477.995940311853
5	0.755511381250	1.955908950000	1.955908950000

Table 2: Values of  $\alpha$  for which  $G_{p^m}$  is  $A_\alpha$ -borderenergetic.

$n$	$\alpha$	$\varepsilon_\alpha(\overline{G}_n)$	$\varepsilon_\alpha(K_n)$
9	0.428571428571	9.142857142857	9.142857142857
27	0.391304347826	31.652173913043	31.652173913043
81	0.380281690141	99.154929577465	99.154929577465
5	0.454545454545	4.363636363636	4.363636363636
25	0.438596491228	26.947368421053	26.947368421053
125	0.420875420875	143.622895622896	143.622895622896
625	0.417501670007	726.957915831663	726.957915831663
49	0.449541284404	52.844036697248	52.844036697248
121	0.463601532567	128.735632183908	128.735632183908

Table 3: Values of  $\alpha$  for which  $\overline{G}_{p^m}$  is  $A_\alpha$ -borderenergetic.

## 6 Conclusion

In this article the  $A_\alpha$ -eigenvalues of unitary addition Cayley graph  $G_n$  when its order is  $p^m$  or even is calculated. When its order is odd, a bound for the  $A_\alpha$ -eigenvalues is found. In addition, the  $A_\alpha$ -energy of  $G_n$  when  $n$  is  $p^m$  or even is computed. For odd  $n$ , bounds for  $A_\alpha$ -energy of  $G_n$  and  $\overline{G}_n$  are estimated. Also the concept of  $A_\alpha$ -borderenergetic and  $A_\alpha$ -hyperenergetic graphs is introduced and found some families of each.

## Acknowledgement

The authors would like to thank the DST, Government of India, for providing support to carry out this work under the scheme 'FIST'(No.SR/FST/MS-I/2019/40). The first author gratefully acknowledges the financial support of University Grants Commission(UGC), India.

## References

- [1] Cui, Shu-Yu, and Gui-Xian Tian. The spectrum and the signless Laplacian spectrum of coranae. *Linear algebra and its applications* 437.7 (2012): 1692-1703.
- [2] Dejter, Italo J., and Reinaldo E. Giudici. On unitary Cayley graphs. *Journal of Combinatorial Mathematics and Combinatorial Computing* 18 (1995): 121-124.
- [3] Ghorbani, M., Deng, B., Hakimi-Nezhaad, M. and Li, X. A survey on borderenergetic graphs. *MATCH Commun Math Comput Chem*, 84, 293-322, (2020).
- [4] Gong, Shicai, Xueliang Li, Guanghui Xu, Ivan Gutman, and Boris Furtula. Borderenergetic graphs. *MATCH Commun. Math. Comput. Chem* 74, no. 2 (2015): 321-332.
- [5] Gutman, Ivan. Hyperenergetic molecular graphs. *Journal of the serbian chemical society* 64, no. 3 (1999): 199-205.
- [6] Ilić, Aleksandar. The energy of unitary Cayley graphs. *Linear Algebra and its Applications*, 431.10 (2009): 1881-1889.
- [7] Jahfar, T. K., V. K. Najiya, and A. V. Chithra. Construction of new families of borderenergetic graphs. *AIP Conference Proceedings*. Vol. 2336. No. 1. AIP Publishing LLC, 2021.
- [8] Naveen Palanivel and AV Chithra. Energy and Laplacian energy of unitary addition Cayley graphs. *Filomat*, 33(11):3599–3613, 2019.
- [9] O. Favaron, M. Maheo, and J.-F. Sacle. Some Eigenvalue Properties in Graphs: Conjectures of Graffiti-II. *Discrete Mathematics*, 111:197–220, 1993.
- [10] Palanivel Naveen, and A. V. Chithra. Some structural properties of unitary addition Cayley graphs. *International Journal of Computer Applications* 121.17 (2015): 1-5.
- [11] Pirzada, Shariefuddin, Bilal A. Rather, Hilal A. Ganie, and Rezwana ul Shaban. On  $\alpha$ -adjacency energy of graphs and Zagreb index. *AKCE International Journal of Graphs and Combinatorics* 18, no. 1 (2021): 39-46.
- [12] Ravindra B Bapat. *Graphs and matrices*, volume 27. Springer, 2010.
- [13] Shanmukha, B., H. N. Shwetha, and N. Manjunath. "Bounds for  $\alpha$ -Adjacency Energy of a Graph." *IOP Conference Series: Materials Science and Engineering*. Vol. 1070. No. 1. IOP Publishing, 2021.

- [14] Sinha, Deepa, Pravin Garg, and Anjali Singh. Some properties of unitary addition Cayley graphs. *Notes on Number Theory and Discrete Mathematics* 17.3 (2011): 49-59.
- [15] Tura, Fernando. L-borderenergetic graphs. *MATCH Commun. Math. Comput. Chem* 77 (2017): 37-44.
- [16] Vaidya, S.K. And Popat, K.M., 2024. Some Borderenergetic and Equienergetic Graphs. *Kragujevac Journal Of Mathematics*, 48(6), 935-949.
- [17] Vladimir Nikiforov. Merging the A-and Q-spectral theories. *Applicable Analysis and Discrete Mathematics*, 11(1):81–107, 2017.
- [18] X. Liu and Z. Zhang. Spectra of subdivision-vertex and subdivision-edge joins of graphs. *Bulletin of the Malaysian Mathematical Sciences Society*, 42 (2019), 15–31.
- [19] Zhang, Fuzhen, ed. *The Schur complement and its applications*. Vol. 4. Springer Science & Business Media, 2006.