

RICCI FLOW UNDER KATO-TYPE CURVATURE LOWER BOUND

MAN-CHUN LEE

ABSTRACT. In this work, we extend the existence theory of non-collapsed Ricci flows from point-wise curvature lower bound to Kato-type curvature lower bound. As an application, we prove that compact three dimensional non-collapsed strong Kato limit space is homeomorphic to a smooth manifold. Moreover, similar result also holds in higher dimension under stronger curvature condition. We also use the Ricci flow smoothing to study stability problem in scalar curvature geometry.

1. INTRODUCTION

There is a long history on the study of compactness of sets of manifolds with uniform lower bound of Ricci curvature. For instance, the Gromov compactness Theorem [20] states that sequence of compact manifolds with uniform Ricci lower bound and diameter upper bound admits a convergent subsequence in the Gromov-Hausdorff topology. Since then, there have been many studies toward understanding the structure of the metric space arising as limits of smooth manifolds, see [10, 11, 12, 13, 14, 17] and the reference therein. The analytic properties such as comparison geometry and heat kernel estimates on manifolds with Ricci lower bound play an important role there.

On the other hand, there are however many interesting scenarios in which uniform Ricci lower bound is missing, especially in the study of Ricci flow. It is then natural and important to consider the case when a uniform Ricci lower bound is further relaxed to bound in a weaker sense. In [35], Petersen and Wei generalized the classical fundamental Laplacian and volume comparison for uniform Ricci curvature lower bounds to smallness assumptions of L^p norm of Ric_- for $p > n/2$ where $\text{Ric}_-(x) = \inf\{\alpha \geq 0 : \text{Ric}(g(x)) + \alpha g(x) \geq 0\}$. This opens the door to understanding the structure of limit spaces under small $\|\text{Ric}_-\|_p$ assumption for $p > n/2$, see also [18, 30].

More recently, it was further generalized by Carron [5] and Rose [36] that in the compact case, a Dynkin-type lower bound of Ricci curvature also suffices to obtain the Li-Yau estimates for the heat kernel, building on the idea of Zhang-Zhu [46]. For a complete Riemannian manifold (M^n, g_0) of dimension

Date: April 19, 2023.

2010 Mathematics Subject Classification. Primary 53C44 .

Key words and phrases. Kato-type lower bound, Ricci flow smoothing, stability.

$n \geq 2$, we denote

$$\kappa_t(M^n, g_0) = \sup_{x \in M} \int_0^t \int_M H_{g_0}(x, y, s) (\text{Ric})_-(y) d\text{vol}_{g_0}(y) ds$$

where H_{g_0} is the heat kernel, that is $H(\cdot, \cdot, t)$ is the kernel of the operator $e^{-t\Delta_{g_0}}$ for any $t > 0$.

Definition 1.1. Let $\{(M_i, g_i)\}_{i=1}^\infty$ be a sequence of compact manifolds¹. We say that $\{(M_i, g_i)\}_{i=1}^\infty$ satisfies

- a uniform Dynkin bound on Ric_- if there exists $T > 0$ such that for all $i \in \mathbb{N}$,

$$\kappa_T(M_i, g_i) \leq \frac{1}{16n};$$

- a uniform Kato bound on Ric_- if there exist a non-decreasing function $f : (0, T] \rightarrow (0, +\infty)$, $T > 0$ such that $f(t) \rightarrow 0$ as $t \rightarrow 0^+$ and for all $t \in (0, T]$ and $i \in \mathbb{N}$,

$$\kappa_t(M_i, g_i) \leq f(t);$$

- a strong uniform Kato bound on Ric_- if there exist a non-decreasing function $f : (0, T] \rightarrow (0, +\infty)$, $T, \Lambda > 0$ such that $f(t) \rightarrow 0$ as $t \rightarrow 0^+$ and for all $t \in (0, T]$ and $i \in \mathbb{N}$,

$$\kappa_t(M_i, g_i) \leq f(t); \quad \int_0^T \frac{\sqrt{f(s)}}{s} ds \leq \Lambda.$$

When $\text{Ric} \geq -K$, it satisfies the uniform Dynkin for $T = (16nK)^{-1}$. In the compact case, it was also shown that a strong uniform Kato bound can be achieved for some small T under suitable L^p bound of Ric_- [38] or suitable Morrey bound [9]. Perhaps more importantly, it was proved by Carron [5] that the set of compact manifolds satisfying a uniform Dynkin bound on Ric_- is pre-compact in the Gromov-Hausdorff topology. When it is strengthened to strong uniform Kato sense, the structure theory of the corresponding volume non-collapsed Gromov-Hausdorff limit has been developed by Carron-Mondello-Tewodrose [6]. This generalizes the earlier works of Cheeger-Colding on Ricci limit spaces. We refer readers to [7, 8, 37, 39] for more interesting and important developments.

In this work, we are interested in the regularity of the Gromov-Hausdorff limits of volume non-collapsed compact manifolds (without boundary) under the strong Kato lower bound on 1-isotropic curvature. To clarify the notion, denote the space of algebraic curvature tensors on \mathbb{R}^n by $\mathcal{C}_B(\mathbb{R}^n)$. For any given $R \in \mathcal{C}_B(\mathbb{R}^n)$, we extend it complex linearly to \mathbb{C}^n . We say that $R \in \mathcal{C}_{\text{PIC}2}$ if for each two complex dimensional subspace Σ of \mathbb{C}^n and orthonormal basis $v, w \in \mathbb{C}^n$ of Σ , we have $R(v, w, \bar{v}, \bar{w}) \geq 0$. If one instead asks for non-negativity of complex sectional curvature only for PIC1 sections, defined to be

¹We might analogously define the notions on complete non-compact case.

those Σ that contain some non-zero vector v whose conjugate \bar{v} is orthogonal to the entire section Σ . The algebraic curvature tensors R with non-negative complex sectional curvature for each such restricted Σ form a cone we denote by C_{PIC_1} . When $n = 3$, it is known that C_{PIC_1} is also the cone of curvature tensors R with $\text{Ric}(R) \geq 0$.

For $n \geq 4$, one can equivalently describe C_{PIC_1} as follows (see [3]): C_{PIC_1} is the cone consisting of curvature tensors R such that for any orthonormal four-frame $\{e_i\}_{i=1}^4 \subset \mathbb{R}^n$ and $\lambda \in [0, 1]$, we have

$$R_{1313} + \lambda^2 R_{1414} + R_{2323} + \lambda^2 R_{2424} - 2\lambda R_{1234} \geq 0.$$

We are interested in studying manifolds where the curvature is bounded from below with respect to C_{PIC_1} . For a Riemannian manifold (M^n, g) with $n \geq 3$ and $x \in M$, we define

$$(\text{IC}_1)_-(x) = \inf\{\alpha \geq 0 : \text{Rm}_{g(x)} + \alpha I \in C_{\text{PIC}_1}\}$$

where I is the curvature tensor defined by $I_{ijkl} = \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk}$. This is the negative part of the lowest eigenvalue of Rm with respect to C_{PIC_1} . This generalizes the notion of Ric_- in the sense that when $n = 3$, $\text{Ric}_- = (n - 1) \cdot (\text{IC}_1)_-$ while when $n \geq 4$, we have $\text{Ric}_- \leq (n - 1) \cdot (\text{IC}_1)_-$.

We follow the spirit of κ_t and define, for all $t > 0$,

$$\kappa_{t, \text{IC}_1}(M^n, g_0) = \sup_{x \in M} \int_0^t \int_M H_{g_0}(x, y, s) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) \, ds.$$

Definition 1.2. Suppose $T > 0$ and $f : (0, T] \rightarrow (0, +\infty)$ is a non-decreasing function such that

$$(1.1) \quad \lim_{t \rightarrow 0} f(t) = 0, \quad \text{and} \quad f(T) \leq \frac{1}{16n(n-1)}.$$

We say that $(M^n, g) \in \mathcal{K}_{\text{IC}_1}(n, f)$ if for all $t \in (0, T]$ we have

$$\kappa_t(M^n, g_0) \leq f(t).$$

If in addition, $\text{Vol}_g(B_g(x_0, \sqrt{T})) \geq vT^{n/2}$ for some $x_0 \in M, v > 0$, then we say that $(M^n, g, x_0) \in \mathcal{K}_{\text{IC}_1}(n, f, v)$.

We will mainly focus on the case when M is compact and the Kato-bound is uniformly strong. Given a sequence of compact manifolds $\{(M_i, g_i)\}_{i=1}^\infty$, we say that $\{(M_i, g_i)\}_{i=1}^\infty$ satisfies a strong uniform IC_1 -Kato if $(M_i, g_i) \in \mathcal{K}_{\text{IC}_1}(n, f)$ where $f : (0, T] \rightarrow (0, +\infty)$ is a non-decreasing function, $T, \Lambda > 0$ such that $f(t) \rightarrow 0$ as $t \rightarrow 0^+$ and satisfies

$$(1.2) \quad \int_0^T \frac{\sqrt{f(s)}}{s} \, ds \leq \Lambda.$$

When $n = 3$, it coincides with the concept of strong Kato bound of Ric_- considered in [6], modulus scaling².

In contrast with the work [6, 7], we intend to strengthen the curvature lower bound from Ricci curvature to 1-isotropic curvature (which two notions coincide when $n = 3$). This is largely motivated by the works of Bamler-Cabezas-Wilking [2], Simon-Topping [32], Lai [25] and Lee-Tam [26] on obtaining the regularity of the Gromov-Hausdorff limit using Ricci flow.

The following is the main technical result which says that such manifolds can be regularized by the Ricci flow uniformly. This generalizes the earlier work of Bamler-Cabezas-Wilking [2] on the theory of Ricci flows with curvature bounded from below in point-wise sense initially.

Theorem 1.1. *Suppose (M^n, g_0, x_0) , $n \geq 3$ is a pointed compact manifold which is inside $\mathcal{K}_{\text{IC}_1}(n, f, v)$ for some non-decreasing function f , T and Λ satisfying (1.2) and $\text{diam}(M, g_0) \leq D$ for some $D > 0$, then there exist S, α, \tilde{v} depending only on n, f, v, D, T, Λ and a solution $g(t), t \in [0, S]$ to the Ricci flow such that for all $(x, t) \in M \times (0, S]$,*

- (a) $|\text{Rm}(x, t)| \leq \alpha t^{-1}$;
- (b) $\text{Vol}_{g(t)} \left(B_{g(t)}(x, \sqrt{T}) \right) \geq \tilde{v} T^{n/2}$;
- (c) $\text{inj}(g(x, t)) \geq \sqrt{\alpha^{-1}t}$;
- (d) $\text{Rm}(g(x, t)) + \alpha(1 + t^{-1}f(\alpha t)) \cdot \text{I} \in \text{C}_{\text{PIC}_1}$.

A similar result also holds in the Kähler case under strong uniform Kato lower bound on bisectional curvature, see Theorem 5.1.

Remark 1.1. Using the partial Ricci flow approach in [25] (see also [32, 26]), the result also holds in the complete non-compact case *without* bounded curvature assumption if suitable *static* heat kernel estimates hold. In the presence of uniform Ricci lower bound, this usually follows from Li-Yau estimates. In the compact case, it has been obtained by Carron [5] and Rose [36] under the Kato-type lower bound of Ricci curvature. The comparison geometry in the complete non-compact case seems out of reach at the moment due to the absence of exhaustion function with good control. We will therefore only focus on the compact case in this work.

As an application, we prove that the Gromov-Hausdorff limit of sequence of compact manifolds satisfying the strong uniform IC1-Kato bound and a uniform diameter upper bound is homeomorphic to a smooth manifold.

Corollary 1.1. *Suppose $(M_i^n, g_{i,0})$ is a sequence of compact manifold such that*

- (a) $(M_i, g_{i,0}, x_i) \in \mathcal{K}_{\text{IC}_1}(n, f, v)$ for some $x_i \in M_i, v > 0$ and a non-decreasing function f satisfying (1.2);

²there will be a normalization constant $n - 1$ since the positivity of IC_1 is defined using language of cone.

(b) $\text{diam}(M_i, g_{i,0}) \leq D$ for some $D > 0$.

Then there exist a smooth compact manifold M_∞ and a distance function d_∞ on M_∞ so that $(M_i, g_{i,0})$ sub-converges to (M_∞, d_∞) in the measured Gromov-Hausdorff sense. Moreover, there exists a Ricci flow $g_\infty(t), t \in (0, S]$ on M_∞ such that

$$\lim_{t \rightarrow 0} d_{GH}((M_\infty, d_\infty), (M_\infty, g_\infty(t))) = 0.$$

Furthermore, if in addition $(M_i, g_{i,0}, x_i) \in \mathcal{K}_{IC_1}(n, f_i, v)$ for some decreasing function f_i in which $f_i(T) \rightarrow 0$, then $\text{Rm}(g_\infty(t)) \in \mathcal{C}_{\text{PIC}_1}$.

The final part of the result also holds if $\mathcal{C}_{\text{PIC}_1}$ is replaced by other curvature cone in Lemma 3.1. Moreover, it particularly infers that if a compact manifold supports a metric with sufficiently small negative lower bound of IC_1 in the strong Kato sense relative to n, D, v , then it also admits a smooth metric whose curvature operator lies in $\mathcal{C}_{\text{PIC}_1}$ via a contradiction argument.

Another application is to understand the stability of metrics on torus with almost non-negative scalar curvature. When $M = \mathbb{T}^n$, the celebrated work of Schoen and Yau [41, 42], Gromov and Lawson [19] stated that metrics with non-negative scalar curvature $\mathcal{R} \geq 0$ must be flat. In [21], Gromov asked if a sequence of metrics g_i with $\mathcal{R}(g_i) \geq -i^{-1}$ on \mathbb{T}^n will sub-converge to a flat metric in some appropriate weak sense. For related works, we refer interested readers to the survey paper of Sormani [43] for a comprehensive discussion. In this regard, we show that if in addition the 1-isotropic curvature is bounded from below in the strong Kato sense, then the sequence will sub-converge to the flat torus in the measured Gromov-Hausdorff sense.

Corollary 1.2. *Suppose $(M_i^n, g_{i,0})$ is a sequence of compact manifolds such that*

- (a) $(M_i, g_{i,0}, x_i) \in \mathcal{K}_{\text{IC}_1}(n, f, v)$ for some $x_i \in M_i, v > 0$ and a non-decreasing function f satisfying (2.2);
- (b) $\text{diam}(M_i, g_{i,0}) \leq D$ for some $D > 0$;
- (c) $\sigma(M_i) \leq 0$ (equivalently, M_i does not admit PSC metric);
- (d) $\mathcal{R}(g_i) \geq -i^{-1}$.

Then after passing to subsequent, $(M_i^n, g_{i,0})$ converges to a compact Ricci flat manifold (M_∞, g_∞) in the measured Gromov-Hausdorff sense. If in addition $b_1(M_i) = n$ for all i , then (M_∞, g_∞) is a flat torus.

Acknowledgement: The author would like to thank C. Rose for his interest and D. Tewodrose for comments on improving the manuscript. The work was partially supported by Hong Kong RGC grant (Early Career Scheme) of Hong Kong No. 24304222, a direct grant of CUHK and a NSFC grant.

2. THE STRONG KATO BOUND AND IMPLICATIONS

In the following, compact manifolds are referred those without boundary. In this section, we will collect some important properties from [6] for compact

manifolds (M^n, g) satisfying the strong Kato bound. The properties will play crucial role in obtaining estimates of Ricci flows.

Since $C_{\text{PIC1}} \subseteq C_{\text{Ric}}$ for $n \geq 3$, it is clear that for all $t \in (0, +\infty)$,

$$\kappa_t(M^n, g_0) \leq (n-1)\kappa_{t, \text{IC1}}(M^n, g_0).$$

Assume $T > 0$ and $f : (0, T] \rightarrow (0, +\infty)$ such that f is non-decreasing and satisfies

$$(2.1) \quad \lim_{t \rightarrow 0} f(t) = 0, \quad \text{and} \quad f(T) \leq \frac{1}{16n}.$$

We say that $(M^n, g) \in \mathcal{K}(n, f)$ if for all $t \in (0, T]$ we have $\kappa_t(M^n, g) \leq f(t)$. Similarly, we define $\mathcal{K}(n, f, v)$ to be the class of pointed compact manifolds (M, g, x_0) such that the volume $\text{Vol}_g(B_g(x_0, \sqrt{T})) \geq vT^{n/2}$. The following Proposition says that the heat kernel will behave in a similar way as the Euclidean one under the Kato control. This is originated in [5]

Proposition 2.1. [7, Proposition 2.6] *There exists $\gamma(n) \geq 1$ such that if $(M^n, g) \in \mathcal{K}(n, f)$, then for all $x, y \in M$ and $t \in (0, T)$,*

$$\frac{\gamma^{-1}}{\text{Vol}_g(B_g(x, r))} \cdot \exp\left(-\frac{\gamma d^2(x, y)}{t}\right) \leq H_{g_0}(x, y, t) \leq \frac{\gamma}{\text{Vol}_g(B_g(x, r))} \cdot \exp\left(-\frac{d^2(x, y)}{5t}\right).$$

Moreover, in this case, it is known that $(M^n, g) \in \mathcal{K}(n, f)$ is a volume doubling space: for any $x \in M$, $0 < s < r \leq \sqrt{T}$,

$$\frac{\text{Vol}_g(B_g(x, r))}{\text{Vol}_g(B_g(x, s))} \leq C \left(\frac{r}{s}\right)^{e^2 n}.$$

If in addition f satisfies a stronger integrability condition:

$$(2.2) \quad \int_0^T \frac{\sqrt{f(s)}}{s} ds \leq \Lambda$$

for some $\Lambda > 0$. As pointed out in [6], f satisfying (2.2) automatically implies that f satisfies (2.1) by shrinking T . It was first shown by Carron [5] that under the stronger assumption (2.2), we can strength the conclusion to Ahlfors n -regular.

Proposition 2.2. [6, Proposition 2.14 & Corollary 2.15] *Suppose $(M^n, g) \in \mathcal{K}(n, f)$ and f satisfies (2.2) for some $\Lambda > 0$. Then there exists $C_n > 0$ such that for all $x \in M$ and $0 < s < r \leq \sqrt{T}$,*

$$\text{Vol}_g(B_g(x, r)) \leq C_n^{\Lambda+1} r^n \quad \text{and} \quad \frac{\text{Vol}_g(B_g(x, r))}{\text{Vol}_g(B_g(x, s))} \leq C_n^{\Lambda+1} \left(\frac{r}{s}\right)^n.$$

Moreover, for any $x, y \in M$ and $0 \leq r \leq \sqrt{T}$,

$$\frac{\text{Vol}_g(B_g(x, \sqrt{T}))}{T^{n/2}} \leq C_n^{(\Lambda+1)T^{-1/2}d(x,y)} \cdot \frac{\text{Vol}_g(B_g(y, r))}{r^n}.$$

We are primarily interested in the non-collapsed case in this work. We end this section by observing that the non-collapsed strong Kato assumption can be re-phased as an estimate in an integral form.

Lemma 2.1. *Suppose $(M^n, g, x_0) \in \mathcal{K}_{\text{IC}_1}(n, f)$ for some f satisfying (2.2), then there exists $C_1(n, \Lambda, T) > 0$ such that for all $(x, t) \in M \times (0, T]$,*

$$\int_M \frac{1}{t^{n/2}} \exp\left(-\frac{C_1 d_{g_0}^2(x, y)}{t}\right) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) \leq \frac{C_1 f(t)}{t}.$$

Proof. Since $\kappa_t(M, g) \leq (n-1)\kappa_{t, \text{IC}_1}$, we have $\mathcal{K}_{\text{IC}_1}(n, f) \subseteq \mathcal{K}(n, f)$ for a possibly larger Λ . By proposition 2.1, Proposition 2.2 and the volume non-collapsing, there exist $C_0(n, \Lambda, T), \gamma(n) > 0$ such that for all $(x, t) \in M \times (0, T]$,

$$\begin{aligned} & \int_0^t \int_M H(x, y, s) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) ds \\ (2.3) \quad & \geq \int_0^t \int_M \frac{\gamma}{V_g(x, \sqrt{s})} \exp\left(-\frac{\gamma d_{g_0}^2(x, y)}{s}\right) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) ds \\ & \geq \frac{\gamma}{C_n^{\Lambda+1}} \int_0^t \int_M \frac{1}{s^{n/2}} \exp\left(-\frac{\gamma d_{g_0}^2(x, y)}{s}\right) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) ds. \end{aligned}$$

Combines this with the assumption of Kato-type bound, we have for all $(x, t) \in M \times (0, T]$,

$$\begin{aligned} (2.4) \quad f(t) & \geq \frac{\gamma}{C_n^{\Lambda+1}} \int_{t/2}^t \int_M \frac{1}{s^{n/2}} \exp\left(-\frac{\gamma d_{g_0}^2(x, y)}{s}\right) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y) ds \\ & \geq \frac{t\gamma}{2C_n^{\Lambda+1}} \int_M \frac{1}{t^{n/2}} \exp\left(-\frac{2\gamma d_{g_0}^2(x, y)}{t}\right) (\text{IC}_1)_-(y) \, d\text{vol}_{g_0}(y). \end{aligned}$$

This completes the proof by re-arranging the inequality. \square

3. RICCI FLOW SMOOTHING AND ESTIMATES

The idea of this work is to mollify the metric g_0 using the Ricci flow. The Ricci flow is a one parameter family of metrics $g(t)$ satisfying

$$(3.1) \quad \partial_t g(t) = -2\text{Ric}(g(t)), \quad g(0) = g_0.$$

In the coming few sections, we will focus on obtaining estimates of Ricci flows.

If (M^n, g_0) is a compact manifold, it was shown by Hamilton [40] that the Ricci flow admits a short-time solution $g(t)$. It is by-now known to be extremely powerful in regularizing metrics even in case without bounded curvature, for instances see the works [22, 33, 25, 32, 24] and the reference therein. Although we are primarily interested in the compact case, we will state our result in this section using a localized form so that it also applies to the non-compact case.

It is a general philosophy that the Ricci flow will tend to be positively curved as it evolves, for instances see the list in Lemma 3.1 and the recent work of

Brendle [4]. The following Proposition illustrates its relation to curvature estimates in the non-collapsed case. This is a slight generalization of the curvature estimates in [25, 26, 31].

Proposition 3.1. *For any $n \geq 3$, $v_0, L > 0$, there exist $\tilde{S}(n, v_0, L), a(n, v_0, L) > 0$ such that the following holds. Suppose $(M^n, g(t)), t \in [0, S]$ is a smooth solution to the Ricci flow (not necessarily complete) such that for some $x_0 \in M$, we have*

- (a) $B_{g(t)}(x_0, 1) \Subset M$ for all $t \in [0, S]$;
- (b) for all $x \in B_{g(t)}(x_0, \frac{3}{4}), t \in [0, S]$ and $r \in (0, \frac{1}{4}\sqrt{L^{-1}t} \wedge \frac{1}{4})$,

$$\text{Vol}_{g(t)}(B_{g(t)}(x, r)) \geq v_0 r^n.$$

- (c) for all $x \in B_{g(t)}(x_0, 1)$ and $t \in (0, S]$, we have either
 - (i) $n = 3$ or;
 - (ii) $\text{IC}_1(g(t)) \geq -Lt^{-1}$ or;
 - (iii) $\text{OB}(g(t))^3 \geq -Lt^{-1}$ when n is even and $g(t)$ is Kähler.

Then for $t \in (0, \tilde{S} \wedge S]$ and $x \in B_{g(t)}(x_0, \frac{1}{2})$, we have

$$|\text{Rm}(x, t)| \leq at^{-1}.$$

Proof. The proof of the curvature estimate uses the Perelman-inspired point-picking argument from [31, Lemma 2.1]. We illustrate the case of (ii) and will point out the modification needed for (i) and (iii).

Suppose the conclusion is false for some $n, v_0, L > 0$. Then for any $a_k \rightarrow +\infty$, we can find a sequence of manifold M_k^n , Ricci flows $g_k(t), t \in [0, S_k]$ and $p_k \in M_k$ satisfying the hypotheses but so the curvature estimate fails with $a = a_k$ in an arbitrarily short time. We might assume $a_k S_k \rightarrow 0$. Using the smoothness of the flow, we can choose $t_k \in (0, S_k]$ such that

- (i) $B_{g_k(t)}(x_k, 1) \Subset M_k$ for $t \in [0, t_k]$;
- (ii) $\text{Vol}_{g_k(t)}(B_{g_k(t)}(x, r)) \geq v_0 r^n$ for $t \in [0, t_k], x \in B_{g_k(t)}(x_k, \frac{3}{4})$ and $r \in (0, \frac{1}{4}\sqrt{L^{-1}t}]$;
- (iii) $\text{IC}_1(g_k(t)) \geq -Lt^{-1}$ on $B_{g_k(t)}(x_k, 1)$ for $t \in [0, t_k]$;
- (iv) $|\text{Rm}_{g_k}(x, t)| < a_k t^{-1}$ for $t \in [0, t_k]$ and $x \in B_{g_k(t)}(x_k, \frac{1}{2})$;
- (v) $|\text{Rm}_{g_k}(z_k, t_k)| = a_k t_k^{-1}$ for some $z_k \in \overline{B_{g_k(t_k)}(x_k, \frac{1}{2})}$.

By (v) and the fact that $a_k t_k \rightarrow 0$, [31, Lemma 5.1] implies that for sufficiently large $k \in \mathbb{N}$, we can find $\beta(n) > 0, \tilde{t}_k \in (0, t_k]$ and $\tilde{x}_k \in B_{g_k(\tilde{t}_k)}(x_k, \frac{3}{4} - \frac{1}{2}\beta\sqrt{a_k \tilde{t}_k})$ such that

$$(3.2) \quad |\text{Rm}_{g_k}(x, t)| \leq 4|\text{Rm}_{g_k}(\tilde{x}_k, \tilde{t}_k)| = 4Q_k$$

whenever $d_{g_k(\tilde{t}_k)}(x, \tilde{x}_k) < \frac{1}{8}\beta a_k Q_k^{-1/2}$ and $\tilde{t}_k - \frac{1}{8}a_k Q_k^{-1} \leq t \leq \tilde{t}_k$ where $\tilde{t}_k Q_k \geq a_k \rightarrow +\infty$. We note here that from the proof of [31, Lemma 5.1], for $(x, t) \in$

³OB denotes the orthogonal bisectional curvature in the Kähler case

$\frac{1}{8}\beta a_k Q_k^{-1/2} \times [\tilde{t}_k - \frac{1}{8}a_k Q_k^{-1}, \tilde{t}_k]$, we have $d_{g_k(t)}(x, x_k) < 1$ so that the conditions applied.

We now consider the parabolic rescaled Ricci flows $\tilde{g}_k(t) = Q_k g(t_k + Q_k^{-1}t)$, $t \in [-\frac{1}{8}a_k, 0]$ so that

- (a) $B_{\tilde{g}_k(0)}(\tilde{x}_k, \frac{1}{8}\beta a_k) \Subset M_k$;
- (b) $|\text{Rm}_{\tilde{g}_k(t)}| \leq 4$ on $B_{\tilde{g}_k(0)}(\tilde{x}_k, \frac{1}{8}\beta a_k) \times [-\frac{1}{8}a_k, 0]$;
- (c) $|\text{Rm}_{\tilde{g}_k}(\tilde{x}_k, 0)| = 1$;
- (d) $\text{Vol}_{\tilde{g}_k(0)}(B_{g_k(t)}(\tilde{x}_k, r)) \geq v_0 r^n$ for $r \in (0, \frac{1}{4}\sqrt{L^{-1}Q_k t_k}]$;
- (e) $\text{IC}_1(\tilde{g}_k(t)) \geq -\frac{L}{t + Q_k \tilde{t}_k}$ on $B_{\tilde{g}_k(0)}(\tilde{x}_k, \frac{1}{8}\beta a_k) \times [-\frac{1}{8}a_k, 0]$.

By (b), (d) and the result of Cheeger, Gromov and Taylor [15], the injectivity radius of $\tilde{g}_k(0)$ at \tilde{x}_k is bounded from below uniformly. Together with the curvature estimates from (b), we may apply Hamilton's compactness theorem [23] to conclude that, $(M_k, \tilde{g}_k(t), \tilde{x}_k)$ sub-converges in the Cheeger-Gromov sense to $(M_\infty, \tilde{g}_\infty(t), \tilde{x}_\infty)$ which is a complete non-flat ancient solution to the Ricci flow with bounded curvature. By (e), $\text{Rm}(\tilde{g}_\infty(t)) \in \text{C}_{\text{PIC}_1}$ for all $t \leq 0$. Moreover, (d) implies that it is of Euclidean volume growth. But this contradicts with [2, Lemma 4.2]. This completes the proof.

In case of (iii), each $g_k(t)$ is in addition Kähler and the conclusion (e) is replaced by $\text{OB}(\tilde{g}_k(t)) \geq -\frac{L}{t + Q_k \tilde{t}_k}$ instead so that the limiting Ricci flow $\tilde{g}_\infty(t)$ is Kähler, non-flat with Euclidean volume growth and has non-negative bounded holomorphic orthogonal bisectional curvature. This contradicts with [29, Proposition 6.1].

In case of (i), the Ricci curvature of $\tilde{g}_k(t)$ is not almost non-negative but we might instead apply Hamilton-Ivey estimates (for instances see [16, Proposition 9.8]) on the limiting Ricci flow $\tilde{g}_\infty(t)$ to conclude that $\tilde{g}_\infty(t)$ has non-negative sectional curvature and Euclidean volume growth. The non-flatness contradicts with Perelman's Theorem [34, Section 11]. \square

Thanks to Proposition 3.1, obtaining curvature estimates is equivalent to establishing the persistence of volume non-collapsing and curvature lower bound. To obtain a the curvature lower bound along the flow, we will need a version of the local maximum principle which allow the initial data to be merely unbounded. This is a modified version of the local maximum principle in [24, 27].

Proposition 3.2. *Suppose $g(t)$ is a smooth solution to the Ricci flow on $M \times [0, S]$ (not necessarily) and there exist $\alpha, r > 0$ so that all $t \in [0, S]$,*

- (a) $|\text{Rm}(g(t))| \leq \alpha t^{-1}$;
- (b) $\text{inj}(g(t)) \geq \sqrt{\alpha^{-1}t}$;
- (c) *either*
 - (i) M is compact or;
 - (ii) $B_{g_0}(x_0, 2r) \Subset B_{g(t)}(x_0, 4r) \subseteq B_{g(t)}(x_0, 16r) \Subset M$ for some $x_0 \in M$.

If φ is a continuous non-negative function on $M \times [0, S]$ with $\varphi(0) = \varphi_0$ such that

- (i) $\varphi(t) \leq \alpha t^{-1}$ on $M \times (0, S]$;
- (ii) φ satisfies

$$\left(\frac{\partial}{\partial t} - \Delta_{g(t)} \right) \varphi \leq \mathcal{R}\varphi + K\varphi^2$$

in the sense of barrier for some constant $K \geq 0$;

- (iii) There exist $\Lambda, C_1 > 0$ and a non-decreasing $f : (0, r^2] \rightarrow (0, +\infty)$ satisfying $\int_0^{r^2} s^{-1} f(s) ds \leq \Lambda$ so that for all $x \in B_{g_0}(x_0, r)$ and $t \in (0, r^2]$,

$$\int_{B_{g_0}(x_0, 2r)} \frac{1}{t^{n/2}} \exp\left(-\frac{C_1}{t} d_{g_0}^2(x, y)\right) \varphi_0(y) \, d\text{vol}_{y, g_0} \leq \frac{C_1 f(t)}{t}.$$

Then there exist $\tilde{\Lambda}(n, \alpha, f, \Lambda, C_1), \tilde{S}(K, n, \alpha, f, \Lambda, C_1) > 0$ so that for all $t \in (0, \tilde{S}r^2 \wedge S]$,

$$\varphi(x_0, t) \leq \tilde{\Lambda} \left(\frac{f(\tilde{\Lambda}t)}{t} + r^{-2} \right).$$

Proof. By considering $\tilde{g}(t) = r^{-2}g(r^2t)$, $\tilde{f}(t) = f(r^2t)$ and $\tilde{\varphi}(t) = r^2\varphi(r^2t)$, we might assume $r = 1$. For each $x \in B_{g_0}(x_0, 1)$, we define

$$\rho(x) = \sup \{r \in (0, 1) : B_{g_0}(x, r) \Subset B_{g_0}(x_0, 1)\}$$

Let $\tilde{\Lambda}, S$ be constants to be determined as well as a differentiable function $\mathfrak{F} : (0, +\infty) \rightarrow (0, +\infty)$ satisfying $\lim_{t \rightarrow 0^+} \mathfrak{F}(t) = 0$ and $\lim_{t \rightarrow 0^+} \mathfrak{F}'(t) = +\infty$. We will specify the choices in the proof.

We want to show that if $\tilde{\Lambda}$ is chosen to be large enough, \tilde{S} is sufficiently small and $\mathfrak{F}(t)$ is chosen appropriately, then for all $(x, t) \in B_{g_0}(x_0, 1) \times (0, S \wedge \tilde{S}]$,

$$\varphi(x, t) < \tilde{\Lambda} \left(\mathfrak{F}'(t) + \frac{1}{\rho^2(x)} \right).$$

Since \mathfrak{F}' is unbounded as $t \rightarrow 0^+$, there exists $t_1 \in (0, S \wedge \tilde{S}]$ such that the conclusion holds on $(0, t_1)$. If $t_1 = S \wedge \tilde{S}$, then we are done. Otherwise, there exists $x_1 \in B_{g_0}(x_0, 1)$ so that $\varphi(x_1, t_1) = \tilde{\Lambda} (\mathfrak{F}'(t_1) + \rho^{-2}(x_1))$. We denote $\rho_1 = \rho(x_1)$ so that for all $z \in B_{g_0}(x_1, \frac{1}{2}\rho_1)$, $\rho(z) \geq \frac{1}{2}\rho_1$. In particular, for all $(z, t) \in B_{g_0}(x_1, \frac{1}{2}\rho_1) \times (0, t_1]$,

$$\varphi(z, t) \leq \tilde{\Lambda} \left(\mathfrak{F}'(t) + \frac{4}{\rho^2(x)} \right).$$

Thus, if we define

$$\tilde{\varphi}(z, t) = e^{-4K\tilde{\Lambda}(\mathfrak{F}(t) + t\rho_1^{-2})} \varphi(z, t),$$

then it satisfies $\left(\frac{\partial}{\partial t} - \Delta_{g(t)} \right) \tilde{\varphi} \leq \mathcal{R}\tilde{\varphi}$ on $B_{g_0}(x_1, \frac{1}{2}\rho_1) \times (0, t_1]$.

By shrinking \tilde{S} , we might assume $t_1 \leq 1$. We might then apply [27, Proposition 4.1] to the rescaled Ricci flow $\tilde{g}(t) = t_1^{-1}g(t_1t), t \in [0, 1]$ with $\Omega = B_{g_0}(x_0, 2) = B_{\tilde{g}(0)}(x_0, 2t_1^{-1/2})$. The same also holds if M is compact, see [27, Remark 4.1]. By rescaling back, we obtain a Dirichlet heat kernel $G(x, t; y, s)$ on $\Omega \times \Omega \times [0, t_1] \times [0, t_1]$ to the operator $\partial_t - \Delta_{g(t)} - \mathcal{R}$ which satisfies

$$(3.3) \quad G(x, t; y, s) \leq \frac{C_2}{(t-s)^{n/2}} \exp\left(-\frac{d_{g(s)}^2(x, y)}{C_2(t-s)}\right)$$

for some $C_2(n, \alpha) > 0$. Consider the function

$$u(x, t) = \int_{\Omega} G(x, t; y, 0) \varphi_0(y) d\text{vol}_{g_0}(y)$$

on $\Omega \times [0, t_1]$ which satisfies $(\frac{\partial}{\partial t} - \Delta_{g(t)})u = \mathcal{R}u$. If $\tilde{S} \leq \min\{1, C_2^{-1}C_1\}$, then the heat kernel estimate (3.3) shows that for $x \in B_{g_0}(x_0, 1)$ and $t \in [0, t_1]$,

$$(3.4) \quad \begin{aligned} u(x, t) &= \int_{\Omega} G(x, t; y, 0) \varphi_0(y) d\text{vol}_{g_0}(y) \\ &\leq \int_{B_{g_0}(x_0, 2)} \varphi_0(y) \cdot \frac{C_2}{t^{n/2}} \exp\left(-\frac{d_{g_0}^2(x, y)}{C_2 t}\right) d\text{vol}_{g_0}(y) \\ &\leq C_2 C_1^{-1} \frac{f(C_2 C_1^{-1} t)}{t}. \end{aligned}$$

Thanks to the assumption of f , we have $f(t) \leq \Lambda(\log(t^{-1}))^{-1}$. Hence by further shrinking $\tilde{S} \leq \min\{C_2^{-1}C_1, 1, \exp(-\Lambda C_2 C_1^{-1})\}$, we have $u \leq t^{-1}$ for $t \in (0, t_1]$. By [27, Theorem 1.1], we deduce that there exists $T_1(n, \alpha) > 0$ such that if $t_1 \leq \tilde{S} \wedge S \wedge T_1 \rho_1^2$, then

$$(3.5) \quad \tilde{\varphi}(x_1, t_1) \leq u(x_1, t_1) + 16(t_1 \rho_1^{-2}) \cdot \rho_1^{-2}$$

We choose

$$\mathfrak{F}(t) = C_2 C_1^{-1} \int_0^{C_2 C_1^{-1} t} \left(\frac{f(s)}{s} + \varepsilon s^{-1/2} \right) ds$$

for $\varepsilon > 0$ so that $\mathfrak{F}'(t) \rightarrow +\infty$ as $t \rightarrow 0^+$. Moreover, $\lim_{t \rightarrow 0^+} \mathfrak{F}(t) = 0$ follows from absolute continuity of integrable function and assumption on f .

On the other hand, using assumption (i),

$$\frac{\tilde{\Lambda}}{\rho_1^2} < \tilde{\Lambda} \left(\mathfrak{F}'(t_1) + \frac{1}{\rho_1^2} \right) \leq \frac{\alpha}{t_1}$$

giving $t_1 \leq \alpha \tilde{\Lambda}^{-1} \rho_1^2$ and hence

$$(3.6) \quad \begin{aligned} &4\tilde{\Lambda} e^{-4K\tilde{\Lambda}\mathfrak{F}(t_1) - 4K\alpha} \cdot (\mathfrak{F}'(t_1) + \rho_1^{-2}) \\ &\leq \tilde{\varphi}(x_1, t_1) \leq u(x_1, t_1) + 16\alpha \tilde{\Lambda}^{-1} \rho_1^{-2} \\ &\leq \mathfrak{F}'(t_1) + 16\alpha \tilde{\Lambda}^{-1} \rho_1^{-2} \end{aligned}$$

By choosing $\tilde{\Lambda}(n, \alpha, K) > 1$ sufficiently large so that $\tilde{\Lambda}^2 e^{-4\alpha K} > 16$, we see that $t_1 > S_2$ for some $S_2(K, \tilde{\Lambda}, C_2, C_1) > 0$ which is independent of $\varepsilon \rightarrow 0$. By choosing $\tilde{S} = \frac{1}{2} \min\{S_2, C_2^{-1}C_1, 1, \exp(-\Lambda C_2 C_1^{-1})\}$, we see that the conclusion holds on $B_{g_0}(x_0, 1) \times (0, S \wedge \tilde{S}]$. This completes the proof by evaluating at x_0 . \square

We end this section by pointing out that the lowest eigenvalues of Rm with respect to various curvature cones satisfy the evolution equation required in Proposition 3.2.

Lemma 3.1. *Let $g(t)$ be a smooth solution to the Ricci flow. Suppose \mathcal{C} is one of the following curvature cone:*

- (i) *non-negative curvature operator;*
- (ii) *2-non-negative curvature operator;*
- (iii) *weakly PIC₂;*
- (iv) *weakly PIC₁;*
- (v) *non-negative bisectional curvature in case $g(t)$ is Kähler ;*
- (vi) *non-negative orthogonal bisectional curvature OB in case $g(t)$ is Kähler;*
- (vii) *non-negative Ricci curvature if $n = 3$,*

then the function

$$\varphi(x, t) = \inf\{s > 0 : \text{Rm}(x, t) + s \cdot \text{Id} \in \mathcal{C}\}$$

is non-negative continuous and satisfies $(\frac{\partial}{\partial t} - \Delta_{g(t)})\varphi \leq \mathcal{R}\varphi + c_n\varphi^2$ in the sense of barrier for some dimensional constant $c_n \geq 0$.

Proof. When $n \geq 4$, the case of (i)-(v) follows from [2] while (vi) is proved in [29]. It remains to prove (vii) which is equivalent to (iv) when $n = 3$. By [40], if we use φ to denote the negative part of the Ricci lower bound when $n = 3$, then whenever $\varphi \geq 0$,

$$(3.7) \quad \left(\frac{\partial}{\partial t} - \Delta_{g(t)}\right)\varphi = -(\mu - \nu)^2 + \varphi(\mu + \nu)$$

where $\mu \geq \nu \geq -\varphi$ denotes the eigenvalues of $\text{Ric}(x, t)$. Since $\mu + \nu - \varphi = \mathcal{R}$, results follows with $c_3 = 1$. \square

4. VOLUME NON-COLLAPSING ALONG FLOW

In view of Proposition 3.1, the volume non-collapsing along the flow plays an important role to avoid curvature blowup. In this section, we will show that under the mildly integrable curvature lower bound, if the initial metric is volume non-collapsed, then so does the flow for a uniform short time. We rely heavily the idea of Simon-Topping [31] which considered the case when the Ricci curvature is uniformly bounded from below along the flow.

We start with a distance distortion under a weaker condition. Namely, $\text{Ric}(g(t)) \geq -\phi$ for some $\phi \in L^1([0, S]) \cap C_{loc}^0((0, S])$ along the flow. In practice, it will be sufficient to take $\phi(t) = t^{-1}(-\log t)^{1+\frac{1}{2}}$.

Lemma 4.1. *Suppose $(M^n, g(t)), t \in [0, S]$ is a smooth solution to the Ricci flow (not necessarily complete) such that for some $x_0 \in M$, we have $B_{g(t)}(x_0, 5) \Subset M$ for all $t \in [0, S]$. Then the following holds.*

- (a) *If the Ricci curvature satisfies $\text{Ric}(g(x, t)) \geq -\phi(t)$ for some non-negative function $\phi(t) \in L^1([0, S]) \cap C_{loc}^0((0, S])$ and for all $x \in B_{g(t)}(x_0, 5)$, $t \in (0, S]$, then there exists $\tilde{S} > 0$ depending only on ϕ such that for any $x, y \in B_{g(0)}(x_0, 1)$ and $0 \leq s < t \leq S \wedge \tilde{S}$, we have*

$$d_{g(t)}(x, y) \leq \exp\left(\int_s^t \phi(z) dz\right) \cdot d_{g(s)}(x, y).$$

- (b) *If $|\text{Rm}(x, t)| \leq \alpha t^{-1}$ on $B_{g(t)}(x_0, 5)$ for $t \in (0, S]$, then for all $0 \leq s \leq t \leq S$,*

$$B_{g(s)}(x_0, 1 - \beta_n \sqrt{\alpha s}) \supset B_{g(t)}(x_0, 1 - \beta_n \sqrt{\alpha t}).$$

Proof. We first prove (a) following the idea in [31, Lemma 3.1]. Define

$$S_1 = \sup\{s \in [0, S] : B_{g_0}(x_0, 1) \subset B_{g(t)}(x_0, 5)\}.$$

Hence, for $t \in (0, S_1]$, we have Ricci lower bound $\phi(t)$. In particular, for all $x \in B_{g_0}(x_0, 1)$ and $t \in (0, S_1]$,

$$(4.1) \quad d_{g(t)}(x, x_0) \leq \exp\left(\int_0^t \phi(s) ds\right) \cdot d_{g_0}(x, x_0).$$

Hence, there exists $\tilde{S}(\phi) > 0$ such that $S_1 \geq S \wedge \tilde{S}$. The conclusion follows using the Ricci lower bound again. The property (b) follows directly from [31, Corollary 3.3]. This completes the proof. \square

The next result shows that the volume non-collapsing will be preserved in the scale of \sqrt{t} under the scaling invariant curvature bound and L^1 (in time) Ricci lower bound.

Proposition 4.1. *Suppose $(M, g(t))$ is a Ricci flow for $t \in [0, S]$ (not necessarily complete) such that $g(0) = g_0$. If for some $x_0 \in M$, $B_{g(t)}(x_0, 10) \Subset M$ for all $t \in [0, S]$ and for some $v_0, \alpha > 0$, we have*

- (i) $v_0^{-1} r^n \geq \text{Vol}_{g_0}(B_{g_0}(x, r)) \geq v_0 r^n$ for $r \in (0, 1]$ and $x \in B_{g_0}(x_0, 5)$;
- (ii) $|\text{Rm}(x, t)| \leq \alpha t^{-1}$ on $B_{g(t)}(x_0, 10)$ for $t \in (0, S]$;
- (iii) for all $x \in B_{g(t)}(x_0, 10)$ and $t \in (0, S]$, the Ricci curvature satisfies

$$\text{Ric}(g(x, t)) \geq -\phi(t)$$

for some non-negative decreasing function $\phi(t) \in L^1([0, S]) \cap C_{loc}^0((0, S])$.

Then there exist $\tilde{\varepsilon}_0(n, v_0, \phi)$ and $\tilde{S}(n, v_0, \alpha, \phi) > 0$ such that for all $t \in (0, \tilde{S} \wedge S]$ and $x \in B_{g(t)}(x_0, 1)$,

$$\text{Vol}_{g(t)}\left(B_{g(t)}(x, \sqrt{t})\right) \geq \tilde{\varepsilon}_0 t^{n/2}.$$

Remark 4.1. In the presence of uniform Ricci lower bound, the Ahlfors n -regularity, i.e. assumption (i), follows easily from volume lower bound of $B_{g_0}(x_0, 10)$ and volume comparison theorem.

Proof. The proof is almost identical to that of [31, Lemma 11.1]. We only give a sketch and point out the necessary modifications. We first note that since $\phi \in L^1([0, S])$, for any $\varepsilon > 0$, there exists S_ε such that if $0 < t < S_\varepsilon$,

$$(4.2) \quad \varepsilon \geq \int_t^{2t} \phi(s) ds \geq t\phi(t).$$

Hence, as $t \rightarrow 0^+$, the Ricci lower bound behaves as

$$(4.3) \quad \text{Ric}(g(t)) \geq -\phi(t) \geq -o(1)t^{-1}.$$

This serves as the crucial replacement of the uniform Ricci lower bound assumption in [31, Lemma 11.1].

With this replacement in mind, we can carry out the exact same argument in the proof of [31, Lemma 11.1] to obtain the following. If the conclusion fails, then for any sufficiently small $\tilde{\varepsilon}_0$, there exist a sequence of n -manifold M_i , a sequence of point $x_i \in M_i$, a sequence of Ricci flows $\tilde{g}_i(t)$ on $M_i \times [0, \tilde{t}_i]$ with \tilde{t}_i decreases to 0, $t_i \in (0, \tilde{t}_i]$ and a sequence of points $y_i \in B_{\tilde{g}_i(t_i)}(x_i, 1)$ such that

- (i) $B_{\tilde{g}_i(t)}(y_i, 7) \Subset M_i$ for all $t \in [0, t_i]$;
- (ii) $\text{Ric}(\tilde{g}_i(t)) \geq -\phi(t)$ on $B_{\tilde{g}_i(t)}(y_i, 7)$ for all $t \in [0, t_i]$;
- (iii) $|\text{Rm}(\tilde{g}_i(t))| \leq \alpha t^{-1}$ on $B_{\tilde{g}_i(t)}(y_i, 7)$ for all $t \in [0, t_i]$;
- (iv) $v_0^{-1}r^n \geq \text{Vol}_{\tilde{g}_i(0)}(y, r) \geq v_0r^n$ for all $r \in (0, 1)$ and $y \in B_{\tilde{g}_i(0)}(y_i, 1)$;
- (v) $\text{Vol}_{\tilde{g}_i(t_i)}(y_i, \sqrt{t_i}) = \tilde{\varepsilon}_0(\sqrt{t_i})^n$;
- (vi) $\text{Vol}_{\tilde{g}_i(t)}(y, \sqrt{t}) \geq 2^{-n-1}\tilde{\varepsilon}_0(\sqrt{t})^n$ for $t \in [0, t_i]$ and all $y \in B_{\tilde{g}_i(t)}\left(y_i, c_n\alpha^{-1}\sqrt{\frac{t_i}{t}}\right)$;

We remark here that since $t_i \rightarrow 0$, [31, (11.15)] now follows from (4.3) when i is sufficiently large. Now consider the rescaled Ricci flow $g_i(t) = t_i^{-1}\tilde{g}_i(t_i t)$, $t \in [0, 1]$ so that the smooth sub-sequential Cheeger-Gromov limit $(M_\infty, g_\infty(t), y_\infty)$ of $(M_i, g_i(t), y_i)$, $t \in (0, 1]$ satisfies

- (I) $\text{Ric}(g_\infty(t)) \geq 0$ for all $t \in (0, 1]$;
- (II) $|\text{Rm}(g_\infty(t))| \leq \alpha t^{-1}$ for all $t \in (0, 1]$;
- (III) $\text{Vol}_{g_\infty(1)}(B_{g_\infty(1)}(y_\infty, 1)) = \tilde{\varepsilon}_0$;
- (IV) $\text{Vol}_{g_\infty(t)}(B_{g_\infty(t)}(y, \sqrt{t})) \geq 2^{-n-1}\tilde{\varepsilon}_0 t^{n/2}$ for all $t \in (0, 1]$ and $y \in M_\infty$;
- (V) For any $\hat{R} > 2r > 0$ and $z_i \in B_{g_i(0)}(y_i, \hat{R})$,

$$\limsup_{i \rightarrow +\infty} \text{Vol}_{\tilde{g}_i(0)}(B_{\tilde{g}_i(0)}(z_i, r)) \leq v_0^{-1}r^n.$$

Here the property (I) follows from (4.3) while (V) follows from the initial Ahlfors n -regularity assumption. Moreover, using the same proof where [31, Lemma 3.4] is now replaced by Lemma 4.1, the **Claim 1** in [31, page 513] also holds for $g_\infty(1)$ in the sense that if we have a ball $B_{g_\infty(1)}(y, L)$ for $y \in M_\infty$ and $L > 0$, that is covered by N balls of radius $r \geq R(n, \alpha)$, then we must have

$N \geq \eta(n) \cdot v_0^2 L^n r^{-n}$. Now we are in position to apply [31, Lemma 11.2] as in the proof of [31, Lemma 11.1] to derive contradiction when $\tilde{\varepsilon}_0$ is chosen to be too small. \square

Finally, we show that the volume non-collapsing is indeed preserved in a large scale for a uniform short time.

Proposition 4.2. *Under the assumption of Proposition 4.1, there exist $\tilde{v}(n, v_0, \phi)$ and $\tilde{S}(n, v_0, \alpha, \phi) > 0$ such that for all $t \in (0, S \wedge \tilde{S}]$ and $x \in B_{g(t)}(x_0, 1)$,*

$$\text{Vol}_{g(t)}(B_{g(t)}(x_0, 1)) \geq \tilde{v}.$$

Proof. Since $B_{g(t)}(x_0, 1)$ is compactly contained inside M , we might take finitely many $\{B_{g(t)}(x_i, \beta_n \sqrt{\alpha t})\}_{i=1}^N$ such that $B_{g(t)}(x_i, \beta_n \sqrt{\alpha t}) \cap B_{g(t)}(x_j, \beta_n \sqrt{\alpha t}) = \emptyset$ for $i \neq j$ and $B_{g(t)}(x_i, \beta_n \sqrt{\alpha t}) \subseteq B_{g(t)}(x_0, 1)$ while $\{B_{g(t)}(x_i, 4\beta_n \sqrt{\alpha t})\}_{i=1}^N$ is a cover of $B_{g(t)}(x_0, 1)$. Therefore, Proposition 4.1 implies

$$(4.4) \quad \text{Vol}_{g(t)}(B_{g(t)}(x_0, 1)) \geq \sum_{i=1}^N \text{Vol}_{g(t)}(B_{g(t)}(x_i, \beta_n \sqrt{\alpha t})) \geq N\gamma(\beta_n^2 \alpha t)^{n/2}.$$

It remains to estimate N from below. By Lemma 4.1, we might restrict S so that

$$(4.5) \quad B_{g_0}\left(x_0, \frac{1}{2}\right) \subset B_{g(t)}(x_0, 1) \subset \bigcup_{i=1}^N B_{g(t)}(x_i, 4\beta_n \sqrt{\alpha t}) \subset \bigcup_{i=1}^N B_{g_0}(x_i, 5\beta_n \sqrt{\alpha t}).$$

By measuring the set using $d\text{vol}_{g_0}$ and volume growth assumption of g_0 , we deduce that

$$(4.6) \quad C_0^{-2} v_0 \leq 5^n N (\beta_n \sqrt{\alpha t})^n$$

This completes the proof by substituting it back to (4.4). \square

5. RICCI FLOW SMOOTHING ON COMPACT MANIFOLDS

In this section, we establish a existence theory of Ricci flow and use it to prove the stability.

Proof of Theorem 1.1. By the short-time existence Theorem of Hamilton [40], there exists a maximal solution $g(t)$, $t \in [0, T_{max})$ to the Ricci flow on M . Let α to be a large constant to be determined. For the given α , we let s_0 be the supremum of $s \in [0, T_{max})$ such that for all $(x, t) \in M \times [0, s)$,

- (i) $|\text{Rm}(x, t)| < \alpha t^{-1}$;
- (ii) $\text{inj}(g(x, t)) > \sqrt{\alpha^{-1} t}$.

By the smoothness of flow, we have $s_0 \in (0, T_{max})$. Hence, the condition (i)-(ii) hold on $[0, s_0)$ and there exists $x_0 \in M$ such that either

- (I) $|\text{Rm}(x_0, s_0)| = \alpha s_0^{-1}$; or
- (II) $\text{inj}(g(x_0, s_0)) = \sqrt{\alpha^{-1} s_0}$.

By Lemma 3.1 and Lemma 2.1, $\varphi = (\text{IC}_1)_-$ satisfies the assumptions in Proposition 3.2 with $r = \sqrt{T}$. Hence, it applies to see that there exist $S_1(n, \alpha, f, T, \Lambda), \tilde{\Lambda}(n, \alpha, f, T, \Lambda) > 0$ so that for all $(x, t) \in M \times (0, S_1 \wedge s_0]$,

$$\varphi(x, t) \leq \tilde{\Lambda} \left[\frac{f(\tilde{\Lambda}t)}{t} + 1 \right] \leq \tilde{\Lambda} \left[1 + \frac{1}{t} \left(\frac{\Lambda}{\log(T/\tilde{\Lambda}) + \log t^{-1}} \right)^2 \right]$$

where we have used the fact that f satisfies (2.2):

$$(5.1) \quad \Lambda \geq \int_t^T \frac{\sqrt{f(s)}}{s} ds \geq \sqrt{f(t)} \cdot \log \left(\frac{T}{t} \right).$$

By taking $\phi(t) = t^{-1}(-\log t)^{1+\frac{1}{2}} \in L^1([0, T]) \cap C_{loc}^0((0, T])$, we see that if we shrink S_1 further (depending also on $\tilde{\Lambda}$), then for all $(x, t) \in M \times (0, S_1 \wedge s_0]$,

$$(5.2) \quad \varphi(x_0, t) \leq \tilde{\Lambda} \left[1 + \frac{1}{t} \left(\frac{\Lambda}{\log(T/\tilde{\Lambda}) + \log t^{-1}} \right)^2 \right] \leq \phi(t).$$

Together with Proposition 2.2 and assumption on the volume non-collapsing, we might apply Proposition 4.1 to conclude that there exists $\gamma(n, f, T, \Lambda, v) > 0$ such that for all $(x, t) \in M \times (0, S_1 \wedge s_0]$,

$$(5.3) \quad \text{Vol}_{g(t)} \left(B_{g(t)}(x, \sqrt{t}) \right) \geq \gamma t^{n/2}.$$

Moreover, the bound of φ at the same time implies $\text{Ric}(x, t) \geq -(n-1)\phi(t) \geq -t^{-1}$ as $t \rightarrow 0^+$ and hence applying the volume comparison to $t^{-1}g(t)$ implies that for all $(x, t) \in M \times (0, S_1 \wedge s_0]$ and $r \leq \frac{1}{4}\sqrt{t}$,

$$\text{Vol}_{g(t)} \left(B_{g(t)}(x, r) \right) \geq \tilde{v} r^n$$

for some $\tilde{v}(n, f, v, D, T, \Lambda) > 0$ independent of α .

Now we can apply Proposition 3.1 to show that for all $(x, t) \in M \times (0, S_1 \wedge s_0]$,

$$(5.4) \quad |\text{Rm}(x, t)| \leq \beta_0 t^{-1}$$

for some $\beta_0(n, f, v, D, T, \Lambda) > 0$. By combining (5.4), (5.3) and injectivity radius estimate of Cheeger-Gromov-Taylor [15], we conclude that for all $(x, t) \in M \times (0, S_1 \wedge s_0]$, $\text{inj}(x, t) \geq \sqrt{\beta_1^{-1}t}$ for some $\beta_1(n, f, v, D, T, \Lambda, \beta_0) > 0$. Therefore, if we fix $\alpha = \beta_1 + \beta_0$, then we see that $s_0 > S_1$. Hence, the conclusion holds on $(0, S_1]$ for this choice of α and the corresponding $S_1 > 0$. This proves (a) and (c) while (d) follows from (5.2). Finally (b) follows from Proposition 4.2 and covering argument after possibly shrinking S_1 further. This completes the proof. \square

In higher dimension, when g_0 is in addition Kähler, one usually consider the holomorphic curvature which is more complex geometric. We say that the holomorphic bisectional curvature BK of a Kähler curvature tensor R is non-negative (denoted by $R \in \text{C}_{\text{BK}}$) if for any $X, Y \in T^{1,0}M$, we have

$$R(X, \bar{X}, Y, \bar{Y}) \geq 0.$$

If one instead ask for the non-negativity only for $X, Y \in T^{1,0}M$ such that $g(X, \bar{Y}) = 0$, then we say that the orthogonal bisectional curvature OB is non-negative (denoted by $R \in C_{OB}$). One might then analogously define the class of compact Kähler manifolds satisfying the strong Kato-type lower bound on BK (or OB). Define for $t > 0$,

$$\kappa_{t, \text{BK}}(M^n, g_0) = \sup_{x \in M} \int_0^t \int_M H_{g_0}(x, y, s) (\text{BK})_-(y) \, d\text{vol}_{g_0}(y) \, ds$$

where $\text{BK}_-(x) = \inf\{\alpha \geq 0 : \text{Rm}_{g(x)} + \alpha B \in C_{\text{BK}}\}$ and B is the Kähler curvature tensor given by $B(X, \bar{Y}, Z, \bar{W}) = g(X, \bar{Y})g(Z, \bar{W}) + g(X, \bar{W})g(Z, \bar{Y})$. In this case, the same proof of Theorem 1.1 yields the following.

Theorem 5.1. *Suppose (M^n, g_0, x_0) is a pointed compact Kähler manifold with $\dim_{\mathbb{C}}(M) = n \geq 2$. Suppose $(M^n, g_0, x_0) \in \mathcal{K}_{OB}(n, f, v) \cap \mathcal{K}(n, f, v)$ for some non-decreasing function f , T and Λ satisfying (1.2) and $\text{diam}(M, g_0) \leq D$ for some $D > 0$, then there exist S, α, \tilde{v} depending only on n, f, v, D, T, Λ and a solution $g(t), t \in [0, S]$ to the Kähler-Ricci flow such that for all $(x, t) \in M \times (0, S]$,*

- (a) $|\text{Rm}(x, t)| \leq \alpha t^{-1}$;
- (b) $\text{Vol}_{g(t)}(B_{g(t)}(x, \sqrt{T})) \geq \tilde{v} T^{n/2}$;
- (c) $\text{inj}(g(x, t)) \geq \sqrt{\alpha^{-1}t}$;
- (d) $\text{Rm}(g(x, t)) + \alpha(1 + t^{-1}f(\alpha t)) \cdot B \in C_{\text{BK}}$.

When a sequence of compact manifold (M_i^n, g_i) satisfy a uniform Dynkin-type lower bound of Ricci, it is already shown by Carron [5] that the sequence admits a Gromov-Hausdorff limit after passing to subsequence. The structure theory was further investigated by Carron-Mondello-Tewodrose [6, 7] in the non-collapsed case when the Dynkin-type lower bound of Ricci is strengthened to be a strong Kato-type lower bound. We now shows that in dimension three, the limit will be Gromov-Hausdorff close and homeomorphic to a smooth manifold. Moreover, the result also holds in higher dimension if we require the strong Kato bound on 1-isotropic curvature.

Proof of Corollary 1.1. The proof is almost identical to that of [2, Corollary 4]. We include it for reader's convenience. By Theorem 1.1, there exist $\alpha, S > 0$ and $\phi \in L^1([0, S]) \cap C_{loc}^0((0, S))$ such that for each $i \in \mathbb{N}$, we can find a Ricci flow $g_i(t), t \in [0, S]$ such that $g_i(0) = g_{i,0}$ and for all $t \in (0, S]$,

- (a) $|\text{Rm}(g_i(t))| \leq \alpha t^{-1}$;
- (b) $\text{inj}(g_i(t)) \geq \sqrt{\alpha^{-1}t}$;
- (c) $\text{IC}_1(g_i(t)) \geq -\phi(t)$.

It follows from Hamilton's compactness [23] that after passing to subsequence, there exists a smooth Ricci flow $(M_\infty, g_\infty(t), x_\infty), t \in (0, S]$ such that $(M_i, g_i(t), x_i)$ converges to $(M_\infty, g_\infty(t), x_\infty)$ in the C^∞ Cheeger-Gromov sense.

By Lemma 4.1, for all i and $0 < s \leq t \leq S$

$$\exp\left(-\int_s^t \phi(z) dz\right) \cdot d_{g_i(t)} \leq d_{g_i(s)} \leq d_{g_i(t)} + C(\sqrt{t} - \sqrt{s}).$$

If we first let $i \rightarrow +\infty$ and followed by $s \rightarrow 0$, we see that M_∞ is compact since $\text{diam}(M_i, g_{i,0}) \leq D$ and $d_{g_\infty(s)} \rightarrow d_\infty$ for some distance function d_∞ on M_∞ . By interchanging the order of taking limit, it is easy to see that $d_{g_{i,0}}$ converges to d_∞ modulus diffeomorphism. This proved the Gromov-Hausdorff convergence. It follows from [6, Theorem A] that the convergence is also in the measured Gromov-Hausdorff sense.

Suppose now $(M_i, g_{i,0}, x_i) \in \mathcal{K}_{\text{IC}_1}(n, f_i, v)$ so that $f_i(T)$ decreases to 0. We want to show that $g_\infty(t)$ satisfies a stronger curvature estimate.

Fix any $x_i \in M$. If $r \geq D$ where $\text{diam}(M_i, g_{i,0}) \leq D$, then for all $x \in B_{g_{i,0}}(x_i, r) = M_i$ and $t \in (0, r^2 \wedge T]$,

$$(5.5) \quad \begin{aligned} & \int_{B_{g_{i,0}}(x_i, 2r)} \frac{1}{t^{n/2}} \exp\left(-\frac{C_1}{t} d_{g_{i,0}}^2(x, y)\right) \varphi_{i,0}(y) d\text{vol}_{g_{i,0}}(y) \\ &= \int_{M_i} \frac{1}{t^{n/2}} \exp\left(-\frac{C_1}{t} d_{g_{i,0}}^2(x, y)\right) \varphi_{i,0}(y) d\text{vol}_{g_{i,0}}(y) \leq \frac{C_1 f_i(t)}{t} \end{aligned}$$

by Lemma 2.1 and assumption. While if $t \in [T, r^2]$, then the diameter bound yields

$$(5.6) \quad \begin{aligned} & \int_{B_{g_{i,0}}(x_i, 2r)} \frac{1}{t^{n/2}} \exp\left(-\frac{C_1}{t} d_{g_{i,0}}^2(x, y)\right) \varphi_{i,0}(y) d\text{vol}_{g_{i,0}}(y) \\ &\leq e^{C_1 D^2(T^{-1}-t^{-1})} \left(\frac{T}{t}\right)^{n/2} \int_{M_i} \frac{1}{T^{n/2}} \exp\left(-\frac{C_1}{T} d_{g_{i,0}}^2(x, y)\right) \varphi_{i,0}(y) d\text{vol}_{g_{i,0}}(y) \\ &\leq C_1 e^{C_1 D^2(T^{-1}-t^{-1})} \frac{f_i(T)}{T}. \end{aligned}$$

For $t \in [T, r^2]$, we extend $f_i(t)$ by defining

$$f_i(t) = t \cdot e^{C_1 D^2(T^{-1}-t^{-1})} \frac{f_i(T)}{T}.$$

Since $f_i(T) \rightarrow 0$ as $i \rightarrow +\infty$, f_i satisfies the assumption (iii) in Proposition 3.2 for $r \rightarrow +\infty$ in the sense that for any $r \geq D$, there exists $N \in \mathbb{N}$ so that for all $i > N$, Proposition 3.2 applies on $B_{g_0}(x_i, r)$ so that

$$(\text{IC}_1)_-(g_i(x_i, t)) \leq \tilde{\Lambda} \left(\frac{f_i(\tilde{\Lambda}t)}{t} + \frac{1}{r^2} \right)$$

for some uniform $\tilde{\Lambda} > 0$. Since x_i is arbitrary in M_i , by letting $i \rightarrow +\infty$ for $t \in (0, S]$ and followed by letting $r \rightarrow +\infty$, we see that $\text{Rm}(g_\infty(t)) \in \mathcal{C}_{\text{PIC}_1}$ on M_∞ for $t \in (0, S]$. This completes the proof. \square

As another application, we obtain a stability in view of Geroch conjecture. This is motivated by the work of [1] where compactness under uniform L^p curvature bound for $p = 3$ is studied when $n = 3$. Since a $L^p, p > \frac{3}{2}$ lower bound on Ric implies the uniform strong Kato bound of Ric [38], together with the regularity theory of Ricci flow (for instances, see [44, 45]), we can view Corollary 1.2 as a generalization.

Proof of Corollary 1.2. By Corollary 1.1 it suffices to identify the Gromov-Hausdorff limit. Recall that there exists $(M_\infty, g_\infty(t), x_\infty), t \in (0, S]$ such that the Ricci flow $(M_i^n, g_i(t), x_i)$ with $g_i(0) = g_{i,0}$ converges to $(M_\infty, g_\infty(t), x_\infty)$ in the smooth Cheeger-Gromov sense after passing to subsequent. By maximum principle, we have $\mathcal{R}(g_i(t)) \geq -i^{-1}$ for all i and hence $\mathcal{R}(g_\infty(t)) \geq 0$. Since M_∞ is compact, the smooth convergence implies that M_∞ is diffeomorphic to M_i for i sufficiently large. In particular, $\sigma(M_\infty) \leq 0$ and hence $g_\infty(t)$ is Ricci flat for $t \in (0, S]$ by strong maximum principle. Thus, $g_\infty = g_\infty(S)$ is desired limit.

If $b_1(M_i) = n$ for all i , we also have $b_1(M_\infty) = b_1(M_i) = n$ for i large. Since $g_\infty(t)$ is of Ricci flat, it is the standard torus by the classical Bochner theorem. This completes the proof. \square

REFERENCES

- [1] Allen, B.; Bryden, E.; Kazaras, D., *Stability of the positive mass theorem and torus rigidity theorems under integral curvature bounds*, arXiv:2210.04340
- [2] Bamler, R. H. ; Cabezas-Rivas, E.; Wilking, B., *The Ricci flow under almost nonnegative curvature conditions*, Invent. Math. 217 (2019) 95–126
- [3] Brendle, S., *Ricci Flow and the Sphere Theorem*, Graduate Studies in Mathematics vol. 111, American Mathematical Society (2010)
- [4] Brendle, S., *Ricci flow with surgery on manifolds with positive isotropic curvature*. Ann. of Math. (2) 190 (2019), no. 2, 465–559.
- [5] Carron, G. , *Geometric inequalities for manifolds with Ricci curvature in the Kato class*. Ann. Inst. Fourier (Grenoble), 69(7):3095–3167, 2019.
- [6] Carron, G.; Mondello, I. ; Tewodrose, D., *Limits of manifolds with a Kato bound on the Ricci curvature*, preprint, arXiv:2102.05940.
- [7] Carron, G.; Mondello, I. ; Tewodrose, D., *Limits of manifolds with a Kato bound on the Ricci curvature. II*, preprint, arXiv:2205.01956.
- [8] Carron, G.; Mondello, I. ; Tewodrose, D., *Torus stability under Kato bounds on the Ricci curvature*. J. Lond. Math. Soc. (2) 107 (2023), no. 3, 943–972.
- [9] Carron, G.; Rose, C. , *Geometric and spectral estimates based on spectral Ricci curvature assumptions*. J. Reine Angew. Math., 772:121–145, 2021.
- [10] Cheeger, J. ; Colding, T. H. , *On the structure of spaces with Ricci curvature bounded below. I*. J. Differential Geom., 46(3):406–480, 1997.
- [11] Cheeger, J. ; Colding, T. H. , *On the structure of spaces with Ricci curvature bounded below. II*. J. Differential Geom., 54(1):13–35, 2000.
- [12] Cheeger, J. ; Colding, T. H. , *On the structure of spaces with Ricci curvature bounded below. III*. J. Differential Geom., 54(1):37–74, 2000.
- [13] Cheeger, J.; Naber, A. , *Regularity of Einstein manifolds and the codimension 4 conjecture*. Ann. of Math. (2), 182(3):1093–1165, 2015.

- [14] Cheeger, J.; Jiang, W.; Naber, A. , *Rectifiability of singular sets of noncollapsed limit spaces with Ricci curvature bounded below*. Ann. of Math. (2), 193(2):407–538, 2021.
- [15] Cheeger, J.; Gromov, M.; Taylor, M. , *Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds*, J. Differential Geom. 17 (1982) 15–53
- [16] Chow, B. ; Knopf, D., *The Ricci flow: an introduction*. Mathematical Surveys and Monographs, 110. American Mathematical Society, Providence, RI, 2004.
- [17] Colding, T. H., *Ricci curvature and volume convergence*. Ann. of Math. (2), 145(3):477–501, 1997.
- [18] Dai, X.;Wei, G.; Zhang, Z. , *Local Sobolev constant estimate for integral Ricci curvature bounds*. Adv. Math., 325:1–33, 2018.
- [19] Gromov, M.; Lawson, H. B., Jr., *Spin and scalar curvature in the presence of a fundamental group. I*, Ann. of Math. (2) 111 (1980), no. 2, 209–230.
- [20] Gromov, M., *Metric structures for Riemannian and non-Riemannian spaces*. Based on the 1981 French original. With appendices by M. Katz, P. Pansu and S. Semmes. Translated from the French by Sean Michael Bates. Progress in Mathematics, 152. Birkhauser Boston, Inc., Boston, MA, 1999.
- [21] Gromov, M., *Dirac and Plateau billiards in domains with corners*, Cent. Eur. J. Math. 12 (2014), no. 8, 1109–1156.
- [22] Huang, S.; Wang, B. ,*Ricci flow smoothing for locally collapsing manifolds*. Calc. Var. Partial Differential Equations 61 (2022), no. 2, Paper No. 64, 32 pp.
- [23] Hamilton R.-S., *A compactness property for solutions of the Ricci flow*, American J. Math. 117 (1995) 545–572.
- [24] Hochard, R., *Theorèmes d’existence en temps court du flot de Ricci pour des variétés non-complètes, non-effondrées, à courbure minorée*, PhD thesis, Université de Bordeaux (2019) Available at <https://hal.inria.fr/tel-02092609/>
- [25] Lai, Y., *Ricci flow under local almost nonnegative curvature conditions*, Adv. Math. 343 (2019) 353–392
- [26] Lee, M.-C.; Tam, L.-F., *Kähler manifolds with almost non-negative curvature*, Geom. Topol. 25–4 (2021), 1979–2015.
- [27] Lee, M.-C.; Tam, L.-F., *Some local maximum principles along Ricci flows*. Canad. J. Math. 74 (2022), no. 2, 329–348.
- [28] Lee, M.-C.; Topping, P.M., *Three-manifolds with non-negatively pinched Ricci curvature*. arXiv:2204.00504
- [29] Li, X. ; Ni, L., *Kähler-Ricci Shrinkers and ancient solutions with nonnegative orthogonal bisectional curvature*. J. Math. Pure Appl. 138(2020), 28–45.
- [30] Ma, Y.; Wang, B., *Ricci curvature integrals, local functionals, and the Ricci flow*, arXiv:2109.02449v1.
- [31] Simon, M.; Topping, P. M. , *Local control on the geometry in 3D Ricci flow*, J. Differential Geom. 122 (2022), no. 3, 467–518.
- [32] Simon, M.; Topping, P. M. , *Local mollification of Riemannian metrics using Ricci flow, and Ricci limit spaces*. Geom. Topol. 25 (2021), no. 2, 913–948.
- [33] Wang, B., *The local entropy along Ricci flow -Part B: the pseudo-locality theorems*, arXiv:2010.09981.
- [34] Perelman, G.,*The entropy formula for the Ricci flow and its geometric applications*. <http://arXiv.org/abs/math/0211159v1> (2002)
- [35] Petersen, P.; Wei, G. , *Relative volume comparison with integral curvature bounds*. Geom. Funct. Anal., 7(6):1031–1045, 1997.
- [36] Rose C., *Li-Yau gradient estimate for compact manifolds with negative part of Ricci curvature in the Kato class*. Ann. Global Anal. Geom. 55 (2019), no. 3, 443–449.

- [37] Rose, C., *Almost positive Ricci curvature in Kato sense—an extension of Myers’ theorem*. Math. Res. Lett. 28 (2021), no. 6, 1841–1849.
- [38] Rose C.; Stollmann, P., *The Kato class on compact manifolds with integral bounds on the negative part of Ricci curvature*. Proc. Amer. Math. Soc., 145(5):2199–2210, 2017.
- [39] Rose, C; Wei, G. , *Eigenvalue estimates for Kato-type Ricci curvature conditions*. Anal. PDE 15 (2022), no. 7, 1703–1724.
- [40] Hamilton, R. S., *Three-manifolds with positive Ricci curvature*, J. Differential Geom. 17 (1982), no. 2, 255–306.
- [41] Schoen, R.; Yau, S.-T., *Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature*, Ann. of Math. (2) 110 (1979), no. 1, 127–142.
- [42] Schoen, R.; Yau, S.-T., *On the structure of manifolds with positive scalar curvature*, Manuscripta Math. 28 (1979), no. 1-3, 159–183.
- [43] Sormani, C., *Conjectures on Convergence and Scalar Curvature*, preprint, arXiv:2103.10093.
- [44] Yang, D., *Convergence of Riemannian manifolds with integral bounds on curvature. I*. Ann. Sci. École Norm. Sup. (4) 25 (1992), no. 1, 77–105.
- [45] Yang, D., *Convergence of Riemannian manifolds with integral bounds on curvature. II*. Ann. Sci. École Norm. Sup. (4) 25 (1992), no. 2, 179–199.
- [46] Zhang, Qi S.; Zhu, M. , *Li-Yau gradient bounds on compact manifolds under nearly optimal curvature conditions*. J. Funct. Anal., 275(2):478–515, 2018.

(Man-Chun Lee) DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T., HONG KONG

Email address: mclee@math.cuhk.edu.hk