Dagger categories via anti-involutions and positivity

Luuk Stehouwer and Jan Steinebrunner

April 7, 2023

Abstract

Dagger categories are an essential tool for categorical descriptions of quantum physics, but pose a challenge to category theorists as their definition is in tension with the "principle of equivalence" that lies at the heart of category theory. In this note we propose the alternative, coherent definition of an "involutive category with a notion of positivity" and show that the 2-category formed by these is biequivalent to the 2-category of dagger categories.

Contents

1	Dagger categories	1
2	Anti-involutive categories	4
3	Hermitian fixed points and Hermitian completion	8
4	Indefinite dagger categories	14
5	Choosing positive Hermitian structures	17
References		22

1 Dagger categories

Hilbert spaces play an important role in the mathematical study of physical systems and in particular in the notion of unitary topological quantum field theory. In the context of unitary TFTs it is especially important to understand Hilbert spaces from a categorical perspective.

When considering the category of finite dimensional Hilbert spaces and bounded operators $\operatorname{Hilb}^{\mathrm{fd}}$ one is faced with a fundamental problem: the forgetful functor $\operatorname{Hilb}^{\mathrm{fd}} \to \operatorname{Vect}_{\mathbb{C}}^{\mathrm{fd}}$ is an equivalence of categories. It is essentially surjective because every finite-dimensional vector space admits a Hilbert space structure and it is fully faithful because every linear map between finite dimensional Hilbert spaces is bounded. We conclude that in this framework, category theory cannot tell apart Hilbert spaces and vector spaces. To resolve this, we need remember how to take the adjoint $A^* \colon H' \to H$ of an operator $A \colon H \to H'$. In other words, we should think of Hilb^{fd} as a dagger category:

A dagger category is a category \mathcal{C} equipped with a functor $\dagger: \mathcal{C}^{\text{op}} \to \mathcal{C}$ satisfying $\dagger \circ \dagger^{\text{op}} = \text{Id}_{\mathcal{C}}$ and $\dagger(x) = x$ for all objects $x \in \mathcal{C}$. A dagger functor $F: (\mathcal{C}, \dagger) \to (\mathcal{D}, \ddagger)$ is a functor $F: \mathcal{C} \to \mathcal{D}$ such that $F(f^{\dagger}) = F(f)^{\ddagger}$ holds for all morphisms $f: x \to y$ in \mathcal{C} .

While dagger categories are key to categorical approaches to quantum physics, they also come with an inherent difficulty: the condition $\dagger(x) = x$ does not play well with many concepts in category theory. Dagger categories are hence sometimes humorously referred to as an "evil" concept [HL+15], as they behave differently from what one might expect from usual category theory. For instance, dagger structures cannot be transported along equivalences of categories.

However, there is still a well-behaved "dagger category theory" obtained by requiring all coherence isomorphisms to be *unitary*. A morphism $u: x \to y$ in a dagger category (\mathcal{C}, \dagger) is called unitary if $u^{\dagger}: x \to y$ is an inverse to u, i.e. if $u \circ u^{\dagger} = \mathrm{id}_y$ and $u^{\dagger} \circ u = \mathrm{id}_x$. There also is a notion of isometries: these are morphisms $i: x \to y$ satisfying only $i^{\dagger} \circ i = \mathrm{id}_x$.

We can define a 2-category †Cat of dagger categories as follows. Objects are dagger categories, morphisms are dagger functors, and 2-morphisms are natural transformations $\alpha: F \to G$ such that each $\alpha_x: Fx \to Gx$ is an isometry. Requiring that the natural transformations are isometries ensures that all invertible 2-morphisms are unitary, and hence the 2-category recovers the appropriate notion of equivalence of dagger categories:

Lemma 1.1 ([Vic11, Lemma 5.1]). We say that dagger functor $F : \mathcal{C} \to \mathcal{D}$ is a dagger equivalence if it satisfies the following equivalent conditions:

- F is an equivalence in the 2-category †Cat. (i.e. there is a dagger functor G: D → C such that F ∘ G and G ∘ F are unitarily naturally isomorphic to the respective identity functors.)
- *F* is fully faithful and surjective up to unitaries. (i.e. for each $d \in D$ there is a $c \in C$ such that F(c) is unitarily isomorphic to d.)

The purpose of this note is to relate dagger categories to their coherent analogue: anti-involutive categories, and to precisely describe which information is lost in the process. There has been plenty of previous work on several notions of involutions on categories, mostly about covariant (sometimes opmonoidal) involutions. A partial list includes [BM09] [Jac12] [Egg11] [BSW19] [HP20] [Yau20] [SCC21] [HPT23]. We define an anti-involutive category to be a category \mathcal{C} equipped with a functor $d: \mathcal{C}^{\text{op}} \to \mathcal{C}$ that squares to the identity functor up to chosen higher coherence.¹ Abstractly, the 2-category ICat of

¹This is sometimes called a category with duality.

anti-involutive categories may be thought of as the homotopy fixed point category of the involution $\mathcal{C} \mapsto \mathcal{C}^{\text{op}}$ on the 2-category Cat. Any dagger category gives rise to an anti-involutive category with trivial higher coherence, and this defines a 2-functor T: \dagger Cat \rightarrow ICat. However, we will see that anti-involutive categories only suffice to capture the behaviour of "indefinite" dagger categories (Definition 4.3).

Theorem 1.2. There is a 2-adjunction

 $T: \dagger Cat \rightleftharpoons ICat : Herm$

and it restricts to an equivalence between the full 2-subcategory of indefinite complete dagger categories and the full 2-subcategory of those anti-involutive categories where each object admits at least one fixed point structure.

To fully capture dagger categories, we additionally introduce the concept of a "positivity notion" P on an anti-involutive category (Definition 5.4), which one can think of as specifying a collection of Hermitian pairings on the category that we prefer to call positive. We then define PCat to be the 2-category of anti-involutive categories equipped with a positivity notion. This approach is in part motivated by LeFanu Lumsdaine's mathoverflow answer [HL+15], which suggests to encode dagger categories by keeping track of a coherent involution and "unitary fixed point data". Our main theorem states that these indeed form an equivalent notion to dagger categories.

Theorem 1.3. There the above adjunction lifts to a biequivalence of 2-categories

 $\dagger Cat \simeq PCat$

that commutes with the forgetful functors to Cat.

One of the key uses of this theorem is that it allows us to compute categories of dagger functors from ordinary functor categories together with information about the anti-involutions and the positivity notions.

Theorem 1.4. Let (\mathcal{C}, \dagger) and (\mathcal{D}, \dagger) be two dagger categories. Then $F \mapsto \dagger_{\mathcal{D}} \circ F \circ \dagger_{\mathcal{C}}$ defines an anti-involution on category of all (not necessarily dagger) functors. The inclusion of the dagger functors into the fixed points

$$\operatorname{Fun}^{\dagger}((\mathcal{C},\dagger),(\mathcal{D},\dagger)) \hookrightarrow (\operatorname{Fun}(\mathcal{C},\mathcal{D}))^{\operatorname{fix}}$$

is fully faithful and its essential image consists of those functors that preserve the positivity notions.

We will now briefly sketch why this is particularly useful for studying unitary topological field theory, mainly following ideas of [FH21]. Let $C = \text{Bord}_{d,d-1}$ be a bordism category that admits the structure of a dagger category. In Atiyah's original definition of topological field theories [Ati88] his Hermitian axiom required the state spaces to come with Hilbert space pairings in such

a way that simultaneously reversing in- and output and orientation-reversing bordisms amounts to taking adjoints of operators. This can be encoded in categorical language by defining unitary TFTs as symmetric monoidal dagger functors $\text{Bord}_{d,d-1} \rightarrow \text{Hilb}$. The above suggests a possible strategy for computing unitary TFTs:²

- 1. First, compute the groupoid of ordinary TFTs $\operatorname{Fun}^{\otimes}(\operatorname{Bord}_{d,d-1},\operatorname{Vect}_{\mathbb{C}})$.
- 2. Next, determine the anti-involution on this groupoid given by conjugating a functor with the respective anti-involutions on both categories.
- 3. Compute the fixed points of this anti-involution: This is the groupoid of involutive functors.
- 4. Determine which of the involutive functors preserve the positivity notion.

Moreover, in rigid symmetric monoidal categories a dual functor provides an equivalence between anti-involutions and covariant involutions. We can think of the covariant involutions (orientation reversal on $\text{Bord}_{d,d-1}$ and complex conjugation on $\text{Vect}_{\mathbb{C}}$) as reflections in the sense of [FH21], and then step 2 and 3 correspond to computing TFTs with reflection structures.

Acknowledgements

The first author would like to thank Theo Johnson-Freyd, Lukas Müller, David Reutter, Stephan Stolz, and Peter Teichner for fruitful discussions on dagger categories, especially in relationship to unitary TFTs. The first author expresses his gratitude to the Max Planck Institute for Mathematics in Bonn for its stimulating research environment leading to this work.

The second author would like to thank André Henriques for introducing him to dagger categories, and Dave Penneys and David Reutter for many enlightening conversations on this topic. The second author is supported by the ERC grant no. 772960, and would like to thank the Copenhagen Centre for Geometry and Topology for their hospitality.

2 Anti-involutive categories

As a first approximation to a more categorically well-behaved version of dagger categories, we can weaken the condition that $\dagger: \mathcal{C} \to \mathcal{C}^{op}$ squares to the identity functor on the nose and we also no longer require it to be the identity on objects. Instead, we give a natural isomorphism $\dagger \circ \dagger^{op} \cong \mathrm{Id}_{\mathcal{C}}$ satisfying some compatibility conditions. Here given a functor $F: \mathcal{C} \to \mathcal{D}$, we denoted the canonical induced functor $\mathcal{C}^{op} \to \mathcal{D}^{op}$ by F^{op} , but we will often abuse notation and write it as F.

A category with anti-involution is exactly a fixed point for the $\mathbb{Z}/2$ -action on the bicategory of categories given by $\mathcal{C} \mapsto \mathcal{C}^{op}$, see [Hes17, section 2.2] or [MS23, Appendix A.2]. This results in the following concrete definition:

 $^{^{2}\}mathrm{As}$ it stands this strategy computes arbitrary functors, not symmetric monoidal ones.

Definition 2.1. For C a category, an *anti-involution* is a functor $d : C \to C^{op}$ and a natural isomorphism $\eta : \mathrm{id}_{\mathcal{C}} \Rightarrow d^{op} \circ d$ such that $\eta_{d(c)} : d(c) \to ddd(c)$ and $d(\eta_c) : ddd(c) \to d(c)$ are inverses.³ We call the triple (C, d, η) an *anti-involutive* category.

Example 2.2. If (\mathcal{C}, \dagger) is a dagger category, define an anti-involution as $d := \dagger$. Since dc = c for every object c of \mathcal{C} , we can set $\eta_c = \mathrm{id}_c$, which gives an anti-involutive category since $\mathrm{id}_c^{\dagger} = \mathrm{id}_c$.

Definition 2.3. An involutive functor $F : (\mathcal{C}_1, d_1, \eta_1) \to (\mathcal{C}_2, d_2, \eta_2)$ consists of a functor $F : \mathcal{C}_1 \to \mathcal{C}_2$ and a natural isomorphism $\varphi : F^{\text{op}} \circ d_1 \cong d_2 \circ F$ such that the following square commutes for all $x \in \mathcal{C}_1$:

The composition of involutive functors $(F : C_1 \to C_2, \varphi) \circ (G : C_2 \to C_3, \psi)$ is defined to come equipped with the natural transformation

$$F \circ G \circ d_1(x) \xrightarrow{F(\psi_x)} F \circ d_2 \circ G(x) \xrightarrow{\varphi_{G(x)}} d_3 \circ F \circ G(x)$$

which is easily shown to satisfy the required condition. An involutive natural transformation $\alpha: (F: \mathcal{C}_1 \to \mathcal{C}_2, \varphi) \Rightarrow (G: \mathcal{C}_1 \to \mathcal{C}_2, \psi)$ is a natural transformation $\alpha: F \Rightarrow G$ such that the following square commutes for all $x \in \mathcal{C}_1$:

$$\begin{array}{ccc} (F \circ d_1)(x) & \xrightarrow{\alpha_{d_1(x)}} & (G \circ d_1)(x) \\ & & \downarrow^{\varphi_x} & & \downarrow^{\psi_x} \\ (d_2 \circ F)(x) & \xleftarrow{d_2(\alpha_x)} & (d_2 \circ G)(x) \end{array}$$

The composition of involutive natural transformations is involutive. Let ICat denote the 2-category of anti-involutive categories, involutive functors, and involutive natural transformations.

Note that similar to isometric natural transformations for dagger categories, an involutive natural transformation α_c admits a left inverse, but not necessarily a right inverse.

Remark 2.4. The observation in example 2.2 that every dagger category is canonically an anti-involutive category extends to give a 2-functor T: \dagger Cat \rightarrow ICat. More precisely, if $F: (\mathcal{C}, \dagger) \rightarrow (\mathcal{D}, \dagger)$ is a dagger functor, we can take $\varphi_c =$ $\mathrm{id}_{F(c)}$. The condition that this defines a natural transformation $F \circ \dagger \Rightarrow \dagger \circ F$

³In fact we could require that $d^{\text{op}} \circ d = \text{id}_{\mathcal{C}}$ and $\eta = \text{id}_{\text{id}_{\mathcal{C}}}$. This would lead to a biequivalent 2-category. However, we will not need this strictification.

is equivalent to F being a dagger functor. The remaining condition between η and φ is satisfied, since all morphisms involved are the identity.

Finally, let $\alpha: F \Rightarrow F'$ be a natural transformation. Then α is an involutive natural transformation between the induced involutive functors if and only if α_c is an isometry for all objects c, which is how we defined 2-morphisms in †Cat. We recall that that α_c need not be invertible, but it is invertible if and only if it is unitary. Clearly these constructions preserve composition of functors and both horizontal and vertical composition of natural transformations.

Lemma 2.5. An involutive functor (F, φ) : $(\mathcal{C}, d, \rho) \to (\mathcal{D}, d, \eta)$ is an equivalence in ICat (i.e. it has an involutive inverse up to involutive natural transformation) if and only if the underlying functor $F: \mathcal{C} \to \mathcal{D}$ is an equivalence of categories.

Proof. The only if direction holds because if (G, ψ) is an involutive inverse functor, then G is an inverse of F up to natural isomorphism.


For the if direction, pick some $G: \mathcal{D} \to \mathcal{C}$ and natural transformations $\alpha: F \circ G \cong \operatorname{Id}_{\mathcal{D}}$ and $\beta: G \circ F \cong \operatorname{Id}_{\mathcal{C}}$. Recall that without loss of generality, we can assume this is an adjoint equivalence, i.e. α and β satisfy the snake identities. It suffices to provide the data ψ that makes G into an involutive functor and show that α and β become involutive natural transformations. Define ψ at an object y of \mathcal{D} as

$$Gdy \xrightarrow{Gd\alpha_y} GdFGy \xrightarrow{G\varphi_{Gy}^{-1}} GFdGy \xrightarrow{\beta_{dGy}} dGy.$$

By definition of being an involutive functor, we have to show the diagram


$$\begin{array}{cccc} Gy & \xrightarrow{G\rho_y} & Gd^2y & \xrightarrow{Gd\alpha_{dy}} & GdFGdy & \leftarrow_{G\varphi_{Gdy}} & GFdGdy \\ & & & & & \downarrow^{\beta_{dGdy}} \\ d^2Gy & \xrightarrow{d\beta_{dGy}} & dGFdGy & \leftarrow_{dG\varphi_{Gy}} & dGdFGy & \xrightarrow{dGd\alpha_y} & dGdy \end{array}$$

commutes. For this, first note that




commutes. Indeed, the left upper triangle commutes by the snake identity, the right upper square commutes by naturality of β , the left lower square commutes by naturality of α and the lower right square commutes because F is an involutive functor. Replacing the morphisms $G\rho_y$ and η_{Gy} in the first diagram by this

second diagram leads us to conclude that it suffices to show that the following diagram commutes. We omitted the choice of input object y in \mathcal{D} from the notation for reasons of space.



Every quadrilateral in the diagram commutes by the interchange law and the upper two bent arrows are equal by the snake identity. We are led to conclude that (G, ψ) is an involutive functor.

It remains to show that α and β are involutive natural transformations. Writing out the definition of the involutive structure on $F \circ G$ this entails that for α we have to show that the diagram



commutes. The two bent arrows are equal by the snake identity and the other two parts commute by the interchange law. The proof that β is involutive is analoguous.

The following example shows that the underlying anti-involution of a dagger category does not preserve enough information.

Example 2.6. Let $\operatorname{Herm}_{\mathbb{C}}$ denote the category where objects are finite dimensional complex vector spaces with a non-degenerate sesquilinear form such that

$$\langle v, w \rangle = \langle w, v \rangle$$

and morphisms are all linear maps. In other words, these are Hermitian vector spaces that are not necessarily positive definite inner product spaces. It becomes

a dagger category when the dagger is defined by taking the adjoint with respect to the pairing.

The category of finite dimensional Hilbert spaces is a full dagger subcategory Hilb \subset Herm_{\mathbb{C}} characterised by the condition that the sesquilinar form be positive definite. This inclusion is not a dagger equivalence, as it is not surjective up to unitaries. Indeed, objects in Herm_{\mathbb{C}} are classified, up to unitary isomorphism, by their signature (p, q) and the fully subcategory only contains those of signature (p, 0).

However, the inclusion $F: \text{Hilb} \to \text{Herm}_{\mathbb{C}}$ is an equivalence of anti-involutive categories. It is fully faithful and essentially surjective because every finite-dimensional vector space admits some Hilbert space structure. By Lemma 2.5, this is an equivalence of anti-involutive categories.

Concretely, we could construct a (highly noncanonical) inverse of this equivalence as follows. Pick for every finite-dimensional Hermitian vector space $(V, \langle ., . \rangle)$ a basis $\alpha_V : V \cong \mathbb{C}^n$ once and for all. Define the functor $G : \operatorname{Herm}_{\mathbb{C}} \to$ Hilb on objects by $G(V, \langle ., . \rangle) = (\mathbb{C}^n, \langle ., . \rangle_{st})$ where $\langle ., . \rangle_{st}$ is the standard Hilbert space structure. On morphisms we set $G(f : V_1 \to V_2) := \alpha_{V_2}^{-1} \circ f \circ \alpha_{V_1}$. There is an associated canonical natural isomorphism $\alpha : \operatorname{id}_{\operatorname{Herm}_{\mathbb{C}}} \Longrightarrow F \circ G$ given by $\alpha(V, \langle ., . \rangle) = \alpha_V : (V, \langle ., . \rangle) \to (\mathbb{C}^n, \langle ., . \rangle_{st})$. Now, G is not a dagger functor since α_V is in general not unitary. But even though the anti-involutions d on both categories are the identity on objects, we can use the recipe in the above lemma to equip G with a non-trivial structure of an involutive functor:

$$\varphi_V \colon G(dV) = G(V) = \mathbb{C}^n \xrightarrow{\alpha_V^+} V \xrightarrow{\alpha_V} \mathbb{C}^n = G(V) = dG(V)$$

Then the condition that φ has to satisfy for G to be an involutive functor boils down to $\varphi_V^{\dagger} = \varphi_V$, which is easy to check. Hence (G, φ) is an involutive inverse of the involutive functor F.

3 Hermitian fixed points and Hermitian completion

In the last section we proposed the notion of an anti-involutive category as a better-behaved analogue of the notion of a dagger category so that every dagger category has an underlying anti-involutive category. However, in example 2.6 we found that there are important examples of dagger categories that are equivalent as anti-involutive categories but not as dagger categories. Heuristically, the example gives us the idea that the dagger category of finite-dimensional Hilbert spaces is not equivalent to the dagger category of finite-dimensional Hermitian vector spaces because in the former fewer Hermitian structures are allowed. Therefore we study an abstraction of the notion of a Hermitian structure as follows.

Definition 3.1. A *Hermitian fixed point* in a category with anti-involution is

an isomorphism $h\colon c\to dc$ such that

$$c \xrightarrow{\eta_c} d^2 c \xrightarrow{dh} dc$$

commutes. The *adjoint* $f^{\dagger}: c_2 \to c_1$ of a morphism $f: c_1 \to c_2$ with respect to Hermitian fixed points $h_1: c_1 \to dc_1$ and $h_2: c_2 \to dc_2$ is the composition

$$c_2 \xrightarrow{h_2} dc_2 \xrightarrow{df} dc_1 \xrightarrow{h_1^{-1}} c_1.$$

Example 3.2. Take $C = \operatorname{Vect}_{\mathbb{C}}^{\operatorname{fd}}$ to be the category of finite-dimensional vector spaces. Recall that the complex conjugate \overline{V} of a vector space V is defined to be the same abelian group but with complex conjugate scalar multiplication. This extends to a functor $\overline{(.)} : \operatorname{Vect}_{\mathbb{C}}^{\operatorname{fd}} \to \operatorname{Vect}_{\mathbb{C}}^{\operatorname{fd}}$. Set $d = \overline{(.)}^*$ so that there is an obvious η given by the evaluation map. It is straightforward to check that η satisfies $\eta_{\overline{V}^*} = \overline{\eta_V}^*$. A Hermitian fixed point consists of a vector space V and an isomorphism $V \to \overline{V}^*$ satisfying a condition. Such an isomorphism is equivalent to the Hermiticity axiom

$$\langle v, w \rangle = \overline{\langle w, v \rangle}$$

The adjoint is given by the usual adjoint of a linear map.

Hermitian fixed points naturally form a category C^{fix} in which morphisms $f: (c_1, h_1) \to (c_2, h_2)$ are morphisms $c_1 \to c_2$ satisfying the compatibility relation

$$\begin{array}{ccc} c_1 & \xrightarrow{J} & c_2 \\ \downarrow h_1 & & \downarrow h \\ dc_1 & \xleftarrow{df} & dc_2 \end{array}$$

Let $f: (c_1, h_1) \to (c_2, h_2)$ be a morphism in \mathcal{C}^{fix} . Note that the condition f has to satisfy exactly says that f^{\dagger} is a left inverse of f. Therefore \mathcal{C}^{fix} is exactly the wide subcategory of isometries of the dagger category $\text{Herm}(\mathcal{C})$ that we shall define now. The construction is closely related to the 'unitary core of a \dagger -isomix category' [SCC21, Definition 5.12]. One could think of $\text{Herm}(\mathcal{C})$ as the 'co-free dagger category on an anti-involutive category'. This idea is made precise by the 2-adjunction that we will establish in theorem 4.9.

Definition 3.3. The Hermitian completion Herm C of the anti-involutive category (C, d, η) is the category in which objects are Hermitian fixed points (c, h) and morphisms $(c_1, h_1) \rightarrow (c_2, h_2)$ are simply given by morphisms $f: c_1 \rightarrow c_2$.

Lemma 3.4. The adjoint on the category $\operatorname{Herm} \mathcal{C}$ makes it into a dagger category.

Proof. Let (c,h) be an object of \mathcal{C} with Hermitian structure $h: c \to dc$. We have that

$$\mathrm{id}_c^\dagger = h^{-1} \circ d(\mathrm{id}_c) \circ h = h^{-1} \circ \mathrm{id}_{dc} \circ h = \mathrm{id}_c$$

If $f: (c_1, h_1) \to (c_2, h_2)$ and $g: (c_2, h_2) \to (c_3, h_3)$, then $(g \circ f)^{\dagger} = f^{\dagger} \circ g^{\dagger}$ follows from the fact that

$$c_3 \stackrel{h_3}{\cong} dc_3 \stackrel{dg}{\longrightarrow} dc_2 \stackrel{h_2^{-1}}{\cong} c_2 \stackrel{h_2}{\cong} dc_2 \stackrel{df}{\longrightarrow} dc_1 \stackrel{h_1}{\cong} c_1$$

is equal to

$$a_3 \stackrel{h_3}{\cong} dc_3 \stackrel{d(g \circ f)}{\longrightarrow} dc_1 \stackrel{h_1}{\cong} c_1$$

by functoriality of d. Now $f^{\dagger\dagger}$ is the composition

$$c_1 \stackrel{h_1}{\cong} dc_1 \stackrel{dh_1^{-1}}{\cong} d^2c_1 \stackrel{d^2f}{\longrightarrow} d^2c_2 \stackrel{dh_2}{\cong} dc_2 \stackrel{h_2^{-1}}{\cong} c_2.$$

Using the fixed point property of a Hermitian structure, this composition is equal to

$$c_1 \stackrel{\eta_{c_1}}{\cong} d^2 c_1 \stackrel{d^2 f}{\longrightarrow} d^2 c_2 \stackrel{\eta_{c_2}^{-1}}{\cong} c_2$$

By naturality of η this composition is equal to f.

Example 3.5. The Hermitian completion of $(\mathcal{C} = \operatorname{Vect}_{\mathbb{C}}^{\mathrm{fd}}, d = \overline{(.)}^*)$ is the dagger category of Hermitian vector spaces we considered in example 2.6. So Herm $\operatorname{Vect}_{\mathbb{C}}^{\mathrm{fd}} = \operatorname{Herm}_{\mathbb{C}}$

Remark 3.6. Unlike for finite-dimensional vector spaces, $\overline{(.)}^*$ does not define an anti-involution on infinite-dimensional vector spaces. Indeed, even though there is still a well-defined bidual map $\eta: V \to \overline{V}^{**}$, it is only injective but not surjective. Hence the dagger category of all Hilbert spaces can not be constructed in a similar fashion as the last example. It would be interesting to study a weakened version of anti-involutive categories in which η is not necessarily an isomorphism and Hermitian fixed points $h: c \to dc$ are not necessarily isomorphisms. The technical disadvantage of such a theory would be that we might have to restrict the morphisms in the Hermitian completion to those that admit an adjoint, for example the bounded operators for Hilbert spaces. An alternative approach would be to work with a certain category of topological vector spaces and use a continuous linear dual.

Example 3.7. Let $TC \in ICat$ be a dagger category C seen as an anti-involutive category. The Hermitian completion Herm(TC) concretely consists of pairs (c, τ) , where $\tau: c \to c$ is invertible and self-adjoint. For $f: (c_1, \tau_1) \to (c_2, \tau_2)$ the new adjoint * on the Hermitian completion is defined as $f^* = \tau_2 \circ f^{\dagger} \circ \tau_1^{-1}$. For example, starting with the dagger category of Hilbert spaces, new objects are triples (V, (-, -), A) consisting of a Hilbert space and a self-adjoint invertible linear operator on V. The adjoints of morphisms between such objects are defined using the Hermitian pairing (-, A-) on V. The resulting dagger category is unitarily equivalent to the dagger category of Hermitian vector spaces.

Example 3.8. Again take $\mathcal{C} = \operatorname{Vect}_{\mathbb{C}}^{\operatorname{fd}}$ to be the category of finite-dimensional vector spaces. Now define d to be the dual $(-)^* \colon \mathcal{C} \to \mathcal{C}^{op}$ and $\eta \colon V \to V^{**}$ the evaluation map. Then a Hermitian fixed point on a vector space V is equivalent to a nondegenerate symmetric bilinear form on V. More generally, we could take \mathcal{C} to be finite-dimensional complex representations of a finite group G. Since for a general G-representation V, there is no G-equivariant isomorphism $V \cong V^*$ there are representations that do not admit the structure of a Hermitian fixed point at all.

Lemma 3.9. Let $(\mathcal{C}, d_{\mathcal{C}}, \eta_{\mathcal{C}}), (\mathcal{D}, d_{\mathcal{D}}, \eta_{\mathcal{D}})$ be two anti-involutive categories. Then there is an anti-involutive structure on the category $\operatorname{Fun}(\mathcal{C}, \mathcal{D})$ of functors between them, such that the category of $\mathbb{Z}/2$ -fixed points $\operatorname{Fun}(\mathcal{C}, \mathcal{D})^{\operatorname{fix}}$ is the category $\operatorname{Hom}_{\operatorname{ICat}}(\mathcal{C}, \mathcal{D})$ of 1-morphisms in ICat.

Proof. The functor category $\operatorname{Fun}(\mathcal{C}, \mathcal{D})$ becomes an anti-involutive category via

$$dF := d_{\mathcal{D}} \circ F \circ d_{\mathcal{C}}$$

Namely, we can define the anti-involution on natural transformations $\alpha \colon F_1 \Rightarrow F_2$ between functors $F_1, F_2 \colon \mathcal{C} \to \mathcal{D}$ as the whiskering

$$d\alpha := \mathrm{id}_{d_{\mathcal{D}}} \bullet \alpha \bullet \mathrm{id}_{d_{\mathcal{C}}},$$

where we denoted horizontal composition of natural transformations with •. This defines a functor $\operatorname{Fun}(\mathcal{C}, \mathcal{D}) \to \operatorname{Fun}(\mathcal{C}, \mathcal{D})$. Define the natural transformation η : $\operatorname{id}_{\operatorname{Fun}(\mathcal{C},\mathcal{D})} \Rightarrow d^2$ on $F \in \operatorname{Fun}(\mathcal{C},\mathcal{D})$ by

$$F \xrightarrow{\eta_{\mathcal{D}} \bullet \mathrm{id}_F \bullet \eta_{\mathcal{C}}} d_{\mathcal{D}}^2 F d_{\mathcal{C}}^2,$$

which is natural by the exchange law. Finally, we have to show that $\eta_{dF} = d\eta_F^{-1}$ for all $F \in \operatorname{Fun}(\mathcal{C}, \mathcal{D})$. This amounts to showing that $\eta_{\mathcal{D}} \bullet \operatorname{id}_{d_{\mathcal{D}}Fd_{\mathcal{C}}} \bullet \eta_{\mathcal{C}}$ is the inverse of $\operatorname{id}_{d_{\mathcal{C}}} \bullet \eta_{\mathcal{D}} \bullet \operatorname{id}_{F} \bullet \eta_{\mathcal{C}} \bullet \operatorname{id}_{d_{\mathcal{D}}}$. This holds because, since $\eta_{\mathcal{D}}$ is part of an anti-involution it satisfies that $\eta_{\mathcal{D}} \bullet \operatorname{id}_{d_{\mathcal{D}}}$ is inverse to $\operatorname{id}_{d_{\mathcal{C}}} \bullet \eta_{\mathcal{D}}$ and similarly for $\eta_{\mathcal{C}}$.

A Hermitian fixed point in $\operatorname{Fun}(\mathcal{C}, \mathcal{D})$ is equivalent to an involutive functor. Indeed, let $\psi \colon F \Rightarrow d_{\mathcal{D}}Fd_{\mathcal{C}}$ be a Hermitian fixed point on F. Writing out the condition results in the commutative diagram

$$Fc \xrightarrow{\psi_c} d_{\mathcal{D}}Fd_{\mathcal{C}}c$$

$$F((\eta_{\mathcal{C}})_c) \downarrow \qquad \qquad \uparrow^{d_{\mathcal{D}}\psi_{d_{\mathcal{C}}}} fd_{\mathcal{D}}\psi_{d_{\mathcal{C}}}$$

$$Fd_{\mathcal{C}}^2c \xrightarrow{(\eta_{\mathcal{D}})_{Fd_{\mathcal{C}}^2c}} d_{\mathcal{D}}^2Fd_{\mathcal{C}}^2c$$

for every object c. A diagram chase shows that under mapping ψ to the composition φ defined by

$$\varphi \colon Fd_{\mathcal{C}} \xrightarrow{\psi \bullet \mathrm{id}_{d_{\mathcal{D}}}} d_{\mathcal{D}}Fd_{\mathcal{C}}^2 \xrightarrow{\mathrm{id}_{d_{\mathcal{D}}F} \bullet \eta_{\mathcal{C}}^{-1}} d_{\mathcal{D}}F$$

this becomes the condition that (F, φ) is an involutive functor. So we see that Herm Fun $(\mathcal{C}, \mathcal{D})$ is the category with objects involutive functors and as morphisms all natural transformations. A natural transformation is involutive if and only if it is an isometry in Herm Fun $(\mathcal{C}, \mathcal{D})$.

Example 3.10. Let $h: c \to dc$ be a Hermitian fixed point. Then $(dh)^{-1}: dc \to d^2c$ is a Hermitian fixed point structure on dc. Indeed, taking d of the diagram saying that h is a fixed point and using that $d\eta_c = \eta_{dc}^{-1}$ yields

$$dc \xrightarrow{\eta_{dc}} d^3c \xleftarrow{d^2h} d^2c$$

Using that $(d^2h)^{-1} = d(dh^{-1})$, this diagram indeed expresses the fact that $(dh)^{-1}: dc \to d^2c$ is a Hermitian fixed point. Note that by construction $h: c \to dc$ is a unitary isomorphism between the objects (c, h) and $(dc, (dh)^{-1})$ in the dagger category Herm \mathcal{C} .

Remark 3.11. We expect the discussion above to be closely related to [Egg11, section 6] as follows. This reference considers covariant op-monoidal involutions which in certain rigid monoidal categories should be related to monoidal anti-involutions after composing with a choice of dual functor. Their notion of Hermitian sesquilinear pairing [Egg11, Definition 6.2] should be related to our notion of Hermitian fixed point and [Egg11, Lemma 6.3] should be related to our Hermitian completion.

Definition 3.12. We extend the construction of definition 3.3 to a 2-functor

Herm: ICat
$$\longrightarrow \dagger$$
Cat

as follows. For an involutive functor $(F, \varphi) \colon (\mathcal{C}, d, \eta) \to (\mathcal{D}, d, \rho)$ we define

Herm $F: \operatorname{Herm} \mathcal{C} \to \operatorname{Herm} \mathcal{D}$

on objects by Herm $F(c, h) = (F(c), h_F := \varphi_c \circ F(h))$, and on morphisms by Herm F(f) = F(f). For an involutive natural transformation $\alpha : (F, \varphi) \to (F', \varphi')$ we define Herm $\alpha :$ Herm $F \to$ Herm F' by (Herm $\alpha)_c := \alpha_c$.

Lemma 3.13. The above yields a well-defined 2-functor.

Proof. We have already checked that $\operatorname{Herm}(\mathcal{C}, d, \eta)$ is indeed a dagger category, so next we need to verify that $\operatorname{Herm}(F, \varphi)$ is a dagger functor. First, note that $h_F := \varphi_c \circ F(h) \colon F(c) \to F(dc) \cong dF(c)$ is indeed a hermitian structure because

of the diagram:

Here the triangle commutes because h is a hermitian fixed point, the trapezoid commutes because φ is part of an involutive functor, and the rectangle commutes because φ is a natural transformation.

Herm F is certainly functorial seeing as morphisms in Herm C are simply composed by composing them in C. To show it is a dagger functor, let

$$f\colon (c_1,h)\to (c_2,h')$$

be a morphism in $\operatorname{Herm} \mathcal{C}$. Then

Herm
$$F(f^{\dagger}) = F(c_2 \xrightarrow{h'} dc_2 \xrightarrow{df} dc_1 \xrightarrow{h^{-1}} c_1) = F(h)^{-1}F(df)F(h')$$

Recall that since F is involutive, the diagram

commutes. Looking at the definition of h_F, h'_F , we obtain

Herm
$$F(f^{\dagger}) = \operatorname{Herm} F(f)^{\dagger}$$
.

To conclude that Herm is a 1-functor we need to check that for two composable involutive functors (F_1, φ_1) and (F_2, φ_2) we have that $\operatorname{Herm}(F_2) \circ \operatorname{Herm}(F_1) =$ $\operatorname{Herm}(F_2 \circ F_1)$. It will suffice to check that both sides do the same on an object (c, h). The two resulting hermitian structures on $F_2F_1(c)$ are

These are indeed the same: the right rectangle commutes because of how the coherence isomorphism φ_{12} of the composite functor is defined.

Finally, we need to consider the effect of Herm on 2-morphisms. Here all there is to check that $\text{Herm}(\alpha)$ is indeed an isometry. This follows from the diagram:

$$\begin{array}{ccc} F(c) & \xrightarrow{\alpha_c} & F'(c) \\ F(h) & & \downarrow F'(h) \\ F(dc) & \xrightarrow{\alpha_{dc}} & F'(dc) \\ \varphi_c & & \downarrow \varphi'_c \\ dF(c) & \xleftarrow{d\alpha_c} & dF'(c) \end{array}$$

The squares commute because α is a natural transformation and because α is involutive with respect to (F, φ) and (F', φ') . The vertical composites are the hermitian structures h_F and h'_F , and therefore the diagram shows that α_c is a one-sided inverse to $\alpha_c^{\dagger} = h_F^{-1} \circ d(\alpha_c) \circ h_F$.

4 Indefinite dagger categories

The 2-functors Herm and T are not inverses of each other for two reasons:

- 1. The anti-involutive category $T(\mathcal{C}, \dagger)$ has the property that every object admits at least one hermitian fixed point structure. This is not true for every anti-involutive category, for instance the discrete category $\mathbb{Z}/2$ with the non-trivial swap, and therefore T is not surjective up to equivalence.
- 2. There exist dagger categories that are not unitarily equivalent, but become equivalent as anti-involutive categories after applying T.

However, we will still be able to show that Herm and T restrict to a biequvialence between certain full 2-subcategories. On the side of the anti-involutive categories we make the following restriction, motivated by point 1 above:

Definition 4.1. Let $C^{\exists fix}$ denote the full subcategory of the anti-involutive category C on the objects c that admit some Hermitian fixed point $h: c \to dc$. This is again an anti-involutive category, also see example 3.10. Let $\operatorname{ICat}^{\exists fix} \subset \operatorname{ICat}$ denote the full 2-subcategory on those anti-involutive categories in which every object admits some Hermitian fixed point structure.

To find the correct property on the dagger category side, we note:

Example 4.2. Consider the dagger category $\operatorname{Herm}_{\mathbb{C}}$ as a category with antiinvolution. Its Hermitian completion is again dagger-equivalent to $\operatorname{Herm}_{\mathbb{C}}$. However, for Hilb it is instead $\operatorname{Herm}_{\mathbb{C}}$ which is not dagger-equivalent to Hilb.

It turns out that the following is the essential property that $\operatorname{Herm}_{\mathbb{C}}$ has and Hilb lacks. In analogy with $\operatorname{Herm}_{\mathbb{C}}$, we think of such dagger categories as containing 'all Hermitian forms, even all the indefinite ones'. **Definition 4.3.** We say that a dagger category \mathcal{D} is *indefinite* if for any object $x \in \mathcal{D}$ and any self-adjoint automorphism $a = a^{\dagger} : x \cong x$ there is another object $y \in \mathcal{D}$ and an isomorphism $f : x \cong y$ such that $a = f^{\dagger} \circ f$. We let $\dagger \operatorname{Cat}^{\operatorname{indef}} \subset \dagger \operatorname{Cat}$ denote the full sub-2-category on the indefinite complete dagger categories.

In the sense of Remark 5.9 this definition can be thought of as saying that in an indefinite dagger category all self-adjoint automorphisms must be "postive".

Lemma 4.4. For any anti-involutive category (C, d, η) the dagger category Herm(C) is indefinite.

Proof. A self-adjoint automorphism is an isomorphism $a \colon (c, h) \to (c, h)$ such that

$$a = a^{\dagger} = h^{-1} \circ d(a) \circ h.$$

We need to find an isomorphism $f: (c, h) \to (c', h')$ such that

$$a \stackrel{?}{=} f^{\dagger} \circ f = h^{-1} \circ d(f) \circ h' \circ f.$$

Indeed, this can always be achieved by setting c' = c, $f = id_c$, and $h' = h \circ a$. It just remains to check that h' is indeed a valid hermitian form on c. For this we consider

$$d(h') \circ \eta_c = d(a) \circ d(h) \circ \eta_c = d(a) \circ h = h \circ a = h'.$$

We now begin to construct the unit and counit for the adjunction between T and Herm.

Definition 4.5. For every anti-involutive category (\mathcal{C}, d, η) we define an involutive functor

$$(K_{\mathcal{C}}, \varphi_{\mathcal{C}}) \colon \mathrm{T}(\mathrm{Herm}(\mathcal{C}, d, \eta)) \longrightarrow (\mathcal{C}, d, \eta)$$

by letting $K_{\mathcal{C}}$ be the functor $(c,h) \mapsto c$ and $f \mapsto f$, and letting $\varphi_{\mathcal{C}} \colon K_{\mathcal{C}}^{\mathrm{op}} \circ \dagger_{\mathrm{T}(\mathrm{Herm}(\mathcal{C}))} \cong d \circ K_{\mathcal{C}}$ be the natural transformation given by

$$\varphi_{(c,h)} := (h \colon c \to d(c)).$$

Lemma 4.6. The involutive functor $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ is well-defined, natural in \mathcal{C} , and it is an equivalence of anti-involutive categories onto the full subcategory $\mathcal{C}^{\exists \text{fix}} \subset \mathcal{C}$.

Proof. To check that φ is indeed a natural transformation we need to consider for each morphism $f: (c_1, h_1) \to (c_2, h_2)$ the square:

$$\begin{array}{ccc} c_1 & \xrightarrow{\varphi_{(c_1,h_1)}=h_1} & d(c_1) \\ f^{\dagger} & & \uparrow \\ c_2 & \xrightarrow{\varphi_{(c_2,h_2)}=h_2} & d(c_2). \end{array}$$

This indeed commutes by the definition of f^{\dagger} . This natural transformation further has to satisfy that for each (c, h) the square

commutes. Upon closer inspection this is exactly the triangle that commutes because h is a Hermitian fixed point.

It follows from the construction that $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ is natural in \mathcal{C} . Moreover, $K_{\mathcal{C}}$ is certainly fully faithful and essentially surjective onto the subcategory of \mathcal{C} that admit a Hermitian fixed point, so by Lemma 2.5 $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ is an equivalence in ICat.

Definition 4.7. For every dagger category (\mathcal{D}, \dagger) we define a dagger functor

$$U_{\mathcal{D}} \colon \mathcal{D} \longrightarrow \operatorname{Herm}(T(\mathcal{D}))$$

by sending x to (x, id) and $f: x \to y$ to $f: (x, id) \to (y, id)$.

The construction of $U_{\mathcal{D}}$ is well-defined and natural in \mathcal{D} . Moreover, $U_{\mathcal{D}}$ is always fully faithful and essentially surjective. However, the more subtle question is when U is surjective up to unitaries.

Lemma 4.8. The functor $U_{\mathcal{D}}$ is an equivalence of dagger categories if and only if \mathcal{D} is indefinite.

Proof. As noted before $U_{\mathcal{D}}: \mathcal{D} \longrightarrow \operatorname{Herm}(T(\mathcal{D}))$ is always an equivalence of categories, so by Lemma 1.1 we only need to check when it is surjective up to unitaries. Suppose $(y,h) \in \operatorname{Herm}(T(\mathcal{D}))$ is an object that is unitarily isomorphic to some object (x, id_x) in the essential image. Then we have an isomorphism $f: (y,h) \to (x, \operatorname{id}_x)$ satisfying $\operatorname{id}_{(y,h)} = f^* \circ f$. (Here we write * for the dagger on $\operatorname{Herm}(T(\mathcal{D}))$ to distinguish it from the dagger \dagger on \mathcal{D} .) Spelling out the definition we see that $\operatorname{id}_{(y,h)} = f^* \circ f = (h^{-1} \circ f^{\dagger} \circ \operatorname{id}_x) \circ f$, or equivalently $h = f^{\dagger} \circ f$. Therefore $U_{\mathcal{D}}$ is surjective up to unitaries if and only if every self-adjoint automorphism h can be written as $f^{\dagger} \circ f$ with f invertible, i.e. if and only if \mathcal{D} is indefinite.

Recall that a 2-adjunction is a Cat_1 -enriched adjunction, i.e. an adjunction for which the unit and counit satisfy the triangle identities *strictly*. [KS06]

Theorem 4.9. The functors U and K exhibit a 2-adjunction:

 $T \colon \dagger Cat \rightleftarrows ICat : Herm$

and this restricts to a biequivalence between $ICat^{\exists fix}$ and the full sub-2-category $\dagger Cat^{indef}$ on the indefinite complete dagger categories.

Proof. To establish the 2-adjunction $T \dashv$ Herm with unit U and counit K we need to check the triangle identities. The first identity concerns for each $(\mathcal{C}, d, \eta) \in$ ICat the composite functor

$$\operatorname{Herm}(\mathcal{C}) \xrightarrow{U_{\operatorname{Herm}(\mathcal{C})}} \operatorname{Herm}(\operatorname{T}(\operatorname{Herm}(\mathcal{C}))) \xrightarrow{\operatorname{Herm}(K_{\mathcal{C}},\varphi_{\mathcal{C}})} \operatorname{Herm}(\mathcal{C})$$

The first functor sends (x, h) to ((x, h), id) and the second functor sends this to $(x, (\varphi_{\mathcal{C}})_{(x,h)} \circ K_{\mathcal{C}}(id)) = (x, h \circ id) = (x, h)$. By construction the composite functor also the identity on morphisms.

The second identity concerns for each $(\mathcal{D}, \dagger) \in \dagger$ Cat the composite functor

$$T(\mathcal{D}) \xrightarrow{T(U_{\mathcal{C}})} T(Herm(T(\mathcal{D}))) \xrightarrow{K_{T(\mathcal{D})}, \varphi_{T(\mathcal{D})}} T(\mathcal{D}).$$

The first functor sends x to (x, id_x) and the second functor sends this to x. On morphisms the composite is also the identity. It remains to check that the involutive data of the composite functor is trivial. For the first functor this holds by definition. For the second functor we have $\varphi_{T(\mathcal{D})}(x, h) = h$, but since we are applying this to the object (x, id_x) , it is also trivial.

Finally, we would like to show that this adjunction restricts to a biequivalence between $\dagger \operatorname{Cat}^{\operatorname{indef}}$ and $\operatorname{ICat}^{\exists \operatorname{fix}}$. The adjunction does restrict because $\operatorname{Herm}(\mathcal{C})$ is always indefinite and $\operatorname{T}(\mathcal{D})$ always has fixed-point structures. The restriction is a biequivalence by lemma 4.8 and lemma 4.6, which state that on these subcategories the unit and counit become equivalences.


5 Choosing positive Hermitian structures

Our next goal is to obtain dagger categories that are not indefinite from categories with anti-involution. To achieve this we will restrict the collection of 'allowed' Hermitian fixed points on the Hermitian completion to a smaller class of 'positive' Hermitian fixed points. This will yield a smaller dagger subcategory for which the underlying category with anti-involution is equivalent. For example, to get the dagger category Hilb^{fd} we take the Hermitian completion of Vect^{fd} and then restrict to the subclass of Hermitian fixed points that are positive definite as Hermitian pairings.

So let (\mathcal{C}, d, η) be a category with anti-involution. For P any subset of the collection of all Hermitian fixed points in \mathcal{C} , let $\mathcal{C}_P \subseteq \text{Herm}\mathcal{C}$ denote the full subcategory on all $(c, h) \in P$. Here P stands for 'positive' to remind us of the typical situation in vector spaces in which we wanted to restrict the Hermitian fixed points to the positive definite ones to obtain the dagger category of Hilbert spaces. The dagger from Herm \mathcal{C} restricts to a dagger on \mathcal{C}_P .

We are interested in understanding how many dagger categories we can get by this procedure that are not unitarily equivalent. For this, first note that if $P \subseteq P'$, inclusion $\mathcal{C}_P \to \mathcal{C}_{P'}$ defines a dagger functor, which is fully faithful. However, even when $P \neq P'$ this inclusion can still be a unitary equivalence. Namely, we will show that adding *transfers* of Hermitian fixed points to P does not change the unitary equivalence class of \mathcal{C}_P : **Definition 5.1.** Given a Hermitian fixed point $h: c \to dc$ and an isomorphism $g: c' \to c$, the *transfer* of h by g is the Hermitian fixed point defined on c' by $d(g) \circ h \circ g$.

Note that this is indeed a Hermitian fixed point because the following diagram commutes



Lemma 5.2. Two objects $(c,h), (c',h') \in \text{Herm } C$ are unitarily isomorphic if and only if h' is a transfer of h.

Proof. An isomorphism $\alpha \colon (c', h') \to (c, h)$ is unitary if and only if

$$\alpha^{-1} = \alpha^{\dagger} \stackrel{\text{defn}}{=} h'^{-1} \circ d\alpha \circ h.$$

This happens if and only if $h' = d\alpha \circ h \circ \alpha$.

Definition 5.3. Given a category C, let $\pi_0(C)$ denote the collection of isomorphism classes of objects. If C is additionally a dagger category, let $\pi_0^U(C)$ denote the collection of unitary isomorphism classes of objects.

We can rephrase the above lemma by saying that $\pi_0^U(\text{Herm }\mathcal{C})$ is the collection of Hermitian fixed points (c, h) modulo transfer. Note that a dagger functor is unitarily essentially surjective if and only if it is surjective on π_0^U . In particular, if P is a collection of Hermitian fixed points and P' is the closure of P under transfers, then $\mathcal{C}_P \to \mathcal{C}_{P'}$ is unitarily essentially surjective and hence an equivalence of dagger categories. Therefore we can assume without loss of generality that P is closed under transfers.

Now let $P_c \subseteq P$ denote the subset of Hermitian fixed points on the object c. Then we will want to require that $P_c \neq \emptyset$, so that every object has 'some positive Hermitian structure'. This will additionally ensure that $C_P \to \text{Herm} C$ is essentially surjective.

This discussion motivates us to make the following definition.

Definition 5.4. Let (\mathcal{C}, d, η) be a category with anti-involution. A *notion of* positivity on \mathcal{C} is a collection of subsets

$$P = \{P_c \subset \operatorname{Hom}_{\mathcal{C}}(c, d(c)) : c \in \operatorname{obj} \mathcal{C}\}$$

such that:

• each P_c is non-empty,

- each $(h: c \to d(c)) \in P_c$ is a Hermitian fixed point,
- *P* is closed under transfer.

Remark 5.5. A necessary and sufficient condition for an anti-involutive category to admit a notion of positivity is that every object admits some Hermitian fixed point structure.

Example 5.6. If C is a category with anti-involution in which every object admits some Hermitian structure, we can take P to consist of all Hermitian fixed points. This is a notion of positivity and $C_P = \text{Herm } C$.

Corollary 5.7. Positivity notions on an anti-involutive category C are in bijection with subsets $[P] \subset \pi_0^U(\text{Herm } C)$ such that the composite

$$[P] \subset \pi_0^U(\operatorname{Herm} \mathcal{C}) \to \pi_0(\mathcal{C})$$

is surjective.

Proof. It follows immediately by the lemma above that a subset $[P] \subset \pi_0^U(\operatorname{Herm} \mathcal{C})$ is equivalent to a choice of Hermitian fixed points on some collection of objects of \mathcal{C} that is additionally closed under transfer. The condition that the given composite is surjective is equivalent to requiring that for every object c there exists an isomorphic object c' and a Hermitian fixed point $h: c' \to dc'$ such that $(c', h: c' \to dc') \in P_{c'}$. In case such (c', h) exists, we also get that $P_c \neq \emptyset$ by transferring h to c. Conversely it is clear that the desired composite is surjective if $P_c \neq \emptyset$ for all c.

Example 5.8. Recall that if (\mathcal{D}, \dagger) is a dagger category, a Hermitian fixed point on the anti-involutive category $T(\mathcal{D}, \dagger)$ is the same as a self-adjoint automorphism $h : h : c \to c^{\dagger} = c$. There is a canonical positivity structure on $T(\mathcal{D}, \dagger)$, which is defined by

 $P_c := \{h: c \to c \mid \text{ There is an automorphism } a: c \to c \text{ with } h = a^{\dagger} \circ a \}.$

Remark 5.9. An endomorphism $h: c \to c$ in a dagger category (or C^* -category) is called *positive* if there is an *endomorphism* $e: c \to c$ with $h = e^{\dagger} \circ e$. The set $P_c \subset \hom_{\mathcal{C}}(c,c)$ from example 5.8 is contained in the set of positive automorphisms. However, it is not true in general that every positive automorphism is in P_c : it might happen that some automorphism $h: c \to c$ can be written $h = a^{\dagger} \circ a$ for $a: c \to c$ some endomorphism, but that a cannot be chosen to be invertible.

Example 5.10. We study the case of finite-dimensional vector spaces with $d = \overline{(.)}^*$ as before. Note that $\pi_0^U(\operatorname{Herm}(\operatorname{Vect}_{\mathbb{C}}^{\operatorname{fd}})) = \mathbb{N} \times \mathbb{N}$ given by the signature of the corresponding Hermitian pairing. Here the signature of (V, (., .)) is the pair (p, q) so that there exists an orthonormal basis $\{e_1, \ldots, e_{p+q}\}$ of V with

$$(e_i, e_i) = 1$$
 $(e_j, e_j) = -1$

for $i \leq p$ and j > p. The forgetful map $\pi_0^U(\operatorname{Herm}(\operatorname{Vect}^{\operatorname{fd}}_{\mathbb{C}})) \to \pi_0(\operatorname{Vect}^{\operatorname{fd}}_{\mathbb{C}}) = \mathbb{N}$ is addition.

We provide some examples of notions of positivity on this category with anti-involution. One example is to take P_V to be the collection of positive definite Hermitian inner products on V. This is a notion of positivity because every finite-dimensional vector space admits a positive definite inner product. The resulting dagger category C_P is the dagger category of finite-dimensional Hilbert spaces. Another example is to take P_V to consist of all Hermitian inner products in which case C_P is the dagger category of finite-dimensional vector spaces with arbitrary Hermitian inner products.

There are also many more unusual notions of positivity on $\operatorname{Vect}_{\mathbb{C}}^{\mathrm{fd}}$. Namely, for every dimension d we can separately specify a nonempty collection of signatures $(p,q) \in \mathbb{N} \times \mathbb{N}$ such that p + q = d we allow for Hermitian forms. Any such choice gives a dagger category \mathcal{C}_P and two different choices are not dagger equivalent compatibly with the map to $\operatorname{Herm}(\operatorname{Vect}_{\mathbb{C}}^{\mathrm{fd}})$. Note that it would be reasonable to restrict the allowed notions of positivity further by requiring compatibility with tensor products or direct sums, but we will not pursue this further here.

Observation 5.11. Given an involutive functor $(F, \varphi) \colon \mathcal{C} \to \mathcal{D}$ and notions of positivity P on \mathcal{C} and Q on \mathcal{D} the following are equivalent:

1. For all $(h: c \to d(c)) \in P_c$, we have

Herm $F(c,h) = (\varphi_c \circ F(h) \colon F(c) \to d(F(c))) \in Q_{F(c)}.$


2. The map $\pi_0^U(\operatorname{Herm} \mathcal{C}) \to \pi_0^U(\operatorname{Herm} \mathcal{D})$ induced by F sends [P] to a subset of [Q].

Definition 5.12. The 2-category PCat has as objects anti-involutive categories equipped with a notion of positivity. Morphisms are involutive functors that intertwine the positivity notions in the sense of equivalent conditions in observation 5.11. The 2-morphisms are the same as in ICat.

Remark 5.13. Note that the forgetful functor $PCat \rightarrow ICat$ is well-behaved: Positivity notions can be transported along equivalences of categories and they can be restricted along fully faithful functors. Therefore the forgetful functor has lifts for equivalences. Moreover, if we restrict to fully faithful functors as morphisms in both 2-categories, then the functor $PCat^{\rm ff} \rightarrow ICat^{\rm ff}$ is equivalent to the Grothendieck construction of the functor $(ICat^{\rm ff})^{\rm op} \rightarrow PoSet$ that sends an anti-involutive category to its poset of possible notions of positivity.

We can also easily characterise the equivalences in the 2-category PCat. An involutive functor $(F, \varphi) : (\mathcal{C}, P) \to (\mathcal{D}, Q)$ is an equivalence in PCat, if and only if F is an equivalence of categories (and hence (F, φ) is an equivalence in ICat by lemma 2.5), and moreover the induced map of sets $[P] \to [Q]$ is surjective. (This map is automatically injective since F is fully faithful.) The latter condition says that every positive Hermitian fixed point in (\mathcal{D}, Q) is (up to transfer) of the form $\varphi_c \circ F(h)$ for $h: c \to d(c)$ a positive Hermitian fixed point in (\mathcal{C}, P) .

Theorem 5.14. Equipping $T(\mathcal{D}, \dagger)$ with the notion of positivity from example 5.8 defines a lift T_p :



This 2-functor T_p is a biequivalence.

Proof. We define an inverse 2-functor

$$\operatorname{Herm}_P\colon\operatorname{PCat}\to\operatorname{\dagger}\operatorname{Cat}$$

by declaring $\operatorname{Herm}_P(\mathcal{C}, d, \eta, P) \subset \operatorname{Herm}(\mathcal{C}, d, \eta)$ to be the full sub-†-category on those hermitian fixed points $(h: c \to d(c))$ where $h \in P_c$. In other words, Herm_P is the subcategory of hermitian fixed points which are positive. This is a well-defined 2-functor because Herm is and because 1-morphisms in PCat preserve the notions of positivity by definition.

The functor $U: \mathcal{D} \to \operatorname{Herm}(\operatorname{T}(\mathcal{D}))$ from definition 4.7 that sends x to (x, id_x) , restricts to a functor $U_P: \mathcal{D} \to \operatorname{Herm}_P(\operatorname{T}_P(\mathcal{D}))$ since (x, id_x) is always a positive hermitian fixed point in $\operatorname{T}_P(\mathcal{D})$. Therefore this defines a natural transformation $U: \operatorname{id}_{\dagger\operatorname{Cat}} \to \operatorname{Herm}_P \circ \operatorname{T}_P$ and as observed below definition 4.7 the dagger functor

$$U_P \colon \mathcal{D} \to \operatorname{Herm}_P(\mathcal{T}_P(\mathcal{D}))$$

is always fully faithful (and essentially surjective). We would like to show that it is surjective up to isometry. Let $(h: x \to x)$ be some object in $\operatorname{Herm}_P(\operatorname{T}_P(\mathcal{D}))$. Since h is positive we can write it as $h = a \circ a^{\dagger}$ for some positive Hermitian fixed point $a: x \to x$ in $\operatorname{T}_P(\mathcal{D})$. In other words, h is the transfer of (id: $x \to x$) along a. By lemma 5.2 this means that there is a unitary isomorphism $(x, \operatorname{id}_x) \cong (x, h)$. Therefore U_P is a dagger equivalence and hence an equivalence in the 2-category \dagger Cat.

Finally, consider the involutive functor

$$(K_{\mathcal{C}}, \varphi_{\mathcal{C}}) \colon \mathrm{T}(\mathrm{Herm}(\mathcal{C}, d, \eta)) \longrightarrow (\mathcal{C}, d, \eta)$$

that we constructed naturally for all $(\mathcal{C}, d, \eta) \in \text{ICat}$ in 4.5. Given a positivity notion P on \mathcal{C} , we also get a positivity notion on $\mathcal{T}_P(\text{Herm}_P(\mathcal{C}, d, \eta))$. We would like to show that $K_{\mathcal{C}}$ preserves positivity notions. A positive Hermitian fixed point in $\mathcal{T}_P(\text{Herm}_P(\mathcal{C}, d, \eta))$ is of the form $p = a^{\dagger} \circ a$ for some automorphism $a: (x, h) \to (x, h)$. Here $h: x \to d(x)$ is a positive hermitian fixed point. Using the definition of the dagger in $\text{Herm}(\mathcal{C})$ we can write this as $p = (h^{-1} \circ d(a) \circ h) \circ a$. In order to show that $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ respects the positivity notions we use condition (1) of observation 5.11, which says that $\varphi_{(x,h)} \circ K_{\mathcal{C}}(p)$ must be positive. Using $\varphi_{(x,h)} = h: x \to d(x)$ we see that

$$\varphi_{(x,h)} \circ K_{\mathcal{C}}(p) = h \circ (h^{-1} \circ d(a) \circ h \circ a) = d(a) \circ h \circ a.$$

This is the transfer of h along a and since h was positive, so is this. Therefore $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ defines a natural morphism

$$T_P(\operatorname{Herm}_P(\mathcal{C}, d, \eta, P)) \longrightarrow (\mathcal{C}, d, \eta, P)$$

of anti-involutive categories with notions of positivity. We already observed in lemma 4.6 that $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$ is an equivalence in ICat. For it to also be an equivalence in PCat we need to check that every positive fixed point in $(\mathcal{C}, d, \eta, P)$ is hit (up to transfer) by $(K_{\mathcal{C}}, \varphi_{\mathcal{C}})$. We saw above that every morphism $d(a) \circ h \circ a$ can be written as $\varphi_{(x,h)} \circ F(p)$. Setting $a = \mathrm{id}_x$ we see that indeed every positive fixed point h can be hit by this.

Corollary 5.15. Let (\mathcal{C}, \dagger) and (\mathcal{D}, \dagger) be two dagger categories. Then $F \mapsto \dagger_{\mathcal{D}} \circ F \circ \dagger_{\mathcal{C}}$ defines an anti-involution on the category of all functors $\mathcal{C} \to \mathcal{D}$. The inclusion of the dagger functors into the fixed points

$$\operatorname{Fun}^{\dagger}((\mathcal{C},\dagger),(\mathcal{D},\dagger)) \hookrightarrow (\operatorname{Fun}(\mathcal{C},\mathcal{D}))^{\operatorname{fix}}$$

is fully faithful and its essential image consists of those functors that preserve the positivity notions.

Proof. We have seen in 3.9 that given two anti-involutive categories $(\mathcal{C}, d_{\mathcal{C}}), (\mathcal{D}, d_{\mathcal{D}})$ there is an anti-involution d on Fun $(\mathcal{C}, \mathcal{D})$ given by the expression $F \mapsto d_{\mathcal{D}} \circ F \circ d_{\mathcal{C}}$. Its fixed points are the category of which objects are involutive functors and morphisms are involutive natural transformations. Specializing to the case where the anti-involutive categories come from dagger categories, we see that

$$(\operatorname{Fun}(\mathcal{C}, \mathcal{D}))^{\operatorname{fix}} = \operatorname{Hom}_{\operatorname{ICat}}(T\mathcal{C}, T\mathcal{D}).$$

The corollary now follows directly from the main theorem.

References

- [Ati88] Michael Atiyah. "Topological quantum field theory". In: *Publications Mathématiques de l'IHÉS* 68 (1988), pp. 175–186.
- [BM09] Edwin Beggs and Shahn Majid. "Bar categories and star operations". In: Algebras and Representation Theory 12.2-5 (2009), pp. 103–152.
- [BSW19] Marco Benini, Alexander Schenkel, and Lukas Woike. "Involutive categories, colored *-operads and quantum field theory". In: *Theory* and Applications of Categories 34.2 (2019), pp. 13–57.
- [Egg11] Jeffrey Egger. "On involutive monoidal categories". In: Theory and Applications of Categories 25.14 (2011), pp. 368–393.
- [FH21] Daniel Freed and Michael Hopkins. "Reflection positivity and invertible topological phases". In: Geometry & Topology 25.3 (2021), pp. 1165–1330.

- [Hes17] Jan Hesse. "Group Actions on Bicategories and Topological Quantum Field Theories". PhD thesis. Staats-und Universitätsbibliothek Hamburg Carl von Ossietzky, 2017.
- [HL+15] Andre Henriques, Peter LeFanu Lumsdaine, et al. Are dagger categories truly evil? https://mathoverflow.net/questions/220032/are-dagger-categories-trul 2015.
- [HP20] André Henriques and David Penneys. *Representations of fusion categories and their commutants.* Available at arXiv:2004.08271. 2020.
- [HPT23] André Henriques, David Penneys, and James Tener. Unitary anchored planar algebras. Available at arXiv:2301.11114. 2023.
- [Jac12] Bart Jacobs. "Involutive categories and monoids, with a GNS-correspondence". In: *Foundations of Physics* 42 (2012), pp. 874–895.
- [KS06] Max Kelly and Ross Street. "Review of the elements of 2-categories". In: Category Seminar: Proceedings Sydney Category Theory Seminar 1972/1973. Springer. 2006, pp. 75–103.
- [MS23] Lukas Müller and Luuk Stehouwer. "Reflection Structures and Spin Statistics in Low Dimensions". In: *arXiv preprint arXiv:2301.06664* (2023).
- [SCC21] Priyaa Srinivasan, Cole Comfort, and Robin Cockett. "Dagger linear logic for categorical quantum mechanics". In: Logical Methods in Computer Science 17 (2021).
- [Vic11] Jamie Vicary. "Completeness of †-categories and the complex numbers". In: J. Math. Phys. 52 (2011), pp. 82–104.
- [Yau20] Donald Yau. Involutive Category Theory. Springer, 2020.