Ricci flow on Finsler manifolds

B. Bidabad^{*} and M. K. Sedaghat

Abstract

This paper investigates the short-time existence and uniqueness of Ricci flow solutions on Finsler manifolds. The main results of this paper are theorems demonstrating the short-time existence of the flow solution for n-dimensional Finsler manifolds and the uniqueness of the solution for isotropic Finsler manifolds. Two examples are also presented to illustrate the results.

Keywords: Ricci flow, Ricci-DeTurck flow, parabolic differential equation, Finsler space. **AMS subject classification**: 53C60, 53B40.

1 Introduction

One of the primary aims of geometric flows is to produce canonical geometric structures by deforming general initial metrics to these canonical structures. The Ricci flow theory and its applications have become a rapidly developing branch of mathematics, with its most remarkable achievement being G. Perelman's proof of Thurston's geometrization conjecture. Hamilton's Ricci flow, introduced in 1982, is a geometric flow with numerous applications in physics and real-world problems. The Ricci flow conformally deforms the Riemannian metric to its induced curvature, such that the curvature tensors evolve by a system of diffusion equations that distributes the curvature uniformly over the manifold. Hamilton demonstrated that, on a closed manifold, there is a unique solution to the Ricci flow over a sufficiently short-time, and that on a compact three-manifold with an initial

^{*}The corresponding author bidabad@aut.ac.ir; behroz.bidabad@math.univ-toulouse.fr

metric of positive Ricci curvature, the Ricci flow converges, after re-scaling to keep constant volume, to a metric of positive constant sectional curvature, proving that the manifold is diffeomorphic to the three-sphere \mathbb{S}^3 or a quotient of the three-sphere \mathbb{S}^3 by a linear group of isometries.

The Ricci flow has demonstrated its great potential in geometry and physics, solving various problems that cannot easily be addressed by alternative methods. In discrete geometry, the discrete Ricci flow is a powerful tool for computing metrics with prescribed Gaussian curvatures on general surfaces. Examples of applications related to the discrete Ricci flow include global conformal parameterizations in computer graphics and analysis of medical imaging, surface matching, and manifold splines.

In Finsler geometry, a natural generalization of Riemannian geometry, the problem of constructing the Finslerian Ricci flow raises several new conceptual and fundamental issues regarding the compatibility of geometrical and physical objects and their optimal configurations. A fundamental step in studying any system of evolutionary partial differential equations is to show the short-time existence and uniqueness of solutions. Several joint works have studied the evolution of a family of Finsler metrics along the Ricci flow, demonstrating that Finsler Ricci flow exists in a short time and converges to a limit metric (see [9]). S. Lakzian has further applied the Finsler–Ricci flow to Harnack estimates of the heat equation, using the tools presented by S. Ohta and K. Sturm in [17].

The authors of the present work have shown the existence and uniqueness of solutions to the Ricci flow on Finsler surfaces in [6], extending the work of R. S. Hamilton on Riemannian surfaces [15]. These results have numerous applications in general relativity, though too many to list here. However, in dimensions n > 2, the uniqueness of solutions is no longer valid, [8]. In [7], it was further shown that the Finsler Ricci flow preserves the positivity of the reduced *hh*-curvature on finite time, and the evolution of the Ricci scalar is a parabolic-type equation. Additionally, if the initial Finsler metric has positive flag curvature, then the flag curvature, as well as the Ricci scalar, remain positive as long as the solution exists.

In the present work, we investigate the Ricci flow on closed *n*-dimensional Finsler manifolds, with n > 2, and prove the short-time existence of solutions. Intuitively, since the Ricci flow system of equations is only weakly parabolic, the standard theory of parabolic equations cannot be used to prove its short-time existence and uniqueness. Following the procedure outlined by D. DeTurck in the Riemannian space [13], we introduce the Finslerian Ricci-DeTurck flow by Eq. (32) and study the solutions. We then find a solution to the original Ricci flow by pulling back the solution of the Ricci-DeTurck flow using suitable diffeomorphisms and prove the existence of short-time solutions.

Theorem 1. Let M be a compact Finsler manifold. Given any initial Finsler structure F_0 ,

there exists a real number T > 0 and a smooth one-parameter family of Finsler structures $\tilde{F}(t)$, for all $t \in [0,T)$, such that $\tilde{F}(t)$ is a solution to the Finslerian Ricci-DeTurck flow with $\tilde{F}(0) = F_0$.

Then, a solution to the original Ricci flow is obtained by pulling back the solution to the Ricci-DeTurck flow via appropriate diffeomorphisms. This result leads to

Theorem 2. Let M be a compact Finsler manifold. Given any initial Finsler structure F_0 , there exists a real number T > 0 and a smooth one-parameter family of Finsler structures F(t), for all $t \in [0,T)$ such that F(t) is a solution to the Finslerian Ricci flow and $F(0) = F_0$.

The uniqueness of the solution to the Ricci flow can be demonstrated by the requirement that the Finsler manifolds be isotropic for the solution to be unique.

Theorem 3. Let M be a compact isotropic Finsler manifold. Given any initial Finsler structure F_0 , there exists a real number T > 0 and a smooth one-parameter family of Finsler structures F(t), for all $t \in [0,T)$, such that F(t) is a unique solution to the Finslerian Ricci flow and $F(0) = F_0$.

2 Preliminaries and notations

2.1 Chern connection; A global approach

Let M be a real n-dimensional smooth manifold, and denote by TM the tangent bundle of tangent vectors, by $\pi : TM_0 \longrightarrow M$ the fiber bundle of non-zero tangent vectors and by $\pi^*TM \longrightarrow TM_0$ the pullback tangent bundle. Let F be a Finsler structure on TM_0 , and g the related Finslerian metric [3]. A *Finsler manifold* is denoted here by the pair (M, F). Any point of TM_0 is denoted by z = (x, y), where $x = \pi z \in M$ and $y \in T_xM$.

Let us denote by TTM_0 , the tangent bundle of TM_0 and by ρ , the canonical linear mapping $\rho: TTM_0 \longrightarrow \pi^*TM$, where, $\rho = \pi_*$. For all $z \in TM_0$, V_zTM is the set of all vertical vectors at z, that is, the set of vectors which are tangent to the fiber through z. Consider the decomposition $TTM_0 = HTM \oplus VTM$, which allows to uniquely represent a vector field $\hat{X} \in \mathcal{X}(TM_0)$ as the sum of the horizontal and vertical parts namely, $\hat{X} = H\hat{X} + V\hat{X}$. The corresponding basis is denoted here by $\{\frac{\delta}{\delta x^i}, \frac{\partial}{\partial y^i}\}$, where, $\frac{\delta}{\delta x^i} := \frac{\partial}{\partial x^i} N_i^j \frac{\partial}{\partial y^j}$, $N_i^j = \frac{1}{2} \frac{\partial G^j}{\partial y^i}$ and G^i are the spray coefficients defined by $G^i = \frac{1}{4}g^{ih}(\frac{\partial^2 F^2}{\partial y^h \partial x^j}y^j - \frac{\partial F^2}{\partial x^h})$. We denote the formal Christoffel symbols by $\gamma_{jk}^i = \frac{1}{2}g^{ih}(\partial_j g_{hk} + \partial_k g_{jh} - \partial_h g_{jk})$ where, $\partial_k = \frac{\partial}{\partial x^k}$. The dual bases are denoted by $\{dx^i, \delta y^i\}$ where, $\delta y^i := dy^i + N_j^i dx^j$. Let us denote a global representation of the Chern connection by $\nabla : \mathcal{X}(TM_0) \times \Gamma(\pi^*TM) \longrightarrow$ $\Gamma(\pi^*TM)$. Consider the linear mapping $\mu: TTM_0 \longrightarrow \pi^*TM$, defined by $\mu(\hat{X}) = \nabla_{\hat{X}} \mathbf{y}$ where, $\hat{X} \in TTM_0$ and $\mathbf{y} = y^i \frac{\partial}{\partial x^i}$ is the canonical section of π^*TM .

The connection 1-forms of Chern connection in these bases are given by $\omega_j^i = \Gamma_{jk}^i dx^k$ where, $\Gamma_{jk}^i = \frac{1}{2}g^{ih}(\delta_j g_{hk} + \delta_k g_{jh} - \delta_h g_{jk})$ and $\delta_k = \frac{\delta}{\delta x^k}$. In the sequel, all the vector fields on TM_0 are decorated with a hat and denoted by \hat{X} , \hat{Y} and \hat{Z} , and the corresponding sections of π^*TM by $X = \rho(\hat{X})$, $Y = \rho(\hat{Y})$ and $Z = \rho(\hat{Z})$, respectively unless otherwise specified. The torsion freeness and almost metric compatibility of the Chern connection are given by

$$\nabla_{\hat{X}}Y - \nabla_{\hat{Y}}X = \rho[\hat{X}, \hat{Y}],\tag{1}$$

$$(\nabla_{\hat{z}}g)(X,Y) = 2C(\mu(\hat{Z}),X,Y), \tag{2}$$

respectively, where C is the Cartan tensor with the components $C_{ijk} = \frac{\partial g_{ij}}{\partial y^k}$. In local coordinates on TM the Chern horizontal and vertical covariant derivatives of an arbitrary (1,2) tensor field S on π^*TM with the components $S^i_{ik}(x,y)$ on TM are given by

$$\begin{aligned} \nabla_l S^i_{jk} &:= \delta_l S^i_{jk} - S^i_{sk} \Gamma^s_{jl} - S^i_{js} \Gamma^s_{kl} + S^s_{jk} \Gamma^i_{sl}, \\ \dot{\nabla}_l S^i_{jk} &:= \dot{\partial}_l S^i_{jk}, \end{aligned}$$

where, $\nabla_l := \nabla_{\frac{\delta}{\delta x^l}}$ and $\dot{\nabla}_l := \nabla_{\frac{\partial}{\partial y^l}}$. Horizontal metric compatibility of the Chern connection in local coordinates translates into $\nabla_l g_{jk} = 0$, see [3, p. 45]. The local *Chern hh-curvature* tensor is given by

$$R_{j\ kl}^{\ i} = \delta_k \Gamma_{jl}^i - \delta_l \Gamma_{jk}^i + \Gamma_{hk}^i \Gamma_{jl}^h - \Gamma_{hl}^i \Gamma_{jk}^h, \tag{3}$$

see [3, p. 52]. In the isotropic case, the contracted hh-curvature tensor $R_{j\ il}^{i}$ is symmetric in two indices j and l. In fact, by means of the symmetry of $R_{j\ il}^{i}$ in Cartan connection in the isotropic case, see [1, p. 152], and the relation between the Cartan and Chern hhcurvature tensors, one can see that in the isotropic case, $R_{j\ il}^{i}$ is symmetric in j and l. More intuitively, it can be part of a symmetric quadratic form.

The reduced hh-curvature tensor is a connection free tensor field which is also referred to as the *Riemann curvature* by certain authors. In local coordinates on TM, the components of the reduced hh-curvature tensor are given by $R^i_k := \frac{1}{F^2} y^j R^i_{j\ km} y^m$, which are entirely expressed in terms of x and y derivatives of spray coefficients G^i as follows

$$R^{i}_{\ k} := \frac{1}{F^{2}} \left(2 \frac{\partial G^{i}}{\partial x^{k}} - \frac{\partial^{2} G^{i}}{\partial x^{j} \partial y^{k}} y^{j} + 2G^{j} \frac{\partial^{2} G^{i}}{\partial y^{j} \partial y^{k}} - \frac{\partial G^{i}}{\partial y^{j}} \frac{\partial G^{j}}{\partial y^{k}} \right), \tag{4}$$

see [3, p. 66].

Let (x, y) be an element of TM and $P(y, X) \subset T_x(M)$ a 2-plane generated by the vectors y and X in $T_x(M)$. Then the flag curvature K(x, y, X) with respect to the plane

P(y,X) at a point $x\in M$ is defined by

$$K(x,y,X) := \frac{g(R(X,y)y,X)}{g(X,X)g(y,y) - g(X,y)^2}$$

where R(X, y)y is the *hh*-curvature tensor. If K is independent of X, then the Finsler manifold (M, F) is called *isotropic* or *space of scalar curvature*. If K has no dependence on x or y, then the Finsler manifold is said to be of *constant curvature*.

2.2 Lie derivatives of Finsler metrics

The Lie derivative of an arbitrary Finslerian (0, 2) tensor field $\mathcal{T} = \mathcal{T}_{jk}(x, y) dx^j \otimes dx^k$ on $\otimes^2 \pi^* TM$ with respect to an arbitrary vector field \hat{V} on TM_0 is given by

$$(\mathcal{L}_{\hat{V}}\mathcal{T})(X,Y) = \hat{V}(\mathcal{T}(X,Y)) - \mathcal{T}(\rho[\hat{V},\hat{X}],Y) - \mathcal{T}(X,\rho[\hat{V},\hat{Y}]),$$

where, $\rho(\hat{X}) = X$, $\rho(\hat{Y}) = Y$ and $\hat{X}, \hat{Y} \in T_z T M_0$, see [5]. The Lie derivative of a Finsler metric g in direction of an arbitrary vector field \hat{V} on $T M_0$ is given by

$$(\mathcal{L}_{\hat{V}}g)(X,Y) = \hat{V}(g(X,Y)) - g(\rho[\hat{V},\hat{X}],Y) - g(X,\rho[\hat{V},\hat{Y}]).$$

Using the torsion freeness of Chern connection defined by (1), the Lie derivative of the Finsler metric g becomes

$$\begin{aligned} (\mathcal{L}_{\hat{V}}g)(X,Y) &= \hat{V}(g(X,Y)) - g(\nabla_{\hat{V}}X - \nabla_{\hat{X}}V,Y) - g(X,\nabla_{\hat{V}}Y - \nabla_{\hat{Y}}V) \\ &= \hat{V}(g(X,Y)) - g(\nabla_{\hat{V}}X,Y) + g(\nabla_{\hat{X}}V,Y) \\ &- g(X,\nabla_{\hat{V}}Y) + g(X,\nabla_{\hat{Y}}V). \end{aligned}$$
(5)

By the almost g-compatibility of Chern connection defined by (2), we have

$$2C(\mu(\hat{V}), X, Y) = (\nabla_{\hat{V}}g)(X, Y) = \hat{V}(g(X, Y)) - g(\nabla_{\hat{V}}X, Y) - g(X, \nabla_{\hat{V}}Y).$$

Therefore,

$$\hat{V}(g(X,Y)) = 2C(\mu(\hat{V}), X, Y) + g(\nabla_{\hat{V}}X, Y) + g(X, \nabla_{\hat{V}}Y).$$
(6)

Plugging the equation (6) in (5) we obtain

$$(\mathcal{L}_{\hat{V}}g)(X,Y) = 2C(\mu(\hat{V}),X,Y) + g(\nabla_{\hat{X}}V,Y) + g(X,\nabla_{\hat{Y}}V).$$
(7)

Replacing X and Y by the canonical section $\mathbf{y} = y^i \frac{\partial}{\partial x^i}$ in (7) we obtain

$$(\mathcal{L}_{\hat{V}}g)(\mathbf{y},\mathbf{y}) = 2C(\mu(\hat{V}),\mathbf{y},\mathbf{y}) + g(\nabla_{\hat{\mathbf{y}}}V,\mathbf{y}) + g(\mathbf{y},\nabla_{\hat{\mathbf{y}}}V),$$

where, $\hat{\mathbf{y}} = y^i \frac{\delta}{\delta x^i}$. Using $C(\mu(\hat{V}), \mathbf{y}, \mathbf{y}) = 0$, see [3, p. 23], and the symmetric property of $g(\nabla_{\hat{\mathbf{y}}} V, \mathbf{y})$ one arrives at

$$(\mathcal{L}_{\hat{V}}g)(\mathbf{y},\mathbf{y}) = 2g(\mathbf{y},\nabla_{\hat{\mathbf{y}}}V),\tag{8}$$

where, $V = v^i \frac{\partial}{\partial x^i}$ is a section of π^*TM . In local coordinates, the above equation translates into

$$y^i y^j \mathcal{L}_{\hat{V}} g_{ij} = 2y^i y^j g_{ik} \nabla_j v^k$$

Using $\nabla_j g_{ik} = 0$, we obtain

$$y^{i}y^{j}\mathcal{L}_{\hat{V}}g_{ij} = 2y^{i}y^{j}\nabla_{j}v_{i}, \qquad (9)$$

where, $v_i = g_{ik} v^k$.

2.3 A geometric setup on *SM*

Let (M, F) be a Finsler manifold and SM the quotient of TM_0 under the following equivalence relation: $(x, y) \sim (x, \tilde{y})$ if and only if y, \tilde{y} are positive multiples of each other. In other words, SM is the bundle of all directions or rays, and is called the (projective) sphere bundle. The local coordinates $x^1, ..., x^n$ on M induce global coordinates $y^1, ..., y^n$ on each fiber T_xM , through the expansion $y = y^i \frac{\partial}{\partial x^i}$. Therefore $(x^i; y^i)$ is a coordinate system on SM, where the coordinates y^i are regarded as homogeneous coordinates in the projective space. Using the canonical projection $p: SM \longrightarrow M$, one can pull the tangent bundle TM back to p^*TM which is a vector bundle with the fiber dimension n over the (2n-1)-manifold SM. The vector bundle p^*TM has a global section $l := \frac{y^i}{F(y)} \frac{\partial}{\partial x^i}$ and a natural Riemannian metric which we here denote by $g := g_{ij}(x, y)dx^i \otimes dx^j$.

Let $\{e_a = u_a^i \frac{\partial}{\partial x^i}\}$ be a g-orthonormal frame for p^*TM with $e_n := l$ and $\{\omega^a = v_i^a dx^i\}$ be its coframe; thus $\omega^a(e_b) = \delta_b^a$. It is clear that $\omega^n = \frac{\partial F}{\partial y^i} dx^i$ which is a global section of p^*T^*M . Also we have $\frac{\partial}{\partial x^i} = v_i^a e_a$ and $dx^i = u_a^i \omega^a$. A basic relation between (u_a^i) and (v_i^a) is given by $v_i^a u_b^i = \delta_b^a$ and $u_a^i v_j^a = \delta_j^i$. For convenience, we shall also regard the e_a 's and ω^a 's as local vector fields and 1-forms, respectively on SM. All p^*TM related indices are raised and lowered with the metric g, see [4]. Let

$$\begin{aligned} \hat{e}_a &= u_a^i \frac{\delta}{\delta x^i}, \quad \hat{e}_{n+a} &= u_a^i F \frac{\partial}{\partial y^i}, \\ \omega^a &= v_a^i dx^i, \quad \omega^{n+a} &= v_a^i \frac{\delta y^i}{F}. \end{aligned}$$

Also, recall that the Latin indices run over the range 1, ..., n and the Greek indices run from 1 to n-1. It can be shown that $\{\hat{e}_a, \hat{e}_{n+\alpha}\}$ is a local basis for the tangent bundle TSM and $\{\omega^a, \omega^{n+\alpha}\}$ is a local basis for the cotangent bundle T^*SM . Tangent vectors on SM which are annihilated by all $\{\omega^{n+\alpha}\}$'s form the horizontal sub-bundle HSM of TSM. The fibers of HSM are n-dimensional. On the other hand, let $VSM := \bigcup_{x \in M} T(S_xM)$ be the vertical sub-bundle of TSM; its fibers are n-1 dimensional. The decomposition $TSM = HSM \oplus VSM$ holds because HSM and VSM are direct summands. The sphere bundle $SM \subset TM$ is a (2n - 1)-dimensional Riemannian manifold equipped with the induced Riemannian metric

$$\hat{g} := \delta_{ab} \omega^a \otimes \omega^b + \delta_{\alpha\beta} \omega^{n+\alpha} \otimes \omega^{n+\beta}.$$

In particular, HSM and VSM are orthogonal with respect to \hat{g} .

2.4 Integrability condition Lemma

Here, we first recall that not all Riemannian metrics on the fibers of the pulled-back bundle come from a Finsler structure. Eventually, any arbitrary symmetric positive-definite (0, 2)tensor field $g_{ij}(x, y)$ does not arise from a Finsler structure F(x, y). Intuitively, to ensure that $g_{ij}(x, y)$ are components of a Finsler structure, the essential integrability criterion is the total symmetry of $(g_{ij})_{y^k} := \frac{\partial g_{ij}}{\partial y^k}$ on all the three indices i, j, and k. In fact, $g_{ij}(x, y)$ follows from a Finsler structure F(x, y) if and only if $(g_{ij})_{y^k}$ is totally symmetric in its three indices, see [2, p. 56]. Symmetry of $(g_{ij})_{y^k}$ on all the three indices i, j, and k is known in the literature as *integrability condition*. Moreover, we must ensure the integrability criterion is satisfied in every step along with the Ricci flow. To this end we consider a general evolution equation given by

$$\frac{\partial}{\partial t}g(t) = \omega(t), \quad g(0) := g_0, \tag{10}$$

where, $\omega(t) := \omega(t, x, y)$ is a family of symmetric (0, 2)-tensors on π^*TM , zero-homogenous with respect to y and $(\omega_{ij})_{y_k} = \frac{\partial \omega_{ij}}{\partial y^k}$ be totally symmetric in the three indices i, j and k. The following Lemma establishes the integrability condition, see also [9, p. 749].

Lemma 2.1. Let g(t) be a solution to the evolution equation (10). There is a family of Finsler structures F(t) on TM such that,

$$g_{ij}(t) = \frac{1}{2} \frac{\partial^2 F(t)}{\partial y^i \partial y^j}.$$
(11)

Proof. Let M be a compact differential manifold, F(t) a family of smooth 1-parameter Finsler structures on TM_0 and g(t) the Hessian matrix of F(t) which defines a scalar product on π^*TM for every t. Let g(t) be a solution to the evolution equation (10). We have

$$g(t) = g(0) + \int_0^t \omega(\tau) d\tau, \qquad \forall \tau \in [0, t).$$
(12)

We show that the metric g(t) satisfies the integrability condition, or equivalently there is a Finsler structure F(t) on TM_0 satisfying (11). For this purpose, we multiply g_{ij} by y^i and y^j in (12),

$$y^i y^j g_{ij}(t) = y^i y^j g_{ij}(0) + \int_0^t y^i y^j \omega_{ij}(\tau) d\tau.$$

By means of the initial condition $y^i y^j g_{ij}(0) = F^2(0)$, we get

$$y^{i}y^{j}g_{ij}(t) = F^{2}(0) + \int_{0}^{t} y^{i}y^{j}\omega_{ij}(\tau)d\tau.$$
(13)

By positive definiteness assumption of g_{ij} , we put $F = (y^i y^j g_{ij})^{\frac{1}{2}}$. Twice vertical derivatives of (13) yields

$$\frac{1}{2}\frac{\partial^2 F^2}{\partial y^k \partial y^l} = g_{kl}(0) + \frac{1}{2} \int_0^t \frac{\partial^2}{\partial y^k \partial y^l} (y^i y^j \omega_{ij}(\tau)) d\tau.$$
(14)

On the other hand, by straightforward calculation we have

$$\frac{1}{2}\frac{\partial^2}{\partial y^k \partial y^l}(y^i y^j \omega_{ij}(\tau)) = \frac{1}{2}\frac{\partial^2 \omega_{ij}(\tau)}{\partial y^k \partial y^l}y^i y^j + \left(\frac{\partial \omega_{ik}(\tau)}{\partial y^l} - \frac{\partial \omega_{il}(\tau)}{\partial y^k}\right)y^i + \omega_{kl}(\tau), \quad (15)$$

for all $\tau \in [0, t)$. Since $(\omega_{ij})_{y_k}$ is totally symmetric in three indices i, j and k, we obtain

$$\frac{1}{2}\frac{\partial^2 \omega_{ij}(\tau)}{\partial y^k \partial y^l} y^i y^j = 0, \quad y^i \frac{\partial \omega_{ik}(\tau)}{\partial y^l} = 0, \quad y^i \frac{\partial \omega_{il}(\tau)}{\partial y^k} = 0.$$

Therefore, (15) is reduced to

$$\frac{1}{2}\frac{\partial^2}{\partial y^k \partial y^l} (y^i y^j \omega_{ij}(\tau)) = \omega_{kl}(\tau), \tag{16}$$

for all $\tau \in [0, t)$. Finally, replacing (16) in (14) we get

$$\frac{1}{2}\frac{\partial^2 F^2}{\partial y^k \partial y^l} = g_{kl}(0) + \int_o^t \omega_{kl}(\tau) d\tau = g_{kl}.$$

Therefore, every $g_{ij}(t)$ on the fibers of the pulled-back bundle, arises from a Finsler structure, which completes the proof.

Remark 2.1. Let g(t) be a solution to the evolution equation (10). The verification of the integrability of $g_{ij}(t)$ can also be done more simply. To verify that $(g_{ij})_{y_k} = (g_{ik})_{y_j}$, consider $(g_{ij})_{y_k} - (g_{ik})_{y_j}$, take the partial derivative ∂_t of the both terms and insert it in the evolution equation (10). Therefore, from the total symmetry of $(\omega_{ij})_{y_k}$ in i, j and kwe have

$$\partial_t \Big((g_{ij})_{y_k} - (g_{ik})_{y_j} \Big) = (w_{ij})_{y_k} - (w_{ik})_{y_j} = 0.$$

Thus $(g_{ij})_{y_k} - (g_{ik})_{y_j}$ does not depend on t. At t = 0, $g_{ij}(0)$ is the fundamental tensor of the initial data F_0 , so $(g_{ij})_{y_k} - (g_{ik})_{y_j} = 0$ at t = 0. The same is true for all t.

3 Semi-linear parabolic equations on SM

Recall that a quasi-linear system is a system of partial differential equations where, the derivatives of the principal order terms occur only linearly and coefficients can depend on the derivatives of the lower order terms. It is called *semi-linear* if it is quasi-linear and coefficients of the principal order terms depend only on the independent variables, but not on the solution, see [18, p. 45]. Let M be an n-dimensional smooth manifold and $u: M \longrightarrow \mathbb{R}$ a smooth function on M. A *semi-linear strictly parabolic* equation is a PDE of the form

$$\frac{\partial u}{\partial t} = a^{ij}(x,t)\frac{\partial^2 u}{\partial x^i\partial x^j} + h(x,t,u,\frac{\partial u}{\partial x^i}),$$

where, a^{ij} and h are smooth functions on M and for some constant $\lambda > 0$ we have the parabolic assumption

$$a^{ij}\xi_i\xi_j \ge \lambda \parallel \xi \parallel^2, \quad 0 \ne \xi \in \chi(M),$$

that is, all eigenvalues of $A = (a^{ij})_{2 \times 2}$ have positive signs or equivalently, A is positive definite.

Definition 3.1. Let M be an n-dimensional smooth manifold and $\phi : SM \longrightarrow \mathbb{R}$ a smooth function on the sphere bundle SM. Consider the following strictly parabolic semi-linear equation on SM;

$$\frac{\partial \phi}{\partial t} = G^{\scriptscriptstyle AB}(x, y, t) \hat{e}_{\scriptscriptstyle A} \hat{e}_{\scriptscriptstyle B} \phi + h(x, y, t, \rho, \hat{e}_{\scriptscriptstyle A} \phi), \qquad A, B = 1, 2, ..., (2n-1),$$

where, \hat{e}_A is a local frame for the tangent bundle TSM and stand for partial derivatives on SM. Here, G^{AB} and h are smooth functions on SM and $G = (G^{AB})$ is positive definite.

More precisely, a strictly parabolic semi-linear system on SM can be written in the following form.

$$\frac{\partial \phi}{\partial t} = p^{ab}(x, y, t) \hat{e}_a \hat{e}_b \phi + q^{\alpha\beta}(x, y, t) \hat{e}_{n+\alpha} \hat{e}_{n+\beta} \phi + m^{a\alpha}(x, y, t) \hat{e}_a \hat{e}_{n+\beta} \phi + \text{lower order terms},$$
(17)

where the Latin indices a, b, ... and the Greek indices $\alpha, \beta, ...$ run over the range 1, ..., nand 1, ..., (n - 1), respectively and the matrix

$$G = \left(\begin{array}{c|c} P & \frac{1}{2}M \\ \hline \frac{1}{2}M^t & Q \end{array}\right)_{(2n-1)\times(2n-1)},$$

is positive definite where, $P = (p^{ab})_{n \times n}, Q = (q^{\alpha\beta})_{(n-1) \times (n-1)}, M = (m^{a\alpha})_{n \times (n-1)}.$

Lemma 3.1. Let (M, F) be a Finsler manifold and $\phi : TM \longrightarrow \mathbb{R}$ a smooth function with zero homogeneity on the tangent bundle TM. The semi-linear differential equation

$$\frac{\partial\phi}{\partial t} = g^{ij} \frac{\delta^2 \phi}{\delta x^i \delta x^j} + F^2 g^{ij} \frac{\partial^2 \phi}{\partial y^i \partial y^j} + lower \ order \ terms, \tag{18}$$

is a strictly parabolic equation on SM.

Proof. Let us denote again by ϕ the restriction of ϕ on SM. By definition, we have

$$\begin{split} \hat{e}_a \phi &= u_a^i \frac{\delta \phi}{\delta x^i}, \\ \hat{e}_b \hat{e}_a \phi &= u_a^i u_b^j \frac{\delta^2 \phi}{\delta x^i \delta x^j} + \hat{e}_b(u_a^i) (\frac{\delta \phi}{\delta x^i}). \end{split}$$

Multiplying both sides of the above equation by g^{ab} we get

$$g^{ab}\hat{e}_b\hat{e}_a\phi = g^{ab}u^i_a u^j_b \frac{\delta^2 \phi}{\delta x^i \delta x^j} + g^{ab}\hat{e}_b(u^i_a)(\frac{\delta \phi}{\delta x^i})$$
$$= g^{ij}\frac{\delta^2 \phi}{\delta x^i \delta x^j} + g^{ab}\hat{e}_b(u^i_a)(\frac{\delta \phi}{\delta x^i}),$$

where, $g_{ab} = g_{ij}u_a^i u_b^j$. Denoting $B^c := v_i^c g^{ab} \hat{e}_b(u_a^i)$ we can rewrite the expression $g^{ij} \frac{\delta^2 \phi}{\delta x^i \delta x^j}$ on SM with respect to \hat{e}_a 's as follows

$$g^{ab}\hat{e}_b\hat{e}_a\phi - B^c\hat{e}_c\phi = g^{ij}\frac{\delta^2\phi}{\delta x^i\delta x^j}.$$
(19)

By definition, we also have

$$\begin{split} \hat{e}_{n+\alpha}\phi &= Fu_{\alpha}^{i}\frac{\partial\phi}{\partial y^{i}},\\ \hat{e}_{n+\beta}\hat{e}_{n+\alpha}\phi &= \hat{e}_{n+\beta}(Fu_{\alpha}^{i}\frac{\partial\phi}{\partial y^{i}})\\ &= F^{2}u_{\beta}^{j}u_{\alpha}^{i}\frac{\partial^{2}\phi}{\partial y^{j}\partial y^{i}} + F(\hat{e}_{n+\beta}u_{\alpha}^{i})(\frac{\partial\phi}{\partial y^{i}}) + Fu_{\beta}^{j}(\frac{\partial F}{\partial y^{j}})u_{\alpha}^{i}\frac{\partial\phi}{\partial y^{i}}. \end{split}$$

Using the fact $u_{\beta}^{j} \frac{\partial F}{\partial y^{j}} = 0$, see [1, p. 161], we have

$$\hat{e}_{n+\beta}\hat{e}_{n+\alpha}\phi = F^2 u^j_\beta u^i_\alpha \frac{\partial^2 \phi}{\partial y^j \partial y^i} + F(\hat{e}_{n+\beta}u^i_\alpha)(\frac{\partial \phi}{\partial y^i}).$$

Multiplying both sides by $g^{\alpha\beta}$ and taking into account $g^{\alpha\beta}u^i_{\alpha}u^j_{\beta}=g^{ij}-y^iy^j$ we get

$$g^{\alpha\beta}\hat{e}_{n+\beta}\hat{e}_{n+\alpha}\phi = F^2 g^{ij} \frac{\partial^2 \phi}{\partial y^j \partial y^i} + F g^{\alpha\beta} (\hat{e}_{n+\beta} u^i_{\alpha}) \frac{\partial \phi}{\partial y^i},$$

where, $g_{\alpha\beta} = g_{ij} u^i_{\alpha} u^j_{\beta}$. If we denote $D^{\gamma} := v^{\gamma}_i F g^{\alpha\beta} \hat{e}_{n+\beta} u^i_{\alpha}$ we can rewrite the expression $F^2 g^{ij} \frac{\partial^2 \phi}{\partial y^j \partial y^i}$ on SM with respect to $\hat{e}_{n+\alpha}$'s as follows

$$g^{\alpha\beta}\hat{e}_{n+\beta}\hat{e}_{n+\alpha}\phi - D^{\gamma}\hat{e}_{n+\gamma}\phi = F^2 g^{ij} \frac{\partial^2 \phi}{\partial y^j \partial y^i}.$$
 (20)

Thus the principal order terms $g^{ij} \frac{\delta^2 \phi}{\delta x^i \delta x^j}$ and $F^2 g^{ij} \frac{\partial^2 \phi}{\partial y^i \partial y^j}$ convert to $g^{ab} \hat{e}_b \hat{e}_a \phi - B^c \hat{e}_c \phi$ and $g^{\alpha\beta} \hat{e}_{n+\beta} \hat{e}_{n+\alpha} \phi - D^{\gamma} \hat{e}_{n+\gamma} \phi$ on SM. On the other hand, the order of the lower order terms in (18) do not change after rewriting them in terms of the basis $\{\hat{e}_a, \hat{e}_{n+\alpha}\}$ on SM. Therefore (18) on SM is written as

$$\frac{\partial \phi}{\partial t} = g^{ab} \hat{e}_b \hat{e}_a \phi + g^{\alpha\beta} \hat{e}_{n+\beta} \hat{e}_{n+\alpha} \phi - B^c \hat{e}_c \phi - D^{\gamma} \hat{e}_{n+\gamma} \phi + \text{lower order terms.}$$
(21)

Using the fact that g is positive definite, the coefficients

$$G = \left(\begin{array}{c|c} g^{ab} & 0\\ \hline 0 & g^{\alpha\beta} \end{array}\right)_{(2n-1)\times(2n-1)}$$

of principal order terms of (21) is positive definite on SM. Therefore, by virtue of (17) the differential equation (21) is a semi-linear strictly parabolic equation on SM.

4 Setting up an operator and a vector field on *SM*

Let (M, F) and (N, \overline{F}) be the two *n*-dimensional Finsler manifolds with the corresponding metric tensors g and h, respectively. We decorate with a bar the geometric objects on N. Let (x^i, y^i) and $(\overline{x}^i, \overline{y}^i)$ be the local coordinate systems on TM and TN, respectively. Considering that c is a geodesic on (M, F), the natural lift of c on TM is given by

$$\tilde{c}: t \in I \longrightarrow \tilde{c}(t) = (x^i(t), \frac{dx^i}{dt}(t)) \in TM.$$

It is well known that \tilde{c} is a *horizontal curve* on TM, that is to say its tangent vector field is a horizontal vector field with the components $\dot{\tilde{c}}(t) = \frac{dx^i}{dt} \frac{\delta}{\delta x^i} \in HTM$. Let us consider a diffeomorphism φ between the tangent bundles TM and TN, such that $\bar{c}(t) := (\varphi \circ \tilde{c})(t)$ is a horizontal curve on TN.

$$\varphi: TM \longrightarrow TN,$$

$$(x^{i}, y^{i}) \mapsto \varphi(x^{i}, y^{i}) = (\varphi^{\alpha}(x^{i}, y^{i})),$$

$$(22)$$

where, the components φ^{α} of φ are $(\varphi^{\alpha}(x^{i}, y^{i})) =: (\varphi^{j}(x^{i}, y^{i}), \varphi^{n+j}(x^{i}, y^{i}))$, and the Latin indices i, j, ... run over the range 1, ..., n and the Greek indice α run over the range 1, ..., 2n. Throughout this section, φ takes a horizontal curve on TM into a horizontal curve on TN.

Denote by Γ^i_{jk} and $\bar{\Gamma}^i_{jk}$ the coefficients of the Chern connection on (M, F) and (N, \bar{F}) , respectively. Clearly

$$\bar{\nabla}_{\dot{c}}\dot{\bar{c}} = \bar{\nabla}_{\dot{c}}\frac{d\bar{x}^j}{dt}\frac{\delta}{\delta\bar{x}^j} = \frac{d^2\bar{x}^j}{dt^2}\frac{\delta}{\delta\bar{x}^j} + \frac{d\bar{x}^j}{dt}\bar{\nabla}_{\dot{c}}\frac{\delta}{\delta\bar{x}^j} = (\frac{d^2\bar{x}^i}{dt^2} + \frac{d\bar{x}^j}{dt}\frac{d\bar{x}^k}{dt}\bar{\Gamma}^i_{jk})\frac{\delta}{\delta\bar{x}^i}, \quad (23)$$

where $\dot{\bar{c}}(t) := \frac{d\bar{x}^j}{dt} \frac{\delta}{\delta \bar{x}^j}$ is the tangent vector to the horizontal curve $\bar{c}(t)$ on TN. On the other hand

$$\frac{d\bar{x}^i}{dt} = \frac{\delta\varphi^i}{\delta x^p} \frac{dx^p}{dt}, \quad \frac{d^2\bar{x}^i}{dt^2} = \frac{\delta^2\varphi^i}{\delta x^p \delta x^q} \frac{dx^p}{dt} \frac{dx^q}{dt} + \frac{\delta\varphi^i}{\delta x^p} \frac{d^2x^p}{dt^2}.$$

Replacing the last equations in (23) leads to

$$\bar{\nabla}_{\dot{c}}\dot{\bar{c}} = \left(\frac{\delta^2\varphi^i}{\delta x^p\delta x^q}\frac{dx^p}{dt}\frac{dx^q}{dt} + \frac{\delta\varphi^i}{\delta x^h}\frac{d^2x^h}{dt^2} + \bar{\Gamma}^i_{jk}\frac{\delta\varphi^j}{\delta x^p}\frac{\delta\varphi^k}{\delta x^q}\frac{dx^p}{dt}\frac{dx^q}{dt}\right)\frac{\delta}{\delta\bar{x}^i},\tag{24}$$

where, all the indices i, j, p, q, \dots , run over the range $1, \dots, n$. The geodesic c on (M, F), satisfies

$$\frac{d^2x^h}{dt^2} + \Gamma^h_{pq} \frac{dx^p}{dt} \frac{dx^q}{dt} = 0.$$
(25)

Substituting $\frac{d^2x^h}{dt^2}$ from the last equation in (24), leads to

$$\bar{\nabla}_{\dot{\bar{c}}}\dot{\bar{c}} = \frac{dx^p}{dt}\frac{dx^q}{dt}\left(\frac{\delta^2\varphi^i}{\delta x^p\delta x^q} - \frac{\delta\varphi^i}{\delta x^h}\Gamma^h_{pq} + \bar{\Gamma}^i_{jk}\frac{\delta\varphi^j}{\delta x^p}\frac{\delta\varphi^k}{\delta x^q}\right)\frac{\delta}{\delta\bar{x}^i}.$$

Next, let

$$A^{i}_{pq} := \frac{\delta^{2}\varphi^{i}}{\delta x^{p}\delta x^{q}} - \frac{\delta\varphi^{i}}{\delta x^{h}}\Gamma^{h}_{pq} + \bar{\Gamma}^{i}_{jk}\frac{\delta\varphi^{j}}{\delta x^{p}}\frac{\delta\varphi^{k}}{\delta x^{q}} + F^{2}\frac{\partial^{2}\varphi^{i}}{\partial y^{p}\partial y^{q}}$$

Contracting A_{pq}^i with g^{pq} leads to the following operators

$$(\Phi_{g,h}\varphi)^{i} := g^{pq} \left(\frac{\delta^{2}\varphi^{i}}{\delta x^{p} \delta x^{q}} + F^{2} \frac{\partial^{2}\varphi^{i}}{\partial y^{p} \partial y^{q}} - \frac{\delta\varphi^{i}}{\delta x^{h}} \Gamma^{h}_{pq} + \bar{\Gamma}^{i}_{jk} \frac{\delta\varphi^{j}}{\delta x^{p}} \frac{\delta\varphi^{k}}{\delta x^{q}}\right),$$
(26)

where, $(\Phi_{g,h}\varphi)^i = g^{pq}A^i_{pq}$ and i = 1, ..., n. For the indices greater than "n" we consider the following operator.

$$(\Phi_{g,h}\varphi)^{n+i} := g^{pq} \left(\frac{\delta^2 \varphi^{n+i}}{\delta x^p \delta x^q} + F^2 \frac{\partial^2 \varphi^{n+i}}{\partial y^p \partial y^q} + \frac{\partial \varphi^{n+i}}{\partial y^k} \frac{\delta N_q^k}{\delta x^p} \right), \tag{27}$$

where, $i = 1, \dots, n$. Summarizing the above definitions we have

$$(\Phi_{g,h}\varphi)^{\alpha} = \begin{cases} (\Phi_{g,h}\varphi)^i & \alpha = i\\ (\Phi_{g,h}\varphi)^{n+i} & \alpha = n+i, \end{cases}$$

where, i = 1, ..., n. Next, we show the operator $(\Phi_{g,h}\varphi)^{\alpha}$ is invariant under all diffeomorphisms on TM. Our approach to defining the above operator on SM and the following key lemma is an extension of that presented in [11].

Lemma 4.1. Let (M, F) and (N, F) be two n-dimensional Finsler manifolds with the corresponding metric tensors g and h, respectively. If ψ is a diffeomorphism from TM to itself, then it leaves invariant the operator $(\Phi_{g,h}\varphi)^{\alpha}$, in the sense that

$$(\Phi_{\psi^*(g),h}\psi^*\varphi)^{\alpha}\mid_{(\tilde{x},\tilde{y})}=(\Phi_{g,h}\varphi)^{\alpha}\mid_{(x,y)}$$

where, $\tilde{x}^i = \psi^* x^i$, $\tilde{y}^i = \psi^* y^i$ and $\alpha = 1, \cdots, 2n$.

Proof. Let (x^i, y^i) and (\bar{x}^i, \bar{y}^i) be the two local coordinate systems on TM and TN, respectively. Letting $\tilde{x}^i = \psi^* x^i$, $\tilde{y}^i = \psi^* y^i$, and considering the diffeomorphism φ given in (22) for $\alpha = i$, we have

$$\begin{split} (\Phi_{g,h}\varphi)^{i}|_{(x,y)} &= g^{pq}(x,y) \Big(\frac{\delta^{2}\varphi^{i}}{\delta x^{p}\delta x^{q}}(x,y) + F^{2}(x,y) \frac{\partial^{2}\varphi^{i}}{\partial y^{p}\partial y^{q}}(x,y) - \frac{\delta\varphi^{i}}{\delta x^{k}}(x,y)\Gamma_{pq}^{k}(x,y) \\ &+ \bar{\Gamma}_{jk}^{i}(\bar{x},\bar{y}) \frac{\delta\varphi^{j}}{\delta x^{p}}(x,y) \frac{\delta\varphi^{k}}{\delta x^{q}}(x,y) \Big). \end{split}$$

Using the diffeomorphism ψ and the pull-back ψ^* we have

$$\begin{split} (\Phi_{g,h}\varphi)^{i} \mid_{(x,y)} &= g^{pq}(\psi(\tilde{x},\tilde{y})) \Big(\frac{\delta^{2}\varphi^{i}}{\delta x^{p}\delta x^{q}}(\psi(\tilde{x},\tilde{y})) + F^{2}(\psi(\tilde{x},\tilde{y})) \frac{\partial^{2}\varphi^{i}}{\partial y^{p}\partial y^{q}}(\psi(\tilde{x},\tilde{y})) \\ &\quad - \frac{\delta\varphi^{i}}{\delta x^{k}}(\psi(\tilde{x},\tilde{y}))\Gamma^{k}_{pq}(\psi(\tilde{x},\tilde{y})) + \bar{\Gamma}^{i}_{jk}(\bar{x},\bar{y}) \frac{\delta\varphi^{j}}{\delta x^{p}}(\psi(\tilde{x},\tilde{y})) \frac{\delta\varphi^{k}}{\delta x^{q}}(\psi(\tilde{x},\tilde{y})) \Big) \\ &= (\psi^{*}g)^{pq}(\tilde{x},\tilde{y}) \Big(\frac{\delta^{2}(\psi^{*}\varphi)^{i}}{\delta \tilde{x}^{p}\delta \tilde{x}^{q}}(\tilde{x},\tilde{y}) + (\psi^{*}F^{2})(\tilde{x},\tilde{y}) \frac{\partial^{2}(\psi^{*}\varphi)^{i}}{\partial \tilde{y}^{p}\partial \tilde{y}^{q}}(\tilde{x},\tilde{y}) \\ &\quad - \frac{\delta(\psi^{*}\varphi)^{i}}{\delta \tilde{x}^{k}}(\tilde{x},\tilde{y})\Gamma(\psi^{*}g)^{k}_{pq}(\tilde{x},\tilde{y}) + \bar{\Gamma}^{i}_{jk}(\bar{x},\bar{y}) \frac{\delta(\psi^{*}\varphi)^{j}}{\delta \tilde{x}^{p}}(\tilde{x},\tilde{y}) \frac{\delta(\psi^{*}\varphi)^{k}}{\delta \tilde{x}^{q}}(\tilde{x},\tilde{y}) \Big) \\ &= (\Phi_{\psi^{*}(g),h}\psi^{*}\varphi)^{i} \mid_{(\tilde{x},\tilde{y})} . \end{split}$$

Similarly, for $\alpha = n + i$, one can show that

$$(\Phi_{g,h}\varphi)^{n+i}\mid_{(x,y)}=(\Phi_{\psi^*(g),h}\psi^*\varphi)^{n+i}\mid_{(\tilde{x},\tilde{y})},$$

where, $i = 1, \dots, n$. This completes the proof.

Remark 4.1. Let (M, F) and (N, \overline{F}) be two n-dimensional Finsler manifolds with the corresponding metric tensors g and h, respectively. Let $\varphi : TM \longrightarrow TN$, $\varphi(x^i, y^i) = (\varphi^{\alpha}(x^i, y^i)), \alpha = 1, ..., 2n$, be a diffeomorphism that takes a horizontal curve to a horizontal curve. Given $\varphi_0 : TM \longrightarrow TN$, we consider the following evolution equation

$$\frac{\partial}{\partial t}\varphi^{\alpha} = (\Phi_{g,h}\varphi)^{\alpha}, \qquad \varphi_{(0)} = \varphi_0.$$
(28)

By restricting φ^{α} 's to SM and using Lemma 3.1, one can see that (28) is a strictly parabolic system. Hence, there is a unique solution for (28) in a short-time.

Corollary 4.1. Let (M, \tilde{F}) and (N, \bar{F}) be two n-dimensional Finsler manifolds with corresponding metric tensors \tilde{g} and h, respectively. Let N = M and φ be the identity map $\varphi = Id: TM \longrightarrow TM, \ \varphi(x^i, y^i) = (x^i, y^i)$, then we have

$$(\Phi_{\tilde{g},h}Id)^{\alpha} = (\Phi_{\tilde{g},h}Id)^{i} = \tilde{g}^{pq}(-\tilde{\Gamma}^{i}_{pq} + \bar{\Gamma}^{i}_{pq}), \quad \alpha = i$$

where, $i = 1, \dots, n$ and $\tilde{\Gamma}^{i}_{pq}, \bar{\Gamma}^{i}_{pq}$ are the coefficients of horizontal covariant derivatives of the Chern connection with respect to the metrics \tilde{g} and h, respectively.

Let ξ be a vector field on TM with the components

$$\xi := (\Phi_{\tilde{g},h}Id)^i \frac{\partial}{\partial x^i} = \tilde{g}^{pq} (-\tilde{\Gamma}^i_{pq} + \bar{\Gamma}^i_{pq}) \frac{\partial}{\partial x^i}.$$
(29)

Using the fact that the difference between any two connections is a tensor field, ξ is a globally well-defined vector field. It is easy to verify that the components of ξ are homogeneous of degree zero on y. Thus ξ can be considered as a vector field on SM.

5 Ricci-DeTurck flow on Finsler manifolds

Here we study the Ricci-DeTurck flow and its existence of solution. More intuitively, there are several well-known definitions of the Ricci tensor in Finsler geometry. For instance, H. Akbar-Zadeh has considered two Ricci tensors on Finsler manifolds in his works, namely, one is defined by $Ric_{ij} := [\frac{1}{2}F^2\mathcal{R}ic]_{y^iy^j}$ where, $\mathcal{R}ic$ is the *Ricci scalar* defined by $\mathcal{R}ic := g^{ik}R_{ik} = R^i_i$ and R^i_k are defined by (4), see [3, p. 192]. Another Ricci tensor is defined by $Rc_{ij} := \frac{1}{2}(\mathsf{R}_{ij} + \mathsf{R}_{ji})$, where R_{ij} is the trace of *hh*-curvature defined by $\mathsf{R}_{ij} = R^i_{i\,lj}$. The difference between these two Ricci tensors is the additional term $\frac{1}{2}y^k \frac{\partial \mathsf{R}_{jk}}{\partial y^i}$ that appeared in the first definition. More precisely, we have $Ric_{ij} - Rc_{ij} = \frac{1}{2}y^k \frac{\partial \mathsf{R}_{jk}}{\partial y^i}$.

In [2], D. Bao, based on the first definition of the Ricci tensor, considered the following evolutionary differential equation as a *Ricci flow* equation in Finsler geometry:

$$\frac{\partial}{\partial t}g_{jk}(t) = -2Ric_{jk}, \qquad g_{(t=0)} = g_0, \tag{30}$$

where, $g_{jk}(t)$ are a family of Finslerian metrics defined on $\pi^*TM \times [0, T)$. Contracting (30) with $y^j y^k$, via Euler's theorem, leads to $\frac{\partial}{\partial t}F^2 = -2F^2\mathcal{R}ic$. That is,

$$\frac{\partial}{\partial t}\log F(t) = -\mathcal{R}ic, \qquad F(t=0) := F_0,$$
(31)

where, F_0 is the initial Finsler structure, see more details on [2]. Here and everywhere in the present work we consider the first Akbar-Zadeh's definition of Ricci tensor and the related Ricci flow (31). One of the advantages of the Ricci quantity Ric_{ij} , used in the present work is its independence with respect to the choice of Cartan, Berwald or Chern connections.

Definition 5.1. Let M be a compact manifold with a fixed background Finsler structure \overline{F} and the related Finsler metric h. Assume that for all $t \in [0, T)$, $\widetilde{F}(t)$ is a one-parameter family of Finsler structures defined on TM and $\widetilde{g}(t)$ is the tensor metric related to $\widetilde{F}(t)$. We say that $\widetilde{F}(t)$ is a solution to the Finslerian Ricci-DeTurck flow if

$$\frac{\partial}{\partial t}\tilde{F}^{2}(t) = -2\tilde{F}^{2}(t)\mathcal{R}ic(\tilde{g}(t)) - \mathcal{L}_{\xi}\tilde{F}^{2}(t), \qquad (32)$$

where, \mathcal{L}_{ξ} is the Lie derivative with respect to the vector field $\xi = (\Phi_{\tilde{g}(t),h}Id)^i \frac{\partial}{\partial x^i}$ on SM as mentioned previously in (29).

The following theorem shows that the Ricci-DeTurck flow (32) is well defined and has a solution in a short-time interval.

Proof of Theorem 1. Let M be a compact manifold with a fixed background Finsler structure \overline{F} and the related Finsler metric h. Here, all the indices run over the range 1, ..., n. The Ricci-DeTurck flow (32) can be written in the following form

$$y^{p}y^{q}\frac{\partial}{\partial t}\tilde{g}_{pq}(t) = -2\tilde{F}^{2}(t)\mathcal{R}ic(\tilde{g}(t)) - \mathcal{L}_{\xi}y^{p}y^{q}\tilde{g}_{pq}(t),$$
(33)

where, $\tilde{g}(t)$ is the metric tensor related to $\tilde{F}(t)$. Also we have

$$\mathcal{L}_{\xi}(y^{p}y^{q}\tilde{g}_{pq}) = y^{p}y^{q}\mathcal{L}_{\xi}\tilde{g}_{pq} + 2y^{p}\tilde{g}_{pq}\mathcal{L}_{\xi}y^{q}.$$

Therefore, (33) becomes

$$y^{p}y^{q}\frac{\partial}{\partial t}\tilde{g}_{pq}(t) = -2\tilde{F}^{2}(t)\mathcal{R}ic(\tilde{g}(t)) - y^{p}y^{q}\mathcal{L}_{\xi}\tilde{g}_{pq} - 2y^{p}\tilde{g}_{pq}\mathcal{L}_{\xi}y^{q}.$$
(34)

By means of the Lie derivative formula (9) along ξ we have

$$y^p y^q \mathcal{L}_{\xi} \tilde{g}_{pq} = 2y^p y^q \nabla_p \xi_q, \tag{35}$$

where, ∇_p is the horizontal covariant derivative in Chern connection. The *h*-metric compatibility of the Chern connection, yields

$$\nabla_p \xi_q = (\nabla_p \tilde{g}_{ql} \xi^l) = \tilde{g}_{ql} (\nabla_p \xi^l).$$

As mentioned earlier, if we denote the coefficients of horizontal covariant derivatives of Chern connection with respect to the metric tensors h and \tilde{g} by $\Gamma(h)$ and $\Gamma(\tilde{g})$ respectively, then by the definition of ξ given in (29), the above equation becomes

$$\begin{aligned} \nabla_p \xi_q &= \tilde{g}_{ql} (\delta_p \xi^l + \Gamma(\tilde{g})^l_{pw} \xi^w) \\ &= \tilde{g}_{ql} [\delta_p (\tilde{g}^{mn} (\Gamma(h)^l_{mn} - \Gamma(\tilde{g})^l_{mn}))] + \tilde{g}_{ql} \Gamma(\tilde{g})^l_{pw} \xi^w \\ &= \tilde{g}_{ql} [(\delta_p \tilde{g}^{mn}) (\Gamma(h)^l_{mn} - \Gamma(\tilde{g})^l_{mn}) + \tilde{g}^{mn} \delta_p (\Gamma(h)^l_{mn}) - \tilde{g}^{mn} \delta_p (\Gamma(\tilde{g})^l_{mn})] \\ &+ \tilde{g}_{ql} \Gamma(\tilde{g})^l_{pw} \xi^w. \end{aligned}$$

According to the horizontal derivative of $\Gamma(\tilde{g})_{mn}^l$ in the last equation, we get

$$\begin{aligned} \nabla_p \xi_q &= \frac{1}{2} \tilde{g}^{mn} (\delta_p \delta_q \tilde{g}_{mn} - \delta_p \delta_n \tilde{g}_{qm} - \delta_p \delta_m \tilde{g}_{qn}) - \frac{1}{2} \tilde{g}_{ql} \tilde{g}^{mn} (\delta_p \tilde{g}^{ls}) (\delta_n \tilde{g}_{sm} - \delta_s \tilde{g}_{mn} + \delta_m \tilde{g}_{ns}) \\ &+ \tilde{g}_{ql} (\delta_p \tilde{g}^{mn}) (\Gamma(h)_{mn}^l - \Gamma(\tilde{g})_{mn}^l) + \tilde{g}_{ql} \tilde{g}^{mn} \delta_p (\Gamma(h)_{mn}^l) + \tilde{g}_{ql} \Gamma(\tilde{g})_{pw}^l \xi^w. \end{aligned}$$

Using the last equation, (35) is written

$$y^{p}y^{q}\mathcal{L}_{\xi}\tilde{g}_{pq} = y^{p}y^{q}\tilde{g}^{mn}(\delta_{p}\delta_{q}\tilde{g}_{mn} - \delta_{p}\delta_{n}\tilde{g}_{qm} - \delta_{p}\delta_{m}\tilde{g}_{qn}) - y^{p}y^{q}\tilde{g}_{ql}\tilde{g}^{mn}(\delta_{p}\tilde{g}^{ls})(\delta_{n}\tilde{g}_{sm} - \delta_{s}\tilde{g}_{mn} + \delta_{m}\tilde{g}_{ns}) + 2y^{p}y^{q}\tilde{g}_{ql}(\delta_{p}\tilde{g}^{mn})(\Gamma(h)^{l}_{mn} - \Gamma(\tilde{g})^{l}_{mn}) + 2y^{p}y^{q}\tilde{g}_{ql}\tilde{g}^{mn}\delta_{p}(\Gamma(h)^{l}_{mn}) + 2y^{p}y^{q}\tilde{g}_{ql}\Gamma(\tilde{g})^{l}_{pw}\xi^{w}.$$
(36)

Also we have

$$-2\tilde{F}^{2}\mathcal{R}ic(\tilde{g}) = -2\tilde{F}^{2}R^{n}_{\ n} = -2\tilde{F}^{2}l^{q}R^{\ n}_{q\ np}l^{p},\tag{37}$$

where, $R_{q np}^{n}$ are the components of the *hh*-curvature tensor of Chern connection and $l^{q} = \frac{y^{q}}{F}$ are the components of the Liouville vector field. Replacing (3) in (37) and using the definition of $\Gamma(\tilde{g})$, yields

$$\begin{split} -2\tilde{F}^2 \mathcal{R}ic(\tilde{g}) &= -2\tilde{F}^2 l^q R_{q\ np}^n l^p \\ &= -2y^p y^q (\delta_n \Gamma_{qp}^n(\tilde{g}) - \delta_p \Gamma_{qn}^n(\tilde{g}) + \Gamma_{mn}^n(\tilde{g}) \Gamma_{qp}^m(\tilde{g}) - \Gamma_{mp}^n(\tilde{g}) \Gamma_{qn}^m(\tilde{g})) \\ &= -2y^p y^q [\delta_n (\frac{1}{2} \tilde{g}^{mn} (\delta_q \tilde{g}_{mp} + \delta_p \tilde{g}_{mq} - \delta_m \tilde{g}_{pq}))] \\ &+ 2y^p y^q [\delta_p (\frac{1}{2} \tilde{g}^{mn} (\delta_q \tilde{g}_{mn} + \delta_n \tilde{g}_{qm} - \delta_m \tilde{g}_{qn}))] \\ &- 2y^p y^q (\Gamma_{mn}^n(\tilde{g}) \Gamma_{qp}^m(\tilde{g}) - \Gamma_{mp}^n(\tilde{g}) \Gamma_{qn}^m(\tilde{g})). \end{split}$$

By applying the δ_p derivative we have

$$-2\tilde{F}^{2}\mathcal{R}ic(\tilde{g}) = y^{p}y^{q}\tilde{g}^{mn}(\delta_{p}\delta_{q}\tilde{g}_{mn} + \delta_{n}\delta_{m}\tilde{g}_{pq} - \delta_{p}\delta_{m}\tilde{g}_{qn} - \delta_{n}\delta_{q}\tilde{g}_{mp}) - y^{p}y^{q}(\delta_{n}\tilde{g}^{nm})(\delta_{q}\tilde{g}_{mp} + \delta_{p}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{pq}) + y^{p}y^{q}(\delta_{p}\tilde{g}^{mn})(\delta_{q}\tilde{g}_{mn} + \delta_{n}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{qn}) - 2y^{p}y^{q}(\Gamma_{mn}^{n}(\tilde{g})\Gamma_{qp}^{m}(\tilde{g}) - \Gamma_{mp}^{n}(\tilde{g})\Gamma_{qn}^{m}(\tilde{g})).$$
(38)

Substituting (36) and (38) in (34), we obtain

$$y^{p}y^{q}\frac{\partial}{\partial t}\tilde{g}_{pq}(t) = y^{p}y^{q}\tilde{g}^{mn}\delta_{n}\delta_{m}\tilde{g}_{pq}$$

$$(39)$$

$$- y^{p}y^{q}(\delta_{n}\tilde{g}^{nm})(\delta_{q}\tilde{g}_{mp} + \delta_{p}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{pq})$$

$$+ y^{p}y^{q}(\delta_{p}\tilde{g}^{mn})(\delta_{q}\tilde{g}_{mn} + \delta_{n}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{qn})$$

$$- 2y^{p}y^{q}(\Gamma_{mn}^{n}(\tilde{g})\Gamma_{qp}^{m}(\tilde{g}) - \Gamma_{mp}^{n}(\tilde{g})\Gamma_{qn}^{m}(\tilde{g}))$$

$$+ y^{p}y^{q}\tilde{g}_{ql}\tilde{g}^{mn}(\delta_{p}\tilde{g}^{ls})(\delta_{n}\tilde{g}_{sm} - \delta_{s}\tilde{g}_{mn} + \delta_{m}\tilde{g}_{ns})$$

$$- 2y^{p}y^{q}\tilde{g}_{ql}(\delta_{p}\tilde{g}^{mn})(\Gamma(h)_{mn}^{l} - \Gamma(\tilde{g})_{mn}^{l})$$

$$- 2y^{p}y^{q}\tilde{g}_{ql}\tilde{g}^{mn}\delta_{p}(\Gamma(h)_{mn}^{l}) - 2y^{p}y^{q}\tilde{g}_{ql}\Gamma(\tilde{g})_{pw}^{l}\xi^{w}$$

$$- 2y^{p}\tilde{g}_{pq}\mathcal{L}_{\xi}y^{q}.$$

Using Euler's theorem yields

$$y^{p}y^{q}\frac{\partial^{2}\tilde{g}_{pq}}{\partial y^{n}\partial y^{m}} = \frac{\partial^{2}}{\partial y^{n}\partial y^{m}}(y^{p}y^{q}\tilde{g}_{pq}) - 2\tilde{g}_{nm} = 0.$$

$$\tag{40}$$

In order to get a strictly parabolic system, by virtue of (40) we add the zero term $\tilde{F}^2 y^p y^q \tilde{g}^{mn} \frac{\partial^2 \tilde{g}_{pq}}{\partial y^n \partial y^m} = 0$ to the right hand side of (39). Therefore, we have

$$y^{p}y^{q}\left(\frac{\partial}{\partial t}\tilde{g}_{pq}(t) - \tilde{g}^{mn}\delta_{n}\delta_{m}\tilde{g}_{pq} - \tilde{F}^{2}\tilde{g}^{mn}\frac{\partial^{2}\tilde{g}_{pq}}{\partial y^{n}\partial y^{m}} + (\delta_{n}\tilde{g}^{nm})(\delta_{q}\tilde{g}_{mp} + \delta_{p}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{pq}) - (\delta_{p}\tilde{g}^{mn})(\delta_{q}\tilde{g}_{mn} + \delta_{n}\tilde{g}_{qm} - \delta_{m}\tilde{g}_{qn}) + 2(\Gamma_{mn}^{n}(\tilde{g})\Gamma_{qp}^{m}(\tilde{g}) - \Gamma_{mp}^{n}(\tilde{g})\Gamma_{qn}^{m}(\tilde{g})) - \tilde{g}_{ql}\tilde{g}^{mn}(\delta_{p}\tilde{g}^{ls})(\delta_{n}\tilde{g}_{sm} - \delta_{s}\tilde{g}_{mn} + \delta_{m}\tilde{g}_{ns}) + 2\tilde{g}_{ql}(\delta_{p}\tilde{g}^{mn})(\Gamma(h)_{mn}^{l} - \Gamma(\tilde{g})_{mn}^{l}) + 2\tilde{g}_{ql}\tilde{g}^{mn}\delta_{p}(\Gamma(h)_{mn}^{l}) - 2y^{p}y^{q}\tilde{g}_{ql}\Gamma(\tilde{g})_{pw}^{l}\xi^{w} + 2\frac{l_{q}}{F}\tilde{g}_{np}\mathcal{L}_{\xi}y^{n}\right) = 0.$$

On the other hand, applying twice the vector field $\frac{\delta}{\delta x^n}$ on the components of metric tensor

 \tilde{g}_{pq} yields

$$\begin{split} \delta_n \delta_m \tilde{g}_{pq} = & \frac{\partial^2 \tilde{g}_{pq}}{\partial x^n \partial x^m} - \frac{\partial N_m^k}{\partial x^n} \frac{\partial \tilde{g}_{pq}}{\partial y^k} - N_m^k \frac{\partial^2 \tilde{g}_{pq}}{\partial x^n \partial y^k} - N_n^l \frac{\partial^2 \tilde{g}_{pq}}{\partial y^l \partial x^m} \\ &+ N_n^l \frac{\partial N_m^k}{\partial y^l} \frac{\tilde{g}_{pq}}{\partial y^k} + N_n^l N_m^k \frac{\partial^2 \tilde{g}_{pq}}{\partial y^k \partial y^l}. \end{split}$$

Convecting the last equation with $y^p y^q$ and using (40) we have

$$y^p y^q \delta_n \delta_m \tilde{g}_{pq} = y^p y^q \frac{\partial^2 \tilde{g}_{pq}}{\partial x^n \partial x^m}$$

Remark that, in the term $y^p y^q \tilde{g}^{mn} \delta_n \delta_m \tilde{g}_{pq}$ in (39), there is no term containing derivatives of \tilde{g} except $y^p y^q \frac{\partial^2 \tilde{g}_{pq}}{\partial x^n \partial x^m}$. One can rewrite (41) as follows

$$y^{p}y^{q}\left(\frac{\partial}{\partial t}\tilde{g}_{pq}(t) - \tilde{g}^{mn}\delta_{n}\delta_{m}\tilde{g}_{pq} - \tilde{F}^{2}\tilde{g}^{mn}\frac{\partial^{2}\tilde{g}_{pq}}{\partial y^{n}\partial y^{m}} + \text{lower order terms}\right) = 0.$$
(42)

Let us set the term between the parentheses in (42) equal to zero, that is,

$$\frac{\partial}{\partial t}\tilde{g}_{pq}(t) - \tilde{g}^{mn}\delta_n\delta_m\tilde{g}_{pq} - \tilde{F}^2\tilde{g}^{mn}\frac{\partial^2\tilde{g}_{pq}}{\partial y^n\partial y^m} + \text{lower order terms} = 0$$
(43)

By restricting the metric tensor \tilde{g} on p^*TM and using Lemma 3.1 we can rewrite (43) in terms of the basis $\{\hat{e}_a, \hat{e}_{n+\alpha}\}$ on SM as follows

$$\frac{\partial}{\partial t}\tilde{g}_{pq} = \tilde{g}^{ab}\hat{e}_b\hat{e}_a\tilde{g}_{pq} + \tilde{g}^{\alpha\beta}\hat{e}_{n+\beta}\hat{e}_{n+\alpha}\tilde{g}_{pq} - B^c\hat{e}_c\tilde{g}_{pq} - D^\gamma\hat{e}_{n+\gamma}\tilde{g}_{pq} + \text{lower order terms} = 0, \qquad (44)$$

where, $B^c := v_i^c \tilde{g}^{ab} \hat{e}_b(u_a^i)$ and $D^{\gamma} := v_i^{\gamma} \tilde{F} \tilde{g}^{\alpha\beta} \hat{e}_{n+\beta} u_{\alpha}^i$ as mentioned in Lemma 3.1.

By hypothesis, M is compact and so is the sphere bundle SM. It is well known that the metric tensor \tilde{g}^{mn} remains positive definite along the Ricci flow, see [9], Corollary 3.7. Since the coefficients of the principal (second) order terms of (44) are positive definite, by Definition 3.1, it is a strictly parabolic semi-linear system on SM. Therefore, the standard existence and uniqueness theorem for parabolic systems on compact domains implies that, (44) has a unique solution on SM. The equation (44) is a special case of the general flow equation (10) and $\tilde{g}(t)$ is a solution. Therefore, by means of Lemma 2.1, $\tilde{g}(t)$ satisfies the integrability condition or equivalently, there exists a Finsler structure $\tilde{F}(t)$ on TMsuch that $\tilde{g}_{ij} = \frac{1}{2} \frac{\partial^2 \tilde{F}}{\partial y^i \partial y^j}$. Hence, \tilde{g} is a Finsler metric and determines a Finsler structure $\tilde{F}^2 := \tilde{g}_{pq} y^p y^p$ which is a solution to the Finsler Ricci-DeTurck flow. This completes the proof of Theorem 1.

Remark 5.1. In Theorem 1, as it is mentioned earlier in the preliminaries, if the Finsler manifold M is isotropic, then $R_{q np}^n$ is symmetric with respect to the indices p and q.

Therefore, by means of symmetry of $\mathcal{L}_{\xi}(y^p y^q \tilde{g}_{pq})$ on the indices p and q, we conclude that the term in the brackets in (42) is symmetric on the indices p and q. If a symmetric bilinear form vanishes on the diagonal, then by the polarization identity it vanishes identically. Hence, according to (42) we have

$$\frac{\partial}{\partial t}\tilde{g}_{pq}(t) - \tilde{g}^{mn}\delta_n\delta_m\tilde{g}_{pq} - \tilde{F}^2\tilde{g}^{mn}\frac{\partial^2\tilde{g}_{pq}}{\partial y^n\partial y^m} + lower \ order \ terms = 0.$$
(45)

As shown in the proof of Theorem 1, by restricting the metric tensor \tilde{g} on p^*TM and using Lemma 3.1 we can rewrite (43) on SM as (44) which is a strictly parabolic semi-linear system on SM. Therefore, the standard existence and uniqueness theorem for parabolic systems on compact domains implies that, (44) has a unique solution on SM. Hence, Finsler structure $\tilde{F}^2 := \tilde{g}_{pq} y^p y^p$ is a unique solution to the Finsler Ricci-DeTurck flow. Therefore, in the isotropic case, the solution to the (32) is unique.

6 Short-time solution to the Ricci flow on Finsler manifolds

In this section, we will prove that there is a one-to-one correspondence between the solutions of the Ricci flow and the Ricci-DeTurck flow on Finsler manifolds. Here, we recall some results which will be used in the sequel.

Lemma A. [12, p. 82] Let $\{X_t : 0 \le t < T \le \infty\}$ be a continuous time-dependent family of vector fields on a compact manifold M. There exists a one-parameter family of diffeomorphisms $\{\varphi_t : M \longrightarrow M; 0 \le t < T \le \infty\}$ defined on the same time interval such that

$$\begin{cases} \frac{\partial}{\partial t}\varphi_t(x) = X_t[\varphi_t(x)]\\ \varphi_0(x) = x, \end{cases}$$

for all $x \in M$ and $t \in [0, T)$.

Remark 6.1. Let M be a compact Finsler manifold. According to Lemma A, there exists a unique one-parameter family of diffeomorphisms $\tilde{\varphi}_t$ on SM, such that

$$\begin{cases} \frac{\partial}{\partial t}\varphi_t(z) = \xi(\varphi_t(z), t), \\ \varphi_0 = Id_{SM}, \end{cases}$$

where, $z = (x, [y]) \in SM$ and $t \in [0, T)$.

Remark 6.2. Let \tilde{g}_{pq} be a solution to the Ricci-DeTurck flow and φ_t the one-parameter global group of diffeomorphisms generating the vector field ξ . Since ξ is a vector field on

SM, then φ_t are homogeneous of degree zero. The zero-homogeneity of \tilde{g}_{pq} implies that $\varphi_t^*(\tilde{g}_{pq})$ is also homogeneous of degree zero. That is,

$$(\varphi_t^* \tilde{g}_{pq})(x, \lambda y) = \tilde{g}_{pq}(\varphi_t(x, \lambda y)) = \tilde{g}_{pq}(\varphi_t(x, y)) = (\varphi_t^* \tilde{g}_{pq})(x, y).$$

Using the fact that \tilde{g}_{pq} is positive definite and φ_t^* are diffeomorphisms, $\varphi_t^*(\tilde{g}_{pq})$ is also positive definite. As well $\varphi_t^*(\tilde{g}_{pq})$ is symmetric. More intuitively,

$$(\varphi_t^*\tilde{g})(X,Y)=g(\varphi_{t_*}(X),\varphi_{t_*}(Y))=g(\varphi_{t_*}(Y),\varphi_{t_*}(X))=(\varphi_t^*\tilde{g})(Y,X).$$

Therefore, $\varphi_t^*(\tilde{g}_{pq})$ determines a Finsler structure as follows

$$F^2 := g_{pq} \tilde{y}^p \tilde{y}^q,$$

where, $g_{pq} := \varphi_t^*(\tilde{g}_{pq})$ and $\varphi_t^* y^p := \tilde{y}^p$.

Lemma 6.1. Let φ_t be a global one-parameter group of diffeomorphisms corresponding to the vector field ξ and $(\gamma_{jk}^i)_{\tilde{g}}$ and $(G^i)_{\tilde{g}}$ are the Christoffel symbols and spray coefficients related to the Finsler metric \tilde{g} , respectively. Then we have

$$\varphi_t^*((\gamma_{jk}^i)_{\tilde{g}}) = (\gamma_{jk}^i)_{\varphi_t^*(\tilde{g})},\tag{46}$$

$$\varphi_t^*(G_{\tilde{g}}^i) = G_{\varphi_t^*(\tilde{g})}^i, \tag{47}$$

where, $(\gamma_{jk}^i)_{\tilde{g}} = \tilde{g}^{is} \frac{1}{2} (\frac{\partial \tilde{g}_{sj}}{\partial x^k} - \frac{\partial \tilde{g}_{jk}}{\partial x^s} + \frac{\partial \tilde{g}_{ks}}{\partial x^j})$ and $G^i_{\tilde{g}} = \frac{1}{2} (\gamma^i_{jk})_{\tilde{g}} y^j y^k$.

Proof. Let us denote $\varphi_t^* x^i := \tilde{x}^i$ and $\varphi_t^* y^i := \tilde{y}^i$. By definition, we have

$$\begin{aligned} \varphi_t^*((\gamma_{jk}^i)_{\tilde{g}}) &= \varphi_t^*(\tilde{g}^{is}\frac{1}{2}(\frac{\partial \tilde{g}_{sj}}{\partial x^k} - \frac{\partial \tilde{g}_{jk}}{\partial x^s} + \frac{\partial \tilde{g}_{ks}}{\partial x^j})) \\ &= \varphi_t^*(\tilde{g}^{is})\varphi_t^*(\frac{1}{2}(\frac{\partial \tilde{g}_{sj}}{\partial x^k} - \frac{\partial \tilde{g}_{jk}}{\partial x^s} + \frac{\partial \tilde{g}_{ks}}{\partial x^j})) \\ &= \varphi_t^*(\tilde{g}^{is})\frac{1}{2}(\frac{\partial \varphi_t^*(\tilde{g}_{sj})}{\partial \tilde{x}^k} - \frac{\partial \varphi_t^*(\tilde{g}_{jk})}{\partial \tilde{x}^s} + \frac{\partial \varphi_t^*(\tilde{g}_{ks})}{\partial \tilde{x}^j})) \\ &= (\gamma_{ik}^i)_{\varphi_t^*(\tilde{g})}. \end{aligned}$$

Next, by means of (46) we have

$$\begin{split} \varphi_t^*(G_{\tilde{g}}^i) &= \varphi_t^*(\frac{1}{2}(\gamma_{jk}^i)_{\tilde{g}}y^jy^k) = \frac{1}{2}\varphi_t^*((\gamma_{jk}^i)_{\tilde{g}})\varphi_t^*y^j\varphi_t^*y^k \\ &= \frac{1}{2}(\gamma_{jk}^i)_{\varphi_t^*(\tilde{g})}\tilde{y}^j\tilde{y}^k = G_{\varphi_t^*(\tilde{g})}^i. \end{split}$$

This completes the proof.

Lemma 6.2. Let φ_t be a global one-parameter group of diffeomorphisms generating the vector field ξ and $\mathcal{R}ic_{\tilde{g}}$ the Ricci scalar related to the Finsler metric \tilde{g} , then we have

$$\varphi_t^*(\mathcal{R}ic_{\tilde{g}}) = \mathcal{R}ic_{\varphi_t^*(\tilde{g})}$$

Proof. Let us consider the *reduced hh-curvature tensor* R^i_k which is expressed entirely in terms of the x and y derivatives of the spray coefficients $G^i_{\bar{a}}$.

$$(R^i_{\ k})_{\tilde{g}} := \frac{1}{\tilde{F}^2} (2\frac{\partial G^i_{\tilde{g}}}{\partial x^k} - \frac{\partial^2 G^i_{\tilde{g}}}{\partial x^j \partial y^k} y^j + 2G^j_{\tilde{g}} \frac{\partial^2 G^i_{\tilde{g}}}{\partial y^j \partial y^k} - \frac{\partial G^i_{\tilde{g}}}{\partial y^j} \frac{\partial G^j_{\tilde{g}}}{\partial y^k}).$$

Therefore, we have

$$\begin{split} \varphi_t^*((R^i{}_k)_{\tilde{g}}) &= \varphi_t^*(\frac{1}{\tilde{F}^2}(2\frac{\partial G^i_{\tilde{g}}}{\partial x^k} - \frac{\partial^2 G^i_{\tilde{g}}}{\partial x^j \partial y^k}y^j + 2G^j_{\tilde{g}}\frac{\partial^2 G^i_{\tilde{g}}}{\partial y^j \partial y^k} - \frac{\partial G^i_{\tilde{g}}}{\partial y^j}\frac{\partial G^j_{\tilde{g}}}{\partial y^k})) \\ &= \varphi_t^*(\frac{1}{\tilde{F}^2})\varphi_t^*(2\frac{\partial G^i_{\tilde{g}}}{\partial x^k} - \frac{\partial^2 G^i_{\tilde{g}}}{\partial x^j \partial y^k}y^j + 2G^j_{\tilde{g}}\frac{\partial^2 G^i_{\tilde{g}}}{\partial y^j \partial y^k} - \frac{\partial G^i_{\tilde{g}}}{\partial y^j}\frac{\partial G^j_{\tilde{g}}}{\partial y^j}) \end{split}$$

Thus, we get

$$\begin{split} \varphi_t^*((R^i_k)_{\tilde{g}}) = & \frac{1}{\varphi_t^*(\tilde{F}^2)} (2 \frac{\partial(\varphi_t^*(G^i_{\tilde{g}}))}{\partial \tilde{x}^k} - \frac{\partial^2(\varphi_t^*(G^i_{\tilde{g}}))}{\partial \tilde{x}^j \partial \tilde{y}^k} \tilde{y}^j \\ &+ 2\varphi_t^*(G^j_{\tilde{g}}) \frac{\partial^2(\varphi_t^*(G^i_{\tilde{g}}))}{\partial \tilde{y}^j \partial \tilde{y}^k} - \frac{\partial(\varphi_t^*(G^i_{\tilde{g}}))}{\partial \tilde{y}^j} \frac{\partial(\varphi_t^*(G^j_{\tilde{g}}))}{\partial \tilde{y}^k}) \end{split}$$

Letting i = k in the last equation and using (47) implies

$$\varphi_t^*(\mathcal{R}ic_{\tilde{g}}) = \mathcal{R}ic_{\varphi_t^*(\tilde{g})},$$

as we have claimed.

Now we are in a position to prove the following proposition.

Proposition 6.1. Fix a compact Finsler manifold (M, \overline{F}) with the related Finsler metric tensor h. Let $\widetilde{F}(t)$ be a family of solutions to the Ricci-DeTurck flow

$$\frac{\partial}{\partial t}\tilde{F}^2(t) = -2\tilde{F}^2(t)\mathcal{R}ic(\tilde{g}(t)) - \mathcal{L}_{\xi}\tilde{F}^2(t), \qquad (48)$$

where, $\xi = (\Phi_{\tilde{g}(t),h}Id)^i \frac{\partial}{\partial x^i}$ and $t \in [0,T)$. Moreover, let φ_t be a one-parameter family of diffeomorphisms satisfying

$$\frac{\partial}{\partial t}\varphi_t(z) = \xi(\varphi_t(z), t),$$

for $z \in SM$ and $t \in [0,T)$. Then the Finsler structures F(t) form a solution to the Finslerian Ricci flow (31) where, F(t) is defined by

$$F^2(t) := g_{pq} \tilde{y}^p \tilde{y}^q = \varphi_t^*(\tilde{F}^2(t)),$$

where, $g_{pq} := \varphi_t^*(\tilde{g}_{pq})$ and $\varphi_t^* y^p := \tilde{y}^p$.

Proof. In order to show that F(t) forms a solution to the Finslerian Ricci flow (31) we need to show that $\frac{\partial}{\partial t}(\log F(t)) = -\mathcal{R}ic$. Derivation of $F^2(t) = \varphi_t^*(\tilde{F}^2(t))$ with respect to the parameter t, leads to

$$\frac{\partial}{\partial t}(\log F(t)) = \frac{1}{2} \frac{\frac{\partial}{\partial t}(\varphi_t^*(\tilde{F}^2(t)))}{\varphi_t^*(\tilde{F}^2(t))}.$$
(49)

The term $\frac{\partial}{\partial t}(\varphi_t^*\tilde{F}^2(t))$ becomes

$$\frac{\partial}{\partial t}(\varphi_t^* \tilde{F}^2(t)) = \frac{\partial}{\partial s}(\varphi_{s+t}^*(\tilde{F}^2(s+t)))|_{s=0}$$

$$= \varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \frac{\partial}{\partial s}(\varphi_{s+t}^*(\tilde{F}^2(t)))|_{s=0}$$

$$= \varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \frac{\partial}{\partial s}((\varphi_t^{-1} \circ \varphi_{t+s})^*(\varphi_t^*(\tilde{F}^2(t))))|_{s=0}$$

$$= \varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \mathcal{L}_{\frac{\partial}{\partial s}(\varphi_t^{-1} \circ \varphi_{t+s})|_{s=0}}\varphi_t^*(\tilde{F}^2(t)).$$
(50)

On the other hand, we have

$$\frac{\partial}{\partial s}(\varphi_t^{-1} \circ \varphi_{t+s}) \mid_{s=0} = (\varphi_t^{-1})_*((\frac{\partial}{\partial s}\varphi_{s+t}) \mid_{s=0}) = (\varphi_t^{-1})_*(\xi).$$

Hence, (50) is written

$$\frac{\partial}{\partial t}(\varphi_t^*\tilde{F}^2(t)) = \varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \mathcal{L}_{(\varphi_t^{-1})_*(\xi)}\varphi_t^*(\tilde{F}^2(t)).$$

Replacing the last equation in (49) and using the assumption (48) we get

$$\begin{split} \frac{\partial}{\partial t}(\log F(t)) &= \frac{1}{2} \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \mathcal{L}_{(\varphi_t^{-1})_*(\xi)}\varphi_t^*(\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} \\ &= \frac{1}{2} \frac{\varphi_t^*(-2\tilde{F}^2(t)\mathcal{R}ic(\tilde{F}(t)) - \mathcal{L}_{\xi}\tilde{F}^2(t)) + \mathcal{L}_{(\varphi_t^{-1})_*(\xi)}\varphi_t^*(\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} \\ &= \frac{1}{2} \frac{\varphi_t^*(-2\tilde{F}^2(t)\mathcal{R}ic(\tilde{F}(t))) - \varphi_t^*(\mathcal{L}_{\xi}\tilde{F}^2(t))) + \mathcal{L}_{(\varphi_t^{-1})_*(\xi)}\varphi_t^*(\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} \\ &= \frac{1}{2} \frac{-2\varphi_t^*(\tilde{F}^2(t))\varphi_t^*(\mathcal{R}ic(\tilde{F}(t)))}{\varphi_t^*(\tilde{F}^2(t))}. \end{split}$$

By virtue of Lemma 6.2 we have

$$\frac{\partial}{\partial t}(\log F(t)) = -\varphi_t^*(\mathcal{R}ic(\tilde{F}(t))) = -\mathcal{R}ic_{\varphi_t^*(\tilde{F}(t))} = -\mathcal{R}ic_{F(t)}.$$

Therefore, the Finsler structures F(t) form a solution to the Finslerian Ricci flow. Hence the proof is complete.

Proof of Theorem 2. We check the existence of a solution to the Finslerian Ricci flow. Recall that Theorem 1 asserts that, there exists a solution $\tilde{F}(t)$ to the Finslerian Ricci-DeTurck flow (32) which is defined on some time interval [0, T) and satisfies $\tilde{F}(0) = F_0$. Let φ_t be the solution of the ODE

$$\frac{\partial}{\partial t}\varphi_t(z) = (\Phi_{\tilde{g}(t),h}Id)(\varphi_t(z),t) = \xi(\varphi_t(z),t),$$

with the initial condition $\varphi_0(z) = z$, for $z \in SM$ and $t \in [0, T)$. By Proposition 6.1, the Finsler structures $F^2(t) = \varphi_t^*(\tilde{F}^2(t))$ form a solution to the Finslerian Ricci flow (31) with $F(0) = F_0$. This completes the proof of Theorem 2.

In the following proposition, we assume that there exists a solution to the Finslerian Ricci flow, from which we construct a solution to the Ricci-DeTurck flow.

Proposition 6.2. Consider a fixed compact Finsler manifold (M, \bar{F}) , with the related Finsler metric tensor h. Let F(t), $t \in [0, T)$, be a family of solutions to the Ricci flow and φ_t a one-parameter family of diffeomorphisms on SM deforming under the following flow,

$$\frac{\partial}{\partial t}\varphi_t = \Phi_{g(t),h}\varphi_t$$

Then the Finsler structures $\tilde{F}(t)$ defined by $F^2(t) = \varphi_t^*(\tilde{F}^2(t))$ form a solution to the following Ricci-DeTurck flow

$$\frac{\partial}{\partial t}\tilde{F}^2(t) = -2\tilde{F}^2(t)\mathcal{R}ic(\tilde{g}(t)) - \mathcal{L}_{\xi}\tilde{F}^2(t),$$

where, $\xi = (\Phi_{\tilde{g}(t),h}Id)^i \frac{\partial}{\partial x^i}$ and $\tilde{g}(t)$ is the metric tensor related to $\tilde{F}(t)$. Furthermore, for all $z \in SM$ and $t \in [0,T)$ we have

$$\frac{\partial}{\partial t}\varphi_t(z) = \xi(\varphi_t(z), t)$$

Proof. Using Lemma 4.1 we have

$$\frac{\partial}{\partial t}\varphi_t = \Phi_{g(t),h}\varphi_t = \Phi_{\varphi_t^*(\tilde{g}(t)),h}\varphi_t = \Phi_{\varphi_t^*(\tilde{g}(t)),h}Id \circ \varphi_t$$
$$= \Phi_{\varphi_t^*(\tilde{g}(t)),h}\varphi_t^*Id = \Phi_{\tilde{g}(t),h}Id = \xi,$$

for all $z \in SM$ and $t \in [0,T)$. Having $F^2(t) = \varphi_t^*(\tilde{F}^2(t))$, we obtain

$$\frac{\partial}{\partial t}(\log F(t)) = \frac{1}{2} \frac{\frac{\partial}{\partial t}(\varphi_t^*(\tilde{F}^2(t)))}{\varphi_t^*(\tilde{F}^2(t))}
= \frac{1}{2} \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t)) + \mathcal{L}_{(\varphi_t^{-1})_*(\xi)}\varphi_t^*(\tilde{F}^2(t)))}{\varphi_t^*(\tilde{F}^2(t))}
= \frac{1}{2} \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))}.$$
(51)

By assumption, F(t) form a solution to the Finslerian Ricci flow (31)

$$0 = \frac{\partial}{\partial t} (\log F(t)) + \mathcal{R}ic_{F(t)}.$$
(52)

By means of (51), (52) and Lemma 6.2 we have

$$\begin{split} 0 &= \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} + 2\mathcal{R}ic_{F(t)} \\ &= \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} + 2\mathcal{R}ic_{\varphi_t^*(\tilde{F}(t))} \\ &= \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t))}{\varphi_t^*(\tilde{F}^2(t))} + 2\varphi_t^*(\mathcal{R}ic_{\tilde{F}(t)}) \\ &= \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t)) + 2\varphi_t^*(\tilde{F}^2(t))\varphi_t^*(\mathcal{R}ic_{\tilde{F}(t)})}{\varphi_t^*(\tilde{F}^2(t))} \\ &= \frac{\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t) + 2\tilde{F}^2(t)\mathcal{R}ic_{\tilde{F}(t)})}{\varphi_t^*(\tilde{F}^2(t))}. \end{split}$$

~ ~

Therefore, $\varphi_t^*(\frac{\partial}{\partial t}\tilde{F}^2(t) + \mathcal{L}_{\xi}\tilde{F}^2(t) + 2\tilde{F}^2(t)\mathcal{R}ic_{\tilde{F}(t)}) = 0$, which implies

$$\frac{\partial}{\partial t}\tilde{F}^2(t) = -2\tilde{F}^2(t)\mathcal{R}ic_{\tilde{F}(t)} - \mathcal{L}_{\xi}\tilde{F}^2(t).$$

Hence, $\tilde{F}(t)$ is a solution to the Ricci-DeTurck flow, as we have stated.

Proof of Theorem 3. The existence statement has been proved in Theorem 2 for a general Finsler structure. For uniqueness statement in the isotropic case, assume that $F_1(t)$ and $F_2(t)$ are both solutions to the Finslerian Ricci flow defined on some time interval [0,T) and satisfy $F_1(0) = F_2(0)$ at the point t = 0. We claim $F_1(t) = F_2(t)$ for all $t \in [0,T)$. In order to prove this fact, we argue by contradiction. Suppose that $F_1(t) \neq F_2(t)$ for some $t \in [0,T)$. Let us consider a real number $\tau \in [0,T)$ where $\tau = \inf\{t \in [0,T) : F_1(t) \neq F_2(t)\}$. We will easily see, $F_1(\tau) = F_2(\tau)$. Let φ_t^1 be a solution of the flow

$$\frac{\partial}{\partial t}\varphi_t^1 = \Phi_{g_1(t),h}\varphi_t^1,$$

with the initial condition $\varphi_{\tau}^1 = Id$ and φ_t^2 a solution of the flow

$$\frac{\partial}{\partial t}\varphi_t^2 = \Phi_{g_2(t),h}\varphi_t^2,$$

with the initial condition $\varphi_{\tau}^2 = Id$. It follows from the standard theory of parabolic differential equations that φ_t^1 and φ_t^2 are defined on some time interval $[\tau, \tau + \epsilon)$, where, ϵ is a positive real number. Moreover, if we choose $\epsilon > 0$ small enough, then φ_t^1 and φ_t^2

are diffeomorphisms for all $t \in [\tau, \tau + \epsilon)$. For each $t \in [\tau, \tau + \epsilon)$ we define the two Finsler structures $\tilde{F}_1(t)$ and $\tilde{F}_2(t)$ by $(F_1(t))^2 = (\varphi_t^1)^* (\tilde{F}_1(t))^2$ and $(F_2(t))^2 = (\varphi_t^2)^* (\tilde{F}_2(t))^2$. It follows from Proposition 6.2 that $\tilde{F}_1(t)$ and $\tilde{F}_2(t)$ are the solutions of the Finslerian Ricci-DeTurck flow. Since $\tilde{F}_1(\tau) = \tilde{F}_2(\tau)$, the uniqueness of solution to the Finslerian Ricci-DeTurck flow on isotropic Finsler manifolds mentioned in Remark 5.1 implies that $\tilde{F}_1(t) = \tilde{F}_2(t)$ for all $t \in [\tau, \tau + \epsilon)$. For each $t \in [\tau, \tau + \epsilon)$, we define a vector field ξ on SMby

$$\xi = \Phi_{\tilde{g}_1(t),h} Id = \Phi_{\tilde{g}_2(t),h} Id.$$

By Proposition 6.2, we have

$$\frac{\partial}{\partial t}\varphi_t^1(z) = \xi(\varphi_t^1(z), t),$$

and

$$\frac{\partial}{\partial t}\varphi_t^2(z) = \xi(\varphi_t^2(z), t)$$

for $z \in SM$ and $t \in [\tau, \tau + \epsilon)$. Since $\varphi_{\tau}^1 = \varphi_{\tau}^2 = Id$, it follows that $\varphi_t^1 = \varphi_t^2$ for all $t \in [\tau, \tau + \epsilon)$. Putting these facts together, we conclude that

$$(F_1(t))^2 = (\varphi_t^1)^* (\tilde{F}_1(t))^2 = (\varphi_t^2)^* (\tilde{F}_2(t))^2 = (F_2(t))^2,$$

for all $t \in [\tau, \tau + \epsilon)$. Therefore, $F_1(t) = F_2(t)$ for all $t \in [\tau, \tau + \epsilon)$. This contradicts the definition of τ . Thus the uniqueness holds well. This completes the proof of Theorem 3.

Example 6.1. Let (M, F_0) be an Einstein Randers manifold with dimension n > 2. We are going to obtain a solution to the Ricci flow (31). Using Schur's lemma for Einstein Randers metrics [19, p. 30], the Ricci scalar Ric of F_0 is necessarily constant, that is $\mathcal{R}ic_{F_0} = K$ where, K is a constant. Consider a family of linear scalars $\tau(t)$ defined by

$$\tau(t) := 1 - 2Kt > 0.$$

Define a smooth one-parameter family of Finsler structures on M by

$$F^2(t) := \tau(t) F_0^2.$$

Thus we have

$$\log(F(t)) = \frac{1}{2}\log(\tau(t)F_0^2).$$

Derivative with respect to t yields

$$\frac{\partial}{\partial t}\log(F(t)) = -\frac{K}{\tau(t)} = -\frac{\mathcal{R}ic_{F_0}}{\tau(t)}.$$
(53)

On the other hand, by straight forward computations we have $\frac{1}{\tau(t)} \mathcal{R}ic_{F_0} = \mathcal{R}ic_{\tau(t)^{\frac{1}{2}}F_0}$, for more details see [10, p. 926]. Replacing the last relation in (53) leads to

$$\frac{\partial}{\partial t}\log(F(t)) = -\mathcal{R}ic_{\tau(t)^{\frac{1}{2}}F_0} = -\mathcal{R}ic_{F(t)}.$$

Hence, F(t) is a solution to the Ricci flow equation (31).

Example 6.2. Let (M, F_0) be a Finsler manifold and $V = V^i(x) \frac{\partial}{\partial x^i}$ a vector field on M. The triple (M, F_0, V) is called a Finslerian Ricci soliton if there exists a constant λ such that

$$2Ric_{ij} + \mathcal{L}_{\hat{V}}g_{ij} = 2\lambda g_{ij},\tag{54}$$

where, g_{ij} is the Hessian related to the Finsler structure F_0 , \hat{V} the complete lift of V and $\lambda \in \mathbb{R}$. By multiplying $y^i y^j$ in both sides of (54), we obtain

$$2F_0^2 \mathcal{R}ic + \mathcal{L}_{\hat{V}}F_0^2 = 2\lambda F_0^2.$$

Depending on the sign of λ , a Finslerian Ricci soliton is called shrinking $(\lambda > 0)$, steady $(\lambda = 0)$, or expanding $(\lambda < 0)$. Suppose (M, F_0, V) is a compact Finslerian Ricci soliton. For each point $p \in M$, we denote by $\varphi_t(p)$ the unique solution of the ordinary differential equation (ODE)

$$\frac{\partial}{\partial t}\varphi_t(p) = \frac{1}{1 - 2\lambda t} V|_{\varphi_t(p)},$$

with the initial condition $\varphi_0(p) = p$. This defines a one-parameter family of diffeomorphisms on M. One can define a smooth one-parameter family of Finsler structures on TM as follow

$$F^2(t) = (1 - 2\lambda t)\tilde{\varphi}_t^*(F_0^2),$$

where, $\tilde{\varphi}_t^*$ is the one-parameter group on TM associated to the complete lift $\frac{1}{1-2\lambda t}\hat{V}$. Then F(t) form a solution to the Ricci flow equation (31), see [10, p. 928].

In conclusion, this paper has studied the short-time existence and uniqueness of Ricci flow solutions on isotropic Finsler manifolds with dimension n. We have shown that the short-time existence of the solution can be established, and if the Finsler manifold is isotropic, the solution is unique. The results of this paper highlight the versatility of the Ricci flow, as it can be applied to various areas of geometry and physics, as well as computer graphics, medical imaging, and surface matching.

References

 H. Akbar-Zadeh, Initiation to global Finslerian geometry, Vol. 68. Elsevier Science, 2006.

- [2] D. Bao, On two curvature-driven problems in Riemann-Finsler geometry: In memory of Makoto Matsumoto, Advanced studies in pure mathematics, Vol. 48, Mathematical Society, Japan, Tokyo, (2007), 19-71.
- [3] D. Bao, S. Chern, Z. Shen, An introduction to Riemann-Finsler Geometry. Graduate Texts in Mathematics, Vol. 200, Springer, 2000.
- [4] D. Bao and Z. Shen, On the volume of unit tangent spheres in a Finsler manifold, Results in Mathematics, Vol. 26, (1994), pp. 1-17.
- [5] B. Bidabad, P. Joharinad, Conformal vector fields on Finsler spaces, Differential Geometry and its Applications, 31, (2013), 33-40.
- [6] B. Bidabad, M.K. Sedaghat, *Ricci flow on Finsler surfaces*, Journal of Geometry and Physics, **129**, (2018), 238-254.
- [7] B. Bidabad, M.K. Sedaghat, On evolution of Finsler Ricci scalar, J. Korean Math. Soc., 55, No. 3, (2018), 749-761.
- B. Bidabad, M.K. Sedaghat, Hamilton's Ricci Flow on Finsler Spaces, https://doi.org/10.48550/arXiv.1508.02893, 1-19.
- B. Bidabad, M. Yar Ahmadi, Convergence of Finslerian metrics under Ricci flow, Sci. China. Math, Vol. 59, (2016), 741-750.
- [10] B. Bidabad, M. Yar Ahmadi, On quasi-Einstein Finler spaces, Bull. Iranian Math. Soc., Vol. 40, No. 4, (2014), 921-930.
- [11] S. Brendle, Ricci Flow and the Sphere Theorem, Vol. III, Graduate Studies in Mathematics, AMS, 2010.
- [12] B. Chow, D. Knopf, *The Ricci Flow: An Introduction*, Mathematical Survays and Monographs, Vol. **110**, AMS, Providence, RI, 2004.
- [13] D. M. DeTurck, Deforming metrics in the direction of their Ricci tensors, J. Differential Geom. 18(1) (1983), 157–162.
- [14] R. S. Hamilton, *Three manifolds with positive Ricci curvature*, J. Differential Geom. 17, (1982), 255-306.
- [15] R. S. Hamilton, *The Ricci flow on surfaces*, Math. and General Relativity, Contemporary Math. **71**, (1988), 237-262.
- [16] S. Lakzian, Differential Harnack estimates for positive solutions to heat equation under Finsler-Ricci flow, Pacific Journal of Mathematics, 278, N. 2, (2015), 447-462.

- [17] S. Ohta, K. Sturm, Bochner-Weitzenböck formula and Li-Yau estimates on Finsler manifolds, Adv. Math., 252, (2014), 429-448.
- [18] M. Renardy, R. C. Rogers, An introduction to partial differential equations, Texts in applied mathematics, second edition, Springer 2004.
- [19] C. Robles, *Einstein metrics of Randers type*, Ph.D. thesis, University of British Columbia, Canada, 2003.

Behroz Bidabad^{1,2}, Maral Khadem Sedaghat¹,

 $bidabad@aut.ac.ir; m_sedaghat@aut.ac.ir$

1. Department of Mathematics and Computer Science,

Amirkabir University of Technology (Tehran Polytechnic),

Hafez Ave., 15914 Tehran, Iran.

2. Institut de Mathematique de Toulouse (IMT), Universite' Paul Sabatier, 118 route de Narbonne - F-31062 Toulouse, France.

behroz.bidabad@math.univ-toulouse.fr