

DLR-KMS correspondence on lattice spin systems

N. Drago^{*a} and C. J. F. van de Ven^{†b}

^aDipartimento di Matematica, Università di Trento and INFN-TIFPA
and INdAM, Via Sommarive 14, I-38123 Povo, Italy

^bJulius Maximilian University of Würzburg, Department of
Mathematics Chair of Mathematics X (Mathematical Physics),
Emil-Fischer-Straße 31, 97074 Würzburg, Germany

July 12, 2023

Abstract

The Dobrushin-Lanford-Ruelle (DLR) condition [13, 22] and the classical Kubo-Martin-Schwinger (KMS) condition [18] are considered in the context of classical lattice systems. In particular, we prove that these conditions are equivalent for the case of a lattice spin system with values in a compact symplectic manifold by showing that infinite volume Gibbs states are in bijection with KMS states.

1 Introduction

Identifying the proper notion of thermal equilibrium in an infinite system is of paramount importance in statistical mechanics [26]. A common approach to this problem is to consider a finite size approximation of the system under consideration, where the notion of thermal equilibrium is captured by the renowned Gibbs state [16], *cf.* Equation (4). A subsequent limit procedure, where the size of the system diverges, leads to states which are identified (a posteriori) as those describing thermal equilibrium in the infinite system. While being a convenient and fruitful approach, this method leaves open the problem of identifying thermal equilibrium states directly on the infinite system. Such possibility is essential because it leads to a proper notion of phase transitions, independently on the chosen order parameters [16, §3].

For lattice spin systems, the Dobrushin-Lanford-Ruelle (DLR) condition provides a satisfactory solution to the stated problem [10, 11, 12, 13, 22, 25]. Therein, thermal equilibrium states are identified with probability measures

^{*}nicolo.drago@unitn.it

[†]christiaan.vandeven@mathematik.uni-wuerzburg.de

which reduce to the Gibbs state when conditioned on the complement of a finite region, see Section 2 for the precise definition. States abiding by the DLR condition are usually called infinite-volume Gibbs states. The main ingredients of this approach are the lattice formulation of the system of interest together with the notion of Gibbs states on a local region of such lattice [19].

Another conceptually clear yet different in spirit condition to pinpoint thermal equilibrium states is the renowned Kubo-Martin-Schwinger (KMS) condition. The quantum version of this condition, usually formulated in the C^* -algebraic setting, identifies thermal equilibrium in terms of an analytic condition on the correlations of the state [9, 20]. A classical version of this condition is at disposal and has been investigated in different contexts ranging from systems of infinitely many particles on \mathbb{R}^d [1, 2, 17, 18] to Poisson geometry [5, 8, 15]. The classical KMS condition fixes the expectation value of the state of interest on Poisson brackets, *cf.* Section 3 for the precise definition. Thus, it requires a Poisson structure on the system under investigation.

At this juncture a natural question arises, namely if the DLR and the (classical) KMS conditions agree whenever the system under investigation is described in terms of a lattice structure endowed with a Poisson bracket. A positive answer for systems of particles on \mathbb{R}^d has been found in [2] and similar positive results have been obtained on the quantum side [9, Thm. 6.2.18].

In this paper we deal with the DLR-KMS correspondence for the case of a lattice system whose configuration space $\Omega = M^{\mathbb{Z}^d}$ is the space of functions $\mathbb{Z}^d \rightarrow M$ with values in a compact symplectic manifold M . This provides a natural setting where both the DLR and KMS approaches apply, leading to the natural question of whether the DLR-KMS correspondence holds true also in this setting. The main result of this paper is the following theorem, which provides a positive answer to this question.

THEOREM 1: Let $\mathcal{G}(\Phi)$ be the convex set of infinite volume Gibbs states on $\Omega = M^{\mathbb{Z}^d}$ —*cf.* Section 3 and Equation (6). Let $\mathcal{K}(\Phi)$ be the convex set of X^Φ -KMS states defined as per Equation (14), *cf.* Section 3. Then $\mathcal{G}(\Phi) = \mathcal{K}(\Phi)$. \diamond

Aside from its relevance in the discussion on the notion of thermal equilibrium, Theorem 1 is relevant for its connection with strict deformation quantization (SDQ) [3, 4, 6]. The latter provides a mathematically sound description of the classical limit of quantum theories and it is particularly suitable for the analysis of the semi-classical limit of quantum states. In view of the recent results in this framework [14, 23, 24, 27, 28], Theorem 1 stands as a technical result which may lead to a deeper understanding of the quantum-to-classical limit of thermal equilibrium states. We plan to address this question in a future investigation.

From a technical point of view this paper profits from the strategy applied in [2] together with the results obtained in [8, 15]. As a matter of fact the proof of $\mathcal{G}(\Phi) \subseteq \mathcal{K}(\Phi)$ is a direct computation, *cf.* Proposition 5. Instead, the converse inclusion requires a more refined analysis of the conditional probability $\varphi_\Lambda(\cdot|\eta)$ of a given KMS state φ with respect to the complement Λ^c of a finite region $\Lambda \Subset \mathbb{Z}^d$. This is where the results of [8, 15] apply, ensuring that such conditional probabilities $\varphi_\Lambda(\cdot|\eta)$ coincide with the local Gibbs state $\varphi_\Lambda^\Phi(\cdot|\eta)$.

The paper is organized as follows: Section 2 is devoted to a brief introduction to the lattice spin system of interest together with the precise definition of the DLR condition. Similarly, Section 3 considers the classical KMS condition in the same setting. Finally Section 4 proves Theorem 1, *cf.* Proposition 5-6.

2 DLR approach on continuous lattice system

In this section we briefly recollect a few crucial results and definitions from continuous lattice systems, see [16]. In what follows we will consider a configuration space over the lattice \mathbb{Z}^d with values in a symplectic manifold M . The differential structure of M is needed for the formulation of the classical KMS condition, *cf.* Section 3.

Let (M, ς) be a compact, connected symplectic manifold with symplectic form ς . For technical convenience we shall assume that M is metrizable and denote with d_M a complete metric on M whose induced topology coincides with the one of M . We will denote by $\mu_M := c_M \varsigma^m / m!$ the induced volume form on M , where $\dim M = 2m$ while the normalization constant $c_M > 0$ is chosen so that $\int_M \mu_M = 1$. For $f, g \in C^\infty(M)$ we denote by $\{f, g\}$ the induced Poisson bracket between f and g and set $X_f := \{ \cdot, f \}$ the Hamiltonian vector field associated with f .

The configuration space of interest is $\Omega := M^{\mathbb{Z}^d}$ which is compact in the product topology. Moreover, Ω is also metrizable once we set

$$d_\Omega(\omega, \eta) := \sum_{i \in \mathbb{Z}^d} 2^{-\|i\|_\infty} \frac{d_M(\omega_i, \eta_i)}{1 + d_M(\omega_i, \eta_i)},$$

where $\|i\|_\infty := \sup_{k \in \{1, \dots, d\}} |i_k|$.

If $\Lambda \subset \mathbb{Z}^d$ we set $\Omega_\Lambda := M^\Lambda$ and denote by

$$\pi_\Lambda: \Omega \rightarrow \Omega_\Lambda \quad \pi_\Lambda(\omega) := \omega_\Lambda := \omega|_\Lambda,$$

the projection over Ω_Λ . With a standard slight abuse of notation we will set, given $\omega_\Lambda \in \Omega_\Lambda$ and $\omega_{\Lambda^c} \in \Omega_{\Lambda^c}$, $\Lambda \subset \mathbb{Z}^d$, $\omega_\Lambda \omega_{\Lambda^c} := \omega_\Lambda \otimes \omega_{\Lambda^c} \in \Omega$. For a given $\Lambda \subset \mathbb{Z}^d$ we will write $\Lambda \Subset \mathbb{Z}^d$ if $|\Lambda| < +\infty$: Notice that in this latter

case $C(\Omega_\Lambda) \simeq C(M^{|\Lambda|})$ where $M^{|\Lambda|}$ is again a connected compact symplectic manifold with symplectic form $\varsigma_\Lambda := \oplus_{i \in \Lambda} \varsigma$.

The algebra of observables over the configuration space Ω is identified with the commutative C^* -algebra $C(\Omega)$ equipped with the supremum norm $\|f\|_\infty := \sup_{\omega \in \Omega} |f(\omega)|$. This space is best described in terms of (continuous) local functions. In particular, $f \in C(\Omega)$ is called **(continuous) local function**, denoted $f \in C_{\text{LOC}}(\Omega)$, if there exists $\Lambda \Subset \mathbb{Z}^d$ and $f_\Lambda \in C(\Omega_\Lambda) \simeq C(M^{|\Lambda|})$ such that $f(\omega) = f_\Lambda(\omega_\Lambda)$ for all $\omega \in \Omega$. Then, for all $f \in C(\Omega)$ there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of local functions such that $\lim_{n \rightarrow \infty} \|f - f_n\|_\infty = 0$ [16, Lem. 6.21] (*e.g.* one considers $\eta \in \Omega$ and a sequence $(\Lambda_n)_{n \in \mathbb{N}}$ of increasing subsets $\Lambda_n \subset \Lambda_{n+1} \Subset \mathbb{Z}^d$ such that $\cup_n \Lambda_n = \Omega$ and set $f_n(\omega) := f(\omega_{\Lambda_n} \eta_{\Lambda_n^c})$ for all $n \in \mathbb{N}$). For this reason, $C(\Omega)$ is usually referred to as the C^* -algebra of **(continuous) quasi-local functions**.

Per definition, a **state** on $C(\Omega)$ is a normalized linear, positive functional $\varphi: C(\Omega) \rightarrow \mathbb{C}$. By the Riesz-Markov-Kakutani theorem every state φ on $C(\Omega)$ is completely described by a Radon probability measure $\mu \in \mathcal{P}(\Omega, \mathcal{F})$ on (Ω, \mathcal{F}) —here \mathcal{F} denotes the Borel σ -algebra over Ω . In particular $\varphi \equiv \varphi_\mu$ and

$$\varphi_\mu(f) = \int_{\Omega} f d\mu \quad \forall f \in C(\Omega). \quad (1)$$

For later convenience we recall the definition of the σ -algebra \mathcal{F}_Λ over Ω of events occurring in $\Lambda \subset \mathbb{Z}^d$:

$$\mathcal{F}_\Lambda := \sigma \left(\bigcup_{\substack{\Lambda' \Subset \mathbb{Z}^d \\ \Lambda' \subset \Lambda}} \{ \pi_{\Lambda'}^{-1}(E) \mid E \in \mathcal{B}_{\Lambda'} \} \right),$$

where $\sigma(A)$ is the σ -algebra generated by the collection A while, if $\Lambda \Subset \mathbb{Z}^d$, \mathcal{B}_Λ denotes the Borel σ -algebra over $\otimes_{i \in \Lambda} M \simeq M^{|\Lambda|}$. Notice that

$$\mathcal{F}_\Lambda = \{ \pi_\Lambda^{-1}(E) \mid E \in \mathcal{B}_\Lambda \} \quad \forall \Lambda \Subset \mathbb{Z}^d,$$

while $\mathcal{F}_{\mathbb{Z}^d}$ coincides with the Borel σ -algebra \mathcal{F} over Ω . A function $f: \Omega \rightarrow \mathbb{C}$ is \mathcal{F}_Λ -measurable if and only if f is Λ -**local**, namely there exists a measurable function $f_\Lambda: \Omega_\Lambda \rightarrow \mathbb{C}$ such that $f(\omega) = f_\Lambda(\omega_\Lambda)$ for all $\omega \in \Omega$ [16, Lem. 6.3].

Among all possible states φ_μ on $C(\Omega)$ we will be interested in the infinite volume Gibbs states. The definition of the latter requires to introduce a **potential**, namely a collection $\Phi = \{ \Phi_\Lambda \}_{\Lambda \Subset \mathbb{Z}^d}$, where $\Phi_\Lambda \in C(\Omega)$ is \mathcal{F}_Λ -measurable for all $\Lambda \Subset \mathbb{Z}^d$. In what follows we will assume the following technical assumptions on Φ :

(I) For all $\Lambda \Subset \mathbb{Z}^d$, $\Phi_\Lambda \in C^1(\Omega_\Lambda) \simeq C^1(M^{|\Lambda|})$.

(II) For all $i \in \mathbb{Z}^d$

$$\sum_{\substack{\Lambda \Subset \mathbb{Z}^d \\ i \in \Lambda}} \|\Phi_\Lambda\|_{C^1(\Omega_\Lambda)} < +\infty, \quad (2)$$

where $\|\Phi_\Lambda\|_{C^1(M_\Lambda)}$ denotes the supremum of Φ_Λ and $d\Phi_\Lambda$ over Ω_Λ , thus it coincides with the norm of the C^* -algebra $C^1(M^{|\Lambda|})$.

Within these assumptions we can define

$$H_\Lambda^\Phi := \sum_{\substack{\Lambda' \in \mathbb{Z}^d \\ \Lambda' \cap \Lambda \neq \emptyset}} \Phi_{\Lambda'}, \quad (3)$$

which is a well-defined element in $C(\Omega)$ on account of assumption (II). For a fixed $\eta \in \Omega$ one then introduces the probability measure $\mu_\Lambda^\Phi(\cdot|\eta)$ on (Ω, \mathcal{F}) defined by

$$\mu_\Lambda^\Phi(A|\eta) := \frac{1}{Z_{\eta, \Lambda}^\Phi} \int_{\Omega_\Lambda} 1_A(\omega_\Lambda \eta_{\Lambda^c}) e^{-H_\Lambda^\Phi(\omega_\Lambda \eta_{\Lambda^c})} d\mu_{M^{|\Lambda|}}(\omega_\Lambda) \quad \forall A \in \mathcal{F}, \quad (4)$$

where $\mu_{M^{|\Lambda|}} := \otimes_{i \in \Lambda} \mu_M$ denotes the (normalized) Liouville volume form over $M^{|\Lambda|}$ while $Z_{\eta, \Lambda}^\Phi$ is a normalization constant —we have implicitly identified $\Omega_\Lambda \simeq M^{|\Lambda|}$. We will refer to $\mu_\Lambda^\Phi(\cdot|\eta)$ as the **Λ -Gibbs measure** associated with the potential Φ and with boundary condition η (at fixed inverse temperature $\beta = 1$). We will denote by $\varphi_\Lambda^\Phi(\cdot|\eta)$ the associated state on $C(\Omega)$ which will be referred to as the **Λ -Gibbs state** associated with Φ and η .

REMARK 2: The assumption on the potential Φ are slightly stronger than those usually imposed [16, §6]. As a matter of fact, one usually requires

$$\sum_{\substack{\Lambda \in \mathbb{Z}^d \\ i \in \Lambda}} \|\Phi_\Lambda\|_{C(\Omega_\Lambda)} < +\infty,$$

which ensures the well-definiteness of H_Λ^Φ as per Equation (3). The stronger assumptions (I)-(II) are necessary for discussing the Poisson bracket on Ω —*cf.* Section 3. \diamond

The collection $\mu^\Phi := \{\mu_\Lambda^\Phi\}_{\Lambda \in \mathbb{Z}^d}$ is called **Gibbs specification** and enjoys the following properties:

- (a) For all $\Lambda \in \mathbb{Z}^d$ and $\eta \in \Omega$, $\mu_\Lambda^\Phi(\cdot|\eta)$ is a probability measure over (Ω, \mathcal{F}) ;
- (b) For all $\Lambda \in \mathbb{Z}^d$ and $A \in \mathcal{F}$, the function $\Omega \ni \eta \mapsto \mu_\Lambda^\Phi(A|\eta)$ is \mathcal{F}_{Λ^c} -measurable.

Properties (a)-(b) can be summarized by saying that μ_Λ^Φ is a **probability kernel** from \mathcal{F}_{Λ^c} to \mathcal{F} [21, §5] (also known as transition measures [7, Def. 10.7.1]).

- (c) For all $\Lambda \in \mathbb{Z}^d$ and $A \in \mathcal{F}_{\Lambda^c}$, it holds $\mu_\Lambda^\Phi(A|\eta) = 1_A(\eta)$. Probability kernels abiding by this assumption are called **proper**.

- (d) The family of probability kernels μ^Φ is **compatible**, namely, for all $\Lambda_1 \subset \Lambda_2 \Subset \mathbb{Z}^d$ it holds $\mu_{\Lambda_2}^\Phi \mu_{\Lambda_1}^\Phi = \mu_{\Lambda_2}^\Phi$, where

$$\mu_{\Lambda_2}^\Phi \mu_{\Lambda_1}^\Phi : \mathcal{F} \times \Omega \ni (A, \eta) \mapsto \int_{\Omega} \mu_{\Lambda_1}^\Phi(A|\omega) d\mu_{\Lambda_2}^\Phi(\omega|\eta).$$

For any probability measure $\mu \in \mathcal{P}(\Omega, \mathcal{F})$ and $\Lambda \Subset \mathbb{Z}^d$ one may introduce a new probability measure $\mu \mu_\Lambda^\Phi \in \mathcal{P}(\Omega, \mathcal{F})$ defined by

$$\mu \mu_\Lambda^\Phi(A) := \int_{\Omega} \mu_\Lambda^\Phi(A|\eta) d\mu(\eta). \quad (5)$$

A state φ_μ is called **infinite volume Gibbs state**, denoted $\varphi_\mu \in \mathcal{G}(\Phi)$, if the associated Radon probability measure μ fulfils

$$\mu \mu_\Lambda^\Phi = \mu \quad \forall \Lambda \Subset \mathbb{Z}^d. \quad (6)$$

Any $\varphi_\mu \in \mathcal{G}(\Phi)$ is interpreted as a state in thermal equilibrium (at fixed inverse temperature $\beta = 1$) with respect to the formal Hamiltonian $H_{\mathbb{Z}^d} := \sum_{\Lambda \Subset \mathbb{Z}^d} \Phi_\Lambda$. This condition allows to characterize thermal equilibrium states on the whole configuration space Ω , without resorting to a finite configuration space Ω_Λ , $\Lambda \Subset \mathbb{Z}^d$. This approach provides a rather neat definition of phase transition: A **phase transition** occurs whenever $\mathcal{G}(\Phi)$ is not a singleton. In the physical jargon this is equivalent to the existence of different equilibrium states in the infinite volume system.

The compatibility condition (6) with the Gibbs specification reflect the property that locally, *i.e.* on Ω_Λ for $\Lambda \Subset \mathbb{Z}^d$ with fixed boundary conditions on Λ^c , all equilibrium states coincide with the Gibbs measure (4). In physical terms, there are no phase transitions in a finite system.

As we shall see in the next section, in the current situation the symplectic structure on M allows for a different characterization of thermal equilibrium states by means of the (classical) KMS condition [17, 18]. The goal of this paper is to prove that these two characterizations of thermal states coincide.

REMARK 3: Condition (6) can be interpreted by saying that, for all $\Lambda \Subset \mathbb{Z}^d$, $\mu_\Lambda^\Phi(\cdot|\eta)$ is a regular conditional measure for μ with respect to \mathcal{F}_{Λ^c} [7, Def. 10.4.1]. For later convenience we stress that any probability measure $\mu \in \mathcal{P}(\Omega, \mathcal{F})$ arising from a state φ_μ on $C(\Omega)$ has a **regular conditional measure** μ_Λ with respect to \mathcal{F}_{Λ^c} for all $\Lambda \Subset \mathbb{Z}^d$. This follows from [7, Cor. 10.4.7], together with the observation that: (a) since M is metrizable, Ω is second-countable (and also a **Polish space** because it is metrizable), therefore, \mathcal{F} is countably generated; (b) since Ω is compact, the Riesz-Markov-Kakutani theorem ensures that the probability measure μ associated with any state φ_μ is Radon, thus it has an approximating compact class (made by compact subsets of Ω). Thus, for all $\Lambda \Subset \mathbb{Z}^d$ there exists

$$\mu_\Lambda : \mathcal{F} \times \Omega \ni (A, \eta) \mapsto \mu_\Lambda(A|\eta) \in [0, 1],$$

such that $\mu_\Lambda(\cdot|\eta) \in \mathcal{P}(\Omega, \mathcal{F})$ for μ -almost all $\eta \in \Omega$ and $\mu_\Lambda(A|\cdot)$ is \mathcal{F}_{Λ^c} -measurable, moreover,

$$\mu(A \cap B) = \int_B \mu_\Lambda(A|\eta) d\mu(\eta) \quad \forall A \in \mathcal{F}, \forall B \in \mathcal{F}_{\Lambda^c}. \quad (7)$$

Thus $\varphi_\mu \in \mathcal{G}(\Phi)$ if and only if $\mu_\Lambda(\cdot|\eta) = \mu_\Lambda^\Phi(\cdot|\eta)$ for μ -almost all $\eta \in \Omega$ and for all $\Lambda \Subset \mathbb{Z}^d$. It is worth observing that, since \mathcal{F} is countably generated, $\mu_\Lambda(\cdot|\eta)$ is unique for all $\eta \in \Omega$ up to a μ -null set [7, Lem. 10.4.3]. Moreover, since $\mu_\Lambda(\cdot|\eta)$ is μ -integrable, a density argument shows that, for all $f \in C(\Omega)$,

$$\varphi_\mu(f) = \int_\Omega \varphi_\Lambda(f|\eta) d\mu(\eta),$$

where $\varphi_\Lambda(\cdot|\eta)$ denotes the state on $C(\Omega)$ associated with $\mu_\Lambda(\cdot|\eta)$. In particular, if $f, g \in C(\Omega)$ and g is \mathcal{F}_{Λ^c} -measurable, $\Lambda \Subset \mathbb{Z}_+^d$, then

$$\varphi_\mu(fg) = \int_\Omega \varphi_\Lambda(f|\eta)g(\eta) d\mu(\eta). \quad (8)$$

Finally, it is worth observing that, as a consequence of [7, Cor. 10.4.10] $\varphi_\Lambda(\cdot|\eta)$ is concentrated on $\pi_{\Lambda^c}^{-1}(\eta_{\Lambda^c})$ for all $\Lambda \Subset \mathbb{Z}^d$ and μ -almost all $\eta \in \Omega$. Notice that $\pi_{\Lambda^c}^{-1}(\eta_{\Lambda^c}) \in \mathcal{F}_{\Lambda^c}$ because $\pi_{\Lambda^c}^{-1}(\eta_{\Lambda^c}) = \bigcap_{n \in \mathbb{N}} \pi_{\Lambda_n \setminus \Lambda}^{-1}(\eta_{\Lambda_n \setminus \Lambda^c})$ where $(\Lambda_n)_{n \in \mathbb{N}}$ is an increasing sequence of finite subsets $\Lambda_n \Subset \mathbb{Z}^d$ such that $\bigcup_{n \in \mathbb{N}} \Lambda_n = \mathbb{Z}^d$. \diamond

3 Classical KMS condition on lattice systems

This section is devoted to describe the classical KMS condition on the lattice system $\Omega = M^{\mathbb{Z}^d}$. This condition is used to select a particular class of states φ_μ by means of the Poisson structure carried by M [1, 2, 17, 18]. This is parallel to the notion of infinite volume Gibbs states (6) which is instead based on the Λ -local Gibbs measure (4).

To set the stage we recall that, for all $\Lambda \Subset \mathbb{Z}^d$, $\Omega_\Lambda \simeq M^{|\Lambda|}$ is a compact, connected symplectic manifold with $\varsigma_\Lambda := \oplus_{i \in \Lambda} \varsigma$. For $f, g \in C^\infty(\Omega_\Lambda)$ we denote by $\{f, g\}_\Lambda$ the associated Poisson bracket.

REMARK 4: For later convenience we recall the following property of the state φ_{Ω_Λ} on $C(\Omega_\Lambda) \simeq C(M^{|\Lambda|})$ induced by the normalized Liouville measure $\mu_{M^{|\Lambda|}}$ on $M^{|\Lambda|}$. Actually, φ_{Ω_Λ} is the unique state invariant under all Hamiltonian vector fields. In fact, if ψ_Λ is a state on $C(\Omega_\Lambda)$ such that

$$\psi_\Lambda(\{f, g\}_\Lambda) = 0 \quad \forall f, g \in C^\infty(\Omega_\Lambda),$$

then $\psi_\Lambda = \varphi_{\Omega_\Lambda}$ *i.e.*

$$\psi_\Lambda(f) = \int_{\Omega_\Lambda} f(\omega_\Lambda) d\mu_{M^{|\Lambda|}}(\omega_\Lambda).$$

The proof of this result can be found in [2, Lem. 1] as well as in [8, Cor. 2.7]. It is based on a localization process in coordinated charts equipped with Darboux coordinates —so that one can write $\partial_a f = \{f, x^a\}_\Lambda$ — together with the observation that the Lebesgue measure on \mathbb{R}^d is the unique (up to multiplicative constants) measure vanishing on derivatives of compactly supported functions [8, Lem. 2.6]. \diamond

The KMS condition requires to endow $C(\Omega)$ with a Poisson bracket. We recall that a **Poisson bracket** $\{ , \}$ on $C(\Omega)$ is a skew-symmetric bilinear map $\{ , \} : D \times D \rightarrow D$ defined on a dense subspace D of $C(\Omega)$ and fulfilling the **Jacobi identity**

$$\{\{f, g\}, h\} = \{\{f, h\}, g\} + \{f, \{g, h\}\} \quad \forall f, g, h \in D.$$

In what follows we will consider $D = C_{\text{LOC}}^\infty(\Omega)$ where $C_{\text{LOC}}^\infty(\Omega)$ is the algebra of smooth local functions: $f \in C_{\text{LOC}}^\infty(\Omega)$ if there exists $\Lambda \Subset \mathbb{Z}^d$ and $f_\Lambda \in C^\infty(\Omega_\Lambda)$ such that $f(\omega) = f_\Lambda(\omega_\Lambda)$.

To define a Poisson bracket on $C(\Omega)$ we notice that, $\{f, g\}_\Lambda$ makes sense also for $f, g \in C^\infty(\Omega_{\Lambda'})$ for all $\Lambda \subset \Lambda' \Subset \mathbb{Z}^d$. As a matter of fact, if $\Lambda_1 \subset \Lambda_2 \Subset \mathbb{Z}^d$, denoting $\pi_1^2 : \Omega_{\Lambda_2} \rightarrow \Omega_{\Lambda_1}$ the projection $\omega_{\Lambda_2} = \omega_{\Lambda_1} \omega_{\Lambda_2 \setminus \Lambda_1} \mapsto \omega_{\Lambda_1}$, we find

$$\{(\pi_1^2)^* f, g\}_{\Lambda_2} = \{(\pi_1^2)^* f, g\}_{\Lambda_1}, \quad (9)$$

for all $f \in C^\infty(\Omega_{\Lambda_1})$ and $g \in C^\infty(\Omega_{\Lambda_2})$ where $(\pi_1^2)^* : C(\Omega_{\Lambda_1}) \rightarrow C(\Omega_{\Lambda_2})$ —in what follows we will not write π_1^2 since its use will be clear from the context. The Poisson bracket $\{ , \}$ on $C(\Omega)$ is defined by

$$\{f, g\} := \{f, g\}_\Lambda, \quad (10)$$

where $f, g \in C_{\text{LOC}}^\infty(\Omega)$ while $\Lambda \Subset \mathbb{Z}^d$ is such that $f, g \in C^\infty(\Omega_\Lambda)$. On account of Equation (9) the value of $\{f, g\}_\Lambda$ does not change if we enlarge Λ . By direct inspection $\{ , \}$ defines a Poisson bracket on $C(\Omega)$.

The next ingredient for stating the KMS condition is the choice of a vector field X on $C(\Omega)$ which plays the role of the infinitesimal generator of the dynamics on Ω . Notice that, despite this interpretation, the (classical) KMS condition does not rely on the existence of a dynamics integrating X . We now consider the vector field

$$X_\Lambda^\Phi : C_{\text{LOC}}^\infty(\Omega) \rightarrow C(\Omega) \quad X_\Lambda^\Phi(f) := \{f, H_\Lambda^\Phi\} = \sum_{\substack{\Lambda' \Subset \mathbb{Z}^d \\ \Lambda' \cap \Lambda \neq \emptyset}} \{f, \Phi_{\Lambda'}\}_{\Lambda'}, \quad (11)$$

where the series converges on account of assumption (II). Notice that, $\{f, H_\Lambda^\Phi\}$ is only continuous on Ω , moreover, it is not a local function.

By direct inspection we also have, for all $\Lambda'' \subseteq \Lambda \Subset \mathbb{Z}^d$ and $f \in C^\infty(\Omega_{\Lambda''})$,

$$\{f, H_\Lambda^\Phi\} = \sum_{\substack{\Lambda' \Subset \mathbb{Z}^d \\ \Lambda' \cap \Lambda \neq \emptyset}} \{f, \Phi_{\Lambda'}\}_{\Lambda' \cap \Lambda''} = \sum_{\substack{\Lambda' \Subset \mathbb{Z}^d \\ \Lambda' \cap \Lambda'' \neq \emptyset}} \{f, \Phi_{\Lambda'}\}_{\Lambda' \cap \Lambda''} = \{f, H_{\Lambda''}^\Phi\}. \quad (12)$$

Equation (12) implies that, if f is Λ'' -local and $\Lambda'' \subseteq \Lambda \in \mathbb{Z}^d$, then $X_\Lambda^\Phi(f)$ does not change if we enlarge Λ . This allows to introduce a global vector field X^Φ defined by

$$X^\Phi: C_{\text{loc}}^\infty(\Omega) \rightarrow C(\Omega) \quad X^\Phi(f) = \{f, H_\Lambda^\Phi\} \quad \forall f \in C^\infty(\Omega_\Lambda). \quad (13)$$

Loosely speaking, one may think of X^Φ as the Hamiltonian vector field associated to the formal Hamiltonian $H = \sum_{\Lambda \in \mathbb{Z}^d} \Phi_\Lambda$.

We can finally introduce the KMS condition associated with the vector field X^Φ and the Poisson bracket $\{, \}$. The latter condition select a particular subclass of states φ_μ on $C(\Omega)$ with a constraint on the expectation value of Poisson brackets.

In more details, a state φ_μ on $C(\Omega)$ is called **X^Φ -KMS state** if

$$\varphi_\mu(\{f, g\}) = \varphi_\mu(gX^\Phi(f)) \quad \forall f, g \in C_{\text{loc}}^\infty(\Omega). \quad (14)$$

We denote by $\mathcal{K}(\Phi)$ the convex set of KMS states.

4 Proof of the classical DLR-KMS correspondence

This section is devoted to the proof of Theorem 1 which asserts the equivalence between the DLR condition (6) and the KMS condition (14). The argument profits of the one presented in [2, Lem. 2].

To lighten the presentation, the proof of Theorem 1 is divided in two propositions.

PROPOSITION 5: It holds $\mathcal{G}(\Phi) \subseteq \mathcal{K}(\Phi)$, namely every infinite volume Gibbs state is also a X^Φ -KMS state. \diamond

Proof. Let $\varphi_\mu \in \mathcal{G}(\Phi)$, $\Lambda \in \mathbb{Z}^d$ and $f, g \in C^\infty(\Omega_\Lambda)$. Equation (6) implies

$$\varphi_\mu(\{f, g\}) = \int \varphi_\Lambda^\Phi(\{f, g\}_\Lambda | \eta) d\mu(\eta).$$

By direct inspection we also have

$$\begin{aligned} \varphi_\Lambda^\Phi(\{f, g\}_\Lambda | \eta) &:= \frac{1}{Z_{\eta, \Lambda}^\Phi} \int_{\Omega_\Lambda} \{f, g\}_\Lambda(\omega_\Lambda \eta_{\Lambda^c}) e^{-H_\Lambda^\Phi(\omega_\Lambda \eta_{\Lambda^c})} d\mu_{M|\Lambda|}(\omega_\Lambda) \\ &= \frac{1}{Z_{\eta, \Lambda}^\Phi} \int_{\Omega_\Lambda} \{f, g e^{-H_\Lambda^\Phi}\}_\Lambda(\omega_\Lambda \eta_{\Lambda^c}) d\mu_{M|\Lambda|}(\omega_\Lambda) \\ &\quad + \frac{1}{Z_{\eta, \Lambda}^\Phi} \int_{\Omega_\Lambda} g \{f, H_\Lambda^\Phi\}_\Lambda(\omega_\Lambda \eta_{\Lambda^c}) e^{-H_\Lambda^\Phi(\omega_\Lambda \eta_{\Lambda^c})} d\mu_{M|\Lambda|}(\omega_\Lambda) \\ &= \varphi_\Lambda^\Phi(g \{f, H_\Lambda^\Phi\}_\Lambda | \eta) = \varphi_\Lambda^\Phi(gX^\Phi(f) | \eta), \end{aligned}$$

where in the last line we used that $\varphi_{\Omega_\Lambda}(\{h_1, h_2\}_\Lambda) = 0$ for all $h_1, h_2 \in C^1(\Omega_\Lambda)$. Overall we have

$$\varphi_\mu(\{f, g\}) = \int_\Omega \varphi_\Lambda^\Phi(gX^\Phi(f) | \eta) d\mu(\eta) = \varphi_\mu(gX^\Phi(f)),$$

therefore, $\varphi_\mu \in \mathcal{K}(\Phi)$. \square

PROPOSITION 6: It holds $\mathcal{K}(\Phi) \subseteq \mathcal{G}(\Phi)$, namely every X^Φ -KMS state is an infinite volume Gibbs state. \diamond

Proof. Using Equation (7) it is enough to prove that $\varphi_\Lambda(\cdot|\eta) = \varphi_\Lambda^\Phi(\cdot|\eta)$ for μ -almost all $\eta \in \Omega$ and all $\Lambda \Subset \mathbb{Z}^d$. We recall that for each $f \in C(\Omega)$, the function $\eta \mapsto \varphi_\Lambda(f|\eta)$ is \mathcal{F}_{Λ^c} -measurable, thus, it only depends on η_{Λ^c} . Moreover, $\varphi_\Lambda(\cdot|\eta)$ is concentrated on $\pi_{\Lambda^c}^{-1}(\eta_{\Lambda^c})$ μ -almost all $\eta \in \Omega$.

Let $\Lambda \Subset \mathbb{Z}^d$, $\Lambda' \Subset \Lambda^c$ and consider $f \in C^\infty(\Omega_\Lambda)$, $h \in C^\infty(\Omega_{\Lambda'})$, $g \in C_{\text{LOC}}^\infty(\Omega)$. Then Equation (9) implies that

$$h\{f, g\} = \{f, gh\},$$

since $\{f, h\} = 0$. Moreover, Equation (8) and the KMS condition (14) imply

$$\begin{aligned} \int_{\Omega} \varphi_\Lambda(\{f, g\}|\eta)h(\eta)d\mu(\eta) &= \varphi_\mu(h\{f, g\}) = \varphi_\mu(\{f, gh\}) \\ &= \varphi_\mu(ghX^\Phi(f)) = \int_{\Omega} \varphi_\Lambda(g\{f, H_\Lambda^\Phi\}|\eta)h(\eta)d\mu(\eta). \end{aligned} \quad (15)$$

The arbitrariness of $h \in C^\infty(\Omega_{\Lambda'}) \subset C(\Omega_{\Lambda^c})$ and of $\Lambda' \Subset \Lambda^c$ entails that, by an approximation argument,

$$\int_{\Omega} [\varphi_\Lambda(\{f, g\}_\Lambda|\eta) - \varphi_\Lambda(g\{f, H_\Lambda^\Phi\}_\Lambda|\eta)]h(\eta)d\mu(\eta) = 0 \quad \forall h \in C(\Omega_{\Lambda^c}).$$

Since $\eta \mapsto \varphi_\Lambda(f|\eta)$ in \mathcal{F}_{Λ^c} -measurable, it follows that there exists $N_{f,g} \in \mathcal{F}_{\Lambda^c} \subset \mathcal{F}$ with $\mu(N_{f,g}) = 0$ such that

$$\varphi_\Lambda(\{f, g\}_\Lambda|\eta) = \varphi_\Lambda(g\{f, H_\Lambda^\Phi\}_\Lambda|\eta) \quad \forall \eta \in \Omega \setminus N_{f,g}. \quad (16)$$

To proceed, we need to cope with the f, g -dependence of $N_{f,g}$. However, since $C(\Omega_\Lambda)$ is separable, we may choose $N_{f,g}$ independently on f, g : Indeed, it suffices to consider Equation (16) for f, g on a countable dense set of $C^\infty(\Omega_\Lambda)$. This leads to countably many μ -null sets whose union leads to the μ -null set N of interest.

We now prove that condition (16) implies that $\varphi_\Lambda(\cdot|\eta) = \varphi_\Lambda^\Phi(\cdot|\eta)$ for all $\eta \in \Omega \setminus N$. For each $\eta \in \Omega \setminus N$ we consider the linear positive functional ψ_Λ on $C(\Omega_\Lambda)$ defined by

$$\psi_\Lambda(f) := \varphi_\Lambda(fe^{H_\Lambda^\Phi}|\eta) \quad \forall f \in C(\Omega_\Lambda).$$

By direct inspection we have, for all $f, g \in C^\infty(\Omega_\Lambda)$,

$$\begin{aligned} \psi_\Lambda(\{f, g\}_\Lambda) &= \varphi_\Lambda(\{f, g\}_\Lambda e^{H_\Lambda^\Phi}) \\ &= \varphi_\Lambda(\{f, ge^{H_\Lambda^\Phi}\}_\Lambda) - \varphi_\Lambda(g\{f, H_\Lambda^\Phi\}_\Lambda e^{H_\Lambda^\Phi}) = 0, \end{aligned}$$

where we applied Equation (16).

Thus, ψ_Λ is a linear, positive functional on $C(\Omega_\Lambda) \simeq C(M^{|\Lambda|})$ which is invariant under all Hamiltonian vector fields. Remark 4 implies that $\widehat{\psi}_\Lambda = \varphi_{\Omega_\Lambda}$ where $\widehat{\psi}_\Lambda(f) = \psi_\Lambda(f)/\psi_\Lambda(1)$. Finally, since $\varphi_\Lambda(\cdot|\eta)$ is concentrated on $\pi_{\Lambda^c}^{-1}(\eta_{\Lambda^c})$ we have

$$\varphi_\Lambda(f|\eta) = \widehat{\psi}_\Lambda(f_\eta e^{-H_{\Lambda,\eta}^\Phi}) = \varphi_\Lambda^\Phi(f|\eta),$$

where $f_\eta e^{-H_{\Lambda,\eta}^\Phi} \in C(\Omega_\Lambda)$ is defined by $(f_\eta e^{-H_{\Lambda,\eta}^\Phi})(\omega_\Lambda) := (f e^{-H_\Lambda^\Phi})(\omega_\Lambda \eta_{\Lambda^c})$. \square

Acknowledgements. We are grateful to prof. Aernout van Enter for his suggestions and helpful discussions on this project. We also thank Klaas Landsman, Rodrigo Bissacot, Teun van Nuland, Y. Velenik and Stefan Waldmann for their fruitful comments. C. J. F. van de Ven is supported by a postdoctoral fellowship granted by the Alexander von Humboldt Foundation (Germany). N.D. acknowledges the support of the GNFM-INdAM Progetto Giovani *Non-linear sigma models and the Lorentzian Wetterich equation*.

Data availability statement. Data sharing is not applicable to this article as no new data were created or analysed in this study.

Conflict of interest statement. The authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter discussed in this manuscript.

References

- [1] Aizenman M., Gallavotti G., Goldstein S., Lebowitz J.L., *Stability and equilibrium states of infinite classical systems*, Commun.Math. Phys. 48, 1-14 (1976).
- [2] Aizenman M., Goldstein S., Gruber C., Lebowitz J.L., Martin P., *On the equivalence between KMS-states and equilibrium states for classical systems*, Commun. Math. Phys. 53, 209-220 (1977).
- [3] Bayen F., Flato M., Fronsdal C., Lichnerowicz A., Sternheimer D., *Deformation Theory and Quantization. 1. Deformations of Symplectic Structures*, Annals Phys. 111, 61 (1978).
- [4] Bayen F., Flato M., Fronsdal C., Lichnerowicz A., Sternheimer D., *Deformation Theory and Quantization. 2. Physical Applications*, Annals Phys. 111-151 (1978).

- [5] Basart H., Flato M., Lichnerowicz A., Sternheimer D., *Deformation theory applied to quantization and statistical mechanics*, Lett Math Phys 8, 483-494 (1984).
- [6] Berezin F.A., *General concept of quantization*, Commun. Math. Phys. 40, 153-174 (1975).
- [7] Bogachev V.I., *Measure Theory Vol. I-II*, Springer Berlin, Heidelberg (2007).
- [8] Bordermann M., Römer H., Waldmann S., *A Remark on Formal KMS States in Deformation Quantization*, Letters in Mathematical Physics, Vol. 45, 49-61 (1998).
- [9] Bratteli O., Robinson D.W., *Operator Algebras and Quantum Statistical Mechanics. Vol. II: Equilibrium States, Models in Statistical Mechanics*, Springer Berlin, Heidelberg (1981).
- [10] Dobrushin R.L., *The description of a random field by means of conditional probabilities and conditions of its regularity*, Theor. Prob. Appl. 13, 197-224 (1968).
- [11] Dobrushin R.L., *Gibbsian random fields for lattice systems with pairwise interactions*, Funct. Anal. Appl. 2, 292-301 (1968).
- [12] Dobrushin R.L., *The problem of uniqueness of a Gibbs random field and the problem of phase transition*, Funct. Anal. Appl. 2, 302-312 (1968).
- [13] Dobrushin R.L., *Prescribing a system of random variables by conditional distributions*, Th. Prob. Appl. 17, 582-600 (1970).
- [14] Drago N., van de Ven C. J. F., *Strict deformation quantization and local spin interactions*, arXiv:2210.10697 [math-ph] (2022).
- [15] Drago N., Waldmann S., *Classical KMS Functionals and Phase Transitions in Poisson Geometry*, arXiv:2107.04399 [math-ph] (2021).
- [16] Friedli S., Velenik Y., *Statistical mechanics of lattice systems: A concrete mathematical introduction*, Cambridge University Press (2017).
- [17] Gallavotti G., Pulvirenti M., *Classical KMS Condition and Tomita-Takesaki Theory*, Comm. Math. Phys. 46, Number 1, 1-9 (1976).
- [18] Gallavotti G., Verboven E., *On the classical KMS boundary condition*, Nuov Cim B, 28, 274-286 (1975).
- [19] Georgii, H.-O. *Gibbs Measures and Phase Transitions*, Berlin, New York: De Gruyter (2011).

- [20] Haag R., Hugenholtz N. M., Winnink M., *On the equilibrium states in quantum statistical mechanics*, Comm. Math. Phys. 5:215-236 (1967).
- [21] Kallenberg O., *Foundations of Modern Probability*, Springer Nature Switzerland AG (2021).
- [22] Lanford O.E., Ruelle. D., *Observables at infinity and states with short range correlations in statistical mechanics*, Comm. Math. Phys., 13: 194-215, (1969).
- [23] Moretti V., van de Ven C.J.F., *The classical limit of Schrödinger operators in the framework of Berezin quantization and spontaneous symmetry breaking as emergent phenomenon*, International Journal of Geometric Methods in Modern Physics, Vol. 19, Iss. 01 (2022).
- [24] Murro S., van de Ven C. J. F., *Injective tensor products in strict deformation quantization* Math Phys Anal Geom 25, 2 (2022).
- [25] Ruelle D., *A variational formulation of equilibrium statistical mechanics and the Gibbs phase rule*, Comm. Math. Phys. 5, 324-329 (1967).
- [26] Ruelle D., *Statistical mechanics: rigorous results*, New York: Benjamin (1969).
- [27] van de Ven C. J. F., *The classical limit and spontaneous symmetry breaking in algebraic quantum theory*, Expositiones Mathematicae Vol. 40, Iss. 3, 543-571 (2022).
- [28] van de Ven C. J. F., *KMS states and their classical limit*, arXiv:2211.01755 [math-ph] (2022).