

COMPACT KÄHLER 3-FOLDS WITH NEF ANTI-CANONICAL BUNDLE

SHIN-ICHI MATSUMURA AND XIAOJUN WU

ABSTRACT. In this paper, we prove that a non-projective compact Kähler 3-fold with nef anti-canonical bundle is, up to a finite étale cover, one of the following: a manifold with vanishing first Chern class; the product of a K3 surface and the projective line; or a projective space bundle over a 2-dimensional torus. This result extends Cao-Höring's structure theorem for projective manifolds to compact Kähler manifolds in dimension 3. For the proof, we investigate the Minimal Model Program for compact Kähler 3-folds with nef anti-canonical bundles by using the positivity of direct image sheaves, \mathbb{Q} -conic bundles, and orbifold vector bundles.

CONTENTS

1. Introduction	2
1.1. Background and the main results	2
1.2. Strategy of the proof of Theorem 1.3	3
Notation and Conventions	4
Acknowledgment	4
2. Preliminary Results	5
2.1. Bott-Chern cohomology groups on normal analytic varieties	5
2.2. Positivity of sheaves on normal analytic varieties	6
2.3. On direct image sheaves for projective morphisms	6
2.4. Hermitian metrics on orbifold vector bundles	12
3. \mathbb{Q} -Conic Bundles and the Minimal Model Program	13
3.1. \mathbb{Q} -conic bundles	13
3.2. Conic bundles	14
3.3. Minimal Model Program	18
4. Proof of the Main Results	23
4.1. On the base of MRC fibrations	23

Date: July 8, 2024, version 0.01.

2020 Mathematics Subject Classification. Primary 32J25, Secondary 53C25, 14E30.

Key words and phrases. Kähler spaces, Structure theorems, Nef anti-canonical bundles, Minimal Model Programs, \mathbb{Q} -conic bundles, Albanese maps, Orbifold structures.

4.2. The case of X being non-simply connected	24
4.3. The case of X being simply connected	26
References	36

1. INTRODUCTION

1.1. Background and the main results. In this paper, we study a structure theorem for compact Kähler manifolds with nef anti-canonical bundle, motivated by the conjecture below. This conjecture serves as a natural extension of the pioneering studies for manifolds with nef tangent bundles [DPS94] and manifolds with non-negative holomorphic bisectional curvatures [HSW81, Mok88]. The studies of manifolds with nef anti-canonical bundles have advanced through interactions with other studies of “non-negatively curved” varieties (e.g., see [Mat20, Mat22a, Mat22b, HIM22] for recent developments).

Conjecture 1.1. *Let X be a compact Kähler manifold with the nef anti-canonical bundle $-K_X$. Then, there exists a fibration $\varphi: X \rightarrow Y$ with the following:*

- $\varphi: X \rightarrow Y$ is a locally constant fibration;
- Y is a compact Kähler manifold with $c_1(Y) = 0$;
- F , which is the fiber of $\varphi: X \rightarrow Y$, is rationally connected.

The notion of locally constant fibrations (e.g., see [MW21, Definition 2.3]) is stronger than that of locally trivial fibrations. However, in this paper, readers unfamiliar with locally constant fibrations may regard them as locally trivial fibrations, except for Proposition 4.17.

Conjecture 1.1 has been addressed by the theory of holonomy groups [CDP15, DPS96] under the stronger assumption that $-K_X$ is semi-positive (i.e., it admits a smooth Hermitian metric with semi-positive curvature). Nevertheless, the nef case presents significantly greater challenges than semi-positive case, just as it was highly non-trivial to generalize the structure theorem from manifolds with non-negative holomorphic bisectional curvatures to those with nef tangent bundles. In the case where X is a projective manifold, Cao-Höring solved the conjecture in [Cao19, CH19] (see [CCM21, MW21, Wan20] for projective klt pairs), but their proofs require an ample line bundle on X ; therefore, we cannot at least directly apply their proofs to compact Kähler manifolds.

This paper aims to solve Conjecture 1.1 in the case of $\dim X = 3$ (see Theorem 1.2). Theorem 1.2 follows directly from Theorem 1.3, as detailed in Proposition 4.17. Thus, this paper focuses on proving Theorem 1.3.

Theorem 1.2. *Conjecture 1.1 is true in the case of $\dim X = 3$.*

Theorem 1.3. *Let X be a non-projective compact Kähler 3-fold with nef anti-canonical bundle. Then X admits a finite étale cover that is one of the following:*

- *a compact Kähler manifold with vanishing first Chern class;*
- *the product of a K3 surface and the projective line \mathbb{P}^1 ;*
- *the projective space bundle $\mathbb{P}(\mathcal{F})$ of a numerically flat vector bundle \mathcal{F} of rank 2 over a 2-dimensional (compact complex) torus.*

1.2. Strategy of the proof of Theorem 1.3. In this subsection, we outline the proof of Theorem 1.3. Let X be a non-projective compact Kähler 3-fold with nef anti-canonical bundle and let $\varphi: X \dashrightarrow R(X)$ be an MRC (maximally rationally connected) fibration of X (see [KoMM92, Cam92] for MRC fibrations). The candidates of X in Theorem 1.3 are determined by $\varphi: X \dashrightarrow R(X)$.

We initially verify that it suffices to consider the case of $\dim R(X) = 2$. In the case of $\dim R(X) = 0$, the manifold X is rationally connected, and hence projective. In the case of $\dim R(X) = 1$, a rationally connected fiber F of $\varphi: X \dashrightarrow R(X)$ has no non-zero holomorphic differential forms. As a result, we obtain $h^2(X, \mathcal{O}_X) = h^0(X, \Omega_X^2) = 0$ from $\dim R(X) = 1$, implying that X is projective. In the case of $\dim R(X) = 3$, the manifold X is non-uniruled; hence K_X is pseudo-effective. This follows from [BDPP13] for projective manifolds of any dimension and from [Bru06, Corollary 1.2] for compact Kähler manifolds of dimension ≤ 3 . Consequently, the nefness of $-K_X$ implies that $c_1(X) = c_1(K_X) = 0$.

We now revisit Cao-Höring's proof [CH19], which shows that a projective manifold X admits a locally constant MRC fibration. For simplicity, we suppose that $\varphi: X \dashrightarrow R(X)$ is an everywhere-defined holomorphic map onto a smooth projective variety $R(X)$. The essence of the proof involves constructing a φ -ample line bundle B on X such that the direct image sheaf $\varphi_*(pB) := \varphi_*\mathcal{O}_X(pB)$ is weakly positively curved and satisfies that $c_1(\varphi_*(pB)) = 0$ for $1 \ll p \in \mathbb{Z}$ (see Subsections 2.2 and 2.3 for details). Then, Simpson's result [Sim92] confirms that $\varphi_*(pB)$ admits a flat connection, which implies that $X \rightarrow R(X)$ is a locally constant fibration.

Now, let us return to the case where X is non-projective. A significant challenge arises in this context: X might not have a φ -ample line bundle, even if $\varphi: X \dashrightarrow R(X)$ is a holomorphic map. Our idea to overcome this challenge is to apply the Minimal Model Program (MMP) for compact Kähler 3-folds, as developed in [HP15a, HP15b, HP16]. Note that the MMP approach has been previously treated in [BP04, PS98]. By running the MMP, we can find $X \dashrightarrow X' \rightarrow Z$, where $X \dashrightarrow X'$ is a composition of divisorial contractions and flips and $\varphi: X' \rightarrow S$ is an MFS (Mori fiber space) (see Theorem 3.6 for details). An advantage of running the MMP is that $-K_{X'}$ is φ -ample by construction; thus, we can expect that Cao-Höring's proof works for $\varphi: X' \rightarrow S$.

In considering $\varphi: X' \rightarrow S$ instead of $\varphi: X \dashrightarrow R(X)$, we face the new difficulties compared to Cao-Höring’s argument: The first difficulty is that $-K_{X'}$ is not necessarily nef. To overcome this difficulty, based on an observation in [EIM23], we focus on the fact that the non-nef locus of $-K_{X'}$ is not dominant over S (see Subsection 3.3), which enables us to treat our situation as in the case where $-K_{X'}$ is nef. Thus, we can construct a φ -ample line bundle B on X' such that $\varphi_*(pB)$ is weakly positively curved and satisfies that $c_1(\varphi_*(pB)) = 0$. The second difficulty is that S may have singularities, which prevents us from obtaining a flat connection on $\varphi_*(pB)$. To overcome this difficulty, we observe that $\varphi: X' \rightarrow S$ is a toroidal \mathbb{Q} -conic bundle and $\varphi_*(pB)$ is an orbifold vector bundle on S , which enables us to obtain a flat connection.

To implement the above ideas, we actually need to divide the situation into two cases. In Subsection 4.2, we consider the case where X is not simply connected. In this case, we focus on the Albanese map $\alpha: X \rightarrow A(X)$ after taking a finite étale cover of X (cf. [NW]). Each step in the MMP contracts rational curves and $A(X)$ has no rational curve; hence we can find a morphism $\beta: S \rightarrow A(X)$. Then, comparing S to $A(X)$, we show that the MFS $\varphi: X' \rightarrow S$ is actually a conic bundle. Then, by studying conic bundles in the non-projective setting, we deduce that $\varphi: X' \rightarrow S$ is a projective space bundle and $X \dashrightarrow X'$ is isomorphic. In Subsection 4.3, we treat the remaining case where X is simply connected. In this case, by applying the theory of orbifold vector bundles, we show that after taking the base change by an appropriate finite quasi-étale cover, the MFS $\varphi: X' \rightarrow S$ is a locally constant fibration, which implies that $X \dashrightarrow X'$ is isomorphic.

Notation and Conventions. We use the terms “Cartier divisors,” “invertible sheaves,” and “line bundles” interchangeably, and adopt the additive notation for tensor products (e.g., $L + M := L \otimes M$ for line bundles L and M). Additionally, we use the terms “locally free sheaves” and “vector bundles” interchangeably, and often refer to singular Hermitian metrics simply as “metrics.” The term “fibrations” refers to a proper surjective morphism with connected fibers, the term “analytic varieties” refers to an irreducible and reduced complex analytic space, the term “Kähler spaces” refers to a normal analytic variety admitting a Kähler form (i.e., a smooth positive $(1, 1)$ -form on X with local potential).

Acknowledgment. The authors would like to thank Prof. Patrick Graf for providing a simpler proof of Lemma 4.15 than the original one in a draft of this paper. They also express their gratitude to an anonymous referee who carefully reviewed the draft, offered numerous constructive suggestions, and pointed out an error in Subsection 4.3 of the draft. The first author was partially supported by Grant-in-Aid for Scientific Research (B) #21H00976 and Fostering Joint International Research (A) #19KK0342 from Japan Society for the Promotion of Science. The second author was supported by

DFG Projekt “Singuläre hermitianische Metriken für Vektorbündel und Erweiterung kanonischer Abschnitte,” managed by Prof. Mihai Păun.

2. PRELIMINARY RESULTS

2.1. Bott-Chern cohomology groups on normal analytic varieties. In this subsection, following [BEG13], we review Bott-Chern cohomology groups and positive currents on normal analytic varieties.

Let X be a normal analytic variety. A pluriharmonic function on X can be locally written as the real part of a holomorphic function, in other words, the kernel of the $\partial\bar{\partial}$ -operator acting on the sheaf of distributions of bidegree $(0,0)$ coincides with the sheaf $\mathbb{R}\mathcal{O}_X$ of real parts of holomorphic functions (e.g., see [BEG13, Lemma 4.6.1]). Then, the *Bott-Chern cohomology group* of X is defined by

$$H_{BC}^{1,1}(X, \mathbb{C}) := H^1(X, \mathbb{R}\mathcal{O}_X).$$

The first Chern class $c_1(L) \in H_{BC}^{1,1}(X, \mathbb{C})$ of a line bundle L on X is defined by the Bott-Chern cohomology class of $(\sqrt{-1}/2\pi)\Theta_h(L)$, where $\Theta_h(L)$ denotes the Chern curvature of a smooth Hermitian metric h on L . Note that $c_1(L)$ does not depend on the choice of smooth Hermitian metrics and the first Chern class of \mathbb{Q} -Cartier divisors can also be defined by linearity. There exists the natural morphism $H_{BC}^{1,1}(X, \mathbb{C}) \rightarrow H^2(X, \mathbb{R})$ such that the first Chern class of line bundles coincides with the topological definition, where $H^2(X, \mathbb{R})$ denotes the singular cohomology of X (e.g., see [BEG13, Page 230]).

The proposition below is often used as an extension theorem for positive currents representing Bott-Chern cohomology classes (see [Dem85] for currents on analytic varieties). The lemma below is a generalization of the support theorem to analytic varieties.

Proposition 2.1 ([BEG13, Proposition 4.6.3]). *Let $\alpha \in H_{BC}^{1,1}(X, \mathbb{C})$ be a Bott-Chern cohomology class on a normal analytic variety X , and let T be a positive current on X_{reg} representing the restricted class $\alpha|_{X_{\text{reg}}} \in H_{BC}^{1,1}(X_{\text{reg}}, \mathbb{C})$. Then, the current T is uniquely extended to the positive current with local potential on X representing $\alpha \in H_{BC}^{1,1}(X, \mathbb{C})$.*

Lemma 2.2. *Let X be an analytic variety, and let T_1, T_2 be d -closed positive currents of bidimension (p, p) (without assuming that they admit local potentials). If the support of the difference $T := T_1 - T_2$ is contained in a Zariski closed subset $A \subset X$ of dimension $< p$, then we have $T = 0$.*

Proof of Lemma 2.2. The statement is local in X ; therefore, we may assume that there exists an embedding $i: X \rightarrow B \subset \mathbb{C}^N$ of X into an open set $B \subset \mathbb{C}^N$. Since the pushforward i_*T is a normal current (e.g., see [Dem, Chap. III, Section 2.C] for the definition of normal currents), the support theorem for smooth varieties shows that $i_*T = 0$, which implies that $T = 0$. \square

2.2. Positivity of sheaves on normal analytic varieties. In this subsection, following [HPS18, PT18, Mat22a], we briefly review singular Hermitian metrics on torsion-free sheaves on normal analytic varieties.

Let \mathcal{E} be a torsion-free coherent sheaf on a normal analytic variety X . A *singular Hermitian metric* h on \mathcal{E} is a possibly singular Hermitian metric on the vector bundle $\mathcal{E}|_{X_0}$ (see [HPS18, Definition 17.1] and [PT18, Definition 2.2.1] for metrics on vector bundles). Here $\mathcal{E}|_{X_0}$ is the restriction of \mathcal{E} to $X_0 := X_{\text{reg}} \cap X_{\mathcal{E}}$, where X_{reg} is the non-singular locus of X and $X_{\mathcal{E}}$ is the maximal locally free locus of \mathcal{E} . Note that $X_0 \subset X$ is a Zariski open set with $\text{codim}(X \setminus X_0) \geq 2$. For a smooth $(1, 1)$ -form θ on X with local potential, we write

$$\sqrt{-1}\Theta_h \geq \theta \otimes \text{id on } X$$

if the function $\log |e|_{h^*} - f$ is psh for any local section e of \mathcal{E}^* , where f is a local potential of θ (i.e., $\theta = \sqrt{-1}\partial\bar{\partial}f$) and h^* is the induced metric on the dual sheaf $\mathcal{E}^* := \text{Hom}(\mathcal{E}, \mathcal{O}_X)$. The plurisubharmonicity can be extended through a Zariski closed set of codimension ≥ 2 ; therefore it suffices to check the plurisubharmonicity on an open set of X_0 .

Definition 2.3. Let X be a Kähler space with a Kähler form ω_X and θ be a $(1, 1)$ -form on X with local potential. Let \mathcal{E} be a torsion-free sheaf on X

- (1) \mathcal{E} is said to be *θ -positively curved* if \mathcal{E} admits a singular Hermitian metric h such that $\sqrt{-1}\Theta_h \geq \theta \otimes \text{id}$ on X .
- (2) \mathcal{E} is said to be *θ -weakly positively curved* if \mathcal{E} admits singular Hermitian metrics $\{h_\varepsilon\}_{\varepsilon>0}$ such that $\sqrt{-1}\Theta_{h_\varepsilon} \geq (\theta - \varepsilon\omega_X) \otimes \text{id}$ on X .
- (3) \mathcal{E} is simply said to be *positively curved* or *weakly positive curved* in the case of $\theta = 0$.

When X is compact, the notion of weakly positively curved sheaves does not depend on the choice of ω_X and is stronger than the notion of pseudo-effective sheaves in the sense of [Mat23, Definition 2.1].

2.3. On direct image sheaves for projective morphisms. This subsection aims to prove Theorem 2.6. For this purpose, we prepare the following proposition:

Proposition 2.4 (cf. [CH19, 2.8 Proposition], [CCM21, Theorem 2.2 (1)]). *Let $\varphi: X \rightarrow Y$ be a fibration between (not necessarily compact) Kähler manifolds X and Y with the Kähler forms ω_X and ω_Y . Let L be a line bundle on X and θ be a $(1, 1)$ -form on Y with local potential. Assume the following conditions:*

- (a) *The non-nef locus of $-K_{X/Y}$ is not dominant over Y in the following sense: $-K_{X/Y}$ has singular Hermitian metrics $\{g_\delta\}_{\delta>0}$ such that $\sqrt{-1}\Theta_{g_\delta} \geq -\delta\omega_X$ holds on X and $\{x \in X \mid \nu(g_\delta, x) > 0\}$ is not dominant over Y , where $\nu(g_\delta, x)$ is the Lelong number of a local potential of g_δ at x ;*

- (b) L is a φ -big line bundle in the following sense: L has a singular Hermitian metric g such that $\sqrt{-1}\Theta_g + \varphi^*\omega_Y \geq \omega_X$ holds on X ;
- (c) L is $\varphi^*\theta$ -weakly positively curved in the following sense: L has singular Hermitian metrics $\{h_{\delta'}\}_{\delta'>0}$ such that $\sqrt{-1}\Theta_{h_{\delta'}} \geq \varphi^*\theta - \delta'\omega_X$ on X .

Then, we have:

- (1) The direct image sheaf $\varphi_*(-mK_{X/Y} + L)$ is $((1 - \varepsilon)\theta - \varepsilon\omega_Y)$ -positively curved for any $m \in \mathbb{Z}_+$ and $\varepsilon > 0$.
- (2) If we further assume that $\omega_Y \geq \theta$ holds, then $\varphi_*(-mK_{X/Y} + L)$ is θ -weakly positively curved. In particular, if L is a pseudo-effective line bundle (i.e., positively curved), then $\varphi_*(-mK_{X/Y} + L)$ is weakly positively curved.

Remark 2.5. Throughout this paper, for a line bundle M on X , the notation $\varphi_*(M)$ denotes the direct image sheaf $\varphi_*(\mathcal{O}_X(M))$ of the invertible sheaf $\mathcal{O}_X(M)$.

Proof. We apply the positivity of direct image sheaves [PT18, Theorem 5.1.2] (see also [HPS18]) to construct the desired singular Hermitian metrics on $\mathcal{W}_m := \varphi_*(-mK_{X/Y} + L)$. Note that [PT18, Theorem 5.1.2] is stated for projective fibrations, but in fact it is valid for Kähler fibrations (see [Wan21, Proposition 2.5, Theorem 2.6]). To this end, we consider the following decomposition of $-mK_{X/Y} + L$:

$$-mK_{X/Y} + L = kK_{X/Y} \underbrace{- (m+k)K_{X/Y}}_{\text{with } g_\delta^{m+k}} + \underbrace{L}_{\text{with } g^\varepsilon \cdot h_{\delta'}^{1-\varepsilon}}.$$

Furthermore, we define the metric G on $-(m+k)K_{X/Y} + L$ by

$$G := g_\delta^{m+k} \cdot g^\varepsilon \cdot h_{\delta'}^{1-\varepsilon}.$$

We can easily confirm that the multiplier ideal sheaf $\mathcal{J}(G^{1/k})$ satisfies that

$$(2.1) \quad \mathcal{J}(G^{1/k})|_{X_y} = \mathcal{J}(g_\delta^{1+(m/k)} \cdot g^{\varepsilon/k} \cdot h_{\delta'}^{(1-\varepsilon)/k})|_{X_y} = \mathcal{O}_{X_y}$$

for a very general fiber X_y and a sufficiently large $k \gg 1$. Indeed, since Condition (a) implies that

$$\nu(g_\delta^{1+(m/k)} \cdot g^{\varepsilon/k} \cdot h_{\delta'}^{(1-\varepsilon)/k}, x) \leq \nu(g^{1/k} \cdot h_{\delta'}^{1/k}, x) < 1 \text{ for any } x \in X_y,$$

we see that

$$\mathcal{O}_{X_y} = \mathcal{J}(G^{1/k}|_{X_y}) \subset \mathcal{J}(G^{1/k})|_{X_y} \subset \mathcal{O}_{X_y}$$

by Skoda's lemma and the restriction formula. This indicates that the support of $\mathcal{O}_X/\mathcal{J}(G^{1/k})$ is not dominant over Y . Hence, the natural inclusion

$$(2.2) \quad \varphi_*((-mK_{X/Y} + L) \otimes \mathcal{J}(G^{1/k})) \rightarrow \varphi_*(-mK_{X/Y} + L)$$

is generically surjective. Meanwhile, by the construction of metrics, we can easily see that

$$\begin{aligned}\sqrt{-1}\Theta_G &\geq -\delta(m+k)\omega_X + \varepsilon\omega_X - \varepsilon\varphi^*\omega_Y + (1-\varepsilon)\varphi^*\theta - (1-\varepsilon)\delta'\omega_X \\ &\geq (\varepsilon - \delta(m+k) - \delta')\omega_X - \varepsilon\varphi^*\omega_Y + (1-\varepsilon)\varphi^*\theta.\end{aligned}$$

For a given $\varepsilon > 0$, after taking $\delta' = \delta'(\varepsilon) > 0$ with $\delta' < (1/2)\varepsilon$, we fix a sufficiently large $k = k(\delta') = k(\varepsilon)$ satisfying (2.1). Furthermore, we take $1 \gg \delta = \delta(m, k, \varepsilon) > 0$ so that $\delta(m+k) < (1/2)\varepsilon$. Then, the right-hand side is bounded from below by $\varphi^*(-\varepsilon\omega_Y + (1-\varepsilon)\theta)$. From this curvature estimate and (2.2), we can deduce that the sheaf $\varphi_*(-mK_{X/Y} + L)$ has the desired singular Hermitian metrics in Conclusion (1), by applying the positivity of direct image sheaves (see [PT18, Theorem 5.1.2] or [Wan21, Proposition 2.5, Theorem 2.6]) to $Ge^{-\varphi^*f}$, where f is a local potential of $-\varepsilon\omega_Y + (1-\varepsilon)\theta$. Conclusion (2) directly follows from $-\varepsilon\omega_Y + (1-\varepsilon)\theta \geq -2\varepsilon\omega_Y + \theta$. \square

Theorem 2.6. *Let $\varphi: X \rightarrow Y$ be an equi-dimensional fibration between compact Kähler spaces X and Y . Let $Y_0 \subset Y$ be a Zariski open set with $\text{codim}(Y \setminus Y_0) \geq 2$ such that $X_0 := \varphi^{-1}(Y_0)$ and Y_0 are smooth and that $\varphi_0 := \varphi|_{X_0}: X_0 \rightarrow Y_0$ is a smooth fibration. Let L be a line bundle on X . Assume the following conditions:*

- (a) $-K_X$ is \mathbb{Q} -Cartier and the non-nef locus of $-K_X$ is not dominant over Y in the sense of Proposition 2.4 (a);
- (b) $-K_Y$ is \mathbb{Q} -Cartier and numerically trivial;
- (c) L is a pseudo-effective (i.e., positively curved) and φ -ample line bundle on X ;
- (d) For any $p \in \mathbb{Z}_+$ with $\varphi_*(pL) \neq 0$, the line bundle $\det(\varphi_*(pL))|_{Y_0}$ has a smooth Hermitian metric g_p such that $\eta_p := \sqrt{-1}\Theta_{g_p} \geq -\omega_Y$ holds on Y_0 for some Kähler form ω_Y .

Let r be the rank of $\varphi_*(L)$ and p be a sufficiently large integer with $p/r \in \mathbb{Z}_+$. Define the sheaf \mathcal{V}_p on Y by

$$\mathcal{V}_p := \varphi_*(pL) \otimes \left(\frac{p}{r} \det \varphi_*(L) \right)^*.$$

Then, both \mathcal{V}_p and $(\det \mathcal{V}_p)^*$ are weakly positively curved.

Remark 2.7. The determinant sheaf $\det \varphi_*(L) := (\Lambda^r \varphi_*(L))^{**}$ is a reflexive sheaf of rank 1, but not necessarily invertible when Y has singularities; therefore, precisely speaking, the notation $(p/r) \det \varphi_*(L)$ should be replaced with $((\det \varphi_*(L))^{\otimes (p/r)})^{**}$. Nevertheless, we mainly handle only $\det \varphi_*(L)|_{Y_0}$, which is a line bundle on Y_0 ; hence, this notation does not cause confusion.

We apply this theorem to the case where the sheaf $\varphi_*(pL)$ is an orbifold vector bundle. Condition (d), which may appear as a technical requirement, is automatically satisfied in this case.

Proof. For the proof, employing L instead of ample line bundles, we adopt the argument in [Cao19, CH19]. However, the original argument in [Cao19, CH19] is not so easy, and our proof requires solving several technical issues. Therefore, we write a detailed proof for the reader's convenience.

We may assume that $\varphi_*(pL)$ is locally free on Y_0 by removing the non-locally free locus of $\varphi_*(pL)$ from Y_0 . We initially prove Claim 2.8 and Claim 2.9.

Claim 2.8. $\varphi_*(pL)$ is weakly positively curved on Y for any $p \in \mathbb{Z}_+$.

Proof. The assumptions of Proposition 2.4 for $\theta = 0$ are satisfied from Conditions (a), (b), (c) of Theorem 2.6. Hence, by applying Proposition 2.4 to $\varphi_0 = \varphi|_{X_0}: X_0 \rightarrow Y_0$, we can obtain singular Hermitian metrics $\{H_\varepsilon\}_{\varepsilon>0}$ on $\varphi_*(pL)|_{Y_0}$ such that $\sqrt{-1}\Theta_{H_\varepsilon} \geq -\varepsilon\omega_Y \otimes \text{id}$ holds on Y_0 for some Kähler form ω_Y . Then, by $\text{codim}(Y \setminus Y_0) \geq 2$, the metrics H_ε can be automatically extended to Y , where we implicitly used that ω_Y is a Kähler form defined on Y (not only on Y_0). This means that $\varphi_*(pL)$ is weakly positively curved on Y . \square

Claim 2.9. Let r_p be the rank of $\varphi_*(pL)$. Then, the sheaf

$$r_p pL \otimes (\varphi^* \det \varphi_*(pL))^*$$

is weakly positively curved on X .

Proof. The basic strategy is the same as in [Cao19, Proposition 3.15], but some different arguments are required because of the lack of ample line bundles. For simplicity of the notation, we assume that $p = 1$ by replacing L with pL . Fix a Kähler form ω_Y on Y with Condition (d) of Theorem 2.6. We may assume that

$$\eta_1 + (1/r)\omega_Y = \sqrt{-1}\Theta_{g_1} + (1/r)\omega_Y$$

is a Kähler form by replacing ω_Y with $k\omega_Y$, where $r := r_1$ and $k \gg 1$. Furthermore, we take a Kähler form ω_X on X such that $\omega_X \geq \varphi^*\omega_Y$. Note that this condition is preserved by replacing ω_X and ω_Y with $k\omega_X$ and $k\omega_Y$. Since L is a φ -ample line bundle on X , we can take a smooth Hermitian metric g on L such that $\sqrt{-1}\Theta_g + \varphi^*\omega_Y \geq c\omega_X$ holds for some constant $1 \gg c > 0$ by replacing ω_X and ω_Y with $k\omega_X$ and $k\omega_Y$.

Let Z be the r -times fiber product of $\varphi: X \rightarrow Y$ with the i -th projection $\text{pr}_i: Z \rightarrow X$ and the natural morphism $\psi: Z \rightarrow Y$:

$$\begin{array}{ccc} Z := X \times_Y \cdots \times_Y X & \xrightarrow{\text{pr}_j} & X \\ \text{pr}_i \downarrow & \searrow \psi & \downarrow \varphi \\ X & \xrightarrow{\varphi} & Y. \end{array}$$

Set

$$L_r := \sum_{i=1}^r \text{pr}_i^* L \text{ and } L' := L_r \otimes (\psi^* \det \varphi_*(L))^*.$$

To apply Proposition 2.4 to

$$\psi_0 = \psi|_{Z_0}: Z_0 := \psi^{-1}(Y_0) \rightarrow Y_0 \text{ equipped with } L'|_{Z_0} \text{ and } \theta := \eta_1$$

we examine the non-nef locus of $-K_{Z/Y}$ and metrics on L' .

By Conditions (a), (b) of Proposition 2.4, we obtain singular Hermitian metrics $\{g_\delta\}_{\delta>0}$ on $-K_{X/Y}|_{X_0} = -K_{X_0/Y_0}$ such that $\sqrt{-1}\Theta_{g_\delta} \geq -\delta\omega_X$ holds on X_0 and the upper-level set $\{x \in X_0 \mid \nu(g_\delta, x) > 0\}$ of Lelong numbers is not dominant over Y_0 . Since $\psi: Z \rightarrow Y$ is a smooth fibration over Y_0 , we have

$$K_{Z_0} = \sum_{i=1}^r \text{pr}_i^* K_{X_0} \text{ on } Z_0.$$

Hence, we obtain the metric $G_\delta := \sum_{i=1}^r \text{pr}_i^* g_\delta$ on $-K_{Z_0/Y_0}$. By construction, the upper-level set $\{x \in Z_0 \mid \nu(G_\delta, x) > 0\}$ of Lelong numbers is not dominant over Y_0 and the curvature current $\sqrt{-1}\Theta_{g_\delta}$ satisfies that

$$\sqrt{-1}\Theta_{g_\delta} \geq -\delta \sum_{i=1}^r \text{pr}_i^* \omega_X \text{ on } Z_0.$$

This indicates that $-K_{Z_0/Y_0}$ satisfies Condition (a) of Proposition 2.4 for the Kähler form $\sum_{i=1}^r \text{pr}_i^* \omega_X$ on Z_0 . Consider the smooth Hermitian metric

$$G := \left(\sum_{i=1}^r \text{pr}_i^* g \right) \cdot (\psi^* g_1)^{-1} \text{ on } L' = \left(\sum_{i=1}^r \text{pr}_i^* L \right) \otimes (\psi^* \det \varphi_*(L))^*.$$

Recall that g_1 is the smooth Hermitian metric on $\det \varphi_* L|_{Y_0}$ in Condition (d). Then, we obtain that

$$\sqrt{-1}\Theta_G(L') + \sum_{i=1}^r \text{pr}_i^* \varphi^* \left(\omega_Y + \frac{1}{r} \eta_1 \right) \geq \sum_{i=1}^r \text{pr}_i^* c \omega_X \text{ on } Z_0.$$

Here we used $\psi^* = \text{pr}_i^* \circ \varphi^*$ for any $1 \leq i \leq r$. Since $\omega_Y + (1/r)\eta_1$ is a Kähler form on Y_0 , the line bundle $L'|_{Z_0}$ satisfies Condition (b) of Proposition 2.4. On the other hand, there exists the non-zero natural morphism

$$\det \varphi_*(L) \rightarrow (\varphi_*(L))^{\otimes r} \cong \psi_*(L_r) \text{ on } Y_0,$$

which shows that $h^0(Z_0, L') \neq 0$ by the definition of L' . In particular, the line bundle $L'|_{Z_0}$ satisfies Condition (c) of Proposition 2.4 for $\theta = 0$ (and $\delta' = 0$). The above arguments enable us to apply Proposition 2.4, and then we obtain singular Hermitian metrics $\{H_\varepsilon\}_{\varepsilon>0}$ on $\psi_*(L')|_{Y_0}$ such that $\sqrt{-1}\Theta_{H_\varepsilon} \geq -\varepsilon\omega_Y \otimes \text{id}$ on Y_0 .

We finally prove the desired conclusion using the metrics on L' induced by H_ε . Let us consider the natural morphism

$$(2.3) \quad \psi^*\psi_*(L') \rightarrow L' \text{ on } Z_0,$$

which is generically surjective by $h^0(Z_0, L') \neq 0$. Let G_ε be the metric on $L'|_{Z_0}$ induced by ψ^*H_ε and the above morphism. The diagonal subset Δ of the fiber product Z_0 is identified with X_0 . Note that $L'|_\Delta \cong rL \otimes (\varphi^* \det \varphi_* L)^*$ holds under this identification and (2.3) is not identically zero on Δ . By construction, the metric $G_\varepsilon|_\Delta$ on $L'|_\Delta \cong rL \otimes (\varphi^* \det \varphi_* L)^*$ is well-defined (i.e., $G_\varepsilon|_\Delta \neq \infty$) and

$$\sqrt{-1}\Theta_{G_\varepsilon|_\Delta} \geq -\varepsilon\psi^*\omega_Y|_\Delta \geq -\varepsilon\omega_X \text{ holds on } \Delta \cong X_0.$$

Note that the well-definedness follows since G_ε is constructed by the pull-back ψ^*H_ε . This curvature condition can be extended to X by $\text{codim}(X \setminus X_0) \geq 2$. Here, we used that $\varphi: X \rightarrow Y$ has equi-dimensional fibers. \square

We finish the proof of Theorem 2.6. Let p be an integer with $p/r \in \mathbb{Z}_+$. By Claim 2.9 and Condition (d), there exist singular Hermitian metrics $\{g_{\delta'}\}_{\delta' > 0}$ on $L|_{X_0}$ such that

$$\sqrt{-1}\Theta_{g_{\delta'}}(L) \geq \varphi^*\left(\frac{1}{pr_p}\eta_p\right) - \delta'\omega_X \text{ on } X_0.$$

Let us apply Proposition 2.4 to $\theta := (1/pr_p)\eta_p$. Then, since η_p is the curvature of $\det \varphi_*(pL)$, we see that

$$(2.4) \quad \varphi_*L \otimes \left(\frac{1}{pr_p} \det \varphi_*(pL)\right)^* \text{ is weakly positively curved}$$

on Y_0 (with respect to ω_Y); hence it is weakly positively curved on Y since ω_Y is defined on Y . The determinant sheaf

$$\det \varphi_*L \otimes \left(\frac{r}{pr_p} \det \varphi_*(pL)\right)^* = \det \varphi_*L - \frac{r}{pr_p} \det \varphi_*(pL)$$

is also weakly positively curved on Y . Here, we use the additive notation on the left-hand side, which is justified on Y_0 (see Remark 2.7). This implies that

$$\begin{aligned} (\det \mathcal{V}_p)^* &= -\det \varphi_*(pL) + \frac{r_p p}{r} \det \varphi_*L \\ &\geq_w -\det \varphi_*(pL) + \frac{r_p p}{r} \cdot \frac{r}{pr_p} \det \varphi_*(pL) = 0 \text{ on } Y_0 \end{aligned}$$

is weakly positively curved, where the notation \geq_w denotes the difference is weakly positively curved. On the other hand, since $(p/r)L$ is sufficiently φ -ample, the natural morphism

$$\mathcal{W}_p := \text{Sym}^p(\varphi_*L) \otimes \left(\frac{p}{r} \det \varphi_*L\right)^* \rightarrow \varphi_*(pL) \otimes \left(\frac{p}{r} \det \varphi_*L\right)^* = \mathcal{V}_p$$

is generically surjective for $p \gg 1$. The sheaf \mathcal{W}_p can be written as the p -th symmetric tensor of (2.4) of $p = 1$; therefore \mathcal{W}_p is weakly positively curved. By the above morphism, we see that \mathcal{V}_p is also weakly positively curved. \square

2.4. Hermitian metrics on orbifold vector bundles. In this subsection, following the discussions in [MM07, Wu], we review some basic facts on orbifold structures.

Let (X, ω) be a compact Kähler space with quotient singularities. The space X can be regarded as a Kähler orbifold (see [MM07, Section 5.4], [Wu, Definition 1] for the precise definition); in particular, there exists an open cover $\mathcal{U} := \{U_\alpha\}_{\alpha \in A}$ of X and a Galois quasi-étale cover $\pi_\alpha: \tilde{U}_\alpha \rightarrow \tilde{U}_\alpha/G_\alpha \cong U_\alpha$, where \tilde{U}_α is a smooth variety and G_α is a finite group. Recall that quasi-étale covers are defined as finite surjective morphisms that are étale in codimension 1. The smooth variety \tilde{U}_α is called a *local smooth ramified cover*.

A reflexive sheaf \mathcal{F} on X is called an *orbifold vector bundle* if $\{(\pi_\alpha^* \mathcal{F})^{**}\}_\alpha$ is locally free for any $\alpha \in A$. The complex orbifold, denoted by F , is determined by the quotients $\{(\pi_\alpha^* \mathcal{F})^{**}/G_\alpha\}_\alpha$ of the total space (as a vector bundle) of $(\pi_\alpha^* \mathcal{F})^{**}$, which has the orbifold morphism $F \rightarrow X$. Furthermore, the projective space orbifold bundle $p: \mathbb{P}(F) \rightarrow X$ and the tautological orbifold line bundle $\mathcal{O}_{\mathbb{P}(F)}(1)$ can be also defined. A singular Hermitian metric h on \mathcal{F} in the sense of Subsection 2.2 determines the *orbifold metric*, that is, the family $\{h_\alpha\}_\alpha$ of a (possibly singular) metric h_α (defined by the pullback of h) on $(\pi_\alpha^* \mathcal{F})^{**}$ compatible with the orbifold structure. The orbifold metric is said to be *smooth* if h_α is smooth for any α . Then, we can define the notation of positivity for \mathcal{F} as in the case of locally free sheaves, and generalize [DPS94, Theorem 1.18] to orbifold vector bundles. We emphasize that orbifold vector bundles themselves are sheaves, but all the metrics and positivity are calculated on local smooth ramified covers.

Definition 2.10. Let ω be a Kähler form on a Kähler orbifold X . An orbifold vector bundle \mathcal{F} is said to be *pseudo-effective in the strong sense* if there exists a family $\{h_\varepsilon\}_{\varepsilon > 0}$ of orbifold metrics on $\mathcal{O}_{\mathbb{P}(F)}(1)$ such that

$$\sqrt{-1} \Theta_{h_\varepsilon}(\mathcal{O}_{\mathbb{P}(F)}(1)) + \varepsilon p^* \omega \geq 0 \text{ holds in the sense of currents}$$

and the polar set $\{h_\varepsilon = +\infty\}$ is not dominant over X , where the above inequality means that it holds on each local smooth ramified cover of $\mathbb{P}(F)$. The orbifold vector bundle \mathcal{F} is said to be *nef* if h_ε in the above condition can be chosen as a smooth orbifold metric.

The orbifold vector bundle \mathcal{F} is said to be *numerically flat* (resp. *Hermitian flat*) if both \mathcal{F} , \mathcal{F}^* are nef (resp. \mathcal{F} admits an orbifold metric whose curvature vanishes over each local smooth ramified cover).

Theorem 2.11 ([Wu, Corollary 2, Theorem D]). *Let \mathcal{F} be an orbifold vector bundle on a compact Kähler orbifold X .*

(1) If \mathcal{F} is weakly positively curved as a sheaf on X (see Definition 2.3) and satisfies that $c_1(\mathcal{F}) = 0$, then \mathcal{F} is a numerically flat orbifold vector bundle.

(2) The orbifold vector bundle \mathcal{F} is numerically flat if and only if there exists a filtration of orbifold vector bundles

$$(2.5) \quad 0 =: \mathcal{F}_0 \subset \mathcal{F}_1 \subset \cdots \subset \mathcal{F}_{m-1} \subset \mathcal{F}_m := \mathcal{F}$$

such that each quotient $\mathcal{F}_k/\mathcal{F}_{k-1}$ is a Hermitian flat orbifold vector bundle.

Proof. Let $\{h_\varepsilon\}_{\varepsilon>0}$ be singular Hermitian metrics on \mathcal{F} satisfying the definition of weakly positively curved sheaves (see Definition 2.3). We can easily see that the metrics on $\mathcal{O}_{\mathbb{P}(F)}(1)$ induced by $\{h_\varepsilon\}_{\varepsilon>0}$ satisfies the definition of pseudo-effective orbifold vector bundles in the strong sense. By [Wu, Lemma 1], a Kähler form on X defines an orbifold Kähler form modulo dd^c -exact forms with continuous potential. In particular, we may assume that ω in Definition 2.10 is an orbifold Kähler form. Thus, Conclusion (1) follows from [Wu, Corollary 2]. Conclusion (2) is a direct consequence of [Wu, Theorem D]. \square

Corollary 2.12. *Let \mathcal{F} be a numerically flat orbifold vector bundle on a compact Kähler orbifold X . Assume that $\pi_1(X_{\text{reg}}) = \{\text{id}\}$ and $H^1(X_{\text{reg}}, \mathcal{O}_X) = 0$. Then, the sheaf \mathcal{F} is a trivial vector bundle on X .*

Proof. Take the filtration as in Theorem 2.11. The quotient sheaf $\mathcal{F}_k/\mathcal{F}_{k-1}$ is a Hermitian flat vector bundle on X_{reg} (but not necessarily on X); therefore, it is induced by a GL-representation of the (topological) fundamental group $\pi_1(X_{\text{reg}})$, which indicates that $\mathcal{F}_k/\mathcal{F}_{k-1}$ is a trivial vector bundle. The extension class of $0 \rightarrow \mathcal{F}_k \rightarrow \mathcal{F} \rightarrow \mathcal{F}_k/\mathcal{F}_{k-1} \rightarrow 0$ (which is an exact sequence of vector bundles on X_{reg}) is trivial by $H^1(X_{\text{reg}}, \mathcal{O}_X) = 0$. This indicates that \mathcal{F} is trivial on X_{reg} , and thus it is trivial on X by reflexivity. \square

3. \mathbb{Q} -CONIC BUNDLES AND THE MINIMAL MODEL PROGRAM

\mathbb{Q} -conic bundles naturally appear as an outcome of the MMP in our situation (see Subsection 3.3 for details). For this reason, we respectively study \mathbb{Q} -conic bundles and conic bundles in Subsections 3.1 and 3.2. Furthermore, we clarify what the nefness of anti-canonical bundles brings to the geometry of \mathbb{Q} -conic bundles.

3.1. \mathbb{Q} -conic bundles. In this subsection, following [MP08a, MP08b, Pro07], we summarize the basic properties of \mathbb{Q} -conic bundles. We first review the definition of \mathbb{Q} -conic bundles.

Definition 3.1. (1) Let X and S be normal analytic varieties. A fibration $\varphi: X \rightarrow S$ is called a *\mathbb{Q} -conic bundle* if it satisfies following conditions:

- X has terminal singularities;
- $\varphi: X \rightarrow S$ is equi-dimensional and of relative dimension 1;

- $-K_X$ is φ -ample.

Throughout this paper, except for Subsection 3.2, we consider only a \mathbb{Q} -conic bundle $\varphi: X \rightarrow S$ with $\dim X = 3$.

(2) The *discriminant divisor* Δ of a \mathbb{Q} -conic bundle $\varphi: X \rightarrow S$ is defined by the union of divisorial components of the non-smooth locus $\{s \in S \mid \varphi \text{ is not a smooth fibration at } s\}$.

(3) A \mathbb{Q} -conic bundle $\varphi: X \rightarrow S$ is said to be *toroidal* at $s \in S$ with respect to $\mu_m := \mathbb{Z}/m\mathbb{Z}$ if X is isomorphic to the quotient $\mathbb{P}^1 \times \mathbb{C}^2/\mu_m$ over a neighborhood of s . Here the μ_m -action is defined by

$$(t; z_1, z_2) \rightarrow (\varepsilon^b t; \varepsilon z_1, \varepsilon^{-1} z_2),$$

where b is an integer with $\gcd(m, b) = 1$ and ε is the primitive m -th root of unity. In this case, the singularities of X are cyclic quotient singularities of types $(1/m)(b, 1, -1)$ and $(1/m)(-b, 1, -1)$; furthermore, the singularities of the base $S \cong \mathbb{C}^2/\mu_m$ are the cyclic quotient of type A_{m-1} .

A \mathbb{Q} -conic bundle $\varphi: X \rightarrow S$ with $\dim X = 3$ can be explicitly described locally over S . The following corollary is a direct consequence of the classification [Pro18, Corollary 10.85].

Lemma 3.2 ([Pro18, Corollary 10.85]). *Let $\varphi: X \rightarrow S$ be a \mathbb{Q} -conic bundle with $\dim X = 3$ and $\Delta \subset S$ be the discriminant divisor. Then $s \notin \Delta$ if and only if $\varphi: X \rightarrow S$ is toroidal at s .*

3.2. Conic bundles. In this subsection, we review conic bundles as presented in [Sar82] and extend certain properties to the non-projective case (see Propositions 3.4 and 3.5). We derive Proposition 3.5 from Proposition 3.4. Proposition 3.4 is proved by the same argument as in [Sar82] even in the non-projective case; hence we omit the detailed proof.

Definition 3.3. A \mathbb{Q} -conic bundle $\varphi: X \rightarrow S$ is called a *conic bundle* if X and S are smooth.

Proposition 3.4. *Let $\varphi: X \rightarrow S$ be a conic bundle. Then we have:*

- (1) $-K_X$ is φ -very ample; $E := \varphi_*(-K_X)$ is a locally free sheaf of rank 3; $E = \varphi_*(-K_X)$ defines an embedding of $\varphi: X \rightarrow S$ into $p: \mathbb{P}(E) \rightarrow S$:

$$\begin{array}{ccc} X & \xrightarrow{\quad} & \mathbb{P}(E) \\ & \searrow \varphi & \swarrow p \\ & & S \end{array}$$

Furthermore, the scheme-theoretic fiber X_s at a point $s \in S$ is a (possibly reducible or non-reduced) conic on the projective plane $\mathbb{P}(E_s)(\cong \mathbb{P}^2)$.

(2) $X \subset \mathbb{P}(E)$ can be written as the zero locus of a section

$$\sigma \in H^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(2) \otimes p^*(-\det E - K_S)).$$

(3) We identify $\mathbb{P}(E)$ with $\mathbb{P}^2 \times U$ over a small open subset $U \subset S$ with a coordinate z . Then, the embedding $X \subset \mathbb{P}(E) = \mathbb{P}^2 \times U$ over U can be written as

$$X = \{([x_0 : x_1 : x_2], z) \in \mathbb{P}^2 \times U \mid \sum_{0 \leq i, j \leq 2} a_{i,j}(z)x_i x_j = 0\},$$

where $a_{i,j} \in \mathcal{O}_S(U)$. Furthermore, the discriminant divisor Δ coincides with the non-smooth locus $\{s \in S \mid \varphi \text{ is not smooth at } s\}$ and can be described as

$$\Delta = \{z \in S \mid \det[a_{i,j}(z)]_{i,j=0}^2 = 0\}.$$

(4) For a given point $s \in U$, by changing a coordinate z of U and a basis of E , we can assume that $a_{i,j} = 0$ for $i \neq j$. Furthermore, we can assume that

- $a_{i,i} \in \mathcal{O}_Y^*(U)$ for any $0 \leq i \leq 2$ when X_s is smooth;
- $a_{i,i} \in \mathcal{O}_Y^*(U)$ for any $1 \leq i \leq 2$ and $\text{mult}_s(a_{0,0}) = 1$ when X_s is reduced and reducible.

(5) The discriminant divisor Δ is normal crossing in codimension 2 in S . Furthermore, the fiber X_s of a general point $s \in \Delta$ is reduced and reducible.

(6) $c_1(\Delta) = -c_1(\det E) - 3c_1(K_S)$ holds.

Proof. Points (1) and (2) are due to [Sar82, 1.5]. Point (3) is due to [Sar82, Proposition 1.7]. Point (4) is due to [Sar82, Proposition 1.8, Point 5]. Point (6) is due to [Sar82, Definition 1.6]. The divisor Δ is reduced by [Sar82, Corollary 1.9]. The first statement of Point (5) follows from the proof of [Sar82, Proposition 1.8, Point 5] (see also [Pro18, 3.3.3]). The second statement of Point (5) follows from [Sar82, Proposition 1.8, Point 4]. \square

Proposition 3.5. *Let $\varphi: X \rightarrow S$ be a conic bundle. Then, we have*

$$\varphi_*(c_1(K_X)^2) = -4c_1(K_S) - c_1(\Delta).$$

Proof. The proposition is proved in the projective case (see [Mi83, 4.11]). We extend the discussion of [Mi83, 4.11] to the non-projective case. By Proposition 3.4 (6), the conclusion is equivalent to

$$\varphi_*(c_1(K_X)^2) = \frac{4}{3}c_1(\det E) + \frac{1}{3}c_1(\Delta).$$

We fix a smooth Hermitian metric h on $E = \varphi_*(-K_X)$ and use the same notation h to denote the induced metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$. Then, by the adjunction formula and Proposition 3.4 (2), the conclusion is equivalent to the following formula:

$$(3.1) \quad p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X]) = \frac{4}{3}c_1(\det E, \det h) + \frac{1}{3}[\Delta],$$

where $[X], [\Delta]$ are the integration currents and $c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h), c_1(\det E, \det h)$ are the Chern curvatures divided by 2π .

We first prove the desired formula on $S \setminus \Delta$. For this purpose, we summarize some formulas for the curvatures of vector bundles (e.g., see [Dem, Section 15.C, Chap. V]). Set $n = \dim S$ and $r := \text{rank } E (= 3)$. For a given $s \in S \setminus \Delta$, we take a local frame $(e_\lambda)_{\lambda=1}^r$ of E giving an orthonormal basis of E_s at $s \in S$, and then write the Chern curvature of E as

$$(3.2) \quad \Theta_h(E) = \sum_{1 \leq j, k \leq n, 1 \leq \lambda, \mu \leq r} c_{jk\lambda\mu} dz^j \wedge d\bar{z}^k \otimes e_\lambda^* \otimes e_\mu,$$

where $(z_j)_{j=1}^n$ is a local coordinate of S . Let $[x] \in \mathbb{P}(E_s)$ be the point represented by a vector $\sum_{\lambda=1}^r x_\lambda e_\lambda^* \in E_s^*$ with $\sum_{\lambda=1}^r |x_\lambda|^2 = 1$. Then, the curvature of $\mathcal{O}_{\mathbb{P}(E)}(1)$ at $[x]$ can be written as

$$(3.3) \quad \Theta_h(\mathcal{O}_{\mathbb{P}(E)}(1))_{[x]} = \sum_{1 \leq j, k \leq n, 1 \leq \lambda, \mu \leq r} c_{jk\lambda\mu} x_\lambda \bar{x}_\mu dz^j \wedge d\bar{z}^k + \sum_{1 \leq \lambda \leq r-1} d\xi_\lambda \wedge d\bar{\xi}_\lambda,$$

where $(\xi_\lambda)_{\lambda=1}^{r-1}$ is the coordinate of $\mathbb{P}(E)$ induced by unitary coordinates on the hyperplane $(\mathbb{C}x)^\perp \subset E_s^*$. We identify $\mathbb{P}(E)$ with $\mathbb{P}^{r-1} \times U$ over an open neighborhood $U \subset S$ of s , and regard the Fubini-Study form Ω on \mathbb{P}^{r-1} as the $(1, 1)$ -form on $\mathbb{P}(E)$. Then, we can easily check that

$$\frac{\sqrt{-1}}{2\pi} \Theta_h(\mathcal{O}_{\mathbb{P}(E)}(1))_{[x]} = \Omega - \frac{\sqrt{-1}}{2\pi} \frac{\langle p^* \Theta_{h^*}(E^*)x, x \rangle}{|x|^2} \quad \text{for any } [x] \in p^{-1}(s).$$

The pushforward of smooth forms can be described as a fiber integration near $s \in S \setminus \Delta$ (e.g., see [GPR94, Theorem 1.14, Proposition 2.15, Chap. II]). Hence, we obtain

$$\begin{aligned} p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X])_s &= \int_{X_s} c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \\ &= -2 \cdot \frac{\sqrt{-1}}{2\pi} \int_{X_s} \Omega \wedge \frac{\langle p^* \Theta_{h^*}(E^*)x, x \rangle}{|x|^2}, \end{aligned}$$

where X_s is the fiber of $\varphi: X \rightarrow S$ at $s \in S$.

We consider the special case of $X_s = \{x_0^2 + x_1^2 + x_2^2 = 0\}$. We can easily see that

$$\int_{X_s} \frac{x_\lambda \bar{x}_\mu}{|x|^2} \Omega = \frac{2}{3} \cdot \delta_{\lambda\mu},$$

where $\delta_{\lambda\mu}$ is the Kronecker delta. Hence, by the formula (3.2), we obtain

$$\begin{aligned} \int_{X_s} c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 &= \frac{4}{3} \frac{\sqrt{-1}}{2\pi} \sum_{\lambda} c_{jk\lambda\lambda} dz^j \wedge d\bar{z}^k \\ &= \frac{4}{3} c_1(\det E, \det h). \end{aligned}$$

The general case can be reduced to the special case. Indeed, by Proposition 3.4 (4), we may assume that $X_s = \{C_0x_0^2 + C_1x_1^2 + C_2x_2^2 = 0\}$, where C_i is a non-zero constant. Hence, we see that X_s is cohomologous to $\{x_0^2 + x_1^2 + x_2^2 = 0\}$ in $\mathbb{P}(E_s)$. By the Stokes formula, we can conclude that

$$\begin{aligned} \int_{X_s} c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 &= \int_{\{x_0^2+x_1^2+x_2^2=0\}} c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \\ &= \frac{4}{3} c_1(\det E, \det h). \end{aligned}$$

The Fubini theorem shows that

$$\int_S \varphi_* (c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2) \wedge \alpha = \int_{S \setminus \Delta} \alpha \wedge \int_{X_s} c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2$$

for any smooth form α with $\text{Supp } \alpha \Subset S \setminus \Delta$, which implies that

$$p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X]) = \frac{4}{3} c_1(\det E, \det h) \text{ on } S \setminus \Delta.$$

We finally prove the desired formula (3.1) on S . The pushforward $p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X])$ is a normal current; hence, by applying the support theorem (see Lemma 2.2), we can find $c_i \in \mathbb{R}$ such that

$$\mathbb{1}_{\Delta} p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X]) = \sum_i c_i [\Delta_i],$$

where $\{\Delta_i\}_{i \in I}$ are the irreducible components of Δ . We show that $c_i = 1/3$ for any i by regarding c_i as the generic Lelong number along Δ_i . Note that c_i is independent of the choice of metric h on E . Indeed, let $h' = he^{-\psi}$ be another smooth Hermitian metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ with some $\psi \in C^\infty(\mathbb{P}(E))$. Then, we have

$$p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h')^2 \wedge [X]) - p_*(c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)^2 \wedge [X]) = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial}F,$$

where F is a function defined by

$$F(s) = \int_{X_s} \psi \cdot (c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h') + c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h)) \text{ for } s \in S.$$

To check that the Lelong number is independent of the choice of h , it suffices to show that F is continuous over any small 1-dimensional disc passing through a general point of Δ , but this follows from the theory of cycle spaces (e.g., see [Bar78, Corollaire 1]).

To finish the proof, for a general point $s \in \Delta_i$, we construct a smooth Hermitian metric h on E (e.g., by a partition of unity) such that h is flat on a neighborhood of s . Then, by $c_1(\mathcal{O}_{\mathbb{P}(E)}(1), h) = \Omega$, it suffices to show that

$$p_*(\Omega^2 \wedge [X]) = \frac{1}{3}[\Delta_i]$$

on a neighborhood U of s . In the following, we locally approximate $X \rightarrow S$ with projective conic bundles $\{X_N \rightarrow S\}_{N=1}^\infty$, and prove the desired equality by using [Mi83, 4.11]. (The proposition itself does not seem to be directly derivable from [Mi83, 4.11], even when considering the following approximate argument.) Let $z_0x_0^2 + f_1x_1^2 + f_2x_2^2$ be a local defining function of $X \subset \mathbb{P}(E)$, where $f_1, f_2 \in \mathcal{O}^*(U)$. Let us regard $U \subset S$ as an open subset in \mathbb{P}^n . Then, we can find $g_N \in H^0(\mathbb{P}^n \times \mathbb{P}^2, \mathcal{O}(N) \boxtimes \mathcal{O}(2))$ such that

$$[X \cap \mathbb{P}(E)] = \lim_{N \rightarrow \infty} [X_N \cap \mathbb{P}(E)] \text{ over } U, \text{ where } X_N := \{g_N = 0\}$$

by using the polynomial approximation of f_1, f_2 (e.g., we can use the Taylor expansion). By the Bertini theorem, a general member in $H^0(\mathbb{P}^n \times \mathbb{P}^2, \mathcal{O}(N) \boxtimes \mathcal{O}(2))$ determines a conic bundle over \mathbb{P}^n . Thus, by replacing g_N with the general member, we may assume that $X_N = \{g_N = 0\} \rightarrow \mathbb{P}^n$ is a conic bundle with discriminant divisor Δ_N . Since $X_N \rightarrow \mathbb{P}^n$ is a projective conic bundle, we have

$$p_*(\Omega^2 \wedge [X_N \cap \mathbb{P}(E)]) = \frac{1}{3}[\Delta_N] \text{ over } U$$

by [Mi83, 4.11]. Consequently, by taking the limit as $N \rightarrow \infty$ in the space of currents, we obtain the desired conclusion. Indeed, the pushforward $p_*(\Omega^2 \wedge \bullet)$ defines a continuous map from the space of currents on $U \times \mathbb{P}^2$ to that of U , which implies that the left-hand side converges to $p_*(\Omega^2 \wedge [X \cap \mathbb{P}(E)])$. On the other hand, the right-hand side converges to $[\Delta_i]$ by construction. \square

3.3. Minimal Model Program. In this subsection, we review the MMP for Kähler 3-folds developed in [HP15a, HP15b, HP16], and observe what the MMP brings to Conjecture 1.1 (see Corollary 3.11).

Theorem 3.6 ([HP15b]). *Let X be a \mathbb{Q} -factorial compact Kähler space of dimension 3 with terminal singularities. Assume that $\dim R(X) = 2$, where $R(X)$ is the base of an MRC fibration $X \dashrightarrow R(X)$ of X . Then, we have:*

(1) X is bimeromorphic to an MFS (Mori fiber space); more precisely, there exist

a bimeromorphic map $\pi: X \dashrightarrow X'$ and an MFS $\varphi: X' \rightarrow S$ such that

(a) $X \dashrightarrow X'$ is obtained from the composition of divisorial contractions and flips;

- (b) X' is a \mathbb{Q} -factorial compact Kähler space with terminal singularities;
- (c) S is a \mathbb{Q} -factorial compact Kähler space of dimension 2 with klt singularities;
- (d) S is non-uniruled and K_S is pseudo-effective;
- (e) $-K_{X'}$ is φ -ample and the relative Picard number $\rho(X'/S)$ is 1;
- (f) $\varphi: X' \rightarrow S$ is equi-dimensional and of relative dimension 1.

(2) The outcome $X \dashrightarrow X' \rightarrow S$ of the MMP factors through the Albanese map $\alpha: X \rightarrow A(X)$, that is, there exists the morphism $\beta: S \rightarrow A(X)$ with the diagram:

$$\begin{array}{ccc} X & \xrightarrow{\pi} & X' \\ \alpha \downarrow & & \downarrow \varphi \\ A(X) & \xleftarrow{\beta} & S. \end{array}$$

Proof. By running the MMP in [HP15b] for the initial variety X , we can find a birational map $\pi: X \dashrightarrow X'$ such that either $K_{X'}$ is nef or there exists an MFS $\varphi: X' \rightarrow S$.

We can easily exclude the case where $K_{X'}$ is nef. Indeed, by [Bru06, Corollary 1.2], we see that Y is non-uniruled if and only if K_Y is pseudo-effective, where Y is a compact Kähler space of dimension ≤ 3 with terminal singularities. Hence, by noting that X' has terminal singularities, if $K_{X'}$ is nef, the variety X' is non-uniruled, which contradicts $\dim R(X) = 2$. Hence, except for (c) and (d), the other properties follow from [HP15b].

The outcome $X \dashrightarrow X' \rightarrow S$ gives one of MRC fibrations of X , which implies that S is a non-uniruled surface by $\dim R(X) = 2$. The non-uniruledness shows that K_S is pseudo-effective. Indeed, for a minimal resolution $\pi: \bar{S} \rightarrow S$ of S , we have $\pi^*K_S = K_{\bar{S}} + E$ for some effective exceptional \mathbb{Q} -divisor E . Since \bar{S} is non-uniruled, we see that $K_{\bar{S}}$ is pseudo-effective; hence so is K_S .

As in the case of the projective case, all the steps of the MMP (i.e., divisorial contractions, flips, MFSs) are obtained from contractions of rational curves. Thus, since the torus $A(X)$ has no rational curve, the outcome $X \dashrightarrow X' \rightarrow S$ factors through the Albanese map $X \rightarrow A(X)$. \square

Let us briefly examine the positivity of $-K_{X'}$. Suppose that we start from X with the nef anti-canonical bundle $-K_X$, and obtain a birational map $\pi: X \dashrightarrow X'$ in Theorem 3.6. We might expect that $-K_{X'}$ is still nef “outside the exceptional locus” of $\pi: X \dashrightarrow X'$, but proving this is not straightforward. In fact, after taking a smooth form $T_\varepsilon \in c_1(-K_X)$ such that $T_\varepsilon \geq -\varepsilon\omega$, we can obtain $\pi_*T_\varepsilon \geq -\varepsilon\pi_*\omega$ on X' , where ω is a Kähler form on X . However, it is unclear how the current $\pi_*\omega$ relates to a Kähler form on X' . For this reason, we prepare Lemma 3.8 to compare Kähler forms on X to those on X' .

Setting 3.7. Before stating Lemma 3.8, we fix the notation. Assume that X in Theorem 3.6 is smooth and $-K_X$ is nef. The bimeromorphic map $X \dashrightarrow X'$ is decomposed as follows:

$$(3.4) \quad X =: X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_N := X',$$

where each bimeromorphic map $\pi_i: X_i \dashrightarrow X_{i+1}$ is a divisorial contraction or flip. Let \bar{X} be a compact Kähler manifold with a bimeromorphic morphism $p_i: \bar{X} \rightarrow X_i$ that resolves the indeterminacy locus of π_i (when π_i is a flip). Depending on whether π_i is a divisorial contraction or flip, we obtain the following diagrams:

Note that Z_i and X_i are Kähler spaces by [HP15b, Theorem 3.15]. The varieties X_i are biholomorphic to each other on a non-empty Zariski open set. This open set, regarded as an open subset of all X_i 's, is called a *biholomorphic locus*. Note that the complement of the biholomorphic locus in $X_N = X'$ is of codimension ≥ 2 .

Lemma 3.8. *We consider Setting 3.7. Then, for any $0 \leq i \leq N$, there exists a Kähler form ω_i on X_i such that the Bott-Chern class*

$$\{p_{0*}(p_{i+1}^*\omega_{i+1} - p_i^*\omega_i)\} + O(E, K_X)$$

is represented by a positive current that is smooth on the biholomorphic locus of $X \dashrightarrow X'$, where $O(E, K_X)$ is a linear combination of the first Chern classes of K_X and the exceptional divisors $p_{0}p_i^*E_i$. Here E_i denotes the exceptional divisor of $\pi_i: X_i \rightarrow X_{i+1}$ (when it is a divisorial contraction). In particular, the Bott-Chern cohomology class $\{p_{0*}p_i^*\omega_i - \omega_0\} + O(E, K_X)$ is represented by a positive current that is smooth on the biholomorphic locus of $X \dashrightarrow X'$.*

Remark 3.9. The proof of Lemma 3.8 works for Kähler spaces of any dimension, provided that divisorial contractions or flips $\pi_i: X_i \dashrightarrow X_{i+1}$ exist. In the projective case, Mori's cone theorem and Kawamata's base-point-free theorem contract the extremal ray whose intersection number with the canonical divisor is strictly negative. This contraction is given by the Stein factorization of the Iitaka fibration of a suitably chosen line bundle. As a result, the outcomes of MMP are always projective varieties. It is conjectured that a Kähler variant of Mori's cone theorem would similarly contract Kähler spaces to Kähler spaces, which is verified up to dimension 3 (see [HP16]).

Proof. Fix a Kähler form ω_N on X_N . Assuming ω_{i+1} has been constructed, we proceed to inductively construct ω_i .

We initially consider the case where $\pi_i: X_i \rightarrow X_{i+1}$ is a divisorial contraction with the exceptional divisor E_i . Since $-E_i$ is π_i -ample, we can take a smooth form $\theta_i \in c_1(E_i)$ such that $\omega_i := \pi_i^* \omega_{i+1} - \varepsilon \theta_i$ is a Kähler form on X_i for $1 \gg \varepsilon > 0$. Then, we see that

$$p_{0*}(p_{i+1}^* \omega_{i+1} - p_i^* \omega_i) = p_{0*} p_i^*(\pi_i^* \omega_{i+1} - \omega_i) = \varepsilon p_{0*} p_i^* \theta_i.$$

The current $p_{0*} p_i^* \theta_i$ represents $c_1(p_{0*} p_i^* E_i) = O(E)$, which finishes the proof.

We now consider the case where $\pi_i: X_i \dashrightarrow X_{i+1}$ is a flip. Fix a Kähler form ω_{Z_i} (up to a rescaling) on Z_i such that $\omega_{i+1} \geq q_{i+}^* \omega_{Z_i}$. Furthermore, since $-K_{X_i}$ is q_i -ample, we can take a smooth form $\eta_i \in c_1(K_{X_i})$ such that $\omega_i := q_i^* \omega_{Z_i} - \varepsilon \eta_i$ is a Kähler form on X_i for $1 \gg \varepsilon > 0$. Then, we can easily see that

$$\begin{aligned} p_{0*}(p_{i+1}^* \omega_{i+1} - p_i^* \omega_i) &= p_{0*}(p_{i+1}^* \omega_{i+1} - p_i^*(q_i^* \omega_{Z_i} - \varepsilon \eta_i)) \\ &= p_{0*}(p_{i+1}^*(\omega_{i+1} - q_{i+}^* \omega_{Z_i}) + \varepsilon p_i^* \eta_i). \end{aligned}$$

The current $\varepsilon p_{0*} p_i^* \eta_i$ represents $\varepsilon c_1(p_{0*} p_i^* K_{X_i}) = O(K_X, E)$ and $p_{0*}(p_{i+1}^*(\omega_{i+1} - q_{i+}^* \omega_{Z_i}))$ is smooth on the biholomorphic locus, which finishes the proof. \square

Later, we will show that $X \dashrightarrow X'$ described in Theorem 3.6 is actually an isomorphism when $-K_X$ is nef. For this purpose, we need the following corollary on the intersection number. Note that when X_i is smooth, some in the following proposition can be simplified by saying that $-K_{X_i}$ is modified nef. However, we do not use this terminology in this paper due to the ambiguity of the definition of modified nefness on singular spaces.

Proposition 3.10. *We consider Setting 3.7. Let ω_i be a Kähler form on X_i . Then, we have:*

(1) *There exists a positive current $T_\varepsilon \in c_1(-K_{X_i}) + \varepsilon \{\omega_i\}$ such that T_ε is smooth on the biholomorphic locus of $X \dashrightarrow X'$.*

(2) *For a surface $V \subset X_i$, we have*

$$(c_1(K_{X_i})^2 \cdot \{\omega_i\}) \geq 0 \text{ and } (c_1(-K_{X_i}) \cdot c_1(V) \cdot \{\omega_i\}) \geq 0.$$

Proof. By the nefness of $-K_X$, there exists a smooth (semi-)positive form $S_\varepsilon \in c_1(-K_X) + \varepsilon \{\omega_0\}$. By Lemma 3.8, we can find a positive current $P \in \{p_{0*} p_i^* \omega_i - \omega_0\} + O(E, K_X)$ such that P is smooth on the biholomorphic locus of $X \dashrightarrow X'$. The current $p_{i*} p_0^*(S_\varepsilon + \varepsilon P)$ (defined on X_i) is positive and represents

$$p_{i*} p_0^*(c_1(-K_X) + \varepsilon p_{0*} p_i^* \{\omega_i\} + \varepsilon O(E, K_X)).$$

The above class coincides with $c_1(-K_{X_i}) + \varepsilon \{\omega_i\} + \varepsilon O(K_{X_i})$ on the biholomorphic locus, which is a Bott-Chern cohomology class on X_i . Hence, by Proposition, 2.1 the positive current $p_{i*} p_0^*(S_\varepsilon + \varepsilon P)$ actually represents $c_1(-K_{X_i}) + \varepsilon \{\omega_i\} + \varepsilon O(K_{X_i})$. Then,

Conclusion (1) easily follows since the $O(K_{X_i})$ -part can be absorbed into the Kähler class and $p_{i*}p_0^*(S_\varepsilon + \varepsilon P)$ is smooth on the biholomorphic locus.

To prove Conclusion (2), we first remark that $c_1(V)$ is well-defined since X_i is \mathbb{Q} -factorial. Let Q be a $(2, 2)$ -class defined by either $c_1(V) \cdot \{\omega_i\}$ or $c_1(-K_{X_i}) \cdot \{\omega_i\}$. In any case, noting that $p_i^*(-K_{X_i})$ and $p_i^*c_1(V)$ are pseudo-effective, the pull-back p_i^*Q can be represented by a positive $(2, 2)$ -current. Hence, since S_ε is a smooth (semi-)positive $(1, 1)$ -form, the intersection number $(\{p_0^*S_\varepsilon\} \cdot p_i^*Q)$ is non-negative. Then, Conclusion (2) follows from

$$\begin{aligned} (c_1(-K_{X_i}) \cdot Q) &= \lim_{\varepsilon \rightarrow 0} ((c_1(-K_{X_i}) + \varepsilon\{\omega_i\} + \varepsilon O(K_{X_i})) \cdot Q) \\ &= \lim_{\varepsilon \rightarrow 0} (\{p_{i*}p_0^*(S_\varepsilon + \varepsilon P)\} \cdot Q) \\ &= \lim_{\varepsilon \rightarrow 0} (\{p_0^*(S_\varepsilon + \varepsilon P)\} \cdot p_i^*Q) \\ &= \lim_{\varepsilon \rightarrow 0} (\{p_0^*S_\varepsilon\} \cdot p_i^*Q) \geq 0. \end{aligned}$$

□

Corollary 3.11. *We consider Setting 3.7 and the MFS $\varphi: X_N = X' \rightarrow S$ in Theorem 3.6. Then, we have:*

(1) $-4c_1(K_S) - c_1(\Delta)$ is pseudo-effective, where Δ is the discriminant divisor of the MFS $\varphi: X \rightarrow S$ (which is a \mathbb{Q} -conic bundle).

(2) $\Delta = 0$, $c_1(K_S) = 0$, and $\kappa(S) = 0$ hold; in particular, $\varphi: X' \rightarrow S$ is toroidal over S and S has only rational double points. Furthermore, when S is smooth, the variety X' is automatically smooth and $\varphi: X' \rightarrow S$ is a (locally trivial) \mathbb{P}^1 -bundle.

Proof. In the proof, we take Zariski open subsets $S_0 \subset S$ with $\text{codim}(S \setminus S_0) \geq 2$. To maintain clarity in notation, we consistently refer to these subsets as S_0 , even though they may vary. Since $-K_{X'}$ is φ -ample by Theorem 3.6 (d), we can take a Kähler form $\omega' \in c_1(-K_{X'}) + \{\varphi^*\omega_S\}$, where ω_S is a fixed Kähler form on S . By Proposition 3.10 (1), there exists a positive current

$$T_\varepsilon \in -c_1(K_{X'}) + \varepsilon\{\omega'\} = -(1 + \varepsilon)c_1(K_{X'}) + \varepsilon\{\varphi^*\omega_S\}$$

such that T_ε is smooth on $\varphi^{-1}(S_0)$. Since X' is smooth in codimension 2, there exists S_0 such that $\varphi|_{X_0}: X_0 := \varphi^{-1}(S_0) \rightarrow S_0$ is a conic bundle. Note that the Bedford-Taylor product T_ε^2 is defined on X'_{reg} . By Proposition 3.5, the pushforward $\varphi_*(T_\varepsilon^2)$ defined on S_0 is a positive current representing the following class on S_0 :

$$\begin{aligned} &\varphi_*((-1 + \varepsilon)c_1(K_{X'}) + \varepsilon\{\varphi^*\omega_S\})^2 \\ (3.5) \quad &= -(1 + \varepsilon)^2(4c_1(K_S) + c_1(\Delta)) - 2\varepsilon(1 + \varepsilon)\varphi_*c_1(K_{X'}) \cdot \{\omega_S\} + \varepsilon^2\varphi_*\{\varphi^*\omega_S^2\} \\ &= -(1 + \varepsilon)^2(4c_1(K_S) + c_1(\Delta)) + 4\varepsilon(1 + \varepsilon) \cdot \{\omega_S\}. \end{aligned}$$

Here, we used that $\varphi_*c_1(K_{X'}) = -2$ and $\varphi_*\{\varphi^*\omega_S^2\} = 0$ hold on S_0 . Proposition 2.1 shows that $\varphi_*(T_\varepsilon^2)$ is actually a positive current on S representing the Bott-Chern cohomology class of the right-hand side. Since the mass measure of $\varphi_*(T_\varepsilon^2)$ is uniformly bounded, we may assume that $\varphi_*(T_\varepsilon^2)$ has the weak limit by weak compactness (see [Dem, (1.14), (1.23) Propositions, Chapter III]). Indeed, the total mass of $\varphi_*(T_\varepsilon^2)$ with respect to ω_S is the intersection number of (3.5) with $\{\omega_S\}$, which is uniformly bounded in $\varepsilon \in [0, 1)$. Note that S has singularities, but we can apply the weak compactness after taking a resolution of singularities of S . Then, the weak limit of $\varphi_*(T_\varepsilon^2)$ is a positive current representing the class

$$\lim_{\varepsilon \rightarrow 0} -(1 + \varepsilon)^2(4c_1(K_S) + c_1(\Delta)) + 4\varepsilon(1 + \varepsilon) \cdot \{\omega_S\} = -4c_1(K_S) - c_1(\Delta).$$

This indicates that $-4c_1(K_S) - c_1(\Delta)$ is pseudo-effective.

Theorem 3.6 (d) shows that $\Delta = 0$ and $c_1(K_S) = 0$ hold. The \mathbb{Q} -conic bundle $\varphi: X' \rightarrow S$ is toroidal by Lemma 3.2 and the singularities of S are rational double points. Therefore, for the minimal resolution $h: \tilde{S} \rightarrow S$, we have $K_{\tilde{S}} = h^*K_S$, which implies that $\kappa(S) = \kappa(\tilde{S}) = 0$. The latter statement of Conclusion (2) is a special case of [AR14, Theorem 5]. \square

Remark 3.12. In the proof, if X' is smooth, the Bedford-Taylor product T_ε^2 can be defined on X' as a positive current representing $(c_1(-K_i) + \varepsilon\{\omega_i\})^2$. This is expected to be true even when X' has singularities, which gives a more direct proof of Proposition 3.10, but we could not prove this expectation. We avoided this difficulty by considering the $(1, 1)$ -current $\varphi_*(T_\varepsilon^2)$ (instead of T_ε^2).

4. PROOF OF THE MAIN RESULTS

This section is devoted to the proof of Theorem 1.3. Throughout this section, let X be a non-projective compact Kähler 3-fold with nef anti-canonical bundle. As explained in Subsection 1.2, we may assume that $\dim R(X) = 2$ for Theorem 1.3. Furthermore, by replacing X with a finite étale cover, we may assume that $\pi_1(X) \cong \mathbb{Z}^{\oplus 2q}$, where q is the irregularity of X . Here we used the fact that $\pi_1(X)$ is almost abelian (see [Pău97, Théorème 2] for the proof based on Monge-Ampère equations and see [Pău17, Theorem 1.4], [Cam95, Theorem 2.2] for the proof based on variations of Kähler-Einstein metrics). We consider the case of $q \neq 0$ in Subsection 4.2 and the case of $q = 0$ in Subsection 4.3.

4.1. On the base of MRC fibrations. Before starting the proof of Theorem 1.3, we determine the smooth minimal base of MRC fibrations of X .

Proposition 4.1. *Let X be a non-projective compact Kähler 3-fold with nef anti-canonical bundle such that $\dim R(X) = 2$, where $R(X)$ denotes the smooth minimal base of MRC fibrations of X . Then, up to a finite étale cover of X , the base $R(X)$ is either a torus or a K3 surface. In particular, the augmented irregularity of X is 2 or 0.*

Proof. By replacing X with a finite étale cover, we assume that $\pi_1(X) \cong \mathbb{Z}^{\oplus 2q}$. In particular, we have $q = \dim A(X)$, where q is the irregularity of X .

Consider the same situation as in Theorem 3.6. Since $X \dashrightarrow X' \rightarrow S$ in Theorem 3.6 is an MRC fibration of X , we obtain the minimal resolution $\gamma: R(X) \rightarrow S$ by noting that $R(X)$ is the smooth minimal base of MRC fibrations and the base of MRC fibrations is uniquely determined up to bimeromorphic models. By Corollary 3.11, the variety S has cyclic quotient singularities of type A_{m-1} (see Definition 3.1), and thus, we have

$$c_1(K_{R(X)}) = c_1(\gamma^* K_S) = 0.$$

The classification of surfaces (e.g., see [BHPV04]) implies that $R(X)$ is a (2-dimensional) torus or a K3 surface. Note that the possibilities of Enriques surfaces and hyperelliptic surfaces are excluded by the non-projectivity of $R(X)$.

We consider the case of $q = \dim A(X) > 0$. The Albanese map $\alpha: X \rightarrow A(X)$ is surjective by [Pău17, Theorem 1.4], and thus, so is $\beta: S \rightarrow A(X)$ in Theorem 3.6. By pulling back 1-forms on $A(X)$ using $\gamma: R(X) \rightarrow S$ and $\beta: S \rightarrow A(X)$, we can see that $R(X)$ is a torus, which indicates that $q \geq 2$. In the case of $q = \dim A(X) = 3$, the manifold X is non-uniruled, which contradicts $\dim R(X) = 2$. Thus, we can conclude that $q = 2$ and $R(X)$ is a torus.

We consider the remaining case of $q = \dim A(X) = 0$. In this case, the manifold X is simply connected, and thus, so is $R(X)$ by $\pi_1(X) \cong \pi_1(R(X))$ (see [Kol93, Theorem 5.2] and [BC15, Theorem 4.1]), which indicates that $R(X)$ is a K3 surface. \square

4.2. The case of X being non-simply connected. In this subsection, we prove Theorem 1.3 under assuming that $\pi_1(X) \cong \mathbb{Z}^{\oplus 2q}$, $q \neq 0$, and $\dim R(X) = 2$.

Theorem 4.2. *Consider the same situation as in the beginning of Subsection 4.2. Then, up to a finite étale cover of X , there exists a numerically flat vector bundle \mathcal{F} on $A(X)$ such that X is isomorphic to the projective space bundle $\mathbb{P}(\mathcal{F})$ over $A(X)$.*

Proof. We initially show that $\beta: S \rightarrow A(X)$ in Theorem 3.6 is actually isomorphic. Note that $A(X)$ is a 2-dimensional torus by Proposition 4.1 and $\beta: S \rightarrow A(X)$ is surjective by [Pău17, Theorem 1.4]. Then, we have $K_S = \beta^* K_{A(X)} + E$ for some effective divisor E supported in the ramified locus of $\beta: S \rightarrow A(X)$. By $c_1(K_{A(X)}) = 0$ and $c_1(K_S) = 0$ (see Corollary 3.11), we deduce that $E = 0$, which indicates that $\beta: S \rightarrow A(X)$ is étale and S is a torus. We can see that $\beta: S \rightarrow A(X)$ is isomorphic by the universal property of Albanese maps.

By Corollary 3.11 (2), the MFS $\varphi: X' \rightarrow S \cong A(X)$ in Theorem 3.6 is a \mathbb{P}^1 -bundle; hence, up to a finite étale cover of X' , there exists a vector bundle \mathcal{F} on $A(X)$ such that $X' \cong \mathbb{P}(\mathcal{F})$ and $\det(\mathcal{F}) = \mathcal{O}_{A(X)}$ by [CP91, Lemma 7.4] (see [Ele82] for more information on Brauer groups). We show that \mathcal{F} can be chosen to be numerically flat. We emphasize

that the following discussion works only when X' and S are smooth. By the formula

$$-K_{X'} = \mathcal{O}_{\mathbb{P}(\mathcal{F})}(2) - \varphi^* \det \mathcal{F} = \mathcal{O}_{\mathbb{P}(\mathcal{F})}(2),$$

it suffices to show that $-K_{X'}$ is nef. By applying the regularization theorem [Dem92, Proposition 3.7] to the current in Proposition 3.10 (1), we obtain positive currents $T_\varepsilon \in c_1(-K_{X'}) + \varepsilon\{\omega_{X'}\}$ with analytic singularities such that the singular locus of T_ε is not dominant over S . By the proof of Corollary 3.11, we have $\varphi_*(c_1(-K_{X'})^2) = 0$ and $\lim_{\varepsilon \rightarrow 0} \varphi_*(T_\varepsilon^2) = 0$; hence, the Lelong number of T_ε uniformly converges to 0 on X' by [Wu22b, Lemma 15]. Thus, we can conclude that $-K_{X'}$ is nef by the regularization with smooth forms (see [Dem92, Theorem 1.1], cf. the end of proof of [Wu22b, Theorem 4] and [Wu22b, Corollary 6]).

We finally show that $X \dashrightarrow X'$ is actually an isomorphism. To achieve this, we focus on the final step $X_{N-1} \dashrightarrow X_N = X'$ of the MMP (see (3.4)) and divide the proof into three subsequent claims.

Claim 4.3. $X_{N-1} \dashrightarrow X_N = X'$ cannot be a flip.

Proof. Every rational curve $R \subset X'$ is contained in a fiber of $\varphi: X' \cong \mathbb{P}(\mathcal{F}) \rightarrow S \cong A(X)$. Consequently, the intersection number $(R \cdot c_1(-K_{X'})) = \deg(-K_{\mathbb{P}^1})$ is positive. This implies that X' has no $K_{X'}$ -positive rational curves; therefore $X_{N-1} \dashrightarrow X_N = X'$ cannot be a flip since a flip generates a $K_{X'}$ -positive rational curve. \square

We consider the possible divisorial contraction $\pi := \pi_{N-1}: X_{N-1} \rightarrow X_N = X'$ with the exceptional divisor E .

Claim 4.4. $X_{N-1} \dashrightarrow X_N = X'$ cannot be a divisorial contraction contracting E to the one point.

Proof. Since X_{N-1} has terminal singularities, we have

$$(4.1) \quad -K_{X_{N-1}} = \pi^*(-K_{X_N}) - aE \quad \text{for some } a > 0.$$

We can confirm that the equality $c_1(K_{X_N})^2 = 0$ is satisfied. Indeed, the condition $c_1(K_{X_N})^2 = 0$ remains invariant under finite étale covers, which permits us to assume that $X' = X_N = \mathbb{P}(\mathcal{F})$ with a numerically flat vector bundle \mathcal{F} . Then, the desired equality $c_1(K_{X_N})^2 = 0$ can be obtained from $K_{X_N} = \mathcal{O}_{\mathbb{P}(\mathcal{F})}(-2)$ and

$$c_1(\mathcal{O}_{\mathbb{P}(\mathcal{F})}(1))^2 = \varphi^*(c_1(\mathcal{F}))c_1(\mathcal{O}_{\mathbb{P}(\mathcal{F})}(1)) - \varphi^*(c_2(\mathcal{F})) = 0.$$

By noting that E is contracted to the one point, we can see that $c_1(-K_{X_{N-1}})^2 = a^2c_1(E)^2$ by $c_1(K_{X_N})^2 = 0$. Then, since $\mathcal{O}_E(-E)$ is ample, we obtain

$$(c_1(K_{X_{N-1}})^2 \cdot \{\omega_{X_{N-1}}\}) = a^2(c_1(E)|_E \cdot \{\omega_{N-1}\}|_E) < 0.$$

This contradicts Proposition 3.10 (2). \square

Claim 4.5. $X_{N-1} \dashrightarrow X_N = X'$ cannot be a divisorial contraction contracting E to the curve $C := \pi(E)$.

Proof. Assume that there exists a surface $V \subset X_N$ such that $K_{X_N}|_V$ is numerically trivial and $\bar{V} \cap E$ is an effective curve, where $\bar{V} \subset X_{N-1}$ is the strict transform of V . Then, by (4.1), we can see that

$$(c_1(-K_{X_{N-1}}) \cdot c_1(\bar{V}) \cdot \omega_{N-1}) = a(c_1(-E) \cdot c_1(\bar{V}) \cdot \omega_{N-1}) < 0.$$

This contradicts Proposition 3.10 (2).

To find such a surface V , we show that C is a fiber of $\varphi: X_N = X' \rightarrow A(X)$, which follows since $X \rightarrow A(X)$ is a smooth fibration outside a Zariski closed set of $A(X)$ of codimension ≥ 2 by [Cao13, Proposition 1.6]. Indeed, otherwise, the image $\varphi(C)$ is a curve in $A(X)$. Then, a fiber of $X_{N-1} \rightarrow X' \rightarrow A(X)$ at a general point $p \in \varphi(C)$ has at least two irreducible components: the strict transform of the fiber F of $\varphi: X' \rightarrow A(X)$ at p and the inverse image of $F \cap C$. This contradicts that $X \rightarrow A(X)$ is smooth in codimension 1. Meanwhile, by [DPS94, Theorem 1.18], the numerically flat vector bundle \mathcal{F} is constructed from a GL-representation of $\pi_1(S)$. Furthermore, since $\pi_1(S)$ is abelian, this representation is the direct sum of 1-dimensional representations. Hence, there exist Hermitian flat line bundles L_1, L_2 on S such that

$$0 \rightarrow L_1 \rightarrow \mathcal{F} \rightarrow L_2 \rightarrow 0.$$

Define the surface $V \subset X' = X_N$ by the image of $\mathbb{P}(L_2) \subset \mathbb{P}(\mathcal{F})$ via $\mathbb{P}(\mathcal{F}) \rightarrow X' = X_N$. Then, we can easily see that $K_{X_N}|_V$ is numerically trivial and the intersection $\bar{V} \cap E$ is the fiber $\pi: X_{N-1} \rightarrow X' = X_N$ at the non-empty 0-dimensional varieties $V \cap C$. \square

This finishes the proof of Theorem 1.3 under assuming that $\pi_1(X) \cong \mathbb{Z}^{\oplus 2q}$, $q \neq 0$, and $\dim R(X) = 2$. \square

Remark 4.6. The final case (where E is contracted to the curve C) can be actually excluded by another approach without using [Cao13, Proposition 1.6] (see the final step of the proof of Theorem 4.8). However, this approach explained in Theorem 4.8 is quite complex, and thus we provide a more straightforward proof here using [Cao13, Proposition 1.6].

4.3. The case of X being simply connected. In this subsection, we prove Theorem 1.3 under assuming that $\pi_1(X) = \{\text{id}\}$ and $\dim R(X) = 2$. Compared to Subsection 4.2, a notable challenge is that S may have singularities, which prevents us from employing the same argument as in Theorem 4.2. The following example helps us to understand this difficulty.

Example 4.7. For a 2-dimensional torus A , we consider

$$X' := (\mathbb{P}^1 \times A)/\mu_2 \rightarrow S = A/\mu_2,$$

where $\mu_2 := \mathbb{Z}/2\mathbb{Z}$ acts on $\mathbb{P}^1 \times A$ by $(-1) \cdot (t, z_1, z_2) = (-t, -z_1, -z_2)$. Both S and X' are simply connected and $\varphi: X' \rightarrow S$ is a \mathbb{Q} -conic bundle such that $-K_{X'}$ is nef. Nevertheless, the fibration $\varphi: X' \rightarrow S$ is not even locally trivial.

In fact, the above example never appears as an outcome of the MMP of X with nef anti-canonical bundle. More precisely, the following theorem can be proved:

Theorem 4.8. *Consider the same situation as in the beginning of Subsection 4.3. Then, the manifold X is isomorphic to the product of a K3 surface and the projective line \mathbb{P}^1 .*

Proof. By [Cam04, Corollary 6.7], there exists a quasi-étale cover $\tau: S^\dagger \rightarrow S$ such that S^\dagger is either a torus or a normal K3 surface (i.e., a normal surface whose minimal resolution is a K3 surface) and $\tau: S^\dagger \rightarrow S$ is an orbifold morphism (i.e., it can be locally described as $\tau: U^\dagger/G^\dagger \rightarrow U/G$ induced by a morphism $\hat{\tau}: U^\dagger \rightarrow U$ and a group homomorphism $\rho: G^\dagger \rightarrow G$ such that $\hat{\tau}(gz) = \rho(g)\hat{\tau}(z)$ holds for any $g \in G^\dagger$ and $z \in U^\dagger$, where U^\dagger (resp. U) is a local smooth ramified cover with the linear action of the finite group G^\dagger (resp. G .) Note that any normal K3 surface is simply connected. We consider the base change:

$$\begin{array}{ccc} X^\dagger & \xrightarrow{\nu} & X' \\ \downarrow \varphi^\dagger & & \downarrow \varphi \\ S^\dagger & \xrightarrow{\tau} & S. \end{array}$$

The fibration $\varphi: X' \rightarrow S$ satisfies the assumptions of Theorem 2.6 when we set $Y_0 := S_{\text{reg}}$ and $L := -K_{X'}$. Indeed, Conditions (a), (b), (c) are confirmed by Proposition 3.10, Corollary 3.11, and Theorem 3.6. Furthermore, Condition (d) can be confirmed by orbifold structures as follows: Indeed, the reflexive sheaf \mathcal{V}_p defined by $L = -K_{X'}$ as in Theorem 2.6 is an orbifold vector bundle on the orbifold S . By [Wu, Lemma 1], there exists a continuous function ψ on S whose pull-back on each local smooth ramified cover is smooth such that $\omega_S + \sqrt{-1}\partial\bar{\partial}\psi$ defines an orbifold Kähler form on S . For a smooth orbifold metric g on $\det \mathcal{V}_p$, we take $C \gg 1$ such that $\sqrt{-1}\Theta_g + C(\omega_S + \sqrt{-1}\partial\bar{\partial}\psi)$ is positive on each local smooth ramified cover. Then $ge^{-C\psi}$ is a metric satisfying Condition (d). Consequently, Theorems 2.6 and 2.11 show that \mathcal{V}_p is a numerical flat orbifold vector bundle on S . In the same way, we deduce that the fibration $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ also satisfies the assumptions of Theorem 2.6 when we set $Y_0 := S^\dagger \setminus \tau^{-1}(S_{\text{sing}})$ and $L = -K_{X^\dagger}$, by noting that $\nu: X^\dagger \rightarrow X'$ is a quasi-étale cover and $-K_{X^\dagger}$ is φ^\dagger -ample by the construction of $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$. Consequently, the sheaf \mathcal{V}_p^\dagger defined by $L = -K_{X^\dagger}$ is also a numerical flat orbifold bundle on S^\dagger .

We divide our situation into Case 1 and Case 2, and respectively prove Claim 4.9 and Claim 4.10.

Case 1: The case where S^\dagger is a (2-dimensional) torus.

Case 2: The case where S^\dagger is a normal K3 surface.

Claim 4.9. *In Case 1, up to a finite étale cover of S^\dagger , the variety X^\dagger is isomorphic to the projective space bundle $\mathbb{P}(\mathcal{F})$ of a numerically flat vector bundle \mathcal{F} over the torus S^\dagger . Moreover, the vector bundle \mathcal{F} admits a filtration by Hermitian flat line bundles:*

$$0 \rightarrow L_1 \rightarrow \mathcal{F} \rightarrow L_2 \rightarrow 0.$$

Proof. Since S^\dagger is smooth, the sheaf \mathcal{V}_p^\dagger defined as above is a numerical flat vector bundle on the torus S^\dagger by [Wu22b, Main Theorem]. Then, [MW21, Proposition 2.5, Remark 2.6 (b)] shows that $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ is locally constant. In particular, the variety X^\dagger is smooth and $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ is a \mathbb{P}^1 -bundle. Then, the first conclusion follows from the same argument as in Theorem 4.2.

The flat vector bundle \mathcal{F} is constructed from a GL-representation of the fundamental group $\pi_1(S^\dagger)$. Thus, since $\pi_1(S^\dagger)$ is abelian, the vector bundle \mathcal{F} admits the desired filtration. \square

Claim 4.10. *In Case 2, the variety X^\dagger is isomorphic to the product $S^\dagger \times \mathbb{P}^1$.*

Proof. We initially show that $\pi_1(X_{\text{reg}}^\dagger) = \{\text{id}\}$ by applying the Van Kampen theorem. Note that any normal K3 surface is simply connected. The variety S^\dagger is the universal cover in the sense of orbifolds (see [Cam04, Définition 5.3]); hence, we obtain $\pi_1(S_{\text{reg}}^\dagger) \cong \pi_1(S^\dagger) = \{\text{id}\}$. Since $\varphi: X' \rightarrow S$ is a smooth \mathbb{P}^1 -bundle on S_{reg} (which is preserved under the base change), we deduce that $\pi_1((\varphi^\dagger)^{-1}(S_{\text{reg}}^\dagger)) = \{\text{id}\}$. Near a singular point in S_{sing} at which $\tau: S^\dagger \rightarrow S$ is not étale, both $\varphi: X' \rightarrow S$ and $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ can be locally described as follows:

$$\begin{array}{ccc} (\mathbb{P}^1 \times \mathbb{C}^2)/\mu_{m^\dagger} & \xrightarrow{\nu} & (\mathbb{P}^1 \times \mathbb{C}^2)/\mu_m \\ \downarrow \varphi^\dagger & & \downarrow \varphi \\ \mathbb{C}^2/\mu_{m^\dagger} & \xrightarrow{\tau} & \mathbb{C}^2/\mu_m. \end{array}$$

Here the action of $\varepsilon \in \mu_{m^\dagger}$ is given by $\varepsilon \cdot (t, z_1, z_2) = (\varepsilon^b t, \varepsilon z_1, \varepsilon^{-1} z_2)$ and τ is given by $\mathbb{C}^2/\mu_{m^\dagger} \rightarrow \mathbb{C}^2/\mu_m$ induced by the inclusion $\mu_{m^\dagger} \rightarrow \mu_m$. To apply the Van Kampen theorem, we regard X_{reg}^\dagger as the union of $(\varphi^\dagger)^{-1}(S_{\text{reg}}^\dagger)$ and open neighborhoods V_i of $(\varphi^\dagger)^{-1}(S_{\text{sing}}^\dagger)$ in X_{reg}^\dagger . Then, we can see that

- V_i is homeomorphic to $(\mathbb{P}^1 \times \mathbb{C}^2 \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\})/\mu_{m^\dagger}$,
- $(\varphi^\dagger)^{-1}(S_{\text{reg}}^\dagger) \cap V_i$ is homeomorphic to $(\mathbb{P}^1 \times (\mathbb{C}^2 \setminus \{\mathbf{0}\}) \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\})/\mu_{m^\dagger}$,

where $\mathbf{0} := (0, 0) \in \mathbb{C}^2$. Consider the homomorphism induced by the natural inclusion:

$$\pi_1(\mathbb{P}^1 \times (\mathbb{C}^2 \setminus \{\mathbf{0}\}) \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\})/\mu_{m^\dagger} \rightarrow \pi_1(\mathbb{P}^1 \times \mathbb{C}^2 \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\})/\mu_{m^\dagger}.$$

Since $\mathbb{P}^1 \times (\mathbb{C}^2 \setminus \{\mathbf{0}\}) \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\}$ and $\mathbb{P}^1 \times \mathbb{C}^2 \setminus \{(0, \mathbf{0}), (\infty, \mathbf{0})\}$ are the universal covers respectively, the above homomorphism can be regarded as the identity map of μ_{m^\dagger} . Therefore, the Van Kampen theorem shows that $\pi_1(X_{\text{reg}}^\dagger) = \{\text{id}\}$.

The sheaf \mathcal{V}_p^\dagger defined by $L := -K_{X^\dagger}$ is a numerical flat orbifold vector bundle on S^\dagger . Since $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ is an orbifold morphism, the sheaf $(\varphi^{\dagger*}\mathcal{V}_p^\dagger)^{**}$ is also a numerically flat orbifold vector bundle. By Lemma 4.15 (which is proved later), we have

$$H^1(X_{\text{reg}}^\dagger, \mathcal{O}_{X_{\text{reg}}^\dagger}) \cong H^1(X^\dagger, \mathcal{O}_{X^\dagger}) = 0.$$

Then, Corollary 2.12 shows that $(\varphi^{\dagger*}\mathcal{V}_p^\dagger)^{**}$ is a trivial vector bundle on X^\dagger . Note that the analogue of Lemma 4.15 in the 2-dimensional case is false, which is the reason to consider the pull-back of \mathcal{V}_p^\dagger to X^\dagger . Since \mathcal{V}_p^\dagger is locally free on S_{reg}^\dagger , the projection formula indicates that $\mathcal{V}_p^\dagger \cong \varphi_*^\dagger((\varphi^{\dagger*}\mathcal{V}_p^\dagger)^{**})$ holds on S_{reg}^\dagger ; hence \mathcal{V}_p^\dagger is a trivial vector bundle on S_{reg}^\dagger . By [MW21, Proposition 2.5] and $\pi_1(S_{\text{reg}}^\dagger) = \{\text{id}\}$, we conclude that $\varphi^\dagger: X^\dagger \rightarrow S^\dagger$ gives the product structure $S_{\text{reg}}^\dagger \times \mathbb{P}^1$ over S_{reg}^\dagger . Precisely speaking, we need to check Condition (2) in [MW21, Proposition 2.5], that is, the natural morphism $\text{Sym}^m \mathcal{V}_p^\dagger \rightarrow \mathcal{V}_{pm}^\dagger$ on S_{reg}^\dagger is a morphism of local systems. In our case, this is automatically satisfied since the reflexive hulls of $\text{Sym}^m \mathcal{V}_p^\dagger$ and \mathcal{V}_{pm}^\dagger are trivial vector bundles on S^\dagger (cf. [MW21, Remark 2.6 (b)]).

If the meromorphic map $S^\dagger \times \mathbb{P}^1 \dashrightarrow X^\dagger$ (obtained from the above product structure over S^\dagger) fails to be an isomorphism, the fiber \mathbb{P}^1 at a singular point of S^\dagger would be contractible by [Kol91, Proposition 2.1.13]. However, such a contraction does not exist. Therefore, we can conclude that $X^\dagger = S^\dagger \times \mathbb{P}^1$. \square

We finally show that $\pi: X \dashrightarrow X'$ is actually isomorphic. This finishes the proof of Theorem 4.8. Indeed, the MFS $\varphi: X' \rightarrow S$ is a conic bundle by noting that $\varphi: X' \rightarrow S$ is toroidal at any point. The fibration $\varphi: X \cong X' \rightarrow S$ is a locally constant \mathbb{P}^1 -bundle by the same argument as in Theorem 4.2. Meanwhile, since $\pi_1(X) \cong \pi_1(S)$ holds by [Kol93, Theorem 5.2] and [BC15, Theorem 4.1], the base S is simply connected, which implies that $X \cong X'$ is the product of S and \mathbb{P}^1 .

To verify that $\pi: X \dashrightarrow X'$ is isomorphic, we divide our situation into the four subsequent claims as in the proof of Theorem 4.2.

Claim 4.11. *In both Case 1 and Case 2, the final step $X_{N-1} \dashrightarrow X_N = X'$ of the MMP cannot be a flip.*

Proof. The same strategy as in Theorem 4.2 works. Let R be a rational curve on X' . Let d be the degree of the restriction $\nu^{-1}(R) \rightarrow R$ of $\nu: X^\dagger \rightarrow X'$. Then, we see that

$$(c_1(K_{X'}) \cdot R) = \frac{1}{d}(c_1(K_{X^\dagger}) \cdot \nu^{-1}(R)) \leq 0.$$

This implies that $\pi: X_{N-1} \dashrightarrow X_N = X'$ cannot be a flip. \square

Claim 4.12. *In both Case 1 and Case 2, the final step $\pi: X_{N-1} \dashrightarrow X_N = X'$ cannot contract a surface to the one point*

Proof. Let d be the degree of $\nu: X^\dagger \rightarrow X'$. In both Case 1 and Case 2, we have $K_{X'}^2 = (1/d)\nu_*K_{X^\dagger}^2 = 0$ since $K_{X^\dagger}^2 = 0$ holds and $\nu: X^\dagger \rightarrow X'$ is an orbifold morphism. Then, by the same argument as in Claim 4.4, we obtain a contradiction. \square

Claim 4.13. *In Case 2, the final step $\pi: X_{N-1} \dashrightarrow X_N = X'$ cannot contract a surface E to a curve C .*

Proof. In Case 2, for a general $t \in \mathbb{P}^1$, the surface $V := \nu(S^\dagger \times \{t\}) \subset X'$ such that $K_{X_N}|_V$ is numerically trivial and $\bar{V} \cap E$ is an effective curve, where \bar{V} is the strict transform of V . Then, by the same argument as in the first paragraph of Claim 4.5 we obtain a contradiction. \square

Claim 4.14. *In Case 1, the final step $\pi: X_{N-1} \dashrightarrow X_N = X'$ cannot contract a surface E to a curve C .*

Proof. We initially consider the case where C intersects with the image $V := \nu(\mathbb{P}(L_2))$. In this case, we can see that $K_{X_N}|_V$ is numerically trivial and $\bar{V} \cap E$ is an effective curve, where $\bar{V} \subset X_{N-1}$ is the strict transform of V . Hence, the same argument as in the first paragraph of Claim 4.5 yields a contradiction. Thus, we may assume that C does not intersect with $\nu(\mathbb{P}(L_2))$.

We will prove that after the base change by $\nu: X^\dagger \rightarrow X$, the contraction $\pi_{N-1}: X_{N-1} \rightarrow X_N = X'$ coincides with the blow-up of $X_N = X'$ along C , which will lead to a contradiction. To confirm this, we consider the following diagram:

$$\begin{array}{ccccc} \mathbb{P}(\mathcal{F}|_{T^\dagger}) & \hookrightarrow & \mathbb{P}(\mathcal{F}) = X^\dagger & \xrightarrow{\nu} & X' = X_N & \xleftarrow{\pi := \pi_{N-1}} & X_{N-1} \\ \downarrow \varphi^\dagger & & \downarrow \varphi^\dagger & & \downarrow \varphi & & \\ T^\dagger := \varphi^\dagger(C^\dagger) & \hookrightarrow & S^\dagger & \xrightarrow{\tau} & S = S^\dagger/G, & & \end{array}$$

where $C^\dagger := \nu^{-1}(C)$.

We now prove that T^\dagger is the disjoint union of elliptic curves. Consider the irreducible decomposition $T^\dagger = \cup_i T_i$. By [Uen75, Lemma 10.8], there exists an abelian variety in S^\dagger containing T_i . Since S^\dagger is a non-projective torus of dimension 2, the curve T_i must be an elliptic curve. If $T_i \cap T_j \neq \emptyset$ for some $i \neq j$, there exists $g \in G$ such that $T_j = gT_i$ holds by $\tau(T_i) = \varphi \circ \nu(C^\dagger) = \tau(T_j)$. Then, up to a translation of T_i and gT_i , we can see that $gT_i \rightarrow S^\dagger \rightarrow S^\dagger/T_i$ is an isogeny. By [BL99, Proposition 6.1, Chap I], up to an isogeny of S^\dagger , the torus S^\dagger is the product of elliptic curves. In particular S^\dagger is projective, which contradicts the assumption that S^\dagger is a non-projective torus.

Subsequently, we prove that C^\dagger is also the disjoint union of elliptic curves. For simplicity, we assume that T^\dagger is connected (equivalently, T^\dagger is irreducible). Note that, in the general case, applying the same argument to each connected component of T^\dagger yields the desired conclusion. To this end, we write the divisor C^\dagger on $\mathbb{P}(\mathcal{F}|_{T^\dagger})$ as

$$(4.2) \quad C^\dagger = \mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(a) + \varphi^{\dagger*} D$$

for some $a \in \mathbb{Z}$ and some line bundle D on T^\dagger , and prove that $a > 0$ and D is numerically trivial. By considering the intersection number with a general fiber of $\mathbb{P}(\mathcal{F}|_{T^\dagger}) \rightarrow T^\dagger$, we can see that $a \geq 0$ since $\mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(1)$ is nef. Meanwhile, by the exact sequence in Claim 4.9, we can see that $c_1(\mathcal{O}_{\mathbb{P}(\mathcal{F})}(1)) = c_1(\mathbb{P}(L_2))$, where $\mathbb{P}(L_2)$ is regarded as a divisor on $\mathbb{P}(\mathcal{F})$. Hence, by the assumption that $C \cap \nu(\mathbb{P}(L_2)) = \emptyset$, we have

$$(4.3) \quad (C^\dagger \cdot \mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(1)) = (C^\dagger \cdot \mathcal{O}_{\mathbb{P}(\mathcal{F})}(1)) = 0.$$

By using (4.3) and $(\mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(1) \cdot \mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(1)) = 0$, we deduce that $c_1(D) = 0$.

We consider two cases, depending on whether the following exact sequence, obtained from restricting the exact sequence in Claim 4.9, splits or not:

$$0 \rightarrow L_1|_{T^\dagger} \rightarrow \mathcal{F}|_{T^\dagger} \rightarrow L_2|_{T^\dagger} \rightarrow 0.$$

We first consider the case where this sequence splits on the elliptic curve T^\dagger . Take a lattice Λ of \mathbb{C} such that $T^\dagger \cong \mathbb{C}/\Lambda$. Then, since $L_1|_{T^\dagger}$, $L_2|_{T^\dagger}$, and D are numerically trivial line bundles, we can take the corresponding unitary representations

$$\rho_1, \rho_2, \rho_D: \Lambda \rightarrow \mathrm{U}(1).$$

Take a section

$$s \in H^0(\mathbb{P}(\mathcal{F}|_{T^\dagger}), \mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(a) + \varphi^{\dagger*} D) \cong H^0(T^\dagger, \mathrm{Sym}^a(L_1|_{T^\dagger} \oplus L_2|_{T^\dagger}) \otimes D)$$

whose divisor $\mathrm{div}(s)$ coincides with C^\dagger . By pulling back s to the universal cover $\mathbb{C} \times \mathbb{P}^1$, the section s can be identified with an element

$$\sum_{i,j \geq 0, i+j=a} f_{ij}(z) u^i v^j \in \mathcal{O}(\mathbb{C})[u, v]$$

satisfying that

$$f_{ij}(z + \lambda) = f_{ij}(z) \rho_1(\lambda)^i \rho_2(\lambda)^j \rho_D(\lambda) \text{ for any } \lambda \in \Lambda,$$

where $(z, [u : v])$ is a coordinate of $\mathbb{C} \times \mathbb{P}^1$. The Liouville theorem shows that f_{ij} are constant functions on \mathbb{C} . This indicates that C^\dagger is smooth and is the disjoint union of elliptic curves. Indeed, the inverse image of C^\dagger in $\mathbb{C} \times \mathbb{P}^1$ can be written as

$$\{(z, [u : v]) \in \mathbb{C} \times \mathbb{P}^1 \mid \sum_{i,j \geq 0, i+j=a} f_{ij} u^i v^j = 0\},$$

and C^\dagger is the quotient of this inverse image.

We consider the remaining case where the above exact sequence does not split. In this case, by considering the extension class, we can see that $L_1|_{T^\dagger} = L_2|_{T^\dagger}$ and that $\mathbb{P}(\mathcal{F}|_{T^\dagger})$ is the same as in [DPS94, Example 1.7]. By [DPS94, Example 1.7], we can see that $c_1(\mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(a))$ contains the only positive current associated with an effective curve whose support is $\mathbb{P}(L_2|_{T^\dagger})$. This indicates that $C^\dagger = \mathbb{P}(L_2|_{T^\dagger})$ by $c_1(C^\dagger) = c_1(\mathcal{O}_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}(a))$, and thus C^\dagger is an elliptic curve.

Consider the normalization X_{N-1}^\dagger of the fiber product $X^\dagger \times_{X'} X_{N-1}$:

$$\begin{array}{ccc} X_{N-1}^\dagger & \xrightarrow{\nu^\dagger} & X_{N-1} \\ \downarrow \pi^\dagger & & \downarrow \pi := \pi_{N-1} \\ X^\dagger & \xrightarrow{\nu} & X', \end{array}$$

We prove that

$$\pi^\dagger: (X_{N-1}^\dagger, E^\dagger := \nu^{\dagger*} E) \rightarrow (X^\dagger, C^\dagger)$$

satisfies all the conditions of Lemma 4.16 (which is proved later) to conclude that it coincides with the blow-up of X^\dagger along C^\dagger . Condition (0) of Lemma 4.16 is satisfied since C^\dagger and X^\dagger are smooth. Additionally, Condition (3) is also satisfied since X_{N-1} has isolated singularities and $\nu: X^\dagger \rightarrow X'$ is ramified along only X'_{sing} (which are finitely many points). As was the projective case, the contraction $\pi: X_{N-1} \rightarrow X'$ satisfies Conditions (1), (2) and both $-E$ and $-K_{X_{N-1}}$ are π -ample (see [HP15a, Lemma 7.8] for (2) and [HP15a, HP15b, HP16]). By noting that $\nu: X^\dagger \rightarrow X'$ is ramified along only X'_{sing} , we can see that $\pi^\dagger: (X_{N-1}^\dagger, E^\dagger) \rightarrow (X^\dagger, C^\dagger)$ also satisfies Conditions (1), (2) and both $-E^\dagger = -\nu^{\dagger*} E$ and $-K_{X_{N-1}^\dagger} = -\nu^{\dagger*} K_{X_{N-1}}$ on X_{N-1}^\dagger are π^\dagger -ample outside $\nu^{-1}(X'_{\text{sing}})$. To check that $-E^\dagger = -\nu^{\dagger*} E$ is π^\dagger -nef, we take a fiber F of $X_{N-1}^\dagger \rightarrow X^\dagger$. Since $\nu^\dagger(F)$ is a curve contracted by $\pi: X_{N-1} \rightarrow X' = X_N$, we have

$$(-E^\dagger \cdot F) = (-\nu^{\dagger*} E \cdot F) = (-E \cdot \nu^\dagger(F)) > 0$$

which implies that $-E^\dagger$ is π^\dagger -nef. Hence, Conditions (4), (5) are satisfied. Lemma 4.16 shows that $\pi^\dagger: (X_{N-1}^\dagger, E^\dagger) \rightarrow (X^\dagger, C^\dagger)$ is actually the blow-up of X^\dagger along C^\dagger .

Now, let us compute intersection numbers and derive a contradiction. By the formula $K_{X_{N-1}^\dagger} = \pi^{\dagger*} K_{X^\dagger} + E^\dagger$, we have

$$(K_{X_{N-1}^\dagger}^2 \cdot \omega) = (\pi^{\dagger*} K_{X^\dagger}^2 \cdot \omega) + 2(\pi^{\dagger*} K_{X^\dagger} \cdot E^\dagger \cdot \omega) + ((E^\dagger)^2 \cdot \omega),$$

where ω is a Kähler form on X_{N-1}^\dagger . The left-hand side is non-negative. Indeed, for $T_\varepsilon \in c_1(-K_{X_{N-1}}) + \varepsilon\{\omega_{N-1}\}$ in Proposition 3.10 (1), the pull-back $\nu^{\dagger*}(T_\varepsilon)$ represents $c_1(-K_{X_{N-1}^\dagger}) + \nu^{\dagger*}\varepsilon\{\omega_{N-1}\}$ and is smooth outside a Zariski closed set of codimension

≥ 2 since $\nu^\dagger: X_{N-1}^\dagger \rightarrow X_{N-1}$ is quasi-étale. Thus, by noting that X_{N-1}^\dagger is smooth, we see that the Monge-Ampère operator $\nu^{\dagger*}(T_\varepsilon)^2$ is well-defined. This implies that $(K_{X_{N-1}^\dagger}^2 \cdot \omega) \geq 0$. Meanwhile, the right-hand side is negative. Indeed, the first term is zero by $K_{X^\dagger}^2 = 0$. Furthermore, the second term is also zero since $K_{X^\dagger}|_{C^\dagger}$ is numerically trivial by

$$K_{X^\dagger}|_{C^\dagger} = (K_{X^\dagger}|_{\mathbb{P}(\mathcal{F}|_{T^\dagger})})|_{C^\dagger} = K_{\mathbb{P}(\mathcal{F}|_{T^\dagger})}|_{C^\dagger} = 0.$$

Since $\mathcal{O}(-E^\dagger)|_{E^\dagger} = \mathcal{O}_{\mathbb{P}(N_{C^\dagger/X^\dagger})}(1)$ holds and N_{C^\dagger/X^\dagger} is numerically flat, we can see that for $\pi^\dagger|_{E^\dagger}: E^\dagger \rightarrow C^\dagger$,

$$\begin{aligned} ((E^\dagger)^2 \cdot \omega) &= (\mathcal{O}_{\mathbb{P}(N_{C^\dagger/X^\dagger})}(-1) \cdot \omega|_{E^\dagger}) \\ &= - \int_{C^\dagger} (\pi^\dagger|_{E^\dagger})_*(\omega|_{E^\dagger}) < 0. \end{aligned}$$

This is a contradiction. \square

This finishes the proof of Theorem 4.8, and thus completes the proof of Theorem 1.3 under assuming that $\pi_1(X) = \{\text{id}\}$ and $\dim R(X) = 2$. \square

In the following, we give the proofs of the two lemmas used in the proof. Lemm 4.15 is an easy variant of [Wu22a, Lemma 4] and Lemma 4.16 is a Kähler counterpart of [Tzi03, Proposition 1.2].

Lemma 4.15. *Let X be a (not necessarily compact) analytic variety of dimension 3 and let E be a vector bundle on X . Assume that X has isolated cyclic quotient singularities: more precisely, near a singular point, there exists $m \in \mathbb{Z}^+$ such that $X \cong \mathbb{D}/\mu_m$, where \mathbb{D} is a polydisc centered at the origin and the action of μ_m is given by*

$$\varepsilon \cdot (z_0, z_1, z_2) = (\varepsilon^{p_0} z_0, \varepsilon^{p_1} z_1, \varepsilon^{p_2} z_2) \text{ for } \varepsilon \in \mu_m$$

for some $p_i \in \mathbb{Z}$ with isolated singular point. Then, the morphism $H^1(X, E) \rightarrow H^1(X_{\text{reg}}, E)$ induced by the restriction morphism is isomorphic.

Proof. The germ of X at any point x is Cohen-Macaulay. In particular, the depth of $\mathcal{O}_{X,x}$ (i.e., the length of maximal regular sequences contained in the maximal ideal) is equal to the Krull dimension of $\mathcal{O}_{X,x}$. Since E is locally free, the depth of E_x at any point x is 3. By [ST06, Theorem 1.14], the morphism $H^1(X, E) \rightarrow H^1(X_{\text{reg}}, E)$ is isomorphic. \square

Lemma 4.16. *Let $f: (Y, E) \rightarrow (X, C)$ be a divisorial contraction between normal Kähler spaces X and Y of dimension 3 such that E is an irreducible surface and $f(E) = C$ is a curve. Assume the following conditions:*

- (0) $f(E) = C$ is a smooth curve and X is smooth near C .
- (1) $f: Y \rightarrow X$ is a projective morphism.

- (2) *A general fiber of $f|_E: E \rightarrow C$ is smooth, connected, and contained in the regular locus of E .*
- (3) *The singular locus of Y is not dominant over C .*
- (4) *$-E$ is a f -nef \mathbb{Q} -Cartier divisor on Y and f -ample outside finitely many points of X .*
- (5) *$-K_Y$ is a \mathbb{Q} -Cartier divisor on Y and f -ample outside finitely many points of X .*

Then, the contraction $f: Y \rightarrow X$ coincides with the blow-up $p: \text{Bl}_C X \rightarrow X$ of X along C .

Proof. The problem is local in X . Thus, we may assume that X is smooth. We first reduce the problem to show that $f: Y \rightarrow X$ coincides with $p: \text{Bl}_C X \rightarrow X$ outside finitely many points of X . If $f: Y \rightarrow X$ coincides with the blow-up outside finitely many points of X , then we can find a bimeromorphic map

$$\pi: Y \dashrightarrow \text{Bl}_C X \text{ over } C \text{ that is isomorphic in codimension } 1$$

by noting that $f: E \rightarrow C$ is an equidimensional fibration. We conclude that $\pi: Y \dashrightarrow \text{Bl}_C X$ is actually an isomorphism by checking the assumptions of [KM98, Remark 6.37, Lemma 6.39] are satisfied. For any f -ample Cartier divisor A on Y , the Weil divisor $\pi_* A$ corresponding to A via $\pi: Y \dashrightarrow \text{Bl}_C X$ is a Cartier divisor on $\text{Bl}_C X$ (since $\text{Bl}_C X$ is smooth). For the p -ample exceptional divisor $-B$ on $\text{Bl}_C X$, the Weil divisor $\pi_*^{-1} B$ is $-E$, which is f -nef and \mathbb{Q} -Cartier by assumption. Thus, [KM98, Lemma 6.39] shows that $\pi: Y \dashrightarrow \text{Bl}_C X$ is an isomorphism.

We finally show that $f: Y \rightarrow X$ coincides with the blow-up of C outside finitely many points of X . By Condition (4), it is sufficient to show that for any $d \geq 0$,

$$f_* \mathcal{O}(-dE) = \mathcal{I}_C^d$$

over the Zariski open set where $-E$ is f -ample. We may assume that Y is smooth by Condition (3). Furthermore, we may assume that the restriction $f|_E: E \rightarrow C$ (on a non-empty Zariski open set of E) is a smooth morphism between smooth spaces. Since $\mathcal{O}_E(-dE) = K_E - K_Y|_E - (d+1)E|_E$ holds and $-K_Y$ is f -ample, the relative Kodaira vanishing theorem shows that

$$R^i f_*(\mathcal{O}_E(-dE)) = 0 \text{ for any } i > 0, d \geq 0,$$

which finishes the proof by [Mor82, Lemma (3.32)]. □

We finally check that Theorem 1.2 follows from Theorem 1.3 by using almost the same arguments as in [CH19, Theorem 1.4].

Proposition 4.17. *Theorem 1.2 follows from Theorem 1.3.*

Proof. Let X be a non-projective compact Kähler 3-fold with nef anti-canonical bundle. We first show that X admits a locally trivial MRC fibration $X \rightarrow Y$ onto a smooth surface with $c_1(Y) = 0$. The manifold X admits a finite étale cover X' , one of the lists in Theorem 1.3. If $c_1(X') = 0$, then $c_1(X) = 0$; hence we may assume that X' admits a non-trivial locally constant MRC fibration. Let $\mathcal{F} \subset T_X$ be the (unique) saturated integrable subsheaf such that the \mathcal{F} -leaf through a very general point $x \in X$ is a fiber of the MRC-fibration. By Theorem 1.3, the sheaf \mathcal{F} is a regular foliation that is invariant under passing to finite étale covers. Hence, by [Hor07, Corollary 2.11], there exists a smooth morphism $\varphi: X \rightarrow Y$ such that $T_{X/Y} = \mathcal{F}$. Since the general fiber is \mathbb{P}^1 , the fibration $\varphi: X \rightarrow Y$ is locally trivial by the Firscher-Grauert theorem. By Proposition 3.5, the line bundle $-K_Y$ is pseudo-effective. Since Y is not uniruled, we have $c_1(Y) = 0$.

The fibration $\varphi: X \rightarrow Y$ is the MFS of X ; hence $-K_X$ is φ -ample line bundle. By Theorem 2.6, since Y is smooth, we can find a φ -ample line bundle B such that $\varphi_*(pB)$ is numerically flat for $1 \ll p \in \mathbb{Z}_+$. Note that the numerical flatness follows from [Wu22b, Main Theorem]. Then, we see that $\varphi: X \rightarrow Y$ is locally constant by [MW21, Proposition 2.5]. Since Y is a compact Kähler, Condition (2) in [MW21, Proposition 2.5] is automatically satisfied (see [MW21, Remark 2.6 (b)]). \square

REFERENCES

- [AR14] C. Araujo, J.J. Ramón-Marí, *Flat deformations of \mathbb{P}^n* , Bull. Braz. Math. Soc. (N.S.) **45** (2014), no. 3, 371–383.
- [Bar78] D. Barlet, *Convexité de l'espace des cycles*, Bull. Soc. Math. France **106** (1978), 373–397.
- [BC15] Y. Brunebarbe, F. Campana, *Fundamental group and pluridifferentials on compact Kähler manifolds*, Mosc. Math. J. **16** (2016), no. 4, 651–658.
- [BDPP13] S. Boucksom, J.-P. Demailly, M. Păun, T. Peternell, *The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension*, J. Algebraic Geom. **22** (2013), no. 2, 201–248.
- [BEG13] S. Boucksom, P. Eyssidieux, V. Guedj, *An introduction to the Kähler-Ricci flow*, Lecture Notes in Mathematics, 2086. Springer, Cham, 2013. viii+333 pp.
- [BHPV04] W. Barth, K. Hulek, C. Peters, A. Van de Ven, *Compact complex surfaces*, A Series of Modern Surveys in Mathematics, **4**. Springer-Verlag, Berlin, 2004. xii+436 pp.
- [BL99] C. Birkenhake, H. Lange, *Complex Tori*, Progr. Math., **177** Birkhäuser Boston, Inc., Boston, MA, 1999. xvi+251 pp.
- [Bru06] M. Brunella, *A positivity property for foliations on compact Kähler manifolds*, Internat. J. Math. **17** (2006), no. 1, 35–43.
- [BP04] T. Bauer, T. Peternell. *Nef reduction and anticanonical bundles*, Asian J. Math. **8**, No. 2, 315–352 (2004).
- [Cam92] F. Campana, *Connexité rationnelle des variétés de Fano*, Ann. Sci. École Norm. Sup. (4) **25** (1992), no. 5, 539–545.
- [Cam95] F. Campana, *Remarques sur les groupes de Kähler nilpotents*, Ann. Sci. École Norm. Sup. (4) **28** (1995), no. 3, 307–316.
- [Cam04] F. Campana, *Orbifolds à première classe de Chern nulle*, The Fano Conference, 339–351. Univ.Torino, Turin, 2004.
- [Cao13] J. Cao, *A remark on compact Kähler manifolds with nef anticanonical bundles and applications*, available at [arXiv:1305.4397v2](https://arxiv.org/abs/1305.4397v2).
- [Cao19] J. Cao, *Albanese maps of projective manifolds with nef anticanonical bundles*, Ann. Sci. Éc. Norm. Supér. (4), **52** (2019), no. 5, 1137–1154.
- [CCM21] F. Campana, J. Cao, S. Matsumura, *Projective klt pairs with nef anti-canonical divisor*, Algebr. Geom. **8** (2021), no. 4, 430–464.
- [CDP15] F. Campana, J.-P. Demailly, T. Peternell, *Rationally connected manifolds and semipositivity of the Ricci curvature*, Recent advances in algebraic geometry, 71–91, London Math. Soc. Lecture Note Ser., **417**, Cambridge Univ. Press, Cambridge, 2015.
- [CH19] J. Cao, A. Höring, *A decomposition theorem for projective manifolds with nef anticanonical bundle*, J. Algebraic Geom. **28** (2019), 567–597.
- [CP91] F. Campana, T. Peternell, *Projective manifolds whose tangent bundles are numerically effective*, Math. Ann. **289** (1991), no. 1, 169–187.
- [Dem85] J.-P. Demailly, *Mesures de Monge-Ampère et caractérisation géométrique des variétés algébriques affines*, Mém. Soc. Math. France (N.S.) No. 19 (1985), 124 pp.
- [Dem92] J.-P. Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. **1** (1992), no. 3, 361–409.
- [Dem] Jean-Pierre Demailly, *Complex analytic and differential geometry*, available at <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.

- [DPS94] J.-P. Demailly, T. Peternell, M. Schneider, *Compact complex manifolds with numerically effective tangent bundles*, J. Algebraic Geom. **3** (1994), no. 2, 295–345.
- [DPS96] J.-P. Demailly, T. Peternell, M. Schneider, *Compact Kähler manifolds with Hermitian semi-positive anticanonical bundle*, Compositio Math. **101** (1996), no. 2, 217–224.
- [EIM23] S. Ejiri, M. Iwai, S. Matsumura, *On asymptotic base loci of relative anti-canonical divisors of algebraic fiber spaces*, J. Algebraic Geom. **32** (2023), no. 3, 477–517.
- [Ele82] G. Elencwajg, *The Brauer groups in complex geometry*, Brauer groups in ring theory and algebraic geometry (Wilrijk, 1981), pp. 222–230, Lecture Notes in Math., **917**, Springer, Berlin-New York, 1982.
- [Fuj88] A. Fujiki, *Finite automorphism groups of complex tori of dimension two*, Publ. Res. Inst. Math. Sci. **24** (1988), no. 1, 1–97.
- [GPR94] H. Grauert, Th. Peternell, R. Remmert, *Several Complex Variables VII*, Sheaf-Theoretical Methods in Complex Analysis, 1994, Encyclopaedia of Mathematical Sciences (EMS, volume 74).
- [HIM22] G. Hosono, M. Iwai, S. Matsumura, *On projective manifolds with pseudo-effective tangent bundle*, J. Inst. Math. Jussieu **21** (2022), no. 5, 1801–1830.
- [Hor07] A. Höring, *Uniruled varieties with split tangent bundle*, Math. Z. **256** (2007), no. 3, 465–479.
- [HP15a] A. Höring, T. Peternell, *Bimeromorphic geometry of Kähler threefolds*, Algebraic geometry: Salt Lake City 2015, 381–402, Proc. Sympos. Pure Math., 97.1, Amer. Math. Soc., Providence, RI, 2018.
- [HP15b] A. Höring, T. Peternell, *Mori fibre spaces for Kähler threefolds*, J. Math. Sci. Univ. Tokyo **22** (2015), no. 1, 219–246.
- [HP16] A. Höring, T. Peternell, *Minimal models for Kähler threefolds*, Invent. Math. **203** (2016), no. 1, 217–264.
- [HPS18] C. Hacon, M. Popa, C. Schnell, *Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Păun*, Local and global methods in algebraic geometry, 143–195, Contemp. Math., **712**, Amer. Math. Soc., Providence, RI, 2018.
- [HSW81] A. Howard, B. Smyth, H. Wu, *On compact Kähler manifolds of nonnegative bisectional curvature I and II*, Acta Math. **147** (1981), no. 1-2, 51–70.
- [Kol91] J. Kollár, *Flips, flops, minimal models, etc.*, Surveys in differential geometry (Cambridge, MA, 1990), 113–199, Lehigh Univ., Bethlehem, PA, 1991.
- [Kol93] J. Kollár, *Shafarevich maps and plurigenera of algebraic varieties*, Invent. Math. **113** (1993), no. 1, 177–215.
- [KM98] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti. Translated from the 1998 Japanese original. Cambridge Tracts in Mathematics, 134. Cambridge University Press, Cambridge, 1998. viii+254 pp.
- [KoMM92] J. Kollár, Y. Miyaoka, S. Mori, *Rationally connected varieties*, J. Algebraic Geom. **1** (1992), no. 3, 429–448.
- [Mat20] S. Matsumura, *On the image of MRC fibrations of projective manifolds with semi-positive holomorphic sectional curvature*, Pure Appl. Math. Q. **16**, No. 5 (2020), pp. 1443–1463.
- [Mat22a] S. Matsumura, *On projective manifolds with semi-positive holomorphic sectional curvature*, Amer. J. Math. **144** (2022), no. 3, 747–777.
- [Mat22b] S. Matsumura, *Open problems on structure of positively curved projective varieties*, Ann. Fac. Sci. Toulouse Math. (6) **31** (2022), no. 3, 1011–1029.
- [Mat23] S. Matsumura, *On the minimal model program for projective varieties with pseudo-effective tangent sheaf*, Épijournal Géom. Algébrique **7** (2023), Art. 22, 13 pp.

- [Mi83] M. Miyanishi, *Algebraic methods in the theory of algebraic threefolds –surrounding the works of Iskovskikh, Mori and Sarkisov*, Algebraic varieties and analytic varieties (Tokyo, 1981), 69–99, Adv. Stud. Pure Math., **1**, North-Holland, Amsterdam, 1983.
- [MM07] X. Ma, G. Marinescu, *Holomorphic Morse inequalities and Bergman kernels*, Progress in Mathematics, 254. Birkhäuser Verlag, Basel, 2007. xiv+422 pp.
- [Mok88] N. Mok, *The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature*, J. Differential Geom. **27** (1988), no. 2, 179–214.
- [Mor82] S. Mori, *Threefolds whose canonical bundles are not numerically effective*, Algebraic threefolds (Varenna, 1981), pp. 155–189 Lecture Notes in Math., **947** Springer-Verlag, Berlin-New York, 1982.
- [MP08a] S. Mori, Y. Prokhorov, *On \mathbb{Q} -conic bundles*, Publ. Res. Inst. Math. Sci. **44** (2008), no. 2, 315–369.
- [MP08b] S. Mori, Y. Prokhorov, *On \mathbb{Q} -conic bundles, II*. Publ. Res. Inst. Math. Sci. **44** (2008), no. 3, 955–971.
- [MW21] S. Matsumura, J. Wang, *Structure theorem for projective klt pairs with nef anti-canonical divisor*, available at [arXiv:2105.14308v2](https://arxiv.org/abs/2105.14308v2).
- [NW] P. Naumann, X. Wu, *Albanese map for Kähler manifolds with nef anticanonical bundle*, available at [arXiv:2310.06695](https://arxiv.org/abs/2310.06695).
- [Pău97] M. Păun, *Sur le groupe fondamental des variétés kählériennes compactes à classe de Ricci numériquement effective*, C. R. Acad. Sci. Paris Sér. I Math. **324** (1997), no. 11, 1249–1254.
- [Pău17] M. Păun, *Relative adjoint transcendental classes and Albanese map of compact Kähler manifolds with nef Ricci curvature*, Higher dimensional algebraic geometry – in honour of Professor Yujiro Kawamata’s sixtieth birthday, 335–356. Adv. Stud. Pure Math., **74**, Mathematical Society of Japan, Tokyo, 2017
- [Pro07] Y. Prokhorov, *The degree of \mathbb{Q} -Fano threefolds*, Mat. Sb. **198** (2007), no. 11, 153–174; translation in Sb. Math. **198** (2007), no. 11-12, 1683–1702.
- [Pro18] Y. Prokhorov, *The rationality problem for conic bundles*, Uspekhi Mat. Nauk **73** (2018), no. 3(441), 3–88; translation in Russian Math. Surveys **73** (2018), no. 3, 375–456.
- [PS98] T. Peternell, F. Serrano, *Threefolds with nef anticanonical bundles*, Collect. Math. **49**, No. 2-3, 465–517 (1998).
- [PT18] M. Păun, S. Takayama, *Positivity of twisted relative pluricanonical bundles and their direct images*, J. Algebraic Geom. **27** (2018), 211–272.
- [Sar82] V. G. Sarkisov, *On conic bundle structures*, Izv. Akad. Nauk SSSR Ser. Mat. **46** (1982), no. 2, 371–408, 432.
- [Sim92] C.-T. Simpson, *Higgs bundles and local systems*, Inst. Hautes Études Sci. Publ. Math. No. **75** (1992), 5–95.
- [ST06] Y.-T. Siu, G. Trautmann, *Gap-Sheaves and Extension of Coherent Analytic Subsheaves*, Lecture Notes in Math., Vol. **172** Springer-Verlag, Berlin-New York, 1971. v+172 pp.
- [Tzi03] N. Tziolas, *Terminal 3-fold Divisorial Contractions of a Surface to a Curve I*, Compositio Math. **139** (2003), no.3, 239–261.
- [Uen75] K. Ueno *Classification Theory of Algebraic Varieties and Compact Complex Spaces* Lecture Notes in Math., vol. 439, Springer-Verlag (1975).
- [Wan21] J. Wang, *On the Itaka Conjecture $C_{n,m}$ for Kähler Fibre Spaces*, Ann. Fac. Sci. Toulouse Math. (6) **30** (2021), no. 4, 813–897.
- [Wu22a] X. Wu, *A study of nefness in higher codimension*, Bull. Soc. Math. France **150** (2022), no. 1, 209–249.

- [Wu22b] X. Wu, *Strongly pseudo-effective and numerically flat reflexive sheaves*, J. Geom. Anal. **32** (2022), no. 4, Paper No. 124, 61 pp.
- [Wu] X. Wu, *On compact Kähler orbifold*, available at [arXiv:2302.11914v1](https://arxiv.org/abs/2302.11914).
- [Wan20] J. Wang, *Structure of projective varieties with nef anticanonical divisor: the case of log terminal singularities*, Math. Ann. **384** (2022), no.1-2, 47–100.

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, 6-3, ARAMAKI AZA-AOBA, AOBA-KU, SENDAI 980-8578, JAPAN.

Email address: `mshinichi-math@tohoku.ac.jp`

Email address: `mshinichi0@gmail.com`

LABORATOIRE MATHÉMATIQUES ET INTERACTIONS J.A. DIEUDONNÉ, UNIVERSITÉ CÔTE D'AZUR, NICE, 06108, FRANCE

Email address: `xiaojun.wu@uni-bayreuth.de`

Email address: `xiaojun.wu@univ-cotedazur.fr`