

Spectral representation of Matsubara n -point functions: Exact kernel functions and applications

Johannes Halbinger, Benedikt Schneider and Björn Sbierski

Department of Physics and Arnold Sommerfeld Center for Theoretical Physics (ASC),

Ludwig-Maximilians-Universität München, Theresienstr. 37, München D-80333, Germany

Munich Center for Quantum Science and Technology (MCQST), Schellingstr. 4, D-80799 München, Germany

(Dated: April 11, 2023)

In the field of quantum many-body physics, the spectral (or Lehmann) representation simplifies the calculation of Matsubara n -point correlation functions if the eigensystem of a Hamiltonian is known. It is expressed via a universal kernel function and a system- and correlator-specific product of matrix elements. Here we provide the kernel functions in full generality, for arbitrary n , arbitrary combinations of bosonic or fermionic operators and an arbitrary number of anomalous terms. As an application, we consider bosonic 3- and 4-point correlation functions for the fermionic Hubbard atom and a free spin, respectively.

I. INTRODUCTION AND DEFINITIONS

Multi-point correlation functions of n quantum mechanical operators, also known as n -point functions, are a central concept in the study of quantum many-body systems and field theory [1]. They generalize the well-known 2-point functions, which, for the example of electrons in the solid state, are routinely measured by scanning tunneling spectroscopy, angle-resolved photon emission spectroscopy [2]. For magnetic systems, the 2-point spin correlators can be probed in a neutron scattering experiment. Higher order correlation functions with $n = 3, 4, 5, \dots$ can for example be measured in non-linear response settings [3]. In the emerging field of cold atomic quantum simulation, (equal-time) n -point functions are even directly accessible [4].

On the theoretical side the study of higher order correlation functions gains traction as well. One motivation is the existence of exact relations between correlation functions of different order n [5, 6]. Although these exact relations can usually not be solved exactly, they form a valuable starting point for further approximations like the parquet approximation [7]. Thus even if the 4-point correlator (or, in that context, its essential part, the one-line irreducible vertex [1]) might not be the primary quantity of interest in a calculation, it appears as a building block of the method. Another example is the functional renormalization group method (fRG) in vertex expansion [8, 9] which rewrites the many body problem as a hierarchy of differential equations for the vertices that interpolate between a simple solvable starting point and the physical theory [10]. Whereas experiments measure correlation functions in real time (or frequency), in theory one often is concerned with the related but conceptually simpler versions depending on imaginary time [1]. In the following, we will focus on these Matsubara correlation functions, which, nevertheless feature an intricate time- or frequency dependence.

Whereas the above theoretical methods usually provide only an approximation for the n -point functions, an important task is to calculate these objects exactly which should be possible for simple quantum many body systems. We consider systems simple if they are amenable to exact diagonalization (ED), i.e. feature a small enough Hilbert space like few-site clusters of interacting quantum spins or fermions. Also impurity systems, where interactions only act locally, can be approximately diagonalized using the numerical renormalization group [11].

Knowing the exact n -point functions for simple systems is important for benchmark testing newly developed methods before deploying them to harder problems. Moreover, n -point functions for simple systems often serve as the starting point of further approximations like in the spin-fRG [12–14] or appear intrinsically in a method like in diagrammatic extensions of dynamical mean field theory [15] with its auxiliary impurity problems. Another pursuit enabled by the availability of exact n -point functions is to interpret the wealth of information encoded in these objects, in particular in their rich frequency structure. For example, Ref. [16] studied the fingerprints of local moment formation and Kondo screening in quantum impurity models.

In this work we complete the task to calculate exact n -point functions by generalizing the spectral (or Lehmann) representation [1, 17] for Matsubara n -point correlation functions to arbitrary n . We assume that a set of eigenstates and -energies is given. Following pioneering work of Refs. [18–21] we split the problem of calculating imaginary frequency correlators into the computation of a universal kernel function and a system- and correlator-specific part (called partial spectral function in Ref. [21]). We provide the kernel functions in full generality for an arbitrary number n of bosonic or fermionic frequencies. Previously, these kernel functions were known exactly only up to the 3-point case [18], for the fermionic 4-point case [19–21] or for the general n -point case [21] but disregarding anomalous contributions to the sum that the kernel function consists of. These anomalous contributions, which are at the heart of the complexity of Matsubara n -point functions occur when certain sums of eigenenergies and frequencies

vanish individually, see the anti-diagonal rays in Fig. 1(c). Physically, they correspond to long-term memory effects, are related to non-ergodicity and, in the case of bosonic two-point functions reflect the difference between static isothermal susceptibilities and the zero-frequency limit of the dynamical Kubo response function [22, 23].

The structure of the paper is as follows: In Sec. II we define the Matsubara n -point function $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_{n-1})$ and review some of its properties. The spectral representation is derived in Sec. III with Eq. (15) being the central equation written in terms of the kernel function $K_n(\Omega_1, \dots, \Omega_{n-1})$. Our main result is an exact closed-form expression of this most general kernel function in Sec. IV. Examples for $n = 2, 3, 4, 5$ are given in Sec. V where we also discuss simplifications for the purely fermionic case. We continue with applications to two particular systems relevant in the field of condensed matter theory: In Sec. VI, we consider the Hubbard atom and the free spin, for which we compute n -point functions not previously available in the literature. We conclude in Sec. VII.

II. DEFINITION OF MATSUBARA n -POINT FUNCTION $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n)$

We consider a set of $n = 2, 3, 4, \dots$ operators $\{A_1, A_2, \dots, A_n\}$ defined on the Hilbert space of a quantum many-body Hamiltonian H . The operators can be fermionic, bosonic or a combination of both types with the restriction that there is an even number of fermionic operators. As an example, $A_1 = d^\dagger d \equiv n$, $A_2 = d$, $A_3 = d^\dagger$ where d^\dagger and d are canonical fermionic creation and annihilation operators. A subset of operators is called bosonic if they create a closed algebra under the commutation operation. They are called fermionic if the algebra is closed under anti-commutation, see Sec. 1 of Ref. [24]. Spin operators are thus bosonic.

We define the imaginary time-ordered n -point correlation functions for imaginary times $\tau_k \in [0, \beta]$, [25, 26],

$$G_{A_1 A_2 \dots A_n}(\tau_1, \tau_2, \dots, \tau_n) \equiv \langle \mathcal{T} A_1(\tau_1) A_2(\tau_2) \dots A_n(\tau_n) \rangle, \quad (1)$$

where $A_k(\tau_k) = e^{\tau_k H} A_k e^{-\tau_k H}$ denotes Heisenberg time evolution. Here and in the following, $k = 1, 2, \dots, n$. The expectation value is calculated as $\langle \dots \rangle = \text{tr}[\rho \dots]$ with $\rho = \exp(-\beta H)/Z$ the thermal density operator at temperature $\beta = 1/T$ and $Z = \text{tr} \exp(-\beta H)$ is the partition function. Note that other conventions for the n -point function differing by a prefactor are also used in the literature, e.g. Ref. [21] multiplies with $(-1)^{n-1}$. In Eq. (1), the imaginary time ordering operator \mathcal{T} orders the string of Heisenberg operators,

$$\mathcal{T} A_1(\tau_1) A_2(\tau_2) \dots A_n(\tau_n) \equiv \zeta(p) A_{p(1)}(\tau_{p(1)}) A_{p(2)}(\tau_{p(2)}) \dots A_{p(n)}(\tau_{p(n)}), \quad (2)$$

where p is the permutation $p \in \mathcal{S}_n$ such that $\tau_{p(1)} > \tau_{p(2)} > \dots > \tau_{p(n)}$ and the sign $\zeta(p)$ is -1 if the operator string $A_{p(1)} A_{p(2)} \dots A_{p(n)}$ differs from $A_1 A_2 \dots A_n$ by an odd number of transpositions of fermionic operators, otherwise it is $+1$. The special case $n = 2$ with $\zeta(12) = 1$ and $\zeta(21) = \zeta$ ($\zeta = 1$ for $A_{1,2}$ bosonic, $\zeta = -1$ for $A_{1,2}$ fermionic) simplifies to

$$\mathcal{T} A_1(\tau_1) A_2(\tau_2) = \begin{cases} A_1(\tau_1) A_2(\tau_2) & : \tau_1 > \tau_2, \\ \zeta A_2(\tau_2) A_1(\tau_1) & : \tau_2 > \tau_1. \end{cases} \quad (3)$$




Figure 1. (a) Ordering convention for imaginary times in Eq. (9). (b) Eigenstates and energies of the Hubbard atom. (c) Matsubara correlation function $G_{S^x S^y S^z}(\omega_1, \omega_2)$ with $\omega_j = 2\pi m_j/\beta$, $j = 1, 2$ for the Hubbard atom (34) at $\beta = 10$, $h = 0.1$, $\epsilon = -2$, $U = 2$, see Eq. (44). The sharp anti-diagonal ray $\propto \delta_{\omega_1 + \omega_2, 0}$ represents an anomalous term of order $a = 1$. The other broadened rays become sharp and anomalous for $h \rightarrow 0$, see Eq. (48).

Imaginary time-ordered correlation functions (1) fulfill certain properties which we review in the following, see e.g. [26] for a more extensive discussion. First, they are invariant under translation of all time arguments,

$$G_{A_1 A_2 \dots A_n}(\tau_1, \tau_2, \dots, \tau_n) = G_{A_1 A_2 \dots A_n}(\tau_1 + \tau, \tau_2 + \tau, \dots, \tau_n + \tau), \quad (4)$$

where $\tau \in \mathbb{R}$ such that $\tau_k + \tau \in [0, \beta]$. They also fulfill periodic or anti-periodic boundary conditions for the individual arguments τ_k ,

$$G_{A_1 \dots A_n}(\tau_1, \dots, \tau_k = 0, \dots, \tau_n) = \zeta_k G_{A_1 \dots A_n}(\tau_1, \dots, \tau_k = \beta, \dots, \tau_n) \quad (5)$$

where $\zeta_k = +1$ if A_k is from the bosonic subset of operators or $\zeta_k = -1$ if it is from the fermionic one. This motivates the use of a Fourier transformation,

$$G_{A_1 \dots A_n}(\tau_1, \dots, \tau_n) \equiv \beta^{-n} \sum_{\omega_1, \dots, \omega_n} e^{-i(\omega_1 \tau_1 + \dots + \omega_n \tau_n)} G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n), \quad (6)$$

$$G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n) = \int_0^\beta d\tau_1 \dots \int_0^\beta d\tau_n e^{+i(\omega_1 \tau_1 + \dots + \omega_n \tau_n)} G_{A_1 \dots A_n}(\tau_1, \dots, \tau_n), \quad (7)$$

where $\omega_k = 2\pi m_k / \beta$ or $\omega_k = 2\pi(m_k + 1/2) / \beta$ with $m_k \in \mathbb{Z}$ are bosonic or fermionic Matsubara frequencies, respectively, and the \sum_{ω_k} is short for $\sum_{m_k \in \mathbb{Z}}$. Note that fermionic Matsubara frequencies are necessarily nonzero, a property that will become important later. As we are not mixing with real-frequency formalisms, we will not write the imaginary unit in front of Matsubara frequencies in the arguments of $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n)$. Again, note that in the literature, different conventions for the Fourier transformation of n -point functions are in use, in particular some authors pick different signs in the exponent of Eq. (7) for fermionic creation and annihilation operators or depending on operator positions.

Time translational invariance (4) implies frequency conservation at the left hand side of Eq. (7),

$$G_{A_1 \dots A_n}(\omega_1, \dots, \omega_{n-1}, \omega_n) \equiv \beta \delta_{0, \omega_1 + \dots + \omega_n} G_{A_1 \dots A_n}(\omega_1, \dots, \omega_{n-1}), \quad (8)$$

where on the right hand side we skipped the n -th frequency entry in the argument list of G . Note that we do not use a new symbol for the correlation function when we pull out the factor β and the Kronecker delta function.

III. SPECTRAL REPRESENTATION OF $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n)$

The integrals involved in the Fourier transformation (7) create all $n!$ different orderings of the time arguments τ_k . As in Refs. [19, 21] it is thus convenient to use a sum over all $n!$ permutations $p \in S_n$ and let a product of $n-1$ step-functions θ with $\theta(x) = 1$ for $x > 0$ and 0 otherwise filter out the unique ordering for which $\beta > \tau_{p(1)} > \tau_{p(2)} > \dots > \tau_{p(n-1)} > \tau_{p(n)} > 0$, see Fig. 1(a),

$$G_{A_1 \dots A_n}(\tau_1, \dots, \tau_n) = \sum_{p \in S_n} \zeta(p) \left[\prod_{i=1}^{n-1} \theta(\tau_{p(i)} - \tau_{p(i+1)}) \right] \langle A_{p(1)}(\tau_{p(1)}) A_{p(2)}(\tau_{p(2)}) \dots A_{p(n)}(\tau_{p(n)}) \rangle. \quad (9)$$

To expose explicitly the time dependence of the Heisenberg operators, we insert n times the basis of eigenstates and -energies of the many-body Hamiltonian H . Instead of the familiar notation $|j_1\rangle, |j_2\rangle, \dots$ and E_{j_1}, E_{j_2}, \dots we employ $|\underline{1}\rangle, |\underline{2}\rangle, \dots$ and $E_{\underline{1}}, E_{\underline{2}}, \dots$ for compressed notation and denote operator matrix elements as $A^{\underline{1}\underline{2}} = \langle \underline{1} | A | \underline{2} \rangle$. We obtain

$$G_{A_1 \dots A_n}(\tau_1, \dots, \tau_n) = \sum_{p \in S_n} \zeta(p) \left[\prod_{i=1}^{n-1} \theta(\tau_{p(i)} - \tau_{p(i+1)}) \right] \times \frac{1}{Z} \sum_{\underline{1} \dots \underline{n}} e^{-\beta E_{\underline{1}}} e^{\tau_{p(1)} E_{\underline{1}}} A_{p(1)}^{\underline{1}\underline{2}} e^{(-\tau_{p(1)} + \tau_{p(2)}) E_{\underline{2}}} A_{p(2)}^{\underline{2}\underline{3}} e^{(-\tau_{p(2)} + \tau_{p(3)}) E_{\underline{3}}} \dots e^{(-\tau_{p(n-1)} + \tau_{p(n)}) E_{\underline{n}}} A_{p(n)}^{\underline{n}\underline{1}} e^{-\tau_{p(n)} E_{\underline{1}}}, \quad (10)$$

and apply the Fourier transform according to the definition (7),

$$G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n) = \frac{1}{Z} \sum_{p \in S_n} \zeta(p) \sum_{\underline{1} \dots \underline{n}} e^{-\beta E_{\underline{1}}} A_{p(1)}^{\underline{1}\underline{2}} A_{p(2)}^{\underline{2}\underline{3}} \dots A_{p(n)}^{\underline{n}\underline{1}} \times \left[\int_0^\beta d\tau_{p(1)} e^{\Omega_{p(1)}^{\underline{1}\underline{2}} \tau_{p(1)}} \right] \left[\int_0^{\tau_{p(1)}} d\tau_{p(2)} e^{\Omega_{p(2)}^{\underline{2}\underline{3}} \tau_{p(2)}} \right] \dots \left[\int_0^{\tau_{p(n-2)}} d\tau_{p(n-1)} e^{\Omega_{p(n-1)}^{\underline{n-1}\underline{n}} \tau_{p(n-1)}} \right] \left[\int_0^{\tau_{p(n-1)}} d\tau_{p(n)} e^{\Omega_{p(n)}^{\underline{n}\underline{1}} \tau_{p(n)}} \right], \quad (11)$$

where we defined

$$\Omega_k^{ab} \equiv i\omega_k + E_{\underline{a}} - E_{\underline{b}} \in \mathbb{C}. \quad (12)$$

In Eq. (11), the first line carries all the system- and correlator dependent information whereas the second line can be regarded as a universal kernel function defined for general $\{\Omega_k\}_k$ with $\Omega_k \in \mathbb{C}$ which is probed at system- and correlator dependent points (12) in the complex plane. Upon renaming the τ -integration variables $\tau_{p(k)} \rightarrow \tau_k$, this kernel function is written as follows:

$$\mathcal{K}_n(\Omega_1, \dots, \Omega_n) \equiv \left[\int_0^\beta d\tau_1 e^{\Omega_1 \tau_1} \right] \left[\int_0^{\tau_1} d\tau_2 e^{\Omega_2 \tau_2} \right] \dots \left[\int_0^{\tau_{n-2}} d\tau_{n-1} e^{\Omega_{n-1} \tau_{n-1}} \right] \left[\int_0^{\tau_{n-1}} d\tau_n e^{\Omega_n \tau_n} \right] \quad (13)$$

$$\equiv \beta \delta_{0, \Omega_1 + \Omega_2 + \dots + \Omega_n} K_n(\Omega_1, \dots, \Omega_{n-1}) + R_n(\Omega_1, \dots, \Omega_n). \quad (14)$$

In the second line we split \mathcal{K}_n into a part K_n proportional to $\beta \delta_{0, \Omega_1 + \Omega_2 + \dots + \Omega_n}$ and the rest R_n . We dropped Ω_n from the argument list of K_n which can be reconstructed from $\{\Omega_1, \dots, \Omega_{n-1}\}$.

Finally, we express $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n)$ of Eq. (11) using the kernel \mathcal{K}_n so that the general $\Omega_k \in \mathbb{C}$ get replaced by Ω_k^{ab} of Eq. (12) for which $\Omega_{p(1)}^{12} + \Omega_{p(2)}^{23} + \dots + \Omega_{p(n)}^{n1} = i(\omega_1 + \omega_2 + \dots + \omega_n)$ since the $E_{\underline{k}}$ cancel pairwise. Due to Eq. (8), the terms proportional to R_n are guaranteed to cancel when summed over permutations $p \in S_n$ and only the terms proportional to K_n remain. We drop the $\beta \delta_{0, \omega_1 + \omega_2 + \dots + \omega_n}$ from both sides [c.f. Eq. (8)] and find the spectral representation of the n -point correlation function in the Matsubara formalism,

$$G_{A_1 \dots A_n}(\omega_1, \dots, \omega_{n-1}) = \frac{1}{Z} \sum_{p \in S_n} \zeta(p) \sum_{\underline{1} \dots \underline{n}} e^{-\beta E_{\underline{1}}} A_{p(1)}^{12} A_{p(2)}^{23} \dots A_{p(n)}^{n1} \times K_n(\Omega_{p(1)}^{12}, \Omega_{p(2)}^{23}, \dots, \Omega_{p(n-1)}^{n-1n}). \quad (15)$$

Equivalent expressions were derived in the literature before [18–21]. However, kernel functions K_n were previously only known for small n or only approximately, restricting to a certain low order of anomalous terms, see the discussion in Sec. V. We define an anomalous term of order $a = 1, 2, \dots, n-1$ as a summand contributing to $K_n(\Omega_1, \dots, \Omega_{n-1})$ that contains a product of a Kronecker delta functions $\delta_{0,x}$ where x is a sum of a subset of $\{\Omega_1, \dots, \Omega_{n-1}\}$. As can be seen in Fig. 1(c), these anomalous terms are qualitatively important features of $G_{A_1 \dots A_n}(\omega_1, \dots, \omega_n)$.

In the next section, we present a simple but yet exact expression for general $K_n(\Omega_1, \dots, \Omega_{n-1})$. Readers not interested in the derivation can directly skip to the result in Eq. (25) or its explicit form for $n = 2, 3, 4, 5$ in Sec. (V).

IV. GENERAL KERNEL FUNCTION $K_n(\Omega_1, \dots, \Omega_{n-1})$

Assuming the eigensystem and matrix elements entering Eq. (15) are known, the remaining task is to find expressions for the kernel function $K_n(\Omega_1, \dots, \Omega_{n-1})$ defined via Eqns. (13) and (14) as the part of $\mathcal{K}_n(\Omega_1, \Omega_2, \dots, \Omega_n)$ multiplying $\beta \delta_{0, \Omega_1 + \Omega_2 + \dots + \Omega_n}$. To facilitate the presentation in this section, in Eq. (13) we rename the integration variables $\tau_k \rightarrow \tau_{n-k+1}$ and define new arguments $z_{n-j+1} = \Omega_j$ for $j = 1, 2, \dots, n-1$,

$$K_n(\Omega_1 = z_n, \Omega_2 = z_{n-1}, \dots, \Omega_n = z_1) = \left[\int_0^\beta d\tau_n e^{z_n \tau_n} \right] \left[\int_0^{\tau_n} d\tau_{n-1} e^{z_{n-1} \tau_{n-1}} \right] \dots \left[\int_0^{\tau_3} d\tau_2 e^{z_2 \tau_2} \right] \left[\int_0^{\tau_2} d\tau_1 \underbrace{e^{z_1 \tau_1}}_{\equiv h_1(\tau_1)} \right] \quad (16)$$

$$\underbrace{\hspace{10em}}_{\equiv h_2(\tau_2)} \\ = \beta \delta_{0, z_1 + z_2 + \dots + z_n} K_n(z_n, z_{n-1}, \dots, z_2) + R(z_n, z_{n-1}, \dots, z_1). \quad (17)$$

As indicated in Eq. (16), we call $h_k(\tau_k)$ the integrand for the $\int_0^{\tau_{k+1}} d\tau_k$ integral for $k = 1, 2, \dots, n$. At $k = 1$ this integrand is given by $h_1(\tau_1) = e^{z_1 \tau_1}$ and we will find h_k for $k = 2, 3, \dots, n$ iteratively. For $z \in \mathbb{C}$, we define the abbreviations $\delta_z \equiv \delta_{0,z}$ and $\Delta_z \equiv (1 - \delta_z)/z$ and consider the integral (for $p = 0, 1, 2, \dots$ and $\tilde{\tau} \geq 0$, proof by partial integration and induction)

$$\int_0^{\tilde{\tau}} d\tau \tau^p e^{z\tau} = \left[\frac{\tilde{\tau}^{p+1}}{p+1} \delta_z + p! (-1)^p \Delta_z^{1+p} \sum_{l=0}^p \frac{(-1)^l}{l!} \Delta_z^{-l} \tilde{\tau}^l \right] e^{z\tilde{\tau}} - p! (-1)^p \Delta_z^{p+1}. \quad (18)$$

Recall that we are only interested in the contribution $K_n(z_n, z_{n-1}, \dots, z_2)$ that fulfills frequency conservation, see Eq. (17). The $\delta_{z_1+z_2+\dots+z_n}$ in front of this term arises from the final τ_n integration of $h_n(\tau_n) \propto e^{(z_1+z_2+\dots+z_n)\tau_n}$ via the first term in Eq. (18). This however requires that all z_k remain in the exponent during the iterative integrations which is violated by the last term in the general integral (18) (which comes from the lower boundary of the integral). All terms in \mathcal{K}_n that stem from this last term in Eq. (18) thus contribute to R and can be dropped in the following [21]. Note however, that it is straightforward to generalize our approach and keep these terms if the full \mathcal{K}_n is desired.

To define the iterative procedure to solve the n -fold integral in Eq. (16), we make the ansatz

$$h_k(\tau_k) = \sum_{l=0}^{k-1} f_k(l) \tau_k^l e^{(z_k+z_{k-1}+\dots+z_1)\tau_k}, \quad (19)$$

which follows from the form of the integral (18) and our decision to disregard the R terms. The ansatz (19) is parameterized by the numbers $f_k(l)$ with $l = 0, 1, \dots, k-1$. These numbers have to be determined iteratively starting from $f_{k=1}(l=0) = 1$, read off from $h_1(\tau_1) = e^{z_1\tau_1}$, c.f. Eq. (16). Iteration rules to obtain the $f_k(l)$ from $f_{k-1}(l)$ are easily derived from Eqns. (16), (18) and (19). We obtain

$$f_k(l) = \sum_{p=0}^{k-1} \tilde{M}_{k-1}(l, p) f_{k-1}(p) \quad (20)$$

which can be understood as a matrix-vector product of $\mathbf{f}_{k-1} = (f_{k-1}(0), f_{k-1}(1), \dots, f_{k-1}(k-2))^T$ with the $k \times (k-1)$ -matrix

$$\tilde{M}_{k-1}(l, p) = \frac{p!}{l!} \left[\delta_{l,p+1} \tilde{\delta}_{k-1} + \theta(p-l+1/2) (-1)^{l+p} \tilde{\Delta}_{k-1}^{1+p-l} \right], \quad (21)$$

where $\tilde{\Delta}_k \equiv \Delta_{z_k+\dots+z_2+z_1}$, $\tilde{\delta}_k \equiv \delta_{z_k+\dots+z_2+z_1}$. The tilde on top of the $\tilde{\delta}_k$ and $\tilde{\Delta}_k$ signals the presence of a sum of z_j in the arguments (below we will define related quantities without tilde for the sum of Ω_j). Note that the first (second) term in brackets of Eq. (21) comes from the first (second) term in square brackets of Eq. (18).

The next step is to find $K_n(z_n, z_{n-1}, \dots, z_2)$. This requires to do the integral $\int_0^\beta d\tau_n h_n(\tau_n)$ which can be again expressed via Eq. (18) but with the replacement $\tilde{\tau} \rightarrow \beta$. Only the first term provides a $\beta \delta_{z_1+z_2+\dots+z_n}$ and is thus identified with K_n . We find:

$$K_n(z_n, z_{n-1}, \dots, z_2) = \sum_{l=0}^{n-1} \frac{\beta^l f_n(l)}{l+1}. \quad (22)$$

The argument z_1 that the right hand side of Eq. (22) depends on is to be replaced by $z_1 = -z_2 - z_3 - \dots - z_n$ in line with the arguments in $K_n(z_n, z_{n-1}, \dots, z_2)$. Then, to conform with Eq. (15), we switch back to $\Omega_j = z_{n-j+1}$ for $j = 1, 2, \dots, n-1$. This amounts to replacing the terms $\tilde{\delta}_j$ and $\tilde{\Delta}_j$ that appear in $f_n(l)$ as follows:

$$\tilde{\delta}_j = \delta_{z_j+\dots+z_2+z_1} = \delta_{\Omega_1+\Omega_2+\dots+\Omega_{n-j}} \equiv \delta_{n-j}, \quad (23)$$

$$-\tilde{\Delta}_j = -\Delta_{z_j+\dots+z_2+z_1} = \Delta_{\Omega_1+\Omega_2+\dots+\Omega_{n-j}} \equiv \Delta_{n-j}. \quad (24)$$

Finally, we can express Eq. (22) using a product of $n-1$ matrices \tilde{M} multiplying the initial length-1 vector with entry $f_1(0) = 1$. Transferring to the Ω -notation by using Eqns. (23) and (24), we obtain

$$K_n(\Omega_1, \dots, \Omega_{n-1}) = \sum_{i_{n-1}=0}^{n-1} \sum_{i_{n-2}=0}^{n-2} \dots \sum_{i_2=0}^2 \sum_{i_1=0}^1 \frac{\beta^{i_{n-1}}}{i_{n-1}+1} M_1(i_{n-1}, i_{n-2}) M_2(i_{n-2}, i_{n-3}) \dots M_{n-2}(i_2, i_1) M_{n-1}(i_1, 0) \quad (25)$$

with

$$M_j(l, p) \equiv \frac{p!}{l!} \left[\delta_{l,p+1} \delta_j - \theta(p-l+1/2) \Delta_j^{1+p-l} \right]. \quad (26)$$

The closed form expression (25) of the universal kernel, to be used in the spectral representation (15), is our main result.

#anom.	$K_2(\Omega_1)$	$K_3(\Omega_1, \Omega_2)$	$K_4(\Omega_1, \Omega_2, \Omega_3)$	$K_5(\Omega_1, \Omega_2, \Omega_3, \Omega_4)$	factor for entire row
$a = 0$	$-\Delta_1$	$+\Delta_1\Delta_2$	$-\Delta_1\Delta_2\Delta_3$	$+\Delta_1\Delta_2\Delta_3\Delta_4$	1
$a = 1$	$+\delta_1$	$-\delta_1\Delta_2$	$+\delta_1\Delta_2\Delta_3$	$-\delta_1\Delta_2\Delta_3\Delta_4$	$\frac{\beta}{2}$
		$-\Delta_1\delta_2$	$+\Delta_1\delta_2\Delta_3$	$-\Delta_1\delta_2\Delta_3\Delta_4$	$\frac{\beta}{2} + \Delta_1$
			$+\Delta_1\Delta_2\delta_3$	$-\Delta_1\Delta_2\delta_3\Delta_4$	$\frac{\beta}{2} + \Delta_1 + \Delta_2$
				$-\Delta_1\Delta_2\Delta_3\delta_4$	$\frac{\beta}{2} + \Delta_1 + \Delta_2 + \Delta_3$
$a = 2$		$+\delta_1\delta_2$	$-\delta_1\delta_2\Delta_3$	$+\delta_1\delta_2\Delta_3\Delta_4$	$\frac{\beta}{2}\frac{\beta}{3}$
			$-\delta_1\Delta_2\delta_3$	$+\delta_1\Delta_2\delta_3\Delta_4$	$\frac{\beta}{2}\left(\frac{\beta}{3} + \Delta_2\right)$
			$-\Delta_1\delta_2\delta_3$	$+\Delta_1\delta_2\delta_3\Delta_4$	$\frac{\beta}{2}\frac{\beta}{3} + \Delta_1\left(\frac{\beta}{2} + \Delta_1\right)$
				$+\delta_1\Delta_2\Delta_3\delta_4$	$\frac{\beta}{2}\left(\frac{\beta}{3} + \Delta_2 + \Delta_3\right)$
				$+\Delta_1\delta_2\Delta_3\delta_4$	$\frac{\beta}{2}\frac{\beta}{3} + (\Delta_1 + \Delta_3)\left(\frac{\beta}{2} + \Delta_1\right)$
				$+\Delta_1\Delta_2\Delta_3\delta_4$	$\frac{\beta}{2}\frac{\beta}{3} + (\Delta_1 + \Delta_2)\left(\frac{\beta}{2} + \Delta_2\right) + \Delta_1^2$
$a = 3$			$+\delta_1\delta_2\delta_3$	$-\delta_1\delta_2\delta_3\Delta_4$	$\frac{\beta}{2}\frac{\beta}{3}\frac{\beta}{4}$
				$-\delta_1\delta_2\Delta_3\delta_4$	$\frac{\beta}{2}\frac{\beta}{3}\left(\frac{\beta}{4} + \Delta_3\right)$
				$-\delta_1\Delta_2\delta_3\delta_4$	$\frac{\beta}{2}\left(\frac{\beta}{3}\frac{\beta}{4} + \Delta_2\left(\frac{\beta}{3} + \Delta_2\right)\right)$
				$-\Delta_1\delta_2\delta_3\delta_4$	$\frac{\beta}{2}\frac{\beta}{3}\frac{\beta}{4} + \Delta_1\left(\frac{\beta}{2}\frac{\beta}{3} + \Delta_1\left(\frac{\beta}{2} + \Delta_1\right)\right)$
$a = 4$			$+\delta_1\delta_2\delta_3\delta_4$	$\frac{\beta}{2}\frac{\beta}{3}\frac{\beta}{4}\frac{\beta}{5}$	

Table I. Universal kernel functions $K_n(\Omega_1, \dots, \Omega_{n-1})$ for $n = 2, 3, 4, 5$ defined in Eq. (14) and calculated from Eq. (25) in Sec. IV. The entries in each column are to be multiplied by the factor from the rightmost column and then summed vertically, see Eqns. (29) and (30) as examples. The symbols δ_j and Δ_j appearing are defined in Eqns. (27) and (28). The rows are organized with respect to the number a of appearances of δ_j , i.e. the order of the anomalous terms.

V. EXAMPLES FOR KERNEL FUNCTION $K_n(\Omega_1, \dots, \Omega_{n-1})$ FOR $n = 2, 3, 4, 5$

While the previous section gives a closed form expression for kernel functions of arbitrary order, we here evaluate the universal kernel functions $K_n(\Omega_1, \dots, \Omega_{n-1})$ defined in Eq. (14) from Eq. (25) for $n = 2, 3, 4, 5$ and show the result in Tab. I. In each column, the kernel function in the top row is obtained by first multiplying the entries with the common factor in the rightmost column and then taking the sum. The symbols δ_j and Δ_j appearing in Tab. I defined for $j = 1, 2, \dots, n-1$ are the same as in Eq. (26) defined by

$$\delta_j \equiv \delta_{\Omega_1 + \Omega_2 + \dots + \Omega_j, 0}, \quad (27)$$

$$\Delta_j \equiv \Delta_{\Omega_1 + \Omega_2 + \dots + \Omega_j} \equiv \frac{1 - \delta_{\Omega_1 + \Omega_2 + \dots + \Omega_j, 0}}{\Omega_1 + \Omega_2 + \dots + \Omega_j}. \quad (28)$$

As an example, for $n = 2$ and $n = 3$ we obtain from Tab. I

$$K_2(\Omega_1) = -\Delta_{\Omega_1} + \frac{\beta}{2}\delta_{\Omega_1}, \quad (29)$$

$$K_3(\Omega_1, \Omega_2) = +\Delta_{\Omega_1}\Delta_{\Omega_1 + \Omega_2} - \frac{\beta}{2}\delta_{\Omega_1}\Delta_{\Omega_2} - \Delta_{\Omega_1}\delta_{\Omega_1 + \Omega_2}\left(\frac{\beta}{2} + \Delta_{\Omega_1}\right) + \delta_{\Omega_1}\delta_{\Omega_2}\frac{\beta}{2}\frac{\beta}{3}, \quad (30)$$

respectively. The rows of Tab. I are organized with respect to the number a of factors δ_l in the summands. Here, $a = 0$ indicates the regular part and $a = 1, 2, \dots, n-1$ indicates anomalous terms. There are $\binom{n-1}{a}$ anomalous terms of order a . Our results are exact and substantially go beyond existing expressions in the literature which are limited to $n \leq 3$ [18] or to fermionic $n = 4$ [19–21] with $a = 0, 1$ (and $a = 2, 3$ guaranteed to vanish, see below) or arbitrary n with $a = 0$ [21]. Alternative expressions for the $n = 3, 4$ kernel functions with $a \leq 1$ were given in [21], but they are consistent with our kernel functions as they yield the same correlation functions, see the Appendix.

In the case of purely fermionic correlators (all A_k fermionic), individual Matsubara frequencies ω_k cannot be zero and thus the $\Omega_k^{a,b} \equiv i\omega_k + E_a - E_b$ of Eq. (12) always have a finite imaginary part and are non-zero, regardless of the eigenenergies. In this case, only sums of an even number of frequencies can be zero, and we can simplify $\delta_1 = \delta_3 = \delta_5 = \dots = 0$. The expressions for the kernels in Tab. I, now denoted by $K_n|_F$ for the fermionic case, simplify to

$$K_2(\Omega_1)|_F = -\Delta_1, \quad (31)$$

$$K_4(\Omega_1, \Omega_2, \Omega_3)|_F = \Delta_1 \Delta_3 \left[\delta_2 \left(\frac{\beta}{2} + \Delta_1 \right) - \Delta_2 \right], \quad (32)$$

$$K_6(\Omega_1, \dots, \Omega_5)|_F = \Delta_1 \Delta_3 \Delta_5 \left\{ -\Delta_2 \Delta_4 - \delta_2 \delta_4 \left[\frac{\beta}{2} \frac{\beta}{3} + (\Delta_1 + \Delta_3) \left(\frac{\beta}{2} + \Delta_1 \right) \right] \right. \\ \left. + \delta_4 \Delta_2 \left(\frac{\beta}{2} + \Delta_1 + \Delta_2 + \Delta_3 \right) + \delta_2 \Delta_4 \left(\frac{\beta}{2} + \Delta_1 \right) \right\}. \quad (33)$$

This concludes the general part of this work. Next, we consider two example systems frequently discussed in the condensed matter theory literature. Using our formalism, we provide analytical forms of correlation functions that to the best of our knowledge were not available before.

VI. APPLICATIONS: HUBBARD ATOM AND FREE SPIN

A. Fermionic Hubbard atom

The Hubbard atom (HA) describes an isolated impurity or otherwise localized system with Hamiltonian

$$H = \epsilon(n_\uparrow + n_\downarrow) + U n_\uparrow n_\downarrow - h(n_\uparrow - n_\downarrow), \quad (34)$$

see Fig. 1(b) for a sketch. The HA corresponds to the limit of vanishing system-bath coupling in the Anderson impurity model (AIM) or vanishing hopping in the Hubbard model (HM). The particle number operators $n_\sigma = d_\sigma^\dagger d_\sigma$ count the number of fermionic particles with spin $\sigma \in \{\uparrow, \downarrow\}$, each contributing an onsite energy ϵ shifted by an external magnetic field h in z -direction. An interaction energy U is associated to double occupation.

Due to its simplicity and the four-dimensional Hilbert space, the correlation functions for the HA can be found analytically using the spectral representation. It is therefore often used for benchmarking [3, 27, 28]. The presence of the interaction term leads to a non-vanishing $n = 4$ one-line irreducible vertex function. The HA serves as an important reference point to study and interpret properties of the AIM and HM beyond the one-particle level, for example divergences of two-line irreducible vertex functions [29–32] and signatures of the local moment formation in generalized susceptibilities [16, 33]. Using the fermionic kernels in Eqns. (31) and (32), we have checked that our formalism reproduces the results for the 2-point and 4-point correlators given in Refs. [19, 26] for half-filling, $\epsilon = -U/2$ and $h = 0$.

Correlation functions including bosonic operators describe the asymptotic behaviour of the $n = 4$ fermion vertex for large frequencies [34] or the interaction of electrons by the exchange of an effective boson [35, 36]. These relations involve correlation functions of two bosonic operators or of one bosonic and two fermionic operators, giving rise to expressions possibly anomalous in at most one frequency argument, i.e. $a \leq 1$.

For the HA, AIM and HM, bosonic correlation functions for $n > 2$ have not been considered thoroughly so far. Only recently, steps in this direction were taken, particularly in the context of non-linear response theory [3]. The response of a system to external perturbation is described by 2- and 3-point correlation functions to linear and second order in the perturbation, respectively. For the HA, physically motivated perturbations are applied to the onsite energy via a term $\delta_\epsilon n$ or via a magnetic field $\delta_h \cdot \mathbf{S}$ with

$$n = n_\uparrow + n_\downarrow, \quad S^x = \frac{1}{2} (d_\uparrow^\dagger d_\downarrow + d_\downarrow^\dagger d_\uparrow), \quad S^y = \frac{-i}{2} (d_\uparrow^\dagger d_\downarrow - d_\downarrow^\dagger d_\uparrow), \quad S^z = \frac{1}{2} (n_\uparrow - n_\downarrow). \quad (35)$$

The resulting changes of the expectation values of the density or magnetization in arbitrary direction are described in second order of the perturbation by the connected parts of the correlation functions $G_{A_1 A_2 A_3}(\tau_1, \tau_2, \tau_3)$ with $A_i \in \{n, S_x, S_y, S_z\}$, where the time-ordered expectation value is evaluated with respect to the unperturbed system (34) and Fourier transformed to the frequencies of interest. These objects have been studied numerically in Ref. [3]. In the following, we give explicit, analytic expressions of the full correlation functions $G_{A_1 A_2 A_3}(\omega_1, \omega_2)$ (i.e. including disconnected parts), for arbitrary parameters ϵ , U and h and for all possible operator combinations using the (bosonic) kernel function K_3 , see Eq. (30). To the best of our knowledge, these expressions have not been reported before.

The eigenstates of the HA Hamiltonian (34) [see Fig. 1(b)] describe an empty ($|0\rangle$), singly occupied ($d_{\uparrow}^{\dagger}|0\rangle = |\uparrow\rangle$, $d_{\downarrow}^{\dagger}|0\rangle = |\downarrow\rangle$) or doubly occupied ($d_{\uparrow}^{\dagger}d_{\downarrow}^{\dagger}|0\rangle = |\uparrow\downarrow\rangle$) impurity with eigenenergies $E_0 = 0$, $E_{\uparrow} = \epsilon - h$, $E_{\downarrow} = \epsilon + h$ and $E_{\uparrow\downarrow} = 2\epsilon + U$, respectively. The partition function is $Z = 1 + e^{-\beta(\epsilon-h)} + e^{-\beta(\epsilon+h)} + e^{-\beta(2\epsilon+U)}$. We define

$$s = \frac{e^{-\beta\epsilon}}{Z} \sinh(\beta h), \quad c = \frac{e^{-\beta\epsilon}}{Z} \cosh(\beta h), \quad (36)$$

and obtain all non-vanishing bosonic 3-point correlation functions (where $\omega_3 = -\omega_1 - \omega_2$):

$$G_{nnn}(\omega_1, \omega_2) = 2\beta^2 \delta_{\omega_1} \delta_{\omega_2} \left(\frac{4e^{-\beta(2\epsilon+U)}}{Z} + c \right), \quad (37)$$

$$G_{nmS^z}(\omega_1, \omega_2) = \beta^2 \delta_{\omega_1} \delta_{\omega_2} s, \quad (38)$$

$$G_{nS^xS^y}(\omega_1, \omega_2) = -\beta \delta_{\omega_1} s \frac{\omega_2}{\omega_2^2 + 4h^2}, \quad (39)$$

$$G_{nS^xS^x}(\omega_1, \omega_2) = G_{nS^yS^y}(\omega_1, \omega_2) = 2\beta \delta_{\omega_1} \frac{h s}{\omega_2^2 + 4h^2}, \quad (40)$$

$$G_{nS^zS^z}(\omega_1, \omega_2) = \frac{\beta^2}{2} \delta_{\omega_1} \delta_{\omega_2} c, \quad (41)$$

$$G_{S^zS^xS^x}(\omega_1, \omega_2) = G_{S^zS^yS^y}(\omega_1, \omega_2) = -s \frac{\omega_2 \omega_3 + 4h^2}{(\omega_2^2 + 4h^2)(\omega_3^2 + 4h^2)} + \beta \delta_{\omega_1} \frac{h c}{\omega_2^2 + 4h^2}, \quad (42)$$

$$G_{S^zS^zS^z}(\omega_1, \omega_2) = \frac{\beta^2}{4} \delta_{\omega_1} \delta_{\omega_2} s, \quad (43)$$

$$G_{S^xS^yS^z}(\omega_1, \omega_2) = 2h s \frac{\omega_1 - \omega_2}{(\omega_1^2 + 4h^2)(\omega_2^2 + 4h^2)} - \frac{\beta}{2} \delta_{\omega_3} c \frac{\omega_1}{\omega_1^2 + 4h^2}. \quad (44)$$

We observe that each conserved quantity, in this case n and S_z , contributes an anomalous term $\propto \delta_{\omega_k}$ in its respective frequency argument ω_k . If an operator A_k is conserved $[H, A_k] = 0$, the basis over which we sum in Eq. (15) can be chosen such that both H and A_k are diagonal, $A_k^{\frac{12}{k}} = A_k^{\frac{1}{k}} \delta_{1,2}$. Thus there will always be vanishing eigenenergy differences leading to the appearance of an anomalous contribution. If the operators in the correlator are additionally commuting with each other, in our case for example $[n, S^z] = 0$, there exists a basis in which all operators and the Hamiltonian are diagonal, giving rise to correlation functions anomalous in all frequency arguments.

In the limit of vanishing field $h \rightarrow 0$, we introduce an additional degeneracy $E_{\uparrow} = E_{\downarrow} = \epsilon$ in the system, potentially resulting in additional anomalous contributions. The corresponding correlation functions can then be obtained by taking appropriate limits, for example

$$\lim_{h \rightarrow 0} \frac{h \sinh(\beta h)}{\omega_k^2 + 4h^2} = \frac{\beta}{4} \delta_{\omega_k}, \quad (45)$$

resulting in

$$G_{nmn}(\omega_1, \omega_2) = \beta^2 \delta_{\omega_1} \delta_{\omega_2} \frac{2(4e^{-\beta(2\epsilon+U)} + e^{-\beta\epsilon})}{Z}, \quad (46)$$

$$G_{nS^{\alpha}S^{\alpha}}(\omega_1, \omega_2) = \beta^2 \delta_{\omega_1} \delta_{\omega_2} \frac{e^{-\beta\epsilon}}{2Z} \quad (\alpha \in \{x, y, z\}), \quad (47)$$

$$G_{S^xS^yS^z}(\omega_1, \omega_2) = \beta \frac{e^{-\beta\epsilon}}{2Z} (-\delta_{\omega_1} \Delta_{\omega_2} + \delta_{\omega_2} \Delta_{\omega_1} - \delta_{\omega_1+\omega_2} \Delta_{\omega_1}), \quad (48)$$

with all other correlation functions vanishing. As already pointed out in Ref. [3], only the last correlation function retains a nontrivial frequency dependence due to non-commuting operators.

B. Free spin S

We now consider correlation functions of a free spin of length S , without a magnetic field so that temperature $1/\beta$ is the only energy scale. The operators $\{S^{\alpha}\}_{\alpha=x,y,z}$ fulfill $S^xS^x + S^yS^y + S^zS^z = S(S+1)$ and the SU(2) algebra $[S^{\alpha_1}, S^{\alpha_2}] = i \sum_{\alpha_3=\{x,y,z\}} \epsilon^{\alpha_1\alpha_2\alpha_3} S^{\alpha_3}$, thus they are bosonic. Since the Hamiltonian vanishes and therefore

$n = 2$	$G_{S^+S^-}(\omega) = G_{S^zS^z}(\omega) = \beta\delta_\omega b_1$
$n = 3$	$G_{S^+S^-S^z}(\omega_1, \omega_2) = \beta b_1(-\delta_{\omega_1}\Delta_{i\omega_2} + \delta_{\omega_2}\Delta_{i\omega_1} + \delta_{\omega_1+\omega_2}\Delta_{i\omega_2}) = -iG_{S^xS^yS^z}(\omega_1, \omega_2)$
$n = 4$	$G_{S^zS^zS^zS^z}(\omega_1, \omega_2, \omega_3) = \delta_{\omega_1}\delta_{\omega_2}\delta_{\omega_3}\beta^3 b_3$
	$G_{S^+S^+S^-S^-}(\omega_1, \omega_2, \omega_3) = \beta b_1[2 \times \delta_{\omega_1}\delta_{\omega_2}\delta_{\omega_3} \times \frac{\beta^2}{5}(3b_1 - \frac{1}{3}) + r]$
	$G_{S^+S^-S^zS^z}(\omega_1, \omega_2, \omega_3) = \beta b_1[1 \times \delta_{\omega_1}\delta_{\omega_2}\delta_{\omega_3} \times \frac{\beta^2}{5}(3b_1 - \frac{1}{3}) - r]$
	$r = \Delta_{i\omega_1}\Delta_{i\omega_2}(\delta_{\omega_1+\omega_3} + \delta_{\omega_2+\omega_3} - \delta_{\omega_3} - \delta_{\omega_4}) - (\delta_{\omega_1}\Delta_{i\omega_2}^2 + \delta_{\omega_2}\Delta_{i\omega_1}^2)(\delta_{\omega_3} + \delta_{\omega_4}) - \Delta_{i\omega_3}\Delta_{i\omega_4}(\delta_{\omega_1} + \delta_{\omega_2})$

Table II. Matsubara correlation functions for a free spin- S up to order $n = 4$. Here, $\omega_4 = -\omega_1 - \omega_2 - \omega_3$.

all eigenenergies are zero, every $\Omega_k^{a,b}$ in the spectral representation (15) can vanish and a proper treatment of all anomalous terms is essential. As the Heisenberg time dependence is trivial, $S^\alpha(\tau) = S^\alpha$, the non-trivial frequency dependence of the correlators, which can be non-vanishing at any order n , derives solely from the action of time ordering.

The correlators are required, for example, as the non-trivial initial condition for the spin-fRG recently suggested by Kopietz et al., Refs. [13, 37–40]. However, for $n > 3$ they are so far only partially available: They are either given for restricted frequency combinations, for the purely classical case $S^{\alpha_1} = S^{\alpha_2} = \dots = S^{\alpha_n}$ where the SU(2) algebra does not matter, or for finite magnetic field via an equation of motion [37] or diagrammatic approach [41, 42].

We define the spin raising and lowering operators,

$$S^\pm = (S^x \pm iS^y)/\sqrt{2}, \quad (49)$$

which have to appear in pairs for a non-vanishing correlator due to spin-rotation symmetry. As for the HA, we do not consider connected correlators in this work for brevity. The classical S^z -correlator can be found from its generating functional with source field h [13],

$$\mathcal{G}(y = \beta h) = \frac{\sinh[y(S + 1/2)]}{(2S + 1) \sinh[y/2]}, \quad (50)$$

$$\langle (S^z)^l \rangle = \lim_{y \rightarrow 0} \partial_y^l \mathcal{G}(y) \equiv b_{l-1}, \quad (51)$$

for example $b_1 = \frac{S}{3}(S + 1)$ and $b_3 = \frac{S}{15}(3S^3 + 6S^2 + 2S - 1)$ and vanishing b_l for even l . For all other correlators, we adapt Eq. (15) for the free spin case,

$$G_{S^{\alpha_1}S^{\alpha_2}\dots S^{\alpha_n}}(\omega_1, \dots, \omega_{n-1}) = \sum_{p \in S_n} \langle S^{\alpha_{p(1)}} S^{\alpha_{p(2)}} \dots S^{\alpha_{p(n)}} \rangle K_n(i\omega_{p(1)}, i\omega_{p(2)}, \dots, i\omega_{p(n-1)}), \quad (52)$$

where we made use of the fact that all eigenenergies are zero and the Heisenberg time evolution is trivial. It is convenient to evaluate the equal-time correlators in Eq. (52) as

$$\langle S^{\alpha_1} S^{\alpha_2} \dots S^{\alpha_n} \rangle = \frac{1}{2S + 1} \sum_{m=-S}^S \langle m | S^{\alpha_1} S^{\alpha_2} \dots S^{\alpha_n} | m \rangle \equiv \frac{1}{2S + 1} \sum_{m=-S}^S \sum_{l=0}^n p_l m^l = p_0 + \sum_{l=2}^n p_l b_{l-1} \quad (53)$$

where in the last step we used Eq. (51). We find the real expansion coefficients $\{p_l\}_{l=0,1,\dots,n}$ iteratively by moving through the string $\alpha_1\alpha_2\dots\alpha_n$ from the right and start from $p_l = \delta_{0,l}$. Based on the S^z eigenstates $\{|m\rangle\}_{m=-S,\dots,S-1,S}$ we obtain the iteration rules from $S^z|m\rangle = m|m\rangle$ and $S^\pm|m\rangle = \sqrt{1/2\sqrt{S(S+1) - m(m \pm 1)}}|m \pm 1\rangle$. We define an auxiliary integer c that keeps track of the intermediate state $|m + c\rangle$, initially $c = 0$. Depending on the α_j that we find in step $j = n, n - 1, \dots, 1$ we take one of the following actions: (i) For $\alpha_j = z$, we update $p_l \leftarrow p_{l-1} + c p_l \forall l$ and leave c unchanged. It is understood that $p_{l < 0} = 0$. (ii) For $\alpha_j = +$, we combine the square-root factor brought by the raising operator with the factor that comes from the necessary $\alpha_{j'} = -$ at another place in the string. We replace $p_l \leftarrow -\frac{1}{2}p_{l-2} - \frac{2c+1}{2}p_{l-1} + (\frac{3}{2}b_1 - c\frac{c+1}{2})p_l \forall l$ and then let $c \leftarrow c + 1$. (iii) For $\alpha_j = -$, we update $c \leftarrow c - 1$ and keep p_l unchanged, $p_l \leftarrow p_l \forall l$.

Our final results for the free spin correlators are reported in Tab. II. We reproduce the known spin correlators for $n = 2, 3$ and determine the non-classical correlators $G_{S^+S^+S^-S^-}$ and $G_{S^+S^-S^zS^z}$ at order $n = 4$ which to the best of our knowledge were not available in the literature [43]. We also confirmed the classical result for $G_{S^zS^zS^zS^z}$ which in our full quantum formalism requires some non-trivial cancellations. To arrive at our results, we used the identity

$$\Delta_{a+b}(\Delta_a + \Delta_b) - \Delta_a\Delta_b = \delta_a\Delta_b^2 + \delta_b\Delta_a^2 - \delta_{a+b}\Delta_a\Delta_b. \quad (54)$$

Note that $G_{S^+S^-S^z} = -iG_{S^xS^yS^z}$, and our result from Tab. (II), if specialized to $S = 1/2$ (where $b_1 = 1/4$) agrees to the zero-field result found for the HA in Eq. (48) up to a factor $2e^{-\beta\epsilon}/Z$. This factor is the expectation value of the projector to the singly-occupied sector in the HA Hilbert space, the latter is implicit in $S^{x,y,z}$ of Eq. (35).

VII. CONCLUSION

In summary, we have provided exact universal kernel functions for the spectral representation of the n -point Matsubara correlator. Our results are an efficient alternative to equation-of-motion approaches which often have difficulties to capture anomalous terms related to conserved or commuting operators. We expect our results to be useful for various benchmarking applications, as starting points for emerging many-body methods and for unraveling the physical interpretation of n -point functions in various settings. Although our work has focused on imaginary frequency (Matsubara) correlators, with analytical expressions now at hand, it is also interesting to study the intricacies of analytical continuation to real frequencies and thus to further explore the connection of Matsubara and Keldysh correlators [44].

ACKNOWLEDGMENTS

We acknowledge useful discussions with Karsten Held, Friedrich Krien, Seung-Sup Lee, Peter Kopietz, Fabian Kugler, Nepomuk Ritz, Georg Rohringer, Andreas Rückriegel. We thank Andreas Rückriegel for sharing unpublished results on 4-point free spin correlators and pointing out further simplifications of the analytical expressions. BS and BS are supported by a MCQST-START fellowship. We acknowledge funding from the International Max Planck Research School for Quantum Science and Technology (IMPRSQST) for JH, from the Deutsche Forschungsgemeinschaft under Germany's Excellence Strategy EXC-2111 (Project No. 390814868), and from the Munich Quantum Valley, supported by the Bavarian state government with funds from the Hightech Agenda Bayern Plus.

APPENDIX: EQUIVALENCE TO CONVENTION OF REF. [21]

In Ref. [21] by Kugler, Lee and von Delft (KLD), only regular ($a = 0$) and anomalous terms of order $a = 1$ have been considered for $n = 3, 4$. The corresponding kernel functions were derived from only $(n - 1)!$ permutations by setting $\tau_n = 0$ and $\tau_{i \neq n} > 0$, but still applied to all $n!$ permutations to obtain the correlation functions. For $n = 3$, the resulting kernel function (Eq. (45) in Ref. [21]) reads

$$K_{3,\text{KLD}}(\Omega_1, \Omega_2) = \Delta_1\Delta_2 - \Delta_1\delta_2\frac{1}{2}(\beta + \Delta_1) - \delta_1\Delta_2\frac{1}{2}(\beta + \Delta_2). \quad (55)$$

This can be compared to the corresponding kernel function for $n = 3$ found in our Eq. (30) truncated to $a \leq 1$,

$$K_3^{a \leq 1}(\Omega_1, \Omega_2) = \Delta_1\Delta_2 - \Delta_1\delta_2\left(\frac{\beta}{2} + \Delta_1\right) - \frac{\beta}{2}\delta_1\Delta_2. \quad (56)$$

Both approaches are equally valid and should yield the same correlation functions (consistently discarding terms with $a = 2$), yet the kernel functions are obviously different. To resolve this issue, we define the difference of the kernel functions

$$K_{3,\text{diff}}(\Omega_1, \Omega_2) = K_{3,\text{KLD}}(\Omega_1, \Omega_2) - K_3^{a \leq 1}(\Omega_1, \Omega_2) = \frac{1}{2}(\Delta_1^2\delta_2 - \delta_1\Delta_2^2) \quad (57)$$

and show that the corresponding contributions to the correlation function vanishes when summed over cyclically

related permutations $p = 123, 231, 312$. These contributions are given by

$$\begin{aligned}
& \frac{1}{Z} \sum_{p=123,231,312} \zeta(p) \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_{p(1)}^{12} A_{p(2)}^{23} A_{p(3)}^{31} K_{3,\text{diff}}(\Omega_{p(1)}^{12}, \Omega_{p(2)}^{23}) \\
&= \frac{\zeta(123)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_1^{12} A_2^{23} A_3^{31} \left(\frac{(1 - \delta_{\omega_1} \delta_{E_{\underline{1}} - E_{\underline{2}}}) \delta_{\omega_1 + \omega_2} \delta_{E_{\underline{1}} - E_{\underline{3}}}}{(i\omega_1 + E_{\underline{1}} - E_{\underline{2}})^2} - \frac{\delta_{\omega_1} \delta_{E_{\underline{1}} - E_{\underline{2}}} (1 - \delta_{\omega_1 + \omega_2} \delta_{E_{\underline{1}} - E_{\underline{3}}})}{(i\omega_1 + i\omega_2 + E_{\underline{1}} - E_{\underline{3}})^2} \right) \\
&+ \frac{\zeta(231)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_2^{12} A_3^{23} A_1^{31} \left(\frac{(1 - \delta_{\omega_2} \delta_{E_{\underline{1}} - E_{\underline{2}}}) \delta_{\omega_2 + \omega_3} \delta_{E_{\underline{1}} - E_{\underline{3}}}}{(i\omega_2 + E_{\underline{1}} - E_{\underline{2}})^2} - \frac{\delta_{\omega_2} \delta_{E_{\underline{1}} - E_{\underline{2}}} (1 - \delta_{\omega_2 + \omega_3} \delta_{E_{\underline{1}} - E_{\underline{3}}})}{(i\omega_2 + i\omega_3 + E_{\underline{1}} - E_{\underline{3}})^2} \right) \\
&+ \frac{\zeta(312)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_3^{12} A_1^{23} A_2^{31} \left(\frac{(1 - \delta_{\omega_3} \delta_{E_{\underline{1}} - E_{\underline{2}}}) \delta_{\omega_3 + \omega_1} \delta_{E_{\underline{1}} - E_{\underline{3}}}}{(i\omega_3 + E_{\underline{1}} - E_{\underline{2}})^2} - \frac{\delta_{\omega_3} \delta_{E_{\underline{1}} - E_{\underline{2}}} (1 - \delta_{\omega_3 + \omega_1} \delta_{E_{\underline{1}} - E_{\underline{3}}})}{(i\omega_3 + i\omega_1 + E_{\underline{1}} - E_{\underline{3}})^2} \right). \tag{58}
\end{aligned}$$

Considering the second term of permutation $p = 312$ and renaming the summation variables $\underline{2} \rightarrow \underline{1}, \underline{3} \rightarrow \underline{2}, \underline{1} \rightarrow \underline{3}$ yields

$$\begin{aligned}
& - \frac{\zeta(312)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_3^{12} A_1^{23} A_2^{31} \frac{\delta_{\omega_3} \delta_{E_{\underline{1}} - E_{\underline{2}}} (1 - \delta_{\omega_3 + \omega_1} \delta_{E_{\underline{1}} - E_{\underline{3}}})}{i(\omega_3 + \omega_1) + E_{\underline{1}} - E_{\underline{3}}} \\
&= - \frac{\zeta(312)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{3}}} A_1^{12} A_2^{23} A_3^{31} \frac{\delta_{\omega_3} \delta_{E_{\underline{3}} - E_{\underline{1}}} (1 - \delta_{\omega_3 + \omega_1} \delta_{E_{\underline{3}} - E_{\underline{2}}})}{(i\omega_3 + i\omega_1 + E_{\underline{3}} - E_{\underline{2}})^2} \\
&= - \frac{\zeta(123)}{2Z} \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_1^{12} A_2^{23} A_3^{31} \frac{\delta_{\omega_1 + \omega_2} \delta_{E_{\underline{1}} - E_{\underline{3}}} (1 - \delta_{\omega_1} \delta_{E_{\underline{1}} - E_{\underline{2}}})}{(i\omega_1 + E_{\underline{1}} - E_{\underline{2}})^2}, \tag{59}
\end{aligned}$$

where we used $\omega_3 = -\omega_1 - \omega_2$ and the fact that δ_{ω_3} enforces the third operator to be bosonic, such that $\zeta(312) = \zeta(123)$. This term exactly cancels the first contribution of permutation $p = 123$ in (58). Repeating similar steps for the remaining terms, we find that the the second term of $p = 123$ and the first term of $p = 231$ as well as the second term of $p = 231$ and the first term of $p = 312$ cancel, leading to

$$\frac{1}{Z} \sum_{p \in \{123, 231, 312\}} \zeta(p) \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_{p(1)}^{12} A_{p(2)}^{23} A_{p(3)}^{31} K_{3,\text{diff}}(\Omega_{p(1)}^{12}, \Omega_{p(2)}^{23}) = 0. \tag{60}$$

Similarly, summing over the second set of cyclicly related permutations $p = 132, 213, 321$ leads to a vanishing result, leading to the conclusion that

$$\frac{1}{Z} \sum_{p \in S_3} \zeta(p) \sum_{\underline{123}} e^{-\beta E_{\underline{1}}} A_{p(1)}^{12} A_{p(2)}^{23} A_{p(3)}^{31} K_{3,\text{diff}}(\Omega_{p(1)}^{12}, \Omega_{p(2)}^{23}) = 0. \tag{61}$$

Thus we have shown that both kernel functions in Eqns. (55) and (56) are equivalent as they yield the same correlation functions after summing over all permutations. The same statement holds true for case of $n = 4$ and $a = 1$.

-
- [1] J. W. Negele and H. Orland. *Quantum Many-Particle Systems*. Westview, 1988.
 - [2] H. Bruus and K. Flensberg. *Many-Body Quantum Theory in Condensed Matter Physics*. Oxford Graduate Texts, 2004.
 - [3] P. Kappl, F. Krien, C. Watzenböck, and K. Held. Non-linear responses and three-particle correlators in correlated electron systems exemplified by the Anderson impurity model. (arXiv:2212.11877), December 2022. doi:10.48550/arXiv.2212.11877.
 - [4] G. Semeghini, H. Levine, A. Keesling, S. Ebadi, T. T. Wang, D. Bluvstein, R. Verresen, H. Pichler, M. Kalinowski, R. Samajdar, A. Omran, S. Sachdev, A. Vishwanath, M. Greiner, V. Vuletić, and M. D. Lukin. Probing topological spin liquids on a programmable quantum simulator. *Science*, 374(6572):1242–1247, December 2021. doi:10.1126/science.abi8794.
 - [5] L. Hedin. New Method for Calculating the One-Particle Green's Function with Application to the Electron-Gas Problem. *Phys. Rev.*, 139(3A):A796–A823, August 1965. doi:10.1103/PhysRev.139.A796.
 - [6] N.E. Bickers. *Self-Consistent Many-Body Theory for Condensed Matter Systems*. Springer.
 - [7] B. Roulet, J. Gavoret, and P. Nozières. Singularities in the X-Ray Absorption and Emission of Metals. I. First-Order Parquet Calculation. *Phys. Rev.*, 178(3):1072–1083, February 1969. doi:10.1103/PhysRev.178.1072.
 - [8] P. Kopietz, L. Bartosch, and F. Schütz. *Introduction to the Functional Renormalization Group*. Springer, 2010.

- [9] W. Metzner, M. Salmhofer, C. Honerkamp, V. Meden, and K. Schönhammer. Functional renormalization group approach to correlated fermion systems. *Rev. Mod. Phys.*, 84:299, March 2012. doi:10.1103/RevModPhys.84.299.
- [10] C. Wetterich. Exact evolution equation for the effective potential. *Physics Letters B*, 301(1):90–94, February 1993. ISSN 0370-2693. doi:10.1016/0370-2693(93)90726-X.
- [11] S.-S. Lee, F. Kugler, and J. von Delft. Computing Local Multipoint Correlators Using the Numerical Renormalization Group. *Phys. Rev. X*, 11(4):041007, October 2021. doi:10.1103/PhysRevX.11.041007.
- [12] J. Reuther and R. Thomale. Cluster functional renormalization group. *Phys. Rev. B*, 89(2):024412, January 2014. doi:10.1103/PhysRevB.89.024412.
- [13] J. Krieg and P. Kopietz. Exact renormalization group for quantum spin systems. *Phys. Rev. B*, 99(6):060403, February 2019. doi:10.1103/PhysRevB.99.060403.
- [14] A. Rückriegel, J. Arnold, R. Goll, and P. Kopietz. Spin functional renormalization group for dimerized quantum spin systems. *Phys. Rev. B*, 105(22):224406, June 2022. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.105.224406.
- [15] G. Rohringer, H. Hafermann, A. Toschi, A. A. Katanin, A. E. Antipov, M. I. Katsnelson, A. I. Lichtenstein, A. N. Rubtsov, and K. Held. Diagrammatic routes to nonlocal correlations beyond dynamical mean field theory. *Rev. Mod. Phys.*, 90(2):025003, May 2018. doi:10.1103/RevModPhys.90.025003.
- [16] P. Chalupa, T. Schäfer, M. Reitner, D. Springer, S. Andergassen, and A. Toschi. Fingerprints of the Local Moment Formation and its Kondo Screening in the Generalized Susceptibilities of Many-Electron Problems. *Phys. Rev. Lett.*, 126(5):056403, February 2021. doi:10.1103/PhysRevLett.126.056403.
- [17] H. Lehmann. Über Eigenschaften von Ausbreitungsfunktionen und Renormierungskonstanten quantisierter Felder. *Nuovo Cim*, 11(4):342–357, April 1954. ISSN 1827-6121. doi:10.1007/BF02783624.
- [18] A.M. Shvaika. On the spectral relations for multitime correlation functions. 9(3(47)):447–458, 2006.
- [19] H. Hafermann, C. Jung, S. Brener, M. I. Katsnelson, A. N. Rubtsov, and A. I. Lichtenstein. Superperturbation solver for quantum impurity models. *Europhys. Lett.*, 85(2):27007, January 2009. ISSN 0295-5075, 1286-4854. doi:10.1209/0295-5075/85/27007.
- [20] A.M. Shvaika. Spectral properties of four-time fermionic Green's functions. 19(3):33004, 2016. doi:10.5488/CMP.19.33004.
- [21] F. B. Kugler, S.-S. B. Lee, and J. von Delft. Multipoint Correlation Functions: Spectral Representation and Numerical Evaluation. *Phys. Rev. X*, 11(4):041006, October 2021. doi:10.1103/PhysRevX.11.041006.
- [22] P. C. Kwok and T. D. Schultz. Correlation functions and Green functions: Zero-frequency anomalies. *J. Phys. C: Solid State Phys.*, 2(7):1196, July 1969. ISSN 0022-3719. doi:10.1088/0022-3719/2/7/312.
- [23] C. Watzenböck, M. Fellinger, K. Held, and A. Toschi. Long-term memory magnetic correlations in the Hubbard model: A dynamical mean-field theory analysis. *SciPost Phys.*, 12(6):184, June 2022. ISSN 2542-4653. doi:10.21468/SciPostPhys.12.6.184.
- [24] A. Tsvelik. *Quantum Field Theory in Condensed Matter Physics*. Cambridge U. Press, 2007.
- [25] G. Rohringer, A. Valli, and A. Toschi. Local electronic correlation at the two-particle level. *Phys. Rev. B*, 86(12):125114, September 2012. doi:10.1103/PhysRevB.86.125114.
- [26] G. Rohringer. *New Routes towards a Theoretical Treatment of Nonlocal Electronic Correlations*. Thesis, Technische Universität Wien, 2013.
- [27] F. Krien and A. Valli. Parquetlike equations for the Hedin three-leg vertex. *Phys. Rev. B*, 100(24):245147, December 2019. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.100.245147.
- [28] F. Krien, A. Kauch, and K. Held. Tiling with triangles: Parquet and G W γ methods unified. *Phys. Rev. Research*, 3(1):013149, February 2021. ISSN 2643-1564. doi:10.1103/PhysRevResearch.3.013149.
- [29] T. Schäfer, S. Ciuchi, M. Wallerberger, P. Thunström, O. Gunnarsson, G. Sangiovanni, G. Rohringer, and A. Toschi. Nonperturbative landscape of the Mott-Hubbard transition: Multiple divergence lines around the critical endpoint. *Phys. Rev. B*, 94(23):235108, December 2016. doi:10.1103/PhysRevB.94.235108.
- [30] P. Thunström, O. Gunnarsson, Sergio Ciuchi, and G. Rohringer. Analytical investigation of singularities in two-particle irreducible vertex functions of the Hubbard atom. *Phys. Rev. B*, 98(23):235107, December 2018. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.98.235107.
- [31] P. Chalupa, P. Gunacker, T. Schäfer, K. Held, and A. Toschi. Divergences of the irreducible vertex functions in correlated metallic systems: Insights from the Anderson impurity model. *Phys. Rev. B*, 97(24):245136, June 2018. doi:10.1103/PhysRevB.97.245136.
- [32] M. Pelz, S. Adler, M. Reitner, and A. Toschi. The highly nonperturbative nature of the Mott metal-insulator transition: Two-particle vertex divergences in the coexistence region. (arXiv:2303.01914), March 2023. doi:10.48550/arXiv.2303.01914.
- [33] S. Adler, F. Krien, P. Chalupa-Gantner, G. Sangiovanni, and A. Toschi. Non-perturbative intertwining between spin and charge correlations: A "smoking gun" single-boson-exchange result. (arXiv:2212.09693), December 2022. doi:10.48550/arXiv.2212.09693.
- [34] N. Wentzell, G. Li, A. Tagliavini, C. Taranto, G. Rohringer, K. Held, A. Toschi, and S. Andergassen. High-frequency asymptotics of the vertex function: Diagrammatic parametrization and algorithmic implementation. *Phys. Rev. B*, 102(8):085106, August 2020. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.102.085106.
- [35] F. Krien, A. Valli, and M. Capone. Single-boson exchange decomposition of the vertex function. *Phys. Rev. B*, 100(15):155149, October 2019. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.100.155149.
- [36] M. Gievers, E. Walter, A. Ge, J. von Delft, and F. B. Kugler. Multiloop flow equations for single-boson exchange fRG. *Eur. Phys. J. B*, 95(7):108, July 2022. ISSN 1434-6036. doi:10.1140/epjb/s10051-022-00353-6.
- [37] R. Goll, D. Tarasevych, J. Krieg, and P. Kopietz. Spin functional renormalization group for quantum Heisenberg ferromagnets: Magnetization and magnon damping in two dimensions. *Phys. Rev. B*, 100(17):174424, November 2019.

doi:10.1103/PhysRevB.100.174424.

- [38] R. Goll, A. Rückriegel, and P. Kopietz. Zero-magnon sound in quantum Heisenberg ferromagnets. *Phys. Rev. B*, 102(22):224437, December 2020. doi:10.1103/PhysRevB.102.224437.
- [39] D. Tarasevych and P. Kopietz. Dissipative spin dynamics in hot quantum paramagnets. *Phys. Rev. B*, 104(2):024423, July 2021. doi:10.1103/PhysRevB.104.024423.
- [40] D. Tarasevych, A. Rückriegel, S. Keupert, V. Mitsioannou, and P. Kopietz. Spin functional renormalization group for the J1-J2-J3 quantum Heisenberg model. *Phys. Rev. B*, 106(17):174412, November 2022. ISSN 2469-9950, 2469-9969. doi:10.1103/PhysRevB.106.174412.
- [41] V. G. Vaks and S. A. Pikin. Thermodynamics of an ideal ferromagnetic substance. *Soviet Journal of Experimental and Theoretical Physics*, 26(1):188, 1968.
- [42] V. G. Vaks, A. I. Larkin, and S. A. Pikin. Spin Waves and Correlation Functions in a Ferromagnetic. *Soviet Journal of Experimental and Theoretical Physics*, 26:647, March 1968. ISSN 1063-7761.
- [43] We thank Andreas Rückriegel for sharing unpublished results on 4-point free spin correlators and pointing out further simplifications of the analytical expressions.
- [44] A. Ge, J. Halbinger, S.-S. Lee, J. von Delft, and F. B. Kugler. manuscript in preparation.