A CONTINUED FRACTION-HYPERBOLA BASED ATTACK ON **RSA CRYPTOSYSTEM**

GILDA RECH BANSIMBA, REGIS F. BABINDAMANA, AND BASILE GUY R. BOSSOTO

Abstract.

In this paper we present new arithmetical and algebraic results following the work of Babindamana and al. on hyperbolas and describe in the new results an approach to attacking a RSA-type modulus based on continued fractions, independent and not bounded by the size of the private key d nor the public exponent e compared to Wiener's attack. When successful, this attack is bounded by $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$ with $b = 10^y$, $\alpha_{i3} + \alpha_{j3}$ a non trivial factor of n and α_{j4} such that $(n+1)/(n-1) = \alpha_{i4}/\alpha_{j4}$. The primary goal of this attack is to find a point $X_{\alpha} = (-\alpha_3, \alpha_3 + 1) \in \mathbb{Z}^2_{\star}$ that satisfies $\langle X_{\alpha_3}, P_3 \rangle = 0$ from a convergent of $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$, with $P_3 \in \mathcal{B}_n(x, y)_{|_{x \ge 4n}}$. We finally present some experimental examples. We believe these results constitute a new direction in RSA Cryptanalysis using continued fractions independently

of parameters e and d.

INTRODUCTION

In [1], Babindamana and al. presented arithmetical results on hyperbola and applied them to lattice points with the Fermat factorization Method. In particular, they used results from the hyperbola parametrization $\mathcal{B}_n(x,y)|_{\mathbb{Z}}$ to prove that for the Fermat factorization equation if $(N-6) \equiv 0 \mod 4$, the Fermat's method fails. Otherwise, in terms of cardinality, it has respectively 4, 8, $2(\alpha + 1)$, $(1 - \delta_{2p_i})2^{n+1}$ and $2\prod_{i=1}^{n} (\alpha_i + 1)$ lattice points respectively if N is an odd prime, $N = N_a \times N_b$ with N_a and N_b odd primes, $N = N_a^{\alpha}$ with N_a prime, $N = \prod_{i=1}^{n} p_i$ with p_i distinct primes and $N = \prod_{i=1}^{n} N_i^{\alpha_i}$ with N_i odd primes.

In this paper, we present new arithmetical results for the hyperbola parametrization considered in [1], in particular we prove that given $p, a, \alpha_i, \alpha_j \in \mathbb{Z}_{>0}$ with $gcd(p + \alpha_j)$ $n \pm 1$

$$1, p-1) \neq 1, \text{ if } \gcd(\alpha_i, \alpha_j) = 1 \text{ and } \frac{p+1}{p-1} = \frac{\alpha_i}{\alpha_j} \text{ then } \alpha_i + \alpha_j = p \text{ and}$$
$$\alpha_i^a - \alpha_j^a = \begin{cases} 1 & \text{if } a = 1\\ p & \text{if } a = 2\\ p\delta & \text{if } 2 \mid a \end{cases}$$

Additionally, among other results, given a RSA modulus n, we prove that for all

 $P_i \in \mathcal{B}_n(x,y)_{|x \ge 4n}, \text{ there exists } \alpha_i \in \mathbb{Z}_{\ge 0} \text{ such that}$ $P_i = \left(4n\frac{(\alpha_i+1)^2}{2\alpha_i+1}, 4n(\alpha_i+1) \cdot \frac{\alpha_i}{2\alpha_i+1}\right), \text{ with } \alpha_0 = 0, \alpha_4 = n-1, \alpha_i \text{ an integer}$ sequence and $\{P_2, P_3, P_4\}$ the algebraic subset of $\mathcal{B}_n(x, y)_{|x \ge 4n}$. Finally, we present an attack on those results using continued fractions whose goal is to find X_{α} such

²⁰²⁰ Mathematics Subject Classification. Primary 11Y05, 11Y40, 11T71.

Key words and phrases. Hyperbola, RSA attack, continued fraction, Cryptanalysis.

that $\langle X_{\alpha_3}, P_3 \rangle = 0$. When successful, this attack has a complexity of $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$ with $b = 10^y$, $\alpha_{i3} + \alpha_{j3}$ a non trivial factor of n and α_{j4} such that $(n+1)/(n-1) = \alpha_{i4}/\alpha_{j4}$.

1. New Algebraic results on $\mathcal{B}_N(x,y)|_{x \ge 4N}$

In [2], the authors presented some arithemtical results on hyperbola and in [1], Babindamana and al. presented other arithmetical and algebraic results on the hyperbola $\mathcal{B}_n(x,y)|_{\mathbb{Z}}$ and their application to the lattice points of the Fermat factorization Method.

Recall that $\mathcal{B}_n(x,y)|_{\mathbb{Z}} = \{(x,y) \in \mathbb{Z} \times \mathbb{Z} / y^2 = x^2 - 4nx\}$ and $\mathcal{B}_n(x,y)|_{x \ge 4n} = (x,y)|_{x \ge 4n}$ $\{(x,y) \in \mathbb{Z}_{\geq 4n} \times \mathbb{Z}_{\geq 0} / y^2 = x^2 - 4nx\}.$

In this section, we present new algebraic results on $\mathcal{B}_N(x,y)|_{x\geq 4N}$

Theorem 1.1.

Let $p, a, \alpha_i, \alpha_j \in \mathbb{Z}_{>0}$ with $gcd(p+1, p-1) \neq 1$. If $gcd(\alpha_i, \alpha_j) = 1$ and $\frac{p+1}{p-1} = \frac{\alpha_i}{\alpha_i}$ then $\alpha_i + \alpha_j = p$ and

$$\alpha_i^a - \alpha_j^a = \begin{cases} 1 & if \ a = 1 \\ p & if \ a = 2 \\ p\delta & if \ 2 \mid a \end{cases}$$

Proof 1.2.

- Consider $p \in \mathbb{Z} \setminus \{0, 1\}$ with $gcd(p+1, p-1) \neq 1$, then the rational fraction $\frac{p+1}{p-1}$ is reducible. Now Assume $\frac{p+1}{p-1} = \frac{\alpha_i}{\alpha_j}$ its irreducible expression, where $gcd(\alpha_i, \alpha_j) = 1$. Rearranging the terms in this expression, we have $p\alpha_j + \alpha_j = p\alpha_i - \alpha_i$, then $-p(\alpha_i - \alpha_j) = -(\alpha_i + \alpha_j)$ hence $p = \frac{\alpha_i + \alpha_j}{\alpha_i - \alpha_j}$. Now since by assumption, p is an integer, then $(\alpha_i - \alpha_j) \mid (\alpha_i + \alpha_j)$ which is true only if $\alpha_i - \alpha_j = 1$ since $gcd(p+1, p-1) \neq 1$. Assume $\alpha_i - \alpha_j \neq 1$, $p+1 = \frac{2\alpha_i}{\alpha_i - \alpha_j}$ and $p-1 = \frac{2\alpha_j}{\alpha_i - \alpha_j}$. In this case, $gcd(p+1, p-1) = gcd(\frac{2\alpha_i}{\alpha_i - \alpha_j}, \frac{2\alpha_j}{\alpha_i - \alpha_j})$. Since the greatest common divisor is defined only in an Euclidean domain, then this would imply that divisor is defined only in an Euclidean domain, then this would imply that $(\alpha_i - \alpha_i) \mid 2\alpha_i$ and $(\alpha_i - \alpha_i) \mid 2\alpha_i$, which is absurd since by hypothesis, $gcd(\alpha_i, \alpha_i) = 1$ and by applying the lemma of Gauss, if $(\alpha_i - \alpha_i) \mid 2\alpha_i$ then $(\alpha_i - \alpha_i) \nmid 2\alpha_i$ or conversely if $(\alpha_i - \alpha_i) \mid 2\alpha_i$ then $(\alpha_i - \alpha_i) \nmid 2\alpha_i$, hence in this case, the gcd is not defined. Therefore, this holds only if $\alpha_i - \alpha_j = 1$.
- If a = 1, since p = α_i + α_j/α_i then α_i α_j = 1
 If a = 2, α_i² α_j² = (α_i α_j)(α_i + α_j) = 1 × p = p.
- Considering the Newton's binomial formula $(\alpha_i + \alpha_j)^n = \sum_{k=0}^n \binom{n}{k} \alpha_i^k \alpha_j^{n-k}$, assume 2 | a, then there exists $\beta \in \mathbb{Z}$ such that $a = 2\beta$. In this case

$$\alpha_i^a - \alpha_j^a = \alpha_i^{2\beta} - \alpha_j^{2\beta} = (\alpha_i^2)^\beta - (\alpha_j^2)^\beta$$

$$\begin{split} &= (\alpha_i^2 - \alpha_j^2)(\alpha_i^{2(\beta-1)} + \alpha_i^{2(\beta-2)}\alpha_j^2 + \ \cdots \ + \alpha_i^2\alpha_j^{2(\beta-2)} + \alpha_j^{2(\beta-1)}) \\ &= p(\alpha_i - \alpha_j)\left(\sum_{k=0}^{\beta-1} \alpha_i^{2(\beta-1-k)}\alpha_j^{2k}\right) \\ &= p\left(\sum_{k=0}^{\beta-1} \alpha_i^{2(\beta-1-k)}\alpha_j^{2k}\right) \\ &\text{Setting } \delta = \left(\sum_{k=0}^{\beta-1} \alpha_i^{2(\beta-1-k)}\alpha_j^{2k}\right), \text{ we have } \alpha_i^a - \alpha_j^a = p\delta \end{split}$$

Theorem 1.3.

If δ is a perfect square of a prime, then $Card\left(\mathcal{B}_{\delta}(x,y)|_{x\geq 4\delta}\right) = 3.$

Proof 1.4.

 δ is a perfect square with prime t such that $\delta = t^2$. In this case, $Div(\delta) = \{1, t, \delta\}$, in other words, δ has 3 divisors.

From theorem 2 in [1], using injective homomorphisms, we have the trivial injective morphism $f_{1\longrightarrow\delta}$, the trivial automorphism $f_{\delta\longrightarrow\delta}$, and the morphism $f_{t\longrightarrow\delta}$. with $\mathcal{B}_1(x,y)|_{x\geq 4}\{(4,0)\}, \mathcal{B}_t(x,y)|_{x\geq 4t}\{(4,0), ((t+1)^2, t^2-1)\}. \mathcal{B}_1(x,y)|_{x\geq 4} = \{(4,0)\}$ and $\mathcal{B}_t(x,y)|_{x\geq 4t} = \{(4t,0), ((t+1)^2, t^2-1)\}$ given by, for all $i \in Div(\delta)$

$$\begin{array}{cccc} f_{i \to \delta} : & \mathcal{B}_i(x, y)_{|_{x \ge 4i}} & \longrightarrow & \mathcal{B}_\delta(x, y)_{|_{x \ge 4\delta}} \\ & (x, y) & \longmapsto & \left(x \frac{\delta}{i}, y \frac{\delta}{i}\right) \end{array}$$

with $f_{i\to\delta}((4,0)) = (4\delta, 0) \in \mathcal{B}_{\delta}(x,y)|_{x\geq 4\delta}, f_{t\to\delta}((4,0)) = (4\delta, 0) \in \mathcal{B}_{\delta}(x,y)|_{x\geq 4\delta}, f_{t\to\delta}(((t+1)^2, t^2-1)) = (t(t+1)^2, t(t^2-1)) \in \mathcal{B}_{\delta}(x,y)|_{x\geq 4\delta} \text{ and } ((\delta+1)^2, \delta^2-1) \in \mathcal{B}_{\delta}(x,y)|_{x\geq 4\delta}.$

Hence $\in \overline{\mathcal{B}}_{\delta}(x,y)|_{x \ge 4\delta}$ has exactly 3 points.

Theorem 1.5.

- (1) Let *n* be a RSA modulus, α_i an integer sequence and $\{P_0, P_2, P_3, P_4\}$ the algebraic subset of $\mathcal{B}_n(x,y)|_{x\geq 4n}$. There exists $X_{\alpha_i} = (-\alpha_i, \alpha_i + 1) \in \mathbb{Z}^2$, such that $\langle X_{\alpha_i}, P_i \rangle_{i=\overline{0,2\cdots 4}} = 0$ and $2\alpha_i + 1 \in \{1, p, q, n\}$.
- (2) For P_2 and P_3 with $\langle X_{\alpha_2}, P_2 \rangle = 0$ and $\langle X_{\alpha_3}, P_3 \rangle = 0$, then $\frac{x_{P_1}}{y_{P_1}} = \alpha_3 + \alpha_2 + 1$
- $\alpha_3 \alpha_2$ (3) p, q respectively n and $\varphi(n)$ are given by $2\alpha_2 + 1, 2\alpha_3 + 1$ respectively $2\alpha_4 + 1$ and $4\alpha_2\alpha_3$.

$$\forall P_i \in \{P_0, P_1, P_2, P_3, P_4\}, \quad P_i = \begin{cases} x_{P_i} \left(1, \frac{\alpha_3 - \alpha_2}{\alpha_3 + \alpha_2 + 1}\right) & if \ i = 1\\ x_{P_i} \left(1, \frac{\alpha_i}{\alpha_i + 1}\right) & else \end{cases}$$

$$(5)$$

 $P_0 = (16\alpha_2\alpha_3 + 8\alpha_2 + 8\alpha_3 + 4, 0)$

 $P_1 = (4\alpha_2^2 + 8\alpha_2\alpha_3 + 4\alpha_3^2 + 8\alpha_2 + 8\alpha_3 + 4, 4\alpha_2^2 - 4\alpha_3^2 + 4\alpha_2 - 4\alpha_3)$ $P_2 = \left(8\,\alpha_2\alpha_3^2 + 16\,\alpha_2\alpha_3 + 4\,\alpha_3^2 + 8\,\alpha_2 + 8\,\alpha_3 + 4, 8\,\alpha_2\alpha_3^2 + 8\,\alpha_2\alpha_3 + 4\,\alpha_3^2 + 4\,\alpha_3\right)$ $P_3 = \left(8\,\alpha_2^2\alpha_3 + 4\,\alpha_2^2 + 16\,\alpha_2\alpha_3 + 8\,\alpha_2 + 8\,\alpha_3 + 4, 8\,\alpha_2^2\alpha_3 + 4\,\alpha_2^2 + 8\,\alpha_2\alpha_3 + 4\,\alpha_2\right)$ $P_4 = (4 \alpha_4^2 + 8 \alpha_4 + 4, 4 \alpha_4^2 + 4 \alpha_4)$ (6) the $\alpha_{i|_{i=2...4}}$ s as defined in (1) satisfy the following system: $4\alpha_{2}\alpha_{3} + 2(\alpha_{2} + \alpha_{3}) + 1 = n \qquad (a)$ $4(16\alpha_{2}^{3}\alpha_{3}^{3} + 32\alpha_{2}^{3}\alpha_{3}^{2} + 32\alpha_{2}^{2}\alpha_{3}^{3} + 20\alpha_{2}^{3}\alpha_{3} + 68\alpha_{2}^{2}\alpha_{3}^{2} + 20\alpha_{2}\alpha_{3}^{3} - 2\alpha_{2}^{2}\alpha_{3}n - 2\alpha_{2}\alpha_{3}^{2}n + 46\alpha_{2}^{2}\alpha_{3} + 46\alpha_{2}\alpha_{3}^{2} + 4\alpha_{3}^{3} - \alpha_{2}^{2}n - 8\alpha_{2}\alpha_{3}n - \alpha_{3}^{2}n + 10\alpha_{2}^{2} + 34\alpha_{2}\alpha_{3} + 10\alpha_{3}^{2} - 4\alpha_{2}n - 4\alpha_{3}n + n^{2} + 8\alpha_{2} + 8\alpha_{3} - 2n + 2) = n(n+1)^{2} \qquad (b)$ $4(8\alpha_{2}^{2}\alpha_{3}^{2} + 12\alpha_{2}^{2}\alpha_{3} + 12\alpha_{2}\alpha_{3}^{2} + 4\alpha_{2}^{2} + 18\alpha_{2}\alpha_{3} + 4\alpha_{3}^{2} - \alpha_{3}n + 6\alpha_{2} + 6\alpha_{3} - n + 2)(2\alpha_{2}\alpha_{3} + \alpha_{2} + \alpha_{3}) = n(n^{2} - 1) \qquad (c)$ or equivalently, setting $P = \alpha_2 \alpha_3$ and $S = \alpha_2 + \alpha_3$, P and S satisfy the following system: $\begin{cases} 16P^3 + 4S^3 + (32S + 28)P^2 + (20P + 10 - n)S^2 + (34 - 2n)PS + (14 - 6n)P + (8 - 4n)S = \frac{n(n+1)^2}{4} - n^2 + 2n - 2 \qquad (b) \\ [8P^2 + 4S^2 + 12PS + 10P + (6 - n)S - n + 2](2P + S) = \frac{n(n^2 - 1)}{4} \end{cases}$ (7) For all $i \in \{0, 2, 3, 4\}$, $\frac{\alpha_{ai}}{\alpha_{bi}} < \frac{\alpha_{aj}}{\alpha_{bj}} \forall j > i$. (8) More generally, there exists $\alpha_i \in \mathbb{Z}_{\geq 0}$ such that $P_i = \left(4n\frac{(\alpha_i+1)^2}{2\alpha_i+1}, 4n(\alpha_i+1) \cdot \frac{\alpha_i}{2\alpha_i+1}\right) \in \mathcal{B}_n(x,y)|_{x \geq 4n}$ (9) Considering $\mathcal{B}_n(x,y)|_{x \geq 4n}$ for a RSA modulus n, the general term of the sequence α_n is $\alpha_i = \frac{\tau_i - 1}{2} \quad \forall \ \tau_i \in Div(n)$, with $\alpha_0 = 0$ and $\alpha_4 = \frac{n-1}{2}$.

Proof 1.6.

(1) To prove this, let's consider $X_{\alpha_i} = (-\alpha_i, \ \alpha_i + 1) \in \mathbb{Z}^2$ and $P_i = (x_{P_i}, \ y_{P_i})$ with $i \in \{0, 2, 3, 4\}$. The scalar product $\langle X_{\alpha_i}, \ P_i \rangle_{i=\overline{0,2\cdots 4}} = 0$ is equivalent to $-\alpha_i x_{P_i} + (\alpha_i + 1)y_{P_i} = 0$, finally $\alpha_i = \frac{y_{P_i}}{x_{P_i} - y_{P_i}}$. Here without loss of generality, for all $P_i \in \{P_0, \ P_2, \ P_3, \ P_4\}$, from $\mathcal{B}_n(x, y)_{|x \ge 4n}$ structure, $x_{P_i} \neq y_{P_i}$, hence α_i is well defined and

$$\begin{split} X_{\alpha_i} &= \left(-\frac{y_{P_i}}{x_{P_i} - y_{P_i}}, \ \frac{x_{P_i}}{x_{P_i} - y_{P_i}} \right) \text{. By the same,} \\ &2\alpha_i + 1 = \frac{2y_{P_i}}{x_{P_i} - y_{P_i}} + 1 \\ &= \frac{x_{P_i} + y_{P_i}}{x_{P_i} - y_{P_i}} = \frac{(x_{P_i} + y_{P_i})^2}{x_{P_i}^2 - y_{P_i}^2} = \frac{x_{P_i}^2 + y_{P_i}^2 + 2x_{P_i}y_{P_i}}{x_{P_i}^2 - y_{P_i}^2} \end{split}$$

$$= \frac{x_{P_i}^2 + x_{P_i}^2 + 2x_{P_i}y_{P_i} - 4nx_{P_i}}{x_{P_i}^2 - x_{P_i}^2 + 4nx_{P_i}} \text{ since } y_{P_i}^2 = x_{P_i}^2 - 4nx_{P_i}^2$$
$$= \frac{2x_{P_i}^2 + 2x_{P_i}y_{P_i} - 4nx_{P_i}}{4nx_{P_i}}$$
$$= \frac{x_{P_i}}{2n} + \frac{y_{P_i}}{2n} - 1, \text{ since for all } P_i \in \{P_0, P_2, P_3, P_4\},$$
$$P_i \in \{(4n, 0), (p(q+1)^2, p(q^2-1)), (q(p+1)^2, q(p^2-1))\}$$
$$\cup \{((n+1)^2, n^2-1)\}$$

for each P_i , we have:

$$\begin{aligned} \frac{x_{P_i}}{2n} + \frac{y_{P_i}}{2n} - 1 &= \frac{4n+0}{2n} - 1 = 2 - 1 = 1 \quad \in \{1, p, q, n\} \text{ or} \\ \frac{x_{P_i}}{2n} + \frac{y_{P_i}}{2n} - 1 &= \frac{p(q+1)^2 + p(q^2 - 1)}{2n} - 1 = \frac{nq+2n+p+qn-p-2n}{2n} = q \\ &\in \{1, p, q, n\} \text{ or} \\ &= \frac{q(p+1)^2 + q(p^2 - 1)}{2n} - 1 = \frac{np+2n+q+pn-q-2n}{2n} = p \\ &\in \{1, p, q, n\} \text{ or} \\ &= \frac{(n+1)^2 + n^2 - 1}{2n} - 1 = \frac{n^2 + 2n + 1 + n^2 - 1}{2n} = \frac{2n^2}{2n} = n \\ &\text{Hence } 2\alpha_i + 1 \quad \in \{1, p, q, n\} \text{ for all } i \in \{0, 2, 3, 4\}. \end{aligned}$$

(2) Here, under the same conditions as above, consider $2\alpha_2 + 1 = p$ and $2\alpha_3 + 1 = q$. From $\mathcal{B}_n(x, y)_{|x \ge 4n}$ structure,

$$\frac{x_{P_1}}{y_{P_1}} = \frac{(p+q)^2}{p^2 - q^2} = \frac{p+q}{p-q}$$
$$= \frac{2\alpha_2 + 1 + 2\alpha_3 + 1}{2\alpha_2 + 1 - 2\alpha_3 - 1}$$
$$= \frac{\alpha_2 + \alpha_3 + 1}{\alpha_2 - \alpha_3}.$$

(3) This results from (2) for p and q, and for the Euler totient function $\varphi(n) = (p-1)(q-1) = \alpha_2 \alpha_3$.

$$\begin{array}{ll} \text{(4) If } i = 1, \text{ from } (2), \ \frac{x_{P_1}}{y_{P_1}} = \frac{\alpha_2 + \alpha_3 + 1}{\alpha_2 - \alpha_3}, \text{ then } y_{P_1} = \frac{\alpha_3 - \alpha_2}{\alpha_3 + \alpha_2 + 1}, \text{ hence} \\ \left(x_{P_1}, \ \frac{\alpha_3 - \alpha_2}{\alpha_3 + \alpha_2 + 1} x_{P_1} \right). \\ \text{If not } i = 1, \text{ considering the scalar product relation from } (1), \ \langle X_{\alpha_i}, \ P_i \rangle_{i=\overline{2\cdots4}} = \\ 0, \text{ we have } \frac{x_{P_i}}{y_{P_i}} = \frac{\alpha_i + 1}{\alpha_i} \text{ then } y_{P_1} = \frac{\alpha_i}{\alpha_i + 1}. \text{ In this case, for all } i \in \\ \{2, 3, 4\}, \ P_i = (x_{P_i}, \ y_{P_i}) = \left(x_{P_i}, \ \frac{\alpha_i}{\alpha_i + 1} x_{P_i} \right). \text{ This holds for } i = 0. \text{ Hence} \\ \forall \ P_i \in \{P_0, P_1, P_2, P_3, P_4\}, \ P_i = \begin{cases} x_{P_i} \left(1, \ \frac{\alpha_3 - \alpha_2}{\alpha_3 + \alpha_2 + 1} \right) & \text{if } i = 1 \\ x_{P_i} \left(1, \ \frac{\alpha_i}{\alpha_i + 1} \right) & \text{else} \end{cases} \end{array}$$

- 6 GILDA RECH BANSIMBA, REGIS F. BABINDAMANA, AND BASILE GUY R. BOSSOTO
 - (5) From (1), $2\alpha_2 + 1 = p$, $2\alpha_3 + 1 = q$ and $2\alpha_4 + 1 = n$, then substituting these values in $\mathcal{B}_n(x,y)|_{x \ge 4n} =$ $\{(4n, 0), ((p+q)^2, p^2 - q^2), (p(q+1)^2, p(q^2 - 1)), (q(p+1)^2, q(p^2 - 1)), ((n+1)^2, n^2 - 1)\}$ For $P_0 = (4n, 0)$, we get $P_0 = (16\alpha_2\alpha_3 + 8\alpha_2 + 8\alpha_3 + 4, 0)$ For P_1, P_2, P_3 and P_4 , after expansion, we get $P_1 = (4\alpha_2^2 + 8\alpha_2\alpha_3 + 4\alpha_3^2 + 8\alpha_2 + 8\alpha_3 + 4, 4\alpha_2^2 - 4\alpha_3^2 + 4\alpha_2 - 4\alpha_3);$ $P_2 = (8\alpha_2\alpha_3^2 + 16\alpha_2\alpha_3 + 4\alpha_3^2 + 8\alpha_2 + 8\alpha_3 + 4, 8\alpha_2\alpha_3^2 + 8\alpha_2\alpha_3 + 4\alpha_3^2 + 4\alpha_3);$ $P_3 = (8\alpha_2^2\alpha_3 + 4\alpha_2^2 + 16\alpha_2\alpha_3 + 8\alpha_2 + 8\alpha_3 + 4, 8\alpha_2^2\alpha_3 + 4\alpha_2^2 + 8\alpha_2\alpha_3 + 4\alpha_2)$ and $P_4 = (4\alpha_4^2 + 8\alpha_4 + 4, 4\alpha_4^2 + 4\alpha_4)$
 - (6) For (a), since $n = (2\alpha_2 + 1)(2\alpha_3 + 1) = 4\alpha_2\alpha_3 + 2(\alpha_2 + \alpha_3) + 1$. Now for (b) and (c), we know from [1], that $P_2 + P_3 = P_4$. Since $(x_{P_2+P_3}, y_{P_2+P_3}) = (x_{P_4}, y_{P_4})$. We set ((b), (c)) = $(x_{P_2+P_3}, y_{P_2+P_3})$. Recall the additive group law on $\mathcal{B}_n(x, y)_{|_{\mathbb{Z}}}$ for points P_2 and P_3 is defined by

$$x_{P_2+P_3} = \frac{1}{2n} \left[(x_{P_2} - 2n)(x_{P_3} - 2n) + y_{P_2}y_{P_3} \right] + 2n$$

$$y_{P_2+P_3} = \frac{1}{2n} \left[y_{P_2}(x_{P_3} - 2n) + y_{P_3}(x_{P_2} - 2n) \right]$$

Then substituting the expressions of P_2 and P_3 from (3) in terms of α'_i s in the system

$$\begin{cases} \frac{1}{2n} \left[(x_{P_2} - 2n)(x_{P_3} - 2n) + y_{P_2}y_{P_3} \right] + 2n = x_{P_4} = (n+1)^2 \\ \frac{1}{2n} \left[y_{P_2}(x_{P_3} - 2n) + y_{P_3}(x_{P_2} - 2n) \right] = y_{P_4} = n^2 - 1 \end{cases}$$

we finally get

.

$$4(16\alpha_{2}^{3}\alpha_{3}^{3} + 32\alpha_{2}^{2}\alpha_{3}^{2} + 32\alpha_{2}^{2}\alpha_{3}^{3} + 20\alpha_{2}^{3}\alpha_{3} + 68\alpha_{2}^{2}\alpha_{3}^{2} + 20\alpha_{2}\alpha_{3}^{3} - 2\alpha_{2}^{2}\alpha_{3}n - 2\alpha_{2}\alpha_{3}^{2}n + 46\alpha_{2}^{2}\alpha_{3} + 46\alpha_{2}\alpha_{3}^{2} + 4\alpha_{3}^{3} - \alpha_{2}^{2}n - 8\alpha_{2}\alpha_{3}n - \alpha_{3}^{2}n + 10\alpha_{2}^{2} + 34\alpha_{2}\alpha_{3} + 10\alpha_{3}^{2} - 4\alpha_{2}n - 4\alpha_{3}n + n^{2} + 8\alpha_{2} + 8\alpha_{3} - 2n + 2) = n(n+1)^{2}$$
(b)
(6.2)

 $10\alpha_{3}^{2} - 4\alpha_{2}n - 4\alpha_{3}n + n^{2} + 8\alpha_{2} + 8\alpha_{3} - 2n + 2) = n(n+1)^{2}$ (b) $4(8\alpha_{2}^{2}\alpha_{3}^{2} + 12\alpha_{2}^{2}\alpha_{3} + 12\alpha_{2}\alpha_{3}^{2} + 4\alpha_{2}^{2} + 18\alpha_{2}\alpha_{3} + 4\alpha_{3}^{2} - \alpha_{2}n - \alpha_{3}n + 6\alpha_{2} + 6\alpha_{3} - n + 2)(2\alpha_{2}\alpha_{3} + \alpha_{2} + \alpha_{3}) = n(n^{2} - 1)$ (c)

setting $P = \alpha_2 \alpha_3$, $S = \alpha_2 + \alpha_3$, and substituting in both the relation (a) and the system (6.2) above, we get:

$$4P + 2S + 1 = n \qquad (a)$$

$$16P^{3} + 4S^{3} + (32S + 28)P^{2} + (20P + 10 - n)S^{2} + (34 - 2n)PS + (14 - 6n)P + (8 - 4n)S = \frac{n(n+1)^{2}}{4} - n^{2} + 2n - 2 \qquad (b)$$

$$[8P^{2} + 4S^{2} + 12PS + 10P + (6 - n)S - n + 2] (2P + S) = \frac{n(n^{2} - 1)}{4} \qquad (c)$$
(7) Here first recall that $(\alpha_{ai}, \alpha_{bi}) = (\alpha_{ai}, \alpha_{ai} + 1)$ and $(\alpha_{aj}, \alpha_{bj}) = (\alpha_{aj}, \alpha_{aj} + 1)$

(7) Here first recall that $(\alpha_{ai}, \alpha_{bi}) = (\alpha_{ai}, \alpha_{ai} + 1)$ and $(\alpha_{aj}, \alpha_{bj}) = (\alpha_{aj}, \alpha_{aj} + 1)$. If j > i, then $\alpha_j > \alpha_i$ considering (1). Now $\alpha_i < \alpha_j$ then $\alpha_i + \alpha_i \alpha_j < \alpha_j + \alpha_i \alpha_j$ since $\alpha_i > 0, \alpha_j > 0$. We have $\alpha_i (\alpha_j + 1) < \alpha_j (\alpha_i + 1)$. Finally $\frac{\alpha_i}{\alpha_i + 1} < \frac{\alpha_j}{\alpha_j + 1}$. Hence $\frac{\alpha_{ai}}{\alpha_{bi}} < \frac{\alpha_{aj}}{\alpha_{bj}}$. A CONTINUED FRACTION-HYPERBOLA BASED ATTACK ON RSA CRYPTOSYSTEM 7

$$\begin{array}{ll} \text{(8) Consider } P_i = \left(4n\frac{(\alpha_i+1)^2}{2\alpha_i+1}, \ 4n(\alpha_i+1)\cdot\frac{\alpha_i}{2\alpha_i+1}\right) \in \mathcal{B}_n(x,y)_{|_{x\geq 4n}}, \text{ since from } (1), 2\alpha_i+1 \in \{1,p,q,n\} \text{ then } \alpha_i \in \left\{0,\frac{p-1}{2},\frac{q-1}{2},\frac{n-1}{2}\right\} \text{ then:} \\ \text{ If } \alpha_i = 0, P_i = \left(4n\frac{1}{1}, \ 4n\cdot(1)\cdot 0\right) = (4n, \ 0) \in \mathcal{B}_n(x,y)_{|_{x\geq 4n}}. \\ \text{ If } \alpha_i = \frac{p-1}{2}, P_i = \left(4n\cdot\frac{\left(\frac{p+1}{2}\right)^2}{p}, \ 4n\cdot\frac{p+1}{2}\cdot\frac{p-1}{2}\right) \\ = \left(4n\cdot\frac{\left(p+1\right)^2}{4p}, \ 4n\cdot\frac{p+1}{2}\cdot\frac{p-1}{2p}\right) \\ = \left(q(p+1)^2, \ q(p^2-1)\right) \in \mathcal{B}_n(x,y)_{|_{x\geq 4n}}. \\ \text{ If } \alpha_i = \frac{q-1}{2}, P_i = \left(4n\cdot\frac{\left(\frac{q+1}{2}\right)^2}{q}, \ 4n\cdot\frac{q+1}{2}\cdot\frac{q-1}{2q}\right) \\ = \left(q(p+1)^2, \ p(q^2-1)\right) \in \mathcal{B}_n(x,y)_{|_{x\geq 4n}}. \\ \text{ If } \alpha_i = \frac{n-1}{2}, P_i = \left(4n\cdot\frac{\left(\frac{n+1}{2}\right)^2}{n}, \ 4n\cdot\frac{n+1}{2}\cdot\frac{n-1}{2n}\right) \\ = \left(qn\cdot\frac{\left(n+1\right)^2}{4n}, \ 4n\cdot\frac{n+1}{2}\cdot\frac{n-1}{2n}\right) \\ = \left((n+1)^2, \ n^2-1\right) \in \mathcal{B}_n(x,y)_{|_{x\geq 4n}}. \end{array}$$

(9) This is a straightforward observation from (1). Since $2\alpha_i + 1 \in \{1, p, q, n\}$, then either $\alpha_i = 0$, $\alpha_i = \frac{p-1}{2}$ or $\alpha_i = \frac{q-1}{2}$ or $\alpha_i = \frac{n-1}{2}$.

Conjecture 1.7.

For all $i \in \{1, 2, 3, 4\}$, $P_i \in \{P_1, P_2, P_3, P_4\}$ the algebraic subset of $\mathcal{B}_n(x, y)|_{x \ge 4n}$. If there exists $(\alpha_{ii}, \alpha_{ji}) \in \mathbb{Z}_{\ge 0}^2$ such that $\frac{x_{P_i}}{y_{P_i}} = \frac{\alpha_{ii}}{\alpha_{ji}}$ with $gcd(\alpha_{ii}, \alpha_{ji}) = 1$ then $\alpha_{i1} + \alpha_{j1} = \alpha_{i3} + \alpha_{j3}$ and $\alpha_{ii} + \alpha_{ji} \in \{p, q, n\}$.

Example 1.8.

Here we consider a sample of 11 different n in the following table, their $\mathcal{B}_n(x,y)|_{x \ge 4n}$ structures with the corresponding $\frac{\alpha_{ii}}{\alpha_{ji}}$:

n	${\mathcal B}_{\boldsymbol{n}}(\boldsymbol{x},\boldsymbol{y}))_{\boldsymbol{x}\geq 4\boldsymbol{n}}$	$\frac{\alpha_{i1}}{\alpha_{j1}}$	$\frac{\alpha_{i2}}{\alpha_{j2}}$	$\frac{\alpha_{i3}}{\alpha_{j3}}$	$rac{lpha_{i4}}{lpha_{j4}}$
15	$\{(60, 0), (64, 16), (80, 40), (108, 72), (256, 224)\}$	6 4	, CI) (C)	a solt
35	$\{(140,0), (144,24), (252,168), (320,240), (1296,1224)\}$	9	σic	2144 0	$\frac{16}{17}$
77	$\{(308,0),(324,72),(704,528),(1008,840),(6084,5928)\}$	6)¢	9 13 [0	<u>س</u> احد	<u>39</u>
143	$\{(572,0),(576,48),(1872,1560),(2156,1848),(20736,20448)\}$	12^{-1}	xo u	21-1C	712
391	$\{(1564,0),(1600,240),(7452,6624),(9792,8976),(153664,152880)\}$	<u>-30</u>	co los	<u>17</u>	<u>196</u> 195
713	$\{(2852,0),(2916,432),(17856,16368),(23552,22080),(509796,508368)\}$	<u>27</u>	12	16-	<u>357</u> 356
14893	(59572, 0), (111556, 76152), (819396, 789048), (4214772, 4184880), (221831236, 221801448)	$\frac{167}{114}$	$\frac{27}{26}$	$\frac{141}{140}$	$\frac{7447}{7446}$
44923	(179692,0),(190096,44472),(759256,7501872),(12174300,12084120),(2018165776,2018075928)	$\frac{218}{51}$	<u>84</u> 83	$\frac{135}{134}$	$\frac{22462}{22461}$
58307	(233228, 0), (242064, 46248), (11720000, 11602800), (17200764, 17083752), (3399822864, 3399706248)	$\frac{246}{47}$	<u>100</u> 99	$\frac{147}{146}$	$\frac{29154}{29153}$
439007603	(1756030412,0), (2065157136, 798996408), (6116692964112, 6115814885880), (13835324622476, 13834446579408), (192727676369820816, 192727675491805608)	$\frac{22722}{8791}$	<u>6966</u> 6965	$\frac{15757}{15756}$	$\frac{219503802}{219503801}$
12100028890	(48400115563041438556, 0), (49139988961662726400, 6029707342921211040), (3720657936997783642938073867293697586729360), (372057939437676), 372065793497757637858076677100, 4751457734679586777100, 47514577346795876766777100, 47514577346795876766777607886074680), (37514577346795876766777100, 4751757746795877766788607468777667878767667876078677776787877676787876767878767678776778787778787787				
760359639	(14641069915723537932	430079469	$\frac{1037458285}{1537458285}$	1967537754	6050014445380179819
	2158145982140929600, 1464106991572353792				

TABLE 1. Table of $\mathcal{B}_n(x,y)|_{x\geq 4n}$ and $\frac{\alpha_{ii}}{\alpha_{ji}}$ for 11 different n values




FIGURE 1. $\frac{\alpha_{ii}}{\alpha_{ji}}$ from a sample of 10 $\mathcal{B}_n(x,y)_{|_{x\geq 4n}}$

From these plots, an observation is that the maximum value of $\frac{\alpha_{ii}}{\alpha_{ji}}$ on $\mathcal{B}_n(x,y)|_{x\geq 4n}$ is obtained with i = 1: the point P_1 , followed by the point with i = 2: the point P_2 then followed by the P_3 and finally by the point P_4 . The sequence $\frac{\alpha_{ii}}{\alpha_{ji}}$ is a decreasing sequence.

Another notice for this sample is that for $i \in \{2, 3, 4\}$, the sequence $\frac{\alpha_i}{\alpha_j}$ seems to converge as n goes to infinity, and $\frac{\alpha_{i2}}{\alpha_{j2}} \simeq \frac{\alpha_{i3}}{\alpha_{j3}} \simeq \frac{\alpha_{i4}}{\alpha_{j4}}$ as n grows.

2. The Attack

In this section, we present the attack. This is a generic attack and only depends on the modulus n and does not depend on the size of the public parameter e or the private exponent d in the cryptosystem setting. This is a new approach compared to Wiener's attack.

Prior to the presentation, let's recall some preliminaries on the RSA cryptosystem and on some non-generic attack related work done so far.

2.1. Recall on RSA Cryptosystem. We denote $\mathcal{P}, \mathcal{C}, \mathcal{K}, \mathcal{E}_k, \mathcal{D}_k$ respectively the Plaintext space, the Ciphertext space, the Key space, the Encryption algorithm and the Decryption algorithm.

Here $\mathcal{P} = \mathcal{C} = \mathbb{Z}/n\mathbb{Z}$, and

 $\begin{aligned} \mathcal{K} &= \left\{ (p,q,n,e,d) \in \mathbb{Z}_+^5, \ p,q \text{ primes } / \ n = pq, \ ed \equiv 1 \mod \varphi(n) \right\}, p,q \text{ are chosen} \\ \text{randomly, } 1 < e < \varphi(n) \ / \ \gcd(e,\varphi(n)) = 1 \text{ and } d \text{ such that } ed \equiv 1 \mod \varphi(n). \\ \text{Public parameters are } (n,e), \text{ secret parameters are } (p,q,d). \text{ The following are the} \\ \text{encryption and decryption algorithms.} \end{aligned}$

$$\mathcal{E}_k : \mathcal{P} \longrightarrow \mathcal{C} ; \qquad \mathcal{D}_k : \mathcal{C} \longrightarrow \mathcal{P} \\ m \longmapsto \mathcal{E}_k(m) = m^e \mod n \qquad \qquad c \longmapsto \mathcal{D}_k(c) = c^d \mod n$$

 $\forall m \in \mathcal{P}, \ \mathcal{E}_k(m) = m^e \mod n = c \text{ and } \mathcal{D}_k(c) = c^d \mod n = m.$ With $\mathcal{D}_k(\mathcal{E}_k) = \mathcal{E}_k(\mathcal{D}_k) = id.$

10 GILDA RECH BANSIMBA, REGIS F. BABINDAMANA, AND BASILE GUY R. BOSSOTO

2.2. Related work on non-generic attack. Most of the non-generic attacks are based on the development of continued fractions [5] and the application of Coppersmith's [8] theorem. From the work of Wiener in [5], it is shown that the secret key d can be recovered in polynomial time for a bound of $d < \frac{1}{3}N^{0.25}$. Following the same perspective, in [6] De Weger proposed another bound $d < \frac{N^{\frac{3}{4}}}{|p-q|}$. Later in [7], Nitaj and al. proposed the improved bound $d < \frac{\sqrt{6\sqrt{2}}}{6}N^{0.25}$. In [4], Bunder and al. proposed a new improved bound and proved that d can be recovered if $d < 2\sqrt{2N^{\frac{1}{4}}}$.

More recently, in [3] Ariffin and al. proposed a new improved short decryption exponent attack for

$$d < \sqrt{\frac{a^{j} + b^{i}}{2}} \left(\frac{N}{e}\right)^{\frac{1}{2}} N^{0.375}.$$

Setting $N_{1} = N - \left\lceil \left(\frac{a^{\frac{j}{i}} + b^{\frac{j}{i}}}{(2ab)^{\frac{j}{2i}}} + \frac{a^{\frac{1}{j}} + b^{\frac{1}{j}}}{(2ab)^{\frac{1}{2j}}}\right) \sqrt{N} \right\rceil + 1$, they showed that d can be

recovered in polynomial time from the convergents of the continued fraction of $\frac{\epsilon}{N_1}$. All of these attacks are bounded by the Euler totient function approximation and the size of the secret key d or the size of the public exponent e and are therefore limited when the public parameter e is properly chosen and consequently d is properly chosen.

Consequently, we present a generic attack based on continued fractions. It is independent of public and private keys e and d respectively.

The attack consists on finding $X_{\alpha} = (-\alpha_3, \alpha_3 + 1)$ that satisfies $\langle X_{\alpha_3}, P_3 \rangle = 0$ from a convergent of $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$.

When successful, it retrieves an RSA private key d in $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$. For that, let's consider the following results.

Lemma 2.1.

 $\forall n \text{ RSA type modulus and consider the structure } \mathcal{B}_n(x,y)_{|_{x>4n}}$, then $3\alpha_{j3} < \alpha_{j4}$.

Proof 2.2.

 $\forall n = pq, pq < n+2$, then $q < \frac{pq}{3} < \frac{n+2}{3}$ finally $q < \frac{n+2}{3}$ then 3q < n+2, and 3(q-1) < n-1 hence $3\alpha_{j3} < \alpha_4$.

Theorem 2.3.

Let *n* be a RSA modulus and $\mathcal{B}_n(x,y)_{|x\geq 4n}$ the associated hyperbola structure. There exists $\delta \leq \frac{1+\alpha_{j4}-\alpha_{j3}}{\alpha_{j4}\alpha_{j3}}$ and more precisely $0 < \frac{\alpha_{j4}-\alpha_{j3}}{\alpha_{j4}\alpha_{j3}} < \delta \leq \frac{1+\alpha_{j4}-\alpha_{j3}}{\alpha_{j4}\alpha_{j3}}$ such that $\frac{\alpha_{i3}}{\alpha_{j3}}$ is a convergent of $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$. **Proof 2.4.**

Consider

$$\begin{aligned} 0 < \frac{\alpha_{j4} - \alpha_{j3}}{\alpha_{j4}\alpha_{j3}} < \delta \leq \frac{1 + \alpha_{j4} - \alpha_{j3}}{\alpha_{j4}\alpha_{j3}} \\ \alpha_{j4} - \alpha_{j3} < \delta\alpha_{j4}\alpha_{j3} \leq 1 + \alpha_{j4} - \alpha_{j3} \\ 0 < \delta\alpha_{j4}\alpha_{j3} - (\alpha_{j4} - \alpha_{j3}) \leq 1 \end{aligned}$$

$$\begin{array}{l} \mbox{From Lemma 2.1, } 3\alpha_{j3} < \alpha_{j4} \\ 0 < 3\alpha_{j3} \left[\delta\alpha_{j4}\alpha_{j3} - (\alpha_{j4} - \alpha_{j3}) \right] \leq 3\alpha_{j3} < \alpha_{j4} \\ 0 < 2\alpha_{j3} \left[\delta\alpha_{j4}\alpha_{j3} - (\alpha_{j4} - \alpha_{j3}) \right] < \alpha_{j4} \\ 0 < 2\alpha_{j3} \left[\alpha_{j4}\alpha_{j3} - \alpha_{j4}\alpha_{j3} + \delta\alpha_{j4}\alpha_{j3} - (\alpha_{j4} - \alpha_{j3}) \right] < \alpha_{j4} \\ 0 < 2\alpha_{j3} \left[\alpha_{j3}(\alpha_{j4} + 1) + \delta\alpha_{j4}\alpha_{j3} - \alpha_{j4}(\alpha_{j3} + 1) \right] < \alpha_{j4} \\ 0 < 2\alpha_{j3} \left[\alpha_{j3}\alpha_{i4} + \delta\alpha_{j4}\alpha_{j3} - \alpha_{j4}\alpha_{i3} \right] < \alpha_{j4} \\ 0 < (\alpha_{j3}\alpha_{i4} + \delta\alpha_{j4}\alpha_{j3} - \alpha_{j4}\alpha_{i3}) < \frac{\alpha_{j4}}{2\alpha_{j3}} = \frac{\alpha_{j4}\alpha_{j3}}{2\alpha_{j3}^2} \\ \mbox{Dividing the whole expression by } \alpha_{j4}\alpha_{j3}, \mbox{ we have:} \end{array}$$

$$0 < \left|\frac{\alpha_{i4}}{\alpha_{j4}} + \delta - \frac{\alpha_{i3}}{\alpha_{j3}}\right| < \frac{1}{2\alpha_j^2}$$

Now applying the Legendre's theorem [9], $\frac{\alpha_{i3}}{\alpha_{j3}}$ is a convergent of $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$. ie there exists $k \leq n$ such that $[a_0, a_1, \ldots, a_k] = \frac{\alpha_{i3}}{\alpha_{j3}}$ with $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta = [a_0, a_1, \ldots, a_n]$.

Remark 2.5.

We consider the following notation for the continued fraction of $x, x = [a_0, a_1, \dots, a_n] =$ $a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{\cdot}}}.$

The k^{th} convergent of x is given by $[a_0, a_1, \ldots, a_k]$ with $k \leq n$, we denote it $\frac{p_k}{q_k} = [a_0, a_1, \dots, a_k]$ and the sequences (p_k) and (q_k) are computed using the recursive formulas:

$$p_{-2} = q_{-1} = 0, \quad p_k = a_k p_{k-1} + p_{k-2} \text{ for all } k \ge 0,$$

 $p_{-1} = q_{-2} = 1, \quad q_k = a_k q_{k-1} + q_{k-2} \text{ for all } k \ge 0.$

Discussion:

Theorem 2.3 gives an important result allowing us to attack any RSA modulus nusing the continued fraction method in a "parameters e and d"-independent way in contrast to existing related attacks both being constrained and depending on parameters e and d.

parameters e and a. From our results, finding δ such that $\delta \leq \frac{1 + \alpha_{j4} - \alpha_{j3}}{\alpha_{j4}\alpha_{j3}}$ is the crutial point. We approach this by approximating the value of $\frac{1}{\alpha_{j3}}$ since by rearranging the terms, we have $\delta \leq \frac{1 + \alpha_{j4} - \alpha_{j3}}{\alpha_{j4}\alpha_{j3}} = \frac{1 + \alpha_{j4}}{\alpha_{j4}\alpha_{j3}} - \frac{1}{\alpha_{j4}} = \frac{1 + \alpha_{j4}}{\alpha_{j4}} \cdot \frac{1}{\alpha_{j3}} - \frac{1}{\alpha_{j4}} > 0.$

Knowing that $0 < \frac{1}{\alpha_{i3}} < 0.5$, it is not difficult to see by induction and from a simple empirical observation (see table 2) that the decimal part of $\frac{1}{\alpha_{i3}}$ has exactly \sharp (number of digits of α_{j3}) - 1) zeros.

TABLE 2 .	Some	values	of α_{j3}	and	$\frac{1}{\alpha_{j3}}$
-------------	------	--------	------------------	-----	-------------------------

α_{j3}	2	3	5	6	11	15	140	146	15756	1967537754	
$\frac{1}{\alpha_{i3}}$	0.5	0.3	0.2	0.16	0.09	00_6	0. 00 714	0. 00 68	0. 0000 635	0. 000000000 508	
<i>a</i> _{j3}						1	2	2	4	9	

Another observation is that if α_{i4} or α_{j4} has x bits then α_{i3} has $\frac{x}{2}$ bits. Equivalently, if α_{j4} has k decimal digits in base 10 then α_{j3} has $\lfloor \frac{k}{2} \rfloor + 1$ decimal digits.

Finally $\frac{1}{\alpha_{j3}}$ has in its decimal part $\lfloor \frac{k}{2} \rfloor$ number of zeros, where k is number of decimal digits of α_{j4} . This information helps us to approximate $\frac{1}{\alpha_{i3}}$.

Therefore the following algorithm.

2.3. Attack algorithm. Here we present the algorithm. Its complexity analysis is given by proposition 2.8

Algorithm 1: ATTACK ALGORITHM

Input: $n, P_4 \in \mathcal{B}_n(x,y)|_{x \ge 4n}, e$: RSA public exponent **Output:** p and q non trivial factors of n**1 Step 1**: Retrieve factors of nInitialize bound $b = 10^y$, $y \in \{2, 3, 4, 5, 6, 7, 8, 9, ...\}$. Initialize $k = \sharp$ decimal digits of nInitialize $k = \sharp$ decimal argues of nInitialize $\frac{\alpha_{i4}}{\alpha_{j4}} = \frac{x_{P_4}}{y_{P_4}}$ while $i \leq b$ do $\delta \leftarrow \frac{1 + \alpha_{j4}}{\alpha_{j4}} \cdot \frac{i}{10^{\lfloor \frac{k}{2} \rfloor + 1 + \sharp \text{decimal of i}}} - \frac{1}{\alpha_{j4}}$ if $\delta > 0$ then Compute convergents $\frac{p_k}{q_k} = \frac{\alpha_{i3}}{\alpha_{j3}}$ of $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$ for $\frac{\alpha_{i3}}{\alpha_{j3}} \in \text{convergents of } \frac{\alpha_{i4}}{\alpha_{j4}} + \delta$ do if $gcd(n, \alpha_{i3} + \alpha_{j3}) \neq 1$, *n* then return $\alpha_{i3} + \alpha_{j3}$, gcd $(n, \alpha_{i3} + \alpha_{j3})$ end if end for end if end while 2 Step 2: Retrieve d : RSA private key from e $\psi \leftarrow (\alpha_{i3} + \alpha_{j3} - 1) \left(\gcd(\alpha_{i3} + \alpha_{j3}) - 1 \right)$ solve using extended Euclidean algorithm $ed + k\psi = 1$ return d

Example 2.6.

RSA modulus	Size (bits)	δ	$\frac{\alpha_{i3}}{\alpha_{j3}'}$	$\gcd(\alpha_{i3}^{\prime}+\alpha_{j3}^{\prime},\ n)$
14893	17	$\frac{7}{103}$	$\frac{141}{140}$	281
439007603	30	$\frac{\frac{10^3}{969}}{5 \cdot 10^7}$	$\frac{122935301}{122932918}$	13931

A CONTINUED FRACTION-HYPERBOLA BASED ATTACK ON RSA CRYPTOSYSTEM 13

Remark 2.7.

This attack can be parallelized in the following way: In the while loop, while iterating on different values of i up to b, discretize the interval in different steps. That said, assume for example we want to distribute the work on 2 CPUs, then: in CPU₁ go from 0 to b with step 2: $0, 2, 4, \cdots$, up to b. in CPU₂ go from 1 to b with step 2: $1, 3, 5, \cdots$, up to b - 1.

Proposition 2.8.

This algorithm 1 has a complexity in $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$.

Proof 2.9.

step 1, initializations are done in constant time $\mathcal{O}(1)$, $\frac{\alpha_{i4}}{\alpha_{j4}} + \delta$ has $\mathcal{O}(\log \alpha_{j4})$ convergents and the computation of the greatest common divisor of n and $\alpha_{i3} + \alpha_{j3}$ has complexity in $\mathcal{O}(\alpha_{i3} + \alpha_{j3})$ since min $(n, \alpha_{i3} + \alpha_{j3}) = \alpha_{i3} + \alpha_{j3}$, thus a total of $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$.

Step 2 has complexity in $\mathcal{O}(\log e)$ since $1 < e < \psi$ implies $\min(e, \psi) = e$. Now $\max(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}), \log e) = b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3})$, hence $\mathcal{O}(\text{step } 1 + \text{step } 2) = \mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$.

CONCLUSION

We have presented new arithmetical and algebraic results following the work of Babindamana and al. in [1] on hyperbolas and describe an approach to attacking a RSA-type modulus based on continued fractions in the new results, different from Wiener's attack approach which depends on the parameters e and d. When successful, this attack has a complexity in $\mathcal{O}(b \log \alpha_{j4} \log (\alpha_{i3} + \alpha_{j3}))$.

The particularity of this attack is that it is independent and not bounded by the size of the private key d nor the public exponent e compared to Weiner's attack and its improvements. As future work, this method should be thoroughly investigated to find the best bounds for δ , cases of modulus with faster $\frac{\alpha'_{i3}}{\alpha'_{j3}}$ retrivial and reduce b for a polynomial time overall complexity.

References

- Regis F. Babindamana, Gilda R. Bansimba, Basile G. R. Bossoto (2022). "Lattice Points on the Fermat Factorization Method", Journal of Mathematics, vol. 2022, Article ID 6360264.
- [2] Gilda R. Bansimba, Regis F. Babindamana, Basile G. R. Bossoto (2021). Some Arithmetical properties on Hyperbola, JP Journal of Algebra, Number Theory and Applications, Vol. 50, no. 1, P. 45-100.
- [3] Abubakar, S. I., Ariffin, M. R. K. and Asbullah, M. A. (2019). A New Improved Bound for Short Decryption Exponent on RSA Modulus N = pq using Wiener's Method, Malaysian Journal of Mathematical Sciences 13(S) April: 89–99
- [4] Bunder, M. and Tonien, J. (2017). A New Attack on the RSA Cryptosystem Based on Continued Fractions, Malaysian Journal of Mathematical Sciences 11(S) August: 45 - 57 (2017).
- [5] Wiener, M. J. (1990). Cryptanalysis of short rsa secret exponents, IEEE Transactions on Information theory, 36(3):553-558.
- [6] De Weger, B. (2002). Cryptanalysis of rsa with small prime difference. Applicable Algebra in Engineering, Communication and Computing, 13(1):17–28.
- [7] Nitaj, A. (2013). Diophantine and lattice cryptanalysis of the rsa cryptosystem. In Artificial Intelligence, Evolutionary Computing and Metaheuristics, pages 139–168. Springer.

- 14 $\,$ GILDA RECH BANSIMBA, REGIS F. BABINDAMANA, AND BASILE GUY R. BOSSOTO
- [8] Don Coppersmith (1997), Small Solutions to Polynomial Equations, and Low Exponent RSA Vulnerabilities. Journal of Cryptology, 10(4), 233-260.
- [9] S. Lang (1966), Introduction to Diophantine Approximations, Addison-Wesley Pub. Co.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITÉ MARIEN NGOUABI, FACULTÉ DES SCIENCES ET TECHNIQUES, BP: 69, BRAZZAVILLE, CONGO

Current address: Department of Mathematics and Computer Science, Université Marien Ngouabi, Faculté des Sciences et Techniques, BP: 69, Brazzaville, Congo

Email address: bansimbagilda@gmail.com

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITÉ MARIEN NGOUABI, FACULTÉ DES SCIENCES ET TECHNIQUES, BP: 69, BRAZZAVILLE, CONGO Email address: regis.babindamana@umng.cg

Department of Mathematics Université Marien Ngouabi, Faculté des Sciences et Techniques, BP: 69, Brazzaville, Congo

Email address: basile.bossoto@umng.cg