

Prophet Inequalities: Separating Random Order from Order Selection

Giordano Giambartolomei[†], Frederik Mallmann-Trenn[†] and Raimundo Saona^{††}

Abstract

Prophet inequalities are a central object of study in optimal stopping theory. A gambler is sent values online, sampled from an instance of independent distributions, in an adversarial, random or selected order, depending on the model. When observing each value, the gambler either accepts it as a reward or irrevocably rejects it and proceeds to observe the next value. The goal of the gambler, who cannot see the future, is maximising the expected value of the reward while competing against the expectation of a prophet (the offline maximum). In other words, one seeks to maximise the gambler-to-prophet ratio of the expectations.

The model, in which the gambler selects the arrival order first, and then observes the values, is known as Order Selection. In this model a ratio of 0.7251 has been proved to be attainable for any instance. In very recent work, this has been improved up to 0.7258. If the gambler chooses the arrival order (uniformly) at random, we obtain the Random Order model. The worst case ratio over all possible instances has been extensively studied for at least 40 years. In the recent work aforementioned, through simulations, this ratio has been shown to be at most 0.7254 for the Random Order model, thus establishing for the first time that carefully choosing the order, instead of simply taking it at random, benefits the gambler. We give an alternative, more rigorous proof of this fact, by showing mathematically that in the Random Order model, no algorithm can achieve a ratio larger than 0.7235. This sets a new state-of-the-art hardness for this model, and establishes more formally that there is a real benefit in choosing the order.

1 Introduction

Prophet inequalities are a central object of study in optimal stopping theory. A gambler is sent nonnegative values online, sampled from an instance of uniformly bounded independent random variables $\{V_i\}$ with known distributions $\{\mathcal{D}_i\}$, in adversarial, random or selected order, depending on the particular model. When observing each value, the gambler either accepts it as a reward, or irrevocably rejects it and proceeds with observing the next value. The goal of the gambler, who cannot see the future, is to maximise the expected value of the reward while competing against the expectation of a prophet (out of metaphor, the offline maximum or supremum, depending on whether the instance is finite or not). In other words, one seeks to maximise the gambler-to-prophet ratio of the expectations.

The gambler represents any online decision maker, such as an algorithm or stopping rule. Probabilistically, we will refer to it as a *stopping time* τ : being online implies not being able to see the future, thus the gambler will always stop at a time τ such that the event $\{\tau = i\}$ depends, informally speaking, only on the first i values observed.

[†]King's College London.

^{††}Institute of Science and Technology Austria.

Due to the online nature of prophet inequalities, some terminology from *competitive analysis* is usually borrowed. The decision maker, determining the stopping time τ , will be often referred to as an (online) algorithm. Somewhat informally, it is common to refer to a worst-case gambler-to-prophet ratio of the algorithm (that is a ratio known to be achievable for any given instance by the algorithm) as *competitive ratio*.

Let $c \in [0, 1]$ be a real constant. An algorithm is said to be c -competitive if it has a competitive ratio of c , meaning that it can attain at least a gambler-to-prophet ratio of c for any instance, but it does not necessarily mean that c is the highest possible ratio. An upper bound on any algorithm's highest possible competitive ratio for a given instance will be called *hardness* of the instance. Saying that a prophet inequality model is c -hard or has a hardness of c , means that there is a c -hard instance for that problem, but it does not necessarily mean that c is the lowest hardness possible amongst all instances for that model.

A hardness for a model is said to be *tight* (or *optimal*) when it is matched by the competitive ratio of an algorithm solving it. Similarly, the competitive ratio of an algorithm solving a model is tight (or optimal) when it is matched by a hardness known for that model. Often, when determining a hardness or a competitive ratio, numerical computations are involved. Thus tightness can be used in a broad sense. For two models A and B , we say that A *beats* B if the hardness of B is strictly less than the competitive ratio of an algorithm solving A . When A beats B or B beats A , we say that A and B are *separated*.

1.1 Models of prophet inequalities

To date, several models and corresponding extensions of prophet inequalities have appeared in the literature. In this section we first introduce the classical model, and then describe three relaxations of it, briefly reviewing the state-of-the-art concerning their hardness and the competitiveness of known algorithms. More details are provided in Section 1.5.

1.1.1 The classical Prophet Inequality

The very first model of prophet inequality, typically referred to as (adversarial) Prophet Inequality (PI), is due to [23, 24]. The given instance is composed of countably many independent random variables $\{V_i\}$ with known distributions $\{\mathcal{D}_i\}$ in a fixed given order, usually referred to as *adversarial* order. The general dynamics previously described is followed: the gambler observes in an online fashion the sequence of sampled values, and makes irrevocable decisions to accept a value and stop, or continue observing, with the goal of maximising the expected value of the reward, while competing against the expectation of a prophet, which represents the offline supremum. More precisely, the goal is to maximise the gambler-to-prophet ratio

$$\frac{\mathbb{E} V_\tau}{\mathbb{E} \sup_i V_i},$$

where τ is the stopping time associated to the gambler's stopping rule.

In [24], it was shown that the $1/2$ -hardness of PI (shown by Garlin [23]) is tight. Later it was shown that a competitive ratio of $1/2$ can be attained even by a *single threshold* algorithm [30]. The classical PI has subsequently been relaxed giving more power to the gambler, which leads to larger competitive ratios.

1.1.2 IID Prophet Inequality, Order Selection and Random Order models

In this section we first explain our working assumptions and some basic notation, and then review the three main models of prophet inequalities related to our work.

We will restrict ourselves to *finite* instances, that is, we will always work with a size $n \in \mathbb{N}$ fixed and with non-negative random variables V_1, \dots, V_n having distributions $\mathcal{D}_1, \dots, \mathcal{D}_n$. Therefore, it is natural that, if the gambler gets to the last stage, any value observed will have to be accepted. Thus the stopping time τ of the gambler will belong to the class of stopping times valued in $[n] := \{k \in \mathbb{N} : k \leq n\} = \{1, 2, \dots, n\}$. We denote this class as C^n . Also, we denote the set of permutations of $[n]$ by S_n , and for any $\pi \in S_n$, we will adopt the *inline* notation $\pi = (\pi_1, \dots, \pi_n)$.

For readability, the following exact values will be denoted by their fourth decimal approximation, and viceversa, whenever using these decimal values, we mean the corresponding exact constants:

- $1 - 1/e \approx 0.6321$;
- $1/\beta \approx 0.7451$, where $\beta \in \mathbb{R}$ is the unique solution to $\int_0^1 \frac{dx}{x(1-\log x)+\beta-1} = 1$;
- $\sqrt{3} - 1 \approx 0.7321$.

The following three variants of PI are related to our work.

- **IID PI** is a specialised case where the random variables are assumed independent and identically distributed (iid). The hardness of this problem has been shown to be 0.7451 in [22]. That this is tight follows from a 0.7451-competitive *quantile strategy* devised in [9]. Remarkably, the quantile thresholds do not depend on the distribution.
- **Order Selection** (OS, also known as *free order*) is a variant where the gambler is allowed to *select* the arrival order $\pi \in S_n$ first, and then observe the values sampled from $V_{\pi_1}, \dots, V_{\pi_n}$, thus seeking to maximise

$$\frac{\mathbb{E} V_{\pi_\tau}}{\mathbb{E} \max_{i \in [n]} V_i},$$

over π and τ . This version, introduced in [19], is also central in the study of revenue maximisation in Posted Price Mechanisms [13]. After various improvements (see Section 1.5) on the competitive ratio of 0.6321 shown in [7], the state-of-the-art for OS is a 0.7251-competitive algorithm exploiting *continuous time arrival design* introduced in [28]. The 0.7451-hardness of OS follows directly from the hardness of IID PI.

- **Random Order** (RO, also known as *prophet secretary*) is a variant where random variables are shown in a uniform random order π to the gambler, who observes the values sampled from $V_{\pi_1}, \dots, V_{\pi_n}$ and seeks to maximise

$$\frac{\mathbb{E} V_{\pi_\tau}}{\mathbb{E} \max_{i \in [n]} V_i},$$

over τ only. One can equivalently see this model as arising from the OS setting, by saying that the gambler chooses the order π uniformly at random in S_n . After various improvements (see Section 1.5) on the initial approximate competitive ratio of 0.6321 achieved in [15], where the model was introduced (note that here *approximate* means that the competitive ratio is approached as the size of the instance $n \rightarrow \infty$), the state-of-the-art is a 0.6697-competitive multi-threshold algorithms known as *blind strategies* introduced in [12]. The state-of-the-art hardness for RO is 0.7321, also shown in [12].

1.2 Applications to pricing

Prophet inequalities are closely related to Posted Price Mechanisms (PPMs), which are an attractive alternative to implementing auctions and are often used in online sales [13]. The way these mechanisms work is as follows. Suppose a seller has an item to sell. Customers arrive one at a time and the seller proposes to each customer a take-it-or-leave-it price. The first customer accepting the offer pays the price and takes the item [11].

If a seller faces buyers with private information about their willingness to pay and there are no further transaction costs, an auction is optimal [13, 18, 27, 29]. However, auctions can have high transaction costs. They take time and require communication with multiple buyers. There are several circumstances under which price posting may be preferable, being much simpler, yet efficient enough [13, 11].

Since PPMs are suboptimal, it is important to know the ratio between PPMs and the optimal auction (Myerson’s auction) [11]. This ratio can be studied from the point of view of prophet inequalities: designing PPMs can be translated into designing algorithms for prophet inequality models [17, 7] and viceversa [11]. In particular, OS readily connects to ordering potential buyers in the PPM setting, while RO can be seen as taking potential buyers uniformly at random. Therefore, it is important to know if ordering the potential buyers can lead to improved performance, compared to a uniform random order. Since the introduction of OS 40 years ago, there has been significant effort spent on finding how far the benefits of ordering go: for PPMs showing that ordering yields benefits is equivalent to separating RO from OS.

1.3 Our contributions

Our main contribution consists in improving on the hardness of RO, to the extent of separating it from OS, through rigorous mathematical arguments. The new hardness of 0.7235 follows from an asymptotic analysis of the optimal algorithm applied to a particular class of instances.

1.3.1 The hard instance

We consider the following class of instances consisting of n iid random variables and a constant.

Definition 1.1. *Let $a, b, p \in \mathbb{R}$ be positive constants and fix $n \in \mathbb{N}$ large enough, so as to have a well defined random variable*

$$V \sim \begin{cases} n, & \text{w.p. } 1/n^2 \\ b, & \text{w.p. } p/n \\ 0, & \text{w.p. } 1 - p/n - 1/n^2. \end{cases}$$

Furthermore assume that $a < 1 < b$, $\log(1 + pb) < p$ and that the following conditions hold:

$$\frac{1 + bp}{1 + (b - a)p} \log \frac{1 + bp}{1 + (b - a)p} \leq ap, \quad \text{I}$$

$$(2 - p)(b - a) < 1, \quad \text{II}$$

$$\frac{1 - (2 - p)(b - a)}{1 + p(b - a)} < 1 + \frac{1}{p} \log \frac{1 + p(b - a)}{1 + pb}, \quad \text{III}$$

$$2 + pb[1 - p(b - a)] \geq 0, \quad \text{IV}$$

$$\frac{bp[1 + (b - a)p]}{(1 + pb) \log(1 + pb)} < 1. \quad \text{V}$$

We define the instance $\{V_1, V_2, \dots, V_{n+1}\}$, which consists of n iid random variables distributed as V and the constant a , that is

$$V_i \sim \begin{cases} V, & 1 \leq i \leq n \\ a, & i = n + 1. \end{cases} \quad \text{H}$$

The hard instance, belonging to this class, has parameters $a = 0.789$, $b = 1.24$, $p = 0.421$.

1.3.2 Main results

To establish a hardness for RO, it is enough to show a uniform upper bound on the competitive ratio of all stopping rules, that is an upper bound, holding for some given instance and uniformly for all stopping rules τ , on

$$\frac{\mathbb{E} V_{\pi_\tau}}{\mathbb{E} \max_i V_i}.$$

This is done by upper-bounding the expected reward $\mathbb{E} V_{\pi_T}$ of an *optimal* stopping rule T , which is a stopping rule (existing by *backward induction*) maximising the expected reward.

Theorem 1.1. *Let $a = 0.789$, $b = 1.24$, $p = 0.421$. Then for Instance H, the optimal stopping time $T \in C^{n+1}$ is such that*

$$\limsup_{n \rightarrow \infty} \frac{\mathbb{E} V_{\pi_T}}{\mathbb{E} \max_{i \in [n+1]} V_i} < 0.7235.$$

As a consequence, RO is 0.7235-hard. Therefore OS is separated from RO, and the former beats the latter.

1.3.3 Our techniques

To show separation we rely on an innovative asymptotic analysis of the optimal algorithm's acceptance thresholds, computed via backward induction, in a random arrival order setting, so as to obtain upper bounds on the competitive ratio of the optimal algorithm.

Future applications of these analytic techniques are not limited to the hard instance studied, nor to RO. In fact, this instance is found by analysing a much larger class of instances, which can be characterised as a collection of a positive constant and n iid random variables (we will informally refer to n as the size of the instance), supported on a set of three nonnegative values:

- a size-dependent value always accepted by the optimal algorithm, which gets larger but also disproportionately more unlikely as n grows;
- a null value, which is never accepted until the final step, and which gets increasingly likely as n grows;
- a positive value, which gets more unlikely (but to a lesser extent than the size-dependent value) as n grows, and whose acceptance depends on the past values and the stage the algorithm is at; we thus refer to this value as *nontrivial*.

This class of instances, which we analyse as the size grows, is thus fairly general and broadly speaking it contains the 0.7321-hard instance studied in [12, §7], which is composed of n iid random variables taking value n with probability $1/n^2$ and zero otherwise, and an additional positive constant a , which is a nontrivial value for the optimal algorithm. Our class is obtained by adding to the iid distributions an increasingly unlikely point mass at another nontrivial value.

The techniques used in the analysis can be further extended towards future work.

- They can be applied to a larger class of similar instances. For example, n iid random variables with more than one nontrivial value could be used: simulations suggest that adding one more nontrivial value is likely to yield better hardness, with the analysis still relatively feasible.
- They can be helpful in providing a rigorous foundation for the computational search of hard instances. Indeed, once our analytic strategy is executed, the problem is reduced to an optimization with respect to the parameters of the instance, which could be either approached mathematically or numerically. If approached numerically, it would typically be simple enough to be carried out via theoretical bounds on the approximation errors, thus keeping under control the propagation of machine precision errors.
- They could offer new insight into other separation problems, through suitable adaptation. There is no lack of open questions to attempt in this sense: it is not known yet whether OS is separated from IID PI, nor whether undisclosed RO is separated from disclosed RO, as we are going to explain in the following section.

1.4 Previous bounds on Random Order

To fully understand the significance of previous bounds on RO, it is best to distinguish between two versions of the mode implicitly appearing in the literature (thus the terminology adopted here is nonstandard).

- **Undisclosed RO.** Also known as *anonymous* RO, this is the original version, enforcing *undisclosed* uniform random order, as per the definition given in [15]: at time i , only the realised value of V_{π_i} is sent to the gambler, who in general will never know the label π_i of the distribution it has been sampled from (in particular cases the *a priori* knowledge of the distributions and the history of the probed values allow the gambler to deduce it). The approximate 0.6321-competitive multi-threshold algorithm designed in [15] is still the state-of-the-art for this model, and it is not known whether it is 0.6321-hard or not.
- **Disclosed RO.** Also known as *personalised* RO, this version, whose introduction is motivated by PPMs, where prices can be personalised, allows for a *disclosed* random order: at time i , the realised value V_{π_i} is sent to the gambler together with the label π_i , revealing what distribution the sample observed comes from. Thus the gambler acquires information about the order, as it observes more values. All the algorithms surpassing 0.6321-competitiveness for RO implicitly refer to this model, which is therefore the most commonly referred to in the more recent literature. It is also a more natural point of view when seeing RO within the OS framework (where every selected label π_i is clearly known in advance).

For the gambler, undisclosed RO is no easier than disclosed RO, since in the undisclosed model less information is provided.¹ However, it is not known if there is a real benefit in disclosing the order, that is if the two versions of RO are separated. Although in undisclosed RO the algorithm

¹Very recently, following the preprint version of this paper, a similar terminology has been introduced in [6], although referring to different models: *order-aware* and *order-unaware* RO. To avoid confusion, we stress that the *order-aware* model does not refer to disclosed RO; rather, it is a model of RO, where the algorithm is partially offline, due to knowing both the past and the future labels π_i . Equivalently, the order is known in advance not only by the prophet, but by the gambler as well. Our analysis is not concerned with this newly introduced variant of RO, which was absent from the literature and from [5] (the preprint version of [6]) to which only we had access at the time of writing the present work. However, our analytic techniques could easily be extended to study the optimal algorithm for this variant on Instance H. More explanations about the present work and [6] are in place. In [6, §1.1] it is claimed, without proof, that the *result* we obtain for Instance H does not apply to order-aware RO. Although we restrict ourselves to the study of online models of RO, we address the fact that, theoretically, this is actually an open question. Thus if they refer to our separation result (and not the numerical upper bound presented), their

does not know what distribution the value shown comes from *in general*, for specific instances it is possible to deduce this information, fully or partially, from the *a priori* knowledge of the distributions given and the history of the probed values. In particular, for some instances, disclosed and undisclosed RO can be equivalent. This is precisely the case for Instance H, as the support of all distributions is given as *a priori* knowledge to the algorithm: values $0, b, n$ are coming from $V_1, \dots, V_n \sim V$, so probing them at step k means that $\pi_k(\omega) \in [n]$, while probing a at step k means that $\pi_k(\omega) = n + 1$. Thus the order oblivious stopping rule has, for Instance H, a way of extracting partial information about π from the history of the probed values. Although partial, distributionally the inferred information is equivalent to the one that the disclosed RO model would allow: indeed besides V_{n+1} , all other distributions are iid and knowing also in which order they arrived in the past offers no advantage in expectation. Therefore, the distinction between the two RO models does not affect the analysis in the present work. For simplicity, being its set-up more straightforward from a measure-theoretic point of view, we will adopt the simpler framework of undisclosed RO, and the hardness achieved will nonetheless hold also for disclosed RO.

For the gambler, RO is no easier than OS: a uniform random permutation can always be selected. Therefore, the competitive ratio achievable by an algorithm with OS is at least the one achievable with RO. However, it is not known if the gambler can benefit from selecting the order, that is if RO and OS are separated.

Since its introduction in [15], the only non trivial bound for RO was the 0.7451-hardness of the IID PI model (which can be seen as a special case) proved in [22] almost 30 years earlier. Unlike the design of algorithms for RO, which made a relatively quick progress (see Section 1.5), it was not until 6 years later that the 0.7321-hard instance previously described was found in [12, §7]. On the other hand, in [28] it was proved that the competitive ratio for OS is at least 0.7251. Refining the techniques of [28], this ratio was recently improved in [6] to a lower bound of 0.7258. These mathematical arguments alone fall short of proving that RO and OS are separated.

In [6, §6] a brute force simulation improving numerically the hardness of RO also appeared. The simulated hardness is strong enough to show that RO is indeed separated from OS. In the simulation, a dynamic program is run on a 26-parameters instance, yielding a 0.7254-hardness for RO, which is less than 0.7258, the state-of-the-art competitive ratio of OS. From a mathematical standpoint, the simulation could, in principle, be extrapolated into a rigorously proven upper bound through formal error analysis, but this has not been carried out. Furthermore, the simulated upper bound of 0.7254 for RO differs from the improved lower bound for OS only by 0.0004. A study of the propagation of errors in the simulations involved would be auspicious. On the other hand, the simulated instance does not seem formally tractable, and it is complicated enough that the simulation does not lend itself to an easy error analysis. This makes it still desirable to find new arguments that establish the separation of the two models more formally. In this paper we provide an alternative mathematical proof, based on a simpler instance, yielding a significantly improved state-of-the-art hardness for RO of 0.7235. At the same time this shows rigorously the separation, and provides a more straightforward argument, since it comes without the need for

claim should be interpreted as conjecture. Clearly, the value of 0.7235 for Instance H is very likely to grow in the order-aware model. However, simulations show that it is unlikely to grow by much. Indeed, this instance is expected to improve on the state-of-the-art hardness of 0.7321 also for order-aware RO. It is also open, although less likely in our opinion, the question of whether Instance H separates order-aware RO from OS. This could be resolved through simulations and proper error analysis, which we do not present, to focus on analytical bounds. We conclude by adding that, in [6, §1.1], the authors refer to the present work as using *the same idea* as theirs. Indeed, both works generalise, in different and independent ways, the hard instance presented in [12]. Apart from this common origin of the hard instance, this work shares very little with theirs. Techniques and methodologies are very different. We conduct a theoretical and analytic study, while they use a simulation-based analysis, of the optimal algorithm for RO.

further improving on the competitive ratio of 0.7251 for OS.

1.5 Related work

In this section, we briefly review some of the vast literature on prophet inequalities, with an emphasis on IID PI, OS and RO. There is in fact a variety of extensions of prophet inequalities to combinatorial structures such as matroids, matchings and combinatorial auctions: the reader interested in these aspects is better served by surveys such as [10], which contains many recent developments, as well as [26] for an economic point of view (especially regarding PPMs); [21] is also an interesting survey, containing classical results concerning infinite instances. In Sections 1.1.2 and 1.4 we already explored the literature on hardness results, so we will use this section to review some of the developments on lower bounds, that is the progress made on the competitive ratio of the various algorithms designed for these models.

We mentioned that an optimal algorithm can be found by backward induction. However, it is of little practical use on general instances, thus the research has focused on designing similar (that is, threshold-based) but simpler algorithms.

- **IID PI.** The first to surpass the $1/2$ barrier of PI was a 0.6321-competitive algorithm based on complicated recursive functions designed in [20]; 35 years later an approximate single threshold 0.7380-competitive algorithm was found in [1], after which the tight optimal competitive ratio of 0.7451 was finally attained as reviewed in Section 1.1.2.
- **OS.** The first to surpass the $1/2$ barrier of PI were certain PPMs proved to be 0.6321-competitive in [7] (recall that PPMs and prophet inequalities are equivalent). After 8 years, further improvement came from a 0.6346-competitive algorithm actually designed for disclosed RO (recall that RO is no easier than OS) in [3]. Shortly after, a 0.6541-competitive PPM appeared in [4]. Further improvement came successively from another algorithm designed for disclosed RO: the already mentioned 0.6697-competitive algorithm introduced in [12]. Thus we reached the state-of-the-art of the 0.7251-competitive algorithm designed in [28], which has been recently polished so as to be 0.7258-competitive in [6]. There are also special cases for which algorithms can achieve an even better competitive ratio: in [1] it was shown that when each distribution in the instance occurs m times, a 0.7380-competitive algorithm can be designed for m large enough. At this point, one might wonder why no progress has been made on the hardness of RO. This is due to the fact that it requires finding the optimal ordering, and in [2] this has been shown to be NP-hard, even under a special case where all the distributions have 3-point support, and where the highest and lowest points of the support are the same for all the distributions.
- **RO.** We already mentioned that, as of yet, no algorithm, which has surpassed the initial approximate 0.6321-competitiveness established in [15], has been designed for undisclosed RO. In fact the algorithms in [3, 12], mentioned in the previous point, are designed and directly analysed for continuous distributions. This analysis is compatible with undisclosed RO; however, if point masses are allowed, the analysis is extended through a standard argument exploiting stochastic tie breaking, which requires revealing the distributional identity of the random variable from which the samples probed come from. Hence stochastic tie breaking can only be performed with disclosed RO. In this sense, not only personalisation has been used to go beyond 0.6321-competitiveness, but it has been relied upon also to design the 0.6321-competitive algorithms for RO, both in [9] (multi-threshold) and [14] (single-threshold). As a result, the question of whether personalisation is actually necessary to even fully achieve 0.6321 remains

open for undisclosed RO. One interesting fact about RO is that 0.7451-competitiveness can be recovered by simply removing a constant number of random variables: more specifically in [25] it is shown that given any instance $\mathbf{V} = \{V_1, \dots, V_n\}$, for any $\varepsilon > 0$ we can find an instance \mathbf{V}' of $n' = n'(\varepsilon)$ random variables (note that n' is independent of n) obtained by removing a constant number (only dependent on ε) of random variables from \mathbf{V} , such that there is a stopping rule $\tau \in C^{n'}$ ensuring

$$\frac{\mathbb{E} V'_\tau}{\mathbb{E} \max_{i \in [n']} V'_i} \geq 0.7451 - \varepsilon.$$

Removing random variables is not the only way to push beyond the 0.7235-hardness we prove: *large markets* hypothesis are also very helpful. For example, in [1] it was shown that, if each distribution in an instance of size n occurs at least $\Theta(\log n)$ times, a 0.7380-competitive algorithm can be designed.

1.6 Organisation of the paper

In Section 2, we provide some background on optimal stopping theory for finite horizon random arrival order problems and apply it to RO and Instance H. In Section 2.1, the existence of an optimal stopping rule for this setting is shown, by deriving backward induction acceptance thresholds. We state the explicit characterisation of these thresholds for Instance H, and describe the optimal stopping rule through its acceptance times in Section 2.2, which is the starting point of our analysis. In Section 2.3, we derive sharp asymptotic estimates for two of the acceptance times and determine the asymptotic ordering of all acceptance times (Lemma 2.1).

In Section 3, we first compute the prophet's expectation (Lemma 3.1) and the optimal algorithm's expectation (Proposition 3.1) and derive sharp asymptotic estimates. Then we analyse the ratio, so as to reduce the problem of upper-bounding it, to an optimisation problem, solving which the upper bound of 0.7235 on the competitive ratio of the optimal algorithm is derived, yielding the separation between OS and RO (Theorem 1.1).

The reader can find in the Appendix:

- some elements of optimal stopping theory and the corresponding measure theoretic background for the reader unfamiliar with the topics (Appendices A and B);
- the more technical of the proofs (Appendices C to E);
- formal error analysis and reference to both the shared code used to determine the numerical value of our estimate of the gambler-to-prophet ratio of the optimal algorithm, and the shared code that suggests tightness of the estimate (Appendix F).

2 Preliminaries

In this section, we adapt classical tools from finite horizon optimal stopping theory to RO, with the goal of describing an optimal stopping rule for Instance H in terms of acceptance times, which we asymptotically estimate, as the size of the instance grows.

2.1 Optimal stopping rule for Random Order

The classical theory of optimal stopping generally studies fixed order stochastic processes. The transition to random order for finite horizon is quite natural, although it requires some prelimi-

nary measure-theoretic remarks. In this section, we clarify this foundation so that even a reader unfamiliar with the topic understands the construction of an optimal stopping rule for RO.

2.1.1 Finite horizon optimal stopping for random order processes

By (finite horizon) random order (stochastic) process we mean a random vector whose components are permuted uniformly at random. In [8, Theorem 3.2], a standard backward induction result for (finite horizon) optimal stopping is shown. This formulation does not assume independence of the components of the random vector $X = (V_1, \dots, V_{n+1})$, where $n+1$ is the finite horizon we will work with (from now on omitted), but it does assume that the order is fixed. Nonetheless, the theorem can be naturally extended to random order processes, by making use of a filtration that accounts for the randomly permuted order of arrival of the $\{V_i\}$.

A random order process is formally obtained by replacing the indices of the vector's components with those of a uniform random permutation of $[n+1]$, $\pi = (\pi_1, \dots, \pi_{n+1})$, obtaining a randomly permuted random vector $X^\pi := (V_{\pi_1}, \dots, V_{\pi_{n+1}})$. The natural filtration of this process is similarly obtained by formally starting with $\mathcal{F}_0 := \{\emptyset, \Omega\}$, and for all $k \in [n+1]$, letting $\mathcal{F}_k := \sigma(V_{\pi_1}, \dots, V_{\pi_k})$, where $\sigma(V_{\pi_1}, \dots, V_{\pi_k})$ denotes the smallest σ -algebra providing all information regarding both sources of randomness: the random order and the random values, up to time k . Given a stopping time τ , that is, a random variable such that $\{\tau = k\} \in \mathcal{F}_k$, we can stop the process X^π , and the stopped value is that of the random variable V_{π_τ} . All rigorous measure-theoretic constructions, supported by a suitable probability space $(\Omega, \mathcal{F}, \mathbb{P})$, are given in Appendix A. For Instance H we will be able to avoid the measure-theoretic formalism, so we will not rely directly on most of those constructions, included merely to provide a solid background to the natural extension of [8, Theorem 3.2], from which we will derive the formalism used in this paper.

Let C_k^{n+1} be the set of stopping rules τ that never stop before time k , that is such that $k \leq \tau \leq n+1$. Note that we previously denoted $C^{n+1} := C_1^{n+1}$. For ease of notation, in this section we omit the reference to the finite horizon and simply denote these classes as C_k . We will also denote as $\mathbb{E}_{\mathcal{F}_k}$ the conditional expectation with respect to the σ -algebra \mathcal{F}_k and omit the reference to the permutation from X^π , thus simply denoted as X . We conclude by providing the heuristics of how to optimally stop X , so as to maximise the expected value obtained by the gambler.

Backward induction is a strategy for the gambler based on the following premise: *take a value only if it is larger than the expected future reward*. Recall that in our framework the rewards come from the observed values $V_{\pi_1}, \dots, V_{\pi_{n+1}}$. Then this strategy defines backwards random rewards $\gamma_{n+1}, \dots, \gamma_2, \gamma_1$ and stopping times s_{n+1}, \dots, s_1, s_2 . For each $k \in [n+1]$, consider the case where the gambler rejected the first $k-1$ values. Then, the value $\mathbb{E} \gamma_k$ represents the expected reward that the gambler obtains using backward induction, characterised through the stopping times s_k , which implement the stopping rule of accepting the first value V_{π_l} that exceeds the expected future reward (conditionally on the rejected values) $\mathbb{E}_{\mathcal{F}_l} \gamma_{l+1}$ for $l \geq k$. Theorem A.1 shows that the expected reward of the stopping rule s_1 is optimal. Rigorous definitions and statements are provided in Appendix A.

2.1.2 Applications to Instance H

In this section, we consider the special case of Theorem A.1 for Instance H, leading to a more elementary formalism, sufficient to describe an optimal stopping rule, so as to make explicit computations more accessible.

Recall that, for Instance H, all components of $X = (V_{\pi_1}, \dots, V_{\pi_{n+1}})$ are independent and the

process takes values in the finite state space

$$\mathcal{S}_n := \{(x_1, \dots, x_{n+1}) : \exists! i \in [n+1], x_i = a, \forall j \neq i, x_j \in \{0, b, n\}\}.$$

Our random order process can be defined, through the standard construction developed in Appendix A, starting from the product space

$$(\mathcal{S}_{n+1} \times \mathcal{S}_n, 2^{\mathcal{S}_{n+1}} \otimes 2^{\mathcal{S}_n}, \mu \otimes \nu),$$

where:

- $\nu := \underbrace{\mathcal{D} \otimes \dots \otimes \mathcal{D}}_n \otimes \delta_a$;
- \mathcal{D} is the law of V ;
- δ_a is the Dirac measure at a ;
- $\mu \equiv \frac{1}{(n+1)!}$ is the uniform probability law on \mathcal{S}_{n+1} ;
- \otimes denotes the standard construction of the product measure and σ -algebra.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the supporting probability space and let \mathbb{E} denote the corresponding expectation. We can replace conditioning on the random order filtration with elementary conditioning on all the possible values of the random variables V_1, \dots, V_{n+1} in all the possible arrangements without loss of generality, due to the σ -algebras \mathcal{F}_k being generated by $\{(X_1 = x_1, \dots, X_k = x_k) : x \in \mathcal{S}_n\}$ for all $k \in [n+1]$.

Let us adopt the same notation as for Theorem A.1, and restate it in this discrete setting. For every $x \in \mathcal{S}_n$, define backwards $\gamma_{n+1}, \gamma_n, \dots, \gamma_1$ as functions of x , by setting

$$\gamma_{n+1}(x_1, \dots, x_{n+1}) := x_{n+1}$$

and for all $l = n, \dots, 1$,

$$\gamma_l(x_1, \dots, x_l) := \max\{x_l, \bar{\gamma}_l(x_1, \dots, x_l)\},$$

where

$$\bar{\gamma}_l(x_1, \dots, x_l) := \mathbb{E}[\gamma_{l+1}(x_1, \dots, x_l, X_{l+1}) | X_1 = x_1, \dots, X_l = x_l].$$

For each $k = n+1, \dots, 1$, let

$$s_k := \inf\{l \geq k : x_l = \gamma_l(x_1, \dots, x_l)\}$$

and denote $T := s_1$. Then, the following is a restatement of Theorem A.1 for discrete instances. In particular, it applies to Instance H.

Theorem 2.1. *Consider $X := (V_{\pi_1}, \dots, V_{\pi_{n+1}})$, the backward induction values $\gamma_{n+1}, \gamma_n, \dots, \gamma_1$ and the stopping times s_{n+1}, s_n, \dots, s_1 as previously defined. Then for any $k \in [n+1]$, $s_k \in C_k$ and for any realization (x_1, \dots, x_k) , the value $\gamma_k(x_1, \dots, x_k)$ is the optimal value the gambler can obtain from time k . More formally, for all stopping rules $\tau \in C_k$ we have that*

$$\mathbb{E}(X_{s_k} | X_1 = x_1, \dots, X_k = x_k) = \gamma_k(x_1, \dots, x_k) \geq \mathbb{E}(X_\tau | X_1 = x_1, \dots, X_k = x_k).$$

Thus for all stopping rules $\tau \in C_k$,

$$\mathbb{E} X_{s_k} = \mathbb{E} \gamma_k(X_1, \dots, X_k) \geq \mathbb{E} X_\tau.$$

In particular, for all stopping rules $\tau \in C_1$,

$$\mathbb{E} V_{\pi_T} := \mathbb{E} X_T = \mathbb{E} \gamma_1(X_1) \geq \mathbb{E} X_\tau := \mathbb{E} V_{\pi_\tau}.$$

Thus $\mathbb{E} \gamma_1(X_1) = \sup_{\tau \in C_1} \mathbb{E} V_{\pi_\tau}$.

Recall that Instance [H](#) has the following property: with undisclosed RO the stopping rule, which at any time $k \in [n+1]$ is not given information about the past values of the random ordering π_1, \dots, π_k , will nonetheless be able to infer, step by step, whether the values probed came from V or not, by simply observing if a has already been probed or not. Moreover, from the fact that

$$\mathbb{P}(\pi_{k+1} = n+1 | X_1 = x_1, \dots, X_k = x_k) = \begin{cases} 0, & \exists 1 \leq i \leq k : x_i = a \\ \frac{1}{n-k+1}, & \text{otherwise,} \end{cases}$$

since $\{\pi_{k+1} = n+1\} = \{X_{k+1} = a\}$, we can compute the quantities in [Theorem 2.1](#) as follows.

Remark 2.1. For Instance [H](#),

$$\bar{\gamma}_n(x_1, \dots, x_n) = \begin{cases} \mathbb{E} V, & \exists 1 \leq i \leq n : x_i = a \\ a, & \text{otherwise,} \end{cases}$$

where $\mathbb{E} V = \frac{1+bp}{n}$, and for all $1 \leq k < n$,

$$\bar{\gamma}_k(x_1, \dots, x_k) = \begin{cases} \mathbb{E} \gamma_{k+1}(x_1, \dots, x_k, V), & \exists 1 \leq i \leq k : x_i = a \\ \frac{\gamma_{k+1}(x_1, \dots, x_k, a)}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \mathbb{E} \gamma_{k+1}(x_1, \dots, x_k, V), & \text{otherwise,} \end{cases}$$

where

$$\mathbb{E} \gamma_{k+1}(x_1, \dots, x_k, V) = \frac{\gamma_{k+1}(x_1, \dots, x_k, n)}{n^2} + \frac{p\gamma_{k+1}(x_1, \dots, x_k, b)}{n} + \gamma_{k+1}(x_1, \dots, x_k, 0) \left(1 - \frac{p}{n} - \frac{1}{n^2}\right).$$

2.2 Acceptance times

Computing the expected value of the optimal algorithm $\mathbb{E} V_{\pi_T}$ explicitly as the size of the instance grows is usually infeasible: in [Appendix B](#) additional comments regarding this matter are provided; moreover, we also provide some background on the fact that there is no loss of generality in T being a nonrandomised stopping rule. Nonetheless, [Theorem 2.1](#) enables us to infer features of the optimal stopping rule T on Instance [H](#), yielding asymptotic estimates of $\mathbb{E} V_{\pi_T}$ very close to its true value. In particular T can be described in terms of *acceptance times*: for each value $0, a, b, n$, there is a time before which the value is never accepted (that is, if it were probed, the algorithm would not stop with it as reward) and such that, from that time onward, the value would always be accepted (that is, if it were probed, the algorithm would stop with it as reward). In this section, we rigorously define these acceptance times. They are deterministic ([Definitions 2.1 to 2.3](#)) and can be computed using simple iterative equations ([Remark 2.5](#)). This leads to the asymptotic estimates and order restrictions in the next section ([Lemma 2.1](#)).

The acceptance times for the values 0 and n are trivial: if n is probed, the optimal algorithm always stops; if 0 is probed, it does not stop unless it is in the last step.

Remark 2.2. For all $x \in \mathcal{S}_n$ and $k \in [n]$, we have $0 < \bar{\gamma}_k(x_1, \dots, x_k) \leq n$. Thus the acceptance time for n is 1 and the acceptance time for 0 is $n + 1$.

The acceptance times for the values a and b are nontrivial. Moreover, the earliest time that b may be accepted is random (and thus of no use in the argument of Proposition 3.1). Therefore, we define two acceptance times for the value b , depending on whether the value a has already been probed or not. The existence of acceptance times follows from the monotonicity of the future expectation, which can be inferred from the nested maxima in their formulation, preceding Theorem 2.1.

Remark 2.3. For every $x \in \mathcal{S}_n$ fixed, we have that for every $k \in [n - 1]$, $\bar{\gamma}_k(x_1, \dots, x_k) \geq \bar{\gamma}_{k+1}(x_1, \dots, x_{k+1})$.

To define the acceptance times for a and b , we exploit the fact that, informally speaking, the thresholds are deterministic given the information regarding whether a has already been revealed or not. More formally, for $i \in [n]$, denote the event that a is probed at time i by

$$\Omega_i := \{\pi_i = n + 1\} = \{X_i = a\},$$

and denote the corresponding conditional expectation $\mathbb{E}_i(\cdot) := \mathbb{E}(\cdot \mid \Omega_i)$ and conditional probability $\mathbb{P}_i(\cdot) := \mathbb{P}(\cdot \mid \Omega_i)$. For every $k \in [n + 1]$, denote the event that the value a is probed by time k as

$$\Pi_k := \bigcup_{i=1}^k \Omega_i = \{\exists 1 \leq i \leq k : \pi_i = n + 1\}$$

and denote its generated σ -algebra by $\sigma(\Pi_k)$. An immediate consequence of Remark 2.1 is the following.

Remark 2.4. In the measure-theoretic notation of Theorem A.1, for every $k \in [n]$,

$$\bar{\gamma}_k(X_1, \dots, X_k) := \mathbb{E}[\gamma_{k+1}(X_1, \dots, X_{k+1}) \mid X_1, \dots, X_k]$$

is $\sigma(\Pi_k)$ -measurable. Since for every $k \in [n]$, any $x \in \mathcal{S}_n$ and $\omega \in \{X_1 = x_1, \dots, X_k = x_k\}$, $\bar{\gamma}_k(X_1, \dots, X_k)(\omega) = \bar{\gamma}_k(x_1, \dots, x_k)$, avoiding measure theoretic terminology, we can equivalently state the $\sigma(\Pi_k)$ -measurability in the notation of Theorem 2.1 by saying that for every $k \in [n]$ and for any $x \in \mathcal{S}_n$, $\bar{\gamma}_k(x_1, \dots, x_k)$ is determined solely by the time step k and whether there is $i \in [k]$ such that $x_i = a$ or not.

Thus we can further simplify the expected future rewards. Recall that the optimal stopping rule T is such that it accepts the currently probed value only if it is larger than the expected future reward. In general the future reward depends on all values that the gambler has already seen in the past up to the present. By Remark 2.4, for Instance H, the future reward can be computed solely based on the time stage k and whether the value a has been probed or not. The future rewards under each of these scenarios are then represented as follows. For every $k \in [n]$ and $x \in \mathcal{S}_n$,

$$\begin{aligned} \phi_k^{n+1} &:= \mathbb{E}(V_{\pi_T} \mid T > k, \Pi_k) = \bar{\gamma}_k(x_1, \dots, x_i = a, \dots, x_k) \\ \bar{\phi}_k^{n+1} &:= \mathbb{E}(V_{\pi_T} \mid T > k, \Pi_k^c) = \bar{\gamma}_k(x_1 \neq a, \dots, x_k \neq a). \end{aligned}$$

As usual, we will omit the time horizon from the notation and denote the expected future reward when a has been already probed at time k simply by ϕ_k and the expected future reward when a has not been already probed yet at time k simply by $\bar{\phi}_k$. Furthermore we will often denote $x \vee y := \max\{x, y\}$.

Remark 2.5. By Remark 2.1 it follows that, for Instance H , we have

$$\begin{aligned}\phi_n &= \mathbb{E} V, \\ \bar{\phi}_n &= a,\end{aligned}$$

where $\mathbb{E} V = \frac{1+bp}{n}$, and for all $k \in [n-1]$,

$$\begin{aligned}\phi_k &= \mathbb{E}(V \vee \phi_{k+1}), \\ \bar{\phi}_k &= \frac{a \vee \phi_{k+1}}{n+1-k} + \left(1 - \frac{1}{n+1-k}\right) \mathbb{E}(V \vee \bar{\phi}_{k+1}).\end{aligned}$$

We first introduce the acceptance times for a .

Definition 2.1. Denote the earliest (deterministic) time that the value a would be accepted if probed by

$$j_n := \inf\{k \in [n+1] : a \geq \phi_k\}.$$

From Remark 2.3, it follows that, for all times after j_n , the optimal stopping rule will always accept the value a if probed, and stop. Moreover, for all large enough n , we have that $j_n \leq n$ since $\phi_n = (1+b)/n \leq a$.

Next, we introduce the two acceptance times for the value b , relative to whether a has already been probed or not.

Definition 2.2. Denote the earliest (deterministic) time that the value b would be accepted if probed, given that the value a has already been probed, by

$$k_n := \inf\{k \in [n+1] : b \geq \phi_k\}.$$

From Remark 2.3, it follows that, if the value a has already been probed, then, for all times after k_n , the optimal stopping rule will accept the value b if probed, and stop. Moreover, for all large enough n , we have that $k_n \leq n$ since $\phi_n = (1+b)/n \leq b$.

Definition 2.3. Denote the earliest (deterministic) time that the value b would be accepted if probed, given that the value a has not been probed, by

$$\bar{k}_n := \inf\{k \in [n+1] : b \geq \bar{\phi}_k\}.$$

From Remark 2.3, it follows that, if the value a has not been probed, then, for all times after \bar{k}_n , the optimal stopping rule will accept the value b if probed, and stop. Moreover, for all large enough n , we have that $\bar{k}_n \leq n$, since $a < b$ and, at time n , not having probed the value a implies that $\pi_{n+1}(\omega) = n+1$, that is $x_{n+1} = a$.

2.3 Asymptotic estimates for the acceptance times

In this section we derive sharp asymptotic estimates on j_n , k_n , and the relative order of j_n , k_n and \bar{k}_n . These results are at the heart of the asymptotics of $\mathbb{E} V_{\pi_T}$. The full proof is given in Appendix C. We provide here its main ideas, and a graphical representation for the hard instance in Figure 1.

Lemma 2.1. As $n \rightarrow \infty$, the optimal stopping rule for Instance H is such that we have the following.

- a) $j_n \sim n \left(1 + \frac{1}{p} \log \frac{1+(b-a)p}{1+bp}\right)$.
- b) $k_n \sim n \left(1 + \frac{1}{p} \log \frac{1}{1+bp}\right)$.
- c) $k_n \leq j_n$. Informally, the gambler accepts the value a later than the value b after seeing the value a .
- d) $k_n \leq \bar{k}_n$. Informally, the gambler accepts the value b not having seen the value a later than the value b after seeing the value a .
- e) $\bar{k}_n \leq j_n$. Informally, the gambler accepts the value a later than the value b not having seen the value a .

Idea of the proof.

- a) By iterating the formulas in Remark 2.5 we obtain a close expression for $\{\phi_k\}$, which can be turned into a sharp asymptotics for j_n as $n \rightarrow \infty$, since it is the smallest k such that $a \geq \phi_k$.
- b) Since k_n is the smallest k such that $b \geq \phi_k$, the asymptotics for k_n is obtained similarly to the one for j_n in Lemma 2.1 (a).
- c) Follows from the fact that, as $n \rightarrow \infty$, the limit of j_n/n obtained in Lemma 2.1 (a) is larger than the limit of k_n/n obtained in Lemma 2.1 (b) since $a < b$.
- d) Assume by contradiction that $\bar{k}_n < k_n$ infinitely often as $n \rightarrow \infty$. For all such n this implies that $\bar{\phi}_{k_n-1} \leq b$. We will reach a contradiction with this fact as follows. First, through Remark 2.5, we derive an iterative lower bound on $\bar{\phi}_k$ for all $k \geq k_n - 1$, for all such n . By exploiting the asymptotics obtained in Lemma 2.1 (b), we derive further asymptotic estimates, which show that $\bar{\phi}_{k_n} > b$ as $n \rightarrow \infty$, which yields the contradiction sought.
- e) Assume by contradiction that $j_n < \bar{k}_n$ infinitely often as $n \rightarrow \infty$. We first derive, through Remark 2.5, an iterative upper bound on $\{\bar{\phi}_k\}$ for all $k \geq j_n$, for all such n . Then we consider the earliest time that b is greater than this upper bound, and denote it as k_n^* . This deterministic time has the following properties, as $n \rightarrow \infty$.
 - $\bar{k}_n \leq k_n^*$.
 - $k_n^* \sim \gamma n$, where $0 < \gamma = \gamma(a, b, p) < 1$ is found explicitly.

Using the value of γ and Lemma 2.1 (a), we have that, $k_n^* < j_n$ for all sufficiently large n . This is a contradiction with the hypothesis that $j_n < \bar{k}_n \leq k_n^*$ for infinitely many values of n .


□

3 Random Order is 0.7235-hard

We start by calculating a sharp asymptotic estimate for the prophet's expectation. The computation is given in Appendix D.

Lemma 3.1. *For Instance H as $n \rightarrow \infty$,*

$$\mathbb{E} \max_{i \in [n+1]} V_i = 1 + b(1 - e^{-p}) + ae^{-p} + \mathcal{O}\left(\frac{1}{n}\right).$$



(a) Future rewards $\{\phi_k\}$ (blue) and $\{\bar{\phi}_k\}$ (amber)

(b) Zoom in at the intersection $\phi_k \approx b$

Figure 1: Simulation of the dynamic program (reference to the code shared is in Appendix F) for Instance H with $a = 0.789$, $b = 1.24$, $p = 0.421$, $n = 10^6$. Figure 1a shows the sequences $\{\phi_k\}$ (blue), $\{\bar{\phi}_k\}$ (amber), and the values a (green), b (red). Figure 1b shows a zoom on the intersection where $\phi_k \approx b$. Informally, the abscissa of the intercept of the blue dotted curve with the red line corresponds to the smallest acceptance time $k_{10^6} \approx 2253$; the abscissa of the intercept of the amber dotted curve with the red line is the second largest acceptance time $\bar{k}_{10^6} \approx 211231$; the abscissa of the intercept of the blue dotted curve with the green line is the largest acceptance time $j_{10^6} \approx 415187$, as per Lemma 2.1 (c, d, e).

Next we derive sharp asymptotic estimates for the expectation of the optimal algorithm, relying on the eventual reciprocal ordering being $k_n \leq \bar{k}_n \leq j_n$ by Lemma 2.1 (c, d, e), and exploiting the partitioning $\{\Omega_i\}$ of the probability space, through the law of total expectation. The proof is given in Appendix E.

Proposition 3.1. For Instance H as $n \rightarrow \infty$,

$$\mathbb{E} V_{\pi_T} = q_{a,b,p}(\lambda_n, \mu_n, \nu_n) + \mathcal{O}\left(\frac{1}{n}\right),$$

where $\lambda_n := j_n/n$, $\mu_n := k_n/n$, $\nu_n := \bar{k}_n/n$ and $q_{a,b,p}(\lambda, \mu, \nu)$ is a multivariate exponential quadratic in the variables λ, μ, ν and parameters a, b, p defined as

$$\begin{aligned} q_{a,b,p}(\lambda, \mu, \nu) &= \frac{\mu^2}{2} - \frac{\nu^2}{2} + \nu + \frac{1}{p} + b + \left(\frac{1}{p} - \mu\right) \left(\frac{1}{p} + b\right) e^{p(\mu-1)} \\ &\quad + \left[\left(\frac{1}{p} + b\right) (\nu - \lambda) - \frac{a}{p}\right] e^{p(\nu-1)} - \frac{1}{p} \left(\frac{1}{p} + b - a\right) e^{p(\nu-\lambda)}. \end{aligned}$$

Lastly we prove the 0.7235-hardness and the separation.

Proof of Theorem 1.1. By Proposition 3.1 and Lemma 2.1 (a, b), letting

$$\lambda_* := 1 + \frac{1}{p} \log \frac{1 + (b-a)p}{1 + bp}$$

and

$$\mu_* := 1 + \frac{1}{p} \log \frac{1}{1 + bp},$$

it follows that

$$\mathbb{E} V_{\pi_T} = q_{a,b,p}(\lambda_*, \mu_*, \nu_n) + \mathcal{O}(1).$$

For simplicity we will omit the reference to the parameters a , b , p , λ_* and μ_* in the notation of the exponential quadratic. The dependence on n is implicit in the notation: recall that in $\mathbb{E} V_{\pi_T}$, $T \in C^{n+1}$. Since as $n \rightarrow \infty$, $\mu_n \rightarrow \mu_*$, $\mu_n \leq \nu_n \leq \lambda_n$ and $\lambda_n \rightarrow \lambda_*$,

$$\mathbb{E} V_{\pi_T} \leq \max_{\nu \in [\mu_*, \lambda_*]} q(\nu) + \mathcal{O}(1).$$

By Lemma 3.1, as $n \rightarrow \infty$

$$\frac{\mathbb{E} V_{\pi_T}}{\mathbb{E} \max_{i \in [n+1]} V_i} \leq \frac{\max_{\nu \in [\mu_*, \lambda_*]} q(\nu)}{1 + b(1 - e^{-p}) + ae^{-p}} + \mathcal{O}(1).$$

Denote as $m(a, b, p) := \max_{\nu \in [\mu_*, \lambda_*]} q(\nu)$. Then

$$\limsup_{n \rightarrow \infty} \frac{\mathbb{E} V_{\pi_T}}{\max_{i \in [n+1]} V_i} \leq \frac{m(a, b, p)}{1 + b(1 - e^{-p}) + ae^{-p}} := M(a, b, p). \quad (3.1)$$

We start by computing $m(a, b, p)$. Then the maximisation is performed on $\mu_* \leq \nu \leq \lambda_*$. Note that

$$q'(\nu) = 1 - \nu + \left[\left(\frac{1}{p} + b \right) + (1 + bp)(\nu - \lambda_*) - a \right] e^{p(\nu-1)} - \left(\frac{1}{p} + b - a \right) e^{p(\nu-\lambda_*)},$$

and since

$$\left(\frac{1}{p} + b - a \right) e^{p(\nu-\lambda_*)} = \left(\frac{1}{p} + b \right) e^{p(\nu-1)},$$

it follows that

$$\begin{aligned} q'(\nu) &= 1 - \nu + [(1 + bp)(\nu - \lambda_*) - a] e^{p(\nu-1)}, \\ q''(\nu) &= -1 + \{1 + p[b - a + (1 + bp)(\nu - \lambda_*)]\} e^{p(\nu-1)}, \end{aligned}$$

and

$$q'''(\nu) = p\{2 + p[2b - a + (1 + bp)(\nu - \lambda_*)]\} e^{p(\nu-1)} > 0,$$

since

$$0 \geq \nu - \lambda_* \geq \mu_* - \lambda_* = -\frac{1}{p} \log[1 + p(b - a)]$$

for all $\mu_* \leq \nu \leq \lambda_*$ implies that

$$2 + p[2b - a + (1 + bp)(\nu - \lambda_*)] \geq 2 + p \left[2b - a - \frac{1 + bp}{p} \log[1 + p(b - a)] \right] \geq 2 + pb[1 - p(b - a)] > 0,$$

where we used $\log(1 + x) \leq x$ in the second last inequality and Condition IV in the last inequality. Thus $q'(\nu)$ is convex in the interval considered for maximisation. Since

$$q'(\mu_*) = \frac{1}{p} \log \left(\frac{1 + bp}{1 + (b - a)p} \right) - \frac{a}{1 + bp}$$

and

$$q'(\lambda_*) = \frac{1}{p} \log \left(\frac{1 + bp}{1 + (b - a)p} \right) - a \frac{1 + (b - a)p}{1 + bp},$$

we can conclude that $q'(\mu_*) > q'(\lambda_*)$. Since by Condition I we have $q'(\lambda_*) \leq 0$, by convexity we have only two possibilities (depending on a , b , p).




Figure 2: Root ν_* of $q'(\nu)$ in $[\mu_*, \lambda_*]$

- $q'(\mu_*) \leq 0$: then $q'(\nu) \leq 0$ for all $\mu_* \leq \nu \leq \lambda_*$; in this case $q(\nu)$ is nonincreasing and therefore $q(\mu_*) = \max_{\nu \in [\mu_*, \lambda_*]} q(\nu)$, hence

$$m(a, b, p) = q(\mu_*). \quad (3.2)$$

Plugging (3.2) into the definition of $M(a, b, p)$ in (3.1) yields

$$M(a, b, p) = \frac{q(\mu_*)}{1 + b(1 - e^{-p}) + ae^{-p}};$$

- $q'(\mu_*) > 0$: then $q'(\nu_*) = 0$ for some $\mu_* < \nu_* < \lambda_*$ and $q(\nu)$ is not monotonic: it increases until it reaches its maximum at ν_* and then decreases. We find ν_* numerically, since it is a nontrivial zero of the exponential linear polynomial $q'(\nu)$. The properties of the function $q'(\nu)$ in the interval $[\mu_*, \lambda_*]$ allow for numerical methods, such as *bisection*, to determine ν_* with arbitrary accuracy up to machine precision (we will adopt the Python method for scalar root finding from the *scipy* library, in the *scipy.optimize* package, *bisect*, with a tolerance of roughly 10^{-13}), since $q'(\nu)$ is smooth and has opposite signs at the ends of the interval (see Figure 2 for a representation). Hence we numerically determine $q(\nu_*) = \max_{\nu \in [\mu_*, \lambda_*]} q(\nu)$, thus

$$m(a, b, p) = q(\nu_*). \quad (3.3)$$

Plugging (3.3) into the definition of $M(a, b, p)$ in (3.1) yields

$$M(a, b, p) = \frac{q(\nu_*)}{1 + b(1 - e^{-p}) + ae^{-p}}.$$

Fix $a = 0.789$, $b = 1.24$, $p = 0.421$. It is possible to determine numerically that we are in the second nonmonotonic scenario and

$$M(a, b, p) \approx 0.72349 < 0.7235.$$

The complete list of correct decimals produced are shared in Appendix F, where we also show that the error committed is at most 10^{-12} .

The competitive ratio of the optimal algorithm solving undisclosed RO has been shown to be less than 0.7235, since for Instance [H](#) it attains a gambler-to-prophet ratio lower than this value, for all n large enough. Thus no algorithm can achieve a better competitive ratio and RO is 0.7235-hard. Recall that since for Instance [H](#) undisclosed RO and disclosed RO are equivalent, the hardness of 0.7235 applies to both models. The separation from OS follows from the algorithm designed in [\[28\]](#), which ensures a competitive ratio for OS of $0.7251 > 0.7235$. \square

As a concluding remark, we would like to observe that the sharpness of the methods used to estimate the gambler-to-prophet ratio of the optimal algorithm T on Instance [H](#) can be checked through simulations. In order to get an estimate of how close to the optimal limiting value $M(a, b, p) \approx 0.72349$ may be, we can perform simulations of the dynamic program (the reference to the shared code is in Appendix [F](#)) producing a gambler-to-prophet ratio of approximately 0.72354 with $n = 10^4$, 0.72349 with $n = 10^5$ and $n = 10^6$. This suggests that the competitive ratio obtained through our mathematical analysis is close to optimal for Instance [H](#).

Acknowledgements

This research was partially supported by the EPSRC grant EP/W005573/1, the ERC CoG 863818 (ForM-SMArt) grant, and the ANID Chile grant ACT210005. We would like to thank José Correa and Bruno Zilotto for their precious advice, and Mona Mohammadi and Roodabeh Safavi for early conversations.

References

- [1] M. Abolhasani, S. Ehsani, H. Esfandiari, M. T. Hajiaghayi, R. Kleinberg, and B. Lucier. Beating $1 - 1/e$ for ordered prophets. In *STOC*, 66-71, ACM, 2017.
- [2] S. Agrawal, J. Sethuraman, and X. Zhang. On optimal ordering in the optimal stopping problem. In *EC*, 187-188, ACM, 2020.
- [3] Y. Azar, A. Chiplunkar, and H. Kaplan, Prophet secretary: surpassing the $1 - 1/e$ barrier. In *EC*, 303-318, ACM, 2018.
- [4] H. Beyhaghi, N. Golrezaei, R. P. Leme, M. Pal, and B. Sivan. Improved approximations for posted price and second price mechanisms. *Oper. Res.*, 69(6):1805-1822, 2021.
- [5] A. Bubna and A. Chiplunkar. Prophet Inequality: order selection beats random order. arXiv:2211.04145v1, *Preprint*, 2022.
- [6] A. Bubna and A. Chiplunkar. Prophet inequality: order selection beats random order. In *EC*, 302-336, ACM, 2023.
- [7] S. Chawla, J. D. Hartline, D. L. Malec, and B. Sivan. Multi-parameter mechanism design and sequential posted pricing. In *STOC*, 311-320, ACM, 2010.
- [8] Y. S. Chow, H. Robbins, and D. Siegmund. Great expectations: the theory of optimal stopping. Houghton Mifflin, 1971.
- [9] J. Correa, P. Foncea, R. Hoeksma, R. Oosterwijk, and T. Vredeveld. Posted price mechanisms for a random stream of customers. In *EC*, 169-189, ACM, 2017.
- [10] J. Correa, P. Foncea, R. Hoeksma, R. Oosterwijk, and T. Vredeveld. Recent developments in prophet inequalities. *SIGecom Exch.*, 17(1):61-70, 2018.
- [11] J. Correa, P. Foncea, D. Pizzarro, and V. Verdugo. From pricing to prophets, and back! *Oper. Res. Lett.*, 47:26-29, 2019.
- [12] J. Correa, R. Saona, and B. Ziliotto. Prophet secretary through blind strategies. *Math. Program.*, 190(1):483-521, 2021.
- [13] L. Einav, C. Farronato, J. Levin, and N. Sundaresan. Auctions versus posted prices in online markets. *J. Political Econ.*, 126(1):178-215, 2018.
- [14] S. Eshani, M. Hajiaghayi, M. Kesselheim, and S. Singla. Prophet secretary for combinatorial auctions and matroids. In *SODA*, 700-714, SIAM, 2018.
- [15] H. Esfandiari, M. Hajiaghayi, V. Liaghat, and M. Monemizadeh. Prophet secretary. *SIAM J. Discrete Math.*, 31(3):1685-1701, 2017.
- [16] T. S. Ferguson. Mathematical statistics: a decision theoretic approach. Academic Press, 1967.
- [17] M. T. Hajiaghayi, R. Kleinberg, and T. Sandholm. Automated online mechanism design and prophet inequalities. In *AAAI*, 58-65, 2007.
- [18] M. Harris and R. M. Townsend. Resource allocation under asymmetric information. *Econometrica*, 49(1):33-64, 1981.

- [19] T. P. Hill. Prophet inequalities and order selection in optimal stopping problems. *Proc. Am. Math. Soc.*, 88(1):131-137, 1983.
- [20] T. P. Hill and R. P. Kertz. Comparisons of stop rules and supremum expectations of i.i.d. random variables. *Ann. Probab.*, 10(2):336-345, 1982.
- [21] T. P. Hill and R. P. Kertz. A survey of prophet inequalities in optimal stopping theory. *Contemp. Math*, 125:191-207, 1992.
- [22] R. P. Kertz. Stop rule and supremum expectations of i.i.d. random variables: a complete comparison by conjugate duality. *J. Multivar. Anal.*, 19(1):88-112, 1986.
- [23] U. Krengel and L. Sucheston. Semiamarts and finite values. *Bull. Amer. Math. Soc.*, 83(4):745-747, 1977.
- [24] U. Krengel and L. Sucheston, On semiamarts, amarts, and processes with finite value. *Adv. Probab. Related Topics*, 4:197–266, 1978.
- [25] A. Liu, R. P. Leme, M. Pál, J. Schneider, and B. Sivan. Variable decomposition for prophet inequalities and optimal ordering. In *EC*, 692, ACM, 2021.
- [26] B. Lucier. An economic view of prophet inequalities. *SIGecom Exch.*, 16(1):24-47, 2017.
- [27] R. B. Myerson. Optimal auction design. *Math. Oper. Res.*, 6(1):58–73, 1981.
- [28] B. Peng and Z. G. Tang. Order selection prophet inequality: from threshold optimization to arrival time design. In *FOCS*, 171-178, IEEE, 2022.
- [29] J. G. Riley and W. F. Samuelson. Optimal auctions. *Am. Econ. Rev.*, 71(3):381–92, 1981.
- [30] E. Samuel-Cahn. Comparisons of threshold stop rule and maximum for independent nonnegative random variables. *Ann. Probab.*, 12(4):1213-1216, 1984.

A Supplements to Section 2.1.1

In this section we provide the details of the rigorous measure-theoretic construction for finite random order processes, so as to extend [8, Theorem 3.2] to them.

Recall that S_{n+1} denotes the set of permutations of $[n+1]$ and $\mathbb{R}_+^{n+1} := \{x \in \mathbb{R}^{n+1} : x_i \geq 0, \forall 1 \leq i \leq n+1\}$. We denote by $2^{S_{n+1}}$ the σ -algebra of subsets of S_{n+1} and by $\mathcal{B}(\mathbb{R}_+^{n+1})$ the Borel σ -algebra on \mathbb{R}_+^{n+1} . Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space supporting jointly π and $X = (V_1, \dots, V_{n+1})$, with (π, X) valued in the product measure space

$$(S_{n+1} \times \mathbb{R}_+^{n+1}, 2^{S_{n+1}} \otimes \mathcal{B}(\mathbb{R}_+^{n+1}), \mu \otimes \nu),$$

where:

- for every $\pi \in S_{n+1}$, $\mu(\pi) \equiv \frac{1}{(n+1)!}$ is the uniform probability law on the permutations;
- ν is the law of X , that is the joint law of $\{V_1, \dots, V_{n+1}\}$;
- \otimes denotes the standard product measure and σ -algebra.

Note that a uniform random permutation can be seen as a stochastic process $\pi = (\pi_1, \dots, \pi_n)$. This is a very natural thing to do, since the uniform law on S_{n+1} can be equivalently derived from the conditional laws

$$\begin{aligned} \pi_1 &\sim \text{Unif}([n+1]). \\ (\pi_2 | \pi_1) &\sim \text{Unif}([n+1] \setminus \{\pi_1\}), \\ &\vdots \\ (\pi_n | \pi_{n-1}, \dots, \pi_1) &\sim \text{Unif}([n+1] \setminus \{\pi_1, \dots, \pi_{n-1}\}), \\ (\pi_{n+1} | \pi_n, \dots, \pi_1) &\sim \delta_{[n+1] \setminus \{\pi_1, \dots, \pi_n\}}, \end{aligned}$$

where $\text{Unif}(\cdot)$ denotes the discrete uniform distribution and δ_x is the Dirac distribution centered at x . As a finite stochastic process, π is the measurable map

$$\begin{aligned} \pi : \Omega \times [n+1] &\longrightarrow [n+1] \\ (\omega, i) &\mapsto \pi_i(\omega) \end{aligned}$$

and can be composed with the vector $X = (V_1, \dots, V_{n+1})$, which can also be seen as a finite stochastic process with the joint laws previously given, that is a measurable map

$$\begin{aligned} X : \Omega \times [n+1] &\longrightarrow \mathbb{R}_+ \\ (\omega, i) &\mapsto V_i(\omega). \end{aligned}$$

The composition needs to be done by exploiting the extended *graph map* associated with π , which is measurable and will be denoted as

$$\begin{aligned} \tilde{\pi} : \Omega \times [n+1] &\longrightarrow \Omega \times [n+1] \\ (\omega, i) &\mapsto (\omega, \pi_i(\omega)). \end{aligned}$$

Due to the measurability of all maps involved (which clearly follows from the time index set being discrete), the result of composing X and $\tilde{\pi}$ is the randomly permuted finite stochastic process

$X^\pi := X \circ \tilde{\pi}$ and corresponds, by going back to the usual random vector point of view, to the randomly permuted random vector of components $X_i^\pi = V_{\pi_i}$ for all $i \in [n+1]$.

For every $B \in \mathcal{B}(\mathbb{R}_+)$, $V_{\pi_k} \in B$ if and only if for every $k \in [n+1]$, $\pi_k = i$ and $V_i \in B$. Therefore

$$\{V_{\pi_k} \in B\} = \bigcap_{i=1}^{n+1} (\{\pi_k = i\} \cap \{V_i \in B\}).$$

Equivalently

$$V_{\pi_k}^{-1}(B) = \bigcap_{i=1}^{n+1} (\pi_k^{-1}(i) \cap V_k^{-1}(B)),$$

and we can thus define the σ -algebra generated by V_{π_k} as

$$\sigma(V_{\pi_k}) := \sigma(\{F \in \mathcal{F} : F = V_{\pi_k}^{-1}(B), B \in \mathcal{B}(\mathbb{R}_+)\}).$$

A filtration that accounts for the random order arrival of a general sequence of random variables V_1, \dots, V_{n+1} with undisclosed RO can be constructed by considering \mathcal{F}_k^{n+1} (which will be denoted simply as \mathcal{F}_k , by omitting the finite horizon) defined as follows: $\mathcal{F}_0 := \{\emptyset, \Omega\}$, and for all $k \in [n+1]$, $\mathcal{F}_k := \sigma(V_{\pi_1}, \dots, V_{\pi_k})$. In disclosed RO one would have to interleave with π , that is $\mathcal{F}_k := \sigma(V_{\pi_1}, \pi_1, \dots, V_{\pi_k}, \pi_k)$. Since, due to the features of Instance [H](#), it will be enough to work with undisclosed RO, we adopt the simpler filtration.

Given a stopping time τ , that is a random variable $\tau : \Omega \rightarrow [n+1]$ such that $\{\tau = k\} \in \mathcal{F}_k$, we can also define the stopped version of the process X^π . We simply consider first the stopped process $\pi^\tau := \{\pi_{i \wedge \tau}, i \in [n+1]\}$ and then define $X^{\pi^\tau} := X \circ \tilde{\pi}^\tau$, where $\tilde{\pi}^\tau : (\omega, i) \mapsto (\omega, \pi_{i \wedge \tau}(\omega))$, and therefore $X_i^{\pi^\tau} = V_{\pi_{i \wedge \tau}}$, so that the stopped values of X^{π^τ} is V_{π_τ} . We will abuse the notation and simply refer to X^π as X . Thus $X_i := X_i^\pi = V_{\pi_i}$ for all $i \in [n+1]$ and the conditional expectation on the filtration $\mathbb{E}_{\mathcal{F}_k}(\cdot)$ can be equivalently denoted as $\mathbb{E}(\cdot | X_1, \dots, X_k)$. To construct the optimal stopping rule we define

$$\gamma_{n+1} := V_{\pi_{n+1}}$$

and, for all $l \in [n]$, we define

$$\gamma_l := \max\{V_{\pi_l}, \mathbb{E}_{\mathcal{F}_l} \gamma_{l+1}\}.$$

We also define, for each $k \in [n+1]$,

$$s_k := \inf\{l \geq k : V_{\pi_l} = \gamma_l\}.$$

Then we have the following result.

Theorem A.1 (Extension of [\[8, Theorem 3.2\]](#)). *For a given instance V_1, V_2, \dots, V_{n+1} denote the random arrival order process $X := (V_{\pi_1}, \dots, V_{\pi_{n+1}})$, and consider the backward induction values $\gamma_{n+1}, \gamma_n, \dots, \gamma_1$ and stopping times s_{n+1}, s_n, \dots, s_1 as previously defined. Then for all $k \in [n+1]$, $s_k \in C_k$ and*

$$\mathbb{E}_{\mathcal{F}_k} V_{\pi_{s_k}} = \gamma_k \geq \mathbb{E}_{\mathcal{F}_k} V_{\pi_\tau}$$

for all stopping rules $\tau \in C_k$. Taking expectations yields

$$\mathbb{E} V_{\pi_{s_k}} = \mathbb{E} \gamma_k \geq \mathbb{E} V_{\pi_\tau}$$

for all stopping rules $\tau \in C_k$. Thus $\mathbb{E} \gamma_k = \sup_{\tau \in C_k} \mathbb{E} V_{\pi_\tau}$.

B Supplements to Section 2.1.2

In this section we describe a stopping rule for the discrete setting more explicitly, slightly adapting [16, §7.1] to random order processes, in order to show how to compute the reward of the optimal stopping rule directly, and provide evidence for the claim that it is not a feasible option in our setting. We also comment on the fact that there is no loss of generality in working only with nonrandomised stopping rules, such as those yielded by backward induction.

Recall from Appendix A that the distribution of the random variable X_i , denoting the i -th component of the random order process $X = (V_{\pi_1}, \dots, V_{\pi_{n+1}})$, is known jointly with the distribution of π_i , since by our model, regardless of whether π_i is disclosed or not,

$$(X_i | \pi_i = \sigma_i) \sim V_{\sigma_i},$$

denoting the values taken by X_i as x_i and those taken by π_i as σ_i .

A (possibly randomised) stopping rule for a sequence of deterministic reward functions $\{y_1(x_1), \dots, y_{n+1}(x_1, \dots, x_{n+1})\}$ can be described as a sequence $\psi = (\psi_1(x_1), \dots, \psi_{n+1}(x_1, \dots, x_{n+1}))$ where $\psi_i = \psi_i(x_1, \dots, x_i)$ is the probability of stopping at step i given that i observations have been taken, namely $X_1 = x_1, \dots, X_i = x_i$ (for non-randomised stopping rules $\psi_i \in \{0, 1\}$ for all $i \in [n+1]$, which is the case of the optimal stopping rule constructed in Theorem 2.1). The stopping rule ψ and the observations of X up to time i only, determine the stopping time $0 \leq \tau \leq n+1$. More formally the conditional probability mass function of τ given $X = x$ can be denoted as $\psi = \{\psi_1, \dots, \psi_{n+1}\}$ where

$$\psi_i(x_1, \dots, x_i) = \mathbb{P}(\tau = i | X_1 = x_1, \dots, X_i = x_i) = \mathbb{P}(\tau = i | X = x).$$

Note that the last equality encodes conditional independence of $\{\tau = i\}$, given the past up to the present, from future observations X_{i+1}, \dots, X_{n+1} , ensuring that it is a stopping time (the randomised case can be reduced to a stopping time by enlarging the probability space, as per the concluding remark of this section). By the assumptions of the model, we also have that

$$\begin{aligned} \psi_i(x_1, \dots, x_i) &= \mathbb{P}(\tau = i | \pi_1 = \sigma_1, X_1 = x_1, \dots, \pi_i = \sigma_i, X_i = x_i) \\ &= \mathbb{P}(\tau = i | \pi = \sigma, X = x) = \mathbb{P}(\tau = i | X_1 = x_1, \dots, X_i = x_i), \end{aligned}$$

since the decision to stop is reached while unaware of the arrival order of the distributions in the past, that is, more precisely, the stopping rule, conditionally on the observed values, is independent of the permutation (the last equality need not hold in general in disclosed RO). The conditional probability mass function ψ determines the law of the stopping time τ . Within this framework, optimal stopping is choosing a stopping rule ψ determining a stopping time T that yields maximal expectation of the stopped reward sequence $\mathbb{E} y_T := \mathbb{E} V_{\pi_T}$. Since in our case $y_i(x_1, \dots, x_i) := x_i$, this expectation is calculated by exploiting $y_T = X_T = \sum_{i=1}^{n+1} X_i \mathbb{1}_{\{T=i\}}$, yielding

$$\begin{aligned} \mathbb{E} X_T &= \mathbb{E}_{\pi, X} \mathbb{E}(X_T | \pi, X) = \mathbb{E}_{\pi, X} \sum_{i=1}^{n+1} X_i \mathbb{P}(T = i | \pi, X) = \mathbb{E}_{\pi, X} \sum_{i=1}^{n+1} X_i \psi_i(X_1, \dots, X_i) \\ &:= \mathbb{E} \sum_{i=1}^{n+1} V_{\pi_i} \psi_i(V_{\pi_1}, \dots, V_{\pi_i}), \end{aligned}$$

where the abuse of notation $\mathbb{E}_{\pi, X}(\cdot)$ does not stand for conditioning, but for the joint laws with respect to which integration is carried, upon change of variables. Such notation is no longer

necessary in the last step, where all sources of randomness are explicitly stated in the integrands' notation. To clarify, all the measures being discrete, turns all integrations into summations:

$$\begin{aligned}
& \mathbb{E}_{\pi, X} \sum_{i=1}^{n+1} X_i \mathbb{P}(T = i | \pi, X) \\
&= \sum_{\sigma \in S_{n+1}} \sum_{x \in \mathcal{S}_n} \sum_{i=1}^{n+1} x_i \mathbb{P}(T = i | \pi = \sigma, X = x) \mathbb{P}(\pi = \sigma, X = x) \\
&= \sum_{\sigma \in S_{n+1}} \sum_{x \in \mathcal{S}_n} \sum_{i=1}^{n+1} x_i \mathbb{P}(T = i | X = x) \mathbb{P}(X = x | \pi = \sigma) \mathbb{P}(\pi = \sigma) \\
&= \sum_{\sigma \in S_{n+1}} \frac{1}{(n+1)!} \sum_{x \in \mathcal{S}_n} \mathbb{P}(V_{\sigma_1} = x_1) \dots \mathbb{P}(V_{\sigma_{n+1}} = x_{n+1}) \sum_{i=1}^{n+1} x_i \psi_i(x_1, \dots, x_i).
\end{aligned}$$

We will not use the abuse of notation aforementioned outside this section.

We conclude this section with a comment on the optimal stopping rule characterised in Theorem 2.1: backward induction yields a non-randomised optimal stopping rule T , but this comes with no loss of generality, because for any randomised stopping rule there is a non-randomised equivalent one (in expectation), and it is therefore sufficient to work with an optimal non-randomised stopping rule. To see the equivalence in our case, start with a possibly randomised stopping rule ψ . An equivalent non-randomised stopping rule is then yielded by enlarging the product space with $[0, 1]^{n+1}$ endowing this factor with the product uniform measure on the unit interval, and considering the interleave process $(\pi, X, U) := \{\pi_1, X_1, U_1, \dots, \pi_{n+1}, X_{n+1}, U_{n+1}\}$ with $\{U_i\}$ being iid copies of $U \sim \text{Unif}[0, 1]$. With respect to the filtration incorporating the uniform random variables interleaved, that is $\mathcal{F}_k = \sigma(V_{\pi_1}, U_1, \dots, V_{\pi_k}, U_k)$, the randomized stopping rule yields a stopping time τ . Define the nonrandomised stopping rule $\tilde{\psi}$ as

$$\tilde{\psi}_i(X_1, U_1, \dots, X_i, U_i) := \mathbb{1}_{\{U_i < \psi_i(X_1, \dots, X_i)\}},$$

maintaining the same reward sequence. By exploiting the independence of the uniform random variables introduced, the two stopping rules are equivalent, since if we denote by $\tilde{\tau}$ the stopping time yielded by $\tilde{\psi}$, then by the independence of U_i it follows that

$$\begin{aligned}
\mathbb{E} V_{\pi_{\tilde{\tau}}} &= \mathbb{E}_{\pi, X, U} \sum_{i=1}^{n+1} X_i \tilde{\psi}_i(X_1, U_1, \dots, X_i, U_i) = \mathbb{E}_{\pi, X} \sum_{i=1}^{n+1} X_i \mathbb{E}_U \mathbb{1}_{\{U_i < \psi_i(X_1, \dots, X_i)\}} \\
&= \mathbb{E}_{\pi, X} \sum_{i=1}^{n+1} X_i \psi_i(X_1, \dots, X_i) = \mathbb{E} V_{\pi_{\tau}}.
\end{aligned}$$

C Asymptotic analysis of the acceptance times

In this section we prove the asymptotic results concerning the acceptance times of the optimal algorithm.

Proof of Lemma 2.1.

a) Recall that j_n is the smallest i such that $a \geq \phi_i$, as per Definition 2.1. Since for all n large enough, $j_n \in [n]$, we consider $i \in [n]$, and in order to derive the sharp asymptotic estimate of the claim, we start by calculating explicitly ϕ_i by deriving $\phi_n, \dots, \phi_{i+1}$ via backward induction.

Step 1. Note that since $a < b$, for all $k \geq j_n$, $\phi_k < b$. In fact by Remark 2.5, $\phi_{k-1} = \mathbb{E}(V \vee \phi_k) \geq \phi_k$ for all $k \in [n]$ and $\phi_{j_n} < a < b$. By Remark 2.5 for $k = n$, as $n \rightarrow \infty$,

$$\phi_n = \frac{1+bp}{n}, \quad (\text{C.1})$$

and for all $i+1 \leq k < n+1$ we will iterate (C.1) through Remarks 2.1 and 2.5, that is through

$$\phi_{k-1} = \mathbb{E}(V \vee \phi_k) = \frac{1}{n} + \frac{p(b \vee \phi_k)}{n} + \left(1 - \frac{p}{n} - \frac{1}{n^2}\right) \phi_k = \frac{1+pb}{n} + \left(1 - \frac{p}{n} - \frac{1}{n^2}\right) \phi_k, \quad (\text{C.2})$$

which follows by expanding the expectation of the maximum as in Remark 2.1 and using $0 < \phi_k \leq n$ as observed in Remark 2.2, along with the aforementioned fact, that for all $i+1 \leq k < n+1$, $b \vee \phi_k = b$. One more iteration will suffice to clarify what the induction hypothesis should be. Consider $k = n$ in (C.2), then we have that

$$\phi_{n-1} = \frac{1+pb}{n} + \left(1 - \frac{p}{n} - \frac{1}{n^2}\right) \frac{1+pb}{n} = \frac{1+pb}{n} \left[1 + \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)\right].$$

The induction hypothesis is therefore that for $i+1 \leq k < n+1$ and n large enough

$$\phi_k = \frac{1+bp}{n} \sum_{j=0}^{n-k} \left(1 - \frac{p}{n} - \frac{1}{n}\right)^j. \quad (\text{C.3})$$

By assuming (C.3) and using (C.2), it follows that

$$\phi_{k-1} = \frac{1+pb}{n} + \left(1 - \frac{p}{n} - \frac{1}{n^2}\right) \left[\frac{1+bp}{n} \sum_{j=0}^{n-k} \left(1 - \frac{p}{n} - \frac{1}{n}\right)^j \right] = \frac{1+bp}{n} \sum_{j=0}^{n-k+1} \left(1 - \frac{p}{n} - \frac{1}{n}\right)^j,$$

and the induction step is complete.

Step 2. Having shown (C.3) for all $i \leq k \leq n$, we take $k = i$, so as to obtain

$$\phi_i = \frac{1+bp}{n} \sum_{j=0}^{n-i} \left(1 - \frac{p}{n} - \frac{1}{n}\right)^j = \frac{1+bp}{n} \left[\frac{1 - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{n-i+1}}{\frac{p}{n} + \frac{1}{n^2}} \right]. \quad (\text{C.4})$$

Thus

$$\phi_i = \frac{1+bp}{p} \left[1 - \left(1 - \frac{p}{n}\right)^{n-i+1} \right] + \mathcal{O}\left(\frac{1}{n}\right), \quad (\text{C.5})$$

having used

$$\frac{\frac{1+bp}{n}}{\frac{p}{n} + \frac{1}{n^2}} = \frac{1+bp}{p + \frac{1}{n}} = \frac{1+bp}{p} \left(1 + \mathcal{O}\left(\frac{1}{n}\right) \right) \quad (\text{C.6})$$

and, upon factorising

$$1 - \frac{p}{n} - \frac{1}{n^2} = \left(1 - \frac{p}{n}\right) \left(1 - \frac{1 + \mathcal{O}(1)}{n^2}\right), \quad (\text{C.7})$$

having used

$$\left(1 - \frac{1 + \mathcal{O}(1)}{n^2}\right)^{n-i+1} = e^{-(n-i+1)\frac{1+\mathcal{O}(1)}{n^2} + \mathcal{O}\left(\frac{1}{n^3}\right)} = 1 + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{C.8})$$

Plugging (C.5) into Definition 2.1, we observe that since j_n is the smallest i such that $a \geq \phi_i$, one obtains equivalently that j_n is the smallest $i \in [n]$ such that

$$a \geq \frac{1+bp}{p} \left[1 - \left(1 - \frac{p}{n}\right)^{n-i+1} \right] + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{C.9})$$

Step 3. We rearrange (C.9) into

$$i \geq n + 1 - \frac{\log\left(1 - \frac{pa}{1+bp} + \mathcal{O}\left(\frac{1}{n}\right)\right)}{\log\left(1 - \frac{p}{n}\right)} \quad (\text{C.10})$$

and then use the Taylor expansion of the logarithms as $n \rightarrow \infty$, yielding

$$\begin{aligned} \frac{\log\left(1 - \frac{pa}{1+bp} + \mathcal{O}\left(\frac{1}{n}\right)\right)}{\log\left(1 - \frac{p}{n}\right)} &= \frac{\log\left(1 - \frac{pa}{1+bp} + \mathcal{O}\left(\frac{1}{n}\right)\right)}{-\frac{p}{n} + \mathcal{O}\left(\frac{1}{n^2}\right)} = -\frac{n \log\left(\frac{1+(b-a)p}{1+bp}\right) + \mathcal{O}\left(\frac{1}{n}\right)}{1 + \mathcal{O}\left(\frac{1}{n}\right)} \\ &= -\frac{n}{p} \log\left(\frac{1+(b-a)p}{1+bp}\right) + \mathcal{O}(1). \end{aligned}$$

Plugging the expansion into (C.10) yields

$$i \geq n + 1 + \frac{n}{p} \log\left(\frac{1+(b-a)p}{1+bp}\right) + \mathcal{O}(1),$$

from which it follows that j_n is the smallest $i \in [n]$ such that, as $n \rightarrow \infty$,

$$i \geq n \left[1 + \frac{1}{p} \log\left(\frac{1+(b-a)p}{1+bp}\right) \right] + \mathcal{O}(1).$$

Thus by the standard asymptotics of the ceiling function, as $n \rightarrow \infty$, we have that

$$j_n = \left\lceil n \left[1 + \frac{1}{p} \log\left(\frac{1+(b-a)p}{1+bp}\right) \right] + \mathcal{O}(1) \right\rceil \sim n \left[1 + \frac{1}{p} \log\left(\frac{1+(b-a)p}{1+bp}\right) \right].$$

Note that since $0 < a < 1 < b$, the coefficient of n is positive by the inequality

$$\log(1+x) \geq \frac{x}{1+x}$$

for all $x > -1$, which implies, by taking

$$x = -\frac{ap}{1+bp},$$

that

$$\log\left(\frac{1+(b-a)p}{1+bp}\right) = \log\left(1 - \frac{ap}{1+bp}\right) > \frac{-\frac{ap}{1+bp}}{1 - \frac{ap}{1+bp}} = -\frac{ap}{1+(b-a)p} > -p,$$

yielding the claim.

b) Replacing a with b in (C.9) shows that, by Definition 2.2, k_n is the smallest $2 \leq i \leq n$ such that, as $n \rightarrow \infty$,

$$i \geq n \left[1 + \frac{1}{p} \log\left(\frac{1}{1+bp}\right) \right] + \mathcal{O}(1),$$

which yields Lemma 2.1 (b) through a similar concluding argument as that of Lemma 2.1 (a). Note that the coefficient of n is positive by the condition $\log(1+pb) < p$ on the parameters of Instance H.

c) Noting that for Instance H we have $b > a > 0$ and $p > 0$, it follows that $1 < 1+(b-a)p$, and therefore Lemma 2.1 (a, b) immediately imply that eventually $k_n \leq j_n$, since

$$1 + \frac{1}{p} \log \frac{1}{1+bp} < 1 + \frac{1}{p} \log \frac{1+(b-a)p}{1+bp}.$$

d) To show that $\bar{k}_n \geq k_n$ as $n \rightarrow \infty$, assume by contradiction that there exists a subsequence $\{n_l\}$ such that $\bar{k}_{n_l} < k_{n_l}$. For simplicity, we relabel the indices with n . Thus our hypothesis is that for infinitely many n , $\bar{k}_n < k_n$. By Definition 2.3, \bar{k}_n is the smallest k such that $\bar{\phi}_k \leq b$, so it follows that for all n considered,

$$\bar{\phi}_{\bar{k}_n-1} \leq b. \quad (\text{C.11})$$

Our strategy will be to derive a contradiction with (C.11) thanks to an iterative lower bound on $\bar{\phi}_{\bar{k}_n-1}$.

Step 1. First of all note that by Remark 2.3, also for all $k \geq k_n$, $\bar{\phi}_k \leq b$. In this first step we exploit this fact to derive a lower bound on $\bar{\phi}_k$ for all $k \geq k_n - 1$. By Remarks 2.1 and 2.5 and the facts aforementioned, we have that

$$\begin{aligned} \bar{\phi}_k &\geq \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left(\frac{1}{n} + p \frac{b \vee \bar{\phi}_{k+1}}{n} + \bar{\phi}_{k+1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)\right) \\ &= \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left(\frac{1+bp}{n} + \bar{\phi}_{k+1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)\right) \end{aligned} \quad (\text{C.12})$$

Then by (C.12) applied with $k = n - 1$, using $\bar{\phi}_n = a$, we derive

$$\bar{\phi}_{n-1} \geq a + \frac{1}{2} \left[\frac{1+(b-a)p}{n} - \frac{a}{n^2} \right]. \quad (\text{C.13})$$

Iterating from (C.13) via (C.12) we obtain, by induction, that for all $k \geq k_n - 1$,

$$\bar{\phi}_k = a + \left(\frac{1+(b-a)p}{n} - \frac{a}{n^2} \right) \sum_{j=0}^{n-k-1} \frac{n-k-j}{n+1-k} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j. \quad (\text{C.14})$$

In fact if (C.14) is true for any $k_n < k < n + 1$, by (C.12) applied to $k - 1$ we obtain

$$\begin{aligned} \bar{\phi}_{k-1} &= \frac{a}{n-k+2} + \left(1 - \frac{1}{n-k+2}\right) \left[\frac{1+pb}{n} + a \left(1 - \frac{p}{n} - \frac{1}{n^2}\right) \right. \\ &\quad \left. + \left(\frac{1+(b-a)p}{n} - \frac{a}{n^2} \right) \sum_{j=0}^{n-k-1} \frac{n-k-j}{n+1-k} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{j+1} \right] \\ &= a + \left(\frac{1+(b-a)p}{n} - \frac{a}{n^2} \right) \frac{n-k+1}{n-k+2} \left[1 + \sum_{j=1}^{n-k} \frac{n-k-(j-1)}{n-(k-1)} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \right] \\ &= a + \left(\frac{1+(b-a)p}{n} - \frac{a}{n^2} \right) \sum_{j=0}^{n-k} \frac{n-(k-1)-j}{n+1-(k-1)} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j. \end{aligned}$$

Step 2. In this step we derive a sharp asymptotic estimate for (C.14). Let $q = q(n, p) := 1 - p/n - 1/n^2$. The summation term can be rewritten as

$$\begin{aligned} S_{k,n} &:= \sum_{j=0}^{n-k-1} \frac{n+1-k-(j+1)}{n+1-k} q^j = \sum_{j=0}^{n-k-1} q^j - \frac{1}{n+1-k} \sum_{j=0}^{n-k-1} (j+1)q^j = \\ &= \frac{1-q^{n-k}}{1-q} - \frac{1}{n+1-k} \frac{d}{dq} \left(\sum_{j=0}^{n-k} q^j \right) = \frac{1-q^{n-k}}{1-q} - \frac{1}{n+1-k} \frac{d}{dq} \left(\frac{1-q^{n-k+1}}{1-q} \right) = \\ &= \frac{1-q^{n-k}}{1-q} - \frac{1}{n+1-k} \frac{1-q^{n-k}[(n-k)(1-q)+1]}{(1-q)^2}, \end{aligned}$$

so by factorising q as in (C.7), and exploiting

$$\left(1 - \frac{p}{n}\right)^n = e^{-p} + \mathcal{O}\left(\frac{1}{n}\right), \quad (\text{C.15})$$

we have that for all $k \geq k_n - 1$,

$$\begin{aligned} S_{k,n} &= \frac{1 - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{n-k}}{\frac{p}{n} + \frac{1}{n^2}} - \frac{1}{n+1-k} \frac{1 - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{n-k} \left[(n-k) \left(\frac{p}{n} + \frac{1}{n^2}\right) + 1\right]}{\left(\frac{p}{n} + \frac{1}{n^2}\right)^2} \\ &= \frac{n}{p} \left[1 - e^{-p\left(1 - \frac{k}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right)\right] - \frac{n}{p^2 \left(1 - \frac{k}{n}\right)} \left\{1 - e^{-p\left(1 - \frac{k}{n}\right)} \left[1 + p \left(1 - \frac{k}{n}\right)\right] + \mathcal{O}\left(\frac{1}{n}\right)\right\} \\ &= \frac{n}{p} \left[1 - \frac{1 - e^{-p\left(1 - \frac{k}{n}\right)}}{p \left(1 - \frac{k}{n}\right)} + \mathcal{O}\left(\frac{1}{n \left(1 - \frac{k}{n}\right)}\right)\right]. \end{aligned}$$

Recall that by Lemma 2.1 (b),

$$\frac{k_n}{n} \rightarrow 1 - \frac{\log(1+bp)}{p}.$$

For $k = k_n - 1$, the error term

$$\mathcal{O}\left(\frac{1}{n \left(1 - \frac{k}{n}\right)}\right) = \mathcal{O}\left(\frac{1}{n}\right).$$

Step 3. We plug $S_{k_n-1,n}$ into (C.14), for $k = k_n$, and we obtain

$$\begin{aligned} \bar{\phi}_{k_n-1} &\geq a + \left(\frac{1 + (b-a)p}{p} - \frac{a}{pn}\right) \left[1 - \frac{1 - e^{-p\left(1 - \frac{k_n-1}{n}\right)}}{p \left(1 - \frac{k_n-1}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right)\right] \\ &= a + \frac{1 + (b-a)p}{p} - \frac{1 + (b-a)p}{p} \frac{1 - e^{-p\left(1 - \frac{k_n}{n}\right)}}{p \left(1 - \frac{k_n}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) \\ &= b + \frac{1}{p} - \frac{1 + (b-a)p}{p} \frac{1 - e^{-p\left(1 - \frac{k_n}{n}\right)}}{p \left(1 - \frac{k_n}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right). \end{aligned}$$

Applying Lemma 2.1 (b) yields

$$\begin{aligned} \bar{\phi}_{k_n-1} &\geq b + \frac{1}{p} - \frac{1 + (b-a)p}{p} \frac{1 - e^{-p\left(1 - \frac{k_n}{n}\right)}}{p \left(1 - \frac{k_n}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) \\ &= b + \frac{1}{p} - \frac{1 + (b-a)p}{p} \frac{1 - \frac{1}{1+bp}}{\log(1+pb)} + \mathcal{O}(1) = b + \frac{1}{p} - \frac{b[1 + (b-a)p]}{(1+pb) \log(1+pb)} + \mathcal{O}(1). \end{aligned}$$

By Condition V, which ensures that

$$\frac{1}{p} - \frac{b[1 + (b-a)p]}{(1+pb) \log(1+pb)} > 0,$$

and n being arbitrarily large, we have that $\bar{\phi}_{k_n-1} > b$, which contradicts (C.11). The assumption that there are infinitely many n such that $\bar{k}_n < k_n$ is therefore false, meaning that for all n large enough, $k_n \leq \bar{k}_n$.

e) To show that $\bar{k}_n \leq j_n$ as $n \rightarrow \infty$, assume by contradiction that there exists a subsequence $\{n_l\}$ such that $\bar{k}_{n_l} > j_{n_l}$ as $l \rightarrow \infty$. For simplicity, relabel n_l as n , thus starting the argument, without loss of generality, with the assumption by contradiction, that $\bar{k}_n > j_n$ for infinitely many n . The overall strategy will be the following: under the assumption by contradiction we derive an upper bound on \bar{k}_n , which we recall to be the smallest integer k such that $b \geq \bar{\phi}_k$ as per Definition 2.3. Thanks to this upper bound we will show that $\bar{k}_n/j_n < 1$, contradicting the assumption that $\bar{k}_n/j_n > 1$. To obtain such an upper bound on \bar{k}_n we will find first a suitable upper bound on $\bar{\phi}_{\bar{k}_n}$.

Step 1. Since for all $\bar{k}_n \leq k \leq n$, we have that $k \geq j_n$, not only we know that $\bar{\phi}_k \leq b$, but we also know that $\phi_k \leq a$, and therefore by Remarks 2.1 and 2.5 we obtain that

$$\bar{\phi}_k \leq \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left[\frac{1+bp}{n} + \bar{\phi}_{k+1} \left(1 - \frac{p}{n}\right) \right]. \quad (\text{C.16})$$

Recall that $\bar{\phi}_n = a$. Then by (C.16) it follows that

$$\bar{\phi}_{n-1} \leq \frac{a}{2} + \frac{1}{2} \left[\frac{1+bp}{n} + a \left(1 - \frac{p}{n}\right) \right] = a + \frac{1+(b-a)p}{2n}. \quad (\text{C.17})$$

The induction hypothesis will be that for any $\bar{k}_n \leq k \leq n-1$,

$$\bar{\phi}_{k+1} \leq a + \frac{n-k-1}{2} \frac{1+(b-a)p}{n}. \quad (\text{C.18})$$

Then by (C.16) and (C.18) it follows that

$$\begin{aligned} \bar{\phi}_k &\leq \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left[\frac{1+bp}{n} + \left(a + \frac{n-k-1}{2} \frac{1+(b-a)p}{n}\right) \left(1 - \frac{p}{n}\right) \right] \\ &= a + \left(1 - \frac{1}{n-k+1}\right) \left[\frac{n-k+1}{2} \frac{1+(b-a)p}{n} - \frac{p}{n} \frac{n-k-1}{2} \frac{1+(b-a)p}{n} \right] \\ &\leq a + \left(1 - \frac{1}{n-k+1}\right) \frac{n-k+1}{2} \frac{1+(b-a)p}{n} = a + \frac{n-k}{2} \frac{1+(b-a)p}{n}. \end{aligned}$$

By induction on k we obtain that

$$\bar{\phi}_{\bar{k}_n} \leq a + \frac{n-\bar{k}_n}{2} \frac{1+(b-a)p}{n}. \quad (\text{C.19})$$

By a trivial induction argument one can iterate this bound until time j_n ; showing the first step will suffice. Since for all $j_n \leq k < \bar{k}_n$, we have that $b \vee \bar{\phi}_k = \bar{\phi}_k$ and $a \vee \phi_k = a$, by Remarks 2.1 and 2.5 it follows that

$$\begin{aligned} \bar{\phi}_k &\leq \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left[\frac{1}{n} + \frac{p}{n} \bar{\phi}_{k+1} + \bar{\phi}_{k+1} \left(1 - \frac{p}{n}\right) \right] \\ &= \frac{a}{n-k+1} + \left(1 - \frac{1}{n-k+1}\right) \left[\frac{1}{n} + \bar{\phi}_{k+1} \right]. \end{aligned} \quad (\text{C.20})$$

In the induction steps past time \bar{k}_n , (C.20) will take the place of (C.16). We show the first step. By (C.19) and (C.20) and the fact that

$$\frac{1}{n} < \frac{1+(b-a)p}{n},$$

we have that

$$\begin{aligned}\bar{\phi}_{\bar{k}_n-1} &\leq \frac{a}{n - \bar{k}_n + 2} + \left(1 - \frac{1}{n - \bar{k}_n + 2}\right) \left[\frac{1}{n} + a + \frac{n - \bar{k}_n}{2} \frac{1 + (b - a)p}{n}\right] \\ &\leq a + \left(1 - \frac{1}{n - \bar{k}_n + 2}\right) \frac{n - \bar{k}_n + 2}{2} \frac{1 + (b - a)p}{n} = a + \frac{n - \bar{k}_n + 1}{2} \frac{1 + (b - a)p}{n},\end{aligned}$$

The mechanism of this iteration is trivial, due to the cancellation of the fractions carrying over for all k , and therefore what we obtained for the previous steps can be iterated by induction for all successive steps. Thus we have shown that for all $j_n \leq k \leq n - 1$,

$$\bar{\phi}_k \leq a + \frac{n - k}{2} \frac{1 + (b - a)p}{n}. \quad (\text{C.21})$$

Step 2. Consider

$$k_n^* := \inf \left\{ k \geq j_n : b \geq a + \frac{n - k}{2} \frac{1 + (b - a)p}{n} \right\}.$$

Equivalently, k_n^* is the smallest $k \geq j_n$ such that

$$k \geq n \frac{1 - (2 - p)(b - a)}{1 + p(b - a)}.$$

Note that $k_n^* \leq n$ due to $b > a$, and $k_n^* > 0$ due to Condition II. Then, by a similar reasoning as in the conclusion of Lemma 2.1 (a),

$$k_n^* = \left\lceil n \frac{1 - (2 - p)(b - a)}{1 + p(b - a)} \right\rceil \sim n \frac{1 - (2 - p)(b - a)}{1 + p(b - a)}.$$

Note that

$$\bar{k}_n \leq k_n^*, \quad (\text{C.22})$$

since, by (C.21), the earliest $k \geq j_n$ such that $b \geq \bar{\phi}_k$ is smaller than the earliest $k \geq j_n$ such that

$$b \geq a + \frac{n - k}{2} \frac{1 + (b - a)p}{n}.$$

Step 3. Recall that we are assuming that $j_n < \bar{k}_n$. Equivalently, we have that

$$1 + \frac{1}{j_n} \leq \frac{\bar{k}_n}{j_n}.$$

By (C.22), Condition III, and Lemma 2.1 (a), we reach the following contradiction: as $n \rightarrow \infty$,

$$\frac{\bar{k}_n}{j_n} \geq 1 + \frac{1}{j_n} \rightarrow 1 + \frac{1}{1 + \frac{1}{p} \log \frac{1 + p(b - a)}{1 + pb}} > 1$$

and

$$\frac{\bar{k}_n}{j_n} \leq \frac{k_n^*}{j_n} \rightarrow \frac{\frac{1 - (2 - p)(b - a)}{1 + p(b - a)}}{1 + \frac{1}{p} \log \frac{1 + p(b - a)}{1 + pb}} < 1.$$

Thus it must hold that for all n large enough, $\bar{k}_n \leq j_n$. □

D Expectation of the prophet

In this section we compute exact asymptotics for the expectation of the maximum for Instance H.

Proof of Lemma 3.1. Since by assumption $b > a$, for all n large enough

$$\max_{i \in [n+1]} V_i \sim \begin{cases} n, & \text{w.p. } 1 - \left(1 - \frac{1}{n^2}\right)^n, \\ b, & \text{w.p. } \left(1 - \frac{1}{n^2}\right)^n - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^n, \\ a, & \text{w.p. } \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^n. \end{cases}$$

Thus

$$\mathbb{E} \max_{i \in [n+1]} V_i = n \left[1 - \left(1 - \frac{1}{n^2}\right)^n\right] + b \left[\left(1 - \frac{1}{n^2}\right)^n - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^n\right] + a \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^n.$$

By a similar estimate as in (C.8), factorising as in (C.7), we have that

$$\left(1 - \frac{1}{n^2}\right)^n = 1 - \frac{1}{n} + \mathcal{O}\left(\frac{1}{n^3}\right),$$

so that by exploiting (C.15) it follows that

$$\mathbb{E} \max_{i \in [n+1]} V_i = n \left[\frac{1}{n} + \mathcal{O}\left(\frac{1}{n^3}\right)\right] + b \left[1 - \frac{1}{n} + \mathcal{O}\left(\frac{1}{n^3}\right) - e^{-p} + \mathcal{O}\left(\frac{1}{n}\right)\right] + a \left[e^{-p} + \mathcal{O}\left(\frac{1}{n}\right)\right]$$

and the claim follows. \square

E Expectation of the optimal algorithm

In this section we compute a sharp asymptotic estimate for the expectation of the optimal algorithm $\mathbb{E} V_{\pi_T}$. We will do so through the law of total expectation with respect to the partitioning $\{\Omega_1, \dots, \Omega_{n+1}\}$ of the sample space Ω . The role of Lemma 2.1 in this computation can be appreciated from the following. Consider $i < j_n$ for instance: conditionally on Ω_i , the optimal stopping rule does not stop when a or 0 are probed (except for 0 , in the last step), but it stops the first time n is probed, or, at certain times, when b is probed. Thus to determine when b is accepted or not, knowledge of the relative position of \bar{k}_n with respect to k_n and j_n is needed. Thanks to Lemma 2.1 we avoid a lengthy case analysis, since only one ordering is possible: $k_n \leq \bar{k}_n \leq j_n$.

Proof of Proposition 3.1. We start by finding the distribution of $(V_{\pi_T} | \Omega_i)$ for all $i \in [n+1]$ (the abuse of notation denotes the conditional distribution of V_{π_T} given Ω_i), so as to determine, for all n large enough, $\mathbb{E}_i V_{\pi_T}$ for all $i \in [n+1]$. Recall that, conditionally on Ω_i , at every step of the process other than the i th, independently, the algorithm could probe n , b and 0 , with probabilities $1/n^2$, p/n and $1 - p/n - 1/n^2$ respectively. While n is always accepted, 0 is never accepted, until the end, by the optimal algorithm.

- i) If $i < k_n$, conditionally on Ω_i , a is probed before time $k_n \leq j_n$ by Lemma 2.1 (c), so a will be rejected. There are in total n steps, in which n could be probed and accepted. Before time k_n , the optimal algorithm stops with reward n if and only if any of the first $k_n - 2$ values probed is n , because if b were probed before time i , the algorithm would not stop by definition of \bar{k}_n , since $i < k_n \leq \bar{k}_n$ by Lemma 2.1 (d); if b were probed between time i and

k_n , the algorithm would not stop by definition of k_n . Thus the probability of stopping with reward n before time k_n is

$$1 - \left(1 - \frac{1}{n^2}\right)^{k_n-2}.$$

From time k_n onward, at each step, the optimal algorithm stops with reward n if and only if all previous values probed, starting from time k_n , are 0. In fact if after or at time k_n , b were to be probed before n , the algorithm would stop with reward b by definition of $k_n > i$. Thus the probability of stopping with reward n at each step from time k_n onward (there are $n + 2 - k_n$ such steps), is the probability of not probing n in any of the first $k_n - 2$ steps (this guarantees that the algorithm reaches the k_n th step), multiplied by the probability of always probing 0 from time k_n up to the first time n is probed:

$$\frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{k_n-2} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

There are in total n steps, in which b could be probed, but only in $n + 2 - k_n$ of these, from time k_n onward, b would be accepted. From the previous comments on the acceptance of b , at each of these steps, the probability that the optimal algorithm stops with reward b is the probability of not probing n in any of the first $k_n - 2$ steps (this guarantees that the algorithm reaches the k_n th step), multiplied by the probability that all remaining probed values are 0, from time k_n up to the first time b is probed. Thus the probability of stopping with reward b is

$$\frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{k_n-2} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

In conclusion, for every $i < k_n$,

$$(V_{\pi_T} | \Omega_i) \sim \begin{cases} n, & \text{w.p. } 1 - \left(1 - \frac{1}{n^2}\right)^{k_n-2} + \frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{k_n-2} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ b, & \text{w.p. } \frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{k_n-2} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ 0, & \text{otherwise.} \end{cases} \quad (\text{E.1})$$

Since

$$\sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j = \frac{1 - \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{n+2-k_n}}{\frac{p}{n} + \frac{1}{n^2}}$$

and from (C.8) and (C.15) it follows that

$$\left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{k_n-2} = \left(1 - \frac{p}{n}\right)^{k_n} + \mathcal{O}\left(\frac{1}{n}\right) = \left[\left(1 - \frac{p}{n}\right)^n\right]^{\frac{k_n}{n}} + \mathcal{O}\left(\frac{1}{n}\right) = e^{-p\frac{k_n}{n}} + \mathcal{O}\left(\frac{1}{n}\right)$$

and

$$\left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{n+2-k_n} = \frac{e^{p\frac{k_n}{n}}}{e^p} + \mathcal{O}\left(\frac{1}{n}\right) = e^{p\left(\frac{k_n}{n}-1\right)} + \mathcal{O}\left(\frac{1}{n}\right),$$

we conclude that

$$\sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j = \frac{1 - e^{p\left(\frac{k_n}{n}-1\right)} + \mathcal{O}\left(\frac{1}{n}\right)}{\frac{p}{n} + \frac{1}{n^2}} = \frac{n}{p} \left[1 - e^{p\left(\frac{k_n}{n}-1\right)} + \mathcal{O}\left(\frac{1}{n}\right)\right]. \quad (\text{E.2})$$

Thus

$$\frac{p}{n} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j = 1 - e^{p(\frac{k_n}{n}-1)} + \mathcal{O}\left(\frac{1}{n}\right) \quad (\text{E.3})$$

and

$$\frac{1}{n^2} \sum_{j=0}^{n+1-k_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j = \frac{1}{pn} \left(1 - e^{p(\frac{k_n}{n}-1)}\right) + \mathcal{O}\left(\frac{1}{n^2}\right). \quad (\text{E.4})$$

Considering that k_n/n is subunitary and bounded away from 0 and 1 by Lemma 2.1 (b), it also follows that

$$\begin{aligned} \left(1 - \frac{1}{n^2}\right)^{k_n-2} &= \left(1 - \frac{1}{n^2}\right)^{k_n} + \mathcal{O}\left(\frac{1}{n^2}\right) = e^{-\frac{k_n}{n^2} + \mathcal{O}\left(\frac{k_n}{n^4}\right)} + \mathcal{O}\left(\frac{1}{n^2}\right) \\ &= 1 - \frac{k_n}{n^2} + \mathcal{O}\left(\frac{k_n}{n^4}\right) + \mathcal{O}\left[\left(\frac{k_n}{n^2} + \mathcal{O}\left(\frac{k_n}{n^4}\right)\right)^2\right] + \mathcal{O}\left(\frac{1}{n^2}\right) \\ &= 1 - \frac{k_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right). \end{aligned} \quad (\text{E.5})$$

In conclusion plugging (E.3) to (E.5) into (E.1) yields

$$\begin{aligned} \mathbb{E}_i V_{\pi_T} &= \\ &= n \left[\frac{k_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right) \right] + \left[1 - \frac{k_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right) \right] \left[\left(\frac{1}{p} + b \right) \left(1 - e^{p(\frac{k_n}{n}-1)} \right) + \mathcal{O}\left(\frac{1}{n}\right) \right] \\ &= \frac{k_n}{n} + \left(\frac{1}{p} + b \right) \left(1 - e^{p(\frac{k_n}{n}-1)} \right) + \mathcal{O}\left(\frac{1}{n}\right). \end{aligned} \quad (\text{E.6})$$

- ii) If $k_n \leq i < \bar{k}_n$ (assuming that there is any such i ; if not, the empty sum convention used in (E.13) will take care of the case $k_n = \bar{k}_n$) conditionally on Ω_i , a is probed after or at time k_n , strictly before time $\bar{k}_n \leq j_n$ by Lemma 2.1 (d, e), so a will be rejected. There are in total n steps, in which n could be probed and accepted. Before time i , the optimal algorithm stops with reward n if and only if any of the first $i - 1$ values probed is n , because if b were probed before time i , the optimal algorithm would not stop by definition of $\bar{k}_n > i$. Thus the probability of stopping with reward n before time i is

$$1 - \left(1 - \frac{1}{n^2}\right)^{i-1}.$$

From time i onward, at each step, the optimal algorithm stops with reward n if and only if all the previous values probed, starting from time $i + 1$, are 0. In fact if after time $i \geq k_n$, b were to be probed before n , the algorithm would stop with reward b by definition of k_n . Thus the probability of stopping with reward n at each step from time $i + 1$ onward (there are $n + 1 - i$ such steps), is the probability of not probing n in any of the first $i - 1$ steps (which ensures that the algorithm reaches the i th step), multiplied by the probability of always probing 0, from time $i + 1$ up to the first time n is probed:

$$\frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{i-1} \sum_{j=0}^{n-i} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

There are in total n steps, in which b could be probed, but only in $n+1-i$ of these steps, from time $i+1$ onward, b would be accepted, since $k_n \leq i < \bar{k}_n$. From the previous comments on the acceptance of b , at each of these steps, the probability that the optimal algorithm stops with reward b is the probability of not probing n in any of the first $i-1$ steps (which ensures that the algorithm reaches the i th step), multiplied by the probability that all remaining probed values are 0, from time $i+1$ up to the first time b is probed. Thus the probability of stopping with reward b is

$$\frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{i-1} \sum_{j=0}^{n-i} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

In conclusion for every $k_n \leq i < \bar{k}_n$,

$$(V_{\pi_T} | \Omega_i) \sim \begin{cases} n, & \text{w.p. } 1 - \left(1 - \frac{1}{n^2}\right)^{i-1} + \frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{i-1} \sum_{j=0}^{n-i} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ b, & \text{w.p. } \frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{i-1} \sum_{j=0}^{n-i} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ 0, & \text{otherwise.} \end{cases} \quad (\text{E.7})$$

which is an expression similar to that of (E.1), yielding, through analogous methods (which apply since $k_n \leq i < j_n$ by Lemma 2.1 (a, b)),

$$\mathbb{E}_i V_{\pi_T} = \frac{i}{n} + \left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{i}{n}-1\right)}\right) + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{E.8})$$

- iii) If $\bar{k}_n \leq i < j_n$ (assuming that there is any such i ; if not, the empty sum convention used in (E.13) will take care of the case $\bar{k}_n = j_n$) conditionally on Ω_i , a is probed after or at time \bar{k}_n , before time j_n , so a will be rejected. There are in total n steps, in which n could be probed and accepted. Before time \bar{k}_n , the optimal algorithm stops with reward n if and only if any of the first $\bar{k}_n - 1$ values probed is n , because if b were probed before step $\bar{k}_n \leq i$, the algorithm would not stop by definition of \bar{k}_n . Thus the probability of stopping with reward n before time \bar{k}_n is

$$1 - \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1}.$$

From time \bar{k}_n onward, the optimal algorithm stops with reward n if and only if all previous values probed (except for the i th), starting from time k_n , are 0. In fact as of time \bar{k}_n , if b were to be probed before n , the optimal algorithm would always stop with reward b : if probed between time \bar{k}_n (included) and time i , by definition of \bar{k}_n ; from time $i+1 > \bar{k}_n \geq k_n$, by definition of k_n and Lemma 2.1 (d). Thus the probability of stopping with reward n at each step from time \bar{k}_n (there are $n+1-\bar{k}_n$ such steps, due to the exception of the i th step), is the probability of not probing n in any of the first $\bar{k}_n - 1$ steps (which ensures that the algorithm reaches the \bar{k}_n th step) multiplied by the probability of always probing 0, from time \bar{k}_n up to the first time n is probed (skipping time i when appropriate: the actual value of i does not affect the final expression):

$$\frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{n-\bar{k}_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

There are in total n steps, in which b could be probed, but only in $n+1-\bar{k}_n$ of these, from time \bar{k}_n onward, b would be accepted. From the previous comments on the acceptance of

b , at each of these steps, either between time \bar{k}_n and i or from time $i + 1 > \bar{k}_n \geq k_n$, the probability of the optimal algorithm stopping with reward b is the probability of not probing n in any of the first $\bar{k}_n - 1$ steps (which ensures that the algorithm reaches the \bar{k}_n th step) multiplied by the probability that all remaining probed values are 0, from time \bar{k}_n up to the first time b is probed (skipping time i when appropriate: the actual value of i does not affect the final expression):

$$\frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{n - \bar{k}_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

In conclusion for every $\bar{k}_n \leq i < j_n$,

$$(V_{\pi_T} | \Omega_i) \sim \begin{cases} n, & \text{w.p. } 1 - \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} + \frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{n - \bar{k}_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ b, & \text{w.p. } \frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{n - \bar{k}_n} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ 0, & \text{otherwise.} \end{cases} \quad (\text{E.9})$$

This is the same as (E.1), except for formally having $\bar{k}_n + 1$ instead of k_n in the expression. This difference only contributes with $\mathcal{O}(1/n)$ -terms, so we can conclude, by similar methods, that

$$\mathbb{E}_i V_{\pi_T} = \frac{\bar{k}_n}{n} + \left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n} - 1\right)}\right) + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{E.10})$$

- iv) If $i \geq j_n$, conditionally on Ω_i , a is probed after or at time j_n , so a will be accepted and the algorithm will not get past step i . There are in total $i - 1$ steps, in which n could be probed and accepted. Before time \bar{k}_n , the optimal algorithm stops with reward n if and only if any of the first $\bar{k}_n - 1$ values probed is n , because if b were probed before time $\bar{k}_n < i$, the algorithm would not stop by definition of \bar{k}_n . Thus the probability of stopping with reward n before time \bar{k}_n is

$$1 - \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1}.$$

From time \bar{k}_n onward, at each step, the optimal algorithm stops with reward n if and only if all previous values probed, starting from time \bar{k}_n , are 0. In fact as of time \bar{k}_n , if b were to be probed before n , the algorithm would stop with reward b by definition of \bar{k}_n . Thus the probability of stopping with reward n at each step from time \bar{k}_n onward (there are $i - \bar{k}_n$ such steps), is the probability of not probing n in any of the first $\bar{k}_n - 1$ steps (which ensures that the algorithm reaches the \bar{k}_n th step), multiplied by the probability of always probing 0, from time \bar{k}_n up to the first time n is probed:

$$\frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{i - \bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

There are in total $i - 1$ steps, in which b could be probed, but only in $i - \bar{k}_n$ of these, from time \bar{k}_n to time $i - 1$, b would be accepted. From the previous comments on the acceptance of b , at each of these steps, the probability that the optimal algorithm stops with reward b is the probability of not probing n in any of the first $\bar{k}_n - 1$ steps (which ensures that the algorithm reaches the \bar{k}_n th step), multiplied by the probability that all remaining probed values are 0, from time \bar{k}_n up to the first time b is probed. Thus the probability of stopping with b is

$$\frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{i - \bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j.$$

From the above it follows also that the probability of stopping with reward a is the probability of not probing n in any of the first $\bar{k}_n - 1$ steps, multiplied by the probability of always probing 0 in all the remaining $i - \bar{k}_n$ steps, that is

$$\left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{i - \bar{k}_n}.$$

In conclusion for every $i > j_n$,

$$(V_{\pi_T} | \Omega_i) \sim \begin{cases} n, & \text{w.p. } 1 - \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} + \frac{1}{n^2} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{i - \bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ b, & \text{w.p. } \frac{p}{n} \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \sum_{j=0}^{i - \bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^j \\ a, & \text{w.p. } \left(1 - \frac{1}{n^2}\right)^{\bar{k}_n - 1} \left(1 - \frac{p}{n} - \frac{1}{n^2}\right)^{i - \bar{k}_n}, \end{cases} \quad (\text{E.11})$$

and therefore, through the usual estimation methods, by Lemma 2.1(c)

$$\begin{aligned} \mathbb{E}_i V_{\pi_T} &= n \left[\frac{\bar{k}_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right) \right] + \left[1 - \frac{\bar{k}_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right) \right] \left[\left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)}\right) \right] \\ &+ a \left(1 - \frac{\bar{k}_n}{n^2} + \mathcal{O}\left(\frac{1}{n^2}\right)\right) \left(e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) \right) + \mathcal{O}\left(\frac{1}{n}\right) \\ &= \frac{\bar{k}_n}{n} + \left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)}\right) + a e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right). \end{aligned} \quad (\text{E.12})$$

Having exhausted all cases, by the law of total expectation we can compute, by adopting empty sum convention,

$$\begin{aligned} \mathbb{E} V_{\pi_T} &= \frac{1}{n+1} \sum_{i=1}^{k_n-1} \mathbb{E}_i V_{\pi_T} + \frac{1}{n+1} \sum_{i=k_n}^{\bar{k}_n-1} \mathbb{E}_i V_{\pi_T} + \frac{1}{n+1} \sum_{i=\bar{k}_n}^{j_n-1} \mathbb{E}_i V_{\pi_T} + \frac{1}{n+1} \sum_{i=j_n}^{n+1} \mathbb{E}_i V_{\pi_T} \\ &:= S_{1, k_n-1} + S_{k_n, \bar{k}_n-1} + S_{\bar{k}_n, j_n-1} + S_{j_n, n+1}. \end{aligned} \quad (\text{E.13})$$

- i) We start by calculating S_{1, k_n-1} . By (E.6), denoting as $\mu_n := k_n/n$ (note that by Lemma 2.1 (b) this quantity is subunitary and bounded away from zero and one as $n \rightarrow \infty$), we have that

$$\begin{aligned} &\frac{1}{n+1} \sum_{i=1}^{k_n-1} \left[\frac{k_n}{n} + \left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{k_n}{n} - 1\right)}\right) + \mathcal{O}\left(\frac{1}{n}\right) \right] \\ &= \frac{k_n - 1}{n+1} \frac{k_n}{n} + \frac{k_n - 1}{n+1} \left(\frac{1}{p} + b\right) \left(1 - e^{p\left(\frac{k_n}{n} - 1\right)}\right) + \mathcal{O}\left(\frac{1}{n}\right), \end{aligned}$$

so it follows that

$$S_{1, k_n-1} = \mu_n^2 + \mu_n \left(\frac{1}{p} + b\right) \left(1 - e^{p(\mu_n - 1)}\right) + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{E.14})$$

- ii) Next we calculate S_{k_n, \bar{k}_n-1} . By (E.8), denoting $\nu_n := \bar{k}_n/n$ (note that by Lemma 2.1 (a, b, d, e) this quantity is subunitary and bounded away from zero and one as $n \rightarrow \infty$), we have

that

$$\begin{aligned}
& \frac{1}{n+1} \sum_{i=k_n}^{\bar{k}_n-1} \left[\frac{i}{n} + \left(\frac{1}{p} + b \right) \left(1 - e^{p\left(\frac{i}{n}-1\right)} \right) + \mathcal{O}\left(\frac{1}{n}\right) \right] = \frac{\bar{k}_n(\bar{k}_n-1) - k_n(k_n-1)}{2n(n+1)} \\
& + \left(\frac{1}{p} + b \right) \frac{\bar{k}_n - k_n}{n+1} - \frac{1}{n+1} \left(\frac{1}{p} + b \right) \sum_{i=k_n}^{\bar{k}_n-1} e^{p\left(\frac{i}{n}-1\right)} + \mathcal{O}\left(\frac{1}{n}\right) \\
& = \frac{\nu_n^2}{2} - \frac{\mu_n^2}{2} + \left(\frac{1}{p} + b \right) (\nu_n - \mu_n) - \frac{e^{-p}}{n+1} \left(\frac{1}{p} + b \right) \frac{e^{p\mu_n} - e^{p\nu_n}}{1 - e^{\frac{p}{n}}} + \mathcal{O}\left(\frac{1}{n}\right).
\end{aligned}$$

Therefore, since

$$\frac{1}{n+1} \frac{e^{p\mu_n} - e^{p\nu_n}}{1 - e^{\frac{p}{n}}} = \frac{1}{n+1} \frac{e^{p\mu_n} - e^{p\nu_n}}{-\frac{p}{n} + \mathcal{O}\left(\frac{1}{n^2}\right)} = \frac{e^{p\mu_n} - e^{p\nu_n}}{-p + \mathcal{O}\left(\frac{1}{n}\right)} = \frac{e^{p\nu_n} - e^{p\mu_n}}{p} + \mathcal{O}\left(\frac{1}{n}\right),$$

it follows that

$$S_{k_n, \bar{k}_n-1} = \frac{\nu_n^2}{2} - \frac{\mu_n^2}{2} + \left(\frac{1}{p} + b \right) (\nu_n - \mu_n) - \frac{e^{-p}}{p} \left(\frac{1}{p} + b \right) (e^{p\nu_n} - e^{p\mu_n}) + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{E.15})$$

iii) Next we calculate $S_{\bar{k}_n, j_n-1}$. Denoting as $\lambda_n := j_n/n$ (note that by Lemma 2.1 (a) this quantity is subunitary and bounded away from zero and one as $n \rightarrow \infty$), by (E.10) we have that

$$\begin{aligned}
& \frac{1}{n+1} \sum_{i=\bar{k}_n}^{j_n-1} \left[\frac{\bar{k}_n}{n} + \left(\frac{1}{p} + b \right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n}-1\right)} \right) + \mathcal{O}\left(\frac{1}{n}\right) \right] \\
& = \frac{(j_n - \bar{k}_n)\bar{k}_n}{(n+1)n} + \frac{j_n - \bar{k}_n}{n+1} \left(\frac{1}{p} + b \right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n}-1\right)} \right) + \mathcal{O}\left(\frac{1}{n}\right) \\
& = (\lambda_n - \nu_n)\nu_n + (\lambda_n - \nu_n) \left(\frac{1}{p} + b \right) \left(1 - e^{p(\nu_n-1)} \right) + \mathcal{O}\left(\frac{1}{n}\right),
\end{aligned}$$

so it follows that

$$S_{\bar{k}_n, j_n-1} = -\nu_n^2 + \lambda_n \nu_n + (\lambda_n - \nu_n) \left(\frac{1}{p} + b \right) \left(1 - e^{p(\nu_n-1)} \right) + \mathcal{O}\left(\frac{1}{n}\right). \quad (\text{E.16})$$

iv) Finally we compute $S_{j_n, n+1}$. By (E.12) we have that

$$\frac{1}{n+1} \sum_{i=j_n}^{n+1} \left[\frac{\bar{k}_n}{n} + \left(\frac{1}{p} + b \right) \left(1 - e^{p\left(\frac{\bar{k}_n}{n}-\frac{i}{n}\right)} \right) + a e^{p\left(\frac{\bar{k}_n}{n}-\frac{i}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) \right]$$

can be expanded as

$$\begin{aligned}
& \frac{\bar{k}_n(n+2-j_n)}{n(n+1)} + \frac{1}{n+1} \left(\frac{1}{p} + b\right) \sum_{i=j_n}^{n+1} \left(1 - e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)}\right) + \\
& \frac{a}{n+1} \sum_{i=j_n}^{n+1} e^{p\left(\frac{\bar{k}_n}{n} - \frac{i}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) = \nu_n - \lambda_n \nu_n + \left(\frac{1}{p} + b\right) \frac{n+2-j_n}{n+1} - \\
& \frac{1}{n+1} \left(\frac{1}{p} + b - a\right) \sum_{j=j_n-\bar{k}_n}^{n+1-\bar{k}_n} e^{-\frac{pj}{n}} + \mathcal{O}\left(\frac{1}{n}\right) = \nu_n - \lambda_n \nu_n + \left(\frac{1}{p} + b\right) (1 - \lambda_n) - \\
& \frac{1}{n+1} \left(\frac{1}{p} + b - a\right) \left(\frac{e^{-p\frac{j_n-\bar{k}_n}{n}} - e^{-p\frac{n+2-\bar{k}_n}{n}}}{1 - e^{-\frac{p}{n}}}\right) + \mathcal{O}\left(\frac{1}{n}\right) = \nu_n - \lambda_n \nu_n + \\
& \left(\frac{1}{p} + b\right) (1 - \lambda_n) - \frac{1}{n+1} \left(\frac{1}{p} + b - a\right) \frac{e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)} + \mathcal{O}\left(\frac{1}{n}\right)}{1 - e^{-\frac{p}{n}}} + \mathcal{O}\left(\frac{1}{n}\right).
\end{aligned}$$

Therefore, since

$$\begin{aligned}
& \frac{1}{n+1} \frac{e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)} + \mathcal{O}\left(\frac{1}{n}\right)}{1 - e^{-\frac{p}{n}}} = \frac{1}{n+1} \frac{e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)} + \mathcal{O}\left(\frac{1}{n}\right)}{\frac{p}{n} + \mathcal{O}\left(\frac{1}{n^2}\right)} \\
& = \frac{1}{p} \frac{e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)}}{1 + \mathcal{O}\left(\frac{1}{n}\right)} + \mathcal{O}\left(\frac{1}{n}\right) = \frac{1}{p} (e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)}) + \mathcal{O}\left(\frac{1}{n}\right),
\end{aligned}$$

it follows that

$$S_{j_n, n+1} = -\lambda_n \nu_n + \nu_n + \left(\frac{1}{p} + b\right) (1 - \lambda_n) - \frac{1}{p} \left(\frac{1}{p} + b - a\right) (e^{-p(\lambda_n - \nu_n)} - e^{-p(1-\nu_n)}) + \mathcal{O}\left(\frac{1}{n}\right). \tag{E.17}$$

Plugging (E.14) to (E.17) into (E.13) yields, after a few cancellations and collecting of common factors, that

$$\begin{aligned}
\mathbb{E} V_{\pi_T} &= \frac{\mu_n^2}{2} - \frac{\nu_n^2}{2} + \nu_n + \frac{1}{p} + b + \left(\frac{1}{p} - \mu_n\right) \left(\frac{1}{p} + b\right) e^{p(\mu_n - 1)} \\
&+ \left[\left(\frac{1}{p} + b\right) (\nu_n - \lambda_n) - \frac{a}{p}\right] e^{p(\nu_n - 1)} - \frac{1}{p} \left(\frac{1}{p} + b - a\right) e^{p(\nu_n - \lambda_n)} + \mathcal{O}\left(\frac{1}{n}\right) \\
&:= q_{a,b,p}(\lambda_n, \mu_n, \nu_n) + \mathcal{O}\left(\frac{1}{n}\right). \tag{E.18}
\end{aligned}$$

□

F Numerical approximations

Root finding. In this section the code for the computation of $M(a, b, p) \approx 0.7235$ is shared. The full correct list of the decimals is 0.72348603329, and the value provided for the hardness is a rounded-up approximation, following from the parameters set in the bisection method. In the Python method used we set $\text{xtol} = 10^{-13}$ and $\text{rtol} = 10^{-14}$, and the bisection method is implemented such that when it stops, if $\hat{\nu}$ is the root found and ν_* is the true root of $\tilde{q}'(\nu)$, it is guaranteed that

$$|\hat{\nu} - \nu_*| \leq \text{xtol} + |\hat{\nu}| \cdot \text{rtol}.$$

Given that in our case we found $\hat{\nu} \approx 0.211 < 0.3$, this ensures

$$|\hat{\nu} - \nu_*| < 10^{-13} + 0.3 \cdot 10^{-14} < 10^{-12}.$$

Moreover the function $q(\nu)$ has derivative close to zero at $\hat{\nu}$, since ν_* is a stationary point. In particular, for any ζ between $\hat{\nu}$ and ν_* , $|q'(\zeta)| < 1$ (this can be easily verified by direct computation), and therefore by the Lagrange remainder formula for Taylor approximation, we have

$$|q(\hat{\nu}) - \tilde{q}(\nu_*)| \leq |q'(\zeta)| |\hat{\nu} - \nu_*| < |\hat{\nu} - \nu_*| < 10^{-12}.$$

This largely ensures the correctness of the first four significant digits provided for $M(a, b, p)$.

Numerical computations. The Python codes used for numerical computations are available at <https://github.com/AnonSODA24/SODA>.

- The code `bisection.py` approximates the value of the upper bound $M(a, b, p)$ on the gambler-to-prophet ratio of the optimal algorithm for Instance [H](#).
- The code `DP.py` simulates the dynamic program used to produce [Figure 1](#) and an approximation of the gambler-to-prophet ratio of the optimal algorithm for Instance [H](#) with $n = 10^6$, so as to estimate the sharpness of the value obtained for $M(a, b, p)$.