Alon-Tarsi Number of Some Complete Multipartite Graphs

Prajnanaswaroopa S

sntrm4@rediffmail.com

April 11, 2023

ABSTRACT

The Alon-Tarsi number of a polynomial is a parameter related to the exponents of its monomials. For graphs, their Alon-Tarsi number is the Alon-Tarsi number of their graph polynomials. As such, it provides an upper bound on their choice and online choice numbers. In this paper, we obtain the Alon-Tarsi number of some complete multipartite graphs and line graphs of some complete graphs of even order.

1 Introduction

The Combinatorial Nullstellensatz theorem has become a trending method in Algebraic Combinatorics. It is used in a variety of unexpected areas of Combinatorics and Algebra. The theorem, which is an extension, or a generalization of the fundamental theorem of algebra to several variables, is quite useful, primarily in graph theory, number theory, and discrete geometry. Here, we use the method (theorem) in graph theory, specifically for graph colorings/list colorings.

Given a graph G with order and size n and E respectively, we define the average degree of the graph as $a = 2\frac{E}{n}$. Note that, for regular graphs, the average degree of the graph equals their maximum and minimum degrees, hence $a = \Delta$ for such graphs, where Δ is the (maximum) degree of the regular graph. If we give a suitable ordering of the vertices of G in the form x_1, x_2, \ldots, x_n , we define the graph polynomial as the product $P = \prod_{i < j} (x_i - x_j)$ where x_i is adjacent to x_j . Note that the graph polynomial P is homogenous with degree equal to the size of G and maximum exponent of any variable equal to the maximum degree of G. From the original paper regarding Combinatorial Nullstellensatz by Alon([1]) and the crucial paper by Tarsi and Alon([2]), we can define the Alon-Tarsi Number of any polynomial $H = \sum_t c_t y_t$ with $y_t = y_1^{i_1} y_2^{i_2} \ldots y_n^{i_n}$ as $min_t(max_{j_k}(y_1^{i_1}y_2^{i_2} \ldots y_n^{i_n}))$, that is, it is the minimum of the highest exponent of any of the monomials in the polynomial. For graphs, the Alon-Tarsi number can be defined as the Alon-Tarsi number, we say a monomial with the minimum exponent (exponent refers to the maximum over all variables in that monomial) is the Alon-Tarsi monomial. From the references, it is clear that the Alon-Tarsi number of any graph is an upper bound (sometimes strict) on the choice number and hence the chromatic number. From [3] and the primary reference thereof ([4]), the Alon-Tarsi number also upper bounds the online list chromatic number.

One trivial observation from the structure of graph polynomial P is the following.

Lemma 1.1. The Alon-Tarsi Number (ATN) of any graph G is at least $\frac{a}{2}$

Proof. We note that the graph polynomial P of G is homogenous with degree E (size). Therefore, the minimum exponent of any variable will be greater than or equal to $\frac{E}{n}$, where n is the order. By the definition of average degree as given above, the lemma at once follows.

2 Theorems

Proposition 1. Let the vertices of the star graph G with k + 1 vertices and k edges be labeled $x_1, x_2, \ldots, x_{k+1}$, where x_1 is the root vertex, and the rest are pendant vertices. Then, the graph polynomial G has 2^k monomials with precisely $\binom{k}{u}$ monomials having an exponent of x_1 as y.

Proof. The proof proceeds by direct multiplication and observation. The graph polynomial is given by $(x_1 - x_2)(x_1 - x_3) \dots (x_1 - x_{k+1})$. It is clear by observing that any monomial can be formed by first choosing a specified number of x_1 s and the remaining variables in the factors. Also, each of the monomials so formed is distinct. Hence, their Alon-Tarsi number is 1 = 1 = 2, as the Alon-Tarsi monomial, in this case, can be chosen as the monomial $x_2x_3 \dots x_{k+1}$.

Though the following result is already implied by the main result in [2], the approach we use here is relatively straightforward.

Theorem 2.1. The Alon-Tarsi monomial of $G = K_{n,n}$ is of the form $c(x_1x_2ldotsx_n)^{fracn^2} c \neq 0$ for even n. Hence ATN of G is $\frac{n}{2} + 1$.

Proof. In the graph G we have $a = n \implies \frac{a}{2} = \frac{n}{2}$. Therefore, ATN is bounded below by $\frac{n}{2} + 1$. We label all the vertices of one partite set as x_1, x_2, \ldots, x_n and the remaining partite set as $x_{n+1}, x_{n+2}, \ldots, x_{2n}$. Now, the monomial $c(x_1x_2\ldots x_{2n})^{\frac{n}{2}}$ can be formed by taking all the variables $\frac{n}{2}$ times in the product. Observe that all edges are of the form $(x_i - x_j)$, where $i \in \{1, 2, \ldots, n\}$ and $j \in \{n + 1, n + 2, \ldots, 2n\}$. The sign of the product of $x_1^{\frac{n}{2}} x_2^{\frac{n}{2}} \ldots x_n^{\frac{n}{2}}$ are all positive as all the individual signs are positive. Now, the signs of $x_{n+1}^{\frac{n}{2}}, x_{n+2}^{\frac{n}{2}} \ldots x_{2n}^{\frac{n}{2}}$ can be negative. Nevertheless, as the number of variables in each part (and hence the second part) is even, therefore the sign of the product overall will be $(1)^{n\frac{n}{2}}$, which is positive (1). Therefore, the sign of the full product $x_1^{\frac{n}{2}} x_2^{\frac{n}{2}} \ldots x_n^{\frac{n}{2}} x_{n+1}^{\frac{n}{2}}, x_{n+2}^{\frac{n}{2}} \ldots x_{2n}^{\frac{n}{2}}$ is positive (1). Since this will always be the case, the sum of such monomials will be non-zero. Thus, the Alon-Tarsi monomial of G is of the form $c(x_1x_2\ldots x_n)^{fracn2}$ for some non-zero c. Thus, ATN of G attains its lower bound $\frac{n}{2} + 1$.

Corollary 2.1. The ATN of $G = K_{n,n}$ is $1 + \lfloor \frac{n}{2} \rfloor$.

Proof. The proof is immediate once we note the following three observations: 1) The ATN of $K_{n,n}$ for even n is $\frac{n}{2}$. 2) The ATN of G is always $\geq \frac{n}{2}$. 3) G is a subgraph of $K_{n+1,n+1}$. \Box

Theorem 2.2. The ATN of the bipartite graphs $G = K_{m,n}$, m < n with n even and (m+n)|mn is equal to $\frac{mn}{m+n} + 1$.

Proof. In the graph G we have $\frac{a}{2} = \frac{mn}{m+n}$. Therefore, ATN of G is bounded below by $\frac{mn}{m+n} + 1$. We label all the vertices of one part as $x_1, x_2, ldots, x_m$ and the remaining part as $x_{m+1}, x_{m+2}, \ldots, x_n$. The monomial $c(x_1x_2\ldots x_n)^{\frac{mn}{m+n}}$ can be formed by multiplying all the variables $\frac{mn}{m+n}$ times. Observe that the partial product $(x_1x_2\ldots x_n)^{\frac{mn}{m+n}}$ has a positive sign owing to the positivity of the variables. Whereas, the overall sign of $(x_{m+1}x_{m+2}\ldots x_n)^{\frac{mn}{m+n}}$ is equal to $(-1)^{n\frac{mn}{m+n}} = 1$. Therefore, all such monomials are always positive in sign; hence the sum of these monomials will also be non-zero. This, in turn, implies the theorem.

Theorem 2.3. The ATN of regular bipartite graph G with 2n vertices with even n and even degree Δ is $\frac{\Delta}{2}$.

Proof. Again, we label the vertices of G as x_1, x_2, \ldots, x_n for the vertices of one part, and $x_{n+1}, x_{n+2}, \ldots, x_{2n}$ for the remaining vertices. All edges are of the form $(x_i - x_j)$ where i and j are from different parts. This implies we can form a monomial by choosing each of the variables $\frac{\Delta}{2}$ times. As in the previous theorem, the sign of the product $(x_1x_2\ldots x_n)^{\frac{\Delta}{2}}$ is always positive as all the individual variables are. As for the sign of $(x_{n+1}x_{n+2}\ldots x_{2n})^{\frac{Delta}{2}}$, we note that as n is even, the combined sign of this partial product would be $(1)^{n\frac{Delta}{2}} = 1$. Therefore, the full sign of the product monomial $(x_1x_2\ldots x_nx_{n+1}x_{n+2}\ldots x_{2n})^{\frac{\Delta}{2}}$ is always positive (1). Hence, the sum of such monomials will be non-zero; in equivalent words, ATN of G is $\frac{Delta}{2}$.

Theorem 2.4. The ATN of the complete k-partite graph $K_{n,n,\dots(k-times),\dots,n}$ lies between $\left\lceil \frac{(k-1)n}{2} \right\rceil + 1$ and $(k-1) \left\lceil \frac{n}{2} \right\rceil + 1$.

Proof. First, we partition the vertices into k parts (independent sets) I_1, I_2, \ldots, I_k , with each part having n vertices as suggested by the k- partiteness. We denote the vertices ordered as x_1, x_2, \ldots, x_n for the first part; $x_{n+1}, x_{n+2}, \ldots, x_{2n}$ for the second and so on. The edges are now given by $(x_i - x_j)$ where i, j are from different partite sets. We observe that for every partite set, there are exactly (k-1) partite sets with which the induced graph formed will be the complete bipartite graph $K_{n,n}$. Hence, the Alon-Tarsi monomials for each of these induced graphs will be of the form $c_i(x_{i_1}x_{i_2}\ldots x_{i_{2n}})^{\left\lceil \frac{n}{2} \right\rceil}$ for some non-zero c_i and variables i_k s. The product of all such monomials will give us the desired non-zero monomial $(x_1x_2\ldots x_{nk})^{\left\lceil \frac{n}{2} \right\rceil}$. Thus, an upper bound for the ATN of G is $(k-1) \left\lceil \frac{n}{2} \right\rceil + 1$. On the other hand, we have $a = \frac{k(k-1)n^2}{nk}$. Therefore, the lower bound for ATN is $(k-1) \left\lceil \frac{n}{2} \right\rceil + 1$. Hence, the claim follows.

Corollary 2.2. The ATN of the complete k- partite graph $K_{n,n,\dots(k-times),\dots,n}$ for even n is equal to $(k-1)\frac{n}{2}$.

Proof. The proof is immediate from the previous theorem once we know that $\left\lceil \frac{n}{2} \right\rceil = \frac{n}{2}$.

A related result to the next result was proved using elaborate techniques in [5] for complete graphs of prime degree.

Theorem 2.5. The ATN of the line graph of $G = K_n$ for n = 4k, $k \in \mathbb{N}$ is n - 1. Hence, the edge choosability of K_n is n - 1, or are chromatic edge choosable.

Proof. We note that we can factorize the edges of K_n into $\Delta = n - 11$ – factors or perfect matchings. We label the edges (vertices of line graph) as x_1, x_2, \ldots, x_{2k} for the first perfect matching; $x_{2k+1}, x_{2k+2}, \ldots, x_n$ and so on. We also observe that an edge in any perfect matching is adjacent to exactly two edges in any other perfect matching (the adjacencies are determined by the end vertices). Thus, the line graphs of G are n - 1 -partite with any two partite sets inducing a regular bipartite graph of degree 2. Now, the Alon-Tarsi monomial of the induced graph formed by any two partite sets in the line graph is of the form $c_i(x_{i_1}x_{i_2}\ldots x_{i_{2n}})^{\frac{2}{2}} = c_i(x_{i_1}x_{i_2}\ldots x_{i_{2n}})$ for some non-zero c_i and indices i_k s. Since there are (n - 2) such regular bipartite graphs (the induced graphs between any two perfect matchings), we get that the final Alon-Tarsi number should be (n - 2)(1) + 1 = n - 1. Hence, the claim follows.

References

- [1] Alon, N. (1999). Combinatorial nullstellensatz. Combinatorics, Probability and Computing, 8(1-2), 7-29.
- [2] Alon, N., and Tarsi, M. (1992). Colorings and orientations of graphs. Combinatorica, 12, 125-13.
- [3] Carraher, J., Loeb, S., Mahoney, T., Puleo, G. J., Tsai, M. T., and West, D. B. (2014). Three topics in online list coloring. Journal of Combinatorics, 5(1), 115-130.

- [4] Schauz, U. (2009). Mr. paint and mrs. correct. the electronic journal of combinatorics, R77-R77.
- [5] Schauz, U. (2014). Proof of the list edge coloring conjecture for complete graphs of prime degree. The Electronic Journal of Combinatorics, P3-43.
- [6] Zhu, X., and Balakrishnan, R. (2021). Combinatorial Nullstellensatz: With Applications to Graph Colouring. CRC Press.