# On the Nash Equilibria of a Simple Discounted Duel

Athanasios Kehagias

April 12, 2023

#### Abstract

We formulate and study a two-player *static duel* game as a *nonzero-sum discounted* stochastic game. Players  $P_1, P_2$  are standing in place and, in each turn, one or both may shoot at the "other" player. If  $P_n$  shoots at  $P_m$   $(m \neq n)$ , either he hits and kills him (with probability  $p_n$ ) or he misses him and  $P_m$  is unaffected (with probability  $1 - p_n$ ). The process continues until at least one player dies; if nobody ever dies, the game lasts an infinite number of turns. Each player receives unit payoff for each turn in which he remains alive; no payoff is assigned to killing the opponent. We show that the the always-shooting strategy is a NE but, in addition, the game also possesses "cooperative" (i.e., non-shooting) Nash equilibria in both stationary and nonstationary strategies. A certain similarity to the repeated Prisoner's Dilemma is also noted and discussed.

## 1 Introduction

In this paper we study a two-player *static duel* game played in turns. Players  $P_1$ ,  $P_2$  are standing in place and, in each turn, one or both may shoot at the "other" player. If  $P_n$  shoots at  $P_m$  $(m \neq n)$ , either he hits and kills him or he misses him and  $P_m$  is unaffected; the respective probabilities are  $p_n$  and  $1-p_n$ . The process continues until at least one player dies; it is possible that nobody ever dies and the game lasts an infinite number of turns. We formulate the above as a nonzero-sum discounted stochastic game. The game rules and the players' payoff function will be presented in the next section.

Little work has been done on the static duel. Actually, as far as we know, it has only been studied as a preliminary step in the study of the "static *truel*", in which *three* stationary players shoot at each other. Early works on the static truel are [\[11,](#page-17-0) [16,](#page-17-1) [18,](#page-17-2) [19,](#page-17-3) [20\]](#page-17-4) in which the postulated game rules guarantee the existence of *exactly one* survivor ("winner"). A more general analysis appears in [\[17\]](#page-17-5) which considers the possibility of "cooperation" between the players. This idea is further studied in [\[13,](#page-17-6) [14,](#page-17-7) [15,](#page-17-8) [25\]](#page-18-0). Recent papers on the truel include  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$  $[1, 2, 3, 5, 6, 7, 8, 9, 22, 23, 24]$ .

<span id="page-0-0"></span><sup>&</sup>lt;sup>1</sup>Let us also note the existence of an extensive literature on a quite different type of duel games, which essentially are *games of timing* [\[4,](#page-17-15) [10,](#page-17-16) [12\]](#page-17-17). However, this literature is not relevant to the game studied in this paper.

While the above papers focus on various forms of the static truel, we believe that the static duel is interesting in its own right and has not received the attention it deserves. In particular we will show that, under our formulation, the static duel has a certain similarity to the repeated Prisoner's Dilemma and possesses "cooperative" Nash equilibria in nonstationary strategies.

This paper is structured as follows. In Section [2](#page-1-0) we define the game rigorously. In Section [3](#page-3-0) we introduce several stationary and nonstationary strategies and compute their expected payoffs. In Section [4](#page-7-0) we prove that certain pairs of the previously defined strategies are Nash equilibria. In Section [5](#page-15-0) we discuss the obtained results and the connection of the static duel to the repeated Prisoner's Dilemma. Finally, in Section [6](#page-16-2) we summarize our results and propose some future research directions.

#### <span id="page-1-0"></span>2 The Game

The game involves players  $P_1, P_2$  and proceeds at discrete time steps (rounds)  $t \in \{1, 2, ...\}$ . The *state* at time t is

$$
s(t) = (s_1(t), s_2(t)) \in S = \{(1, 1), (1, 0), (0, 1), (0, 0), (\tau, \tau)\}.
$$

For  $n \in \{1, 2\}$ ,  $s_n(t)$  is  $P_n$ 's state at  $t \in \{0, 1, 2, ...\}$  and can be

- 1 : when  $P_n$  is alive,
- 0 : when  $P_n$  dies in the current round,
- $\tau$ : when one or both players have died in a previous round.

 $P_n$ 's action at  $t \in \{1, 2, ...\}$  is  $f_n(t)$ , which can be 1  $(P_n$  is shooting) or 0  $(P_n$  is not shooting). If  $s_n(t-1) \neq 1$ ,  $P_n$  cannot shoot at t and  $f_n(t)$  must equal 0; if  $s_n(t-1) = 1$  then  $f_n(t)$  can be either 0 or 1. When  $f_n(t) = 1$ ,  $s_{-n}(t) = 0$  (i.e.,  $P_{-n}$  dies<sup>[2](#page-1-1)</sup>) with probability  $p_n \in (0,1)$  and  $s_{-n}(t) = 1$  with probability  $1 - p_n$ . We set  $f(t) = (f_1(t), f_2(t))$  and  $\mathbf{p} = (p_1, p_2)$ . Note that we have assumed that  $p_1$ ,  $p_2$  are different from both zero and one.

The game starts at an initial state  $s(0)$ ; obviously, the main case of interest is  $s(0)$  = (1, 1). At times  $t \in \{1, 2, ...\}$  the players choose simultaneously the actions  $f_1(t)$ ,  $f_2(t)$ and the game moves to state  $s(t)$  according to the conditional state transition probability  $Pr(s(t)|s(t-1), f(t))$ . In Figure 1 we present the state transition diagram, in which the action-dependent transition probabilities are written next to the edges; it is easily verified that these probabilities conform to the game rules. The figure shows that the game starting at  $(1, 1)$ , lasts an infinite number of rounds and two possibilities exist.

- 1. The game always stays in (1, 1) (no player is ever killed).
- 2. At some t' the game moves to a state  $s \in \{(1,0), (0,1), (0,0)\}$  (one or both players are killed) and at  $t' + 1$  the game moves to the *terminal* state  $(\tau, \tau)$ , where it stays for ever after.

<span id="page-1-1"></span><sup>&</sup>lt;sup>2</sup>In the sequel we use the standard game theoretic notation by which  $s_{-1} = s_2$ ,  $s_{-2} = s_1$ . The same notation is used for players, actions etc.



Figure 1: State transition diagram of the static duel game.

A finite history is a sequence  $h = s(0) f(1) s(1) ... f(T) s(T)$ ; an infinite history is an  $h = s(0) f(1) s(1) \dots$ . An *admissible* history is one which conforms to the game rules; the set of all admissible finite (resp. infinite) histories is denoted by H (resp.  $H^{\infty}$ ). For every history h and for  $n \in \{1, 2\}$ , we define  $P_n$ 's total payoff function to be

$$
Q_n(h) = \sum_{t=0}^{\infty} \gamma^t q_n(s(t))
$$

where  $\gamma \in (0,1)$  is the discounting factor and  $q_n : S \to \mathbb{R}$  is  $P_n$ 's stage payoff function, defined as follows.

$$
q_1(\tau, \tau) = 0, \t q_2(\tau, \tau) = 0,
$$
  
\n
$$
q_1(1, 1) = 1, \t q_2(1, 1) = 1,
$$
  
\n
$$
q_1(1, 0) = \frac{1}{1-\gamma}, \t q_2(1, 0) = 0,
$$
  
\n
$$
q_1(0, 1) = 0, \t q_2(0, 1) = \frac{1}{1-\gamma},
$$
  
\n
$$
q_1(0, 0) = 0, \t q_2(0, 0) = 0.
$$

stays alive; the payoff  $q_1(1,0) = \frac{1}{1-\gamma}$  incorporates the infinite payoff sequence  $\sum_{t=0}^{\infty} \gamma^t 1 = \frac{1}{1-\gamma}$ The above values indicate that each player receives one payoff unit for every turn in which he (this will result when  $P_1$  kills  $P_2$  and stays alive for the infinite number of subsequent turns). Note that a player receives no direct payoff from killing his opponent, but he has the indirect benefit of removing the possibility of being killed himself.

A strategy for  $P_n$  is a function  $\sigma_n : H \to [0,1]$  which corresponds to every finite history h the probability

$$
x_n = \sigma_n(h) = \Pr(\text{``}P_n \text{ shoots at } P_{-n}\text{''})\,.
$$

A stationary strategy is a  $\sigma_n$  depending only on the current state s, hence we simply write  $x_n = \sigma_n(s)$ . A strategy profile is a vector  $\sigma = (\sigma_1, \sigma_2)$ . We denote the set of all *admissible*  strategies (those which are compatible with the game rules) by  $\Sigma$  and the set of all admissible stationary strategies by  $\Sigma$ .

Given the initial state  $s(0)$  and the strategies  $\sigma_1$  and  $\sigma_2$ , used by  $P_1$  and  $P_2$  respectively, a probability measure is defined on the set of all infinite histories. Since  $\gamma \in (0,1)$ , the total expected payoffs

$$
\forall n \in \{1,2\} : \overline{Q}_n(s(0), \sigma_1, \sigma_2) = \mathbb{E}\left(Q_n(h) \, | s(0), \sigma_1, \sigma_2\right)
$$

are well defined.

We have thus formulated the simultaneous static duel as a discounted stochastic game, which we will denote by  $\Gamma(s(0), \gamma, \mathbf{p})$  or simpl;  $\Gamma(s(0), \gamma)$ , when  $(p_1, p_2)$  is fixed. Our main interest is in the nonzero-sum game  $\Gamma((1,1), \gamma)$ . We assume that  $P_1$  and  $P_2$  attempt to reach a Nash equilibrium (NE), i.e., a strategy profile  $(\widehat{\sigma}_1, \widehat{\sigma}_2)$  such that

$$
\forall n \in \{1,2\} : \forall \sigma_n \in \Sigma : \overline{Q}_n((1,1),\widehat{\sigma}_n,\widehat{\sigma}_{-n}) \geq \overline{Q}_n((1,1),\sigma_n,\widehat{\sigma}_{-n}).
$$

#### <span id="page-3-0"></span>3 Some Basic Strategies and Their Payoffs

In this section we introduce several strategies which we will use in our later exploration of Nash equilibria.

#### 3.1 The Stationary Strategy  $\sigma^S$

When  $P_n$   $(n \in \{1,2\})$  uses a stationary admissible strategy  $\sigma_n$ , we have  $\sigma_n(s) = 0$  for  $s \in$  $\{(1,0), (0,1), (0,0), (\tau, \tau)\}\$ and  $\sigma_n$  is fully specified by the value  $\sigma_n(1, 1) = x_n$ . Hence we will sometimes write  $\overline{Q}_n((1,1), x_1, x_2)$  in place of  $\overline{Q}_n((1,1), \sigma_1, \sigma_2)$ .

Let  $V_1^S(x_1, x_2) = \overline{Q}_1((1,1), x_1, x_2)$ ; for brevity we will also write simply  $V_1^S$ . Then  $V_1^S$ satisfies the equation

<span id="page-3-1"></span>
$$
V_1^S = 1 + \gamma p_1 x_1 (x_2 (1 - p_2) + (1 - x_2)) \frac{1}{1 - \gamma} + \gamma (x_1 (1 - p_1) + (1 - x_1)) (x_2 (1 - p_2) + (1 - x_2)) V_1^S,
$$
\n(1)

obtained by the following reasoning. When the game is in state  $s = (1, 1)$ ,  $P_1$ 's expected payoff is one unit for the current state plus the discounted expected payoff from the subsequent state  $s'$ , for which we have the following possibilities.

- 1.  $s' = (1, 0)$  when  $P_1$  shoots and hits  $P_2$  and  $P_2$  either shoots and misses or does not shoot; the respective probability is  $p_1x_1(x_2(1-p_2) + (1-x_2))$ . The total expected payoff of this case is  $\overline{Q}_1((1,0), x_1, x_2) = \frac{1}{1-\gamma}$ .
- 2.  $s' = (1, 1)$  when each of  $P_1$  and  $P_2$  either shoots and misses or does not shoot; the respective probability is  $(x_1 (1 - p_1) + (1 - x_1)) (x_2 (1 - p_2) + (1 - x_2))$ . In this case we have returned to the starting state  $(1, 1)$  and the additional total expected payoff is again  $Q_1((1,1), x_1, x_2).$

3. We also have the possibilities of moving into  $(0, 1)$  and  $(0, 0)$ , but these yield zero payoff to  $P_1$ , so they are not included in  $(1)$ .

Solving  $(1)$  we get, after some algebraic calculations<sup>[3](#page-4-0)</sup>, that

<span id="page-4-1"></span>
$$
V_1^S(x_1, x_2) = \frac{1 - \gamma (1 - p_1 x_1 (1 - p_2 x_2))}{(1 - \gamma) (1 - \gamma (1 - p_1 x_1) (1 - p_2 x_2))}.
$$
\n(2)

In what follows we will often work with the *normalized* total expected payoff. In this case, it is

<span id="page-4-2"></span>
$$
v_1^S(x_1, x_2) = (1 - \gamma) \overline{Q}_1((1, 1), x_1, x_2) = \frac{1 - \gamma (1 - p_1 x_1 (1 - p_2 x_2))}{1 - \gamma (1 - p_1 x_1 (1 - p_2 x_2))}
$$
(3)

Formulas for  $V_2^S(x_1, x_2) = \overline{Q}_2((1, 1), x_1, x_2)$  and  $v_2^S(x_1, x_2) = (1 - \gamma)\overline{Q}_2((1, 1), x_1, x_2)$  can be obtained by interchanging the indices 1 and 2 in [\(2\)](#page-4-1)-[\(3\)](#page-4-2).

#### $3.2$  The Cooperating Strategy  $\sigma^C$

The stationary "cooperating" (the name will be justified in Section [5\)](#page-15-0) strategy  $\sigma^C$  is defined by

 $\sigma^{C}(1,1) = 0$ , which means the player never shoots.

Obviously,  $\sigma^C$  is  $\sigma^S$  with  $x_n = 0$  and  $\overline{Q}_n((1,1), \sigma^C, \sigma^C) = \overline{Q}_n((1,1), 0, 0)$ . Hence we obtain  $V_1^C = \overline{Q}_1((1,1), \sigma^C, \sigma^C) = \overline{Q}_1((1,1), 0, 0)$  by setting  $x_1 = x_2 = 0$  in [\(2\)](#page-4-1). Consequently the expected and normalized expected total payoff are

$$
V_1^C = \frac{1}{1 - \gamma}, \qquad v_1^C = (1 - \gamma) V_1^C = 1
$$

Formulas for  $V_2^C = \overline{Q}_2(\sigma^C, \sigma^C)$ ,  $v_2^c = (1 - \gamma) V_2^C$  are obtained by exchanging the indices 1 and 2 in the above formulas.

#### $3.3$  The Defecting Strategy  $\sigma^D$

The stationary "defecting" (the name will be justified in Section [5\)](#page-15-0) strategy  $\sigma^D$  is defined by

 $\sigma^{D}(1,1) = 1$ , which means the player always shoots with probability one.

Obviously,  $\sigma^D$  is  $\sigma^S$  with  $x_n = 1$  and  $\overline{Q}_n((1,1), \sigma^D, \sigma^D) = \overline{Q}_n((1,1), 1, 1)$ . Hence we obtain  $V_1^D = \overline{Q}_1((1,1), \sigma^D, \sigma^D) = \overline{Q}_1((1,1), 1, 1)$  by setting  $x_1 = x_2 = 1$  in [\(2\)](#page-4-1). We then get the expected and normalized expected total payoff to be

$$
V_1^D = \frac{1 - \gamma (1 - p_1 (1 - p_2))}{(1 - \gamma) (1 - \gamma (1 - p_1) (1 - p_2))}
$$

$$
v_1^D = (1 - \gamma) V_1^D(\gamma) = \frac{1 - \gamma (1 - p_1 (1 - p_2))}{1 - \gamma (1 - p_1) (1 - p_2)}
$$

Formulas for  $V_2^D = \overline{Q}_2(\sigma^D, \sigma^D)$ ,  $v_2^D = (1 - \gamma) V_2^D$  are obtained by exchanging the indices 1 and 2 in the above formulas.

<span id="page-4-0"></span><sup>&</sup>lt;sup>3</sup>The calculations required to obtain the solution have been performed by the computer algebra system Maple and then verified by hand. This is also true of additional (sometimes quite complicated) calculations required in the rest of the paper.

#### $3.4$  The Early-Shooting Strategy  $\sigma^{DC(K)}$

The nonstationary "early-shooting" strategy  $\sigma^{DC(K)}$  is to shoot (with probability one) only at times 1, 2, ..., K, where K is a parameter of the strategy. Let  $V_1^{DC(K)} = Q_1((1,1), \sigma^{DC(K)}, \sigma^{DC(K)})$ ; then  $V_1^{DC(K)}$  $\sum_{1}^{D}C(A)$  satisfies the equation:

<span id="page-5-0"></span>
$$
V_1^{DC(K)} = 1
$$
  
+  $\gamma \left( p_1 (1 - p_2) \frac{1}{1 - \gamma} + (1 - p_1) (1 - p_2) \right)$   
+  $\gamma^2 \left( (1 - p_1) p_1 (1 - p_2)^2 \frac{1}{1 - \gamma} + (1 - p_1)^2 (1 - p_2)^2 \right)$   
+ ...  
+  $\gamma^{K-1} \left( (1 - p_1)^{K-2} p_1 (1 - p_2)^{K-1} \frac{1}{1 - \gamma} + (1 - p_1)^{K-1} (1 - p_2)^{K-1} \right)$   
+  $\gamma^K \left( (1 - p_1)^{K-1} p_1 (1 - p_2)^K \frac{1}{1 - \gamma} + (1 - p_1)^K (1 - p_2)^K V_1^C \right)$  (4)

This equation is justified as follows.

- 1. At time  $t = 0$ ,  $P_1$  receives a payoff of one unit.
- 2. At times  $t = 1, ..., K 1$ , the expected payoffs are the following.
	- (a) With probability  $(1 p_1)^{t-1} p_1 (1 p_2)^t$ ,  $P_1$  misses at times  $t' = 1, ..., t 1$  and succeeds at time  $t'$ , while  $P_2$  misses as times  $t' = 1, ..., t$ . In this case  $P_1$  receives payoff 1  $\frac{1}{1-\gamma}$ .
	- (b) With probability  $(1-p_1)^t (1-p_2)^t$ , both  $P_1$  and  $P_2$  miss at times  $t' = 1, ..., t$ . In this case  $P_1$  receives payoff 1.
	- (c) All other possibilities yield zero payoff, so they are not included in the equation.
- 3. At time  $t = K$ , the expected payoffs are the following.
	- (a) With probability  $(1 p_1)^{K-1} p_1 (1 p_2)^K$ ,  $P_1$  misses at times  $t' = 1, ..., K 1$  and succeeds at time  $t' = K$ , while  $P_2$  misses as times  $t' = 1, ..., K$ . In this case  $P_1$  receives payoff  $\frac{1}{1-\gamma}$ .
	- (b) With probability  $(1 p_1)^K (1 p_2)^K$ , both  $P_1$  and  $P_2$  miss at times  $t' = 1, ..., K$ . In this case both players will never shoot at subsequent times, so we have returned to the starting state  $(1, 1)$  and the additional total expected payoff is  $V_1^C = \overline{Q}_1((1, 1), \sigma^C, \sigma^C)$ .
	- (c) All other possibilities yield zero payoff, so they are not included in the equation.

Substituting in [\(4\)](#page-5-0) the expression for  $V_1^C$  we obtain the following expressions for expected total and normalized expected total payoff:

$$
V_1^{DC(K)} = \frac{1 + \gamma^{K+1} (1 - p_1)^K (1 - p_2)^K p_2 - \gamma (1 - p_1 + p_1 p_2)}{(1 - \gamma) (1 - \gamma (1 - p_1) (1 - p_2))}
$$
  

$$
v_1^{DC(K)} = (1 - \gamma) V_1^{DC(K)} = \frac{1 + \gamma^{K+1} (1 - p_1)^K (1 - p_2)^K p_2 - \gamma (1 - p_1 + p_1 p_2)}{1 - \gamma (1 - p_1) (1 - p_2)}
$$

Formulas for  $V_2^{DC(K)}$  $v_2^{DC(K)}, v_2^{DC(K)}$  $\frac{DC(K)}{2}$  are obtained by exchanging the indices 1 and 2 in the above formulas.

#### 3.5 The Late-Shooting Strategy  $\sigma^{CD(K)}$

The nonstationary "late-shooting" strategy  $\sigma^{CD(K)}$  is to shoot (with probability one) only at times  $K, K+1, ...,$  where  $K$  is a parameter of the strategy. Let  $V_1^{CD(K)} = Q_1((1,1), \sigma^{CD(K)}, \sigma^{CD(K)});$ then  $V_1^{CD(K)}$  $\mathcal{L}^{CD(K)}$  satisfies the equation:

<span id="page-6-0"></span>
$$
V_1^{CD(K)} = 1 + \gamma + \dots + \gamma^{K-1} + \gamma^K \left( p_1 \left( 1 - p_2 \right) \frac{1}{1 - \gamma} + \left( 1 - p_1 \right) \left( 1 - p_2 \right) V_1^D \right) \tag{5}
$$

This equation is justified as follows.

- 1. At times  $t = 0, 1, ..., K 1, P_1$  receives discounted payoff of one unit.
- 2. At time  $t = K$  we have the following possibilities.
	- (a) With probability  $p_1 (1 p_2)$ ,  $P_1$  hits  $P_2$ , while  $P_2$  misses. In this case  $P_1$  receives payoff  $\frac{1}{1-\gamma}$ .
	- (b) With probability  $(1 p_1) (1 p_2)$ , both  $P_1$  and  $P_2$  miss. In this case both  $P_1$  and  $P_2$ revert to strategy  $\sigma^D$  and  $P_1$  receives total expected payoff  $V_1^D = \overline{Q}_1((1,1), \sigma^D, \sigma^D)$ .
	- (c) All other possibilities yield zero payoff, so they are not included in the equation.

Substituting in [\(5\)](#page-6-0) the previously obtained expression for  $V_1^D$  we get the following expressions for expected total and normalized expected total payoff:

$$
V_1^{CD(K)} = \frac{1 - \gamma^K p_2 - \gamma (1 - p_1) (1 - p_2)}{(1 - \gamma) (1 - \gamma (1 - p_1) (1 - p_2))}
$$
  

$$
v_1^{CD(K)} = (1 - \gamma) V_1^{CD(K)} = \frac{1 - \gamma^K p_2 - \gamma (1 - p_1) (1 - p_2)}{1 - \gamma (1 - p_1) (1 - p_2)}
$$

Formulas for  $V_2^{CD(K)}$  $v_2^{CD(K)}, v_2^{CD(K)}$  $_{2}^{\text{U}(K)}$  are obtained by exchanging the indices 1 and 2 in the above formulas.

#### 3.6 The Periodic-Shooting Strategy  $\sigma^{P(M)}$

The nonstationary "periodic-shooting" strategy  $\sigma^{P(M)}$  is to shoot only at times  $M+1, 2M+2, ...,$ where M is strategy parameter. Let  $V_1^{P(M)} = Q_1((1,1), \sigma^M, \sigma^M)$ ; by reasoning similar to that of the previous cases, we see that  $V_1^{P(M)}$  $\sum_{1}^{H(M)}$  satisfies the equation

$$
V_1^{P(M)} = 1 + \gamma + \dots + \gamma^M
$$
  
+  $\gamma^{M+1} \left( p_1 (1 - p_2) \frac{1}{1 - \gamma} + (1 - p_1) (1 - p_2) V_1^{P(M)} \right)$  (6)

Solving [\(6\)](#page-7-1) we get the following expressions for expected total and normalized expected total payoff:

<span id="page-7-1"></span>
$$
V_1^{P(M)} = \frac{1 - \gamma^{M+1} (1 - p_1 (1 - p_2)))}{(1 - \gamma) (1 - \gamma^{M+1} (1 - p_1) (1 - p_2))}
$$

$$
v_1^{P(M)} = \frac{1 - \gamma^{M+1} (1 - p_1 (1 - p_2))}{1 - \gamma^{M+1} (1 - p_1) (1 - p_2)}
$$

Formulas for  $V_2^{P(M)}$  $v_2^{P(M)}, v_2^{P(M)}$  $2^{P(M)}$  are obtained by exchanging the indices 1 and 2 in the above formulas.

#### 3.7 Grim Strategies

Any  $P_n$ 's strategy  $\sigma_n$  can be used to define a corresponding grim strategy  $\tilde{\sigma}_n$  as follows:

 $\widetilde{\sigma}_n$ : As long as  $P_{-n}$  uses  $\sigma_n$ ,  $P_n$  also uses  $\sigma_n$ ; if at the t-th turn  $P_{-n}$  deviates from  $\sigma_n$ ,  $P_n$  uses  $\sigma^D$  in all subsequent turns.

For example, the grim-cooperating strategy  $\tilde{\sigma}^C$  dictates that:  $P_n$  never shoots, as long as  $P_{-n}$ <br>does not shoot gither; if at the t th turn  $P_n$  shoots than starting at the  $(t+1)$  th turn  $P_n$ does not shoot either; if at the t-th turn  $P_{-n}$  shoots then, starting at the  $(t + 1)$ -th turn,  $P_n$ will always shoot with probability one. Similarly we can get:

- 1. the grim-defecting strategy  $\tilde{\sigma}^D$  (it is identical to  $\sigma^D$ ),
- 2. the grim-early-shooting strategy  $\tilde{\sigma}^{DC(K)}$ ,
- 3. the grim-late-shooting strategy  $\tilde{\sigma}^{CD(K)}$ ,
- 4. the grim-periodic-shooting strategy  $\tilde{\sigma}^{P(M)}$ .

## <span id="page-7-0"></span>4 Nash Equilibria

We will now pesent a sequence of propositions; each one indicates that a certain strategy pair is a (stationary or nonstationary) NE; sometimes this will only hold for a certain range of  $\gamma$  and possibly  $p_1, p_2$  values.

**Proposition 4.1** For every  $\gamma \in (0,1)$ , the only stationary NE of  $\Gamma((1,1), \gamma)$  are  $(\sigma^C, \sigma^C)$  and  $(\sigma^D, \sigma^D).$ 

**Proof.** Suppose that  $P_1$  (resp.  $P_2$ ) uses the stationary strategy  $\sigma^S$  with  $\sigma^S(1,1) = x_1$  (resp.  $\sigma^S$  with  $\sigma^S(1,1) = x_2$ ). Then  $P_1$ 's payoff is

$$
V_1^S(x_1,x_2)=\frac{1-\gamma(1-p_1x_1(1-p_2x_2))}{(1-\gamma)(1-\gamma(1-p_1x_1)(1-p_2x_2))}.
$$

Now suppose  $P_1$  switches to  $\sigma_S(1,1) = y_1$ ; his payoff becomes

$$
V_1^S(y_1, x_2) = \frac{1 - \gamma (1 - p_1 y_1 (1 - p_2 x_2))}{(1 - \gamma) (1 - \gamma (1 - p_1 y_1) (1 - p_2 x_2))}.
$$

Let us look at the difference of normalized payoffs

$$
\delta v_1 = (1 - \gamma) (Q_1((1, 1), x_1, x_2) - Q_1((1, 1), y_1, x_2))
$$
  
= 
$$
\frac{1 - \gamma (1 - p_1 x_1 (1 - p_2 x_2))}{1 - \gamma (1 - p_1 x_1) (1 - p_2 x_2)} - \frac{1 - \gamma (1 - p_1 y_1 (1 - p_2 x_2))}{1 - \gamma (1 - p_1 y_1) (1 - p_2 x_2)}
$$
  
= 
$$
\frac{\gamma^2 p_1 p_2 x_2 (x_1 - y_1)(1 - p_2 x_2)}{((1 - \gamma (1 - p_2 x_2)(1 - p_1 y_1))(1 - \gamma (1 - p_2 x_2)(1 - p_1 x_1)))}
$$

Now,  $P_1$  has no incentive to switch from  $x_1$  to  $y_1$  iff  $\delta v_1 \geq 0$  which is equivalent to

$$
\gamma^2 p_1 p_2 x_2 (x_1 - y_1)(1 - p_2 x_2) \ge 0
$$

Similarly,  $P_2$  has no incentive to switch from  $x_2$  to  $y_2$  iff

$$
\gamma^2 p_1 p_2 x_1 (x_2 - y_2)(1 - p_1 x_1) \ge 0
$$

Hence, the following hold for  $n \in \{1,2\}$ .

- 1. If  $(x_1, x_2) = (0, 0), P_n$  has no incentive to change  $x_n$ ;  $(x_1, x_2) = (0, 0)$ ; hence  $(\sigma^C, \sigma^C)$  is a NE.
- 2. If  $(x_1, x_2) = (1, 1), P_n$  has no incentive to change  $x_n$ ;  $(x_1, x_2) = (1, 1)$ ; hence  $(\sigma^D, \sigma^D)$  is a NE.
- 3. If  $(x_1, x_2) \in (0, 1) \times (0, 1)$  then it cannot be a NE, because  $P_n$  has incentive to change (unilaterally) from  $x_n$  to 1.

This completes the proof. ■

Now we will start looking at NE obtained from combinations of grim strategies.

**Proposition 4.2** For every  $\gamma \in (0,1)$ ,  $(\tilde{\sigma}^C, \tilde{\sigma}^C)$  is a NE of  $\Gamma((1,1), \gamma)$ .

**Proof.** Suppose that both  $P_1$  and  $P_2$  use  $\tilde{\sigma}^C$ . Then  $P_1$ 's payoff is

$$
V_1^C = Q_1\left(\widetilde{\sigma}^C, \widetilde{\sigma}^C\right) = Q_1\left(\sigma^C, \sigma^C\right) = \frac{1}{1-\gamma}.
$$

Now suppose  $P_1$  deviates from  $\tilde{\sigma}^C$ . It suffices to examine the case in which  $P_1$  deviates at  $t = 1$ ; furthermore after P deviates (i.e. starting at  $t = 2$ ) P will switch to  $\sigma^D$  and P has no  $t = 1$ ; furthermore, after  $P_1$  deviates (i.e., starting at  $t = 2$ )  $P_2$  will switch to  $\sigma^D$  and  $P_1$  has no incentive to not shoot at any  $t \geq 2$ . <sup>[4](#page-9-0)</sup> Hence  $P_1$  is essentially using the strategy  $\sigma_1 = \sigma^D$  and his total expected payoff will then be

$$
V_1 = Q_1 \left( \sigma^D, \tilde{\sigma}^C \right) = 1 + \gamma \left( p_1 \frac{1}{1 - \gamma} + (1 - p_1) \left( 1 + \gamma Q_1 \left( \sigma^D, \sigma^D \right) \right) \right)
$$
  
=  $1 + \gamma \left( p_1 \frac{1}{1 - \gamma} + (1 - p_1) \left( 1 + \gamma \frac{1 - \gamma \left( 1 - p_1 \left( 1 - p_2 \right) \right)}{\left( 1 - \gamma \left( 1 - \gamma \right) \left( 1 - \gamma \left( 1 - p_1 \right) \left( 1 - p_2 \right) \right)} \right) \right)$   
=  $\frac{1 - p_2 (1 - p_1) \gamma^3 - (1 - p_2) (1 - p_1) \gamma}{(1 - \gamma) \left( 1 - p_1 \right) (1 - p_2) \gamma}$ 

Now

$$
(1 - \gamma) \left(Q_1\left(\tilde{\sigma}^C, \tilde{\sigma}^C\right) - Q_1\left(\sigma^D, \tilde{\sigma}^C\right)\right) = (1 - \gamma) \left(V_1^C - V_1\right)
$$

$$
= 1 - \frac{1 - p_2(1 - p_1)\gamma^3 - (1 - p_2)(1 - p_1)\gamma}{1 - \gamma(1 - p_1)(1 - p_2)}
$$

$$
= \frac{\gamma^3 p_2 (1 - p_1)}{1 - \gamma(1 - p_1)(1 - p_2)} > 0
$$

Hence  $P_1$  has no incentive to deviate from  $\tilde{\sigma}^C$ . The same can be proved for  $P_2$ . Consequently  $(\tilde{\sigma}^C, \tilde{\sigma}^C)$  is a NF  $(\tilde{\sigma}^C, \tilde{\sigma}^C)$  is a NE.

In the next proposition the strategy profile is a NE only for "large enough"  $\gamma$ .

**Proposition 4.3** There exist some  $\gamma_0 \in (0,1)$  such that: for all  $\gamma \in (\gamma_0,1)$ , and for all  $K \in \mathbb{N}$ ,  $(\widetilde{\sigma}^{DC(K)}, \widetilde{\sigma}^{DC(K)})$  is a NE of  $\Gamma((1,1), \gamma)$ .

**Proof.** Recall that, when both players use  $\tilde{\sigma}^{DC(K)}$ ,  $P_1$  receives payoff

$$
V_1^{DC(K)} = Q_1((1,1), \tilde{\sigma}^{DC(K)}, \tilde{\sigma}^{DC(K)}) = \frac{1 + \gamma^{K+1}(1 - p_1)^K (1 - p_2)^K p_2 - \gamma (1 - p_1 + p_1 p_2)}{(1 - \gamma)(1 - \gamma (1 - p_1)(1 - p_2))}
$$

Let us show that  $P_1$  has no incentive to use a deviating strategy  $\sigma_1$ .

<span id="page-9-0"></span><sup>&</sup>lt;sup>4</sup>This is a consequence of the following fact, which we will often use in the remainder of the paper. If  $P_n$ starts using a stationary strategy  $\sigma_n$  at some time t, then  $P_{-n}$ 's best response is also a stationary strategy. This is the case because, for a fixed stationary  $\sigma_n$ ,  $P_{-n}$  has to solve a *Markov Decision Process*, for which the optimal strategy is stationary. For more details see [\[21\]](#page-17-18).

1. Case I: Let us first consider strategies which deviate at times  $t \in \{K+1, K+2, ...\}$ ; i.e., they shoot after the game has entered the no-shooting phase. We actually need to consider only  $\sigma_1$  which will shoot at  $t = K + 1$  and with probability one. In this case

$$
Q_{1} ((1,1), \tilde{\sigma}^{DC(K)}, \tilde{\sigma}^{DC(K)}) = A + \gamma^{K+1} Q_{1} ((1,1), \tilde{\sigma}^{DC(K)}, \tilde{\sigma}^{DC(K)})
$$
  
\n
$$
= A + \gamma^{K+1} Q_{1} ((1,1), \sigma^{C}, \sigma^{C}) = A + \gamma^{K+1} V_{1}^{C}
$$
  
\n
$$
Q_{1} ((1,1), \sigma^{1}, \tilde{\sigma}^{DC(K)}) = A + \gamma^{K+1} Q_{1} ((1,1), \sigma^{1}, \tilde{\sigma}^{DC(K)})
$$
  
\n
$$
= A + \gamma^{K+1} Q_{1} ((1,1), \sigma^{1}, \tilde{\sigma}^{DC(K)})
$$

where A is the expected payoff summed over times  $t \in \{0, ..., K\}$  and is the same for both strategies used by P<sub>1</sub>. Now, for the usual reasons, P<sub>1</sub> will keep shooting at  $t \in$  ${K + 2, K + 3, ...}$  and we will have

$$
V_1 = Q_1 ((1,1), \sigma^1, \widetilde{\sigma}^{DC(K)}) = p_1 \frac{1}{1-\gamma} + (1-p_1) (1 + \gamma Q_1 (\sigma^D, \sigma^D))
$$
  
=  $p_1 \frac{1}{1-\gamma} + (1-p_1) (1 + \gamma \frac{1-\gamma(1-p_1(1-p_2))}{(1-\gamma)(1-\gamma(1-p_1)(1-p_2))})$   
=  $\frac{1-p_2(1-p_1)\gamma^2 - (1-p_2)(1-p_1)\gamma}{(1-\gamma)(1-\gamma(1-p_1)(1-p_2))}$ 

Then we have

$$
(1 - \gamma) (Q_1 ((1, 1), \tilde{\sigma}^C, \tilde{\sigma}^C) - Q_1 ((1, 1), \sigma^D, \tilde{\sigma}^C))
$$
  
=  $(1 - \gamma) (V_1^C - V_1)$   
=  $1 - \frac{1 - p_2 (1 - p_1) \gamma^2 - (1 - p_2) (1 - p_1) \gamma}{1 - \gamma (1 - p_1) (1 - p_2)}$   
=  $\frac{p_2 (1 - p_1) \gamma^2}{1 - \gamma (1 - p_1) (1 - p_2)} > 0$ 

Hence  $P_1$  has no incentive to shoot at  $t > K$ .

2. Case II: Let us next consider strategies which deviate at times  $t \in \{1, 2, ..., K\}$ , i.e., they do not shoot during the shooting phase. Again, after the first deviation  $P_1$  has no incentive to not shoot. So we only need to consider strategies  $\sigma_1$  which (a) do not shoot at some  $t = L \in \{1, 2, ..., K\}$  and (b) shoot at all  $t \in \{1, 2, ..., L - 1, L + 1, ...\}$ . Then, by the usual arguments,

$$
V_{1} = Q_{1} ((1, 1), \sigma_{1}, \tilde{\sigma}^{DC(K)})
$$
  
\n
$$
= 1
$$
  
\n
$$
+ \gamma \left( p_{1} (1 - p_{2}) \frac{1}{1 - \gamma} + (1 - p_{1}) (1 - p_{2}) \right)
$$
  
\n
$$
+ \gamma^{2} \left( (1 - p_{1}) p_{1} (1 - p_{2})^{2} \frac{1}{1 - \gamma} + (1 - p_{1})^{2} (1 - p_{2})^{2} \right)
$$
  
\n+ ...  
\n
$$
+ \gamma^{L-1} \left( (1 - p_{1})^{L-2} p_{1} (1 - p_{2})^{L-1} \frac{1}{1 - \gamma} + (1 - p_{1})^{L-1} (1 - p_{2})^{L-1} \right)
$$
  
\n
$$
+ \gamma^{L} \left( (1 - p_{1})^{L-1} p_{1} (1 - p_{2})^{L} \frac{1}{1 - \gamma} + (1 - p_{1})^{L} (1 - p_{2})^{L} (1 + \gamma V_{1}^{D}) \right)
$$
  
\n
$$
= 1 + \sum_{k=1}^{L-1} \gamma^{k} \left( (1 - p_{1})^{k-1} p_{1} (1 - p_{2})^{k} \frac{1}{1 - \gamma} + (1 - p_{1})^{k} (1 - p_{2})^{k} \right)
$$
  
\n
$$
+ \gamma^{L} (1 - p_{1})^{L-1} (1 - p_{2})^{L} p_{1} \frac{1}{1 - \gamma}
$$
  
\n
$$
+ \gamma^{L} (1 - p_{1})^{L} (1 - p_{2})^{L} \left( 1 + \gamma \frac{1 - \gamma (1 - p_{1} (1 - p_{2}))}{(1 - \gamma) (1 - \gamma (1 - p_{1}) (1 - p_{2}))} \right)
$$

Let

$$
\delta v_1(\gamma) = (1 - \gamma) \left( Q_1 \left( (1, 1), \widetilde{\sigma}^{DC(K)}, \widetilde{\sigma}^{DC(K)} \right) - Q_1 \left( (1, 1), \sigma_1, \widetilde{\sigma}^{DC(K)} \right) \right) = V_1^{DC(K)} - V_1.
$$

Note that  $\delta v_1(\gamma)$  is well defined and continuous for all  $\gamma \in [0,1]$ , because the factor  $(1 - \gamma)$ cancels the  $(1 - \gamma)$  factor in the denominator of

$$
Q_1((1,1), \widetilde{\sigma}^{DC(K)}, \widetilde{\sigma}^{DC(K)}) - Q_1((1,1), \sigma_1, \widetilde{\sigma}^{DC(K)})
$$
.

After a considerable amount of algebra<sup>[5](#page-11-0)</sup> we find that

$$
\delta v_1(1) = \frac{\left(1 - p_1\right)^K \left(1 - p_2\right)^K p_2}{p_1 + \left(1 - p_1\right) p_2} > 0.
$$

Since  $\delta v_1(\gamma)$  is continuous, there will exist some  $\gamma_0 \in (0,1)$  such that  $\delta v_1(\gamma)$  will be positive for every  $\gamma \in (\gamma_0, 1)$  and for every  $K \in \mathbb{N}$ . Hence, for such values,  $P_1$  has no incentive to deviate during the shooting phase.

Putting together Cases I and II we see that  $P_1$  has no incentive to deviate from  $\tilde{\sigma}^{DC(K)}$ . The same is proved, similarly, for  $P_2$ . Hence  $(\tilde{\sigma}^{DC(K)}, \tilde{\sigma}^{DC(K)})$  is a NE.

Next we present a negative result: mutual late shooting is not a NE.

<span id="page-11-0"></span><sup>5</sup>Using Maple once again.

**Proposition 4.4** For every  $\gamma \in (0,1)$  and every  $K \in \mathbb{N}$ ,  $(\tilde{\sigma}^{CD(K)}, \tilde{\sigma}^{CD(K)})$  is not a NE of  $\Gamma ((1,1), \gamma)$ .

Proof. Recall that

$$
V_1^{CD(K)} = Q_1\left(\tilde{\sigma}^{CD(K)}, \tilde{\sigma}^{CD(K)}\right) = \frac{1 - \gamma^K p_2 - \gamma(1 - p_1)(1 - p_2)}{(1 - \gamma)(1 - \gamma(1 - p_1)(1 - p_2))}
$$

We just need to show that  $P_1$  has one profitable deviating strategy  $\sigma_1$ . Let  $\sigma_1$  be: do not shoot at  $t \in \{1, 2, ..., K-2\}$ , shoot at  $t \in \{K-1, K, ...\}$ ; in other words start shooting one turn before the shooting phase starts. Then, by the usual arguments,  $P_1$ 's payoff is

$$
V_1 = Q_1((1,1), \sigma_1, \widetilde{\sigma}^{CD(K)}) = \sum_{k=0}^{K-2} \gamma^k + \gamma^{K-1} \left( p_1 \frac{1}{1-\gamma} + (1-p_1)(1+\gamma V_D) \right)
$$
  
= 
$$
\frac{\left( -\gamma^{K+1} p_2 (1-p_1) + 1 - \gamma (1-p_2)(1-p_1) \right)}{(1-\gamma)(1-\gamma(1-p_2)(1-p_1))}
$$

By appropriate substitutions and algebraic calculations, we get

$$
\delta v_1 = (1 - \gamma) \left( Q_1 \left( \tilde{\sigma}^{CD(K)}, \tilde{\sigma}^{CD(K)} \right) - Q_1 \left( \sigma_1, \tilde{\sigma}^{CD(K)} \right) \right)
$$

$$
= -\frac{p_2 \gamma^K \left( 1 - \gamma \left( 1 - p_1 \right) \right)}{(1 - \gamma (1 - p_2)(1 - p_1))} < 0
$$

Hence  $P_1$  has incentive to switch to  $\sigma_1$  and  $(\tilde{\sigma}^{CD(K)}, \tilde{\sigma}^{CD(K)})$  is not a NE.

**Proposition 4.5** For every  $M \in \mathbb{N}$ , there exists a  $\delta_M > 0$  such that: if

$$
\gamma_M = \frac{9}{10},
$$
  
\n
$$
p_M = \frac{1 - e^{-M}}{10},
$$
  
\n
$$
I_M = (\gamma_M - \delta_M, \gamma_M + \delta_M) \times (p_M - \delta_M, p_M + \delta_M) \times (p_M - \delta_M, p_M + \delta_M),
$$

then  $(\tilde{\sigma}^{P(M)}, \tilde{\sigma}^{P(M)})$  is a NE of  $\Gamma((1,1), \gamma, \mathbf{p})$  for every  $(\gamma, p_1, p_2) \in I_M$ .

Proof. Recall that

$$
V^{P(M)} = Q_1 \left( \tilde{\sigma}^{P(M)}, \tilde{\sigma}^{P(M)} \right) = \frac{1 - \gamma^{M+1} (1 - p_1 (1 - p_2))}{(1 - \gamma) (1 - \gamma^{M+1} (1 - p_1) (1 - p_2))}
$$

We will prove that, for every  $(\gamma, p_1, p_2) \in I_M$ ,  $P_1$  has no incentive to deviate from  $\tilde{\sigma}^{P(M)}$  (the proof for P is identical) proof for  $P_2$  is identical).

Suppose that  $P_1$  uses some strategy  $\sigma_1$  by which he shoots at  $P_2$  at some  $t \neq i \cdot (M + 1)$ . For the usual reasons, it suffices to consider strategies by which  $P_1$  shoots in the first period and with probability one. So suppose that  $P_1$  abstains for all  $t \in (1, ..., K)$  and then shoots at  $P_2$  at some  $t' = K + 1 \leq M$ . Then the following two possibilities exist.

- 1. With probability  $p_1$ :  $P_2$  is killed and  $P_1$  receives payoff  $\frac{1}{1-\gamma}$ .
- 2. With probability  $1-p_1: P_2$  is missed,  $P_1$  receives payoff one and for all subsequent rounds  $P_2$  will always shoot at  $P_1$  with probability one. In this case  $P_1$ 's best response at time  $t'' > t'$  is to always shoot at  $P_2$  with probability one; hence, starting at the  $(t' + 1)$ -th round, both players use the  $\sigma^D$  strategy. The total expected payoff received by  $P_1$  in this case is  $Q_1(\sigma^D, \sigma^D)$ .

Hence, assuming  $P_1$  will first shoot at  $t = K + 1 \in \{1, ..., M\}$ , by the above reasoning  $P_1$ 's expected total payoff will be

$$
V_1 = Q_1 \left( \sigma_1, \tilde{\sigma}^{P(M)} \right) = \left( \sum_{k=0}^K \gamma^k \right) + \gamma^{K+1} \left( p_1 \frac{1}{1 - \gamma} + (1 - p_1) Q_1 \left( \sigma^D, \sigma^D \right) \right)
$$

Substituting the  $Q_1(\tilde{\sigma}^{P(M)}, \tilde{\sigma}^{P(M)})$  and  $Q_1(\sigma^D, \sigma^D)$  values and performing a considerable amount of algebra we get of algebra we get

$$
\delta v_1(\gamma, p_1, p_2) = (1 - \gamma) \left( Q_1 \left( \tilde{\sigma}^{P(M)}, \tilde{\sigma}^{P(M)} \right) - Q_1 \left( \sigma_1, \tilde{\sigma}^{P(M)} \right) \right)
$$
  
= 
$$
\frac{p_2 \gamma^{K+2} \left( -(1 - p_1)^2 (1 - p_2) \gamma^{M+1} + (1 - p_1) (1 - p_2) \gamma^{M-K} - \gamma^{M-K-1} + 1 - p_1 \right)}{(1 - \gamma^{M+1} (1 - p_1) (1 - p_2)) (1 - \gamma (1 - p_1) (1 - p_2))}
$$

Setting  $p_1 = p_2 = p$  we get

$$
\delta v_1(\gamma, p, p) = \frac{p\gamma^{K+2} \left(-\left(1-p\right)^3 \gamma^{M+1} + \left(1-p\right)^2 \gamma^{M-K} - \gamma^{M-K-1} + 1-p\right)}{\left(1-\gamma^{M+1} \left(1-p\right)^2\right) \left(1-\gamma \left(1-p\right)^2\right)}
$$

The sign of  $\delta v_1(\gamma, p, p)$  is the same as that of

$$
f_{M,K}(\gamma, p) = -(1-p)^3 \gamma^{M+K+3} + (1-p)^2 \gamma^{M+2} - \gamma^{M+1} + (1-p) \gamma^{K+2}
$$
  
=  $f_{1,M,K}(\gamma, p) + f_{2,M,K}(\gamma, p)$ 

with

$$
f_{M,K,1}(\gamma, p) = (1 - p)^2 \gamma^{M+2} - (1 - p)^3 \gamma^{M+K+3}
$$
  

$$
f_{M,K,2}(\gamma, p) = -\gamma^{M+1} + (1 - p) \gamma^{K+2}
$$

Now we consider the following cases.

1. **Case I:**  $K \leq M - 2$ . Then  $M - K - 1 \geq 1$ . For all M and  $K \in \{1, ..., M - 2\}$  we have  $(1-p)^2 > (1-p)^3$  and  $\gamma^{M+2} > \gamma^{M+K+3}$ 

hence we will always have  $f_{1,M,K}(\gamma, p) > 0$ . To also have  $f_{M,K,2}(\gamma, p) > 0$  for a specific K, it suffices that

$$
\gamma^{M-K-1} < 1 - p \Leftrightarrow \gamma < (1 - p)^{\frac{1}{M-K-1}}
$$

To have  $f_{M,K,2}(\gamma, p) > 0$  for all  $K \in \{1, ..., M-2\}$ , it suffices that

<span id="page-14-0"></span>
$$
\gamma < (1-p)^{\frac{1}{M-2}}\tag{7}
$$

In other words, for all  $M$  we have:

$$
\gamma \in \left(0, (1-p)^{\frac{1}{M-2}}\right) \Rightarrow (\forall K \in \{1, ..., M-2\} : f_{M,K}(\gamma, p) > 0)
$$
  

$$
\gamma \in \left(0, (1-p)^{\frac{1}{M-2}}\right) \Rightarrow (\forall K \in \{1, ..., M-2\} : \delta v_1(\gamma, p, p) > 0)
$$

2. Case II:  $K = M - 1$ . In this case we want

$$
-(1-p)^{3} \gamma^{M+K+3} + (1-p)^{2} \gamma^{M+2} - \gamma^{M+1} + (1-p) \gamma^{K+2} > 0 \Leftrightarrow
$$
  

$$
-(1-p)^{3} \gamma^{M+M-1+3} + (1-p)^{2} \gamma^{M+2} - \gamma^{M+1} + (1-p) \gamma^{M-1+2} > 0 \Leftrightarrow
$$
  

$$
-(1-p)^{3} \gamma^{2M+2} + (1-p)^{2} \gamma^{M+2} - \gamma^{M+1} + (1-p) \gamma^{M+1} > 0 \Leftrightarrow
$$
  

$$
-(1-p)^{3} \gamma^{M+1} + (1-p)^{2} \gamma - 1 + (1-p) > 0 \Leftrightarrow
$$
  

$$
-(1-p)^{3} \gamma^{M+1} + (1-p)^{2} \gamma - p > 0 \Leftrightarrow
$$
  

$$
-(1-p)^{3} \gamma^{M+1} + \gamma p^{2} - (2\gamma + 1) p + \gamma > 0
$$

Let us define the function

$$
\overline{f}_{M}(\gamma, p) = -(1-p)^{3} \gamma^{M+1} + \gamma p^{2} - (2\gamma + 1) p + \gamma
$$

By continuity, in a sufficiently small neighborhood of  $(\gamma_M, p_M) = \left(\frac{9}{10}, \frac{1-e^{-M}}{10}\right)$ , the sign of  $\overline{f}_M\left(\gamma,p\right)$  will be the same as that of

$$
h_1(M) = \overline{f}_M(\gamma_M, p_M) = -\frac{\left(9 + e^{-M}\right)^3 \left(\frac{9}{10}\right)^{M+1}}{1000} + \frac{9\left(1 - e^{-M}\right)^2}{1000} + \frac{31}{50} + \frac{7e^{-M}}{25}
$$

and it suffices to show that  $h_1(M) > 0$  for all M. To this end we first note that

$$
h_1(1) = -\frac{\left(9 + e^{-1}\right)^3 \left(\frac{9}{10}\right)^2}{1000} + \frac{9 \left(1 - e^{-1}\right)^2}{1000} + \frac{31}{50} + \frac{7e^{-1}}{25} = 0.60703... > 0
$$

Also, letting

$$
h_2\left(M\right) = -\frac{\left(9 + e^{-M}\right)^3 \left(\frac{9}{10}\right)^{M+1}}{1000} + \frac{31}{50}
$$

we have

$$
\forall M: h_1(M) > h_2(M).
$$

Now,  $h_2(M)$  is strictly increasing in M and  $h_2(2) = 0.064222...$ . Consequently

$$
\forall M \ge 2 : h_1(M) > h_2(M) > h_2(2) > 0
$$

Hence finally we have

$$
\forall M \geq 1 : \overline{f}_M(\gamma_M, p_M) = h_1(M) > 0.
$$

Now, to have

$$
\forall M, \forall K \in \{1, ..., M-1\} : f_{M,K}(\gamma_M, p_M) > 0
$$

we must ensure that [\(7\)](#page-14-0) holds for  $(\gamma, p) = (\gamma_M, p_M)$ . In other words, we want  $\gamma_M < (1 - p_M)^{\frac{1}{M-2}}$ or, equivalently,

$$
\frac{9}{10} < \left(1 - \frac{1 - e^{-M}}{10}\right)^{\frac{1}{M-2}} = \left(\frac{9}{10} + \frac{e^{-M}}{10}\right)^{\frac{1}{M-2}}.
$$

This holds: since for all  $M \in \mathbb{N}$  we have  $\frac{9}{10} + \frac{e^{-M}}{10} < 1$ , we also have

$$
\frac{9}{10} < \left(\frac{9}{10} + \frac{e^{-M}}{10}\right) < \left(\frac{9}{10} + \frac{e^{-M}}{10}\right)^{\frac{1}{M-2}}.
$$

In short we have shown that

$$
\forall M, \forall K \in \{1, ..., M - 1\} : f_{M,K}(\gamma_M, p_M) > 0
$$
  

$$
\forall M, \forall K \in \{1, ..., M - 1\} : \delta v_1(\gamma_M, p_M, p_M) > 0
$$

For all M and K,  $\delta v_1(\gamma, p_1, p_2)$  is a continuous function. Hence, for all M, there exists some  $\delta_M > 0$  such that

$$
\forall K \in \{1, ..., M - 1\}, \forall (\gamma, p_1, p_2) \in I_M : \delta v_1 (\gamma, p_1, p_2) > 0
$$

which shows that  $P_1$  has no incentive to deviate from  $\tilde{\sigma}^{P(M)}$ . The same argument can be applied<br>to  $P$  Hones for every  $(\alpha, p, p) \in L$ ,  $(\tilde{\sigma}^{P(M)} | \tilde{\sigma}^{P(M)})$  is a NF of  $\Gamma((1, 1), \alpha, p)$ to  $P_2$ . Hence, for every  $(\gamma, p_1, p_2) \in I_M$ ,  $(\widetilde{\sigma}^{P(M)}, \widetilde{\sigma}^{P(M)})$  is a NE of  $\Gamma((1, 1), \gamma, \mathbf{p})$ .

### <span id="page-15-0"></span>5 Some Additional Remarks

Let us now justify our terms "cooperating" and "defecting" strategy. From the results of Section [3,](#page-3-0) for  $n \in \{1, 2\}$ , we have

$$
\overline{Q}_{n}((1,1), \sigma^{C}, \sigma^{C}) = \frac{1}{1-\gamma}
$$

$$
\overline{Q}_{n}((1,1), \sigma^{D}, \sigma^{D}) = \frac{1-\gamma(1-p_{n}(1-p_{-n}))}{(1-\gamma)(1-\gamma(1-p_{1})(1-p_{2}))}.
$$

It follows that

$$
\overline{Q}_{n}\left(\left(1,1\right),\sigma^{C},\sigma^{C}\right)-\overline{Q}_{n}\left(\left(1,1\right),\sigma^{D},\sigma^{D}\right)=\frac{\gamma p_{-n}}{\left(1-\gamma\right)\left(1-\gamma\left(1-p_{1}\right)\left(1-p_{2}\right)\right)}>0.
$$

In short, just like in PD, it is more profitable for both players to not-shoot rather than shoot. Because in our formulation there is no direct profit from killing the opponent, both  $(\sigma^C, \sigma^C)$ and  $(\sigma^D, \sigma^D)$  are NE; however, for both players,  $(\sigma^C, \sigma^C)$  is more profitable NE than  $(\sigma^D, \sigma^D)$ . This is the reason for calling  $(\sigma^C, \sigma^C)$  a cooperating, and  $(\sigma^D, \sigma^D)$  a defecting strategy.

All this may be surprising, since one would expect that, in a duel, each player's goal will be to eliminate his opponent. It may be supposed that the higher profitability of  $(\sigma^C, \sigma^C)$ 

follows from our choice of not assigning any direct payoff to killing one's opponent. But this is not true. Even with a positive "killing payoff",  $\overline{Q}_n((1,1), \sigma^C, \sigma^C)$  can still be greater than  $\overline{Q}_n((1,1), \sigma^D, \sigma^D)$ , provided  $\gamma$  is sufficiently close to one<sup>[6](#page-16-3)</sup>. The reason for the superiority of  $(\sigma^C, \sigma^C)$  is this: if a positive payoff is assigned to survival, this, compounded over an infinite number of turns, can always outweigh the killing payoff. Hence our model can be understood as a more "pacifist" version than the usual duel model[7](#page-16-4) .

Let us now compare our static duel to the PD. In both the PD and the duel, cooperation is more profitable than defection. While  $(\sigma^C, \sigma^C)$  is not a NE in PD,  $(\sigma^D, \sigma^D)$  is a NE in both of them. However, both the duel and the repeated version of PD, possess several NE in grim strategies; the common characteristic of all such equilibria is that they promote cooperation or, in other words, punish defection (shooting). In fact, similarly to the case of repeated PD, it might be possible to prove a "Folk Theorem" for the static duel as well; namely that every feasible and individually rational payoff is a NE for  $\gamma$  sufficiently close to one. We intend to study this question in the future.

## <span id="page-16-2"></span>6 Conclusion

We have formulated the simultaneous shooting static duel as a discounted stochastic game. We have shown that it has two Nash equilibria in stationary strategies, namely the "always-shooting" and the "never-shooting" strategies; in addition several nonstationary, "cooperation-promoting" Nash equilibria also exist. In the future we intend to extend the study of the static duel in several directions.

First, we want to extend our study and obtain similar results for two variants: (a) the case of non-zero killing payoff and (b) the case of terminal-only payoffs. In addition, we want to formulate and study a version of the static duel in which each player wants to kill his opponent in the shortest possible time.

Secondly, we hope to prove a form of "Folk Theorem", namely that every every feasible and individually rational payoff is a NE for  $\gamma$  sufficiently close to one.

Finally, we want to formulate the static *Nuel* (i.e., the duel-like game which involves N players shooting at each other) as a discounted stochastic game and extend our results for this case.

## References

- <span id="page-16-0"></span>[1] Amengual, P. and Toral, R. "Distribution of winners in truel games". AIP Conference Proceedings (2005), pp. 128–141.
- <span id="page-16-1"></span>[2] Amengual, P. and Toral, R. "A Markov chain analysis of truels". Proceedings of the 8th Granada Seminar on Computational Physics (2005).

<span id="page-16-3"></span><sup>6</sup>This, as well as additional results regarding the positive killing payoff case, will be reported in a future publication.

<span id="page-16-4"></span><sup>7</sup>This point has also been raised by Donald Knuth in the context of the truel [\[17\]](#page-17-5). For example, he remarks that "a player who passes is guaranteeing that his opponent has no reason to shoot back, as far as the opponent's survival is concerned ".

- <span id="page-17-9"></span>[3] Amengual, P. and Toral, R. "Truels, or survival of the weakest." Computing in Science.
- <span id="page-17-15"></span>[4] Barron, E. N. Game theory: an introduction. John Wiley & Sons (2013).
- <span id="page-17-10"></span>[5] Bossert, W., Brams S.J. and Kilgour, D.M. "Cooperative vs non-cooperative truels: little agreement, but does that matter?." Games and Economic Behavior, vol. 40 (2002), pp. 185-202.
- <span id="page-17-11"></span>[6] Brams, S.J., and Kilgour, D.M. "The truel". Mathematics Magazine, vol. 70 (1997), pp. 315-326.
- <span id="page-17-12"></span>[7] Brams, S.J., and Kilgour, D.M. "Games that End in a Bang or a Whimper". preprint, CV Starr Center for Applied Economics (2001).
- <span id="page-17-13"></span>[8] Brams, S.J., Kilgour, D.M. and Dawson, B. "Truels and the Future." Math Horizons, vol. 10 (2003), pp. 5-8.
- <span id="page-17-14"></span>[9] Dorraki, M., Allison, A. and Abbott, D.. "Truels and strategies for survival." Scientific Reports, vol. 9 (2019), pp. 1-7.
- <span id="page-17-16"></span>[10] Dresher, M. Games of strategy: theory and applications. Rand Corp. (1961).
- <span id="page-17-0"></span>[11] Gardner, M. : New Mathematical Puzzles and Diversions. (1966), pp. 42-49.
- <span id="page-17-17"></span>[12] Karlin, S., Mathematical Methods and Theory in Games, Programming, and Economics, vol. 2, Addison- Wesley (1959).
- <span id="page-17-6"></span>[13] Kilgour, D. M. "The simultaneous truel." *International Journal of Game Theory*, vol. 1 (1971), pp. 229-242.
- <span id="page-17-7"></span>[14] Kilgour, D. M. "The sequential truel." International Journal of Game Theory, vol. 4 (1975), pp. 151-174.
- <span id="page-17-8"></span>[15] Kilgour, D. M. "Equilibrium points of infinite sequential truels." International Journal of Game Theory, vol. 6 (1977), pp. 167-180.
- <span id="page-17-1"></span>[16] Kinnaird, C. Encyclopedia of puzzles and pastimes. Secaucus (1946).
- <span id="page-17-5"></span>[17] Knuth, D.E. "The Triel: A New Solution." Journal of Recreational Mathematics, vol. 6 (1972), pp. 1-7.
- <span id="page-17-2"></span>[18] Larsen, H.D. "A Dart Game." American Mathematical Monthly, (1948) pp. 640-41.
- <span id="page-17-3"></span>[19] Mosteller, F. Fifty challenging problems in probability with solutions. Courier Corporation (1987).
- <span id="page-17-4"></span>[20] Shubik, M. "Does the Fittest Necessarily Survive?", Readings in Game Theory and Political Behavior, Doubleday.
- <span id="page-17-18"></span>[21] Sobel, M.J. "Noncooperative stochastic games." The Annals of Mathematical Statistics, vol. 42 (1971), pp. 1930-1935.
- <span id="page-18-1"></span>[22] Toral, R., and Amengual, P.. "Distribution of winners in truel games." AIP Conference Proceedings, vol. 779. American Institute of Physics (2005).
- <span id="page-18-2"></span>[23] Wegener, M. and Mutlu, E.. "The good, the bad, the well-connected." International Journal of Game Theory, vol. 50 (2021), pp. 759-771.
- <span id="page-18-3"></span>[24] Xu, X. "Game of the truel." Synthese, vol. 185 (2012), pp. 19-25.
- <span id="page-18-0"></span>[25] Zeephongsekul, P. "Nash Equilibrium Points Of Stochastic N-Uels." Recent Developments in Mathematical Programming. CRC Press (1991), pp. 425-452.