

# SPECTRAL ANALYSIS OF AN OPEN $q$ -DIFFERENCE TODA CHAIN WITH TWO-SIDED BOUNDARY INTERACTIONS ON THE FINITE INTEGER LATTICE

JAN FELIPE VAN DIEJEN

ABSTRACT. A quantum  $n$ -particle model consisting of an open  $q$ -difference Toda chain with two-sided boundary interactions is placed on a finite integer lattice. The spectrum and eigenbasis are computed by establishing the equivalence with a previously studied  $q$ -boson model from which the quantum integrability is inherited. Specifically, the  $q$ -boson-Toda correspondence in question yields Bethe Ansatz eigenfunctions in terms of hyperoctahedral Hall-Littlewood polynomials and provides the pertinent solutions of the Bethe Ansatz equations via the global minima of corresponding Yang-Yang type Morse functions.

## 1. INTRODUCTION

The relativistic Toda chain is an ubiquitous one-dimensional  $n$ -particle model introduced by Ruijsenaars that is integrable both at the level of classical and quantum mechanics [R90]. In the case of an open chain, integrable perturbations at the boundary were implemented via the boundary Yang-Baxter equation [KT96, S90]. At the quantum level, the hamiltonian of the relativistic Toda chain is given by a ( $q$ -)difference operator. Quantum groups connect the difference operator at issue to the quantum  $K$ -theory of flag manifolds [BF05, GL03] and provide a natural representation-theoretical habitat for the construction of its eigenfunctions [E99, S00].

When considering the quantum dynamics on an integer lattice the eigenvalue problem for the  $q$ -difference Toda chain can be solved in terms of  $q$ -Whittaker functions that arise as a parameter specialization of the Macdonald polynomials, both in the case of particles moving on an infinite lattice [GLO11] and in the case of particles moving on a finite periodic lattice [DP16]. From the perspective of integrable probability, such particle models are of interest in connection with the  $q$ -Whittaker process [BC14]. Representation-theoretical constructions for the pertinent  $q$ -Whittaker functions can be found in [DK18, DKT17, FJM09].

This note addresses the spectral problem for an open  $n$ -particle  $q$ -difference Toda chain on the finite lattice  $\{0, 1, 2, \dots, m\}$  that is endowed with two-parameter boundary interactions on both ends. The model could be thought of as a finite discrete and  $q$ -deformed counterpart of Sklyanin's open quantum Toda chain with general two-sided boundary perturbations governed by Morse potentials [S88]. In the limit  $m \rightarrow \infty$  the pertinent  $q$ -difference Toda hamiltonian was diagonalized in

---

*Date:* January 2023.

*2010 Mathematics Subject Classification.* Primary: 33D52; Secondary: 05E05, 81Q35, 81Q80, 81U15, 82B23.

*Key words and phrases.*  $q$ -difference Toda chain, Bethe Ansatz, Hall-Littlewood polynomials.

terms of hyperoctahedral  $q$ -Whittaker functions that arise in turn through a parameter specialization of the Macdonald-Koornwinder polynomials [DE15]. Here it will be shown that for finite  $m$  an explicit eigenbasis can be constructed from Bethe Ansatz wave functions given by Macdonald's hyperoctahedral Hall-Littlewood polynomials [M00]. The main idea is to exploit an equivalence between  $q$ -difference Toda chains and  $q$ -boson models pointed out in [DP16]. By establishing a version of this equivalence in the current situation of an open chain with boundary perturbations, our  $q$ -difference Toda hamiltonian is mapped to the hamiltonian of a  $q$ -boson model previously diagonalized in [DEZ18]. The upshot is that the commuting quantum integrals and the Bethe Ansatz eigenfunctions for the  $q$ -difference Toda chain can in this approach be retrieved directly from those in [DEZ18] for the corresponding  $q$ -boson model.

The material is organized as follows. Section 2 describes the hamiltonian of our  $q$ -difference Toda chain and verifies its self-adjointness. Section 3 establishes the equivalence with the  $q$ -boson model from [DEZ18] and therewith retrieves the corresponding Bethe Ansatz wave functions in terms of hyperoctahedral Hall-Littlewood polynomials. The Bethe Ansatz equations of interest are of a convex type studied in wider generality in [DE19], which entails an explicit description of the spectrum via the global minima of associated Yang-Yang-type Morse functions detailed in Section 4. The presentation closes in Section 5 with a description of the spectral analysis for the  $q$ -difference Toda chain in the degenerate limit  $q \rightarrow 1$ .

## 2. OPEN $q$ -DIFFERENCE TODA CHAIN WITH BOUNDARY INTERACTIONS

**2.1. Quantum hamiltonian.** Given  $m, n \in \mathbb{N}$ , the  $q$ -difference Toda chain under consideration describes the quantum dynamics of  $n$  interacting particles hopping over the finite integer lattice  $\{0, 1, 2, \dots, m\}$ . The positions of these particles are encoded by a partition  $\mu = (\mu_1, \mu_2, \dots, \mu_n)$  in the configuration space

$$\Lambda^{(n,m)} = \{\mu \in \mathbb{Z}^n \mid m \geq \mu_1 \geq \mu_2 \geq \dots \geq \mu_n \geq 0\}. \quad (2.1)$$

The dynamics is governed in turn by the following quantum hamiltonian

$$\begin{aligned} H &= \beta_+(1 - q^{m-\mu_1}) + \beta_-(1 - q^{\mu_n}) \\ &+ \sum_{1 \leq i \leq n} (1 - \alpha_+ q^{m-\mu_i-1})^{\delta_{i-1}} (1 - q^{\mu_{i-1}-\mu_i}) T_i \\ &+ \sum_{1 \leq i \leq n} (1 - \alpha_- q^{\mu_n-1})^{\delta_{n-i}} (1 - q^{\mu_i-\mu_{i+1}}) T_i^{-1}, \end{aligned} \quad (2.2)$$

with

$$\delta_i = \begin{cases} 1 & \text{if } i = 0, \\ 0 & \text{if } i \neq 0, \end{cases} \quad \begin{cases} \mu_0 \equiv m, \\ \mu_{n+1} \equiv 0. \end{cases}$$

Here  $T_i$  and  $T_i^{-1}$  denote hopping operators that act on  $n$ -particle wave functions  $\psi$  via a unit translation of the  $i$ th particle to the left and to the right, respectively:

$$(T_i^\epsilon \psi)(\mu_1, \dots, \mu_n) = \psi(\mu_1, \dots, \mu_{i-1}, \mu_i + \epsilon, \mu_{i+1}, \dots, \mu_n) \quad (\epsilon \in \{1, -1\}).$$

The action of  $H$  (2.2) on wave functions  $\psi : \Lambda^{(n,m)} \rightarrow \mathbb{C}$  is well-defined in the sense that the coefficient of  $(T_i^\epsilon \psi)(\mu_1, \dots, \mu_n)$  in  $(H\psi)(\mu_1, \dots, \mu_n)$  vanishes for any  $(\mu_1, \dots, \mu_n) \in \Lambda^{(n,m)}$  such that  $(\mu_1, \dots, \mu_{i-1}, \mu_i + \epsilon, \mu_{i+1}, \dots, \mu_n) \notin \Lambda^{(n,m)}$ . Notice also that the convention in the second brace below Eq. (2.2) can be interpreted as representing the positions of two additional particles fixed at the lattice end-points

0 and  $m$ , respectively. The parameter  $q \in (-1, 1) \setminus \{0\}$  denotes a scale parameter of the model governing the nearest neighbour interaction between the particles whereas the parameters  $\alpha_{\pm} \in (-1, 1)$  and  $\beta_{\pm} \in \mathbb{R}$  represent coupling constants regulating additional interactions at the boundary of the chain.

Our main goal is to solve the spectral problem for the  $q$ -difference Toda hamiltonian  $H$  (2.2) in the  $\binom{n+m}{n}$ -dimensional Hilbert space  $\ell^2(\Lambda^{(n,m)}, \Delta)$  of functions  $\psi : \Lambda^{(n,m)} \rightarrow \mathbb{C}$ , endowed with an inner product

$$\langle \psi, \phi \rangle_{\Delta} = \sum_{\mu \in \Lambda^{(n,m)}} \psi(\mu) \overline{\phi(\mu)} \Delta_{\mu} \quad \left( \psi, \phi \in \ell^2(\Lambda^{(n,m)}, \Delta) \right) \quad (2.3a)$$

determined by positive weights given by perturbed  $q$ -multinomials on  $\Lambda^{(n,m)}$ :

$$\Delta_{\mu} = \frac{(q; q)_m}{(\alpha_+; q)_{m-\mu_1} (\alpha_-; q)_{\mu_n} \prod_{0 \leq i \leq n} (q; q)_{\mu_i - \mu_{i+1}}} \quad (2.3b)$$

( $\mu \in \Lambda^{(n,m)}$ ). Here we have employed the standard  $q$ -shifted factorial

$$(a; q)_l = \begin{cases} 1 & \text{if } l = 0 \\ (1-a)(1-aq) \cdots (1-aq^{l-1}) & \text{if } l = 1, 2, 3, \dots \end{cases}$$

**Proposition 2.1** (Self-adjointness). *For  $\alpha_{\pm} \in (-1, 1)$ ,  $\beta_{\pm} \in \mathbb{R}$ , and  $q \in (-1, 1) \setminus \{0\}$ , the  $q$ -difference Toda hamiltonian  $H$  (2.2) is self-adjoint in  $\ell^2(\Lambda^{(n,m)}, \Delta)$ , i.e.*

$$\forall \psi, \phi \in \ell^2(\Lambda^{(n,m)}, \Delta) : \quad \langle H\psi, \phi \rangle_{\Delta} = \langle \psi, H\phi \rangle_{\Delta}.$$

*Remark 2.2.* For  $m \rightarrow \infty$ , the  $q$ -difference Toda hamiltonian  $H$  (2.2) was diagonalized in [DE15, Section 7] in terms of a unitary eigenfunction transform with a  $q$ -Whittaker kernel built from a parameter specialization of the Macdonald-Koornwinder polynomials.

**2.2. Proof of Proposition 2.1.** The action of  $H$  (2.2) on  $\psi \in \ell^2(\Lambda^{(n,m)}, \Delta)$  is of the form

$$\begin{aligned} (H\psi)(\mu) &= (\beta_+(1 - q^{m-\mu_1}) + \beta_-(1 - q^{\mu_n}))\psi(\mu) \\ &+ \sum_{\substack{1 \leq i \leq n \\ \mu + e_i \in \Lambda^{(n,m)}}} (1 - \alpha_+ q^{m-\mu_1-1})^{\delta_{i-1}} (1 - q^{\mu_{i-1}-\mu_i})\psi(\mu + e_i) \\ &+ \sum_{\substack{1 \leq i \leq n \\ \mu - e_i \in \Lambda^{(n,m)}}} (1 - \alpha_- q^{\mu_n-1})^{\delta_{n-i}} (1 - q^{\mu_i-\mu_{i+1}})\psi(\mu - e_i), \end{aligned}$$

where the vectors  $e_1, \dots, e_n$  represent the standard unit basis for  $\mathbb{Z}^n$ .

Since all coefficients of the difference operator in question are real, the asserted symmetry  $\langle H\psi, \phi \rangle_{\Delta} = \langle \psi, H\phi \rangle_{\Delta}$  is immediate from the following bilinear identity

$\forall \psi, \phi \in \ell^2(\Lambda^{(n,m)}, \Delta)$ :

$$\begin{aligned} & \sum_{\mu \in \Lambda^{(n,m)}} \left( \sum_{\substack{1 \leq i \leq n \\ \mu + e_i \in \Lambda^{(n,m)}}} (1 - \alpha_+ q^{m-\mu_1-1})^{\delta_{i-1}} (1 - q^{\mu_i-1-\mu_i}) \psi(\mu + e_i) \right) \phi(\mu) \Delta_\mu \\ \stackrel{(i)}{=} & \sum_{\tilde{\mu} \in \Lambda^{(n,m)}} \sum_{\substack{1 \leq i \leq n \\ \tilde{\mu} - e_i \in \Lambda^{(n,m)}}} (1 - \alpha_+ q^{m-\tilde{\mu}_1})^{\delta_{i-1}} (1 - q^{\tilde{\mu}_i-1-\tilde{\mu}_i+1}) \psi(\tilde{\mu}) \phi(\tilde{\mu} - e_i) \Delta_{\tilde{\mu} - e_i} \\ \stackrel{(ii)}{=} & \sum_{\tilde{\mu} \in \Lambda^{(n,m)}} \psi(\tilde{\mu}) \left( \sum_{\substack{1 \leq i \leq n \\ \tilde{\mu} - e_i \in \Lambda^{(n,m)}}} (1 - \alpha_- q^{\tilde{\mu}_n-1})^{\delta_{n-i}} (1 - q^{\tilde{\mu}_i-\tilde{\mu}_i+1}) \phi(\tilde{\mu} - e_i) \right) \Delta_{\tilde{\mu}}. \end{aligned}$$

Step (i) hinges on the substitution  $\mu = \tilde{\mu} - e_i$ , which for a given  $i \in \{1, \dots, n\}$  determines a bijection from the subset  $\{\tilde{\mu} \in \Lambda^{(n,m)} \mid \tilde{\mu} - e_i \in \Lambda^{(n,m)}\}$  onto the subset  $\{\mu \in \Lambda^{(n,m)} \mid \mu + e_i \in \Lambda^{(n,m)}\}$ . Step (ii) uses the elementary recurrence

$$(1 - \alpha_+ q^{m-\tilde{\mu}_1})^{\delta_{i-1}} (1 - q^{\tilde{\mu}_i-1-\tilde{\mu}_i+1}) \Delta_{\tilde{\mu} - e_i} = (1 - \alpha_- q^{\tilde{\mu}_n-1})^{\delta_{n-i}} (1 - q^{\tilde{\mu}_i-\tilde{\mu}_i+1}) \Delta_{\tilde{\mu}}$$

for  $\tilde{\mu} \in \Lambda^{(n,m)}$  such that  $\tilde{\mu} - e_i \in \Lambda^{(n,m)}$  (and the convention  $\tilde{\mu}_0 \equiv m$ ,  $\tilde{\mu}_{n+1} \equiv 0$ ).

### 3. EIGENFUNCTIONS

**3.1. Bethe Ansatz.** While Proposition 2.1 implies that the existence of an orthogonal eigenbasis diagonalizing  $H$  (2.2) in  $\ell^2(\Lambda^{(n,m)}, \Delta)$  is evident from the spectral theorem for self-adjoint operators in finite dimension, the aim here is to provide an *explicit* eigenbasis given by Bethe Ansatz wave functions in the spirit of [D21] for  $n = 1$ .

To this end let us recall that for any  $\lambda = (\lambda_1, \dots, \lambda_m) \in \Lambda^{(m,n)}$  and  $\xi = (\xi_1, \dots, \xi_m)$  belonging to

$$\mathbb{R}_{\text{reg}}^m = \{\xi \in \mathbb{R}^m \mid 2\xi_j, \xi_j - \xi_k, \xi_j + \xi_k \notin 2\pi\mathbb{Z}, \forall 1 \leq j \neq k \leq m\}, \quad (3.1)$$

Macdonald's hyperoctahedral Hall-Littlewood polynomial (associated with the root system  $BC_m$ ) is given by [M00, §10]

$$R_\lambda(\xi_1, \dots, \xi_m) = \quad (3.2a)$$

$$\sum_{\substack{\sigma \in S_m \\ \epsilon \in \{1, -1\}^m}} C(\epsilon_1 \xi_{\sigma(1)}, \dots, \epsilon_m \xi_{\sigma(m)}) \exp(i\epsilon_1 \xi_{\sigma(1)} \lambda_1 + \dots + i\epsilon_n \xi_{\sigma(m)} \lambda_m)$$

with

$$\begin{aligned} C(\xi_1, \dots, \xi_m) &= \prod_{1 \leq j \leq m} \frac{(1 - \beta_+ e^{-i\xi_j} + \alpha_+ e^{-2i\xi_j})}{1 - e^{-2i\xi_j}} \\ &\times \prod_{1 \leq j < k \leq m} \left( \frac{1 - qe^{-i(\xi_j - \xi_k)}}{1 - e^{-i(\xi_j - \xi_k)}} \right) \left( \frac{1 - qe^{-i(\xi_j + \xi_k)}}{1 - e^{-i(\xi_j + \xi_k)}} \right), \end{aligned} \quad (3.2b)$$

where the summation is over all permutations  $\sigma = (\sigma(1) \ \sigma(2) \ \dots \ \sigma(m))$  of the symmetric group  $S_m$  and all sign configurations  $\epsilon = (\epsilon_1, \dots, \epsilon_m) \in \{1, -1\}^m$ .

For any  $\lambda \in \Lambda^{(m,n)}$  and  $0 \leq i \leq n$  we denote the multiplicity of  $i$  in  $\lambda$  by

$$m_i(\lambda) = |\{1 \leq j \leq m \mid \lambda_j = i\}|. \quad (3.3a)$$

Additionally, for  $\mu \in \Lambda^{(n,m)}$  we write

$$\mu' = (0^{m-\mu_1} 1^{\mu_1-\mu_2} 2^{\mu_2-\mu_3} \dots (n-1)^{\mu_{n-1}-\mu_n} n^{\mu_n}) \quad (3.3b)$$

for its conjugate partition  $\mu' \in \Lambda^{(m,n)}$  (i.e. ‘with the columns and rows swapped’). In other words,  $\mu' = (\mu'_1, \mu'_2, \dots, \mu'_m)$  is the (unique) partition in  $\Lambda^{(m,n)}$  such that  $m_i(\mu') = \mu_i - \mu_{i+1}$  for  $i = 0, \dots, n$  (where, recall,  $\mu_0 \equiv m$  and  $\mu_{n+1} \equiv 0$ ). Notice in this connection that  $|\Lambda^{(n,m)}| = |\Lambda^{(m,n)}| = \frac{(n+m)!}{n!m!}$  and that the mapping  $\mu \rightarrow \mu'$  (3.3b) defines a bijection from  $\Lambda^{(n,m)}$  onto  $\Lambda^{(m,n)}$ .

From now on it will moreover be assumed (unless explicitly stated otherwise) that the boundary parameters  $\alpha_{\pm}, \beta_{\pm}$  have values such that the roots  $p_{\pm}, q_{\pm}$  of the two quadratic polynomials  $x^2 - \beta_{\pm}x + \alpha_{\pm}$  belong to the interval  $(-1, 1) \setminus \{0\}$ :

$$\boxed{\alpha_{\pm} = p_{\pm}q_{\pm} \quad \text{and} \quad \beta_{\pm} = p_{\pm} + q_{\pm} \quad \text{with} \quad q_{\pm}, p_{\pm} \in (-1, 1) \setminus \{0\}} \quad (3.4a)$$

or equivalently

$$\boxed{0 < \alpha_{\pm}^2 < 1 \quad \text{and} \quad 4\alpha_{\pm} \leq \beta_{\pm}^2 \leq (1 + \alpha_{\pm})^2.} \quad (3.4b)$$

**Theorem 3.1** (Bethe Ansatz Wave Function). *Let  $q \in (-1, 1) \setminus \{0\}$  and let the boundary parameters  $\alpha_{\pm}, \beta_{\pm}$  belong to the domain specified in Eqs. (3.4a), (3.4b). Given  $\xi = (\xi_1, \dots, \xi_m) \in \mathbb{R}_{\text{reg}}^m$  (3.1), we define the wave function  $\psi_{\xi} \in \ell^2(\Lambda^{(n,m)}, \Delta)$  through its values on  $\Lambda^{(n,m)}$  as follows:*

$$\psi_{\xi}(\mu) = R_{\mu'}(\xi_1, \dots, \xi_m) \quad (\mu \in \Lambda^{(n,m)}). \quad (3.5a)$$

The wave function  $\psi_{\xi}$  (3.5a) solves the eigenvalue equation for the  $q$ -difference Toda hamiltonian  $H$  (2.2)

$$H\psi_{\xi} = E(\xi)\psi_{\xi} \quad \text{with} \quad E(\xi) = 2(1-q) \sum_{1 \leq j \leq m} \cos(\xi_j), \quad (3.5b)$$

provided the spectral parameter  $\xi \in \mathbb{R}_{\text{reg}}^m$  satisfies the algebraic system of Bethe Ansatz equations

$$e^{2in\xi_j} = \frac{(1 - \beta_+ e^{i\xi_j} + \alpha_+ e^{2i\xi_j})(1 - \beta_- e^{i\xi_j} + \alpha_- e^{2i\xi_j})}{(e^{2i\xi_j} - \beta_+ e^{i\xi_j} + \alpha_+)(e^{2i\xi_j} - \beta_- e^{i\xi_j} + \alpha_-)} \quad (3.5c)$$

$$\times \prod_{\substack{1 \leq k \leq m \\ k \neq j}} \frac{(1 - qe^{i(\xi_j - \xi_k)})(1 - qe^{i(\xi_j + \xi_k)})}{(e^{i(\xi_j - \xi_k)} - q)(e^{i(\xi_j + \xi_k)} - q)} \quad \text{for } j = 1, \dots, m.$$

*Remark 3.2.* The hyperoctahedral Hall-Littlewood polynomial  $R_{\lambda}(\xi_1, \dots, \xi_m)$ ,  $\lambda \in \Lambda^{(m,n)}$  is in fact a symmetric polynomial in  $\cos(\xi_1), \dots, \cos(\xi_m)$  of total degree  $\lambda_1 + \lambda_2 + \dots + \lambda_n$ . Hence, it is clear that the Bethe Ansatz wave function  $\psi_{\xi}$  in Theorem 3.1 extends smoothly in the spectral parameter  $\xi$  from values in  $\mathbb{R}_{\text{reg}}^m$  to values in  $\mathbb{R}^m$ .

**3.2. Proof of Theorem 3.1.** In [DEZ18], Propositions 8.4, 9.2 and (the proof of) Theorem 9.5 imply that the hyperoctahedral Hall-Littlewood polynomials satisfy

the following recurrence for  $\lambda \in \Lambda^{(m,n)}$ :

$$\begin{aligned}
E(\xi)R_\lambda(\xi) &= (\beta_+(1 - q^{\mathfrak{m}_0(\lambda)}) + \beta_-(1 - q^{\mathfrak{m}_n(\lambda)}))R_\lambda(\xi) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda + e_j \in \Lambda^{(m,n)}}} (1 - \alpha_+ q^{\mathfrak{m}_0(\lambda)-1})^{\delta_{\lambda_j}} (1 - q^{\mathfrak{m}_{\lambda_j}(\lambda)})R_{\lambda+e_j}(\xi) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda - e_j \in \Lambda^{(m,n)}}} (1 - \alpha_- q^{\mathfrak{m}_n(\lambda)-1})^{\delta_{n-\lambda_j}} (1 - q^{\mathfrak{m}_{\lambda_j}(\lambda)})R_{\lambda-e_j}(\xi),
\end{aligned} \tag{3.6}$$

which is on-shell in the sense that the relation holds provided  $\xi = (\xi_1, \dots, \xi_m)$  satisfies the Bethe Ansatz equations (3.5c). Substituting  $\lambda = \mu'$  with  $\mu \in \Lambda^{(n,m)}$  leads us via Eq. (3.3b) to the recurrence relation

$$\begin{aligned}
E(\xi)R_{\mu'}(\xi) &= (\beta_+(1 - q^{m-\mu_1}) + \beta_-(1 - q^{\mu_n}))R_{\mu'}(\xi) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \mu' + e_j \in \Lambda^{(m,n)}}} (1 - \alpha_+ q^{m-\mu_1-1})^{\delta_{\mu'_j}} (1 - q^{\mathfrak{m}_{\mu'_j}(\mu')})R_{\mu'+e_j}(\xi) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \mu' - e_j \in \Lambda^{(m,n)}}} (1 - \alpha_- q^{\mu_n-1})^{\delta_{n-\mu'_j}} (1 - q^{\mathfrak{m}_{\mu'_j}(\mu')})R_{\mu'-e_j}(\xi).
\end{aligned}$$

We now observe that for any  $\mu \in \Lambda^{(n,m)}$  and  $j \in \{1, \dots, m\}$ :

$$\mu' + e_j \in \Lambda^{(m,n)} \iff \mu' + e_j = (\mu + e_i)' \quad \text{with } i = \mu'_j + 1 \in \{1, \dots, n\}$$

and

$$\mu' - e_j \in \Lambda^{(m,n)} \iff \mu' - e_j = (\mu - e_i)' \quad \text{with } i = \mu'_j \in \{1, \dots, n\}.$$

The recurrence of interest can thus be rewritten in the form

$$\begin{aligned}
E(\xi)R_{\mu'}(\xi) &= (\beta_+(1 - q^{m-\mu_1}) + \beta_-(1 - q^{\mu_n}))R_{\mu'}(\xi) \\
&+ \sum_{\substack{1 \leq i \leq n \\ \mu + e_i \in \Lambda^{(n,m)}}} (1 - \alpha_+ q^{m-\mu_1-1})^{\delta_{i-1}} (1 - q^{\mu_{i-1}-\mu_i})R_{(\mu+e_i)'}(\xi) \\
&+ \sum_{\substack{1 \leq i \leq n \\ \mu - e_i \in \Lambda^{(n,m)}}} (1 - \alpha_- q^{\mu_n-1})^{\delta_{n-i}} (1 - q^{\mu_i-\mu_{i+1}})R_{(\mu-e_i)'}(\xi),
\end{aligned}$$

where we have employed once more Eq. (3.3b).

#### 4. SPECTRAL ANALYSIS

**4.1. Solutions for the Bethe Ansatz equations.** The following system of transcendental equations provides a logarithmic form of the Bethe Ansatz equations in Theorem 3.1:

$$\begin{aligned}
2n\xi_j + v_{p_+}(\xi_j) + v_{q_+}(\xi_j) + v_{p_-}(\xi_j) + v_{q_-}(\xi_j) \\
+ \sum_{\substack{1 \leq k \leq m \\ k \neq j}} (v_q(\xi_j + \xi_k) + v_q(\xi_j - \xi_k)) &= 2\pi(m + 1 - j + \kappa_j),
\end{aligned} \tag{4.1a}$$

with  $j = 1, \dots, m$ ,  $\kappa \in \Lambda^{(m,n)}$  and

$$v_a(z) = \int_0^z \frac{(1-a^2) dx}{1-2a \cos(x) + a^2} = i \log \left( \frac{1 - ae^{iz}}{e^{iz} - a} \right) \quad (-1 < a < 1). \quad (4.1b)$$

Indeed, upon multiplying Eq. (4.1a) by  $i$  ( $= \sqrt{-1}$ ) and applying the exponential function on both sides it is readily seen that any of its solutions gives rise to a solution of the Bethe Ansatz equations (3.5c) (where—recall—the boundary parameters  $\alpha_{\pm}$ ,  $\beta_{\pm}$  and  $p_{\pm}$ ,  $q_{\pm}$  are related via Eq. (3.4a)).

For any  $\kappa \in \Lambda^{(m,n)}$ , the system in Eqs. (4.1a), (4.1b) describes the critical point of a Yang-Yang type Morse function:

$$V_{\kappa}(\xi_1, \dots, \xi_m) = \sum_{1 \leq j < k \leq m} \left( \int_0^{\xi_j + \xi_k} v_q(x) dx + \int_0^{\xi_j - \xi_k} v_q(x) dx \right) + \quad (4.2)$$

$$\sum_{1 \leq j \leq m} \left( n \xi_j^2 - 2\pi(m+1-j+\kappa_j)\xi_j + \int_0^{\xi_j} (v_{p_+}(x) + v_{q_+}(x) + v_{p_-}(x) + v_{q_-}(x)) dx \right).$$

The function  $V_{\kappa}(\xi_1, \dots, \xi_m)$  belongs to a wider class of smooth, strictly convex and radially unbounded Morse functions studied in [DE19, Section 3]. The upshot is that via a Yang-Yang type analysis, one arrives at  $\binom{m+n}{n}$  solutions for the Bethe Ansatz equations given by the respective minima of  $V_{\kappa}(\xi_1, \dots, \xi_m)$ ,  $\kappa \in \Lambda^{(m,n)}$  (cf. [DEZ18, Remark 3.5]).

**Proposition 4.1** (Solutions for the Bethe Ansatz Equations). *Let  $q \in (-1, 1) \setminus \{0\}$  and let the boundary parameters  $\alpha_{\pm}$ ,  $\beta_{\pm}$  belong to the domain specified in Eqs. (3.4a), (3.4b).*

(i) *For any  $\kappa \in \Lambda^{(m,n)}$ , the logarithmic form of the Bethe Ansatz equations in Eqs. (4.1a), (4.1b) has a unique solution  $\boxed{\xi_{\kappa} \in \mathbb{R}^m}$  given by the global minimum of the strictly convex radially unbounded Morse function  $V_{\kappa}(\xi_1, \dots, \xi_m)$  (4.2).*

(ii) *The global minima  $\xi_{\kappa}$ ,  $\kappa \in \Lambda^{(m,n)}$  in part (i) are all distinct and located within the open alcove*

$$\mathbb{A}^m = \{(\xi_1, \xi_2, \dots, \xi_m) \in \mathbb{R}^m \mid \pi > \xi_1 > \xi_2 > \dots > \xi_m > 0\} \subset \mathbb{R}_{\text{reg}}^m. \quad (4.3a)$$

Moreover, at a global minimum  $\xi = \xi_{\kappa}$  the following estimates are fulfilled:

$$\frac{\pi(m+1-j+\kappa_j)}{n+K_+} \leq \xi_j \leq \frac{\pi(m+1-j+\kappa_j)}{n+K_-} \quad (4.3b)$$

(for  $1 \leq j \leq m$ ), and

$$\frac{\pi(k-j+\kappa_j-\kappa_k)}{n+K_+} \leq \xi_j - \xi_k \leq \frac{\pi(k-j+\kappa_j-\kappa_k)}{n+K_-} \quad (4.3c)$$

(for  $1 \leq j < k \leq m$ ), where

$$K_{\pm} = (m-1) \left( \frac{1+|q|}{1-|q|} \right)^{\pm 1} + \quad (4.3d)$$

$$\frac{1}{2} \left( \left( \frac{1+|p_+|}{1-|p_+|} \right)^{\pm 1} + \left( \frac{1+|q_+|}{1-|q_+|} \right)^{\pm 1} + \left( \frac{1+|p_-|}{1-|p_-|} \right)^{\pm 1} + \left( \frac{1-|q_-|}{1-|q_-|} \right)^{\pm 1} \right).$$

*Proof.* The assertions of this proposition follow by applying [DE19, Propositions 3.1 and 3.2] to the Bethe Ansatz equations of Theorem 3.1 (cf. [DEZ18, Remark 3.5]).  $\square$

*Remark 4.2.* From a mostly academic perspective, Proposition 4.1 invites us to compute  $\xi_\kappa$  via the gradient flow of the pertinent Morse function:

$$\frac{d\xi_j}{dt} + \partial_{\xi_j} V_\kappa(\xi_1, \dots, \xi_m) = 0, \quad j = 1, \dots, m, \quad (4.4)$$

which gives rise to the following system of differential equations (cf. Eq. (4.1a))

$$\begin{aligned} \frac{d\xi_j}{dt} + 2n\xi_j + v_{p_+}(\xi_j) + v_{q_+}(\xi_j) + v_{p_-}(\xi_j) + v_{q_-}(\xi_j) \\ + \sum_{\substack{1 \leq k \leq m \\ k \neq j}} \left( v_q(\xi_j + \xi_k) + v_q(\xi_j - \xi_k) \right) = 2\pi(m+1-j+\kappa_j), \end{aligned} \quad (4.5)$$

$j = 1, \dots, m$ . For  $n = 0$  and  $\kappa = (0^m)$ , the corresponding gradient flow was analyzed in [D19] and seen to converge exponentially fast to the roots of the Askey-Wilson polynomial  $p_m(\cos \vartheta; p_+, q_+ p_-, q_- | q)$  [KLS10, Equation (14.1.1)] located within the interval of orthogonality  $0 < \vartheta < \pi$ . A minor variation of [D19, Theorem 2] reveals that in our present setting the equilibrium  $\xi_\kappa$  of the gradient system in Eq. (4.5) remains globally exponentially stable, i.e. for any initial condition  $\xi_\kappa(0)$  the unique solution  $\xi_\kappa(t)$ ,  $t \geq 0$  of the gradient system converges exponentially fast to the equilibrium  $\xi_\kappa$ . More specifically, by slightly adapting the analysis in [D19, Section 4] one readily deduces that for any  $0 < \varepsilon < 2(n + \kappa_-)$  (= a lower bound for the eigenvalues of the hessian of  $V_\kappa(\xi_1, \dots, \xi_m)$ ), there exists a constant  $C_\varepsilon > 0$  such that

$$\forall t \geq 0: \quad \|\xi_\kappa(t) - \xi_\kappa\|_\infty \leq C_\varepsilon e^{-\varepsilon t}, \quad (4.6)$$

where  $\|\xi\|_\infty \equiv \max_{1 \leq j \leq m} |\xi_j|$ . Apart from  $\varepsilon$ , the actual value of the constant  $C_\varepsilon$  in the uniform estimate of the error term will depend on the choice of the initial condition  $\xi_\kappa(0) \in \mathbb{R}^m$ , as well as on  $\kappa \in \Lambda^{(m,n)}$  and  $q, p_\pm, q_\pm \in (-1, 1)$  (cf. Remark 4.4 below). Notice that the  $q$ -difference Toda hamiltonian  $H$  (2.2) degenerates to a discrete Laplacian on  $\Lambda^{(n,m)}$  in the symplectic Schur limit  $\alpha_\pm, \beta_\pm, q \rightarrow 0$ . At this elementary point in the parameters space, one has that (cf. Eq. (4.3b))

$$\xi_\kappa \rightarrow \left( \frac{\pi(m+\kappa_1)}{m+n+1}, \frac{\pi(m-1+\kappa_2)}{m+n+1}, \dots, \frac{\pi(m+1-j+\kappa_j)}{m+n+1}, \dots, \frac{\pi(1+\kappa_m)}{m+n+1} \right),$$

which serves as a convenient initial condition for the gradient flow (4.5). Indeed, at this particular value of the spectral parameter the bounds in Eqs. (4.3b) and (4.3c) are fulfilled for any  $p_\pm, q_\pm, q \in (-1, 1)$ .

**4.2. Spectrum and eigenbasis.** By combining Theorem 3.1 and Proposition 4.1, an eigenbasis of Bethe Ansatz wave functions for the  $q$ -difference Toda hamiltonian  $H$  (2.2) in the Hilbert space  $\ell(\Lambda^{(n,m)}, \Delta)$  is found together with the corresponding eigenvalues.

**Theorem 4.3** (Spectrum and Eigenbasis). *Let  $q \in (-1, 1) \setminus \{0\}$  and let the boundary parameters  $\alpha_\pm, \beta_\pm$  belong to the domain specified in Eqs. (3.4a), (3.4b). For any  $\xi \in \mathbb{R}_{\text{reg}}^m$  and  $\kappa \in \Lambda^{(m,n)}$ , the function  $\psi_\xi : \Lambda^{(n,m)} \rightarrow \mathbb{R}$  refers to the Hall-Littlewood Bethe Ansatz wave function from Theorem 3.1 and  $\xi_\kappa \in \mathbb{A}^m \subset \mathbb{R}_{\text{reg}}^m$  denotes the unique global minimum of  $V_\kappa(\xi_1, \dots, \xi_m)$  (4.2) detailed in Proposition 4.1.*

(i) *The spectrum of the  $q$ -difference Toda hamiltonian  $H$  (2.2) in the Hilbert space  $\ell(\Lambda^{(n,m)}, \Delta)$  consists of the eigenvalues  $E(\xi_\kappa)$ ,  $\kappa \in \Lambda^{(m,n)}$ , where  $E(\xi)$  is given by Eq. (3.5b).*



(ii) The corresponding Bethe Ansatz wave functions  $\psi_{\xi_\kappa}$ ,  $\kappa \in \Lambda^{(m,n)}$  constitute an eigenbasis for  $H$  (2.2) in  $\ell(\Lambda^{(n,m)}, \Delta)$  such that

$$H\psi_{\xi_\kappa} = E(\xi_\kappa)\psi_{\xi_\kappa} \quad (\kappa \in \Lambda^{(m,n)}). \quad (4.7)$$

*Proof.* It is clear from Theorem 3.1 and Proposition 4.1 that for any  $\kappa \in \Lambda^{(m,n)}$  the Bethe Ansatz wave function  $\psi_{\xi_\kappa}$  solves the eigenvalue equation (4.7). Moreover, the value of the Bethe Ansatz wave functions at the origin is given by Macdonald's three-parameter Poincaré series for the root system  $BC_m$  [M00, §10]:

$$\begin{aligned} \psi_\xi(0^n) &= R_{(0^m)}(\xi_1, \dots, \xi_m) = \sum_{\substack{\sigma \in S_m \\ \epsilon \in \{1, -1\}^m}} C(\epsilon_1 \xi_{\sigma(1)}, \dots, \epsilon_m \xi_{\sigma(m)}) \\ &= \frac{(\alpha_+; q)_m (q; q)_m}{(1-q)^m} \neq 0 \end{aligned}$$

(cf. also [DEZ18, Remark 3.4]). Hence, for any  $\kappa \in \Lambda^{(m,n)}$  the wave function  $\psi_{\xi_\kappa}$  constitutes a proper (i.e. nontrivial) eigenfunction of the  $q$ -difference Toda hamiltonian  $H$  with eigenvalue  $E(\xi_\kappa)$ . To confirm the completeness of the Bethe Ansatz it remains to check that the wave functions in question indeed form a basis for  $\ell(\Lambda^{(n,m)}, \Delta)$ , or equivalently, that the hyperoctahedral Hall-Littlewood polynomials  $R_{\mu'}(\xi_1, \dots, \xi_m)$ ,  $\mu \in \Lambda^{(n,m)}$  are linearly independent as functions on the Bethe spectrum  $\{\xi_\kappa \mid \kappa \in \Lambda^{(m,n)}\}$ . This independence is immediate from the second part of [DEZ18, Theorem 3.1].  $\square$

*Remark 4.4.* The  $q$ -difference Toda hamiltonian  $H$ , the positive weights  $\Delta_\mu$ , and the Bethe Ansatz wave function  $\psi_\xi$  clearly extend smoothly in the parameters to the domain  $q, p_\pm, q_\pm \in (-1, 1)$ . With the aid of the implicit function theorem, it is seen that the same is true for the solutions  $\xi_\kappa$ ,  $\kappa \in \Lambda^{(n,m)}$  of the (logarithmic) Bethe Ansatz equation in Proposition 4.1 (cf. [DEZ18, Remark 3.6]). Indeed, the Morse function  $V_\kappa(\xi_1, \dots, \xi_m)$  extends smoothly in the parameters and remains strictly convex. Hence, for  $\kappa \in \Lambda^{(m,n)}$  the Bethe Ansatz wave function  $\psi_{\xi_\kappa}$  constitutes in fact an eigenfunction of  $H$  (2.2) in  $\ell(\Lambda^{(n,m)}, \Delta)$  with eigenvalue  $E(\xi_\kappa)$  (3.5b) for any  $q, p_\pm, q_\pm \in (-1, 1)$  (i.e. even if one or more of the parameters in question vanish).

*Remark 4.5.* The self-adjointness of  $H$  (2.2) in Proposition 2.1 implies that

$$\forall \kappa, \nu \in \Lambda^{(m,n)} : \quad \langle \psi_{\xi_\kappa}, \psi_{\xi_\nu} \rangle_\Delta = 0 \quad \text{if } \kappa \neq \nu, \quad (4.8a)$$

provided  $E(\xi_\kappa) \neq E(\xi_\nu)$ . Rewritten in terms of the (real-valued) hyperoctahedral Hall-Littlewood polynomials  $R_\lambda(\xi) = R_\lambda(\xi_1, \dots, \xi_m)$  one obtains that in this situation:

$$\sum_{\lambda \in \Lambda^{(m,n)}} R_\lambda(\xi_\kappa) R_\lambda(\xi_\nu) \Delta'_\lambda = 0 \quad \text{if } \kappa \neq \nu, \quad (4.8b)$$

with

$$\Delta'_\lambda = \frac{(q; q)_m}{(\alpha_+; q)_{m_0(\lambda)} (\alpha_-; q)_{m_n(\lambda)} \prod_{0 \leq i \leq n} (q; q)_{m_i(\lambda)}} \quad (4.8c)$$

(so  $\Delta'_{\mu'} = \Delta_\mu$  for  $\mu \in \Lambda^{(n,m)}$ ). To date the latter orthogonality relation has been checked directly without the proviso regarding the nondegeneracy of the corresponding eigenvalues of  $H$  in the following four cases:

- (a) if  $q = 0$  and  $p_\pm, q_\pm \in (-1, 1)$ , cf. [DE18a, Theorem 3.1];
- (b) if  $0 < q < 1$ ,  $\alpha_\pm = 0$  and  $\beta_\pm \in (-1, 1)$ , cf. [DE17, Section 11.4];

- (c) if  $q, p_{\pm}, q_{\pm} \in (-1, 1)$  and  $n \geq 2m$ , cf. [D22, Theorem 4.2];
- (d) if  $q, p_{\pm}, q_{\pm} \in (-1, 1)$  and  $n = 1$ , cf. [D22, Section 5.7].

In view of Remark 4.4, within these four subdomains the statements concerning the spectrum and completeness formulated in parts (i) and (ii) of Theorem 4.3 therefore persist with the corresponding Bethe Ansatz eigenbasis being orthogonal in  $\ell(\Lambda^{(n,m)}, \Delta)$  (as expected).

*Remark 4.6.* In [DEZ18], Eq. (3.6) is interpreted as the eigenvalue equation for a hamiltonian of an  $m$ -particle  $q$ -boson model on the lattice  $\{0, 1, \dots, n\}$ . The quantum integrability of this  $m$ -particle  $q$ -boson hamiltonian, which is thus given explicitly by the difference operator acting at the RHS of Eq. (3.6), was established for  $\alpha_+ = \alpha_- = 0$  in [DE17] (using the quantum inverse scattering method) and for general boundary parameters in [DEZ18, Section 8] (using representations of the double affine Hecke algebra of type  $C^{\vee}C$  at the critical level  $q = 0$ ). As detailed explicitly for the hamiltonian in the proof of Theorem 3.1, the mapping  $\mu \rightarrow \mu'$  from  $\Lambda^{(n,m)}$  onto  $\Lambda^{(m,n)}$  allows us to pull back the commuting quantum integrals for the  $q$ -boson model from  $\ell(\Lambda^{(m,n)}, \Delta')$  to  $\ell(\Lambda^{(n,m)}, \Delta)$ . This maps the commuting quantum integrals in question to an algebra of commuting difference operators in  $\ell(\Lambda^{(n,m)}, \Delta)$  containing  $H$  (2.2). Since [DEZ18, Theorem 9.5] guarantees that the latter algebra of commuting difference operators is (Harish-Chandra-)isomorphic to the algebra of complex functions on the joint spectrum  $\{\xi_{\kappa} \mid \kappa \in \Lambda^{(m,n)}\} \subset \mathbb{A}^m$ , this establishes the quantum integrability of our  $q$ -difference Toda hamiltonian  $H$  (2.2).

## 5. THE LIMIT $q \rightarrow 1$

**5.1. Quantum hamiltonian.** Upon dividing out an overall scaling factor  $1 - q$ , the  $q$ -difference Toda hamiltonian  $H$  (2.2) degenerates for  $q \rightarrow 1$  to an elementary difference operator  $\tilde{H}$  with linear coefficients:

$$\begin{aligned} \tilde{H} &= \beta_+(m - \mu_1) + \beta_-(\mu_n) + \\ &\sum_{1 \leq i \leq n} \left( (1 - \alpha_+)^{\delta_{i-1}} (\mu_{i-1} - \mu_i) T_i + (1 - \alpha_-)^{\delta_{n-i}} (\mu_i - \mu_{i+1}) T_i^{-1} \right). \end{aligned} \quad (5.1)$$

From Proposition 2.1 it follows that—assuming  $\alpha_{\pm} \in (-1, 1)$  and  $\beta_{\pm} \in \mathbb{R}$ —this limiting quantum hamiltonian is self-adjoint in a Hilbert space  $\ell(\Lambda^{(n,m)}, \tilde{\Delta})$  governed by the weights of a two-parameter multinomial distribution on the partitions  $\Lambda^{(n,m)}$ :

$$\begin{aligned} \tilde{\Delta}_{\mu} &= \lim_{q \rightarrow 1} \Delta_{\mu} \\ &= \frac{m!}{(1 - \alpha_+)^{m - \mu_1} (1 - \alpha_-)^{\mu_n} \prod_{0 \leq i \leq n} (\mu_i - \mu_{i+1})!} \\ &= \mathcal{N} \cdot \frac{m! \prod_{0 \leq i \leq n} \rho_i^{\mu_i - \mu_{i+1}}}{\prod_{0 \leq i \leq n} (\mu_i - \mu_{i+1})!}, \end{aligned} \quad (5.2a)$$

with

$$\rho_i = \begin{cases} \frac{(1 - \alpha_+)^{-1}}{(n-1) + (1 - \alpha_+)^{-1} + (1 - \alpha_-)^{-1}} & \text{if } i = 0, \\ \frac{1}{(n-1) + (1 - \alpha_+)^{-1} + (1 - \alpha_-)^{-1}} & \text{if } 0 < i < n, \\ \frac{(1 - \alpha_-)^{-1}}{(n-1) + (1 - \alpha_+)^{-1} + (1 - \alpha_-)^{-1}} & \text{if } i = n, \end{cases} \quad (5.2b)$$

and

$$\mathcal{N} = \sum_{\mu \in \Lambda^{(n,m)}} \tilde{\Delta}_\mu = ((n-1) + (1-\alpha_+)^{-1} + (1-\alpha_-)^{-1})^m. \quad (5.2c)$$

Notice in particular that  $\rho_0 = \rho_2 = \dots = \rho_n = \frac{1}{n+1}$  if  $\alpha_+ = \alpha_- = 0$ .

**5.2. Bethe Ansatz.** The Bethe Ansatz wave function  $\psi_\xi$  (3.5a) degenerates in the limit  $q \rightarrow 1$  to a wave function  $\tilde{\psi}_\xi : \Lambda^{(n,m)} \rightarrow \mathbb{C}$  with values

$$\tilde{\psi}_\xi(\mu) = \tilde{R}_{\mu'}(\xi_1, \dots, \xi_m) \quad (5.3a)$$

that separate in terms of univariate ( $BC_1$ -type) Hall-Littlewood polynomials. Specifically, for any  $\lambda \in \Lambda^{(m,n)}$  and  $\xi \in \mathbb{R}_{\text{reg}}^m$  one has that

$$\begin{aligned} \tilde{R}_\lambda(\xi_1, \dots, \xi_m) &= \lim_{q \rightarrow 1} R_\lambda(\xi_1, \dots, \xi_m) \\ &= \sum_{\sigma \in S_m} R_{\lambda_1}(\xi_{\sigma(1)}) R_{\lambda_2}(\xi_{\sigma(2)}) \cdots R_{\lambda_m}(\xi_{\sigma(m)}) \end{aligned} \quad (5.3b)$$

with

$$\begin{aligned} R_l(\vartheta) &= \frac{(1 - \beta_+ e^{-i\vartheta} + \alpha_+ e^{-2i\vartheta})}{1 - e^{-2i\vartheta}} \exp(il\vartheta) + \\ &\quad \frac{(1 - \beta_+ e^{i\vartheta} + \alpha_+ e^{2i\vartheta})}{1 - e^{2i\vartheta}} \exp(-il\vartheta) \end{aligned} \quad (5.3c)$$

( $l \in \{0, \dots, n\}$ ,  $\vartheta \notin \pi\mathbb{Z}$ ).

Heuristically, for the Bethe Ansatz wave function  $\tilde{\psi}_\xi$  (5.3a)–(5.3c) to solve the  $q \rightarrow 1$  eigenvalue equation

$$\tilde{H}\tilde{\psi}_\xi = \tilde{E}(\xi)\tilde{\psi}_\xi \quad \text{with} \quad \tilde{E}(\xi) = 2 \sum_{1 \leq j \leq m} \cos(\xi_j), \quad (5.4a)$$

one expects the spectral parameter  $\xi$  to be required to satisfy the following decoupled system of Bethe Ansatz equations arising from Eq. (3.5c) in the limit  $q \rightarrow 1$ :

$$e^{2in\xi_j} = \frac{(1 - \beta_+ e^{i\xi_j} + \alpha_+ e^{2i\xi_j})(1 - \beta_- e^{i\xi_j} + \alpha_- e^{2i\xi_j})}{(e^{2i\xi_j} - \beta_+ e^{i\xi_j} + \alpha_+)(e^{2i\xi_j} - \beta_- e^{i\xi_j} + \alpha_-)}, \quad (5.4b)$$

$j = 1, \dots, m$ .

**5.3. Solutions for the Bethe Ansatz equations.** It is illuminating to emphasize that for  $p_+, q_+, p_-, q_- \in (-1, 1)$  the decoupled Bethe Ansatz equations in Eq. (5.4b) can be conveniently solved in terms of the roots of the Askey-Wilson polynomial  $p_{n+1}(\cos \vartheta; p_+, q_+, p_-, q_- | q)$  [KLS10, Equation (14.1.1)] at  $q = 0$ . Indeed, it is clear from the orthogonality relation [KLS10, Equation (14.1.2)] that at  $q = 0$  the Askey-Wilson polynomials fall within a well-known class of orthogonal polynomials studied by Bernstein and Szegő [S75, Chapter 2.6]. The classical theory of Bernstein and Szegő tells us, moreover, that the polynomials in question can be written explicitly

as follows (cf. e.g. [DE18b, Section 4.3]):

$$\begin{aligned} & \frac{p_{n+1}(\cos \vartheta; p_+, q_+ p_-, q_- | 0)}{(p_+ q_+ p_- q_- q^n; q)_{n+1}} = \\ & \frac{\prod_{\epsilon=\pm} (1 - p_\epsilon e^{-i\vartheta})(1 - q_\epsilon e^{-i\vartheta})}{1 - e^{-2i\vartheta}} \exp(i(n+1)\vartheta) \\ & + \frac{\prod_{\epsilon=\pm} (1 - p_\epsilon e^{i\vartheta})(1 - q_\epsilon e^{i\vartheta})}{1 - e^{2i\vartheta}} \exp(-i(n+1)\vartheta) \end{aligned} \quad (5.5)$$

( $\vartheta \notin \pi\mathbb{Z}$ ). This explicit formula reveals in particular that the roots

$$0 < \vartheta_0 < \vartheta_1 < \dots < \vartheta_n < \pi \quad (5.6)$$

of  $p_{n+1}(\cos \vartheta; p_+, q_+ p_-, q_- | 0)$  solve the Bethe Ansatz equation in Eq. (5.4b) (with  $\xi_j$  replaced by  $\vartheta$ ). More specifically, the root  $\vartheta_{\mathbf{k}}$  corresponds to the solution of the associated logarithmic Bethe Ansatz equation (cf. Remark 4.2 above)

$$2n\vartheta + v_{p_+}(\vartheta) + v_{q_+}(\vartheta) + v_{p_-}(\vartheta) + v_{q_-}(\vartheta) = 2\pi(\mathbf{k} + 1), \quad (5.7)$$

with  $\mathbf{k} \in \{0, 1, \dots, n\}$ .

The upshot is that to any  $\kappa = (\kappa_1, \dots, \kappa_m) \in \Lambda^{(m,n)}$ , we can now attach a solution  $\tilde{\xi}_\kappa$  of the decoupled system of Bethe Ansatz equations in Eq. (5.4b) by forming the following vector of  $q = 0$  Askey-Wilson roots:

$$\tilde{\xi}_\kappa = (\vartheta_{\kappa_1}, \vartheta_{\kappa_2}, \dots, \vartheta_{\kappa_m}) \in \{\xi \in \mathbb{R}^m \mid \pi > \xi_1 \geq \xi_2 \geq \dots \geq \xi_m > 0\}. \quad (5.8)$$

Notice that  $\tilde{\xi}_\kappa$  encodes the unique global minimum of the decoupled Morse function

$$\begin{aligned} \tilde{V}_\kappa(\xi_1, \dots, \xi_m) = & \\ & \sum_{1 \leq j \leq m} \left( n\xi_j^2 - 2\pi(\kappa_j + 1)\xi_j + \int_0^{\xi_j} (v_{p_+}(x) + v_{q_+}(x) + v_{p_-}(x) + v_{q_-}(x)) dx \right). \end{aligned} \quad (5.9)$$

**5.4. Spectrum and eigenbasis for  $q \rightarrow 1$ .** When tying the above observations together, one is led to the following  $q \rightarrow 1$  counterpart of Theorem 4.3.

**Theorem 5.1** (Spectrum and Eigenbasis for  $q \rightarrow 1$ ). *Let the boundary parameters  $\alpha_\pm, \beta_\pm$  be of the form in Eq. (3.4a) with  $p_\pm, q_\pm \in (-1, 1)$ . For any  $\xi \in \{\xi \in \mathbb{R}^m \mid \xi_j \notin \pi\mathbb{Z}, \forall 1 \leq j \leq m\}$  and  $\kappa \in \Lambda^{(m,n)}$ , the function  $\tilde{\psi}_\xi : \Lambda^{(n,m)} \rightarrow \mathbb{R}$  refers to the  $q = 1$  Hall-Littlewood Bethe Ansatz wave function in Eq. (5.3a)–(5.3c) and  $\tilde{\xi}_\kappa$  denotes the solution in Eq. (5.8) of the decoupled Bethe Ansatz equations (5.4b).*

(i) *The spectrum of  $\tilde{H}$  (5.1) in the Hilbert space  $\ell(\Lambda^{(n,m)}, \tilde{\Delta})$  consists of the eigenvalues  $\tilde{E}(\tilde{\xi}_\kappa)$ ,  $\kappa \in \Lambda^{(m,n)}$ , where  $\tilde{E}(\xi)$  is given by Eq. (5.4a).*

(ii) *The corresponding Bethe Ansatz wave functions  $\tilde{\psi}_{\tilde{\xi}_\kappa}$ ,  $\kappa \in \Lambda^{(m,n)}$  constitute an orthogonal eigenbasis for  $\tilde{H}$  (5.1) in  $\ell(\Lambda^{(n,m)}, \tilde{\Delta})$  such that*

$$\tilde{H}\tilde{\psi}_{\tilde{\xi}_\kappa} = \tilde{E}(\tilde{\xi}_\kappa)\tilde{\psi}_{\tilde{\xi}_\kappa} \quad (\kappa \in \Lambda^{(m,n)}). \quad (5.10)$$

*Remark 5.2.* Systematic studies of orthogonal polynomials associated with the multinomial distribution give rise to multivariate generalizations of the Krawtchouk polynomials [DG14, GLZ13, GR11, I12, I20, MT04]. For the particular instance of the two-parameter multinomial distribution in Eqs. (5.2a)–(5.2c), Theorem 5.1 suggests an intriguing link to the  $q = 1$  Hall-Littlewood Bethe Ansatz wave function  $\tilde{\psi}_{\tilde{\xi}_\kappa}$ . When  $n = 1$  this link is actually well-understood in the literature as a relation

between classical univariate Krawtchouk polynomials and elementary symmetric polynomials, cf. e.g. [D21, Equation (5.14)].

**5.5. Proof of Theorem 5.1.** In order to establish the claims of the theorem in full rigor avoiding tricky formal limits, let us first check that the Bethe Ansatz wave functions  $\tilde{\psi}_{\tilde{\xi}_\kappa}$ ,  $\kappa \in \Lambda^{(m,n)}$  are indeed orthogonal in the Hilbert space  $\ell(\Lambda^{(n,m)}, \tilde{\Delta})$  (i.e. with the weights  $\tilde{\Delta}_\mu$  replacing  $\Delta_\mu$ ):

$$\forall \kappa, \nu \in \Lambda^{(m,n)} : \quad \langle \tilde{\psi}_{\tilde{\xi}_\kappa}, \tilde{\psi}_{\tilde{\xi}_\nu} \rangle_{\tilde{\Delta}} = 0 \quad \text{if } \kappa \neq \nu. \quad (5.11)$$

Rewritten in terms of the  $q = 1$  hyperoctahedral Hall-Littlewood polynomials, the inner product on the LHS of Eq. (5.11) becomes (cf. Eqs. (4.8b), (4.8c)):

$$\sum_{\lambda \in \Lambda^{(m,n)}} \frac{\tilde{R}_\lambda(\tilde{\xi}_\kappa) \tilde{R}_\lambda(\tilde{\xi}_\nu) m!}{(1 - \alpha_+)^{m_0(\lambda)} (1 - \alpha_-)^{m_n(\lambda)} \prod_{0 \leq i \leq n} m_i(\lambda)!}, \quad (5.12)$$

with  $\tilde{R}_\lambda(\xi)$  and  $\tilde{\xi}_\kappa$  taken from Eqs. (5.3b) and (5.8), respectively. In particular, if  $m = 1$  then the sum in Eq. (5.12) is of the form

$$\sum_{0 \leq \lambda \leq n} \frac{R_\lambda(\vartheta_\kappa) R_\lambda(\vartheta_\nu)}{(1 - \alpha_+)^{\delta_\lambda} (1 - \alpha_-)^{\delta_{n-\lambda}}} \quad (5.13)$$

with  $0 \leq \kappa \neq \nu \leq n$ , where  $\vartheta_0, \dots, \vartheta_n$  refer to the  $q = 0$  Askey-Wilson roots from Eq. (5.6). Since the sum in Eq. (5.13) coincides with that of the inner product in Case (a) of Remark 4.5 (specialized to  $m = 1$ ), in this simplest situation the asserted orthogonality is immediate from the remark in question.

If on the other hand  $m > 1$ , then the inner product in Eq. (5.12) decomposes into a sum of contributions of the form

$$\begin{aligned} & \sum_{\lambda \in \Lambda^{(m,n)}} \frac{R_{\lambda_1}(\vartheta_{\mathbf{k}_1}) \cdots R_{\lambda_m}(\vartheta_{\mathbf{k}_m}) R_{\lambda_1}(\vartheta_{\mathbf{n}_1}) \cdots R_{\lambda_m}(\vartheta_{\mathbf{n}_m}) m!}{(1 - \alpha_+)^{m_0(\lambda)} (1 - \alpha_-)^{m_n(\lambda)} \prod_{0 \leq i \leq n} m_i(\lambda)!} \\ &= \sum_{0 \leq \lambda_1, \dots, \lambda_m \leq n} \frac{R_{\lambda_1}(\vartheta_{\mathbf{k}_1}) \cdots R_{\lambda_m}(\vartheta_{\mathbf{k}_m}) R_{\lambda_1}(\vartheta_{\mathbf{n}_1}) \cdots R_{\lambda_m}(\vartheta_{\mathbf{n}_m})}{\prod_{1 \leq j \leq m} (1 - \alpha_+)^{\delta_{\lambda_j}} (1 - \alpha_-)^{\delta_{n-\lambda_j}}} \\ &= \prod_{1 \leq j \leq m} \sum_{0 \leq \lambda_j \leq n} \frac{R_{\lambda_j}(\vartheta_{\mathbf{k}_j}) R_{\lambda_j}(\vartheta_{\mathbf{n}_j})}{(1 - \alpha_+)^{\delta_{\lambda_j}} (1 - \alpha_-)^{\delta_{n-\lambda_j}}}, \end{aligned}$$

where the  $m$ -tuple  $(\mathbf{k}_1, \mathbf{k}_2, \dots, \mathbf{k}_m)$  denotes a reordering  $(\kappa_{\sigma(1)}, \kappa_{\sigma(2)}, \dots, \kappa_{\sigma(m)})$  of  $\kappa \in \Lambda^{(m,n)}$  and the  $m$ -tuple  $(\mathbf{n}_1, \mathbf{n}_2, \dots, \mathbf{n}_m)$  denotes a reordering of  $\nu \in \Lambda^{(m,n)}$  (not necessarily stemming from the same permutation  $\sigma \in S_m$ ). From the orthogonality for  $m = 1$  it is now clear that all such contributions vanish, unless  $\mathbf{k}_j = \mathbf{n}_j$  for  $j = 1, \dots, m$ , i.e., except when  $\kappa = \nu$  and both reorderings coincide.

It remains to verify that the eigenvalue equation in Eq. (5.4a) is satisfied at  $\xi = \tilde{\xi}_\kappa$  for  $\kappa \in \Lambda^{(m,n)}$ . Rewritten in terms of  $\lambda = \mu' \in \Lambda^{(m,n)}$ , this eigenvalue

equation reads explicitly (cf. Eq. (3.6)):

$$\begin{aligned}
\tilde{E}(\tilde{\xi}_\kappa)\tilde{R}_\lambda(\tilde{\xi}_\kappa) &= (\beta_+ m_0(\lambda) + \beta_- m_n(\lambda))\tilde{R}_\lambda(\tilde{\xi}_\kappa) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda + e_j \in \Lambda^{(m,n)}}} (1 - \alpha_+)^{\delta_{\lambda_j}} m_{\lambda_j}(\lambda) \tilde{R}_{\lambda + e_j}(\tilde{\xi}_\kappa) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda - e_j \in \Lambda^{(m,n)}}} (1 - \alpha_-)^{\delta_{n - \lambda_j}} m_{\lambda_j}(\lambda) \tilde{R}_{\lambda - e_j}(\tilde{\xi}_\kappa).
\end{aligned} \tag{5.14}$$

For instance, if  $m = 1$  then Eq. (5.14) simplifies to

$$\begin{aligned}
2 \cos(\vartheta_\kappa) R_\lambda(\vartheta_\kappa) &= (\beta_+ \delta_\lambda + \beta_- \delta_{n-\lambda}) R_\lambda(\vartheta_\kappa) + \\
(1 - \alpha_+)^{\delta_\lambda} (1 - \delta_{n-\lambda}) R_{\lambda+1}(\vartheta_\kappa) &+ (1 - \alpha_-)^{\delta_{n-\lambda}} (1 - \delta_\lambda) R_{\lambda-1}(\vartheta_\kappa),
\end{aligned} \tag{5.15}$$

where  $0 \leq \kappa, \lambda \leq n$ . Apart from a missing overall factor  $(1 - q)$  on both sides (which was actually divided out at the start of Subsection 5.1), Eq. (5.15) coincides precisely with the  $m = 1$  specialization of the eigenvalue equation from Theorem 4.3 at  $\mu = \lambda'$  (which agrees with the observation that the dependence on  $q$  drops out when  $m = 1$ ). Upon recalling Remark 4.4, this settles the validity of Eq. (5.15) for the full parameter regime  $p_\pm, q_\pm \in (-1, 1)$ .

Moreover, by virtue of Eq. (5.15) one has more generally that for  $m > 1$  and any  $\sigma \in S_m$ :

$$\begin{aligned}
&2 \sum_{1 \leq j \leq m} \cos(\vartheta_{\kappa_j}) \prod_{1 \leq l \leq m} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) = \\
&\sum_{1 \leq j \leq m} (\beta_+ \delta_{\lambda_j} + \beta_- \delta_{n-\lambda_j}) \prod_{1 \leq l \leq m} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) \\
&+ \sum_{1 \leq j \leq m} (1 - \alpha_+)^{\delta_{\lambda_j}} (1 - \delta_{n-\lambda_j}) R_{\lambda_j+1}(\vartheta_{\kappa_{\sigma(j)}}) \prod_{\substack{1 \leq l \leq m \\ l \neq j}} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) \\
&+ \sum_{1 \leq j \leq m} (1 - \alpha_-)^{\delta_{n-\lambda_j}} (1 - \delta_{\lambda_j}) R_{\lambda_j-1}(\vartheta_{\kappa_{\sigma(j)}}) \prod_{\substack{1 \leq l \leq m \\ l \neq j}} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) \\
&= (\beta_+ m_0(\lambda) + \beta_- m_n(\lambda)) \prod_{1 \leq l \leq m} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda + e_j \in \Lambda^{(m,n)}}} (1 - \alpha_+)^{\delta_{\lambda_j}} m_{\lambda_j}(\lambda) R_{\lambda_j+1}(\vartheta_{\kappa_{\sigma(j)}}) \prod_{\substack{1 \leq l \leq m \\ l \neq j}} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}) \\
&+ \sum_{\substack{1 \leq j \leq m \\ \lambda - e_j \in \Lambda^{(m,n)}}} (1 - \alpha_-)^{\delta_{n-\lambda_j}} m_{\lambda_j}(\lambda) R_{\lambda_j-1}(\vartheta_{\kappa_{\sigma(j)}}) \prod_{\substack{1 \leq l \leq m \\ l \neq j}} R_{\lambda_l}(\vartheta_{\kappa_{\sigma(l)}}),
\end{aligned}$$

which entails Eq. (5.14) through symmetrization by summing over all  $\sigma \in S_m$  on both sides.

#### ACKNOWLEDGEMENTS

This work was supported in part by the *Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT)* Grant # 1210015.

## REFERENCES

- BC14. A. Borodin and I. Corwin, Macdonald processes, *Probab. Theory Related Fields* **158** (2014), 225–400.
- BF05. A. Braverman and M. Finkelberg, Finite difference quantum Toda lattice via equivariant  $K$ -theory, *Transform. Groups* **10** (2005), 363–386.
- DG14. P. Diaconis and R. Griffiths, An introduction to multivariate Krawtchouk polynomials and their applications, *J. Statist. Plann. Inference* **154** (2014), 39–53.
- D19. J.F. van Diejen, Gradient system for the roots of the Askey-Wilson polynomial, *Proc. Amer. Math. Soc.* **147** (2019), 5239–5249.
- D21. J.F. van Diejen,  $q$ -deformation of the Kac-Sylvester tridiagonal matrix, *Proc. Amer. Math. Soc.* **149** (2021), 2291–2304.
- D22. J.F. van Diejen, Harmonic analysis of boxed hyperoctahedral Hall-Littlewood polynomials, *J. Funct. Anal.* **282** (2022), no. 1, Paper No. 109256.
- DE15. J.F. van Diejen and E. Emsiz, Integrable boundary interactions for Ruijsenaars’ difference Toda chain, *Comm. Math. Phys.* **337** (2015), 171–189.
- DE17. J.F. van Diejen and E. Emsiz, Orthogonality of Bethe Ansatz eigenfunctions for the Laplacian on a hyperoctahedral Weyl alcove, *Comm. Math. Phys.* **350** (2017), 1017–1067.
- DE18a. J.F. van Diejen and E. Emsiz, Discrete Fourier transform associated with generalized Schur polynomials, *Proc. Amer. Math. Soc.* **146** (2018), 3459–3472.
- DE18b. J.F. van Diejen and E. Emsiz, Quadrature rules from finite orthogonality relations for Bernstein-Szegő polynomials, *Proc. Amer. Math. Soc.* **146** (2018), 5333–5347.
- DE19. J.F. van Diejen and E. Emsiz, Solutions of convex Bethe Ansatz equations and the zeros of (basic) hypergeometric orthogonal polynomials, *Lett. Math. Phys.* **109** (2019), 89–112.
- DEZ18. J.F. van Diejen, E. Emsiz, and I.N. Zurrián, Completeness of the Bethe Ansatz for an open  $q$ -boson system with integrable boundary interactions, *Ann. Henri Poincaré* **19** (2018), 1349–1384.
- DK18. P. Di Francesco and R. Kedem, Difference equations for graded characters from quantum cluster algebra, *Transform. Groups* **23** (2018), 391–424.
- DKT17. P. Di Francesco, R. Kedem, and B. Turmunkh, A path model for Whittaker vectors, *J. Phys. A* **50** (2017), no. 25, 255201.
- DP16. A. Duval and V. Pasquier,  $q$ -bosons, Toda lattice, Pieri rules and Baxter  $q$ -operator, *J. Phys. A* **49** (2016), no. 15, 154006.
- E99. P. Etingof, Whittaker functions on quantum groups and  $q$ -deformed Toda operators, in: *Differential Topology, Infinite-Dimensional Lie Algebras, and Applications*, A. Ashtashkevich and S. Tabachnikov (eds.), Amer. Math. Soc. Transl. Ser. 2, Vol.194, Amer. Math. Soc., Providence, RI, 1999, p. 9–25.
- FJM09. B. Feigin, E. Feigin, M. Jimbo, T. Miwa, and E. Mukhin, Fermionic formulas for eigenfunctions of the difference Toda Hamiltonian, *Lett. Math. Phys.* **88** (2009), 39–77.
- GLZ13. V.X. Genest, L. Vinet, and A. Zhedanov, The multivariate Krawtchouk polynomials as matrix elements of the rotation group representations on oscillator states, *J. Phys. A* **46** (2013), no. 50, 505203.
- GLO11. A. Gerasimov, D. Lebedev, and S. Oblezin, On  $q$ -deformed  $gl_{l+1}$ -Whittaker function III, *Lett. Math. Phys.* **97** (2011), 1–24.
- GL03. A. Givental and Y.-P. Lee, Quantum  $K$ -theory on flag manifolds, finite-difference Toda lattices and quantum groups, *Invent. Math.* **151** (2003), 193–219.
- GR11. F.A. Grünbaum and M. Rahman, A system of multivariable Krawtchouk polynomials and a probabilistic application, *SIGMA Symmetry Integrability Geom. Methods Appl.* **7** (2011), 119.
- I12. P. Iliev, A Lie-theoretic interpretation of multivariate hypergeometric polynomials, *Compos. Math.* **148** (2012), no. 3, 991–1002.
- I20. P. Iliev and Y. Xu, Hahn polynomials on polyhedra and quantum integrability, *Adv. Math.* **364** (2020), 107032.

- KLS10. R. Koekoek, P.A. Lesky, and R.F. Swarttouw, *Hypergeometric Orthogonal Polynomials and their  $q$ -Analogues*, Springer Monographs in Mathematics. Springer-Verlag, Berlin-Heidelberg, 2010.
- KT96. V.B. Kuznetsov and A.V. Tsyganov, Quantum relativistic Toda chains, *J. Math. Sci.* **80** (1996), 1802–1810.
- M00. I.G. Macdonald, Orthogonal polynomials associated with root systems, *Sém. Lothar. Combin.* **45** (2000/01), Art. B45a.
- MT04. H. Mizukawa and H. Tanaka,  $(n + 1, m + 1)$ -hypergeometric functions associated to character algebras, *Proc. Amer. Math. Soc.* **132** (2004), 2613–2618.
- R90. S.N.M. Ruijsenaars, Relativistic Toda systems, *Comm. Math. Phys.* **133** (1990), 217–247.
- S00. A. Sevostyanov, Quantum deformation of Whittaker modules and the Toda lattice, *Duke Math. J.* **105** (2000), 211–238.
- S88. E.K. Sklyanin, Boundary conditions for integrable quantum systems, *J. Phys. A* **21** (1988), 2375–2389.
- S90. Y.B. Suris, Discrete time generalized Toda lattices: complete integrability and relation with relativistic Toda lattices, *Phys. Lett. A* **145** (1990), 113–119.
- S75. G. Szegő, *Orthogonal Polynomials*, Fourth Edition, American Mathematical Society, Colloquium Publications, vol. XXIII, American Mathematical Society, Providence, R.I., 1975.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD DE TALCA, CASILLA 747, TALCA, CHILE  
Email address: diejen@inst-mat.otalca.cl