# Foundations for an Abstract Proof Theory in the Context of Horn Rules

Tim S. Lyon<sup>1</sup> and Piotr Ostropolski-Nalewaja<sup>1,2</sup> <sup>1</sup>Computational Logic Group, TU Dresden <sup>2</sup>University of Wrocław

Abstract—We introduce a novel, logic-independent framework for the study of sequent-style proof systems, which covers a number of proof-theoretic formalisms and concrete proof systems that appear in the literature. In particular, we introduce a generalized form of sequents, dubbed g-sequents, which are taken to be binary graphs of typical, Gentzen-style sequents. We then define a variety of *inference rule types* as sets of operations that act over such objects, and define abstract (sequent) calculi as pairs consisting of a set of g-sequents together with a finite set of operations. Our approach permits an analysis of how certain inference rule types interact in a general setting, demonstrating under what conditions rules of a specific type can be permuted with or simulated by others, and being applicable to any sequentstyle proof system that fits within our framework. We then leverage our permutation and simulation results to establish generic calculus and proof transformation algorithms, which show that every abstract calculus can be effectively transformed into a lattice of polynomially equivalent abstract calculi. We determine the complexity of computing this lattice and compute the relative sizes of proofs and sequents within distinct calculi of a lattice. We recognize that top elements in lattices correspond to nested sequent systems, while bottom elements correspond to labeled sequent systems, and observe that top and bottom elements coincide with many known (cut-free) nested and labeled sequent systems for logics characterized by Horn properties.

## I. INTRODUCTION

Proof calculi are indispensable tools in the theory and application of logics, serving as engines that facilitate reasoning within a given logical paradigm. Of particular importance are sequent-style calculi, which were first introduced by Gentzen in the 1930s [1], [2]. Gentzen's sequent systems consist of inference rules, which operate over formulae called sequents, i.e. formulae of the form  $\varphi_1, \ldots, \varphi_n \Rightarrow \psi_1, \ldots, \psi_k$  with  $\varphi_i$ and  $\psi_i$  logical formulae, being used to derive theorems of a specified logic. A crucial feature of Gentzen's sequent calculi is that they exhibit the so-called *sub-formula property*, meaning every formula occurring in the premise of an inference rule is a sub-formula of one occurring in the conclusion of the rule. This feature, and the sequent formalism more generally, proved to be fruitful from both a theoretical and practical standpoint, being used to supply proof systems for a wide array of logics [3], [4], [5], [6], to discover new logics [7], to establish properties of logics (e.g. interpolation [8]), and to automate reasoning with logics [9], [10].

Yet, the discovery of new, expressive logics (e.g. the modal logic S5 and bi-intuitionistic logic) led to the realization that the sequent formalism was *too strict* as sequent calculi exhibiting the sub-formula property remained elusive [11],

[12]. In response, a variety of formalisms extending Gentzen's traditional sequent formalism were introduced to recapture the sub-formula property; e.g. hypersequents were introduced as multisets of sequents [13], [14], 2-sequents/linear nested sequents were introduced as lines of sequents [15], [16], nested sequents were introduced as trees of sequents [17], [18], and labeled sequents were introduced [19], [20], being interpretable as binary graphs of sequents [21], [22].

Such proof systems have found a wide array of applications for diverse classes of logics, being used in the design of interpolant construction algorithms [23], [24], [25], in writing decision algorithms (with counter-model extraction) [26], [27], [28], and have been applied in knowledge integration scenarios [29]. Nevertheless, it was found that differing formalisms possessed distinct advantages over one another; e.g. nested calculi were found to be suitable for writing proof-search and decision algorithms [28], [29], whereas labeled calculi were found to admit algorithmic construction [30].

Naturally, the arrival of new sequent-based formalisms gave rise to questions concerning their relationships: How are calculi in one formalism transformed into 'deductively equivalent' calculi in another? What are the relative sizes of proofs and sequents in one formalism compared to another? Under what conditions are proofs transformable between formalisms and what are the complexity bounds thereof? Such questions have typically been investigated in restricted concrete settings, focusing on specific sequent-based calculi for known classes of logics [21], [31], [32], [16], [33]. In contrast, we propose an alternative methodology, designing a novel *abstract framework* for the general study of structural sequent calculi.<sup>1</sup>

In particular, we formulate calculi as pairs consisting of (1) a set of objects called *generalized sequents* accompanied by (2) a finite set of inference rules. We therefore shift our attention from proof systems for logics, and instead, focus on proof systems in and of themselves, yielding a *logic independent* approach for studying the properties of, and relationships between, sequent-style systems. Due to its generality, our framework subsumes the various sequent-based formalisms mentioned above, and our results hold for any logic or sequent-style system that can be viewed as an object in our framework. Specifically, we accomplish the following:

• We generalize the notion of *sequent* to a graph of Gentzen-

<sup>&</sup>lt;sup>1</sup>Due to the duality exhibited between sequent systems and tableaux [34], our work is also applicable to the latter.

style sequents, referred to as *g-sequents*, which cover various kinds of sequents (e.g. labeled, nested, linear nested) that commonly appear in proof-theoretic works.

• We generalize inference rules to select *inference rule types* that operate over g-sequents, revealing the critical components that constitute an inference rule. These inference rule types subsume standard inference rules for sequent-style systems.

• We define a generic notion of calculus, so that our results hold generally for all sequent-style calculi that can be viewed as objects within our framework.

• Our abstract calculi include structural rules that facilitate reasoning with Horn properties. Therefore, a sizable number of (non-)classical logics, semantically characterized by 'Horn' frame conditions, and their accompanying sequent-based systems are subsumed by our work. Examples of logics with proof systems covered by our framework can be viewed in Figure 1.

• We define proof transformation notions (e.g. *permutation* and *simulation*) as well as explain how to strengthen or weaken certain rules (via notions called *absorption* and *fracturing*), which are used to provide generic calculus and derivation transformation algorithms and to compute complexity bounds thereof. This work contributes to a better understanding of how structural rules are eliminated from proofs, and how reachability and propagation rules [35], [22] arise from this process.

• We discover that every abstract calculus exists within a finite lattice of 'polynomially equivalent' abstract calculi, which we show how to compute. We observe that the top and bottom of a lattice is one of two calculus types, which we call *implicit* and *explicit* calculi, respectively. When we instantiate a lattice with known sequent-style systems, we find that nested calculi [17], [18] serve as the top element, whereas labeled calculi [19], [20] serve as the bottom element, establishing that nested and labeled calculi are *dual*.

Our abstract approach has explanatory value, yielding deep insights into the nature of, and connection between, sequentbased systems. Moreover, we provide a widely applicable toolkit for the manipulation of proofs and proof systems.

This paper is organized as follows: In Section II, we explain how our framework was designed by abstracting general underlying patterns appearing in calculi, considering various inference rule types and proof manipulation techniques. In Section III, we define our framework, and then put it to use in Section IV to establish a large number of permutation and simulation relationships between various inference rule types. In Section V, we demonstrate how abstract calculi can be converted into lattices of polynomially equivalent calculi. Finally, in Section VI, we discuss how nested and labeled systems can be identified with top and bottom elements of lattices, and discuss avenues for future research. Due to space constraints, we defer all proofs to the appendix.

#### II. OVERVIEW OF OUR APPROACH

We now turn our attention toward explaining our abstract proof-theoretic framework, arising from the study of numerous calculi, and the formalization of underlying patterns. While

Logics	Labeled/Explicit	Nested/Implicit
Bi-intuitionistic Logic	[36]	[33]
(Deontic) STIT Logics	[37], [22]	[27], [22]
FO Modal Logics	[38], [39]	[39]
FO Intuitionistic Logics	[40], [22]	[41], [22]
Provability Logic	[38]	[42]
Intuitionistic Modal Logics	[19]	[43], [44]
Tense/Grammar Logics	[45], [22]	[46], [28]

Fig. 1. Logics and associated sequent-style calculi covered by our abstract framework. Citations to papers with (cut-free) labeled and nested sequent systems thereof are provided in the second and third columns.

it is infeasible to thoroughly describe this lengthy process, one can provide a narrated walk-through of this work for a single (labeled) sequent calculus. This is the purpose of this section. It is meant to make the reader adopt a certain mindset by following a concrete example foreshadowing our general framework defined in Section III. It is worth noting, that while this section provides a description of the aforementioned process, Section III reflects its entirety by providing the end result. Thus, it might be worthwhile to consult the exact definitions in the following section, if the reader wishes to.

We have chosen a fragment of the labeled sequent calculus G3I [38] for propositional intuitionistic logic to use as our running example. We denote this fragment by G3l', and define it to be the set of rules shown in Figure  $2.^2$  This calculus avoids the unnecessary complexities of other sequent-based systems, while also possessing revelatory attributes that justify concepts later defined within our framework.

#### A. The Structure of Sequents and Inference Rules

As mentioned in Section I, generalizations of Gentzenstyle sequents (we hitherto refer to *Gentzen-style sequents* as *sequents*) take various forms, typically being types of graphs with sequents as vertices. These may take the form of general graphs [19], polytrees [21], trees [18], lines [15], or points, yielding standard sequents [1], [2]; e.g. the labeled sequents used in G3l' take the form of graphs, as explained below.

Labeled sequents in G3I' are objects of the form  $\mathcal{R}, \Gamma \vdash \Delta$ , where  $\mathcal{R}$  is a set of *relational atoms* (or, *edges*) of the form wEu and  $\Gamma, \Delta$  are multisets of *labeled formulae* of the form  $w: \varphi$ , which use a set  $\{w, u, v, \ldots\}$  of *labels*. In G3I', labeled formulae employ formulae from the language  $\mathcal{L}$  of propositional intuitionistic logic, generated via the following grammar in BNF:  $\varphi ::= p \mid \bot \mid \varphi \lor \varphi \mid \varphi \land \varphi \mid \varphi \supset \varphi$ , where p ranges over a set of propositional variables.



We observe that each labeled sequent can be viewed as a binary graph of sequents, obtained by depicting all labels as vertices, all relational atoms as edges, and all labeled formulae as sequents labeling nodes (cf. [22]). For example, the graph shown to the left

 $^{2}$ We employ a slight variation of the notation used for labeled sequents in [38] to better fit within the notation of our framework.

$$\frac{\overline{\mathcal{R}, wEu, \Gamma, w: p \vdash u: p, \Delta}}{\mathcal{R}, wEu, \Gamma, w: \varphi \vdash \Delta} (id) \quad \frac{\mathcal{R}, \Gamma, w: \varphi \vdash \Delta}{\mathcal{R}, \Gamma, w: \varphi \lor \psi \vdash \Delta} (\lor_L) \\
\frac{\mathcal{R}, wEu, \Gamma, w: \varphi \supset \psi \vdash u: \varphi, \Delta}{\mathcal{R}, wEu, \Gamma, w: \varphi \supset \psi \vdash \Delta} (\supset_L) \quad \frac{\mathcal{R}, wEw, \Gamma \vdash \Delta}{\mathcal{R}, \Gamma \vdash \Delta} (ref) \\
\frac{\overline{\mathcal{R}, wEu, uEv, wEv, \Gamma \vdash \Delta}}{\mathcal{R}, wEu, uEv, \Gamma \vdash \Delta} (tra) \quad \frac{\mathcal{R}, wEu, \Gamma, u: \varphi \vdash u: \psi, \Delta}{\mathcal{R}, \Gamma \vdash w: \varphi \supset \psi, \Delta} (\supset_R)^{\dagger}$$

Fig. 2. Some inference rules from the labeled calculus G3I for propositional intuitionistic logic [38]. We let G3I' denote the collection of the above rules. The side condition  $\dagger$  stipulates that the rule is applicable only if u is fresh, i.e. u does not occur in the surrounding context  $\mathcal{R}, \Gamma, \Delta$ .

corresponds to the labeled sequent  $\mathcal{R}, \Gamma \vdash \Delta$ , given that we let  $\mathcal{R} = wEw, wEz, wEv, wEu, uEv, \Gamma = w : \varphi, u : \chi, z : \psi$ , and  $\Delta = w : \psi, z : \chi, z : \xi$ . Every labeled sequent can be rewritten in an equivalent form  $\mathcal{R} \vdash \Sigma$ , where  $\mathcal{R}$  is a set of relational atoms as before, but  $\Sigma$  is a set of prefixed sequents; e.g. the labeled sequent  $\mathcal{R}, \Gamma \vdash \Delta$  depicted above, can we written as  $\mathcal{R} \vdash \Sigma$ , where

$$\Sigma = w : (\varphi \Rightarrow \psi), z : (\psi \Rightarrow \chi, \xi), u : (\chi \Rightarrow \emptyset), v : (\emptyset \Rightarrow \emptyset).$$

We view this perspective of labeled (or, graphical) sequents as beneficial for a couple reasons: (1) The internal structure of a sequent is *logic-dependent*; e.g. certain intuitionistic logics may restrict the succedent to at most one formula [1], [2] or certain sub-structural logics may employ sequences of formulae as opposed to (multi)sets in the antecedent or succedent [7]. As we are interested in providing a generic framework that studies the graphical properties of 'generalized or graphical sequents' and their associated proof systems, we may simply view sequents as types of labels. We therefore define the notion of a generalized sequent (g-sequent for short) as a graph of 'sequents' without specifying the internal structure of such sequents, yielding a logic-independent study of sequentstyle systems, as presented in Section III. (2) As mentioned above, various sequent-style formalisms beget proof systems that operate over certain types of graphs of sequents. Thus, our notion of g-sequent captures all such formalisms uniformly as restricting the g-sequents used yields a certain formalism.

We also find it important to clarify the connection between our notation and the notation typically used in labeled sequents, as this demonstrates how various sequent-style systems can be viewed as objects within our framework. Thus, we will view labeled sequents in the examples below as binary graphs of sequents. Moreover, we remark that a variety of works clarify how generalized versions of sequents correspond to graphs of sequents and how labeled sequents subsume such formalisms [21], [32], [47], [22], letting one view various proof systems as systems within our formalism.

Since we have adopted the view that labeled sequents are graphs of sequents, we reinterpret the labeled sequents and inference rules of G3l' in light of this perspective. We now take a labeled sequent to be an object of the form  $\mathcal{R} \vdash \Sigma$ such that  $\mathcal{R}$  is a set of edges (i.e. relational atoms) as before and  $\Sigma$  is a set of *prefixed sequents*, which are of the form  $w: (X \Rightarrow Y)$  with  $X = \varphi_1, \ldots, \varphi_n$  and  $Y = \psi_1, \ldots, \psi_k$ multisets of intuitionistic formulae. We rewrite the inference rules of G3I' in this notation and find certain commonalities among sets of rules in G3I' that give rise to various notions of *inference rule types*, of which we now elaborate.

**Initial Rules.** In our new notation, the (id) rule takes the form shown below.

$$\mathcal{R}, wEu \vdash \Sigma, w : (X, p \Rightarrow Y), u : (X' \Rightarrow p, Y')$$
<sup>(id)</sup>

We ask: what are the features of an inference rule that make it initial? Obviously, they are free of premises, and dictate what is taken to be axiomatic. Second, we observe that such rules may rely on the existence of relational data; e.g. in the (id) rule above, an edge wEu connecting one sequent of a certain type to another sequent of a certian type must be present. That is, (*id*) is subject to a *structural constraint*, which we may formalize as a labeled graph of the form  $C = (\{w, u\}, \{(w, u)\}, L)$  such that L(w, u) = E. One may verify that an instance of (id) satisfies such a constraint in the sense that any instance of (id) can be 'pattern matched' to such a constraint (with the edge (w, u) being associated with wEu). Third, we notice that although structural constraints appear to be critical features of initial (or, inference) rules, such objects are not enough to clearly express the operation of (id). We also require that prefixed sequents satisfy a certain relation, which we refer to as a sequent constraint. For example, in the (id) rule above, a relation R is additionally required, where  $R(S_1, S_2)$  holds with  $S_1 = X_1 \Rightarrow Y_1$  and  $S_2 = X_2 \Rightarrow Y_2$  iff  $p \in X_1$  and  $p \in Y_2$ .

**Local Rules.** We define a *local rule* to be an inference rule that only operates on sequents at a specific label. For example, if we rewrite  $(\vee_L)$  in our notation, the rule becomes:

$$\frac{\mathcal{R} \vdash \Sigma, w : (X, \varphi \Rightarrow Y) \quad \mathcal{R} \vdash \Sigma, w : (X, \psi \Rightarrow Y)}{\mathcal{R} \vdash \Sigma, w : (X, \varphi \lor \psi \Rightarrow Y)} (\lor_L)$$

Observe that this rule is local in the sense that it only manipulates data occurring in sequents at the label w. As in the initial rule case above, we recognize that a sequent constraint is required to fully specify the operation of the  $(\vee_L)$  rule: for sequents  $S_1 = X_1 \Rightarrow Y_1$ ,  $S_2 = X_2 \Rightarrow Y_2$ , and  $S_3 = X_3 \Rightarrow Y_3$ , we define  $R(S_1, S_2, S_3)$  iff  $\varphi \in X_1$ ,  $\psi \in X_2$ , and  $\varphi \lor \psi \in X_3$ . Such a relation must hold in any application of this rule for it to qualify as a valid rule application.

**Expansion Rules.** We classify *expansion rules* as inference rules that bottom-up introduce an edge (in  $\mathcal{R}$ ) to a fresh label,

thus expanding the graphical structure of the conclusion. The  $(\supset_R)$  rule serves as an example of an expansion rule, which takes the following form when rewritten in our notation:

$$\frac{\mathcal{R}, wEu \vdash \Sigma, w: (X \Rightarrow Y), u: (\varphi \Rightarrow \psi)}{\mathcal{R} \vdash \Sigma, w: (X \Rightarrow Y, \varphi \supset \psi)} (\supset_R)$$

Similar to the case of the  $(\lor_L)$  rule above, we observe that a sequent constraint must hold, specifying how the sequents at w and u in the premise relate to each other and the sequent at w in the conclusion.

**Transmission Rules.** We take a *transmission rule* to be an inference rule that updates two sequents connected by a single edge. The  $(\supset_L)$  rule serves as an example of a transmission rule, which takes the following form in our notation:

$$\begin{aligned}
\mathcal{G}_1 &= \mathcal{R}, \ wEu \vdash \Sigma, w : (X, \varphi \supset \psi \Rightarrow Y), u : (X' \Rightarrow \varphi, Y') \\
\mathcal{G}_2 &= \mathcal{R}, \ wEu \vdash \Sigma, w : (X, \varphi \supset \psi \Rightarrow Y), u : (X', \psi \Rightarrow Y') \\
\frac{\mathcal{G}_1 \qquad \mathcal{G}_2}{\mathcal{R}, \ wEu \vdash \Sigma, \ w : (X, \varphi \supset \psi \Rightarrow Y), \ u : (X' \Rightarrow Y')} (\supset_L)
\end{aligned}$$

Similar to the case of the (id) rule, the use of an edge (viz. wEu) in updating the sequents at w and u implies that structural constraints must be enforced on the left and right premises. In addition, a sequent constraint is required to fully specify that operation of the above rule, relating the sequents at w and u in the left and right premises with the sequents at w and u in the conclusion.

**Horn Rules.** In our setting we consider *Horn rules* to be inference rules that encode a Horn property, stipulating that if a certain sequence of edges exist in the conclusion of the rule, then a single type of edge must occur in the premise (cf. [20], [44]). Such rules serve as types of *structural rules* [21] or *relational rules* [20] existing in the literature. The (ref) and (tra) rules stand as examples of Horn rules, which take the following form in our notation:

$$\frac{\mathcal{R}, wEw \vdash \Sigma}{\mathcal{R} \vdash \Sigma} (ref) \quad \frac{\mathcal{R}, wEu, uEv, wEv \vdash \Sigma}{\mathcal{R}, wEu, uEv \vdash \Sigma} (tra)$$

The (ref) rule encodes reflexivity, by adding a single 'loop' (i.e. wEw) to the premise, whereas the the (tra) rule encodes transitivity, requiring a sequence of two edges (i.e. wEu, uEv), and connecting w to v via a single edge (i.e. wEv) in the premise. Both rules encode types of Horn conditions, and we note that their functionality can be specified without the use of constraints. As we discuss below, Horn rules can be 'absorbed' into the constraints associated with initial and transmission rules, producing new inference rules.

## B. Calculus Transformation and Rule Trading

Permutations arguments are at the heart of proof theory; e.g. Gentzen's celebrated cut-elimination theorem shows how the cut rule can be eliminated via permutations, yielding a proof exhibiting the sub-formula property [1], [2]. Likewise, simulations between sets of inference rules is of critical importance as they can be used to establish the 'relative strength' of proof systems and to establish the relative sizes of proofs. In Section IV, we will define these notions, using them to confirm a broad set of general relationships between rule types within our framework, and assisting us in writing generic algorithms (with complexity bounds) that transform calculi and their associated proofs.

We now exemplify simulations and permutations in the context of G3l'. In particular, we look at how initial, transmission, and Horn rules relate to one another. This investigation will demonstrate the connection between structural constraints and Horn rules, justifying their presence in our framework.

We begin by studying simulations between the initial rule (id) and the Horn rules (ref) and (tra), and look at the cases where the explicit edge wEu in (id) is 'active' in applications of (ref) and (tra). The first case yields a derivation of the following form:

$$\frac{\hline \mathcal{R}, wEw \vdash \Sigma, w: (X, p \Rightarrow p, Y)}{\mathcal{R} \vdash \Sigma, w: (X, p \Rightarrow p, Y)} \stackrel{(id)}{(ref)}$$

where as the second case yields a derivation of the form:

$$\frac{\mathcal{R}, wEu, uEv, wEv \vdash \Sigma'}{\mathcal{R}, wEu, uEv \vdash \Sigma'} (ta)$$

with  $\Sigma' = \Sigma, w : (X, p \Rightarrow Y), v : (X' \Rightarrow p, Y')$ . We observe that the conclusion in the (ref) case is similar to an instance of (id). However, whereas (id) requires the existence of prefixed sequents of the form  $w : (X, p \Rightarrow Y)$  and  $u : (X' \Rightarrow p, Y')$ connected by a single edge wEu, the conclusion of (ref)*identifies* these two sequents as  $w : (X, p \Rightarrow p, Y)$  and omits the occurrence of an edge. In the (tra) case, the conclusion of (tra) contains two prefixed sequents like (id), but with these two prefixed sequents connected by a path of edges wEu, uEv. Taking this into account, we recognize that we could simulate such derivations with a stronger form of (id)that absorbs the functionality of the (ref) and (tra) rules:

$$\mathcal{R} \vdash \Sigma, w : (X, p \Rightarrow Y), u : (X' \Rightarrow p, Y')$$
<sup>(id)</sup>

where (id)' is subject to the side condition that a path  $wEv_1, \ldots, v_{n-1}Eu$  of relational atoms of length 0 (meaning w = u) or greater exists between w and u. We can formalize this requirement as a structural constraint of the form  $C = (\{w, u\}, \{(w, u)\}, L)$  with  $L(w, u) \in \{\varepsilon, E, EE, \ldots\}$ , where  $\varepsilon$  is the empty string (meaning w = u), E is treated as a character, and each  $EE \cdots E$  is a word. Moreover, we require the same sequent relation to be enforced on (id)' just as it was with (id). We can take the conclusion of (ref) (in the derivation above) to be an instance of (id)' where  $L(w, u) = \varepsilon$ , the conclusion of a typical (id) rule to be an instance of (id)' where L(w, u) = E, and the conclusion of (tra) to be an instance of (id)' where L(w, u) = EE.

One can indeed show that any labeled sequent derivable by (id) followed by applications of (ref) or (tra) can be simulated by (id)' and vice-versa [40], [22]. Furthermore, this example justifies the inclusion of constraints in our framework as it shows that constraints can be modified, generating stronger inference rules, and forging new derivations that simulate others, effectively yielding new types of calculi.

We also observe a similar behavior when applying (ref)and (tra) to the transmission rule  $(\supset_L)$ . Let us consider applying the (ref) rule after an instance of  $(\supset_L)$  such that the relational atom 'active' in the latter is removed by (ref). We then have a derivation of the following form:

$$\begin{split} \mathcal{G}_1 &= \mathcal{R}, \, wEw \, \vdash \, \Sigma, \, w: (X, \varphi \supset \psi \Rightarrow \varphi, Y) \\ \mathcal{G}_2 &= \mathcal{R}, \, wEw \, \vdash \, \Sigma, \, w: (X, \varphi \supset \psi, \psi \Rightarrow Y) \\ \frac{\mathcal{G}_1 \quad \mathcal{G}_2}{\mathcal{R}, \, wEw \, \vdash \, \Sigma, \, w: (X, \varphi \supset \psi \Rightarrow Y)} \, \stackrel{(\supset_L)}{(ref)} \end{split}$$

Similar to the (*id*) case above, whereas  $(\supset_L)$  acts on prefixed sequents at w and u, separated by a single edge wEu, (ref)requires the identification of these two prefixed sequents as  $w: (X, \varphi \supset \psi \Rightarrow Y)$ . An investigation of applying (tra)to an instance of  $(\supset_L)$  would exhibit behavior as in the *(id)* case as well, where the two prefixed sequents are connected via a chain of relational atoms greater than one. We could therefore modify the constraints imposed on  $(\supset_L)$ , enforcing a constraint family  $C = (C_1, C_2)$  with  $C_1$  applied to the left premise and  $C_2$  applied to the right premise. Specifically, we let  $C_i = (\{w, u\}, \{(w, u)\}, L_i)$  such that  $i \in \{1, 2\}$  and  $L_i(w,u) \in \{\varepsilon, E, EE, \ldots\}$ . (NB. Although  $C_1 = C_2$  in this example, we allow constraints within a constraint family to differ in the general setting.) This constraint family can be imposed to define a new rule  $(\supset_L)'$ , which operates like  $(\supset_L)$ , but applies between sequents connected via a chain of relational atoms of length zero or greater. Using this modified rule, we find that the above derivation can be simulated by an application of (ref) followed by an application of  $(\supset_L)'$ , yielding a type of permutation, as shown below.

$$\mathcal{D}_{1} = \frac{\mathcal{R}, wEw \vdash \Sigma, w: (X, \varphi \supset \psi \Rightarrow \varphi, Y)}{\mathcal{R} \vdash \Sigma, w: (X, \varphi \supset \psi \Rightarrow \varphi, Y)} (ref)$$
$$\mathcal{D}_{1} = \frac{\mathcal{R}, wEw \vdash \Sigma, w: (X, \varphi \supset \psi, \psi \Rightarrow Y)}{\mathcal{R} \vdash \Sigma, w: (X, \varphi \supset \psi, \psi \Rightarrow Y)} (ref)$$
$$\frac{\mathcal{D}_{1} \quad \mathcal{D}_{2}}{\mathcal{R} \vdash \Sigma, w: (X, \varphi \supset \psi \Rightarrow Y)} (\supset_{L})'$$

If we replace (id) and  $(\supset_L)$  by (id)' and  $(\supset_L)'$  in G3l', we find that (ref) and (tra) can be permuted upward in any given derivation and ultimately eliminated [48], [40]. Rules such as (id)' and  $(\supset_L)'$  have been referred to as *reachability* (and in certain cases, *propagation*) rules [46], [22], and form a crucial component of our framework. Such rules witness the importance of structural constraints, and as we will show in Section IV, the interplay between constraints, reachability rules, and Horn rules uncover a wealth of permutation and simulation relationships between classes of inference rule types. Ultimately, in Section V, such rules will play a vital role, helping us identify spaces of polynomially equivalent calculi.

# III. ABSTRACT SEQUENT CALCULI

We now present our framework, and introduce the notion of a *generalized sequent* (or, *g-sequent* for short). These objects are graphs whose edges are labeled with characters and vertices are labeled with sequents. As we are in a general setting, and are interested in (the interaction between) inference rules that operate on such graphs, we do not discuss the internal structure of sequents, and thus, a *sequent* is merely a label in our context. The use of such objects is motivated by more expressive sequent systems, such as labeled calculi [19], [20] and nested calculi [17], [18], which can be seen as systems that reason over graphs of sequents, as previously explained.

After the introduction of g-sequents, we define various types of inference rules that typically appear in sequent-style proof systems. Certain classes of inference rules (e.g. reachability rules) require the use of *languages* in their constraints. We will use restricted versions of semi-Thue systems [49] to generate such languages and to facilitate our proof-theoretic study in subsequent sections.

#### A. Generalized Sequents

We let  $S = \{S_1, S_2, S_3, ...\}$  be a countably infinite set of *sequents*, which are denoted by S and annotated versions thereof. As sequents are taken to be atomic entities in our framework, we do not describe their internal structure. We let  $U = \{w, u, v, ...\}$  be the *universe*, whose entities are denoted by  $w, u, v, ...\}$  be the *universe*, whose entities are denoted by w, u, v, ... (potentially annotated), and which serve as vertices in the various graphs we define. Below, we define *gsequents* relative to a non-empty, finite set  $E = \{a, b, c, ...\}$  of *edge types*, which are used to index the edges of a g-sequent.

**Definition 1** (Generalized Sequent). A generalized sequent (g-sequent) is defined to be a tuple  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{L})$  such that

- $\mathcal{V} \subseteq \mathcal{U}$  is a (potentially empty) set of vertices;
- $\mathcal{E} = \{\mathcal{E}_a \mid a \in E\}$  with  $\mathcal{E}_a \subseteq \mathcal{V} \times \mathcal{V}$  for each  $a \in E$ ;
- $\mathcal{L}: \mathcal{V} \to S.$

We use  $\mathcal{G}$  (possibly annotated) to denote g-sequents, and let  $\mathfrak{G}(E)$  be the set of all g-sequents defined relative to a set E of edge types. For a g-sequent  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{L})$ , we let  $\mathcal{U}(\mathcal{G}) = \mathcal{V}$ .

As proof systems are concerned with the manipulation of syntactic entities via inference rules, we employ a more standard 'sequent-style' notation for g-sequents in our technical work. In particular, we use the equivalent notation  $\Gamma \vdash \Delta$  to denote a g-sequent  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{L})$ , where the *antecedent*  $\Gamma$  is a set of *edge atoms* of the form  $w\mathcal{E}_a u$  and the *succedent*  $\Delta$  is a set of *prefixed sequents* of the form w : S such that (1) for each  $a \in E$ ,  $w\mathcal{E}_a u \in \Gamma$  *iff*  $(w, u) \in \mathcal{E}_a$ , and (2)  $w : S \in \Delta$  *iff*  $\mathcal{L}(w) = S$ . We define the *size* of a g-sequent  $\mathcal{G} = \Gamma \vdash \Delta = (\mathcal{V}, \mathcal{E}, \mathcal{L})$  to be  $s(\mathcal{G}) = |\Gamma| + |\Delta| = |\mathcal{E}| + |\mathcal{V}|$ . Also, we let  $PS = \mathcal{U} \times S$  denote the set of prefixed sequents.

To improve intuition concerning g-sequents and their representations, we provide examples in Figure 3. We also specify a special subclass of g-sequents (whose importance will be discussed in Sections V and VI) referred to as *polytree gsequents*. A polytree g-sequent is a g-sequent that is (1)



Fig. 3. We give two examples of g-sequents:  $\mathcal{G} = (\mathcal{V}, \{\mathcal{E}_a, \mathcal{E}_b, \mathcal{E}_c\}, \mathcal{L})$  is shown left with  $\mathcal{V} = \{u_1, u_2, u_3, u_4\}, \mathcal{E}_a = \{(u_1, u_3), (u_2, u_4)\}, \mathcal{E}_b = \{(u_2, u_3)\}, \mathcal{E}_c = \{(u_1, u_2), (u_3, u_4), (u_4, u_4)\}, \text{ and for } i \in [4], \mathcal{L}(u_i) = S_i. \Gamma \vdash \Delta \text{ is shown right with } \Gamma = w_1 \mathcal{E}_c w_2, w_1 \mathcal{E}_b w_3, w_4 \mathcal{E}_b w_2, w_5 \mathcal{E}_a w_2 \text{ and } \Delta = w_1 : S_1, w_2 : S_2, w_3 : S_3, w_4 : S_4, w_5 : S_5.$ 

connected, and is (2) free of (un)directed cycles. Observe that the g-sequent shown right in Figure 3 is a polytree g-sequent.

#### B. E-Systems and Propagation

To control the functionality of certain inference rules, we make use of a restricted version of semi-Thue systems [49] that rewrite single edge types to strings of thereof.

Given a set A we define the set  $A^*$  of *strings* over A to be the set of finite sequences of elements of A including the *empty* string  $\varepsilon$ . We denote strings with (possibly annotated) letters s, t, r. A production rule is defined to be an object of the form  $p = s \longrightarrow s'$  such that  $s, s' \in A^*$ . We often use p and annotated versions thereof to denote production rules. A semi-Thue system is defined to be a finite set G of production rules. Semi-Thue systems permit us to derive strings via repeated applications of production rules. Given a semi-Thue system G over A, and a pair of strings  $t, t' \in A^*$  we write  $t \longrightarrow_{\mathbf{G}} t'$  iff there exists a rule  $s \longrightarrow s' \in \mathbf{G}$  such that s is a sub-string of t, and t' can be obtained from t by replacing some occurrence of s in t by s'. A **G**-derivation of a string  $t \in A^*$  from a string  $s \in A^*$ , denoted  $s \longrightarrow^*_{\mathbf{G}} t$ , is defined accordingly: (1)  $s \longrightarrow^*_{\mathbf{G}} t$ s, (2) if  $s \longrightarrow^*_{\mathbf{G}} r$  and  $r \longrightarrow_{\mathbf{G}} t$ , then  $s \longrightarrow^*_{\mathbf{G}} t$ . We define the *length* of a G-derivation of a string  $t \in A^*$  from a string  $s \in A^*$  to be the minimal number of rule applications used to derive t from s. The *language* of a string  $s \in A^*$  relative to a semi-Thue system **G** is defined as:  $\mathbf{G}(s) = \{t \mid s \longrightarrow_{\mathbf{G}}^{*} t\}.$ 

Let  $\overline{\mathbf{E}}$  be the following set  $\{\overline{a} \mid a \in \mathbf{E}\}$ . For a production rule of the form  $p = x \longrightarrow y_1 \cdots y_n$  with  $x, y_1, \ldots, y_n \in \mathbf{E} \cup \overline{\mathbf{E}}$ , we define  $\overline{p} = \overline{x} \longrightarrow \overline{y}_n \cdots \overline{y}_1$ , where  $\overline{z} = z$  for  $z \in \mathbf{E} \cup \overline{\mathbf{E}}$ . We define an E-system to be a semi-Thue system  $\mathbf{G}$  over  $\mathbf{E} \cup \overline{\mathbf{E}}$ satisfying: (1) for every rule  $s \longrightarrow t \in \mathbf{G}$  we have |s| = 1, and (2)  $s \longrightarrow t \in \mathbf{G}$  iff  $\overline{s} \longrightarrow \overline{t} \in \mathbf{G}$ . A production pair from  $\mathbf{G}$  is defined be a pair  $(p, \overline{p})$  such that  $p, \overline{p} \in \mathbf{G}$ . We define  $P(\mathbf{G})$  to be the set of all production pairs in  $\mathbf{G}$ . For a set Pof production pairs, we let  $\mathbf{G}(P)$  be the set of all rules found in a production pair of P.

Given a g-sequent  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{L})$ , two vertices  $u, w \in \mathcal{V}$ , and an element  $a \in \mathbf{E}$  we write  $\mathcal{G} \models u \stackrel{a}{\rightarrow} w$  *iff*  $(u, w) \in \mathcal{E}_a$ , and  $\mathcal{G} \models u \stackrel{\overline{a}}{\rightarrow} w$  *iff*  $(w, u) \in \mathcal{E}_a$ . Moreover, given a string  $xs \in (\mathbf{E} \cup \overline{\mathbf{E}})^*$  where  $x \in \mathbf{E} \cup \overline{\mathbf{E}}$ , we inductively define  $\mathcal{G} \models$   $u \underset{\sim}{xs} w$  as  $\exists_{v \in \mathcal{V}} \mathcal{G} \models u \underset{\sim}{x} v$  and  $\mathcal{G} \models v \underset{\sim}{s} w'$ , and  $\mathcal{G} \models u \underset{\sim}{x} v$  and  $\mathcal{G} \models v \underset{\sim}{s} w'$ . Additionally, when  $\mathcal{G}$  is clear from the context we may simply write  $u \underset{\sim}{s} w$  to express  $\mathcal{G} \models u \underset{\sim}{s} w$ . Finally, given a language  $\mathscr{L}$  (of some E-system) we use  $u \underset{\sim}{s} w$  iff there is a string  $s \in \mathscr{L}$  such that  $u \underset{\sim}{s} w$ .

#### C. Rules and Abstract Systems

Let us move on to discuss operations over g-sequents. We will define such operations in the format of inference rules as this will let us view our work more clearly in the context of structural proof theory. First, we define *structural constraints*, which are used to specify classes of g-sequents that are permitted to appear in specific inference rules.

**Definition 2** (Structural Constraint). Let E be a set of edge types. We define a *structural constraint* C to be a finite labeled tree (V, E, L) such that  $V \subseteq U, E \subseteq V \times V$ , and if  $(w, u) \in E$ , then  $L(w, u) = \mathbf{G}(a)$  for  $a \in \mathbf{E}$  and  $\mathbf{G}$  an E-system.

We define a *constraint family* to be a finite sequence  $C = (C_1, \ldots, C_n)$  of constraints, and we say that an E-system **G** participates in a constraint C = (V, E, L) iff there exists an  $(w, u) \in E$  and  $a \in E$  such that  $L(w, u) = \mathbf{G}(a)$ . Likewise, we say that an E-system **G** participates in a constraint family C iff there exists a constraint C in C such that **G** participates in C. We let  $\mathbf{G}(C) = \mathbf{G}_1 \cup \cdots \cup \mathbf{G}_n$  such that  $\mathbf{G}_1, \ldots, \mathbf{G}_n$  are all E-systems participating in C, and define the *size* of a constraint C as:  $|C| = |\mathbf{G}|$ .

**Definition 3** (Constraint Satisfaction). Let C = (V, E, L) be a constraint,  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{L})$  a g-sequent, and suppose  $V \subseteq \mathcal{V}$ . We say that  $\mathcal{G}$  satisfies C iff the following condition holds: if  $L(w, u) = \mathbf{G}(a)$ , then  $\mathcal{G} \models w \overset{\mathbf{G}(a)}{\leadsto} u$ .

**Definition 4** (Sequent Constraint). We define a *sequent constraint* R to be an (n + 1)-ary relation such that:

$$R \subseteq \underbrace{\mathbf{S} \times \cdots \times \mathbf{S}}_{n} \times 2^{\mathsf{PS}}.$$

We say that  $S_1, \ldots, S_n \in S$  and  $\Delta \subseteq PS$  satisfy R iff there exists a tuple  $(S_{k_1}, \ldots, S_{k_n}, \Delta) \in R$  with  $k_i, k_j \in \{1, \ldots, n\}$  and  $k_i \neq k_j$  for each  $1 \leq i < j \leq n$ ,

As certain inference rules in the literature are *context* dependent, e.g. the  $L\exists$  rule of Fitting [41], sequent constraints must take the entire succedent  $\Delta$  of a g-sequent into account in inference rule applications. This explains the presence of  $\Delta$  in sequent constraints. Note that we will hitherto refer to structural constraints as *constraints* more simply, while referring to sequent constraints as *sequent constraints*.

We now specify certain classes of inference rules, which will be collected together in finite sets to define our abstract calculi later on. Note that in the formulation of the inference rules below, for any constraint C = (V, E, L) parameterizing an inference rule, the vertices in V are always assumed to occur in the g-sequents of the rule, as specified by Definition 3. For inference rules with multiple premises, we use  $i \in [n]$  to mean  $1 \le i \le n$ .

**Initial Rule.** We define an *initial rule* to be an operation of the following form: i(C, R)

$$\Gamma \vdash \Delta i(C, F)$$

satisfying two conditions: (1) the g-sequent  $\Gamma \vdash \Delta$  satisfies the constraint C = (V, E, L), and (2) if  $\Gamma \vdash \Delta := (\mathcal{V}, \mathcal{E}, \mathcal{L})$ , then  $\mathcal{L}(w_1), \ldots, \mathcal{L}(w_n)$ , and  $\Delta' = (\Delta \setminus \{w_i : \mathcal{L}(w_i) \mid i \in [n]\})$ satisfy R, where  $V = \{w_1, \ldots, w_n\}$ . Examples of initial rules include init<sub>2</sub> in [24] and  $(\perp L)$  in [19].

Local Rule. We define a *local rule* to have the following form:

$$\frac{\{\Gamma \vdash \Delta, w : S_i\}_{i \in [n]}}{\Gamma \vdash \Delta, w : S} l(R)$$

such that (1)  $S_1, \ldots, S_n, S$  and  $\Delta$  satisfy R. Examples of local rules include  $(\neg \rightarrow)$  in [17] and CUT in [7].

Expansion Rule. We define an expansion rule as:

$$\frac{\Gamma, \Sigma \vdash \Delta, w : S_1, u : S_2}{\Gamma \vdash \Delta, w : S} e(R)$$

where (1)  $S_1$ ,  $S_2$ , S,  $\Delta$  satisfy the sequent constraint R, and (2)  $\mathcal{U}(\Gamma \vdash \Delta) \cap \mathcal{U}(\Sigma) = \{w\}$  with  $\Sigma \in \{w\mathcal{E}_a u, u\mathcal{E}_a w \mid a \in E\}$ . Examples of such rules are  $\Box R$  in [20] and [a] in [28].

Forward Horn Rule. If  $s = a_1 \cdots a_n \in (E \cup \overline{E})^*$ , then we define  $w\mathcal{E}_s u = w\mathcal{E}_{a_1}v_1, \ldots, v_{n-1}\mathcal{E}_{a_1}u$ , where  $w\mathcal{E}_{\overline{a}}u := u\mathcal{E}_a w$  and  $w\mathcal{E}_{\varepsilon}u = (w = u)$ . We define a *forward Horn rule* to be an operation of the form shown below left, which takes the form shown below right when  $s = \varepsilon$ .

$$\frac{\Gamma, w\mathcal{E}_s u, w\mathcal{E}_a u \vdash \Delta}{\Gamma, w\mathcal{E}_s u \vdash \Delta} h_f \quad \frac{\Gamma, w\mathcal{E}_a w \vdash \Delta}{\Gamma \vdash \Delta} h_f$$

For a production rule  $p = a \longrightarrow s$ , we define the singleton set  $H(p, \overline{p})$  to be the set containing the forward Horn rule above left, which takes the form above right when  $s = \varepsilon$ .

**Backward Horn Rule.** We define a *backward Horn rule* to be an operation of the form shown below left, which takes the form shown below right when  $s = \varepsilon$ .

$$\frac{\Gamma, w\mathcal{E}_s u, u\mathcal{E}_a w \vdash \Delta}{\Gamma, w\mathcal{E}_s u \vdash \Delta} h_b \quad \frac{\Gamma, w\mathcal{E}_a w \vdash \Delta}{\Gamma \vdash \Delta} h_b$$

For a production rule  $p = \overline{a} \longrightarrow s$ , we define the singleton set  $H(p,\overline{p})$  to be the set containing the backward Horn rule shown above left, which takes the form shown above right when  $s = \varepsilon$ . We define a *Horn rule* to be either a forward or backward Horn rule, and for a set P of production pairs, we let  $H(P) = \bigcup_{(p,\overline{p})\in P} H(p,\overline{p})$ . Examples of Horn rules include  $\chi_B$  in [19] and (Path) in [21].

We remark that Horn rules encode (universally closed) relational properties of the form  $w\mathcal{E}_s u \to w\mathcal{E}_x u$  with  $x \in E \cup \overline{E}$ , covering standard frame conditions for tense logics [46], first-order intuitionistic logics [50], and even agency logics [27].

**Reachability and Transmission Rules.** We define a *reachability rule* to be an operation of the following form:

$$\frac{\{\Gamma \vdash \Delta, w : S_i, u : S'_i\}_{i \in [n]}}{\Gamma \vdash \Delta, w : S, u : S'} r(\mathcal{C}, R)$$

such that (1)  $\mathcal{C} = (C_1, \ldots, C_n)$  and each  $C_i$  is of the form  $(\{w, u\}, \{(w, u)\}, L_i)$  with  $L_i(w, u) = \mathbf{G}_i(a_i)$  for some  $a_i \in \mathbf{E}$  and E-system  $\mathbf{G}_i$ , (2) the  $i^{th}$  premise satisfies  $C_i$ , and (3)  $S_1, S'_1, \ldots, S_n, S'_n, S, S'$  and  $\Delta$  satisfy R. Examples of reachability rules include  $\operatorname{Prop}(\mathbf{P})$  in [46] and  $(\forall_i^n)$  in [40].

We define a *transmission rule*  $t(\mathcal{C}, R)$  (as discussed in the previous section) to be a special instance of a reachability rule satisfying conditions (1) and (2) above, but where for every constraint  $C_i$ ,  $L_i(w, u) = \{a\}$  for some  $a \in E$ . Examples of transmission rules include  $\Diamond^\circ$  in [43] and Lift in [24].

We refer to any inference rule of the above form as either an *inference rule* or *rule*, more generally, and use  $\rho$ ,  $\sigma$ ,  $\tau$ , ... (potentially annotated) to denote them. For those inference rules parameterized by a constraint C or constraint family C, we say that an E-system *participates in the rule iff* the Esystem participates in the constraint C or constraint family C. Let us now define the notion of an *abstract calculus*.

**Definition 5** (Abstract Calculus). Let E be a set of edge types. We define an *abstract (sequent) calculus* (over E) to be an ordered pair  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  with  $\mathfrak{G}(E)$  the set of g-sequents defined relative to E and  $\mathfrak{R}$  a finite collection of inference rules. We use  $\mathfrak{A}, \mathfrak{B}, \mathfrak{C}, \ldots$  (occasionally annotated) to denote abstract calculi and define S(E) to be the collection of all abstract calculi over E. Furthermore, for an abstract calculus  $\mathfrak{A} = (\mathfrak{G}(E_1), \mathfrak{R}_1)$  and  $\mathfrak{B} = (\mathfrak{G}(E_2), \mathfrak{R}_2)$ , we say that  $\mathfrak{B}$  is an *extension* of  $\mathfrak{A}$ , and write  $\mathfrak{A} \subseteq \mathfrak{B}$ , *iff*  $E_1 \subseteq E_2$  and  $\mathfrak{R}_1 \subseteq \mathfrak{R}_2$ .

Given a set R of rules, we define a *derivation*  $\mathcal{D}$  to be any sequence of applications of rules in R to g-sequents in  $\mathfrak{G}(E)$ . If a g-sequent  $\mathcal{G}$  occurs in a derivation  $\mathcal{D}$ , then we write  $\mathcal{G} \in \mathcal{D}$ to indicate this. The *quantity* of a derivation  $\mathcal{D}$  is defined as  $q(\mathcal{D}) = |\{\mathcal{G} \in \mathfrak{G}(E) \mid \mathcal{G} \in \mathcal{D}\}|$  and the *size* of a derivation  $\mathcal{D}$ is defined to be  $s(\mathcal{D}) = \max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{D}\} \times q(\mathcal{D})$ .

A proof  $\mathcal{P}$  is defined to be a derivation beginning with applications of initial rules, and a *complete proof* is any proof ending with a g-sequent of the form w : S. Finally, a *polytree proof* is defined to be a proof such that every g-sequent occurring in the proof is a polytree g-sequent.

Two abstract calculi  $\mathfrak{A}, \mathfrak{B} \in \mathbb{S}(E)$  are defined to be *polynomially equivalent*, written  $\mathfrak{A} \dashv_p \mathfrak{B}$ , when a proof of a gsequent  $\mathcal{G}$  exists in  $\mathfrak{A}$  *iff* a proof  $\mathcal{P}'$  of  $\mathcal{G}$  exists in  $\mathfrak{B}$ , and there there exist PTIME functions f and g such that  $f(\mathcal{P}) = \mathcal{P}'$ and  $g(\mathcal{P}') = \mathcal{P}$ . We also lift specific set-theoretic operations to abstract calculi: for a set R of rules and an abstract calculus  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$ , we define  $\mathfrak{A} \setminus R := (\mathfrak{G}(E), \mathfrak{R} \setminus R)$  and  $\mathfrak{A} \cup R := (\mathfrak{G}(E), \mathfrak{R} \cup R)$ . Last, we let  $H(\mathfrak{A})$  denote the set of Horn rules in an abstract calculus  $\mathfrak{A}$ , and remark that H and annotated versions thereof will be exclusively used to denote sets of Horn rules throughout the remainder of the paper.

**Definition 6.** Let  $\rho \in \{i(C, R), r(\mathcal{C}, R)\}$ . We define the grammar  $\mathbf{G}(\rho)$  of  $\rho$  accordingly:  $p, \overline{p} \in \mathbf{G}(\rho)$  iff there exists an E-system  $\mathbf{G}$  that participates in  $\rho$  such that  $p, \overline{p} \in \mathbf{G}$ . For  $\rho \in \{l(R), e(R)\}$ , we define  $\mathbf{G}(\rho) = \emptyset$ . For a Horn rule  $h_f$  or  $h_b$ , we define the grammar  $\mathbf{G}(h_f) = \{a \longrightarrow s, \overline{a} \longrightarrow \overline{s}\}$  and  $\mathbf{G}(h_b) = \{\overline{a} \longrightarrow s, a \longrightarrow \overline{s}\}$ , respectively. Given a set of

rules  $R = \{\rho_1, \dots, \rho_n\}$ , we let  $G(R) = G(\rho_1) \cup \dots \cup G(\rho_n)$ . For an abstract calculus  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R}), G(\mathfrak{A}) = G(\mathfrak{R})$ .

Similarly, for a set R of rules, we define the set of production pairs of R as  $P(R) = P(\mathbf{G}(R))$ , and for an abstract calculus  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$ , we let  $P(\mathfrak{A}) = P(\mathbf{G}(\mathfrak{R}))$ .

#### IV. PERMUTATIONS AND SIMULATIONS

We now put our framework to use, studying the relationships between inference rules. A novel feature of our approach concerns the manipulation of constraints with E-systems via two operations, referred to as *absorb* and *fracture*. The absorb operation increases the expressiveness of constraints, whereas the fracture operation decreases the expressiveness of constraints. Increasing the expressiveness of a constraint in a rule secures new permutability relationships, while decreasing the expressiveness of a constraint, yields a weaker version that requires additional rules to recover its original functionality.

## A. Permutation and Absorption

We begin by defining the absorb operation, which 'adds' an E-system to the constraint of rule. We note that this operation only affects rules parameterized with constraints that associate E-systems with the edges of a constraint, namely, the i(C, R) and r(C, R) rules (see Section III). As local, expansion, and Horn rules omit the use of constraints entirely, such inference rules would be unaffected by the absorb operation, and thus, we disregard the absorb operation in these cases.

**Definition 7** (Absorb). We define the *absorb* operation between a constraint C = (V, E, L) and a E-system **G** denoted  $C \oplus \mathbf{G}$ , as the constraint (V, E, L') such that for each  $(w, u) \in$  $E, L'(w, u) = (\mathbf{G}' \cup \mathbf{G})(a)$  *iff*  $L(w, u) = \mathbf{G}'(a)$ . For a constraint family  $\mathcal{C} = (C_1, \ldots, C_n), \mathcal{C} \oplus \mathbf{G} = (C_1 \oplus \mathbf{G}, \ldots, C_n \oplus$ **G**). We lift the absorb operation from constraints to initial and reachability rules as follows:  $i(C, R) \oplus \mathbf{G} = i(C \oplus \mathbf{G}, R)$  and  $r(\mathcal{C}, R) \oplus \mathbf{G} = r(\mathcal{C} \oplus \mathbf{G}, R)$ .

We now define the notion of *permutation* in our setting, clarifying what it means for two rule sets to be permutable with one another.

**Definition 8** (Permutation). Let E be a set of edge types, and  $R_1$  and  $R_2$  be two sets of rules. We say that  $R_1$  *permutes above*  $R_2$ , written  $R_1 \nearrow R_2$ , *iff* for any g-sequents  $\mathcal{G}, \mathcal{G}_1, \ldots, \mathcal{G}_n \in \mathfrak{G}(E)$ , if  $\mathcal{G}$  can be derived via an application of a rule  $\sigma \in R_2$  followed by an application of a rule  $\rho \in R_1$  from  $\mathcal{G}_1, \ldots, \mathcal{G}_n$ , then  $\mathcal{G}$  can be derived via an application of  $\rho$  followed by an application of  $\sigma$  from  $\mathcal{G}_1, \ldots, \mathcal{G}_n$ .

If  $R_1 \nearrow R_2$  and  $R_2 \nearrow R_1$ , then we say that  $R_1$  and  $R_2$  are *permutable* with one another, and write  $R_1 \rightleftharpoons R_2$ . We note that when  $R_1$  or  $R_2$  is a singleton (i.e. a single rule  $\rho$ ), we simply write the rule name  $\rho$  in the notation defined above.

We now present a sequence of permutation results. In particular, we find that Horn rules are permutable with local rules (Theorem 9), Horn rules can always be permuted above expansion rules (Theorem 10), and Horn rules are always permutable with reachability rules, given that such Horn rules have been absorbed into their constraints (Theorem 11). For the remainder of the section, we fix a set E of edge types, and consider relationships between rules that participate in abstract calculi of S(E), unless specified otherwise.

**Theorem 9.** If l(R) is a local rule and H is a set of Horn rules, then  $l(R) \rightleftharpoons$  H.

**Theorem 10.** If e(R) is an expansion rule and H is a set of Horn rules, then H  $\nearrow e(R)$ .

**Theorem 11.** If  $r(\mathcal{C}, R)$  is a reachability rule, and H is a set of Horn rules, then  $r(\mathcal{C}, R) \oplus \mathbf{G}(H) \rightleftharpoons H$ .

## B. Simulation and Fracturing

We now introduce the fracture operation, which under certain conditions, functions as the inverse of the absorb operation, thus weakening constraints on initial and reachability rules. Subsequently, we define the notion of simulation, and show how weakened variants of initial and reachability rules can be simulated with the help of Horn rules.

**Definition 12** (Fracture). We define the *fracture* operation between a constraint C = (V, E, L) and a E-system **G**, denoted  $C \ominus \mathbf{G}$ , to be the constraint (V, E, L') such that for each  $(w, u) \in E$ ,  $(\mathbf{G}' \setminus \mathbf{G})(a) = L'(w, u)$  iff  $\mathbf{G}'(a) = L(w, u)$ . For a constraint family  $\mathcal{C} = (C_1, \ldots, C_n)$ , we let  $\mathcal{C} \ominus \mathbf{G} =$  $(C_1 \ominus \mathbf{G}, \ldots, C_n \ominus \mathbf{G})$ . We lift the fracture operation from constraints to initial rules and reachability rules as follows:  $i(C, R) \ominus \mathbf{G} = i(C \ominus \mathbf{G}, R)$  and  $r(\mathcal{C}, R) \ominus \mathbf{G} = r(\mathcal{C} \ominus \mathbf{G}, R)$ .

It is straightforward to verify that absorbing a grammar into a transmission rule (discussed in Sections II and III), yields a reachability rule, and that 'fracturing' the grammar  $\mathbf{G}(r(\mathcal{C}, R))$  from a reachability rule  $r(\mathcal{C}, R)$ , gives a transmission rule.

**Proposition 13.** Let t(C, R) be a transmission rule, r(C, R) be a reachability rule, and H be a non-empty set of Horn rules.

- 1)  $t(\mathcal{C}, R) \oplus \mathbf{G}(H)$  is a reachability rule;
- 2)  $r(\mathcal{C}, R) \ominus \mathbf{G}(r(\mathcal{C}, R))$  is a transmission rule.

Moreover, one can confirm that under certain conditions, the absorb and fracture operations are inverses of one another, and exhibit the following properties.

**Lemma 14.** Let  $\rho \in \{i(C, R), r(\mathcal{C}, R)\}$  and H be a set of Horn rules. Then,

- 1)  $(\rho \oplus \mathbf{G}(\mathbf{H})) \ominus \mathbf{G}(\mathbf{H}) = \rho \ominus \mathbf{G}(\mathbf{H});$
- 2) if  $\mathbf{G}(\mathrm{H}) \cap \mathbf{G}(\rho) = \emptyset$ , then  $(\rho \oplus \mathbf{G}(\mathrm{H})) \ominus \mathbf{G}(\mathrm{H}) = \rho$ ;
- 3)  $(\rho \ominus \mathbf{G}(\mathbf{H})) \oplus \mathbf{G}(\mathbf{H}) = \rho \oplus \mathbf{G}(\mathbf{H});$
- 4) if  $\mathbf{G}(\mathbf{H}) \subseteq \mathbf{G}(\rho)$ , then  $(\rho \ominus \mathbf{G}(\mathbf{H})) \oplus \mathbf{G}(\mathbf{H}) = \rho$ .

We now define the simulation and bi-simulation relation between rule sets and abstract calculi. In the sequel, we state a variety of useful properties concerning such relations.

**Definition 15** (Simulation). Let E be a set of edge types, and  $R_1$  and  $R_2$  two sets of rules. We say that  $R_2$  *simulates*  $R_1$ , written  $R_1 \preceq R_2$ , *iff* for any g-sequents  $\mathcal{G}, \mathcal{G}_1, \ldots, \mathcal{G}_n \in \mathfrak{G}(E)$ ,

if  $\mathcal{G}$  is derivable from  $\mathcal{G}_1, \ldots, \mathcal{G}_n$  with  $R_1$ , then  $\mathcal{G}$  derivable from  $\mathcal{G}_1, \ldots, \mathcal{G}_n$  with  $R_2$ . If  $R_1 \leq R_2$  and  $R_2 \leq R_1$ , then we say that  $R_1$  and  $R_2$  *bi-simulate* each other and write  $R_1 \simeq R_2$ .

Let  $\mathfrak{A}_1 = (\mathfrak{G}(\mathsf{E}), \mathfrak{R}_1)$  and  $\mathfrak{A}_2 = (\mathfrak{G}(\mathsf{E}), \mathfrak{R}_2)$  be two abstract calculi. We say that  $\mathfrak{A}_2$  *simulates*  $\mathfrak{A}_1$ , and write  $\mathfrak{A}_1 \leq \mathfrak{A}_2$ , *iff*  $\mathfrak{R}_1 \leq \mathfrak{R}_2$ . We say that  $\mathfrak{A}_1$  *bi-simulates*  $\mathfrak{A}_2$ , and write  $\mathfrak{A}_1 \simeq \mathfrak{A}_2$ , *iff*  $\mathfrak{A}_1 \leq \mathfrak{A}_2$  and  $\mathfrak{A}_2 \leq \mathfrak{A}_1$ .

It is a basic exercise to establish the following properties:

**Lemma 16.** Let E and E' be sets of edge types,  $\mathfrak{A} \in \mathbb{S}(E)$  with  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$ , and  $\mathfrak{B} \in \mathbb{S}(E')$ . Then,

- 1)  $\leq$  is a pre-order over  $\mathbb{S}(E)$ ;
- 2)  $\simeq$  is an equivalence relation over  $\mathbb{S}(\mathbb{E})$ ;
- 3) if  $R_1 \subseteq \mathfrak{R}$  and  $R_1 \preceq R_2$  with  $R_2$  a set of rules, then  $\mathfrak{A} \preceq (\mathfrak{G}(\mathsf{E}), (\mathfrak{R} \setminus R_1) \cup R_2);$
- 4) if  $\mathfrak{A} \subseteq \mathfrak{B}$ , then  $\mathfrak{A} \preceq \mathfrak{B}$ .

To discuss simulations and properties thereof, we require the use of *ordered rule sets*. In essence, an ordered rule set is a set  $R = R_1 \cup \cdots \cup R_n$  of rules such that any derivation constructed with R must proceed in a certain order, being obtained by applying at least 0 or 1 rule applications from  $R_1$ , followed by at least 0 or 1 rule applications from  $R_2$ , etc.

**Definition 17** (Ordered Rule Sets). Let  $R_1, \ldots, R_n$  be sets of rules. We define  $\triangleright^{i_1}R_1 \cdots \triangleright^{i_n} R_n$  with  $i_j \in \{0, 1\}$  to be an *ordered rule set* such that any derivation constructed with the rules in  $R_1 \cup \cdots \cup R_n$  must proceed by first applying  $i_1$  or more applications of rules from  $R_1$ , followed by  $i_2$  or more applications of rules from  $R_2$ , etc. When a set of rules in an ordered rule set is a singleton  $\{\rho\}$ , we will simply write  $\rho$ .

The following lemma is straightforward to establish:

**Lemma 18.** Let  $R_1$  and  $R_2$  be two set of rules. Then,

- if  $\mathbf{R}_1 \nearrow \mathbf{R}_2$ , then  $(\triangleright^1 \mathbf{R}_2 \triangleright^1 \mathbf{R}_1) \preceq (\triangleright^1 \mathbf{R}_1 \triangleright^1 \mathbf{R}_2)$ ;
- for  $i \in \{0, 1\}$ ,  $\triangleright^1 \mathbf{R}_1 \triangleright^i \mathbf{R}_2 \preceq \triangleright^0 \mathbf{R}_1 \triangleright^i \mathbf{R}_2$ ;
- for  $i \in \{0,1\}$ ,  $\triangleright^i \mathbf{R}_1 \triangleright^1 \mathbf{R}_2 \preceq \triangleright^i \mathbf{R}_1 \triangleright^0 \mathbf{R}_2$ .

As stated in the lemma below, we find that applying the absorb operation to an initial or reachability rule  $\rho$  strengthens the rule in the sense that  $\rho \oplus \mathbf{G}(\mathbf{H})$  can simulate  $\rho$  for a set  $\mathbf{H}$  of Horn rules, and conversely, we find that the fracture operation weakens an initial or reachability rule. Moreover, the absorb operation satisfies a monotonicity property relative to the subset relation over Horn rules, while the fracture operation satisfies an antitonicity property, as expressed by the fourth claim of the following lemma.

**Lemma 19.** Let H and H' be two sets of Horn rules,  $H \subseteq H'$ ,  $\rho \in \{i(C, R), r(C, R)\}$ , and  $i, j \in \{0, 1\}$ . Then,

1)  $\rho \preceq \rho \oplus \mathbf{G}(\mathbf{H})$  and  $\rho \ominus \mathbf{G}(\mathbf{H}) \preceq \rho$ ; 2)  $(\triangleright^i \rho \triangleright^j \mathbf{H}) \preceq (\triangleright^i \rho \oplus \mathbf{G}(\mathbf{H}) \triangleright^j \mathbf{H})$ ; 3)  $(\triangleright^i \rho \ominus \mathbf{G}(\mathbf{H}) \triangleright^j \mathbf{H}) \preceq (\triangleright^i \rho \triangleright^j \mathbf{H})$ ; 4)  $\rho \oplus \mathbf{G}(\mathbf{H}) \preceq \rho \oplus \mathbf{G}(\mathbf{H}')$  and  $\rho \ominus \mathbf{G}(\mathbf{H}') \preceq \rho \ominus \mathbf{G}(\mathbf{H})$ .

The following theorem is crucial for our generic algorithms in Section V. The theorem states that any derivation consisting of initial rules followed by applications of Horn rules can be simulated by initial rules under absorption.

**Theorem 20.** If i(C, R) is an initial rule and H is a set of Horn rules, then  $(\triangleright^1 i(C, R) \triangleright^0 H) \preceq i(C, R) \oplus \mathbf{G}(H)$ .

Let us turn our attention toward investigating simulations in the presence of fracturing. When fracturing an initial or reachability rule  $\rho$  with an E-system  $\mathbf{G}(\mathbf{H})$  for  $\mathbf{H}$  a set of Horn rules, we find that  $\rho$  can be simulated by  $\rho \ominus \mathbf{G}(\mathbf{H})$  along with applications of other inference rules. Yet, it so happens that *dependencies* between Horn rules in  $\mathbf{H}$  are of importance when considering simulations in this context. Intuitively, one Horn rule  $h_1$  depends on another Horn rule  $h_2$  when an application of  $h_2$  produces a g-sequent  $\mathcal{G}$  such that  $h_1$  becomes applicable to it. For the interested reader, we have provided an example of Horn rules and dependencies (which are captured by the following notion of a *dependency graph*) in Example 49 of Section V. We now define these dependencies:

**Definition 21** (Dependency Graph). Let **G** be an E-system with  $(p, \overline{p})$  and  $(p', \overline{p}')$  distinct propagation pairs in  $P(\mathbf{G})$  such that  $p = x \longrightarrow s$  and  $p' = y \longrightarrow t$  with  $x, y \in \mathbf{E} \cup \overline{\mathbf{E}}$ . We say that  $(p', \overline{p}')$  depends on  $(p, \overline{p})$ , written  $(p, \overline{p}) \sqsubset (p', \overline{p}')$ , iff s or  $\overline{s}$  is of the form  $s_1ys_2$ . We define the dependency graph of **G** to be the pair  $\mathsf{DG}(\mathbf{G}) = (P, \sqsubseteq)$  such that  $P = P(\mathbf{G})$  and  $\sqsubseteq$  is the reflexive-transitive closure of  $\sqsubset$ .

For a set H of Horn rules, we define  $DG(H) = (H, \sqsubseteq')$ such that for  $h, h' \in H$ ,  $h \sqsubseteq' h'$  iff for  $(p, \overline{p}) \in P(h)$  and  $(p', \overline{p}') \in P(h')$ ,  $(p, \overline{p}) \sqsubseteq (p', \overline{p}')$  in  $DG(G(H)) = (P, \sqsubseteq)$ . To capture dependency graphs over E-systems and Horn rules in a uniform notation, we may denote them by  $DG = (V, \sqsubseteq)$ .

Of critical importance in dependency graphs is the notion of a *fracturable set*. In essence, for a dependency graph  $DG = (V, \sqsubseteq)$ , a fracturable set is a set  $V' \subseteq V$  of vertices such that every vertex  $v \in V'$  'sees' only vertices in V'.

**Definition 22** (Fracturable Set). Given a dependency graph  $DG = (V, \sqsubseteq)$  we say that a subset V' of V is *fracturable* when there are no  $\sqsubseteq$ -edges from V' to  $V \setminus V'$ , and we define a subset V'' of V to be *anti-fracturable iff* there exists a fracturable subset V' and  $V'' = V \setminus V'$ . Given any  $V' \subseteq V$  of a dependency graph  $DG = (V, \sqsubseteq)$  of an E-system or a set of Horn rules H, we let G(V') denote  $\bigcup_{(p,\overline{p})\in V'}\{p,\overline{p}\}$  and  $\bigcup_{h\in V'} G(h)$ , respectively. For a set H of Horn rules we say that a subset H' is *(anti-)fracturable iff* 

The following properties of (anti-)fracturable subsets are useful and follow from the above definition.

**Lemma 23.** Let  $DG = (V, \sqsubseteq)$  be a dependency graph. Then,

- 1) V and  $\emptyset$  are both (anti-)fracturable subsets of V;
- 2) if both V' and V'' are (anti-)fracturable subsets of V and  $V \setminus V'$  respectively, then  $V' \cup V''$  is (anti-)fracturable.

**Theorem 24.** If H is a set of Horn rules with H' a fracturable subset of DG(H), then  $H \setminus H' \nearrow H'$ .

**Definition 25** (Saturation). Let H be a set of Horn rules with  $h \in H$ . If h is of the form shown below left, we define the *inverse*  $\overline{h}$  of h to be the rule of the form shown below right:

$$\frac{\mathcal{G}}{\mathcal{G}'}h \qquad \frac{\mathcal{G}'}{\mathcal{G}}\overline{h}$$

We write  $\overline{h}(\mathcal{G}) = \mathcal{G}'$  to mean that an application of  $\overline{h}$  to  $\mathcal{G}$  produces  $\mathcal{G}'$ . A g-sequent  $\mathcal{G}$  is defined to be  $\overline{h}$ -saturated iff every application of  $\overline{h}$  to  $\mathcal{G}$  produces  $\mathcal{G}$ . We say that  $\overline{h}(\mathcal{G})$  is permissible iff  $\overline{h}(\mathcal{G})$  produces a g-sequent  $\mathcal{G}' \neq \mathcal{G}$ . We define  $\overline{\mathrm{H}} = \{\overline{h} \mid h \in \mathrm{H}\}$  and define a g-sequent  $\mathcal{G}$  to be  $\overline{\mathrm{H}}$ -saturated iff it is  $\overline{h}$ -saturated for every  $\overline{h} \in \overline{\mathrm{H}}$ .

We now provide a sequence of results of that will ultimately be used to show under what conditions initial and reachability rules can be simulated with 'weaker' variants along with applications of Horn rules. In what follows, we let  $\overline{H}(\mathcal{G})$  denote the  $\overline{H}$ -saturated g-sequent obtained by repeatedly applying all permissible applications of rules in  $\overline{H}$  to  $\mathcal{G}$ .

**Lemma 26.** If  $\mathcal{G}$  is a g-sequent and H is a finite set of Horn rules, then  $\overline{H}(\mathcal{G})$  is computable in PTIME and is  $\overline{H}$ -saturated.

**Observation 27.** Let H be a set of Horn rules,  $\mathcal{G} = \Gamma \vdash \Delta$  be a g-sequent, and  $\overline{H}(\mathcal{G}) = \Gamma' \vdash \Delta'$ . Then,  $\mathcal{U}(\mathcal{G}) = \mathcal{U}(\overline{H}(\mathcal{G}))$ ,  $\Gamma \subseteq \Gamma'$ , and  $\Delta = \Delta'$ .

**Lemma 28.** Let H be a set of Horn rules,  $\mathcal{G}$  be a g-sequent, and  $\mathbf{G} = \mathbf{G}(\mathrm{H})$ . For any  $s \in (\mathbf{E} \cup \overline{\mathbf{E}})^*$  and  $u, w \in \mathcal{U}(\mathcal{G})$  if  $\mathcal{G} \models w \underset{\sim}{\sim} u$ , then for every string t such that  $t \longrightarrow_{\mathbf{G}}^* s$ , we have  $\overline{\mathrm{H}}(\mathcal{G}) \models w \underset{\sim}{\leftarrow} u$ .

We introduce the *path weakening* rule (denoted pw) in the lemma below, which serves as a restricted form of the conventional weakening rule (cf. WrL in [20]), only permitting the introduction of single edges between vertices in a gsequent. By standard arguments, one can establish that pwis eliminable from any proof.

**Lemma 29.** The path weakening rule pw (shown below) is eliminable from any proof:

$$\frac{\Gamma\vdash\Delta}{\Gamma,\Sigma\vdash\Delta}pw$$

where  $\Sigma \in \{ w \mathcal{E}_a u \mid a \in \mathbf{E} \text{ and } w, u \in \mathcal{U}(\Gamma \vdash \Delta) \}.$ 

The two lemmas below follow from Observation 27 with the latter lemma also relying on Lemma 28, and are used to establish the two subsequent theorems.

**Lemma 30.** If H is a set of Horn rules, then  $\overline{H} \leq pw$ .

**Lemma 31.** Let  $\mathcal{G}$  be a g-sequent, C be a constraint, and  $H := H(\mathbf{G}(C))$ . If H' is a fracturable subset of H and  $\mathcal{G}$  satisfies C, then  $\overline{H'}(\mathcal{G})$  satisfies  $C \ominus \mathbf{G}(H')$ .

**Theorem 32.** Let i(C, R) be an initial rule with the set  $H := H(\mathbf{G}(i(C, R)))$ . If H' is a fracturable subset of H, then  $i(C, R) \preceq \triangleright^1 i(C, R) \ominus \mathbf{G}(H') \triangleright^0 H'$ .

**Theorem 33.** Let  $r(\mathcal{C}, R)$  be a reachability rule and let  $H := H(\mathbf{G}(r(\mathcal{C}, R)))$ . If H' is a fracturable subset of H, then  $r(\mathcal{C}, R) \preceq \triangleright^0 pw \triangleright^1 r(\mathcal{C}, R) \ominus \mathbf{G}(H') \triangleright^0 H'$ .

## V. GENERIC TRANSFORMATION ALGORITHMS

We now put our permutation and simulation results to work, yielding generic transformations that establish the polynomial equivalence between abstract calculi within certain spaces. What we discover is rather remarkable: we find that every abstract calculus sits within two isomorphic lattices (dubbed upward and downward spaces), with the top and bottom elements exhibiting unique properties. We call calculi that sit at that the top of a lattice *implicit* and call calculi that sit at the bottom of a lattice explicit. Implicit calculi perform Horn reasoning with constraints only, whereas explicit calculi perform such reasoning with Horn rules. In Section VI, we observe that for a diverse number of known logics their (cutfree) nested sequent calculi (cf. [17], [18]) are implicit, while their (cut-free) labeled sequent calculi are explicit (cf. [19], [20]), establishing that such proof systems are *dual* to one another.

#### A. Upward Spaces/Lattices and Explicit Calculi

We first investigate the *upward space* of an arbitrary abstract calculus, which is defined by taking an abstract calculus  $\mathfrak{A}$  and strengthening its rules with  $\mathbf{G}(\mathbf{H})$  for  $\mathbf{H}$  a specific set of Horn rules, while simultaneously removing  $\mathbf{H}$  from  $\mathfrak{A}$ . This operation permits us to define a partial order over the produced abstract calculi, which can be viewed as a complete lattice (see Theorem 36 below). We will utilize these lattices later to identify spaces of polynomially equivalent calculi. First, let us lift  $\oplus$  and  $\ominus$  to the level of calculi.

**Definition 34** (Calculus Absorption and Fracturing). Let  $\mathfrak{A} = (\mathfrak{G}(\mathbb{E}), \mathfrak{R})$  be an abstract calculus, **G** be an E-system, and  $\odot \in \{\oplus, \ominus\}$ . We define  $\mathfrak{A} \odot \mathbf{G}$  to be the same as  $\mathfrak{A}$ , but where each initial rule i(C, R) and reachability rule  $r(\mathcal{C}, R)$  in  $\mathfrak{R}$  is replaced by  $i(C, R) \odot \mathbf{G}$  and  $r(\mathcal{C}, R) \odot \mathbf{G}$ , respectively.

**Definition 35** (Upward Space). Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus. We define the *upward space*  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  inductively: (1)  $\mathfrak{A} \in \mathbf{S}$ , (2) if  $\mathfrak{B} \in \mathbf{S}$  with  $\mathsf{DG}(H) = (H, \sqsubseteq)$  the dependency graph of  $H := H(\mathfrak{B})$ , then for an anti-fracturable subset  $H' \subseteq H$ ,  $\mathfrak{C} = ((\mathfrak{B} \oplus \mathbf{G}(H')) \setminus H') \in \mathbf{S}$  and  $\mathfrak{B} \notin \mathfrak{C}$ .

**Theorem 36.** Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its upward space. Then, the  $\leqslant$  relation is a connected partial-order. Moreover, for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , if we take  $\mathfrak{B} \wedge \mathfrak{C} = \inf{\{\mathfrak{B}, \mathfrak{C}\}}$  and  $\mathfrak{B} \vee \mathfrak{C} = \sup{\{\mathfrak{B}, \mathfrak{C}\}}$  under  $\leqslant$ , then  $(\mathbf{S}, \wedge, \vee)$  is a complete lattice with  $\mathfrak{A} = \bot$ .

Given an abstract calculus  $\mathfrak{A}$  and its upward space  $\mathbb{U}(\mathfrak{A})$ , we find that any proof from  $\mathfrak{A}$  may be transformed along the partial order of  $\mathbb{U}(\mathfrak{A})$  in PTIME. We also establish that proofs may shrink when transformed along the partial order of  $\mathbb{U}(\mathfrak{A})$ .

**Theorem 37** (Up the Upward Space). Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its upward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{A}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{B}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') \leq s(\mathcal{P})$ .

```
Algorithm: IMPLICATE(\mathfrak{A})

Input: An explicit calculus \mathfrak{A}.

Output: The upward space \mathbb{U}(\mathfrak{A}).

Set \mathbb{U}(\mathfrak{A}) := (\mathbf{S}, \leqslant), \mathbf{S} := \{\mathfrak{A}\}, \text{ and } (\leqslant) := \emptyset;

While \mathbb{U}(\mathfrak{A}) grows;

For each \gtrless-maximal element \mathfrak{B} of \mathbb{U}(\mathfrak{A});

For each \gtrless-maximal element \mathfrak{B} of \mathbb{U}(\mathfrak{A});

For each anti-fracturable subset H' in DG(H(\mathfrak{B}));

Add f(\mathfrak{B}, \mathrm{H'}) to S and \mathfrak{B} \leqslant f(\mathfrak{B}, \mathrm{H'}) to (\gtrless);

Return \mathbb{U}(\mathfrak{A}).
```

Fig. 4. The IMPLICATE algorithm takes an explicit calculus as input and computes its upward space.

To transform proofs 'down' the partial order of an upward space  $\mathbb{U}(\mathfrak{A})$  into a proof of  $\mathfrak{A}$  requires an additional condition, namely, the initial and reachability rules of  $\mathfrak{A}$  must satisfy a certain set of equations. This gives rise to the notion of an *explicit calculus*. Intuitively, an explicit calculus is one where all initial and reachability rules are parameterized with minimal constraints, i.e. constraints C such that |C| = 0. This has the effect that if a proof utilizes Horn rules, then such rules cannot be eliminated from the proof as they cannot be 'mimicked' by other rules of the calculus.

**Definition 38** (Explicit Calculus). Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus. We define  $\mathfrak{A}$  to be *explicit iff* for every initial i(C, R) and reachability rule  $r(\mathcal{C}, R)$  in  $\mathfrak{R}$ , the following hold:

$$i(C, R) \ominus \mathbf{G}(\mathfrak{A}) = i(C, R) \text{ and } r(\mathcal{C}, R) \ominus \mathbf{G}(\mathfrak{A}) = r(\mathcal{C}, R).$$

**Theorem 39** (Down the Upward Space). Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$ be an explicit calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \ll)$  its upward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{B}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{A}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

#### B. Downward Spaces/Lattices and Implicit Calculi

Above, we investigated the upward spaces of calculi obtained by strengthening initial and reachability rules via absorbtion. Conversely, we obtain *downward spaces* by weakening initial and reachability rules via fracturing.

**Definition 40** (Downward Space). Let  $\mathfrak{A} = (\mathfrak{G}(\mathsf{E}), \mathfrak{R})$  be an abstract calculus. We define the *downward space*  $\mathbb{D}(\mathfrak{A}) =$  $(\mathbf{S}, \leqslant)$  inductively as follows: (1)  $\mathfrak{A} \in \mathbf{S}$ , (2) if  $\mathfrak{B} \in \mathbf{S}$  with  $\mathrm{H} := \mathrm{H}(\mathfrak{B})$  and  $\mathrm{DG}(\mathbf{G}) = (P, \sqsubseteq)$  the dependency graph of  $\mathbf{G} := \mathbf{G}(\mathfrak{B} \setminus \mathrm{H})$ , then for a fracturable subset  $P' \subseteq P$ ,  $\mathfrak{C} = ((\mathfrak{B} \ominus \mathbf{G}(P')) \cup \mathrm{H}(\mathbf{G}(P'))) \in \mathbf{S}$  and  $\mathfrak{C} \leqslant \mathfrak{B}$ .

As with upward spaces, we obtain that downward spaces are partially ordered sets, which can be viewed as lattices.

**Theorem 41.** Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its downward space. Then, the  $\leqslant$  relation is a connected partial order. Moreover, for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , if we take  $\mathfrak{B} \land \mathfrak{C} = \inf{\{\mathfrak{B}, \mathfrak{C}\}}$  and  $\mathfrak{B} \lor \mathfrak{C} = \sup{\{\mathfrak{B}, \mathfrak{C}\}}$  under  $\leqslant$ , then  $(\mathbf{S}, \land, \lor)$  is a complete lattice with  $\mathfrak{A} = \top$ .

Given an abstract calculus  $\mathfrak{A}$ , we can translate proofs 'down' the partial order of  $\mathbb{D}(\mathfrak{A})$  in PTIME, similar to Theorem 39.



Fig. 5. The EXPLICATE algorithm takes an implicit calculus as input and computes its downward space.

**Theorem 42** (Down the Downward Space). Let  $\mathfrak{A} = (\mathfrak{G}(\mathbb{E}), \mathfrak{R})$  be a calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its downward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{A}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{B}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

We find that transforming proofs 'up' the partial order of a downward space  $\mathbb{D}(\mathfrak{A})$  requires that  $\mathfrak{A}$  is of a specific form. Namely, we find that such proofs can be transformed when  $\mathfrak{A}$  is an *implicit calculus*, defined below.

**Definition 43** (Implicit Calculus). Let  $\mathfrak{A} = (\mathfrak{G}(\mathsf{E}), \mathfrak{R})$  be an abstract calculus. We define  $\mathfrak{A}$  to be *implicit iff*  $\mathfrak{A}$  does not contain any Horn rules, and every initial rule i(C, R)and reachability rule  $r(\mathcal{C}, R)$  in  $\mathfrak{R}$ , satisfies the following equations:

$$i(C, R) \oplus \mathbf{G}(\mathfrak{A}) = i(C, R) \text{ and } r(\mathcal{C}, R) \oplus \mathbf{G}(\mathfrak{A}) = r(\mathcal{C}, R).$$

**Theorem 44** (Up the Downward Space). Let  $\mathfrak{A} = (\mathfrak{G}(\mathbb{E}), \mathfrak{R})$ be an implicit calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its downward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{B}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{A}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') \leq s(\mathcal{P})$ .

Beyond their use in the theorem above, another interesting feature of implicit calculi is that every *complete proof* employs only g-sequents of a polytree shape.<sup>3</sup> Such calculi are reminiscent of nested sequent systems [17], [18], and later on, we will identify a sizable number of nested sequent systems appearing in the literature with implicit calculi.

**Theorem 45.** If  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  is an implicit calculus, then every complete proof is a polytree proof.

#### C. Generic Calculus Transformations

An interesting and new discovery that arises from our framework is that certain abstract calculi participate in lattices of polynomially equivalent calculi. These lattices can be identified by transforming an explicit calculus into its upward space, or by transforming an implicit calculus into its downward space. We provide two calculus transformation algorithms IMPLICATE and EXPLICATE, which take an abstract calculus as input, and compute its upward or downward space,

 $<sup>^{3}</sup>$ See Section III for the definition of complete proofs and polytree g-sequents.

effectively generating a lattice of polynomially equivalent calculi. To state these algorithms we employ the following notation:

**Definition 46.** Let  $\mathfrak{A}$  be an abstract calculus with H a set of Horn rules and *P* a set of production pairs. We define:

$$f(\mathfrak{A},\mathrm{H}):=(\mathfrak{A}\oplus\mathbf{G}(\mathrm{H}))\backslash\mathrm{H}\quad g(\mathfrak{A},P):=(\mathfrak{A}\ominus\mathbf{G}(P))\cup\mathrm{H}(P)$$

Our first calculus transformation algorithm IMPLICATE is presented in Figure 4. The algorithm is named 'IMPLICATE' as it successively computes better approximations of the implicit calculus  $\mathfrak{B}$  that is polynomially equivalent to the input. It is straightforward to verify that IMPLICATE terminates as every execution of the while-loop strictly reduces the finite set of Horn rules associated with all  $\ll$ -maximal calculi. The following relies on Theorems 37 and 39, and we remark that  $\top$  can be obtained from  $\mathfrak{A}$  in PTIME by computing  $\top := f(\mathfrak{A}, H(\mathfrak{A})).$ 

**Theorem 47.** Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an explicit calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \gtrless)$  its upward space. Then,

- 1)  $\mathbb{U}(\mathfrak{A}) = \text{IMPLICATE}(\mathfrak{A});$
- 2)  $\mathbb{U}(\mathfrak{A})$  is computable from  $\mathfrak{A}$  in EXPTIME;
- 3) If  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , then  $\mathfrak{B} \dashv \vdash_p \mathfrak{C}$ ;
- 4)  $\top$  is computable from  $\mathfrak{A}$  in PTIME;
- 5)  $\top$  is the only implicit calculus in  $\mathbb{U}(\mathfrak{A})$ .

Our second calculus transformation algorithm EXPLICATE is displayed in Figure 5. The algorithm is named 'EXPLICATE' since it successively computes better approximations of the explicit calculus polynomially equivalent to the input. We note that EXPLICATE depends on sets of production pairs, and observe that EXPLICATE terminates since each grammar  $G(\mathfrak{B} \setminus H(\mathfrak{B}))$  strictly decreases for each  $\leqslant$ -minimal calculus  $\mathfrak{B}$  after each execution of the while-loop.

The following theorem is similar to Theorem 47, but relies on Theorem 42 and 44 to establish the polynomial equivalence of all abstract calculi in the downward space. Moreover, we remark that  $\perp$  can be obtained from the input  $\mathfrak{A}$  in PTIME by computing  $\perp := g(\mathfrak{A}, P(\mathfrak{A})).$ 

**Theorem 48.** Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an implicit calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its downward space. Then,

- 1)  $\mathbb{D}(\mathfrak{A}) = \text{EXPLICATE}(\mathfrak{A});$
- 2)  $\mathbb{D}(\mathfrak{A})$  is computable from  $\mathfrak{A}$  in EXPTIME;
- 3) If  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , then  $\mathfrak{B} \dashv \vdash_p \mathfrak{C}$ ;
- 4)  $\perp$  is computable from  $\mathfrak{A}$  in PTIME;
- 5)  $\perp$  is the only explicit calculus in  $\mathbb{D}(\mathfrak{A})$ .

**Example 49.** To demonstrate the functionality of our two calculus transformation algorithms, we consider an example with an explicit calculus  $\mathfrak{A}$  consisting of the following rules, and where |C| = 0.

$$\frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta} i(C, R) = \frac{\Gamma, w\mathcal{E}_{c}u, u\mathcal{E}_{c}v, w\mathcal{E}_{a}v \vdash \Delta}{\Gamma, w\mathcal{E}_{c}u, u\mathcal{E}_{c}v \vdash \Delta} h_{1}$$

$$\frac{\Gamma, w\mathcal{E}_{a}w \vdash \Delta}{\Gamma \vdash \Delta} h_{2} = \frac{\Gamma, w\mathcal{E}_{a}u, u\mathcal{E}_{b}w \vdash \Delta}{\Gamma, w\mathcal{E}_{a}u \vdash \Delta} h_{3}$$



Fig. 6. The lattice above left corresponds to a computation of IMPLICATE( $\mathfrak{A}$ ), where  $\mathfrak{A}$  is defined as in Example 49. The lattice shown above right corresponds to a computation of EXPLICATE( $\mathfrak{B}$ ), where  $\mathfrak{B} = f(\mathfrak{A}, \{h_1, h_2, h_3\})$ .

We let  $H = \{h_1, h_2, h_3\}$  and observe that in the dependency graph DG(H) = (H,  $\sqsubseteq$ ),  $h_3 \sqsubseteq h_1$  and  $h_3 \sqsubseteq h_2$ . In this setting, IMPLICATE( $\mathfrak{A}$ ) constructs the lattice shown on the left of Figure 6, eventually yielding the implicit calculus  $f(\mathfrak{A}, \{h_1, h_2, h_3\}) = \mathfrak{B}$  at the top. Furthermore, if we let  $P(H) = P_1 \cup P_2 \cup P_3$  such that  $P_i = P(h_i) = \{(p_i, \overline{p}_i)\}$ , then in DG(P) = (P,  $\sqsubseteq'$ ),  $(p_3, \overline{p}_3) \sqsubseteq (p_1, \overline{p}_1)$  and  $(p_3, \overline{p}_3) \sqsubseteq$  $(p_2, \overline{p}_2)$ . If we run IMPLICATE( $\mathfrak{B}$ ), we obtain the lattice shown on the right of Figure 6.

**Theorem 50.** Let  $\mathfrak{A}$  be an explicit calculus,  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$ ,  $\mathfrak{B}$  be an implicit calculus, and  $\mathbb{D}(\mathfrak{B}) = (\mathbf{S}', \leqslant)$ . If either  $\mathfrak{B} \in \mathbf{S}$  or  $\mathfrak{A} \in \mathbf{S}'$ , then  $\mathbf{S} = \mathbf{S}'$ ,  $\mathbb{U}(\mathfrak{A}) \cong \mathbb{D}(\mathfrak{B})$ , and for any  $\mathfrak{C}, \mathfrak{D} \in \mathbf{S} = \mathbf{S}'$ ,  $\mathbf{G}(\mathfrak{C}) = \mathbf{G}(\mathfrak{D})$ .

#### VI. CONCLUDING REMARKS

Our novel framework for the study of sequent-style systems yields rich insights into the nature of numerous concrete proof systems that appear in the literature. As shown in Figure 1, diverse classes of logics with applications in computer science [51], [52], mathematics [42], and philosophy [53], [50] admit sequent-style calculi that can be viewed as instances of abstract calculi. More specifically, if we observe the labeled sequent systems of the logics in Figure 1 (operating over graphs of Gentzen-style sequents [21], [22]), we discover that such systems are instances of explicit calculi. In addition, due to a unique property of implicit calculi, namely, that every complete proof is a polytree proof (Theorem 45), one can identify nested sequents [17], [18], [21]) with implicit calculi.

When viewed through the lens of our theory, we thus find that labeled and nested sequent systems exist within a spectrum (or, lattice) of polynomially equivalent calculi. We therefore get the polynomial equivalence of nested and labeled calculi for free via our general results. Our theory also suggests that labeled and nested systems are *dual* to one another with labeled systems serving as the bottom of a lattice and nested systems serving as the top. This reveals that labeled and nested systems tend to come in pairs. Our work also confirms that nested sequent systems generally admit shorter proofs with syntactically simpler sequents than labeled systems as witnessed by Theorem 37 and 44. There are various avenues for future research. We could provide a more fine-grained treatment, taking the internal structure of sequents into account. Doing so, we could investigate the inter-workings of cut-elimination in a broad setting. Second, we could generalize the types of structural rules considered in our framework, moving beyond Horn rules. In fact, as discovered in [27], [22], certain proof systems utilizing disjunctive properties admit transformations similar to those in Section V. Examining these cases and incorporating them into our framework seems promising. Third, we could extend our methods to pinpoint how other types of proof systems arise (e.g. linear nested sequents), identifying the spaces these calculi exist within and uncovering transformations that navigate them.

## REFERENCES

- G. Gentzen, "Untersuchungen über das logische schließen. i," Mathematische zeitschrift, vol. 39, no. 1, pp. 176–210, 1935.
- [2] —, "Untersuchungen über das logische schließen. ii," Mathematische Zeitschrift, vol. 39, no. 1, pp. 405–431, 1935.
- [3] G. Corsi, "A cut-free calculus for dummett's lc quantified," *Mathematical Logic Quarterly*, vol. 35, no. 4, pp. 289–301, 1989.
- [4] H. B. Curry, "The elimination theorem when modality is present," *The Journal of Symbolic Logic*, vol. 17, no. 4, pp. 249–265, 1952. [Online]. Available: http://www.jstor.org/stable/2266613
- [5] C. Rauszer, "An algebraic and kripke-style approach to a certain extension of intuitionistic logic, dissertations mathematicae," 1980.
- [6] N. Tennant, "Natural deduction and sequent calculus for intuitionistic relevant logic," *The Journal of symbolic logic*, vol. 52, no. 3, pp. 665– 680, 1987.
- [7] J.-Y. Girard, "Linear logic," *Theoretical computer science*, vol. 50, no. 1, pp. 1–101, 1987.
- [8] S. Maehara, "On the interpolation theorem of craig," Sûgaku, vol. 12, no. 4, pp. 235–237, 1960.
- [9] R. Dyckhoff, "Contraction-free sequent calculi for intuitionistic logic," *The Journal of Symbolic Logic*, vol. 57, no. 3, pp. 795–807, 1992.
- [10] J. K. Slaney, "Minlog: A minimal logic theorem prover," in Automated Deduction - CADE-14, 14th International Conference on Automated Deduction, Townsville, North Queensland, Australia, July 13-17, 1997, Proceedings, ser. Lecture Notes in Computer Science, W. McCune, Ed., vol. 1249. Springer, 1997, pp. 268–271. [Online]. Available: https://doi.org/10.1007/3-540-63104-6\_27
- [11] L. Buisman and R. Goré, "A cut-free sequent calculus for bi-intuitionistic logic," in Automated Reasoning with Analytic Tableaux and Related Methods, N. Olivetti, Ed. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 90–106.
- [12] H. Wansing, "Sequent calculi for normal modal propositional logics," *Journal of Logic and Computation*, vol. 4, no. 2, pp. 125–142, 1994.
- [13] A. Avron, "The method of hypersequents in the proof theory of propositional non-classical logics," in *Logic: From Foundations to Applications: European Logic Colloquium*, W. Hodges, M. Hyland, C. Steinhorn, and J. Truss, Eds. USA: Clarendon Press, 1996, p. 1–32.
  [14] G. Pottinger, "Uniform, cut-free formulations of t, s4 and s5," *Journal*
- [14] G. Pottinger, "Uniform, cut-free formulations of t, s4 and s5," *Journal of Symbolic Logic*, vol. 48, no. 3, p. 900, 1983.
- [15] A. Masini, "2-sequent calculus: A proof theory of modalities," Annals of Pure and Applied Logic, vol. 58, no. 3, pp. 229–246, 1992.
- [16] B. Lellmann, "Linear nested sequents, 2-sequents and hypersequents," in Automated Reasoning with Analytic Tableaux and Related Methods, ser. Lecture Notes in Computer Science, H. De Nivelle, Ed., vol. 9323. Cham: Springer International Publishing, 2015, pp. 135–150.
- [17] R. A. Bull, "Cut elimination for propositional dynamic logic without \*," Z. Math. Logik Grundlag. Math., vol. 38, no. 2, pp. 85–100, 1992.
- [18] R. Kashima, "Cut-free sequent calculi for some tense logics," *Studia Logica*, vol. 53, no. 1, pp. 119–135, 1994.
- [19] A. K. Simpson, "The proof theory and semantics of intuitionistic modal logic," Ph.D. dissertation, University of Edinburgh. College of Science and Engineering. School of Informatics, 1994.
- [20] L. Viganò, Labelled Non-Classical Logics. Springer Science & Business Media, 2000.

- [21] A. Ciabattoni, T. Lyon, R. Ramanayake, and A. Tiu, "Display to labelled proofs and back again for tense logics," ACM Transactions on Computational Logic, vol. 22, no. 3, p. 1–31, 2021.
- [22] T. Lyon, "Refining labelled systems for modal and constructive logics with applications," Ph.D. dissertation, Technische Universität Wien, 2021.
- [23] R. Kuznets, "Proving craig and lyndon interpolation using labelled sequent calculi," in *Logics in Artificial Intelligence*, ser. Lecture Notes in Computer Science, L. Michael and A. Kakas, Eds., vol. 10021. Cham: Springer International Publishing, 2016, pp. 320–335.
- [24] R. Kuznets and B. Lellmann, "Interpolation for intermediate logics via hyper- and linear nested sequents," in Advances in Modal Logic 12, proceedings of the 12th conference on "Advances in Modal Logic," held in Bern, Switzerland, August 27-31, 2018, G. Bezhanishvili, G. D'Agostino, G. Metcalfe, and T. Studer, Eds. College Publications, 2018, pp. 473–492. [Online]. Available: http://www.aiml.net/volumes/volume12/Kuznets-Lellmann.pdf
- [25] T. Lyon, A. Tiu, R. Goré, and R. Clouston, "Syntactic interpolation for tense logics and bi-intuitionistic logic via nested sequents," in 28th EACSL Annual Conference on Computer Science Logic, CSL 2020, January 13-16, 2020, Barcelona, Spain, ser. LIPIcs, M. Fernández and A. Muscholl, Eds., vol. 152. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020, pp. 28:1–28:16.
- [26] K. Brünnler, "Deep sequent systems for modal logic," Arch. Math. Log., vol. 48, no. 6, pp. 551–577, 2009. [Online]. Available: https://doi.org/10.1007/s00153-009-0137-3
- [27] T. Lyon and K. van Berkel, "Automating agential reasoning: Proofcalculi and syntactic decidability for stit logics," in *PRIMA 2019: Principles and Practice of Multi-Agent Systems - 22nd International Conference, Turin, Italy, October 28-31, 2019, Proceedings*, ser. Lecture Notes in Computer Science, M. Baldoni, M. Dastani, B. Liao, Y. Sakurai, and R. Zalila Wenkstern, Eds., vol. 11873. Cham: Springer International Publishing, 2019, pp. 202–218.
- [28] A. Tiu, E. Ianovski, and R. Goré, "Grammar logics in nested sequent calculus: Proof theory and decision procedures," in Advances in Modal Logic 9, papers from the ninth conference on "Advances in Modal Logic," held in Copenhagen, Denmark, 22-25 August 2012, T. Bolander, T. Braüner, S. Ghilardi, and L. S. Moss, Eds. College Publications, 2012, pp. 516–537. [Online]. Available: http://www.aiml.net/volumes/volume9/Tiu-Ianovski-Gore.pdf
- [29] T. S. Lyon and L. Gómez Álvarez, "Automating Reasoning with Standpoint Logic via Nested Sequents," in *Proceedings of* the 19th International Conference on Principles of Knowledge Representation and Reasoning, 8 2022, pp. 257–266. [Online]. Available: https://doi.org/10.24963/kr.2022/26
- [30] A. Ciabattoni, P. Maffezioli, and L. Spendier, "Hypersequent and labelled calculi for intermediate logics," in *Automated Reasoning with Analytic Tableaux and Related Methods*, ser. Lecture Notes in Computer Science, D. Galmiche and D. Larchey-Wendling, Eds., vol. 8123. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013, pp. 81–96.
- [31] D. Galmiche, M. Marti, and D. Méry, "Relating Labelled and Label-Free Bunched Calculi in BI Logic," in 28th International Conference on Automated Reasoning with Anamytic Tableaux and Related Methods, TABLEAUX 2019, ser. 28Th Int.Conference on Automated Reasoning with Analytic Tableaux and Related Methods, Tableaux 2019, vol. 11714. Londres, United Kingdom: Springer, 2019, pp. 130–146.
- [32] R. Goré and R. Ramanayake, "Labelled tree sequents, tree hypersequents and nested (deep) sequents," in Advances in Modal Logic 9, papers from the ninth conference on "Advances in Modal Logic," held in Copenhagen, Denmark, 22-25 August 2012, T. Bolander, T. Braüner, S. Ghilardi, and L. S. Moss, Eds. College Publications, 2012, pp. 279– 299. [Online]. Available: http://www.aiml.net/volumes/volume9/Gore-Ramanayake.pdf
- [33] L. Pinto and T. Uustalu, "A proof-theoretic study of bi-intuitionistic propositional sequent calculus," *Journal of Logic and Computation*, vol. 28, no. 1, pp. 165–202, 2018.
- [34] M. Fitting, "Prefixed tableaus and nested sequents," Annals of Pure and Applied Logic, vol. 163, no. 3, pp. 291–313, 2012. [Online]. Available: https://www.sciencedirect.com/science/article/pii/S0168007211001266
- [35] M. A. Castilho, L. F. del Cerro, O. Gasquet, and A. Herzig, "Modal tableaux with propagation rules and structural rules," *Fundamenta Informaticae*, vol. 32, no. 3, 4, pp. 281–297, 1997.
- [36] L. Pinto and T. Uustalu, "Proof search and counter-model construction for bi-intuitionistic propositional logic with labelled sequents," in Auto-

*mated Reasoning with Analytic Tableaux and Related Methods*, M. Giese and A. Waaler, Eds. Berlin, Heidelberg: Springer Berlin Heidelberg, 2009, pp. 295–309.

- [37] K. van Berkel and T. Lyon, "Cut-free calculi and relational semantics for temporal stit logics," in *Logics in Artificial Intelligence*, F. Calimeri, N. Leone, and M. Manna, Eds. Cham: Springer International Publishing, 2019, pp. 803–819.
- [38] S. Negri and J. Von Plato, Proof analysis: a contribution to HIlbert's last problem. Cambridge University Press, 2011.
- [39] T. S. Lyon, "Nested sequents for first-order modal logics via reachability rules," arXiv, p. submitted, 2022. [Online]. Available: https://arxiv.org/abs/2210.00789
- [40] T. Lyon, "On the correspondence between nested calculi and semantic systems for intuitionistic logics," *Journal of Logic and Computation*, vol. 31, no. 1, pp. 213–265, 12 2020.
- [41] M. Fitting, "Nested sequents for intuitionistic logics," Notre Dame Journal of Formal Logic, vol. 55, no. 1, pp. 41–61, 2014.
- [42] F. Poggiolesi, "A purely syntactic and cut-free sequent calculus for the modal logic of provability," *The Review of Symbolic Logic*, vol. 2, no. 4, p. 593–611, 2009.
- [43] L. Straßburger, "Cut elimination in nested sequents for intuitionistic modal logics," in *Foundations of Software Science and Computation Structures*, ser. Lecture Notes in Computer Science, F. Pfenning, Ed., vol. 7794. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013, pp. 209–224.
- [44] T. S. Lyon, "Nested sequents for intuitionistic modal logics via structural refinement," in *Automated Reasoning with Analytic Tableaux and Related Methods*, A. Das and S. Negri, Eds. Cham: Springer International Publishing, 2021, pp. 409–427.
- [45] B. Boretti, "Proof analysis in temporal logic," Ph.D. dissertation, University of Milan, 2008.
- [46] R. Goré, L. Postniece, and A. Tiu, "On the correspondence between display postulates and deep inference in nested sequent calculi for tense logics," *Log. Methods Comput. Sci.*, vol. 7, no. 2, 2011. [Online]. Available: https://doi.org/10.2168/LMCS-7(2:8)2011
- [47] B. Lellmann and E. Pimentel, "Proof search in nested sequent calculi," in Logic for Programming, Artificial Intelligence, and Reasoning - 20th International Conference, LPAR-20 2015, Suva, Fiji, November 24-28, 2015, Proceedings, ser. Lecture Notes in Computer Science, M. Davis, A. Fehnker, A. McIver, and A. Voronkov, Eds., vol. 9450. Berlin, Heidelberg: Springer-Verlag, 2015, p. 558–574.
- [48] E. Pimentel, "A semantical view of proof systems," in Logic, Language, Information, and Computation - 25th International Workshop, WoLLIC 2018, Bogota, Colombia, July 24-27, 2018, Proceedings, ser. Lecture Notes in Computer Science, L. S. Moss, R. J. G. B. de Queiroz, and M. Martínez, Eds., vol. 10944. Springer, 2018, pp. 61–76.
- [49] E. L. Post, "Recursive unsolvability of a problem of Thue," *The Journal of Symbolic Logic*, vol. 12, no. 1, pp. 1–11, 1947.
- [50] M. C. Fitting and R. L. Mendelsohn, *First-Order Modal Logic*. Kluwer Academic Publishers, 1998.
- [51] L. F. del Cerro and M. Penttonen, "Grammar logics," *Logique et Analyse*, vol. 31, no. 121/122, pp. 123–134, 1988.
- [52] D. Galmiche and Y. Salhi, "Tree-sequent calculi and decision procedures for intuitionistic modal logics," *Journal of Logic and Computation*, vol. 28, no. 5, pp. 967–989, 06 2015.
- [53] K. van Berkel and T. Lyon, "The varieties of ought-implies-can and deontic stit logic," in *Deontic Logic and Normative Systems: 15th International Conference (DEON2020/2021, Munich)*, F. Liu, A. Marra, P. Portner, and F. V. D. Putte, Eds. College Publications, 2021.

#### Appendix

## A. Proofs for Section IV

Theorem 9. If l(R) is a local rule and H is a set of Horn rules, then  $l(R) \rightleftharpoons H$ .

*Proof:* Consider any  $\Gamma$ ,  $\Gamma'$ ,  $\Delta$ ,  $\Delta'$ , and  $\{\Delta_i\}_{i \in [n]}$ . Observe that the applicability of any local rule l(R) to a set of g-sequents does not depend on the edges thereof, that is, the derivation shown below left is a valid application of l(R) *iff* the derivation shown below right is:

$$\frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\Gamma \vdash \Delta} l(R) \quad \frac{\{\Gamma' \vdash \Delta_i\}_{i \in [n]}}{\Gamma' \vdash \Delta} l(R)$$

Analogously, the application of any rule  $\rho \in H$  does not depend on the labeling of vertices in g-sequents, that is, the derivation shown below left is a valid application of  $\rho$  *iff* the derivation shown below right is:

$$\frac{\Gamma \vdash \Delta}{\Gamma' \vdash \Delta} \rho \quad \frac{\Gamma \vdash \Delta'}{\Gamma' \vdash \Delta'} \rho$$

Thus, for any  $\rho \in H$ , we have a derivation of the form shown below left *iff* if we have a derivation of the form shown below right, where  $\rho \times n$  indicates that  $\rho$  is applied n times.

$$\frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\Gamma' \vdash \Delta} \rho l(R) = \frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\{\Gamma' \vdash \Delta_i\}_{i \in [n]}} \rho \times n$$
$$\frac{\{\Gamma' \vdash \Delta_i\}_{i \in [n]}}{\Gamma' \vdash \Delta} l(R)$$

Theorem 10. If e(R) is an expansion rule and H is a set of Horn rules, then H  $\nearrow e(R)$ .

*Proof:* We want to prove that for any rule  $\rho \in H$ , if we are given a derivation of the form shown below left, then we can swap the applications of the rules, yielding a derivation as shown below right. We let  $\Sigma \in \{w\mathcal{E}_a u, u\mathcal{E}_a w \mid a \in E\}$ .

$$\frac{\Gamma, \Sigma \vdash \Delta}{\Gamma' \vdash \Delta'} \rho e(R) \quad \frac{\Gamma, \Sigma \vdash \Delta}{\Gamma', \Sigma \vdash \Delta} \rho \\ \frac{\Gamma', \Sigma \vdash \Delta}{\Gamma' \vdash \Delta'} e(R)$$

To prove the claim, we must show that  $\mathcal{G} = \Gamma', \Sigma \vdash \Delta$ satisfies the conditions required to apply the e(R) rule: (1) In the derivation above left, observe that  $\rho$  will only affect an edge between vertices  $v, z \in \mathcal{U}(\Gamma)$  and that  $\mathcal{U}(\Gamma \vdash \Delta) \cap \mathcal{U}(\Sigma) = \{w\}$  holds due to the side condition on e(R). Therefore,  $\mathcal{U}(\Gamma' \vdash \Delta) \cap \mathcal{U}(\Sigma) = \{w\}$  as  $\mathcal{U}(\Gamma' \vdash \Delta) =$  $\mathcal{U}(\Gamma \vdash \Delta)$ . (2) The sequent constraint R will continue to hold for  $\mathcal{G}$  as well since  $\Delta$  is left unchanged. Therefore,  $\mathcal{G}$ satisfies the requirements for e(R) to be applied, showing that the derivation above right is permitted.

Theorem 11. If  $r(\mathcal{C}, R)$  is a reachability rule, and H is a set of Horn rules, then  $r(\mathcal{C}, R) \oplus \mathbf{G}(H) \rightleftharpoons H$ .

*Proof:* Let  $\rho \in H$  and  $\mathbf{G}(\rho) = \{a \longrightarrow s, \overline{a} \longrightarrow \overline{s}\}$ . We will only consider the case of  $h_f$ , as the case for  $h_b$  is analogous. We let  $\hat{\mathcal{C}} = \mathcal{C} \oplus \mathbf{G}(H)$  be the constraint of  $\sigma = r(\mathcal{C}, R) \oplus \mathbf{G}(H)$ . Our aim is to prove that we have a derivation of the form shown below top *iff* we have a derivation of the form shown below bottom.

$$\frac{\{\Gamma, w\mathcal{E}_{s}u, w\mathcal{E}_{a}u \vdash \Delta_{i}\}_{i \in [n]}}{\Gamma, w\mathcal{E}_{s}u, w\mathcal{E}_{a}u \vdash \Delta} \sigma \\
\frac{\{\Gamma, w\mathcal{E}_{s}u, w\mathcal{E}_{a}u \vdash \Delta_{i}\}_{i \in [n]}}{\{\Gamma, w\mathcal{E}_{s}u \vdash \Delta_{i}\}_{i \in [n]}} h_{f} \times m \\
\frac{\{\Gamma, w\mathcal{E}_{s}u \vdash \Delta_{i}\}_{i \in [n]}}{\Gamma, w\mathcal{E}_{s}u \vdash \Delta} \sigma$$

We note that the right-to-left direction is trivial, and thus, we focus on the left-to-right direction. In particular, we want to argue that if a derivation of the form shown below left is a

valid application of  $\sigma$ , then the derivation below right is a valid application of  $\sigma$ , where the premises are the g-sequents  $\mathcal{G}_i$  and  $\mathcal{G}'_i$ , respectively.

$$\underbrace{\{\overline{\Gamma, w\mathcal{E}_s u, w\mathcal{E}_a u \vdash \Delta_i}\}_{i \in [n]}}_{\Gamma, w\mathcal{E}_s u, w\mathcal{E}_a u \vdash \Delta} \sigma \xrightarrow{\frac{\mathcal{G}'_i}{\{\Gamma, w\mathcal{E}_s u \vdash \Delta_i\}_{i \in [n]}}}_{\Gamma, w\mathcal{E}_s u \vdash \Delta} \sigma$$

Let  $C = (C_1, \ldots, C_n)$  such that  $C_i = (\{v, v'\}, \{(v, v')\}, L_i)$ with  $L_i(v, v') = \mathbf{G}_i(a_i)$ . Furthermore, we let  $C_i \oplus \mathbf{G}(\mathbf{H}) = (\{v, v'\}, \{(v, v')\}, L'_i)$ , where  $L'_i(v, v') = \mathbf{G}'_i(a_i)$  and  $\mathbf{G}'_i = \mathbf{G}_i \cup \mathbf{G}(\mathbf{H})$  as dictated by Definition 7. Assume that  $L_i(v, v')$ uses  $u \stackrel{a}{\longrightarrow} w$  or  $u \stackrel{a}{\longrightarrow} w$ , as otherwise the case is trivial. Since  $\{a \longrightarrow s, \overline{a} \longrightarrow \overline{s}\} = \mathbf{G}(h_f) \subseteq \mathbf{G}(\mathbf{H})$ , anytime  $u \stackrel{a}{\longrightarrow} w$  or  $u \stackrel{a}{\longrightarrow} w$  is used in the propagation path witnessing satisfaction of  $L_i(v, v')$  in  $\mathcal{G}_i$  we can use  $u \stackrel{s}{\longrightarrow} w$  or  $u \stackrel{\overline{s}}{\longrightarrow} w$  in  $\mathcal{G}'_i$  instead. Thus,  $\mathcal{G}'_i \models v \stackrel{\mathbf{G}'_i(a_i)}{\longrightarrow} v'$  for  $i \in [n]$ , showing that  $h_f$  can be applied first, and  $\sigma$  second.

Theorem 20. If i(C, R) is an initial rule and H is a set of Horn rules, then  $(\triangleright^1 i(C, R) \triangleright^0 H) \preceq i(C, R) \oplus \mathbf{G}(H)$ .

*Proof:* We consider w.l.o.g. a forward Horn rule  $h_f \in H$  as the case for a backward Horn rule is similar. Let us assume that we have an instance of i(C, R) with C = (V, E, L), followed by an application of  $h_f$ , as shown below.

$$\frac{\Gamma, w\mathcal{E}_s u, w\mathcal{E}_a u \vdash \Delta}{\Gamma, w\mathcal{E}_s u \vdash \Delta} \frac{i(C, R)}{h_f}$$

We let  $\mathcal{G} = \Gamma, w\mathcal{E}_s u, w\mathcal{E}_a u \vdash \Delta = (\mathcal{V}, \mathcal{E}, \mathcal{L})$  and let  $\mathcal{G}' = \Gamma, w\mathcal{E}_s u \vdash \Delta = (\mathcal{V}, \mathcal{E}', \mathcal{L})$ . Furthermore, let  $i(\hat{C}, R) = i(C, R) \oplus \mathbf{G}(\mathbf{H})$  and  $\hat{C} = C \oplus \mathbf{G}(\mathbf{H}) = (V, E, L')$  as in Definition 7. Since i(C, R) is an initial rule, we know that  $\mathcal{G}$  satisfies C. We aim to show that  $\mathcal{G}'$  satisfies  $\hat{C}$ , showing that  $i(\hat{C}, R)$  can simulate  $\triangleright^1 i(C, R) \triangleright^0 \mathbf{H}$ .

First, observe that the only difference between  $\mathcal{E}$  and  $\mathcal{E}'$  is that  $\mathcal{E}'_a = \mathcal{E}_a \setminus \{(w, u)\}$  for  $a \in \mathsf{E}, \mathcal{E}_a \in \mathcal{E}$ , and  $\mathcal{E}'_a \in \mathcal{E}'$ . Let us now argue that  $\mathcal{G}'$  satisfies  $\hat{C}$ . We must show for any  $(v, z) \in \mathcal{E}'$ , if  $L'(v, z) = \mathbf{G}'(b)$ , then  $\exists r \in \mathbf{G}'(b)$  such that  $v \stackrel{r}{\rightharpoondown} z$ . Therefore, let  $(v, z) \in \mathcal{E}'$  and suppose that  $L'(v, z) = \mathbf{G}'(b)$  with  $\mathbf{G}' = \mathbf{G} \cup \mathbf{G}(\mathsf{H})$ . Due to the fact that  $\mathcal{G}$  satisfies C, we know that  $\exists t \in \mathbf{G}(b)$  such that  $v \stackrel{t}{\leadsto} z$ . We now use the path  $v \stackrel{t}{\leadsto} z$  to find a new path  $v \stackrel{r}{\leadsto} z$  in  $\mathcal{G}'$ . Since  $w\mathcal{E}_s u$  occurs in  $\mathcal{G}'$ , there exists a path  $w \stackrel{s}{\Longrightarrow} u$  and its converse  $u \stackrel{s}{\Longrightarrow} w$  in  $\mathcal{G}'$ . To find  $v \stackrel{r}{\leadsto} z$ , we take  $v \stackrel{t}{\longleftrightarrow} z$  and replace each occurrence of  $w \stackrel{a}{\Longrightarrow} u$  by  $w \stackrel{s}{\Longrightarrow} u$ , as well as replace each occurrence of  $u \stackrel{a}{\longrightarrow} w$  by  $u \stackrel{s}{\Longrightarrow} w$ . Hence,  $\mathcal{G}' \models v \stackrel{r}{\longleftrightarrow} z$ .

Last, we need to show that  $r \in (\mathbf{G} \cup \mathbf{G}(\mathbf{H}))(b)$ . To do this, we first recognize that  $\{a \longrightarrow s, \overline{a} \longrightarrow \overline{s}\} = \mathbf{G}(h_f) \subseteq \mathbf{G}(\mathbf{H})$ . Because  $t \in \mathbf{G}(b)$  and  $\mathbf{G}(b) \subseteq (\mathbf{G} \cup \mathbf{G}(\mathbf{H}))(b)$ , there exists a derivation  $b \longrightarrow^*_{\mathbf{G} \cup \mathbf{G}(\mathbf{H})} t$ . By applying the production rule  $a \longrightarrow s$  to each occurrence of a in t and the production rule  $\overline{a} \longrightarrow \overline{s}$  to each occurrence of  $\overline{a}$  in t, we obtain a derivation  $b \longrightarrow^*_{\mathbf{G} \cup \mathbf{G}(\mathbf{H})} r$ , showing the desired claim. Hence, we have verified that  $\mathcal{G}'$  is an instance of  $i(\hat{C}, R)$ .

Theorem 24. If H is a set of Horn rules with H' a fracturable subset of DG(H), then  $H \setminus H' \nearrow H'$ .

*Proof:* We consider the case of a backward Horn rule  $h_b \in H \setminus H'$  as the forward case is analogous, and argue by contraposition. Suppose that  $h_b$  cannot be permuted above a rule  $h' \in H'$  (we assume w.l.o.g. that h' is a forward Horn rule) in a derivation, which occurs *iff* a derivation of the form shown below exists, where  $w \ \mathcal{E}_{s\overline{b}s'} u, w\mathcal{E}_a u$  and  $v\mathcal{E}_t v', v'\mathcal{E}_b v$  are 'active' in h' and  $h_b$ , respectively.

$$\frac{w \mathcal{E}_{s\overline{b}s'} u}{\Gamma, v\mathcal{E}_t v', \underbrace{w\mathcal{E}_s v, v'\mathcal{E}_b v, v'\mathcal{E}_{s'} u}_{\Gamma, w\mathcal{E}_s v, v'\mathcal{E}_{s'} u, v\mathcal{E}_t v', v'\mathcal{E}_b v \vdash \Delta} h_b} h'$$

However, from the above, we can conclude that  $h' \sqsubseteq h_b$  in DG(H), showing that H' is not a fracturable subset. *Lemma* 26. If  $\mathcal{G}$  is a g-sequent and H is a finite set of Horn rules, then  $\overline{\mathrm{H}}(\mathcal{G})$  is computable in PTIME and is  $\overline{\mathrm{H}}$ -saturated.

*Proof:* It should be clear, from the definition of  $\overline{\mathrm{H}}(\mathcal{G})$ , that saturating  $\mathcal{G}$  under applications of rules from  $\overline{\mathrm{H}}$  builds a bottom-up derivation of the following form:

$$\frac{\overline{\mathrm{H}}(\mathcal{G})}{\frac{\vdots}{\mathcal{G}}} h_1$$

where  $h_1, \ldots, h_n \in H$ . Observe the following: (1) any bottomup application of a Horn rule builds the upper g-sequent from the lower by adding a single edge. (2) As only permissible rules can be applied, the height of this derivation is bound by the maximal number of edges in a graph, which is quadratic in the number of its vertices. (3) Given a Horn rule h verifying if a sequent  $\mathcal{G}$  is  $\overline{h}$ -saturated can be done in PTIME, as checking the existence of a path labeled with a given string between any two vertices of a graph can be done in PTIME by standard algorithmic techniques. (4)  $\overline{\mathrm{H}}(\mathcal{G})$  is obviously  $\overline{\mathrm{H}}$ -saturated by its definition.

*Lemma* 28. Let H be a set of Horn rules,  $\mathcal{G}$  be a g-sequent, and  $\mathbf{G} = \mathbf{G}(\mathrm{H})$ . For any  $s \in (\mathbf{E} \cup \overline{\mathbf{E}})^*$  and  $u, w \in \mathcal{U}(\mathcal{G})$  if  $\mathcal{G} \models w \underset{\sim}{\sim} u$ , then for every string t such that  $t \longrightarrow_{\mathbf{G}}^* s$ , we have  $\overline{\mathrm{H}}(\mathcal{G}) \models w \underset{\sim}{\sim} u$ .

**Proof:** Assume towards contradiction that lemma does not hold. Let H,  $\mathcal{G}$ , w, u, s, and t form a counterexample. Note that w.l.o.g we can assume  $t \longrightarrow_{\mathbf{G}} s$ , meaning s is derivable from t in one step. As  $t \longrightarrow_{\mathbf{G}} s$  we know that (1) there exists a rule  $h \in \mathbf{H}$  (we may assume w.l.o.g. that  $h = h_f$ ) such that  $\mathbf{G}(h) = \{a \longrightarrow r, \overline{a} \longrightarrow \overline{r}\}, (2) t$ is of the form  $t_1 a t_2$  (or  $t_1 \overline{a} t_2$ ), and (3) s is of the form  $t_1 r t_2$  (or  $t_1 \overline{r} t_2$ ). We will consider the case of  $t = t_1 a t_2$ , as the latter is analogous. Let  $w', u' \in \mathcal{U}(\mathcal{G})$  be such that:  $\overline{\mathrm{H}}(\mathcal{G}) \models w \stackrel{t}{\longrightarrow} w', w' \stackrel{r}{\longrightarrow} u', u' \stackrel{t}{\longrightarrow} u$ , but not  $\overline{\mathrm{H}}(\mathcal{G}) \models w \stackrel{t}{\longrightarrow} u$ . Let  $\overline{\mathrm{H}}(\mathcal{G}) = \Gamma \vdash \Delta$  and consider the following:

$$\frac{\Gamma, \, w' \mathcal{E}_a u' \, \vdash \, \Delta}{\Gamma \, \vdash \, \Delta} \, h$$

Note, that when viewed bottom-up, this is a permissible application of  $\overline{h}$ , which leads to a contradiction as  $\overline{\mathrm{H}}(\mathcal{G})$  is  $\overline{\mathrm{H}}$ -saturated.

*Lemma* 29. The path weakening rule pw (shown below) is eliminable from any proof:

$$\frac{\Gamma \vdash \Delta}{\Gamma, \Sigma \vdash \Delta} pw$$

where  $\Sigma \in \{ w \mathcal{E}_a u \mid a \in \mathsf{E} \text{ and } w, u \in \mathcal{U}(\Gamma \vdash \Delta) \}.$ 

*Proof:* We will show that: (1) i(C, R) followed by pw can be simulated by i(C, R) alone, (2)  $pw \nearrow l(R)$ , (3)  $pw \nearrow e(R)$ , (4)  $pw \nearrow r(C, R)$ , and (5) that for any derivation consisting of a Horn rule h followed by a weakening rule  $\rho$  one of the following holds: (5a)  $\rho$  can be permuted above h, or (5b) both applications can be removed. Using these facts, the result follows by induction on the quantity of the given proof.

(1) Observe that if we have a derivation of the form shown below left, then we have a derivation of the form shown below right, as the satisfaction of C and R cannot be falsified by the addition of edges.

$$\frac{\overline{\Gamma \vdash \Delta}}{\Gamma, \Sigma \vdash \Delta} \frac{i(C, R)}{pw} \qquad \overline{\Gamma, \Sigma \vdash \Delta} i(C, R)$$

(2) Assume we have a derivation of the form shown below left. As applicability of a local rule does not depend on the set of the edges of a g-sequent, any derivation of the form shown below right is valid as well.

$$\frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\frac{\Gamma \vdash \Delta}{\Gamma, \Sigma \vdash \Delta} pw} l(R) \qquad \frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\{\Gamma, \Sigma \vdash \Delta_i\}_{i \in [n]}} pw \times n$$

(3) Assume we have a derivation of the form shown below left, and assume w.l.o.g. that  $\mathcal{U}(\Gamma \vdash \Delta) \cap \{w\mathcal{E}_b u\} = \{w\}$ . Note as  $u' \neq u$  and  $w' \neq u$  in the derivation below left, we may permute pw above e(R), obtaining the derivation shown below right.

$$\frac{\Gamma, w\mathcal{E}_b u \vdash \Delta}{\Gamma \vdash \Delta'} e(R) = \frac{\Gamma, w\mathcal{E}_b u \vdash \Delta}{\Gamma, w\mathcal{E}_b u, w'\mathcal{E}_a u' \vdash \Delta} pw$$

$$\frac{\Gamma, w\mathcal{E}_b u, w'\mathcal{E}_a u' \vdash \Delta}{\Gamma', w'\mathcal{E}_a u' \vdash \Delta'} e(R)$$

(4) Assume we have a derivation of the form shown below left. As satisfaction of C and R cannot be falsified by the addition of edges, we may permute pw above r(C, R), as shown below right.

$$\frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\frac{\Gamma \vdash \Delta}{\Gamma, \Sigma \vdash \Delta} pw} r(\mathcal{C}, R) \qquad \frac{\{\Gamma \vdash \Delta_i\}_{i \in [n]}}{\{\Gamma, \Sigma \vdash \Delta_i\}_{i \in [n]}} pw \times n}{\frac{\{\Gamma, \Sigma \vdash \Delta_i\}_{i \in [n]}}{\Gamma, \Sigma \vdash \Delta} r(\mathcal{C}, R)}$$

(5) We will consider only  $h_f$  case as the  $h_b$  case is analogous. Assume we have a derivation of the form shown below left. If  $\Sigma = \Sigma'$  then both applications can be removed from any derivation; otherwise, we may permute pw above  $h_f$  as shown below right.

$$\frac{\frac{\Gamma, \Sigma \vdash \Delta}{\Gamma, \Sigma' \vdash \Delta} h_f}{\Gamma, \Sigma' \vdash \Delta} pw \qquad \frac{\frac{\Gamma, \Sigma \vdash \Delta}{\Gamma, \Sigma, \Sigma' \vdash \Delta} pw}{\Gamma, \Sigma' \vdash \Delta} h_f$$

Theorem 32. Let i(C, R) be an initial rule with the set  $H := H(\mathbf{G}(i(C, R)))$ . If H' is a fracturable subset of H, then  $i(C, R) \preceq \rhd^1 i(C, R) \ominus \mathbf{G}(H') \rhd^0 H'$ .

*Proof:* Suppose we have an application of i(C, R), yielding the g-sequent  $\mathcal{G}$ . By Lemma 31, we know that the g-sequent  $\mathcal{G}' = \overline{\mathrm{H}'}(\mathcal{G})$  satisfies  $C \ominus \mathbf{G}(\mathrm{H}')$ , and thus, serves as an instance of  $i(C, R) \ominus \mathbf{G}(\mathrm{H}')$ . As  $\overline{\mathrm{H}'}(\mathcal{G})$  is  $\overline{\mathrm{H}'}$ -saturated, we may apply the rules from  $\mathrm{H}'$  to  $\mathcal{G}'$  to derive  $\mathcal{G}$ .

Theorem 33. Let  $r(\mathcal{C}, R)$  be a reachability rule and let  $H := H(\mathbf{G}(r(\mathcal{C}, R)))$ . If H' is a fracturable subset of H, then  $r(\mathcal{C}, R) \preceq \triangleright^0 pw \triangleright^1 r(\mathcal{C}, R) \ominus \mathbf{G}(H') \triangleright^0 H'$ .

*Proof:* Suppose that we have an application of  $r(\mathcal{C}, R)$  as shown below left, where  $\mathcal{C} = (C_1, \ldots, C_n)$ . By Lemma 30, we can derive the g-sequent  $\mathcal{G}'_i = \overline{\mathrm{H}'}(\mathcal{G}_i)$  from  $\mathcal{G}_i$  for each  $i \in [n]$  via some number k of applications of pw. By Lemma 31, we know that each g-sequent  $\mathcal{G}'_i$  satisfies the constraint  $C_i \ominus \mathbf{G}(\mathrm{H}')$ , and thus, we may apply  $r(\mathcal{C}, R) \ominus \mathbf{G}(\mathrm{H}')$ to derive  $\overline{\mathrm{H}'}(\mathcal{G})$ . As  $\overline{\mathrm{H}'}(\mathcal{G})$  is  $\overline{\mathrm{H}'}$ -saturated, we can derive  $\mathcal{G}$ by subsequent applications of H', as shown below right.

$$\frac{\{\mathcal{G}_i\}_{i\in[n]}}{\mathcal{G}}r(\mathcal{C},R) \qquad \frac{\frac{\{\mathcal{G}_i\}_{i\in[n]}}{|\{\overline{\mathrm{H}'}(\mathcal{G}_i)\}_{i\in[n]}}pw \times k}{\frac{|\overline{\mathrm{H}'}(\mathcal{G})|_{i\in[n]}}{\mathcal{G}}H'}r(\mathcal{C},R)\ominus \mathbf{G}(\mathrm{H}')$$

## B. Proofs for Section V

Theorem 36. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \ll)$  its upward space. Then, the  $\ll$  relation is a connected partial-order. Moreover, for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , if we take  $\mathfrak{B} \wedge \mathfrak{C} = \inf{\mathfrak{B}, \mathfrak{C}}$  and  $\mathfrak{B} \vee \mathfrak{C} = \sup{\mathfrak{B}, \mathfrak{C}}$  under  $\ll$ , then  $(\mathbf{S}, \wedge, \vee)$  is a complete lattice with  $\mathfrak{A} = \bot$ .

*Proof:* Let  $\mathfrak{A} = (\mathfrak{G}(\mathsf{E}), \mathfrak{R})$  be an abstract calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its upward space. We first prove that  $\leqslant$  is a partial order. By Lemma 23, we know that  $\emptyset$  is an anti-fracturable subset in any dependency graph. Observe that for any  $\mathfrak{B} \in \mathbf{S}, \mathfrak{B} = (\mathfrak{B} \oplus \mathbf{G}(\emptyset)) \setminus \emptyset$ , showing that  $\mathfrak{B} \notin \mathfrak{B}$ , i.e.  $\gtrless$  is reflexive.

Second, assume that  $\mathfrak{B} \leq \mathfrak{C}$  and  $\mathfrak{C} \leq \mathfrak{D}$  for  $\mathfrak{B}, \mathfrak{C}, \mathfrak{D} \in$ S. Then, there exist anti-fracturable subsets  $H'_1$  of  $H(\mathfrak{B})$  and  $H'_2$  of  $H(\mathfrak{C})$  such that  $\mathfrak{C} = (\mathfrak{B} \oplus \mathbf{G}(H'_1)) \setminus H'_1$  and  $\mathfrak{D} = (\mathfrak{C} \oplus \mathbf{G}(H'_2)) \setminus H'_2$ . Observe that  $\mathfrak{D} = (\mathfrak{B} \oplus \mathbf{G}(H'_1 \cup H'_2)) \setminus (H'_1 \cup H'_2)$ . Therefore, to complete the proof of the case, we need to show that  $H'_1 \cup H'_2$  is anti-fracturable. Observe that the sets  $H(\mathfrak{B}) \setminus H'_1 = H(\mathfrak{C})$  and  $H(\mathfrak{C}) \setminus H'_2$  are fracturable, meaning  $(H(\mathfrak{B}) \setminus H'_1) \setminus H'_2 = H(\mathfrak{B}) \setminus (H'_1 \cup H'_2)$  is fracturable; consequently,  $H'_1 \cup H'_2$  is anti-fracturable, showing that  $\leq$  is transitive.

Third, let us assume that  $\mathfrak{B} \ll \mathfrak{C}$  and  $\mathfrak{C} \ll \mathfrak{B}$  for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ . Then, there exist anti-fracturable subsets  $H'_1$  of  $H(\mathfrak{B})$  and  $H'_2$ of  $H(\mathfrak{C})$  such that  $\mathfrak{C} = (\mathfrak{B} \oplus \mathbf{G}(H'_1)) \setminus H'_1$  and  $\mathfrak{B} = (\mathfrak{C} \oplus \mathbf{G}(H'_2)) \setminus H'_2$ . Plugging either equation into the other shows that  $H'_1 = \emptyset = H'_2$ , implying that  $\mathfrak{C} = \mathfrak{B}$ .

Last, one can show that for any  $\mathfrak{B} \in \mathbf{S}$ ,  $\mathfrak{A} \notin \mathfrak{B}$ , that is,  $\mathbb{U}(\mathfrak{A})$  is connected. Moreover, as every partially ordered set

can be viewed as a lattice, we can view  $\mathbb{U}(\mathfrak{A})$  as a lattice, and by the previous sentence,  $\mathfrak{A}$  is the bottom element.

Theorem 37. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its upward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{A}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{B}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') \leq s(\mathcal{P})$ .

**Proof:** Let  $\mathfrak{A}$  be an abstract calculus,  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \boldsymbol{\leqslant})$  its upward space, and  $\mathfrak{B} \in \mathbf{S}$ . By Theorem 36, we know that  $\mathfrak{A} \boldsymbol{\leqslant} \mathfrak{B}$ , and hence, there exists a set H' of Horn rules such that H'  $\subseteq$  H = H( $\mathfrak{A}$ ) is an anti-fracturable subset in DG(H) which witnesses that the relation  $\mathfrak{A} \boldsymbol{\leqslant} \mathfrak{B}$  holds. Let us take a proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{A}$  and consider the topmost application of a Horn rule in H'. We assume w.l.o.g. that the Horn rule is  $h_f$  and argue by induction on the quantity of the proof that  $h_f$  can be permuted upward and eliminated entirely.

*Base case.* Suppose we have an application of i(C, R) followed by an application of  $h_f$ . By Lemma 18, we know that  $\triangleright^1 i(C, R) \triangleright^1 h_f \preceq \triangleright^1 i(C, R) \triangleright^0 h_f$ , and by Theorem 20, we know that  $\triangleright^1 i(C, R) \triangleright^0 h_f \preceq i(C, R) \oplus \mathbf{G}(h_f)$ . By Lemma 19,  $i(C, R) \oplus \mathbf{G}(h_f) \preceq i(C, R) \oplus \mathbf{G}(H')$  since  $h_f \in \mathbf{H}'$ . As simulations are transitive (Lemma 16), we know that  $\triangleright^1 i(C, R) \triangleright^1 h_f \preceq i(C, R) \oplus \mathbf{G}(H')$ . We can therefore replace the application of  $i(C, R) \oplus \mathbf{G}(H')$ .

Inductive step. Suppose we have an application of a local, expansion, reachability, or Horn rule  $\rho$  in  $\mathfrak{B}$  followed by an application of  $h_f$ . By Theorem 9 (local case), 10 (expansion case), 11 (reachability case), and 24 (Horn rule case) we know that  $h_f$  can be permuted above  $\rho$ , and thus by the induction hypothesis,  $h_f$  is elimimable from the proof altogether. Note that by definition H' is an anti-fracturable subset of H( $\mathfrak{A}$ ), and thus, Theorem 24 is applicable.

We repeat the above algorithm by successively considering each topmost occurrence of a horn rule in H' until all such rules are eliminated, giving a PTIME procedure for computing the output proof  $\mathcal{P}'$  of  $\mathfrak{B}$ . Moreover,  $\max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}'\} \leq \max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}\}$  since the elimination of Horn rules removes edges from g-sequents in  $\mathcal{P}$ , giving potentially smaller g-sequents in  $\mathcal{P}'$ . Additionally,  $|\{\mathcal{G} \in \mathfrak{G}(E) \mid \mathcal{G} \in \mathcal{P}'\}| \leq |\{\mathcal{G} \in \mathfrak{G}(E) \mid \mathcal{G} \in \mathcal{P}\}|$  since Horn rules are 'absorbed' into the initial rules. Thus, we have that  $s(\mathcal{P}') \leq s(\mathcal{P})$ .  $\blacksquare$ *Theorem* 39. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an explicit calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its upward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{B}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{A}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

*Proof:* Let  $\mathfrak{A}$  be an explicit calculus,  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \boldsymbol{\leqslant})$  its upward space, and  $\mathfrak{B} \in \mathbf{S}$ . By Theorem 36, we know that  $\mathfrak{A} \boldsymbol{\leqslant} \mathfrak{B}$ , and hence, there exists a set H' of Horn rules such that  $\mathrm{H}' \subseteq \mathrm{H} = \mathrm{H}(\mathfrak{A})$  is an anti-fracturable subset in DG(H) and which witnesses that the relation  $\mathfrak{A} \boldsymbol{\leqslant} \mathfrak{B}$  holds.

Let us first consider any initial rule  $i(\hat{C}, R)$  or reachability rule  $r(\hat{C}, R)$  in  $\mathfrak{B}$ . We know that each rule was obtained from an initial rule i(C, R) and reachability rule  $r(\mathcal{C}, R)$  in  $\mathfrak{A}$  such that  $\hat{C} = C \oplus \mathbf{G}(\mathbf{H}')$  and  $\hat{C} = \mathcal{C} \oplus \mathbf{G}(\mathbf{H}')$ . Since  $\mathfrak{A}$  is explicit, we know that  $\mathbf{G}(\rho) \cap \mathbf{G}(\mathbf{H}') = \emptyset$  for  $\rho \in \{i(C, R), r(\mathcal{C}, R)\}$ , and thus, by Lemma 14  $(\rho \oplus \mathbf{G}(\mathbf{H}')) \oplus \mathbf{G}(\mathbf{H}') = \rho$ . This fact is required to complete the remainder of the proof.

Let us now suppose that we have a proof  $\mathcal{P}$  of  $\mathcal{G}$  in  $\mathfrak{B}$ . By Theorem 32 and 33, we can replace every application of an initial rule  $i(\hat{C}, R)$  and reachability rule  $r(\hat{C}, R)$  from  $\mathfrak{B}$  by a derivation from  $\triangleright^1 i(\hat{C}, R) \ominus \mathbf{G}(\mathbf{H}') \triangleright^0 \mathbf{H}'$  and  $\triangleright^0 pw \triangleright^1 r(\hat{C}, R) \ominus$  $\mathbf{G}(\mathbf{H}') \triangleright^0 \mathbf{H}'$  as  $\mathbf{H}'$  is a fracturable subset of  $\mathbf{H}'$  by Lemma 23. As mentioned above,  $i(\hat{C}, R) \ominus \mathbf{G}(\mathbf{H}')$  and  $r(\hat{C}, R) \ominus \mathbf{G}(\mathbf{H}')$  are initial and reachability rules in  $\mathfrak{A}$ . Furthermore, by Lemma 29, we know that pw is eliminable from any proof, and thus, by eliminating all occurrences of pw, we obtain a proof  $\mathcal{P}'$  that is a proof in  $\mathfrak{A}$ .

The above yields a PTIME algorithm for transforming  $\mathcal{P}$  into  $\mathcal{P}'$ , and observe that in the worst-case

$$\max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}'\} = \mathcal{O}(\max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}\}^2)$$

as applications of pw could transform g-sequents from  $\mathcal{P}$  into at most complete graphs with quadratically more edges. Moreover, for each application of a rule in  $\mathcal{P}$ , we could have (in the worst-case) at most quadratically many applications of rules in  $\mathcal{P}'$  as Horn rules can only be applied at most quadratically many times (by what was said above) until the edges from a g-sequent are completely removed. Therefore,  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

*Theorem* 41. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an abstract calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its downward space. Then, the  $\leqslant$  relation is a connected partial order. Moreover, for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , if we take  $\mathfrak{B} \land \mathfrak{C} = \inf{\{\mathfrak{B}, \mathfrak{C}\}}$  and  $\mathfrak{B} \lor \mathfrak{C} = \sup{\{\mathfrak{B}, \mathfrak{C}\}}$  under  $\leqslant$ , then  $(\mathbf{S}, \land, \lor)$  is a complete lattice with  $\mathfrak{A} = \top$ .

*Proof:* Let  $\mathfrak{A} = (\mathfrak{G}(\mathbb{E}), \mathfrak{R})$  be an abstract calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$  its downward space. We first prove that  $\leqslant$  is a partial order. By Lemma 23, we know that  $\emptyset$  is a fracturable subset in DG(G( $\mathfrak{B}\setminus H$ )) with  $\mathfrak{B}$  an abstract calculus and H := H(\mathfrak{B}), implying that for any  $\mathfrak{B} \in \mathbf{S}, \mathfrak{B} = (\mathfrak{B} \ominus \emptyset) \cup H(\emptyset)$ , showing that  $\mathfrak{B} \leqslant \mathfrak{B}$ , i.e.  $\leqslant$  is reflexive.

Second, assume that  $\mathfrak{B} \leq \mathfrak{C}$  and  $\mathfrak{C} \leq \mathfrak{D}$  for  $\mathfrak{B}, \mathfrak{C}, \mathfrak{D} \in \mathbf{S}$ . Then, there exist fracturable subsets  $P'_1$  of  $\mathbf{G}(\mathfrak{C} \setminus H_1)$  with  $H_1 := \mathrm{H}(\mathfrak{C})$  and  $P'_2$  of  $\mathbf{G}(\mathfrak{D} \setminus H_2)$  with  $H_2 := \mathrm{H}(\mathfrak{D})$  such that  $\mathfrak{B} = (\mathfrak{C} \ominus \mathbf{G}(P'_1)) \cup \mathrm{H}(P'_1)$  and  $\mathfrak{C} = (\mathfrak{D} \ominus \mathbf{G}(P'_2)) \cup \mathrm{H}(P'_2)$ . Observe that  $\mathfrak{B} = (\mathfrak{D} \ominus \mathbf{G}(P'_1 \cup P'_2)) \cup \mathrm{H}(P'_1 \cup P'_2)$ . Therefore, to complete the proof of the case, we need to show that  $P'_1 \cup P'_2$  is fracturable. We know that  $P'_1 \subseteq \mathbf{G}(\mathfrak{C} \setminus H_1) \subseteq \mathbf{G}(\mathfrak{O} \setminus H_2)$  as  $\mathbf{G}(\mathfrak{C} \setminus H_1) = \mathbf{G}(\mathfrak{D} \setminus H_2) \setminus \mathbf{G}(P'_2)$ . Since  $P'_2$  is fracturable in  $\mathbf{G}(\mathfrak{D} \setminus H_2)$  and  $P'_1 \subseteq \mathbf{G}(\mathfrak{O} \setminus H_2)$ , we know that  $P'_2$  does not depend on any pair in  $P'_1$  in the dependency graph of  $\mathbf{G}(\mathfrak{O} \setminus H_2)$ . Furthermore, a pair in  $P'_1$  may depend on a pair in  $P'_2$ , but cannot depend on any pair in  $\mathbf{G}(\mathfrak{O} \setminus H_2) \setminus \mathbf{G}(P'_2)$ . Thus,  $P'_1 \cup P'_2$  is a fracturable subset of  $\mathbf{G}(\mathfrak{O} \setminus H_2)$ , showing that  $\mathfrak{B} \leq \mathfrak{D}$ , i.e.  $\leq$  is transitive.

Third, let us assume that  $\mathfrak{B} \leq \mathfrak{C}$  and  $\mathfrak{C} \leq \mathfrak{B}$  for  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ . Then, there exist fracturable subsets  $P'_1$  of  $\mathbf{G}(\mathfrak{B} \setminus \mathrm{H}_1)$  with  $\mathrm{H}_1 := \mathrm{H}(\mathfrak{B})$  and  $P'_2$  of  $\mathbf{G}(\mathfrak{C} \setminus \mathrm{H}_2)$  such that  $\mathfrak{C} = (\mathfrak{B} \ominus \mathbf{G}(P'_1)) \cup \mathrm{H}(P_1)$  and  $\mathfrak{B} = (\mathfrak{C} \ominus \mathbf{G}(P'_2)) \cup \mathrm{H}(P_2)$ . Plugging either equation into the other shows that  $P'_1 = \emptyset = P'_2$ , implying that  $\mathfrak{C} = \mathfrak{B}$ .

Finally, one can show that for any  $\mathfrak{B} \in \mathbf{S}$ ,  $\mathfrak{B} \leq \mathfrak{A}$ , that is,  $\mathbb{U}(\mathfrak{A})$  is connected, and since every partially ordered set can be viewed as a lattice, we can view  $\mathbb{D}(\mathfrak{A})$  as a lattice with  $\mathfrak{A}$  the top element.

Theorem 42. Let  $\mathfrak{A} = (\mathfrak{G}(\mathsf{E}), \mathfrak{R})$  be an abstract calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its downward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{A}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{B}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

*Proof:* Let  $\mathfrak{A}$  be an abstract calculus,  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its downward space, and  $\mathfrak{B} \in \mathbf{S}$ . By Theorem 41, we know that  $\mathfrak{B} \leq \mathfrak{A}$  and hence there exists a fracturable subset P' in the dependency graph  $\mathsf{DG}(\mathbf{G}(\mathfrak{A} \setminus \mathrm{H})) = (P, \sqsubseteq)$  with  $\mathrm{H} := \mathrm{H}(\mathfrak{A})$ , witnessing that the relation  $\mathfrak{B} \leq \mathfrak{A}$  holds.

Let us suppose that we have a proof  $\mathcal{P}$  of  $\mathcal{G}$  in  $\mathfrak{A}$ . By Theorem 32 and 33, we can replace every application of an initial rule i(C, R) and reachability rule  $r(\mathcal{C}, R)$  from  $\mathfrak{A}$  by a derivation from  $\triangleright^1 i(C, R) \ominus \mathbf{G}(P') \triangleright^0 \mathrm{H}(\mathbf{G}(P'))$  and  $\triangleright^0 pw \triangleright^1$  $r(\mathcal{C}, R) \ominus \mathbf{G}(P') \triangleright^0 \mathrm{H}(\mathbf{G}(P'))$ . By Lemma 29, we know that pw is eliminable from any proof, and thus, by eliminating all occurrences of pw, we obtain a proof  $\mathcal{P}'$  that is a proof in  $\mathfrak{A}$ .

The above yields a PTIME algorithm for transforming  $\mathcal{P}$  into  $\mathcal{P}'$ , and a complexity analysis similar to the one in Theorem 42 shows that  $s(\mathcal{P}') = \mathcal{O}(s(\mathcal{P})^2)$ .

Theorem 44. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an implicit calculus with  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its downward space. For any abstract calculus  $\mathfrak{B} \in \mathbf{S}$  and proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{B}$ , there exists a proof  $\mathcal{P}'$  of  $\mathcal{G}$  in  $\mathfrak{A}$  such that  $\mathcal{P}'$  is computable from  $\mathcal{P}$  in PTIME with  $s(\mathcal{P}') \leq s(\mathcal{P})$ .

*Proof:* Let  $\mathfrak{A}$  be an implicit calculus,  $\mathbb{D}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its downward space, and  $\mathfrak{B} \in \mathbf{S}$ . By Theorem 41, we know that  $\mathfrak{B} \leq \mathfrak{A}$ , and hence, there exists a fracturable subset P' of  $\mathsf{DG}(\mathbf{G}(\mathfrak{A}))$  (note that  $\mathrm{H}(\mathfrak{A}) = \emptyset$  as  $\mathfrak{A}$  is implicit), which witnesses that the relation  $\mathfrak{B} \leq \mathfrak{A}$  holds.

Let us first consider any initial rule  $i(\check{C}, R)$  or reachability rule  $r(\check{C}, R)$  in  $\mathfrak{B}$ . We know that each rule was obtained from an initial rule i(C, R) and reachability rule  $r(\mathcal{C}, R)$  in  $\mathfrak{A}$  such that  $\check{C} = C \ominus \mathbf{G}(P')$  and  $\check{C} = \mathcal{C} \ominus \mathbf{G}(P')$ . Since  $\mathfrak{A}$  is implicit, we know that  $\mathbf{G}(P') \subseteq \mathbf{G}(\rho)$  for  $\rho \in \{i(C, R), r(\mathcal{C}, R)\}$ , and thus, by Lemma 14 and the fact that  $\mathbf{G}(P') = \mathbf{G}(\mathrm{H}(P'))$ , we have  $(\rho \ominus \mathbf{G}(P')) \oplus \mathbf{G}(P') = \rho$ . This fact is required to complete the remainder of the proof.

Let us take a proof  $\mathcal{P}$  of a g-sequent  $\mathcal{G}$  in  $\mathfrak{B}$  and consider the topmost application of a Horn rule in  $\mathrm{H}' := \mathrm{H}(\mathbf{G}(P'))$ . We assume w.l.o.g. that the Horn rule is  $h_f$  and argue by induction on the quantity of the proof that  $h_f$  can be permuted upward and eliminated entirely.

*Base case.* Suppose we have an application of  $i(\check{C}, R)$ followed by an application of  $h_f$ . By Lemma 18, we know that  $\triangleright^1 i(\check{C}, R) \triangleright^1 h_f \preceq \triangleright^1 i(\check{C}, R) \triangleright^0 h_f$ , and by Theorem 20, we know that  $\triangleright^1 i(\check{C}, R) \triangleright^0 h_f \preceq i(\check{C}, R) \oplus \mathbf{G}(h_f)$ . By Lemma 19,  $i(\check{C}, R) \oplus \mathbf{G}(h_f) \preceq i(\check{C}, R) \oplus \mathbf{G}(H')$  since  $h_f \in H'$ . As simulations are transitive (Lemma 16), we know that  $\triangleright^1 i(C, R) \triangleright^1 h_f \leq i(C, R) \oplus \mathbf{G}(\mathbf{H}')$ . We can therefore replace the application of  $i(\check{C}, R)$  followed by  $h_f$  by a single application of  $i(\check{C}, R) \oplus \mathbf{G}(\mathbf{H}')$ , which is an initial rule in  $\mathfrak{A}$ since  $(\rho \ominus \mathbf{G}(P')) \oplus \mathbf{G}(P') = \rho$  for every initial rule  $\rho$  in  $\mathfrak{A}$ .

Inductive step. Suppose we have an application of a local, expansion, or reachability  $\rho$  in  $\mathfrak{B}$  followed by an application of  $h_f$ . In the case that  $\rho$  is a reachability rule, replace  $r(\check{C}, R)$ by  $r(\check{C}, R) \oplus \mathbf{G}(\mathbf{H}') = r(\mathcal{C}, R)$ , which is a reachability rule in  $\mathfrak{A}$  since  $(\rho \ominus \mathbf{G}(P')) \oplus \mathbf{G}(P') = \rho$  for every reachability rule  $\rho$ in  $\mathfrak{A}$ . By Theorem 9 (local case), 10 (expansion case), and 11 (reachability case), we know that  $h_f$  can be permuted above  $\rho$ , and thus by the induction hypothesis,  $h_f$  is eliminable from the proof altogether. Note that we need not consider permutations of  $h_f$  above Horn rules as  $h_f$  is a topmost application of a Horn rule.

We repeat the above algorithm by successively considering each topmost occurrence of a horn rule in H' until all such rules are eliminated, giving a PTIME procedure for computing the output proof  $\mathcal{P}'$  of  $\mathfrak{A}$ . Moreover,  $\max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}'\} \leq \max\{s(\mathcal{G}) \mid \mathcal{G} \in \mathcal{P}\}$  since the elimination of Horn rules removes edges from g-sequents in  $\mathcal{P}$ , giving potentially smaller g-sequents in  $\mathcal{P}'$ . Additionally,  $|\{\mathcal{G} \in \mathfrak{G}(E) \mid \mathcal{G} \in \mathcal{P}'\}| \leq |\{\mathcal{G} \in \mathfrak{G}(E) \mid \mathcal{G} \in \mathcal{P}\}|$  since Horn rules are 'absorbed' into the initial rules. Thus, we have that  $s(\mathcal{P}') \leq s(\mathcal{P})$ .  $\blacksquare$ *Theorem* 45. If  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  is an implicit calculus, then every complete proof is a polytree proof.

**Proof:** Consider a complete proof of a g-sequent  $\vdash w : S$  in an implicit calculus  $\mathfrak{A}$ . Observe that  $\vdash w : S$  is a polytree g-sequent, and any bottom-up application of l(R), e(R), or  $r(\mathcal{C}, R)$  will yield polytree g-sequents as the premises; hence, the entire proof will consist of polytree g-sequents.

Theorem 47. Let  $\mathfrak{A} = (\mathfrak{G}(E), \mathfrak{R})$  be an explicit calculus with  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leq)$  its upward space. Then,

- 1)  $\mathbb{U}(\mathfrak{A}) = \text{IMPLICATE}(\mathfrak{A});$
- 2)  $\mathbb{U}(\mathfrak{A})$  is computable from  $\mathfrak{A}$  in EXPTIME;
- 3) If  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ , then  $\mathfrak{B} \twoheadrightarrow_p \mathfrak{C}$ ;
- 4)  $\top$  is computable from  $\mathfrak{A}$  in PTIME;
- 5)  $\top$  is the only implicit calculus in  $\mathbb{U}(\mathfrak{A})$ .

*Proof:* As (1) is straightforward to confirm, we argue claims (2)–(5) in turn:

(2) During the execution of IMPLICATE the algorithm considers all anti-fracturable subsets of any calculus  $\mathfrak{B} \in \mathbf{S}$ , which may be exponential in the size of the input. Therefore, IMPLICATE has a worst-case complexity of EXPTIME.

(3) Suppose  $\mathfrak{B}, \mathfrak{C} \in \mathbf{S}$ . Then, by Theorem 37 and 39, we know that  $\mathfrak{B} \dashv \vdash_p \mathfrak{A}$  and  $\mathfrak{A} \dashv \vdash_p \mathfrak{C}$ , which implies that  $\mathfrak{B} \dashv \vdash_p \mathfrak{C}$ .

(4) By taking  $H := H(\mathfrak{A})$  and defining  $\mathfrak{B} := f(\mathfrak{A}, H)$ , we obtain the calculus  $\top$ . This procedure can be performed in PTIME in the size of  $\mathfrak{A}$ .

(5) Since  $\top$  is free of Horn rules and  $\mathfrak{A}$  is explicit,  $\top$  satisfies the properties of an implicit calculus by Definition 43. Moreover, for every calculus  $\mathfrak{B} \in \mathbf{S}$  such that  $\mathfrak{B} \neq \top$  and  $\mathfrak{B} \notin \top$ , it must be the case that  $\mathfrak{B}$  contains Horn rules by the definition of  $\notin$ .

Theorem 50. Let  $\mathfrak{A}$  be an explicit calculus,  $\mathbb{U}(\mathfrak{A}) = (\mathbf{S}, \leqslant)$ ,  $\mathfrak{B}$  be an implicit calculus, and  $\mathbb{D}(\mathfrak{B}) = (\mathbf{S}', \leqslant)$ . If either  $\mathfrak{B} \in \mathbf{S}$  or  $\mathfrak{A} \in \mathbf{S}'$ , then  $\mathbf{S} = \mathbf{S}'$ ,  $\mathbb{U}(\mathfrak{A}) \cong \mathbb{D}(\mathfrak{B})$ , and for any  $\mathfrak{C}, \mathfrak{D} \in \mathbf{S} = \mathbf{S}'$ ,  $\mathbf{G}(\mathfrak{C}) = \mathbf{G}(\mathfrak{D})$ .

*Proof*: Assume w.l.o.g that  $\mathfrak{B} \in \mathbf{S}$ . As  $\mathbb{U}(\mathfrak{A})$  contains only one implicit calculus (Theorem 47) we know that  $\mathfrak{B}$  is the top element of  $\mathbb{U}(\mathfrak{A})$ . Let  $\mathrm{H} := \mathrm{H}(\mathfrak{A})$ , and observe that for any anti-fracturable subset  $\mathrm{H}'$  of  $\mathrm{H}$ ,  $f(\mathfrak{A}, \mathrm{H}') = g(\mathfrak{B}, P(\mathrm{H} \setminus \mathrm{H}'))$ . We define  $h(f(\mathfrak{A}, \mathrm{H}')) = g(\mathfrak{B}, P(\mathrm{H} \setminus \mathrm{H}'))$ , and note that hserves as an isomorphism between  $\mathbb{U}(\mathfrak{A})$  and  $\mathbb{D}(\mathfrak{A})$ , meaning  $\mathbb{U}(\mathfrak{A}) \cong \mathbb{D}(\mathfrak{B})$ . The fact that for any  $\mathfrak{C}, \mathfrak{D} \in \mathbf{S} = \mathbf{S}', \mathbf{G}(\mathfrak{C}) =$  $\mathbf{G}(\mathfrak{D})$  follows from the fact that all Horn rules of  $\mathfrak{A}$  either occur explicitly in any calculus of  $\mathbb{U}(\mathfrak{A})$  or were absorbed into initial and reachability rules. ■