Liouvillian exceptional points in continuous variable system

B. A. Tay^a

^aDepartment of Foundation in Engineering, Faculty of Science and Engineering, University of Nottingham Malaysia, Jalan Broga, 43500 Semenyih, Selangor, Malaysia

Abstract

The Liouvillian exceptional points for a quantum Markovian master equation of an oscillator in a generic environment are obtained. They occur at the points when the modified frequency of the oscillator vanishes, whereby the eigenvalues of the Liouvillian become real. In a generic system there are two parameters that modify the oscillator's natural frequency. One of the parameters can be the damping rate. The exceptional point then corresponds to critical damping of the oscillator. This situation is illustrated by the Caldeira–Leggett (CL) equation and the Markovian limit of the Hu–Paz–Zhang (HPZ) equation. The other parameter changes the oscillator's effective mass whereby the exceptional point is reached in the limit of extremely heavy oscillator. This situation is illustrated by a modified form of the Kossakowski–Lindblad (KL) equation. The eigenfunctions coalesce at the exceptional points and break into subspaces labelled by a natural number *N*. In each of the *N*-subspace, there is a (*N* + 1)-fold degeneracy and the Liouvillian has a Jordan block structure of order- $(N + 1)$. We obtain the explicit form of the generalized eigenvectors for a few Liouvillians. Because of the degeneracies, there is a freedom of choice in the generalized eigenfunctions. This freedom manifests itself as an invariance in the Jordan block structure under a similarity transformation whose form is obtained. We compare the relaxation of the first excited state of an oscillator in the underdamped region, critically damped region which corresponds to the exceptional point, and overdamped region using the generalized eigenvectors of the CL equation.

1. Introduction

In open quantum systems, the generators governing the evolution of a system are intrinsically dissipative and hence non-Hermitian. At certain choices of parameters of the system, the eigenvalues of the generator and the corresponding eigenvectors could coalesce. The points in the parameter space at which the eigenvalues coalesce are called exceptional points [1, 2]. The exceptional points are different from degeneracy of eigenvalues in conservative systems. Whereas the generators of evolution in conservative systems can be diagonalized at degenerate eigenvalues, those in dissipative systems can at best be brought into a Jordan canonical form. In scattering theory, the coalescence of eigenvalues manifests itself as double or higher order poles in the complex energy plane [3, 4, 5, 6] that lead to resonances.

The occurrence of exceptional points are ubiquitous in dissipative systems. It was observed in microwave cavity [7], optical cavity [8], atom–cavity [9], electronics [10], photonics [11, 12], driven superconducting qubits [13] and etc. It was found that at the exceptional points, the enhanced gain or loss of the system, or a balance between them could potentially lead to novel applications. This had been exploited to create enhanced sensors [14, 15], nonreciprocal topological energy transfer [16], revival of lasing [17], and etc. There were also connections between exceptional points and the symmetry breaking in PT symmetric Hamiltonians [18] as well as the onset of quantum phase transitions [19].

Recently, there are increasing interests in extending the understanding of exceptional points physics from the level of Hamiltonian to the level of Liouvillian, i.e., the evolution generator of quantum master equation. The Liouvillian of quantum master equation is a non-Hermitian superoperator [20]. Along this line, the exceptional points of the

Email address: BuangAnn.Tay@nottingham.edu.my (B. A. Tay)

Preprint submitted to Physica A 12 April 2023

Liouvillians of finite-level systems were explored in Ref. [21] and from the perspective of quantum jumps in Refs. [22, 23, 24].

In this work we focus on the Liouvillian of an infinite-level or continuous variable system. Utilizing the results obtained recently on the spectrum of the Liouvillian of a quantum oscillator in a generic environment [25], we obtain the exceptional points of the Liouvillian at arbitrary order and the generalized eigenfunctions from the Jordan block structure of the Liouvilian.

We first discuss the degeneracy in the Kossakowski–Lindblad (KL) equation in 2. Then, the approach to exceptional points in a generic quantum master equation are elucidated with two examples in 3. The generalized eigenvectors are obtained in 4. This is followed by a discussion on the invariance of the Jordan block structure of the reduced dynamics under similarity transformation in 5. We illustrate the results by comparing the evolution of a quantum state in three regions of damping in 6. A conclusion then follows. Some of the details of the calculations are presented in the appendices.

2. Degeneracy in Kossakowski–Lindblad (KL) equation

We consider a quantum Markovian master equation, $\partial \rho / \partial t = -K\rho$, for the density operator ρ of a harmonic oscillator. The effect of the environment is encoded in the Liouvillian *K*. We start by considering the Kossakowski– Lindblad (KL) equation [26, 27] (also known as the Gorini–Kossakowski–Sudarshan–Lindblad or simply the Lindblad equation) which is widely used to describe the influence of environment on an oscillator in many fields [28, 29, 30, 31], such as in quantum information, quantum optics, condensed matter, energy transfer in molecular systems, and etc. Its Liouvillian is usually written in terms of the creation and annihilation operators of a harmonic oscillator as

$$
K_{\text{KL}}\rho = i\omega_0[a^\dagger a,\rho] - \frac{\gamma}{2}\left(b + \frac{1}{2}\right)(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a) - \frac{\gamma}{2}\left(b - \frac{1}{2}\right)(2a^\dagger \rho a - a a^\dagger \rho - \rho a a^\dagger),\tag{1}
$$

where γ is a relaxation rate and *b* is related to the temperature *T* of the environment by $b = \frac{1}{2} \coth[\hbar \omega/(2kT)]$. At absolute zero when $b = 1/2$ the last group of operators in K_{W} o drops out absolute zero when $b = 1/2$, the last group of operators in $K_{KL}\rho$ drops out.

To continue our discussion, we specialize in the position coordinates. We first introduce the dimensionless position coordinate $x \equiv \sqrt{m\omega_0/\hbar} q$, where q is the position coordinate with a dimension of length. In the Liouville space, we
have the bra-space \mathbb{R}^n and the ket-space \mathbb{R}^n . The density function in the position coo have the bra-space $|\tilde{x}\rangle$ and the ket-space $\langle x|$. The density function in the position coordinates is denoted by $\langle x|\rho|\tilde{x}\rangle$. It is customary to introduce the centre and relative coordinates

$$
Q = \frac{1}{2}(x + \tilde{x}), \qquad r \equiv x - \tilde{x}, \tag{2}
$$

respectively, and write the density function as

$$
\langle x|\rho|\tilde{x}\rangle = \left\langle Q + \frac{r}{2}|\rho|Q - \frac{r}{2}\right\rangle \equiv \rho(Q, r). \tag{3}
$$

The annihilation operator in the position coordinates is $a = (\hat{x} + i\hat{p})/$
the dimensionless position and momentum operators. Then √ $2 = (\hat{x} + \partial/\partial \hat{x})$ √ 2, where \hat{x} and $\hat{p} = -i\partial/\partial \hat{x}$ are the dimensionless position and momentum operators. Then,

$$
\langle x|a\rho|\tilde{x}\rangle = \frac{1}{\sqrt{2}}\left(x + \frac{\partial}{\partial x}\right)\langle x|\rho|\tilde{x}\rangle = \frac{1}{\sqrt{2}}\left(Q + \frac{r}{2} + \frac{1}{2}\frac{\partial}{\partial Q} + \frac{\partial}{\partial r}\right)\rho(Q, r). \tag{4}
$$

Similarly,

$$
\langle x|\rho a|\tilde{x}\rangle = \frac{1}{\sqrt{2}} \left(\tilde{x} - \frac{\partial}{\partial \tilde{x}}\right) \langle x|\rho|\tilde{x}\rangle = \frac{1}{\sqrt{2}} \left(Q - \frac{r}{2} - \frac{1}{2}\frac{\partial}{\partial Q} + \frac{\partial}{\partial r}\right) \rho(Q, r).
$$
 (5)

Eqs. (4) and (5) are defined under a trace, see 5.1. In the process of getting Eq. (5), an integration by parts is carried out with an assumption that $\langle x|\rho|\tilde{x}\rangle$ vanishes fast enough at the infinity. $\langle x|\rho|\tilde{x}\rangle$ with the form of a Gaussian multiplying a polynomial is a useful example of functions satisfying this requirement. The corresponding expressions of creation operator in the position coordinates can be obtained similarly by using $a^{\dagger} = (\hat{x} - i\hat{p})/\sqrt{2} = (\hat{x} - \partial/\partial \hat{x})\sqrt{2}$.

In this way, in the position coordinates the Liouvillian K_{KL} can be simplified to

$$
K_{\text{KL}}(Q,r) = i\omega \left(-\frac{\partial^2}{\partial Q \partial r} + Qr \right) - \frac{\gamma}{2} \left(\frac{\partial}{\partial Q} Q - r \frac{\partial}{\partial r} \right) - b\frac{\gamma}{2} \left(\frac{\partial^2}{\partial Q^2} - r^2 \right). \tag{6}
$$

The eigenvalue equation

$$
K_{\text{KL}}f_{mn}^{\pm} = \lambda_{mn}^{\pm}f_{mn}^{\pm}, \qquad 0 \le n \le m = 0, 1, 2, \dots,
$$
 (7)

has the solution [32, 33, 34]

$$
\lambda_{mn}^{\pm} = \pm in\omega + (2m - n)\frac{\gamma}{2}, \qquad 0 \le n \le m = 0, 1, 2, \dots
$$
 (8)

The eigenfunctions $f_{mn}^{\pm}(Q, r)$ can be decomposed into two parts [33],

$$
f_{mn}^{\pm}(Q,r) = \Pi_{mn}^{\pm}(Q,r)f_{00}(Q,r), \qquad 0 \le n \le m = 0, 1, 2, \dots,
$$
 (9)

where

$$
f_{00}(Q,r) = \frac{1}{\sqrt{2\pi b}} e^{-Q^2/(2b) - br^2/2}
$$
\n(10)

 $\sqrt{2\pi b}$
is a stationary state, whereas $\Pi_{mn}^{\pm}(Q, r)$ is a polynomial in *Q* and *r*. Its expression is given in Appendix A for the
convenience of the reader. In the following discussion, we will omit the coordinates convenience of the reader. In the following discussion, we will omit the coordinates dependence on the functions when confusion does not arise.

Fig. 1(a) gives the positions of the eigenvalues (8) on a complex plane. As ω decrease to zero, two or more eigenvalues such as λ_{mn}^{\pm} and $\lambda_{mn'}^{\pm}$, are degenerate whenever the condition $2m' - n' = 2m - n$ is met. Both the λ_{mn}^{\pm} and λ_{mn}^{\pm} and λ_{mn}^{\pm} and λ_{mn}^{\pm} and λ_{mn}^{\pm} and λ_{mn}^{\pm} and λ_{mn} ϵ_{m} and the contention of the label eigenvalues become real. At this point, it is convenient to introduce the label $\frac{1}{mn}$ move vertically towards the real horizontal axis, see Fig. 1(b), and meet there at $\omega = 0$, see Fig. 1(c), at which the

$$
N \equiv 2m - n, \qquad N = 0, 1, 2, 3, \dots
$$
 (11)

to write the eigenvalue as

$$
\lambda_N \equiv N \frac{\gamma}{2} \,. \tag{12}
$$

Each *N* form a subspace of eigenvectors with a $(N + 1)$ -fold degeneracy. Note that for even *N*, there is already an existing eigenvalue $\lambda_{N/2,0}$ on the real axis. When ω becomes imaginary, the eigenvalues remain real. The degeneracies that occur at $\omega = 0$ are lifted and the eigenvalues split from each other along the horizontal axis, see Fig. 1(d). The separation between two nearest eigenvalues of the same *^N*-series is 2|ω|. In analogy to a damped oscillator in classical mechanics [37], Fig. 1(a) and Fig. 1(b) represent the case of an underdamped oscillator, Fig. 1(c) a critically damped oscillator, and Fig. 1(d) an overdamped oscillator.

For the KL equation, the eigenfunctions of degenerate eigenvectors remain distinct. As a result, the structure of the eigenvalue equation remains intact. This is because the unitary part of the Liouvillian *K*⁰ (the terms on the righthand-side (RHS) of Eq. (6) containing ω), and the rest of its dissipative part K_d containing γ are separable, i.e., they separately satisfy the eigenvalue equation

$$
K_0 f_{mn}^{\pm} = \pm in\omega f_{mn}^{\pm} , \qquad K_0 f_{mn}^{\pm} = (2m - n) \frac{\gamma}{2} f_{mn}^{\pm} . \tag{13}
$$

Consequently, the dynamics of the unitary and dissipative part do not affect each other in the limit of zero ω .

We illustrate this with the $m = 1$ subspace. Taking the limit $\omega \to 0$, there are two degenerate eigenvalues $j_{11}^+ = \lambda_{11}^- = \gamma/2$. The eigenfunctions remain linearly independent,

$$
f_{11}^{\pm} = \mp i \left(\frac{Q}{\sqrt{2b}} \pm \sqrt{\frac{b}{2}} r \right) f_{00}
$$
 (14)

Consequently, there is a 2-fold degeneracy in this subspace

$$
K_{\text{KL}} f_{11}^{\pm} = \frac{\gamma}{2} f_{11}^{\pm} \,. \tag{15}
$$

We note that f_{11}^+ and f_{11}^- remain orthogonal to each other [33].

Figure 1: The change in the positions of eigenvalues λ_{mn}^{\pm} (8) on a complex plane as real ω reduces to 0 (the exceptional point) and then turns imaginary Horizontal and vertical axis are scaled in units of $\alpha/2$ imaginary. Horizontal and vertical axis are scaled in units of $\gamma/2$ and ω , respectively. (a) The eigenvalues away from the exceptional point. Dashed lines connect eigenvalues with the same $m = (N + n)/2$. The corresponding eigenvalue is given above the cross. The position of λ_{mn}^- is a mirror
reflection of λ_{mn}^+ along the horizontal axis. (b) As ω decreases the reflection of λ_{mn}^+ along the horizontal axis. (b) As ω decreases, the eigenvalues move vertically towards the horizontal axis from both the above and below of the axis. (c) At the exceptional point, the eigenvalues and below of the axis. (c) At the exceptional point, the eigenvalues λ_N (12) coalesce on the horizontal axis and become real. They are labelled by circles with a cross. They have a $(N + 1)$ -fold degeneracy. (d) When ω turns imaginary, the degeneracies are removed. The eigenvalues split away from each other along the horizontal axis. The distance between two nearest eigenvalues of the same *^N*-subspace is 2|ω|. The eigenvalues that correspond to the crosses are given in the respective order up to $N = 4$ for illustration.

3. Exceptional points in quantum master equation

In previous works [35, 36], it was shown that the Liouvillian of a generic quantum Markovian master equation has the form

$$
K = 2\omega_0 i L_0 + h_1 i M_1 + h_2 i M_2 + \gamma (O_0 - I/2) + g_0 O_+ + g_1 L_{1+} + g_2 L_{2+}.
$$
\n(16)

The h_i and g_i are constant parameters, whereas γ is a damping or relaxation constant of the system. The expressions of the seven operators in the position coordinates are given in Appendix A. We note that K_{KL} (1) is obtained by choosing $g_0 = -2\gamma b$ and zero for the rest of the parameters. *K* can be related to the Liouvillian of the KL equation by a similarity transformation $K = S K_{KL} S^{-1}$ [25], provided that the modified frequency of system,

$$
\omega = \sqrt{\omega_0^2 - \frac{h_1^2}{4} - \frac{h_2^2}{4}},\tag{17}
$$

satisfies the condition

$$
\omega^2 + \frac{\gamma^2}{4} \neq 0. \tag{18}
$$

Then, the eigenvalue of the transformed system is also given by Eq. (8), with eigenfunction $S f_{mn}^{\pm}$.

In elementary mechanics, there are three regions of harmonic motion of an oscillator under a friction force [37, 38], i.e., underdamping, critical damping and overdamping that correspond to real $\omega > 0$, $\omega = 0$ and ω imaginary, respectively. The consideration carries over into the reduced dynamics of a quantum oscillator. From the discussion in the previous section, degeneracy occurs when $\omega = 0$ at critical damping. For a given natural frequency ω_0 , points of degeneracy lie on a circle of radius $2\omega_0$ in the (h_1, h_2) -parameter space

$$
h_1^2 + h_2^2 = 4\omega_0^2.
$$
 (19)

We will consider two independent ways in the approach to $\omega = 0$, (1) from $h_2 \rightarrow 2\omega_0$ with $h_1 = 0$ in the Caldeira– Leggett (CL) equation, and (2) from $h_1 \rightarrow 2\omega_0$ with $h_2 = 0$ in the modified Kossakowski–Lindblad (KL) equation. For generic quantum master equation, the approach to zero ω exhibits behaviour that is distinct from the KL equation.

3.1. Caldeira–Leggett (CL) equation

We first consider the Caldeira–Leggett (CL) equation [39] with $h_2 = -\gamma$, $g_0 = g_1 = -2\gamma \bar{b}$ in Eq. (16), whereas the rest of the parameters are zero. We will used "bar" to denote quantities related to the CL equation. In terms of the creation and annihilation operators of harmonic oscillator, the CL equation is written as

$$
\bar{K}_{\text{CL}} = i\omega_0[a^{\dagger}a,\rho] + i\frac{\gamma}{2}[\hat{x},\{\hat{p},\rho\}] + \gamma\bar{b}[\hat{x},[\hat{x},\rho]],
$$
\n(20)

where $\{\cdot,\cdot\}$ denotes an anti-commutator bracket. In the high temperature limit, the second term with the anti-commutator bracket can be dropped, as is usually done in the studies on environmental decoherence [40, 41].

In the position coordinates, the Liouvillian of the CL equation is

$$
\bar{K}_{\text{CL}}(\bar{Q}, \bar{r}) = i\omega_0 \left(-\frac{1}{2} \frac{\partial^2}{\partial \bar{Q} \partial \bar{r}} + 2 \bar{Q} \bar{r} \right) + \gamma \bar{r} \frac{\partial}{\partial \bar{r}} + 2\gamma \bar{r}^2, \tag{21}
$$

where \bar{b} is absorbed into the coordinates

$$
\bar{Q} \equiv \frac{Q}{\sqrt{2\bar{b}}}, \qquad \bar{r} \equiv \sqrt{\frac{\bar{b}}{2}}r. \tag{22}
$$

 \bar{K}_{CL} is similarly related to K_{KL} by $\bar{K}_{\text{CL}} = S_2 K_{\text{KL}} S_2^{-1}$ [25], where

$$
S_2 = e^{i\phi M_1} e^{\eta L_2}.
$$
 (23)

The stationary state is

$$
\bar{f}_{00}(\bar{Q}, \bar{r}) = S_2 f_{00}(Q, r) = \frac{1}{\sqrt{\pi}} e^{-\bar{Q}^2 - \bar{r}^2}.
$$
\n(24)

The corresponding function $\bar{\Pi}^{\pm}_{mn}$ in Eq. (9) can be obtain in a similar way from f^{\pm}_{mn} . We refer the reader to Ref. [25] for the details. A few \bar{f}^{\pm}_{mn} are listed in Appendix B.1.

The damped oscillator has the modified frequency

$$
\omega_2 \equiv \sqrt{\omega_0^2 - \gamma^2/4} \,. \tag{25}
$$

For the approach of ω_2 to zero at critical damping, we consider the limit $\gamma \to 2\omega_0$ from the side $\gamma < 2\omega_0$ so that ω_2 remains real. In this limit, the eigenvalues are still given by Eq. (12). However, small denominator could develop in the eigenfunctions leading to divergence. As an example, we illustrate this with the eigenfunction \bar{f}_{11}^{\pm} under the $N = 1$ subspace of the CL equation. In Appendix B.1, we expand \bar{f}_{11}^{\pm} in powers of the parameter

$$
\delta_2 \equiv 1 - \frac{\gamma}{2\omega_0} \,. \tag{26}
$$

In the limit $\omega_0 \rightarrow \gamma/2$, the most singular term in the series is

$$
\bar{f}_{11}^{\pm} = \frac{1}{\sqrt{\delta_2}} \frac{1 \mp i}{2} (\bar{Q} - i\bar{r}) \bar{f}_{00} + O(\delta_2^0), \qquad (27)
$$

which has a square root singularity. Eigenfunctions with higher order *N* will have increasingly higher power of square root singularity. The singularity shows that this is not a case of simple degeneracy. Furthermore, notice that both eigenvectors are the same up to an overall constant, i.e., they are no longer orthogonal. $\omega_2 = 0$ is called the exceptional point to distinguish it from ordinary degeneracy.

Another Liouvillian with a similar behaviour is the Markovian limit of the HPZ equation [42, 43] which is related to the CL equation by a similarity transformation. Owing to its similarity with the CL equation, it approaches the exceptional point in a similar way with the CL equation. We present the results for the HPZ equation in Appendix E.

3.2. Modified Kossakowski–Lindblad (KL) equation

We consider another model with a different approach to the exceptional points. We add a $h_1 i M_1$ term (A.7) to K_{KL} (1) to get a modified KL equation. The Liouvillian becomes

$$
\tilde{K}_{\text{mKL}}(\tilde{Q}, \tilde{r}) = -\frac{i}{2} \left(\omega_0 - \frac{h_1}{2} \right) \frac{\partial^2}{\partial \tilde{Q} \partial \tilde{r}} + 2i \left(\omega_0 + \frac{h_1}{2} \right) \tilde{Q} \tilde{r} - \frac{\gamma}{2} \left(\frac{\partial}{\partial \tilde{Q}} \tilde{Q} - \tilde{r} \frac{\partial}{\partial \tilde{r}} + \frac{1}{2} \frac{\partial^2}{\partial \tilde{Q}^2} - 2\tilde{r}^2 \right),\tag{28}
$$

where we define

$$
\tilde{Q} \equiv \frac{Q}{\sqrt{2\tilde{b}}}, \qquad \tilde{r} \equiv \sqrt{\frac{\tilde{b}}{2}}r,\tag{29}
$$

and use "tilde" to denote quantities related to the modified KL equation.

Due to the interaction between the oscillator and the environment, the oscillator acquires an effective mass

$$
\tilde{m} \equiv m \left(1 - \frac{h_1}{2\omega_0} \right)^{-1} . \tag{30}
$$

The frequency of the oscillator is modified to

$$
\omega_1 \equiv \sqrt{\omega_0^2 - h_1^2/4} \,. \tag{31}
$$

We approach the limit of heavy oscillator $h_1 \rightarrow 2\omega_0$ from the side $h_1 < 2\omega_0$, so that ω_1 is real.

 \tilde{K} is similarly related to K_{KL} through

$$
S_1 = e^{\eta_0 L_{1+}} e^{\psi i M_2} e^{\eta_2 i L_{2+}}.
$$
\n(32)

Using the method discussed in Ref. [25], we apply the transformation to the stationary state of the KL equation to get the stationary state of \tilde{K}_{mKL} ,

$$
\tilde{f}_{00}(\tilde{Q},\tilde{r}) = S_1 f_{00} = \frac{1}{\sqrt{\pi}} \sqrt{\frac{\omega_1^2 + \gamma^2/4}{\omega_0(\omega_0 - h_1/2) + \gamma^2/4}} \exp\left[-\frac{(\omega_1^2 + \gamma^2/4)\tilde{Q}^2 + \gamma h_1 i \tilde{Q}\tilde{r}/2 + (\omega_0^2 + \gamma^2/4)\tilde{r}^2}{\omega_0(\omega_0 - h_1/2) + \gamma^2/4} \right].
$$
 (33)

The corresponding transformed polynomial $\tilde{\Pi}^{\pm}_{mn} = S_1 \Pi^{\pm}_{mn} S_1^{-1}$, cf. Eq. (9), can be worked out in the same way. A few of them can be found in Appendix B.2.

In the case of the modified KL equation, we show in Appendix B.2 that in the limit $h_1 \rightarrow 2\omega_0$, two of the eigenfunctions in the $N = 1$ subspace go into

$$
\tilde{f}_{11}^{\pm} = -\sqrt{2}\tilde{\omega}ir\tilde{f}_{00} + O\left(\sqrt{\delta_1}\right),\tag{34}
$$

where

$$
\tilde{\omega} \equiv \frac{2\omega_0}{\gamma} \,. \tag{35}
$$

In contrast to the eigenfunctions of \bar{K}_{CL} which develop small denominator, the eigenfunctions of \tilde{K}_{mKL} contain increasing power of the square root of a small numerator

$$
\delta_1 \equiv 1 - \frac{h_1}{2\omega_0} \,. \tag{36}
$$

Furthermore, both the \tilde{f}_{11}^+ and \tilde{f}_{11}^- are identical in this limit. This again implies that we lose a copy of linearly independent vector. Hence, $\omega_1 = 0$ is the exceptional point for the modified KL equation.

It is known in scattering theory that whenever the *S* -matrix has a pole of order *N*, its Hamiltonian when extended beyond the Hilbert space reveals a Jordan block structure of order *N* [4, 5]. On the level of density operator, we have seen that a change of parameters causes the eigenvalues to coalesce. At the exceptional points, the usual eigenvalue equation is no longer capable of describing the reduced dynamics. The eigenvalue equation has to be generalized to a Jordan block structure,

$$
K_E F_N^{(0)} = \lambda_N F_N^{(0)},\tag{37a}
$$

$$
K_E F_N^{(z)} = \lambda_N \left(F_N^{(z)} + F_N^{(z-1)} \right), \qquad z = 1, 2, ..., N,
$$
 (37b)

for an exceptional point of order *N*+1. In the following section we will construct higher-order generalized eigenvectors by solving the Jordan block structure (37a)–(37b) in a systematic way. There, we realize that Eqs. (27) and (34) give the zeroth-order vector $F_N^{(0)}$ up to a normalization constant.

We note that in the literature, the coefficient of $F_N^{(z-1)}$ in Eq. (37b) is usually chosen as 1. The discrepancy is due to a different definition in the normalization constant of $F_N^{(z)}$. The definition we adopt render all $F_N^{(z)}$ dimensionless. We will discuss the normalization of the generalized eigenfunctions in 5.1.

4. Generalized eigenvectors of quantum master equation

In this section we construct the generalized eigenvectors of the Liouvillian of CL and modified KL at the exceptional points. Similar to the eigenvectors of K_{KL} (9), we decompose the generalized eigenvectors into two parts,

$$
F_N^{(z)} = \Phi_N^{(z)} F_0^{(0)},\tag{38}
$$

where $F_0^{(0)}$ $\phi_0^{(0)}$ is the stationary state and $\Phi_0^{(0)}$ $\phi_0^{(0)} = 1$. $\Phi_N^{(z)}$ is a polynomial in the coordinates *Q* and *ir* with real coefficients. In Eq. (38) and the equations to follow, we omit the coordinates dependence on the functions to simplify the expressions.

Before we construct the eigenvectors, we discuss the hermiticity requirement on the density operator $\rho^{\dagger} = \rho$.
he coordinate representation, this requirement is translated into $\langle x|\sigma^{\dagger}|\tilde{x}\rangle = \langle \tilde{x}|\sigma|x\rangle^* = \langle x|\sigma^{\$ In the coordinate representation, this requirement is translated into $\langle x|\rho^{\dagger}|\tilde{x}\rangle = \langle \tilde{x}|\rho|x\rangle^* = \langle x|\rho|\tilde{x}\rangle$, or equivalently, $\rho^*(Q - r) = \rho(Q, r)$. In the language of superpoperator, a superpoperator with this proper symmetric [44, 36]. We can put a function in an explicitly adjoint-symmetric form by writing the function as a ^{*}(*Q*, -*r*) = ρ (*Q*,*r*). In the language of superoperator, a superoperator with this property is also called adjoint-
wmmetric [*AA*, **36]** We can put a function in an explicitly adjoint-symmetric form by writing t polynomial in powers of *Q* and *ir*. The coefficients in the polynomials are then real numbers.

We find that the $\Phi_N^{(z)}$ can be decomposed into a combination of lower order generalized eigenvectors multiplying terms that are linear in the coordinates. We start with two ansatz, which we call the diagonal and parallel representation of the generalized eigenvector,

Diagonal:
$$
\Phi_N^{(z)} = (u_N^{(z)}Q + v_N^{(z)}ir)\Phi_{N-1}^{(z)} + w_N^{(z)}\Phi_{N-2}^{(z)}, \qquad z \neq N,
$$
(39a)

Parallel:
$$
\Phi_N^{(z)} = (p_N^{(z)}Q + q_N^{(z)}ir)\Phi_{N-1}^{(z-1)} + s_N^{(z)}\Phi_{N-2}^{(z-2)}, \qquad z \neq 0,
$$
 (39b)

where *u*, *v*, *w*, *p*, *q*, *s* are real constant. By definition, we set $\Phi_N^{(z)} = 0$ whenever $z < 0$ or $z > N$ or $N < 0$. We note that additional terms, such as $x_N^{(z)} \Phi_{N-2}^{(z-1)}$ might need to be added to Eqs. (39a)–(39b) for different choices of generalized eigenvectors, see Appendix C for the details.

We summarize the procedure here. The details are left to Appendix C and Appendix D.m We start with the subspace $N = 1$ by applying Eq. (37a) to the lowest order generalized eigenvector $F_1^{(0)}$ using the diagonal representation (39a). By comparing the coefficients of the polynomial in *Q* and *ir* on both sides of the equation, we solve for their (39a). coefficients. We next apply Eq. (37b) to $F_1^{(1)}$ $_1^{(1)}$ using the parallel representation and solve for the coefficients. Then we move on to the next subspace of generalized eigenvectors with $N = 2$. In this way we can determine the coefficients of the polynomials from one order to the next for all *N*.

z	$N = 0$	$N = 1$	$N = 2$	$N = 3$
0	1	$\bar{Q} - i\bar{r}$	$(\bar{Q} - i\bar{r})^2 - \frac{1}{2}$	$(\bar{Q} - i\bar{r})^3 - \frac{3}{2}(\bar{Q} - i\bar{r})$
1	$-i\bar{r}$	$-2i\bar{r}(\bar{Q} - i\bar{r})$	$-3i\bar{r}((\bar{Q} - i\bar{r})^2 - \frac{1}{2})$	
2	$2(i\bar{r})^2$	$\frac{9}{2}(i\bar{r})^2(\bar{Q} - i\bar{r})$		
3	$-\frac{9}{2}(i\bar{r})^3$			

Table 1: The first few series of $\bar{\Phi}_N^{(z)}$.

4.1. Caldeira–Leggett (CL) equation

The exceptional points for \bar{K}_{CL} is reached by increasing the damping to $\gamma = 2\omega_0$ where the modified frequency ω_2
ishes. At the exceptional point \bar{K}_{CL} (21) turns into vanishes. At the exceptional point, \bar{K}_{CL} (21) turns into

$$
\bar{K}_E(\bar{Q}, \bar{r}) \equiv i \frac{\gamma}{2} \left(-\frac{1}{2} \frac{\partial^2}{\partial \bar{Q} \partial \bar{r}} + 2 \bar{Q} \bar{r} \right) + \gamma \bar{r} \frac{\partial}{\partial \bar{r}} + 2 \gamma \bar{r}^2.
$$
\n(40)

The details in obtaining the coefficients in Eqs. (39a)–(39b) are worked out in Appendix C. The lowest order generalized eigenvector $\bar{F}_0^{(0)}$ $_{0}^{(0)}$ is identical with \bar{f}_{00} (24), and $\bar{\Phi}_{0}^{(0)}$ $\binom{00}{0}$ = 1. The higher order generalized eigenvectors can be summarized as

$$
\text{Diagonal}: \qquad \frac{\bar{\Phi}_N^{(z)}}{N^z} = (\bar{Q} - i\bar{r}) \frac{\bar{\Phi}_{N-1}^{(z)}}{(N-1)^z} - \frac{N-1-z}{2} \frac{\bar{\Phi}_{N-2}^{(z)}}{(N-2)^z}, \qquad z \neq N, \tag{41a}
$$

Parallel :
$$
\frac{\bar{\Phi}_{N}^{(z)}}{N^{z}} = \frac{1}{z}(-i\bar{r}) \frac{\bar{\Phi}_{N-1}^{(z-1)}}{(N-1)^{z-1}}, \qquad z \neq 0,
$$
 (41b)

Recall that we set $\bar{\Phi}_N^{(z)} = 0$ whenever $z < 0$ or $z > N$ or $N < 0$. A few examples are listed in 4.1. In the overlapping
region of the two equations i.e., when $z = 1, 2, 3$ $N = 1$, we can show that the two sets of equa region of the two equations, i.e., when $z = 1, 2, 3, \ldots, N-1$, we can show that the two sets of equation (41a) and (41b) are identical using a proof by induction. The details are presented in Appendix C 2 and Appendix C 3. are identical using a proof by induction. The details are presented in Appendix C.2 and Appendix C.3.

This set of generalized eigenvectors is not unique. The non-uniqueness arises from the degeneracies. It reveals itself as a freedom in the choice of the coefficients in the diagonal and parallel representation. Examples are given in Appendix C.1. We will elaborate this freedom of choice in 5.2.

4.2. Modified Kossakowski–Lindblad (KL) equation

In the second approach to the exceptional point in \tilde{K}_{mKL} , we increase the effective mass of the oscillator until the modified frequency ω_1 vanishes in the limit $h_1 \to 2\omega_0$. In this limit, the Liouvillian \tilde{K}_{mKL} (28) can be cast into the form

$$
\tilde{K}_E(\tilde{Q}, \tilde{r}) = 4i\omega_0 \tilde{Q}\tilde{r} - \frac{\gamma}{2} \left(\frac{1}{2} \frac{\partial^2}{\partial \tilde{Q}^2} + \tilde{Q} \frac{\partial}{\partial \tilde{Q}} - \tilde{r} \frac{\partial}{\partial \tilde{r}} - 2\tilde{r}^2 + 1 \right).
$$
\n(42)

The zeroth-order generalized eigenvector \tilde{F}_0^0 is obtained from the stationary state in Eq. (33) by setting $h_1 = 2\omega_0$. It can be simplified to

$$
\tilde{F}_0^{(0)} = \frac{1}{\sqrt{\pi}} e^{-\tilde{Q}^2 - 2\tilde{\omega} i \tilde{Q} \tilde{r} - (1 + \tilde{\omega}^2) \tilde{r}^2},\tag{43}
$$

where $\tilde{\omega}$ is already defined in Eq. (35).

\mathcal{Z}	$N=0$	$N=1$	$N=2$	$N=3$
0		i $\tilde{\omega}$ r̃	$(i\tilde{\omega}\tilde{r})^2$	$(i\tilde{\omega}\bar{r})^3$
1		$\tilde{Q}/2$	$i\tilde{\omega}\tilde{r}\tilde{Q}$	$rac{3}{2}(i\tilde{\omega}\tilde{r})^2\tilde{Q}$
2			$rac{1}{2}\tilde{Q}^2-\frac{1}{4}$	$\frac{9}{8}(i\tilde{\omega}\tilde{r})\left(\tilde{Q}^2-\frac{1}{2}\right)$
3				$\frac{9}{16}\tilde{Q}\left(\tilde{Q}^2-\frac{3}{2}\right)$

Table 2: The first few series of $\tilde{\Phi}_N^{(z)}$.

In Appendix D, we construct the higher order generalized eigenvectors in increasing the procedure described in the paragraph after Eqs. (39a)–(39b). The results are

$$
\text{Diagonal}: \quad \frac{\tilde{\Phi}_N^{(z)}}{N^z} = i\tilde{\omega}\tilde{r}\frac{\tilde{\Phi}_{N-1}^{(z)}}{(N-1)^z}, \quad z \neq N, \tag{44a}
$$

Parallel :
$$
\frac{\tilde{\Phi}_N^{(z)}}{N^z} = \frac{\tilde{Q}}{2z} \frac{\tilde{\Phi}_{N-1}^{(z-1)}}{(N-1)^{z-1}} - \frac{1}{8z} \frac{\tilde{\Phi}_{N-2}^{(z-2)}}{(N-2)^{z-2}}, \qquad z \neq 0.
$$
 (44b)

Again, we set $\tilde{\Phi}_N^{(z)} = 0$ whenever $z < 0$ or $z > N$ or $N < 0$. The generalized eigenvectors from the first few subspaces are given in 4.2.

We can also prove by induction that in the overlapping region of the equations, $z = 1, 2, \ldots, N - 1$, both equations are identical. The set of generalized eigenvectors again are not unique. Its origin will be discussed in the next section.

5. Freedom in the basis of generalized eigenvectors

5.1. Normalization of generalized eigenvectors

Let us discuss the normalization of generalized eigenvectors. The trace of a density operator in the coordinate representation is defined as (by setting $\tilde{x} = x$ and integrate over *x*)

$$
\text{tr}\rho = \int_{-\infty}^{\infty} \rho(x, r = 0) \, dx = 1 \,. \tag{45}
$$

We note that except for the stationary state $F_0^{(0)}$ $_{0}^{(0)}$, all decaying vectors have zero trace [33, 45],

$$
\int_{-\infty}^{\infty} F_N^{(z)}(x, r = 0) dx = \delta_{N0} \delta_{z0}.
$$
\n(46)

From Eqs. (37a)–(37b), we deduce that once we fix the overall constant of $F_N^{(0)}$, the overall constant of higher order vectors in the same series are also fixed.

We mentioned at the end of 3 that in the literature the coefficient of $F_N^{(z-1)}$ in Eq. (37b) chosen as 1. We can bring it into the form of Eq. (37b), or vice versa, by a redefinition of the overall constant in $F_N^{(z)}$. In the system we consider, λ_N is real and non-zero. We can redefine $F_N^{(z)}$ as $F_N'^{(z)} \equiv F_N^{(z)}/\lambda_N^z$. Then Eq. (37a) r becomes

$$
KF_N^{\prime(z)} = \lambda_N F_N^{\prime(z)} + F_N^{\prime(z-1)},\tag{47}
$$

which is the usual way of writing the generalized eigenvalue equation in the literature. By adopting the normalization implied by Eqs. (37a)–(37b), all the $F_N^{(z)}$ are dimensionless. We can also simplify the expressions when we discuss the invariance of the Liouvillain at the exceptional point in the next section.

5.2. Freedom in generalized eigenvectors

As already mentioned in the last section, the set of generalized eigenvectors we have constructed are not unique because of degeneracies. Equally valid sets of generalized eigenvectors satisfying Eqs. (37a)–(37b) can be generated from the known set of vectors by exploiting this freedom.

(1) Linear sum of two series of generalized eigenvectors

Given two series of generalized eigenvectors of the same order, $F_N^{(z)}$ and $G_N^{(z)}$ that satisfy Eqs. (37a)–(37b), any linear combination of them with constant coefficients,

$$
H_N^{(z)} = c_1 F_N^{(z)} + c_2 G_N^{(z)}, \tag{48}
$$

are also the generalized eigenvectors of K_E . Note that the stationary states are always the same, $F_0^{(0)}$ $G_0^{(0)} = G_0^{(0)}$ $\binom{0}{0}$ = $H_0^{(0)}$,(v)
0 When we start with $F_N^{(z)}$ and $G_N^{(z)}$ that are adjoint-symmetric, cf. the discussion at the beginning of 4, we require c_1 and *c*₂ to be real so that $H_N^{(z)}$ is also adjoint-symmetric. We can then set $c_1 + c_2 = 1$ to fix the overall constant of $H_N^{(z)}$. *(2) Invariance of K^E at the exceptional points*

It is convenient to adopt a matrix representation to discuss the invariance of *K^E* under a similarity transformation. We write K_E in the subspace of generalized eigenvector of order N (37a)–(37b) as

$$
\mathbf{K}_N = \lambda_N \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 1 & 1 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 1 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 1 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 & 1 \end{pmatrix} . \tag{49}
$$

The generalized eigenvectors are

$$
\boldsymbol{F}_{N}^{(0)} \equiv \begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}, \quad \boldsymbol{F}_{N}^{(1)} \equiv \begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 1 \\ 0 \end{pmatrix}, \quad \cdots, \quad \boldsymbol{F}_{N}^{(N-1)} \equiv \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \boldsymbol{F}_{N}^{(N)} \equiv \begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}.
$$
 (50)

 K_N is then invariant under a similarity transformation

$$
\boldsymbol{T}_N \cdot \boldsymbol{K}_N \cdot \boldsymbol{T}_N^{-1} = \boldsymbol{K}_N \,, \tag{51}
$$

where the matrix T_N and its inverse are, respectively,

$$
\boldsymbol{T}_{N} = \begin{pmatrix} c_{0} & 0 & 0 & \cdots & 0 & 0 \\ c_{1} & c_{0} & 0 & \cdots & 0 & 0 \\ c_{2} & c_{1} & c_{0} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ c_{N-1} & c_{N-2} & c_{N-3} & \cdots & c_{0} & 0 \\ c_{N} & c_{N-1} & c_{N-2} & \cdots & c_{1} & c_{0} \end{pmatrix}, \qquad \boldsymbol{T}_{N}^{-1} = \begin{pmatrix} d_{0} & 0 & 0 & \cdots & 0 & 0 \\ d_{1} & d_{0} & 0 & \cdots & 0 & 0 \\ d_{2} & d_{1} & d_{0} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ d_{N-1} & d_{N-2} & d_{N-3} & \cdots & d_{0} & 0 \\ d_{N} & d_{N-1} & d_{N-2} & \cdots & d_{1} & d_{0} \end{pmatrix}.
$$
 (52)

The c_i and d_i are real coefficients and $c_0 \neq 0$. They are related by

$$
d_0 = \frac{1}{c_0},\tag{53}
$$

$$
d_z = -\frac{1}{c_0} \sum_{i=0}^{z-1} c_{z-i} d_i, \qquad z = 1, 2, 3, \dots, N. \tag{54}
$$

We note that the form (52) is preserved by $(T_N)^n$, where *n* is a natural number. The invariance in Eq. (51) implies that

$$
\boldsymbol{G}_{N}^{(z)} \equiv \boldsymbol{T}_{N} \cdot \boldsymbol{F}_{N}^{(z)}, \qquad \boldsymbol{H}_{N}^{(z)} \equiv \boldsymbol{T}_{N}^{-1} \cdot \boldsymbol{F}_{N}^{(z)}
$$
(55)

are also generalized eigenvectors of K_N satisfying Eqs. (37a)–(37b). The generalized eigenvectors generated from $F_N^{(z)}$ are

$$
G_N^{(z)} = \sum_{i=0}^{z} c_i F_N^{(z-i)}, \qquad H_N^{(z)} = \sum_{i=0}^{z} d_i F_N^{(z-i)} \qquad z = 0, 1, 2, ..., N,
$$
 (56)

*c*⁰ fixes the overall constant in the series of the generalized eigenvectors mentioned at the beginning of 5.2. Once the overall constant of $F_N^{(0)}$ is fixed, the overall constant of other generalized eigenvectors in the series then follows.

As an illustration of this freedom, we list a few interesting cases for the subspace $N = 3$.

 $G_3^{(1)}$

1. If we set $c_i = 1$ for all *i*, then $d_0 = 1, d_1 = -1$, and $d_2 = d_3 = 0$. The following series of $G_N^{(z)}$ and $H_N^{(z)}$ are produced,

$$
G_3^{(0)} = F_3^{(0)}, \qquad H_3^{(0)} = F_3^{(0)}, \qquad (57a)
$$

$$
G_3^{(1)} = F_3^{(1)} + F_3^{(0)}, \qquad H_3^{(1)} = F_3^{(1)} - F_3^{(0)}, \qquad (57b)
$$

$$
G_3^{(2)} = F_3^{(2)} + F_3^{(1)} + F_3^{(0)}, \qquad H_3^{(2)} = F_3^{(2)} - F_3^{(1)}.
$$
 (57c)

2. Setting $c_0 = 1$, $c_1 = c_3 = 0$, and $c_2 = c$, we get

$$
G_3^{(0)} = F_3^{(0)}, \qquad H_3^{(0)} = F_3^{(0)}, \qquad (58a)
$$

$$
H_3^{(1)} = F_3^{(1)}, \t H_3^{(1)} = F_3^{(1)}, \t (58b)
$$

$$
G_3^{(2)} = F_3^{(2)} + cF_3^{(0)}, \t H_3^{(2)} = F_3^{(2)} - cF_3^{(0)},
$$

\n
$$
G_3^{(3)} = F_3^{(3)} + cF_3^{(1)}, \t H_3^{(3)} = F_3^{(3)} - cF_3^{(1)}.
$$
\n(58c)

Notice that in this series, the even-order $(F_3^{(0)})$ $F_3^{(0)}$, $F_3^{(2)}$ $S_3^{(2)}$) and the odd-order ($F_3^{(1)}$) $F_3^{(1)}, F_3^{(3)}$ $3^{(3)}$) of generalized eigenvectors are mixed in separate subspace.

3. Setting $c_0 = 1$, $c_1 = c_2 = 0$ and $c_3 = d$, we can add arbitrary copies of $F_3^{(0)}$ $t_3^{(0)}$ to the highest order vector in a series. Then only the generalized eigenvector of highest order is modified,

$$
G_3^{(z)} = F_3^{(z)}, \t H_3^{(z)} = F_3^{(z)}, \t z = 0, 1, 2, (59a)
$$

\n
$$
G_3^{(3)} = F_3^{(3)} + dF_3^{(0)}, \t H_3^{(3)} = F_3^{(3)} - dF_3^{(0)}.
$$
\n(59b)

We can also apply the freedom to the generalized eigenvectors $\bar{\Phi}_N^{(z)}$ and $\tilde{\Phi}_N^{(z)}$ listed in ?? and ??, respectively. As shown in Appendix C.1, different choices of coefficients produce different series of generalized eigenvectors $\bar{\Psi}_{N}^{(z)}$ for the Liouvillian of the CL equation, \bar{K}_{E} . A few of them are listed in ??. In fact, the $\bar{\Phi}_{N}^{(z)}$ and $\Psi_{N}^{(z)}$ are connected by a similarity transformation $\Psi_N = \bar{T}_N \cdot \bar{\Phi}_N$, where for the first few subspaces $N = 1, 2, 3$, they are

$$
\bar{T}_1 = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \qquad \bar{T}_2 = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 2 & -2 & 1 \end{pmatrix}, \qquad \bar{T}_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -3 & 1 & 0 & 0 \\ \frac{9}{2} & -3 & 1 & 0 \\ -\frac{9}{2} & \frac{9}{2} & -3 & 1 \end{pmatrix}, \tag{60}
$$

respectively.

An alternate series of generalized eigenvectors of the Liouvillian of the modified KL equation, $\tilde{\Psi}_N^{(z)}$ using a different choice of coefficients chosen in Appendix D, is listed in ??. It happens that the $\tilde{\Phi}_{N}^{(z)}$ and $\tilde{\Psi}_{N}^{(z)}$ are related by the inverse of the similarity transformation in Eq. (60), $\tilde{\Psi}_N = \tilde{T}_N \cdot \tilde{\Phi}_N$, where $\tilde{T}_N = \bar{T}_N^{-1}$. The first few \tilde{T}_N are

$$
\tilde{\boldsymbol{T}}_1 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \qquad \tilde{\boldsymbol{T}}_2 = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 2 & 2 & 1 \end{pmatrix}, \qquad \tilde{\boldsymbol{T}}_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ \frac{9}{2} & 3 & 1 & 0 \\ \frac{9}{2} & \frac{9}{2} & 3 & 1 \end{pmatrix}.
$$
 (61)

Table 3: The first few series of $\Psi_N^{(z)}$.

Table 4: The first few series of $\tilde{\Psi}_{N}^{(z)}$.

6. Evolution of quantum state at the exceptional points

6.1. Deviation from purely exponential decay

We know that as friction increases, the oscillatory motion of a classical harmonic oscillator gradually vanishes, i.e., its underdamped motion goes into critical damping and then followed by overdamping [37, 38]. At critical damping, the oscillator relaxes to its equilibrium position asymptotically with a time dependence of *e*^{−γ*t*/2} and *te*^{−γ*t*/2}. The generalized eigenvectors exhibit a similar time dependence at the exceptional point. For example, this behaviour was also observed in Ref. [4] for second-order resonance pole in Friedrichs model in which *te*[−]γ*^t* was called a secular term. Similar behaviours are also obtained in collective spin models, such as in Refs. [46, 47].

The solutions to a non-Hermitian eigenvalue problem can be worked out using the method discussed in Ref. [48]. Let ϕ_N be an initial state consisting of the combination of generalized eigenvectors from the *N*-subspace with real coefficients $\phi_N = c_0 F^{(0)} + c_1 F^{(1)} + \cdots + c_N F^{(N)}$ In matrix form it is coefficients $\phi_N = c_0 F_N^{(0)} + c_1 F_N^{(1)} + \cdots + c_N F_N^{(N)}$. In matrix form, it is

$$
\boldsymbol{\phi}_N(0) = \begin{pmatrix} c_N \\ c_{N-1} \\ \vdots \\ c_1 \\ c_0 \end{pmatrix} . \tag{62}
$$

The solution to the matrix equation $\frac{\partial}{\partial t} \phi_N = -K_N \cdot \phi_N$ is [48]

$$
\phi_N(t) = e^{-\lambda_N t} \begin{pmatrix}\n1 & & & & & \\
-\lambda_N t & & & & & \\
\vdots & & \ddots & & & \\
\frac{(-\lambda_N t)^{N-2}}{(N-2)!} & \cdots & -\lambda_N t & 1 & 0 & 0 \\
\frac{(-\lambda_N t)^{N-1}}{(N-1)!} & \cdots & \frac{(\lambda_N t)^2}{2!} & -\lambda_N t & 1 & 0 \\
\frac{(-\lambda_N t)^N}{N!} & \cdots & -\frac{(\lambda_N t)^3}{3!} & \frac{(\lambda_N t)^2}{2!} & -\lambda_N t & 1\n\end{pmatrix} \cdot \phi_N(0).
$$
\n(63)

For instance, in the $N = 2$ subspace of the oscillator model we consider, where $\lambda_2 = \gamma$, a state that starts initially as $\phi_2(0) = c_2 F^{(0)} + c_1 F^{(1)} + c_2 F^{(2)}$ evolves into $\phi_2(0) = c_0 F_2^{(0)}$ $c_2^{(0)}$ + $c_1F_2^{(1)}$ $c_2^{(1)}$ + $c_2F_2^{(2)}$ $2^{(2)}$ evolves into

$$
\phi_2(t) = e^{-\gamma t} \left[\left(c_0 - \gamma t c_1 + \frac{\gamma^2 t^2}{2} c_2 \right) F_2^{(0)} + (c_1 - \gamma t c_2) F_2^{(1)} + c_2 F_2^{(2)} \right] \tag{64a}
$$

$$
= e^{-\gamma t} \phi_2(0) - \gamma t e^{-\gamma t} \left(c_1 F_2^{(0)} + c_2 F_2^{(1)} \right) + \frac{(\gamma t)^2}{2} e^{-\gamma t} c_2 F_2^{(0)}.
$$
 (64b)

We find that deviations from the exponential decay of the initial state reveal themselves through polynomial terms in *t n* .

6.2. Relaxation of an excited state under Caldeira–Leggett (CL) equation

In this section we follow the relaxation of the first excited state of a harmonic oscillator to the stationary state under the CL equation. The first excited state has the wave function $\phi_1(x) = (2/\sqrt{\pi})^{1/2} x \exp(-x^2/2)$ in dimensionless position coordinate. Its density function is $\Xi(x, \tilde{x}) = \phi_1^*(x)\phi_1(\tilde{x})$. It decomposes into the eigenfunction of the KL equation $f^{\pm}(Q, r)$ as (with $h = 1/2$) [49] equation $f_{mn}^{\pm}(Q, r)$ as (with $b = 1/2$) [49]

$$
\Xi(Q,r) = f_{00}(Q,r) - 2f_{10}(Q,r) = \frac{2}{\sqrt{\pi}} \left(Q^2 - \frac{r^2}{4}\right) e^{-Q^2 - r^2/4}.
$$
\n(65)

This state has a two-peak probability distribution function $\Xi(x, 0)$ along the diagonal (*r* = 0), cf. Fig. 2(a). It has two
troughs along the off-diagonal (*O* = 0), which is the coherence or interference component of t troughs along the off-diagonal $(Q = 0)$, which is the coherence or interference component of the state, cf. Fig. 2(e). We can think of it as a superposition between two stationary wave packets giving rise to interference effect. Because this state does not contain f_{mn}^{\pm} component with non-zero *n*, the wave packet does not execute oscillating motion around the equilibrium position while it relaxes to the stationary state. We consider three regions of damping below.

(a) Underdamped region, $\gamma = \omega_0$ (real ω).

Using the transformation S_2 , we obtain the transformed state of Ξ in the CL system. For simplicity, we take $\gamma = \omega_0$. Then substituting $\delta_2 = 1 - \gamma/(2\omega_0) = 1/2$ into \bar{f}_{10} (B.6) yields $S_2 f_{10}(Q, r) = \bar{f}_{10}(\bar{Q}, \bar{r}) = \bar{\Pi}_{10}\bar{f}_{00}$, where \bar{f}_{20} is the stationary state already given in Eq. (24), and the polynomial \bar{f}_{00} is the stationary state already given in Eq. (24), and the polynomial part is

$$
\bar{\Pi}_{10}(\bar{Q}, \bar{r}) = \frac{2}{3} \left(1 - 2\bar{Q}^2 + 2i\bar{Q}\bar{r} + 2\bar{r}^2 \right).
$$
 (66)

Though a similarity transformation preserves the trace and the hermiticity (or adjoint-symmetry) of the density function $\Xi(Q, r)$ [36], it does not guarantee the positivity of the transformed state $\bar{f}_{00} - 2\bar{f}_{10}$. We have to modify the coefficient of \bar{f}_{10} to −3/2 to ensure its positive definiteness,

$$
\bar{\Xi}_{U}(0) = \bar{f}_{00} - \frac{3}{2}\bar{f}_{10} = \left(1 - \frac{3}{2}\bar{\Pi}_{10}\right)\bar{f}_{00}.
$$
\n(67)

Notice that the trace or normalization is not altered by a change in the coefficient in front of the decaying state \bar{f}_{10} which has zero trace. The time evolution is (with $\lambda_{10} = \gamma = \omega_0$)

$$
\bar{\Xi}_U(t) = e^{-\bar{K}t}\bar{\Xi}(0) = \left(1 - \frac{3}{2}e^{-\omega_0 t}\bar{\Pi}_{10}\right)\bar{f}_{00}.
$$
\n(68)

Figure 2: Top and bottom row show the evolution of the probability distributions of $\bar{\Xi}_U(t)$ and $\bar{\Xi}_O(t)$ along the \bar{Q} -axis with $\bar{r} = 0$, and their coherence components along the F-axis with $\bar{Q} = 0$ respect coherence components along the \bar{r} -axis with $\bar{Q} = 0$, respectively, for three different damping at different time, (a) and (e) at $\omega_0 t = 0$, (b) and (f) at $\omega_0 t = 0.5$, (c) and (g) at $\omega_0 t = 1$, (d) and (h) at $\omega_0 t = 3$. Dashed, solid and dot-dashed lines denote underdamping ($\gamma = \omega_0$), critical damping ($\gamma = 2\omega_0$), which corresponds to the exceptional point, and overdamping ($\gamma = 3\omega_0$).

(b) Critically damped region, $\gamma = 2\omega_0$ (at the exceptional point $\omega = 0$).

When approaching the exceptional points through $\gamma \to 2\omega_0$, the dynamics is now determined by \bar{K}_E . Therefore, we need to write $\bar{\Xi}$ in the basis of the generalized eigenvectors $\bar{\Phi}_{N}^{(z)}$ listed in 5.2. We find that

$$
\bar{\Pi}_{10} = -\frac{4}{3} \left(\bar{\Phi}_2^{(0)} - \frac{1}{2} \bar{\Phi}_2^{(1)} + \frac{1}{2} \bar{\Phi}_2^{(2)} \right).
$$
 (69)

Consequently, we start with (cf. Eq. (67))

$$
\bar{\Xi}_E(0) = \bar{\Xi}_U(0) = \left[1 + 2\left(\bar{\Phi}_2^{(0)} - \frac{1}{2}\bar{\Phi}_2^{(1)} + \frac{1}{2}\bar{\Phi}_2^{(2)}\right)\right]\bar{f}_{00}
$$
\n(70)

as the initial state for time evolution at the exceptional point. Using Eq. (64a) and $\lambda_2 = \gamma = 2\omega_0$, the evolution of $\bar{\Xi}_E$ is

$$
\bar{\Xi}_E(t) = e^{-\bar{K}_E t} \bar{\Xi}_E(0) = \left(1 + 2e^{-2\omega_0 t} \left[(1 + \omega_0 t + \omega_0^2 t^2) \bar{\Phi}_2^{(0)} - \frac{1}{2} (1 + 2\omega_0 t) \bar{\Phi}_2^{(1)} + \frac{1}{2} \bar{\Phi}_2^{(2)} \right] \right) \bar{f}_{00} \,. \tag{71}
$$

(c) Overdamped region, $\gamma = 3\omega_0$ (imaginary ω).

When the relaxation rate is greater than $2\omega_0$, the modified frequency of the oscillator becomes imaginary, $\omega = i\sqrt{5}\omega_0/2$. The oscillator now experiences overdamning. Substituting $\delta_0 = -1/2$ into the eigenfunctions $i\sqrt{5\omega_0/2}$. The oscillator now experiences overdamping. Substituting $\delta_2 = -1/2$ into the eigenfunctions in
Fas (B.6)–(B.7), we find that the decomposition of the initial condition involves also eigenfunctions from th Eqs. (B.6)–(B.7), we find that the decomposition of the initial condition involves also eigenfunctions from the $N = 2$ subspace. We will label the eigenfunctions of \bar{K}_{CL} in the overdamped region by $\bar{\Omega}_{mn}^{\pm} \bar{f}_{00}$. $\bar{\Pi}_{10}(\bar{Q}, \bar{r})$ can then be decomposed into them as be decomposed into them as √

$$
\bar{\Pi}_{10} = \frac{1}{3}\bar{\Omega}_{10} + \frac{2\sqrt{2}}{3}i\left(\bar{\Omega}_{22}^{+} - \bar{\Omega}_{22}^{-}\right),\tag{72}
$$

where

$$
\bar{\Omega}_{10} = -\frac{2}{5}(1 - 2\bar{Q}^2 + 6i\bar{Q}\bar{r} + 2\bar{r}^2),\tag{73a}
$$

$$
\bar{\Omega}_{22}^{\pm} = \pm \frac{i}{5\sqrt{2}} \left(3 - 6\bar{Q}^2 + 8i\bar{Q}\bar{r} + 6\bar{r}^2 \pm \sqrt{5}(1 - 2\bar{Q}^2 - 2\bar{r}^2) \right). \tag{73b}
$$

The expressions of $\bar{\Omega}_{mn}^{\pm}$ can be obtained by setting $\delta_2 = -1/2$ in Eqs. (B.6) and (B.7). The eigenvalues are $\lambda_{10} = 3\omega_0$ and $\lambda_{22}^{\pm} = (3 \mp \sqrt{5})\omega_0$. Using the initial state

$$
\bar{\Xi}_O(0) = \bar{\Xi}(0) = \left(1 - \frac{1}{2}\bar{\Omega}_{10} - \sqrt{2}i\left(\bar{\Omega}_{22}^+ - \bar{\Omega}_{22}^-\right)\right)\bar{f}_{00},\tag{74}
$$

the state evolves into

$$
\bar{\Xi}_O(t) = \left(1 - e^{-3\omega_0 t} \left[\frac{1}{2}\bar{\Omega}_{10} + \sqrt{2}i \left(e^{\sqrt{5}\omega_0 t} \bar{\Omega}_{22}^+ - e^{-\sqrt{5}\omega_0 t} \bar{\Omega}_{22}^-\right)\right]\right) \bar{f}_{00} \,. \tag{75}
$$

In Fig. 2, the evolution of $\bar{\Xi}_U(t)$, $\bar{\Xi}_E(t)$ and $\bar{\Xi}_O(t)$ are plotted on the same graph at different instants. They are labelled by dashed lines, solid lines and dot-dashed lines, respectively. It was reported in dissipative collective spin models [46, 47] that there is a slow down of relaxation at the exceptional point. In the continuous variable model we consider, we find that there is indeed a slow down in the relaxation of the probability distribution function $\bar{\Xi}(Q, 0)$ when it passes from the underdamping region across the critical damping (the exceptional point) region into the overdamping region, as can be observed from the plots on the first row of Fig. 2. The polynomial terms in *t* (71) that was introduced at the exceptional point tend to slow down the relaxation at the initial stage of the evolution.

However, the relaxation of the coherence component $\bar{\Xi}(0, r)$ exhibits an opposite behaviour. From the plots on the second row of Fig. 2, we observe that at the initial stage, the decoherence is slowest in the underdamped oscillator. This is because the coherence component selects the off-diagonal components $(Q = 0)$ from the density function. For example, the \bar{Q}^2 and \bar{r}^2 terms in $\bar{\Xi}_U(\bar{Q}, \bar{r})$ and $\bar{\Xi}_O(\bar{Q}, \bar{r})$ have opposite signs. This results in different kinds of behavior in their relaxation in their relaxation.

There is also contribution from the exponential decay $exp(-\gamma t)$ that tends to speed up the relaxation when the damping rate γ increases from underdamping, across critical damping into overdamping region. In the final stage of the evolution when all the fast decaying components become negligible, we are left with the slowest decaying component $\exp(-(3 - \sqrt{5})\omega_0 t)\overline{\Omega}_{22}^+ \approx \exp(-0.76\omega_0 t)\overline{\Omega}_{22}^+$ from the overdamped region (75). That is why in the last
figure on each row of Fig. 2 we see that the overdamped oscillator is left behind by the underdamped figure on each row of Fig. 2 we see that the overdamped oscillator is left behind by the underdamped and critically damped oscillator which have almost arrived at their stationary state.

In the case of the modified KL equation, because the stationary state (33) depends on the parameter h_1 which decides the transition to the exceptional point, the evolution of a state at and away from the exceptional point cannot be compared as easily as in the case of CL equation. Therefore, we do not discuss it here. Nonetheless, when purely Gaussian states are considered, it is possible to obtain the exact time evolution of the generic models using the results from Ref. [50]. We will present the results elsewhere.

7. Conclusion

In this work we obtain the spectrum of the quantum Markovian Liouvillian for an oscillator in a generic environment at the exceptional points and clarify the structure of its eigenspace. The exceptional points occur when the modified frequency of the oscillator vanishes, which corresponds to critical damping of the oscillator. For a given natural frequency, the exceptional points lie on a circle in a two-parameter space. All orders of degeneracy up to infinity occur simultaneously. Consequently, the eigenspace also breaks into an infinite number of subspaces, each one has its own Jordan block structure. The degeneracy allows freedom in the choice of generalized eigenvectors. The freedom reveals itself as the invariance in the Jordan block structure of the Liouvillian under a similarity transformation whose form is obtained. In the Kossakowski–Lindblad equation whose natural frequency is not modified, the eigenfunctions do not coalesce so that it remains a case of ordinary degeneracy.

Acknowledgments

This work is supported by the Ministry of Higher Education Malaysia (MOHE) under the Fundamental Research Grant Scheme (FRGS), Grant No. FRGS/1/2020/STG07/UNIM/02/01.

Appendix A. Expressions of functions and operators in position coordinates

The explicit form of the polynomial $\Pi^{\pm}_{mn}(Q, r)$ in Eq. (9) is [49, 33]

$$
\Pi_{mn}^{\pm}(Q,r) = \sum_{\mu=0}^{m-n} \sum_{\nu=0}^{\mu} \sum_{\sigma=0}^{n} c_{mn}^{\pm \mu \nu \sigma} \left(\frac{Q}{\sqrt{2b}} \right)^{2(\mu-\nu)+n-\sigma} H_{2\nu+\sigma} \left(\sqrt{\frac{b}{2}} r \right), \tag{A.1}
$$

in which $H_{2v+\sigma}$ denotes the Hermite polynomials. The coefficients are

$$
c_{mn}^{\pm\mu\nu\sigma} = (\pm 1)^{n+\sigma} \frac{(-1)^{\mu+\nu}}{i^{n}2^{2\nu+\sigma}\mu!} \sqrt{\frac{(m-n)!}{m!}} {m \choose n+\mu} {\mu \choose \nu} {n \choose \sigma}.
$$
 (A.2)

For example,

$$
\Pi_{11}^{\pm}(Q,r) = \mp i \left(\frac{Q}{\sqrt{2b}} \pm \sqrt{\frac{b}{2}} r \right),\tag{A.3}
$$

$$
\Pi_{10}(Q,r) = \frac{1}{2} - \frac{Q^2}{2b} + \frac{b}{2}r^2,
$$
\n(A.4)

$$
\Pi_{22}^{\pm}(Q,r) = \frac{1}{\sqrt{2}} \left(\frac{1}{2} - \frac{Q^2}{2b} \mp Qr - \frac{b}{2}r^2 \right). \tag{A.5}
$$

In terms of the annihilation and creation operators of harmonic oscillator as well as in the position coordinates, the seven operators in Eq. (16) are

$$
iL_0 \rho = i \left[a^\dagger a, \rho \right] \doteq \frac{i}{2} \left(-\frac{\partial^2}{\partial Q \partial r} + Qr \right) \rho, \tag{A.6}
$$

$$
iM_1 \rho = \frac{i}{4} ([a^{\dagger} a^{\dagger}, \rho] + [aa, \rho]) \doteq \frac{i}{2} \left(\frac{\partial^2}{\partial Q \partial r} + Qr \right) \rho, \tag{A.7}
$$

$$
iM_2 \rho = \frac{1}{4} ([a^{\dagger} a^{\dagger}, \rho] - [aa, \rho]) \doteq -\frac{1}{2} \left(\frac{\partial}{\partial Q} Q + r \frac{\partial}{\partial r} \right),
$$
 (A.8)

$$
(O_0 - I/2)\rho = \frac{1}{4} ([a^{\dagger}, \{a, \rho\}] - [a, \{a^{\dagger}, \rho\}]) \doteq -\frac{1}{2} \left(\frac{\partial}{\partial Q} Q - r \frac{\partial}{\partial r}\right) \rho, \tag{A.9}
$$

$$
O_{+}\rho = -\frac{1}{2}[a,[a^{\dagger},\rho]] \doteq \frac{1}{4}\left(\frac{\partial^2}{\partial Q^2} - r^2\right)\rho,
$$
\n(A.10)

$$
L_{1+}\rho = -\frac{1}{4} ([a^{\dagger}, [a^{\dagger}, \rho]] + [a, [a, \rho]]) \doteq -\frac{1}{4} \left(\frac{\partial^2}{\partial Q^2} + r^2 \right) \rho, \tag{A.11}
$$

$$
L_{2+}\rho = \frac{i}{4} \left([a^{\dagger}, [a^{\dagger}, \rho]] - [a, [a, \rho]] \right) \doteq -\frac{i}{2} r \frac{\partial}{\partial Q} \rho, \tag{A.12}
$$

where we use $\dot{=}$ to denote the coordinate representation of an operator.

Appendix B. Dominant term under similarity transformation

Appendix B.1. Caldeira–Leggett (CL) equation

The Liouvillian of the CL equation is related to that of the KL equation by a similarity transformation S_2 (23) through $K_{CL} = S_2 K_{KL} S_2^{-1}$ [36, 25]. The parameters in S_2 are $\eta = -2b \tanh \phi$ and $\sinh \phi = -\gamma/(2\omega_2)$. The coefficients of the two Liouvillians are related by $\omega_0 = \omega_0 \cosh \phi$ and $\bar{b} = b/\cosh \phi$ of the two Liouvillians are related by $\omega_0 = \omega_2 \cosh \phi$ and $\bar{b} = b/\cosh \phi$.

Applying a similarity transformation S_2 (23) to the coordinates yields

$$
\hat{Q} = S_2 \frac{Q}{\sqrt{2b}} S_2^{-1} = \frac{1}{\sqrt{\cosh \phi}} \left[\cosh(\phi/2) \bar{Q} + \sinh(\phi/2) \frac{i}{2} \frac{\partial}{\partial \bar{r}} \right] + \frac{\sinh \phi}{\sqrt{\cosh \phi}} \left[\cosh(\phi/2) i\bar{r} - \sinh(\phi/2) \frac{1}{2} \frac{\partial}{\partial \bar{Q}} \right], \quad (B.1)
$$

$$
i\hat{r} = S_2 \sqrt{\frac{b}{2}} i r S_2^{-1} = \sqrt{\cosh \phi} \left[\cosh(\phi/2) i\bar{r} - \sinh(\phi/2) \frac{1}{2} \frac{\partial}{\partial \bar{Q}} \right], \quad (B.2)
$$

where

$$
\cosh \phi = \frac{\omega_0}{\omega_2}, \qquad \cosh(\phi/2) = \frac{1}{2\sqrt{\omega_2}} \left(\sqrt{\omega_0 - \gamma/2} + \sqrt{\omega_0 + \gamma/2} \right), \tag{B.3}
$$

$$
\sinh \phi = -\frac{\gamma}{2\omega_2}, \quad \sinh(\phi/2) = \frac{1}{2\sqrt{\omega_2}} \left(\sqrt{\omega_0 - \gamma/2} - \sqrt{\omega_0 + \gamma/2} \right). \tag{B.4}
$$

Using the stationary state \bar{f}_{00} (24) of \bar{K}_{CL} , we obtain

$$
\bar{f}_{11}^{\pm} = S_2 f_{11}^{\pm} = \mp i(\hat{Q} \pm \hat{r}) \bar{f}_{00} = \frac{1 \pm i}{2 \sqrt{\delta_2}} (\bar{Q} - i\bar{r}) \bar{f}_{00} - \frac{1 \pm i}{2 \sqrt{2 - \delta_2}} (\bar{Q} + i\bar{r}) \bar{f}_{00},
$$
\n(B.5)

$$
\bar{f}_{10} = S_2 f_{10} = \left(\frac{1}{2} - \hat{Q}^2 + \hat{r}^2\right) \bar{f}_{00} = \frac{1 - 2\bar{Q}^2 + 4i\bar{Q}\bar{r}(1 - \delta_2) + 2\bar{r}^2}{2\delta_2(2 - \delta_2)} \bar{f}_{00},
$$
\n(B.6)

$$
\bar{f}_{22}^{\pm} = S_2 f_{22}^{\pm} = \frac{1}{2 \sqrt{2}} \left(1 - 2 \hat{Q}^2 \mp 4 \hat{Q} \hat{r} - 2 \hat{r}^2 \right) \bar{f}_{00}
$$
\n
$$
= \frac{1}{2 \sqrt{2} \delta_2 (2 - \delta_2)} \left(\pm i \left[(1 - \delta_2)(1 - 2 \hat{Q}^2 + 2 \hat{r}^2) + 4i \bar{Q} \bar{r} \right] + \sqrt{\delta_2 (2 - \delta_2)} (1 - 2 \bar{Q}^2 - 2 \bar{r}^2) \right) \bar{f}_{00},
$$
\n(B.7)

where $\delta_2 \equiv 1 - \gamma/(2\omega_0)$ (26). Note that $\bar{\Pi}^{\pm}_{mn}$ can be extracted from $\bar{f}^{\pm}_{mn} = \bar{\Pi}^{\pm}_{mn} \bar{f}_{00}$. We further note that \bar{f}_{11}^{\pm} belong to the $N - 1$ subspace, whereas \bar{f}_{22} and \bar{f}^{\pm}_{21} ar the *N* = 1 subspace, whereas \bar{f}_{10} and \bar{f}_{22}^{\pm} are in the *N* = 2 subspace.

Appendix B.2. Modified Kossakowski–Lindblad (KL) equation

The modified KL equation can be obtained from the KL equation via a similarity transformation $S_1(32)$, \tilde{K}_{mKL} = $S_1 K_{\text{KL}} S_1^{-1}$ [36, 25]. The parameters in S_1 are given by $\tanh \psi = -h_1/(2\omega_0)$, $\eta_0 = -\eta_2 \gamma/h_1$ and $\eta_2 = -2\gamma bh_1/(4\omega_0 \omega_1)$, where ω_1 is the modified frequency (31). The coefficient in the equation is relate where ω_1 is the modified frequency (31). The coefficient in the equation is related by $\bar{b} = b(\omega_1^2 + \gamma^2/4)/(\omega_0\omega_1)$.
Applying a similarity transformation S₁ (32) to the coordinates produces the following results Applying a similarity transformation S_1 (32) to the coordinates produces the following results,

$$
\hat{Q} = S_1 \frac{Q}{\sqrt{2b}} S_1^{-1} = \frac{1}{\omega_1} \sqrt{\frac{\omega_0 + h_1/2}{\omega_0(\omega_1^2 + \gamma^2/4)}} \left[\left(\omega_1^2 + \frac{\gamma^2}{4} \right) \tilde{Q} + \frac{\gamma^2}{4} \frac{1}{2} \frac{\partial}{\partial \tilde{Q}} + \frac{\gamma}{2} \frac{h_1}{2} i \tilde{r} \right],
$$
\n(B.8)

$$
i\hat{\vec{r}} = S_1 \sqrt{\frac{b}{2}} i r S_1^{-1} = \sqrt{\frac{\omega_0 (\omega_0 + h_1/2)}{\omega_1^2 + \gamma^2/4}} i \tilde{r}.
$$
 (B.9)

Making use of the stationary state \tilde{f}_{00} (33) of \tilde{K}_{mKL} , we obtain

$$
\tilde{f}_{11}^{\pm} = S_1 f_{11}^{\pm} = \mp i(\hat{Q} \pm \hat{r}) \tilde{f}_{00}
$$
\n
$$
= \frac{-\tilde{\omega}}{\sqrt{1 + \delta_1 (2 - \delta_1)\tilde{\omega}^2}} \left[\sqrt{2 - \delta_1} i \tilde{r} \pm i \frac{\sqrt{\delta_1}}{1 + \delta_1 \tilde{\omega}^2} \left(\tilde{Q} + i \tilde{\omega} \tilde{r} + \delta_1 \left[(2 - \delta_1) \tilde{\omega}^2 \tilde{Q} - i \tilde{\omega} \tilde{r} \right] \right) \right] \tilde{f}_{00}, \qquad (B.10)
$$
\n
$$
\tilde{f}_{10} = S_1 f_{10} = \left(\frac{1}{2} - \hat{Q}^2 + \hat{r}^2 \right) \tilde{f}_{00}
$$
\n
$$
= \frac{\tilde{\omega}^2}{2(1 + \delta_1 \tilde{\omega}^2)^2} \left(4\tilde{r}^2 + \delta_1 \left[1 - 2\tilde{Q}^2 - 4\tilde{\omega} i \tilde{Q} \tilde{r} - 2(1 - \tilde{\omega}^2) \tilde{r}^2 \right] + \delta_1^2 \left[\tilde{\omega}^2 (1 - 4\tilde{Q}^2) + 4\tilde{\omega} i \tilde{Q} \tilde{r} \right] + 2\delta_1^3 \tilde{\omega}^2 \tilde{Q}^2 \right) \tilde{f}_{00}, \qquad (B.11)
$$

$$
\tilde{f}_{22}^{\pm} = S_1 f_{22}^{\pm} = \frac{1}{2 \sqrt{2}} \left(1 - 2 \hat{Q}^2 \mp 4 \hat{Q} \hat{r} - 2 \hat{r}^2 \right) \tilde{f}_{00}
$$
\n
$$
= \frac{1}{2 \sqrt{2}} \left[\frac{-2(2 - \delta_1) \tilde{\omega}^2 \tilde{r}^2}{1 + \delta_1 (2 - \delta_1) \tilde{\omega}^2} + \frac{\delta_1 \tilde{\omega}^2}{1 + \delta_1 \tilde{\omega}^2} - \frac{2 \delta_1 \tilde{\omega}^2}{(1 + \delta_1 \tilde{\omega}^2)^2} \left(\left[1 + \delta_1 (2 - \delta_1) \right] \tilde{Q}^2 + 2 \tilde{\omega} (1 - \delta_1) i \tilde{Q} \tilde{r} - \frac{\tilde{\omega}^2 (1 - \delta_1)^2 \tilde{r}^2}{1 + \delta_1 (2 - \delta_1) \tilde{\omega}^2} \right) \right]
$$
\n
$$
\pm i \frac{\sqrt{\delta_1 (2 - \delta_1)} \tilde{\omega}^2}{1 + \delta_1 \tilde{\omega}^2} \left(4i \tilde{Q} \tilde{r} - \frac{4(1 - \delta_1) \tilde{\omega} \tilde{r}^2}{1 + \delta_1 (2 - \delta_1) \tilde{\omega}^2} \right) \Big] \tilde{f}_{00} \,, \tag{B.12}
$$

where $\tilde{\omega} \equiv 2\omega_0/\gamma$ (35) and $\delta_1 \equiv 1 - h_1/(2\omega_0)$ (36).

Appendix C. Generalized eigenvectors of Caldeira–Leggett (CL) equation

In this appendix, we will work out the first few series of the generalized eigenvectors of \bar{K}_E (40) for the CL equation using Eqs. (39a)–(39b), and the Jordan block structure of the reduced dynamics Eqs. (37a)–(37b) at the exceptional points.

Appendix C.1. First few series of generalized eigenvectors

Starting from Eq. (39a) with $\bar{\Phi}_0^{(0)}$ $_{0}^{(0)} = 1$, in the $N = 1$ subspace we have $\bar{\Phi}_{1}^{(0)}$ $u_1^{(0)} = u_1^{(0)} \bar{Q} + v_1^{(0)}$ $i_1^{(0)}$ *ir*. From the discussion in 5.2, once we choose the overall constant of the lowest order generalized eigenvector $F_N^{(0)}$, the overall constants of other vectors are also fixed. We fix the overall constant by choosing $u_1^{(0)} = 1$. Substituting $\bar{\Phi}_1^{(0)}$ into Eq. (39a), we γ_1 – 1. Substituting \mathcal{L}_1 solve the equation to get $v_1^{(0)}$ $\bar{\Phi}^{(0)}_1 = -1$. Therefore, $\bar{\Phi}^{(0)}_1$ $1_{1}^{(0)} = \overline{Q} - i\overline{r}$. We continue by using Eq. (39b) *z* = *N* = 1. Substituting $\bar{\Phi}_1^{(1)}$ $u_1^{(1)} = u_1^{(1)}\bar{Q} + v_1^{(1)}$ $i_1^{(1)}$ *ir* into Eq. (39b) yield $p_1^{(1)}$ $q_1^{(1)} + q_1^{(1)}$ $1_1^{(1)} = -1$. We can now make different choices of $p_1^{(1)}$ $q_1^{(1)}$ and $q_1^{(1)}$ $1^{(1)}$. This is the freedom stemming from the discussion in 5.2. For example, we could choose

$$
p_1^{(1)} = 0, \qquad q_1^{(1)} = -1, \qquad \bar{\Phi}_1^{(1)} = -i\bar{r}, \tag{C.1a}
$$

$$
p_1^{(1)} = -1 \,, \qquad q_1^{(1)} = 0 \,, \qquad \bar{\Psi}_1^{(1)} = -\bar{Q} \,, \tag{C.1b}
$$

or many others. All this choices produce valid series of generalized eigenvectors.

In the $N = 2$ subspace, we continue with $\bar{\Phi}_{2}^{(0)}$ $v_2^{(0)} = (u_2^{(0)}\bar{Q} + v_2^{(0)}$ $\bar q_2^{(0)}$ i $\bar r) \bar \Phi_1^{(0)}$ $y_1^{(0)} + w_2^{(0)} \bar{\Phi}_0^{(0)}$ $_{0}^{(0)}$. Substituting it into Eq. (39a) we get the condition $u_2^{(0)}$ $v_2^{(0)} + v_2^{(0)}$ $y_2^{(0)} = 0$ and $w_2^{(0)}$ $v_2^{(0)} = (v_2^{(0)}$ ⁽⁰⁾ − 1)/4. We fix the overall constant by choosing $u_2^{(0)}$
(\overline{Q} = ¹²)² + 1/2. We applies with the shairs Fr. (C₁)² $2^{(0)} = 1$. Then $v_2^{(0)}$ $y_2^{(0)} = -1$ and $w_2^{(0)}$ $\bar{\Phi}_2^{(0)} = -1/2$, leading to $\bar{\Phi}_2^{(0)}$ $v_2^{(0)} = -1$ and $w_2^{(0)} = -1/2$, leading to $\bar{\Phi}_2^{(0)} = (\bar{Q} - i\bar{r})^2 - 1/2$. We continue with the choice Eq. (C.1a). The next vector in this series is $\bar{\Phi}_2^{(1)} = (u_2^{(1)}\bar{Q} + v_2^{(1)}i\bar{r})\bar{\Phi}_1^{(1)}$. Upon substituting $y_2^{(1)} = (u_2^{(1)}\overline{Q} + v_2^{(1)})$ $_{2}^{(1)}$ i $\bar{r})\bar{\Phi}_{1}^{(1)}$ $1^{(1)}$. Upon substituting this choice of $\bar{\Phi}_2^{(1)}$ $2^{(1)}$ into Eq. (39a) and simplify the expression, we find that $u_2^{(1)}$ $v_2^{(1)} = 2, v_2^{(1)}$ $_2^{(1)} = -2$. This means $\bar{\Phi}_2^{(1)}$ $\Omega_2^{(1)} = -2i\bar{r}(\bar{Q} - i\bar{r})$. The last vector in this order is $\bar{\Phi}_{2}^{(2)}$ $Q_2^{(2)} = (p_2^{(2)}\bar{Q} + q_2^{(2)})$ $^{(2)}_{2}$ *ir*) $\bar{\Phi}^{(1)}_{1}$ $s_1^{(1)} + s_2^{(2)} \bar{\Phi}_0^{(0)}$ ⁽⁰⁾. Applying Eq. (37b) onto it, we obtain $p_2^{(2)}$ $q_2^{(2)} = 0, q_2^{(2)}$ $s_2^{(2)} = -2, s_2^{(2)}$ $p_2^{(2)} = -p_2^{(2)}$ $\Phi_2^{(2)} = (p_2^{(2)}\bar{Q} + q_2^{(2)}i\bar{r})\Phi_1^{(1)} + s_2^{(2)}\Phi_0^{(0)}$. Applying Eq. (37b) onto it, we obtain $p_2^{(2)} = 0$, $q_2^{(2)} = -2$, $s_2^{(2)} = -p_2^{(2)}/4 = 0$. We then conclude that $\Phi_2^{(2)} = 2(i\bar{r})^2$. $2^{(2)} = 2(i\bar{r})^2$.

We continue in this way for $N = 3$ and summarize the results in the 4.1. We can continue in this manner to generate all the high order generalized eigenvectors. The results can be generalized to Eqs. (41a)–(41b).

Other set of generalized eigenvectors can be generated using the freedom discussed in 5.2. For example, if we instead continue with Eq. (C.1b), we will get the series of generalized eigenvectors of \bar{K}_E summarized in 5.2. However, for this series of generalized eigenvectors, additional terms, $x_N^{(z)} \overline{\Phi}_N^{(z)}$ and $x_N^{(z)} \overline{\Phi}_N^{(z)}$, should be added to Eqs. (39a) and (39b), respectively.

Appendix C.2. Proof of the equality of diagonal and parallel representation of $\bar{\Phi}_{N}^{(z)}$

In this section, we prove by induction that the diagonal representation (41a) and parallel representation (41b) of $\bar{\Phi}_{N}^{(z)}$ are indeed identical in the overlapping region, i.e., when $z = 1, 2, ..., N - 1$.
First we check that Eq. (41a) and (41b) are indeed identical for $N = 2$ and 3

First, we check that Eq. (41a) and (41b) are indeed identical for $N = 2$ and 3. Then assuming that the right hand side (RHS) of Eqs. (??) are equal for $\bar{\Phi}_N^{(z)}$ in the subspace $N = m - 1$ and *m*, we will show that they are also equal in the subspace $N = m + 1$.

We start by simplifying the expression obtained from the RHS of the diagonal representation (41a) for $N = m + 1$, and show that it equals Eq. (41b) for $N = m + 1$. The RHS of Eq. (41a) for $N = m + 1$ is

$$
(\bar{Q} - i\bar{r}) \frac{\bar{\Phi}_{m}^{(z)}}{m^{z}} - \frac{m - z}{2} \frac{\bar{\Phi}_{m-1}^{(z)}}{(m-1)^{z}} = (\bar{Q} - i\bar{r}) \frac{-i\bar{r}}{z} \frac{\bar{\Phi}_{m-1}^{(z-1)}}{(m-1)^{z-1}} - \frac{m - z}{2} \frac{-i\bar{r}}{z} \frac{\bar{\Phi}_{m-2}^{(z-1)}}{(m-2)^{z-1}}
$$

$$
= \frac{-i\bar{r}}{z} \left((\bar{Q} - i\bar{r}) \frac{\bar{\Phi}_{m-1}^{(z-1)}}{(m-1)^{z-1}} - \frac{(m-1) - (z-1)}{2} \frac{\bar{\Phi}_{m-2}^{(z-1)}}{(m-2)^{z-1}} \right)
$$

$$
= \frac{-i\bar{r}}{z} \frac{\bar{\Phi}_{m}^{(z-1)}}{m^{z-1}}, \qquad z = 1, 2, ..., m,
$$
 (C.2)

which is the RHS of Eq. (41b) for $N = m + 1$. In the first equality, we have made used of Eq. (41b) twice to replace $\bar{\Phi}_m^{(z)}$ and $\bar{\Phi}_{m-1}^{(z)}$, which are assumed to be correct and are valid for $z \neq 0$. To obtain the last equality, we use Eq. (41a) Φ_m and Φ_{m-1} , which are assumed to be correct and are vand for $z \neq 0$. To obtain the last equality, we use Eq. (+1a) for *N* = *m* and for *z* = 1 to replace the expression in the bracket by $\bar{\Phi}_m^{(z-1)}/m^{z-1}$ since by definition we set $\bar{\Phi}_{m-1}^{(m)} = 0$. Therefore, Eqs. (41a) and (41b) are equal in the overlapping region.

A ppendix $C.3$. Proof that $\bar{\Phi}_{m}^{(z)}$ are generalized eigenvectors of \bar{K}_{E}

We will show that $\bar{\Phi}_N^{(z)}$ in Eq. (38) satisfies Eqs. (37a) and (37b). We start with $z = 0$. We know that $\bar{\Phi}_N^{(0)}$ for $N = 0, 1, 2, 3$ satisfy Eqs. (37a)–(37b) by construction. We prove the rest by induction. We first assume that Eq. (37a)–(37b) are correct for arbitrary $N = m$, then we deduce that they are correct for $N = m + 1$ too. First, using Eq. (41a), we have

$$
\bar{F}_N^{(0)} = \bar{\Phi}_N^{(0)} \bar{F}_0^{(0)} = (\bar{Q} - i\bar{r}) F_{N-1}^{(0)} - \frac{N-1}{2} \bar{F}_{N-2}^{(0)}.
$$
\n(C.3)

Then, acting \bar{K}_E (40) on it, we obtain

$$
\bar{K}_{E}\bar{F}_{m}^{(0)} = (\bar{Q} - i\bar{r})\bar{K}_{E}\bar{F}_{m-1}^{(0)} - \frac{m-1}{2}\bar{K}_{E}\bar{F}_{m-2}^{(0)} - \frac{\gamma}{4} \left(\frac{\partial \bar{F}_{m-1}^{(0)}}{\partial \bar{Q}} + i \frac{\partial \bar{F}_{m-1}^{(0)}}{\partial \bar{r}} + 4i\bar{r}F_{m-1}^{(0)} \right)
$$
(C.4)

Using $\bar{K}_E \bar{F}_{m-1}^{(0)} = \lambda_{m-1} \bar{F}_{m-1}^{(0)}$, $\lambda_{m-1} = (m-1)\gamma/2$ and a similar expression for $N = m-2$ on the RHS of Eq. (C.4), we put the RHS of Eq. (C.4), we put the RHS of Eq. $(C.4)$ in the following form,

$$
\bar{K}_{E}\bar{F}_{m}^{(0)} = \lambda_{m} \left((\bar{Q} - i\bar{r}) F_{m-1}^{(0)} - \frac{m-1}{2} \bar{F}_{m-2}^{(0)} \right) - \frac{\gamma}{2} \left(\frac{1}{2} \frac{\partial \bar{F}_{m-1}^{(0)}}{\partial \bar{Q}} + \frac{i}{2} \frac{\partial \bar{F}_{m-1}^{(0)}}{\partial \bar{r}} + \bar{F}_{m}^{(0)} + 2i\bar{r} F_{m-1}^{(0)} - \frac{m-1}{2} \bar{F}_{m-2}^{(0)} \right). \tag{C.5}
$$

Notice that the first bracket equals $F_m^{(0)}$. Therefore, we are left to show that the rest of the terms in the second bracket vanishes. That is, defining

$$
D_N(\bar{Q}, \bar{r}) \equiv \frac{1}{2} \frac{\partial \bar{F}_{N-1}^{(0)}}{\partial \bar{Q}} + \frac{i}{2} \frac{\partial \bar{F}_{N-1}^{(0)}}{\partial \bar{r}}, \qquad (C.6)
$$

we want to show that

$$
D_N(\bar{Q}, \bar{r}) = -\bar{F}_N^{(0)} - 2i\bar{r}\bar{F}_{N-1}^{(0)} + \frac{N-1}{2}\bar{F}_{N-2}^{(0)}, \qquad N \ge 2.
$$
 (C.7)

Let us prove Eq. (C.7) by induction, i.e., we show that given Eq. (C.7) is true for D_m and D_{m-1} , then it is also true for D_{m+1} .

We first verify that Eq. $(C.7)$ is indeed true for $N = 2, 3$ by direct calculation. Then we substitute the expression of $\bar{F}_m^{(0)}$ (C.3) into the RHS of Eq. (C.6) for $N = m + 1$. After carrying out differentiation, we simplify it to

$$
D_{m+1}(\bar{Q}, \bar{r}) = \frac{1}{2} \frac{\partial \bar{F}_m^{(0)}}{\partial \bar{Q}} + \frac{i}{2} \frac{\partial \bar{F}_m^{(0)}}{\partial \bar{r}} = F_{m-1}^{(0)} + (\bar{Q} - i\bar{r}) D_m(\bar{Q}, \bar{r}) - \frac{m-1}{2} D_{m-1}(\bar{Q}, \bar{r}).
$$
 (C.8)

Assuming that Eq. (C.7) is true for $N = m - 1$ and m , we substitute them into D_m and D_{m-1} in Eq. (C.8). After simplifying the expression we get

$$
D_{m+1}(\bar{Q}, \bar{r}) = -\bar{F}_{m+1}^{(0)} - 2i\bar{r}\bar{F}_m^{(0)} + \frac{m}{2}\bar{F}_{m-1}^{(0)},
$$
\n(C.9)

which is Eq. (C.7) for $N = m + 1$. This proves that Eq. (41a) indeed satisfies Eq. (37a).

We then prove that Eq. (41b), or $\bar{\Phi}_N^{(z)}$ for $z \neq 0$, satisfy Eq. (37b) by induction. We know that Eq. (41b) satisfy Eq. (37b) for $N = 1, 2$ by construction. Then we assume that Eq. (37b) is true for $N = m - 1$, and show that it is true
for $N = m$ too. We start with Eq. (41b) by multiplying it with $\bar{F}_0^{(0)}$ to get $\int_0^{(0)}$ to get

$$
\frac{F_m^{(z)}}{m^z} = -\frac{i\bar{r}}{z} \frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}.
$$
\n(C.10)

Then we act \bar{K}_E (40) on it to get

$$
\bar{K}_{E}\left(\frac{\bar{F}_{m}^{(z)}}{m^{z}}\right) = \frac{-i\bar{r}}{z}\bar{K}_{E}\left(\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}\right) - \gamma \frac{i\bar{r}}{z}\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}} - \frac{\gamma}{4z}\frac{\partial}{\partial\bar{Q}}\left(\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}\right). \tag{C.11}
$$

Let us calculate the first term on the RHS of Eq. (C.11) using the of the generalized eigenvalue equation (37b), and then use Eq. (41b) to substitute $\bar{F}_{m-1}^{(z-1)}$ $\bar{F}_{m-1}^{(z-1)}$ and $\bar{F}_{m-1}^{(z-2)}$ $\sum_{m=1}^{(z-2)}$ in the expression. The result is

$$
\frac{-i\bar{r}}{z}\bar{K}_{E}\left(\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}\right) = \lambda_{m-1}\frac{-i\bar{r}}{z}\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}} + \lambda_{m-1}\frac{-i\bar{r}}{z}\frac{\bar{F}_{m-1}^{(z-2)}}{(m-1)^{z-2}} = \lambda_{m}\left(\frac{\bar{F}_{m}^{(z)}}{m^{z}} + \frac{\bar{F}_{m}^{(z-1)}}{m^{z}}\right) - \frac{\gamma}{2}\left(\frac{\bar{F}_{m}^{(z)}}{m^{z}} + \frac{1}{z}\frac{\bar{F}_{m}^{(z-1)}}{m^{z-1}}\right), (C.12)
$$

where we use Eq. (C.10) and $\lambda_{m-1} = \lambda_m - \gamma/2$ to get the last equality. Notice that the two terms in the first bracket on the RHS of the last equality equal $\bar{K}_E(\bar{F}_{m}^{(z)}/(\lambda_N m^z))$, i.e., the left hand side (LHS) of Eq. (C.11). Substituting Eqs. (C.12) back into Eq. (C.11), we are left to show that Eqs. (C.12) back into Eq. (C.11), we are left to show that

$$
\frac{1}{2z}\frac{\partial}{\partial \bar{Q}}\left(\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}\right) = -\frac{\bar{F}_{m}^{(z)}}{m^{z}} - \frac{1}{z}\frac{\bar{F}_{m}^{(z-1)}}{m^{z-1}} - \frac{2i\bar{r}}{z}\frac{\bar{F}_{m-1}^{(z-1)}}{(m-1)^{z-1}}.
$$
(C.13)

Denoting the LHS of Eq. (C.13) by

$$
D'_{N,z}(\bar{Q},\bar{r}) = \frac{1}{2z} \frac{\partial}{\partial \bar{Q}} \left(\frac{\bar{F}_{N-1}^{(z-1)}}{(N-1)^{z-1}} \right),\tag{C.14}
$$

we assume that if Eq. (C.13) were true for $N = m$ and $z \neq 0$, then it is also true for $N = m + 1$ for $z \neq 0$. We calculate $D'_{m+1,z}(\tilde{Q}, \tilde{r})$ by substituting $\bar{F}_m^{(z-1)}$ using Eq. (41b) for $N = m$ and for $z \to z - 1$ to get

$$
D'_{m+1,z}(\bar{Q},\bar{r}) = \frac{1}{2z} \frac{\partial}{\partial \bar{Q}} \left(\frac{\bar{F}_m^{(z-1)}}{m^{z-1}} \right) = \frac{-i\bar{r}}{z} \frac{1}{2(z-1)} \frac{\partial}{\partial \bar{Q}} \left(\frac{\bar{F}_{m-1}^{(z-2)}}{(m-1)^{z-2}} \right). \tag{C.15}
$$

Then using Eq. (C.13) for $z \to z - 1$ to substitute the term on the RHS of Eq. (C.15), we finally obtain

$$
D'_{m+1,z}(\bar{Q},\bar{r}) = \frac{-i\bar{r}}{z} \left(-\frac{\bar{F}_m^{(z-1)}}{m^{z-1}} - \frac{1}{z-1} \frac{\bar{F}_m^{(z-2)}}{m^{z-2}} - \frac{2i\bar{r}}{z-1} \frac{\bar{F}_{m-1}^{(z-2)}}{(m-1)^{z-2}} \right) = -\frac{\bar{F}_{m+1}^{(z)}}{(m+1)^z} - \frac{1}{z} \frac{\bar{F}_{m+1}^{(z-1)}}{(m+1)^{z-1}} - \frac{2i\bar{r}}{z} \frac{\bar{F}_m^{(z-1)}}{m^{z-1}} \tag{C.16}
$$

where we repeatedly use Eq. (41b) to replace $\bar{F}_m^{(z-1)}$, $\bar{F}_m^{(z-2)}$ and $\bar{F}_{m-1}^{(z-2)}$
last equality. Now notice that Eq. (C 16) is Eq. (C 13) for $N - m +$ $\bar{F}_{m-1}^{(z-2)}$ by $\bar{F}_{m+1}^{(z)}$ $\bar{F}_{m+1}^{(z-1)}$, $\bar{F}_{m+1}^{(z-1)}$
 \bar{z} + 0 as w $\bar{F}_{m+1}^{(z-1)}$ and $\bar{F}_{m}^{(z-1)}$, respectively, to get the last equality. Now notice that Eq. (C.16) is Eq. (C.13) for $N = m + 1$ for $z \neq 0$ as we set up to prove. This completes the proof that Eqs. (41a)–(41b) satisfy Eqs. (37a)–(37b) for \bar{K}_E .

Appendix D. Generalized eigenvectors of modified Kossakowski–Lindblad (KL) equation

In this appendix, we will work out the first few series of the generalized eigenvectors of the modified KL equation with the Liouvillian \tilde{K}_E (42) at the exceptional point, using the similar method in the last section. $E(E(42))$ at the exceptional point, using the similar method in the last section.

We start with $\tilde{\Phi}_0^{(0)}$ $\phi_0^{(0)} = 1$. From Eq. (39a), the next order vector is $\tilde{\Phi}_1^{(0)}$ $u_1^{(0)} = u_1^{(0)} \tilde{Q} + v_1^{(0)}$ $\tilde{\Phi}_1^{(0)}$ *ir*̃. Substituting $\tilde{\Phi}_1^{(0)}$ $\int_1^{(0)}$ into Eq. (37b), we obtain the condition $u_1^{(0)}$ $v_1^{(0)} = 0$ only. There is no requirement on $v_1^{(0)}$ $v_1^{(0)}$. We choose $v_1^{(0)}$ $\tilde{\omega}_1^{(0)} = \tilde{\omega}$, where $\tilde{\omega} \equiv 2\omega_0/\gamma$ (35). Other choices will give different set of generalized eigenvectors. This ambiguity is attributed to the freedom discussed in 5.2. Then $\tilde{\Phi}_1^{(0)} = i\tilde{\omega}\tilde{r}$. We then use Eq. (39b) to get the n to the freedom discussed in 5.2. Then $\Phi_1^{(0)} = i\tilde{\omega}\tilde{r}$. We then use Eq. (39b) to get the next vector in this series $\tilde{\Phi}_1^{(1)} = p_1^{(1)}\tilde{Q} + q_1^{(1)}\tilde{r}$. Acting \tilde{K}_E on $\tilde{\Phi}_1^{(1)}$, we obtain $p_1^{(1)} =$ $p_1^{(1)} = p_1^{(1)}\tilde{Q} + q_1^{(1)}$ $i_1^{(1)}\tilde{r}$. Acting \tilde{K}_E on $\tilde{\Phi}_1^{(1)}$ $_1^{(1)}$, we obtain $p_1^{(1)}$ $q_1^{(1)} = 1/2$, with no constraint on $q_1^{(1)}$ $1^{(1)}$. The simplest choice is $q_1^{(1)}$ $_{1}^{(1)} = 0$. As a result, $\tilde{\Phi}_{1}^{(1)}$ $\frac{1}{1} = \frac{\tilde{Q}}{2}$ $\frac{Q}{2}$.

Continuing on to $N = 2$, applying Eq. (37a) on $\tilde{\Phi}_{2}^{(0)}$ $v_2^{(0)} = (u_2^{(0)}\tilde{Q} + v_2^{(0)}$ $j_2^{(0)}$ i \tilde{r}) $\tilde{\Phi}_1^{(0)}$ $w_1^{(0)} + w_2^{(0)} \tilde{\Phi}_0^{(0)}$ $u_0^{(0)}$ yield $u_2^{(0)}$ $v_2^{(0)} = 0$ and $w_2^{(0)}$ $\frac{1}{2}^{(0)}_{...}=0.$ Choosing the free coefficient $v_2^{(0)}$ $\tilde{\Phi}_2^{(0)} = \tilde{\omega}$, we get $\tilde{\Phi}_2^{(0)}$ $\tilde{Q}^{(0)} = (i\tilde{\omega}\tilde{r})^2$. The next generalized eigenvector is $\tilde{\Phi}^{(1)}_2$ $u_2^{(1)} = (u_2^{(1)}\tilde{Q} +$ $v_2^{(1)}$ $_2^{(1)}$ i \tilde{r}) $\tilde{\Phi}_1^{(1)}$ $_1^{(1)}$. Substituting in $\tilde{\Phi}_1^{(1)}$ $_1^{(1)}$, Eq. (37b) then produces the condition $u_2^{(1)}$ $v_2^{(1)} = 0$ and $v_2^{(1)}$ $\tilde{\Phi}_2^{(1)} = 2\tilde{\omega}$ to yield $\tilde{\Phi}_2^{(1)}$ $v_2^{(1)}$ *i* \tilde{p} $\tilde{\Phi}_1^{(1)}$. Substituting in $\tilde{\Phi}_1^{(1)}$, Eq. (37b) then produces the condition $u_2^{(1)} = 0$ and $v_2^{(1)} = 2\tilde{\omega}$ to yield $\tilde{\Phi}_2^{(1)} = i\tilde{\omega}\tilde{r}\tilde{Q}$.
The last generalized eigenvector in thi $q_2^{(2)} = (p_2^{(2)}\tilde{Q} + q_2^{(2)})$ $^{(2)}_{2}$ *ir*) $\tilde{\Phi}^{(1)}_{1}$ $s_1^{(1)} + s_2^{(2)}$ $\tilde{\Phi}_0^{(0)}$ $_{0}^{(0)}$, see Eq. (39b). Applying Eq. (37b)

for $z = 2$ then gives $p_2^{(2)}$ $q_2^{(2)} = 1, q_2^{(2)}$ $s_2^{(2)} = 0$ and $s_2^{(2)}$ $p_2^{(2)} = -p_2^{(2)}$ $\tilde{\Phi}_2^{(2)}/4 = -1/4$. Hence, $\tilde{\Phi}_2^{(2)}$ $\zeta_2^{(2)} = \frac{1}{2}\tilde{Q}^2 - \frac{1}{4}$. Continuing in this way, we obtain the generalized eigenvectors for $N = 3$. We can continue in this way to generate higher order generalized eigenvectors. The results can be generalized to Eqs. (44a)–(44b). The first few series of the generalized eigenvectors are summarized in 4.2.

The proof that Eqs. (44a) and (44b) are equal in the overlapping region and that they lead to generalized eigenvectors (38) of \tilde{K}_E that satisfy Eqs. (37a)–(37b) can be proved by induction, similar to the corresponding proof for the generalized eigenvectors of \bar{K}_E in Appendix C. We omit the proof here.

If we start with a different choice of the coefficients, such as $q_1^{(1)}$ If we start with a different choice of the coefficients, such as $q_1^{(1)} = \tilde{\omega}$, we will get a different series of generalized eigenvectors. However, we note that additional terms, such as $x_N^{(z)} \bar{\Phi}_N^{(z)}$ and $x_N^{(z)}$ Eqs. (39a) and (39b), respectively, for this series of generalized eigenvectors. The results are listed in 5.2.

Appendix E. Hu–Paz–Zhang equation

The Liouvillian of the Markovian limit of the HPZ equation [51] differs from that of \bar{K}_{CL} by an extra term, $K'_{\text{H}PZ} = \bar{K}_{\text{CL}} - dL_{2+}$. It can be obtained from \bar{K}_{CL} by a similarity transformation $K'_{\text{H}PZ} = S_3 \bar{K}_{\text{CL}} S_3^{-1}$, where

$$
S_3 = e^{\zeta L_{1+}}.
$$
 (E.1)

In terms of the following coordinates,

$$
Q_{+} \equiv \frac{Q}{\sqrt{2b_{+}}}, \qquad r_{-} \equiv \sqrt{\frac{b_{-}}{2}}r, \tag{E.2}
$$

where

$$
b_{+} \equiv b_{\text{HPZ}} + \frac{d}{2\omega_{0}} = \bar{b} + \frac{d}{4\omega_{0}}, \qquad b_{-} \equiv b_{\text{HPZ}} = \bar{b} - \frac{d}{4\omega_{0}}.
$$
 (E.3)

The stationary state is

$$
f'_{00}(Q_+, r_-) \equiv S_3 \bar{f}_{00}(\bar{Q}, \bar{r}) = \frac{1}{\sqrt{\pi}} e^{-Q_+^2 - r_-^2}.
$$
 (E.4)

We refer the readers to Ref. [25] for the expressions of $\Phi_{mn}^{\prime\pm}$, cf. Eq. (38).

Appendix E.1. Hu–Paz–Zhang (HPZ) equation

In the position coordinates (E.2), the Markovian limit of the HPZ equation has the Liouvillian

$$
K'_{\rm HPZ}(Q_+,r_-) = i\omega_0 \left(-\frac{1}{2} \sqrt{\frac{b_-}{b_+}} \frac{\partial^2}{\partial Q_+ \partial r_-} + 2 \sqrt{\frac{b_+}{b_-}} Q_+ r_- \right) + \gamma \bar{r} \frac{\partial}{\partial \bar{r}} + 2 \gamma \bar{r}^2 + i\omega_0 \frac{b_+ - b_-}{\sqrt{b_+ b_-}} r_- \frac{\partial}{\partial Q_+}.
$$
 (E.5)

The transformed coordinate operators are

$$
\hat{Q}_{+} \equiv S_{3} \bar{Q} S_{3}^{-1} = \frac{1}{\sqrt{2b_{+}(b_{+} + b_{-})}} \left[2b_{+} Q_{+} + \frac{1}{2} (b_{+} - b_{-}) \frac{\partial}{\partial Q_{+}} \right],
$$
\n(E.6a)

$$
\hat{r}_{-} \equiv S_{3}\bar{r}S_{3}^{-1} = \sqrt{\frac{b_{+} + b_{-}}{2b_{-}}}r_{-}.
$$
\n(E.6b)

 K'_{HPZ} has the same exceptional point as \bar{K}_{CL} , that is at $\gamma = 2\omega_0$. We replace this into Eq. (E.5) to get the Liouvillian \mathbb{R}^2 directly from those of K' directly from those of of HPZ equation at the exceptional point K_E' . We can obtain the generalized eigenvectors of K_E' directly from those of \bar{K}_E by similarity transformation. They are

$$
F_N^{'(z)} \equiv S_3 \bar{F}_N^{(z)}, \tag{E.7}
$$

with the stationary state $F_0^{(0)}$ $f_0^{(0)} = f_{00}$ (E.4). The higher order vectors can be generated using

$$
S_3 \bar{Q} \bar{F}_0^{(0)} = \hat{Q}_+ F_0^{'(0)} = \sqrt{\frac{b_+ + b_-}{2b_+}} Q_+ F_0^{'(0)}, \tag{E.8}
$$

$$
S_3(i\bar{r})\bar{F}_0^{(0)} = i\hat{r}_-F_0'^{(0)} = \sqrt{\frac{b_+ + b_-}{2b_-}}ir_-F_0'^{(0)}.
$$
 (E.9)

Applying S_3 to Eqs. (41a) and (41b), the results are

Diagonal:
$$
\frac{\Phi_N^{\prime(z)}}{N^z} = \sqrt{\frac{b_+ + b_-}{2b_+}} \left(Q_+ - \sqrt{\frac{b_+}{b_-}} i r_- \right) \frac{\Phi_{N-1}^{\prime(z)}}{(N-1)^z} - \frac{N-1-z}{2} \frac{\Phi_{N-2}^{\prime(z)}}{(N-2)^z}, \qquad z \neq N, \qquad (E.10a)
$$

Parallel:
$$
\frac{\Phi_N^{\prime(z)}}{N^z} = \sqrt{\frac{b_+ + b_-}{2b_-}} \frac{1}{z} (-ir_-) \frac{\Phi_{N-1}^{\prime(z-1)}}{(N-1)^{z-1}},
$$
 $z \neq 0.$ (E.10b)

As usual, we set $\Phi_{N}^{\prime}(z) = 0$, whenever $z < 0$, $z > N$ or $N < 0$. We can then construct the whole series of generalized eigenfunctions successively starting with $\Phi_0^{\prime(0)}$ $\binom{00}{0} = 1.$

References

- [1] W. D. Heiss, The physics of exceptional points, J. Phys. A: Math. Theor. 45 (44) (2012) 444016. doi:10.1088/1751-8113/45/44/444016.
- [2] M. Berry, Physics of nonhermitian degeneracies, Czech. J. Phys, 54 (10) (2004) 1039–1047. doi:10.1023/B:CJOP.0000044002.05657.04.
- [3] A. Mondragon, E. Hernandez, Degeneracy and crossing of resonance energy surfaces, J. Phys. A: Math. Gen. 26 (20) (1993) 5595. doi:10.1088/0305-4470/26/20/039.
- [4] G. Bhamathi, E. C. G. Sudarshan, Double resonances and jordan block spectra, Int. J. Mod. Phys. B 10 (13-14) (1996) 1531–1544. doi:10.1142/S0217979296000635.
- [5] A. Bohm, M. Loewe, S. Maxson, P. Patuleanu, C. Püntmann, M. Gadella, Gamow-jordan vectors and non-reducible density operators from higher-order s-matrix poles, J. Math. Phys. 38 (1997) 6072–6100. doi:10.1063/1.532203.
- [6] C. Ferise, P. del Hougne, S. Félix, V. Pagneux, M. Davy, Exceptional points of *pt*-symmetric reflectionless states in complex scattering systems, Phys. Rev. Lett. 128 (2022) 203904. doi:10.1103/PhysRevLett.128.203904.
- [7] C. Dembowski, H.-D. Graf, H. L. Harney, A. Heine, W. D. Heiss, H. Rehfeld, A. Richter, Experimental observation of the topological ¨ structure of exceptional points, Phys. Rev. Lett. 86 (2001) 787–790. doi:10.1103/PhysRevLett.86.787.
- [8] S.-B. Lee, J. Yang, S. Moon, S.-Y. Lee, J.-B. Shim, S. W. Kim, J.-H. Lee, K. An, Observation of an exceptional point in a chaotic optical microcavity, Phys. Rev. Lett. 103 (2009) 134101. doi:10.1103/PhysRevLett.103.134101.
- [9] Y. Choi, S. Kang, S. Lim, W. Kim, J.-R. Kim, J.-H. Lee, K. An, Quasieigenstate coalescence in an atom-cavity quantum composite, Phys. Rev. Lett. 104 (2010) 153601. doi:10.1103/PhysRevLett.104.153601.
- [10] J. Schindler, A. Li, M. C. Zheng, F. M. Ellis, T. Kottos, Experimental study of active lrc circuits with PT symmetries, Phys. Rev. A 84 (2011) 040101. doi:10.1103/PhysRevA.84.040101.
- [11] S. K. Ozdemir, S. Rotter, F. Nori, L. Yang, Parity-time symmetry and exceptional points in photonics, Nat. Mat. 18 (2019) 783–798. ¨ doi:10.1038/s41563-019-0304-9.
- [12] M. A. Quiroz-Juárez, A. Perez-Leija, K. Tschernig, B. M. Rodríguez-Lara, O. S. M. na Loaiza, K. Busch, Y. N. Joglekar, R. de J. León-Montiel, Exceptional points of any order in a single, lossy waveguide beam splitter by photon-number-resolved detection, Photon. Res. 7 (8) (2019) 862–867. doi:10.1364/PRJ.7.000862.
- [13] W. Chen, M. Abbasi, Y. N. Joglekar, K. W. Murch, Quantum jumps in the non-hermitian dynamics of a superconducting qubit, Phys. Rev. Lett. 127 (2021) 140504. doi:10.1103/PhysRevLett.127.140504.
- [14] J. Wiersig, Robustness of exceptional-point-based sensors against parametric noise: The role of hamiltonian and liouvillian degeneracies, Phys. Rev. A 101 (2020) 053846. doi:10.1103/PhysRevA.101.053846.
- [15] J. Yuan, L. Geng, J. Huang, Q. Guo, J. Yang, G. Hu, X. Zhou, Exceptional points induced by time-varying mass to enhance the sensitivity of defect detection, Phys. Rev. Appl. 18 (2022) 064055. doi:10.1103/PhysRevApplied.18.064055.
- [16] H. Xu, D. Mason, L. Jiang, J. G. E. Harris, Topological energy transfer in an optomechanical system with exceptional points, Nature 537 (2016) 80–83. doi:10.1038/nature18604.
- [17] B. Peng, S. K. Ozdemir, S. Rotter, H. Yilmaz, M. Liertzer, F. Monifi, C. M. Bender, F. Nori, L. Yang, Loss-induced suppression and revival ¨ of lasing, Science 346 (2014) 328–332. doi:https://doi.org/10.1126/science.1258004.
- [18] A. Guo, G. J. Salamo, D. Duchesne, R. Morandotti, M. Volatier-Ravat, V. Aimez, G. A. Siviloglou, D. N. Christodoulides, Observation of PT-symmetry breaking in complex optical potentials, Phys. Rev. Lett. 103 (2009) 093902. doi:10.1103/PhysRevLett.103.093902.
- [19] W. D. Heiss, F. G. Scholtz, H. B. Geyer, The large n behaviour of the lipkin model and exceptional points, J. Phys. A Math. Gen. 38 (9) (2005) 1843. doi:10.1088/0305-4470/38/9/002.
- [20] T. Petrosky, I. Prigogine, The liouville space extension of quantum mechanics, Adv. Chem. Phys. 99 (1997) 1.
- [21] N. Hatano, Exceptional points of the lindblad operator of a two-level system, Mol. Phys. 117 (15-16) (2019) 2121–2127. doi:10.1080/00268976.2019.1593535.
- [22] F. Minganti, A. Miranowicz, R. W. Chhajlany, F. Nori, Quantum exceptional points of non-hermitian hamiltonians and liouvillians: The effects of quantum jumps, Phys. Rev. A 100 (2019) 062131. doi:10.1103/PhysRevA.100.062131.
- [23] F. Minganti, A. Miranowicz, R. W. Chhajlany, I. I. Arkhipov, F. Nori, Hybrid-liouvillian formalism connecting exceptional points of non-hermitian hamiltonians and liouvillians via postselection of quantum trajectories, Phys. Rev. A 101 (2020) 062112. doi:10.1103/PhysRevA.101.062112.
- [24] I. I. Arkhipov, A. Miranowicz, F. Minganti, F. Nori, Liouvillian exceptional points of any order in dissipative linear bosonic systems: Coherence functions and switching between PT and anti- PT symmetries, Phys. Rev. A 102 (2020) 033715. doi:10.1103/PhysRevA.102.033715.
- [25] B. A. Tay, Eigenvalues of the liouvillian of quadratic master equation for a harmonic oscillator, Physica A 556 (2020) 124768. doi:https://doi.org/10.1016/j.physa.2020.124768.
- [26] V. Gorini, A. Kossakowski, E. C. G. Sudarshan, Completely positive dynamical semigroups of *n*-level systems, J. Math. Phys. 17 (1976) 821–825. doi:10.1063/1.522979.
- [27] G. Lindblad, On the generators of quantum dynamical semigroups, Commun. Math. Phys. 48 (1976) 119–130. doi:10.1007/BF01608499.
- [28] M. A. Nielsen, I. L. Chuang, Quantum Computation and Quantum Information, Cambridge, New York, 2000.
- [29] C. W. Gardiner, P. Zoller, Quantum Noise, 3rd Edition, Springer, Berlin, 2004.
- [30] H.-P. Breuer, F. Petruccione, The Theory of Open Quantum Systems, Oxford, New York, 2002.
- [31] V. May, O. Kuhn, Charge and Energy Transfer Dynamics in Molecular Systems, 3rd Edition, Wiley-VCH, Weinheim, 2011. ¨
- [32] H.-J. Briegel, B.-G. Englert, Quantum optical master equations: The use of damping bases, Phys. Rev. A 47 (4) (1993) 3311–3329. doi:10.1103/PhysRevA.47.3311.
- [33] B. A. Tay, T. Petrosky, Biorthonormal eigenbasis of a markovian master equation for the quantum brownian motion, J. Math. Phys. 49 (11) (2008) 113301. doi:DOI:10.1063/1.3005968.
- [34] D. Honda, H. Nakazato, M. Yoshida, Spectral resolution of the liouvillian of the lindblad master equation for a harmonic oscillator, J. Math. Phys. 51 (7) (2010) 072107. doi:DOI:10.1063/1.3442363.
- [35] P. Talkner, Gauss markov process of a quantum oscillator, Z. Phy. B: Cond. Matt. 41 (4) (1981) 365–374. doi:10.1007/BF01307328. URL http://dx.doi.org/10.1007/BF01307328
- [36] B. A. Tay, Symmetry of bilinear master equations for a quantum oscillator, Physica A 468 (2017) 578–589. doi:http://dx.doi.org/10.1016/j.physa.2016.10.067.
- [37] K. R. Symon, Mechanics, 3rd Edition, Addison-Wesley, New York, 1971.
- [38] D. Heiss, Circling exceptional points, Nat. Phys. 12 (2016) 823-824. doi:10.1038/nphys3864.
- [39] A. Caldeira, A. Leggett, Path integral approach to quantum brownian motion, Physica A 121 (1983) 587–616. doi:https://doi.org/10.1016/0378-4371(83)90013-4.
- [40] E. Joos, H. D. Zeh, C. Kiefer, D. J. W. Giulini, J. Kupsch, I. O. Stamatescu, Decoherence and the Appearance of a Classical World in Quantum Theory, 2nd Edition, Springer, Berlin, 2003.
- [41] P. J. Dodd, J. J. Halliwell, Disentanglement and decoherence by open system dynamics, Phys. Rev. A 69 (5) (2004) 052105. doi:10.1103/PhysRevA.69.052105.
- [42] B. L. Hu, J. P. Paz, Y. Zhang, Quantum brownian motion in a general environment: Exact master equation with nonlocal dissipation and colored noise, Phys. Rev. D 45 (1992) 2843–2861. doi:10.1103/PhysRevD.45.2843.
- [43] B. A. Tay, T. Petrosky, Thermal symmetry of the markovian master equation, Phys. Rev. A (2007) 042102doi:10.1103/PhysRevA.76.042102.
- [44] I. Prigogine, C. George, F. Henin, L. Rosenfeld, A unified formulation of dynamics and thermodynamics, Chem. Scr. 4 (1973) 5–32.
- [45] F. Minganti, A. Biella, N. Bartolo, C. Ciuti, Spectral theory of liouvillians for dissipative phase transitions, Phys. Rev. A 98 (2018) 042118. doi:10.1103/PhysRevA.98.042118.
- [46] P. W. Claeys, A. Lamacraft, Dissipative dynamics in open xxz richardson-gaudin models, Phys. Rev. Res. 4 (2022) 013033. doi:10.1103/PhysRevResearch.4.013033.
- [47] A. Rubio-García, A. L. Corps, A. Relaño, R. A. Molina, F. Pérez-Bernal, J. E. García-Ramos, J. Dukelsky, Exceptional spectral phase in a dissipative collective spin model, Phys. Rev. A 106 (2022) L010201. doi:10.1103/PhysRevA.106.L010201. URL https://link.aps.org/doi/10.1103/PhysRevA.106.L010201
- [48] S. H.Weintraub, Jordan Canonical Form: Theory and Practice, Morgan & Claypool, San Rafael, 2009.
- [49] B. A. Tay, Coherence and Decoherence Processes of a Harmonic Oscillator Coupled with Finite Temperature Field Exact Eigenbasis Solution of Kossakowski-Lindblad's Equation, Ph.D. Dissertation, University of Texas at Austin, 2004.
- [50] B. A. Tay, Solutions of generic bilinear master equations for a quantum oscillator—positive and factorized conditions on stationary states, Physica A 477 (2017) 42–64. doi:http://dx.doi.org/10.1016/j.physa.2017.02.020.
- [51] B. A. Tay, G. Ordonez, Exact markovian kinetic equation for a quantum brownian oscillator, Phys. Rev. E 73 (2006) 016120. doi:10.1103/PhysRevE.73.016120.