

# The semiclassical limit from the Pauli-Poiswell to the Euler-Poiswell system by WKB methods

Changhe Yang<sup>a</sup>, Norbert J. Mauser<sup>b</sup>, and Jakob Möller<sup>b</sup>

<sup>a</sup>Computing and Mathematical Sciences (CMS) Department, California Institute of Technology, 91125 Pasadena, CA, USA

<sup>b</sup>Research Platform MMM "Mathematics-Magnetism-Materials" c/o Fak. Mathematik, Univ. Wien, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria

June 12, 2023

## Abstract

The self-consistent Pauli-Poiswell equation for 2-spinors is the first order in  $1/c$  semi-relativistic approximation of the Dirac-Maxwell equation for 4-spinors coupled to the self-consistent electromagnetic fields generated by the density and current density of a fast moving electric charge. It consists of a vector-valued magnetic Schrödinger equation with an extra term coupling spin and magnetic field via the Pauli matrices coupled to 1+3 Poisson type equations as the magnetostatic approximation of Maxwell's equations. The Pauli-Poiswell equation is a consistent  $O(1/c)$  model that keeps both relativistic effects magnetism and spin which are both absent in the non-relativistic Schrödinger-Poisson equation and inconsistent in the magnetic Schrödinger-Maxwell equation.

We present the mathematically rigorous semiclassical limit  $\hbar \rightarrow 0$  of the Pauli-Poiswell equation towards the magnetic Euler-Poiswell equation. We use WKB analysis which is valid locally in time only. A key step is to obtain an a priori energy estimate for which we have to take into account the Poisson equations for the magnetic potential with the current as source term. Additionally we obtain the weak convergence of the monokinetic Wigner transform and strong convergence of the density and the current density. We also prove local wellposedness of the Euler-Poiswell equation which is global unless a finite time blow-up occurs.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Notations . . . . .	5
1.2	Outline . . . . .	6
<b>2</b>	<b>Statement of the main result</b>	<b>6</b>
2.1	WKB Ansatz and formal limit to the Euler-Poiswell equation . . . . .	6

2.2	Wigner transform and Pauli-Wigner equation . . . . .	11
2.3	Main result . . . . .	13
<b>3</b>	<b>A priori estimates</b>	<b>15</b>
3.1	Estimate of $E_s(t)$ . . . . .	16
3.2	Estimates for the Poisson equations . . . . .	19
3.3	Estimate of $\varepsilon\ a\ _{H^s}$ in $E_s^1(t)$ . . . . .	22
3.3.1	Motivation of Lemma 5 . . . . .	23
3.3.2	Proof of Lemma 5 . . . . .	25
3.4	Proof of Proposition 1 . . . . .	30
<b>4</b>	<b>Existence and Uniqueness of Solutions</b>	<b>32</b>
<b>5</b>	<b>Behavior at Blowup Time</b>	<b>43</b>
<b>6</b>	<b>Semiclassical limit</b>	<b>44</b>
<b>A</b>	<b>Some inequalities</b>	<b>49</b>

## 1 Introduction

We deal with the "semiclassical" limit of vanishing Planck constant for nonlinear Schrödinger equations. In particular we deal with the self-consistent Pauli-Poiswell equation as a relativistic extension of the Schrödinger-Poisson equation in  $3 - d$  (e.g in 3 space dimensions) which keeps magnetic and spin effects. There are 2 main techniques for such semiclassical limits:

a) *Wigner transforms* (see [8] for an overview) that yield global (in time) limits towards Vlasov type equations, see e.g. Lions-Paul [21] and Markowich-Mauser [22] for the limit from the Schrödinger-Poisson equation to the Vlasov-Poisson equation for the *mixed state* case in  $3 - d$  and Zhang-Zheng-Mauser [37] for the *pure state* case which is possible in  $1 - d$  only with appropriate non unique measures valued solutions of the Vlasov-Poisson equation.

b) *WKB methods* (after Wentzel, Kramers and Brillouin) that yield local (in time) limits to Euler type equations which correspond to monokinetic Vlasov equations for pure states. For the WKB analysis of the Schrödinger-Poisson equation and its convergence towards the Euler-Poisson equation we refer to the work of Zhang [35, 36] and Alazard and Carles [2, 5]. Grenier [13] proved the semiclassical limit of the cubic NLS and recently, Gui and Zhang [14] proved the semiclassical limit of the Gross-Pitaevskii equation using WKB analysis.

For a unified presentation of these approaches and extensions to multi-valued WKB see e.g. [31].

The analysis for the Pauli-Poiswell equation is much harder than for the Schrödinger-Poisson equation. Already for the linear Pauli equation (i.e. with given "external" potentials) the analysis is much more complicated than for the (magnetic) Schrödinger equation because of the existence of zero modes due to the presence of the Stern-Gerlach term involving the magnetic field  $B = \nabla \times A$ , see e.g. [7], where even the case of constant magnetic fields shows hard technical challenges. At the core of the technical difficulties, besides the Stern-Gerlach term, is

the "advective" term  $A \cdot \nabla$  in the magnetic Schrödinger operator and the nonlinear coupling of the magnetic vector potential  $A$  via the 3 additional Poisson type equations for the 3 components of the magnetic potential with the current density as source term.

In the present work we deal with case b) , i.e. the semiclassical limit via WKB methods. The Wigner method case a) is dealt with in [24] and the existence and uniqueness analysis of the Pauli-Poisson system is dealt with in [11]. The related simplified *Pauli-Poisson equation* was presented in [26] where an external magnetic field is applied that is much stronger than the magnetic self-interaction that can hence be neglected, while the electric field is kept self-consistent by a mere coupling to the Poisson equation for  $V^{\hbar,c}$ . The semiclassical limit of the Pauli-Poisson equation to the Vlasov-Poisson equation with Lorentz force by the Wigner method a) was proven in [25] (with some additional details in [27]), which is a first true extension of the results for the Schrödinger-Poisson equation 30 years ago.

The Pauli-Poisson equation for a 2-spinor  $\psi^{\hbar,c}(x, t) \in (L^2(\mathbb{R}^3 \times \mathbb{R}, \mathbb{C}))^2$  and the electromagnetic potentials  $A^{\hbar,c}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  and  $V^{\hbar,c}: \mathbb{R}^3 \rightarrow \mathbb{R}$  depending on two parameters  $\hbar, c$  reads (cf. [23])

$$i\hbar\partial_t\psi^{\hbar,c} = -\frac{1}{2}(\hbar\nabla - \frac{i}{c}A^{\hbar,c})^2\psi^{\hbar,c} + V^{\hbar,c}\psi^{\hbar,c} - \frac{1}{2}\frac{\hbar}{c}(\boldsymbol{\sigma} \cdot B^{\hbar,c})\psi^{\hbar,c}, \quad (1.1)$$

$$-\Delta V^{\hbar,c} = \rho^{\hbar,c} := |\psi^{\hbar,c}|^2, \quad (1.2)$$

$$-\Delta A^{\hbar,c} = \frac{1}{c}J^{\hbar,c}. \quad (1.3)$$

where the Pauli current density is given by

$$J^{\hbar,c}(\psi^{\hbar,c}, A^{\hbar,c}) = \text{Im}(\overline{\psi^{\hbar,c}}(\hbar\nabla - \frac{i}{c}A^{\hbar,c})\psi^{\hbar,c}) - \hbar\nabla \times (\overline{\psi^{\hbar,c}}\boldsymbol{\sigma}\psi^{\hbar,c}), \quad (1.4)$$

with initial data

$$\psi^{\hbar,c}(x, 0) = \psi^{\hbar,c,0}(x) \in (L^2(\mathbb{R}^3))^2. \quad (1.5)$$

We denote the Pauli Hamiltonian by

$$H = -\frac{1}{2}(\hbar\nabla - \frac{i}{c}A^{\hbar,c})^2 + V^{\hbar,c} - \frac{1}{2}\frac{\hbar}{c}(\boldsymbol{\sigma} \cdot B^{\hbar,c}) \quad (1.6)$$

where  $B^{\hbar,c} := \nabla \times A^{\hbar,c}$  is the magnetic field and  $|\psi^{\hbar,c}|^2 := |\psi_1^{\hbar,c}|^2 + |\psi_2^{\hbar,c}|^2$  the scalar charge density. Note that we work in Lorenz gauge, which is consistent in  $O(1/c)$ ,

$$\text{div}A^{\hbar,c} + \frac{1}{c}\partial_t V^{\hbar,c} = 0. \quad (1.7)$$

The Stern-Gerlach term is  $\boldsymbol{\sigma} \cdot B^{\hbar,c} := \sum_{k=1}^3 \sigma_k B_k^{\hbar,c}$  where the  $\sigma_k$  are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1.8)$$

The expressions  $\overline{\psi^{\hbar,c}}\nabla\psi^{\hbar,c}$  and  $\overline{\psi^{\hbar,c}}\boldsymbol{\sigma}\psi^{\hbar,c}$  are to be understood as the vectors with components

$$\left(\overline{\psi_1^{\hbar,c}}, \overline{\psi_2^{\hbar,c}}\right)\partial_k \begin{pmatrix} \psi_1^{\hbar,c} \\ \psi_2^{\hbar,c} \end{pmatrix}, \quad \left(\overline{\psi_1^{\hbar,c}}, \overline{\psi_2^{\hbar,c}}\right)\sigma_k \begin{pmatrix} \psi_1^{\hbar,c} \\ \psi_2^{\hbar,c} \end{pmatrix},$$

for  $k = 1, 2, 3$ . Note that using the Pauli vector identity  $(a \cdot \boldsymbol{\sigma})(b \cdot \boldsymbol{\sigma}) = (a \cdot b)I + i(a \times b) \cdot \boldsymbol{\sigma}$  the Pauli Hamiltonian can be rewritten as

$$H = -\frac{1}{2}(\boldsymbol{\sigma} \cdot (\hbar\nabla - \frac{i}{c}A^{\hbar,c}))^2 + V^{\hbar,c}, \quad (1.9)$$

and we can also rewrite  $J^{\hbar,c}$  as

$$J^{\hbar,c} = \text{Re} \left( \overline{\psi^{\hbar,c}} \boldsymbol{\sigma} (\boldsymbol{\sigma} \cdot (-i\hbar\nabla - \frac{1}{c}A^{\hbar,c})) \psi^{\hbar,c} \right). \quad (1.10)$$

which is to be understood as the 3-vector with components

$$J_k^{\hbar,c} = \text{Re} \left( \overline{\psi^{\hbar,c}} \sigma_k (\boldsymbol{\sigma} \cdot (-i\hbar\nabla - \frac{1}{c}A^{\hbar,c})) \psi^{\hbar,c} \right).$$

Since we are interested in the semiclassical ( $\hbar \rightarrow 0$ ) limit with  $c$  fixed we will omit the  $c$  superscript in the sequel. The self-consistent Pauli-Poisson equation (a portmanteau of *Poisson* and *Maxwell* coined in [23]), arises as the semi-relativistic approximation at  $O(1/c)$  of the fully relativistic Dirac-Maxwell equation for a 4-spinor. The Poisson equations for  $A^{\hbar}$  and  $V^{\hbar}$  are the magnetostatic approximation of Maxwell's equation by keeping terms of order  $O(1/c)$ . The two components of the Pauli equation describe the two spin states of a charged spin-1/2-particle, in particular a fermion, whereas the Poisson equations describe the electrodynamic self-interaction of a fast moving particle with the electromagnetic field that it generates itself due to the finite speed of light.

We use a scaling where the dimensionless „semiclassical parameter“  $\varepsilon$  is proportional to  $\hbar$  and (1.1)-(1.3) reads

$$i\varepsilon\partial_t\psi^\varepsilon = -\frac{1}{2}(\varepsilon\nabla - iA^\varepsilon)^2\psi^\varepsilon + V^\varepsilon\psi^\varepsilon - \frac{1}{2}\varepsilon(\boldsymbol{\sigma} \cdot B^\varepsilon)\psi^\varepsilon, \quad (1.11)$$

$$-\Delta V^\varepsilon = \rho^\varepsilon := |\psi^\varepsilon|^2, \quad (1.12)$$

$$-\Delta A^\varepsilon = J^\varepsilon. \quad (1.13)$$

where the Pauli current density is given by

$$J^\varepsilon(\psi^\varepsilon, A^\varepsilon) = \text{Im}(\overline{\psi^\varepsilon}(\varepsilon\nabla - iA^\varepsilon)\psi^\varepsilon) - \varepsilon\nabla \times (\overline{\psi^\varepsilon}\boldsymbol{\sigma}\psi^\varepsilon). \quad (1.14)$$

Since we work in  $d = 3$  dimensions the Poisson equation for  $V^\varepsilon$  is equivalent to the Hartree nonlinearity where the density is convoluted with  $|x|^{-1}$ , i.e. the fundamental solution of the Poisson equation in three dimensions:

$$V^\varepsilon[\psi^\varepsilon] = (-\Delta)^{-1}\rho^\varepsilon = \frac{1}{4\pi|x|} * \rho^\varepsilon = \frac{1}{4\pi|x|} * |\psi^\varepsilon|^2 \quad (1.15)$$

The current density becomes

$$J^\varepsilon = \frac{i\varepsilon}{2} \sum_{j=1}^2 (\overline{\psi_j^\varepsilon} \nabla \psi_j^\varepsilon - \psi_j^\varepsilon \nabla \overline{\psi_j^\varepsilon}) - \rho A^\varepsilon - \frac{\varepsilon}{2} \nabla \times (\overline{\psi^\varepsilon} \boldsymbol{\sigma} \psi^\varepsilon). \quad (1.16)$$

Note that  $\rho$  and  $J$  satisfy the continuity equation

$$\partial_t \rho^\varepsilon - \operatorname{div}(J^\varepsilon) = 0. \quad (1.17)$$

**Remark 1.** Let us briefly discuss the gauge of the Pauli-Poiswell equation (1.1)-(1.5). In the derivation in [23] the authors use Lorenz gauge

$$\operatorname{div} A^\varepsilon + \frac{1}{c} \partial_t V^\varepsilon = 0, \quad (1.18)$$

where  $\varepsilon = 1/c$ , in order to transform Maxwell's equations for the fields into a system of wave equations for the potentials

$$\square A^\varepsilon = -\frac{1}{c} J^\varepsilon, \quad \square V^\varepsilon = -\rho, \quad \square = \frac{1}{c^2} \partial_t^2 - \Delta.$$

They perform an asymptotic expansion up to first order in  $\varepsilon = 1/c$  and arrive at (1.1)-(1.5). This means that using the Lorenz gauge for the Pauli-Poiswell equation is consistent at first order.

## 1.1 Notations

Let  $C_{M_1, \dots, M_n}$  denote a constant which only depends on some constants  $M_1, \dots, M_n$ . Let

$$A \lesssim_{M_1, \dots, M_n} B,$$

denote  $|A| \leq C_{M_1, \dots, M_n} |B|$ . Also, we omit the dependence on the regularity parameter  $s$  and use  $\lesssim$  instead of  $\lesssim_s$  for simplicity, where  $s$  is the required regularity in  $H^s$ . For a  $m \times n$  matrix  $A$  and a  $n$ -dimensional vector  $x$  we use summation convention, i.e.

$$A_i^j x_j := \sum_{j=1}^n A_{ij} x_j.$$

For two vector-valued functions  $f, g : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , if  $f$  is differentiable, we use  $g \cdot \nabla f$  to denote the vector  $(\partial^j f_i g_j)_{1 \leq i \leq d}$ . Notice that here we use summation convention. For  $f \in H^s(\Omega)$  and a vector  $\alpha = (\alpha_1, \dots, \alpha_s)$  with  $\forall 1 \leq i \leq s, \alpha_i \in \{1, 2, 3\}$ , we denote its weak derivative by

$$\partial^\alpha f(x) := \partial^{\alpha_1} \dots \partial^{\alpha_s} f(x).$$

Let  $V, W$  be vector-valued functions with components  $V_j, W_j$ . By  $V \otimes W$  we denote the matrix

$$(V \otimes W)_{ij} = V_i W_j.$$

For a vector  $\alpha$ , let  $n(\alpha)$  denote the length of  $\alpha$ . For a matrix  $X \in \mathbb{R}^{m \times n}$ , we will define the magnitude, the  $L^2$  norm, the  $H^s$  norm and the  $W^{s,\infty}$  norm as

$$\begin{aligned} |X|^2 &= \sum_{1 \leq i \leq m, 1 \leq j \leq n} |X_{ij}|^2, & \|X\|_{L^2} &= \|(|X|)\|_{L^2}, \\ \|X\|_{H^s} &= \sum_{n(\alpha) < s} \|\partial^\alpha X\|_{L^2}, & \|X\|_{W^{s,\infty}} &= \sup_{\substack{n(\alpha) < s, \\ 1 \leq i \leq m, \\ 1 \leq j \leq n}} \|\partial^\alpha X_{ij}\|_{L^\infty}. \end{aligned}$$

For vectors  $v \in \mathbb{R}^d$  we can define the respective norms regarding  $v$  as an  $1 \times d$  matrix.

## 1.2 Outline

In Section 2 we introduce the WKB ansatz and transform the Pauli-Poiswell equation into a system with unknowns  $a^\varepsilon$  and  $u^\varepsilon$  instead of  $\psi^\varepsilon$  where  $a^\varepsilon$  is the amplitude and  $u^\varepsilon$  is the velocity, i.e. the gradient of the phase  $S^\varepsilon$  in the WKB ansatz (2.2). Then we introduce the Wigner transform and Wigner measures. Finally we state our main results: The local wellposedness and blow up alternative of the Vlasov-Poiswell and Euler-Poiswell equations are stated in Theorem 1. The semiclassical limit from the Vlasov-Poiswell equation to the Euler-Poiswell equation is stated in Theorem 2.

In section 3 we will derive an a priori estimate (Proposition 1) which allows us to bound the  $H^s$  norm of  $u^\varepsilon$  and the  $H^{s-1}$  norm of  $a^\varepsilon$ . First we obtain an estimate for the charge (Lemma 2) and combine it with an estimate for the potentials  $A^\varepsilon$  and  $V^\varepsilon$  (Lemma 3). This gives an estimate involving  $\varepsilon \|a^\varepsilon\|_{H^s}$  (Lemma 4). Using energy conservation (Lemma 5) we arrive at an estimate independent of  $\varepsilon$ . The a priori estimate is the key estimate for Theorems 1 and Theorem 2.

We will then provide existence and uniqueness for both equations using the a priori estimate and a contraction argument. Based on these results we are able to prove the semiclassical limit of the Pauli-Poiswell-WKB equation (2.5) towards an Euler-Poiswell equation in Section 6. Some technical lemmas which we need in the analysis will be introduced in Appendix A.

## 2 Statement of the main result

### 2.1 WKB Ansatz and formal limit to the Euler-Poiswell equation

The WKB ansatz consists of assuming that the initial data  $\psi^{\varepsilon,0}$  of the Pauli-Poiswell equation are of the form

$$\psi_j^{\varepsilon,0}(x) = a_j^{\varepsilon,0}(x) e^{\frac{i}{\varepsilon} S^{\varepsilon,0}(x)}, \quad j = 1, 2, \quad (2.1)$$

where  $a_j^{\varepsilon,0}$  are the *initial amplitudes* of the components of the 2-spinor. The *initial phase*  $S^{\varepsilon,0}$ , which we choose to be the same for both components, is real-valued. The gradient of  $S^{\varepsilon,0}$  is the *initial velocity*  $u^{\varepsilon,0} := \nabla S^{\varepsilon,0}$ . One then expects that at least for short times the solution  $\psi^\varepsilon$  will be of the form

$$\psi_j^\varepsilon(x, t) = a_j^\varepsilon(x, t) e^{\frac{i}{\varepsilon} S^\varepsilon(x, t)}. \quad (2.2)$$

where  $a_j^\varepsilon$  are the *amplitudes* and  $S^\varepsilon$  is the *phase*. The *velocity*  $u^\varepsilon$  is defined as the gradient of the phase,  $u^\varepsilon := \nabla S^\varepsilon$ . We can also write (2.2) in spinor notation, i.e. by writing  $\psi^\varepsilon = (\psi_1^\varepsilon, \psi_2^\varepsilon)^T$  and  $a^\varepsilon = (a_1^\varepsilon, a_2^\varepsilon)^T$  we have

$$\psi^\varepsilon(x, t) = a^\varepsilon(x, t) e^{\frac{i}{\varepsilon} S^\varepsilon(x, t)}. \quad (2.3)$$

**Remark 2.** Note that the two components of  $\psi^{\varepsilon, 0}$  in the ansatz (2.1) have the same phase  $S^{\varepsilon, 0}$ . If one would choose more complicated initial data with two different phases  $S_j^{\varepsilon, 0}$ ,  $j = 1, 2$ , then also the solution  $\psi^\varepsilon$  would have two phases and oscillatory cross terms like  $\exp(\frac{i}{\varepsilon}(S_1^\varepsilon - S_2^\varepsilon))$ , would appear in (2.4).

Note that when dealing with matrix-valued Hamiltonians one usually employs the "multi-component WKB" ansatz (2.1) by taking a vector-valued amplitude and a scalar phase which is a sound mathematical choice that is also usually chosen by physicists (see e.g. [19], [32]).

Substituting  $\psi^\varepsilon$  given by (2.2) into (1.11) yields

$$\begin{aligned} i\varepsilon(\partial_t a_j^\varepsilon + \frac{i}{\varepsilon} a_j^\varepsilon \partial_t S^\varepsilon) e^{\frac{i}{\varepsilon} S^\varepsilon} &= -\frac{\varepsilon^2}{2}(\Delta a_j^\varepsilon + \frac{2i}{\varepsilon} \nabla a_j^\varepsilon \cdot \nabla S^\varepsilon + \frac{i}{\varepsilon} a_j^\varepsilon \Delta S^\varepsilon - \frac{1}{\varepsilon^2} |\nabla S^\varepsilon|^2 a_j^\varepsilon) e^{\frac{i}{\varepsilon} S^\varepsilon} \\ &\quad - i\varepsilon A^\varepsilon \cdot (\nabla a_j^\varepsilon + \frac{i}{\varepsilon} a_j^\varepsilon \nabla S^\varepsilon) e^{\frac{i}{\varepsilon} S^\varepsilon} - \frac{i\varepsilon}{2} \operatorname{div} A^\varepsilon a_j^\varepsilon e^{\frac{i}{\varepsilon} S^\varepsilon} + (\frac{|A^\varepsilon|^2}{2} + V^\varepsilon) a_j^\varepsilon e^{\frac{i}{\varepsilon} S^\varepsilon} \\ &\quad - \frac{\varepsilon}{2} \mathring{\mathbf{B}}_j^i a_i^\varepsilon e^{\frac{i}{\varepsilon} S^\varepsilon}. \end{aligned}$$

Here,  $\mathbf{B}^\varepsilon = \boldsymbol{\sigma} \cdot B^\varepsilon = \sum_{k=1}^3 \sigma_k B_k^\varepsilon$  is a  $2 \times 2$  matrix and we use summation convention for  $\mathring{\mathbf{B}}_j^i a_i$ . Rewrite the equation above as

$$\begin{aligned} &a_j \left( \partial_t S^\varepsilon + \frac{1}{2} |\nabla S^\varepsilon|^2 + (\frac{|A^\varepsilon|^2}{2} + V^\varepsilon) + A^\varepsilon \cdot \nabla S^\varepsilon \right) \\ &- i\varepsilon \left( \partial_t a_j^\varepsilon - \frac{i}{2} \mathring{\mathbf{B}}_j^i a_i^\varepsilon - \frac{i}{2} (\varepsilon \Delta a_j^\varepsilon + 2i \nabla a_j^\varepsilon \cdot \nabla S^\varepsilon + i a_j^\varepsilon \Delta S^\varepsilon) + \frac{1}{2} \operatorname{div} A^\varepsilon a_j^\varepsilon + A^\varepsilon \cdot \nabla a_j^\varepsilon \right) = 0. \end{aligned}$$

Then we have the following system:

$$\begin{aligned} \partial_t a_j^\varepsilon + A^\varepsilon \cdot \nabla a_j^\varepsilon + \nabla a_j^\varepsilon \cdot \nabla S^\varepsilon + \frac{1}{2} a_j^\varepsilon (\Delta S^\varepsilon + \operatorname{div} A^\varepsilon) &= \frac{i\varepsilon}{2} \Delta a_j^\varepsilon + \frac{i}{2} \mathring{\mathbf{B}}_j^i a_i^\varepsilon, \\ \partial_t S^\varepsilon + \frac{1}{2} |\nabla S^\varepsilon|^2 + A^\varepsilon \cdot \nabla S^\varepsilon + (\frac{|A^\varepsilon|^2}{2} + V^\varepsilon) &= 0, \\ a_j^\varepsilon(x, 0) &= a_j^{\varepsilon, 0}(x), \\ S^\varepsilon(x, 0) &= S^{\varepsilon, 0}(x). \end{aligned} \quad (2.4)$$

Let now  $u^{\varepsilon, 0} = \nabla S^{\varepsilon, 0}$  and  $u^\varepsilon = \nabla S^\varepsilon$  be the initial velocity and velocity, respectively. By differentiating the second equation in (2.4) we obtain the *Pauli-Poisswell-WKB equation*

$$\begin{aligned} \partial_t a_j^\varepsilon + (A^\varepsilon + u^\varepsilon) \cdot \nabla a_j^\varepsilon + \frac{1}{2} a_j^\varepsilon \operatorname{div}(u^\varepsilon + A^\varepsilon) &= \frac{i\varepsilon}{2} \Delta a_j^\varepsilon + \frac{i}{2} \mathring{\mathbf{B}}_j^i a_i^\varepsilon \\ \partial_t u^\varepsilon + (A^\varepsilon + u^\varepsilon) \cdot \nabla u^\varepsilon + u^\varepsilon \cdot \nabla A^\varepsilon + \nabla \left( \frac{|A^\varepsilon|^2}{2} + V^\varepsilon \right) &= 0 \\ a_j^\varepsilon(x, 0) &= a_j^{\varepsilon, 0}(x) \\ u^\varepsilon(x, 0) &= u^{\varepsilon, 0}(x). \end{aligned} \quad (2.5)$$

where

$$-\Delta V^\varepsilon = \rho^\varepsilon, \quad (2.6)$$

$$-\Delta A^\varepsilon = \varepsilon w^\varepsilon - \varepsilon v^\varepsilon - \rho^\varepsilon u^\varepsilon - \rho^\varepsilon A^\varepsilon, \quad (2.7)$$

with

$$\rho^\varepsilon := |a^\varepsilon|^2 \quad (2.8)$$

$$v^\varepsilon := \frac{1}{2} \nabla \times (\overline{a^\varepsilon} \sigma_1 a^\varepsilon, \overline{a^\varepsilon} \sigma_2 a^\varepsilon, \overline{a^\varepsilon} \sigma_3 a^\varepsilon)^T, \quad (2.9)$$

$$w^\varepsilon := \frac{i}{2} \sum_{j=1}^2 (\overline{a_j^\varepsilon} \nabla a_j^\varepsilon - a_j^\varepsilon \nabla \overline{a_j^\varepsilon}). \quad (2.10)$$

Then  $J$  is given by

$$J^\varepsilon = \varepsilon w^\varepsilon - \varepsilon v^\varepsilon - \rho^\varepsilon (u^\varepsilon + A^\varepsilon). \quad (2.11)$$

There is a one-to-one match between each solution  $\psi^\varepsilon$  of the original Pauli-Poiswell equation (1.11) (up to a constant  $e^{i\theta}$ ) and each solution  $(a^\varepsilon, u^\varepsilon)$  of (2.5). Given  $(a^\varepsilon, u^\varepsilon)$  solving (2.5), we can construct  $\psi^\varepsilon$  solving (1.11) by (2.2). Inversely, if one has  $\psi^\varepsilon$  solving (1.11) one needs to calculate  $V^\varepsilon$  and  $A^\varepsilon$ . Then  $u^\varepsilon$  is obtained through the second equation in (2.5) and then  $a^\varepsilon$  through the first equation in (2.5). This is where existence and uniqueness of solutions to (2.5) are needed. However this approach is only local in time since (2.5) has the following characteristics

$$\dot{x} = u^\varepsilon + A^\varepsilon, \quad x(0) = x_0.$$

In general the characteristics can intersect in finite time which means that caustics appear (cf. [31]).

**Remark 3.** Note that in the WKB ansatz we choose a complex-valued amplitude as opposed to the Madelung transform where one uses a real-valued amplitude. This additional degree of freedom allows us to avoid singular quantum pressure in the equation for the velocity  $u^\varepsilon$ . Instead one obtains a skew-symmetric term in the transport equation (due to the term  $(i/\varepsilon)/2\Delta a_j^\varepsilon$ ). For more details on the Madelung transform and its connection to the hydrodynamic formulation of nonlinear Schrödinger equations we refer to the survey in [3].

Suppose  $a^0$  and  $u^0$  are the limit of  $a^{\varepsilon,0}$  and  $u^{\varepsilon,0}$  in a sense to be specified later. We can formally pass to the limit  $\varepsilon \rightarrow 0$  in system (2.5) and obtain the following *pressureless Euler-Poiswell equation*,

$$\begin{aligned} \partial_t a_j + (A + u) \cdot \nabla a_j + \frac{1}{2} a_j \operatorname{div}(u + A) &= \frac{i}{2} \mathbf{B}_j^i a_i, \\ \partial_t u + (A + u) \cdot \nabla u + u \cdot \nabla A + \nabla \left( \frac{|A|^2}{2} + V \right) &= 0, \\ a_j(x, 0) &= a_j^0(x), \\ u(x, 0) &= u^0(x). \end{aligned} \quad (2.12)$$



where

$$-\Delta V = \rho, \quad (2.13)$$

$$-\Delta A = -\rho u - \rho A, \quad (2.14)$$

and

$$\rho = |a|^2.$$

Multiplying the first equation in (2.12) by  $\bar{a}_j$  summing over  $j = 1, 2$ , and taking the real part yields

$$\begin{aligned} \partial_t \rho + \nabla \cdot (\rho(u + A)) &= 0, \\ \partial_t u + (A + u) \cdot \nabla u + u \cdot \nabla A + \nabla \left( \frac{|A|^2}{2} + V \right) &= 0, \\ a_j(x, 0) &= a_j^0(x), \\ u(x, 0) &= u^0(x). \end{aligned} \quad (2.15)$$

and

$$-\Delta V = \rho, \quad (2.16)$$

$$-\Delta A = -\rho u - \rho A, \quad (2.17)$$

since  $\text{Re}(i\bar{a}(\boldsymbol{\sigma} \cdot B)a) = 0$ .

The Euler-Poisson equation (2.15) reduces to the *repulsive pressureless Euler-Poisson equation* when there is no magnetic field present, i.e.  $A = 0$ :

$$\begin{aligned} \partial_t \rho + \nabla \cdot (\rho u) &= 0 \\ \partial_t (\rho u) + \nabla \cdot (\rho u \otimes u) &= -\rho \nabla V, \\ \rho(x, 0) &= \rho^0(x), \\ u(x, 0) &= u^0(x). \end{aligned} \quad (2.18)$$

where

$$-\Delta V = \rho, \quad \rho = |a|^2. \quad (2.19)$$

The Euler-Poisson equation arises as the semiclassical limit of the Schrödinger-Poisson equation with WKB initial data as was shown in [2]. Recently, Golse and Paul [12] showed that the bosonic  $N$ -particle Schrödinger dynamics with Coulomb interaction converge to the Euler-Poisson equation in the combined semiclassical and mean field limit  $\hbar + 1/N \rightarrow 0$  if the first marginal of the initial data has monokinetic Wigner measure (for the definition cf Section 2.2) which is equivalent to the WKB formulation.

Depending on the sign in front of the term  $\rho \nabla V$  on the LHS of the second equation in (2.18) the Euler-Poisson equation is called *repulsive* (with a negative sign) or *attractive* (with a positive sign). In other words, the Euler-Poisson equation is repulsive if the sign on the LHS of

the second equation in (2.18) and the sign on the RHS of the Poisson equation (2.19) are the same and attractive if they are opposite.

In the repulsive case the Euler-Poisson equation is a non-relativistic self-consistent model for the self-interaction with the electric field, coupled via the Poisson equation (2.19)), for a system (plasma) of charged particles of charge of the same sign (e.g. electrons). This is because the Coulomb interaction is repulsive for charges of the same sign.

In the attractive case the Euler-Poisson equation is a self-consistent model for non-relativistic gravitational interaction, e.g. gaseous stars, which is attractive.

The analytical properties of the Euler-Poisson equation depend heavily on the sign. The repulsive case is less delicate than the attractive case: The dispersive nature of the electric field may prevent the formation of singularities [17]. In 3d Guo and Pausader [15, 17] proved the global existence of small smooth irrotational flows. Recently, Hadžić and Jang constructed global solutions for both the gravitational and electrostatic Euler-Poisson equation in [18].

However it is also possible to have blow-up even for the repulsive case, e.g. [34] or even non-existence, e.g. [30] for the 1d case. Further references can also be found in [6].

The fully relativistic self-consistent interaction with the electromagnetic field is modeled by the *pressureless Euler-Maxwell equation*, studied in [10, 16]. Following the notation in [10] it is given by

$$\begin{aligned}
\partial_t \rho + \nabla \cdot (\rho u) &= 0 \\
\partial_t (\rho u) + \nabla \cdot (\rho u \otimes u) &= \rho(E + u \times B), \\
\partial_t E - \nabla \times B &= \rho u, \quad \partial_t B + \nabla \times E = 0, \\
\nabla \cdot E &= -\rho, \quad \nabla \cdot B = 0. \\
(\rho, u, E, B)(x, 0) &= (\rho^0(x), u^0(x), E^0(x), B^0(x))
\end{aligned} \tag{2.20}$$

The non-relativistic limit  $c \rightarrow \infty$  of the Euler-Maxwell equation (2.20) to the Euler-Poisson equation was shown in [29, 33]. In the Euler-Maxwell case there is no distinction between repulsive or attractive interaction since the force described by Maxwell's equations is given by the Lorentz force  $F_L = q(E + u \times B)$  (for the non-relativistic Euler-Poisson equation it reduces to the repulsive electrostatic force  $F_{el} = qE$ ). Similar to the Euler-Poisson equation the Euler-Maxwell equation is in fact more stable than the pure Euler system without self-interaction and smooth solutions can exist globally without shock formation [16].

The Euler-Poiswell equation (2.15)-(2.17) is the  $O(1/c)$  semi-relativistic approximation of the Euler-Maxwell equation which is natural since the Pauli-Poiswell equation is the semi-relativistic  $O(1/c)$  approximation of the Dirac-Maxwell equation and the Euler-Maxwell equation is the semiclassical limit of the Dirac-Maxwell equation. Therefore the Euler-Poiswell equation, with the repulsive sign in all Poisson type equations, is the correct semi-relativistic self-consistent  $O(1/c)$  model of a system (plasma) of charged particles with self-interaction with the electromagnetic field. Similar to the Euler-Maxwell equation the Euler-Poiswell equation describes the interaction with the electromagnetic field via the Lorentz force.

Our new local existence result in Theorem 1.a for the Euler-Poiswell equation is based on the local wellposedness of the Pauli-Poiswell-WKB equation (2.5).

## 2.2 Wigner transform and Pauli-Wigner equation

The Wigner transform  $f^\varepsilon(x, \xi, t)$  of  $\psi^\varepsilon(x, t)$  is defined by

$$f^\varepsilon(x, \xi, t) := (2\pi\varepsilon)^{-3} \int_{\mathbb{R}^3} e^{-i\xi \cdot y} \psi^\varepsilon(x + \frac{\varepsilon y}{2}, t) \overline{\psi^\varepsilon(x - \frac{\varepsilon y}{2}, t)} dy. \quad (2.21)$$

Note that some authors use the opposite sign in the exponential or a different normalization. More generally, we define the *Wigner matrix*  $F^\varepsilon$  (cf. [8]) as the Wigner transform of  $\psi^\varepsilon \otimes \psi^\varepsilon$ , i.e.

$$F^\varepsilon(x, \xi, t) = (2\pi\varepsilon)^{-3} \int_{\mathbb{R}^3} e^{-i\xi \cdot y} \psi^\varepsilon(x + \frac{\varepsilon y}{2}, t) \otimes \overline{\psi^\varepsilon(x - \frac{\varepsilon y}{2}, t)} dy,$$

Note that

$$f^\varepsilon = \text{Tr } F^\varepsilon \quad (2.22)$$

where  $\text{Tr } F^\varepsilon$  denotes the trace of the  $2 \times 2$  matrix  $F^\varepsilon$ . A simple calculation shows that

$$\rho^\varepsilon = \int_{\mathbb{R}^3_\xi} f^\varepsilon d\xi. \quad (2.23)$$

The Pauli current density  $J^\varepsilon$  can be written as a first order moment of the Wigner transform, more precisely:

$$J^\varepsilon = \int \text{Tr}(\text{Re}(\sigma \cdot (\xi - A^\varepsilon) F^\varepsilon)) d\xi \quad (2.24)$$

where  $\text{Tr}$  denotes the  $2 \times 2$  matrix trace. After a straightforward calculation one obtains the self-consistent Pauli-Wigner equation for the Wigner matrix  $F^\varepsilon$ ,

$$\begin{aligned} \partial_t F^\varepsilon + \xi \cdot \nabla_x F^\varepsilon - \mathcal{F}_y[\beta[A^\varepsilon]] *_\xi \nabla_x F^\varepsilon - i \mathcal{F}_y[\delta[A^\varepsilon]] *_\xi (\xi F^\varepsilon) \\ + \frac{1}{2} \theta[(A^\varepsilon)^2] F^\varepsilon - \frac{\varepsilon}{2} \theta[\sigma \cdot B^\varepsilon] F^\varepsilon + \theta[V^\varepsilon] F^\varepsilon = 0, \\ -\Delta V^\varepsilon = \rho^\varepsilon, \\ -\Delta A^\varepsilon = J^\varepsilon \\ F^\varepsilon(x, \xi, 0) = F^{\varepsilon,0}(x, \xi), \end{aligned} \quad (2.25)$$

where  $\theta[\cdot]$  is the pseudo-differential operator defined by

$$(\theta[\cdot] f^\varepsilon)(x, \xi, t) := \frac{i}{(2\pi)^3} \int_{\mathbb{R}^6} \delta[\cdot](x, y, t) f^\varepsilon(x, \eta, t) e^{-i(\xi - \eta) \cdot y} d\eta dy. \quad (2.26)$$

where

$$\beta[g] := \frac{1}{2} (g(x + \frac{\varepsilon y}{2}) + g(x - \frac{\varepsilon y}{2})),$$

and

$$\delta[g] := \frac{1}{\varepsilon} (g(x + \frac{\varepsilon y}{2}) - g(x - \frac{\varepsilon y}{2})).$$

The initial Wigner function  $f^{\varepsilon,0}$  is the Wigner transform of the initial wave function  $\psi^{\varepsilon,0}$ , i.e.

$$f^{\varepsilon,0}(x) = (2\pi\varepsilon)^{-3} \int_{\mathbb{R}^3} e^{-i\xi \cdot y} \psi^{\varepsilon,0}\left(x + \frac{\varepsilon y}{2}\right) \overline{\psi^{\varepsilon,0}\left(x - \frac{\varepsilon y}{2}\right)} dy. \quad (2.27)$$

The Wigner transform can attain negative values in general which corresponds to the uncertainty principle for conjugate variables position-momentum in quantum mechanics. Thus it can only be regarded as a "quasi-probability density" since it has the right moments (2.23) and (2.24). Formally passing  $\varepsilon \rightarrow 0$  in (2.25) yields

$$\partial_t F + p \cdot \nabla_x F - (\nabla_x V + p \times B) \cdot \nabla_p F = 0, \quad (2.28)$$

(where  $B = \nabla \times A$  and  $p = \xi + A$ ) and

$$-\Delta V(x, t) = \rho(x, t), \quad \rho(x, t) = \int_{\mathbb{R}^3_p} f(x, p, t) dp, \quad (2.29)$$

$$-\Delta A(x, t) = J(x, t), \quad J(x, t) = \int_{\mathbb{R}^3_p} p f(x, p, t) dp, \quad (2.30)$$

verifying the initial condition

$$f(x, p, 0) = f^0. \quad (2.31)$$

If  $\{\psi^\varepsilon\}$  is a bounded family in  $(L^2(\mathbb{R}^3))^2$  then  $f^\varepsilon$  has a weak limit  $f$  in  $\mathcal{A}'$  which is the dual to the test function space (in fact an algebra)  $\mathcal{A}$  defined by

$$\mathcal{A} := \{\phi \in C_0(\mathbb{R}_x^3 \times \mathbb{R}_\xi^3) : \mathcal{F}_\xi[\phi](x, \eta) \in L^1(\mathbb{R}_\eta^3, C_0(\mathbb{R}_x^3))\} \quad (2.32)$$

This space is specifically tailored to the Wigner transform and was introduced in [21]. One can also show (cf. [8]) that  $f^\varepsilon$  converges weakly to  $f$  as a tempered distribution, i.e. one can also work in  $\mathcal{S}'$  instead of  $\mathcal{A}'$ . In any case one then shows that the limit  $f$  is a non-negative Radon measure, called *Wigner measure* and can be interpreted as a classical phase-space density obeying the Vlasov-Poisson equation (2.28)-(2.31). It is related to the semiclassical measures introduced in [9]. If we now take the WKB initial data (2.1) and assume that  $\rho^{\varepsilon,0} = |a^{\varepsilon,0}|^2$  and  $u^{\varepsilon,0}$  converge to  $\rho^0$  and  $u^0$  in a sense to be clarified later we observe that the corresponding Wigner measure is

$$f_0(x, p, t) = \rho^0(x, t) \delta(p - u^0(x, t)), \quad (2.33)$$

If we use this as initial data for (2.28) we can expect that for short times,  $f(x, p, t)$  is of the form

$$f(x, p, t) = \rho(x, t) \delta(p - u(x, t)). \quad (2.34)$$

It will be shown that  $f^\varepsilon$  converges weakly in  $\mathcal{A}'$  to (2.34). We will closely follow an argument by Zhang [35]. This will be a consequence of our analysis of the energy of the Pauli-Poisson-WKB equation (2.5).

### 2.3 Main result

Define the normed space  $X^s$  by

$$X^s := H^{s-1}(\Omega, (\mathbb{C})^2) \times H^s(\Omega, \mathbb{R}^3), \quad \|(a, u)\|_{X^s} := \|a\|_{H^{s-1}} + \|u\|_{H^s} \quad (2.35)$$

We also frequently write  $\|(a, u)\|_{X^s}$  instead of  $\|(a, u)(t)\|_{X^s}$  for simplicity.

**Remark 4.** In this paper we only consider the three dimensional case. However we can generalize our results to cases in higher dimensions. In  $d \geq 4$  we just need to replace the assumption  $s > 7/2$  by  $s > d/2 + 2$  (due to Sobolev's embedding) and all the results still hold true. However note that the Poisson equation for  $V^\varepsilon$  is then no longer equivalent to the Hartree term.

The following two theorems are our main result. The first theorem is about the wellposedness and blow up of the Euler-Poisswell equation (2.12) and the Pauli-Poisswell-WKB equation (2.5). Theorem 1.a will be proved in Proposition 2, Theorem 1.b will be proved in Proposition 3.

#### Theorem 1.a. Local wellposedness of Euler-Poisswell and Pauli-Poisswell

Let  $\psi^{\varepsilon,0} \in (L^2(\mathbb{R}^3), \mathbb{C})^2$  be an initial 2-spinor of the form (2.1), i.e.

$$\psi_j^{\varepsilon,0}(x) = a_j^{\varepsilon,0}(x) e^{\frac{i}{\varepsilon} S^{\varepsilon,0}(x)}, \quad j = 1, 2.$$

. Let  $s > 7/2$  and let

$$\|u^{\varepsilon,0}\|_{H^s} + \|a^{\varepsilon,0}\|_{H^s} + \|u^0\|_{H^s} + \|a^0\|_{H^s} \leq Q,$$

where  $Q > 0$  is independent of  $\varepsilon$ . Assume that  $(a^{\varepsilon,0}, u^{\varepsilon,0})$  converges to  $(a^0, u^0)$  in  $X^{s-2}$ . Then:

- (i) There exists a unique local solution  $(a, u) \in C([0, T^0], H^{s-1}) \times C([0, T^0], H^s)$  to the Euler-Poisswell equation (2.12) with initial data  $(a^0, u^0) \in H^s \times H^{s-1}$ . Here  $T^0 > 0$  is the maximal (possibly infinite) time of existence.
- (ii) There exists a unique local solution  $(a^\varepsilon, u^\varepsilon) \in C([0, T^\varepsilon], H^s) \times C([0, T^\varepsilon], H^s)$  to the Pauli-Poisswell-WKB equation (2.5) with initial data  $(a^{\varepsilon,0}, u^{\varepsilon,0}) \in H^s \times H^s$ . Here  $T^\varepsilon > 0$  is the maximal (possibly infinite) time of existence.
- (iii) There exists  $\varepsilon_0 > 0$  such that  $T^\varepsilon$  has a uniform positive lower bound  $T$  for  $0 < \varepsilon < \varepsilon_0$ . In other words, there exist  $\varepsilon_0 > 0$  and  $T > 0$ , such that

$$\inf_{0 < \varepsilon < \varepsilon_0} T^\varepsilon > T. \quad (2.36)$$

#### Theorem 1.b. Blow up alternative for Euler-Poisswell and Pauli-Poisswell

Under the assumptions of Theorem 1.a, either the time of existence is global, or finite time blow up occurs. In the latter case it holds that:

- (i) If  $T^0$  is finite, then the solution  $(a, u)$  of (2.12) blows up at  $T^0$  such that

$$\lim_{t \rightarrow T^0-} (\|a(t)\|_{L^\infty} + \|u(t)\|_{W^{1,\infty}}) = \infty.$$

(ii) If  $T^\varepsilon$  is finite, then the solution  $(a^\varepsilon, u^\varepsilon)$  of (2.5) blows up at  $T^\varepsilon$  such that

$$\lim_{t \rightarrow T^\varepsilon -} (\|a^\varepsilon(t)\|_{H^1} + \|a^\varepsilon(t)\|_{W^{1,\infty}} + \|a^\varepsilon(t)\|_{W^{2,3}} + \|u^\varepsilon(t)\|_{W^{1,\infty}}) = \infty.$$

Furthermore, although  $\|u^\varepsilon(t)\|_{W^{1,\infty}} + \|a^\varepsilon(t)\|_{L^\infty}$  may not blow up at  $T^\varepsilon$  like in the Euler-Poisson equation, we can still show the following nearly singular behavior: For  $\varepsilon$  small enough, there exists a  $K = K(\varepsilon)$ , such that

$$\limsup_{t \rightarrow T^\varepsilon -} \|u^\varepsilon(t)\|_{W^{1,\infty}} + \|a^\varepsilon(t)\|_{L^\infty} \geq K.$$

Here,  $K$  goes to infinity as  $\varepsilon$  goes to zero.

The second result is about the semiclassical limit.

**Theorem 2. Semiclassical limit of Pauli-Poisson-WKB**

Let  $T^0$  and  $T^\varepsilon$  be the maximal times of existence of the Euler-Poisson equation (2.12) and the Pauli-Poisson-WKB equation (2.5). Under the assumptions of Theorem 1.a we have:

(i) Suppose that  $T < T^0$  is a lower bound of  $T^\varepsilon$  satisfying (2.36). We have the following semiclassical limit

$$\|(a^\varepsilon - a, u^\varepsilon - u)\|_{L^\infty([0,T], X^{s-2})} \xrightarrow{\varepsilon \rightarrow 0} 0. \quad (2.37)$$

where  $X^s$  is defined by (2.35).

(ii) For all  $t < T$  the Wigner transform  $f^\varepsilon$  satisfies

$$f^\varepsilon(x, \xi, t) \xrightarrow{\varepsilon \rightarrow 0} f(x, p, t) = \rho(x, t) \delta(p - u(x, t)) \text{ in } \mathcal{A}'$$

where  $\mathcal{A}'$  is the dual of  $\mathcal{A}$  defined by (2.32). In particular,  $f$  is the solution of (2.28) with initial data

$$f(0, x, \xi) = \rho^0(x, t) \delta(\xi - u^0(x, t)). \quad (2.38)$$

(iii) For the density  $\rho^\varepsilon$  and the current density  $J^\varepsilon$  it holds that

$$\begin{aligned} \rho^\varepsilon &= |a^\varepsilon|^2 \xrightarrow{\varepsilon \rightarrow 0} \rho = |a|^2 && \text{in } L^\infty([0, T], H^{s-3}(\mathbb{R}^3)), \\ J^\varepsilon &= \varepsilon w^\varepsilon - \varepsilon v^\varepsilon - \rho^\varepsilon(u^\varepsilon + A^\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} J = -(\rho u + \rho A) && \text{in } L^\infty([0, T], H^{s-3}(\mathbb{R}^3)), \end{aligned}$$

where  $A$  is given by  $-\Delta A = -(\rho u + \rho A)$ .

(iv) If  $s \geq 6$ , then the asymptotic behavior of the maximal time of existence  $T^\varepsilon$  satisfies

$$\liminf_{t \rightarrow T^\varepsilon -} T^\varepsilon \geq T^0. \quad (2.39)$$

Theorem 2 will be proved in Section 6.

**Remark 5.** As mentioned before the WKB ansatz only works locally in time. In order to obtain global results one can use the Wigner transform which we introduced in 2.2. In this paper we work with a special type of initial data for the Wigner function which is adapted to the WKB ansatz, i.e. the monokinetic initial data (cf. (2.33)) which leads to a monokinetic Wigner measure (2.34) obeying the Vlasov-Poisson equation in the distributional sense for a finite time  $T$ . For general initial data with a certain amount of regularity there is a downside: convergence is usually only weak and the limit equation only holds in the sense of distributions. For the linear case a general survey can be found in [8]. The semiclassical limit of the Schrödinger-Poisson equation has been proven in [21] and [22]. The global-in-time semiclassical limit of the Pauli-Poisson system using Wigner methods is ongoing work [24].

**Remark 6.** We draw our inspiration from three different papers in which the semiclassical limit of different nonlinear Schrödinger equations was discussed. In [13], Grenier showed the semiclassical limit for the nonlinear Schrödinger equation with  $A^\varepsilon \equiv 0$  and the nonlinearity given by  $f(|\psi|^2)\psi$  with  $f \in C^\infty(\mathbb{R}_+, \mathbb{R})$ . In [35], Zhang established the semiclassical limit of the Wigner-Poisson equation and in [2], Alazard and Carles showed the semiclassical limit of the Schrödinger-Poisson equation with doping profile in  $d \geq 3$  dimensions.

### 3 A priori estimates

In this section we will prove an a priori estimate for (2.5). We will omit the  $\varepsilon$ -superscript in this section in order to maintain readability. For

$$a \in L^\infty([0, T], H^s(\Omega, \mathbb{C})), \quad u \in W^{1, \infty}([0, T], H^s(\Omega, \mathbb{R}^3))$$

we define the following functionals. Let  $\mu, \mu_1, \mu_2 > 0$ .

$$E_s(t) := \|(a, u)(t)\|_{X^s}, \tag{3.1}$$

$$E_s^\mu(t) := E_s(t) + \mu \varepsilon \|a(t)\|_{H^s}, \tag{3.2}$$

$$E_s^{\mu_1, \mu_2}(t) := E_s^{\mu_1}(t) + \mu_2 \|\partial_t u(t)\|_{H^{s-1}}, \tag{3.3}$$

$$M(t) := 1 + \|u(t)\|_{W^{1, \infty}} + \|a(t)\|_{L^\infty} + \varepsilon \|a(t)\|_{H^1} + \varepsilon \|a(t)\|_{W^{1, \infty}} + \varepsilon \|a(t)\|_{W^{2, 3}}, \tag{3.4}$$

$$N(t) := \sup_{0 \leq s \leq t} M(s). \tag{3.5}$$

Note that by Sobolev's inequality, when  $s > 7/2$ , we have

$$M(t) \lesssim E_s^\mu(t). \tag{3.6}$$

**Proposition 1.** *Suppose  $u$  and  $a$  solve (2.5). Then we have the following estimate*

$$E_s^1(t) \leq CN(t)^{2s+3} E_s^1(0) e^{CN(t)^{2s+3}t}. \tag{3.7}$$

$E_s^1(t)$  is defined by (3.2) with  $\mu = 1$ .  $C$  is some constant that only depends on  $s$ .

**Remark 7.** This proposition is very important for the following two reasons. First, it allows us to prove local wellposedness as well as establish the semiclassical limit  $\varepsilon \rightarrow 0$  of (2.5). In fact, the proof of Theorem 1.a is following the spirit of the proof of Proposition 1. Second, this proposition leads to the second part of Theorem 1.b simply by a bootstrap argument. (See Section 5).

The remainder of this section is devoted to the proof of this proposition which we split into four steps. The first step (Section 3.1) is based on charge conservation (i.e. the conservation of  $\|a\|_2$ ) and is similar to the proof of Lemma 2.2 in [35]. It yields the following estimate

$$\frac{d}{dt}E_s(t) \lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}) + \|\nabla V\|_{H^s}.$$

If  $\|A\|_{W^{1,\infty}}$ ,  $\|\nabla A\|_{H^s}$  and  $\|\nabla V\|_{H^s}$  would not appear on the RHS one could use Gronwall's inequality directly. Therefore we have to bound these terms: In the second step (Section 3.2) we will bound  $\|A\|_{W^{1,\infty}}$ ,  $\|\nabla A\|_{H^s}$  and  $\|\nabla V\|_{H^s}$  by  $M(t)$ ,  $E_s(t)$  and  $\varepsilon\|a\|_{H^s}$ . This is achieved by some standard elliptic estimates. We obtain

$$\frac{d}{dt}E_s(t) \lesssim M(t)^{2s+1}E_s^1(t).$$

in Lemma 4. Since  $E_s^1(t) = E_s(t) + \varepsilon\|a(t)\|_{H^s}$  contains  $\|a(t)\|_{H^s}$ , in Section 3.3 we will prove a bound for  $\varepsilon\|a\|_{H^s}$ . Combining it with Lemma 4, we can close the argument and prove Proposition 1.

### 3.1 Estimate of $E_s(t)$

In this part we will use charge conservation to obtain a first estimate of  $E_s(t)$ . By multiplying the first equation in (2.5) by  $\bar{a}_j$  it is easy to show that

$$\partial_t|a_j|^2 + (u + A) \cdot \nabla|a_j|^2 + |a_j|^2 \operatorname{div}(u + A) = \frac{i\varepsilon}{2}(\Delta a_j \bar{a}_j - \Delta \bar{a}_j a_j) + \frac{1}{2} \operatorname{Im}(\mathbf{B}_i^j a_i \bar{a}_i).$$

Sum with respect to  $j$  and obtain

$$\partial_t \rho + (u + A) \cdot \nabla \rho + \rho \operatorname{div}(u + A) = \frac{i\varepsilon}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i). \quad (3.8)$$

Integrating over  $\Omega$  and by parts yields

$$\frac{d}{dt}\|a\|_{L^2}^2 = 0, \quad (3.9)$$

which is the conservation of charge. In general we have  $\|a\|_{L^2} = 1$ . Combining (3.9) with (3.4), interpolation and Hölder's inequality yields the estimate

$$\|a\|_{L^p} \lesssim \|a\|_{L^2} + \|a\|_{L^\infty} \lesssim M(t), \quad 2 \leq p \leq +\infty, \quad (3.10)$$

where  $M(t)$  is defined by (3.4).



**Remark 8.** There are two key observations concerning the estimate above. The first point is that although we are not able to obtain good estimates for  $\nabla\rho$  we can use integration by parts to transfer the derivative from  $\rho$  to  $(u + A)$ . The second point is that the term  $(i\varepsilon/2)\Delta a$  cancels due to Green's second identity since  $\Delta a$  has constant coefficients.

Since we need higher regularity for  $a$  later we need to extend the argument above to  $\partial_x^\alpha a_j$  and  $\partial_x^\alpha u$  which can be done by similar reasoning. To clarify the analogy between the charge conservation and our estimate for  $\partial_x^\alpha a_j$ , we first show the following lemma, which is inspired by the two observations in Remark 8.

**Lemma 1.** *Consider the following equation,*

$$\partial_t a + (A + u) \cdot \nabla a = \frac{i\varepsilon}{2} \Delta a + F,$$

with  $a$  a complex-valued,  $u$  real vector-valued and  $F$  is a source term. Then,

$$\frac{d}{dt} \|a\|_{L^2}^2 = \int_{\Omega} \operatorname{div}(u + A)|a|^2 + 2 \int_{\Omega} \operatorname{Re}(F \cdot \bar{a}), \quad (3.11)$$

and

$$\frac{d}{dt} \|a\|_{L^2} \lesssim (M(t) + \|A\|_{W^{1,\infty}}) \|a\|_{L^2} + \|F\|_{L^2}. \quad (3.12)$$

hold. If the term  $(i\varepsilon/2)\Delta a$  is absent, the two conclusions still hold true.

*Proof.* Similar to (3.8), it is easy to see that

$$\partial_t |a|^2 + (u + A) \cdot \nabla |a|^2 = \frac{i\varepsilon}{2} (\Delta a \bar{a} - \Delta \bar{a} a) + 2 \operatorname{Re}(F \bar{a}).$$

Integrating on  $\Omega$  and by parts, we obtain (3.11). Again, by integrating by parts, we get rid of the terms  $\nabla a$  and  $\Delta a$ , for which we do not have good estimates. Noticing that

$$\begin{aligned} \frac{d}{dt} \|a\|_{L^2}^2 &= 2 \|a\|_{L^2} \frac{d}{dt} \|a\|_{L^2}, \\ \left| \int_{\Omega} \operatorname{Re}(F \cdot \bar{a}) \right| &\leq \|a\|_{L^2} \|F\|_{L^2}, \\ \left| \int_{\Omega} \operatorname{div}(u + A)|a|^2 \right| &\leq (\|\nabla A\|_{L^\infty} + \|u\|_{W^{1,\infty}}) \|a\|_{L^2}^2 \\ &\leq (M(t) + \|A\|_{W^{1,\infty}}) \|a\|_{L^2}^2. \end{aligned}$$

(3.12) immediately follows. □

Applying Lemma 1 to system (2.5), we have the following estimate.

**Lemma 2.** *Suppose  $u$  and  $a$  solve (2.5). Then we have the following estimate*

$$\frac{d}{dt} E_s(t) \lesssim (M(t) + \|A\|_{W^{1,\infty}}) (E_s(t) + \|\nabla A\|_{H^s}) + \|\nabla V\|_{H^s}. \quad (3.13)$$

*Proof.* Apply  $\partial_x^\alpha$  to the first equation of (2.5) for  $1 \leq n(\alpha) \leq s-1$  and find

$$\partial_t \partial_x^\alpha a_i + (u + A) \cdot \nabla \partial_x^\alpha a_i = \frac{i\varepsilon}{2} \Delta \partial_x^\alpha a_i - \mathbf{R}_i^c, \quad (3.14)$$

with

$$\begin{aligned} \mathbf{R}_i^c &= (\operatorname{div} \partial_x^\alpha((u + A)a_i) - (u + A) \cdot \nabla \partial_x^\alpha a_i) - \frac{1}{2} \partial_x^\alpha (a_i \operatorname{div}(u + A)) - \frac{i}{2} \partial_x^\alpha (\mathbf{B}_i^j a_j) \\ &:= \sum_{j=1}^3 F_j \end{aligned} \quad (3.15)$$

for  $i = 1, 2$ . It is easy to see that  $\mathbf{R}^c$  does not contain  $s$ -th order derivative of  $a_i$ . Then due to the Kato-Ponce inequality (Lemma 10), we have the following estimate for  $F_1$ :

$$\begin{aligned} \|F_1\|_2 &= \|\operatorname{div} \partial_x^\alpha((u + A)a_i) - (u + A) \cdot \nabla \partial_x^\alpha a_i\|_2 \\ &\lesssim \|a\|_{\dot{H}^{s-1}} \|\nabla(u + A)\|_{L^\infty} + \|a\|_{L^\infty} \|u + A\|_{\dot{H}^s}, \\ &\lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}). \end{aligned}$$

For  $F_2$  we have

$$\begin{aligned} \|F_2\|_2 &= \|\partial_x^\alpha (a_i \operatorname{div}(u + A))\|_2 \\ &\lesssim \|a\|_{\dot{H}^{s-1}} \|\nabla(u + A)\|_{L^\infty} + \|a\|_{L^\infty} \|u + A\|_{\dot{H}^s} \\ &\lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}). \end{aligned}$$

For  $F_3$  we have

$$\begin{aligned} \|F_3\|_2 &= \|\partial_x^\alpha (\mathbf{B}_i^j a_j)\|_2 \\ &\lesssim \|\mathbf{B}\|_{L^\infty} \|a\|_{H^{s-1}} + \|a\|_{L^\infty} \|\mathbf{B}\|_{H^{s-1}} \\ &\lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}). \end{aligned}$$

Here, we used the fact  $1 \leq n(\alpha) \leq s-1$ . In the estimate for the last term, we also used the following fact,

$$\|\mathbf{B}\|_{H^{s-1}} \lesssim \|\nabla A\|_{H^{s-1}}, \quad \|\mathbf{B}\|_{L^\infty} \lesssim \|\nabla A\|_{L^\infty}. \quad (3.16)$$

It follows that for  $\mathbf{R}_i^c$ , we have

$$\|\mathbf{R}_i^c\|_{L^2} \lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}). \quad (3.17)$$

For (3.14), we can apply estimate (3.12) in Lemma 1 to obtain

$$\frac{d}{dt} \|\partial_x^\alpha a_i\|_{L^2} \lesssim (M(t) + \|A\|_{W^{1,\infty}}) \|\partial_x^\alpha a_i\|_{L^2} + \|\mathbf{R}_i^c\|_{L^2},$$

Adding up with respect to  $i = 1, 2$  and plugging in (3.17), it follows that

$$\frac{d}{dt} \|\partial_x^\alpha a\|_{L^2} \lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}).$$

Summing for all  $1 \leq |\alpha| \leq s-1$  yields

$$\frac{d}{dt} \|a\|_{H^{s-1}} \lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}). \quad (3.18)$$

The estimate for  $\|u\|_{H^s}$  is similar and we obtain

$$\frac{d}{dt} \|u\|_{H^s} \lesssim (M(t) + \|A\|_{W^{1,\infty}})(E_s(t) + \|\nabla A\|_{H^s}) + \|\nabla V\|_{H^s}. \quad (3.19)$$

Combining (3.18) and (3.19) finishes the proof.  $\square$

### 3.2 Estimates for the Poisson equations

In (3.13) we need to eliminate  $\|\nabla A\|_{H^s}$  and  $\|\nabla V\|_{H^s}$  in order to use Gronwall's inequality. Thus in this part, we will estimate  $\|\nabla A\|_{H^s}$  and  $\|\nabla V\|_{H^s}$ . Recall that  $V$  and  $A$  are given by (2.6) and (2.7). The result is following.

**Lemma 3.** *Let  $s \geq 0$ . For  $V$  and  $A$  given by (2.6) and (2.7), we have the following estimates,*

$$\|\nabla V\|_{H^s} \lesssim M(t) \|a\|_{H^{s-1}}, \quad (3.20)$$

$$\|\nabla A\|_{H^s} \lesssim M(t)^{2s} E_s^1(t), \quad (3.21)$$

$$\|A\|_{W^{1,\infty}} \lesssim M(t)^5. \quad (3.22)$$

where  $E_s^1(t)$  is given by (3.2) with  $\mu = 1$  and  $M(t)$  is given by (3.4).

**Remark 9.** Unfortunately, we have no estimate for  $\|A\|_{L^2}$ , which means we have to avoid using  $\|A\|_{L^2}$  to bound other terms in the analysis.

*Proof.* We divide the proof into three steps.

*Step 1: Estimate of  $\|\nabla V\|_{H^s}$ .* By definition, we have

$$\|\nabla V\|_{H^s} = \|\nabla V\|_{L^2} + \|\nabla^2 V\|_{H^{s-1}}$$

For  $\|\nabla V\|_{L^2}$ , we need the help of Sobolev inequality and then use the  $W^{2,p}$  estimate for the Poisson equation. For  $\|\nabla^2 V\|_{H^{s-1}}$ , we use the  $W^{2,p}$  estimate directly. Sobolev's inequality gives

$$\|\nabla V\|_{L^2} \lesssim \|\nabla^2 V\|_{L^{6/5}}.$$

By Lemma 11 and Hölder's inequality, we have

$$\|\nabla^2 V\|_{L^{6/5}} \lesssim \| |a|^2 \|_{L^{6/5}} \lesssim (\|a\|_{L^2}^{5/6} \|a\|_{L^\infty}^{1/6})^2 \lesssim M(t) \|a\|_{L^2}. \quad (3.23)$$

Applying Lemma 11 to (2.6), we get

$$\|\nabla^2 V\|_{H^{s-1}} \lesssim \| |a|^2 \|_{H^{s-1}} \lesssim \|a\|_{H^{s-1}} \|a\|_{L^\infty} \lesssim M(t) \|a\|_{H^{s-1}}. \quad (3.24)$$

In the second step we used Lemma 10. From (3.2) and (3.23), we get

$$\|\nabla V\|_{L^2} \lesssim M(t)\|a\|_{L^2}.$$

Combining the equation above and (3.24), we get (3.20).

*Step 2: Estimate of  $\|A\|_{W^{1,\infty}}$ .* By Lemma 9, we know that

$$\|A\|_{L^\infty} \lesssim \|A\|_{L^6}^{\alpha_1} \|A\|_{\dot{W}^{2,6}}^{1-\alpha_1}, \quad \|\nabla A\|_{L^\infty} \lesssim \|A\|_{L^6}^{\alpha_2} \|A\|_{\dot{W}^{2,6}}^{1-\alpha_2}.$$

For some  $\alpha_1, \alpha_2 \in [0, 1]$ . Then  $\|A\|_{W^{1,\infty}}$  is bounded by

$$\|A\|_{W^{1,\infty}} \lesssim \|A\|_{L^6} + \|A\|_{\dot{W}^{2,6}}$$

From (1.13), (2.11) and Lemma 11, we know that

$$\begin{aligned} \|A\|_{\dot{W}^{2,6}} &\lesssim \|J\|_{L^6} \\ &\lesssim \varepsilon\|w\|_{L^6} + \varepsilon\|v\|_{L^6} + \|\rho u\|_{L^6} + \|\rho A\|_{L^6}. \end{aligned}$$

Recalling (2.9) and (2.10), with the help of Hölder inequality, it follows that

$$\begin{aligned} \|A\|_{\dot{W}^{2,6}} &\lesssim \varepsilon\|a\|_{L^6} \|a\|_{W^{1,\infty}} + \|\rho\|_{L^6} \|u\|_{L^\infty} + \|\rho\|_{L^\infty} \|A\|_{L^6} \\ &= \varepsilon\|a\|_{L^6} \|a\|_{W^{1,\infty}} + \|a\|_{L^{12}}^2 \|u\|_{L^\infty} + \|a\|_{L^\infty}^2 \|A\|_{L^6} \\ &\lesssim M(t)^2(1 + \|A\|_{L^6}). \end{aligned}$$

In the last step, we use (3.10) to bound the norms of  $a$  and  $u$ . By the Sobolev inequality,

$$\|A\|_{L^6} \lesssim \|\nabla A\|_{L^2}, \quad (3.25)$$

which leads to

$$\|\nabla A\|_{L^2} \lesssim \|\varepsilon(v - w) + \rho u\|_{L^{6/5}}.$$

Thus  $\|A\|_{W^{1,\infty}}$  is bounded by

$$\|A\|_{W^{1,\infty}} \lesssim M(t)^2(1 + \|\varepsilon(v - w) + \rho u\|_{L^{6/5}}).$$

Since

$$\begin{aligned} \|\varepsilon(v - w) + \rho u\|_{L^{6/5}} &\leq \|\rho u\|_{L^{6/5}} + \varepsilon\|v - w\|_{L^{6/5}} \\ &\lesssim \|a\|_{L^{6/5}}^2 \|u\|_{L^\infty} + \varepsilon\|a\|_{L^3} \|\nabla a\|_{L^2} \\ &\lesssim M(t)^3. \end{aligned}$$

we can conclude.

*Step 3: Estimate of  $\|\nabla A\|_{H^s}$ .* Multiplying (1.13) by  $A$  and integrating by parts yields

$$\begin{aligned} \|\nabla A\|_{L^2}^2 &= \int_{\Omega} (\varepsilon(v - w) + \rho(u - A))A dx \leq \int_{\Omega} (\varepsilon(v - w) + \rho u)A dx \\ &\lesssim \|\varepsilon(v - w) + \rho u\|_{L^{6/5}} \|A\|_{L^6}. \end{aligned}$$

Recalling the Sobolev inequality (3.25) and the definition of  $v$  and  $w$  via (2.9) and (2.10), by Hölder inequality, we can get

$$\begin{aligned}\|\varepsilon(v-w) + \rho u\|_{L^{6/5}} &\leq \|\rho u\|_{L^{6/5}} + \varepsilon\|v-w\|_{L^{6/5}} \\ &\lesssim \| |a|^2 \|_{L^3} \|u\|_{L^2} + \varepsilon \|a\|_{L^3} \|\nabla a\|_{L^2} \\ &\lesssim M(t)^2 E_s^1(t).\end{aligned}$$

In the last step we used (3.10). It follows that

$$\|\nabla A\|_{L^2} \lesssim M(t)^2 E_s^1(t). \quad (3.26)$$

We obtain the estimate of  $\|\nabla A\|_{L^2}$ . The next step is to estimate  $\|\nabla A\|_{H^s}$ . For  $s \geq 1$ , thanks to Lemma 11, we know that

$$\begin{aligned}\|\nabla A\|_{H^s} &\lesssim \|\nabla^2 A\|_{H^{s-1}} + \|\nabla A\|_{L^2} \\ &\lesssim \varepsilon\|v-w\|_{H^{s-1}} + \|\rho u\|_{H^{s-1}} + \|\rho A\|_{H^{s-1}} + \|\nabla A\|_{L^2}.\end{aligned}$$

$\|\nabla A\|_{L^2}$  is bounded by (3.26).  $\varepsilon\|v-w\|_{H^{s-1}}$  and  $\|\rho u\|_{H^{s-1}}$  are bounded by

$$\begin{aligned}\varepsilon\|v-w\|_{H^{s-1}} &\lesssim \varepsilon\|a\|_{L^\infty} \|\nabla a\|_{H^{s-1}} \\ &\lesssim M(t) E_s^1(t),\end{aligned} \quad (3.27)$$

and

$$\begin{aligned}\|\rho u\|_{H^{s-1}} &\lesssim \|\rho\|_{L^\infty} \|u\|_{H^{s-1}} + \|u\|_{L^\infty} \|\rho\|_{H^{s-1}} \\ &\lesssim \|a\|_{L^\infty}^2 \|u\|_{H^{s-1}} + \|u\|_{L^\infty} \|a\|_{L^\infty} \|a\|_{H^{s-1}} \\ &\lesssim M(t)^2 E_s^1(t).\end{aligned} \quad (3.28)$$

We still need to estimate  $\|\rho A\|_{H^{s-1}}$  and the key point is that we should avoid  $\|A\|_{L^2}$ , for which we have no upper bounds. Since

$$\begin{aligned}\|\rho A\|_{H^{s-1}} &\lesssim \sum_{n(\alpha)+n(\beta)\leq s-1, n(\beta)\geq 1} \|\partial^\alpha \rho \partial^\beta A\|_{L^2} + \sum_{n(\alpha)=s-1} \|\partial^\alpha \rho A\|_{L^2} \\ &\lesssim \|\rho \nabla A\|_{H^{s-2}} + \sum_{n(\alpha)=s-1} \|\partial^\alpha \rho\|_{L^2} \|A\|_{L^\infty},\end{aligned}$$

by Lemma 10,  $\|\rho A\|_{H^{s-1}}$  is given by

$$\begin{aligned}\|\rho A\|_{H^{s-1}} &\lesssim \|\nabla A\|_{H^{s-2}} \|\rho\|_{L^\infty} + \|\nabla A\|_{L^\infty} \|\rho\|_{H^{s-2}} + \|\rho\|_{H^{s-1}} \|A\|_{L^\infty} \\ &\lesssim \|\nabla A\|_{H^{s-2}} \|a\|_{L^\infty}^2 + \|A\|_{W^{1,\infty}} \|a\|_{L^\infty} \|a\|_{H^{s-1}}.\end{aligned}$$

By the definition of the  $H^s$  norm via the Fourier transform and Hölder's inequality, we obtain

$$\|\nabla A\|_{H^{s-2}} \leq \|\langle \xi \rangle^{s-2} \widehat{\nabla A}^{\frac{s-2}{s-1}}\|_{\frac{2(s-1)}{s-2}} \|\widehat{\nabla A}^{\frac{1}{s-1}}\|_{2(s-1)} \lesssim \|\nabla A\|_{H^{s-1}}^{\frac{s-2}{s-1}} \|\nabla A\|_2^{\frac{1}{s-1}}.$$

Then we get

$$\|\rho A\|_{H^{s-1}} \lesssim \|\nabla A\|_{H^{s-1}}^{\frac{s-2}{s-1}} \|\nabla A\|_{L^2}^{\frac{1}{s-1}} \|a\|_{L^\infty}^2 + \|A\|_{W^{1,\infty}} \|a\|_{L^\infty} \|a\|_{H^{s-1}}. \quad (3.29)$$

Plugging (3.27),(3.28) and (3.29) into (3.2) yields

$$\begin{aligned} \|\nabla A\|_{H^s} &\lesssim \varepsilon \|v - w\|_{H^{s-1}} + \|\rho u\|_{H^{s-1}} + \|\nabla A\|_{L^2} \\ &+ \|\nabla A\|_{H^{s-1}}^{\frac{s-2}{s-1}} \|\nabla A\|_{L^2}^{\frac{1}{s-1}} \|a\|_{L^\infty}^2 + \|A\|_{W^{1,\infty}} \|a\|_{L^\infty} \|a\|_{H^{s-1}} \\ &\lesssim \varepsilon \|v - w\|_{H^{s-1}} + \|\rho u\|_{H^{s-1}} + (1 + \|a\|_{L^\infty}^{2(s-1)}) \|\nabla A\|_{L^2} + \|A\|_{W^{1,\infty}} \|a\|_{L^\infty} \|a\|_{H^{s-1}} \\ &\lesssim M(t) (\|A\|_{W^{1,\infty}} + M(t)) E_s^1(t) + M(t)^{2s-2} \|\nabla A\|_{L^2}. \end{aligned}$$

In the second step we used the following fact (which is just Young's inequality for products),

$$C_1 \|\nabla A\|_{H^{s-1}}^{\frac{s-2}{s-1}} (\|\nabla A\|_{L^2} \|a\|_{L^\infty}^{2(s-1)})^{\frac{1}{s-1}} \leq \frac{1}{2} \|\nabla A\|_{H^{s-1}} + C_2 \|\nabla A\|_{L^2} \|a\|_{L^\infty}^{2(s-1)}.$$

Moreover, we absorbed  $(1/2)\|\nabla A\|_{H^{s-1}}$  into the LHS by the trivial estimate  $\|\nabla A\|_{H^{s-1}} \leq \|\nabla A\|_{H^s}$ . Then we can apply (3.26) and get

$$\|\nabla A\|_{H^s} \lesssim M(t) (\|A\|_{W^{1,\infty}} + M(t)) E_s^1(t) + M(t)^{2s} E_s^1(t). \quad (3.30)$$

To finish the proof of (3.21), we only need (3.22).  $\square$

Combining Lemma 2 and Lemma 3 leads to

**Lemma 4.** *Suppose  $u$  and  $a$  solve (2.5). Then we have the following estimate*

$$\frac{d}{dt} E_s(t) \lesssim M(t)^{2s+1} E_s^1(t), \quad (3.31)$$

where  $E_s(t)$  is defined by (3.1).

### 3.3 Estimate of $\varepsilon \|a\|_{H^s}$ in $E_s^1(t)$

To apply Gronwall's inequality to (3.31) we need to estimate the term  $\varepsilon \|a\|_{H^s}$  in  $E_s^1(t)$ . We have the following lemma.

**Lemma 5.** *Suppose  $S$  and  $a$  solve (2.4). Then we have the following estimate. For any vector  $\alpha$  with  $n(\alpha) = s$ ,*

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim M(t)^{2s+2} E_s^1(t). \quad (3.32)$$

where  $E_s^1(t)$  is defined by (3.2).

Lemma 5 in combination with Lemma 4 will yield Proposition 1 which will be shown in the last part of this section. In the following section we will motivate Lemma 5 by an analogous result for the Schrödinger-Poisson equation.

### 3.3.1 Motivation of Lemma 5

Consider the case  $A \equiv 0$ . Then (1.11) becomes

$$i\varepsilon\partial_t\psi = -\frac{\varepsilon^2}{2}\Delta\psi + V\psi, \quad (3.33)$$

where  $\psi = (\psi_1, \psi_2)$ . By abuse of notation we use  $\psi$  and all other original notations for the respective quantities of the Schrödinger-Poisson equation. For the Schrödinger-Poisson equation the WKB equation (2.5) becomes

$$\begin{aligned} \partial_t a_j + u \cdot \nabla a_j + \frac{1}{2}a_j \operatorname{div} u &= \frac{i\varepsilon}{2}\Delta a_j \\ \partial_t u + u \cdot \nabla u + \nabla V &= 0 \\ a_j(x, 0) &= a_j^{\varepsilon, 0}(x), \\ u(x, 0) &= u^{\varepsilon, 0}(x). \end{aligned} \quad (3.34)$$

While  $\rho$  and  $w$  are still defined by (1.12) and (2.10), the current density  $J$  becomes

$$J = -\operatorname{Re}(\bar{\psi}(-i\varepsilon\nabla)\psi),$$

or equivalently

$$J = \varepsilon w - \rho u. \quad (3.35)$$

Then (3.8) becomes

$$\partial_t \rho + \operatorname{div}(\rho u) = \frac{i\varepsilon}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i). \quad (3.36)$$

Then with the help of energy conservation, we will show the following lemma.

**Lemma 6.** *Suppose  $u$  and  $a$  solve (3.34). Then we have the following equation*

$$-2\varepsilon \int_{\Omega} u \cdot \partial_t w dx + \varepsilon^2 \frac{d}{dt} \|\nabla a\|_{L^2}^2 = 0. \quad (3.37)$$

**Remark 10.** To see why Lemma 6 motivates Lemma 5, replace  $u = \nabla S$  by  $\partial^\alpha S$  and  $\nabla a$  by  $\partial^\alpha a$  on the LHS of (3.37) where  $\alpha$  is some vector satisfying  $n(\alpha) = s$ . Recall that  $w$  is defined by (2.10). Then  $w$  should be replaced by  $\frac{i}{2} \sum_{j=1}^2 (\bar{a}_j \partial^\alpha a_j - a_j \partial^\alpha \bar{a}_j)$ . The LHS of (3.37) becomes

$$\varepsilon^2 \frac{d}{dt} \|\partial^\alpha a\|_{L^2}^2 - i\varepsilon \int_{\Omega} \partial^\alpha S \cdot \partial_t \sum_{j=1}^2 (\bar{a}_j \partial^\alpha a_j - a_j \partial^\alpha \bar{a}_j) dx. \quad (3.38)$$

This is exactly the LHS of (3.32) in Lemma 5. In fact, inspired by Lemma 6, we tried to estimate (3.38) and finally obtained Lemma 5. The difficulty is now that for the Pauli-Poisson equation,  $A \neq 0$  and we have to modify our calculations in the proof of Lemma 6.

*Proof.* The proof is split into three steps. First, we show that the energy of the Schrödinger-Poisson equation (3.33) is conserved and obtain (3.39). Then we will use the WKB system (2.5) to obtain a similar conclusion (3.40). Finally we combine (3.39) and (3.40) and finish the proof.

*Step 1.* First let us show energy conservation. Let

$$E(t) = \| -i\varepsilon\nabla\psi \|_{L^2}^2 + \|\nabla V\|_{L^2}^2,$$

denote the energy of the Schrödinger-Poisson equation (3.33). Multiplying (3.33) with  $\partial_t\bar{\psi}$  and taking the real part, we will get

$$2\operatorname{Re}\left(\partial_t\bar{\psi}\left[\frac{1}{2}(-i\varepsilon\nabla)^2 + V\right]\psi\right) = 0.$$

Using  $2\operatorname{Re}(V\partial_t\bar{\psi}\psi) = V\partial_t|\psi|^2$  yields

$$0 = \operatorname{Re}(\partial_t\bar{\psi}(-i\varepsilon\nabla)^2\psi) + V\partial_t|\psi|^2.$$

Now,  $-\Delta V = |\psi|^2$  and integration by parts yields

$$\begin{aligned} 0 &= \int_{\Omega} \operatorname{Re}(\partial_t\bar{\psi}(-i\varepsilon\nabla)^2\psi)dx + \int_{\Omega} V\partial_t|\psi|^2dx \\ &= \operatorname{Re} \int_{\Omega} \partial_t(-i\varepsilon\nabla\psi)^*(-i\varepsilon\nabla\psi)dx + \int_{\Omega} \partial_t(\nabla V) \cdot \nabla V dx \\ &= \frac{1}{2}\partial_t \left( \int_{\Omega} |\varepsilon\nabla\psi|^2 dx + \int_{\Omega} |\nabla V|^2 dx \right) \\ &= \frac{1}{2}\partial_t E(t). \end{aligned}$$

We can plug our ansatz (2.2) into  $E(t)$  and it follows that

$$E(t) = \int_{\Omega} \rho|u|^2 dx + \|\nabla V\|_{L^2}^2 + 2\varepsilon \operatorname{Im} \int_{\Omega} \bar{a}(u \cdot \nabla)a dx + \varepsilon^2 \|\nabla a\|_{L^2}^2.$$

Recall  $w$  defined by (2.10). Then  $(d/dt)E(t) = 0$  becomes

$$\frac{d}{dt} \int_{\Omega} \rho|u|^2 dx + \frac{d}{dt} \|\nabla V\|_{L^2}^2 - 2\varepsilon \frac{d}{dt} \int_{\Omega} u \cdot w dx + \varepsilon^2 \frac{d}{dt} \|\nabla a\|_{L^2}^2 = 0. \quad (3.39)$$

*Step 2.* Now we consider the WKB system (3.34) and try to get another form of energy conservation. From (3.34), we know that

$$\partial_t|u|^2 + u \cdot \nabla|u|^2 + 2u \cdot \nabla V = 0.$$

Together with (3.36), we get

$$\partial_t\rho|u|^2 + \operatorname{div}(|u|^2\rho u) + 2\rho u \cdot \nabla V = \frac{i\varepsilon}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i) |u|^2.$$



Recalling the expression of  $J$  (3.35), we can rewrite the equation above as

$$\partial_t \rho |u|^2 + \operatorname{div}(|u|^2 \rho u) - 2J \cdot \nabla V + 2\varepsilon w \cdot \nabla V = \frac{i\varepsilon}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i) |u|^2.$$

Integrating on  $\Omega$ , it follows that

$$\partial_t \int_{\Omega} \rho |u|^2 dx - 2 \int_{\Omega} J \cdot \nabla V dx = \int_{\Omega} \left( -2\varepsilon w \cdot \nabla V + \frac{i\varepsilon}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i) |u|^2 \right) dx.$$

By (1.17), we obtain

$$\int_{\Omega} J \cdot \nabla V dx = - \int_{\Omega} \operatorname{div}(J)V dx = - \int_{\Omega} \partial_t \rho V dx = \int_{\Omega} \partial_t (\Delta V)V dx = - \frac{d}{dt} \|\nabla V\|_{L^2}^2.$$

Then (3.40) becomes

$$\frac{d}{dt} \int_{\Omega} \rho |u|^2 dx + \frac{d}{dt} \|\nabla V\|_{L^2}^2 = \varepsilon \int_{\Omega} \left( -2w \cdot \nabla V + \frac{i}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i) |u|^2 \right) dx. \quad (3.40)$$

This equation could be seen as an analogy of the energy conservation.

*Step 3.* Combining (3.40) with (3.39), we can easily see that

$$\varepsilon \int_{\Omega} \left( -2w \cdot \nabla V + \frac{i}{2} \sum_i (\Delta a_i \bar{a}_i - \Delta \bar{a}_i a_i) |u|^2 \right) dx - 2\varepsilon \frac{d}{dt} \int_{\Omega} u \cdot w dx + \varepsilon^2 \frac{d}{dt} \|\nabla a\|_{L^2}^2 = 0.$$

Integrating by parts, we will get

$$-2\varepsilon \int_{\Omega} w \cdot \left( \nabla V + \frac{1}{2} \nabla |u|^2 \right) dx - 2\varepsilon \frac{d}{dt} \int_{\Omega} u \cdot w dx + \varepsilon^2 \frac{d}{dt} \|\nabla a\|_{L^2}^2 = 0.$$

Since

$$\nabla V + \frac{1}{2} \nabla |u + A|^2 = -\partial_t u,$$

it follows that

$$2\varepsilon \int_{\Omega} w \cdot \partial_t u dx - 2\varepsilon \frac{d}{dt} \int_{\Omega} u \cdot w dx + \varepsilon^2 \frac{d}{dt} \|\nabla a\|_{L^2}^2 = 0.$$

This immediately leads to (3.37) and finishes the proof.  $\square$

### 3.3.2 Proof of Lemma 5

Now we return to the original system (1.11), which means that  $A \neq 0$ .

*Proof of Lemma 5.* We need to estimate two terms in Lemma 5 which we denote by  $I_1$  and  $I_2$ :

$$I_1 = -i\varepsilon \int_{\Omega} \partial^\alpha S \cdot \partial_t \sum_{j=1}^2 (\bar{a}_j \partial^\alpha a_j - a_j \partial^\alpha \bar{a}_j) dx,$$

and

$$I_2 = \varepsilon^2 \frac{d}{dt} \|\partial^\alpha a\|_{L^2}^2.$$

It turns out that when estimating  $I_1$  and  $I_2$  separately, two terms will cancel each other when adding  $I_1$  and  $I_2$ . This leads to an estimate for  $\|\partial^\alpha a\|_{L^2}^2$ .

**Estimate of  $I_2$ .** First, we rewrite (3.14) as

$$\partial_t \partial_x^\alpha a_i + (u + A) \cdot \nabla \partial_x^\alpha a_i + \frac{1}{2} a_i \partial_x^\alpha \operatorname{div}(u + A) + \mathbf{R}_a = \frac{i\varepsilon}{2} \Delta \partial_x^\alpha a_i, \quad (3.41)$$

with

$$\begin{aligned} \mathbf{R}_a &= (\partial_x^\alpha((u + A) \cdot \nabla a_i) - (u + A) \cdot \nabla \partial_x^\alpha a_i) - \frac{i}{2} \partial_x^\alpha (\mathbf{B}_i^j a_j) \\ &\quad + \frac{1}{2} (\partial_x^\alpha (a_i \operatorname{div}(u + A)) - a_i \partial_x^\alpha \operatorname{div}(u + A)). \end{aligned} \quad (3.42)$$

Applying (3.11) in Lemma 1 leads to

$$\begin{aligned} &\frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 + \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(a_i \partial_x^\alpha \operatorname{div}(u + A) \partial_x^\alpha \bar{a}_i) dx \\ &= \sum_{i=1}^2 \left( \int_{\Omega} |\partial_x^\alpha a_i|^2 \operatorname{div}(u + A) dx - \int_{\Omega} \operatorname{Re}(\mathbf{R}_a \partial_x^\alpha \bar{a}_i) dx \right). \end{aligned}$$

Using Lemma 10 and (3.16), (3.21) gives the following estimates. For the second term on the LHS of the equation above we have

$$\begin{aligned} \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(a_i \partial_x^\alpha \operatorname{div}(A) \partial_x^\alpha \bar{a}_i) dx &\lesssim \|a\|_{L^\infty} \|\partial_x^\alpha a\|_{L^2} \|\partial_x^\alpha \operatorname{div}(u + A)\|_{L^2} \\ &\lesssim M(t) \|a\|_{H^s} \|\nabla A\|_{H^s}, \\ &\lesssim \frac{M(t)^{2s+1}}{\varepsilon} E_s^1(t)^2. \end{aligned}$$

For the first term on the RHS we have

$$\begin{aligned} \left| \int_{\Omega} |\partial_x^\alpha a_i|^2 \operatorname{div}(u + A) dx \right| &\lesssim \|\partial_x^\alpha a\|_{L^2}^2 \|u + A\|_{W^{1,\infty}} \\ &\lesssim \frac{M(t)^5}{\varepsilon^2} E_s^1(t)^2. \end{aligned}$$

For the second term on the RHS we have to consider  $\mathbf{R}_a$ , cf. (3.42). We can write  $\int_{\Omega} \operatorname{Re}(\mathbf{R}_a \partial_x^\alpha \bar{a}_i) dx$  as

$$\begin{aligned} \int_{\Omega} \operatorname{Re}(\mathbf{R}_a \partial_x^\alpha \bar{a}_i) dx &= \int_{\Omega} \operatorname{Re}((\partial_x^\alpha((u+A) \cdot \nabla a_i) - (u+A) \cdot \nabla \partial_x^\alpha a_j) \partial_x^\alpha \bar{a}_i) dx \\ &\quad + \frac{1}{2} \int_{\Omega} \operatorname{Im}(\partial_x^\alpha (\mathbf{B}_i^j a_j) \partial_x^\alpha \bar{a}_i) dx \\ &\quad + \int_{\Omega} \operatorname{Re}((\partial_x^\alpha (a_i \operatorname{div}(u+A)) - a_i \partial_x^\alpha \operatorname{div}(u+A)) \partial_x^\alpha \bar{a}_i) dx \\ &:= \sum_{j=1}^3 F_j \end{aligned}$$

The estimate for  $F_1$  is given by

$$\begin{aligned} |F_1| &= \left| \int_{\Omega} \operatorname{Re}((\partial_x^\alpha((u+A) \cdot \nabla a_i) - (u+A) \cdot \nabla \partial_x^\alpha a_j) \partial_x^\alpha \bar{a}_i) dx \right| \\ &\lesssim (\|a\|_{H^s} \|u+A\|_{W^{1,\infty}} + \|u+A\|_{H^s} \|\nabla a\|_{L^\infty}) \|\partial_x^\alpha a\|_{L^2} \\ &\lesssim \frac{M(t)^{2s+1}}{\varepsilon^2} E_s^1(t)^2. \end{aligned}$$

For  $F_2$  we have

$$\begin{aligned} |F_2| &= \left| \int_{\Omega} \operatorname{Im}(\partial_x^\alpha (\mathbf{B}_i^j a_j) \partial_x^\alpha \bar{a}_i) dx \right| \\ &\lesssim (\|a\|_{H^s} \|\mathbf{B}\|_{L^\infty} + \|\mathbf{B}\|_{H^s} \|a\|_{L^\infty}) \|\partial_x^\alpha a\|_{L^2} \\ &\lesssim (\|a\|_{H^s} \|\nabla A\|_{L^\infty} + \|\nabla A\|_{H^s} \|a\|_{L^\infty}) \|\partial_x^\alpha a\|_{L^2} \\ &\lesssim \frac{M(t)^{2s+1}}{\varepsilon} E_s^1(t)^2. \end{aligned}$$

And for  $F_3$  we have

$$\begin{aligned} |F_3| &= \left| \int_{\Omega} \operatorname{Re}((\partial_x^\alpha (a_i \operatorname{div}(u+A)) - a_i \partial_x^\alpha \operatorname{div}(u+A)) \partial_x^\alpha \bar{a}_i) dx \right| \\ &\lesssim (\|a\|_{H^s} \|u+A\|_{W^{1,\infty}} + \|u+A\|_{H^s} \|\nabla a\|_{L^\infty}) \|\partial_x^\alpha a\|_{L^2} \\ &\lesssim \frac{M(t)^{2s+1}}{\varepsilon^2} E_s^1(t)^2, \end{aligned}$$

Applying the estimates above and multiplying by  $\varepsilon^2$  yields

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 + \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(a_i \partial_x^\alpha \operatorname{div}(u) \partial_x^\alpha \bar{a}_i) dx \lesssim M(t)^{2s+1} E_s^1(t)^2. \quad (3.43)$$

Since the term

$$\sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(a_i \partial_x^\alpha \operatorname{div}(u) \partial_x^\alpha \bar{a}_i) dx$$

contains a  $(s + 1)$ -th order derivative of  $u$ , we cannot apply Lemma 10. Otherwise, we will get  $\|u\|_{H^{s+1}}$ . But if we add (3.43) to the estimate of  $I_1$ , which we will show in the next step, this tricky term will cancel with another term in the estimate of  $I_1$ .

**Estimate of  $I_1$ .** Multiplying (3.41) by  $\bar{a}_i$ , we get

$$\bar{a}_i \partial_t \partial_x^\alpha a_i + \bar{a}_i (u + A) \cdot \nabla \partial_x^\alpha a_i + \frac{1}{2} |a_i|^2 \partial_x^\alpha \operatorname{div}(u + A) + \bar{a}_i \mathbf{R}_a = \frac{i\varepsilon}{2} \bar{a}_i \Delta \partial_x^\alpha a_i.$$

Multiplying the first equation in (2.5) by  $\partial_x^\alpha \bar{a}_i$  and complex conjugation yield

$$\partial_x^\alpha a_i \partial_t \bar{a}_i + \partial_x^\alpha a_i (A + u) \cdot \nabla \bar{a}_i + \frac{1}{2} \bar{a}_i \partial_x^\alpha a_i \operatorname{div}(u + A) = -\frac{i\varepsilon}{2} \Delta \bar{a}_i \partial_x^\alpha a_i - \frac{i}{2} \mathbf{B}_i^j \bar{a}_j \partial_x^\alpha a_i.$$

Adding the two equations above together yields

$$\partial_t (\bar{a}_i \partial_x^\alpha a_i) + (u + A) \cdot \nabla (\bar{a}_i \partial_x^\alpha a_i) + \frac{1}{2} |a_i|^2 \partial_x^\alpha \operatorname{div}(u + A) + \mathbf{R}_b = \frac{i\varepsilon}{2} \bar{a}_i \Delta \partial_x^\alpha a_i - \frac{i\varepsilon}{2} \Delta \bar{a}_i \partial_x^\alpha a_i,$$

where  $\mathbf{R}_b$  is defined as

$$\mathbf{R}_b := \bar{a}_i \mathbf{R}_a + \frac{1}{2} \bar{a}_i \partial_x^\alpha a_i \operatorname{div}(u + A) + \frac{i}{2} \mathbf{B}_i^j \bar{a}_j \partial_x^\alpha a_i.$$

Recall that  $\mathbf{R}_a$  was defined as

$$\begin{aligned} \mathbf{R}_a &= (\partial_x^\alpha ((u + A) \cdot \nabla a_i) - (u + A) \cdot \nabla \partial_x^\alpha a_i) - \frac{i}{2} \partial_x^\alpha (\mathbf{B}_i^j a_j) \\ &\quad + \frac{1}{2} (\partial_x^\alpha (a_i \operatorname{div}(u + A)) - a_i \partial_x^\alpha \operatorname{div}(u + A)). \end{aligned}$$

Now take the imaginary part and multiply with  $\partial_x^\alpha S$ :

$$\begin{aligned} &\partial_x^\alpha S \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) + \partial_x^\alpha S (u + A) \cdot \nabla \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) + \partial_x^\alpha S \operatorname{Im}(\mathbf{R}_b) \\ &= -\frac{\varepsilon}{2} \partial_x^\alpha S \operatorname{Re}(\bar{a}_i \Delta \partial_x^\alpha a_i - \Delta \bar{a}_i \partial_x^\alpha a_i). \end{aligned}$$

Integrate by parts:

$$\begin{aligned} &\int_\Omega \partial_x^\alpha S \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx - \int_\Omega \operatorname{div}(\partial_x^\alpha S (u + A)) \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx + \int_\Omega \partial_x^\alpha S \operatorname{Im}(\mathbf{R}_b) dx \\ &= -\frac{\varepsilon}{2} \int_\Omega \partial_x^\alpha S \operatorname{Re}(\bar{a}_i \Delta \partial_x^\alpha a_i - \Delta \bar{a}_i \partial_x^\alpha a_i) dx \\ &= -\frac{\varepsilon}{2} \int_\Omega \Delta \partial_x^\alpha S \operatorname{Re}(\bar{a}_i \partial_x^\alpha a_i) dx - \varepsilon \int_\Omega \nabla \partial_x^\alpha S \cdot \operatorname{Re}(\nabla \bar{a}_i \partial_x^\alpha a_i) dx \\ &= -\frac{\varepsilon}{2} \int_\Omega \partial_x^\alpha \operatorname{div}(u) \operatorname{Re}(\bar{a}_i \partial_x^\alpha a_i) dx - \varepsilon \int_\Omega \partial_x^\alpha u \cdot \operatorname{Re}(\nabla \bar{a}_i \partial_x^\alpha a_i) dx. \end{aligned}$$

The first term on the LHS is the term we want to estimate. The first term on the RHS is the term that will cancel with the tricky term in  $I_2$ . That means we have to estimate the remaining terms. For the second term on the LHS we have

$$\begin{aligned} \left| \int_{\Omega} \operatorname{div}(\partial_x^\alpha S(u+A)) \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \right| &\lesssim \|\partial_x^\alpha a_i\|_2 \|a_i\|_\infty \|\operatorname{div}(\partial_x^\alpha S(u+A))\|_2 \\ &\lesssim \|\partial_x^\alpha a_i\|_2 \|a_i\|_\infty \|\partial_x^\alpha S\|_{H^1} \|u+A\|_{W^{1,\infty}} \\ &\lesssim \frac{M(t)^6}{\varepsilon} E_s^1(t)^2. \end{aligned}$$

The next five inequalities give the estimate for the third term on the LHS, i.e.  $\int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\mathbf{R}_b) dx$ .

$$\begin{aligned} \left| \int_{\Omega} \partial_x^\alpha S \operatorname{Re}(\bar{a}_i \partial_x^\alpha (\mathbf{B}_i^k a_k) - \mathbf{B}_i^k \bar{a}_k \partial_x^\alpha a_i) dx \right| &\lesssim \|\partial_x^\alpha S\|_2 (\|a_i\|_\infty \|\partial_x^\alpha (\mathbf{B}_i^k a_k)\|_2 + \|\partial_x^\alpha a_i\|_2 \|\mathbf{B}_i^k a_k\|_\infty) \\ &\lesssim \|u\|_{H^s} \|a_i\|_\infty (\|\mathbf{B}\|_{H^s} \|a_i\|_\infty + \|a_i\|_{H^s} \|\mathbf{B}\|_\infty) \\ &\lesssim \|u\|_{H^s} \|a_i\|_\infty (\|\nabla A\|_{H^s} \|a_i\|_\infty + \|a_i\|_{H^s} \|\nabla A\|_\infty) \\ &\lesssim M(t)^{2s+2} \|u\|_{H^s} (\|u\|_{H^s} + \|a_i\|_{H^s}) \\ &\lesssim \frac{M(t)^{2s+2}}{\varepsilon} E_s^1(t)^2. \end{aligned}$$

$$\begin{aligned} \left| \int_{\Omega} \partial_x^\alpha S \operatorname{Re}(\Delta \bar{a}_i \partial_x^\alpha a_i) dx \right| &\lesssim \|\Delta a_i\|_{L^3} \|\partial_x^\alpha S\|_{L^6} \|\partial_x^\alpha a_i\|_2 \\ &\lesssim \|\Delta a_i\|_{L^3} \|\nabla \partial_x^\alpha S\|_2 \|\partial_x^\alpha a_i\|_2 \\ &\lesssim \frac{M(t)}{\varepsilon^2} E_s^1(t)^2. \end{aligned}$$

$$\begin{aligned} \left| \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) \operatorname{div}(u+A) dx \right| &\lesssim \|\operatorname{div}(u+A)\|_\infty \|a_i\|_\infty \|\partial_x^\alpha a_i\|_2 \|\partial_x^\alpha S\|_2 \\ &\lesssim \frac{M(t)^6}{\varepsilon} E_s^1(t)^2, \end{aligned}$$

$$\begin{aligned} &\left| \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i (\partial_x^\alpha (a_i \operatorname{div}(u+A)) - a_i \partial_x^\alpha \operatorname{div}(u+A))) dx \right| \\ &\lesssim \|\partial_x^\alpha S\|_2 \|a_i\|_\infty \|\partial_x^\alpha (a_i \operatorname{div}(u+A)) - a_i \partial_x^\alpha \operatorname{div}(u+A)\|_2 \\ &\lesssim \|u\|_{H^s} (\|a_i\|_{W^{1,\infty}} \|\operatorname{div}(u+A)\|_{H^{s-1}} + \|\operatorname{div}(u+A)\|_\infty \|a_i\|_{H^s}) \\ &\lesssim \frac{M(t)}{\varepsilon} \|u\|_{H^s} (\|u\|_{H^s} + \|\nabla A\|_{H^{s-1}}) + \varepsilon \|a_i\|_{H^s} \\ &\lesssim \frac{M(t)^{2s+1}}{\varepsilon} E_s^1(t)^2. \end{aligned}$$

$$\begin{aligned}
& \left| \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i (\partial_x^\alpha ((u+A) \cdot \nabla a_i) - (u+A) \cdot \nabla \partial_x^\alpha a_j)) dx \right| \\
& \lesssim \|\partial_x^\alpha S\|_2 \|a_i\|_\infty \|\partial_x^\alpha ((u+A) \cdot \nabla a_i) - (u+A) \cdot \nabla \partial_x^\alpha a_j\|_2 \\
& \lesssim \|\partial_x^\alpha S\|_2 (\|\nabla(u+A)\|_\infty \|\nabla a_i\|_{\dot{H}^{s-1}} + \|\nabla a_i\|_\infty \|u+A\|_{\dot{H}^s}) \\
& \lesssim \frac{M(t)^{2s+1}}{\varepsilon} E_s^1(t)^2.
\end{aligned}$$

For the last term on the RHS we have

$$\begin{aligned}
\left| \int_{\Omega} \partial_x^\alpha u \cdot \operatorname{Re}(\nabla \bar{a}_i \partial_x^\alpha a_i) dx \right| & \lesssim \|u\|_{H^s} \|\nabla a_i\|_\infty \|a_i\|_{H^s} \\
& \lesssim \frac{M(t)}{\varepsilon} \|u\|_{H^s} \|a\|_{H^s} \\
& \lesssim \frac{M(t)}{\varepsilon^2} E_s^1(t)^2.
\end{aligned}$$

Combining all estimates yields

$$-2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx - \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \operatorname{div}(u) \operatorname{Re}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim M(t)^{2s+2} E_s^1(t)^2. \quad (3.44)$$

Notice that the second term of the equation above will cancel with the second term of (3.43). We can use this observation to finish the estimate of  $\|a\|_{H^s}^2$ .

**Combining  $I_1$  and  $I_2$ .** Adding (3.43) and (3.44) immediately yields (3.32).  $\square$

### 3.4 Proof of Proposition 1

Now we will use Lemma 4 and Lemma 5 to derive the a priori estimate in Proposition 1.

*Proof of Proposition 1.* Notice that

$$\begin{aligned}
& \varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \\
& = \varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx + 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_t \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx.
\end{aligned}$$

First we estimate the third term. By the second equation in (2.4), we have

$$\begin{aligned}
2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_t \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx & \lesssim \varepsilon \|\partial_t \partial_x^\alpha S\|_{L^2} \|\partial_x^\alpha a\|_{L^2} \|a\|_{L^\infty} \\
& \lesssim \varepsilon \|\partial_t u\|_{H^{s-1}} \|a\|_{H^s} \|a\|_{L^\infty}.
\end{aligned}$$

Using

$$\begin{aligned}
\|\partial_t u\|_{H^{s-1}} &\lesssim \|(A+u) \cdot \nabla u\|_{H^{s-1}} + \|u \cdot \nabla A\|_{H^{s-1}} + \|\nabla |A|^2\|_{H^{s-1}} + \|\nabla V\|_{H^{s-1}} \\
&\lesssim (\|u\|_{L^\infty} + \|A\|_{L^\infty})(\|u\|_{H^s} + \|A\|_{\dot{H}^s}) + \|\nabla V\|_{H^{s-1}} \\
&\lesssim M(t)^{2s+1} E_s^1(t).
\end{aligned} \tag{3.45}$$

yields

$$2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_t \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim M(t)^{2s+2} E_s^1(t)^2.$$

Now we use Lemma 5 and rewrite (3.32) as

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_2^2 - 2\varepsilon \frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim M(t)^{2s+2} E_s^1(t)^2.$$

Integrating from 0 to  $t$ , we get

$$\begin{aligned}
&\varepsilon^2 \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \\
&- \varepsilon^2 \|\partial_x^\alpha a^{\varepsilon,0}\|_{L^2}^2 + 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S^{\varepsilon,0} \operatorname{Im}(\bar{a}_i^{\varepsilon,0} \partial_x^\alpha a_i^{\varepsilon,0}) dx \lesssim \int_0^t M(r)^{2s+2} E_s^1(r)^2 dr.
\end{aligned} \tag{3.46}$$

It is easy to see that

$$\begin{aligned}
\left| \int_{\Omega} \partial_x^\alpha S \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \right| &\lesssim \|a\|_{L^\infty} \|\partial_x^\alpha S\|_{L^2} \|\partial_x^\alpha a\|_{L^2} \\
&\lesssim M(t) \|u\|_{H^{s-1}} \|a\|_{H^s} \\
&\lesssim M(t) (E_s(t)^2 + E_s(t) \|a\|_{\dot{H}^s}),
\end{aligned}$$

and similarly that

$$\begin{aligned}
\varepsilon^2 \|\partial_x^\alpha a^{\varepsilon,0}\|_{L^2}^2 + \varepsilon \left| \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha S^{\varepsilon,0} \operatorname{Im}(\bar{a}_i^{\varepsilon,0} \partial_x^\alpha a_i^{\varepsilon,0}) dx \right| &\lesssim \varepsilon \|u^{\varepsilon,0}\|_{H^{s-1}} \|a^{\varepsilon,0}\|_{H^s} \|a^{\varepsilon,0}\|_{L^\infty} + \varepsilon^2 \|a^{\varepsilon,0}\|_{H^s}^2 \\
&\lesssim M(0) E_s^1(0)^2.
\end{aligned}$$

Plugging the two estimate above into (3.46), it follows that

$$\varepsilon^2 \|\partial_x^\alpha a\|_{L^2}^2 \lesssim M(0) E_s^1(0)^2 + \varepsilon M(t) E_s(t) \|a\|_{\dot{H}^s} + M(t) E_s(t)^2 + \int_0^t M(r)^{2s+2} E_s^1(r)^2 dr.$$

Summing over all  $\alpha$  such that  $n(\alpha) = s$ , we get

$$\varepsilon^2 \|a\|_{\dot{H}^s}^2 \lesssim M(0) E_s^1(0)^2 + \varepsilon M(t) E_s(t) \|a\|_{\dot{H}^s} + M(t) E_s(t)^2 + \int_0^t M(r)^{2s+2} E_s^1(r)^2 dr. \tag{3.47}$$

Noticing that for any constant  $C > 0$ , we have

$$C\varepsilon M(t)E_s(t)\|a\|_{\dot{H}^s} \leq \frac{1}{2}\varepsilon^2\|a\|_{\dot{H}^s}^2 + \frac{1}{2}C^2M(t)^2E_s(t)^2,$$

it follows that

$$\varepsilon^2\|a\|_{\dot{H}^s}^2 \lesssim M(0)E_s^1(0)^2 + M(t)^2E_s(t)^2 + \int_0^t M(r)^{2s+2}E_s^1(r)^2dr. \quad (3.48)$$

From Lemma 4, i.e. (3.31), it follows that

$$\frac{d}{dt}E_s(t)^2 \lesssim M(t)^{2s+1}E_s^1(t)^2.$$

The integral form of this inequality is

$$E_s(t)^2 \lesssim M(0)^{2s+1}E_s^1(0)^2 + \int_0^t M(r)^{2s+1}E_s^1(r)^2dr. \quad (3.49)$$

Plugging (3.49) into (3.48), we get

$$\varepsilon^2\|a\|_{\dot{H}^s}^2 \lesssim M(t)^2M(0)^{2s+1}E_s^1(0)^2 + M(t)^2 \int_0^t M(r)^{2s+1}E_s^1(r)^2dr. \quad (3.50)$$

Adding (3.49) with (3.50) and recalling the definition of  $N(t)$  given by (3.5), we get

$$E_s^1(t)^2 \lesssim N(t)^{2s+3} \left( E_s^1(0)^2 + \int_0^t E_s^1(r)^2dr \right).$$

By the Gronwall inequality we can conclude that (3.7) holds. This finishes the proof.  $\square$

## 4 Existence and Uniqueness of Solutions

In this section we discuss existence and uniqueness of solutions for the Pauli-Poiswell-WKB system (2.5), i.e. Theorem 1.a. Proposition 2 is equivalent to the last two parts in Theorem 1.a. By the same argument, we can easily prove the first part in Theorem 1.a, i.e. the local wellposedness of the Euler-Poiswell equation (2.12).

We use a fixed point argument to show existence and uniqueness in the space  $Y_{\mu_1, \mu_2}^{T, s}$  defined by (4.1) for some subtly selected  $\mu_1, \mu_2$ . For the fixed point argument we need that the solution map  $\Psi$  is an involution on a suitable subset of  $Y_{\mu_1, \mu_2}^{T, s}$  and that it is a contraction on that set. The first part is shown in Lemma 7, the second part in Lemma 8.

**Proposition 2.** *Under the assumptions of Theorem 1.a there exists a unique solution  $(a^\varepsilon, u^\varepsilon)$  of (2.5), such that  $a^\varepsilon \in L^\infty([0, T], H^s)$ ,  $u^\varepsilon \in W^{1, \infty}([0, T], H^s)$  for some  $T > 0$  that only depends on  $Q, s$ . In particular,  $T$  is independent of  $\varepsilon$ .*



We will close the argument the space  $Y_{\mu_1, \mu_2}^{T, s}$ :

$$Y_{\mu_1, \mu_2}^{T, s} := L^\infty([0, T], H^s(\Omega, \mathbb{C})) \times W^{1, \infty}([0, T], H^s(\Omega, \mathbb{R}^3)), \quad (4.1)$$

equipped with the following norm:

$$\begin{aligned} \|(a, u)\|_{Y_{\mu_1, \mu_2}^{T, s}} &:= \sup_{t \in [0, T]} E_s^{\mu_1, \mu_2}(t) \\ &= \sup_{t \in [0, T]} (\|(a, u)(t)\|_{X^s} + \mu_1 \varepsilon \|a(t)\|_{H^s} + \mu_2 \|\partial_t u(t)\|_{H^{s-1}}), \end{aligned}$$

where  $0 < \mu_1, \mu_2 < 1$ . In order to improve readability we will simply write  $\|(a, u)\|_Y$ . Define the operator  $\Psi$  as:

$$\Psi_{a^\varepsilon, 0, u^\varepsilon, 0}^\varepsilon : Y_{\mu_1, \mu_2}^{T, s} \rightarrow Y_{\mu_1, \mu_2}^{T, s}, \quad (\tilde{a}, \tilde{u}) \mapsto (a, u),$$

where  $(a, u)$  is the solution at time  $t$  of the following linear system

$$\begin{aligned} \partial_t a_j + (\tilde{A} + \tilde{u}) \cdot \nabla a_j + \frac{1}{2} a_j \operatorname{div}(\tilde{u} + \tilde{A}) &= \frac{i\varepsilon}{2} \Delta a_j + \frac{i}{2} \tilde{\mathbf{B}}_j^i a_i, \\ \partial_t u + (\tilde{A} + \tilde{u}) \cdot \nabla u + u \cdot \nabla \tilde{A} + \nabla \left( \frac{|\tilde{A}|^2}{2} + \tilde{V} \right) &= 0, \\ a_j(x, 0) &= a_j^{\varepsilon, 0}(x) \\ u(x, 0) &= u^{\varepsilon, 0}(x). \end{aligned} \quad (4.2)$$

$$-\Delta \tilde{V} = \tilde{\rho}, \quad (4.3)$$

$$-\Delta \tilde{A} = \varepsilon \tilde{w} - \varepsilon \tilde{v} - \tilde{\rho} \tilde{u} - \rho \tilde{A}, \quad (4.4)$$

where

$$\tilde{\rho} = |\tilde{a}_1|^2 + |\tilde{a}_2|^2, \quad (4.5)$$

and

$$\begin{aligned} \tilde{v} &= \frac{1}{2} \nabla \times (\tilde{a}^* \sigma_1 \tilde{a}, \tilde{a}^* \sigma_2 \tilde{a}, \tilde{a}^* \sigma_3 \tilde{a})^T, \\ \tilde{w} &= \frac{i}{2} \sum_{j=1}^2 (\tilde{a}_j \nabla \tilde{a}_j - \tilde{a}_j \nabla \tilde{a}_j). \end{aligned}$$

with  $\tilde{\mathbf{B}}$  defined similarly. Let  $\mathcal{B} = \mathcal{B}_{s, \mu_1, \mu_2, T, L}$  denote the ball of radius  $L$  in  $Y$ :

$$\mathcal{B}_{s, \mu_1, \mu_2, T, L} := \left\{ (a, u) : \|(a, u)\|_{Y_{\mu_1, \mu_2}^{T, s}} \leq L \right\}.$$

*Proof of Proposition 2.* Lemma 7 shows that  $\Psi$  maps  $\mathcal{B}$  into itself and Lemma 8 shows that it is indeed a contraction. Then we can conclude by Banach's fixed point theorem.  $\square$

**Lemma 7.** Let  $(a^{\varepsilon,0}, u^{\varepsilon,0})$  satisfy the assumptions of Theorem 1.a. Then there exists  $L > 0$  only depending on  $Q$  and  $s$  and  $\mu_0 > 0$  such that for all  $\mu_1, \mu_2 \in (0, \mu_0)$  there is a  $T_0 = T_0(\mu_1, \mu_2, L, s) > 0$  independent of  $\varepsilon$  such that for all  $T \in [0, T_0)$

$$\Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon(\mathcal{B}_{s, \mu_1, \mu_2, T, L}) \subset \mathcal{B}_{s, \mu_1, \mu_2, T, L}.$$

In other words,  $\Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon$  maps  $\mathcal{B}_{s, \mu_1, \mu_2, T, L}$  into itself.

**Lemma 8.** Under the assumptions of Theorem 1.a and for any  $L > 0$ , there exist  $\mu_{10} = \mu_{10}(L, s) > 0$  and  $\mu_{20} = \mu_{20}(\mu_1, L, s) > 0$  such that for all  $0 < \mu_1 < \mu_{10}$  and  $0 < \mu_2 < \mu_{20}$  there exists  $T_0 = T_0(\mu_1, \mu_2, L, s) > 0$  independent of  $\varepsilon$  such that for all  $T \in [0, T_0)$  and  $(\tilde{a}^1, \tilde{u}^1), (\tilde{a}^2, \tilde{u}^2) \in \mathcal{B}_{s, \mu_1, \mu_2, T, L}$  we have

$$\|\Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon(\tilde{a}^1, \tilde{u}^1) - \Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon(\tilde{a}^2, \tilde{u}^2)\|_{Y_{\mu_1, \mu_2}^{T, s}} \leq \tau \|(\tilde{a}^1, \tilde{u}^1) - (\tilde{a}^2, \tilde{u}^2)\|_{Y_{\mu_1, \mu_2}^{T, s}}. \quad (4.6)$$

Here,  $\tau \in (0, 1)$  is a constant.

*Proof of Lemma 7.* We need to show that for  $(\tilde{a}, \tilde{u})$  such that  $\|(\tilde{a}, \tilde{u})\|_Y \leq L$  the image  $\Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon(\tilde{a}, \tilde{u})$  also satisfies  $\|\Psi_{a^{\varepsilon,0}, u^{\varepsilon,0}}^\varepsilon(\tilde{a}, \tilde{u})\|_Y \leq L$ . We have

$$\partial_t \partial_x^\alpha a_i + (\tilde{u} + \tilde{A}) \cdot \nabla \partial_x^\alpha a_i = \frac{i\varepsilon}{2} \Delta \partial_x^\alpha a_i - \tilde{\mathbf{R}}_i^c,$$

with

$$\tilde{\mathbf{R}}_i^c = (\operatorname{div} \partial_x^\alpha ((\tilde{u} + \tilde{A}) a_i) - (\tilde{u} + \tilde{A}) \cdot \nabla \partial_x^\alpha a_i) - \frac{1}{2} \partial_x^\alpha (a_i \operatorname{div}(\tilde{u} + \tilde{A})) - \frac{i}{2} \partial_x^\alpha (\tilde{\mathbf{B}}_i^j a_j).$$

*Step 1: Estimate for  $\|(a, u)\|_{X^s}$ .* Let us first look at the  $X^s$ -norm of  $(a, u)$  which we have to estimate in terms of the  $Y$ -norm of  $(\tilde{a}, \tilde{u})$ . Similarly to (3.18), we can apply Lemma 1 and prove that

$$\begin{aligned} \frac{d}{dt} \|a\|_{H^{s-1}} &\lesssim (M(\tilde{a}, \tilde{u})(t) + \|\tilde{A}\|_{W^{1, \infty}}) \|(a, u)\|_{X^s} + M(a, u)(t) (\|(\tilde{a}, \tilde{u})\|_{X^s} + \|\nabla \tilde{A}\|_{H^s}) \\ &\lesssim (E_s^1(\tilde{a}, \tilde{u})(t) + \|\nabla \tilde{A}\|_{H^s}) \|(a, u)\|_{X^s}. \end{aligned}$$

The second inequality follows from (3.6) and Sobolev embedding. Similarly to (3.19), we can prove that

$$\begin{aligned} \frac{d}{dt} \|u\|_{H^s} &\lesssim (\|\nabla \tilde{A}\|_{H^s} + \|\tilde{u}\|_{H^s}) \|u\|_{L^\infty} + \|u\|_{H^s} (\|\tilde{u}\|_{W^{1, \infty}} + \|\tilde{A}\|_{W^{1, \infty}}) \\ &\quad + \|\nabla \tilde{A}\|_{H^s}^2 + \|\nabla \tilde{V}\|_{H^s} \\ &\lesssim (\|\nabla \tilde{A}\|_{H^s} + E_s^1(\tilde{a}, \tilde{u})(t)) \|(a, u)\|_{X^s} + \|\nabla \tilde{A}\|_{H^s}^2 + \|\nabla \tilde{V}\|_{H^s} \end{aligned}$$

Now we have to estimate the potential terms. Due to Lemma 3, we know that

$$\begin{aligned} \|\nabla \tilde{V}\|_{H^s} &\lesssim M(\tilde{a}, \tilde{u})(t) \|\tilde{a}\|_{H^{s-1}} \lesssim \|(\tilde{a}, \tilde{u})\|_Y^2, \\ \|\nabla \tilde{A}\|_{H^s} &\lesssim M(\tilde{a}, \tilde{u})(t)^{2s} E_s^1(\tilde{a}, \tilde{u})(t) \lesssim \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1}}{\mu_1}, \\ \|\tilde{A}\|_{W^{1, \infty}} &\lesssim M(\tilde{a}, \tilde{u})(t)^5 \lesssim \|(\tilde{a}, \tilde{u})\|_Y^5 \end{aligned}$$

Here we used (3.6) and the fact that

$$E_s^1(\tilde{a}, \tilde{u})(t) \lesssim \frac{\|(\tilde{a}, \tilde{u})\|_Y}{\mu_1}, \quad \forall 0 < t < T.$$

Now we get a first estimate for  $\|(a, u)\|_{X^s}$ :

$$\frac{d}{dt} \|(a, u)\|_{X^s} \lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} \right) \|(a, u)\|_{X^s} + \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1}{\mu_1^2} \right).$$

Multiply both sides with  $\|(a, u)\|_{X^s}$  and get

$$\begin{aligned} \frac{d}{dt} \|(a, u)\|_{X^s}^2 &\lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} \right) \|(a, u)\|_{X^s}^2 + \|(a, u)\|_{X^s} \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1}{\mu_1^2} \right) \\ &\lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} \right) \|(a, u)\|_{X^s}^2 + \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{8s+4} + 1}{\mu_1^4} \right). \end{aligned}$$

Writing the estimate in integral form and using the assumptions of Theorem 1.a we obtain

$$\|(a, u)\|_{X^s}^2 \lesssim Q^2 + \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} \right) \int_0^t \|(a, u)(r)\|_{X^s}^2 dr + t \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{8s+4} + 1}{\mu_1^4} \right). \quad (4.7)$$

*Step 2: Estimate for  $\|a\|_{H^s}$ .* Applying the proof of (3.43) and (3.44) to system (4.2) yields

$$\begin{aligned} &\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 + \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(a_i \partial_x^\alpha \operatorname{div}(\tilde{u}) \partial_x^\alpha \bar{a}_i) dx \\ &\lesssim (\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1) (\varepsilon \|a\|_{H^s} + \|(a, u)\|_{X^s})^2 \\ &\lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1^2} \right) E_s^{\mu_1}(a, u)(t)^2, \end{aligned}$$

and

$$\begin{aligned} &-2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx - \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \operatorname{div}(\tilde{u}) \operatorname{Re}(\bar{a}_i \partial_x^\alpha a_i) dx \\ &\lesssim (\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1) (\varepsilon \|a\|_{H^s} + \|(a, u)\|_{X^s})^2 \\ &\lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1^2} \right) E_s^{\mu_1}(a, u)(t)^2. \end{aligned}$$

In the last step of both estimates, we used

$$E_s^1(a, u)(t) \lesssim \frac{E_s^{\mu_1}(a, u)(t)}{\mu_1}.$$

Adding both estimates leads to the following estimate

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1}{\mu_1^2} \right) E_s^{\mu_1}(a, u)(t)^2,$$

which is the analogue of (3.32). Since

$$\frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx = \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \partial_t \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx + \sum_{i=1}^2 \int_{\Omega} \partial_t \partial_x^\alpha \tilde{S} \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx$$

and

$$\begin{aligned} \varepsilon \left| \sum_{i=1}^2 \int_{\Omega} \partial_t \partial_x^\alpha \tilde{S} \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \right| &\lesssim \varepsilon \|\partial_t \partial_x^\alpha \tilde{S}\|_{L^2} \|\partial_x^\alpha a\|_{L^2} \|a\|_{L^\infty} \\ &\lesssim \varepsilon \|\partial_t \tilde{u}\|_{H^{s-1}} \|a\|_{H^s} \|a\|_{L^\infty} \\ &\lesssim \frac{1}{\mu_1 \mu_2} \|(\tilde{a}, \tilde{u})\|_Y E_s^{\mu_1}(a, u)(t)^2, \end{aligned}$$

we have

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx \lesssim \frac{(\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1) E_s^{\mu_1}(a, u)(t)^2}{(\mu_1 \wedge \mu_2) \mu_1}.$$

Integrating from 0 to  $t$  yields

$$\begin{aligned} &\varepsilon^2 \|\partial_x^\alpha a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\bar{a}_i \partial_x^\alpha a_i) dx - \varepsilon^2 \|\partial_x^\alpha a^{\varepsilon,0}\|_{L^2}^2 \\ &+ 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S}(0, t) \operatorname{Im}(\bar{a}_i^{\varepsilon,0} \partial_x^\alpha a_i^{\varepsilon,0}) dx \lesssim \frac{1}{(\mu_1 \wedge \mu_2) \mu_1} (\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1) \int_0^t E_s^{\mu_1}(a, u)(r)^2 dr. \end{aligned}$$

Similar to (3.47), we can use

$$\begin{aligned} \varepsilon^2 \|a\|_{H^s}^2 &\lesssim (\varepsilon \|(a^{\varepsilon,0}, u^{\varepsilon,0})\|_{X^s} \|(\tilde{a}, \tilde{u})(0)\|_{X^s} \|a^{\varepsilon,0}\|_{H^s} + \varepsilon^2 \|a^{\varepsilon,0}\|_{H^s}^2) \\ &+ \varepsilon \|a\|_{H^s} \|(\tilde{a}, \tilde{u})\|_Y \|(a, u)\|_{X^s} + \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1}{(\mu_1 \wedge \mu_2) \mu_1} \right) \int_0^t E_s^{\mu_1}(a, u)(r)^2 dr. \end{aligned}$$

Due to the assumption of Theorem 1.a we know that

$$\begin{aligned} \varepsilon^2 \|a\|_{H^s}^2 &\lesssim Q^2(\|(\tilde{a}, \tilde{u})(0)\|_{X^s} + 1) + \varepsilon \|a\|_{H^s} \|(\tilde{a}, \tilde{u})\|_Y \|(a, u)\|_{X^s} \\ &+ \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1}{(\mu_1 \wedge \mu_2) \mu_1} \right) \int_0^t E_s^{\mu_1, \mu_2}(a, u)(r)^2 dr. \end{aligned} \tag{4.8}$$

Step 3: Estimate for  $\|\partial_t u\|_{H^{s-1}}$ . For  $\partial_t u$ , a similar estimate to (3.45) is given by

$$\begin{aligned}
\|\partial_t u\|_{H^{s-1}} &\lesssim (\|\tilde{u}\|_{L^\infty} + \|\tilde{A}\|_{L^\infty})(\|u\|_{H^s} + \|\tilde{A}\|_{\dot{H}^s}) + \|u\|_{L^\infty}(\|\tilde{u}\|_{H^s} + \|\tilde{A}\|_{\dot{H}^s}) + \|\nabla \tilde{V}\|_{H^{s-1}} \\
&\lesssim (\|\tilde{u}\|_{H^s} + \|\tilde{A}\|_{\dot{H}^s} + \|\tilde{A}\|_{L^\infty})(\|u\|_{H^s} + \|\tilde{A}\|_{\dot{H}^s}) + \|(\tilde{a}, \tilde{u})\|_Y^2 \\
&\lesssim (\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1) (\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1 + \|(a, u)\|_{X^s}). \tag{4.9}
\end{aligned}$$

Step 4: Estimate for  $\|(a, u)\|_Y$ . Since  $\|(a, u)\|_Y = \sup_t E_s^{\mu_1, \mu_2}(a, u)(t)$  we have to combine (4.7), (4.8) and (4.9):

$$\begin{aligned}
E_s^{\mu_1, \mu_2}(a, u)(t)^2 &\lesssim \|(a, u)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|a\|_{H^s}^2 + \mu_2^2 \|\partial_t u\|_{H^{s-1}}^2 \\
&\lesssim Q^2(\mu_1^2 \|(\tilde{a}, \tilde{u})(0)\|_{X^s} + 1) + t \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{8s+4} + 1}{\mu_1^4} \right) \\
&\quad + \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} + \frac{\mu_1 (\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1)}{(\mu_1 \wedge \mu_2)} \right) \int_0^t E_s^{\mu_1, \mu_2}(a, u)(r)^2 dr \\
&\quad + \mu_1^2 \varepsilon \|a\|_{H^s} \|(\tilde{a}, \tilde{u})\|_Y \|(a, u)\|_{X^s} \\
&\quad + \mu_2^2 (\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1) (\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1 + \|(a, u)\|_{X^s}^2).
\end{aligned}$$

Using

$$\frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+1} + 1}{\mu_1} + \frac{\mu_1 (\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1)}{(\mu_1 \wedge \mu_2)} \lesssim \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1}{\mu_1 \wedge \mu_2},$$

yields

$$\begin{aligned}
E_s^{\mu_1, \mu_2}(a, u)(t)^2 &\lesssim Q^2(\mu_1^2 \|(\tilde{a}, \tilde{u})(0)\|_{X^s} + 1) + t \left( \frac{\|(\tilde{a}, \tilde{u})\|_Y^{8s+4} + 1}{\mu_1^4} \right) \\
&\quad + \frac{\|(\tilde{a}, \tilde{u})\|_Y^{2s+2} + 1}{\mu_1 \wedge \mu_2} \int_0^t E_s^{\mu_1, \mu_2}(a, u)(r)^2 dr + \mu_1^2 \varepsilon \|a\|_{H^s} \|(\tilde{a}, \tilde{u})\|_Y \|(a, u)\|_{X^s} \\
&\quad + \mu_2^2 (\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1) (\|(\tilde{a}, \tilde{u})\|_Y^{4s+2} + 1 + \|(a, u)\|_{X^s}^2).
\end{aligned}$$

Now suppose that  $\|(\tilde{a}, \tilde{u})\|_Y \leq L$ . Then the estimate above leads to

$$\begin{aligned}
E_s^{\mu_1, \mu_2}(a, u)(t)^2 &\leq C_s \left( Q^2(\mu_1^2 L + 1) + \frac{t(L^{8s+4} + 1)}{\mu_1^4} + \left( \frac{L^{2s+2} + 1}{\mu_1 \wedge \mu_2} \right) \int_0^t E_s^{\mu_1, \mu_2}(a, u)(r)^2 dr \right. \\
&\quad \left. + \mu_2^2 (L^{4s+2} + 1)^2 + (\mu_1 L + \mu_2^2 (L^{4s+2} + 1)) E_s^{\mu_1, \mu_2}(a, u)(r)^2 \right),
\end{aligned}$$

for some constant  $C_s$ . If we choose  $L$  large enough and  $\mu_1, \mu_2$  small enough such that

$$\begin{aligned}
C_s(\mu_1 L + \mu_2^2 (L^{4s+2} + 1)) &\leq 1/2, \\
C_s(Q^2(\mu_1^2 L + 1) + \mu_2^2 (L^{4s+2} + 1)^2) &\leq L^2/4,
\end{aligned}$$

we can get

$$E_s^{\mu_1, \mu_2}(a, u)(t)^2 \leq \frac{L^2}{2} + 2C_s \left( \left( \frac{L^{2s+2} + 1}{\mu_1 \wedge \mu_2} \right) \int_0^t E_s^{\mu_1, \mu_2}(a, u)(r)^2 dr + \frac{t(L^{8s+4} + 1)}{\mu_1^4} \right).$$

By Gronwall's inequality, we get

$$E_s^{\mu_1, \mu_2}(a, u)(t)^2 \leq \left( \frac{L^2}{2} + \frac{2C(s)(L^{8s+4} + 1)}{\mu^4} t \right) \exp \left( 2C_s \left( \frac{L^{2s+2} + 1}{\mu_1 \wedge \mu_2} \right) t \right).$$

Then there is a  $T$  small enough such that

$$\|(a, u)\|_Y^2 = \sup_{t \in [0, T]} E_s^{\mu_1, \mu_2}(a, u)(t)^2 \leq L^2.$$

It follows that  $\|(a, u)\|_Y \leq L$ . □

*Proof of Lemma 8.* We want to estimate the difference between the image of two solutions  $(\tilde{a}^1, \tilde{u}^1)$  and  $(\tilde{a}^2, \tilde{u}^2)$  to the linearized system. Let

$$(a^j, u^j) = \Psi_{a^{\varepsilon, 0}, u^{\varepsilon, 0}}(\tilde{a}^j, \tilde{u}^j), \quad j = 1, 2.$$

By definition,  $(a^j, u^j)$  satisfies

$$\begin{aligned} \partial_t a_k^j + (\tilde{A}^j + \tilde{u}^j) \cdot \nabla a_k^j + \frac{1}{2} a_k^j \operatorname{div}(\tilde{u}^j + \tilde{A}^j) &= \frac{i\varepsilon}{2} \Delta a_k^j + \frac{i}{2} \tilde{\mathbf{B}}_k^{j, i} a_i^j, \\ \partial_t u^j + (\tilde{A}^j + \tilde{u}^j) \cdot \nabla u^j + u^j \cdot \nabla \tilde{A}^j + \nabla \left( \frac{|\tilde{A}^j|^2}{2} + \tilde{V}^j \right) &= 0, \\ a_k^j(x, 0) &= a_k^{\varepsilon, 0}(x), \quad k = 1, 2 \\ u(x, 0) &= u^{\varepsilon, 0}(x). \end{aligned} \tag{4.10}$$

where  $\tilde{A}^j, \tilde{\mathbf{B}}^j$  and  $\tilde{V}^j$  are defined as above with every variable added with a superscript  $j$ . Define

$$\delta a := a^1 - a^2.$$

The quantities  $\delta u, \delta A, \delta V, \delta \mathbf{B}, \delta \rho$  and  $\delta \tilde{a}, \delta \tilde{u}, \delta \tilde{A}, \delta \tilde{V}, \delta \tilde{\mathbf{B}}, \delta \tilde{\rho}$  are similarly defined. Now subtract (4.10) with  $j = 2$  from (4.2) with  $j = 1$ :

$$\begin{aligned} \partial_t \delta a_j + (\delta \tilde{A} + \delta \tilde{u}) \cdot \nabla a_j^1 + (\tilde{A}^2 + \tilde{u}^2) \cdot \nabla \delta a_j \\ + \frac{1}{2} a_j^1 \operatorname{div}(\delta \tilde{u} + \delta \tilde{A}) + \frac{1}{2} \delta a_j \operatorname{div}(u^2 + A^2) &= \frac{i}{2} \delta \tilde{\mathbf{B}}_j^i a_i^1 + \frac{i}{2} \tilde{\mathbf{B}}_j^{2, i} \delta a_i + \frac{i\varepsilon}{2} \Delta \delta a_j, \\ \partial_t \delta u + (\delta \tilde{A} + \delta \tilde{u}) \cdot \nabla u^1 + (\tilde{A}^2 + \tilde{u}^2) \cdot \nabla \delta u \\ + \delta u \cdot \nabla \tilde{A}^1 + u^2 \cdot \nabla \delta \tilde{A} + \nabla \left( \frac{\delta \tilde{A} \cdot (\tilde{A}^1 + \tilde{A}^2)}{2} + \delta \tilde{V} \right) &= 0, \\ \delta a_j(x, 0) &= 0, \\ \delta u(x, 0) &= 0. \end{aligned} \tag{4.11}$$

Here,  $\delta\tilde{V}$  and  $\delta\tilde{A}$  are given by

$$-\Delta\delta\tilde{V} = \delta\tilde{\rho}, \quad (4.12)$$

$$-\Delta\tilde{A} = \varepsilon\delta\tilde{w} - \varepsilon\delta\tilde{v} - (\delta\tilde{\rho}\tilde{u}^1 - \tilde{\rho}^2\delta\tilde{u}) + \delta\tilde{\rho}\tilde{A}^2 - \tilde{\rho}^1\delta\tilde{A}. \quad (4.13)$$

Again we have to split the proof into multiple steps since we have to estimate the difference in  $Y$ , so we have to estimate  $\|(\delta a, \delta u)\|_{X^s}$ ,  $\mu_1\varepsilon\|\delta a\|_{H^s}$  and  $\mu_2\|\partial_t\delta u\|_{H^{s-1}}$ .

*Step 1.* Apply  $\partial^\alpha$  with  $0 \leq n(\alpha) \leq s-1$  to the first equation in (4.11) which yields

$$\partial_t\partial^\alpha\delta a_j + (\tilde{A}^2 + \tilde{u}^2) \cdot \nabla\partial^\alpha\delta a_j + \mathbf{R}_{d,j} = \frac{i\varepsilon}{2}\Delta\partial^\alpha\delta a_j,$$

where  $\mathbf{R}_{d,j}$  is defined as

$$\begin{aligned} \mathbf{R}_{d,j} &:= \partial^\alpha((\delta\tilde{A} + \delta\tilde{u}) \cdot \nabla a_j^1) + \partial^\alpha((\tilde{A}^2 + \tilde{u}^2) \cdot \nabla\delta a_j) - (\tilde{A}^2 + \tilde{u}^2) \cdot \nabla\partial^\alpha\delta a_j \\ &\quad + \frac{1}{2}\partial^\alpha(a_j^1 \operatorname{div}(\delta\tilde{u} + \delta\tilde{A})) + \frac{1}{2}\partial^\alpha(\delta a_j \operatorname{div}(u^2 + A^2)) - \frac{i}{2}\partial^\alpha(\delta\tilde{\mathbf{B}}_j^i a_i^1) + \frac{i}{2}\partial^\alpha(\tilde{\mathbf{B}}_j^{2,i}\delta a_i) \\ &:= \sum_{j=1}^7 F_j. \end{aligned}$$

Then Lemma 1 implies

$$\begin{aligned} \frac{d}{dt}\|\partial^\alpha\delta a_j\|_{L^2} &\lesssim (M(\tilde{a}_i^2, \tilde{u}^2)(t) + \|\tilde{A}^2\|_{W^{1,\infty}})\|\partial^\alpha\delta a_j\|_{L^2} + \|\mathbf{R}_{d,j}\|_{L^2} \\ &\lesssim_{\mu_1, \mu_2, L} \|\partial^\alpha\delta a_j\|_{L^2} + \|\mathbf{R}_{d,j}\|_{L^2}. \end{aligned}$$

In the second step, we used  $(\tilde{a}^2, \tilde{u}^2) \in \mathcal{B}_{s, \mu_1, \mu_2, T, L}$ . With the help of Lemma 10 and (3.16) we obtain the following estimates. For  $F_1$  in  $\mathbf{R}_{d,j}$  we have the following estimates. If  $n(\alpha) \geq 1$ ,

$$\begin{aligned} \|F_1\|_{L^2} &\lesssim \|\delta\tilde{A} + \delta\tilde{u}\|_{\dot{H}^{n(\alpha)}}\|\nabla a_j^1\|_{L^\infty} + \|a_j^1\|_{H^{s-2}}\|\delta\tilde{A} + \delta\tilde{u}\|_{L^\infty} \\ &\lesssim_{\mu_1, \mu_2, L} \|\delta\tilde{A} + \delta\tilde{u}\|_{\dot{H}^{n(\alpha)}} + \|\delta\tilde{A} + \delta\tilde{u}\|_{L^\infty} \\ &\lesssim_{\mu_1, \mu_2, L} \|\nabla\delta\tilde{A}\|_{H^{s-1}} + \|\delta\tilde{u}\|_{H^s}. \end{aligned}$$

If  $n(\alpha) = 0$  then

$$\begin{aligned} \|F_1\|_{L^2} &\lesssim \|a_j^1\|_{H^1}\|\delta\tilde{A} + \delta\tilde{u}\|_{L^\infty} \\ &\lesssim_{\mu_1, \mu_2, L} \|\nabla\delta\tilde{A}\|_{H^{s-1}} + \|\delta\tilde{u}\|_{H^s}. \end{aligned}$$

Now consider  $F_2 - F_3$ . If  $n(\alpha) \geq 1$ ,

$$\begin{aligned} \|F_2 - F_3\|_{L^2} &\lesssim \|\tilde{A}^2 + \tilde{u}^2\|_{\dot{H}^{n(\alpha)}}\|\nabla\delta a_j\|_{L^\infty} + \|\delta a_j\|_{H^{s-1}}\|\tilde{A}^2 + \tilde{u}^2\|_{W^{1,\infty}} \\ &\lesssim_{\mu_1, \mu_2, L} \|\delta a_j\|_{H^{s-1}}. \end{aligned}$$

And if  $n(\alpha) = 0$  then  $\|F_2 - F_3\|_{L^2} = 0$ . For  $F_4$  we have

$$\begin{aligned}\|F_4\|_{L^2} &\lesssim \|\delta\tilde{u} + \delta\tilde{A}\|_{\dot{H}^{n(\alpha)}} \|\nabla a_j^1\|_{L^\infty} + \|a_j^1\|_{H^{s-1}} \|\delta\tilde{u} + \delta\tilde{A}\|_{W^{1,\infty}} \\ &\lesssim_{\mu_1, \mu_2, L} \|\nabla\delta\tilde{A}\|_{H^{s-1}} + \|\delta\tilde{u}\|_{H^s}.\end{aligned}$$

For  $F_5$  we have

$$\begin{aligned}\|F_5\|_{L^2} &\lesssim \|u^2 + A^2\|_{\dot{H}^{n(\alpha)}} \|\nabla\delta a_j\|_{L^\infty} + \|\delta a_j\|_{H^{s-1}} \|u^2 + A^2\|_{W^{1,\infty}} \\ &\lesssim_{\mu_1, \mu_2, L} \|\delta a_j\|_{H^{s-1}}.\end{aligned}$$

For  $F_6$  and  $F_7$  we have

$$\begin{aligned}\|F_6\|_{L^2} &\lesssim \|\delta\tilde{\mathbf{B}}\|_{H^{s-1}} \|a^1\|_{L^\infty} + \|a^1\|_{H^{s-1}} \|\delta\tilde{\mathbf{B}}\|_{W^{1,\infty}} \\ &\lesssim \|\nabla\delta\tilde{A}\|_{H^{s-1}} \|a^1\|_{L^\infty} + \|a^1\|_{H^{s-1}} \|\nabla\delta\tilde{A}\|_{W^{1,\infty}} \\ &\lesssim_{\mu_1, \mu_2, L} \|\nabla\delta\tilde{A}\|_{H^{s-1}},\end{aligned}$$

and

$$\begin{aligned}\|F_7\|_{L^2} &\lesssim \|\tilde{\mathbf{B}}^2\|_{H^{s-3}} \|\delta a\|_{L^\infty} + \|\delta a\|_{H^{s-1}} \|\tilde{\mathbf{B}}^2\|_{W^{1,\infty}} \\ &\lesssim_{\mu_1, \mu_2, L} \|\delta a\|_{H^{s-1}},\end{aligned}$$

For  $\|\partial^\alpha((\delta\tilde{A} + \delta\tilde{u}) \cdot \nabla a_j^1)\|_{L^2}$  and  $\|\partial^\alpha((\tilde{A}^2 + \tilde{u}^2) \cdot \nabla\delta a_j) - (\tilde{A}^2 + \tilde{u}^2) \cdot \nabla\partial^\alpha\delta a_j\|_{L^2}$ , we used two different estimates in two different cases, i.e.  $n(\alpha) \geq 1$  and  $n(\alpha) = 0$ , because we need to avoid the term  $\|A\|_{L^2}$  (See Remark 9). Collecting the estimates above, similar to (3.18), we can easily prove that

$$\frac{d}{dt} \|\delta a\|_{H^{s-1}} \lesssim_{\mu_1, \mu_2, L} \|\delta a\|_{H^{s-1}} + \|\delta\tilde{u}\|_{H^s} + \|\nabla\delta\tilde{A}\|_{H^{s-1}}.$$

For  $\delta u$ , similar to (3.19), we have

$$\frac{d}{dt} \|\delta u\|_{H^s} \lesssim_{\mu_1, \mu_2, L} \|\delta u\|_{H^s} + \|\delta\tilde{u}\|_{H^s} + \|\nabla\delta\tilde{A}\|_{H^{s-1}} + \|\nabla\delta\tilde{V}\|_{H^s}.$$

For  $\delta\tilde{V}$ , similar to (3.20), we can prove that

$$\|\nabla\delta\tilde{V}\|_{H^s} \lesssim_{\mu_1, \mu_2, L} \|\delta a\|_{H^{s-1}}.$$

For  $\delta A$ , similar to (3.26) and (3.30), from (4.13) we can derive

$$\|\nabla\delta A\|_{L^2} \lesssim_{\mu_1, \mu_2, L} \|\delta u\|_{L^2} + \|\delta a\|_{L^2} + \varepsilon \|\nabla\delta a\|_{L^2},$$

and

$$\begin{aligned}\|\nabla\delta A\|_{H^{s-1}} &\lesssim_{\mu_1, \mu_2, L} \|\delta u\|_{H^s} + \|\delta a\|_{H^{s-1}} + \varepsilon \|\delta a\|_{H^s} + \|\nabla\delta A\|_{L^2} \\ &\lesssim_{\mu_1, \mu_2, L} \|\delta u\|_{H^s} + \|\delta a\|_{H^{s-1}} + \varepsilon \|\delta a\|_{H^s}\end{aligned}$$



It follows that

$$\frac{d}{dt} \|(\delta a, \delta u)\|_{X^s} \lesssim_{\mu_1, \mu_2, L} \|(\delta a, \delta u)\|_{X^s} + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y.$$

The integral form of this estimate is

$$\|(\delta a, \delta u)\|_{X^s} \lesssim_{\mu_1, \mu_2, L} \int_0^t \|(\delta a, \delta u)(r)\|_{X^s} dr + t \|(\delta \tilde{a}, \delta \tilde{u})\|_Y. \quad (4.14)$$

This completes the estimate in  $X^s$ .

*Step 2.* Similarly to (3.43), we can prove that

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha \delta a\|_{L^2}^2 + \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(\delta a_i \partial_x^\alpha \operatorname{div}(\tilde{u}^2) \partial_x^\alpha \delta \bar{a}_i) dx \lesssim_{\mu_1, \mu_2, L} \|(\delta a, \delta u)\|_{X^s}^2 + \|(\delta \tilde{a}, \delta \tilde{u})\|_{Y_{\mu_1, \mu_2}^{T, s}}^2.$$

Similarly to (3.44), we can prove that

$$\begin{aligned} & -2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \partial_t \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx - \varepsilon^2 \sum_{i=1}^2 \int_{\Omega} \operatorname{Re}(\delta a_i \partial_x^\alpha \operatorname{div}(\tilde{u}^2) \partial_x^\alpha \delta \bar{a}_i) dx \\ & \lesssim_{\mu_1, \mu_2, L} \|(\delta a, \delta u)\|_{X^s}^2 + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2. \end{aligned}$$

Then similarly to (3.32), adding the two estimates above yields

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha \delta a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \partial_t \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \lesssim_{\mu_1, \mu_2, L} \|(\delta a, \delta u)\|_{X^s}^2 + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2.$$

It follows that

$$\varepsilon^2 \frac{d}{dt} \|\partial_x^\alpha \delta a\|_{L^2}^2 - 2\varepsilon \frac{d}{dt} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \lesssim_{\mu_1, \mu_2, L} \|(\delta a, \delta u)\|_{X^s}^2 + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2.$$

Integrating from 0 to  $t$  gives

$$\begin{aligned} & \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2 - 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \\ & \lesssim_{\mu_1, \mu_2, L} \int_0^t \|(\delta a, \delta u)(r)\|_{X^s}^2 dr + t \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2. \end{aligned} \quad (4.15)$$

Notice that the second term of the LHS is bounded by

$$\begin{aligned} \left| 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \right| & \leq C_{L, s} \varepsilon \|(\delta a, \delta u)(r, \cdot)\|_{X^s} \|\partial_x^\alpha \delta a\|_{L^2} \\ & \leq \frac{C_{L, s}}{\mu_1} (\|(\delta a, \delta u)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2), \end{aligned}$$

for some constant  $C_{L,s}$ . Then if we choose  $\mu_1$  small enough such that

$$\mu_1 C_{L,s} |\{\alpha : n(\alpha) = s\}| \leq \frac{1}{2},$$

we have the following estimate,

$$\sum_{n(\alpha)=s} \mu_1^2 \left| 2\varepsilon \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \right| \lesssim \frac{1}{2} (\|(\delta a, \delta u)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2).$$

Summing (4.15) with respect to  $\alpha$  yields

$$\begin{aligned} & \varepsilon^2 \|\delta a\|_{H^s}^2 - \frac{1}{2\mu_1^2} (\|(\delta a, \delta u)(r)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2) \\ & \lesssim \sum_{n(\alpha)=s} \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2 - 2\varepsilon \sum_{n(\alpha)=s} \sum_{i=1}^2 \int_{\Omega} \partial_x^\alpha \tilde{S} \operatorname{Im}(\delta \bar{a}_i \partial_x^\alpha \delta a_i) dx \\ & \lesssim_{\mu_1, \mu_2, L} \int_0^t \|(\delta a, \delta u)(r)\|_{X^s}^2 dr + t \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2. \end{aligned}$$

Combine this estimate and (4.14) and it follows that

$$\|(\delta a, \delta u)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2 \lesssim_{\mu_1, \mu_2, L} \int_0^t \|(\delta a, \delta u)(r)\|_{X^s}^2 dr + t \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2. \quad (4.16)$$

*Step 3.* For  $\partial_t \delta u$ , we have the following estimate

$$\|\partial_t \delta u\|_{H^{s-1}} \leq C_{\mu_1, L} (\|(\delta a, \delta u)\|_{X^s} + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y)$$

We can choose  $\mu_2$  small enough, which depending on  $\mu_1$  and  $L$ , such that

$$\mu_2 C_{\mu_1, L} \leq \frac{1}{4}.$$

Then it is obvious that

$$\mu_2^2 \|\partial_t \delta u\|_{H^{s-1}}^2 \leq \frac{1}{4} (\|(\delta a, \delta u)\|_{X^s}^2 + \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2).$$

*Step 4.* Combine this estimate with (4.16) and get

$$\begin{aligned} & \|(\delta a, \delta u)\|_{X^s}^2 + \mu_1^2 \varepsilon^2 \|\partial_x^\alpha \delta a\|_{L^2}^2 + \mu_2^2 \|\partial_t \delta u\|_{H^{s-1}}^2 \\ & \leq \frac{1}{2} \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2 + C_{\mu_1, \mu_2, L} \left( \int_0^t \|(\delta a, \delta u)(r)\|_{X^s}^2 dr + t \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2 \right). \end{aligned}$$

By Gronwall's inequality, we know that

$$\|(\delta a, \delta u)\|_Y^2 \leq \left( \frac{1}{2} + C_{\mu_1, \mu_2, L} T \right) \exp(C_{\mu_1, \mu_2, L} T) \|(\delta \tilde{a}, \delta \tilde{u})\|_Y^2.$$

By choosing  $T$  small enough we can get (4.6). □

## 5 Behavior at Blowup Time

In this section, we are going to prove Theorem 1.b. We only prove the second part of 1.b, i.e. the blow up of the Pauli-Poiswell-WKB equation (2.5). It is equivalent to Proposition 3. Again, by a similar but simpler argument, we easily obtain the first part in Theorem 1.b, i.e. the blow up of the Euler-Poiswell equation (2.12).

**Proposition 3.** *Under the assumptions of Proposition 2 there exists a unique local solution  $(a^\varepsilon, u^\varepsilon)$  to the Pauli-Poiswell-WKB equation (2.5). Suppose the solution is defined on a maximal time interval  $[0, T^*)$ . Then we have*

$$\lim_{t \rightarrow T^*-} M(a^\varepsilon, u^\varepsilon)(t) = +\infty.$$

where  $M$  is defined by (3.4). Furthermore, when  $\varepsilon$  is small enough, there exists a  $K = K_{\varepsilon, Q, T^*}$ , such that

$$\limsup_{t \rightarrow T^*-} \|u^\varepsilon(t)\|_{W^{1,\infty}} + \|a^\varepsilon(t)\|_{L^\infty} > K.$$

Here, for fixed  $Q$  and  $T^*$ ,  $K_{\varepsilon, Q, T^*}$  goes to infinity as  $\varepsilon$  goes to zero.

*Proof.* Suppose that  $T^*$  is the maximal time of existence of the solution. Then  $E_s^1(t)(a^\varepsilon, u^\varepsilon)$  blows up as  $t \uparrow T^*$ . If  $N(a^\varepsilon, u^\varepsilon)(t)$  would remain bounded when  $t \uparrow T^*$ , then by Proposition 1 and the assumption of Theorem 1.a,  $E_s^1(a^\varepsilon, u^\varepsilon)$  would remain bounded when  $t \uparrow T^*$  which is a contradiction. Thus,

$$\lim_{t \uparrow T^*} N(a^\varepsilon, u^\varepsilon)(t) = +\infty.$$

On the other hand, it is obvious that  $N(a^\varepsilon, u^\varepsilon)(t)$  remains bounded when  $t < T^*$ . It follows that

$$\lim_{t \uparrow T^*} M(a^\varepsilon, u^\varepsilon)(t) = +\infty.$$

For the second part we are going to use a bootstrap argument. If for all times

$$\|u^\varepsilon(t)\|_{W^{1,\infty}} + \|a^\varepsilon(\cdot, t)\|_{L^\infty} \leq K, \tag{5.1}$$

for some constant  $K$  to be assigned we also assume that

$$\|a(t)\|_{H^1} + \|a(t)\|_{W^{1,\infty}} + \|a(t)\|_{W^{2,3}} \leq K/\varepsilon, \tag{5.2}$$

for all times. Here we also require that

$$\|a^{\varepsilon,0}\|_{H^1} + \|a^{\varepsilon,0}\|_{W^{1,\infty}} + \|a^{\varepsilon,0}\|_{W^{2,3}} \leq K/\varepsilon. \tag{5.3}$$

Then it follows that for all  $0 < t < T^*$ ,

$$N(a^\varepsilon, u^\varepsilon)(t) \leq 2K + 1.$$

Since  $s > 7/2$ , by the Sobolev embedding, we can immediately get

$$\|a(t)\|_{H^1} + \|a(t)\|_{W^{1,\infty}} + \|a(t)\|_{W^{2,3}} \lesssim \|a\|_{H^s},$$

where the constant only depends on  $s$ . Due to the a priori estimate we know that

$$\begin{aligned} \|a(t)\|_{H^1} + \|a(t)\|_{W^{1,\infty}} + \|a(t)\|_{W^{2,3}} &\lesssim E_s^1(a^\varepsilon, u^\varepsilon)(t) \\ &\lesssim N(a^\varepsilon, u^\varepsilon)(t)^{2s+3} Q e^{CN(a^\varepsilon, u^\varepsilon)(t)^{2s+3}t} \\ &\leq C(2K+1)^{2s+3} Q e^{C(2K+1)^{2s+3}T^*}. \end{aligned} \quad (5.4)$$

where  $C = C_s$ . If we have

$$C(2K+1)^{2s+3} Q e^{C(2K+1)^{2s+3}T^*} < \frac{K}{\varepsilon}, \quad (5.5)$$

then (5.4) is a stronger estimate than (5.2). By the bootstrap argument, we can actually obtain that (5.4) holds true all the time. Then (5.1) implies that  $N(a^\varepsilon, u^\varepsilon)(t)$  is bounded for all times, which contradicts the fact that  $T^*$  is maximal. Thus we only need to show that when  $\varepsilon$  is small enough, there exists  $K = K(\varepsilon, Q, T^*)$  such that (5.5) holds true. And we also require that  $K > CQ$ . It is easy to see that such a  $K$  exists. In fact, we can take

$$K = \frac{1}{2} \left( \left| \log \frac{\sqrt{\varepsilon}}{CT^*} \right|^{1/(2s+3)} - 1 \right).$$

Then the LHS of (5.5) is

$$\frac{CQ \left| \log \frac{\sqrt{\varepsilon}}{CT^*} \right|}{\sqrt{\varepsilon}},$$

and the RHS of (5.5) is

$$\frac{1}{2\varepsilon} \left( \left| \log \frac{\sqrt{\varepsilon}}{CT^*} \right|^{1/(2s+3)} - 1 \right).$$

It is obvious that when  $\varepsilon$  is small enough, (5.5) and (5.3) hold true and  $K$  goes to infinity as  $\varepsilon \rightarrow 0+$ . This finishes the proof.  $\square$

## 6 Semiclassical limit

In this section, we are going to prove our main result, Theorem 2. Note that in equation (2.5) the semiclassical parameter  $\varepsilon$  only appears in the terms  $(i\varepsilon/2)\Delta a_j$  and  $\varepsilon w - \varepsilon v$ . In Proposition 1 established the regularity of  $a$  and  $u$  in  $X^s = H^{s-1} \times H^s$  by the a priori estimate. Now we can directly pass to the semiclassical limit  $\varepsilon \rightarrow 0$  by taking the difference between  $(a^\varepsilon, u^\varepsilon)$  and  $(a, u)$  in  $X^s$ .

*Proof of Theorem 2.* We split the proof into three parts. In the first part we estimate the difference between the solution  $(a^\varepsilon, u^\varepsilon)$  of the Pauli-Poiswell-WKB equation and the solution  $(a, u)$  of the Euler-Poiswell equation. Here we use the machinery developed so far (i.e. charge and energy conservation and the Poisson estimates). In the second part we show the semiclassical limit of the density  $\rho^\varepsilon$ , the current density  $J^\varepsilon$  and the monokinetic Wigner transform  $f^\varepsilon$ . In the

third part we prove the statement about the time of existence in the theorem.

Let  $(a^\varepsilon, u^\varepsilon, A^\varepsilon, V^\varepsilon)$  be the solution of (2.5) and  $(a, u, A, V)$  the solution of (2.12). Let

$$\begin{aligned}\delta a &:= a^\varepsilon - a, & \delta u &:= u^\varepsilon - u, & \delta A &:= A^\varepsilon - A, \\ \delta V &:= V^\varepsilon - V, & \delta \mathbf{B} &:= \mathbf{B}^\varepsilon - \mathbf{B}, & \delta \rho &:= \rho^\varepsilon - \rho.\end{aligned}$$

Subtracting (2.12) from (2.5) yields

$$\begin{aligned}\partial_t \delta a + (\delta A + \delta u) \cdot \nabla a_j^\varepsilon + (A + u) \cdot \nabla \delta a_j \\ + \frac{1}{2} a_j^\varepsilon \operatorname{div}(\delta u + \delta A) + \frac{1}{2} \delta a_j \operatorname{div}(u + A) &= \frac{i}{2} \delta \mathbf{B}_j^i a_i^\varepsilon + \frac{i}{2} \mathbf{B}_j^i \delta a_i + \frac{i\varepsilon}{2} \Delta a_j^\varepsilon, \\ \partial_t \delta u + (\delta A + \delta u) \cdot \nabla u^\varepsilon + (A + u) \cdot \nabla \delta u + \delta u \cdot \nabla A^\varepsilon \\ + u^\varepsilon \cdot \nabla \delta A + \nabla \left( \frac{1}{2} \delta A \cdot (A + A^\varepsilon) + \delta V \right) &= 0, \\ \delta a_j(x, 0) &= 0 \\ \delta u(x, 0) &= 0.\end{aligned}\tag{6.1}$$

Similarly,  $\delta V$  and  $\delta A$  are given by

$$-\Delta \delta V = \delta \rho,\tag{6.2}$$

$$-\Delta \delta A = \varepsilon w^\varepsilon - \varepsilon v^\varepsilon - (\delta \rho u^\varepsilon - \rho \delta u) + \delta \rho A - \rho^\varepsilon \delta A.\tag{6.3}$$

Apply  $\partial^\alpha$  with  $0 \leq n(\alpha) \leq s - 3$  to the first equation in (6.1), we obtain

$$\partial_t \partial^\alpha \delta a_j + (A + u) \cdot \nabla \partial^\alpha \delta a_j + \mathbf{R}_d = 0,$$

with

$$\begin{aligned}\mathbf{R}_d &:= \partial^\alpha ((\delta A + \delta u) \cdot \nabla a_j^\varepsilon) + \partial^\alpha ((A + u) \cdot \nabla \delta a_j) - (A + u) \cdot \nabla \partial^\alpha \delta a_j \\ &+ \frac{1}{2} \partial^\alpha (a_j^\varepsilon \operatorname{div}(\delta u + \delta A)) + \frac{1}{2} \partial^\alpha (\delta a_j \operatorname{div}(u + A)) \\ &- \frac{i}{2} \partial^\alpha (\delta \mathbf{B}_j^i a_i^\varepsilon) + \frac{i}{2} \partial^\alpha (\mathbf{B}_j^i \delta a_i) - \frac{i\varepsilon}{2} \Delta \partial^\alpha a_j^\varepsilon. \\ &:= \sum_{j=1}^8 F_j\end{aligned}$$

Thanks to Lemma 1, we obtain

$$\frac{d}{dt} \|\partial^\alpha \delta a_j\|_{L^2}^2 = \int_{\Omega} \operatorname{div}(A + u) |\partial^\alpha \delta a_j|^2 dx - \int_{\Omega} \operatorname{Re}(\mathbf{R}_d \partial^\alpha \overline{\delta a_j}).$$

*Step 1: Estimate for  $\mathbf{R}_d$ .* With the help of Lemma 10 and (3.16), we get the following estimates. If  $n(\alpha) \geq 1$ ,

$$\begin{aligned}\|F_1\|_{L^2} &= \|\partial^\alpha ((\delta A + \delta u) \cdot \nabla a_j^\varepsilon)\|_{L^2} \lesssim \|\delta A + \delta u\|_{\dot{H}^{n(\alpha)}} \|\nabla a_j^\varepsilon\|_{L^\infty} + \|a_j^\varepsilon\|_{H^{s-2}} \|\delta A + \delta u\|_{L^\infty} \\ &\lesssim_{Q,T} \|\delta A + \delta u\|_{\dot{H}^{n(\alpha)}} + \|\delta A + \delta u\|_{L^\infty} \\ &\lesssim_{Q,T} \|\nabla \delta A\|_{H^{s-3}} + \|\delta u\|_{H^s},\end{aligned}$$

and

$$\begin{aligned}\|F_2 - F_3\|_{L^2} &= \|\partial^\alpha((A+u) \cdot \nabla \delta a_j) - (A+u) \cdot \nabla \partial^\alpha \delta a_j\|_{L^2} \\ &\lesssim \|A+u\|_{\dot{H}^{n(\alpha)}} \|\nabla \delta a_j\|_{L^\infty} + \|\delta a_j\|_{H^{s-3}} \|A+u\|_{W^{1,\infty}} \\ &\lesssim_{Q,T} \|\delta a_j\|_{H^{s-3}}.\end{aligned}$$

If  $n(\alpha) = 0$ ,

$$\begin{aligned}\|F_1\|_{L^2} &= \|\partial^\alpha((\delta A + \delta u) \cdot \nabla a_j^\varepsilon)\|_{L^2} \lesssim \|a_j^\varepsilon\|_{H^1} \|\delta A + \delta u\|_{L^\infty} \\ &\lesssim_{Q,T} \|\nabla \delta A\|_{H^{s-3}} + \|\delta u\|_{H^s},\end{aligned}$$

and

$$\|F_2 - F_3\|_{L^2} = \|\partial^\alpha((A+u) \cdot \nabla \delta a_j) - (A+u) \cdot \nabla \partial^\alpha \delta a_j\|_{L^2} = 0,$$

Moreover,

$$\begin{aligned}\|F_4\|_{L^2} &\lesssim \|\partial^\alpha(a_j^\varepsilon \operatorname{div}(\delta u + \delta A))\|_{L^2} \lesssim \|\delta A + \delta u\|_{\dot{H}^{n(\alpha)}} \|\nabla a_j^\varepsilon\|_{L^\infty} + \|a_j^\varepsilon\|_{H^{s-1}} \|\delta A + \delta u\|_{W^{1,\infty}} \\ &\lesssim_{Q,T} \|\nabla \delta A\|_{H^{s-3}} + \|\delta u\|_{H^s}\end{aligned}$$

For  $F_5$  we have

$$\begin{aligned}\|F_5\|_{L^2} &\lesssim \|\partial^\alpha(\delta a_j \operatorname{div}(u + A))\|_{L^2} \lesssim \|A+u\|_{\dot{H}^{n(\alpha)}} \|\nabla \delta a_j\|_{L^\infty} + \|\delta a_j\|_{H^{s-3}} \|A+u\|_{W^{1,\infty}} \\ &\lesssim_{Q,T} \|\delta a_j\|_{H^{s-3}}.\end{aligned}$$

For  $F_6$  we have

$$\begin{aligned}\|F_6\|_{L^2} &\lesssim \|\partial^\alpha(\delta \mathbf{B}_j^i a_i^\varepsilon)\|_{L^2} \lesssim \|\delta \mathbf{B}\|_{H^{s-3}} \|a^\varepsilon\|_{L^\infty} + \|a^\varepsilon\|_{H^{s-3}} \|\delta \mathbf{B}\|_{W^{1,\infty}} \\ &\lesssim \|\nabla \delta A\|_{H^{s-3}} \|a^\varepsilon\|_{L^\infty} + \|a^\varepsilon\|_{H^{s-3}} \|\nabla \delta A\|_{W^{1,\infty}} \\ &\lesssim_{Q,T} \|\nabla \delta A\|_{H^{s-3}}.\end{aligned}$$

For  $F_7$  we have

$$\begin{aligned}\|F_7\|_{L^2} &\lesssim \|\partial^\alpha(\mathbf{B}_j^i \delta a_i)\|_{L^2} \lesssim \|\mathbf{B}\|_{H^{s-3}} \|\delta a\|_{L^\infty} + \|\delta a\|_{H^{s-3}} \|\mathbf{B}\|_{W^{1,\infty}} \\ &\lesssim_{Q,T} \|\delta a\|_{H^{s-3}},\end{aligned}$$

And for  $F_8$ , by Proposition 2, we have

$$\|F_8\|_{L^2} \lesssim \varepsilon \|\Delta \partial^\alpha a_j^\varepsilon\|_{L^2} \lesssim \varepsilon \|a^\varepsilon\|_{H^{s-1}} \lesssim_{Q,T} \varepsilon.$$

For  $\|\partial^\alpha((\delta A + \delta u) \cdot \nabla a_j^\varepsilon)\|_{L^2}$  and  $\|\partial^\alpha((A+u) \cdot \nabla \delta a_j) - (A+u) \cdot \nabla \partial^\alpha \delta a_j\|_{L^2}$ , we used two different estimates in two different cases, i.e.  $n(\alpha) \geq 1$  and  $n(\alpha) = 0$ , because we need to avoid the term  $\|A\|_{L^2}$  (See Remark 9). Similar to (3.17), we can combine the estimates above together and get

$$\|\mathbf{R}_d\|_{L^2} \lesssim_{Q,T} \|\nabla \delta A\|_{H^{s-3}} + \|\delta u\|_{H^{s-2}} + \|\delta a\|_{H^{s-3}} + \varepsilon.$$

*Step 2: Estimates for  $\|\delta u\|_{H^{s-2}}$  and  $\|\delta a\|_{H^{s-3}}$ .* It is obvious that

$$\left| \int_{\Omega} \operatorname{div}(A + u) |\partial^\alpha \delta a_j|^2 dx \right| \lesssim_{Q,T} \|\delta a\|_{H^{s-3}}^2.$$

Similar to (3.18), it follows that

$$\frac{d}{dt} \|\delta a\|_{H^{s-3}} \lesssim_{Q,T} \|\delta a\|_{H^{s-3}} + \|\delta u\|_{H^{s-2}} + \|\nabla \delta A\|_{H^{s-3}} + \varepsilon. \quad (6.4)$$

Similarly, for  $\delta u$ , we can obtain

$$\frac{d}{dt} \|\delta u\|_{H^{s-2}} \lesssim_{Q,T} \|\delta u\|_{H^{s-2}} + \|\nabla \delta A\|_{H^{s-3}} + \|\nabla \delta V\|_{H^{s-1}}. \quad (6.5)$$

*Step 3: Estimates for  $\delta V$  and  $\delta A$*  For  $\delta V$ , similar to (3.20), we can prove that

$$\|\nabla \delta V\|_{H^{s-1}} \lesssim_{Q,T} \|\delta a\|_{H^{s-3}}. \quad (6.6)$$

For  $\delta A$ , similar to (3.26) and (3.30), we can prove that

$$\begin{aligned} \|\nabla \delta A\|_{L^2} &\lesssim_{Q,T} \|\delta u\|_{L^2} + \|\delta a\|_{L^2} + \varepsilon, \\ \|\nabla \delta A\|_{H^{s-3}} &\lesssim_{Q,T} \varepsilon + \|\delta u\|_{H^{s-3}} + \|\delta a\|_{H^{s-3}} + \|\nabla \delta A\|_{L^2}. \end{aligned}$$

It follows that

$$\|\nabla \delta A\|_{H^{s-3}} \lesssim_{Q,T} \varepsilon + \|\delta u\|_{H^{s-3}} + \|\delta a\|_{H^{s-3}}. \quad (6.7)$$

Combining (6.4), (6.5), (6.6) and (6.7), we finally get

$$\frac{d}{dt} (\|\delta a\|_{H^{s-3}} + \|\delta u\|_{H^{s-2}}) \lesssim_{Q,T} \|\delta a\|_{H^{s-3}} + \|\delta u\|_{H^{s-2}} + \varepsilon.$$

Applying the Gronwall inequality, we immediately get

$$\|\delta a\|_{H^{s-3}} + \|\delta u\|_{H^{s-2}} \lesssim_{Q,T} \|a^{\varepsilon,0} - a^0\|_{H^{s-3}} + \|u^{\varepsilon,0} - u^0\|_{H^{s-2}} + \varepsilon.$$

(2.37) immediately follows from the assumption of Theorem 1.a. This finishes the proof of the first part of Theorem 2.

Now we turn to the second part of Theorem 2, which is the semiclassical limit of the density  $\rho^\varepsilon$ , the current  $J^\varepsilon$  and the Wigner transform  $f^\varepsilon$ . Recall that  $\rho^\varepsilon$  and  $J^\varepsilon$  are given by

$$\begin{aligned} \rho^\varepsilon &= |a^\varepsilon|^2, \\ J^\varepsilon &= \varepsilon w^\varepsilon - \varepsilon v^\varepsilon - \rho^\varepsilon (u^\varepsilon + A^\varepsilon). \end{aligned}$$

Since  $a^\varepsilon$  and  $u^\varepsilon$  are uniformly bounded in  $H^{s-1}$  and  $H^s$  respectively, we can pass to the limit  $\varepsilon \rightarrow 0$ , which yields

$$\begin{aligned} \rho^\varepsilon &\xrightarrow{\varepsilon \rightarrow 0} \rho = |a|^2, \\ \rho^\varepsilon (u^\varepsilon + A^\varepsilon) &\xrightarrow{\varepsilon \rightarrow 0} J = -(\rho u + \rho A), \\ \varepsilon w^\varepsilon - \varepsilon v^\varepsilon &\xrightarrow{\varepsilon \rightarrow 0} 0. \end{aligned}$$

in  $H^{s-3}$ . The semiclassical limit  $J^\varepsilon \rightarrow J$  as  $\varepsilon \rightarrow 0$  immediately follows. Finally, we come to the semiclassical of  $f^\varepsilon$ . First, by [8], for any fixed  $t < T$ , there exists a subsequence  $\{f^\varepsilon\}$  (we still denote it by  $f^\varepsilon$  for convenience) and a non-negative Radon measure  $f$ , such that

$$f^\varepsilon \xrightarrow{\varepsilon \rightarrow 0} f \quad \text{in } \mathcal{A}'.$$

It is easy to see that

$$\|\varepsilon \nabla \psi^\varepsilon\|_{L^2} \leq C,$$

for some constant  $C$  independent of  $\varepsilon$ . Due to [21], it follows that

$$\int_{\mathbb{R}^3} f(x, \xi, t) d\xi = \lim_{\varepsilon \rightarrow 0} \rho^\varepsilon(x, t) = \rho(x, t).$$

It is easy to see that

$$\|(u(x, t) - i\varepsilon \nabla) \psi^\varepsilon\|_{L^2} \xrightarrow{\varepsilon \rightarrow 0} 0.$$

Then using the argument in [35], we know that

$$\int_{\mathbb{R}^3 \times \mathbb{R}^3} |\xi - u(x, t)|^2 f(x, \xi, t) d\xi dx = 0,$$

and that

$$f(x, \xi, t) = \rho(x, t) \delta(\xi - u(x, t)).$$

It is easy to see that  $f$  satisfies (2.28) and (2.38).

The last part of Theorem 2 is proved by a bootstrap argument. For any  $T < T^0$ , there exists  $K = K(T) > 0$ , such that for all  $0 < t < T$ ,

$$\|(a, u)(t)\|_{X^s} + N(a, u)(t) \leq K.$$

Suppose that

$$E_s^1(a^\varepsilon, u^\varepsilon)(t) \leq K', \tag{6.8}$$

for all  $0 < t < T$  with  $K'$  to be determined. By virtue of the first part of Theorem 2, it follows that for all  $0 < t < T$ ,

$$\|(a, u)(t) - (a^\varepsilon, u^\varepsilon)(t)\|_{X^{s-2}} \lesssim_{K, K', T} \varepsilon.$$

Using Sobolev embedding, we obtain

$$\|u - u^\varepsilon\|_{W^{1, \infty}} + \|a - a^\varepsilon\|_{H^1} + \|a - a^\varepsilon\|_{W^{1, \infty}} + \|a - a^\varepsilon\|_{W^{2, 3}} \lesssim_{K, Q, T} \varepsilon.$$

When  $\varepsilon$  is small enough, this estimate yields for all  $0 < t < T$ ,

$$\|N(a^\varepsilon, u^\varepsilon)(T)\|_{X^s} \leq 2K,$$

By Proposition 1 we know that for all  $0 < t < T$ ,

$$E_s^1(a^\varepsilon, u^\varepsilon)(t) \leq C(2K)^{2s+3} Q e^{C(2K)^{2s+3} T}.$$

If we take  $K'$  larger than  $C(2K)^{2s+3} Q e^{C(2K)^{2s+3} T}$ , then the above estimate is sharper than (6.8). By the bootstrap argument, we know that (6.8) holds true and  $(a^\varepsilon, u^\varepsilon)$  does not blow up at  $T$ . It follows that for all  $T < T^0$ ,

$$\liminf_{t \rightarrow T^-} T^\varepsilon \geq T,$$

which immediately leads to (2.39) and finishes the proof.  $\square$



## A Some inequalities

**Lemma 9 (Gagliardo-Nirenberg).** *Suppose  $n, j, m \in \mathbb{N}$ ,  $1 \leq p, q, r \leq \infty$  and  $\alpha \in [0, 1]$  such that*

$$\frac{1}{p} = \frac{j}{n} + \left( \frac{1}{r} - \frac{m}{n} \right) \alpha + \frac{1-\alpha}{q},$$

and

$$\frac{j}{m} \leq \alpha \leq 1.$$

*Suppose that  $u : \mathbb{R}^n \rightarrow \mathbb{R}$  is a function in  $L^q(\mathbb{R}^n)$  with  $D^m u \in L^r(\mathbb{R}^n)$ . Then  $D^j u \in L^p(\mathbb{R}^n)$  and there exists a constant  $C$  depending on  $m, n, j, q, r$  and  $\alpha$  but independent of  $u$  such that*

$$\|D^j u\|_{L^p} \leq C \|D^m u\|_{L^r}^\alpha \|u\|_{L^q}^{1-\alpha}.$$

*The result has two exceptional cases:*

1. *If  $j = 0$ ,  $mr < n$  and  $q = \infty$ , then it is necessary to make the additional assumption that either  $\lim_{|x| \rightarrow \infty} u(x) = 0$  or that  $u \in L^s(\mathbb{R}^n)$  for some  $s < \infty$ .*
2. *If  $1 < r < \infty$  and  $m - j - \frac{n}{r}$  is a non-negative integer, then it is necessary to assume also that  $\alpha \neq 1$ .*

Cf. [4].

**Lemma 10 (Kato-Ponce).** *Assume that  $f, g \in H^n(\mathbb{R}^d)$ . Then for any multi-index  $\alpha$ ,  $|\alpha| = n$ , we have*

$$\begin{aligned} \|\partial^\alpha(fg)\|_{L^2} &\leq C_n (\|f\|_{L^\infty} \|g\|_{\dot{H}^n} + \|g\|_{L^\infty} \|f\|_{\dot{H}^n}), \\ \|\partial^\alpha(fg) - f\partial^\alpha(g)\|_{L^2} &\leq C_n (\|\nabla f\|_{L^\infty} \|g\|_{\dot{H}^{n-1}} + \|g\|_{L^\infty} \|f\|_{\dot{H}^n}). \end{aligned}$$

Cf. Appendix in [20].

**Lemma 11.** *Let  $u \in \mathcal{S}(\mathbb{R}^d)$  solve the Poisson equation  $-\Delta u(x) = f(x)$  for a.e.  $x \in \mathbb{R}^d$ . Then*

$$\sup_{1 \leq i, j \leq d} \left\| \frac{\partial^2 u}{\partial x_i \partial x_j} \right\|_{L^p} \leq C \|f\|_{L^p}$$

for any  $1 < p < \infty$  and

$$\sup_{1 \leq i, j \leq d} \left[ \frac{\partial^2 u}{\partial x_i \partial x_j} \right]_\alpha \leq C [f]_\alpha$$

for any  $0 < \alpha < 1$ . Here  $[\cdot]_\alpha$  denotes the Hölder seminorm and  $C$  depends only on  $p$  and  $d$ .

Cf. Corollary 7.7 in [28].

## Acknowledgement

We acknowledge support from the Austrian Science Fund (FWF) via the SFB project 'Taming Complexity in Partial Differential Equations' No. F 65 and project No. W1450 and by the Vienna Science and Technology Fund (WWTF) project MA16-W1450 and by the Vienna Science and Technology Fund (WWTF) project MA16-066.

C.Y. acknowledges the hospitality and support of the Wolfgang Pauli Institute Vienna which boosted this research. Further C.Y. expresses his gratitude towards Zhennan Zhou from Peking University for continuous support.

## References

- [1] R.A. Adams and J.J.F. Fournier. *Sobolev Spaces*. ISSN. Elsevier Science, (2003).
- [2] T. Alazard and R. Carles. Semi-classical limit of Schrödinger–Poisson equations in space dimension  $n \geq 3$ , *J. Diff. Eq.* **233** (2007) 241–275.
- [3] R. Carles R. Danchin and J.C. Saut. Madelung, Gross–Pitaevskii and Korteweg, *Nonlinearity* **25**(10) (2012) 2843–2873.
- [4] H. Brezis and P. Mironescu. Gagliardo–Nirenberg inequalities and non-inequalities: The full story. *Ann. Inst. H. Poincaré C, Analyse non linéaire* **35** (2018) 1355–1376.
- [5] R. Carles. WKB analysis for nonlinear Schrödinger equations with potential. *Comm. Math. Phys.*, **269**(1) (2007) 195–221.
- [6] S. Engelberg, H. Liu and E. Tadmor. Critical Thresholds in Euler-Poisson Equations. *Ind. Univ. Math. J.* **50**(1) (2001) 109–157.
- [7] L. Erdős and J. P. Solovej. Semiclassical eigenvalue estimates for the Pauli operator with strong non-homogeneous magnetic fields: II. Leading order asymptotic estimates. *Comm. Math. Phys.* **188** (1997) 599–656.
- [8] P. Gérard, P.A. Markowich, N.J. Mauser and F. Poupaud. Homogenization limits and Wigner transforms. *Comm. Pure Appl. Math.* **50** (1997) 321–377.
- [9] P. Gérard. Mesures semi-classiques et ondes de Bloch. *Sém. Éq. dér. part.* **16** (1991) 1–19.
- [10] P. Germain and N. Masmoudi. Global existence for the Euler-Maxwell system. *Ann. Scient. Éc. Norm. Sup.* **4**(47) (2014) 469–503
- [11] P. Germain, N.J. Mauser and J. Möller. Weak and strong solutions of the Pauli-Poiswell equation, *Manuscript* (2023).
- [12] F. Golse and T. Paul. Mean-Field and Classical Limit for the  $N$ -Body Quantum Dynamics with Coulomb Interaction. *Comm. Pure Appl. Math.* **75**, 1332–1376 (2022).

- [13] E. Grenier. Semiclassical limit of the nonlinear Schrödinger equation in small time. *Proc. Am. Math. Soc.*, **126**(2) (1998) 523–530.
- [14] G. Gui and P. Zhang. Semiclassical Limit of Gross–Pitaevskii Equation with Dirichlet Boundary Condition. *SIAM J. Math. Anal.* **54**(1) (2022) 1053–1104.
- [15] Y. Guo. Smooth Irrotational Flows in the Large to the Euler–Poisson System in  $\mathbb{R}^3+1$ . *Comm. Math. Phys.* **195** (1998), 249 – 265.
- [16] Y. Guo, A. Ionescu, and B. Pausader Global solutions of the Euler–Maxwell two-fluid system in 3D. *Ann. Math.* **183**(2) (2016) 377–498.
- [17] Y. Guo and B. Pausader. Global Smooth Ion Dynamics in the Euler-Poisson System. *Comm. Math. Phys.* **303** (2011) 89–125.
- [18] M. Hadzic and J. Juhi Jang. A Class of Global Solutions to the Euler–Poisson System. *Comm. Math. Phys.* **370** (2019) 475–505.
- [19] S. Keppeler. Semiclassical quantisation rules for the Dirac and Pauli equations. *Prog. Theo. Phys.* **304** (2003) 40–71.
- [20] S. Klainerman and A. Majda. Singular limits of quasilinear hyperbolic systems with large parameters and the incompressible limit of compressible fluids. *Comm. Pure Appl. Math.*, **34**(4) (1981) 481–524.
- [21] P.L. Lions and T. Paul. Sur les mesures de Wigner, *Rev. Mat. Ib.* **9**(3) (1993) 553–618.
- [22] P.A. Markowich and N.J. Mauser. The classical limit of a self-consistent quantum-Vlasov equation in 3D, *Math. Mod. Meth. Appl. Sc.* **3**(1) (1993) 109–124.
- [23] N. Masmoudi and N.J. Mauser. The selfconsistent Pauli equation, *Monatshefte Math.*, **123**(1) (2001).
- [24] J. Möller and N.J. Mauser. The semiclassical limit of the Pauli-Poiswell equation by the Wigner method, *Manuscript* (2023).
- [25] J. Möller. The Pauli-Poisson equation and its semiclassical limit, *Submitted*, (2023).
- [26] J. Möller and N.J. Mauser. Nonlinear PDE models in semi-relativistic quantum physics, *Submitted*, (2023).
- [27] J. Möller. (Asymptotic) Analysis of the Pauli-Poiswell equation in semi-relativistic Quantum Physics., *PhD Thesis*, (2023).
- [28] C. Muscalu and W. Schlag. Classical and Multilinear Harmonic Analysis, *Cambridge Studies Adv. Math.* (2013).
- [29] Y. Peng and S. Wang. Convergence of Compressible Euler-Maxwell Equations to Compressible Euler-Poisson Equations. *Chin. Ann. Math.* **28B**(5) (2007), 583-602.

- [30] B. Perthame. Non-existence of global solutions to Euler-Poisson equations for repulsive forces. *Jap. J. Appl. Math.* **7**(2) (1990) 363–367.
- [31] C. Sparber, P. Markowich and N.J. Mauser. Wigner functions versus WKB-methods in multivalued geometrical optics. *Asym. Anal.*, **33**(2) (2003) 153–187
- [32] H. Yamasaki. A New Derivation of Classical Models of the Spinning Electron from the WKB Solutions to the Pauli and Dirac Equations. *Prog. Theo. Phys.* **36**(1) (1966) 72–84.
- [33] J. Yang and S. Wang. Non-relativistic limit of two-fluid Euler-Maxwell equations arising from plasma physics. *Z. Angew. Math. Mech.* **89**(12) (2009) 981 – 994.
- [34] M. Yuen. Blowup for the Euler and Euler–Poisson equations with repulsive forces. *Nonl. Anal.* **74** (2011) 1465-1470.
- [35] P. Zhang. Wigner measure and the semiclassical limit of Schrödinger–Poisson equations. *SIAM J. Math. Anal.* **34**(3) (2003) 700–718.
- [36] P. Zhang. Wigner measure and semiclassical limits of nonlinear Schrödinger equations, *Courant Lecture Notes* **17** (2008).
- [37] P. Zhang, Y. Zheng, and N. J. Mauser. The limit from the Schrödinger-Poisson to the Vlasov-Poisson equations with general data in one dimension. *Comm. Pure Appl. Math.* **55**(5) (2002) 582–632.