

The ground state of the Kondo lattice

Igor N. Karnaukhov¹

¹G.V. Kurdyumov Institute for Metal Physics, 36 Vernadsky Boulevard, 03142 Kyiv, Ukraine
*karnai@yahoo.com

ABSTRACT

The Kondo insulator state (KIS) is among the least understood phase state in condensed matter physics. KIS is the state of the electron liquid in the Kondo lattice at half filling, is studied within the mean field approach. We demonstrate, that Z_2 -field, which is formed by electrons and local moments, leads to the state of the Kondo insulator in a lattice with a double cell. In the ground state, electrons and local moments form singlets; in this case, no spin or charge density waves are realized in a lattice with a double cell. We have shown that a Majorana-type gap spectrum of the quasi-particle excitations is realized in Z_2 -field. The gap in the spectrum decreases with increasing external magnetic field; it closes at a critical value at the insulator-metal phase transition point. Thus, the introduction of the Z_2 -field allows you to answer the key question what is the ground state of KIS.

Introduction

In contrast to the Kondo problem, the exact solution of which was obtained by a weak interaction in continuum approach^{1,2}, the behavior (phase state) of an electron liquid in the Kondo lattice is an unsolved problem of condensed matter physics. The scattering electrons on a local moment with spin flip leads to the Abrikosov-Suhl resonance, a new behavior of an electron liquid in the Kondo problem at low temperatures and magnetic field.

Speaking of the Kondo lattice, we do not know the answers to simple but important questions: what is the ground state of the electron liquid in KIS, why is there a large Fermi surface at conservation of the number of electrons, what is the nature of the charge and spin gaps in the excitation spectrum³⁻⁹. When solving the Kondo lattice problem, it is also necessary to take into account the scattering of electrons by local moments with spin flip (as it takes place in the Kondo problem). The effective Hamiltonian should also not break the symmetry of the model. This is a non-trivial problem that has not yet been solved, so we cannot say anything definitive about what the ground state is implemented in KIS.

However, despite this pessimistic introduction, the purpose of the article is to answer a main question, what is the ground state of the Kondo lattice at half filling. The antiferromagnetic exchange interaction between electrons and local moments leads to a Z_2 -field whose uniform configuration forms a lattice with a double cell in KIS. Note that charge or spin density waves are not realized in a lattice with a double cell; in this sense, this is an unusual phase state.

Model

The Hamiltonian of the spin- $\frac{1}{2}$ Kondo lattice dimension D $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_K$ includes two terms, the first of which is determined by energy of electrons, the second is determined by the contact exchange interaction

of these electrons with moments located at the lattice sites

$$\begin{aligned}\mathcal{H}_0 &= - \sum_{\langle i,j \rangle} \sum_{\sigma=\uparrow,\downarrow} c_{i\sigma}^\dagger c_{j\sigma} - 2h \sum_j (s_j^z + S_j^z), \\ \mathcal{H}_K &= 2 \sum_{j=1}^N [J s_j^z S_j^z + K (s_j^x S_j^x + s_j^y S_j^y)],\end{aligned}\quad (1)$$

where $c_{j\sigma}^\dagger$ and $c_{j\sigma}$ are the fermion operators determined on a lattice site j , $\sigma = \uparrow, \downarrow$ denotes the spin of electron, the hopping integral between the nearest-neighbor lattice sites is equal to one, the spin operators of electrons $s_j^\alpha = \frac{1}{2} c_{j\sigma}^\dagger \sigma_{\sigma\sigma'}^\alpha c_{j\sigma'}$ are determined by the Pauli matrices σ^α ($\alpha = x, y, z$), \mathbf{S}_j is the spin- $\frac{1}{2}$ operator defined on the lattice site j (S is its value), $J \geq 0$ and $K > 0$ are the magnitudes of the exchange interaction ($K = J > 0$ corresponds to an isotropic antiferromagnetic exchange interaction, $J = 0, K > 0$ corresponds to a strong anisotropic interaction), h is an external magnetic field (g -factor is 2, we assume the Bohr magneton is 1), N is the total number of lattice sites. We study the behavior of an electron liquid in the chain (1D) and on the square (2D) and cubic (3D) lattices at half-filling.

The ground-state of the Kondo lattice

We consider the term of the Hamiltonian (1) \mathcal{H}_K in detail in the following form

$$\mathcal{H}_K = \sum_{j=1}^N [2J s_j^z S_j^z + K (s_j^+ S_j^- + s_j^- S_j^+)], \quad (2)$$

where the spin operators redefine via the fermion operators, $s_j^z = \frac{1}{2}(m_{j\uparrow} - m_{j\downarrow})$, $s_j^+ = c_{j\uparrow}^\dagger c_{j\downarrow}$, $s_j^- = c_{j\downarrow}^\dagger c_{j\uparrow}$, here $m_{j\sigma} = c_{j\sigma}^\dagger c_{j\sigma}$, $m_j = m_{j\uparrow} + m_{j\downarrow}$ are the density operators.

We use the following presentation for the \mathcal{H}_K term: $2J s_j^z S_j^z + K (s_j^+ S_j^- + s_j^- S_j^+) = -J (s_j^z - S_j^z)^2 - K (s_j^+ - S_j^+)(s_j^- - S_j^-) + (J - K)(S_j^z)^2 - \frac{1}{4}(J + 2K)m_j^2 + \frac{1}{2}(J + K)m_j + KS(S + 1) \implies 2\Lambda_j (s_j^z - S_j^z) + \lambda_j (c_{j\uparrow}^\dagger c_{j\downarrow} - S_j^+) + \lambda_j^* (c_{j\downarrow}^\dagger c_{j\uparrow} - S_j^-) + 2\mu_j m_j$. Using the Hubbard-Stratanovich transformation we introduce the effective Hamiltonian which is determined by two component $\Lambda - \lambda$ -field. We will study in detail the case $S = \frac{1}{2}$. $(S_j^z)^2$ -operator is conserved and μ -component shifts the Fermi energy, so they can be neglected. We can define an effective Hamiltonian \mathcal{H}_{eff} , which describes the behavior of the electron liquid in the Kondo lattice in the mean field approach

$$\mathcal{H}_{eff} = \mathcal{H}_0 \sum_j [2\Lambda_j (s_j^z - S_j^z) + \lambda_j (c_{j\uparrow}^\dagger c_{j\downarrow} - S_j^+) + \lambda_j^* (c_{j\downarrow}^\dagger c_{j\uparrow} - S_j^-) + \sum_j [\frac{\Lambda_j^2}{J} + \frac{|\lambda_j|^2}{K}]].$$

Let us consider equations for the one-particle wave functions $\psi(\mathbf{j}, \sigma) c_{j\sigma}^\dagger + \phi(\mathbf{j}, \pm\sigma) S_j^\pm$ ($\sigma = \uparrow, \downarrow$) with energy ε , the $\psi(\mathbf{j}, \sigma)$ and $\phi(\mathbf{j}, \sigma)$ amplitudes satisfy the following equations :

$$\begin{aligned}(\varepsilon - \Lambda_j) \psi(\mathbf{j}, \sigma) + \lambda_j \psi(\mathbf{j}, -\sigma) + \sum_{\mathbf{1}} \psi(\mathbf{j}+\mathbf{1}, \sigma) &= 0, \\ (\varepsilon + \Lambda_j) \psi(\mathbf{j}, -\sigma) + \lambda_j^* \psi(\mathbf{j}, \sigma) + \sum_{\mathbf{1}} \psi(\mathbf{j}+\mathbf{1}, -\sigma) &= 0, \\ (\varepsilon + \Lambda_j) \phi(\mathbf{j}, \sigma) - \lambda_j \phi(\mathbf{j}, -\sigma) &= 0, \\ (\varepsilon - \Lambda_j) \phi(\mathbf{j}, -\sigma) - \lambda_j^* \phi(\mathbf{j}, \sigma) &= 0,\end{aligned}\quad (3)$$

where sums over the nearest lattice sites. The variables $\Lambda_j = \pm\Lambda$ and $\lambda_j = \pm\lambda$ are identified with a static two component \mathbb{Z}_2 -field determined on the lattice sites. A configuration of this field, which corresponds to an energy minimum, defines the ground state. The local moments form the flat band

states, with the energies $\varepsilon_S = \pm\sqrt{\Lambda^2 + \lambda^2}$. The moments are located at the lattice sites, their energy is not determined by its spin.

Detailed numerical analysis shows, that an uniform sector with $\Lambda_j = \Lambda$, $\lambda_j = \lambda$ and $\Lambda_{j+1} = -\Lambda$, $\lambda_{j+1} = -\lambda$ corresponds to the ground state of an electron liquid for arbitrary values of J and K ¹⁰⁻¹³. Using Eqs(3) we calculate the energy of quasi-particle excitations wich correspond to this uniform configuration of the Z_2 -field,. The spectrum includes two branches of local moments ε_S and two branches of electrons $\varepsilon_s(\mathbf{k}) = \pm\sqrt{\Lambda^2 + \lambda^2 + |w(\mathbf{k})|^2}$, here $w(\mathbf{k}) = \sum_{\alpha}^D [1 + \exp(ik_{\alpha})]$, $\mathbf{k} = (k_x, k_y, k_z)$ is the wave vector. The solution for the values of the Z_2 -field components satisfies the energy minimum or the saddle point of the action. For isotropic ($J = K > 0$, here $\Lambda = \lambda \neq 0$) and anisotropic ($J = 0, K > 0$, here $\Lambda = 0, \lambda \neq 0$) exchange interaction, a self-consistent equation has the following form at $T = 0K$

$$\frac{2\lambda}{J} = \frac{1}{N} \sum_{\mathbf{k}} \frac{\lambda}{|\varepsilon_s(\mathbf{k})|} + \frac{\lambda}{|\varepsilon_S|}. \quad (4)$$

The behavior of the electron liquid in the case of strongly anisotropic, when $J = 0, K > 0$, and isotropic, when $J = K > 0$, exchange antiferromagnetic interaction will be considered in detail.

Strongly anisotropic exchange interaction $J = 0, K > 0$

For a strongly anisotropic exchange interaction, the Z_2 - field is one-component, since $\Lambda = 0$ and $\lambda \neq 0$. According to the numerical analysis, the solutions $\lambda_j = \lambda$ and $\lambda_{j+1} = -\lambda$ correspond to the minimum energy and for an arbitrary values of the exchange integral K and magnetic field h .

In magnetic field the energies of the quasi-particle excitations transform to $\varepsilon_{s,+}(\mathbf{k}) = \pm\sqrt{\Lambda^2 + \lambda^2 + (h + |w(\mathbf{k})|)^2}$, $\varepsilon_{s,-}(\mathbf{k}) = \pm\sqrt{\Lambda^2 + \lambda^2 + (h - |w(\mathbf{k})|)^2}$, $\varepsilon_S = \pm\sqrt{\Lambda^2 + \lambda^2 + h^2}$. The spectrum is symmetrical with respect to zero energy or chemical potential, which is zero at half filling.

In the electron spectrum the gap opens at $\lambda \neq 0$ and is equal to 2λ at $\Lambda = 0$, according to (4) its value is determined by K and h . Using Eq (4) we numerically calculate λ as function of K and h for the chain Fig1a), square Fig1b) and cubic Fig1c) lattices. Should be note an universal behavior of an electron liquid in KIS, the curves in Figs are similar for an arbitrary dimension. In a weak coupling limit at $K \rightarrow 0$ the last term in Eq (4) dominates, so $\lambda \rightarrow \frac{K}{2}$. We illustrate the spectrum of the quasi-particle excitations in Fig2 a) for the chain and Fig2 b) for square lattice. The magnetic field breaks the spin degeneracy of the spectrum of electrons, spreading the branches of the electron spectrum. The λ - value (or the value of the gap) decreases with magnetic field. A critical value of the magnetic field h_c , at which the gap closes, depends on K -value. Numerical calculations of h_c are shown in Fig 3 (the curves are calculated for different dimension of the model). In magnetic field h_c the phase transition from insulator state to metal state is realized, in other words KIS is stable at $h < h_c$.

Magnetic properties of an electron liquid in KIS are determined by both band electrons and local moments, they determine the uniform configuration of the Z_2 -field. Electrons and local moments form singlet states in the lattice with a double cell, which are not fixed in time. These singlet states (with different spins of electrons and local moments) are degenerated in energy in absence of magnetic field, so the magnetization density $M = \frac{2}{N} \sum_j (s_j^z + S_j^z)$ is zero. The magnetic field does not break this degeration of energy for local moments, so the magnetization is detemined by an electron term $M = \frac{1}{N} \sum_j (m_{j,\uparrow} - m_{j,\downarrow})$:

$$M = \frac{h}{N} \sum_{\mathbf{k}} \frac{1}{|\varepsilon_{s,+}(\mathbf{k})| + |\varepsilon_{s,-}(\mathbf{k})|} \left(\frac{\lambda^2 + h^2 - |W(\mathbf{k})|^2}{|\varepsilon_{s,+}(\mathbf{k})\varepsilon_{s,-}(\mathbf{k})|} + 1 \right). \quad (5)$$

The calculations of the magnetization density M as function of magnetic field and the exchange integral K are presented in Figs 4 for different dimension of the model. Formula for a static magnetic susceptibility




Figure 1. (Color online) λ –value as a function of the exchange integral K and magnetic field h , calculated for the chain a), square b) and cubic c) lattices .




Figure 2. (Color online) The spectrum of quasi-particle excitations of electron liquid (the gap in the spectrum of charge excitations is equal to 2λ) in the chain a) and square lattice b) as a function of the wave vector, calculated for $\lambda = 0.3$, $h = 0.2$ ($K = 0.524$ for chain and $K = 0.597$ for square lattice).




Figure 3. (Color online) Critical value of magnetic field h_c , at which the gap in the quasi-particle spectrum closes, as a function of the exchange integral, calculated for different dimension of the Kondo lattice.




Figure 4. (Color online) Magnetization density as a function of the exchange integral K and magnetic field h , calculated for the chain a), square b) and cubic c) lattices .




Figure 5. (Color online) Static magnetic susceptibility as a function of an exchange integral calculated for different dimension of the model.




Figure 6. (Color online) The excitation energies corresponding to the Z_2 -field configuration with ν "defects" are calculated for various ν (marked with dots, the lines correspond to the excitation branches). Calculations were carried out for a chain with $\lambda = 0.3$, $h = 0.1$, $K = 0.476$.

leads from M at $h \rightarrow 0$ $\chi = \frac{1}{N} \sum_{\mathbf{k}} \frac{\lambda^2}{(\lambda^2 + |W(\mathbf{k})|)^{3/2}}$. The value of a static magnetic susceptibility is calculated a function of the exchange integral for different dimension of the model, calculations are shown in Fig 5, the susceptibility is a monotonic function of K .

An uniform configuration stabilises the phase state with a double cell, a free configuration corresponds to gapless state with higher energy. According to numerical calculations, a total energy of individual ν "defects" is greater than the energy of one "defect" of size ν . One "defect" with $\nu = 1$ corresponds to a Z_2 -field configuration with an incorrect value of λ_j , namely $-\lambda_j$ instead of λ_j , which occurs in a uniform configuration. As an example, we present the calculations of the excitation energies in a chain with "defects" as a function of the defect size ν . Configurations with "defects" in an uniform configuration have energies lying in the gap, a number of excitations increases with increasing ν , so that for $1 < \nu < 5$ only one excitation is split off from the continuous spectrum, for $5 < \nu < 10$, $10 < \nu < 15$ there are 2 and 3 such states, respectively (see in Fig 6). We note, that the lattice with a double cell is formed by an uniform configuration of λ -field, with no spin or charge density waves being realized.

Isotropic exchange interaction $J = K > 0$

As noted above, in the absence of a magnetic field for an isotropic exchange interaction, the λ - and Λ -components of the Z_2 -fields are equal and are solutions of Eq (3). Along with this solution, there are also a number of non-trivial solutions: $\lambda \neq 0$ and $\Lambda = 0$, $\lambda = 0$ and $\Lambda \neq 0$. Three solutions of Eq (3) have the same energy of the ground state, which follows from numerical calculations of the ground state energy as function of J at $h = 0$ for different dimensions of the model. In the absence of an external magnetic field, the phase state of the electron liquid is degenerate. A magnetic field removes this degeneracy, namely, the solution $\lambda \neq 0$ and $\Lambda = 0$ corresponds to a lower energy than solution $\lambda = 0$ and $\Lambda \neq 0$ for an arbitrary value of magnetic field and an isotropic exchange integral. Another nontrivial solution $\lambda \neq 0, \Lambda \neq 0$ not satisfy Eq(3) for arbitrary h . KIS is determined by the XX -exchange interaction (the value of K in Hamiltonian (1)), the ZZ -exchange interaction (the value of J in Hamiltonian (1)) does not participate in the formation of KIS. Scattering processes with spin flip lead to the formation of KIS in the Kondo lattice as it takes place in the Kondo problem.

In the case of an arbitrary spin S , Eq (4) for the amplitudes of the electron wave function does not change. The energies of local moments change, so for $S = 3/2$ $\epsilon_S = \pm \frac{1}{2}(-1 \pm \sqrt{5})\lambda$. As a result, the formula for the magnetization (5) does not depend on the magnetization of local moments, Eqs (3) are transformed taking into account the value of ϵ_S .

Conclusion

We studied the behavior of electron liquid in the spin- $\frac{1}{2}$ Kondo lattice at half-filling for different dimension. Due to antiferromagnetic exchange interaction between band electrons and local moments a static Z_2 - field is formed. An uniform configuration of the Z_2 -field corresponds to ground state of electron liquid in KIS, leads to formation of a lattice with a double cell. The spin or charge density waves are not realized in KIS. In KIS the spectrum of quasi-particle excitations is symmetric about zero energy, as it takes place for the Majorana spectrum. At a critical value of a magnetic field, at which a gap closes, the phase transition to metal state is realized.

Acknowledgments

The author thanks the Weizmann Institute of Science and personally Prof. E. Berg for support.

References

1. Andrei N., K. Furuya K., & Lowenstein J. H. Solution of the Kondo problem. *Rev. Mod. Phys.*, **55**, 331 (1983)
2. Tsvetick, A.M. & Wiegmann, P.B. Exact results in the theory of magnetic alloys. *Advances in Physics* **32**, 453 (1983)
3. Coleman, P. & Andrei, N. Kondo-stabilised spin liquids and heavy fermion superconductivity. *J.Phys.:Condens. Matter.* **1**, 4057 (1989)
4. Yang, W.-W., Zhao, J., Luo, H.-G., & Zhong, Y. Exactly solvable Kondo lattice model in the anisotropic limit. *Phys. Rev. B* **100**, 045148 (2019)
5. Karnaukhov, I.N. Exactly solvable model of a one-dimensional Kondo lattice. *Phys.Rev.B* **56**, R4313 (1997)
6. Karnaukhov, I.N. & Slietsov I.O. Exactly solvable 2D topological Kondo lattice model. *Europhysics Letters* **109**, 57005 (2015)
7. Sykora, S. & Becker, K.W. Heavy fermion properties of the Kondo Lattice model. *Scientific Reports* **3**, 2691 (2013)
8. Lavagna, M. & Pépin, C. The Kondo lattice model. *Acta physica polonica B.* **29**, 3753 (1998)
9. Karnaukhov, I.N. Electron liquid state in the symmetric Anderson lattice. *Scientific Reports* **11**, 5842 (2021).
10. Karnaukhov, I.N. Electron liquid state in the spin- anisotropic Kondo lattice, *Scientific Reports* **12**, 74202 (2022).
11. Karnaukhov, I.N. Gapped electron liquid state in the symmetric Anderson lattice, Kondo insulator state, *Scientific Reports* **12**, 18607 (2022).
12. Lieb, R.H. Flux phase of the half-filled band. *Phys. Rev. Lett.* **73**, 2158–2161 (1994).
13. Kitaev, A.Yu. Anyons in an exactly solved model and beyond. *Annals of Physics* **321**, 2-111 (2006)

Author contributions statement

I.K. is an author of the manuscript

Additional information

The author declares no competing financial interests.

Availability of Data and Materials

All data generated or analysed during this study are included in this published article. Correspondence and requests for materials should be addressed to I.N.K.