

Categorical Quantum Groups and Braided Monoidal 2-Categories

Hank Chen¹ and Florian Girelli¹

¹Department of Applied Mathematics, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, Canada, N2L 3G1

June 13, 2023

Abstract

We develop a systematic *categorification* of the theory of quantum groups/bialgebras following the dimensional ladder, and study their higher-representation theory. By following closely the generalized quantum double construction of Majid, we construct in particular the *2-quantum double* $D(\mathcal{G})$ associated to a 2-bialgebra \mathcal{G} , and prove its duality and factorization properties. We also characterize a notion of (quasi)triangular *2-R-matrix* \mathcal{R} and identify the associated (2-)Yang-Baxter equations, which can be seen as a categorification of the usual notion of *R-matrix* in an ordinary quantum group. The main result we prove in this paper is that the weak 2-representation 2-category $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ of a quasitriangular 2-bialgebra $(\mathcal{G}, \mathcal{T}, \mathcal{R})$ — when monoidally weakened by a Hochschild 3-cocycle \mathcal{T} — forms a braided monoidal 2-category.

Contents

1	Introduction	2
2	Strict 2-bialgebras	4
2.1	Associative 2-algebras	4
2.1.1	Classification of 2-groups and associative 2-algebras	6
2.1.2	Example: group 2-algebras from 2-groups	7
2.2	Associative 2-bialgebras	8
2.2.1	Definition	8
2.2.2	Example: function 2-bialgebras on 2-groups	11
3	Strict 2-quantum doubles and the universal 2-R-matrix	13
3.1	Matched pair of 2-(bi)algebras	13
3.2	Construction of the strict 2-quantum double	15
3.3	Factorizability of 2-bialgebras	17
3.4	Characterization of quantum 2-R-matrices	19
4	Weak 2-bialgebras	21
4.1	Definition	21
4.1.1	Weak 2-algebra homomorphisms	22
4.1.2	Example: weak 2-group algebras	22
4.2	Weak 2-coalgebras	22
4.3	Weak 2-bialgebras	23
5	Weak (skeletal) 2-quantum double	24
5.1	Matched pair of skeletal weak 2-bialgebras	25
5.2	Factorizability of weak 2-bialgebras	25

*hank.chen@uwaterloo.ca

†fgirelli@uwaterloo.ca

6	The fusion 2-category of 2-representations	26
6.1	Weak 2-representations	27
6.1.1	Endomorphism 2-algebra on a 2-vector space	27
6.1.2	Weak 2-representations, weak 2-intertwiners and modifications	28
6.2	Fusion structure on the 2-representations	30
6.2.1	Monoidal structure	30
6.2.2	Naturality and Gray property of the tensor product	31
6.3	Fusion associators	33
7	The braided monoidal 2-category of 2-representations	35
7.1	The braiding maps and their naturality	35
7.2	Braided 2-quasi-bialgebras; the modified hexagon relations	38
7.3	The braiding hexagonator: weak 2-representations of a braided 2-quasi-bialgebra	39
7.4	Main theorem and its proof	41
8	Conclusion	44
A	2-Hopf algebras	46
B	Classical limits of (weak) 2-bialgebras and 2-R-matrices	47
B.1	Lie 2-bialgebras and the 2-classical double	47
B.2	The classical 2- r -matrix	48
B.3	Weak Lie 2-bialgebras	50
C	Module 2-categories and 2-representation theory	51
C.1	Action 2-functors	51
C.2	Higher-representation theory	52

1 Introduction

It is well-known that the algebra of excitations in 3D BF theory (equivalent to a Chern-Simons theory [1, 2, 3]) for a given gauge group is described by its Drinfel'd double quantum group [4, 5, 2], which forms a quasitriangular Hopf algebra [6, 7]. The tensor category of its representations has equipped very important monoidal structures that capture the essential topological properties of the underlying topological field theory. In particular, the universal R -matrix equipped on a Hopf algebra H defines the braiding map $b = \text{flip} \circ R$ on its representation category $\text{Rep}(H)$.

The fact that the modules of such quantum group Hopf algebras are braided means that $\text{Rep}(H)$ furnishes a representation of the Artin braid group [6], which allows quantum invariants of knots and tangles to be developed from the Hopf algebra machinery. This was first pointed out by Witten [8] in which the Jones polynomial knot invariants were recovered from modules of the quantum $U_q(\mathfrak{su}(2))$ symmetry of $SU(2)$ -Chern-Simons theory. Indeed, the deep connection between Hopf algebras and 3d geometry/topology is well-known [9, 10].

The well-known 2d toric code [11] is another fruitful example of such construction. It has been shown to be described by a 3d \mathbb{Z}_2 -gauge BF theory with the underlying Drinfel'd double $D(\mathbb{Z}_2)$ symmetry. Moreover, the theory of non-degenerate fusion categories, modelling 3-dimensional gapped topological phases and their boundary excitations, are now very well-understood [12, 13, 14].

The characterization of quasitriangular Hopf algebras/quantum groups, as well as its R -matrix, plays also a central role in the theory of quantum and classical integrable systems [15, 16, 17, 18]. In particular for 1-dimensional spin chains such as the Toda lattice [16] or the XXX/XXZ/XYZ family of Ising spins [19], obtaining the quantum R -matrix is equivalent to solving the entire model, through the method known as *quantum inverse scattering*.

One might wonder how these different examples extend to a dimension up. For example, one might wonder what is the right tool to characterize the algebra of excitations in a $4d$ BF theory, or any higher-dimensional topological phases in general [20, 21, 22], such as the 4d toric code [23, 24, 25]. In a similar way, given the well-studied integrable systems are typically in 1+1 dimension, one could wonder what would be the relevant structure for an integrable system in 2+1 dimension.

According to the dimensional (or categorical) ladder proposal [26, 27, 28], one way to obtain the relevant structure is through *categorification*. In fact, several constructions already point to the efficiency of using 2-categories (arising from for instance higher-representation theory [29, 30, 31, 32]) to describe a 4-dimensional gapped topological phases, such as the 4d toric code [23, 24, 25].

The goal of this paper is to describe a systemic categorification of the rich and fruitful theory of quantum groups, starting from the theory of higher-dimensional L_∞ -algebras of [33, 34, 35]. Together with the quantization step, which we understand as taking a Lie algebra to a quantum Hopf algebras (cf. Drinfel'd-Jimbo quantization [36, 37, 38] and Kontsevich \star -quantization [39]), we arrive at the following diagram,

$$\begin{array}{ccc}
(\text{Poisson-})\text{Lie algebra} & \xrightarrow{\text{categorify}} & (\text{Poisson}) L_\infty\text{-algebras} \\
\downarrow \text{quantize} & & \downarrow \text{quantize} \\
(\text{Hopf}) \text{ algebras} & \xrightarrow{\text{categorify}} & (\text{Hopf}) A_\infty\text{-algebras}
\end{array} \quad . \quad (1.1)$$

As such, focusing on 4-dimensions, it follows that a **categorical quantum group** living on the bottom-right corner of (1.1) can be modelled as a "2-bialgebra" \mathcal{G} in the A_∞ context. We shall construct such a structure starting from the strict/associative 2-algebras of [40], then consider a weakening of the associativity up to homotopy expressed by a *Hochschild 3-cocycle* \mathcal{T} on \mathcal{G} . Moreover, we will substantiate the diagram (1.1) in Section B, where we prove that we recover the known notions of Lie 2-bialgebras [41, 42, 43] by taking an appropriate "classical limit".

A structure of particular interest that we will also construct is the *2-quantum double*, which is a "categorical Drinfel'd double" generalizing the works of Majid [6, 44, 7, 36]. We will prove some crucial structural theorems about this 2-quantum double construction, such as quasitriangularity and self-duality. We explicitly demonstrate that these proofs run parallel to classical results for ordinary quantum doubles.

Next, we consider higher-representations of the weak 2-bialgebra \mathcal{G} by examining its action on 2-vector spaces. We emphasize that the 2-representation theory based on the Baez-Crans notion of 2-vector spaces [45] $2\text{Vect}_{\text{int}}$, which is equivalently a category internal to Vect [33], is insufficient for our purposes. As such, we shall base our "weak 2-representation theory" on a modified version of $2\text{Vect}_{\text{int}}$, which we shall denote simply by 2Vect , which has the key property that the endomorphism 1-categories on each class of its simple objects have the structure of a 2-term A_∞ -algebra. We denote the resulting 2-category of weak 2-representations by $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$.

This modification is necessary in order for our weak 2-representations to carry higher-homotopical data, which play very important roles in the following main result of this paper.

Theorem 1.1. *The 2-representation 2-category $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ of a weak quasitriangular 2-bialgebra \mathcal{G} forms a braided monoidal 2-category with trivial left-/right-unitors.*

This paper will be dedicated towards the proof of this theorem. We will, in particular, universally characterize the quantum 2- R -matrices on \mathcal{G} which is responsible for the braiding, and provide the **(categorified) Yang-Baxter equations** it satisfies. The theorem is then proven by relating the 2-algebraic structure to the 2-representation theory, and explicitly checking all relevant coherence diagrams [46, 47]. We will further prove in the appendix that the weak 2-representation theory we develop here is capable of reproducing the 2-representation theory of skeletal 2-groups studied in the literature [48, 32, 31, 30].

There is then naturally a forgetful functor $2\text{Rep}(\mathcal{G}) \rightarrow 2\text{Vect}$ into the k -linear semisimple 2-category of *weak* 2-vector spaces 2Vect — and *not* the Baez-Crans 2-category $2\text{Vect}_{\text{int}}$ — in which the 1- and 2-morphisms are represented respectively by certain homotopically-associative cochain maps and cochain homotopies. The precise formulation of this 2-category 2Vect is currently under investigation by one of the authors.

The paper is outlined as follows.

Section 2 constructs a *strict* 2-bialgebra (where associativity is retained) following the definition of a strict 2-algebra/algebra crossed-module in [40]. In Section 2.2, we begin by introducing a *graded* coproduct Δ on \mathcal{G} , and define the appropriate notion for two 2-bialgebras to be *dually paired*. The 2-bialgebra axiom plays a central role in this notion of duality, just as in the case of the usual 1-bialgebra. We also provide a classic example of the function 2-bialgebra on a 2-group, and verify explicitly the 2-bialgebra axioms.

Section 3 delves into the construction of the 2-quantum double $D(\mathcal{G})$ associated to a pair of mutually dual 2-bialgebras $\mathcal{G}, \mathcal{G}^*$. We prove key structural theorems about $D(\mathcal{G})$, such as its factorizability and self-duality. Furthermore, motivated by Majid's construction of the generalized double [44], based on the existence of a quantum R -matrix, we introduce the notion of 2- R -matrix \mathcal{R} and derive explicitly the "2-Yang-Baxter equations" that \mathcal{R} satisfies. We demonstrate in particular how $D(\mathcal{G})$ is in fact naturally equipped with a universal 2- R -matrix.

In Section 4, we weaken our 2-bialgebra construction by introducing a *Hochschild 3-cocycle* that witnesses associativity. We note that we obtain precisely a 2-bialgebra in the A_∞ context, which fits into the diagram (1.1) as desired. We prove that there is once again a well-defined notion of duality in this case by extending the 2-bialgebra axioms to take the weakened associativity/coassociativity into account.

In Section 5, we generalize the 2-quantum double construction, as well as their structural theorems, to the weak case. We specialize to the skeletal case, where the structural t -map is trivial, in order to leverage the construction given in the strict case.

In Section 6, we discuss directly the notion of weak 2-representations $2\text{Rep}(\mathcal{G})$, based on a modified notion of Baez-Crans 2-vector spaces living in 2Vect . We discuss the different key aspects of the fusion 2-category $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, namely the monoidal structure and the braiding. We will in particular prove their naturality from the underlying structures of the 2-bialgebra.

Finally, in Section 7, we prove the main theorem by explicitly checking all the relevant coherence diagrams of a braided monoidal 2-category. We will moreover emphasize how the fusion associators/pentagonators and braiding hexagonators arise from the higher homotopical data attached to a weak quasitriangular 2-bialgebra $(\mathcal{G}, \mathcal{T}, \mathcal{R})$, as well as its weak 2-representation theory $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$.

We collect some supplementary but also important results that we have obtained in the Appendix. In Section A, we provide an appropriate definition of an antipode in the strict case, and examine briefly its properties. In Section B, we prove that under an appropriate "classical limit", the quantum 2-bialgebra and the universal 2- R -matrix that we have defined in fact recovers the Lie 2-bialgebra and the classical 2- r -matrix as studied in [41, 42, 43]. Finally, in Section C, we will prove that the "theory of weak 2-representations" developed in this paper is capable of reproducing the 2-representation theory of skeletal 2-groups studied in the literature [31, 30, 48].

2 Strict 2-bialgebras

Quantum groups are Hopf algebras, hence we expect to define quantum 2-groups as "2-Hopf algebras". Different notions of 2-Hopf algebra have already been previously proposed in [49] and [40].

2.1 Associative 2-algebras

We begin with the following definition, and build up to the definition of an associative 2-algebra in [40].

Definition 2.1. Let $\mathcal{G}_0, \mathcal{G}_{-1}$ denote a pair of associative algebras. We say that \mathcal{G}_{-1} is a \mathcal{G}_0 -**bimodule** if we have a left and a right action¹ \cdot_l, \cdot_r of \mathcal{G}_0 on \mathcal{G}_{-1} which commute.

$$(x'x) \cdot y = x' \cdot (x \cdot y), \quad (x \cdot y) \cdot x' = x \cdot (y \cdot x'), \quad y \cdot (xx') = (y \cdot x) \cdot x' \quad (2.1)$$

for all $y \in \mathcal{G}_{-1}$ and $x', x \in \mathcal{G}_0$.

Equivalently we can demand that the following diagrams are commutative. We note μ_i the multiplication in $\mathcal{G}_i, i = -1, 0$.

The associativity of the multiplication is encoded in the usual diagram.

$$\begin{array}{ccc}
 & \mathcal{G}_i \otimes \mathcal{G}_i \otimes \mathcal{G}_i & \\
 \swarrow \text{id} \otimes \mu_i & & \searrow \mu_i \otimes \text{id} \\
 \mathcal{G}_i \otimes \mathcal{G}_i & & \mathcal{G}_i \otimes \mathcal{G}_i \\
 \searrow \mu_i & & \swarrow \mu_i \\
 & \mathcal{G}_i &
 \end{array} \quad (2.2)$$

The bimodularity conditions (2.1) read as commutative diagrams,

$$\begin{array}{ccc}
 & \mathcal{G}_0 \otimes \mathcal{G}_0 \otimes \mathcal{G}_{-1} & \\
 \swarrow \mu_0 \otimes \text{id} & & \searrow \text{id} \otimes \cdot_l \\
 \mathcal{G}_0 \otimes \mathcal{G}_{-1} & & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \\
 \searrow \cdot_l & & \swarrow \cdot_l \\
 & \mathcal{G}_{-1} &
 \end{array}
 \quad
 \begin{array}{ccc}
 & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \otimes \mathcal{G}_0 & \\
 \swarrow \text{id} \otimes \mu_0 & & \searrow \cdot_r \\
 \mathcal{G}_{-1} \otimes \mathcal{G}_0 & & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\
 \searrow \cdot_r & & \swarrow \cdot_r \\
 & \mathcal{G}_{-1} &
 \end{array} \quad (2.3)$$

¹We will often omit the subscript when there is no ambiguity.

and the bimodule condition, the middle one of (2.1) is

$$\begin{array}{ccc}
 & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_0 & \\
 \swarrow \cdot_l \otimes \text{id} & & \searrow \text{id} \otimes \cdot_r \\
 \mathcal{G}_{-1} \otimes \mathcal{G}_0 & & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \\
 \searrow \cdot_r & & \swarrow \cdot_l \\
 & \mathcal{G}_{-1} &
 \end{array} \tag{2.4}$$

If we introduce a homomorphism t between \mathcal{G}_{-1} and \mathcal{G}_0 , subject to some conditions, then \mathcal{G}_{-1} and \mathcal{G}_0 can be used to define a crossed module of algebras.

Definition 2.2. A crossed-module of (finite dimensional) associative algebras, $\mathcal{G}_0, \mathcal{G}_{-1}$, or an **associative 2-algebra**, is given by an algebra homomorphism $t : \mathcal{G}_{-1} \rightarrow \mathcal{G}_0$ where

1. \mathcal{G}_{-1} is a \mathcal{G}_0 -bimodule,
2. t is **two-sided \mathcal{G}_0 -equivariant**,

$$t(x \cdot y) = xt(y), \quad t(y \cdot x) = t(y)x \tag{2.5}$$

for all $y \in \mathcal{G}_{-1}, x \in \mathcal{G}_0$, and

3. the **Peiffer identity** is satisfied,

$$t(y) \cdot y' = yy' = y \cdot t(y'), \tag{2.6}$$

where $y, y' \in \mathcal{G}_{-1}$.

We call the latter two the *Peiffer conditions*. We denote an associative 2-algebra simply by \mathcal{G} , or by (\mathcal{G}, \cdot) to emphasize the bimodule structure. Let k denote the ground ring of the 2-vector space underlying \mathcal{G} . We call \mathcal{G} **unital** if there exists a unit map $\eta = (\eta_{-1}, \eta_0) : k \rightarrow \mathcal{G}$ such that

$$\eta_{-1}y = y\eta_{-1} = y, \quad \eta_0x = x\eta_0 = x, \quad \eta_0 \cdot y = y \cdot \eta_0 = y, \tag{2.7}$$

for all $y \in \mathcal{G}_{-1}, x \in \mathcal{G}_0$. Moreover, t should respect the units such that $t(\eta_{-1}) = \eta_0$.

Note that one may consider \mathcal{G}_{-1} first as a vector space and *define* its product with the Peiffer identity. This notion is how one may show the bijective correspondence between Lie algebra crossed-modules and 2-term L_∞ -algebras [41, 42]. However, in the skeletal case, since the Peiffer identity is empty, which forces the product on \mathcal{G}_{-1} to be trivial.

Remark 2.1. If $t \neq 0$ were non-trivial then the Peiffer conditions, together with bimodularity, imply that

$$x \cdot (yy') = (x \cdot y)y', \quad y(x \cdot y') = (y \cdot x)y', \quad (yy') \cdot x = y(y' \cdot x)$$

for each $x \in \mathcal{G}_0, y, y' \in \mathcal{G}_{-1}$. This puts strong constraints on the algebra action \cdot , which is not necessarily imposed in the skeletal $t = 0$ case.

Equivalently, we can encode the different conditions defining the 2-algebra in terms of commutative diagrams. The equivariance reads

$$\begin{array}{ccc}
 & \mathcal{G}_0 \otimes \mathcal{G}_{-1} & \\
 \swarrow \cdot_l & \xrightarrow{\text{id} \otimes t} & \searrow \\
 \mathcal{G}_{-1} & & \mathcal{G}_0 \otimes \mathcal{G}_0 \\
 \searrow t & & \swarrow \mu_0 \\
 & \mathcal{G}_0 &
 \end{array}, \quad
 \begin{array}{ccc}
 & \mathcal{G}_{-1} \otimes \mathcal{G}_0 & \\
 \swarrow \cdot_r & \xrightarrow{t \otimes \text{id}} & \searrow \\
 \mathcal{G}_{-1} & & \mathcal{G}_0 \otimes \mathcal{G}_0 \\
 \searrow t & & \swarrow \mu_0 \\
 & \mathcal{G}_0 &
 \end{array} \tag{2.8}$$

and the Peiffer identity is

$$\begin{array}{ccc}
 & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} & \\
 \swarrow t \otimes \text{id} & \xrightarrow{\text{id} \otimes t} & \searrow \\
 \mathcal{G}_0 \otimes \mathcal{G}_{-1} & & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\
 \searrow \cdot_r & \xrightarrow{\mu_{-1}} & \swarrow \cdot_l \\
 & \mathcal{G}_{-1} &
 \end{array} \tag{2.9}$$

Finally the unit map is encoded in the following commutative diagrams.

$$\begin{array}{ccc}
& & \mathcal{G}_i \otimes \mathcal{G}_i \\
& \nearrow \eta_i \otimes \text{id} & \downarrow \mu_i \\
& & \text{id} \otimes \eta_i \\
k \otimes \mathcal{G}_i & \xrightarrow{\cong} & \mathcal{G}_i \xleftarrow{\cong} \mathcal{G}_i \otimes k
\end{array}
\qquad
\begin{array}{ccc}
& & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \\
& \nearrow \eta_0 \otimes \text{id} & \downarrow \cdot r \\
& & \mathcal{G}_{-1} \\
k \otimes \mathcal{G}_{-1} & \xrightarrow{\cong} & \mathcal{G}_{-1}
\end{array}
\qquad
\begin{array}{ccc}
& & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\
& \nearrow \text{id} \otimes \eta_0 & \downarrow \cdot l \\
& & \mathcal{G}_{-1} \\
\mathcal{G}_{-1} \otimes k & \xrightarrow{\cong} & \mathcal{G}_{-1}
\end{array}
\qquad
\begin{array}{ccc}
& & \mathcal{G}_{-1} \\
& \nearrow \eta_{-1} & \downarrow t \\
& & \mathcal{G}_0 \\
k & \xrightarrow{\eta_0} & \mathcal{G}_0
\end{array}
\tag{2.10}$$

2.1.1 Classification of 2-groups and associative 2-algebras

Recall a 2-group is a connected 2-groupoid $[G_{-1}, G_0, \text{pt}]$ [30, 50], or equivalently its loop 1-groupoid $G_{-1} \times G_0 \rightrightarrows G_0$ [31]. These are equivalent to the following crossed-module description [42, 50].

Definition 2.3. The crossed-module model of a **2-group** G is a group homomorphism $t : G_{-1} \rightarrow G_0$ together with an action \triangleright of G_0 on G_{-1} such that the following conditions

$$t(x \triangleright y) = xt(y)x^{-1}, \quad (ty) \triangleright y' = yy'y^{-1} \tag{2.11}$$

are satisfied for each $x \in G_0$ and $y, y' \in G_{-1}$. The first and second conditions are known respectively as the **equivariance** and the **Peiffer identity**.

A **2-group homomorphism**, is a graded map $\phi = (\phi_{-1}, \phi_0) : G \rightarrow G'$ such that

1. $\phi_0 : G_0 \rightarrow G'_0$ and $\phi_{-1} : G_{-1} \rightarrow G'_{-1}$ are group homomorphisms,
2. $\phi_{-1}(x \triangleright y) = (\phi_0 x) \triangleright' (\phi_{-1} y)$ for each $x \in G_0, y \in G_{-1}$, and
3. $\phi_0 t = t' \phi_{-1}$.

We say that two (strict) 2-groups G, G' are elementary equivalent, or quasi-isomorphic, if there exists an invertible 2-group homomorphism between them. The fundamental classification result [51, 52] is that

Theorem 2.1. (Gerstenhaber, attr. Mac-Lane). *2-groups are classified up to quasi-isomorphism by a degree-3 group cohomology class $\tau \in H^3(N, V)$, where $N = \text{coker } t, V = \ker t$.*

τ is also called the **Postnikov class** in the literature [52, 53, 3]. Note $V = \ker t$ must be Abelian due to the Peiffer identities. The tuple (N, V, τ) is known as *Hoàng data* [53, 54], which was proven by Hoàng to classify "Gr-categories".

Similar to the case of 2-groups [33, 40], a 2-algebra homomorphism $f = (f_{-1}, f_0) : \mathcal{G} \rightarrow \mathcal{G}'$ is a graded pair of algebra homomorphisms that respect the underlying bimodule structure, such that

1. $f_0 : \mathcal{G}_0 \rightarrow \mathcal{G}'_0$ and $f_{-1} : \mathcal{G}_{-1} \rightarrow \mathcal{G}'_{-1}$ are algebra homomorphisms,
2. $f_{-1}(x \cdot y) = (f_0 x) \cdot' (f_{-1} y)$ and $f_{-1}(y \cdot x) = (f_{-1} y) \cdot' (f_0 x)$ for each $x \in \mathcal{G}_0, y \in \mathcal{G}_{-1}$, and
3. $f_0 t = t' f_{-1}$.

We say that two 2-algebras are elementary equivalent, or quasi-isomorphic, if there exists an invertible 2-algebra homomorphism between them.

Theorem 2.2. (Gerstenhaber, attr. Wagemann [40]). *Associative 2-algebras are classified up to quasi-isomorphism by a degree-3 Hochschild cohomology class $\mathcal{T} \in HH^3(\mathcal{N}, V)$, where $\mathcal{N} = \text{coker } t$ and $V = \ker t$.*

See [40] for a definition of Hochschild cohomology of an algebra. The Peiffer identity implies that $V \subset Z(\mathcal{G}_{-1})$ is in the *nucleus* of \mathcal{G}_{-1} ; it is in fact a square-free ideal [40]. Note the nucleus is *not* the same as the centre, which have commutative (but non-trivial) multiplication.

2.1.2 Example: group 2-algebras from 2-groups

Let k denote a field of characteristic zero. One example of 2-algebras comes from using a 2-group G . One way to construct a 2-algebra from G is to take the group algebra functor kG [55], and extend the t -map linearly such that $t : kG_{-1} \rightarrow kG_0$ is an algebra map. To form an associative 2-algebra, we need a kG_0 -bimodule structure on kG_{-1} , which can be induced from the group action

$$x \cdot y = x \triangleright y, \quad y \cdot x = x^{-1} \triangleright y,$$

where $x \in kG_0, y \in kG_{-1}$.

However, the subtlety here is that G_0 acts by group automorphism, not algebra automorphism,

$$x \cdot (yy') = (x \cdot y)(x \cdot y') \neq (x \cdot y)y', \quad (2.12)$$

which contradicts the condition listed in *Remark 2.1* if $t \neq 0$ or the group action $\triangleright \neq 0$ were non-trivial.

We are going to present three alternative ways to circumvent this issue, which will rely on different properties of the 2-group G . We will therefore obtain three different resulting 2-algebras kG associated to G (depending on the properties of G).

a) Wagemann's quotienting construction. The first way to guarantee the functoriality of the map $G \mapsto kG$ follows [40]. Essentially, this amounts to quotienting out certain terms such that we can recover the algebra automorphism.

We begin with the group algebra $k(G_{-1} \otimes G_0)$ equipped with the *balanced* algebra structure

$$(y, x) \cdot (y', x') = (yy' + y \cdot x' + x \cdot y', xx'), \quad x, x' \in kG_0, y, y' \in kG_{-1},$$

and let π_0 denote the projection onto kG_0 . Embed $kG_{-1} \hookrightarrow k(G_{-1} \times G_0)$ by $y \mapsto (y, 0)$ as a subalgebra, and define the ideal

$$X = kG_{-1} \cdot D + D \cdot kG_{-1}, \quad \text{with } D = \{(y, x) \in k(G_{-1} \times G_0) \mid t(y) = -x\}.$$

Explicitly, elements in the subalgebra $D \subset k(G_{-1} \times kG_0)$ satisfy $t(y) = -x$, and hence consist of pairs $(y, x) = (y, -t(y))$ parameterized by $y \in kG_{-1}$. The ideal X is then a direct sum of elements of the forms

$$\begin{aligned} (y', 0) \cdot (y, x) &= (y'y - t(y^{-1}) \triangleright y', 0) = (y'y - y^{-1}y'y, 0), \\ (y, x) \cdot (y', 0) &= (yy' - t(y) \triangleright y', 0) = (yy' - yy'y^{-1}, 0) \end{aligned}$$

using the 2-group properties, where $y' \in kG_{-1}, x \in kG_0$ are arbitrary.

We now form the algebra quotient $k(G_{-1} \otimes G_0)/X$, and denote by $\tilde{\pi}_0 : k(G_{-1} \otimes G_0)/X \rightarrow kG_0$ the induced projection map. Explicitly, this quotient consist of pairs $(y, x) \in k(G_{-1} \otimes G_0)$ such that

$$(y' - y^{-1}y')y = 0, \quad y(y' - y'y^{-1}) = 0, \quad y, y' \in kG_{-1}. \quad (2.13)$$

Put $\tilde{t} = t|_{\widetilde{kG_{-1}}}$ as the restriction of the t -map onto the kernel $\widetilde{kG_{-1}} \equiv \ker \tilde{\pi}_0$ — in other words, $\widetilde{kG_{-1}}$ consist of the G_{-1} -invariant elements of kG_{-1} under both left- and right-multiplication. The induced t -map then induces the following 2-vector space

$$kG := \widetilde{kG_{-1}} \xrightarrow{\tilde{t}} kG_0 \quad (2.14)$$

which has an associative 2-algebra structure. For a proof that this construction indeed yields an associative 2-algebra, see Theorem 3.8.3 of [40].

b) 2-group with adjoint action. The condition (2.13) is a very stringent requirement, especially in the $t = \text{id}$ case. As an alternative, we can follow a different route when the 2-group action is given by the adjoint action. In this case, to guarantee the functoriality of the map $G \mapsto kG$, one needs to find bilinear maps $\cdot : G_{-1} \otimes G_0 \oplus G_0 \otimes G_{-1} \rightarrow G_{-1}$ satisfying

$$x \triangleright y = x \cdot y \cdot x^{-1}, \quad x \in G_0, y \in G_{-1}. \quad (2.15)$$

Provided \cdot extends linearly to left- and right-regular algebra representations, one may use it directly to define the kG_0 -bimodule structure of kG_{-1} . The equivariance condition and Peiffer identity

$$\begin{aligned} xt(y \cdot x^{-1}) &= t(x \cdot y)x^{-1} = t(x \cdot y \cdot x^{-1}) = t(x \triangleright y) = xt(y)x^{-1}, \\ yy'y^{-1} &= t(y) \triangleright y' = t(y) \cdot y' \cdot t(y^{-1}). \end{aligned}$$

are directly verified. This construction is closely related to the invertibles functor $\mathcal{G} \mapsto G = \mathcal{G}^\times$, which is left-adjoint to the group algebra functor [40].

If $G = (G_{-1} \xrightarrow{t=\text{id}} G_0)$ is a *trivial* 2-group, the t -map is the identity and there is no group kernel nor cokernel. Hence it is in the trivial class under elementary equivalence [56, 34] of 2-groups — this is why such 2-groups are called trivial. By definition, the group action \triangleright must be the conjugation action. Hence the second approach to associate a 2-algebra to G seems well suited to this case.

The identity t -map extends $\text{id} : kG_{-1} \rightarrow kG_0$ directly to the group algebras, whence we obtain an associative 2-algebra kG . Following (2.15), the kG_0 -bimodule structure of kG_{-1} is given by group multiplication, and *not* conjugation. Similarly to the 2-group, there is no algebra kernel nor cokernel, whence kG is in the trivial class under elementary equivalence of 2-algebras [40]. As such, we shall also call such 2-algebras $A \xrightarrow{\text{id}} A$ trivial.

This construction defines a functor $G_0 \mapsto G$ assigning the group G_0 (resp. the algebra A_0) to the associated trivial 2-group G (resp. the trivial associative 2-algebra A), which commutes with the group algebra functor [40]. We will show in Section 2.2.2 that this functor, which defines an embedding of the category of associative algebras into that of associative 2-algebras, can be extended to the bialgebra/2-bialgebra context.

c) The skeletal case. The skeletal case is peculiar enough we can check what happens to the different above proposals and even propose yet another way to associate a 2-algebra to a skeletal 2-group.

For *skeletal* 2-group, the t -map $t = 1 : G_{-1} \rightarrow G_0$ is the constant map onto to identity $1 \in G_0$. By the (2-group) Peiffer identity, G_{-1} must be Abelian, and the group kernel $\ker t = G_{-1}$ is the entire group G_{-1} . If we take the group algebras and simply extend t linearly, we obtain the augmentation $t : kG_{-1} \rightarrow k \cdot 1 = k$, whose algebra kernel $\ker t = I$ is the augmentation ideal, which is in general distinct from the group algebra kG_{-1} .

This case is in drastic contrast with the trivial case. The Peiffer identity dictates that kG_{-1} is commutative, but *not* nuclear, and hence kG would be non-skeletal. In other words, merely taking the group algebras of the graded components of a 2-group does not preserve skeletality, and hence does not preserve its elementary equivalence class. Since the linearized t -map is non-trivial $t \neq 0$, there are then two ways to see kG as a 2-algebra:

1. By following the quotient prescription (2.14), we must kill the entire degree-(-1) structure, yielding $\tilde{t} : 0 \rightarrow kG_0$. Indeed, the only totally invariant subgroups of an Abelian group under group multiplication by itself is trivial, and $D \cong k(\ker t) = kG_{-1}$ is the entire group algebra. This is the *only* group 2-algebra one can construct associated to a skeletal 2-group G following the prescription by Wagemann [40].
2. Alternatively, we may seek to solve (2.15) with the given group action \triangleright . However, solutions do not exist unless $x \triangleright \in \text{Aut}(G_{-1})$ is *inner* for all $x \in G_0$, which is certainly not always the case. This gives us a broader class of skeletal 2-groups with which we may form the 2-group algebra by linearly extending t .

Yet another alternative construction (which we emphasize is specialized to the skeletal case) is to linearly extend the group action \triangleright but *not* the t -map. Instead, we take the skeletal 2-algebra $t = 0 : kG_{-1} \rightarrow kG_0$, whose trivial t -map allows us to circumvent the entire issue posed by (2.12). This construction $G \mapsto kG$ not only holds for any skeletal 2-group G , but for certain G_{-1}, G_0 it also preserves the classification! This is due to a deep result in [57], which computes an explicit isomorphism between the Hochschild cohomology and group cohomology in certain cases. We will adopt this approach in an accompanying paper [58], which applies the general framework of 2-Hopf algebras that we shall develop here to construct excitations in the Kitaev model.

With either (2.14), (2.15) understood or the skeletal case proposal, we shall neglect the tilde on the t -map and simply denote $kG = kG_{-1} \xrightarrow{t} kG_0$ in the following. Given a 2-group G , while we can construct in effect different 2-algebras kG with the above two proposals, at the end of the day, we will obtain an associative 2-algebra, which will be the starting point of the 2-bialgebra definition.

2.2 Associative 2-bialgebras

2.2.1 Definition

Coassociative 2-coalgebra. We seek a dual notion of an associative 2-algebra **Definition 2.2.** The idea will be to reverse the arrows in the diagrams and swap the degree. Indeed, our duality structure will typically swap degrees. This is a consequence of how "dualization" is defined in homological algebra [41, 3, 42, 43].

Let us consider a pair of vector spaces, $\mathcal{G}_0, \mathcal{G}_{-1}$ with the map $t : \mathcal{G}_{-1} \rightarrow \mathcal{G}_0$. In direct analogy with the 2-cocycle $\delta = \delta_{-1} + \delta_0$ that were introduced to define a classical Lie 2-bialgebra [41, 3], we introduce the *coproduct* maps

$$\Delta_{-1} : \mathcal{G}_{-1} \rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_{-1}, \quad \Delta_0 : \mathcal{G}_0 \rightarrow (\mathcal{G}_{-1} \otimes \mathcal{G}_0) \oplus (\mathcal{G}_0 \otimes \mathcal{G}_{-1}). \quad (2.16)$$

Note that Δ_0 comes in two components, $\Delta_0 = \Delta_0^l + \Delta_0^r$ (we used the graded sum) with

$$\Delta_0^l : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_0, \quad \Delta_0^r : \mathcal{G}_0 \rightarrow \mathcal{G}_0 \otimes \mathcal{G}_{-1}.$$

In the following, we shall use extensively the conventional Sweedler notation

$$\Delta(y, x) \equiv \Delta_{-1}(y) + \Delta_0(x) = y_{(1)} \otimes y_{(2)} + (x_{(1)}^l \otimes x_{(2)}^l + x_{(1)}^r \otimes x_{(2)}^r) \quad (2.17)$$

where $x_{(1)}^l, x_{(2)}^r \in \mathcal{G}_0$ and $y_{(1)}, y_{(2)}, x_{(2)}^l, x_{(1)}^r \in \mathcal{G}_{-1}$.

Now let

$$\Delta'_0 : \mathcal{G}_0 \rightarrow \mathcal{G}_0 \otimes \mathcal{G}_0$$

denote a coproduct in degree-0, such that Δ_{-1}, Δ'_0 are subject to the following **coassociativity** conditions

$$(\text{id} \otimes \Delta_{-1}) \circ \Delta_{-1} = (\Delta_{-1} \otimes \text{id}) \circ \Delta_{-1}, \quad (\text{id} \otimes \Delta'_0) \circ \Delta'_0 = (\Delta'_0 \otimes \text{id}) \circ \Delta'_0, \quad (2.18)$$

which can be obtained by reversing the arrows in (2.2) for the products μ_{-1} and μ_0 . Hence $(\mathcal{G}_{-1}, \Delta_{-1})$ and $(\mathcal{G}_0, \Delta'_0)$ are *coassociative coalgebras* if (2.18) is satisfied [7]. In the following, we shall use the Sweedler notation

$$\Delta'_0(x) = \bar{x}_{(1)} \otimes \bar{x}_{(2)} \in \mathcal{G}_0 \otimes \mathcal{G}_0. \quad (2.19)$$

Definition 2.4. Let $(\mathcal{G}_{-1}, \Delta_{-1})$ and $(\mathcal{G}_0, \Delta'_0)$ denote a pair of coassociative coalgebras with the coactions Δ_0^l and Δ_0^r . We say that \mathcal{G}_0 forms a \mathcal{G}_{-1} -**cobimodule** if the following **cobimodularity conditions**

$$\begin{aligned} (\Delta_{-1} \otimes \text{id}) \circ \Delta_0^l &= (\text{id} \otimes \Delta_0^l) \circ \Delta'_0, \\ (\text{id} \otimes \Delta_{-1}) \circ \Delta_0^r &= (\Delta_0^r \otimes \text{id}) \circ \Delta'_0, \\ (\text{id} \otimes \Delta_0^r) \circ \Delta_0^l &= (\Delta_0^l \otimes \text{id}) \circ \Delta'_0 \end{aligned} \quad (2.20)$$

are satisfied. In terms of commutative diagrams, we have

$$\begin{array}{ccc} \begin{array}{ccc} & \mathcal{G}_0 & \\ \Delta_0^l \swarrow & & \searrow \Delta'_0 \\ \mathcal{G}_{-1} \otimes \mathcal{G}_0 & & \mathcal{G}_0 \otimes \mathcal{G}_0 \\ \text{id} \otimes \Delta'_0 \searrow & & \swarrow \Delta_0^l \otimes \text{id} \\ & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \otimes \mathcal{G}_0 & \end{array} & , & \begin{array}{ccc} & \mathcal{G}_0 & \\ \Delta_0^r \swarrow & & \searrow \Delta'_0 \\ \mathcal{G}_0 \otimes \mathcal{G}_{-1} & & \mathcal{G}_0 \otimes \mathcal{G}_0 \\ \Delta'_0 \otimes \text{id} \searrow & & \swarrow \text{id} \otimes \Delta_0^r \\ & \mathcal{G}_0 \otimes \mathcal{G}_0 \otimes \mathcal{G}_{-1} & \end{array} \\ \\ \begin{array}{ccc} & \mathcal{G}_0 & \\ \Delta_0^l \swarrow & & \searrow \Delta'_0 \\ \mathcal{G}_0 \otimes \mathcal{G}_0 & & \mathcal{G}_0 \otimes \mathcal{G}_0 \\ \text{id} \otimes \Delta_0^l \searrow & & \swarrow \Delta_0^r \otimes \text{id} \\ & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_0 & \end{array} & & \end{array} \quad (2.21)$$

We emphasize again that, upon dualizing the commutative diagrams (2.2)-(2.10), we must also swap the grading: the action $\cdot_l : \mathcal{G}_0 \otimes \mathcal{G}_{-1} \rightarrow \mathcal{G}_{-1}$ is dualized to the coaction component $\Delta_0^l : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_0$ of the coproduct.

Definition 2.5. A **coassociative 2-coalgebra** (\mathcal{G}, Δ) is a coalgebra homomorphism $t : \mathcal{G}_{-1} \rightarrow \mathcal{G}_0$ such that

1. \mathcal{G}_0 is a \mathcal{G}_{-1} -cobimodule,
2. t is **coequivariant**

$$D_t^+ \circ \Delta_{-1} = \Delta_0 \circ t, \quad (2.22)$$

where we have introduced a convenient tensor notation for the induced t -map

$$D_t^\pm := t \otimes 1 \pm 1 \otimes t$$

in terms of the graded sum. This condition is encoded by the following commutative diagram.

$$\begin{array}{ccc} & \mathcal{G}_{-1} & \\ t \swarrow & & \searrow \Delta_{-1} \\ \mathcal{G}_0 & & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} \\ \Delta_0 \searrow & & \swarrow D_t^+ \\ & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \oplus \mathcal{G}_{-1} \otimes \mathcal{G}_0 & \end{array} \quad (2.23)$$

3. the coPeiffer identity

$$(t \otimes \text{id}) \circ \Delta'_0 = \Delta'_0 = (\text{id} \otimes t) \circ \Delta_0^r, \quad (2.24)$$

which in particular means that we must necessarily have

$$D_t^- \Delta_0 = (t \otimes \text{id}) \circ \Delta_0^l - (\text{id} \otimes t) \circ \Delta_0^r = 0.$$

It is encoded in the following commutative diagram

$$\begin{array}{ccccc} & & \mathcal{G}_0 \otimes \mathcal{G}_0 & & \\ & \text{id} \otimes t \nearrow & \uparrow & \nwarrow t \otimes \text{id} & \\ \mathcal{G}_0 \otimes \mathcal{G}_{-1} & & \mathcal{G}_0 & & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\ & \Delta_0^r \searrow & \uparrow \Delta'_0 & \Delta_0^l \searrow & \\ & & \mathcal{G}_0 & & \end{array} \quad (2.25)$$

We call (\mathcal{G}, Δ) **counital** if there is a counit map $\epsilon = (\epsilon_{-1}, \epsilon_0) : \mathcal{G} \rightarrow k$ such that

$$\begin{aligned} \text{id} &= (\text{id} \otimes \epsilon_{-1}) \circ \Delta_{-1}, & \text{id} &= (\epsilon_{-1} \otimes \text{id}) \circ \Delta_{-1}, \\ \text{id} &= (\epsilon_{-1} \otimes \text{id}) \circ \Delta_0^l, & \text{id} &= (\text{id} \otimes \epsilon_{-1}) \circ \Delta_0^r. \end{aligned} \quad (2.26)$$

Moreover, ϵ should respect the t -map such that $\epsilon_0 = \epsilon_{-1} \circ t$.

The counit conditions can be seen as reversing the arrows of the diagrams (2.10) (and also a swap of the grading since we are dualizing).

$$\begin{array}{ccc} & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} & \\ \epsilon_{-1} \otimes \text{id} \swarrow & \uparrow \Delta_{-1} & \searrow \text{id} \otimes \epsilon_{-1} \\ k \otimes \mathcal{G}_{-1} & \xrightarrow{\cong} \mathcal{G}_{-1} & \xleftarrow{\cong} \mathcal{G}_{-1} \otimes k \\ \\ \text{id} \otimes \epsilon_{-1} \swarrow & \mathcal{G}_0 \otimes \mathcal{G}_{-1} & \uparrow \Delta_0^r \\ k \otimes \mathcal{G}_0 & \xrightarrow{\cong} \mathcal{G}_0 & \\ \\ \epsilon_{-1} \otimes \text{id} \swarrow & \mathcal{G}_{-1} \otimes \mathcal{G}_0 & \uparrow \Delta_0^l \\ \mathcal{G}_0 \otimes k & \xrightarrow{\cong} \mathcal{G}_0 & \\ \\ \epsilon_{-1} \swarrow & \mathcal{G}_{-1} & \downarrow t \\ k & \xleftarrow{\epsilon_0} \mathcal{G}_0 & \end{array} \quad (2.27)$$

Note again that in **Definition 2.5**, the coequivariance and coPeiffer identity are treated as constraints between two coalgebras and the coalgebra homomorphism t between them. With these constraints, we can deduce

$$\text{id} = (\epsilon_0 \otimes \text{id}) \circ \Delta'_0 = (\text{id} \otimes \epsilon_0) \circ \Delta'_0 \quad (2.28)$$

from (2.19) and (2.26). In the skeletal $t = 0$ case, the coproducts $\Delta_{-1}, \Delta_0, \Delta'_0$ and the counits $\epsilon_{-1}, \epsilon_0$ are independent, and this condition is separate from (2.26).

Remark 2.2. Similar to the 2-algebra case, if $t \neq 0$ were not trivial, then we could have the following conditions

$$\begin{aligned} (\text{id} \otimes \Delta'_0) \circ \Delta_0^l &= (\Delta_0^l \otimes \text{id}) \circ \Delta'_0, \\ (\Delta'_0 \otimes \text{id}) \circ \Delta_0^r &= (\text{id} \otimes \Delta_0^r) \circ \Delta'_0, \\ (\text{id} \otimes \Delta_0^l) \circ \Delta'_0 &= (\Delta_0^r \otimes \text{id}) \circ \Delta'_0 \end{aligned} \quad (2.29)$$

between the coproducts Δ_0 and Δ'_0 . By making use of the Sweedler notation (2.17), (2.19), these conditions translate to

$$\begin{cases} ty_{(1)} = (ty)_{(1)}^r, & \bar{x}_{(1)} = tx_{(1)}^l = x_{(1)}^r \\ ty_{(2)} = (ty)_{(2)}^l, & \bar{x}_{(2)} = x_{(2)}^l = tx_{(2)}^r \end{cases} \quad (2.30)$$

When combined, they give $\bar{t}y_{(1)} = ty_{(1)}, \bar{t}y_{(2)} = ty_{(2)}$ which will become important later. In the skeletal case, the constraints involving t drop and we would only have $\bar{x}_{(1)} = x_{(1)}^r, \bar{x}_{(2)} = x_{(2)}^l$.

2-bialgebra. Using the Sweedler notations (2.17), (2.19), we state the condition that the coproduct map Δ given in (2.16) preserves the algebra/bimodule structure:

$$\begin{aligned} \Delta_{-1}(x \cdot y) &= \bar{x}_{(1)} \cdot y_{(1)} \otimes \bar{x}_{(2)} \cdot y_{(2)}, & \Delta_{-1}(y \cdot x) &= y_{(1)} \cdot \bar{x}_{(1)} \otimes y_{(2)} \cdot \bar{x}_{(2)}, \\ \Delta_0^l(xx') &= x_{(1)}^l x'_{(1)} \otimes x_{(2)}^l x'_{(2)}, & \Delta_0^r(xx') &= x_{(1)}^r x'_{(1)} \otimes x_{(2)}^r x'_{(2)}. \end{aligned} \quad (2.31)$$

We call these conditions the **2-bialgebra axioms**.

The bialgebra axioms in each degree,

$$\Delta_{-1}(yy') = y_{(1)}y'_{(1)} \otimes y_{(2)}y'_{(2)}, \quad \Delta'_0(xx') = \bar{x}_{(1)}\bar{x}'_{(1)} \otimes \bar{x}_{(2)}\bar{x}'_{(2)},$$

follow directly from (2.31) and the coequivariance and coPeiffer identities (2.22), (2.24); see *Remark 2.2*.

Definition 2.6. The tuple $(\mathcal{G}, \cdot, \Delta)$ is an **associative 2-bialgebra** iff (\mathcal{G}, \cdot) is an associative 2-algebra and (\mathcal{G}, Δ) is a coassociative 2-coalgebra that are mutually compatible, in the sense that the coproduct map Δ satisfies (2.18)-(2.24) and (2.31).

We call $(\mathcal{G}, \cdot, \eta, \Delta, \epsilon)$ **unital** if $(\mathcal{G}, \cdot, \eta)$ and $(\mathcal{G}, \Delta, \epsilon)$ are respectively unital and counital.

The 2-bialgebra axioms are equivalently described in terms of the following commutative diagrams, where we use the swap $\sigma : \mathcal{G}_0 \otimes \mathcal{G}_{-1} \rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_0$ and $\sigma' : \mathcal{G}_{-1} \otimes \mathcal{G}_0 \rightarrow \mathcal{G}_0 \otimes \mathcal{G}_{-1}$,

$$\begin{array}{ccc}
\begin{array}{ccc}
\mathcal{G}_0 \otimes \mathcal{G}_{-1} & \xrightarrow{\Delta'_0 \otimes \Delta_{-1}} & \mathcal{G}_0 \otimes \mathcal{G}_0 \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} \\
\downarrow \cdot_l & & \downarrow \text{id} \otimes \sigma \otimes \text{id} \\
\mathcal{G}_{-1} & \xrightarrow{\Delta_{-1}} & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} \\
\downarrow \Delta_{-1} & & \downarrow \cdot_l \otimes \cdot_l \\
\mathcal{G}_{-1} \otimes \mathcal{G}_{-1} & & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1}
\end{array} & &
\begin{array}{ccc}
\mathcal{G}_{-1} \otimes \mathcal{G}_0 & \xrightarrow{\Delta_{-1} \otimes \Delta'_0} & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_0 \otimes \mathcal{G}_0 \\
\downarrow \cdot_r & & \downarrow \text{id} \otimes \sigma' \otimes \text{id} \\
\mathcal{G}_{-1} & \xrightarrow{\Delta_{-1}} & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\
\downarrow \Delta_{-1} & & \downarrow \cdot_r \otimes \cdot_r \\
\mathcal{G}_{-1} \otimes \mathcal{G}_0 & & \mathcal{G}_{-1} \otimes \mathcal{G}_0
\end{array} \\
\begin{array}{ccc}
\mathcal{G}_0 \otimes \mathcal{G}_0 & \xrightarrow{\Delta'_0 \otimes \Delta'_0} & \mathcal{G}_{-1} \otimes \mathcal{G}_0 \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_0 \\
\downarrow \mu_0 & & \downarrow \text{id} \otimes \sigma \otimes \text{id} \\
\mathcal{G}_0 & \xrightarrow{\Delta'_0} & \mathcal{G}_{-1} \otimes \mathcal{G}_{-1} \\
\downarrow \Delta'_0 & & \downarrow \mu_{-1} \otimes \mu_0 \\
\mathcal{G}_{-1} \otimes \mathcal{G}_0 & & \mathcal{G}_{-1} \otimes \mathcal{G}_0
\end{array} & &
\begin{array}{ccc}
\mathcal{G}_0 \otimes \mathcal{G}_0 & \xrightarrow{\Delta'_0 \otimes \Delta'_0} & \mathcal{G}_0 \otimes \mathcal{G}_{-1} \otimes \mathcal{G}_0 \otimes \mathcal{G}_{-1} \\
\downarrow \mu_0 & & \downarrow \text{id} \otimes \sigma' \otimes \text{id} \\
\mathcal{G}_0 & \xrightarrow{\Delta'_0} & \mathcal{G}_0 \otimes \mathcal{G}_0 \\
\downarrow \Delta'_0 & & \downarrow \mu_0 \otimes \mu_{-1} \\
\mathcal{G}_0 \otimes \mathcal{G}_{-1} & & \mathcal{G}_0 \otimes \mathcal{G}_{-1}
\end{array}
\end{array} \quad (2.32)$$

2.2.2 Example: function 2-bialgebras on 2-groups

Let k denote a field of characteristic zero, and let G denote a (finite) 2-group.

Function 2-coalgebra on a (finite) 2-group. Consider the 2-group 2-algebra kG constructed in section 2.1.2. We denote its linear dual by $k[G] = \text{Hom}_k(kG, k)$, which consist of k -linear functions on kG . This space inherits the *dual* grading of the 2-group 2-algebra kG , in the sense that $k[G_{-1}]$ is in degree-0 while $k[G_0]$ is in degree-(-1). Each element $F \in k[G]$, then admits a decomposition in terms of this grading, $F = \xi \oplus \zeta$, with $\xi \in k[G_0]$, $\zeta \in k[G_{-1}]$, for which the t -map is given by the pullback,

$$k[G] = k[G_0] \xrightarrow{t^*} k[G_{-1}], \quad (t^*\xi)(y) = \xi(ty),$$

where $x \in kG_0, y \in kG_{-1}$. Note this t -map is the one on kG defined in (2.14), which may differ from that of the underlying 2-group G .

The reversal of degree for the function 2-coalgebra allows us to define a grading-odd duality pairing given by the function evaluation

$$\langle (y, x), (\xi, \zeta) \rangle = \text{ev}((y, x) \otimes (\xi, \zeta)) = \xi(x) + \zeta(y),$$

with respect to which the coproduct Δ on $k[G]$ defined in (2.33) is dual to the 2-algebra structure on kG . Conversely, if there is a well-defined coassociative coproduct Δ on kG (ie. satisfying (2.22)-(2.24)), then the evaluation pairing dualizes it to a well-defined associative 2-algebra structure on $k[G]$. This is a guiding principle with which we shall construct the 2-quantum double in the following sections.

The natural coproduct $\Delta^* = \Delta_{-1}^* + \Delta_0^*$ on $k[G]$, is induced by the 2-algebra structure on kG ,

$$\begin{aligned}
\Delta_{-1}^*(\xi) &= \xi_{(1)} \otimes \xi_{(2)} \iff \xi(xx') = \xi_{(1)}(x)\xi_{(2)}(x'), \\
(\Delta_0^*)^l(\zeta) &= \zeta_{(1)}^l \otimes \zeta_{(2)}^l \iff \zeta(x \cdot y) = \zeta_{(1)}^l(x)\zeta_{(2)}^l(y),
\end{aligned}$$

$$(\Delta_0^*)^r \zeta = \zeta_{(1)}^r \otimes \zeta_{(2)}^r \iff \zeta(y \cdot x) = \zeta_{(1)}^r(y) \zeta_{(2)}^r(x), \quad (2.33)$$

where $x, x' \in kG_0$ and $y \in kG_{-1}$. The superscript $*$ encodes the fact we are considering the dual of kG as we will emphasize soon. The induced coproduct $\Delta_0^{*'}$ in degree-0 is induced from the product μ_{-1} ,

$$\Delta_0^{*'}(\zeta) = \bar{\zeta}_{(1)} \otimes \bar{\zeta}_{(2)} \iff \zeta(yy') = \bar{\zeta}_{(1)}(y) \bar{\zeta}_{(2)}(y'),$$

and is also related to the actions, thanks to the Peiffer identity, so that $\Delta_0^{*'} = \frac{1}{2} D_{t^*}^+ \Delta_0$. In a similar way, all the 2-co-algebra axioms are satisfied as we are dualizing all the properties of the 2-algebra kG , hence we are reversing the arrows.

For example, the co-bimodularity is obtained from the bimodularity in kG .

$$\begin{aligned} (\Delta_0^*)^l(t^*\xi)(x, y) &= \xi(t(x \cdot y)) = \xi(xt(y)) = \xi_{(1)}(x) \xi_{(2)}(t(y)) = ((1 \otimes t^*) \Delta_{-1}^*(\xi))(x, y), \\ (\Delta_0^*)^r(t^*\xi)(y, x) &= \xi(t(y \cdot x)) = \xi(t(y)x) = \xi_{(1)}(ty) \xi_{(2)}(x) = ((t^* \otimes 1) \Delta_{-1}^*(\xi))(y, x), \end{aligned}$$

The co-Peiffer identity is obtained from the Peiffer identity

$$\zeta(yy') = \zeta(ty \cdot y') = \zeta(y \cdot ty') \implies (t^* \otimes 1) (\Delta_0^*)^l(\zeta) = (1 \otimes t^*) (\Delta_0^*)^r(\zeta),$$

and so on and so forth.

Moreover, there is also a counit $\epsilon : k[G] \rightarrow k$ given by evaluation at the identity, $\epsilon_{-1}(\xi) = \xi(1)$, $\epsilon_0(\zeta) = \zeta(1)$. Since the t -map respects the identity, we have $\epsilon_{-1}(t^*\xi) = t^*\xi(1) = \xi(1) = \epsilon_0(\xi)$, as desired for a counit.

In order to determine the 2-algebra structure of $k[G]$, we can use the coalgebra structure of kG . For this we are going to consider the specific case $t = \text{id}$ as otherwise we cannot make the example as explicit as one would wish, due to the constraints (2.30).

The trivial 2-algebra. Recall a *trivial* 2-group G has $G_{-1} = G_0$ and the t -map is the identity $t = \text{id}$. This the simplest case one can consider, as its corresponding 2-group 2-algebra kG is obtained by just extending every structure k -linearly. We shall now consider endowing a coproduct on kG from that of kG_0 .

Let Δ denote a coproduct on kG . (2.22), (2.24) dictates that all components $\Delta_{-1}, \Delta_0, \Delta'_0$ of the coproduct are identical, and hence (kG_0, Δ'_0) is a coalgebra iff (kG, Δ) is also one. Indeed, if $\Delta'_0(x) = x_{(1)} \otimes x_{(2)}$ then we must have

$$\Delta_{-1}(y) = y_{(1)} \otimes y_{(2)}, \quad \Delta_0(x) = y_{(1)} \otimes x_{(2)} + x_{(1)} \otimes y_{(2)},$$

where $y_{(1)}, y_{(2)}$ are identical to $x_{(1)}, x_{(2)} \in kG_0$ as elements of the group algebra kG_0 , but with degree-(-1); in other words, we have $t(y_{(1)}) = x_{(1)}$ under $t = \text{id}$. Explicitly, we can choose the group like coproduct

$$\Delta'_0(x) = x \otimes x \quad \Delta_{-1}(y) = y \otimes y, \quad \Delta_0(x) = y \otimes x + x \otimes y, \quad \text{with } ty = y = x$$

We recover then a 2-algebra structure on the graded function space $k[G]$, for which

$$\begin{aligned} \zeta \eta'(y) &= \zeta(y_{(1)}) \zeta'(y_{(1)}) = \zeta(y) \zeta'(y), & \xi \xi'(x) &= \xi(x_{(1)}) \xi'(x_{(2)}) = \xi(x) \xi'(x), \\ (\zeta \cdot \xi)(x) &= \zeta(y_{(1)}) \xi(x_{(2)}) = \zeta(y) \xi(x), & (\xi \cdot \zeta)(x) &= \xi(x_{(1)}) \zeta(y_{(2)}) = \xi(x) \zeta(y). \end{aligned}$$

We have recovered the pointwise products on $k[G_0]$ and $k[G_{-1}]$, the bimodule structure in terms of these pointwise product (since $ty = y = x$). Moreover, one can check that we have the 2-bialgebra axioms (2.31),

$$\begin{aligned} (\zeta \zeta')(x \cdot y) &= (\zeta \zeta')_{(1)}^l(x) (\zeta \zeta')_{(2)}^l(y) = \zeta_{(1)}^l(x_{(1)}) \zeta_{(1)}^l(x_{(2)}) \zeta_{(2)}^l(y_{(1)}) \zeta_{(2)}^l(y_{(2)}) \\ &= (\zeta_{(1)}^l \zeta_{(1)}^l)(x) (\zeta_{(2)}^l \zeta_{(2)}^l)(y), \\ (\zeta \zeta')(y \cdot x) &= (\zeta \zeta')_{(1)}^r(y) (\zeta \zeta')_{(2)}^r(x) = \zeta_{(1)}^r(y_{(1)}) \zeta_{(1)}^r(y_{(2)}) \zeta_{(2)}^r(x_{(1)}) \zeta_{(2)}^r(x_{(2)}) \\ &= (\zeta_{(1)}^r \zeta_{(1)}^r)(x) (\zeta_{(2)}^r \zeta_{(2)}^r)(y) \\ (\zeta \cdot \xi)(xx') &= (\zeta \cdot \xi)_{(1)}(x) (\zeta \cdot \xi)_{(2)}(x') = \zeta_{(1)}(y_{(1)}) \xi_{(1)}(x_{(2)}) \zeta_{(2)}(x'_{(1)}) \xi_{(2)}(x'_{(2)}) \\ &= (\zeta_{(1)} \xi_{(1)})(x) (\zeta_{(2)} \xi_{(2)})(x'); \end{aligned}$$

note $\bar{\zeta} = \zeta$ as the t -map is the identity. Therefore $(k[G], \Delta^*)$ and (kG, Δ) are (unital) associative 2-bialgebras that are mutually dual in a certain sense, with the former being commutative and the latter being cocommutative, when using the group-like coproducts, just as in the 1-group case [44]. We shall rigorously establish this general 2-bialgebra duality in **Proposition 3.1**.

Remark 2.3. In general, given an ordinary 1-bialgebra G , we can form its associated trivial 2-bialgebra $\mathcal{G} = G \xrightarrow{\text{id}} G$ in analogy with the above. Moreover, it is clear that any bialgebra homomorphism $f : G \rightarrow G'$ extends to a 2-bialgebra homomorphism $f \oplus f : \mathcal{G} \rightarrow \mathcal{G}'$, whose graded components are just copies of f . We therefore have an embedding

$$\text{Bialg}_{\text{ass}} \hookrightarrow 2\text{Bialg}_{\text{ass}} \quad (2.34)$$

of the category of associative bialgebras into the category of associative 2-bialgebras. This is a "quantum" version of an analogous result for Lie 2-bialgebras stated in *Remark 3.11* of [59].

3 Strict 2-quantum doubles and the universal 2-R-matrix

In this section, we construct our main example of a strict 2-bialgebra given by the strict 2-quantum doubles which can be seen a categorification of the standard quantum double [7], and the quantization of a classical 2-double [42, 3] of Lie 2-algebras.

The goal for studying (2-)quantum doubles is that, for the ordinary 1-bialgebra H , the *skew-pairing* involved in the construction of the quantum double $D(H, H)$ of Majid [44] provides a characterization of R-matrices on H . Moreover, this construction is *universal* in the sense that any R-matrix on H can be derived this way from $D(H, H)$. We wish to directly categorify Majid's construction, and derive a universal characterization of 2-R-matrices from our construction of a 2-quantum double.

Our strategy will be as follows. Firstly, we consider a pair of dual associative 2-bialgebras. They are dual in the sense that the coalgebra sector is given by the algebra sector of its dual counterpart. We then define a notion of a canonical *coadjoint action* of a 2-bialgebra on its dual. By requesting that the mutually-dual 2-bialgebras act on each other by such coadjoint actions, we are then able to form the 2-quantum double as a 2-bialgebra. We will then also prove a key factorization theorem for 2-quantum doubles.

3.1 Matched pair of 2-(bi)algebras

Dually paired 2-bialgebras Let $(\mathcal{G}, \cdot, \Delta)$ denote a (finite dimensional) 2-bialgebra, and let \mathcal{G}^* denote its linear dual, defined with respect to the following duality evaluation/pairing map²

$$\langle (g, f), (y, x) \rangle = \langle f, y \rangle_{-1} + \langle g, x \rangle_0 \quad (3.1)$$

for each $x \in \mathcal{G}_0, y \in \mathcal{G}_{-1}, f \in \mathcal{G}_{-1}^*, g \in \mathcal{G}_0^*$. Note that the grading is flipped by dualizing the t -map: $\langle t^* \cdot, \cdot \rangle = \langle \cdot, t \cdot \rangle$, whence $t^* : \mathcal{G}_0^* \rightarrow \mathcal{G}_{-1}^*$ and \mathcal{G}^* is skeletal whenever \mathcal{G} is. In the following, we shall denote this pairing also by an evaluation ev.

So far, \mathcal{G}^* merely forms a 2-vector space. By leveraging the duality (3.1), we can induce algebraic structures on \mathcal{G}^* according to the coalgebraic structures (2.16), (2.19) on \mathcal{G} as follows:

$$\begin{aligned} \langle f \otimes f', \Delta_{-1}(y) \rangle &= \langle f f', y \rangle, & \langle g \otimes g', \Delta'_0(x) \rangle &= \langle g g', x \rangle, \\ \langle f \otimes g, \Delta_0^l(x) \rangle &= \langle f \cdot^* g, x \rangle, & \langle g \otimes f, \Delta_0^r(x) \rangle &= \langle g \cdot^* f, x \rangle, \\ \langle \Delta_0^{*l} f, y \otimes y' \rangle &= \langle f, y y' \rangle, & \langle \Delta_{-1}^* g, x \otimes x' \rangle &= \langle g, x x' \rangle, \\ \langle \Delta_0^{*r} f, x \otimes y \rangle &= \langle f, x \cdot_l y \rangle, & \langle \Delta_0^{*l} f, y \otimes x \rangle &= \langle f, y \cdot_r x \rangle. \end{aligned}$$

The conditions (2.22), (2.24), (2.18), (2.20), then ensure that (\mathcal{G}^*, \cdot^*) forms an associative 2-algebra. More is true, in fact, which we now prove in the following.

Proposition 3.1. *Let $\mathcal{G}, \mathcal{G}^*$ be dually paired as in (3.1), then $(\mathcal{G}, \cdot, \Delta)$ is an (unital) associative 2-bialgebra iff $(\mathcal{G}^*, \cdot^*, \Delta^*)$ is an (unital) associative 2-bialgebra.*

Proof. This is a straightforward computation using the pairing (3.1). In particular, the equivariance and Peiffer identity of t^* , as well as the fact that \mathcal{G}_{-1}^* forms a \mathcal{G}_0^* -bimodule, follow directly from dualizing (2.22), (2.24), (2.18), (2.20).

What is non-trivial is (2.31). Define Δ_0^* by dualizing the bimodule structure \cdot of \mathcal{G} , then we have

$$\begin{aligned} \langle (\Delta_0^*)^l(f f'), x \otimes y \rangle &= \langle f \otimes f', \Delta_{-1}(x \cdot y) \rangle, & \langle (\Delta_0^*)^r(f f'), y \otimes x \rangle &= \langle f \otimes f', \Delta_{-1}(y \cdot x) \rangle, \\ \langle (\Delta_{-1}^*)(f \cdot^* g), x \otimes x' \rangle &= \langle f \otimes g, \Delta_0^l(x x') \rangle, & \langle (\Delta_{-1}^*)(g \cdot^* f), x \otimes x' \rangle &= \langle f \otimes g, \Delta_0^r(x x') \rangle. \end{aligned}$$

We now compute using analogues of (2.31) for Δ^* , that

$$\begin{aligned} \langle f_{(1)}^l f_{(1)}'^l \otimes f_{(2)}^l f_{(2)}'^l, x \otimes y \rangle &= \langle (f_{(1)}^l \otimes f_{(1)}'^l) \otimes (f_{(2)}^l \otimes f_{(2)}'^l), (\bar{x}_{(1)} \otimes \bar{x}_{(2)}) \otimes (y_{(1)} \otimes y_{(2)}) \rangle \\ &= \langle (\Delta_0^*)^l(f) \otimes (\Delta_0^*)^l(f'), (\bar{x}_{(1)} \otimes y_{(1)}) \otimes (\bar{x}_{(2)} \otimes y_{(2)}) \rangle \\ &= \langle f \otimes f', (\bar{x}_{(1)} \cdot y_{(1)}) \otimes (\bar{x}_{(2)} \cdot y_{(2)}) \rangle, \\ \langle f_{(1)}^r f_{(1)}'^r \otimes f_{(2)}^r f_{(2)}'^r, y \otimes x \rangle &= \langle f \otimes f', (y_{(1)} \cdot \bar{x}_{(1)}) \otimes (y_{(2)} \cdot \bar{x}_{(2)}) \rangle, \end{aligned}$$

and similarly

$$\begin{aligned} \langle \bar{f}_{(1)} \cdot^* g_{(1)} \otimes \bar{f}_{(2)} \cdot^* g_{(2)}, x \otimes x' \rangle &= \langle (\bar{f}_{(1)} \otimes g_{(1)}) \otimes (\bar{f}_{(2)} \otimes g_{(2)}), (x_{(1)}^l \otimes x_{(2)}^l) \otimes (x_{(1)}'^l \otimes x_{(2)}'^l) \rangle \\ &= \langle \Delta_0^*(f) \otimes \Delta_{-1}^*(g), (x_{(1)}^l \otimes x_{(1)}'^l) \otimes (x_{(2)}^l \otimes x_{(2)}'^l) \rangle, \end{aligned}$$

²We shall drop the subscripts on the pairing forms $\langle \cdot, \cdot \rangle$ when no confusion arises.

$$\begin{aligned}
&= \langle f \otimes g, x_{(1)}^l x_{(1)}'^l \otimes x_{(2)}^l x_{(2)}'^l \rangle, \\
\langle g_{(1)} \cdot^* \bar{f}_{(1)} \otimes g_{(2)} \cdot^* \bar{f}_{(2)}, x \otimes x' \rangle &= \langle g \otimes f, x_{(1)}^r x_{(1)}'^r \otimes x_{(2)}^r x_{(2)}'^r \rangle,
\end{aligned}$$

hence Δ also satisfies (2.31). This proves that $(\mathcal{G}^*, \cdot^*, \Delta^*)$ is an associative 2-bialgebra iff $(\mathcal{G}, \cdot, \Delta)$ also is.

Now consider the units and counits. Given

$$\begin{aligned}
\langle g, \eta x \rangle &= \langle (\eta^* \otimes \text{id}) \circ \Delta_{-1}^*(g), x \rangle, & \langle g, x \eta \rangle &= \langle (\text{id} \otimes \eta^*) \circ \Delta_{-1}^*(g), x \rangle, \\
\langle f, \eta \cdot y \rangle &= \langle (\eta^* \otimes \text{id}) \circ (\Delta_0^*)^l(f), y \rangle, & \langle f, y \cdot \eta \rangle &= \langle (\text{id} \otimes \eta^*) \circ (\Delta_0^*)^r(f), y \rangle,
\end{aligned}$$

we see that η is a unit for (\mathcal{G}, \cdot) (ie. these quantities all vanish) iff η^* is a counit for $(\mathcal{G}^*, \Delta^*)$. Similarly, ϵ is a counit for (\mathcal{G}, Δ) iff ϵ^* is a unit for (\mathcal{G}^*, \cdot^*) . The converse direction is identical. \square

Coadjoint action.

Definition 3.1. The canonical **coadjoint action** of \mathcal{G} on \mathcal{G}^* is specified in terms of three components, $\bar{\triangleright} = ((\triangleright_0, \triangleright_{-1}), \Upsilon)$ given by

$$\begin{aligned}
\triangleright_0 : \mathcal{G}_0 &\rightarrow \text{End } \mathcal{G}_0^*, & \langle g, x x' \rangle &= -\langle x \triangleright_0 g, x' \rangle, \\
\triangleright_{-1} : \mathcal{G}_0 &\rightarrow \text{End } \mathcal{G}_{-1}^*, & \langle f, x \cdot y \rangle &= -\langle x \triangleright_{-1} f, y \rangle, \\
\Upsilon : \mathcal{G}_{-1} &\rightarrow \text{Hom}(\mathcal{G}_{-1}^*, \mathcal{G}_0^*), & \langle f, y \cdot x \rangle &= -\langle \Upsilon_y f, x \rangle.
\end{aligned} \tag{3.2}$$

As we will see when discussing 2-representations in Section 6, the coadjoint action can also be interpreted as a 2-representation.

Analogously, we have the coadjoint back-action $\bar{\triangleleft} = ((\triangleleft_0, \triangleleft_{-1}), \tilde{\Upsilon})$ of \mathcal{G}^* on \mathcal{G} , which we write from the right³. The "bar" notation is used to distinguish $\bar{\triangleright}$ from the group action \triangleright in the case where $\mathcal{G} = kG$ is defined through a 2-group G .

Matched pair. Given the pair of strict 2-bialgebras $(\mathcal{G}, \mathcal{G}^*)$, we allow them to act upon each other by coadjoint actions $\bar{\triangleright}$ and $\bar{\triangleleft}$. In analogy with [7], we impose the following monstrous set of twelve compatibility conditions

$$\begin{aligned}
x \triangleright_{-1} (f f') &= (t x_{(1)}^l \triangleright_0 f_{(1)}^l) \cdot^* ((x_{(2)}^l \triangleleft_{-1} f_{(2)}^l) \triangleright_{-1} f') + (x_{(1)}^r \triangleright_0 f_{(1)}^l) \cdot^* (\Upsilon_{x_{(2)}^l \triangleleft_0 f_{(2)}^l} f'), \\
&\quad + (\Upsilon_{x_{(1)}^l f_{(1)}^r} f') \cdot^* (\Upsilon_{x_{(2)}^l \tilde{\Upsilon}_{f_{(2)}^r}} f') + (x_{(1)}^r \triangleright_{-1} f_{(1)}^r) \cdot^* (\Upsilon_{x_{(2)}^l \triangleleft_0 (t^* f_{(2)}^r)} f'), \\
\Upsilon_y (f f') &= ((t y_{(1)} \triangleright_0 f_{(1)}^l) \cdot^* (\Upsilon_{y_{(2)} \triangleleft_0 f_{(2)}^l} f') + (\Upsilon_{y_{(1)} f_{(1)}^r} f') \cdot^* (\Upsilon_{y_{(2)} \triangleleft_0 (t^* f_{(2)}^r)} f')), \\
x \triangleright_0 (f \cdot^* g) &= (t x_{(1)}^l \triangleright_0 f_{(1)}^l) \cdot^* ((x_{(2)}^l \triangleleft_{-1} f_{(2)}^l) \triangleright_0 g) + (x_{(1)}^r \triangleright_0 f_{(1)}^l) \cdot^* (t x_{(2)}^r \triangleleft_0 f_{(2)}^l \triangleright_0 g) \\
&\quad + (\Upsilon_{x_{(1)}^l f_{(1)}^r} f') \cdot^* (t x_{(2)}^l \tilde{\Upsilon}_{f_{(2)}^r} \triangleright_0 g) + (x_{(1)}^r \triangleright_{-1} f_{(1)}^r) \cdot^* (t x_{(2)}^r \triangleleft_0 (t^* f_{(2)}^r) \triangleright_0 g), \\
t y \triangleright_0 (f \cdot^* g) &= (t y_{(1)} \triangleright_0 f_{(1)}^l) \cdot^* (t y_{(2)} \triangleleft_0 f_{(2)}^l \triangleright_0 g) + (\Upsilon_{y_{(1)} f_{(1)}^r} f') \cdot^* (t y_{(2)} \triangleleft_0 (t^* f_{(2)}^r) \triangleright_0 g), \\
x \triangleright_0 (g \cdot^* f) &= (t x_{(1)}^l \triangleright_0 g_{(1)}) \cdot^* (\Upsilon_{x_{(2)}^l \tilde{\Upsilon}_{g_{(2)}}} f) + (x_{(1)}^r \triangleright_0 g_{(1)}) \cdot^* (\Upsilon_{x_{(2)}^l \triangleleft_0 (t^* g_{(2)})} f) \\
t y \triangleright_0 (g \cdot^* f) &= (t y_{(1)} \triangleright_0 g_{(1)}) \cdot^* (\Upsilon_{y_{(2)} \triangleleft_0 (t^* g_{(2)})} f), \\
(x x') \triangleleft_{-1} f &= (x \tilde{\Upsilon}_{t x_{(1)}^l \triangleright_0 f_{(1)}^l}) \cdot (x_{(2)}^l \triangleleft_{-1} f_{(2)}^l) + (x \tilde{\Upsilon}_{x_{(1)}^r \triangleright_0 f_{(1)}^l}) \cdot (x_{(2)}^r \triangleleft_0 f_{(2)}^l) \\
&\quad + (x \tilde{\Upsilon}_{\Upsilon_{x_{(1)}^l} f_{(1)}^r}) \cdot (x_{(2)}^l \tilde{\Upsilon}_{f_{(2)}^r}) + (x \triangleleft_{-1} (x_{(1)}^r \triangleright_{-1} f_{(1)}^r)) \cdot (x_{(2)}^r \triangleleft_0 (t^* f_{(2)}^r)), \\
(x x') \tilde{\Upsilon}_g &= (x \tilde{\Upsilon}_{t x_{(1)}^l \triangleright_0 g_{(1)}}) \cdot (x_{(2)}^l \tilde{\Upsilon}_{g_{(2)}}) + (x \tilde{\Upsilon}_{x_{(1)}^r \triangleright_0 g_{(1)}}) \cdot (x_{(2)}^r \triangleleft_0 (t^* g_{(2)})), \\
(y \cdot x) \triangleleft_0 f &= (y \triangleleft_0 t^* (t x_{(1)}^l \triangleright_0 f_{(1)}^l)) \cdot (x_{(2)}^l \triangleleft_{-1} f_{(2)}^l) + (y \triangleleft_0 t^* (x_{(1)}^r \triangleright_0 f_{(1)}^r)) \cdot (x_{(2)}^r \triangleleft_0 f_{(2)}^l) \\
&\quad + (y \triangleleft_0 t^* (\Upsilon_{x_{(1)}^l} f_{(1)}^r)) \cdot (x_{(2)}^l \tilde{\Upsilon}_{f_{(2)}^r}) + (y \triangleleft_0 (x_{(1)}^r \triangleright_{-1} f_{(1)}^r)) \cdot (x_{(2)}^r \triangleleft_0 (t^* f_{(2)}^r)), \\
(y \cdot x) \triangleleft_0 t^* g &= (y \triangleleft_0 t^* (t x_{(1)}^l \triangleright_0 g_{(1)})) \cdot (x_{(2)}^l \tilde{\Upsilon}_{g_{(2)}}) + (y \triangleleft_0 t^* (x_{(1)}^r \triangleright_0 g_{(1)})) \cdot (x_{(2)}^r \triangleleft_0 t^* g_{(2)}), \\
(x \cdot y) \triangleleft_0 f &= (x \tilde{\Upsilon}_{t y_{(1)} \triangleright_0 f_{(1)}^l}) \cdot (y_{(2)} \triangleleft_0 f_{(2)}^l) + (x \tilde{\Upsilon}_{\Upsilon_{y_{(1)}} f_{(1)}^r}) \cdot (y_{(2)} \triangleleft_0 (t^* f_{(2)}^r)), \\
(x \cdot y) \triangleleft_0 t^* g &= (x \tilde{\Upsilon}_{t y_{(1)} \triangleright_0 g_{(1)}}) \cdot (y_{(2)} \triangleleft_0 t^* g_{(2)}),
\end{aligned}$$

where we have made use of the Sweedler notation (2.17).

We define a shorthand notation where $z = (y, x) \in \mathcal{G}$, $h = (g, f) \in \mathcal{G}^*$, such that the following

$$z \bar{\triangleright} (h \cdot^* h') = (z_{(1)} \bar{\triangleright} h_{(1)}) \cdot^* ((z_{(2)} \bar{\triangleleft} h_{(2)}) \bar{\triangleright} h'), \tag{3.3}$$

³This means that we have, for instance, $\langle g \cdot^* f, x \rangle = -\langle g, x \triangleleft_{-1} f \rangle$ and $\langle f \cdot^* g, x \rangle = -\langle f, x \tilde{\Upsilon}_g \rangle$.

$$(z \cdot z') \bar{\triangleleft} h = (z \bar{\triangleleft} (z'_{(1)} \bar{\triangleright} h_{(1)})) \cdot (z'_{(2)} \bar{\triangleleft} h_{(2)}) \quad (3.4)$$

encode respectively the first six and last six of the above conditions. We also have the cross relations

$$z_{(1)} \bar{\triangleleft} h_{(1)} \otimes z_{(2)} \bar{\triangleright} h_{(2)} = z_{(2)} \bar{\triangleleft} h_{(2)} \otimes z_{(1)} \bar{\triangleright} h_{(1)}, \quad (3.5)$$

as well as the unity axioms against the unit η and counit ϵ ,

$$z \bar{\triangleright} \eta = \epsilon(z), \quad \eta \bar{\triangleleft} h = \epsilon(h). \quad (3.6)$$

Definition 3.2. We call a tuple $(\mathcal{G}, \mathcal{G}^*)$ of (finite dimensional) 2-bialgebras satisfying (3.3)-(3.6) a **matched pair**.

Remark 3.1. Note that in the skeletal case $t, t^* = 0$, the crossed relations (3.3), (3.4) reduce to just two non-trivial equations. These are given by

$$\begin{aligned} x \triangleright_{-1} (ff') &= (x_{(1)}^r \triangleright_0 f_{(1)}^l) \cdot^* (\Upsilon_{x_{(2)}^r \triangleleft_0 f_{(2)}^l} f') + (\Upsilon_{x_{(1)}^l} f_{(1)}^r) \cdot^* (\Upsilon_{x_{(2)}^l} \tilde{\Upsilon}_{f_{(2)}^r} f') \\ &\equiv (x_{(1)} \bar{\triangleright} f_{(1)}) \cdot^* ((x_{(2)} \bar{\triangleleft} f_{(2)}) \bar{\triangleright} f'), \\ (xx') \triangleleft_{-1} f &= (x \tilde{\Upsilon}_{x_{(1)}^r \triangleright_0 f_{(1)}^l}) \cdot (x_{(2)}^r \triangleleft_0 f_{(2)}^l) + (x \tilde{\Upsilon}_{x_{(1)}^l} f_{(1)}^r) \cdot (x_{(2)}^l \tilde{\Upsilon}_{f_{(2)}^r}) \\ &\equiv (x \bar{\triangleleft} (x'_{(1)} \bar{\triangleright} f_{(1)})) \cdot (x_{(2)} \bar{\triangleleft} f_{(2)}), \end{aligned} \quad (3.7)$$

where we have used a convenient notation for brevity. One may notice that these are precisely the usual crossed relations for a quantum double group (cf. [7]) of a semidirect product 2-bialgebra $\mathcal{G}_{-1} \rtimes \mathcal{G}_0$, where \mathcal{G}_{-1} is nuclear.

3.2 Construction of the strict 2-quantum double

We now begin our construction of the general 2-quantum double given a matched pair $(\mathcal{G}, \mathcal{G}^*)$. We shall explicitly construct its 2-bialgebra structure such that its self-duality is manifest.

2-algebra structure. We consider $D(\mathcal{G})$ defined in terms of the graded components given by

$$D(\mathcal{G})_0 \cong \mathcal{G}_0 \otimes \mathcal{G}_0^* \ni (x, f), \quad D(\mathcal{G})_{-1} \cong \mathcal{G}_{-1} \otimes \mathcal{G}_0^* \ni (y, g),$$

for which we have a "right-moving" semidirect product $\bar{\triangleright} = (\cdot, \bar{\triangleright})$, giving rise to $D(\mathcal{G})_{-1} \bar{\triangleright} D(\mathcal{G})_0$. Similarly, we also have a "left-moving" semidirect product $\bar{\triangleleft} = (\cdot, \bar{\triangleleft})$ giving rise to $D(\mathcal{G})_{-1} \bar{\triangleleft} D(\mathcal{G})_0$. The combined t -map $T = t \otimes t^*$ is equivariant with respect to these semidirect products

$$t^*(x \triangleright_0 g) = x \triangleright_{-1} t^*g, \quad t(y \triangleleft_0 f) = (ty) \triangleleft_{-1} f, \quad (3.8)$$

since the coadjoint action is 2-representation, while the commutativity of (6.1) implies

$$\begin{aligned} (ty) \triangleright_0 g &= \Upsilon_y(t^*g), & y \triangleleft_0 (t^*g) &= (ty) \tilde{\Upsilon}_g, \\ (ty) \triangleright_{-1} f &= t^*(\Upsilon_y f), & x \triangleleft_{-1} (t^*g) &= t(x \tilde{\Upsilon}_g). \end{aligned} \quad (3.9)$$

These are in fact generalizations of the Peiffer identity.

Proposition 3.2. *If $\bar{\triangleright}, \bar{\triangleleft}$ are given by the coadjoint representations (see (6.4)), then (3.9) reproduces the Peiffer identity.*

Proof. This is a direct computation. By the equality in the second row of (3.9), we have

$$\langle f, y \cdot ty' \rangle = -\langle t^* \Upsilon_y f, y' \rangle = -\langle (ty) \triangleright_{-1} f, y' \rangle = \langle f, ty \cdot y' \rangle,$$

giving $ty \cdot y' = y \cdot ty'$. Now by the fact that t is an algebra homomorphism, we have

$$\begin{aligned} \langle (ty) \triangleright_0 g, ty' \rangle &= -\langle g, (ty)(ty') \rangle = -\langle g, t(yy') \rangle, \\ \langle \Upsilon_y(t^*g), ty' \rangle &= -\langle t^*g, y \cdot ty' \rangle = -\langle g, t(y \cdot ty') \rangle, \end{aligned}$$

for which the first row of (3.9) states $yy' = y \cdot ty'$. Altogether yields

$$yy' = y \cdot t(y') = t(y) \cdot y'$$

for any $y, y' \in \mathcal{G}_{-1}$, which is precisely the Peiffer identity on \mathcal{G} . Similarly, if $\bar{\triangleleft}$ is the coadjoint representation then (3.9) reproduces the Peiffer identity on \mathcal{G}^* . \square

In other words, the Peiffer identity in $D(\mathcal{G})$ is *by definition* given as in (3.9). The multiplication between the sectors $\mathcal{G}_{-1}, \mathcal{G}_{-1}^*$ is given by $yg = \Upsilon_y(t^*g)$ and $gy = (ty)\check{\Upsilon}_g$.

Now that we have defined the product of the graded components and the t -map associated to $D(\mathcal{G})$, we can identify the bimodule structure.

We combine the right-moving $\overrightarrow{\times} = (\cdot, \triangleright)$ and left-moving $\overleftarrow{\times} = (\cdot, \triangleleft)$ multiplications on $D(\mathcal{G})$ to form $\hat{\cdot} = \overrightarrow{\times} + \overleftarrow{\times}$,

$$(z, h)\hat{\cdot}(z', h') = (z \cdot z' + z \overleftarrow{\triangleleft} h' + z' \overleftarrow{\triangleleft} h, h \cdot h' + z \overrightarrow{\triangleright} h' + z' \overrightarrow{\triangleright} h), \quad z, z' \in \mathcal{G}, h, h' \in \mathcal{G}^*. \quad (3.10)$$

Since $\hat{\cdot}$ is a combination of the internal 2-algebra structures of $\mathcal{G}, \mathcal{G}^*$ and the 2-representations $\overrightarrow{\triangleright}, \overleftarrow{\triangleleft}$, we have respectively the Peiffer conditions and associativity for $\mathcal{G}, \mathcal{G}^*$, as well as the 2-representation properties (3.8), (3.9) and the matched pair conditions (3.3), (3.4), (3.6). These imply that the map $\hat{\cdot}$

- (i) is associative,
- (ii) makes $D(\mathcal{G})_{-1}$ into a $D(\mathcal{G})_0$ -bimodule,
- (iii) satisfies the Peiffer conditions under $T = t \otimes t^*$.

Hence $(D(\mathcal{G}), \hat{\cdot})$ is a 2-algebra.

2-coalgebra structure. We intend now to construct the coproduct $\Delta_D : D(\mathcal{G}) \rightarrow D(\mathcal{G})^{2\otimes}$. We have to build the components

$$\begin{aligned} \Delta_{D_{-1}} &: D(\mathcal{G})_{-1} \rightarrow D(\mathcal{G})_{-1} \otimes D(\mathcal{G})_{-1} = (\mathcal{G}_{-1} \otimes \mathcal{G}_0^*) \otimes (\mathcal{G}_{-1} \otimes \mathcal{G}_0^*) \\ \Delta_{D_0} &: D(\mathcal{G})_0 \rightarrow (D(\mathcal{G})_{-1} \otimes D(\mathcal{G})_0) \oplus (D(\mathcal{G})_0 \otimes D(\mathcal{G})_{-1}) \end{aligned}$$

We can directly infer some of the components $\Delta_{D_{-1}}$ from the coproducts $\Delta_{-1}, \Delta_{-1}^*$ of $\mathcal{G}, \mathcal{G}^*$. Explicitly, it is defined as

$$\Delta_{D_{-1}}^d = \Delta_{-1} \otimes \Delta_{-1}^*.$$

This coproduct by construction encodes the separate coproducts $\Delta = \Delta_D|_{\mathcal{G}}, \Delta^* = \Delta_D|_{\mathcal{G}^*}$ by restriction and it is consistent with the products of each 2-algebras. These components are diagonal in a sense and we need to introduce some off diagonal contributions,

$$\xi_{-1} : \mathcal{G}_{-1} \rightarrow \mathcal{G}_0^* \otimes \mathcal{G}_{-1}, \quad \zeta_{-1} : \mathcal{G}_0^* \rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_0^*.$$

such that

$$\Delta_{D_{-1}} = (\Delta_D)_{-1}^d + \xi_{-1} \otimes \zeta_{-1}. \quad (3.11)$$

ξ_{-1} and ζ_{-1} can be interpreted as coactions and

are defined as dualized components of the coadjoint actions. Taking as usual $(x, f) \in D(\mathcal{G})_0 \cong \mathcal{G}_0 \otimes \mathcal{G}_{-1}^*$ and $(y, g) \in D(\mathcal{G})_{-1} \cong \mathcal{G}_{-1} \otimes \mathcal{G}_0^*$ we have

$$\langle \xi_{-1}(y), x \otimes f \rangle := \langle y, x \triangleright_{-1} f \rangle, \quad \langle \zeta_{-1}(g), f \otimes x \rangle := \langle g, x \triangleleft_{-1} f \rangle \quad (3.12)$$

These coactions are 2-algebra maps by (3.5), and hence $\Delta_{D_{-1}}$ satisfies (2.31) on $D(\mathcal{G})$.

In a similar way, Δ_{D_0} is also made of several components. We use the components $\Delta_0 : \mathcal{G}_0 \rightarrow (\mathcal{G}_0 \otimes \mathcal{G}_{-1}) \oplus (\mathcal{G}_{-1} \otimes \mathcal{G}_0)$ and $\Delta_0^* : \mathcal{G}_{-1}^* \rightarrow (\mathcal{G}_0^* \otimes \mathcal{G}_{-1}^*) \oplus (\mathcal{G}_{-1}^* \otimes \mathcal{G}_0^*)$ of \mathcal{G} and \mathcal{G}^* to define the "diagonal" contribution,

$$(\Delta_D^l)_0^d := \Delta_0^l \otimes \Delta_0^{*l}, \quad (\Delta_D^r)_0 := \Delta_0^r \otimes \Delta_0^{*r}.$$

Once again, by restriction, one recovers the separate coproducts $\Delta_0^{r,l}$ and $\Delta_0^{*r,l}$ on respectively \mathcal{G} and \mathcal{G}^* .

We also have to recover the mixed terms.

$$\begin{aligned} \xi_0^l : \mathcal{G}_0 &\rightarrow \mathcal{G}_0^* \otimes \mathcal{G}_0, & \xi_0^r : \mathcal{G}_0 &\rightarrow \mathcal{G}_{-1}^* \otimes \mathcal{G}_{-1} \\ \zeta_0^l : \mathcal{G}_{-1}^* &\rightarrow \mathcal{G}_{-1} \otimes \mathcal{G}_{-1}^*, & \zeta_0^r : \mathcal{G}_0 &\rightarrow \mathcal{G}_0 \otimes \mathcal{G}_0^*, \end{aligned}$$

such that

$$(\Delta_D^{r,l})_0 := (\Delta_D^l)_0^d + \xi_0^{r,l} \otimes \zeta_0^{r,l}. \quad (3.13)$$

These mixed terms are again obtained by dualizing the components of the coadjoint actions

$$\langle \xi_0^l(x), x' \otimes g \rangle := \langle x, x' \triangleright_0 g \rangle, \quad \langle \xi_0^r(x), y \otimes f \rangle := \langle x, \Upsilon_y f \rangle,$$

$$\langle \zeta_0^l(f), f' \otimes y \rangle := \langle f, y \triangleleft_{-1} f' \rangle, \quad \langle \zeta_0^r(f), g \otimes x \rangle := \langle f, x \tilde{\Upsilon}_g \rangle. \quad (3.14)$$

Once again, these coactions are 2-algebra maps by (3.5), and hence Δ_D satisfies (2.31) on $D(\mathcal{G})$.

We now need to show that it also satisfies (2.22), (2.24). We do this by leveraging the self-duality $D(\mathcal{G}) \cong D(\mathcal{G})^*$ under the natural non-degenerate self-pairing via (3.1) (cf. [41]),

$$\langle (z, h), (z', h') \rangle = \langle f, y' \rangle + \langle g, x' \rangle + \langle f', y \rangle + \langle g', x \rangle. \quad (3.15)$$

By **Proposition 3.1**, $(D(\mathcal{G}), \cdot)$ is an associative 2-algebra iff $(D(\mathcal{G})^* \cong D(\mathcal{G}), \Delta_D)$ is a coassociative 2-coalgebra, which implies (2.22)-(2.24) for Δ_D .

Definition 3.3. We call the 2-bialgebra

$$\mathcal{G} \overleftrightarrow{\bowtie} \mathcal{G}^* := D(\mathcal{G}) = (D(\mathcal{G})_{-1} \xrightarrow{T} D(\mathcal{G})_0, \hat{\cdot}, \Delta_D)$$

built out of the the matched pair of strict 2-bialgebras $(\mathcal{G}, \mathcal{G}^*)$ with the product, coproduct, and counit given respectively in (3.10), (3.11) and (3.13), (3.14), the **strict 2-quantum double** of \mathcal{G} .

3.3 Factorizability of 2-bialgebras

Conversely, we can determine when a strict 2-bialgebra is actually a strict 2-quantum double, which is given by a *factorizability/splitting* condition. In fact, we prove that any 2-bialgebra that factorizes appropriately into 2-bialgebras will automatically determine a 2-quantum double.

Theorem 3.1. *Suppose a (unital) 2-bialgebra $(\mathcal{K} = \mathcal{K}_{-1} \xrightarrow{T} \mathcal{K}_0, \hat{\cdot})$ factorizes into two (unital) sub-2-bialgebras \mathcal{G}, \mathcal{H} , meaning that there is a span of inclusions,*

$$\mathcal{G} \hookrightarrow \mathcal{K} \hookleftarrow \mathcal{H}, \quad (3.16)$$

such that $\hat{\cdot} \circ (\iota \otimes j)$ is an isomorphism of 2-vector spaces and such that the 2-sub-bialgebras \mathcal{G}, \mathcal{H} are dually paired, with their t -maps satisfying $\langle t_{\mathcal{G}} \cdot, \cdot \rangle = \langle \cdot, t_{\mathcal{H}} \cdot \rangle$. Then $(\mathcal{G}, \mathcal{H})$ is a matched pair and $\mathcal{K} \cong \mathcal{G} \overleftrightarrow{\bowtie} \mathcal{H}$.

Proof. Let $\mathcal{K} = \mathcal{K}_{-1} \xrightarrow{T} \mathcal{K}_0$ be a 2-bialgebra factorizing into two 2-subbialgebras \mathcal{G}, \mathcal{H} , with typical elements $w \in \mathcal{K}_0$ and $e \in \mathcal{K}_{-1}$. Its 2-algebra structure $\hat{\cdot}$ contains a multiplication ww' in \mathcal{K}_0 and a \mathcal{K}_0 -bimodule structure $w\hat{\cdot}e, e\hat{\cdot}w$ on \mathcal{K}_{-1} , which are both associative. Since (3.16) is a span of 2-vector spaces, we have

$$T \circ (\iota_{-1} \otimes j_{-1}) = (\iota_0 \circ t_{\mathcal{G}}) \otimes (j_0 \circ t_{\mathcal{H}}) = (\iota_0 \otimes j_0) \circ (t_{\mathcal{G}} \otimes t_{\mathcal{H}}),$$

where $t_{\mathcal{G}}, t_{\mathcal{H}}$ are the t -maps in \mathcal{G}, \mathcal{H} respectively, and ι_{-1}, ι_0 are the graded components of the inclusion ι ; similarly for j .

We now separate the bimodule structure $\hat{\cdot}$ into components according to the span (3.16),

$$\hat{\cdot} \equiv \hat{\cdot}|_{\text{im}(\iota_0 \otimes j_{-1})}, \quad \hat{\Upsilon} \equiv \hat{\cdot}|_{\text{im}(\iota_{-1} \otimes j_0)},$$

then for $e = \iota_{-1}(y), e' = j_{-1}(g)$ where $y \in \mathcal{G}_{-1}, g \in \mathcal{H}_{-1}$ we have

$$(T\iota_{-1}(y))\hat{\cdot}j_{-1}(g) = \iota_0(t_{\mathcal{G}}y)\hat{\cdot}j_{-1}(g).$$

By the Peiffer identity in \mathcal{K} , this should read as a left-multiplication of y on g . We lift this action along $t_{\mathcal{H}}$ to create a map $\hat{\Upsilon}_y : \mathcal{H}_0 \rightarrow \mathcal{H}_{-1}$, for which $\hat{\Upsilon}_y(t_{\mathcal{H}}g)$ denotes the left-multiplication of y by g . Similarly we have the lift $\hat{\Upsilon}_g : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1}$ of the right-multiplication of g on y .

Provided we identify $\hat{\Upsilon}_{y \otimes g} = \Upsilon_y \otimes \hat{\Upsilon}_g$, the Peiffer conditions in \mathcal{K} are then equivalent to the 2-representation properties (3.8), (3.9). In particular, the multiplication $y \cdot g = \Upsilon_y(t^*g) = (ty)\hat{\Upsilon}_g$ is given by the generalized Peiffer identity as shown in **Proposition 3.2**.

Now we prove that (3.16) is in fact a span of 2-algebras. Due to the linear isomorphism $\hat{\cdot} \circ (\iota \otimes j)$, there exists a tuple of well-defined linear maps $\Psi = (\Psi_0, \Psi_{-1}; \bar{\Psi}) : \mathcal{G} \otimes \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{G}$, called the **braided transposition**, such that

$$\begin{aligned} \iota_0(x) \cdot j_0(f) &= \cdot \circ (j_0 \otimes \iota_0) \circ \Psi_0(x \otimes f), \\ \iota_0(x)\hat{\cdot}j_{-1}(g) &= \hat{\Upsilon} \circ (j_{-1} \otimes \iota_0) \circ \Psi_{-1}^r(x \otimes g), \\ \iota_{-1}(y)\hat{\Upsilon}j_0(f) &= \hat{\cdot} \circ (j_0 \otimes \iota_{-1}) \circ \Psi_{-1}^l(y \otimes f), \\ \iota_{-1}(y) \cdot j_{-1}(g) &= \cdot \circ (j_{-1} \otimes \iota_{-1}) \circ \bar{\Psi}(y \otimes g), \end{aligned}$$

where $\Psi_{-1} = \Psi_{-1}^l + \Psi_{-1}^r$ and $x \in \mathcal{G}_0, y \in \mathcal{G}_{-1}, f \in \mathcal{H}_0, g \in \mathcal{H}_{-1}$. Due to Peiffer conditions on \mathcal{K} , these braiding maps are not independent and must satisfy

$$\begin{aligned} (t_{\mathcal{H}} \otimes 1) \circ \Psi_{-1}^r &= \Psi_0 \circ (1 \otimes t_{\mathcal{H}}), & (1 \otimes t_{\mathcal{G}}) \circ \Psi_{-1}^l &= \Psi_0 \circ (t_{\mathcal{G}} \otimes 1), \\ \Psi_{-1}^r \circ (t_{\mathcal{G}} \otimes 1) &= \bar{\Psi} = \Psi_{-1}^l \circ (1 \otimes t_{\mathcal{H}}). \end{aligned}$$

By collecting all of the graded components of Ψ in accordance with the shorthand notation $z = (y, x) \in \mathcal{G}, h = (g, f) \in \mathcal{H}$, the definition of Ψ can be concisely written as

$$\iota(z) \hat{\cdot} j(h) = \hat{\cdot} \circ (j \otimes \iota) \circ \Psi(z \otimes h), \quad (3.17)$$

and the relations between its components is summarized as

$$T' \circ \Psi_{-1} = \Psi_0 \circ T, \quad \bar{\Psi} = \Psi_{-1} \circ T, \quad (3.18)$$

where $T' = t_{\mathcal{H}} \otimes t_{\mathcal{G}}$ is the t -map of the 2-bialgebra $\mathcal{K}' \cong \mathcal{H} \otimes \mathcal{G}$ with \mathcal{G}, \mathcal{H} swapped in the span (3.16). (3.18) then implies in particular that $\Psi : \mathcal{K} \rightarrow \mathcal{K}'$ is a 2-vector space homomorphism.

We now proceed formally as in the 1-bialgebra case [7, 44]. The associativity in \mathcal{K} is

$$\begin{aligned} (\iota(z) \hat{\cdot} \iota(z')) \hat{\cdot} j(h) &= \iota(z) \hat{\cdot} (\iota(z') \hat{\cdot} j(h)), \\ (\iota(z) \hat{\cdot} j(h)) \hat{\cdot} j(h') &= \iota(z) \hat{\cdot} (j(h) \hat{\cdot} j(h')), \end{aligned}$$

which yields the **2-braiding relations**

$$\begin{aligned} \Psi \circ (\hat{\cdot} \otimes \text{id}) &= (\text{id} \otimes \hat{\cdot}) \circ \Psi_{12} \circ \Psi_{23}, \\ \Psi \circ (\text{id} \otimes \hat{\cdot}) &= (\hat{\cdot} \otimes \text{id}) \circ \Psi_{23} \circ \Psi_{12}. \end{aligned} \quad (3.19)$$

This then allows us to define the actions

$$\begin{aligned} \bar{\triangleright} &= (\text{id} \otimes \epsilon) \circ \Psi : \mathcal{G} \otimes \mathcal{H} \rightarrow \mathcal{H}, \\ \bar{\triangleleft} &= (\epsilon \otimes \text{id}) \circ \Psi : \mathcal{G} \otimes \mathcal{H} \rightarrow \mathcal{G}, \end{aligned}$$

where ϵ denotes the counit map. Applying $\text{id} \otimes \epsilon$ and $\epsilon \otimes \text{id}$ respectively to the first and second equation of (3.19) implies that $\bar{\triangleright}, \bar{\triangleleft}$ respect the semidirect product structures $\mathcal{G}_{-1} \rtimes \mathcal{G}_0, \mathcal{H}_{-1} \rtimes \mathcal{H}_0$, respectively. Together with our above result, (3.16) is in fact a span of 2-algebras.

We now prove that (3.16) is actually a span of 2-bialgebras, which proves the theorem. Applying $\epsilon \otimes \text{id}$ and $\text{id} \otimes \epsilon$ respectively to the first and second (3.19) yields

$$(z \cdot z') \bar{\triangleleft} h = \hat{\cdot} (z \bar{\triangleleft} \Psi(z' \otimes h)), \quad z \bar{\triangleright} (h \cdot h') = \hat{\cdot} (\Psi(z \otimes h) \bar{\triangleright} h'). \quad (3.20)$$

We now take the coproduct $\Delta_{\mathcal{K}} : \mathcal{K} \rightarrow \mathcal{K}^{2\otimes}$ on \mathcal{K} , given in components and Sweedler notation (see (2.16), (2.17)) by

$$(\Delta_{\mathcal{K}})_{-1}(e) = e_{(1)} \otimes e_{(2)}, \quad (\Delta_{\mathcal{K}})_0(w) = w_{(1)}^l \otimes w_{(2)}^l + w_{(1)}^r \otimes w_{(2)}^r;$$

note $w_{(1)}^l, w_{(2)}^r \in \mathcal{K}_{-1}$. With the span (3.16), we can write $w = \iota_0(x)j_0(f), e = \iota_{-1}(y)j_{-1}(g)$ for some appropriate elements x, f, y, g such that

$$\begin{aligned} (\Delta_{\mathcal{K}})_{-1}(y, g) &= (y_{(1)} \otimes g_{(1)}) \otimes (y_{(2)} \otimes g_{(2)}), \\ (\Delta_{\mathcal{K}})_0^l(x, f) &= (x_{(1)}^l \otimes f_{(1)}^l) \otimes (x_{(2)}^l \otimes f_{(2)}^l), \\ (\Delta_{\mathcal{K}})_0^r(x, f) &= (x_{(1)}^r \otimes f_{(1)}^r) \otimes (x_{(2)}^r \otimes f_{(2)}^r). \end{aligned}$$

This then allows us to define coproducts on \mathcal{G}, \mathcal{H} by

$$\begin{aligned} (\Delta_{\mathcal{G}})_{-1}(y) &= y_{(1)} \otimes y_{(2)}, & (\Delta_{\mathcal{G}})_0(x) &= x_{(1)}^l \otimes x_{(2)}^l + x_{(1)}^r \otimes x_{(2)}^r, \\ (\Delta_{\mathcal{H}})_{-1}(g) &= g_{(1)} \otimes g_{(2)}, & (\Delta_{\mathcal{H}})_0(f) &= f_{(1)}^l \otimes f_{(2)}^l + f_{(1)}^r \otimes f_{(2)}^r, \end{aligned}$$

whence $\Delta_{\mathcal{K}} = \Delta_{\mathcal{G} \otimes \mathcal{H}}$, which implies that $\hat{\cdot} \circ (\iota \otimes j)$ and $\hat{\cdot} \circ (j \otimes \iota)$ by construction respects the coproducts.

As such, Ψ is a 2-coalgebra map. In particular, we have

$$\Delta_{\mathcal{K}} \circ \Psi = (\Psi \otimes \Psi) \circ \Delta_{\mathcal{K}'}, \quad (\epsilon \otimes \epsilon) \circ \Psi = \epsilon \otimes \epsilon \quad (3.21)$$

where \mathcal{K}' is the 2-bialgebra with \mathcal{G}, \mathcal{H} swapped in the span (3.16). An application of $\epsilon \otimes \text{id} \otimes \epsilon \otimes \text{id}$ and $\text{id} \otimes \epsilon \otimes \text{id} \otimes \epsilon$ to (3.21) gives

$$\Delta_{\mathcal{G}} \circ \bar{\triangleleft} = (\bar{\triangleleft} \otimes \bar{\triangleleft}) \circ \Delta_{\mathcal{K}}, \quad \Delta_{\mathcal{H}} \circ \bar{\triangleright} = (\bar{\triangleright} \otimes \bar{\triangleright}) \circ \Delta_{\mathcal{K}},$$

which ensures that $\bar{\triangleright}, \bar{\triangleleft}$ are 2-coalgebra maps.

Now applying $\epsilon \otimes \text{id} \otimes \text{id} \otimes \epsilon$ and $\text{id} \otimes \epsilon \otimes \epsilon \otimes \text{id}$ to (3.21) yields

$$\begin{aligned} z_{(1)} \bar{\triangleleft} h_{(1)} \otimes z_{(2)} \bar{\triangleright} h_{(2)} &= \tau \circ \Psi(z \otimes h), \\ z_{(1)} \bar{\triangleright} h_{(1)} \otimes z_{(2)} \bar{\triangleleft} h_{(2)} &= \Psi(z \otimes h). \end{aligned}$$

Using the second equation, together with (3.20), gives (3.6) and

$$\begin{aligned} z \bar{\triangleright} (h \hat{\cdot}_{\mathcal{H}} h') &= \hat{\cdot}((z_{(1)} \bar{\triangleright} h_{(1)} \otimes z_{(2)} \bar{\triangleleft} h_{(2)}) \hat{\cdot} h') = (z_{(1)} \bar{\triangleright} h_{(1)}) \hat{\cdot}_{\mathcal{H}} ((z_{(2)} \bar{\triangleleft} h_{(2)}) \bar{\triangleright} h'), \\ (z \hat{\cdot}_{\mathcal{G}} z') \bar{\triangleleft} h &= \hat{\cdot}(z \bar{\triangleleft} (z'_{(1)} \bar{\triangleright} h_{(1)} \otimes z'_{(2)} \bar{\triangleleft} h_{(2)})) = (z \bar{\triangleleft} (z'_{(1)} \bar{\triangleright} h_{(1)})) \hat{\cdot}_{\mathcal{G}} (z'_{(2)} \bar{\triangleleft} h_{(2)}), \end{aligned}$$

which are precisely the matched pair conditions (3.3), (3.4) for $\hat{\cdot}_{\mathcal{G}} = \cdot, \hat{\cdot}_{\mathcal{H}} = \cdot^*$. On the other hand, using the first equation gives (3.5). Thus (3.16) is a span of 2-bialgebras and so $K \cong \mathcal{G} \bar{\bowtie} \mathcal{H}$. \square

Note that the span (3.16) factorizes the 2-algebra structure on \mathcal{K} into the right- $\bar{\bowtie} = (\hat{\cdot}_{\mathcal{G}}, \bar{\triangleleft})$ and left-moving $\bar{\bowtie} = (\hat{\cdot}_{\mathcal{H}}^{\text{opp}}, \bar{\triangleright}^{\text{opp}})$ 2-algebra structures. In other words, in order to identify \mathcal{K} with a 2-quantum double, we must have [44]

$$\mathcal{K} \cong \mathcal{G} \bar{\bowtie} \mathcal{H} \cong D(\mathcal{G}, \mathcal{H}^{\text{opp}}), \quad (3.22)$$

where \mathcal{H}^{opp} denotes the opposite 2-algebra; see Appendix A.

This is because, as can be seen in (3.14), the back-action $\bar{\triangleleft}$ is written from right to left.

3.4 Characterization of quantum 2-R-matrices

As we have mentioned in the beginning of this section, we wish to leverage the 2-quantum double construction we have given above in order to provide a notion of a quantum R-matrix on a 2-bialgebra \mathcal{G} . More precisely, we shall use the skew-pairing on \mathcal{G} used in forming the 2-quantum double $D(\mathcal{G}, \mathcal{G}) = \mathcal{G} \bar{\bowtie} \mathcal{G}^{\text{opp}}$ in order to provide a definition of the 2-R-matrix on \mathcal{G} . We shall show in Section 7.2 that such a characterization is *universal*, in the sense that our definition of a 2-R-matrix gives rise to a *braiding* on the 2-representations of \mathcal{G} .

Review of the 1-bialgebra case. We first recall the explicit construction of the universal R-matrix for the ordinary 1-bialgebra H . It was noted by Majid (see eg. [44, 36]) that, in forming the quantum double $D(H, H) = H \bowtie H^{\text{opp}}$ as a bicrossed product, the (non-degenerate) skew-pairing which dualizes H with itself satisfies

$$\begin{aligned} \langle x x', g \rangle_{\text{sk}} &= \langle x \otimes x', \Delta(g) \rangle_{\text{sk}} = \langle x, g_{(1)} \rangle_{\text{sk}} \langle x', g_{(2)} \rangle_{\text{sk}}, \\ \langle x, g g' \rangle_{\text{sk}} &= \langle \Delta(x), g' \otimes g \rangle_{\text{sk}} = \langle x_{(1)}, g' \rangle_{\text{sk}} \langle x_{(2)}, g \rangle_{\text{sk}}, \end{aligned}$$

where $x, x' \in H$ and $g, g' \in H^{\text{opp}} \cong H$. If we define this skew-pairing as a functional $\langle \cdot, \cdot \rangle_{\text{sk}} = R^* : H^{2\otimes} \rightarrow k$, then we see that the above conditions translate to

$$R^* \circ (\mu \otimes \text{id}) = R_{13}^* R_{23}^*, \quad R^* \circ (\text{id} \otimes \mu) = R_{13}^* R_{12}^*,$$

which is nothing but the defining properties of a *dual* R-matrix on H . Indeed, together with the property

$$g_{(1)} x_{(1)} R^*(x_{(2)}, g_{(2)}) = R^*(x_{(1)}, g_{(1)}) x_{(2)} g_{(2)}, \quad (3.23)$$

we obtain the (dual) Yang-Baxter equations [44, 7].

In other words, the duality pairing $\langle \cdot, \cdot \rangle_{\text{sk}}$ on the bicrossed product quantum double $D(H, H) = H \bowtie H^{\text{opp}}$ gives rise to a R-matrix R on H , and conversely any R-matrix gives rise to such a duality bilinear form. Moreover, this pairing is non-degenerate iff the corresponding R-matrix is quasitriangular (ie. R is invertible).

(Dual) 2-R-matrix. We now follow an analogous treatment to characterize dual 2-R-matrices of a quasitriangular 2-bialgebra \mathcal{G} . Take the 2-quantum double $D(\mathcal{G}, \mathcal{G})$, whose underlying duality pairing (3.15) is given by a non-degenerate self-duality skew-pairing $\langle \cdot, \cdot \rangle_{\text{sk}} : \mathcal{G} \otimes \mathcal{G} \rightarrow k$. Explicitly, this pairing satisfies

$$\begin{aligned} \langle x \cdot_l y, f \rangle_{\text{sk}} &= \langle x \otimes y, \Delta_0^l(f) \rangle_{\text{sk}}, & \langle y \cdot_r x, f \rangle_{\text{sk}} &= \langle y \otimes x, \Delta_0^r(f) \rangle_{\text{sk}}, \\ \langle x, f \cdot_l g \rangle_{\text{sk}} &= \langle \Delta_0^r(x), g \otimes f \rangle_{\text{sk}}, & \langle x, g \cdot f \rangle_{\text{sk}} &= \langle \Delta_0^l(x), f \otimes g \rangle_{\text{sk}}, \end{aligned} \quad (3.24)$$

and also in addition to the fact that it should respect the t -map $T = t \otimes t$ on $D(\mathcal{G}, \mathcal{G})$,

$$\langle ty, g \rangle_{\text{sk}} = \langle y, tg \rangle_{\text{sk}},$$

where $x, f, f' \in \mathcal{G}_0$ and $y, g \in \mathcal{G}_{-1}$. Writing the skew-pairing in terms of a functional $\mathcal{R}^* : \mathcal{G}^{2\otimes} \rightarrow k$ by

$$\mathcal{R}_l^*(y, f) = \langle y, f \rangle_{\text{sk}}, \quad \mathcal{R}_r^*(x, g) = \langle x, g \rangle_{\text{sk}},$$

we can rewrite (3.24) as

$$\begin{aligned} \mathcal{R}_l^* \circ (\cdot_l \otimes \text{id}) &= (\mathcal{R}_r^*)_{13} (\mathcal{R}_l^*)_{23}, & \mathcal{R}_l^* \circ (\cdot_r \otimes \text{id}) &= (\mathcal{R}_l^*)_{13} (\mathcal{R}_r^*)_{23}, \\ \mathcal{R}_r^* \circ (\text{id} \otimes \cdot_l) &= (\mathcal{R}_l^*)_{13} (\mathcal{R}_r^*)_{12}, & \mathcal{R}_r^* \circ (\text{id} \otimes \cdot_r) &= (\mathcal{R}_r^*)_{13} (\mathcal{R}_l^*)_{12}, \end{aligned}$$

where \cdot_l, \cdot_r denotes respectively the left and right \mathcal{G}_0 -actions on \mathcal{G}_{-1} . We also have the compatibility conditions with the t -map:

$$\mathcal{R}_r^* \circ (\text{id} \otimes t) = \mathcal{R}_l^* \circ (t \otimes \text{id}) \in \mathcal{G}_{-1}^{2\otimes}.$$

By dualizing the above functional \mathcal{R}^* , we are able to characterize the 2-R-matrix \mathcal{R} on \mathcal{G} .

Definition 3.4. A **2-R-matrix** associated to a 2-bialgebra $(\mathcal{G}, \cdot, \Delta)$ is an element $\mathcal{R} \in \mathcal{G} \otimes \mathcal{G}$ consisting of the graded components

$$\mathcal{R}^l \in \mathcal{G}_{-1} \otimes \mathcal{G}_0, \quad \mathcal{R}^r \in \mathcal{G}_0 \otimes \mathcal{G}_{-1}.$$

such that the following identities are satisfied:

1. the **compatibility with the coproduct**

$$\begin{aligned} (\Delta_0^l \otimes \text{id}) \mathcal{R}^r &= \mathcal{R}_{13}^l \cdot_l \mathcal{R}_{23}^r, & (\Delta_0^r \otimes \text{id}) \mathcal{R}^r &= \mathcal{R}_{13}^r \cdot_r \mathcal{R}_{23}^l, \\ (\text{id} \otimes \Delta_0^l) \mathcal{R}^l &= \mathcal{R}_{13}^l \cdot_r \mathcal{R}_{12}^r, & (\text{id} \otimes \Delta_0^r) \mathcal{R}^l &= \mathcal{R}_{13}^r \cdot_l \mathcal{R}_{12}^l, \end{aligned} \quad (3.25)$$

2. the **coproduct permutation identity**

$$\mathcal{R}^r \Delta_0^r(x) = (\sigma \circ \Delta_0^l(x)) \mathcal{R}^r, \quad \mathcal{R}^l \Delta_0^l(x) = (\sigma \circ \Delta_0^r(x)) \mathcal{R}^l \quad (3.26)$$

for each $x \in \mathcal{G}_0$, where $\sigma : \mathcal{G} \otimes \mathcal{G} \rightarrow \mathcal{G} \otimes \mathcal{G}$ is the permutation of tensor factors, and

3. the **equivariance condition**

$$D_t^- \mathcal{R} = 0 \iff (t \otimes \text{id}) \mathcal{R}^l = (\text{id} \otimes t) \mathcal{R}^r \in \mathcal{G}_0^{2\otimes}. \quad (3.27)$$

We call the tuple $(\mathcal{G}, \cdot, \Delta, \mathcal{R})$ a **quasitriangular 2-bialgebra** if $\mathcal{R}^l, \mathcal{R}^r$ are both invertible.

We now derive the categorified notion of the Yang-Baxter equations.

Proposition 3.3. *The 2-R-matrix of a quasitriangular 2-bialgebra $(\mathcal{G}, \cdot, \Delta, \mathcal{R})$ satisfies the 2-Yang-Baxter equations*

$$\begin{aligned} \mathcal{R}_{23}^r (\mathcal{R}_{13}^r \cdot_l \mathcal{R}_{12}^l) &= (\mathcal{R}_{12}^l \cdot_r \mathcal{R}_{13}^r) \mathcal{R}_{23}^r, & (\mathcal{R}_{23}^l \cdot_l \mathcal{R}_{13}^r) \mathcal{R}_{12}^r &= \mathcal{R}_{12}^r (\mathcal{R}_{13}^r \cdot_r \mathcal{R}_{23}^l), \\ \mathcal{R}_{23}^l (\mathcal{R}_{13}^l \cdot_r \mathcal{R}_{12}^r) &= (\mathcal{R}_{12}^r \cdot_l \mathcal{R}_{13}^l) \mathcal{R}_{23}^l, & (\mathcal{R}_{23}^r \cdot_r \mathcal{R}_{13}^l) \mathcal{R}_{12}^l &= \mathcal{R}_{12}^l (\mathcal{R}_{13}^l \cdot_l \mathcal{R}_{23}^r). \end{aligned} \quad (3.28)$$

Proof. Recall that \mathcal{R} is quasitriangular iff $\mathcal{R}^l, \mathcal{R}^r$ are square and invertible. This pairs \mathcal{G} with itself and hence $\dim \mathcal{G}_0 = \dim \mathcal{G}_{-1}$. We calculate $(\text{id} \otimes \sigma \circ \Delta_0^l) \mathcal{R}^l$ and $(\sigma \circ \Delta_0^l \otimes \text{id}) \mathcal{R}^r$, as well as $(\text{id} \otimes \sigma \circ \Delta_0^r) \mathcal{R}^l$ and $(\sigma \circ \Delta_0^r \otimes \text{id}) \mathcal{R}^r$ in two ways. First using (3.25), we have

$$\begin{aligned} (\text{id} \otimes \sigma \circ \Delta_0^l) \mathcal{R}^l &= (\text{id} \otimes \sigma) \mathcal{R}_{13}^l \cdot_r \mathcal{R}_{12}^r = \mathcal{R}_{12}^l \cdot_r \mathcal{R}_{13}^r, & (\sigma \circ \Delta_0^l \otimes \text{id}) \mathcal{R}^r &= (\sigma \otimes \text{id}) \mathcal{R}_{13}^r \cdot_l \mathcal{R}_{23}^r = \mathcal{R}_{23}^l \cdot_l \mathcal{R}_{13}^r, \\ (\text{id} \otimes \sigma \circ \Delta_0^r) \mathcal{R}^l &= (\text{id} \otimes \sigma) \mathcal{R}_{13}^r \cdot_l \mathcal{R}_{12}^l = \mathcal{R}_{12}^r \cdot_l \mathcal{R}_{13}^l, & (\sigma \circ \Delta_0^r \otimes \text{id}) \mathcal{R}^r &= (\sigma \otimes \text{id}) \mathcal{R}_{13}^l \cdot_r \mathcal{R}_{23}^l = \mathcal{R}_{23}^r \cdot_r \mathcal{R}_{13}^l. \end{aligned}$$

On the other hand from (3.26), we have that,

$$\begin{aligned} (\text{id} \otimes \sigma \circ \Delta_0^l) \mathcal{R}^l &= \mathcal{R}_{23}^r ((\text{id} \otimes \Delta_0^r) \mathcal{R}^l) \mathcal{R}_{23}^{r-1} = \mathcal{R}_{23}^r (\mathcal{R}_{13}^r \cdot_l \mathcal{R}_{12}^l) \mathcal{R}_{23}^{r-1} \\ (\sigma \circ \Delta_0^l \otimes \text{id}) \mathcal{R}^r &= \mathcal{R}_{12}^r ((\Delta_0^r \otimes \text{id}) \mathcal{R}^r) \mathcal{R}_{12}^{r-1} = \mathcal{R}_{12}^r (\mathcal{R}_{13}^r \cdot_r \mathcal{R}_{23}^l) \mathcal{R}_{12}^{r-1}, \\ (\text{id} \otimes \sigma \circ \Delta_0^r) \mathcal{R}^l &= \mathcal{R}_{23}^l ((\text{id} \otimes \Delta_0^l) \mathcal{R}^l) \mathcal{R}_{23}^{l-1} = \mathcal{R}_{23}^l (\mathcal{R}_{13}^l \cdot_r \mathcal{R}_{12}^r) \mathcal{R}_{23}^{l-1} \\ (\sigma \circ \Delta_0^r \otimes \text{id}) \mathcal{R}^r &= \mathcal{R}_{12}^l ((\Delta_0^l \otimes \text{id}) \mathcal{R}^r) \mathcal{R}_{12}^{l-1} = \mathcal{R}_{12}^l (\mathcal{R}_{13}^l \cdot_l \mathcal{R}_{23}^r) \mathcal{R}_{12}^{l-1} \end{aligned}$$

Putting each equation with its above counterpart leads to (3.28). \square

Remark 3.2. It is easy to see that, when $\mathcal{G} = D(\mathcal{H})$ is itself the 2-quantum double of a 2-bialgebra \mathcal{H} , then the skew-pairing required in forming the "2-quantum quadruple" $D(\mathcal{G}, \mathcal{G}) = D(D(\mathcal{H}), D(\mathcal{H}))$ splits into two copies the self-pairing form (3.15),

$$\langle [(y, x), (g, f)], [(y', x'), (g', f')] \rangle_{\text{sk}} = \langle (g, f), (y', x') \rangle + \langle (g', f'), (y, x) \rangle.$$

Since (3.15) is non-degenerate, then so is $\langle \cdot, \cdot \rangle_{\text{sk}}$ and the corresponding universal 2-R-matrix $\mathcal{R} \in D(\mathcal{H}, \mathcal{H})$ on $D(\mathcal{H})$ is automatically quasitriangular.

The (dual) 2-R-matrix from factorizability. Due to the factorizability result **Theorem 3.1**, we could have begun our characterization with a general associative 2-bialgebra \mathcal{K} which factorizes into two copies of \mathcal{G} , instead of the 2-quantum double $D(\mathcal{G}, \mathcal{G})$. This introduces the braided transposition $\Psi : \mathcal{G} \otimes \mathcal{G} \rightarrow \mathcal{G} \otimes \mathcal{G}$ given in (3.17) into the definition of the dual 2-R-matrix:

$$\mathcal{R}_l^* = \text{ev}_l \circ \Psi_{-1}^l, \quad \mathcal{R}_r^* = \text{ev}_r \circ \Psi_{-1}^r$$

where $\text{ev} = \text{ev}_l + \text{ev}_r$ is precisely the skew-pairing $\langle \cdot, \cdot \rangle_{\text{sk}}$ that we have introduced previously.

Dualizing this construction then gives

$$\mathcal{R}^l = \Psi_{-1}^l \circ \text{coev}_l, \quad \mathcal{R}^r = \Psi_{-1}^r \circ \text{coev}_r, \quad (3.29)$$

where $\text{coev} = \text{coev}_l + \text{coev}_r : k \rightarrow \mathcal{G} \otimes \mathcal{G}$ is the coevaluation. In other words, we are able to reconstruct the 2-R-matrix from the braided transposition Ψ on the 2-quantum double $\mathcal{K} \cong D(\mathcal{G}, \mathcal{G})$. Indeed, (3.18) gives the equivariance (3.27), and the relation (3.19) implies (3.25).

As mentioned in the proof of **Proposition 3.3**, having a *quasitriangular* structure on \mathcal{G} implies that \mathcal{G} is self-dual. This explains why only Ψ_{-1} appears in the reconstruction of the 2-R-matrix: the degree-0 component Ψ_0 dualizes to that in degree-(-2) $\bar{\Psi}^*$ for the dual $\mathcal{K}^* \cong \mathcal{K}$, which has the same t -map $T = t \otimes t$. As $\bar{\Psi}^*$ is determined by $(\Psi_{-1}^*)^{l,r} = \Psi_{-1}^{r,l}$ per (3.18), the component Ψ_0 is also completely determined by Ψ_{-1} .

Since **Theorem 3.1** implies that $\mathcal{K} \cong D(\mathcal{G}, \mathcal{G})$, this particular construction is isomorphic to the one we have given above directly from $D(\mathcal{G}, \mathcal{G})$. The characterization of the 2-R-matrix, **Definition 3.4**, thus does not depend on whether we induce R from the skew-pairing on $D(\mathcal{G}, \mathcal{G})$ or the braiding transposition Ψ on \mathcal{K} .

4 Weak 2-bialgebras

We now begin our endeavour to weaken the associativity conditions in the above 2-quantum double construction. The idea of *non-associative* 2-algebra has not been developed nearly as much as their associative counterpart, but we shall take inspiration from their Lie 2-algebra counterparts.

4.1 Definition

We provide the notion of a weak 2-algebra by generalizing **Definition 2.2**.

Definition 4.1. A **weak 2-algebra** $(\mathcal{G}, \mathcal{T})$ is a map $t : \mathcal{G}_{-1} \rightarrow \mathcal{G}_0$ between a pair of *not necessarily associative* algebras, together with an **invertible homotopy map** $\mathcal{T} : \mathcal{G}_0^{3\otimes} \rightarrow \mathcal{G}_{-1}$ such that we have the conditions (2.5), (2.6), as well as

1. the **weak 1-associativity**,

$$(xx')x'' - x(x'x'') = t\mathcal{T}(x, x', x''), \quad (yy')y'' - y(y'y'') = \mathcal{T}(ty, ty', ty'')$$

and the **weak bimodularity**,

$$\begin{aligned} x \cdot (x' \cdot y) - (xx') \cdot y &= \mathcal{T}(x, x', ty) & (x \cdot y) \cdot x' - x \cdot (y \cdot x') &= \mathcal{T}(x, ty, x'), \\ (y \cdot x) \cdot x' - y \cdot (xx') &= \mathcal{T}(ty, x, x'), \end{aligned}$$

for each $x, x', x'' \in \mathcal{G}_0$ and $y, y', y'' \in \mathcal{G}_{-1}$,

2. the **Hochschild 3-cocycle condition**,

$$x_1 \cdot \mathcal{T}(x_2, x_3, x_4) + \mathcal{T}(x_1, x_2, x_3) \cdot x_4 = \mathcal{T}(x_1x_2, x_3, x_4) - \mathcal{T}(x_1, x_2x_3, x_4) + \mathcal{T}(x_1, x_2, x_3x_4)$$

for each $x_1, \dots, x_4 \in \mathcal{G}_0$.

We call $(\mathcal{G}, \mathcal{T})$ a unital weak 2-algebra if we have a unit map $\eta : k \rightarrow \mathcal{G}$ that satisfies the usual conditions (2.7), and such that \mathcal{T} is **normalized** — namely it vanishes whenever any of its arguments are 0 or η_0 .

We note here that this structure is precisely the definition of a homotopy A_∞ -algebra \mathcal{G} , together with the Peiffer identity constraint (2.6). The correspondence between the n -nary product $m_n \in \text{Hom}^{n-2}(\mathcal{G}^{n\otimes}, \mathcal{G})$ and the weak 2-algebra structure is given by

$$m_1(-) = t(-), \quad m_2(-, -) = (-, -, -), \quad m_3(-, -, -) = \mathcal{T}(-, -, -),$$

with $m_n = 0$ trivial for $n \geq 4$. Nevertheless, we shall see that the Peiffer identity on \mathcal{G} shall play a very important role.

Similar to **Remark 2.1**, the Peiffer identity implies the further constraints

$$\begin{aligned} (x \cdot y)y' - x \cdot (yy') &= \mathcal{T}(x, ty, ty'), & (y \cdot x)y' - y(x \cdot y') &= \mathcal{T}(ty, x, ty'), \\ y(y' \cdot x) - (yy') \cdot x &= \mathcal{T}(ty, ty', x) \end{aligned}$$

for $t \neq 0$, where $x \in \mathcal{G}_0, y, y' \in \mathcal{G}_{-1}$.

4.1.1 Weak 2-algebra homomorphisms

We define a map between weak 2-algebras $(\mathcal{G}, \mathcal{T}) \rightarrow (\mathcal{G}', \mathcal{T}')$ as a cochain map $F = (F_1, F_0, F_{-1}) : \mathcal{G} \rightarrow \mathcal{G}'$:

$$F_1 : \mathcal{G}_0^{2\otimes} \rightarrow \mathcal{G}'_{-1}, \quad F_0 : \mathcal{G}_0 \rightarrow \mathcal{G}'_0, \quad F_{-1} : \mathcal{G}_{-1} \rightarrow \mathcal{G}'_{-1},$$

such that $t' \circ F_{-1} = F_0 \circ t$ and the following conditions are satisfied,

$$\begin{aligned} t'F_1(x, x') &= F_0(xx') - F_0(x)F_0(x'), \\ F_1(x, ty) &= F_{-1}(x \cdot y) - F_0(x) \cdot' F_{-1}(y), \\ F_1(ty, x) &= F_{-1}(y \cdot x) - F_{-1}(y) \cdot' F_0(x) \\ \mathcal{T}'(F_0(x), F_0(x'), F_0(x'')) - F_{-1}(\mathcal{T}(x, x', x'')) &= F_0(x) \cdot' F_1(x', x'') - F_1(xx', x'') \\ &\quad + F_1(x, x'x'') - F_1(x, x') \cdot' F_0(x''). \end{aligned} \quad (4.1)$$

In other words, F_1 contributes as an "obstruction" for the other components (F_0, F_{-1}) to define a strict 2-algebra homomorphism, but only up to homotopy in the sense that F_1 by definition (see the last equation of (4.1)) gives an explicit trivialization of the Hochschild cohomology class $[\mathcal{T}' \circ F_0] - [F_{-1} \circ \mathcal{T}] = 0$.

It can then be deduced that quasi-isomorphism classes of weak 2-algebras — where $\mathcal{G} \sim \mathcal{G}'$ are said to be quasi-isomorphic iff there exists a weakly invertible cochain map (4.1) between them — is still labeled by Hochschild cohomology classes $\mathcal{T} \in HH^3(\mathcal{N}, V)$, where $\mathcal{N} = \text{coker } t$ and $V = \text{ker } t$. In particular, $(\mathcal{G}, \mathcal{T})$ is always quasi-isomorphic to its skeleton $(\mathcal{N} \xrightarrow{0} V, [\mathcal{T}])$, which is in fact associative.

4.1.2 Example: weak 2-group algebras

Our definition of the weak 2-algebra is less natural in the context of groups, as weakening the associativity in a group G reads

$$(xx')x'' = \tau(x, x', x'') \cdot x(x'x''), \quad x, x', x'' \in G,$$

which does not reproduce our above notion of a weak 2-algebra when we pass to the group algebra kG . There is hence an inherent disconnect between a natural notion of a "weak 2-group" and that of a weak 2-algebra.

Consider a skeletal 2-group G with Hoang data (G_0, G_{-1}, τ) as a categorical group [60]. There is a copy of G_{-1} over each object $x \in G_0$ as the space of endomorphisms on x . Notice here that G_0 is a genuine group with an associative product, but there are distinguished associator isomorphisms valued in G_{-1} ,

$$\tau(x, x', x'') : (xx')x'' \rightarrow x(x'x''), \quad x, x', x'' \in G_0,$$

that represents the Postnikov class $\tau \in H^3(G_0, G_{-1})$. The 3-cocycle condition for τ holds due to the pentagon relation. Note the group G_{-1} is Abelian (G being skeletal) and we use the addition for its product.

We wish to take the same point of view with weak 2-algebras. Suppose $\mathcal{G} : \mathcal{G}_{-1} \xrightarrow{0} \mathcal{G}_0$ is a weak skeletal 2-algebra, then \mathcal{G}_0 is in fact associative, and \mathcal{G}_{-1} is an associative \mathcal{G}_0 -bimodule. The difference with the strict case is that there are now distinguished associator isomorphisms

$$\mathcal{T}(x, x', x'') : (xx')x'' \rightarrow x(x'x''), \quad (4.2)$$

which is given by the data of the homotopy map \mathcal{T} . Unfortunately, the construction of a 2-group algebra kG described in (2.14), (2.15) does *not* preserve the classifying 3-cocycles. This is because that the t -map on kG is the augmentation, and hence kG is classified by $HH^3(kG_0/k, \varepsilon)$ where $\text{im } t \cong k$ while $\varepsilon = \text{ker } t$ is the augmentation ideal.

It is, however, possible to construct a version of the group 2-algebra kG that does give rise to a correspondence of the form

$$H^3(G_0, G_{-1}) \rightarrow HH^3(kG_0, kG_{-1}), \quad \tau \mapsto \mathcal{T},$$

as we have noted at the end of Section 2.1.2. Moreover, for certain skeletal 2-groups, one may even leverage the natural (ring!) isomorphism $HH^*(kN, kN) \cong H^*(N, kN)$ [57] to produce a bijective correspondence between the classifying 3-cocycles. This will be important in an accompanying work [58].

4.2 Weak 2-coalgebras

We begin by defining the notion of a weak 2-coalgebra. Recall that the weakening in **Definition 4.1** concerns only the associativity of the 2-algebra structure. Correspondingly, the weakening of a 2-coalgebra should only concern the coassociativity.

For brevity of notation later, we first rewrite the equations (2.18), (2.20) in a more concise way. Consider coassociativity (2.18); we naturally extend Δ_{-1} to act on tensor products (with alternating sign) such that

$$\Delta_{-1} \circ \Delta_{-1} \equiv (\text{id} \otimes \Delta_{-1}) \circ \Delta_{-1} - (\Delta_{-1} \otimes \text{id}) \circ \Delta_{-1}.$$

Secondly, we recombine $\Delta_0 = \Delta_0^l + \Delta_0^r$ and extend it as well to tensor products, such that

$$\begin{aligned} (\Delta_{-1} + \Delta_0) \circ \Delta_0 &\equiv [(\Delta_{-1} \otimes \text{id}) \circ \Delta_0^l - (\text{id} \otimes \Delta_0^l) \circ \Delta_0^l] \\ &\quad + [(\text{id} \otimes \Delta_{-1}) \circ \Delta_0^r - (\Delta_0^r \otimes \text{id}) \circ \Delta_0^r] \end{aligned}$$

encodes two expressions in (2.20). We extend the t -map to the triple tensor product,

$$D_t = \text{id} \otimes \text{id} \otimes t - \text{id} \otimes t \otimes \text{id} + t \otimes \text{id} \otimes \text{id},$$

such that the equation

$$D_t \circ \Delta_0 \circ \Delta_0 = \Delta_0 \circ D_t \circ \Delta_0$$

encodes all three equations in (2.29). For convenience, we define also the map

$$D_t[2] \equiv t \otimes t \otimes \text{id} - t \otimes \text{id} \otimes t + \text{id} \otimes t \otimes t,$$

which is an extension of two applications of t to the 3-fold tensor product.

Definition 4.2. Let $\Delta_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1}^{3\otimes}$ denote an *invertible* trilinear map. Together with the maps (Δ_{-1}, Δ_0) defined as in (2.16), we say that the tuple $(\mathcal{G}, \Delta = (\Delta_{-1}, \Delta_0, \Delta_1))$ is a **weak 2-coalgebra** iff coequivariance (2.22), coPeiffer identity (2.24), **weak coassociativity**

$$\begin{aligned} \Delta_{-1} \circ \Delta_{-1} &= \Delta_1 \circ t, \\ (\Delta_{-1} + \Delta_0) \circ \Delta_0 &= D_t \circ \Delta_1, \end{aligned} \tag{4.3}$$

and **2-coassociativity**

$$\Delta_1 \circ \Delta_0 = \Delta_{-1} \circ \Delta_1 \tag{4.4}$$

are satisfied. In which case we call Δ_1 the **coassociator** of \mathcal{G} .

We call (\mathcal{G}, Δ) **counital** if it is equipped with a counit $\epsilon : k \rightarrow \mathcal{G}$ satisfying the usual conditions, and $\epsilon \circ \Delta_1 = 0$.

Notice that, provided the coequivariance and the coPeiffer identity are satisfied, applying one more t -map to (4.3) yields

$$\Delta_0' \circ \Delta_0 - \Delta_0 \circ \Delta_0' = D_t[2] \circ \Delta_1, \tag{4.5}$$

which is a monoidal weakening of the condition (2.29). Similarly, applying the t -map yet once more gives a map $\Phi \equiv (t \otimes t \otimes t)\Delta_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_0^{3\otimes}$ that lands only in \mathcal{G}_0 . We write this element multiplicatively such that

$$(\Delta_0' \otimes \text{id}) \circ \Delta_0' = \Phi \circ (\text{id} \otimes \Delta_0') \circ \Delta_0'. \tag{4.6}$$

Recall that, in the skeletal case where $t = 0$, the coproducts $\Delta_{-1}, \Delta_0, \Delta_0'$ are independent and hence (4.6) should also be imposed independently from (4.3).

4.3 Weak 2-bialgebras

Suppose now $(\mathcal{G}, \mathcal{T})$ is a weak 2-algebra equipped with the tuple $\Delta = (\Delta_{-1}, \Delta_0, \Delta_1)$ of linear maps. Recall the Sweedler notation (2.19) for $\Delta_0' : \mathcal{G}_0 \rightarrow \mathcal{G}_0^{2\otimes}$. We use it to state the condition that the coassociator Δ_1 preserves the algebra structure on \mathcal{G} ,

$$\begin{aligned} (\Delta_{-1} \circ \mathcal{T})(x, x', x'') &= \mathcal{T}(\bar{x}_{(1)}, \bar{x}'_{(1)}, \bar{x}''_{(1)}) \otimes \mathcal{T}(\bar{x}_{(2)}, \bar{x}'_{(2)}, \bar{x}''_{(2)}), \\ \Delta_1(xx') &= x_{(1)}x'_{(1)} \otimes x_{(2)}x'_{(2)} \otimes x_{(3)}x'_{(3)}, \end{aligned} \tag{4.7}$$

for $x, x', x'' \in \mathcal{G}_0$. Note that $\bar{x}_{(1)}, \bar{x}_{(2)} \in \mathcal{G}_0$ are not to be confused with the elements $x_{(1)}^{l,r}$ in (2.17).

Definition 4.3. The tuple $(\mathcal{G}, \mathcal{T}, \Delta)$ is a (unital) **weak 2-bialgebra** iff $(\mathcal{G}, \mathcal{T})$ is a weak 2-algebra and (\mathcal{G}, Δ) is a (counital) weak 2-coalgebra. Equivalently, $(\mathcal{G}, \mathcal{T}, \Delta)$ is a weak 2-bialgebra iff the tuple $\Delta = (\Delta_1, \Delta_0, \Delta_{-1})$ satisfies (2.24), (2.22), (4.3)-(4.5), (2.31) and (4.7).

A weak 2-bialgebra $(\mathcal{G}, \mathcal{T}, \Delta)$ is called **quasi-2-bialgebra** if $\mathcal{T} = 0$.

Similar to what we have done for the strict case, we suppose \mathcal{G} is dually paired with its dual 2-algebra through (3.1). The coassociator Δ_1 on \mathcal{G} induces a linear map $\mathcal{T}^* : \mathcal{G}_{-1}^* \rightarrow \mathcal{G}_0^*$ by

$$\langle f \otimes f' \otimes f'', \Delta_1(x) \rangle = \langle \mathcal{T}^*(f, f', f''), x \rangle.$$

Similarly, the Hochschild 3-cocycle \mathcal{T} on \mathcal{G} induces a linear map $\Delta_1^* : \mathcal{G}_{-1}^* \rightarrow (\mathcal{G}_0^*)^{3\otimes}$. We form the tuple $\Delta^* = (\Delta_1^*, \Delta_0^*, \Delta_{-1}^*)$.

Proposition 4.1. *Let $\mathcal{G}, \mathcal{G}^*$ be dually paired, then $(\mathcal{G}, \mathcal{T}, \Delta)$ is a (unital) weak 2-bialgebra iff $(\mathcal{G}^*, \mathcal{T}^*, \Delta^*)$ is a (unital) weak 2-bialgebra.*

Proof. Let $(\mathcal{G}^*, \mathcal{T}^*, \Delta^*)$ be the dual weak 2-bialgebra. By a straightforward computation, it is obvious that the conditions (2.22), (2.24), (4.3), (4.5) on $(\mathcal{G}^*, \Delta^*)$ hold iff imply the Peiffer conditions, weak 1-associativity and weak bimodularity hold for the 2-algebra \mathcal{G} . The fact that $(\Delta_{-1}^*, \Delta_0^*)$ are 2-algebra maps, as well as the units/counits, are treated the same way as in the proof of **Proposition 3.1**.

The two non-trivial identities to check are (4.4), (4.7). We will use the Sweedler notation for Δ_1 , $\Delta_1 x = x_{(1)} \otimes x_{(2)} \otimes x_{(3)}$. Note $x \in \mathcal{G}_0$, but $x_{(i)} \in \mathcal{G}_{-1}$ for $i = 1, 2, 3$.

We begin first with the latter, and check that Δ_1 is a 2-algebra map. We have

$$\langle \Delta_1^*(f f'), x \otimes x' \otimes x'' \rangle = \langle f f', \mathcal{T}(x, x', x'') \rangle = \langle f \otimes f', \Delta_{-1} \mathcal{T}(x, x', x'') \rangle,$$

while (recall Δ_0^* dualizes to the multiplication on \mathcal{G}_0^* , and $f_{(i)} \in \mathcal{G}_{-1}^*$)

$$\begin{aligned} \langle f_{(1)} f'_{(1)} \otimes f_{(2)} f'_{(2)} \otimes f_{(3)} f'_{(3)}, x \otimes x' \otimes x'' \rangle &= \langle \bigotimes_{i=1}^3 f_{(i)} \otimes f'_{(i)}, \Delta_0^*(x) \otimes \Delta_0^*(x') \otimes \Delta_0^*(x'') \rangle \\ &= \langle (f_{(1)} \otimes f'_{(1)}) \otimes (f_{(2)} \otimes f'_{(2)}) \otimes (f_{(3)} \otimes f'_{(3)}), \\ &\quad (\bar{x}_{(1)} \otimes \bar{x}_{(2)}) \otimes (\bar{x}'_{(1)} \otimes \bar{x}'_{(2)}) \otimes (\bar{x}''_{(1)} \otimes \bar{x}''_{(2)}) \rangle \\ &= \langle \Delta_1^*(f) \otimes \Delta_1^*(f'), (\bar{x}_{(1)} \otimes \bar{x}'_{(1)} \otimes \bar{x}''_{(1)}) \otimes (\bar{x}_{(2)} \otimes \bar{x}'_{(2)} \otimes \bar{x}''_{(2)}) \rangle \\ &= \langle f \otimes f', \mathcal{T}(\bar{x}_{(1)}, \bar{x}'_{(1)}, \bar{x}''_{(1)}) \otimes \mathcal{T}(\bar{x}_{(2)}, \bar{x}'_{(2)}, \bar{x}''_{(2)}) \rangle, \end{aligned}$$

meaning that Δ_1 is a 2-algebra map on $(\mathcal{G}, \mathcal{T})$ iff Δ_1^* is also one on $(\mathcal{G}^*, \mathcal{T}^*)$.

We now check (4.5). Fix arbitrary elements $f_1, \dots, f_4 \in \mathcal{G}_{-1}^*$, we compute

$$\begin{aligned} \langle f_1 \cdot \mathcal{T}^*(f_2, f_3, f_4), x \rangle &= \langle f_1 \otimes \mathcal{T}^*(f_2, f_3, f_4), \Delta_0^l(x) \rangle = \langle f_1 \otimes \dots \otimes f_4, (1 \otimes \Delta_1) \circ \Delta_0^l(x) \rangle, \\ \langle \mathcal{T}^*(f_1, f_2, f_3) \cdot f_4, x \rangle &= \langle \mathcal{T}^*(f_1, f_2, f_3) \otimes f_4, \Delta_0^r(x) \rangle = \langle f_1 \otimes \dots \otimes f_4, (\Delta_1 \otimes 1) \circ \Delta_0^r(x) \rangle; \end{aligned}$$

on the other hand, we have

$$\begin{aligned} \langle \mathcal{T}^*(f_1 f_2, f_3, f_4), x \rangle &= \langle f_1 f_2 \otimes f_3 \otimes f_4, \Delta_1(x) \rangle = \langle f_1 \otimes \dots \otimes f_4, (\Delta_{-1} \otimes 1 \otimes 1) \circ \Delta_1(x) \rangle, \\ \langle \mathcal{T}^*(f_1, f_2, f_3 f_4), x \rangle &= \langle f_1 \otimes f_2 \otimes f_3 f_4, \Delta_1(x) \rangle = \langle f_1 \otimes \dots \otimes f_4, (1 \otimes 1 \otimes \Delta_{-1}) \circ \Delta_1(x) \rangle, \\ \langle \mathcal{T}^*(f_1, f_2 f_3, f_4), x \rangle &= \langle f_1 \otimes f_2 f_3 \otimes f_4, \Delta_1(x) \rangle = \langle f_1 \otimes \dots \otimes f_4, (1 \otimes \Delta_{-1} \otimes 1) \circ \Delta_1(x) \rangle. \end{aligned}$$

By summing these up, the Hochschild 3-cocycle condition for \mathcal{T}^* is equivalent to

$$(1 \otimes \Delta_1) \circ \Delta_0^l + (\Delta_1 \otimes 1) \circ \Delta_0^r = \Delta_1 \circ \Delta_0 = \Delta_{-1} \circ \Delta_1.$$

which is exactly (4.5). \square

Given $(\mathcal{G}, \mathcal{G}^*)$ are dually paired 2-bialgebras, we see that a quasi-2-bialgebra $(\mathcal{G}, \mathcal{T} = 0, \Delta)$ encode the same data as a weak but coassociative 2-bialgebra $(\mathcal{G}^*, \mathcal{T}^*, \Delta^*)$, in which $\Delta_1^* = 0$.

5 Weak (skeletal) 2-quantum double

Let $\mathcal{G}, \mathcal{G}^*$ be dually paired (weak) 2-bialgebras. To form its weak 2-quantum double, we require them to act on each other *weakly*. This means, in particular, that the coadjoint actions $\triangleright, \triangleleft$ now come with the additional components

$$\triangleright_1 : \mathcal{G}_0^{2\otimes} \rightarrow \text{Hom}(\mathcal{G}_{-1}^*, \mathcal{G}_0^*), \quad \triangleleft_1 : (\mathcal{G}_{-1}^*)^{2\otimes} \rightarrow \text{Hom}(\mathcal{G}_0, \mathcal{G}_{-1}).$$

This will be justified further in Section 6.1 where we show that the coadjoint action can be interpreted weak representation. More specifically, just like the product and actions in (3.2) contribute to defining dually some (crossed) relations, the cocycle \mathcal{T} should also contribute dually to the adjoint action. This is what \triangleright_1 and \triangleleft_1 stand for, as we will see in (5.1).

To construct non-skeletal weak 2-quantum doubles, one must explicitly keep track of how $\mathcal{T}, \mathcal{T}^*, \triangleright_1, \triangleleft_1$ appear in the crossed-relations (3.3), (3.4), (3.5). This is a Herculean task that we leave to the ambitious reader. We will restrict from now on to the skeletal case when defining the quantum double.

5.1 Matched pair of skeletal weak 2-bialgebras

Though the situation is drastically simplified in the skeletal case $t = 0$, it is now important for us to keep track of the associators. We shall do this by using the notation of (4.2).

The non-trivial crossed relations (3.7), in particular, are attached with the components $\triangleright_1, \triangleleft_1$ of the coadjoint actions,

$$\begin{aligned} (x) \triangleleft_1^{f, f'} & : x \triangleright_{-1} (ff') \xrightarrow{\sim} (x_{(1)} \bar{\triangleright} f_{(1)}) \cdot * ((x_{(2)} \bar{\triangleleft} f_{(2)}) \bar{\triangleright} f'), \\ \triangleright_1^{x, x'}(f) & : (xx') \triangleleft_{-1} f \xrightarrow{\sim} (x \bar{\triangleleft} (x'_{(1)} \bar{\triangleright} f_{(1)})) \cdot (x_{(2)} \bar{\triangleleft} f_{(2)}), \end{aligned}$$

where we have made use of the shorthand notation defined in Remark 3.1. These come together to allow us to define a Hochschild 3-cochain on the 2-quantum double $D(\mathcal{G})$,

$$\mathcal{T}_D : D(\mathcal{G})_0^{3\otimes} \rightarrow D(\mathcal{G})_{-1}, \quad \mathcal{T}_D(w, w', w'') = \begin{cases} \mathcal{T}(x, x', x'') \\ \triangleright_1^{x, x'}(f'') \\ (x) \triangleleft_1^{f', f''} \\ \mathcal{T}^*(f, f', f'') \end{cases}, \quad (5.1)$$

where $w = (x, f) \in D(\mathcal{G})_0$ is a degree-0 element, with $x \in \mathcal{G}_0$ and $f \in \mathcal{G}_{-1}^*$.

Definition 5.1. The pair $(\mathcal{G}, \mathcal{G}^*)$ of mutually paired weak skeletal 2-bialgebras forms a **(skeletal) matched pair** iff, in addition to the compatibility conditions (3.3)-(3.6), the 3-cochain \mathcal{T}_D defined in (5.1) is a Hochschild 3-cocycle on $D(\mathcal{G}) \cong \mathcal{G} \otimes \mathcal{G}^*$.

For arguments contained solely in \mathcal{G}_0 or \mathcal{G}_{-1}^* , this condition merely states the 3-cocycle conditions for $\mathcal{T}, \mathcal{T}^*$, respectively. The other ones mix non-trivially the different components of the 3-cocycle \mathcal{T}_D ,

$$\begin{aligned} x_1 \triangleright_0 (\triangleright_1^{x_2, x_3}(f)) - \mathcal{T}(x_1, x_2, x_3) \triangleleft_0 f & = \triangleright_1^{x_1 x_2, x_3}(f) - \triangleright_1^{x_1, x_2 x_3}(f) + \mathcal{T}(x_1, x_2, x_3 \triangleleft_{-1} f), \\ x_1 \cdot (x_2) \triangleleft_1^{f_1, f_2} - \triangleright_1^{x_1, x_2}(f_1) \cdot * f_2 & = (x_1 x_2) \triangleleft_1^{f_1, f_2} - (x_1) \triangleleft_1^{x_2 \triangleright_{-1} f_1, f_2} + \triangleright_1^{x_1, x_2}(f_1 f_2), \\ x \triangleright_0 \mathcal{T}^*(f_1, f_2, f_3) - ((x) \triangleleft_1^{f_1, f_2}) \triangleleft_0 f_3 & = \mathcal{T}^*(x \triangleright_{-1} f_1, f_2, f_3) - (x) \triangleleft_1^{f_1 f_2, f_3} + (x) \triangleleft_1^{f_1, f_2 f_3}. \end{aligned} \quad (5.2)$$

Then, we construct $D(\mathcal{G})$ as a 2-bialgebra as in Section 3.

Since we are in the skeletal case, it is easy to see from (4.1) that the 2-quantum double is *weakly* self-dual $D(\mathcal{G}) \sim D(\mathcal{G})^*$, where we recall \sim denotes equivalence of 2-algebras under the classification result **Theorem 2.2**. This means that the associated Hochschild 3-cocycles $\mathcal{T}_D, \mathcal{T}_D^*$ are cohomologous, where

$$\mathcal{T}_D^* : D(\mathcal{G})_0^{3\otimes} \rightarrow D(\mathcal{G})_{-1}, \quad \mathcal{T}_D^*(w, w', w'') = \begin{cases} \overset{\circ}{\mathcal{T}}(f, f', f'') \\ \overset{\circ}{\triangleright}_1^{f, f'}(x'') \\ (f) \overset{\circ}{\triangleleft}_1^{x', x''} \\ \overset{\circ}{\mathcal{T}}^*(x, x', x'') \end{cases},$$

denotes the dual of the 3-cocycle \mathcal{T}_D . The "dual" version of (5.2) reads

$$\begin{aligned} f_1 \triangleleft_0 (\overset{\circ}{\triangleright}_1^{f_2, f_3}(x)) - \overset{\circ}{\mathcal{T}}(f_1, f_2, f_3) \triangleright_0 x & = \overset{\circ}{\triangleright}_1^{f_1 f_2, f_3}(x) - \overset{\circ}{\triangleright}_1^{f_1, f_2 f_3}(x) + \overset{\circ}{\mathcal{T}}(f_1, f_2, f_3 \triangleleft_{-1} x), \\ f_1 \cdot * (f_2) \overset{\circ}{\triangleleft}_1^{x_1, x_2} - \overset{\circ}{\triangleright}_1^{f_1, f_2}(x_1) \cdot x_2 & = (f_1 f_2) \overset{\circ}{\triangleleft}_1^{x_1, x_2} - (f_1) \overset{\circ}{\triangleleft}_1^{f_2 \triangleleft_{-1} x_1, x_2} + \overset{\circ}{\triangleright}_1^{f_1, f_2}(x_1 x_2), \\ f \triangleleft_0 \overset{\circ}{\mathcal{T}}^*(x_1, x_2, x_3) - ((f) \overset{\circ}{\triangleleft}_1^{x_1, x_2}) \triangleright_0 x_3 & = \overset{\circ}{\mathcal{T}}^*(f \triangleright_{-1} x_1, x_2, x_3) - (f) \overset{\circ}{\triangleleft}_1^{x_1 x_2, x_3} + (f) \overset{\circ}{\triangleleft}_1^{x_1, x_2 x_3}. \end{aligned} \quad (5.3)$$

It is important to note that the components $\triangleright_1, \triangleleft_1$ do *not* form Hochschild 3-cocycles by themselves, and similarly for the components $\overset{\circ}{\triangleright}_1, \overset{\circ}{\triangleleft}_1$.

5.2 Factorizability of weak 2-bialgebras

We now prove the analogue of **Theorem 3.1**.

Theorem 5.1. *Suppose $(\mathcal{K}, \hat{\cdot}, \mathcal{T}_{\mathcal{K}})$ is a weak 2-bialgebra that weakly factors into two skeletal weak sub-2-bialgebras \mathcal{G}, \mathcal{H} , namely the inclusions in the span (3.16) are weak homomorphisms as defined in (4.1), then $\mathcal{K} \sim D(\mathcal{G})$ are equivalent as 2-bialgebras.*

Recall two weak 2-bialgebras are equivalent when there exists an invertible weak 2-homomorphism (4.1) between them.

Proof. The fact that \mathcal{K} factors into skeletal 2-subalgebras means that it must also be skeletal itself. This allows us to leverage the proof of **Theorem 3.1** to reconstruct the underlying 2-bialgebra structure of $\mathcal{K} \cong D(\mathcal{G})$ as a 2-quantum double.

The subtlety here is that we must now keep track of the 3-cocycle $\mathcal{T}_K : \mathcal{K}_0^{3\otimes} \rightarrow \mathcal{K}_{-1}$ in \mathcal{K} when we, in particular, invoke associativity in the form

$$\begin{aligned}\mathcal{T}_K(\iota_0(x), \iota_0(x'), j_0(f)) &\equiv \triangleright_1^{x, x'}(f) : (\iota_0(x) \hat{\smile} \iota_0(x')) \hat{\smile} j_0(f) \xrightarrow{\sim} \iota_0(x) \hat{\smile} (\iota_0(x') \hat{\smile} j_0(f)), \\ \mathcal{T}_K(\iota_0(x), j_0(f), j_0(f')) &\equiv \triangleleft_1^{f, f'} : (\iota_0(x) \hat{\smile} j_0(f)) \hat{\smile} j_0(f') \xrightarrow{\sim} \iota_0(x) \hat{\smile} (j_0(f) \hat{\smile} j_0(f')).\end{aligned}$$

Now in the skeletal case, the braiding map $\Psi = (\Psi_0, \Psi_{-1}; \bar{\Psi}) : \mathcal{G} \otimes \mathcal{H} \rightarrow \mathcal{H} \otimes \mathcal{G}$ is still defined as in (3.17). However, the components $\triangleright_1, \triangleleft_1$ now give rise to associators

$$\begin{aligned}\triangleright_1 &: \Psi \circ (\hat{\cdot} \otimes \text{id}) \xrightarrow{\sim} (\text{id} \hat{\otimes} \hat{\cdot}) \circ \Psi_{12} \circ \Psi_{23}, \\ \triangleleft_1 &: \Psi \circ (\text{id} \hat{\otimes} \hat{\cdot}) \xrightarrow{\sim} (\hat{\cdot} \otimes \text{id}) \circ \Psi_{23} \circ \Psi_{12}\end{aligned}\tag{5.4}$$

that implement the braiding relations (3.19). These braiding associators satisfy a set of algebraic conditions following from the 3-cocycle condition (5.2) for \mathcal{T}_K .

With the components $\triangleright_1, \triangleleft_1$ as defined above, we now wish to reconstruct the Hochschild 3-cocycles $\mathcal{T}_G, \mathcal{T}_H$ of \mathcal{G}, \mathcal{H} from \mathcal{T}_K . Note this cannot be achieved by just restricting \mathcal{T}_K via the span (3.16), as this does not have the desired codomains. For instance, the restriction $\mathcal{T}_K|_{\text{im } \iota = \mathcal{G}} : G_0^{3\otimes} \rightarrow \mathcal{K}_{-1} \cong \mathcal{G}_{-1} \otimes \mathcal{H}_{-1}$ in general lands in the tensor product, for which only the \mathcal{G}_{-1} -valued component gives the desired 3-cocycle \mathcal{T}_G on \mathcal{G} . Nevertheless, with $\mathcal{T}_G, \mathcal{T}_H$ defined in this way, having the span (3.16) means that the 3-cocycle condition for \mathcal{T}_K implies $(\mathcal{G}, \mathcal{T}_G), (\mathcal{H}, \mathcal{T}_H)$ form a matched pair of weak 2-bialgebras, as in (5.2).

The "undesirable" piece $\tilde{\mathcal{T}}_G$, namely the component of $\mathcal{T}_K|_{\mathcal{G}}$ valued in \mathcal{H}_{-1} , is a Hochschild coboundary. This follows from the definition of the inclusion $\iota = (\iota_{-1}, \iota_0, \iota_1) : \mathcal{G} \hookrightarrow \mathcal{K}$ as a weak homomorphism. Indeed, by projecting the last of (4.1) for ι_1 to \mathcal{H} , the first term $\iota_{-1}(\mathcal{T}_G(x, x', x''))|_{\mathcal{H}} = 0$ vanishes whence

$$\begin{aligned}\tilde{\mathcal{T}}_G &\equiv \mathcal{T}_K(\iota_0(x), \iota_0(x'), \iota_0(x''))|_{\mathcal{H}} \\ &= \iota_0(x) \hat{\smile} \iota_1(x', x'')|_{\mathcal{H}} - \iota_1(x x', x'')|_{\mathcal{H}} + \iota_1(x, x' x'')|_{\mathcal{H}} - \iota_1(x, x') \hat{\smile} \iota_0(x'') \\ &= d_{HH}[\iota_1|_{\mathcal{H}}](x, x', x''),\end{aligned}$$

where d_{HH} is the Hochschild differential [40]. Similar arguments show that $\tilde{\mathcal{T}}_H = d_{HH}[j_1|_{\mathcal{G}}]$ is a Hochschild coboundary as well. This establishes the weak equivalence $\mathcal{K} \sim D(\mathcal{G})$.

The same argument as above, but dualized, is applied to reconstruct $(\Delta_G)_1$ and $(\Delta_H)_1$ from the coassociator $(\Delta_K)_1$. The coassociator conditions (4.3)-(4.5), as well as (4.7), for them follow from those for $(\Delta_K)_1$. \square

Note the coadjoint actions $\triangleright, \triangleleft$ only define genuine algebra representations when $\mathcal{T}, \mathcal{T}^* = 0$ (as in **Theorem 3.1**), or when $t, t^* = 0$. Without skeletality, the braiding transposition Ψ is no longer of the form given in (3.17). Terms like $\triangleright_1^{t, \cdot}, \triangleleft_1^{\cdot, t^*}$ must now appear. This, of course, would modify (3.19) in a complicated and intricate manner.

Remark 5.1. If the components ι_1, j_1 are not required as part of the data for the inclusions ι, j in the span (3.16), then $\mathcal{K} \not\sim D(\mathcal{G})$ in general. In particular, without the component ι_1 trivializing $\tilde{\mathcal{T}}_G$ by (4.1), its (possibly non-trivial) Hochschild class $[\tilde{\mathcal{T}}_G] \in HH^3(\mathcal{K}_0, \mathcal{K}_{-1})$ is in fact an *extra* piece of data in \mathcal{K} that is not in $D(\mathcal{G})$, despite them sharing the same 2-bialgebra structure. Such a factorizable weak 2-bialgebra is still weakly self-dual $\mathcal{K} \sim \mathcal{K}^*$.

In the following, we shall shift gears a bit and study the 2-representation theory of quasitriangular 2-bialgebras, and finally work our way towards proving the main theorem.

6 The fusion 2-category of 2-representations

With the above algebraic machinery in place, we are now ready to discuss the 2-representations of a strict or weak 2-bialgebra \mathcal{G} . In the following, we shall follow the Baez-Crans definition of a 2-vector space and the monoidal 2-category $2\text{Vect}_{\text{int}}$ they form [33, 61].

Definition 6.1. A 2-vector space is a 2-term cochain complex of vector spaces; equivalently, a 2-vector space is a *nuclear* 2-algebra [40], or an Abelian Lie 2-algebra [41, 42].

2-vector spaces of this type form a 2-category $2\text{Vect}_{\text{int}}$ in which the 1-morphisms are cochain maps and 2-morphisms are cochain homotopies. Concretely, let $V = V_{-1} \xrightarrow{\partial} V_0, W = W_{-1} \xrightarrow{\partial'} W_0$ denote two 2-vector spaces. A cochain map $f : V \rightarrow W$ is a collection linear maps $f_{0, -1} : V_{0, -1} \rightarrow W_{0, -1}$ such that

$$\partial' f_{-1} = f_0 \partial.$$

Given two such cochain maps f, g , a cochain homotopy $q : f \Rightarrow g$ is a linear map $q : V_0 \rightarrow W_{-1}$ such that

$$\partial q = f_0 - g_0, \quad q\partial = f_{-1} - g_{-1}.$$

We shall refine these notions to fit the definition of a 2-representation of \mathcal{G} in the following.

6.1 Weak 2-representations

Recall that a representation of an ordinary algebra A on the vector space V is an algebra homomorphism $A \rightarrow \text{End}(V)$. Morally, a 2-representation should therefore be a 2-algebra homomorphism between a 2-algebra \mathcal{G} and a "categorified" notion of the endomorphism algebra $\text{End}(V)$. Correspondingly, a weak 2-representation should be a weak 2-homomorphism as in (4.1) into a "weak endomorphism 2-algebra".

6.1.1 Endomorphism 2-algebra on a 2-vector space

In the strict case, the endomorphisms of a 2-vector space are naturally given in the setting of $2\text{Vect}_{\text{int}}$ — namely $\text{End}(V) = \text{End}_{2\text{Vect}_{\text{int}}}(V)$, which forms an associative 2-algebra $\text{End}(V) = \text{End}(V)_{-1} \xrightarrow{\delta} \text{End}(V)_0$ of linear transformations on a 2-term cochain complex V [45],

$$\begin{aligned} \text{End}(V)_0 &= \{(M, N) \in \text{End}(V_{-1}) \times \text{End}(V_0) \mid \partial M = N\partial\}, \\ \text{End}(V)_{-1} &= \{A \in \text{Hom}(V_0, V_{-1}) \mid (A\partial, \partial A) \in \text{End}(V_{-1}) \times \text{End}(V_0)\}, \end{aligned}$$

equipped with the 2-algebra structure (take $A \in \text{End}(V)_{-1}$, $(M, N) \in \text{End}(V)_0$)

$$\delta : A \mapsto (A\partial, \partial A), \quad (M, N) \cdot A = MA, \quad A \cdot (M, N) = AN.$$

The associativity of matrix multiplication implies that $\text{End}(V)_{-1}$ is clearly a $\text{End}(V)_0$ -bimodule. Moreover, we have the Peiffer conditions (note $A, A' \in \text{End}(V)_{-1}$)

$$\begin{aligned} \delta((M, N) \cdot A) &= (MA\partial, \partial MA) = (MA\partial, N\partial A) = (M, N)\delta(A), \\ \delta(A \cdot (M, N)) &= (AN\partial, \partial AN) = (A\partial M, \partial AN) = \delta(A)(M, N), \\ A * A' &\equiv \delta(A) \cdot A' = A\partial A' = A \cdot \delta(A'), \end{aligned}$$

and hence $\text{End}(V)$ is an associative 2-algebra. Note that none of the matrices here are required to be invertible.

As weak 2-algebras are no longer associative, the above presentation of $\text{End}(V)$ in terms of matrices is no longer sufficient: we require a weaker version of $\text{End}(V)$. Such a notion of the weak endomorphism 2-algebra $\mathfrak{End}(V)$ would still have the same graded structure $\delta : \mathfrak{End}(V)_{-1} \rightarrow \mathfrak{End}(V)_0$ as in the strict case above, but its algebra structure should have its associativity controlled by a Hochschild 3-cocycle \mathfrak{T} , in accordance with **Definition 4.1**.

To begin, we extend the idea of [62] to weak 2-algebras. In essence, we leverage the observation in the strict case that an algebra 2-homomorphism $\mathcal{G} \rightarrow \mathfrak{End}(V)$ is equivalent to a \mathcal{G} -bimodule structure on V . We are going to provide a weak generalization of such a \mathcal{G} -bimodule structure in **Definition 6.2**.

Let 2Alg denote the category of weak 2-algebras $(\mathcal{G}, \mathcal{T})$, which contains the full subcategory $2\text{Alg}_{\text{gass}}$ of *strict* 2-algebras. A 2-vector space $V \in 2\text{Vect} \subset 2\text{Alg}_{\text{gass}} \subset 2\text{Alg}$ fits as a strict 2-algebra with trivial multiplication. We consider \mathcal{G} a weak 2-algebra (as defined in **Definition 4.1**). We then equip the direct sum $\mathcal{G} \oplus V$ with a semidirect product structure,

$$\begin{aligned} (z + u) \cdot (z' + u') &= yy' + x \cdot_l y' + y \cdot_r x' + xx' \\ &\quad + x \triangleright w' + x \triangleright v' + y \succ w' + y \succ v' \\ &\quad + w \triangleleft x' + v \triangleleft x' + w \prec y' + v \prec y', \end{aligned}$$

where we have used the shorthand notation $z = (y, x) \in \mathcal{G}_{-1} \times \mathcal{G}_0 = \mathcal{G}$, $u = (w, v) \in V_{-1} \times V_0 = V$ and where

$$\begin{aligned} \cdot_l : \mathcal{G}_0 \otimes \mathcal{G}_{-1} &\rightarrow \mathcal{G}_{-1}, & \cdot_r : \mathcal{G}_{-1} \otimes \mathcal{G}_0 &\rightarrow \mathcal{G}_{-1}, \\ \triangleright : \mathcal{G}_0 \otimes V &\rightarrow V, & \triangleleft : V \otimes \mathcal{G}_0 &\rightarrow V, \\ \succ : \mathcal{G}_{-1} \otimes V &\rightarrow V, & \prec : V \otimes \mathcal{G}_{-1} &\rightarrow V \end{aligned}$$

are all bilinear maps.

Definition 6.2. We say that V is a \mathcal{G} -bimodule if $(\mathcal{G} \oplus V, \cdot) \in 2\text{Alg}$ is a weak 2-algebra. In other words,

- (i) $(\mathcal{G} \oplus V)_{-1} \equiv \mathcal{G}_{-1} \oplus V_{-1}$ is a weak $(\mathcal{G} \oplus V)_0 := \mathcal{G}_0 \oplus V_0$ -bimodule,

- (ii) the map $t \oplus \partial : \mathcal{G}_{-1} \oplus V_{-1} \rightarrow \mathcal{G}_0 \oplus V_0$ is equivariant with respect to \cdot and satisfies the Peiffer identity⁴,
- (iii) there exists a well-defined trilinear invertible map $(\mathcal{G}_0 \oplus V_0)^{3\otimes} \rightarrow \mathcal{G}_{-1} \oplus V_{-1}$ that satisfies the Hochschild 3-cocycle condition.

To proceed, we will make the crucial assumption that there exists a "universal functor" $\mathfrak{E}\mathfrak{n}\mathfrak{d} : 2\mathfrak{Vect} \rightarrow 2\mathfrak{Alg}$ that characterizes \mathcal{G} -bimodules: a 2-homomorphism $\mathcal{G} \rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ is equivalent to a \mathcal{G} -bimodule structure on V . We call $\mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ the **weak endomorphism 2-algebra** on V , and denote by $\mathfrak{T} : \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)_0^{3\otimes} \rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)_{-1}$ its Hochschild 3-cocycle obtained from the third point of **Definition 6.2**. This motivates our following theory of weak 2-representations.

Remark 6.1. We emphasize that $\mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ is no longer part of the theory of Baez-Crans 2-vector spaces, as the 2-category of Baez-Crans 2-vector spaces is completely strict [33, 63] and hence $\text{End}(V) = \text{End}_{2\mathfrak{Vect}_{\text{int}}}(V)$ cannot carry any non-trivial Hochschild 3-cocycle \mathfrak{T} . In order to not detract from our goal of proving the main **Theorem 1.1**, we take the existence of the object $\mathfrak{E}\mathfrak{n}\mathfrak{d}(V) = \text{End}_{2\mathfrak{Vect}}(V)$ as granted. A rigorous formulation of the setting $2\mathfrak{Vect}$ of "weak 2-vector spaces" is currently under investigation by one of the authors. In Appendix C, we shall establish properties of weak 2-representations and relate it to those in the literature [48, 32, 31, 30].

6.1.2 Weak 2-representations, weak 2-intertwiners and modifications

Definition 6.3. A **weak 2-representation** $(\varrho, \rho) : \mathcal{G} \rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ of \mathcal{G} on V is a homomorphism between weak 2-algebras as in (4.1). In other words, $\rho = (\rho_0, \rho_1)$ is a chain map

$$\begin{array}{ccc} \mathcal{G}_{-1} & \xrightarrow{t} & \mathcal{G}_0 \\ \downarrow \rho_1 & & \downarrow \rho_0 \\ \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)_{-1} & \xrightarrow{\delta} & \mathfrak{E}\mathfrak{n}\mathfrak{d}(V)_0 \end{array} \quad (6.1)$$

which preserves the 2-algebra structures *up to homotopy*,

$$\begin{aligned} \delta\varrho(x, x') &= \rho_0(xx') - \rho_0(x)\rho_0(x'), \\ \varrho(x, ty) &= \rho_1(x \cdot y) - \rho_0(x) \cdot \rho_1(y), \\ \varrho(ty, x) &= \rho_1(y \cdot x) - \rho_1(y) \cdot \rho_0(x), \end{aligned} \quad (6.2)$$

and for which the Hochschild 3-cocycles $\mathcal{T}, \mathfrak{T}$ of respectively \mathcal{G} and $\mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ satisfy the following compatibility conditions

$$\begin{aligned} \rho_1(\mathcal{T}(x, x', x'')) - \mathfrak{T}(\rho_0(x), \rho_0(x'), \rho_0(x'')) &= \rho_0(x) \cdot \varrho(x', x'') - \varrho(xx', x'') \\ &\quad + \varrho(x, x'x'') - \varrho(x, x') \cdot \rho_0(x''), \end{aligned} \quad (6.3)$$

where $x, x', x'' \in \mathcal{G}_0$ and $y \in \mathcal{G}_{-1}$. We require ϱ to be invertible.

We call ρ a **strict 2-representation** if $\varrho = 0$ identically.

As $\mathcal{T}, \mathfrak{T}$ are normalized, ϱ by definition vanishes if any of its arguments are 0 or the unit $\eta_0 \in \mathcal{G}_0$.

Remark 6.2. Due to the classification **Theorem 2.2** of 2-algebras [40], a non-trivial 2-algebra \mathcal{G} with $\mathcal{T} \neq 0$ cannot admit a strict 2-representation. Conversely, however, 2-representations of a strict 2-algebra can still be weak, as (6.3) only states that the *cohomology class* of \mathfrak{T} is trivial, not that it is trivial as a 3-cocycle. However, if we further restrict to the case where V is a strict \mathcal{G} -bimodule (ie. the trilinear map in **Definition 6.2** vanishes), then $\mathfrak{T} = 0$ and $\mathfrak{E}\mathfrak{n}\mathfrak{d}(V)$ is isomorphic to $\text{End}(V)$.

Example: weak coadjoint representation. A very natural example of a 2-representation is achieved by dualizing, using (3.1), the 2-representation $\mathcal{G} \rightarrow \text{End}(\mathcal{G})$ given by the weak 2-algebra structure of \mathcal{G} on itself.

This gives rise to the *coadjoint representation* (cf. [41, 3]) $\bar{\triangleright} = (\triangleright_1, (\triangleright_0, \triangleright_{-1}), \Upsilon) : \mathcal{G} \rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(\mathcal{G}^*)$ of \mathcal{G} on its dual \mathcal{G}^* , given explicitly by

$$\begin{aligned} \triangleright_0 : \mathcal{G}_0 &\rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(\mathcal{G}_0^*), & \langle g, xx' \rangle &= -\langle x \triangleright_0 g, x' \rangle, \\ \triangleright_{-1} : \mathcal{G}_0 &\rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(\mathcal{G}_{-1}^*), & \langle f, x \cdot y \rangle &= -\langle x \triangleright_{-1} f, y \rangle, \\ \Upsilon : \mathcal{G}_{-1} &\rightarrow \text{Hom}(\mathcal{G}_{-1}^*, \mathcal{G}_0^*), & \langle f, y \cdot x \rangle &= -\langle \Upsilon_y f, x \rangle \end{aligned} \quad (6.4)$$

and

$$\triangleright_1 : \mathcal{G}_0^{2\otimes} \rightarrow \mathfrak{E}\mathfrak{n}\mathfrak{d}(\mathcal{G}^*)_{-1} = \text{Hom}(\mathcal{G}_{-1}^*, \mathcal{G}_0^*), \quad \langle f, \mathcal{T}(x, x', x'') \rangle = +\langle \triangleright_1^{x, x'}(f), x'' \rangle. \quad (6.5)$$

⁴The Peiffer identity states $y \succ w = (ty) \triangleright w = y \succ (\partial w)$, and similarly $w \prec y = (\partial w) \prec y = w \prec (ty)$. If we write $y \succ v = \Upsilon_y v$, then we reproduce precisely the 2-representation properties (3.9).

Notice a plus sign occurs here, in contrast with the rest of the components defined in (6.4). This is because we have dualized two elements in \mathcal{G} , instead of one.

Analogously, we have the coadjoint back-action $\bar{\triangleleft} = ((\triangleleft_0, \triangleleft_{-1}), \tilde{\Upsilon})$ of \mathcal{G}^* on \mathcal{G} , which we write from the right⁵. The "bar" notation is used to distinguish $\bar{\triangleleft}$ from the crossed-module action \triangleright in the case where $\mathcal{G} = kG$ is the 2-group algebra defined in (2.14).

Due to (6.2), the components of a weak 2-representation are not genuine algebra representations in general, but only up to homotopy. We have in general that

$$(xx') \triangleright_0 g = x \triangleright_0 (x' \triangleright_0 g) + \triangleright_1^{x,x'}(t^*g), \quad (xx') \triangleright_{-1} f = x \triangleright_{-1} (x' \triangleright_{-1} f) + t^* \triangleright_1^{x,x'}(f),$$

where t^* is the dual t -map on \mathcal{G}^* , and

$$\Upsilon_{x \cdot y} f = x \triangleright_0 (\Upsilon_y f) + \triangleright_1^{x,ty}(f), \quad \Upsilon_{y \cdot x} f = \Upsilon_y(x \triangleright_{-1} f) + \triangleright_1^{ty,x}(f).$$

Of course, these components reduce to genuine strict algebra representations if $\triangleright_1 = 0$ or $t = 0$, which simplifies the situation considerably.

Remark 6.3. Recall the 2-algebra $\mathcal{G} = kG$ associated to the 2-group G in Example 2.1.2 In the skeletal case, we can induce a 2-representation ρ of the 2-algebra from that of the 2-group, by extending it linearly. All 2-representations of kG shall arise this way, in this case.

1- and 2-morphisms on the weak 2-representation 2-category. With Definition 6.3 in hand, we are now ready to define the morphisms on the weak 2-representations. Let $\rho = (\varrho, \rho_0, \rho_1)$ and $\rho' = (\varrho', \rho'_0, \rho'_1)$ denote two weak 2-representations on $V, W \in 2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, respectively.

Definition 6.4. A **weak 2-intertwiner** $i = (I, i_1, i_0) : V \rightarrow W$ consist of a 2-vector space homomorphism $(i_1, i_0) : V \rightarrow W$ together with a collection of invertible cochain homotopies $I_{x,i} : V_0 \rightarrow W_{-1}$ satisfying

$$\partial I_{x,i} = i_0 \circ \rho_0^0(x) - \rho_0^0(x) \circ i_0, \quad I_{x,i} \partial = i_1 \circ \rho_0^1(x) - \rho_0^1(x) \circ i_1$$

for each $x \in \mathcal{G}_0$, as well as

$$I_{ty,i} = i_1 \circ \rho_1(y) - \rho_1(y) \circ i_0$$

for each $y \in \mathcal{G}_{-1}$. Moreover, $I_{\bullet,i}$ trivializes $\varrho - \varrho'$ as a Hochschild 2-cocycle, in the sense that for each $x, x' \in \mathcal{G}_0$,

$$\text{id}_i \otimes \varrho(x, x') - \varrho'(x, x') \otimes \text{id}_i = \text{id}_{\rho_0(x)} \otimes I_{x',i} - I_{xx',i} + I_{x,i} \otimes \text{id}_{\rho_0(x')}, \quad (6.6)$$

where $\text{id}_i : i \Rightarrow i$ denotes the identity cochain homotopy on the intertwiner i .

In other words, a weak 2-intertwiner $i : V \rightarrow W$ is such that the following diagrams

$$\begin{array}{ccc} V_{-1} & \xrightarrow{\partial} & V_0 \\ \downarrow \rho_0^1 & \searrow i_1 & \downarrow i_0 \\ & W_{-1} & \xrightarrow{\rho_0^0} W_0 \\ \downarrow \rho_0^1 & \searrow i_1 & \downarrow i_0 \\ V_{-1} & \xrightarrow{\partial} & V_0 \\ & W_{-1} & \xrightarrow{\rho_0^1} W_0 \end{array}, \quad \begin{array}{ccc} V_0 & \xrightarrow{\rho_1} & V_{-1} \\ \downarrow i_0 & & \downarrow i_1 \\ W_0 & \xrightarrow{\rho'_1} & W_{-1} \end{array}, \quad (6.7)$$

commute up to a natural invertible 2-morphism given by $I_{\bullet,i}$. By definition, we have $I_{0,i} = I_{\eta_0,i} = 0$ where η_0 is the unit of \mathcal{G}_0 .

Now let $i, i' : \rho \rightarrow \rho'$ denote two weak 2-intertwiners, we have the following.

Definition 6.5. A **modification** $\mu : i \Rightarrow i'$ between two weak 2-intertwiners is a \mathcal{G} -equivariant cochain homotopy

$$\begin{array}{ccc} V_{-1} & \xrightarrow{\partial} & V_0 \\ i_1 - i'_1 \downarrow & \mu & \downarrow i_0 - i'_0 \\ W_{-1} & \xrightarrow{\partial'} & W_0 \end{array}, \quad (6.8)$$

where μ intertwines between $\rho_1(y), \rho'_1(y)$ for each $y \in \mathcal{G}_{-1}$, as cochain homotopies. Moreover, μ trivializes $I_{\bullet,i} - I_{\bullet,i'}$ as a Hochschild 1-cocycle, in the sense that

$$I_{x,i} - I_{x,i'} = \text{id}_{\rho_0(x)} \otimes \mu - \mu \quad (6.9)$$

for all $x \in \mathcal{G}_0$, as a relation between cochain homotopies.

⁵This means that we have, for instance, $\langle g \cdot * f, x \rangle = -\langle g, x \triangleleft_{-1} f \rangle$ and $\langle f \cdot * g, x \rangle = -\langle f, x \tilde{\Upsilon} g \rangle$.

We shall denote by $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ the 2-category of weak 2-representations of the weak 2-bialgebra $(\mathcal{G}, \mathcal{T})$, consisting of weak 2-representation (V, ρ) objects, weak 2-intertwiners i as 1-morphisms and modifications μ as 2-morphisms. We will prove in Appendix C that our definition in fact coincides with the higher-representation theory developed in the literature [64, 31, 48, 32], in the case where $\mathcal{G} = kG$ corresponding to a skeletal 2-group G . We devote the remainder of this section to proving that $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ forms a monoidal (in fact, fusion) 2-category.

6.2 Fusion structure on the 2-representations

Recall that vector space cochain complexes come equipped with natural notions of direct sum \oplus , as well as tensor product \otimes , which satisfy the distributive law

$$V \otimes (W \oplus U) = (V \otimes W) \oplus (V \otimes U),$$

where V, W, U are vector space cochains. For 2-vector spaces (or equivalently two-term cochain complexes of vector spaces [33]), the direct sum is given simply by

$$V \oplus W = V_{-1} \oplus W_{-1} \xrightarrow{\partial \oplus \partial'} V_0 \oplus W_0,$$

while the tensor product is given by the following 3-term complex (cf. [41])

$$V \otimes W = \underbrace{V_{-1} \otimes W_{-1}}_{\text{deg}=-2} \xrightarrow{D^+} \underbrace{V_{-1} \otimes W_0 \oplus V_0 \otimes W_{-1}}_{\text{deg}=-1} \xrightarrow{D^-} \underbrace{V_0 \otimes W_0}_{\text{deg}=0}, \quad (6.10)$$

where $D^{\pm} = \pm 1 \otimes \partial' + \partial \otimes 1$ is the tensor extension of the differentials $\partial : V_{-1} \rightarrow V$ and $\partial' : W_{-1} \rightarrow W_0$.

We endow the direct sum and tensor product structure on 2-representations of \mathcal{G} in the same way as above. Note the direct double $\mathcal{G}^{2\oplus}$ and the tensor square $\mathcal{G}^{2\otimes}$ of a strict 2-algebra \mathcal{G} also have the same structure.

Direct sums. For the direct sum 2-representation, this is simply accomplished by extending **Definition 6.3** to a direct sum of 2-algebra homomorphisms

$$(\varrho, \rho) \oplus (\varrho', \rho') = (\varrho \oplus \varrho', \rho \oplus \rho') : \mathcal{G} \oplus \mathcal{G} \rightarrow \mathfrak{End}(V) \oplus \mathfrak{End}(W).$$

In particular, the direct sum $V \oplus W$ of 2-representations of \mathcal{G} is given by the components

$$(\rho \oplus \rho')_0^0 = \rho_0^0 \oplus \rho_0^0, \quad (\rho \oplus \rho')_0^1 = \rho_0^1 \oplus \rho_0^1, \quad (\rho \oplus \rho')_1 = \rho_1 \oplus \rho_1'$$

such that the square (6.1) commutes,

$$(\rho \oplus \rho')_0 \circ (t \oplus t) = (\delta \oplus \delta') \circ (\rho \oplus \rho')_1,$$

where δ, δ' are the differentials of the two 2-algebras $\mathfrak{End}(V), \mathfrak{End}(W)$, respectively. The zero 2-representation under direct sum is of course the trivial complex $0 \rightarrow 0$.

6.2.1 Monoidal structure

As in the 1-bialgebra case, the tensor product of 2-representations is accomplished by precomposing with the coproduct. However, the graded components of the coproduct $\Delta = \Delta_{-1} + \Delta_0$ in (2.16), as well as Δ'_0 in (2.19), allows us to define the tensor product between 2-representations $V \otimes W$

$$\rho_{V \otimes W}(x) = ((\rho_V)_0 \otimes (\rho_W)_0) \circ \Delta'_0(x), \quad x \in \mathcal{G}_0, \quad (6.11)$$

as well as its weak component (cf. **Definition 6.3**)

$$\varrho_{V \otimes W}(x, x') = \varrho_V(\bar{x}_{(1)}, \bar{x}'_{(1)}) \otimes \varrho_W(\bar{x}_{(2)}, \bar{x}'_{(2)}), \quad x, x' \in \mathcal{G}_0.$$

We also have the tensor product between a 2-intertwiner $i : V \rightarrow U$ and a 2-representation

$$\begin{aligned} \rho_{i \otimes W}(x) &= ((\rho_U)_1 \circ i \otimes (\rho_W)_0) \circ \Delta_0^l(x) + (-1)^{\text{deg}}(i \circ (\rho_V)_0 \otimes (\rho_W)_1) \circ \Delta_0^r(x), \\ \rho_{W \otimes i}(x) &= ((\rho_W)_0 \otimes (\rho_U)_1 \circ i) \circ \Delta_0^r(x) + (-1)^{\text{deg}}((\rho_W)_1 \otimes i \circ (\rho_V)_0) \circ \Delta_0^l(x) \end{aligned} \quad (6.12)$$

for each $x \in \mathcal{G}_0$, where the sign depends on the degree of the components in (6.10). Lastly, the tensor product between 2-intertwiners $i : V \rightarrow U, j : W \rightarrow T$ is given by just

$$\rho_{i \otimes j}(y) = ((\rho_U)_1 \circ i \otimes (\rho_T)_1 \circ j + (-1)^{\text{deg}} i \circ (\rho_V)_1 \otimes j \circ (\rho_W)_1) \circ \Delta_{-1}(y) \quad (6.13)$$

for each $y \in \mathcal{G}_{-1}$. This defines the invertible natural 2-morphism $I_{i \otimes j, \bullet}$ (cf. **Definition 6.4**).

The fact that (6.11), (6.12), (6.13) define genuine 2-representations (up to the homotopy ϱ ; cf. **Definition 6.3** and (6.2)), for instance

$$\delta_{\varrho_{V \otimes W}}(x, x') = \rho_{V \otimes W}(xx') - \rho_{V \otimes W}(x)\rho_{V \otimes W}(x'),$$

requires the 2-bialgebra axioms (2.31).

Tensor unit. Now if \mathcal{G} is a unital 2-bialgebra, then there is a tensor unit, denoted by $I \in 2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ given by the ground field complex $k \xrightarrow{1} k$, and a unit 2-intertwiner given by the identity $\text{id}_I : 1 \rightarrow 1$, such that \mathcal{G} acts on them through multiplication of the counit ϵ ,

$$\rho_I(x) = \epsilon_0(x), \quad \rho_{\text{id}_I}(y) = \epsilon_{-1}(y).$$

From (6.2), the corresponding component $\varrho = \text{id}$ for the tensor unit I is clearly the identity 2-morphism. In according with (6.11), (6.12), (6.13), the condition (2.26) then implies that the left- and right-unitor morphisms in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ are all 1- and 2-isomorphisms. For instance, (2.28) implies

$$\rho_{V \otimes 1} = \rho_V = \rho_{1 \otimes V},$$

whence $V \otimes 1, 1 \otimes V$ and V coincides as 2-representations.

Due to this, all coherence diagrams in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ concerning the unitors, such as the homotopy triangle and the zig-zag axioms [22, 46], are trivially satisfied, and hence we will not directly prove them. The conditions (2.26), (2.28) can of course be easily relaxed to give non-trivial coherent unitors, but we shall not consider this here.

6.2.2 Naturality and Gray property of the tensor product

Recall the space $\mathbf{End}(V)_{-1}$ is modelled by cochain homotopies, which can be interpreted as "endomorphisms" on $\mathbf{End}(V)_0$. Using this perspective, we will prove the following key results.

Lemma 6.1. *Let $i : V \rightarrow U$ denote a 2-intertwiner. We have the following diagrams*

$$\begin{array}{ccc} V \otimes W & \xrightarrow{\rho_{V \otimes W}} & V \otimes W \\ \downarrow i & \xrightarrow{\rho_{i \otimes W}} & \downarrow i \\ U \otimes W & \xrightarrow{\rho_{U \otimes W}} & U \otimes W \end{array}, \quad \begin{array}{ccc} W \otimes V & \xrightarrow{\rho_{W \otimes V}} & W \otimes V \\ \downarrow i & \xrightarrow{\rho_{W \otimes i}} & \downarrow i \\ W \otimes U & \xrightarrow{\rho_{W \otimes U}} & W \otimes U \end{array}$$

in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$.

Proof. Let us focus first on the left diagram. The goal is to show that $\rho_{i \otimes W}$ defines a cochain homotopy which fits into the following diagram

$$\begin{array}{ccccc} V_{-1} \otimes W_{-1} & \xrightarrow{D^+} & V_{-1} \otimes W_0 \oplus V_0 \otimes W_{-1} & \xrightarrow{D^-} & V_0 \otimes W_0 \\ \downarrow & \swarrow \rho_{i \otimes W} & \downarrow & \swarrow \rho_{i \otimes W} & \downarrow \\ U_{-1} \otimes W_{-1} & \xrightarrow{D^+} & U_{-1} \otimes W_0 \oplus U_0 \otimes W_{-1} & \xrightarrow{D^-} & U_0 \otimes W_0 \end{array},$$

where the horizontal maps are the differentials given in (6.10), and the vertical maps are various components of $\rho_{V \otimes W} \circ i - i \circ \rho_{V \otimes W}$.

The key is the commutation relation (6.1), which allows us to write

$$\delta(\rho_1(y)) = (\rho_1(y)\partial, \partial\rho_1(y)) = (\rho_0^1(ty), \rho_0^0(ty))$$

for each $y \in \mathcal{G}_{-1}$, as well as the definition (2.19) of Δ'_0 . Directly computing, we have for the rightmost triangle

$$\begin{aligned} D^- \rho_{i \otimes W} &= \partial_U(\rho_U)_1(x_{(1)}^l) \circ i \otimes (\rho_W)_0^0(x_{(2)}^l) - (-1)^{\deg i} i \circ (\rho_V)_0(x_{(1)}^r) \otimes \partial_W(\rho_W)_0^0(x_{(2)}^r) \\ &= (\rho_U)_0^0(tx_{(1)}^l) \circ i \otimes (\rho_W)_0^0(x_{(2)}^l) - i \circ (\rho_V)_0^0(x_{(1)}^r) \otimes (\rho_W)_0^0(tx_{(2)}^r) \\ &= \rho_{U \otimes W} \circ i - i \circ \rho_{V \otimes W} \end{aligned}$$

as maps on $V_0 \otimes W_0$ (with $\deg = 0$), and similarly we have for the leftmost triangle

$$\begin{aligned}\rho_{i \otimes W} D^+ &= (\rho_U)_1(x_{(1)}^l) \partial_U \circ i \otimes (\rho_W)_0^1(x_{(2)}^l) + (-1)^{\deg i} \circ (\rho_V)_0(x_{(1)}^r) \otimes (\rho_W)_0^0(x_{(2)}^r) \partial_W \\ &= (\rho_U)_0^1(tx_{(1)}^l) \circ i \otimes (\rho_W)_0^1(x_{(2)}^l) - i \circ (\rho_V)_0^1(x_{(1)}^r) \otimes (\rho_W)_0^1(tx_{(2)}^r) \\ &= \rho_{U \otimes W} \circ i - i \circ \rho_{V \otimes W}\end{aligned}$$

as maps on $V_{-1} \otimes W_{-1}$ (with $\deg = -1$).

Now consider the middle section. We need to compute

$$\begin{aligned}D^+ \rho_{i \otimes W} &= (\rho_U)_0^1(tx_{(1)}^l) \circ i \otimes (\rho_W)_0^0(x_{(2)}^l) - i \circ (\rho_V)_0^0(x_{(1)}^r) \otimes (\rho_W)_0^1(tx_{(2)}^r), \\ \rho_{i \otimes W} D^- &= (\rho_U)_0^0(tx_{(1)}^l) \circ i \otimes (\rho_W)_0^1(x_{(2)}^l) - i \circ (\rho_V)_0^1(x_{(1)}^r) \otimes (\rho_W)_0^0(tx_{(2)}^r),\end{aligned}$$

and sum them to find

$$\begin{aligned}D^+ \rho_{i \otimes W} + \rho_{i \otimes W} D^- &= \left[(\rho_U)_0^1(tx_{(1)}^l) \otimes (\rho_W)_0^0(x_{(2)}^l) + (\rho_U)_0^0(tx_{(1)}^l) \otimes (\rho_W)_0^1(x_{(2)}^l) \right] \circ i \\ &\quad - i \circ \left[(\rho_V)_0^0(x_{(1)}^r) \otimes (\rho_W)_0^1(tx_{(2)}^r) + (\rho_V)_0^1(x_{(1)}^r) \otimes (\rho_W)_0^0(tx_{(2)}^r) \right] \\ &= \rho_{U \otimes W} \circ i - i \circ \rho_{V \otimes W}\end{aligned}$$

as maps on $V_{-1} \otimes W_0 \oplus V_0 \otimes W_{-1}$. The other diagram is treated identically. \square

We now show that (6.13) is in fact *not* independently defined.

Lemma 6.2. *If $j : W \rightarrow T$ is another 2-intertwiner, then $i \otimes j$ decomposes as $i \otimes j = i \otimes T \circ V \otimes j \cong U \otimes j \circ i \otimes W$. The homotopy $I_{i \otimes j, \bullet} = I_{i \otimes \text{id}_W} * I_{\text{id}_V \otimes j}$ also decomposes accordingly.*

Proof. What we need to show is that $\rho_{i \otimes j} = (\rho_{i \otimes T} * \rho_{V \otimes j}) \circ t = (\rho_{U \otimes j} * \rho_{i \otimes W}) \circ t$ as 2-morphisms. Recall cochain homotopies $q : f \Rightarrow g, p : g \Rightarrow h$ in 2Vect compose by $p * q = p \partial_U q : f \Rightarrow h$, where U is the source 2-vector space of the cochain map g . Indeed, we have

$$\begin{aligned}\partial_W(p * q) &= (\partial_W p) \circ (\partial_U q) = (g_0 - h_0) \circ (f_0 - g_0), \\ (p * q) \partial_V &= (p \partial_U) \circ (q \partial_V) = (g_{-1} - h_{-1}) \circ (f_{-1} - g_{-1})\end{aligned}$$

as desired, where W is the target of h and V is the source of f . Notice this is exactly how elements in $\mathfrak{Cnd}(V)_{-1}$ compose, $A * A' = A \delta A$.

The goal is to prove that $D^\pm \rho_{i \otimes j}(y)$ in fact decomposes as described above for each $y \in \mathcal{G}_{-1}$. This follows from the coequivariance condition (2.22). By direct computation, precomposing (6.12) yields (here we neglect the 2-vector space subscripts for brevity)

$$\begin{aligned}\rho_{i \otimes W} \circ t &= (\rho_1 i \otimes (\rho_0 t) + (-1)^{\deg i} (\rho_0 t) \otimes \rho_1) \circ \Delta_{-1} \\ &= (\rho_1 i \otimes \partial \rho_1 + (-1)^{\deg i} (\partial \rho_1) \otimes \rho_1) \circ \Delta_{-1}, \\ \rho_{U \otimes j} \circ t &= ((\rho_0 t) \otimes \rho_1 j + (-1)^{\deg \rho_1} \rho_1 \otimes i(\rho_0 t)) \circ \Delta_{-1} \\ &= ((\rho_1 \partial) \otimes \rho_1 j + (-1)^{\deg \rho_1} \rho_1 \otimes j(\rho_1 \partial)) \circ \Delta_{-1},\end{aligned}$$

where we have used (6.1) to commute the t -map past the 2-representations to the differential ∂ . Using the Sweeder notation (2.17) for Δ_{-1} , we compute their graded composition to be

$$\begin{aligned}(\rho_{U \otimes j})(ty) * (\rho_{i \otimes W})(ty) &= (\rho_1(y_{(1)}) \partial) \rho_1(y_{(1)}) i \otimes \rho_1(y_{(2)}) j(\partial \rho_1(y_{(2)})) \\ &\quad + (-1)^{\deg \rho_1(y_{(1)})} (\partial \rho_1(y_{(1)})) \otimes j(\rho_1(y_{(2)}) \partial) \rho_1(y_{(2)}) \\ &= (\rho_1(y_{(1)}) * \rho_1(y_{(1)})) i \otimes j(\rho_1(y_{(2)}) * \rho_1(y_{(2)})) \\ &\quad + (-1)^{\deg (\rho_1(y_{(1)}) * \rho_1(y_{(1)}))} (\rho_1(y_{(1)}) * \rho_1(y_{(1)})) i \otimes j(\rho_1(y_{(2)}) * \rho_1(y_{(2)})) \\ &= (\rho_1 i \otimes \rho_1 j + (-1)^{\deg i} \rho_1 \otimes j \rho_1) \circ \Delta_{-1}(y) = \rho_{i \otimes j}(y)\end{aligned}$$

as desired, where we have noted the property $i_{-1}(\rho_V)_1 = (\rho_U)_1 i_0$ of the 2-intertwiners i, j to permute them past the ρ 's. This proves that the 2-algebra homomorphisms $\rho_{i \otimes j} = \rho_{i \otimes T} * \rho_{V \otimes j}$ coincide. A similar argument shows that the 2-algebra homomorphisms $\rho_{i \otimes j} = \rho_{U \otimes j} * \rho_{i \otimes W}$ also coincide.

This is not sufficient to imply that $i \otimes T \circ V \otimes j = U \otimes j \circ i \otimes W$, however. Indeed, the weak component ϱ of the two decomposed 2-representations in general may differ. After some computations, one can show that we have

$$\begin{aligned}\varrho_{(i \otimes T) \circ (V \otimes j)} \circ \Delta_0(x) &= \varrho(tx_{(1)}^l, x_{(1)}^r) \otimes \varrho(x_{(2)}^l, tx_{(2)}^r) + (-1)^{\deg \varrho} \varrho(x_{(1)}^r, tx_{(1)}^l) \otimes \varrho(tx_{(2)}^r, x_{(2)}^l), \\ \varrho_{(U \otimes j) \circ (i \otimes W)} \circ \Delta_0(x) &= \varrho(x_{(1)}^r, tx_{(1)}^l) \otimes \varrho(tx_{(2)}^r, x_{(2)}^l) + (-1)^{\deg \varrho} \varrho(tx_{(1)}^l, x_{(1)}^r) \otimes \varrho(x_{(2)}^l, tx_{(2)}^r).\end{aligned}$$

The difference $\varrho_{(i \otimes T) \circ (V \otimes j)} * \varrho_{(U \otimes j) \circ (i \otimes W)}^{-1}$ between these 2-morphisms is what gives rise to the 2-isomorphism $i \otimes T \circ V \otimes j \cong U \otimes j \circ i \otimes W$. \square

The fact that the tensor product of 1-morphisms decompose into two 2-isomorphic "mixed" tensor products is a signature property of Gray-enriched categories [65, 29]. We call the property that "structures on the 1-morphisms are determined by the mixed structure, together with appropriate coherence 2-isomorphisms" the **Gray property**.

These lemmas are important, as its proof techniques will be used repeatedly in what follows.

6.3 Fusion associators

In this section, we shall focus on the *associator* morphisms attached to the 2-representations in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, as they play a direct role in the main theorem. Other subtleties that show up are studied in detail in Appendix C.

Recall from Section 6.2.1 that the tensor product on $2\text{Rep}(\mathcal{G})$ is given by the coproduct Δ . The associator morphisms a are therefore given by the coassociator $\Delta_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1}^{3\otimes}$ attached to the coproduct in \mathcal{G} , and *not* the Hochschild 3-cocycle \mathcal{T} . However, the data Δ_1, \mathcal{T} are dual to each other by **Proposition 4.1**, hence if \mathcal{G} is self-dual (like the weak (skeletal) 2-quantum double as we constructed in Section 5), they in fact constitute the same data. As such we shall denote the weak 2-representation 2-category by $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$. We shall neglect the tensor product notation \otimes in the following.

We begin by constructing the associator 2-morphism $a_{ijk} : (i \otimes j) \otimes k \Rightarrow i \otimes (j \otimes k)$ on the triple $i : V \rightarrow V', j : W \rightarrow W', k : U \rightarrow U'$ of 2-intertwiners. By (4.3) and the definition (4.3), we see that the following quantity

$$a_{ijk} = ((\rho_{V'})_1 \circ i \otimes (\rho_{W'})_1 \circ j \otimes (\rho_{U'})_1 \circ k + (-1)^{\deg i} i \circ (\rho_V)_1 \otimes j \circ (\rho_W)_1 \otimes k \circ (\rho_U)_1) \circ (\Delta_1 \circ t) \quad (6.14)$$

defines a cochain homotopy that fits into the following equation $\rho_{(ij)k} - \rho_{i(jk)} = a_{ijk}$, which induces a 2-morphism (also denoted by a_{ijk}) between the 2-intertwiners

$$a_{ijk} : (ij)k \Rightarrow i(jk).$$

Secondly, (4.3) implies that the following quantities based on $D_t \Delta_1$,

$$\begin{aligned} a_{Vjk} &= ((\rho_V)_0 \otimes (\rho_{W'})_1 \circ j \otimes (\rho_{U'})_1 \circ k + (-1)^{\deg(\rho_V)_0} (\rho_V)_0 \otimes j \circ (\rho_W)_1 \otimes k \circ (\rho_U)_1) \circ (t \otimes 1 \otimes 1) \Delta_1, \\ a_{iWk} &= ((\rho_{V'})_1 \circ i \otimes (\rho_W)_0 \otimes (\rho_{U'})_1 \circ k + (-1)^{\deg i} i \circ (\rho_V)_1 \otimes (\rho_W)_0 \otimes k \circ (\rho_U)_1) \circ (1 \otimes t \otimes 1) \Delta_1, \\ a_{ijU} &= ((\rho_{V'})_1 \circ i \otimes (\rho_{W'})_1 \circ j \otimes (\rho_U)_0 + (-1)^{\deg i} i \circ (\rho_V)_1 \otimes j \circ (\rho_W)_1 \otimes (\rho_U)_0) \circ (1 \otimes 1 \otimes t) \Delta_1, \end{aligned} \quad (6.15)$$

give rise to the associators for the following tensor products,

$$a_{Vjk} : (Vj)k \Rightarrow V(jk), \quad a_{iWk} : (iW)k \Rightarrow i(WK), \quad a_{ijU} : (ij)U \Rightarrow i(jU)$$

for the mixed tensor products defined by (6.12). Thirdly, (4.4) implies that the following quantities based in $D_t[2]\Delta_1$,

$$\begin{aligned} a_{VWk} &= ((\rho_V)_0 \otimes (\rho_W)_0 \otimes (\rho_{U'})_1 \circ k + (-1)^{\deg(\rho_V)_0} (\rho_V)_0 \otimes (\rho_W)_0 \otimes k \circ (\rho_U)_1) \circ (t \otimes t \otimes 1) \Delta_1, \\ a_{iWU} &= ((\rho_{V'})_1 \circ i \otimes (\rho_W)_0 \otimes (\rho_U)_0 + (-1)^{\deg i} i \circ (\rho_V)_1 \otimes (\rho_W)_0 \otimes (\rho_U)_0) \circ (1 \otimes t \otimes t) \Delta_1, \\ a_{VjU} &= ((\rho_V)_0 \otimes (\rho_{W'})_1 \circ j \otimes (\rho_U)_0 + (-1)^{\deg(\rho_V)_0} (\rho_V)_0 \otimes j \circ (\rho_W)_1 \otimes (\rho_U)_0) \circ (t \otimes 1 \otimes t) \Delta_1, \end{aligned} \quad (6.16)$$

serve as the associators

$$a_{VWk} : (VW)k \Rightarrow V(Wk), \quad a_{VjU} : (Vj)U \Rightarrow V(jU), \quad a_{iWU} : (iW) \Rightarrow i(WU),$$

Notice that these quantities we have defined so far are all cochain homotopies/2-morphisms in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, due to the appearance of ρ_1 in their tensor products.

Lastly, (4.5) allows us to define the associator *1-morphism*,

$$a_{VWU} = ((\rho_V)_0 \otimes (\rho_W)_0 \otimes (\rho_U)_0)(\Phi), \quad (6.17)$$

with $\Phi \equiv (t \otimes t \otimes t) \Delta_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_0^{3\otimes}$, which induces an invertible 1-morphism

$$a_{VWU} : (VW)U \rightarrow V(WU)$$

that intertwines between $\rho_{(V \otimes W) \otimes U}$ and $\rho_{V \otimes (W \otimes U)}$.

The adjoint associator 2-morphism a^\dagger is implemented by minus the corresponding cochain homotopy. For (6.17), however, the adjoint morphism a^\dagger_{VWU} is given by the inverse Φ^{-1} .

The pentagon relation and naturality of the associator. We now prove the following.

Lemma 6.3. *Suppose the 3-cocycle $\mathfrak{T} = 0$ is trivial for the moment. The pentagon relation for the associators arising from (6.14), (6.15), (6.16), (6.17) follows from the 2-coassociativity condition (4.5) for Δ_1 .*

Proof. Consider first (6.14). We precompose (4.5) with t and reconstruct the associators corresponding to each term according to the definition,

$$\begin{aligned} (\text{id} \otimes (\Delta_1 \circ t)) \circ \Delta_{-1} &\rightsquigarrow \text{id}_i \otimes a_{jkl}, & ((\Delta_1 \circ t) \otimes \text{id}) \circ \Delta_{-1} &\rightsquigarrow a_{ijk} \otimes \text{id}_l, & (1 \otimes \Delta_{-1} \otimes 1) \circ \Delta_1 \circ t &\rightsquigarrow a_{i(jk)l}, \\ -(\Delta_{-1} \otimes 1 \otimes 1) \circ \Delta_1 \circ t &\rightsquigarrow a_{(ij)kl}^\dagger, & -(1 \otimes 1 \otimes \Delta_{-1}) \circ \Delta_1 \circ t &\rightsquigarrow a_{ij(kl)}^\dagger, \end{aligned}$$

where $\text{id}_i : i \Rightarrow i$ denotes the identity modification on the 1-intertwiner i . Now note that, by coequivariance (2.22) $D_t \circ \Delta_{-1} = \Delta_0 \circ t$, we have

$$(\text{id} \otimes (\Delta_1 \circ t)) \circ \Delta_{-1} = (\text{id} \otimes \Delta_1) \circ \Delta_0^l \circ t, \quad ((\Delta_1 \circ t) \otimes \text{id}) \circ \Delta_{-1} = (\Delta_1 \otimes \text{id}) \circ \Delta_0^r \circ t,$$

whence the pentagon relation

$$\begin{array}{ccccc} & & & & ((ij)k)l \\ & & & & \nearrow a_{ijk} \text{id}_l \\ (ij)(kl) & \xrightarrow{a_{(ij)kl}^\dagger} & & & (i(jk))l \\ & \nwarrow a_{ij(kl)}^\dagger & & & \searrow a_{i(jk)l} \\ & & i(j(kl)) & \xleftarrow{\text{id}_i a_{jkl}} & i((jk)l) \end{array} \quad (6.18)$$

is equivalently expressed as

$$\begin{aligned} 0 &= (1 \otimes \Delta_{-1} \otimes 1) \circ \Delta_1 \circ t - (\Delta_{-1} \otimes 1 \otimes 1) \circ \Delta_1 \circ t - (1 \otimes 1 \otimes \Delta_{-1}) \circ \Delta_1 \circ t \\ &\quad + (\Delta_1 \otimes 1) \circ \Delta_0^r \circ t + (1 \otimes \Delta_1) \circ \Delta_0^l \circ t \\ &= [-\Delta_{-1} \circ \Delta_1 + \Delta_1 \circ \Delta_0] \circ t, \end{aligned}$$

which is nothing but the 2-coassociativity (4.5) precomposed with t . Now by the coPeiffer identity $\Delta'_0 = D_t \Delta_0$ (2.19), the same argument shows that the pentagon relations for the rest of the associator 2-morphisms (6.15), (6.16) are equivalent to applying the t -map $D_t, D_t[2]$ to (4.5).

Similarly, under the complete t -map $D_t[3] = t \otimes t \otimes t$, the 2-coassociativity condition (4.5) becomes

$$\Delta'_0 \circ \Phi = \Phi \circ \Delta'_0, \quad (6.19)$$

which by (6.11) implies the pentagon relation for the associator 1-morphism (6.17). \square

We examine the case where $\mathfrak{T} \neq 0$ is non-trivial in Appendix C. In particular, we show in **Proposition C.1** that \mathfrak{T} gives rise to the *pentagonator* 2-morphism π in $2\text{Rep}^T(\mathcal{G})$ implementing the pentagon relations (6.18).

Recall from **Proposition 4.1** that, for a self-dual weak 2-bialgebra, (4.5) follows from the 3-cocycle condition for the Hochschild 3-cocycle \mathcal{T} . Thus the entirety of the 2-bialgebra (or 2-Hopf algebra) structure plays a central role, precisely as one would expect in Tannakian duality [6, 55].

Lemma 6.4. *The associator 2-morphism (6.16) fits into diagrams of the form*

$$\begin{array}{ccc} (VW)U & \xrightarrow{a_{VWU}} & V(WU) \\ \downarrow k & \xrightarrow{a_{VWk}} & \downarrow k \\ (VW)U' & \xrightarrow{a_{VWU'}} & V(WU') \end{array} \quad (6.20)$$

together with the associator morphism (6.17). Moreover, the associator 2-morphisms (6.14), (6.15) are completely determined by (6.16), (6.17).

Proof. The first statement follows directly from the definitions, and by using the same argument as in the proofs of **Lemma 6.1**, and also later in **Lemma 7.2**. Similarly, by adapting the proof of **Lemma 6.2**, we see that (6.14), (6.15) admit the following decompositions

$$a_{ijk} = (a_{V'W'k} \cdot a_{ijU}) \circ t = \dots \text{etc.}, \quad D_\delta a_{ijU} = a_{V'jU} \cdot a_{iWU} = \dots \text{etc.},$$

where D_δ is the tensor triple of the t -map δ on $\mathbf{End}(V)$, and "etc." means permutations of the subscripts. This proves the second statement. \square

This naturally property shall become very important later in Section 7.2.

Remark 6.4. Suppose the endomorphism Φ in (6.17) is *inner*, in the sense that it is given by conjugation with an element — also denoted Φ — of $\mathcal{G}_0^{3\otimes}$, then the coassociativity condition becomes

$$(\text{id} \otimes \Delta'_0) \circ \Delta'_0 = \Phi((\Delta'_0 \otimes \text{id}) \circ \Delta'_0) \Phi^{-1},$$

and the 2-coassociativity condition (6.19) becomes

$$((\text{id} \otimes \text{id} \otimes \Delta'_0) \Phi)((\Delta'_0 \otimes \text{id} \otimes \text{id}) \Phi) = (\Phi \otimes \eta_0)((\text{id} \otimes \Delta'_0 \otimes \text{id}) \Phi)(\eta_0 \otimes \Phi),$$

where η_0 is the unit of \mathcal{G}_0 . In other words, $(\mathcal{G}_0, \Delta'_0, \Phi)$ in fact forms a *quasi-bialgebra* [66].

We have established $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ as a monoidal (fusion) 2-category, or equivalently a monoidal bicategory. We now turn to the braiding structure in the following.

7 The braided monoidal 2-category of 2-representations

We now turn to the braiding structure on the weak 2-representations afforded by the 2-R-matrix \mathcal{R} . We shall first examine some of the basic properties of the braiding map in Section 7.1. We will then study how such braiding maps interact with the weakened monoidal structures of the 2-representations in Section 7.2.

Let $(\mathcal{G}, \cdot, \Delta, \mathcal{R})$ denote a strict quasitriangular 2-bialgebra as defined in Section 3.4. Recall that a 2-R-matrix $\mathcal{R} = \mathcal{R}^l + \mathcal{R}^r$ on the 2-bialgebra \mathcal{G} consist of the following components

$$\mathcal{R}^l = \mathcal{R}_{(1)}^l \otimes \mathcal{R}_{(2)}^l \in \mathcal{G}_{-1} \otimes \mathcal{G}_0, \quad \mathcal{R}^r = \mathcal{R}_{(1)}^r \otimes \mathcal{R}_{(2)}^r \in \mathcal{G}_0 \otimes \mathcal{G}_{-1}$$

for which (3.25), (3.26), (3.27) are satisfied. The equivariance condition, (3.27), unambiguously defines an element

$$R = \mathcal{R}_{(1)}^r \otimes t\mathcal{R}_{(2)}^r (\equiv R^r) = t\mathcal{R}_{(1)}^l \otimes \mathcal{R}_{(2)}^l (\equiv R^l) \in \mathcal{G}_0 \otimes \mathcal{G}_0, \quad (7.1)$$

where $t: \mathcal{G}_{-1} \rightarrow \mathcal{G}_0$ is the t -map on \mathcal{G} . Notice by applying the t -map (at every leg in \mathcal{G}_{-1}) to (3.28), we obtain two identical expressions that are equivalent to the usual 1-Yang-Baxter equations

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$$

for the degree-0 R -matrix (7.1).

7.1 The braiding maps and their naturality

We shall use these components to define the braiding b on $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$. Take two 2-representations V, W of \mathcal{G} ; we define the braiding map between V, W by

$$b_{VW}: V \otimes W \rightarrow W \otimes V, \quad b_{VW} = \text{flip} \circ \rho_0(R) \quad (7.2)$$

where $\rho_0 = (\rho_V)_0 \otimes (\rho_W)_0$ on $V \otimes W$, and $R \in \mathcal{G}_0 \otimes \mathcal{G}_0$ is given in (7.1). By (6.11), the braiding between the tensor product 2-representations are then given by

$$b_{V(W \otimes U)} = \text{flip} \circ \rho_0((1 \otimes \Delta'_0)R), \quad b_{(V \otimes W)U} = \text{flip} \circ \rho_0((\Delta'_0 \otimes 1)R).$$

If $W = V$ are the same 2-representations of \mathcal{G} , then we have the *self-braiding* map $b_V = b_{VV}$. On the other hand, we define the *mixed braiding* map between a 1-morphism $i: V \rightarrow U$ and an object W by

$$\begin{aligned} b_{iW} &= \text{flip} \circ [i \circ \rho_{10}(\mathcal{R}^l) + (-1)^{\text{deg}} \rho_{01}(\mathcal{R}^r) \circ i], \\ b_{Wi} &= \text{flip} \circ [i \circ \rho_{01}(\mathcal{R}^r) + (-1)^{\text{deg}} \rho_{10}(\mathcal{R}^l) \circ i], \end{aligned} \quad (7.3)$$

where we have used the shorthand $\rho_{10} = (\rho_V)_1 \otimes (\rho_W)_0$ and $\rho_{01} = (\rho_U)_0 \otimes (\rho_W)_1$. The sign $(-1)^{\deg}$ depends on the degree of the complex $V \otimes W$; more explicitly, b_{iW} gives two maps

$$b_{iW}^1 : V_0 \otimes W_0 \rightarrow (W_{-1} \otimes U_0) \oplus (W_0 \otimes U_{-1}), \quad b_{iW}^2 : (V_{-1} \otimes W_0) \oplus (V_0 \otimes W_{-1}) \rightarrow W_{-1} \otimes U_{-1}$$

on the tensor product $V \otimes W$, the latter of which carries a non-trivial sign $(-1)^{\deg} = -1$; similarly for b_{Wi} . Now in the spirit of **Lemma 6.2**, we shall *define* the braiding maps b_{ij} between two 1-morphisms i, j by the decomposition formula⁶

$$b_{ij} = b_{jU} \cdot b_{Wi} \cong b_{Ti} \cdot b_{jV}, \quad \begin{cases} i : V \rightarrow U \\ j : W \rightarrow T \end{cases}. \quad (7.4)$$

Let $i : V \rightarrow V', j : U \rightarrow U'$ denote any 2-intertwiner. The above definition (7.3), together with (6.12) then allows us to form

$$\begin{aligned} b_{(i \otimes W)j} &= \text{flip}_{(V' \otimes U') \otimes W} \circ [(i \otimes j) \rho_{101}((\Delta_0^l \otimes 1) \mathcal{R}^r) + (-1)^{\deg} \rho_{011}((\Delta_0^r \otimes 1) \mathcal{R}^r) \circ (i \otimes j)], \\ b_{i(W \otimes j)} &= \text{flip}_{W \otimes (V' \otimes U')} \circ [(i \otimes j) \rho_{101}((\text{id} \otimes \Delta_0^r) \mathcal{R}^l) + (-1)^{\deg} \rho_{110}((1 \otimes \Delta_0^l) \mathcal{R}^l) \circ (i \otimes j)]. \end{aligned}$$

By applying strict 2-representations to (3.25), we obtain the following strict *higher hexagon relations*,

$$b_{(i \otimes W)j} = \text{id}_i \otimes b_{Wj} * b_{Wi} \otimes \text{id}_j, \quad b_{i(W \otimes j)} = \text{id}_i \otimes b_{jW} * b_{iW} \otimes \text{id}_j, \quad (7.5)$$

in which the associator isomorphisms a have been suppressed. We will reinstate them later in Section 7.2.

With the definitions (7.2), (7.3) in hand, we now need to prove some very important lemmas.

Lemma 7.1. *The maps b_{VW} and b_{iW}, b_{Wi} are respectively 2-intertwiners and modifications in $2\text{Rep}(\mathcal{G})$ for all 2-representation V, W and 2-intertwiner i iff (3.26) is satisfied.*

Proof. Note for each 2-representation ρ , the flip map, $\text{flip} : V \otimes W \rightarrow W \otimes V$ is a 2-intertwiner between ρ and $\rho' = \rho \circ \sigma$. Moreover, we interpret the cochain homotopy defined by $(\rho_{V \otimes W})_0^1(x)$ for each $x \in \mathcal{G}_0$ as a modification between the action $(\rho_{V \otimes W})_0^0(x)$ and itself, treated as a 2-intertwiner; similarly for ρ' . Therefore, in order for the mixed braiding map b_{iW} to be a modification in $2\text{Rep}(\mathcal{G})$, it must commute with the cochain homotopy $(\rho_{V \otimes W})_0^1(x)$ — namely

$$b_{iW} * (\rho_{V \otimes W})_0^1(x) = (\rho'_{W \otimes V})_0^1(x) * b_{iW},$$

where $*$ denotes the composition of cochain homotopies. With $\rho'_{W \otimes V} = (\rho_W \otimes \rho_V) \circ \sigma \circ \Delta$, this is satisfied by definition (7.2) of b_{iW} iff

$$\mathcal{R}^r \Delta_0^r(x) = \sigma(\Delta_0^l(x)) \mathcal{R}^r, \quad \mathcal{R}^l \Delta_0^l = \sigma(\Delta_0^r(x)) \mathcal{R}^l, \quad (7.6)$$

which is precisely (3.26).

Similarly, in order for the braiding map b_{VW} to be a 2-intertwiner, it must commute with the action $(\rho_{V \otimes W})_0^0(x)$ for each $x \in \mathcal{G}_0$:

$$b_{VW} \circ (\rho_{V \otimes W})_0^0(x) = (\rho'_{W \otimes V})_0^0(x) \circ b_{VW},$$

where \circ denotes the composition of 2-intertwiners.

First if the 2-representation ρ were strict, then this translates to the algebraic condition

$$\sigma \Delta'_0(x) R = R \Delta'_0(x),$$

which in fact follows also from (3.26). To see this, we recall the definitions (7.1) of R and (2.19) of the coproduct Δ'_0 , and simply apply $t \otimes 1$ and $1 \otimes t$ respectively to (3.26). The fact that t is an algebra homomorphism and that $(t \otimes 1) \circ \sigma = \sigma \circ (1 \otimes t)$ proves the statement.

Second, if the 2-representation ρ were weak, then in general the component ϱ gives rise to a possibly non-trivial invertible natural 2-morphism

$$\varrho(\sigma \Delta'_0(x), R) - \varrho(R, \Delta'_0(x)).$$

We will not need this 2-morphism in the following so we shall suppose $I_{b_{VW}, \bullet} = \text{id}$. □

⁶Alternatively, provided there exists a well-defined R-matrix $R_{-1} \in \mathcal{G}_{-1}^{2 \otimes}$ for the degree-(-1) coproduct Δ_{-1} , satisfying $(t \otimes 1 + 1 \otimes t) R_{-1} = \mathcal{R}$, we can define

$$b_{ij} = \text{flip} \circ [(i \otimes j) \circ \rho_{11}(R_{-1}) + (-1)^{\deg} \rho_{11}(R_{-1}) \circ i \otimes j]$$

such that (7.4) follows from the definition of R_{-1} .

Notice this lemma implies that $(\mathcal{G}_0, \Delta'_0, R)$ forms an ordinary quasitriangular 1-bialgebra. We can then leverage the well-known result in the literature [7, 67] that the Yang-Baxter equation for R implies the hexagon relation for the braiding structure b_{VW} at the level of the objects.

Next, we need to prove the naturality of b with respect to the 2-intertwiners $i : V \rightarrow U$. We shall do this via the same technique as **Lemma 6.1**.

Lemma 7.2. *Consider the intertwiners $i : V \rightarrow U$ and $j : U \rightarrow T$. The mixed braiding maps b_{iW}, b_{Wi} fit into the following diagrams*

$$\begin{array}{ccc} V \otimes W & \xrightarrow{b_{VW}} & W \otimes V \\ \downarrow i & \xrightarrow{b_{iW}} & \downarrow i \\ U \otimes W & \xrightarrow{b_{UW}} & W \otimes U \end{array}, \quad \begin{array}{ccc} W \otimes V & \xrightarrow{b_{WV}} & V \otimes W \\ \downarrow i & \xrightarrow{b_{Wi}} & \downarrow i \\ W \otimes U & \xrightarrow{b_{WU}} & U \otimes W \end{array}$$

in $2\text{Rep}^T(\mathcal{G})$. Moreover, given a 2-intertwiner $j : U \rightarrow T$ composable with i , the corresponding braiding 2-morphisms compose as $b_{jW} * b_{iW} = b_{j \circ i, W}$.

$$\begin{array}{ccc} V \otimes W & \xrightarrow{b_{VW}} & W \otimes V \\ \downarrow i & \xrightarrow{b_{iW}} & \downarrow i \\ U \otimes W & \xrightarrow{b_{UW}} & W \otimes U \\ \downarrow j & \xrightarrow{b_{jW}} & \downarrow j \\ T \otimes W & \xrightarrow{b_{TW}} & W \otimes T \end{array} = \begin{array}{ccc} V \otimes W & \xrightarrow{b_{VW}} & W \otimes V \\ \downarrow j \circ i & \xrightarrow{b_{(j \circ i)W}} & \downarrow j \circ i \\ T \otimes W & \xrightarrow{b_{TW}} & W \otimes T \end{array}$$

Proof. For brevity, we shall suppress the subscripts V, U, W on the 2-representations. Recall the two equivalent ways R^r, R^l to express R in (7.1). We can then write

$$b_{UW} \circ i = \text{flip} \circ \rho_0(R^r) \circ i, \quad i \circ b_{VW} = i \circ \text{flip} \circ \rho_0(R^l).$$

Consider the left diagram. As 2-morphisms in $2\text{Rep}(\mathcal{G})$ are given by cochain homotopies, we need to show that the definition (7.3) of the mixed braiding map $b_{iW} = b_{iW}^1 + b_{iW}^2$ fits into the following diagram

$$\begin{array}{ccccc} V_{-1} \otimes W_{-1} & \xrightarrow{D^+} & V_{-1} \otimes W_0 \oplus V_0 \otimes W_{-1} & \xrightarrow{D^-} & V_0 \otimes W_0 \\ \downarrow & \swarrow b_{iW}^2 & \downarrow & \swarrow b_{iW}^1 & \downarrow \\ W_{-1} \otimes U_{-1} & \xrightarrow{D^+} & W_{-1} \otimes U_0 \oplus W_0 \otimes U_{-1} & \xrightarrow{D^-} & W_0 \otimes U_0 \end{array}, \quad (7.7)$$

where the vertical arrows are the various graded components of $b_{UW} \circ i - i \circ b_{VW}$, and the horizontal arrows are the differentials on the three-term tensor product complex (6.10); for instance, the ones at the top row are given by $D^\pm = 1 \otimes \partial_V \pm \partial_V \otimes 1$.

As in **Lemma 6.1**, the key towards this is the commutative square (6.1), which states that for each $y \in \mathcal{G}_{-1}$ we have

$$(\rho_1(y)\partial, \partial\rho_1(y)) = \delta(\rho_1)(y) = (\rho_0)(Ty) = (\rho_0^1(Ty), \rho_0^0(Ty)).$$

Let us examine first the commutative triangle on the ends of (7.7). First, for the right-most triangle, we compute in terms of the components $b_{iW}^{1,2}$ that

$$\begin{aligned} D^- b_{iW}^1 &= (1 \otimes \partial_V - \partial_W \otimes 1) \circ \text{flip} \circ \rho(\mathcal{R}) \\ &= \text{flip} \circ \left[\rho_0^0(\mathcal{R}_{(1)}^r) \otimes \partial_W(\rho_1(\mathcal{R}_{(2)}^r)) \circ i - i \circ i \partial_V(\rho_1(\mathcal{R}_{(1)}^l)) \otimes \rho_0^0(\mathcal{R}_{(2)}^l) \right] \\ &= \text{flip} \circ \left[\rho_0^0(\mathcal{R}_{(1)}^r) \otimes t\mathcal{R}_{(2)}^r \circ i + i \circ \rho_0^0(-t\mathcal{R}_{(1)}^l \otimes \mathcal{R}_{(2)}^l) \right] \\ &= b_{UW} \circ i - i \circ b_{VW} \end{aligned}$$

as maps on $V_0 \otimes W_0$. Similarly for the left-most triangle, we have

$$b_{iW}^2 D^+ = \text{flip} \circ \rho(\mathcal{R}) \circ (1 \otimes \partial_W + \partial_V \otimes 1)$$

$$\begin{aligned}
&= \text{flip} \circ \left[\rho_0^1(\mathcal{R}_{(1)}^r) \otimes (\rho_1(\mathcal{R}_{(2)}^r)) \partial_W \circ i - i \circ (\rho_1(\mathcal{R}_{(1)}^l)) \partial_V \otimes \rho_0^1(\mathcal{R}_{(2)}^l) \right] \\
&= \text{flip} \circ \left[\rho_0^1(\mathcal{R}_{(1)}^r) \otimes t\mathcal{R}_{(2)}^r \circ i - i \circ \rho_0^1(t\mathcal{R}_{(1)}^l) \otimes \mathcal{R}_{(2)}^l \right] \\
&= b_{UW} \circ i - i \circ b_{VW}
\end{aligned}$$

as maps $V_{-1} \otimes W_{-1}$. Note the sign $(-1)^{\text{deg}}$ in (7.3) is non-trivial here as \mathcal{R} acts on the degree-(-1) part of the tensor product $V \otimes W$.

We now turn to the middle section of (7.7). We are required to compute the following,

$$\begin{aligned}
D^+ b_{iW}^2 &= (1 \otimes \partial_V + \partial_W \otimes 1) \circ \text{flip} \circ \rho(\mathcal{R}) \\
&= \text{flip} \circ \left[\rho_0^1(\mathcal{R}_{(1)}^r) \otimes \partial_W(\rho_1(\mathcal{R}_{(2)}^r)) \circ i - i \circ \partial_V(\rho_1(\mathcal{R}_{(1)}^l)) \otimes \rho_0^1(\mathcal{R}_{(2)}^l) \right] \\
&= \text{flip} \circ \left[\rho_0^1(\mathcal{R}_{(1)}^r) \otimes \rho_0^0(t\mathcal{R}_{(2)}^r) \circ i - i \circ \rho_0^0(t\mathcal{R}_{(1)}^l) \otimes \rho_0^1(\mathcal{R}_{(2)}^l) \right], \\
b_{iW}^1 D^- &= \text{flip} \circ \rho(\mathcal{R}) \circ (1 \otimes \partial_W - \partial_V \otimes 1) \\
&= \text{flip} \circ \left[\rho_0^0(\mathcal{R}_{(1)}^r) \otimes \rho_1(\mathcal{R}_{(2)}^r) \partial_W \circ i - i \circ \rho_1(\mathcal{R}_{(1)}^l) \partial_V \otimes \rho_0^0(\mathcal{R}_{(2)}^l) \right] \\
&= \text{flip} \circ \left[\rho_0^0(\mathcal{R}_{(1)}^r) \otimes \rho_0^1(t\mathcal{R}_{(2)}^r) \circ i - i \circ \rho_0^1(t\mathcal{R}_{(1)}^l) \otimes \rho_0^0(\mathcal{R}_{(2)}^l) \right].
\end{aligned}$$

Summing these and rearranging terms gives, as maps on $V_{-1} \otimes W_0 \oplus V_0 \otimes W_{-1}$,

$$\begin{aligned}
&\text{flip} \circ \left[\rho_0^0(\mathcal{R}_{(1)}^r) \otimes \rho_0^1(t\mathcal{R}_{(2)}^r) + \rho_0^1(\mathcal{R}_{(1)}^r) \otimes \rho_0^0(t\mathcal{R}_{(2)}^r) \right] \circ i \\
&\quad - \text{flip} \circ i \circ \left[\rho_0^1(t\mathcal{R}_{(1)}^l) \otimes \rho_0^0(\mathcal{R}_{(2)}^l) + \rho_0^0(t\mathcal{R}_{(1)}^l) \otimes \rho_0^1(\mathcal{R}_{(2)}^l) \right] \\
&\quad = b_{UW} \circ i - i \circ b_{VW}.
\end{aligned} \tag{7.8}$$

The diagram on the right is treated identically, and this establishes the first statement. The second statement directly follows from the fact that $(j \circ i) \circ \rho_V = j \circ \rho_U \circ i = \rho_X \circ (j \circ i)$ for composable 2-intertwiners i, j . \square

In particular, since **Lemma 7.1** proves that b_{VW} is a 1-morphism, we can *iterate* the braiding maps and define $b_{b_{VW}U}$ as a 2-morphism. **Lemma 7.2** then implies that this is a 2-morphism

$$\begin{array}{ccc}
(V \otimes W) \otimes U & \xrightarrow{b_{(VW)U}} & U \otimes (V \otimes W) \\
\downarrow b_{VW} & \xrightarrow{b_{b_{VW}U}} & \downarrow b_{VW} \\
(W \otimes V) \otimes U & \xrightarrow{b_{(WV)U}} & U \otimes (W \otimes V)
\end{array} \tag{7.9}$$

on three 2-representations V, W, U , and similarly for $b_{Vb_{WU}}$. This will be important later in Section 7.4.

Recall the "higher-hexagon relations" (7.5) following directly from the identities (3.25). We shall prove this in the weakened context in Section 7.2.

7.2 Braided 2-quasi-bialgebras; the modified hexagon relations

We now wish to keep track of the interplay between the fusion associators a and the braiding maps b — or, algebraically, the coassociator and the 2- R -matrix — on $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$. We shall do this by revisiting the universal characterization of 2- R -matrices in Section 3.4. In other words, we are prompted to study the *weak* 2-quantum double $D(\mathcal{G}, \mathcal{G})$ and its braided transposition Ψ .

Fix the weak 2-bialgebra \mathcal{G} . Despite the skeletal construction in Section 5, we are able to form $D(\mathcal{G}, \mathcal{G})$ here *without* assuming skeletality, since we know exactly how \mathcal{G} acts on itself by weak 2-representations — in the canonical way according to **Definition 4.1**. This fact also allows us to identify \mathcal{T}_D as merely several copies of the 3-cocycle \mathcal{T} on \mathcal{G} , and in particular the components $\triangleright_1 = \triangleleft_1 = \mathcal{T}$ are equal.

To proceed, we recall two facts we have learned previously.

- The condition (3.25) in the strict case follows from dualizing the braiding relation (3.19) (see (3.29)).
- From (5.4), the braiding relation (3.19) for the braided transposition Ψ is modified by \mathcal{T} whenever we invoke the associativity in $\mathcal{K} \cong D(\mathcal{G}, \mathcal{G})$.

Combining these means that (3.25) is modified by the *dual* of \mathcal{T} — ie. the coassociator Δ_1 — in the weakened case. More explicitly, we have

$$\begin{aligned} D_t \Delta_1(x)_{231} \cdot (1 \otimes \Delta_0) \mathcal{R} \cdot D_t \Delta_1(x)_{123} &= \mathcal{R}_{13} \cdot D_t \Delta_1(x)_{213} \cdot \mathcal{R}_{12}, \\ D_t \Delta_1(x)_{312}^{-1} \cdot (\Delta_0 \otimes 1) \mathcal{R} \cdot D_t \Delta_1(x)_{213}^{-1} &= \mathcal{R}_{13} \cdot D_t \Delta_1(x)_{132}^{-1} \cdot \mathcal{R}_{23}, \end{aligned} \quad (7.10)$$

for each $x \in \mathcal{G}_0$. This bears a striking resemblance to the defining relations of a *braided quasi-bialgebra* [66]; indeed, applying the double- t -map $D_t[2]$ to (7.10) yields, by definition (7.1), (6.17),

$$\Phi_{231}(x)(1 \otimes \Delta'_0) R \Phi_{123}(x) = R_{13} \Phi_{213}(x) R_{12}, \quad \Phi_{312}^{-1}(x)(\Delta'_0 \otimes 1) R \Phi_{213}^{-1}(x) = R_{13} \Phi_{132}^{-1}(x) R_{23}, \quad (7.11)$$

which is precisely a braided quasi-bialgebra structure on $(\mathcal{G}_0, \Delta'_0, R, \Phi)$; see *Remark 6.4*. This motivates the following definition.

Definition 7.1. A **braided 2-quasi-bialgebra**⁷ $(\mathcal{G}, \Delta = (\Delta_1, \Delta_0, \Delta_{-1}), \mathcal{T}, \mathcal{R})$ is a weak 2-bialgebra equipped with a universal 2- R -matrix \mathcal{R} and a coassociator $\Delta_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1}^{3 \otimes}$ such that (7.10), (7.11), (3.26) and (3.27) hold.

Similar to (7.5), by applying *strict* 2-representations $\rho = (\rho_1, \rho_0)$ to (7.10), we obtain:

Lemma 7.3. For each $X \in 2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, we have the decompositions (the **hexagon relations**)

$$\begin{aligned} \begin{cases} b_{(VW)X} = a_{XVW} \circ b_{VX} \circ a_{VXW}^\dagger \circ b_{WX} \circ a_{VWX} \\ b_{V(WX)} = a_{WXV}^\dagger \circ b_{VX} \circ a_{WVX} \circ b_{VW} \circ a_{VWX}^\dagger \end{cases} &\iff (7.11), \\ \begin{cases} b_{(Vj)X} = a_{XVj} * \text{id}_{b_{VX}} * a_{VXj}^\dagger * b_{jX} * a_{VjX} \\ b_{V(jX)} = a_{jXV}^\dagger * \text{id}_{b_{VX}} * a_{jVX} * b_{Vj} * a_{VjX}^\dagger \end{cases} &\iff \text{apply } D_t^+ \text{ to (7.10)}, \\ \begin{cases} b_{(iW)k} = a_{kiW} * b_{ik} * a_{ikW}^\dagger * b_{Wk} * a_{iWk} \\ b_{i(Wk)} = a_{Wki}^\dagger * b_{ik} * a_{Wik} * b_{iW} * a_{iWk}^\dagger \end{cases} &\iff (7.10), \end{aligned} \quad (7.12)$$

as 1-/2-morphisms, and similarly for all the other possible braiding maps on tensor products.

The decomposition formula for b_{ijk} follows from these, as well as the fact that b_{ij}, a_{ijk} are all determined by the mixed braiding/associators.

The 2-morphism $b_{(iW)X}$, for instance, can be expressed in terms of the following composition diagram

$$\begin{array}{ccccccccccc} (VW)X & \xrightarrow{a_{VWX}} & V(WX) & \xrightarrow{b_{WX}} & V(XW) & \xrightarrow{a_{VXW}^\dagger} & (VX)W & \xrightarrow{b_{VX}} & (XV)W & \xrightarrow{a_{XVW}} & X(VW) \\ \downarrow i & & \downarrow i & & \downarrow i & & \downarrow i & & \downarrow i & & \downarrow i \\ (UW)X & \xrightarrow{a_{UWX}} & V(WX) & \xrightarrow{b_{WX}} & U(XW) & \xrightarrow{a_{UXW}^\dagger} & (UX)W & \xrightarrow{b_{UX}} & (XU)W & \xrightarrow{a_{UVW}} & X(UW) \end{array}, \quad (7.13)$$

which has also appeared in [47]. This establishes most of the structural properties of $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ as a braided 2-category, and the final ingredient to introduce is the *hexagonator*.

7.3 The braiding hexagonator: weak 2-representations of a braided 2-quasi-bialgebra

We obtained the decomposition **Lemma 7.3** by applying a strict 2-representation to (7.10). However, as we have noted previously in *Remark 6.2*, 2-representations of a weak 2-bialgebra $(\mathcal{G}, \mathcal{T})$ *cannot* be strict, even when \mathcal{G} is skeletal. As such, we must take into account the additional component $\varrho : \mathcal{G}_0^{2 \otimes} \rightarrow \mathfrak{Cnd}(V)_{-1}$ when deriving the decompositions above (in particular (7.12)).

For the rest of the paper, it suffices to consider the case $t = 0$ or $t = \eta_0$, the constant map to the unit $\eta_0 \in \mathcal{G}_0$. Since ϱ is normalized and the second and third equations in (6.2) involve *pre*-composing ϱ with t , the only non-trivial relation is

$$\rho_0(xx') - \rho_0(x)\rho_0(x') = \delta\varrho(x, x'), \quad x, x' \in \mathcal{G}_0,$$

where we recall that $\delta : \mathfrak{Cnd}(V)_{-1} \rightarrow \mathfrak{Cnd}(V)_0$ is the t -map on the weak endomorphism 2-algebra. Therefore, in order to obtain the decomposition of the form (7.12) from (7.11), we must keep track of the terms involving ϱ that appear. For instance, we have

$$\rho_0^{3 \otimes}(R_{13} \Phi_{213}) - \rho_0^{3 \otimes}(R_{13}) \rho_0^{3 \otimes}(\Phi_{213}) = (\delta\varrho)^{3 \otimes}(R_{13}, \Phi_{213}),$$

⁷Note that a *quasi* 2-bialgebra, as opposed to a 2-quasi-bialgebra here, refers to a weak 2-bialgebra with trivial 3-cocycle $\mathcal{T} = 0$ but non-trivial coassociator Δ_1 .

in which we notice that the second term on the left-hand side is the composition $b_{VU} \circ a_{WVU}$.

More explicitly, translating (7.11) to (7.12) comes at a price given by a cochain homotopy

$$\begin{aligned}
\Omega_{V|WU}(x) &= (\varrho_V \otimes \varrho_W \otimes \varrho_U)(\Phi_{231}(x), (1 \otimes \Delta'_0)R\Phi_{123}(x)) - (\varrho_V \otimes \varrho_W \otimes \varrho_U)(R_{13}, \Phi_{213}(x)R_{12}) \\
&\quad + (\varrho_V \otimes \varrho_W \otimes \varrho_U)((1 \otimes \Delta'_0)R, \Phi_{123}(x)) - (\varrho_V \otimes \varrho_W \otimes \varrho_U)(\Phi_{213}(x), R_{12}) \\
&= (\varrho_V \otimes \varrho_W \otimes \varrho_U)(\Phi_{231}(x)(1 \otimes \Delta'_0)R, \Phi_{123}(x)) - (\varrho_V \otimes \varrho_W \otimes \varrho_U)(R_{13}\Phi_{213}(x), R_{12}) \\
&\quad + (\varrho_V \otimes \varrho_W \otimes \varrho_U)(\Phi_{231}(x), (1 \otimes \Delta'_0)R) - (\varrho_V \otimes \varrho_W \otimes \varrho_U)(R_{13}, \Phi_{213}(x)) \quad (7.14)
\end{aligned}$$

between the two sides of (7.11) for each $x \in \mathcal{G}_0$, and similarly its adjoint $\Omega_{V|WU}^\dagger$. We thus have the following diagrams

$$\begin{array}{ccc}
(V_{-1} \otimes W_{-1}) \otimes U_{-1} & \xrightarrow{\partial^{3\otimes}} & (V_0 \otimes W_0) \otimes U_0 \\
\downarrow & \swarrow \Omega_{V|WU} & \downarrow \\
W_{-1} \otimes (U_{-1} \otimes V_{-1}) & \xrightarrow{\partial^{3\otimes}} & W_0 \otimes (U_0 \otimes V_0) \\
V_{-1} \otimes (W_{-1} \otimes U_{-1}) & \xrightarrow{\partial^{3\otimes}} & V_0 \otimes (W_0 \otimes U_0) \\
\downarrow & \swarrow \Omega_{V|WU}^\dagger & \downarrow \\
(U_{-1} \otimes V_{-1}) \otimes W_{-1} & \xrightarrow{\partial^{3\otimes}} & (U_0 \otimes V_0) \otimes W_0
\end{array} ,$$


where the vertical arrows denote the decomposition (7.12). These diagrams cast Ω, Ω^\dagger as the **hexagonator** 2-morphisms in $2\text{Rep}^\mathcal{T}(\mathcal{G})$:

$$\begin{array}{ccc}
& V(WU) \xrightarrow{b_{V(WU)}} (WU)V & \\
a_{VWU} \nearrow & & \searrow a_{WUV} \\
(VW)U & \xrightarrow{\Omega_{V|WU}} & W(UV) , \\
b_{VW} \searrow & & \nearrow b_{VU} \\
& (WV)U \xrightarrow{a_{WVU}} W(VU) & \\
V(WU) & \xrightarrow{b_{(VW)U}} U(VW) & \\
a_{VWU}^\dagger \nearrow & & \searrow a_{UVW}^\dagger \\
V(WU) & \xrightarrow{\Omega_{V|WU}^\dagger} & (UV)W . \\
b_{WU} \searrow & & \nearrow b_{VU} \\
& V(UW) \xrightarrow{a_{VUW}^\dagger} (VU)W &
\end{array}$$

In other words, the quantities $\Omega_{V|WU}, \Omega_{V|WU}^\dagger$ by definition is an invertible modification implementing the two sides of the decomposition (7.12).

Now by the diagram (7.13), the 2-intertwiners $i : V \rightarrow U$ and their associated mixed braiding maps b_{iW} preserve these hexagon relations. This leads to the naturality of the hexagonator $\Omega_{V|WU}$ with respect to

2-intertwiners such that we have (cf. diagram (2.2) in [47])



and similarly for the adjoint diagrams with Ω^\dagger . The tensor product VX of 2-representations is equipped with the tensor product $\Omega_{VX|WU}$ hexagonator, which are by construction natural with respect to the braiding and associator morphisms we have studied above (cf. diagram (2.4) in [47]).

Remark 7.1. Notice we did not define any associators for the 2-morphisms μ in $2\text{Rep}^\mathcal{T}(\mathcal{G})$. This is because 2-morphisms in a 2-category the tensor product $\mu \otimes \nu = \mu * \nu$ given by composition is *strictly* associative; indeed, such an associator $a_{\mu\nu\lambda} : (\mu\nu)\lambda \Rightarrow \mu(\nu\lambda)$ would have to be a 3-morphism.

By the same token, the hexagon relations involving the mixed braiding maps (ie. the decompositions in **Lemma 7.3** aside from (7.12)), as well as the pentagon relations for the associator 2-morphisms (6.14), (6.15), (6.16), must hold strictly on-the-nose. However, the fact that a_{VWU} is a 1-morphism implies we can have a 2-morphism π , called the **pentagonator**, that implements its pentagon relation. We will show in Appendix C that π is given by the Hochschild 3-cocycle \mathfrak{T} attached to the weak endomorphism 2-algebra $\mathfrak{End}(V)$.

7.4 Main theorem and its proof

We are finally ready to state and prove the main theorem. As earlier, we will often omit the tensor products to lighten the notations.

Theorem 7.1. *The 2-representation 2-category $2\text{Rep}^\mathcal{T}(\mathcal{G})$ of a weak quasitriangular 2-bialgebra \mathcal{G} is a braided monoidal 2-category with trivial left-/right-equivalences $l : 1V \xrightarrow{\sim} V$, $r : V1 \xrightarrow{\sim} V$.*

We will prove this by using algebraic and diagrammatic manipulations that we have outlined throughout the paper, and reproduce all the coherence relations defining a braided monoidal 2-category in [46, 47].

Recall first that, from Section 6.2, we have trivial left- and right-unitors $l : 1V \rightarrow V$, $r : V1 \rightarrow V$, and hence all coherence relations involving them (ie. diagrams (2.5), (2.7)-(2.9) of [47]) are vacuously satisfied.

Braiding on the associator; diagram (2.6) of [47]. Let $V, W, U \in 2\text{Rep}^\mathcal{T}(\mathcal{G})$ be four 2-representations. Consider the mixed braiding 2-morphism $b_{a_{VWU}X}$, which by **Lemma 7.2** fits into a diagram of the form

$$\begin{array}{ccc}
 ((VW)U)X & \xrightarrow{b_{((VW)U)X}} & X((VW)U) \\
 \downarrow a_{VWU} & \xrightarrow{b_{a_{VWU}X}} & \downarrow a_{VWU} \\
 (V(WU))X & \xrightarrow{b_{(V(WU))X}} & X(V(WU))
 \end{array}$$

Lemma 7.3 states that we can in fact decompose the top and bottom 1-morphisms in this diagram, provided we keep in mind the hexagonator Ω, Ω^\dagger (7.14) that appears in doing so. We thus obtain a formula of the form

$$b_{((VW)U)X} \xrightarrow{\Omega_{((VW)U)X}^\dagger} a_{X(VW)U} \circ b_{(VW)X} \circ a_{(VW)XU}^\dagger \circ b_{UX} \circ a_{(VW)UX}$$

$$\begin{aligned} \xrightarrow{\Omega_{V|WX}^\dagger} & a_{X(VW)U} \circ \left[a_{XVW} \circ b_{VX} \circ a_{VXW}^\dagger \circ b_{WX} \circ a_{VWX} \right] \\ & \circ a_{(VW)XU}^\dagger \circ b_{UX} \circ a_{(VW)UX}, \end{aligned} \quad (7.15)$$

and similarly for the bottom 1-morphism $b_{(V(WU))X}$,

$$\begin{aligned} b_{(V(WU))X} & \xrightarrow{\Omega_{V|(WU)X}^\dagger} a_{XV(WU)} \circ b_{VX} \circ a_{VX(WU)}^\dagger \circ b_{(WU)X} \circ a_{V(WU)X} \\ & \xrightarrow{\Omega_{W|UX}^\dagger} a_{XV(WU)} \circ b_{VX} \circ a_{VX(WU)}^\dagger \\ & \circ \left[a_{XWU} \circ b_{WX} \circ a_{WXU}^\dagger \circ b_{UX} \circ a_{WUX} \right] \circ a_{V(WU)X}. \end{aligned} \quad (7.16)$$

Now notice that there are three identical braiding maps that appear in both of these formulas, b_{VX}, b_{WX}, b_{UX} , but they act on objects that differ by an associator: we have $b_{UX} : (VW)(UX) \rightarrow (VW)(XU)$ from (7.15) and $b_{UX} : V(W(UX)) \rightarrow V(W(XU))$ from (7.16), for instance. Such a square is precisely given by the diagram (6.20),

$$\begin{array}{ccc} (VW)(UX) & \xrightarrow{a_{VW(UX)}} & V(W(UX)) \\ \downarrow b_{UX} & \xrightarrow{a_{VWb_{UX}}} & \downarrow b_{UX} \\ (VW)(XU) & \xrightarrow{a_{VW(XU)}} & V(W(XU)) \end{array} ,$$

and similarly for the other braiding maps that occur in both (7.16), (7.15).

Putting this all together, by successively decomposing the braiding maps, we achieve the following diagrammatic expression for $b_{a_{VWU}X}$ (here we only label the 2-morphisms for clarity):

$$\begin{array}{ccccccc} ((VW)U)X & \xrightarrow{\hspace{15em}} & & & & & X((VW)U) \\ \downarrow & & & \xrightarrow{\Omega_{(VW)|UX}^\dagger} & & & \downarrow \\ (VW)(UX) & \longrightarrow (VW)(XU) & \longrightarrow & ((VW)X)U & \xrightarrow{\hspace{15em}} & & (X(VW))U \\ \downarrow \xrightarrow{a_{VWb_{UX}}} \downarrow & & \downarrow & & \downarrow \xrightarrow{\Omega_{V|WX}^\dagger} \downarrow & & \downarrow \\ V(W(UX)) & \longrightarrow V(W(XU)) & \longrightarrow & V((WX)U) & \longrightarrow & V((XW)U) & \longrightarrow ((VX)W)U \\ \downarrow & & \downarrow \xrightarrow{\Omega_{W|UX}^\dagger} \downarrow & & \downarrow \xrightarrow{a_{b_{VX}WU}} \downarrow & & \downarrow \\ V((WU)X) & \xrightarrow{\hspace{15em}} & V(X(WU)) & \longrightarrow & (VX)(WU) & \longrightarrow & (XV)(WU) \\ \downarrow & & \downarrow \xrightarrow{\Omega_{V|(WU)X}^\dagger} \downarrow & & \downarrow & & \downarrow \\ (V(UW))X & \xrightarrow{\hspace{15em}} & & & & & X(V(UW)) \end{array}$$

This is precisely diagram (2.6) in [47], with non-trivial hexagonators.

The iterated braiding map and diagram (2.10) of [47]. Now consider the iterated braiding 2-morphism $b_{Vb_{UW}}$ (7.9). By the same logic as above, we can use the decomposition (7.12) once again on the top and bottom braiding morphisms that appear in the diagram,

$$\begin{aligned} b_{V(UW)} & \xrightarrow{\Omega_{V|UW}} a_{UWV}^\dagger \circ b_{VW} \circ a_{UVW} \circ b_{VU} \circ a_{VUW}^\dagger, \\ b_{V(WU)} & \xrightarrow{\Omega_{V|WU}} a_{WUV}^\dagger \circ b_{VU} \circ a_{WVU} \circ b_{VW} \circ a_{VWU}^\dagger. \end{aligned}$$

We can thus form the composition

$$b_{\Omega_{V|WU}} \equiv \Omega_{V|WU}^{-1} \cdot b_{Vb_{UW}} \cdot \Omega_{V|UW}, \quad (7.17)$$

which fits into a diagram that "pastes" two hexagon diagrams together,

This is precisely (2.10) in [47]. Note that, by construction (7.17), the 2-morphisms $b_{\Omega_V|W\bullet}$ are natural and invertible, and constitutes part of the braiding data $(b_{VW}, b_{\Omega_V|W\bullet})$ on the objects V, W .

The second axiom of [46], or (2.1) in [47]. Let us now focus on (6.3). Recall that it states, for $x_1, x_2, x_3 \in \mathcal{G}_0$, that

$$\begin{aligned} \rho_1(\mathcal{T}(x_1, x_2, x_3)) - \mathfrak{T}(\rho_0(x_1), \rho_0(x_2), \rho_0(x_3)) &= \rho_0(x_1) \cdot \varrho(x_2, x_3) - \varrho(x_1 x_2, x_3) \\ &\quad + \varrho(x_1, x_2 x_3) - \varrho(x_1, x_2) \cdot \rho_0(x_3), \end{aligned}$$

where \mathfrak{T} is the Hochschild 3-cocycle on the weak endomorphism 2-algebra $\mathfrak{E}nd(V)$ of a particularly chosen weak 2-vector space $V \in 2\mathbf{Vect}$. We shall now specialize x_1, \dots, x_3 to the elements in \mathcal{G}_0^3 of (7.11), and let the equation act on V .

By some computations, we see that the right-hand side translates to the composition of 2-morphisms

$$\text{id}_{\text{id}_W} \Omega_{V|UX} * \Omega_{V|W(UX)} * \Omega_{V|(WU)X}^{-1} * (\Omega_{V|WU} \text{id}_{\text{id}_X})^{-1},$$

while on the term $\rho \circ \mathcal{T}$ on the left dualizes to terms of the form $(\rho_V \otimes \dots \otimes \rho_X)(\Delta_1 \circ R - \mathcal{R} \circ D_t^\dagger \Delta_1)$, which translates to

$$a_{Wb_{VU}X}^\dagger * a_{b_{VW}UX} * a_{WU b_{VX}}^\dagger * b_{V a_{WUX}}.$$

Now by leveraging the result **Proposition C.1** in Appendix C, the term $\mathfrak{T} \circ \rho_0^{3\otimes}$ in fact defines the pentagonators π on $2\text{Rep}^\tau(\mathcal{G})$. The left-hand side then acquires also the contribution

$$\pi_{WVUX} * \pi_{WUVX} * \pi_{VWUX}^\dagger * \pi_{WUXV}^\dagger,$$

where $\pi_{WUXV}(x) = \mathfrak{T}((\rho_W)_0(x), (\rho_U)_0(x), (\rho_X)_0(x))(V)$; see (C.1).

Altogether, this gives rise to the equation

$$\begin{aligned} \pi_{WVUX} * \pi_{WUVX} * \text{id}_{\text{id}_W} \Omega_{V|UX} * \Omega_{V|W(UX)} * a_{b_{VW}UX}^\dagger * b_{V a_{WUX}}^\dagger = \\ \pi_{VWUX} * \pi_{WUXV} * \Omega_{V|WU} \text{id}_{\text{id}_X} * \Omega_{V|(WU)X} * a_{Wb_{VU}X}^\dagger * a_{WU b_{VX}}^\dagger \end{aligned}$$

for $V, W, U, X \in 2\text{Rep}^\tau(\mathcal{G})$, which is precisely the second braided axiom in [46] (or equivalently axiom (2.1) in [47]). In [47], this axiom was also captured in a cohomological manner in (3.2) there, but where the adjoint equivalences a_{iVW}, a_{iVW}^\dagger are omitted.

In summary, we find that $2\text{Rep}^\tau(\mathcal{G})$ has the following ingredients:

objects	1-morphisms	2-morphisms
2-representations $(V, b_{V\bullet}, \Omega_{V \bullet\bullet})$	2-intertwiners $(i, b_{i\bullet})$	equivariant cochain homotopies μ

This establishes **Theorem 7.1** and concludes our paper.

8 Conclusion

In this paper, we have given a construction of a categorified notion of a quantum double suited for 2-groups/2-algebras [40]. This was accomplished by naturally lifting Majid's quantum double construction [44, 7] to internal categories, which makes the structural theorems manifest. In particular, we have provided explicit algebraic computations that demonstrate concretely the notion of duality between 2-bialgebras, and how two 2-bialgebras can be "pasted together" through the notion of 2-representations [45]. We have also given examples which demonstrate that the category of bialgebras (and their quantum doubles) embeds into the category of 2-bialgebras (and their 2-quantum doubles), generalizing an analogous statement proven in [41] for the classical case.

By endowing Baez-Crans 2-vector spaces $2\text{Vect}_{\text{int}}$ with higher homotopical data, we described the *weak* 2-representation theory for 2-bialgebras \mathcal{G} . We give a concrete description of the 2-category $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$, which by definition comes with a forgetful 2-functor $2\text{Rep}(\mathcal{G}) \rightarrow 2\text{Vect}$ into the k -linear semisimple fusion 2-category 2Vect of such "deformed" Baez-Crans 2-vector spaces. Though we had provided an operational description of 2Vect — in particular the property that its endomorphism categories are modelled by A_2 -algebras — its precise construction shall appear soon in a later work.

As in the case of 1-Hopf algebras [36], the monoidal structure on $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ is controlled by the coproduct and the coassociator, whence the naturality and Gray property of the tensor product follow from the coequivariance and coPeiffer identities (**Lemmas 6.1, 6.2**). We have also introduced the 2- R -matrix \mathcal{R} of \mathcal{G} , which was defined naturally from the properties of the 2-quantum double $D(\mathcal{G}, \mathcal{G})$. We show that the resulting braiding on $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ is in fact natural and coherent (**Lemmas 7.1, 7.2**), suggesting that our characterization of the 2- R -matrix is *universal*.

Then, together with the structure of weak 2-representations, we identified the pentagonator and hexagonator 2-morphisms and exhibited all the necessary coherence diagrams to prove **Theorem 7.1**. This gives a direct correspondence of the ingredients of a braided monoidal 2-category [46, 47] with those of an underlying weak quasitriangular 2-bialgebra \mathcal{G} .

We also note that **Theorem 7.1** hints towards a (braided) *higher Tannakian reconstruction* (cf. [55]). It should state that, morally, given a sufficiently "nice" braided 2-category (such as the Drinfel'd centre $Z_1(\mathcal{D})$ of a monoidal 2-category \mathcal{D} [47]) and a fibre 2-functor $F : \mathcal{C} \rightarrow 2\text{Vect}$, there is a braided equivalence $\mathcal{C} \simeq 2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ such that the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & 2\text{Vect} \\ & \searrow \sim & \nearrow \\ & 2\text{Rep}^{\mathcal{T}}(\mathcal{G}) & \end{array}$$

commutes. The authors are aware that efforts towards such a result in the semisimple setting are currently being undertaken. Together with this work, it allows to distill all coherences in \mathcal{C} into a sequence of Hochschild cohomological descent equations between the weak 2-bialgebra $(\mathcal{G}, \mathcal{T}, \mathcal{R})$ and the weak 2-endomorphism 2-algebra $\mathfrak{End}(V)$.

Furthermore, as mentioned in the Introduction, such a braided equivalence would also allow us to reconstruct the topological field theory associated to a 4D gapped topological phase, described by a braided 2-category [68]. The work towards this goal, in the case of the 3D toric code as well as its spin- \mathbb{Z}_2 variant, is in [58].

We show in Appendix C that we are able to reproduce the 2-representation 2-category $2\text{Rep}_{\mathcal{G}}$ of a skeletal 2-group G [32], which forms a fusion 2-category used by [31] as a way to construct quantum invariants of 4-manifolds. The notion of 2-bialgebras (or 2-Hopf algebras in Appendix A) we have developed here would then also serve the same purpose, in analogy with the 3-dimensional case with ordinary Hopf algebras as noticed by Witten [8] — or, more generally, with the theory of modular functors by Turaev [69]. We shall say a bit more words about this in the following, as well as some open research directions that we find interesting.

Relations with existing quantum group categorifications. There have been numerous proposals for the notion of "categorified quantum groups" in the literature, such as the trialgebra proposal of Pfeiffer [55], the quantum 2-group of Majid [49], or the quantum groupoid of Lu [70]. It is well-known that group crossed-modules are equivalent to 2-groups and group groupoids [42], and the cat^1 -Hopf algebras of Wagemann [40] coincide with trialgebras in the cocommutative case [55]. It then stands to reason that our definition of *strict* 2-bialgebras may be closely related to all of these alternative formulations.

On the other hand, the content of **Theorem 1.1** has been explored in the context of Hopf categories of Gurski [65] by Neuchl [29]. Hence, we posit that the relationship between our *weak* 2-bialgebras and these Hopf categories should be understood through a "model change" from 2-vector spaces of the "weak Baez-Crans" sense

2Vect to the Kapranov-Voevodsky sense 2Vect^{KV} [71].⁸ This is the main motivating problem for an upcoming work.

Morita duality and sub-symmetry gauging. Note in Appendix C, we proved that (weak) 2-representations as defined in **Definition 6.3** endow a \mathcal{G} -module action 2-functor on the 2-category 2Vect of 2-vector spaces. In particular, we explicitly showed that 2-representation 2-category 2Rep_G of a 2-group G , in the context of [48], can be recovered from our definition $2\text{Rep}(kG)$ by restricting to the skeletal 2-group algebra kG (with trivial Postnikov class $\tau = 0$ and grouplike coproduct). This suggests to us the possibility to generalize the theory of **Morita duality** developed in [32] to general 2-bialgebras.

To be more precise, we first recall that the *Morita dual* $\mathcal{C}_{\mathcal{D}}^*$ of a category \mathcal{C} with respect to a \mathcal{C} -module category \mathcal{D} is defined as

$$\mathcal{C}_{\mathcal{D}}^* = \text{Func}_{\mathcal{C}}(\mathcal{D}, \mathcal{D}),$$

namely it is the category of \mathcal{C} -linear endofunctors (ie. \mathcal{C} -intertwining endomorphisms) of \mathcal{D} . Physically, this Morita duality operation has the interpretation of "gauging" the subsymmetry in \mathcal{C} by condensing out the excitations labeled by \mathcal{D} . The same idea can be applied to the context of 2-categories; in particular, various Morita duals of the 2-representation 2-category 2Rep_G of a (skeletal) 2-group G have been computed in [32], such as

$$\begin{aligned} (2\text{Rep}_G)_{2\text{Rep}_{G_0}}^* &\simeq 2\text{Rep}_{G_{-1} \times G_0}, \\ (2\text{Rep}_G)_{2\text{Vect}}^* &\simeq 2\text{Vect}_G, \\ (2\text{Rep}_G)_{2\text{Vect}_{G_{-1} \times G_0/G_0}}^* &\simeq 2\text{Vect}_{G_{-1} \times G_0}, \end{aligned}$$

where $2\text{Vect}_G = 2\text{Fun}([*, G_{-1}, G_0], 2\text{Vect})$ denotes the 2-category of G -graded 2-vector spaces, obtained by taking the 2-group G as a 2-category with object-set G_0 , 1-morphisms G_{-1} and trivial 2-morphisms (recall in contrast $2\text{Rep}_G = 2\text{Fun}([G_{-1}, G_0, *], 2\text{Vect})$, which is the notion of 2-representations we are using). Notice condensing the trivial phase 2Vect is in fact not a trivial operation: it shifts the grading of the 2-groupoid symmetry G .

In any case, by leveraging the results of Appendix C, it would be possible to study Morita duals of a general 2-bialgebra \mathcal{G} as defined in this paper. It would be interesting to study in particular the various Morita duals of the 2-quantum double symmetry $D(kG)$ associated to a 2-group G with respect to one of its duality sectors kG, kG^* . Such an operation would have the physical interpretation of condensing the excitations labeled by kG in a "fully-gauged theory" equipped with the complete 2-quantum double symmetry $D(kG) \cong kG \bowtie kG^*$, which we expect to have excitations in $2\text{Rep}(kG^*)$ or $2\text{Vect}(kG^*)$ leftover. This point us towards a more complete and systematic understanding of categorified **Kramers-Wannier duality** [32].

Ribbon tensor 2-categories and modular invariants of 4-manifolds. Recall that, in the 1-Hopf algebra case H , the representation category $\text{Rep}(H)$ can be modelled with *ribbon diagrams* [36], which are pictorial presentations for the computations one can do in $\text{Rep}(H)$. This allowed one to deduce quantum invariants of 3-manifolds from the underlying braided fusion category: such as the Turaev-Viro state-sum invariants [72, 73], the multifusion invariants of Cui [60], and the modular invariants of Reshetikhin-Turaev [74].

The above Turaev-Viro state sum model has seen a recent generalization to 4-dimensions, known as the *Douglas-Reutter model* [31]. This is a state sum model which takes as input a fusion spherical 2-category, and outputs a 4-manifold invariant. Similar constructions had also been studied by Mackaay [28, 75, 76], but the *2-ribbon calculus* underlying the Douglas-Reutter model served tantamount importance.

Nevertheless, these invariants are *not* modular, and a 4-dimensional analogue of the Reshetikhin-Turaev invariant is still unknown. For this, one requires the notion of a **ribbon tensor 2-category**, in which both the braiding and the *twist* operations of the underlying fusion 2-category play a central role. Algebraically, we expect such higher ribbon structures to be captured by **ribbon 2-Hopf algebras**.

In Appendix A, we introduced a strict notion of a 2-Hopf algebra equipped with both a 2-R-matrix \mathcal{R} and an *antipode* S . However, to model a general ribbon tensor 2-category, we would require a "weak antipode" S as well as the central ribbon/tortile element ν that arises from \mathcal{R}, S . We shall pursue this line of thinking in a followup work, in an effort to construct the 4-dimensional analogue of the Reshetikhin-Turaev modular invariant.

⁸We emphasize that this model change would not be possible in the strict Baez-Crans sense, as $2\text{Vect}_{\text{int}}$ is known to be distinct from 2Vect^{KV} as 2-categories.

A 2-Hopf algebras

Recall in the usual 1-algebra case that a Hopf algebra H is by definition a bialgebra equipped with an *antipode* $S : H \rightarrow H$ which is an anti-algebra and anti-coalgebra map, satisfying

$$\cdot \circ (\text{id} \otimes S) \circ \Delta = \cdot \circ (S \otimes \text{id}) \circ \Delta = \eta \circ \epsilon,$$

where \cdot, Δ are the product/coproduct on H and η, ϵ are the unit/counit maps. Correspondingly, we shall define a 2-Hopf algebra \mathcal{G} as a 2-bialgebra equipped with an appropriate notion of an antipode $S : \mathcal{G} \rightarrow \mathcal{G}$.

Opposite 2-algebras. Recall that any algebra A comes with an "opposite" algebra A^{opp} , for which the algebra structure is written "backwards"; the multiplication is given by

$$A \otimes A \rightarrow A, \quad x \otimes x' \mapsto x'x.$$

Similarly, an **opposite 2-algebra** \mathcal{G}^{opp} of a (strict) 2-algebra \mathcal{G} consist of opposites $\mathcal{G}_0^{\text{opp}}, \mathcal{G}_{-1}^{\text{opp}}$ of the graded components of \mathcal{G} , and a "swapped" \mathcal{G}_0 -bimodule structure on \mathcal{G}_{-1} :

$$\begin{aligned} \cdot_l^{\text{opp}} : \mathcal{G}_0 \otimes \mathcal{G}_{-1} &\mapsto \mathcal{G}_{-1} & \cdot_r^{\text{opp}} : \mathcal{G}_{-1} \otimes \mathcal{G}_0 &\mapsto \mathcal{G}_{-1} \\ x \otimes y &\mapsto y \cdot x = x \cdot_l^{\text{opp}} y, & y \otimes x &\mapsto x \cdot y = y \cdot_r^{\text{opp}} x. \end{aligned} \quad (\text{A.1})$$

In other words, the left \mathcal{G}_0 -module structure of \mathcal{G}_{-1} is swapped with the right one. It is easy to see that the equivariance property and Peiffer identity for t still take the form (2.5), (2.6).

Antipode and 2-Hopf algebras. In analogy with the 1-algebra case, an **anti-2-algebra map** on \mathcal{G} is equivalent to a 2-algebra map $\mathcal{G} \rightarrow \mathcal{G}^{\text{opp}}$ into the opposite 2-algebra \mathcal{G}^{opp} as defined above. More explicitly, an anti-2-algebra map $\phi : \mathcal{G} \rightarrow \mathcal{G}$ has graded components $\phi_{0,-1} : \mathcal{G}_{0,-1} \rightarrow \mathcal{G}_{0,-1}$ as anti-algebra maps, such that $\phi_0 t = t \phi_{-1}$ and

$$\phi_{-1}(x \cdot y) = \phi_{-1}(y) \cdot \phi_0(x), \quad \phi_{-1}(y \cdot x) = \phi_0(x) \cdot \phi_{-1}(y)$$

for each $x \in \mathcal{G}_0, y \in \mathcal{G}_{-1}$.

This allows us to define an *antipode* $S_0 = (S_0^1, S_0^0) : \mathcal{G} \rightarrow \mathcal{G}$ as an anti-2-algebra map on \mathcal{G} such that

$$\begin{aligned} \cdot \circ (\text{id} \otimes S_0^1) \circ \Delta_{-1} &= \cdot \circ (S_0^1 \otimes \text{id}) \circ \Delta_{-1} = \eta_{-1} \epsilon_{-1}, \\ \cdot \circ (\text{id} \otimes S_0^0) \circ \Delta_0^l &= \cdot \circ (S_0^1 \otimes \text{id}) \circ \Delta_0^l = \eta_{-1} \epsilon_0, \\ \cdot \circ (\text{id} \otimes S_0^1) \circ \Delta_0^r &= \cdot \circ (S_0^0 \otimes \text{id}) \circ \Delta_0^r = \eta_{-1} \epsilon_0, \end{aligned} \quad (\text{A.2})$$

where \cdot, Δ are the 2-bialgebra product/coproduct and η, ϵ are the unit/counit on \mathcal{G} .

Definition A.1. A (strict) **2-Hopf algebra** is a (strict) counital 2-bialgebra $(\mathcal{G}, \cdot, \Delta, \epsilon)$ as in **Definition 2.6** equipped with an antipode S_0 satisfying (A.2).

Note (A.2) directly implies that $(\mathcal{G}_{-1}, \Delta_{-1}, S_0^1)$ forms a Hopf algebra. Furthermore, recalling the definition (2.19) $\Delta'_0 = \frac{1}{2} D_t^+ \Delta_0$ of the coproduct in degree-0, as well as the property that $t \eta_{-1} = \eta_0$, we have

$$\cdot \circ (\text{id} \otimes S_0^0) \circ \Delta'_0 = \cdot \circ (S_0^0 \otimes \text{id}) \circ \Delta'_0 = \eta_0 \circ \epsilon_0$$

from (A.2), as the t -map by definition intertwines between the antipodes $S_0^0 t = t S_0^1$. As such, $(\mathcal{G}_0, \Delta'_0, S_0^0)$ itself forms a Hopf algebra for which the t -map is a Hopf algebra map, given the conditions (2.18), (2.20), (2.22), (2.24) hold. Moreover, the 2-coalgebra compatibility conditions (2.31) implies that $H_{-1} = (\mathcal{G}_{-1}, \Delta_{-1}, S_0^1)$ forms a Hopf bimodule algebra over $H_0 = (\mathcal{G}_0, \Delta'_0, S_0^0)$.

In other words, our definition of a *strict* 2-Hopf algebra gives rise to a crossed-module of Hopf algebras $t : (\mathcal{G}_{-1}, \Delta_{-1}, S_0^1) \rightarrow (\mathcal{G}_0, \Delta'_0, S_0^0)$, which is precisely the definition of a "cat¹-Hopf algebra" of [40]. We are able to go even further, as we are able to introduce the 2-R-matrix, as well as study the monoidal weakening of \mathcal{G} . We shall say a bit more about the former in the following.

Quasitriangular strict 2-Hopf algebras. Now let $(\mathcal{G}, \Delta, \mathcal{R})$ denote a quasitriangular 2-bialgebra, equipped with the 2-R-matrix \mathcal{R} . Recall that an antipode $S : \mathcal{G} \rightarrow \mathcal{G}$ is a anti-2-algebra map on \mathcal{G} such that (A.2) are satisfied. Together with the 2-R-matrix and (3.25), it follows that we must have

$$\begin{aligned} (S_0^1 \otimes \text{id}) \mathcal{R}^l \cdot \mathcal{R}^r &= \eta_{-1}^{2 \otimes}, & (S_0^0 \otimes \text{id}) \mathcal{R}^r \cdot \mathcal{R}^l &= \eta_{-1}^{2 \otimes}, \\ (\text{id} \otimes S_0^1) \mathcal{R}^r \cdot \mathcal{R}^l &= \eta_{-1}^{2 \otimes}, & (\text{id} \otimes S_0^0) \mathcal{R}^l \cdot \mathcal{R}^r &= \eta_{-1}^{2 \otimes}. \end{aligned} \quad (\text{A.3})$$

Note here that, as the skew-pairing $\langle \cdot, \cdot \rangle_{\text{sk}}$ is non-degenerate (the "quasi" in quasitriangular), the 2-R-matrix components $\mathcal{R}^{l,r}$ are square and hence admit uniquely defined inverses as square matrices.

Definition A.2. Such a tuple $(\mathcal{G}, \Delta, \mathcal{R}, S)$ satisfying (A.3) defines a strict **quasitriangular strict 2-Hopf algebra**.

If $\mathcal{G} = kG$ were the 2-group algebra constructed from (2.15), then the algebra action \cdot is simply matrix product, and we recover the usual notion that $(S \otimes \text{id})\mathcal{R}, (\text{id} \otimes S)\mathcal{R} = \mathcal{R}^{-1}$ provided \mathcal{R} is quasitriangular.

To not bloat up this paper anymore than we already have, we conclude this section by noting that the embedding in *Remark 2.34* extends to (quasitriangular) 2-Hopf algebras,

$$\text{HopfAlg} \hookrightarrow 2\text{HopfAlg}, \quad H \mapsto \mathcal{H} = H \xrightarrow{\text{id}} H.$$

The antipode on \mathcal{H} is simply two copies of that S on H , and (A.2), (A.3) is automatically satisfied as the coproduct components $\Delta_{-1} = \Delta_0 = \Delta'_0$ and the R-matrices $\mathcal{R} = R$ all coincide; see Section 2.2.2.

In general, we should consider a weakened form of this 2-Hopf structure, based on weak 2-bialgebras that we have introduced in the main text. In particular, there is a "cochain homotopy" component $S_1 : \mathcal{G}_0 \rightarrow \mathcal{G}_{-1}$ to the antipode which "deforms" the relations (A.2), (A.3). We shall study this structure in more detail in a followup paper.

B Classical limits of (weak) 2-bialgebras and 2- R -matrices

In this section, we prove that the notion of 2-quantum doubles we have defined in Section 3 in the main text reproduces the known notion of 2-Manin triples of Lie 2-bialgebras [41, 3] in the classical limit. We shall also examine the quasi-weak case and show that a weak/quasi-2-bialgebra as defined in **Definition 4.3** reproduces a weak/quasi-Lie 2-bialgebra as defined in [42, 43].

Classical limit and the Lie-ification functor. Given an (associative) algebra $A \in \text{Alg}_{\text{ass}}$, it is well-known [36, 40] that there is a *Lie-ification functor* $\mathcal{L} : \text{Alg}_{\text{ass}} \rightarrow \text{LieAlg}$ that assigns A to its "classical" Lie algebra $\mathfrak{g}(A)$. The Lie bracket is given by the commutator $[X, X'] = XX' - X'X$, where $X \in \mathfrak{g}(A)$ is the image of an element $x \in A$ under \mathcal{L} . The associativity of A implies the Jacobi identity of $[\cdot, \cdot]$; note A only needs to be left-symmetric (not necessarily associative) in order for $\mathfrak{g}(A)$ to enjoy the Jacobi identity [41].

There is a left-adjoint to the Lie-ification functor given by the universal envelope $U : \mathfrak{g} \mapsto U(\mathfrak{g})$, which can be understood as a "quantization" map [44]. There is an analogous result for associative 2-algebras [40].

Lemma B.1. *The Lie-ification functor $\mathcal{L} : 2\text{Alg}_{\text{ass}} \rightarrow \text{Lie}2\text{Alg}$ lifts to associative 2-algebras (see **Definition 2.2**), where $\mathfrak{g}(\mathcal{G}) = \mathcal{L}(\mathcal{G}_{-1}) \xrightarrow{t} \mathcal{L}(\mathcal{G}_0)$ is a Lie 2-algebra with*

$$X \triangleright Y = X \cdot Y - Y \cdot X, \quad X = \mathcal{L}(x), \quad Y = \mathcal{L}(y),$$

where $x \in \mathcal{G}_0, y \in \mathcal{G}_{-1}$. Moreover, the universal envelop functor U also lifts to Lie 2-algebras $U(\mathfrak{g}) = U(\mathfrak{g}_{-1}) \xrightarrow{t} U(\mathfrak{g}_0)$, such that U is left-adjoint to \mathcal{L} .

In the following, we shall write $[\cdot, \cdot] : \mathfrak{g}^{2\wedge} \rightarrow \mathfrak{g}$ as the binary L_2 -bracket on \mathfrak{g} , which consists of the Lie bracket in \mathfrak{g}_0 as well as the action \triangleright of \mathfrak{g}_0 on \mathfrak{g}_{-1} [41, 3].

Note Lie-ification \mathcal{L} is a functor. This means that, in particular, it sends a 2-algebra representation $\rho : \mathcal{G} \rightarrow \text{End}(V)$ on 2-vector space V to a Lie 2-algebra representation $\mathcal{L}(\rho) : \mathfrak{g}(\mathcal{G}) \rightarrow \mathfrak{gl}(V)$ as defined in [41, 45].

B.1 Lie 2-bialgebras and the 2-classical double

We now extend the above lemma to associative 2-quantum doubles. Let $(\mathcal{G}, \cdot, \Delta)$ denote a strict 2-bialgebra as defined in **Definition 3.1**, and let $(\mathcal{G}^*, \cdot^*, \Delta^*)$ denote its dually-paired 2-algebra. We put $\mathfrak{g} = \mathcal{L}(\mathcal{G})$ and $\mathfrak{g}^* = \mathcal{L}(\mathcal{G}^*)$ as the corresponding Lie-ification of these 2-bialgebras.

The Lie-ification procedure can be understood loosely as an "expansion", or linearization, $x \approx 1 + X$ near the identity. Indeed, we have

$$xx' - x'x \approx (1 + X)(1 + X') - (1 + X')(1 + X) \approx [X, X']$$

modulo terms of higher order. We make use of this notion on the coproduct Eq. (2.16), and also perform a skew-symmetrization, in order to define a Lie 2-algebra 2-cochain $\mathcal{L}(\Delta) = \delta = \delta_{-1} + \delta_0$ on \mathfrak{g} ,

$$\begin{aligned} \delta_{-1}(Y) &= Y_{(1)} \wedge 1 + 1 \wedge Y_{(2)}, \\ \delta_0(X) &= \left[X_{(1)}^l - X_{(2)}^r \right] \wedge 1 + 1 \wedge \left[X_{(2)}^l - X_{(1)}^r \right] \\ &\equiv X_{(1)} \wedge 1 + 1 \wedge X_{(2)}, \end{aligned} \tag{B.1}$$

where we have made use of the Sweedler notation Eq. (2.17), and the conventional notation \wedge to denote skew-symmetric tensor products. Note the skew-symmetrization $\mathcal{G}_{-1} \wedge \mathcal{G}_0$ lands as a subspace in $\mathcal{G}_{-1} \otimes \mathcal{G}_0 \oplus \mathcal{G}_0 \otimes \mathcal{G}_{-1}$.

In degree-0, we have of course also the coproduct Δ'_0 defined in Eq. (2.19). It gives rise to a Lie algebra cochain on $\mathcal{L}(\mathcal{G}_0) = \mathfrak{g}_0$ by

$$\delta'_0(X) = \bar{X}_{(1)} \wedge 1 + 1 \wedge \bar{X}_{(2)} = tX_{(1)} \wedge 1 + 1 \wedge X_{(2)},$$

where $X_{(1)}, X_{(2)}$ have been given in Eq. (B.1).

Proposition B.1. *The Lie-ification functor \mathcal{L} sends a strict 2-bialgebra (\mathcal{G}, Δ) to a Lie 2-bialgebra (\mathfrak{g}, δ) .*

Proof. Recall (\mathfrak{g}, δ) is a Lie 2-bialgebra iff δ is a Lie 2-algebra 2-cocycle [41]. Therefore it suffices to show that the 2-cochain defined in Eq. (B.1) is a 2-cocycle. This shall follow from the fact that $(\mathcal{G}, \cdot, \Delta)$ is a 2-bialgebra — namely the coproduct map Δ Eq. (2.16) satisfies Eqs. (2.22), (2.24) and (2.31).

First note that Eqs. (2.22) and (2.24) for the coproduct Δ translates directly to the conditions

$$(t \otimes 1 + 1 \otimes t)\delta_{-1} = \delta_0 \circ t, \quad (t \otimes 1 - 1 \otimes t)\delta_0 = 0$$

for the 2-cochain $\delta = \delta_{-1} + \delta_0$. Now by a direct computation using Eq. (B.1), the condition Eq. (2.31) implies

$$\begin{aligned} \delta_0[X, X'] &= \delta_0(XX') - \delta_0(X'X) \\ &= X_{(1)}X'_{(1)} \wedge 1 + 1 \wedge X_{(2)}X'_{(2)} \\ &\quad - \left(X'_{(1)}X_{(1)} \wedge 1 + 1 \wedge X'_{(2)}X_{(2)} \right) \\ &= [X_{(1)}, X'_{(1)}] \wedge 1 + 1 \wedge [X_{(2)}, X'_{(2)}] \\ &= tX_{(1)} \triangleright X'_{(1)} \wedge 1 + 1 \wedge [X_{(2)}, X'_{(2)}] \\ &= (\bar{X}_{(1)} \triangleright \otimes 1 + 1 \otimes \text{ad}_{X_{(2)}})\delta_0(X') - (\bar{X}'_{(1)} \triangleright \otimes 1 + 1 \otimes \text{ad}_{X'_{(2)}})\delta_0(X), \end{aligned}$$

where we have used the the Peiffer identity and the fact that $\bar{X}_{(1)} = tX_{(1)}$ inherited from the constraints (2.30), and

$$\begin{aligned} \delta_{-1}(X \triangleright Y) &= \delta_{-1}(X \cdot Y) - \delta_{-1}(Y \cdot X) \\ &= \bar{X}_{(1)} \cdot Y_{(1)} \wedge 1 + 1 \wedge \bar{X}_{(2)} \cdot Y_{(2)} \\ &\quad - (Y_{(1)} \cdot \bar{X}_{(1)} \wedge 1 - 1 \wedge Y_{(2)} \cdot \bar{X}_{(1)}) \\ &= (\bar{X}_{(1)} \triangleright Y_{(1)}) \wedge 1 + 1 \wedge (\bar{X}_{(2)} \triangleright Y_{(2)}) \\ &= [X_{(1)}, Y_{(1)}] \wedge 1 + 1 \wedge (X_{(2)} \triangleright Y_{(2)}) \\ &= (\text{ad}_{X_{(1)}} \otimes 1 + 1 \otimes X_{(2)} \triangleright)\delta_{-1}(Y) - (\text{ad}_{Y_{(1)}} \otimes 1 - 1 \otimes Y_{(2)})\delta_0(X), \end{aligned}$$

where $\bar{X}_{(2)} = X_{(2)}$. These are precisely the Lie 2-algebra 2-cocycle conditions for δ [41, 3]. \square

Now the characterization result in [41] states that $(\mathfrak{g}, \mathfrak{g}^*)$ form a matched pair of Lie 2-bialgebras iff δ is a Lie 2-algebra 2-cobacket on \mathfrak{g} , namely δ satisfies the 2-coJacobi identities. For the 2-cocycle $\delta = \mathcal{L}(\Delta)$ defined in Eq. (B.1), this is guaranteed precisely by coassociativity Eqs. (2.22), (2.24). We have therefore the immediate corollary:

Corollary B.1. *Suppose $(\mathcal{G}, \mathcal{G}^*)$ form a matched pair of strict 2-bialgebras. The Lie-ification functor \mathcal{L} sends a 2-quantum double $D(\mathcal{G}) = \mathcal{G} \bowtie \mathcal{G}^*$ to a 2-Manin triple $\mathfrak{d} = \mathfrak{g} \bowtie \mathfrak{g}^*[1]$.*

In other words, our construction of the 2-quantum double $D(\mathcal{G})$ admits the classical 2-Drinfel'd double as a classical limit, which directly categorifies an analogous statement between the general quantum double construction of Majid [44] and the classical Drinfel'd double [37].

B.2 The classical 2- r -matrix

Let us now turn to the classical limit of the 2- R -matrix as defined in Section 3.4. Prior to that, we first describe one of the key properties of the duality pairing on a 2-quantum double, namely its *invariance*. This is expressed by, for instance, Eq. (6.4) in the case of the coadjoint representation. For the skew-pairing $\langle \cdot, \cdot \rangle_{\text{sk}}$ forming the 2-quantum double $D(\mathcal{G}, \mathcal{G}) = \mathcal{G} \bowtie \mathcal{G}^{\text{opp}}$, however, \mathcal{G} acts on \mathcal{G}^{opp} via its underlying (opposite) 2-algebra structure, which means that the skew-pairing satisfies the invariance property

$$\langle xx', g \rangle_{\text{sk}} = -\langle x', g \cdot x \rangle_{\text{sk}}, \quad \langle x \cdot y, f \rangle_{\text{sk}} = -\langle y, fx \rangle_{\text{sk}}, \quad \langle ff', y \rangle_{\text{sk}} = -\langle f', f \cdot y \rangle_{\text{sk}}.$$

Given the adjoint action $\bar{\triangleright} = (\Upsilon, (\triangleright_0, \triangleright_{-1}))$ of \mathcal{G} on \mathcal{G}^{opp} ,

$$x \triangleright_0 g = g \cdot x, \quad x \triangleright_{-1} f = fx, \quad \Upsilon_y f = f \cdot y,$$

this invariance property translates to the following conditions on the 2- R -matrix $\mathcal{R}^{l,r}$,

$$(x \cdot \otimes 1 + 1 \otimes x \triangleright_0) \mathcal{R}^l = 0, \quad (x \cdot \otimes 1) \mathcal{R}^r + (1 \otimes x \triangleright_{-1}) \mathcal{R}^l = 0, \quad (f \cdot \otimes 1 + 1 \otimes f \triangleright_0) \mathcal{R}^r = 0.$$

Consider the first and last conditions with $x = f \in \mathcal{G}_0$. They can be rewritten equivalently as the conditions

$$(x \cdot \otimes 1) \mathcal{R}^l + (1 \otimes x \triangleright_0) \mathcal{R}^r = 0, \quad (x \cdot \otimes 1) \mathcal{R}^r + (1 \otimes x \triangleright_0) \mathcal{R}^l = 0,$$

which together with the second condition may be compactly expressed as, using the graded sum,

$$(x \bar{\triangleright} \otimes 1 + 1 \otimes x \bar{\triangleright})(\mathcal{R} + \sigma(\mathcal{R})) = 0, \quad \forall x \in \mathcal{G}_0, \quad (\text{B.2})$$

where σ is a permutation of the $\mathcal{G}_0, \mathcal{G}_{-1}$ components.

Let us now finally recover the universal classical 2- r -matrix. This is once again accomplished by taking the Lie-ification functor on the universal quantum 2- R -matrix, $\mathfrak{r} = \mathcal{L}(\mathcal{R}) \in \mathfrak{g} \otimes \mathfrak{g}$, whence

$$\mathfrak{g}_{-1} \otimes \mathfrak{g}_0 \ni \mathfrak{r}^r = \mathcal{L}(\mathcal{R}^r), \quad \mathfrak{g}_0 \otimes \mathfrak{g}_{-1} \ni \mathfrak{r}^l = \mathcal{L}(\mathcal{R}^l). \quad (\text{B.3})$$

The equivariance condition Eq. (3.27) clearly implies

$$D_t^- \mathfrak{r} = 0, \quad (\text{B.4})$$

while applying the Lie-ification functor \mathcal{L} to Eq. (B.2) gives

$$[X \otimes 1 + 1 \otimes X, \mathfrak{r} + \sigma(\mathfrak{r})] = 0, \quad X = \mathcal{L}(x) \in \mathfrak{g}_0.$$

Here, we have used the fact that the adjoint action ρ of \mathcal{G} on itself gives rise to the adjoint representation (using the graded Lie bracket) $\mathcal{L}(\bar{\triangleright}) = [\cdot, \cdot]$ of \mathfrak{g} on itself [41].

Finally, we consider the 2-Yang-Baxter equations (3.28). We sum each equation in (3.28) in the total graded complex $\mathcal{G}^{3\otimes}$, and rearrange them to the form

$$\begin{aligned} 0 &= (\mathcal{R}_{23}^r (\mathcal{R}_{13}^r \cdot_l \mathcal{R}_{12}^l) - (\mathcal{R}_{12}^l \cdot_r \mathcal{R}_{13}^r) \mathcal{R}_{23}^r) + (\mathcal{R}_{23}^l \cdot_l \mathcal{R}_{13}^r) \mathcal{R}_{12}^r - \mathcal{R}_{12}^r (\mathcal{R}_{13}^r \cdot_r \mathcal{R}_{23}^l) \\ &+ (\mathcal{R}_{23}^l (\mathcal{R}_{13}^l \cdot_r \mathcal{R}_{12}^r) - (\mathcal{R}_{12}^r \cdot_l \mathcal{R}_{13}^l) \mathcal{R}_{23}^l) + ((\mathcal{R}_{23}^r \cdot_r \mathcal{R}_{13}^l) \mathcal{R}_{12}^l - \mathcal{R}_{12}^l (\mathcal{R}_{13}^l \cdot_l \mathcal{R}_{23}^r)). \end{aligned} \quad (\text{B.5})$$

Applying the Lie-ification functor \mathcal{L} to this equation yields

$$\begin{aligned} 0 &= ([\mathfrak{r}_{13}^r, \mathfrak{r}_{12}^l] + [\mathfrak{r}_{23}^r, \mathfrak{r}_{13}^r] + [\mathfrak{r}_{23}^l, \mathfrak{r}_{12}^l])|_{rrl} + ([\mathfrak{r}_{23}^l, \mathfrak{r}_{13}^r] + [\mathfrak{r}_{23}^r, \mathfrak{r}_{12}^l] + [\mathfrak{r}_{13}^r, \mathfrak{r}_{12}^l])|_{lrr} \\ &+ ([\mathfrak{r}_{13}^l, \mathfrak{r}_{12}^r] + [\mathfrak{r}_{23}^l, \mathfrak{r}_{13}^l] + [\mathfrak{r}_{23}^r, \mathfrak{r}_{12}^r])|_{llr} + ([\mathfrak{r}_{23}^r, \mathfrak{r}_{13}^l] + [\mathfrak{r}_{23}^l, \mathfrak{r}_{12}^r] + [\mathfrak{r}_{13}^l, \mathfrak{r}_{12}^r])|_{rll}, \end{aligned}$$

where the subscripts indicate where each term came from in (B.5).

Consider the two places in which $\mathfrak{r}_{23}^l \mathfrak{r}_{12}^r$ occurs in the above. These terms take the form respectively in Sweedler notation

$$\begin{aligned} \mathfrak{r}_{23}^l \mathfrak{r}_{12}^r|_{lrr} &= \mathfrak{r}_{(1)}^r \eta_0 \otimes \mathfrak{r}_{(1)}^l \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^l \cdot \eta_{-1}, \\ \mathfrak{r}_{23}^l \mathfrak{r}_{12}^r|_{llr} &= \mathfrak{r}_{(1)}^r \cdot \eta_{-1} \otimes \mathfrak{r}_{(1)}^l \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^l \eta_0, \end{aligned}$$

where η_0, η_{-1} are the units in $\mathcal{G}_0, \mathcal{G}_{-1}$. By using the Peiffer identity and the equivariance condition (B.4)

$$(t \mathfrak{r}_{(1)}^l) \otimes \mathfrak{r}_{(2)}^l = (t \otimes 1) \mathfrak{r}^l = (1 \otimes t) \mathfrak{r}^r = \mathfrak{r}_{(1)}^r \otimes (t \mathfrak{r}_{(2)}^r),$$

we can compute that

$$\begin{aligned} \mathfrak{r}_{23}^l \mathfrak{r}_{12}^r|_{llr} &= \mathfrak{r}_{(1)}^r \cdot \eta_{-1} \otimes \mathfrak{r}_{(1)}^l \cdot (t \mathfrak{r}_{(2)}^r) \otimes \mathfrak{r}_{(2)}^l \eta_0 & \mathfrak{r}_{23}^l \mathfrak{r}_{12}^r|_{lrr} &= \mathfrak{r}_{(1)}^r \eta_0 \otimes (t \mathfrak{r}_{(1)}^l) \cdot \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^l \cdot \eta_{-1} \\ &= (t \mathfrak{r}_{(1)}^l) \cdot \eta_{-1} \otimes \mathfrak{r}_{(1)}^l \cdot \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^l \eta_0 & &= \mathfrak{r}_{(1)}^r \eta_0 \otimes \mathfrak{r}_{(1)}^r \cdot \mathfrak{r}_{(2)}^r \otimes (t \mathfrak{r}_{(2)}^l) \cdot \eta_{-1} \\ &= \mathfrak{r}_{(1)}^l \eta_{-1} \otimes \mathfrak{r}_{(1)}^l \cdot_r \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^l \eta_0 & &= \mathfrak{r}_{(1)}^r \eta_0 \otimes \mathfrak{r}_{(1)}^r \cdot_l \mathfrak{r}_{(2)}^r \otimes \mathfrak{r}_{(2)}^r \eta_{-1} \\ &= \mathfrak{r}_{23}^l \cdot_r \mathfrak{r}_{12}^l & &= \mathfrak{r}_{23}^r \cdot_l \mathfrak{r}_{12}^r \end{aligned}$$

As such, we have

$$[\mathfrak{r}_{23}^l, \mathfrak{r}_{12}^r] = [\mathfrak{r}_{23}^l, \mathfrak{r}_{12}^l] = [\mathfrak{r}_{23}^r, \mathfrak{r}_{12}^r],$$

and hence collecting all terms from the above gives

$$\begin{aligned} [\mathfrak{r}_{12}, \mathfrak{r}_{13}] &= [\mathfrak{r}_{12}^r, \mathfrak{r}_{13}^r] + [\mathfrak{r}_{12}^r, \mathfrak{r}_{13}^l] + [\mathfrak{r}_{12}^l, \mathfrak{r}_{13}^r] + [\mathfrak{r}_{12}^l, \mathfrak{r}_{13}^l] \\ [\mathfrak{r}_{13}, \mathfrak{r}_{23}] &= [\mathfrak{r}_{13}^r, \mathfrak{r}_{23}^r] + [\mathfrak{r}_{13}^r, \mathfrak{r}_{23}^l] + [\mathfrak{r}_{13}^l, \mathfrak{r}_{23}^r] + [\mathfrak{r}_{13}^l, \mathfrak{r}_{23}^l] \\ [\mathfrak{r}_{12}, \mathfrak{r}_{23}] &= [\mathfrak{r}_{12}^r, \mathfrak{r}_{23}^r] + [\mathfrak{r}_{12}^r, \mathfrak{r}_{23}^l] + [\mathfrak{r}_{12}^l, \mathfrak{r}_{23}^r] + [\mathfrak{r}_{12}^l, \mathfrak{r}_{23}^l] \end{aligned}$$

This is precisely the 2-graded classical Yang-Baxter equation of [41]

$$[[\mathfrak{r}, \mathfrak{r}]] = [\mathfrak{r}_{12}, \mathfrak{r}_{13}] + [\mathfrak{r}_{13}, \mathfrak{r}_{23}] + [\mathfrak{r}_{12}, \mathfrak{r}_{23}] = 0$$

for the expansion $\mathfrak{r} = \mathcal{L}(\mathcal{R}) = \mathfrak{r}^r + \mathfrak{r}^l$.

Theorem B.1. \mathcal{R} admits \mathfrak{r} as a classical limit: the Lie-ification functor sends the universal 2- R -matrix to a 2-graded classical r -matrix.

In other words, the "quantization" of the classical 2- r -matrix and the associated Lie 2-bialgebra \mathfrak{g} yields a universal 2- R -matrix with the associated quasitriangular 2-bialgebra \mathcal{G} , as expected from (1.1).

B.3 Weak Lie 2-bialgebras

We now prove the weak analogues of the classical limit for 2-bialgebras.

Lemma B.2. The Lie-ification functor $\mathcal{L} : \text{Alg} \rightarrow \text{Lie}$ extends to weak 2-algebras, assigning $(\mathcal{G}, \mathcal{T})$ to a weak Lie 2-algebra $(\mathfrak{g}(\mathcal{G}), \mu_3)$ where the homotopy map μ_3 is the total skew-symmetrization of \mathcal{T} .

Proof. We construct the Lie 2-algebra structure as in **Lemma B.1**. Let $U_3 = \mathcal{L} \circ \mathcal{T} \circ \mathcal{L}$ denote the induced trilinear map on $\mathcal{L}(\mathcal{G})$. We apply \mathcal{L} to the Jacobiator,

$$\begin{aligned} [X, [X', X'']] + [X', [X'', X']] + [X'', [X, X']] &= X(X'X'') - X(X''X') - (X'X'')X + (X''X')X \\ &\quad + X'(X''X) - X'(XX'') - (X''X)X' + (XX'')X' \\ &\quad + X''(XX') - X''(X'X) - (XX')X'' + (X'X)X'' \\ &= tU_3(X, X', X'') - tU_3(X, X'', X') + tU_3(X', X'', X) \\ &\quad - tU_3(X', X, X'') + tU_3(X'', X', X) - tU_3(X'', X, X') \\ &= t(U_3(X, X', X'') - U_3(X, X'', X') + U_3(X', X'', X) \\ &\quad - U_3(X', X, X'') + U_3(X'', X, X')) - U_3(X'', X', X), \end{aligned}$$

where we have used the weak 1-associativity condition for \mathcal{G} . Similarly, we have

$$\begin{aligned} X \triangleright (X' \triangleright Y) - X' \triangleright (X \triangleright Y) - [X, X'] \triangleright Y &= t(U_3(X, X', tY) - U_3(X, tY, X') + U_3(X', tY, X) \\ &\quad - U_3(X', X, tY) + U_3(tY, X, X') - U_3(tY, X', X)), \end{aligned}$$

hence if we define the total skew-symmetrization

$$\begin{aligned} \mu_3(X, X', X'') &\equiv U_3(X, X', X'') - U_3(X, X'', X') + U_3(X', X'', X) \\ &\quad - U_3(X', X, X'') + U_3(X'', X, X') - U_3(X'', X', X), \end{aligned}$$

then weak 1-associativity implies the 2-Jacobi identity on $\mathcal{L}(\mathcal{G})$.

Using the Peiffer conditions on this fact, we see that the weak bimodularity condition also implies the 2-Jacobi identity, with two tY 's inserted in U_3 instead. Similar computations show that the Hochschild 3-cocycle condition for \mathcal{T} implies the Lie 3-cocycle condition for μ_3 .

Finally, let $F : (\mathcal{G}, \mathcal{T}) \rightarrow (\mathcal{G}', \mathcal{T}')$ denote a weak 2-algebra homomorphism as defined in Eq. (4.1). By applying the Lie-ification functor and appropriately skew-symmetrizing $\mathcal{T}, \mathcal{T}'$ and the 2-algebra structure, we recover precisely the definition of a weak 2-algebra map $\mathcal{L}(F) : (\mathfrak{g}, \mu) \rightarrow (\mathfrak{g}', \mu')$ [63]. Thus \mathcal{L} is functorial. \square

Similar to the Lie 2-algebra 2-cocycle Eq. (B.1) defined from the coproduct Δ , we form the classical limit of the coassociator Δ_1 by totally skew-symmetrizing and linearizing it, such that we have the Lie cochain

$$\delta_1(X) = X_{(1)} \wedge 1 \wedge 1 - 1 \wedge X_{(2)} \wedge 1 + 1 \wedge 1 \wedge X_{(3)}, \quad X \in \mathfrak{g}_0 = \mathcal{L}(\mathcal{G}_0). \quad (\text{B.6})$$

It is not hard to see by, for instance, dualizing the computations in the proof of **Lemma B.2**, that the conditions Eqs. (4.3), (4.5) reduce to

$$\delta_{-1} \circ \delta_{-1} = \delta_1 \circ t, \quad \text{cf. Eq. (42) in [42]}$$

$$\begin{aligned}
(\delta_{-1} + \delta_0) \circ \delta_0 &= D_t \circ \delta_1, & \text{cf. Eq. (43) in [42],} \\
\delta_1 \circ \delta_0 &= \delta_{-1} \circ \delta_1, & \text{cf. Eq. (44) in [42].}
\end{aligned}$$

Let $(\mathcal{G}, \mathcal{T}, \Delta_1)$ be a weak 2-bialgebra as given in **Definition 4.3**. The conditions Eq. (4.7) translate directly to

$$\begin{aligned}
\delta_{-1}(\mu_3(X, X', X'')) &= \mu_3(\bar{X}_{(1)}, \bar{X}'_{(1)}, \bar{X}''_{(1)}) \wedge \mu_3(\bar{X}_{(2)}, \bar{X}'_{(2)}, \bar{X}''_{(2)}), \\
\delta_1([X, X']) &= [X_{(1)}, X'_{(1)}] \wedge 1 \wedge 1 - 1 \wedge [X_{(2)}, X'_{(2)}] \wedge 1 + 1 \wedge 1 \wedge [X_{(3)}, X'_{(3)}],
\end{aligned}$$

which are precisely the conditions for a *weak-Lie 2-bialgebra* $(\mathfrak{g}, \mu_3, \delta)$ [43], expressed explicitly. In other words, we have the weak version of **Proposition B.1**:

Proposition B.2. *The Lie-ification functor takes a weak 2-bialgebra $(\mathcal{G}, \mathcal{T}, \Delta)$ to a weak Lie 2-bialgebra $(\mathfrak{g}, \mu, \delta)$, with the 2-cocycle data given as in Eqs. (B.1), (B.6).*

Note that this is a general result, which does not require the skeletality assumption on \mathcal{G} . When $\mathcal{T} = 0$ and $\mu_3 = 0$, we recover the conditions for a quasi-Lie 2-bialgebra studied also in [42].

C Module 2-categories and 2-representation theory

Recall from *Remark 6.1* that the *weak* endomorphism 2-algebra $\mathfrak{E}nd(V)$ of a 2-vector space V does not fit into the 2-category $2\text{Vect}_{\text{int}}$ of Baez-Crans 2-vector spaces. The proper setting for this should be a k -linear semisimple fusion 2-category, which we denote by 2Vect , which has the property that $\text{End}_{2\text{Vect}}(V) = \mathfrak{E}nd(V)$ is an A_2 -algebra as we have described in the main text. This structure is currently under investigation by one of the authors in an upcoming work.

Nonetheless, provided that such an appropriate setting 2Vect exists, we show in this Section that $\mathfrak{E}nd(V)$ can be interpreted in the context of module 2-categories. This relates our 2-representation theory developed here to that developed elsewhere in the literature [30, 31, 48], in particular the formulation given in [32].

We first recall briefly some key aspects of a module 2-category [48, 32]. To be more concrete, let \mathcal{C} denote a semisimple (monoidal) 2-category. A **\mathcal{C} -module 2-category** is a k -linear semisimple 2-category \mathcal{D} with a \mathcal{C} action 2-functor $\triangleright : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{D}$ and a set of adjoint natural equivalences (the *associators*)

$$\alpha_{XY|A} : (X \otimes Y) \triangleright A \rightarrow X \triangleright (Y \triangleright A)$$

for each $X, Y \in \mathcal{C}$ and $A \in \mathcal{D}$, satisfying the module pentagon relations up to a possibly non-trivial *module pentagonator* 2-morphism $\pi_{XYZ|A}$. These pentagonators must satisfy *on the nose* an additional coherence condition, called the *associahedron condition*. The explicit expressions of these conditions can be found in [48, 32].

C.1 Action 2-functors

Consider a 2-bialgebra \mathcal{G} as a connected "weak cat¹-2-algebra" [40]

$$\mathcal{G} = \mathcal{G}_{-1} \otimes \mathcal{G}_0 \rightrightarrows \mathcal{G}_0 \rightrightarrows \text{pt},$$

which is a 2-category with a single object pt , 1-morphisms \mathcal{G}_0 and each 2-Hom space over \mathcal{G}_0 is a copy of \mathcal{G}_{-1} . Evaluating an action 2-functor $\triangleright : \mathcal{G} \times 2\text{Vect} \rightarrow 2\text{Vect}$ on the object V gives precisely a weak 2-representation $\rho : \mathcal{G} \rightarrow \text{End}_{2\text{Vect}}(V) = \mathfrak{E}nd(V)$ of \mathcal{G} on V , as we have defined in the main text.

Proposition C.1. *The collection $\{\rho : \mathcal{G} \rightarrow \mathfrak{E}nd(V) \mid V \in 2\text{Vect}\}$ of weak 2-representations for each $V \in 2\text{Vect}$ makes 2Vect into a k -linear **\mathcal{G} -module 2-category**.*

Proof. The k -linearity is immediate. We first reconstruct the module associator α and pentagonator π of the associated \mathcal{G} -module 2-category V by taking

$$\alpha_{x_1 x_2|V} = \varrho(\rho_0(x_1), \rho_0(x_2))(V), \quad \pi_{x_1 x_2 x_3|V} = \mathfrak{T}(\rho_0(x_1), \rho_0(x_2), \rho_0(x_3))(V) \quad (\text{C.1})$$

in terms of the data of a weak 2-representation $\rho = (\varrho, \rho_0, \rho_1)$, as well as the Hochschild 3-cocycle \mathfrak{T} . What we need to prove is the pentagon relation between α, π , as well as the associahedron condition for π .

We identify the action 2-functor \triangleright as the weak 2-representation ρ such that $x \triangleright V = \rho_0(x)V$ for each $x \in \mathcal{G}_0$. An arrow $x \triangleright V \rightarrow x' \triangleright V$ is therefore expressed as $\rho_1(y)V$, where $y \in \mathcal{G}_{-1}$ is interpreted as a 2-morphism

$x \xrightarrow{y} x'$ between $x, x' = x + ty$ [40, 30], or simply by $\rho_1(y)$. The pentagon relation can therefore be written in the following way:

$$\begin{array}{ccccc}
& & (x_1(x_2x_3)) \triangleright V & & \\
& \nearrow^{\rho_1(\mathcal{T}(x_1, x_2, x_3))} & & \searrow^{\varrho(x_1, x_2, x_3)} & \\
((x_1x_2)x_3) \triangleright V & & \xrightarrow{\pi_{x_1x_2x_3|V}} & & x_1 \triangleright ((x_2x_3) \triangleright V) \\
& \searrow^{\varrho(x_1x_2, x_3)} & & \nearrow^{\rho_0(x_1)\varrho(x_2, x_3)} & \\
& (x_1x_2) \triangleright (x_3 \triangleright V) & \xrightarrow{\varrho(x_1, x_2)\rho_0(x_3)} & & x_1 \triangleright (x_2 \triangleright (x_3 \triangleright V))
\end{array}$$

Rewriting π in terms of the 3-cocycle \mathfrak{T} , we have

$$\begin{aligned}
\mathfrak{T}(\rho_0(x_1), \rho_0(x_2), \rho_0(x_3)) &= -\varrho(x_1x_2, x_3) - \varrho(x_1, x_2)\rho_0(x_3) \\
&\quad + \rho_1(\mathcal{T}(x_1, x_2, x_3)) + \varrho(x_1, x_2x_3) + \rho_0(x_1)\varrho(x_2, x_3),
\end{aligned}$$

which is nothing but the last equation of (4.1). It is then easy to see that the associahedron condition follows from the Hochschild 3-cocycle condition for $\mathcal{T} \circ \rho_0$. \square

Clearly, the collection $\{(V, \rho) \mid V \in 2\text{Vect}, \rho : \mathcal{G} \rightarrow \mathfrak{Cnd}(V)\}$ by **Definition 6.3** constitutes the objects of the 2-representation 2-category $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ that we have been using in the main text. On the other hand, the objects of the 2-representation 2-category 2Rep_{G_0} of an *ordinary* group G_0 as defined in [32] carry the action 2-functor $\triangleright : G_0 \times 2\text{Vect} \rightarrow 2\text{Vect}$, whose pentagonator π is induced by a group 3-cocycle $\lambda \in H^3(G_0, U(1))$.

We may then recover the formulation of 2Rep_{G_0} — at least at the level of the objects — from our formulation by taking the 2-group algebra $\mathcal{G} = kG$ of the 2-group $G = U(1) \xrightarrow{0} G_0$, considered as a 2-groupoid [31, 30]. In the following, we shall consider the 1- and 2-morphisms of this 2-category and prove that we in fact recover the entire 2-representation 2-category 2Rep_G of skeletal 2-groups G .

2-intertwiners. Recall the notion of weak 2-intertwiners that we have given in **Definition 6.4**. By treating V as a \mathcal{G} -module 2-category and taking $\triangleright, \triangleright'$ as the action 2-functors corresponding to the 2-representations ρ, ρ' , we equivalently characterize the cochain homotopy I as a collection of invertible natural transformations $I_{\bullet, i} : i(\bullet \triangleright V) \Rightarrow \bullet \triangleright' i(V)$, such that the following pentagon relation

$$\begin{array}{ccccc}
& & i(\rho_0(xx')V) & & \\
& \swarrow^{\text{id}_i \otimes \varrho(x, x')} & & \searrow^{I_{xx', I}} & \\
i(\rho_0(x)\rho_0(x')V) & & & & \rho'_0(xx') \circ i(V) \\
& \searrow^{I_{x, i} \otimes \text{id}_{\rho_0(x')}} & & \nearrow^{\varrho'(x, x') \otimes \text{id}_i} & \\
& \rho'_0(x) \circ i(\rho_0(x')V) & \xrightarrow{\text{id}_{\rho'_0(x)} \otimes I_{x', i}} & & \rho'_0(x)\rho'_0(x') \circ i(V)
\end{array}$$

follows directly from (6.6) This recovers precisely the notion of a 2-intertwiner as given in [32]. Notice no pentagonator appears here, as this is a relation on the 2-morphisms in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ and hence a pentagonator for it would have to be a 3-morphism.

Modifications. Now let us consider the notion of modifications in $2\text{Rep}^{\mathcal{T}}(\mathcal{G})$ we have defined in **Definition 6.5**. The condition (6.9) is equivalent to the composition of 2-morphisms $(\text{id}_{\rho_0(x)} \mu) \circ I_{x, i} = I_{x, i'} \circ \mu$, which is exactly a module natural transformation as given in [32].

These correspondences are not coincidences. We will prove in the following that we in fact recover the very notion of 2-representations in the skeletal and finite 2-group case, as developed in the literature [48, 32, 31, 30].

C.2 Higher-representation theory

For this section, we specialize to the case of a *finite* skeletal 2-group $G = G_{-1} \xrightarrow{1} G_0$, where $1 = \eta_0$ denotes the group unit in G_0 , and take the corresponding 2-group algebra kG via part c) of **Example 2.1.2**.

Theorem C.1. *Let kG denote the 2-group algebra of a skeletal 2-group with trivial Postnikov class $\tau = 0$ (and hence trivial Hochschild 3-cocycle $\mathcal{T} = 0$ on kG). The 2-category $2\text{Rep}(kG)$ of weak 2-representations defined here coincides with the 2-category of 2-representations 2Rep_G of G as studied in [32].*

Proof. We shall consider the skeletal 2-group G as a connected 2-groupoid, denoted conventionally by $[G_{-1}, G_0, \text{pt}]$. We first recall that a 2-representation $\rho \in 2\text{Rep}_G$, in the sense of [32] in particular, is a 2-functor $\rho : G \rightarrow 2\text{Vect}$ consisting of

- a 2-vector space $\rho(\text{pt}) = V$,
- a 2-endomorphism $\rho(x) \in \text{End}(V)$ for each $x \in G_0$, and
- a natural transformation $\rho(y) : \rho(x) \Rightarrow \rho(x)$ for each $y \in G_{-1}$, where $x \in G_0$ is the source/target of y .

Notice that this is precisely the data of a 2-representation $\rho : kG \rightarrow \text{End}(V)$ as defined in **Definition 6.3**; in particular, the cochain homotopies $\rho_1(y)$ precisely play the role of natural transformations/2-morphisms on V as a category.

Now since $t = 1$ and by definition $\varrho(x, 1) = \varrho(1, x) = 0$, the left-bimodule structure in particular is respected $\rho_1(x \cdot y) = \rho_0(x)\rho_1(y)$ from (6.2). This implies the condition

$$\rho(x \triangleright y)_v = \rho(y)_{\rho(x)v}, \quad \forall x \in G_0, v \in V_0$$

for each $y \in G_{-1}$, as the left-bimodule action \cdot coincides with \triangleright by construction (2.15).

However, in [32] we also have the following data:

1. The composition of elements $x \in G_0$ is preserved only up to an invertible natural transformation

$$p_{x_1, x_2} : \rho(x_1) \circ \rho(x_2) \xrightarrow{\sim} \rho(x_1 x_2),$$

satisfying

$$p_{x_1, x_2 x_3} * (\text{id}_{\rho(x_1)} \circ p_{x_2, x_3}) = p_{x_1 x_2, x_3} * (p_{x_1, x_2} \circ \text{id}_{\rho(x_3)}), \quad (\text{C.2})$$

where $*$ is the composition of 2-morphisms.

2. A 1-morphism $i : \rho \rightarrow \rho'$ between two 2-representations assigns an object $i_{\text{pt}} \in \text{Fun}(\rho(\text{pt}), \rho'(\text{pt}))$ and an invertible natural transformation $i_x : i_{\text{pt}} \circ \rho(x) \xrightarrow{\sim} \rho'(x) \circ i_{\text{pt}}$ to each $x \in G_0$ satisfying

$$(p_{x_1, x_2} \circ \text{id}_{i_{\text{pt}}}) * (\text{id}_{\rho(x_1)} \circ i_{x_2}) * (i_{x_1} \circ \text{id}_{\rho(x_2)}) = i_{x_1 x_2} * (\text{id}_{i_{\text{pt}}} \circ p_{x_1, x_2}), \quad (\text{C.3})$$

as well as the naturality condition

$$i_x * (\text{id}_{i_{\text{pt}}} \circ \rho(y)) = (\rho'(y) \circ \text{id}_{i_{\text{pt}}}) * i_x \quad (\text{C.4})$$

for each $x \in G_0, y \in G_{-1}$.

3. A 2-morphism $\mu : i \Rightarrow i'$ between two 1-morphisms assigns a natural transformation $\mu_* \in \text{Fun}(i_{\text{pt}}, i'_{\text{pt}})$ satisfying

$$(\text{id}_{\rho(x)} \circ \mu_*) * i_x = i'_x * \mu_*. \quad (\text{C.5})$$

Now let $\rho = (\varrho, \rho_0, \rho_1)$ denote a weak 2-representation as defined in **Definition 6.3**. The theorem then follows from the definition once we note the following points.

- We identify the invertible natural transformation p_{x_1, x_2} with $\varrho(\rho_0(x_1), \rho_0(x_2))$ for each $x_1, x_2 \in G_0$; as $\tau = 0$ is trivial, the Hochschild 3-cocycles $\mathcal{T}, \mathfrak{T}$ are both trivial, whence last equation of (4.1) implies (C.2).
- Take $\rho(*) = V$ and $\rho'(*) = W$, a 1-morphisms i_{pt} clearly denote a cochain map $i : V \rightarrow W$. We identify the invertible natural transformation i_x with the cochain homotopy $I_{x, i}$ defined in **Definition 6.4**, whence (C.3) is equivalent to (6.6). Moreover, as G is skeletal $t = 1$, the 2-morphisms $\text{id}_i \circ \rho(y), \rho(y) \circ \text{id}_i$ are self-modifications $\mu : i \Rightarrow i$ on $i = i_{\text{pt}}$, whence (C.4) follows from (6.9).
- We identify the 2-morphism μ_* with a modification μ as defined in **Definition 6.5**. (C.5) then clearly follows also from (6.9).

This completes the proof. □

As a fusion 2-category, 2Rep_G is endowed with the commutative tensor product [48, 32]

$$(\rho \otimes \rho')(*) = \rho(*) \otimes \rho'(*), \quad (\rho \otimes \rho')(x) = \rho(x) \otimes \rho'(x), \quad x \in G_0$$

which corresponds to the case where the coproduct on $\mathcal{G} = kG$ is grouplike cocommutative (see for instance Section 2.2.2) in our framework.

References

- [1] E. Witten, *(2+1)-Dimensional Gravity as an Exactly Soluble System*, *Nucl. Phys.* **B311** (1988) 46.
- [2] P. K. Osei and B. J. Schroers, *Classical r -matrices via semidualisation*, *J. Math. Phys.* **54** (2013) 101702, [[1307.6485](#)].
- [3] H. Chen and F. Girelli, *Categorified drinfel'd double and bf theory: 2-groups in 4d*, *Phys. Rev. D* **106** (Nov, 2022) 105017.
- [4] C. Delcamp, B. Dittrich and A. Riello, *Fusion basis for lattice gauge theory and loop quantum gravity*, *JHEP* **02** (2017) 061, [[1607.08881](#)].
- [5] M. Dupuis, L. Freidel, F. Girelli, A. Osumanu and J. Rennert, *On the origin of the quantum group symmetry in 3d quantum gravity*, [[2006.10105](#)].
- [6] S. Majid, *Tannaka-Krein theorem for quasiHopf algebras and other results*, .
- [7] S. Majid, *Foundations of quantum group theory*. Cambridge University Press, 2011.
- [8] E. Witten, *Gauge theories, vertex models, and quantum groups*, *Nuclear Physics B* **330** (1990) 285–346.
- [9] H. Hopf, *Über die topologie der gruppen-mannigfaltigkeiten und ihre verallgemeinerungen*, *Annals of Mathematics* **42** (1941) 22–52.
- [10] A. Borel, *Sur la cohomologie des espaces fibres principaux et des espaces homogenes de groupes de lie compacts*, *Annals of Mathematics* **57** (1953) 115–207.
- [11] A. Yu. Kitaev, *Fault tolerant quantum computation by anyons*, *Annals Phys.* **303** (2003) 2–30, [[quant-ph/9707021](#)].
- [12] A. Kitaev and L. Kong, *Models for gapped boundaries and domain walls*, *Communications in Mathematical Physics* **313** (jun, 2012) 351–373.
- [13] X. Chen, Z. C. Gu and X. G. Wen, *Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order*, *Phys. Rev. B* **82** (2010) 155138, [[1004.3835](#)].
- [14] L. Kong and X.-G. Wen, *Braided fusion categories, gravitational anomalies, and the mathematical framework for topological orders in any dimensions*, [[1405.5858](#)].
- [15] C. Tomei, *The toda lattice, old and new*, *Journal of Geometric Mechanics* **5** (2013) 511–530.
- [16] M. Adler, *On a Trace Functional for Formal Pseudo-Differential Operators and the Symplectic Structure of the Korteweg-Devries Type Equation.*, *Inventiones Mathematicae* **50** (Jan., 1978) 219.
- [17] D. B. Olivier Babelon and M. Talon, *Introduction to Classical Integrable Systems*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2003, [10.1017/CBO9780511535024](#).
- [18] C. Meusburger, *Poisson–Lie Groups and Gauge Theory*, *Symmetry* **13** (2021) 1324.
- [19] M. Q. Zhang, *How to find the Lax pair from the Yang-Baxter equation*, *Communications in Mathematical Physics* **141** (1991) 523 – 531.
- [20] A. Bullivant, M. Calcada, Z. Kadar, P. Martin and J. Martins, *Topological phases from higher gauge symmetry in 3+1 dimensions*, *Phys. Rev.* **B95** (2017) 155118, [[1606.06639](#)].
- [21] T. Lan and X.-G. Wen, *Classification of 3 + 1D bosonic topological orders (ii): The case when some pointlike excitations are fermions*, *Phys. Rev. X* **9** (Apr, 2019) 021005.
- [22] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang and H. Zheng, *Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry*, *Phys. Rev. Research* **2** (Oct, 2020) 043086.
- [23] A. Hamma, P. Zanardi and X. G. Wen, *String and membrane condensation on 3-D lattices*, *Phys. Rev. B* **72** (2005) 035307, [[cond-mat/0411752](#)].
- [24] L. Kong, Y. Tian and Z.-H. Zhang, *Defects in the 3-dimensional toric code model form a braided fusion 2-category*, *JHEP* **12** (2020) 078, [[2009.06564](#)].
- [25] D. V. Else and C. Nayak, *Cheshire charge in (3+1)-dimensional topological phases*, *Phys. Rev. B* **96** (2017) 045136, [[1702.02148](#)].

- [26] L. Crane and I. Frenkel, *Four-dimensional topological field theory, Hopf categories, and the canonical bases*, *J. Math. Phys.* **35** (1994) 5136–5154, [[hep-th/9405183](#)].
- [27] J. C. Baez and J. Dolan, *Higher dimensional algebra and topological quantum field theory*, *J. Math. Phys.* **36** (1995) 6073–6105, [[q-alg/9503002](#)].
- [28] M. Mackaay, *Finite groups, spherical 2-categories, and 4-manifold invariants*, *Adv. Math.* **153**(2) (2000), 353–390 (2000) , [[math/9903003](#)].
- [29] M. Neuchl, *Representation Theory of Hopf Categories*. Verlag nicht ermittelbar, 1997.
- [30] J. Baez, A. Baratin, L. Freidel and D. Wise, *Infinite-dimensional representations of 2-groups*, *Memoirs of the American Mathematical Society* **219** (2012) 0–0.
- [31] C. L. Douglas and D. J. Reutter, *Fusion 2-categories and a state-sum invariant for 4-manifolds*, 2018. 10.48550/ARXIV.1812.11933.
- [32] C. Delcamp and A. Tiwari, *Higher categorical symmetries and gauging in two-dimensional spin systems*, [[2301.01259](#)].
- [33] J. C. Baez and A. S. Crans, *Higher-Dimensional Algebra VI: Lie 2-Algebras*, *Theor. Appl. Categor.* **12** (2004) 492–528, [[math/0307263](#)].
- [34] J. Baez and U. Schreiber, *Higher gauge theory: 2-connections on 2-bundles*, [[hep-th/0412325](#)].
- [35] T. Nikolaus, U. Schreiber and D. Stevenson, *Principal ∞ -bundles: presentations*, *Journal of Homotopy and Related Structures* **10** (mar, 2014) 565–622.
- [36] J. M. T. Thompson and S. Majid, *Non-commutative geometry and quantum groups*, *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **358** (2000) 89–109, [<https://royalsocietypublishing.org/doi/pdf/10.1098/rsta.2000.0521>].
- [37] M. A. Semenov-Tyan-Shanskii, *Poisson-Lie groups. The quantum duality principle and the twisted quantum double*, *Theoretical and Mathematical Physics* **93** (Nov, 1992) 1292–1307.
- [38] V. G. Drinfeld, *Quantum groups*, *Zap. Nauchn. Semin.* **155** (1986) 18–49.
- [39] M. Kontsevich, *Deformation quantization of Poisson manifolds. 1.*, *Lett. Math. Phys.* **66** (2003) 157–216, [[q-alg/9709040](#)].
- [40] F. Wagemann, *Crossed Modules*. De Gruyter, 2021, doi:10.1515/9783110750959.
- [41] C. Bai, Y. Sheng and C. Zhu, *Lie 2-bialgebras*, *Communications in Mathematical Physics* **320** (apr, 2013) 149–172.
- [42] Z. Chen, M. Stiénon and P. Xu, *Poisson 2-groups*, *J. Diff. Geom.* **94** (2013) 209–240, [[1202.0079](#)].
- [43] Z. Chen, M. Stiénon and P. Xu, *Weak lie 2-bialgebras*, *Journal of Geometry and Physics* **68** (Jun, 2013) 59–68.
- [44] S. Majid, *Some remarks on the quantum double*, *Czech. J. Phys.* **44** (1994) 1059, [[hep-th/9409056](#)].
- [45] C. Angulo, *A cohomology theory for lie 2-algebras and lie 2-groups*, 2018. 10.48550/arxiv.1810.05740.
- [46] N. Gurski, *Loop spaces, and coherence for monoidal and braided monoidal bicategories*, *Advances in Mathematics* **226** (2011) 4225–4265.
- [47] L. Kong, Y. Tian and S. Zhou, *The center of monoidal 2-categories in 3+1d dijkgraaf-witten theory*, *Advances in Mathematics* **360** (2020) 106928.
- [48] C. Delcamp, *Tensor network approach to electromagnetic duality in (3+1)d topological gauge models*, *JHEP* **08** (2022) 149, [[2112.08324](#)].
- [49] S. Majid, *Strict quantum 2-groups*, [[1208.6265](#)].
- [50] J. C. Baez, A. Baratin, L. Freidel and D. K. Wise, *Infinite-Dimensional Representations of 2-Groups*, vol. 1032. American Mathematical Society, 2012, [10.1090/S0065-9266-2012-00652-6](#).
- [51] C. Zhu, T. Lan and X.-G. Wen, *Topological nonlinear σ -model, higher gauge theory, and a systematic construction of (3+1)d topological orders for boson systems*, *Physical Review B* **100** (Jul, 2019) .

- [52] A. Kapustin and R. Thorngren, *Higher symmetry and gapped phases of gauge theories*, [1309.4721].
- [53] J. Ang and A. Prakash, *Higher categorical groups and the classification of topological defects and textures*, 2018.
- [54] N. T. Quang, P. T. Cuc and N. T. Thuy, *Strict gr-categories and applications on classification of extensions of groups of the type of a crossed module*, 2011. 10.48550/ARXIV.1112.2440.
- [55] H. Pfeiffer, *2-groups, trialgebras and their hopf categories of representations*, *Advances in Mathematics* **212** (2007) 62–108, [0411468].
- [56] F. Wagemann, *On lie algebra crossed modules*, *Communications in Algebra* **34** (2006) 1699–1722, [https://doi.org/10.1080/00927870500542705].
- [57] S. F. Siegel and S. J. Witherspoon, *The hochschild cohomology ring of a group algebra*, *Proceedings of the London Mathematical Society* **79** (1999) 131–157, [https://londmathsoc.onlinelibrary.wiley.com/doi/pdf/10.1112/S0024611599011958].
- [58] H. Chen, *2-drinfel'd double symmetry of the 4d kitaev model*, [2305.04729].
- [59] C. Bai, *A unified algebraic approach to the classical yang–baxter equation*, *Journal of Physics A: Mathematical and Theoretical* **40** (Aug, 2007) 11073–11082.
- [60] S. X. Cui and Z. Wang, *State sum invariants of three manifolds from spherical multi-fusion categories*, *Journal of Knot Theory and Its Ramifications* **26** (dec, 2017) 1750104.
- [61] J. C. Baez and J. Huerta, *An Invitation to Higher Gauge Theory*, *Gen. Rel. Grav.* **43** (2011) 2335–2392, [1003.4485].
- [62] R. D. Schafer, *Structure and representation of nonassociative algebras*, *Bulletin of the American Mathematical Society* **61** (1955) 469–484.
- [63] J. C. Baez, D. Stevenson, A. S. Crans and U. Schreiber, *From loop groups to 2-groups*, [math/0504123].
- [64] J. C. Baez and D. K. Wise, *Teleparallel Gravity as a Higher Gauge Theory*, *Commun. Math. Phys.* **333** (2015) 153–186, [1204.4339].
- [65] M. Gurski, *An Algebraic Theory of Tricategories*. University of Chicago, Department of Mathematics, 2006.
- [66] S. Caenepeel, D. Bulacu, F. Panaite and F. Van Oystaeyen, *Quasi-Hopf Algebras: A Categorical Approach*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2019.
- [67] A. Joyal and R. Street, *Braided tensor categories*, *Advances in Mathematics* **102** (Nov., 1993) 20–78.
- [68] T. Johnson-Freyd, *(3+1)D topological orders with only a \mathbb{Z}_2 -charged particle*, [2011.11165].
- [69] V. Turaev and A. Virelizier, *Monoidal Categories and Topological Field Theory*, vol. 322 of *Progress in Mathematics*. Birkhauser, Cham, 2017, 10.1007/978-3-319-49834-8.
- [70] J.-H. LU, *Hopf algebroids and quantum groupoids*, *International Journal of Mathematics* **07** (1996) 47–70, [https://doi.org/10.1142/S0129167X96000050].
- [71] M. M. Kapranov and V. A. Voevodsky, *2-categories and Zamolodchikov tetrahedra equations*, vol. 56 of *Proc. Sympos. Pure Math.*, p. 177–259. Amer. Math. Soc., Providence, RI, 1994.
- [72] V. G. Turaev and O. Y. Viro, *State sum invariants of 3 manifolds and quantum 6j symbols*, *Topology* **31** (1992) 865–902.
- [73] J. W. Barrett and B. W. Westbury, *Invariants of piecewise linear three manifolds*, *Trans. Am. Math. Soc.* **348** (1996) 3997–4022, [hep-th/9311155].
- [74] N. Reshetikhin and V. G. Turaev, *Invariants of three manifolds via link polynomials and quantum groups*, *Invent. Math.* **103** (1991) 547–597.
- [75] M. Mackaay and R. Picken, *2-categories, 4d state-sum models and gerbes*, [math/0104285].
- [76] M. Mackaay, *Spherical 2-categories and 4-manifold invariants*, *Adv. Math.* **143** (1999), 288–348 (1999) , [math/9805030].