

A REFINED BLOCH-WIGNER EXACT SEQUENCE IN CHARACTERISTIC 2

BEHROOZ MIRZAI AND ELVIS TORRES PÉREZ

ABSTRACT. Let A be a local domain of characteristic 2 such that its residue field has more than 64 elements. Then we find an exact relation between the third integral homology of the group $\mathrm{SL}_2(A)$ and Hutchinson's refined Bloch group $\mathcal{RB}(A)$.

INTRODUCTION

The classical Bloch-Wigner exact sequence studies the indecomposable part of the third K -group of a field [4]. The general Bloch-Wigner exact sequence for fields, claims that for any field F we have the exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\mu(F), \mu(F))^{\sim} \rightarrow K_3^{\mathrm{ind}}(F) \rightarrow \mathcal{B}(F) \rightarrow 0.$$

Here $\mathcal{B}(F)$, called the Bloch group of F , is a certain subgroup of the scissors congruence group $\mathcal{P}(A)$ (see Section 1) and $\mathrm{Tor}_1^{\mathbb{Z}}(\mu(F), \mu(F))^{\sim}$ is the unique nontrivial extension of $\mu_2(F)$ by $\mathrm{Tor}_1^{\mathbb{Z}}(\mu(F), \mu(F))$ [18], [6]. This exact sequence can be extended to local domains, where its residue field has more than 9 elements [16].

When F is quadratically closed, we have a natural isomorphism $H_3(\mathrm{SL}_2(F), \mathbb{Z}) \simeq K_3^{\mathrm{ind}}(F)$ [14],[17]. In general we have a natural surjective map

$$H_3(\mathrm{SL}_2(F), \mathbb{Z}) \twoheadrightarrow K_3^{\mathrm{ind}}(F)$$

(see [10]). The indecomposable group $K_3^{\mathrm{ind}}(F)$ has been studied extensively in the literature (see for example [13] or [12]). In many applications in algebraic K -theory and number theory it is important to understand the structure of the group $H_3(\mathrm{SL}_2(F), \mathbb{Z})$. When F is not quadratically closed, the above map has a nontrivial, and often quite large, kernel (see [6], [9]).

In this paper we prove a refined version of the Bloch-Wigner exact sequence over domains of characteristic 2 (see Theorem 6.1 for the general statement). More precisely we show that for any local domain of characteristic 2, where its residue field has more than 64 elements, we have an exact sequence

$$0 \rightarrow \mathrm{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)) \rightarrow H_3(\mathrm{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

This gives a positive answer to a question raised by Coronado and Hutchinson in [3] over such rings and it improves similar results of Hutchinson (see [5],[7]), as it leaves no ambiguity on 2-torsion elements.

The homology groups $H_\bullet(\mathrm{SL}_2(A), \mathbb{Z})$ are naturally $\mathcal{R}_A := \mathbb{Z}[A^\times / (A^\times)^2]$ -modules and this module structure plays a central role in the study of the homology groups $H_3(\mathrm{SL}_2(A), \mathbb{Z})$.

The refined Bloch group of a ring A , introduced by Hutchinson, is a certain subgroup of the the refined scissors congruence group of a A . The refined scissors congruence group $\mathcal{RP}_1(A)$ of A is defined by a presentation analogous to $\mathcal{P}(A)$ but as a module over the group ring \mathcal{R}_A rather than as an abelian group.

In a series of papers [6], [5], [7], [8], [9], Hutchinson extensively studied the homology group $H_3(\mathrm{SL}_2(A), \mathbb{Z})$ when A is a field or a local ring with sufficiently large residue field. Recently it has been proved that the third homology of SL_2 over discrete valuation rings satisfies certain localisation property [11].

Our main result follows from a careful analysis of a spectral sequence which converge to the homology of SL_2 . Our spectral sequence is a variant of a spectral sequence which is studied by Hutchinson in his series of papers and is similar to the one studies for GL_2 in [15]. This variant has certain advantage when it comes to calculation of some differentials.

Notations. In this note all rings are commutative, except probably group rings, and have the unit element 1. For a ring A , let \mathcal{W}_A be the set of $a \in A^\times$ such that $1 - a \in A^\times$. Thus

$$\mathcal{W}_A := \{a \in A : a(1 - a) \in A^\times\}.$$

Let $\mathcal{G}_A := A^\times / (A^\times)^2$ and set $\mathcal{R}_A := \mathbb{Z}[\mathcal{G}_A]$. The element of \mathcal{G}_A represented by $a \in A^\times$ is denoted by $\langle a \rangle$. We set $\langle\langle a \rangle\rangle := \langle a \rangle - 1 \in \mathcal{R}_A$.

1. THE SCISSORS CONGRUENCE GROUPS

Let A be a ring. The *scissors congruence group* $\mathcal{P}(A)$ of A is defined as the quotient of the free abelian group generated by symbols $[a]$, $a \in \mathcal{W}_A$, by the subgroup generated by elements

$$[a] - [b] + \left[\frac{b}{a}\right] - \left[\frac{1 - a^{-1}}{1 - b^{-1}}\right] + \left[\frac{1 - a}{1 - b}\right],$$

where $a, b, a/b \in \mathcal{W}_A$. Let

$$S_{\mathbb{Z}}^2(A^\times) := (A^\times \otimes A^\times) / \langle a \otimes b + b \otimes a : a, b \in A^\times \rangle.$$

The map

$$\lambda : \mathcal{P}(A) \rightarrow S_{\mathbb{Z}}^2(A^\times), \quad [a] \mapsto a \otimes (1 - a),$$

is a well-defined homomorphism and its kernel is called the *Bloch group* of A (see [4], [18], [15]). We denote this kernel by $\mathcal{B}(A)$.

Let $\mathcal{RP}(A)$ be the quotient of the free \mathcal{R}_A -module generated by symbols $[a]$, $a \in \mathcal{W}_A$, by the \mathcal{R}_A -submodule generated by the elements

$$[a] - [b] + \langle a \rangle \left[\frac{b}{a}\right] - \langle a^{-1} - 1 \rangle \left[\frac{1 - a^{-1}}{1 - b^{-1}}\right] + \langle 1 - a \rangle \left[\frac{1 - a}{1 - b}\right],$$

where $a, b, a/b \in \mathcal{W}_A$. We have a natural surjective map $\mathcal{RP}(A) \rightarrow \mathcal{P}(A)$ and from the definition it follows immediately that

$$\mathcal{P}(A) \simeq \mathcal{RP}(A)_{\mathcal{G}_A} = H_0(\mathcal{G}_A, \mathcal{RP}(A)).$$

Let \mathcal{I}_A be the augmentation ideal of the group ring $\mathcal{R}_A = \mathbb{Z}[\mathcal{G}_A]$. By direct computation one can show that the map

$$\lambda_1 : \mathcal{RP}(A) \rightarrow \mathcal{I}_A^2, \quad [a] \mapsto \langle\langle a \rangle\rangle \langle\langle 1 - a \rangle\rangle$$

is a well-defined \mathcal{R}_A -homomorphism. If we consider $S_{\mathbb{Z}}^2(A^\times)$ as a trivial \mathcal{G}_A -module, then

$$\lambda_2 : \mathcal{RP}(A) \rightarrow S_{\mathbb{Z}}^2(A^\times), \quad [a] \mapsto a \otimes (1 - a),$$

is a homomorphism of \mathcal{R}_A -modules. In fact λ_2 is the composite $\mathcal{RP}(A) \rightarrow \mathcal{P}(A) \xrightarrow{\lambda} S_{\mathbb{Z}}^2(A^\times)$.

Hutchinson defined the *refined scissors congruence group* of A as the \mathcal{R}_A -module

$$\mathcal{RP}_1(A) := \ker(\lambda_1 : \mathcal{RP}(A) \rightarrow \mathcal{I}_A^2).$$

He defined the *refined Bloch group* of A as the \mathcal{R}_A -module

$$\mathcal{RB}(A) := \ker(\lambda_2|_{\mathcal{RP}_1(A)} : \mathcal{RP}_1(A) \rightarrow S_{\mathbb{Z}}^2(A^\times))$$

(see [5, page 28], [6, Subsection 2.3]).

2. THE COMPLEX OF UNIMODULAR VECTORS

The bridge between the third homology of $\mathrm{SL}_2(A)$ and the refined scissors congruence group usually is given by a complex built out of unimodular vectors.

Let A be a commutative ring. A (column) vector $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \in A^2$ is said to be unimodular if $u_1A + u_2A = A$. Equivalently, $\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$ is said to be unimodular if there exists a vector $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$ such that $\{\mathbf{u}, \mathbf{v}\}$ is a basis of A^2 .

For any non-negative integer n , let $X_n(A^2)$ be the free abelian group generated by the set of all $(n+1)$ -tuples $(\langle \mathbf{v}_0 \rangle, \dots, \langle \mathbf{v}_n \rangle)$, where every $\mathbf{v}_i \in A^2$ is unimodular and any two distinct vectors $\mathbf{v}_i, \mathbf{v}_j$ are a basis of A^2 . Observe that $\langle \mathbf{v} \rangle \subseteq A^2$ is the line generated by $\mathbf{v} \in A^2$, i.e. $\langle \mathbf{v} \rangle = \mathbf{v}A$.

We consider $X_n(A^2)$ as a left $\mathrm{GL}_2(A)$ -module (respectively left $\mathrm{SL}_2(A)$ -module) in a natural way. If necessary, we convert this action to a right action by the definition $m.g := g^{-1}m$. Let us define the n -th differential operator

$$\partial_n : X_n(A^2) \rightarrow X_{n-1}(A^2), \quad n \geq 1,$$

as an alternating sum of face operators which throws away the i -th component of generators. Let $\partial_{-1} = \epsilon : X_0(A^2) \rightarrow \mathbb{Z}$ be defined by $\sum_i n_i \langle \mathbf{v}_{0,i} \rangle \mapsto \sum_i n_i$. Hence we have the complex

$$X_\bullet(A^2) \xrightarrow{\epsilon} \mathbb{Z} : \dots \rightarrow X_2(A^2) \xrightarrow{\partial_2} X_1(A^2) \xrightarrow{\partial_1} X_0(A^2) \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0.$$

The following is an important case that we should keep in our mind.

Proposition 2.1 (Hutchinson). *Let A be a local ring. If $|A/\mathfrak{m}_A| = k$, then the complex $X_\bullet(A^2) \rightarrow \mathbb{Z}$ is exact in dimension $< k$.*

Proof. See [7, Lemma 3.21]. □

3. THE MAIN SPECTRAL SEQUENCE

Let $Z_i(A^2) = \ker(\partial_i)$. Assume that $X_\bullet(A^2) \rightarrow \mathbb{Z}$ is exact in dimension < 2 . Consider the complex

$$0 \rightarrow Z_1(A^2) \xrightarrow{\text{inc}} X_1(A^2) \xrightarrow{\partial_1} X_0(A^2) \rightarrow 0.$$

Let $F_\bullet \rightarrow \mathbb{Z}$ be a projective resolution of \mathbb{Z} over $\text{SL}_2(A)$. Let $D_{\bullet,\bullet}$ be the double complex

$$D_{\bullet,\bullet} : 0 \rightarrow F_\bullet \otimes_{\text{SL}_2(A)} Z_1(A^2) \xrightarrow{\text{id}_{F_\bullet} \otimes \text{inc}} F_\bullet \otimes_{\text{SL}_2(A)} X_1(A^2) \xrightarrow{\text{id}_{F_\bullet} \otimes \partial_1} F_\bullet \otimes_{\text{SL}_2(A)} X_0(A^2) \rightarrow 0.$$

In our calculations we usually use either the bar resolution $B_\bullet(\text{SL}_2(A)) \rightarrow \mathbb{Z}$ or the standard resolution $C_\bullet(\text{SL}_2(A)) \rightarrow \mathbb{Z}$ [1]. Note that we have the natural morphisms

$$B_\bullet(\text{SL}_2(A)) \rightarrow C_\bullet(\text{SL}_2(A)), \quad [h_1|h_2|\cdots|h_n] \mapsto (1, h_1, h_1h_2, \dots, h_1h_2\cdots h_n),$$

$$C_\bullet(\text{SL}_2(A)) \rightarrow B_\bullet(\text{SL}_2(A)), \quad (g_0, \dots, g_n) \mapsto g_0[g_0^{-1}g_1|g_1^{-1}g_2|\cdots|g_{n-1}^{-1}g_n],$$

which induce the identity on the homology of $\text{SL}_2(A)$ with any coefficient [1].

From the double complex $D_{\bullet,\bullet}$ we obtain the first quadrant spectral sequence

$$E_{p,q}^1 = \begin{cases} H_q(\text{SL}_2(A), X_p(A^2)) & p = 0, 1 \\ H_q(\text{SL}_2(A), Z_1(A^2)) & p = 2 \\ 0 & p > 2 \end{cases} \implies H_{p+q}(\text{SL}_2(A), \mathbb{Z}).$$

The group $\text{SL}_2(A)$ acts transitively on the sets of generators of $X_i(A^2)$ for $i = 0, 1$. Let

$$\infty := \langle \mathbf{e}_1 \rangle, \quad \mathbf{0} := \langle \mathbf{e}_2 \rangle, \quad \mathbf{a} := \langle \mathbf{e}_1 + a\mathbf{e}_2 \rangle, \quad a \in A^\times.$$

We choose (∞) and $(\infty, \mathbf{0})$ as representatives of the orbit of the generators of $X_0(A^2)$ and $X_1(A^2)$ respectively. Therefore

$$X_0 \simeq \text{Ind}_{B(A)}^{\text{SL}_2(A)} \mathbb{Z}, \quad X_1 \simeq \text{Ind}_{T(A)}^{\text{SL}_2(A)} \mathbb{Z},$$

where

$$B(A) := \text{Stab}_{\text{SL}_2(A)}(\infty) = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a \in A^\times, b \in A \right\},$$

$$T(A) := \text{Stab}_{\text{SL}_2(A)}(\infty, \mathbf{0}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} : a \in A^\times \right\}.$$

Note that $T(A) \simeq A^\times$. In our calculations usually we identify $T(A)$ with A^\times . By Shapiro's lemma we have

$$E_{0,q}^1 \simeq H_q(B(A), \mathbb{Z}), \quad E_{1,q}^1 \simeq H_q(T(A), \mathbb{Z}).$$

In particular $E_{0,0}^1 \simeq \mathbb{Z} \simeq E_{1,0}^1$. Moreover

$$d_{1,q}^1 = H_q(\sigma) - H_q(\text{inc}),$$

where $\sigma : T(A) \rightarrow B(A)$ is given by $\sigma(X) = wXw^{-1}$, $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

These implies that $d_{1,0}^1$ is trivial, $d_{1,1}^1$ is induced by the map $T(A) \rightarrow B(A)$ given by $X \mapsto X^{-2}$ and $d_{1,2}^1$ is trivial. Thus

$$\ker(d_{1,1}^1) \simeq \mu_2(A) = \{b \in A^\times : b^2 = 1\}.$$

It is straightforward to check that for any $b \in \mu_2(A)$,

$$d_{2,1}^1([b] \otimes \partial_2(\infty, \mathbf{0}, \mathbf{a})) = b.$$

Therefore $E_{1,1}^2 = 0$.

4. THE REFINED SCISSORS CONGRUENCE GROUP

Assume that the complex $X_\bullet(A^2) \rightarrow \mathbb{Z}$ is exact in dimension < 4 . From the short exact sequence

$$0 \rightarrow Z_2(A^2) \rightarrow X_2(A^2) \rightarrow Z_1(A^2) \rightarrow 0$$

we obtain the long exact sequence

$$(4.1) \quad H_1(\mathrm{SL}_2(A), X_2(A^2)) \rightarrow H_1(\mathrm{SL}_2(A), Z_1(A^2)) \xrightarrow{\delta} H_0(\mathrm{SL}_2(A), Z_2(A^2)) \\ \rightarrow H_0(\mathrm{SL}_2(A), X_2(A^2)) \rightarrow H_0(\mathrm{SL}_2(A), Z_1(A^2)) \rightarrow 0.$$

Choose $(\infty, \mathbf{0}, \mathbf{a})$, $\langle a \rangle \in \mathcal{G}_A$, as representatives of the orbits of the generators of $X_2(A^2)$. Then

$$X_2 \simeq \bigoplus_{\langle a \rangle \in \mathcal{G}_A} \mathrm{Ind}_{\mu_2(A)}^{\mathrm{SL}_2(A)} \mathbb{Z}\langle a \rangle,$$

where $\mu_2(A) = \mathrm{Stab}_{\mathrm{SL}_2(A)}(\infty, \mathbf{0}, \mathbf{a})$. Thus

$$H_q(\mathrm{SL}_2(A), X_2(A^2)) \simeq \bigoplus_{\langle a \rangle \in \mathcal{G}_A} H_q(\mu_2(A), \mathbb{Z}) \simeq \mathcal{R}_A \otimes H_q(\mu_2(A), \mathbb{Z}).$$

From the exact sequence $X_4(A^2) \xrightarrow{\partial_4} X_3(A^2) \xrightarrow{\partial_3} Z_2(A^2) \rightarrow 0$ we obtain the exact sequence

$$X_4(A^2)_{\mathrm{SL}_2(A)} \xrightarrow{\overline{\partial}_4} X_3(A^2)_{\mathrm{SL}_2(A)} \xrightarrow{\overline{\partial}_3} Z_2(A^2)_{\mathrm{SL}_2(A)} \rightarrow 0$$

of \mathcal{R}_A -modules. The orbits of the action of $\mathrm{SL}_2(A)$ on $X_3(A)$ and $X_4(A)$ are represented by

$$\langle a \rangle[x] := (\langle \mathbf{e}_1 \rangle, \langle \mathbf{e}_2 \rangle, \langle \mathbf{e}_1 + a\mathbf{e}_2 \rangle, \langle \mathbf{e}_1 + ax\mathbf{e}_2 \rangle), \quad \langle a \rangle \in \mathcal{G}_A, x \in \mathcal{W}_A,$$

and

$$\langle a \rangle[x, y] := (\langle \mathbf{e}_1 \rangle, \langle \mathbf{e}_2 \rangle, \langle \mathbf{e}_1 + a\mathbf{e}_2 \rangle, \langle \mathbf{e}_1 + ax\mathbf{e}_2 \rangle, \langle \mathbf{e}_1 + ay\mathbf{e}_2 \rangle), \quad \langle a \rangle \in \mathcal{G}_A, x, y, x/y \in \mathcal{W}_A,$$

respectively. Thus $X_3(A^2)_{\mathrm{SL}_2(A)}$ is the free \mathcal{R}_A -module generated by the symbols $[x]$, $x \in \mathcal{W}_A$ and $X_4(A^2)_{\mathrm{SL}_2(A)}$ is the free \mathcal{R}_A -module generated by the symbols $[x, y]$, $x, y, x/y \in \mathcal{W}_A$. It is straightforward to check that

$$\overline{\partial}_4([x, y]) = [x] - [y] + \langle x \rangle \left[\frac{y}{x} \right] - \langle x^{-1} - 1 \rangle \left[\frac{1 - x^{-1}}{1 - y^{-1}} \right] - \langle 1 - x \rangle \left[\frac{1 - x}{1 - y} \right].$$

These imply that

$$H_0(\mathrm{SL}_2(A), Z_2(A^2)) \simeq \mathcal{RP}(A).$$

It is not difficult to see that the composite

$$\mathcal{RP}(A) \simeq H_0(\mathrm{SL}_2(A), Z_2(A^2)) \rightarrow H_0(\mathrm{SL}_2(A), X_2(A^2)) \simeq \mathbb{Z}[\mathcal{G}_A] = \mathcal{R}_A$$

is given by

$$[x] \mapsto \langle\langle x \rangle\rangle \langle\langle 1-x \rangle\rangle.$$

This is the map λ_1 discussed in Section 1.

we obtain the exact sequence

$$(4.2) \quad H_1(\mathrm{SL}_2(A), Z_2(A^2)) \rightarrow \mathcal{R}_A \otimes \mu_2(A) \rightarrow E_{2,1}^1 \rightarrow \mathcal{RP}_1(A) \rightarrow 0.$$

5. THE REFINED BLOCH GROUP

Let A be a ring such that $\mu_2(A) = 1$ (e.g. a domains of characteristic two). Moreover let $X_\bullet(A^2) \rightarrow \mathbb{Z}$ is exact in dimension < 4 . It follows from the exact sequence (4.2) that

$$E_{2,1}^1 = H_1(\mathrm{SL}_2(A), Z_1(A^2)) \simeq \mathcal{RP}_1(A).$$

Since $\ker(d_{1,1}) \simeq \mu_2(A) = 1$, we have $E_{2,1}^2 = E_{2,1}^1$.

The group $B(A)$ sits in the extension

$$1 \rightarrow N(A) \rightarrow B(A) \xrightarrow{\alpha} T(A) \rightarrow 1,$$

where $N(A) := \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}_2(A) \mid b \in A \right\}$ and $\alpha \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$. Note that $N(A) \simeq A$. This extension splits canonically. Let

$$H_2(B(A), \mathbb{Z}) \simeq H_2(T(A), \mathbb{Z}) \oplus \mathcal{S}_2 \simeq (A^\times \wedge A^\times) \oplus \mathcal{S}_2$$

and consider the differential

$$d_{2,1}^2 : \mathcal{RP}_1(A) \simeq E_{2,1}^2 \rightarrow H_2(B(A), \mathbb{Z}) \simeq (A^\times \wedge A^\times) \oplus \mathcal{S}_2.$$

Lemma 5.1. *The kernel of the composite $\mathcal{RP}_1(A) \xrightarrow{d_{2,1}^2} H_2(B(A), \mathbb{Z}) \xrightarrow{\alpha_*} A^\times \wedge A^\times$ is isomorphic with the refined Bloch group $\mathcal{RB}(A)$.*

Proof. From the complex $0 \rightarrow Z_1(A^2) \xrightarrow{\mathrm{inc}} X_1(A^2) \xrightarrow{\partial_1} X_0(A^2) \rightarrow 0$ we obtain the first quadrant spectral sequence

$$\mathcal{E}_{p,q}^1 = \begin{cases} H_q(\mathrm{GL}_2(A), X_p(A^2)) & p = 0, 1 \\ H_q(\mathrm{GL}_2(A), Z_1(A^2)) & p = 2 \\ 0 & p > 2 \end{cases} \implies H_{p+q}(\mathrm{GL}_2(A), \mathbb{Z}).$$

This spectral sequence have been studied in [15]. The inclusion $\mathrm{SL}_2(A) \hookrightarrow \mathrm{GL}_2(A)$ gives us the morphism of spectral sequences

$$\begin{array}{ccc} E_{p,q}^1 & \implies & H_{p+q}(\mathrm{SL}_2(A), \mathbb{Z}) \\ \downarrow & & \downarrow \\ \mathcal{E}_{p,q}^1 & \implies & H_{p+q}(\mathrm{GL}_2(A), \mathbb{Z}). \end{array}$$

From this we obtain the commutative diagram

$$\begin{array}{ccccc}
\mathcal{RP}_1(A) & \xrightarrow{\alpha_* \circ d_{2,1}^2} & H_2(T(A), \mathbb{Z}) & \xrightarrow{\cong} & A^\times \wedge A^\times \\
\downarrow & & \downarrow & & \downarrow \\
\mathcal{P}(A) & \xrightarrow{\beta_* \circ d_{2,1}^2} & \frac{H_2(T_2(A), \mathbb{Z})}{(\sigma_* - \text{inc}_*)(H_2(T_2(A), \mathbb{Z}))} & \xrightarrow{\cong} & (A^\times \wedge A^\times) \oplus \mathcal{S}_{\mathbb{Z}}^2(A),
\end{array}$$

where

$$\begin{aligned}
B_2(A) &:= \text{Stab}_{\text{GL}_2(A)}(\infty) = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} : a, d \in A^\times, b \in A \right\}, \\
T_2(A) &:= \text{Stab}_{\text{GL}_2(A)}(\infty, \mathbf{0}) = \left\{ \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} : a, d \in A^\times \right\},
\end{aligned}$$

and $\beta : B_2(A) \rightarrow T_2(A)$ is defined similar to α . It is straightforward to check that the vertical map on the right is given by

$$a \wedge b \rightarrow (2(a \wedge b), 2(a \otimes b)).$$

Moreover by a tedious computation one can show that the bottom horizontal composite is given by

$$[a] \mapsto (a \wedge (1 - a), -a \otimes (1 - a)),$$

(for details see [16], [15]). The lemma follows from the commutativity of the above diagram. \square

Our calculations suggest that the image of the differential

$$d_{2,1}^2 : \mathcal{RP}_1(A) \simeq E_{2,1}^2 \rightarrow H_2(B(A), \mathbb{Z}) \simeq (A^\times \wedge A^\times) \oplus \mathcal{S}_2.$$

may have a nonzero part in $\mathcal{S}_2 \subseteq H_2(B(A), \mathbb{Z})$ (see [16]). To obtain a good relation between the third homology of SL_2 and the refined Bloch group $\mathcal{RB}(A)$, we consider those rings that the summand \mathcal{S}_n of $H_n(B(A), \mathbb{Z})$ is trivial when $n = 2, 3$. The isomorphism $H_n(T(A), \mathbb{Z}) \xrightarrow{\text{inc}_*} H_n(B(A), \mathbb{Z})$ has been already studied for a certain class of rings. For example we have the following result of Hutchinson.

Proposition 5.2. *Let A be any local integral domain. If the residue field A/\mathfrak{m}_A is finite of order p^d we suppose that $(p - 1)d > 2n$. Then the natural maps $T(A) \rightarrow B(A)$ induces the isomorphism $H_n(T(A), \mathbb{Z}) \simeq H_n(B(A), \mathbb{Z})$.*

Proof. See [7, Proposition 3.19]. \square

One more fact that is needed in the next section is the structure of the homology group $H_3(T(A), \mathbb{Z})$.

Let B be an abelian group. Let $\sigma_1 : \text{Tor}_1^{\mathbb{Z}}(B, B) \rightarrow \text{Tor}_1^{\mathbb{Z}}(B, B)$ be obtained by interchanging the group B . It is not difficult to show that σ_1 is induced by the involution $B \otimes B \rightarrow B \otimes B, a \otimes b \mapsto -b \otimes a$.

Let $\Sigma'_2 = \{1, \sigma'\}$ be the symmetric group of order 2. Consider the following action of Σ'_2 on $\text{Tor}_1^{\mathbb{Z}}(B, B)$:

$$(\sigma', x) \mapsto -\sigma_1(x).$$

Proposition 5.3. *For any abelian group B we have the exact sequence*

$$0 \rightarrow \bigwedge_{\mathbb{Z}}^3 B \rightarrow H_3(B, \mathbb{Z}) \rightarrow \text{Tor}_1^{\mathbb{Z}}(B, B)^{\Sigma'_2} \rightarrow 0,$$

where the right side homomorphism is obtained from the composition

$$H_3(B, \mathbb{Z}) \xrightarrow{\Delta_{B*}} H_3(B \oplus B, \mathbb{Z}) \rightarrow \text{Tor}_1^{\mathbb{Z}}(B, B),$$

Δ_B being the diagonal map $B \rightarrow B \oplus B$, $b \mapsto (b, b)$.

Proof. See [18, Lemma 5.5], [2, Section 6]. □

6. THE REFINED BLOCH-WIGNER EXACT SEQUENCE IN CHARACTERISTIC 2

Let A be a ring which satisfies the following three conditions:

- (1) $\mu_2(A) = 1$,
- (2) $X_{\bullet}(A^2) \rightarrow \mathbb{Z}$ is exact in dimension < 4 ,
- (3) $H_n(T(A), \mathbb{Z}) \simeq H_n(B(A), \mathbb{Z})$ for $n = 2, 3$.

Note that if $\mu_2(A) = 1$, then A is of characteristic 2. The main example that we should keep in our mind is a domain of characteristic 2 such that its residue field has more than 64 elements (see Propositions 2.1 and 5.2). Here is our main result.

Theorem 6.1. *Let A be a ring which satisfies in the conditions (1), (2) and (3). Then we have the refined Bloch-Wigner exact sequence*

$$\text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

Moreover if A is a domain, then we have the exact sequence

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)) \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

Proof. By Lemma 5.1, $E_{2,1}^{\infty} \simeq E_{2,1}^3 \simeq \mathcal{RB}(A)$. We show that the differential

$$d_{2,2}^1 : H_2(\text{SL}_2(A), Z_1(A^2)) \rightarrow A^{\times} \wedge A^{\times}$$

is surjective. For $a \in A^{\times}$, denote $(\infty, \mathbf{0}, \mathbf{a}) \in X_2(A^2)$ by X_a . Let $Y = (\infty, \mathbf{0}) + (\mathbf{0}, \infty) \in Z_1(A^2)$. For $a, b \in A^{\times}$, let

$$\lambda(a, b) \in H_2(\text{SL}_2(A), Z_1(A)) = H_2(B_{\bullet}(\text{SL}_2(A)) \otimes_{\text{SL}_2(A)} Z_1(A))$$

be the element

$$\begin{aligned} \lambda(a, b) := & ([a|b] + [w|ab] - [w|a] - [w|b]) \otimes Y + [wab|wab] \otimes \partial_2(X_{ab}) \\ & - [wa|wa] \otimes \partial_2(X_a) - [wb|wb] \otimes \partial_2(X_b) + [w|w] \otimes \partial_2(X_1). \end{aligned}$$

Recall that $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. We have

$$\begin{aligned} d_{2,2}^1(\lambda(a,b)) &= (w+1)([a|b] + [w|ab] - [w|a] - [w|b]) \otimes (\infty, \mathbf{0}) \\ &\quad + (g_{ab}^{-1} - h_{ab}^{-1} + 1)([wab|wab]) \otimes (\infty, \mathbf{0}) \\ &\quad - (g_a^{-1} - h_a^{-1} + 1)([wa|wa]) \otimes (\infty, \mathbf{0}) \\ &\quad - (g_b^{-1} - h_b^{-1} + 1)([wb|wb]) \otimes (\infty, \mathbf{0}) \\ &\quad + (g_1^{-1} - h_1^{-1} + 1)([w|w]) \otimes (\infty, \mathbf{0}), \end{aligned}$$

where $g_x := \begin{pmatrix} 0 & 1 \\ 1 & x \end{pmatrix}$ and $h_x := \begin{pmatrix} 1 & x^{-1} \\ 0 & 1 \end{pmatrix}$. This element is in $H_2(\mathrm{SL}_2(A), X_1(A^2)) = H_2(B_\bullet(\mathrm{SL}_2(A)) \otimes_{\mathrm{SL}_2(A)} X_1(A))$. The morphisms

$$B_\bullet(\mathrm{SL}_2(A)) \otimes_{\mathrm{SL}_2(A)} X_1(A) \rightarrow B_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z} \rightarrow C_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z},$$

$$[g_1 | \cdots | g_n] \otimes (\infty, \mathbf{0}) \mapsto [g_1 | \cdots | g_n] \otimes 1 \mapsto \otimes(1, g_1, \dots, g_1 \cdots g_n) \otimes 1,$$

induce the isomorphisms

$$H_2(B_\bullet(\mathrm{SL}_2(A)) \otimes_{\mathrm{SL}_2(A)} X_1(A)) \simeq H_2(B_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z}) \simeq H_2(C_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z}).$$

Following these maps we see that $d_{2,2}^1(\lambda(a,b))$ as an element of $H_2(C_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z})$ find the following form

$$\begin{aligned} d_{2,2}^1(\lambda(a,b)) &= \left((w, wa, wab) + (w, 1, ab) - (w, 1, a) - (w, 1, b) \right. \\ &\quad + (1, a, ab) + (1, w, wab) + (1, w, wa) + (1, w, wb) \left. \right) \otimes 1 \\ &\quad + ((g_{ab}^{-1}, g_{ab}^{-1}wab, g_{ab}^{-1}) - (h_{ab}^{-1}, h_{ab}^{-1}wab, h_{ab}^{-1}) + (1, wab, 1)) \otimes 1 \\ &\quad - ((g_a^{-1}, g_a^{-1}wa, g_a^{-1}) - (h_a^{-1}, h_a^{-1}wa, h_a^{-1}) + (1, wa, 1)) \otimes 1 \\ &\quad - ((g_b^{-1}, g_b^{-1}wb, g_b^{-1}) - (h_b^{-1}, h_b^{-1}wb, h_b^{-1}) + (1, wb, 1)) \otimes 1 \\ &\quad + ((g_1^{-1}, g_1^{-1}w, g_1^{-1}) - (h_1^{-1}, h_1^{-1}w, h_1^{-1}) + (1, w, 1)) \left. \right) \otimes 1. \end{aligned}$$

Now we want to find a representative of this element in $C_\bullet(T(A)) \otimes_{T(A)} \mathbb{Z}$ by the isomorphism

$$H_2(C_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z}) \simeq H_2(C_\bullet(A^\times) \otimes_{A^\times} \mathbb{Z}).$$

Let $s : \mathrm{SL}_2(A) \setminus T(A) \rightarrow \mathrm{SL}_2(A)$ be any (set-theoretic) section of the canonical projection $\pi : \mathrm{SL}_2(A) \rightarrow \mathrm{SL}_2(A) \setminus T(A)$. For $g \in \mathrm{SL}_2(A)$, set $\bar{g} = (g)(s \circ \pi(g))^{-1}$. Then the morphism

$$C_\bullet(\mathrm{SL}_2(A)) \otimes_{T(A)} \mathbb{Z} \xrightarrow{s_\bullet} C_\bullet(T(A)) \otimes_{T(A)} \mathbb{Z}, \quad [g_1 | \cdots | g_n] \otimes 1 \mapsto [\bar{g}_1 | \cdots | \bar{g}_n] \otimes 1$$

induces the desired isomorphism. By choosing the section

$$s(T(A) \begin{pmatrix} a & b \\ c & d \end{pmatrix}) = \begin{cases} \begin{pmatrix} 1 & a^{-1}b \\ ac & ad \end{pmatrix} & \text{if } a \neq 0 \\ \begin{pmatrix} 0 & 1 \\ 1 & bd \end{pmatrix} & \text{if } a = 0 \end{cases}$$

We see that $d_{2,2}^1(\lambda(a,b))$ in $H_2(C_\bullet(T(A)) \otimes_{T(A)} \mathbb{Z}) = H_2(C_\bullet(A^\times) \otimes_{A^\times} \mathbb{Z})$ is of the following form

$$d_{2,2}^1(\lambda(a,b)) = \left((1, a^{-1}, (ab)^{-1}) + (1, a, ab) + (1, (ab)^{-1}, 1) - (1, a^{-1}, 1) - (1, b^{-1}, 1) \right) \otimes 1.$$

In $H_2(B_\bullet(A^\times) \otimes_{A^\times} \mathbb{Z})$ this elements corresponds to

$$d_{2,2}^1(\lambda(a,b)) = \left([a^{-1}|b^{-1}] + [a|b] + [a^{-1}b^{-1}|ab] - [a^{-1}|a] - [b^{-1}|b] \right) \otimes 1.$$

Now by adding the following null element in $H_2(A^\times, \mathbb{Z}) = H_2(B_\bullet(A^\times) \otimes_{A^\times} \mathbb{Z})$

$$d_3([a^{-1}|b^{-1}|ab] - [b^{-1}|b|a]) \otimes 1,$$

it is easy to see that

$$d_{2,2}^1(\lambda(a,b)) = ([a|b] - [b|a]) \otimes 1 \in H_2(B_\bullet(A^\times) \otimes_{A^\times} \mathbb{Z}).$$

This shows that $d_{2,2}^1$ is surjective. Therefore $E_{1,2}^2 = 0$.

Now we need to study $E_{0,3}^\infty = E_{0,3}^3$. To do this, first consider the differential

$$d_{1,3}^1 = H_3(\sigma) - H_3(\text{inc}) : H_3(T(A), \mathbb{Z}) \rightarrow H_3(B(A), \mathbb{Z}) \simeq H_3(T(A), \mathbb{Z}).$$

By Proposition 5.3, we have the exact sequence

$$0 \rightarrow \bigwedge_{\mathbb{Z}}^3 A^\times \rightarrow H_3(T(A), \mathbb{Z}) \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} \rightarrow 0.$$

It is straightforward to check that $d_{1,3}^1 |_{\bigwedge_{\mathbb{Z}}^3 A^\times}$ coincides with multiplication by 2. Thus we have the exact sequence

$$\bigwedge_{\mathbb{Z}}^3 A^\times / 2 \rightarrow E_{0,3}^2 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} / \mathcal{T} \rightarrow 0,$$

for some subgroup \mathcal{T} of $\text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2}$. By an easy analysis of the spectral sequence we have the exact sequence

$$E_{0,3}^3 \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

We denote the image of $a \wedge b \wedge c \in \bigwedge_{\mathbb{Z}}^3 A^\times / 2$ in $E_{0,3}^2$ again by $a \wedge b \wedge c$. Since $d_{2,2}^1(\lambda(ab,c) - \lambda(a,c) - \lambda(b,c)) = 0$, we have $\lambda(ab,c) - \lambda(a,c) - \lambda(b,c) \in E_{2,2}^2$. We show that

$$(6.1) \quad d_{2,2}^2(\lambda(ab,c) - \lambda(a,c) - \lambda(b,c)) = -a \wedge b \wedge c \in E_{0,3}^2.$$

This would imply that there is a surjective map $\mathrm{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} \rightarrow E_{0,3}^3$ and therefore we obtain the exact sequence

$$\mathrm{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} \rightarrow H_3(\mathrm{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0,$$

which proof the first claim of the theorem. Now we prove the equality (6.1).

Consider the diagram

$$\begin{array}{ccc} B_3(\mathrm{SL}_2(A)) \otimes X_0(A^2) & \xleftarrow{\mathrm{id}_{B_3} \otimes \partial_1} & B_3(\mathrm{SL}_2(A)) \otimes X_1(A^2) \\ & & \downarrow d_3 \otimes \mathrm{id}_{X_1} \\ & & B_2(\mathrm{SL}_2(A)) \otimes X_0(A^2) \xleftarrow{\mathrm{id}_{B_2} \otimes \mathrm{inc}} B_2(\mathrm{SL}_2(A)) \otimes Z_1(A^2). \end{array}$$

The element $\Lambda(a, b, c) := \lambda(ab, c) - \lambda(a, c) - \lambda(b, c)$ is

$$\begin{aligned} \Lambda(a, b, c) := & ([ab|c] - [a|c] - [b|c] + [w|abc] - [w|ac] - [w|bc] - [w|ab] + [w|a] + [w|b] + [w|c]) \otimes Y \\ & + [wabc|wabc] \otimes \partial_2(X_{abc}) - [wab|wab] \otimes \partial_2(X_{ab}) - [wbc|wbc] \otimes \partial_2(X_{bc}) \\ & - [wac|wac] \otimes \partial_2(X_{ac}) + [wa|wa] \otimes \partial_2(X_a) + [wb|wb] \otimes \partial_2(X_b) \\ & + [wc|wc] \otimes \partial_2(X_c) - [w|w] \otimes \partial_2(X_1). \end{aligned}$$

For an element $z \in A^\times$, consider $[wz|wz] \otimes \partial_2(X_z) \in B_2(\mathrm{SL}_2(A)) \otimes_{\mathrm{SL}_2(A)} Z_1(A^2)$. For the matrices g_z and h_z we have the identities

$$z^{-1}g_{z^{-1}} = g_z z, \quad zh_z = h_{z^{-1}}z, \quad g_z^{-1}w = h_{z^{-1}}, \quad h_z^{-1} = h_z.$$

Using these identities we obtain

$$\begin{aligned} [wz|wz] \otimes \partial_2(X_z) = & [wz|wz] \otimes (\infty, 0) \\ & + (d_3 \otimes \mathrm{id}_{X_1})((g_z^{-1}|wz|wz] - [h_z|wz|wz] + [z^{-1}|g_z^{-1}|wz] - [z|h_z|wz]) \otimes (\infty, \mathbf{0})) \\ & + (d_3 \otimes \mathrm{id}_{X_1})((z|z^{-1}|g_z] - [z^{-1}|z|h_z] + [z^{-1}|z|z^{-1}]) \otimes (\infty, 0)). \end{aligned}$$

If

$$\begin{aligned} \theta_z := & [g_z^{-1}|wz|wz] - [h_z|wz|wz] + [z^{-1}|g_z^{-1}|wz] - [z|h_z|wz] \\ & + [z|z^{-1}|g_z] - [z^{-1}|z|h_z] + [z^{-1}|z|z^{-1}], \end{aligned}$$

then by a direct calculation we have

$$\Lambda(a, b, c) = (d_3 \otimes \mathrm{id}_{X_1})((w|ab|c] - [w|a|c] - [w|b|c]) \otimes Y + (\Phi_{a,b,c} + \Psi_{a,b,c}) \otimes (\infty, \mathbf{0}),$$

where

$$\begin{aligned} \Phi_{a,b,c} = & \theta_{abc} - \theta_{ab} - \theta_{bc} - \theta_{ac} + \theta_a + \theta_b + \theta_c - \theta_1 \\ \Psi_{a,b,c} = & [wab|wab|c] - [wa|wa|c] - [wb|wb|c] + [w|w|c] \\ & + [c|wabc|wabc] - [c|wac|wac] - [c|wbc|wbc] + [c|wc|wc] \\ & + [ab|wab|c] - [a|wa|c] - [a|c|wac] + [ab|c|wabc] - [c|ab|wabc] + [c|a|wac] \\ & - [b|wb|c] - [b|c|wbc] + [c|b|wbc] - [b|a|c] + [b|c|a] - [c|b|a]. \end{aligned}$$

Since $\partial_1(Y) = 0$, we need only to study

$$(\text{id}_{B_3} \otimes \partial_1)((\Phi_{a,b,c} + \Psi_{a,b,c}) \otimes (\infty, \mathbf{0})).$$

Through the maps

$$B_3(\text{SL}_2(A)) \otimes_{\text{SL}_2(A)} X_1(A^2) \xrightarrow{\text{id}_{B_3} \otimes \partial_1} B_3(\text{SL}_2(A)) \otimes_{\text{SL}_2(A)} X_0(A^2)$$

the above element maps to

$$(w\Phi_{a,b,c} - \Phi_{a,b,c}) \otimes (\infty) + (w\Psi_{a,b,c} - \Psi_{a,b,c}) \otimes (\infty).$$

Now consider the composite

$$B_3(\text{SL}_2(A)) \otimes_{\text{SL}_2(A)} X_0(A^2) \rightarrow C_3(\text{SL}_2(A)) \otimes_{\text{SL}_2(A)} X_0(A^2) \rightarrow C_3(\text{SL}_2(A)) \otimes_{B(T)} \mathbb{Z}.$$

Then

$$\begin{aligned} (w\theta_z - \theta_z) \otimes (\infty) \mapsto & \left\{ (w, wg_z^{-1}, wg_z^{-1}wz, wg_z^{-1}) - (w, wh_z, wh_zwz, wh_z) \right. \\ & + (w, wz^{-1}, wz^{-1}g_z^{-1}wz) - (w, wz, wzh_z, wzh_zwz) \\ & + (w, wz, w, wg_z^{-1}) - (w, wz^{-1}, w, wh_z) \\ & + (w, wz^{-1}, w, wz^{-1}) - (1, g_z^{-1}, g_z^{-1}wz, g_z^{-1}) \\ & + (1, h_z, h_zwz, h_z) - (1, z^{-1}, z^{-1}g_z^{-1}, z^{-1}g_z^{-1}wz) \\ & + (1, z, zh_z, zh_zwz) - (1, z, 1, g_z^{-1}) \\ & \left. + (1, z^{-1}, 1, h_z) - (1, z^{-1}, 1, z^{-1}) \right\} \otimes 1 \end{aligned}$$

and

$$\begin{aligned} (w\Psi_{a,b,c} - \Psi_{a,b,c}) \otimes (\infty) \mapsto & \left\{ (w, ab, w, wc) - (w, a, w, wc) - (w, b, w, wc) + (w, 1, w, wc) \right. \\ & + (w, wc, ab, wc) - (w, wc, a, wc) - (w, wc, b, wc) + (w, wc, 1, wc) \\ & + (w, wab, 1, c) - (w, wa, 1, c) - (w, wa, wac, 1) + (w, wab, wabc, 1) \\ & - (w, wc, wabc, 1) + (w, wc, wac, 1) - (w, wb, 1, c) - (w, wb, wbc, 1) \\ & + (w, wc, wbc, 1) - (w, wb, wab, wabc) + (w, wb, wbc, wabc) \\ & - (w, wc, wbc, wabc) - (1, wab, 1, c) + (1, wa, 1, c) + (1, wb, 1, c) \\ & - (1, w, 1, c) - (1, c, wab, c) + (1, c, wa, c) + (1, c, wb, c) - (1, c, w, c) \\ & - (1, ab, w, wc) + (1, a, w, wc) + (1, a, ac, w) - (1, ab, abc, w) \\ & + (1, c, abc, w) - (1, c, ac, w) + (1, b, w, wc) + (1, b, bc, w) \\ & \left. - (1, c, bc, w) + (1, b, ab, abc) - (1, b, bc, abc) + (1, c, bc, abc) \right\} \otimes 1. \end{aligned}$$

Now we want to follow these elements through the maps

$$C_3(\text{SL}_2(A)) \otimes_{B(A)} \mathbb{Z} \xrightarrow{s} C_3(B(A)) \otimes_{B(A)} \mathbb{Z} \rightarrow C_3(T(A)) \otimes_{T(A)} \mathbb{Z} \rightarrow B_3(T(A)) \otimes_{T(A)} \mathbb{Z}$$

where $s : \mathrm{SL}_2(A) \setminus T(A) \rightarrow \mathrm{SL}_2(A)$ is the section discussed in above. It is straightforward to check that modulo $\mathrm{im}(d_4)$ we have

$$(w\theta_z - \theta_z) \otimes (\infty) \mapsto -[z^{-1}|z|z^{-1}] \otimes 1 = [z|z^{-1}|z] \otimes 1$$

Moreover

$$\begin{aligned} (w\Psi_{a,b,c} - \Psi_{a,b,c}) \otimes (\infty) \mapsto & \left\{ [c^{-1}|abc|(abc)^{-1}] - [c^{-1}|ac|(ac)^{-1}] - [c^{-1}|bc|(bc)^{-1}] + [c^{-1}|c|c^{-1}] \right. \\ & - [a^{-1}|c^{-1}|ac] + [(ab)^{-1}|c^{-1}|abc] - [c^{-1}|(ab)^{-1}|abc] + [c^{-1}|a^{-1}|ac] \\ & - [b^{-1}|c^{-1}|bc] + [c^{-1}|b^{-1}|bc] - [b^{-1}|a^{-1}|c^{-1}] + [b^{-1}|c^{-1}|a^{-1}] \\ & - [c^{-1}|b^{-1}|a^{-1}] - [c|(abc)^{-1}|abc] + [c|(ac)^{-1}|ac] + [c|(bc)^{-1}|bc] \\ & - [c|c^{-1}|c] + [a|c|(ac)^{-1}] - [ab|c|(abc)^{-1}] + [c|ab|(abc)^{-1}] \\ & - [c|a|(ac)^{-1}] + [b|c|(bc)^{-1}] - [c|b|(bc)^{-1}] + [b|a|c] - [b|c|a] \\ & \left. + [c|b|a] \right\} \otimes 1. \end{aligned}$$

Combining all these we see that $d_{2,2}^2(\Lambda(a, b, c))$ is the following element of $E_{0,3}^2$

$$\begin{aligned} d_{2,2}^2(\Lambda(a, b, c)) = & \left\{ [c^{-1}|abc|(abc)^{-1}] - [c^{-1}|ac|(ac)^{-1}] - [c^{-1}|bc|(bc)^{-1}] + [c^{-1}|c|c^{-1}] \right. \\ & - [a^{-1}|c^{-1}|ac] + [(ab)^{-1}|c^{-1}|abc] - [c^{-1}|(ab)^{-1}|abc] + [c^{-1}|a^{-1}|ac] \\ & - [b^{-1}|c^{-1}|bc] + [c^{-1}|b^{-1}|bc] - [b^{-1}|a^{-1}|c^{-1}] + [b^{-1}|c^{-1}|a^{-1}] \\ & - [c^{-1}|b^{-1}|a^{-1}] - [c|(abc)^{-1}|abc] + [c|(ac)^{-1}|ac] + [c|(bc)^{-1}|bc] \\ & + [a|c|(ac)^{-1}] - [ab|c|(abc)^{-1}] + [c|ab|(abc)^{-1}] - [c|a|(ac)^{-1}] \\ & + [b|c|(bc)^{-1}] - [c|b|(bc)^{-1}] + [b|a|c] - [b|c|a] + [c|b|a] \\ & + [abc|(abc)^{-1}|abc] - [ab|(ab)^{-1}|ab] - [bc|(bc)^{-1}|bc] \\ & \left. - [ac|(ac)^{-1}|ac] + [a|a^{-1}|a] + [b|b^{-1}|b] \right\} \otimes 1 \end{aligned}$$

By adding the null element

$$\begin{aligned} d_4 \left(\left\{ - [c|c^{-1}|abc|(abc)^{-1}] + [c|c^{-1}|ac|(ac)^{-1}] + [c|c^{-1}|bc|(bc)^{-1}] - [c|c^{-1}|a^{-1}|ac] \right. \right. \\ - [c|c^{-1}|c|c^{-1}] - [c|c^{-1}|b^{-1}|bc] + [c^{-1}|c|(abc)^{-1}|abc] + [ac|a^{-1}|c^{-1}|ac] \\ - [abc|(ab)^{-1}|c^{-1}|abc] + [bc|b^{-1}|c^{-1}|bc] + [c|ab|(ab)^{-1}|ab] - [c|b|b^{-1}|b] - [c|a|a^{-1}|a] \\ - [c|a|b|(ab)^{-1}] - [a|b|c|(abc)^{-1}] + [abc|b^{-1}|a^{-1}|c^{-1}] - [abc|b^{-1}|c^{-1}|a^{-1}] \\ + [a|c|c^{-1}|a^{-1}] - [a|bc|b^{-1}|c^{-1}] + [a|c|b|b^{-1}] + [ac|b|b^{-1}|a^{-1}] - [a|bc|(bc)^{-1}|a^{-1}] \\ \left. \left. - [b|c|(bc)^{-1}|a^{-1}] + [c|c^{-1}|b^{-1}|a^{-1}] - [c|c^{-1}|c|(abc)^{-1}] \right\} \otimes 1 \right) \end{aligned}$$

we see that, modulo $\text{im}(d_4)$,

$$\begin{aligned} d_{2,2}^2(\Lambda(a, b, c)) &= -([a|b|c] + [c|a|b] + [b|c|a] - [b|a|c] - [c|b|a] - [a|c|b]) \otimes 1 \\ &= -a \wedge b \wedge c. \end{aligned}$$

Thus we obtain the desired exact sequence

$$\text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

Now let A be a domain. Since $\mu(A)$ is direct limit of finite cyclic groups, then

$$\text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A))^{\Sigma'_2} = \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)).$$

Let F be the quotient field of A and \overline{F} the algebraic closure of F . It is very easy to see that $\mathcal{RB}(\overline{F}) = \mathcal{B}(\overline{F})$. The classical Bloch-Wigner exact sequence claims that the sequence

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(\overline{F}), \mu(\overline{F})) \rightarrow H_3(\text{SL}_2(\overline{F}), \mathbb{Z}) \rightarrow \mathcal{B}(\overline{F}) \rightarrow 0$$

is exact. Now the final claim follows from the commutative diagram with exact rows

$$\begin{array}{ccccccc} \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)) & \longrightarrow & H_3(\text{SL}_2(A), \mathbb{Z}) & \longrightarrow & \mathcal{RB}(A) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \text{Tor}_1^{\mathbb{Z}}(\mu(\overline{F}), \mu(\overline{F})) & \longrightarrow & H_3(\text{SL}_2(\overline{F}), \mathbb{Z}) & \longrightarrow & \mathcal{B}(\overline{F}) \longrightarrow 0 \end{array}$$

and the fact that the natural map $\text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)) \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(\overline{F}), \mu(\overline{F}))$ is injective. \square

Corollary 6.2. *Let A be a local domain of characteristic 2, where its residue field has more than 2^6 elements. Then we have the refined Bloch-Wigner exact sequence*

$$0 \rightarrow \text{Tor}_1^{\mathbb{Z}}(\mu(A), \mu(A)) \rightarrow H_3(\text{SL}_2(A), \mathbb{Z}) \rightarrow \mathcal{RB}(A) \rightarrow 0.$$

Proof. This follows from Proposition 2.1, Proposition 5.2 and Theorem 6.1. \square

REFERENCES

- [1] Brown, K. S. Cohomology of Groups. Graduate Texts in Mathematics, 87. Springer-Verlag, New York (1994) [4](#)
- [2] L. Breen. On the functorial homology of abelian groups. Journal of Pure and Applied Algebra **142** (1999) 199–237. [8](#)
- [3] Coronado, R. C., Hutchinson, K. Bloch Groups of Rings. <https://arxiv.org/abs/2201.04996> [1](#)
- [4] Dupont, J- L., Sah, C. Scissors congruences. II. J. Pure Appl. Algebra **25** (1982), no. 2, 159–195. [1](#), [2](#)
- [5] Hutchinson, K. A Bloch-Wigner complex for SL_2 . J. K-Theory **12**, no. 1, 15–68 (2013) [1](#), [2](#), [3](#)
- [6] Hutchinson, K. A refined Bloch group and the third homology of SL_2 of a field. J. Pure Appl. Algebra **217**, 2003–2035 (2013) [1](#), [2](#), [3](#)
- [7] Hutchinson, K. The third homology of SL_2 of local rings. Journal of Homotopy and Related Structures **12**, 931–970 (2017) [1](#), [2](#), [3](#), [7](#)
- [8] Hutchinson, K. The third homology of SL_2 of fields with discrete valuation. J. Pure Appl. Algebra **221** (5), 1076–1111 (2017) [2](#)
- [9] Hutchinson, K. The third homology of $\text{SL}_2(\mathbb{Q})$. J. Algebra **570** (2021), 366–396 [1](#), [2](#)
- [10] Hutchinson, K., Tao, L. The third homology of the special linear group of a field. J. Pure Appl. Algebra **213** (2009), no. 9, 1665–1680. [1](#)

- [11] Hutchinson, K., Mirzaii, B., Mokari, F. Y. The homolog of SL_2 of discrete valuation rings. *Advances in Mathematics* **402** (2022) 108313 [2](#)
- [12] Levine, M. The indecomposable K_3 of fields. *Ann. Sci. École Norm. Sup. (4)* **22** (1989), 255–344. [1](#)
- [13] Merkurjev, A. S., Suslin, A. A. The group K_3 for a field. *Math. USSR-Izv.* **36**, no. 3, 541–565 (1991) [1](#)
- [14] Mirzaii, B. Third homology of general linear groups. *J. Algebra* **320** (2008), no. 5, 1851–1877. [1](#)
- [15] Mirzaii, B. Bloch-Wigner theorem over rings with many units. *Math. Z.* **268**, 329–346 (2011). Erratum to: Bloch-Wigner theorem over rings with many units. *Math. Z.* **275**, 653–655 (2013) [2](#), [6](#), [7](#)
- [16] Mirzaii, B. A Bloch-Wigner exact sequence over local rings. *Journal of Algebra* **476**, 459–493 (2017) [1](#), [7](#)
- [17] Sah, C. Homology of classical Lie groups made discrete. III. *J. Pure Appl. Algebra* **56** (1989), no. 3, 269–312. [1](#)
- [18] Suslin, A. A. K_3 of a field and the Bloch group. *Proc. Steklov Inst. Math.* **183**, no. 4, 217–239 (1991) [1](#), [2](#), [8](#)