

On modeling the TSP as a polynomial-sized linear program: Escaping/Side-stepping the “barriers”

Moustapha Diaby

OPIIM Department; University of Connecticut; Storrs, CT 06268

moustapha.diaby@uconn.edu

Mark H. Karwan

Department of Industrial and Systems Engineering; SUNY at Buffalo; Amherst, NY 14260

mkarwan@buffalo.edu

Lei Sun

Department of Industrial and Systems Engineering; SUNY at Buffalo; Amherst, NY 14260

leisun@buffalo.edu

Abstract: In view of the *extended formulations* (EFs) developments, we bring a clarification in this paper to the question of whether it is possible to model an NP-Complete problem as a polynomial-sized linear program. For the sake of simplicity of exposition, we focus on the traveling salesman problem (TSP). We show that a finding that there exists no polynomial-sized EF of “*the* TSP polytope” does not (necessarily) imply that it is “impossible” for a polynomial-sized linear program to solve the TSP optimization problem. We show that under appropriate conditions the TSP optimization problem can be solved without recourse to the traditional city-to-city (“travel leg”) variables, thereby side-stepping/“escaping from” “*the* TSP polytope” and hence, the “barriers.” Some illustrative examples are discussed.

Keywords: Combinatorial Optimization; Computational Complexity; Linear Programming; Linear Assignment Problem; Traveling Salesman Problem.

1 Introduction

A fundamental issue for modern mathematics in general has been the question of decidability/computability (Hilbert [1901]; Turing [1936]). The decidability/computability question was reduced (in essence) to that of the equality/non-equality of the computational complexity classes “P” and “NP” (“P vs. NP” problem) in the early 1970’s (Cook [1971]). Roughly, the “NP” class of problems are such that checking the correctness of a given solution to them (respectively) is “easy.” The “P” class consists of “NP” problems which are known to also be “easy” to solve (i.e., for which polynomial-time algorithms are known). Hence, “P” is a subclass of “NP.” Within the “NP” class, the so-called NP-Complete (NPC; see Garey and Johnson [1979], among others) problems play a central role in the sense that: (1) they are polynomially reducible to each other among themselves; and (2) every other problem in “NP” is polynomially reducible to any of them. The essence of the “P vs. NP” problem is

the question of whether or not “P” is a proper subclass of “NP,” or in other words, whether or not there exists a polynomial time procedure for solving any of the NPC problems.

In the field of mathematical programming, NPC problems are generally analyzed under the umbrella of combinatorial optimization. Hence, with the advent of polynomial-time algorithms for solving linear programs (LPs) in the early 1980’s (Khachiyan [1979]; Karmarkar [1984]), the “P vs. NP” problem was reduced in part (for operations research (OR) in particular) to the question of whether or not any of the NP-Complete combinatorial optimization problems (COPs) can be modeled as a polynomial-sized LP. The challenge in this sense is to develop a set of linear constraints in such a way that the extreme points of the resulting LP are in a one-to-one correspondence with the combinatorial configuration (e.g. “tours,” “partitions,” “covers,” etc.) for the COP at hand. The NPC problem which has been used in research focused on this issue in general has been the traveling salesman problem (TSP). NP-Completeness pertains to decision problems. However, throughout the remainder of this paper, we will use the TSP optimization problem (which is NP-Hard; see Garey and Johnson [1979], among others) instead of the TSP decision problem. Our reasons for doing this are convenience (since the framework of our exposition is a mathematical programming one) and the fact that the TSP optimization problem subsumes the TSP decision problem.

The first polynomial-sized LP models proposed for solving the TSP date back to the mid-1980’s (Swart [1986/1987]). We do not know of any details of these models. However, the consensus of the research communities is that their validities were refuted by a negative result (“barrier”) developed in the early 1990’s (Yannakakis [1991]). The most prominent conclusion of the Yannakakis [1991] work is that “*the* TSP polytope” does not have a polynomial-sized extended formulation (EF) which is symmetric. By “*the* TSP polytope,” we mean the polytope described using the traditional/standard/“natural”/“travel leg”/“city-to-city” variables for the TSP. Since these Yannakakis [1991] developments, making inferences about possible LP models for COPs by analyzing EFs of their *natural* polytopes has become a standard practice. Recently, Fiorini et al. [2015] strengthened the Yannakakis *barrier* into one which no longer has the symmetry caveat. That work has been, in turn, seminal for many of the other recent developments (e.g. Faenza et al. [2021], Aprile [2022], and Kwan et al. [2022]).

The purpose of this paper is not to dispute the validity of the *barriers* themselves. Rather, our objective is to show that the non-existence of a polynomial-sized integral extended formulation of “*the* TSP polytope” does not imply that it is “impossible” for a polynomial-sized linear program to solve the TSP optimization problem. The importance of this comes from the significance of NPC problems for both practice and theory in many fields of study and the fact that, in the literature, there have been assertions based on developments for “*the* TSP polytope” that it is “impossible” to formulate any of these problems as a linear program.

An optimization problem (using mathematical programming) is defined by both its constraints and its objective function. An oddity of the TSP (and incidentally, of many of the other NP-Complete problems as well) is its relation to the linear assignment problem (LAP)/bi-partite matching problem. Specifically, each extreme point of the LAP defined by its linear constraint set can be interpreted as a TSP tour with the assignment of row i to column j (like job i assigned to person j) interpreted as city i being the j^{th} city visited. The difficulty in this abstraction however, is that the cost function of traveling from city to




Figure 1: Illustration of the *TSP Graph*

city cannot be captured using the two-dimensional, *natural* variables traditionally used to state the constraints of the LAP. Hence, the key to being able to exploit this feature is to extend the LAP into a higher dimension in a way that an objective function relating to sets of joint assignments can be expressed. Also, the numbers of variables and constraints must respectfully be polynomial, and the model must have all-integer extreme points and project to the LAP polytope. As far as we know, this possibility is not ruled out by any existing result as it may be able to be accomplished without regard to the traditional city-to-city variables of the TSP but only by using an appropriate cost function on the EF of the LAP.

The plan of the remainder of the paper is as follows. In section 2, we give an overview of our basic notions, definitions, and notations; discuss alternate representations of TSP tours; and finally, briefly overview the existing *barriers* and discuss some simple misconceptions and misinterpretations in relation to them. In section 3, we provide a motivating illustrative example and develop conditions under which “*the TSP polytope*” (and hence, the *barriers*) can be side-stepped/“escaped from” with respect to the goal of developing polynomial-sized LP models for solving the TSP optimization problem. Finally, we offer some conclusions in section 4.

2 Preliminaries

2.1 Basic notions, definitions, and notations

Our discussions will involve two alternate abstractions of the TSP optimization problem. We give overviews of these and their accompanying notations in this section. The graph which underlies the first abstraction (which we label the “traditional abstraction”) is illustrated in Figure 1. Following the terminology commonly used in the literature, we refer to this graph as the “TSP graph.” Each node of the *TSP graph* represents a city, whereas each arc represents a *travel leg*.




Figure 2: Illustration of the TSP Assignment Graph (TSPAG)

The second abstraction we consider in this paper (which we label as the “alternate representation”) is a variant of the *time-dependent* model first proposed by Picard and Queyranne [1978]. In this representation, city 0 always occupies the first and last position in the sequence formed from a given TSP tour. In other words, we assume that every travel begins and ends at city 0. Hence in the sequencing/permutation-finding optimization problem, the “positioning” of city 0 ceases to be a decision variable. In other words, the search for an optimal TSP tour reduces to a search for a permutation over the set of cities, minus city 0. Hence, the graph which underlies this abstraction is correctly illustrated by Figure 2. Observe that this graph is a generic Assignment Problem graph (see Burkard *et al.* [2009]). In the context of job assignments for example, each row of nodes of this graph (which we will refer to as a “level” of the graph) could represent a worker, and each column of nodes (which we refer to as a “stage” of the graph) could represent a task. Hence, we will refer to this graph as the “TSP Assignment Graph (TSPAG).”

The formalization of these graphs above for a TSP with n cities and the general notation which will be used throughout the remainder of the paper are shown below. (For convenience, table-lists of all the notations used in the paper are also provided in the Appendix to this paper.)

Notation 1 (General Notation)

1. n : Number of TSP cities.
2. $m := n - 1$ (Numbers of *levels* and *stages* respectively, of the TSPAG).
3. $C := \{0, \dots, m\}$: Set of cities.
4. $\forall (i, j) \in C^2, d_{ij}$: Cost of traveling from city i to city j .

5. $\ell := n \times m$ (Number of arcs in the *TSP graph*).
6. \mathcal{P}_n : “the TSP polytope” defined over n cities.
7. $L := \{1, \dots, m\}$ (Index set for the *levels* of the TSPAG).
8. $S := \{1, \dots, m\}$ (Index set for the *stages* of the TSPAG).
9. $N := \{[l, s] : “(l, s)”, (\forall l \in L), (\forall s \in S)\}$ (There is no “ceiling” or “floor” meaning attached to “[” and “]”).
10. \mathcal{A}_n : The LAP polytope defined over the TSPAG associated with a n -city TSP.
11. $(A)^t$: Transpose of matrix A .
12. $Ext(B)$: Set of extreme points of polyhedron B .
13. $\pi_x(B)$: Projection of polyhedron B on the space of vector x .
14. \mathbb{R} : Set of real numbers.
15. $\mathbb{R}_{\geq 0}$: Set of nonnegative real numbers.
16. \mathbb{N} : Set of natural numbers (excluding “0”).

2.2 Alternate representations of TSP tours

2.2.1 “The TSP polytope”

The existing negative results (*barriers*) are based on the conventional, standard abstraction which focuses on the arcs of the *TSP graph* (i.e., the *travel legs* of the TSP). The goal in this abstraction is to select a set of arcs of the *TSP graph* which form a Hamiltonian cycle of the nodes of the graph. This is illustrated in Figure 3. Hence, in this abstraction, a TSP tour can be represented in terms of the TSP cities as a $(n + 1)$ -tuple $h = (h_0 = 0, h_1, \dots, h_{n-1}, h_n = 0)$ such that $\{h_1, \dots, h_{n-1}\} = C \setminus \{0\}$. The TSP tour represented by a given tuple h can also be represented in terms of *travel legs* as $\tau(h) := \{(h_k, h_{k+1}) \in h^2, k = 0, \dots, n - 1\}$.

For the purpose of a linear programming (LP) modeling, attach a vector of variables $y \in \{0, 1\}^{n(n-1)}$ to the *travel legs* with $y_{ij} \in ((i, j) \in C^2)$ equal to 1 if and only if there is travel from city i to city j . The TSP tour represented by a given tuple h is then modeled by the y -instance, \bar{y}^h , which is such that, for each $(i, j) \in C^2$, \bar{y}_{ij}^h is equal to 1 if and only if $(i, j) \in \tau(h)$. “The TSP polytope” (which we denote by \mathcal{P}_n) is defined as the polytope in the space of y which has the \bar{y}^h ’s as its extreme points.

Note that city 0 can be duplicated into a *dummy* city n , so that the condition “ $h_n = 0$ ” can be alternatively written as “ $h_n = n$.” Hence, h represents a constrained permutation of the TSP cities, since the “ $h_0 = 0$ ” and “ $h_n = 0$ ” requirements must be enforced explicitly (including through some classic types of inequalities (see Lawler et al. [1988])). Hence, in other words, in an exact EF of \mathcal{P}_n , each extreme point of the model must represent a constrained/restricted permutation of the TSP cities. Hence, such EF would essentially




Figure 3: Illustration of a *Hamiltonian Cycle*

include “side-constraints” in addition to the descriptive constraints of the LAP polytope defined over (an Assignment-graph version of) the *TSP graph*, while -at the same time- being required to be integral. Solutions of this *TSP graph*-based LAP which comprise subtours of the cities must also be excluded from consideration through additional “side-constraints” (such as in the classical Dantzig et. al. [1954] model, or such as relating to the *cut polytope* discussed in Fiorini et al. [2015; section 3.2] for example). We believe that the notorious difficulty involved in coming up with a “small” LP model of the TSP optimization problem in which the *travel leg* variables y are irredundant/non-superfluous (from an optimization perspective) and the constraints set is an EF of \mathcal{P}_n resides principally in these facts.

2.2.2 LAP solution representation of TSP tours

A simplification of the representation described in section 2.2.1 above can be obtained by observing that under the assumption that all travels begin and end at city 0, an m -tuple $p = (p_1, \dots, p_m)$ which is such that $\{p_1, \dots, p_m\} = C \setminus \{0\}$ is sufficient for abstracting a TSP tour. The problem of choosing from among these m -tuples is most naturally modeled by the LAP defined over the TSPAG. To specify this model, attach to the nodes of the TSPAG variables $w \in \{0, 1\}^{m^2}$ with w_{ir} ($i \in L, r \in S$) equal to 1 if and only if i is the r^{th} city visited after city 0. The instance of w, \bar{w}^p , associated with tuple p has set of positive components $\{\bar{w}_{p_1,1}, \dots, \bar{w}_{p_m,m}\}$, and the sequence of the visits in the TSP tour represented by p is $0 \rightarrow p_1 \rightarrow \dots \rightarrow p_m \rightarrow 0$. It is well-known (see Burkhard et al. [2009]) that the polytope which has the \bar{w}^p 's as its extreme points is the LAP/bipartite-matching polytope in the space of w , which we denote by \mathcal{A}_n . Hence, there exists a one-to-one correspondence between the TSP tours and the extreme points of \mathcal{A}_n .

In order for the modeling based on this alternate abstraction of TSP tours to be complete with respect to the task of solving the TSP optimization problem, a judicious extension of \mathcal{A}_n in which TSP tour costs are correctly captured through auxiliary variables must be developed. In section 3 of this paper, we show that such an EF would allow for the solution




Figure 4: Illustration of *TSP paths*

of the TSP optimization problem irrespective of whether it is polynomial-sized or not and of whether it projects to “the TSP polytope” (\mathcal{P}_n) or not.

We will close this section with notions which formalize the representation of TSP tours in terms of the nodes of the TSPAG.

Definition 2 (“TSP paths”)

1. We refer to a set of nodes at consecutive *stages* of the TSPAG with exactly one node at each *stage* in the set involved as a *path* of the TSPAG. In other words, for $(r, s) \in S^2 : s > r$, we refer to $\{[u_p, p] \in N, p = r, \dots, s\}$ as a *path* of the TSPAG.
2. We refer to a *path* of the TSPAG which (simultaneously) spans the *stages* and the *levels* of the TSPAG as a *TSP path* (of the TSPAG). In other words, letting m be the number of *stages* of the TSPAG, we refer to $\{[u_p, p] \in N, p = 1, \dots, m; (p \neq q \iff u_p \neq u_q)\}$ as a *TSP path* (of the TSPAG).

A *TSP path* is illustrated in Figure 4.

Remark 3

1. The *TSP paths* of the TSPAG are in a one-to-one correspondence with the TSP tours of the TSP.
2. Every extreme point of the Linear Assignment Problem (LAP) polytope defined over the TSPAG (\mathcal{A}_n) corresponds to exactly one TSP tour and vice versa.

2.3 Overview of the *barriers*

2.3.1 Basic results

Definition 4 (*Extended Formulation (EF)*; Yannakakis [1991]) A polytope

$$V := \left\{ \begin{pmatrix} u \\ v \end{pmatrix} \in \mathbb{R}^{\mu+\nu} : E^=u + F^=v = g^=, E^{\leq}u + F^{\leq}v \leq g^{\leq} \right\}$$

is an EF of

$$U := \{u \in \mathbb{R}^{\mu} : Au \leq b\}$$

if

$$\pi_u(V) = \left\{ u \in \mathbb{R}^{\mu} : \left(\exists v \in \mathbb{R}^{\nu} : \begin{pmatrix} u \\ v \end{pmatrix} \in V \right) \right\} = U.$$

(where $E^= \in \mathbb{R}^{m^=\times\mu}$, $F^= \in \mathbb{R}^{m^=\times\nu}$, $g^= \in \mathbb{R}^{m^=}$,
 $E^{\leq} \in \mathbb{R}^{m^{\leq}\times\mu}$, $F^{\leq} \in \mathbb{R}^{m^{\leq}\times\nu}$, and $g^{\leq} \in \mathbb{R}^{m^{\leq}}$.)

Definition 5 (“Symmetric Polytope”; Yannakakis [1991]) :

1. A polytope U in the space of *travel leg* variables $u = (u_{ij})$ and auxiliary variables v is symmetric if every given permutation of the TSP nodes can be also extended to the auxiliary variables v so that U remains invariant.
2. A set of linear constraints is called symmetric if its feasible space is.

Theorem 6 (*Barrier #1*; Yannakakis [1991]) “The TSP polytope does not have a *symmetric* EF of subexponential size.”

Theorem 7 (*Barrier #2*; Fiorini et al. [2015]) “The extension complexity of the TSP polytope $\text{TSP}(n)$ is $2^{\Omega(\sqrt{n})}$.”

(This statement of *Barrier #2* can be paraphrased to make it “line up” more directly with Yannakakis’ as: “The TSP polytope does not have an EF of subexponential size.”)

2.3.2 Inferences and Misconceptions

Although the discussions in this section do not have any bearing (or only a peripheral one at best) on the developments in this paper, we believe they highlight the importance of the point of the paper, namely, that the finding of the *barriers* above (Theorems 6-7) does not imply that it is “impossible” for a polynomial-sized LP to solve the TSP optimization problem, as suggested in Fiorini et al. [2015] in particular.

As indicated earlier, the purpose of this paper is not dispute the validity of the *barriers* themselves, but to show that some of the inferences that can be/have been made from them about what is/is not possible in terms of the LP modeling of the TSP optimization problem in general are overscoped. We will use the equivalence between the notions of *extension* and *extended formulation* discussed in Fiorini et al. [2015] for example to provide a rough illustration of this, as follows.

Claim 8 (Reasoning) : Let u be arbitrary variables which are disjoint from the *travel leg* variables, y , of “the TSP polytope” \mathcal{P}_n . Let there be a claim that a linear program (LP) over

$$U := \{u \in \mathbb{R}^\mu : Au \leq b\} \quad (\text{where } \mu \in \mathbb{N})$$

models the (n -city) TSP exactly. One would arrive at a possibly incorrect assessment of this claim by using the following line of reasoning:

- If there exists a linear map $y - Bu = 0$ between $u \in U$ and $y \in \mathcal{P}_n$, then

$$\bar{U} := \left\{ \begin{pmatrix} u \\ y \end{pmatrix} \in \mathbb{R}^{\mu+n(n-1)} : Au \leq b; y - Bu = 0 \right\}$$

is an EF of \mathcal{P}_n , and hence, U cannot be polynomial-sized.

- If there does not exist a linear map $y - Bu = 0$ between $u \in U$ and $y \in \mathcal{P}_n$, then U cannot be an EF of \mathcal{P}_n , and hence, LP cannot solve the TSP.
- Hence, one concludes that the claim cannot be true, which may not be the correct conclusion, as we will develop in this paper.

Remark 9 (Misconceptions) Some reasons why using the *extension*-based EF illustrated above in particular in making inferences about model sizes can lead to overscoped/erroneous conclusions are as follows:

1. The lack of distinction between what is/is not a meaningful EF in the context of making inferences about model sizes;
2. Inferences about solvability for an optimization problem being made without consideration to the expression of the objective function;
3. The significance of superfluousness (“redundancy”) from an optimization problem perspective being overlooked;

4. The implicit presumption that the TSP optimization problem in its natural form (i.e., without reduction to the optimization version of another NPC problem) can be solved only through “*the* TSP polytope”.

We will show in the remainder of this paper that provided a given polytope projects to the LAP polytope, \mathcal{A}_n , and subject to the possibility of expressing TSP tour costs as linear functions of its descriptive variables, an LP over that polytope (or an EF of it) could solve the TSP optimization problem, irrespective of whether it (or any EF of it) projects to “*the* TSP polytope,” \mathcal{P}_n , or not.

3 Side-stepping of the *barriers*

The key to “escaping from” the *barriers* lies in the judicious exploitation of the one-to-one correspondence that exists between the TSP tours and the extreme points of the LAP polytope \mathcal{A}_n (and equivalently, the *TSP paths* of the TSPAG). Because of this one-to-one correspondence, if tour costs can be correctly modeled by a linear function of the auxiliary variables of an EF of \mathcal{A}_n , then the TSP optimization problem can be solved without recourse to “*the* TSP polytope,” using that EF. Our developments to show this consist essentially of examining the *extensions* of the linear assignment problem (LAP) polytope summarized in Table 1.

<p><i>Polytope</i> \mathcal{A}_n :</p> $\sum_{r=1}^m w_{ir} = 1; (\forall i = 1, \dots, m).$ $\sum_{i=1}^m w_{ir} = 1; (\forall r = 1, \dots, m).$ $w_{ir} \geq 0; (\forall i, r = 1, \dots, m).$	<p><i>Polytope</i> Q_0 :</p> $w \in \mathcal{A}_n.$ $\mathbf{A}w + \mathbf{B}x = \mathbf{b}.$ $(\mathbf{A} \in \mathbb{R}^{\alpha \times m^2}; \mathbf{B} \in \mathbb{R}^{\alpha \times \kappa}; \mathbf{b} \in \mathbb{R}^{\alpha}.)$
<p><i>Polytope</i> Q_1 :</p> $w \in \mathcal{A}_n.$ $\mathbf{E}^=w + \mathbf{F}^=y = \mathbf{f}^=.$ $\mathbf{E}^{\leq}w + \mathbf{F}^{\leq}y \leq \mathbf{f}^{\leq}.$ $(\mathbf{E}^= \in \mathbb{R}^{\beta^= \times m^2}; \mathbf{F}^= \in \mathbb{R}^{\beta^= \times \ell};$ $\mathbf{f}^= \in \mathbb{R}^{\beta^=}; \mathbf{E}^{\leq} \in \mathbb{R}^{\beta^{\leq} \times m^2};$ $\mathbf{F}^{\leq} \in \mathbb{R}^{\beta^{\leq} \times \ell}; \mathbf{f}^{\leq} \in \mathbb{R}^{\beta^{\leq}}.)$	<p><i>Polytope</i> Q_2 :</p> $\begin{pmatrix} w \\ x \end{pmatrix} \in Q_0.$ $\mathbf{G}^=w + \mathbf{H}^=y = \mathbf{h}^=.$ $\mathbf{G}^{\leq}w + \mathbf{H}^{\leq}y \leq \mathbf{h}^{\leq}.$ $(\mathbf{G}^= \in \mathbb{R}^{\gamma^= \times m^2}; \mathbf{H}^= \in \mathbb{R}^{\gamma^= \times \ell};$ $\mathbf{h}^= \in \mathbb{R}^{\gamma^=}; \mathbf{G}^{\leq} \in \mathbb{R}^{\gamma^{\leq} \times m^2};$ $\mathbf{H}^{\leq} \in \mathbb{R}^{\gamma^{\leq} \times \ell}; \mathbf{h}^{\leq} \in \mathbb{R}^{\gamma^{\leq}}.)$

Table 1: Statements of the Polytopes

Polytope \mathcal{A}_n is the LAP polytope explicitly stated in the space of the w variables. *Polytope* Q_0 is in the space of w and its “auxiliary” variables x . *Polytope* Q_1 is in the space of w and TSP *travel leg* variables y . *Polytope* Q_2 is in the space of all of the three classes of variables w , x and y .

The optimization problems associated with the polytopes described above (Table 1) are summarized in Table 2, where $c \in \mathbb{R}^{m^2}$ and $\tilde{c} \in \mathbb{R}^{\ell}$. *Problem LAP* is the “standard” linear assignment problem (LAP) stated in the space of w . *Problem LP₀* seeks to minimize a linear function of x over Q_0 . *Problem LP₁* seeks to minimize a linear function of w over Q_1 . *Problem LP₂* also seeks to minimize a linear function of x , but the applicable constraints set is Q_2 .

<p><i>Problem LAP :</i></p> <p>Minimize : $c^t w$</p> <p>Subject to: $w \in \mathcal{A}_n$.</p>	<p><i>Problem LP₀ :</i></p> <p>Minimize : $\tilde{c}^t x$</p> <p>Subject to: $\begin{pmatrix} w \\ x \end{pmatrix} \in Q_0$.</p>
<p><i>Problem LP₁ :</i></p> <p>Minimize : $c^t w$</p> <p>Subject to: $\begin{pmatrix} w \\ y \end{pmatrix} \in Q_1$.</p>	<p><i>Problem LP₂ :</i></p> <p>Minimize : $\tilde{c}^t x$</p> <p>Subject to: $\begin{pmatrix} w \\ x \\ y \end{pmatrix} \in Q_2$.</p>

Table 2: Statements of the Optimization Problems

Observe that because costs are not put on the *travel leg* variables, y , if *Polytope* Q_1 is assumed to be an EF of *Polytope* \mathcal{A}_n , then *Problem LAP* and *Problem LP₁* would be equivalent optimization problems in the sense that they would have the same set of optimal w -variables and the same objective value. Similarly, since costs are not put on the *travel leg* variables, y , in *Problem LP₂*, *Problem LP₂* and *Problem LP₀* would be equivalent optimization problems if Q_2 is an EF of Q_0 .

The equivalence between *Problem LP₂* and *Problem LP₀* is the one that is essential in our argumentation. The equivalence between *Problem LAP* and *Problem LP₁*, however, is particularly easy to use in developing an illustration, as we will do next.

3.1 A motivating illustrative example

The example we discuss has the general form of *Problem LP₁*. The polytope involved is as follows:

Polytope \overline{Q}_1 :

$$\sum_{r=1}^m w_{ir} = 1; (\forall i = 1, \dots, m). \quad (1)$$

$$\sum_{i=1}^m w_{ir} = 1; (\forall r = 1, \dots, m). \quad (2)$$

$$w_{ir} \geq 0; (\forall i, r = 1, \dots, m). \quad (3)$$

$$y_{0,i} - w_{i,1} = 0; (\forall i = 1, \dots, m). \quad (4)$$

$$y_{i,0} - w_{i,m} = 0; (\forall i = 1, \dots, m). \quad (5)$$

$$w_{ir} + w_{j,r+1} - y_{ij} \leq 1; (\forall i, r, j = 1, \dots, m (r \leq m - 1 \wedge i \neq j)). \quad (6)$$

$$\sum_{i=1}^m \sum_{j=1; j \neq i}^m y_{ij} = m - 1. \quad (7)$$

$$y_{ij} \geq 0; (\forall i, j = 0, \dots, m). \quad (8)$$

Constraints (1) – (3) are those of *Polytope* \mathcal{A}_n . They are stated again for the sake of completeness and convenience. The optimization problem associated with this polytope is *Problem LAP*, as shown in Table 2. Constraints (4) – (8) are a special case of the additional/extension constraints of *Polytope* Q_1 . We will denote this special case of Q_1 (i.e., constraints (1) – (8)) and its associated optimization problem by \overline{Q}_1 and \overline{LP}_1 , respectively. In other words, *Problem* \overline{LP}_1 is as follows:

Problem \overline{LP}_1 :

$$\left| \begin{array}{l} \text{Minimize : } \sum_{i=1}^m \sum_{r=1}^m c_{ir} w_{ir} \\ \text{Subject to: (1) – (8).} \end{array} \right.$$

We have the following results.

Lemma 10 \overline{Q}_1 is an EF of \mathcal{A}_n .

Proof. We want to show that

$$\pi_w(\overline{Q}_1) := \left\{ w \in \mathbb{R}^{m^2} : \left(\exists y \in \mathbb{R}^{nm} : \begin{pmatrix} w \\ x \end{pmatrix} \in \overline{Q}_1 \right) \right\} = \mathcal{A}_n.$$

(a) Since the descriptive constraints of \mathcal{A}_n ((1)-(3)) are included in the description of \overline{Q}_1 ,

$$\begin{pmatrix} w \\ x \end{pmatrix} \in \overline{Q}_1 \implies w \in \mathcal{A}_n. \quad (9)$$

(9) implies:

$$\pi_w(\overline{Q}_1) \subseteq \mathcal{A}_n. \quad (10)$$

(b) Let $\widehat{w} \in \text{Ext}(\mathcal{A}_n)$. Let $\widehat{y} \in \mathbb{R}^{nm}$ be specified in terms of \widehat{w} as follows:

$$\forall i \in \{1, \dots, m\}, \widehat{y}_{0,i} = \begin{cases} 1 & \text{if } \widehat{w}_{i,1} = 1 \\ 0 & \text{otherwise} \end{cases} \quad (11)$$

$$\forall i \in \{1, \dots, m\}, \widehat{y}_{i,0} = \begin{cases} 1 & \text{if } \widehat{w}_{i,m} = 1 \\ 0 & \text{otherwise} \end{cases} \quad (12)$$

$$\forall i, j \in \{1, \dots, m\}, \widehat{y}_{ij} = \begin{cases} 1 & \text{if } \exists r \in \{1, \dots, m-1\} : \widehat{w}_{ir} = \widehat{w}_{j,r+1} = 1. \\ 0 & \text{otherwise} \end{cases} \quad (13)$$

Observe that (11)-(13), (1)-(3), and the integrality of \widehat{w} imply:

$$\exists! i \in C : \widehat{y}_{0,i} = 1; \quad (14)$$

$$\exists! i \in C : \widehat{y}_{i,0} = 1; \quad (15)$$

$$\forall r \in \{1, \dots, m-1\}, \exists!(i, j) \in C^2 : \widehat{y}_{ij} = 1. \quad (16)$$

Observe also that these y -variables (i.e., as set according to (11)-(13)) also satisfy (7). Hence, $\begin{pmatrix} \widehat{w} \\ \widehat{y} \end{pmatrix}$ satisfies (1)-(8). In other words, we have:

$$\begin{pmatrix} \widehat{w} \\ \widehat{y} \end{pmatrix} \in \overline{Q}_1. \quad (17)$$

(17) and the arbitrariness of \widehat{w} imply:

$$\widehat{w} \in \text{Ext}(\mathcal{A}_n) \implies \exists \widehat{y} \in \mathbb{R}^{nm} : \begin{pmatrix} \widehat{w} \\ \widehat{y} \end{pmatrix} \in \overline{Q}_1. \quad (18)$$

(18) implies:

$$\text{Conv}(\text{Ext}(\mathcal{A}_n)) = \mathcal{A}_n \subseteq \pi_w(\overline{Q}_1). \quad (19)$$

(c) (10) and (19) imply:

$$\pi_w(\overline{Q}_1) = \mathcal{A}_n.$$

■

Lemma 11 \overline{Q}_1 is not an EF of \mathcal{P}_n . Moreover, \overline{Q}_1 is non-integral.

Proof. First, we will show that:

$$\begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \not\Rightarrow y \in \mathcal{P}_n. \quad (20)$$

The proof is by construction. Consider the special case of \overline{Q}_1 for a 5-city TSP. Consider $\begin{pmatrix} \overline{w} \\ \overline{y} \end{pmatrix} \in \mathbb{R}_{\geq 0}^{m^2+nm}$ which has the following non-zero components:

$$\begin{cases} \overline{w}_{1,1} = \overline{w}_{2,2} = \overline{w}_{4,3} = \overline{w}_{3,4} = 0.25; \\ \overline{w}_{4,1} = \overline{w}_{3,2} = \overline{w}_{1,3} = \overline{w}_{2,4} = 0.75; \\ \overline{y}_{0,1} = \overline{y}_{3,0} = 0.25; \overline{y}_{0,4} = \overline{y}_{2,0} = 0.75; \\ \overline{y}_{3,1} = \overline{y}_{1,2} = 0.50; \overline{y}_{4,3} = 2.0. \end{cases} \quad (21)$$

Observe that $\begin{pmatrix} \overline{w} \\ \overline{y} \end{pmatrix}$ satisfies (1) – (8) for $n = 5$. Also, since every point of “the TSP polytope” has coordinates between 0 and 1, $\overline{y}_{4,3}$ being greater than 1 ($\overline{y}_{4,3} = 2 > 1$), in particular, implies that $\overline{y} \notin \mathcal{P}_5$. Statement (20) follows directly from these observations.

Focusing back on \mathcal{P}_n and \overline{Q}_1 in general, observe that every feasible solution to (1) – (8) is such that its y -vector component is integral if and only if its w -vector component is integral. In other words, it can be observed that the following is true:

$$\begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \implies \left(w \in \{0, 1\}^{m^2} \iff y \in \{0, 1\}^{nm} \right). \quad (22)$$

The combination of the fact that every extreme point of \mathcal{A}_n is an integral w -vector, statement (22), and Remark 3 gives that:

$$\left(\left\{ y \in \{0, 1\}^{nm} : \left(\exists w \in \{0, 1\}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\} \right) \text{ is} \\ \text{in a one-to-one correspondence with the set of TSP tours.} \quad (23)$$

(23) implies:

$$\mathcal{P}_n = \text{Conv} \left\{ y \in \{0, 1\}^{nm} : \left(\exists w \in \{0, 1\}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\}. \quad (24)$$

(24) implies:

$$\mathcal{P}_n \subseteq \left\{ y \in \mathbb{R}^{nm} : \left(\exists w \in \mathbb{R}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\}. \quad (25)$$

From (20), we have:

$$\left\{ y \notin \mathcal{P}_n : \left(\exists w \in \mathbb{R}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\} \neq \emptyset. \quad (26)$$

(25) and (26) imply:

$$\mathcal{P}_n \subset \left\{ y \in \mathbb{R}^{mn} : \left(\exists w \in \mathbb{R}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\}. \quad (27)$$

Hence,

$$\left\{ y \in \mathbb{R}^{mn} : \left(\exists w \in \mathbb{R}^{m^2} : \begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1 \right) \right\} \neq \mathcal{P}_n. \quad (28)$$

Hence, \overline{Q}_1 is not an EF of \mathcal{P}_n .

Finally (thirdly), we will show that \overline{Q}_1 is non-integral. To that effect, let $I(\overline{Q}_1)$ be the set of integral solutions of \overline{Q}_1 . Then, we have the following.

Statements (24) and (27) imply:

$$\overline{Q}_1 \setminus \text{Conv}(I(\overline{Q}_1)) \neq \emptyset. \quad (29)$$

Since every feasible point of a polytope must be a convex combination of extreme points of that polytope (Bazaraa et al. [2010; pp. 70-82]), and since $\begin{pmatrix} \tilde{w} \\ \tilde{y} \end{pmatrix} \in \overline{Q}_1 \setminus \text{Conv}(I(\overline{Q}_1))$ cannot be a convex combination of integral points of \overline{Q}_1 (since $\begin{pmatrix} \tilde{w} \\ \tilde{y} \end{pmatrix} \notin \text{Conv}(I(\overline{Q}_1))$), at least one of the extreme points of \overline{Q}_1 with positive weights in the representation of $\begin{pmatrix} \tilde{w} \\ \tilde{y} \end{pmatrix}$ must be non-integral. Hence, \overline{Q}_1 must have at least one non-integral extreme point. Hence, \overline{Q}_1 is non-integral. ■

The following result is proved more generally in the next section for *Problem LAP* and *Problem LP₁* (Theorem 17 of section 3.2). Hence, its proof is omitted in this section.

Lemma 12 $\begin{pmatrix} w^* \\ y^* \end{pmatrix}$ is optimal for Problem \overline{LP}_1 iff w^* is optimal for *Problem LAP*.

The main take-aways from this section are summarized in the following remark.

Remark 13

1. Polytope \overline{Q}_1 is an EF of the LAP polytope, \mathcal{A}_n .
2. Problem \overline{LP}_1 and Problem *LAP* have the same objective value and the same set of optimal w variables.
3. If the optimum in (2) above is unique, it is integral and must be obtained whether Problem *LAP* or Problem \overline{LP}_1 is used/solved.

4. If there are multiple optima to Problem \overline{LP}_1 (or equivalently to Problem LAP), then:
- (a) There must exist more than one integral extreme point of \overline{Q}_1 (\mathcal{A}_n) that is optimal for *Problem LP_1* (*Problem LAP*);
 - (b) A fractional solution which is a convex combination of the optimal extreme points in (a) above for \overline{Q}_1 (i.e., an interior point of \overline{Q}_1) may be obtained when *Problem LP_1* is solved.
 - (c) The w -component of any fractional solution of \overline{Q}_1 obtained as described in (b) above would be an interior point of \mathcal{A}_n , and would hence be a convex combination of integral w 's. Each of these integral w 's would correspond to exactly one TSP tour.
5. Statements (1)-(4) of Remark 13 above are true, irrespective of whether an EF of *Polytope \overline{Q}_1* projects to “*the TSP polytope,*” \mathcal{P}_n , or not.

The following example illustrates the fact that an LP with feasible set \overline{Q}_1 (or an EF of it) could solve the TSP optimization problem despite the fact that \overline{Q}_1 is not integral and *does not project* to “*the TSP polytope,*” \mathcal{P}_n . This exemplifies the fact that an LP could solve the TSP optimization problem, irrespective of whether it projects to \mathcal{P}_n or not, provided it project to \mathcal{A}_n .

Example 14 Re-consider the special case of \overline{Q}_1 for a 5-city TSP. Consider the solution instance shown in (21). We will show that this solution can be associated with TSP tours by focusing on \overline{w} . To this effect, let \widehat{w}^1 and \widehat{w}^2 be extreme points of \mathcal{A}_5 with positive components as follows:

$$\begin{cases} \widehat{w}_{1,1}^1 = \widehat{w}_{2,2}^1 = \widehat{w}_{4,3}^1 = \widehat{w}_{3,4}^1 = 1; \\ \widehat{w}_{4,1}^2 = \widehat{w}_{3,2}^2 = \widehat{w}_{1,3}^2 = \widehat{w}_{2,4}^2 = 1. \end{cases}$$

Then, by the one-to-one correspondence between TSP tours and the extreme points of \mathcal{A}_5 , there are unique TSP tours associated with \widehat{w}^1 and \widehat{w}^2 , respectively. The tour represented by \widehat{w}^1 , $\tau(\widehat{w}^1)$, is: $0 \rightarrow 1 \rightarrow 2 \rightarrow 4 \rightarrow 3 \rightarrow 0$. Similarly, the tour, $\tau(\widehat{w}^2)$, represented by \widehat{w}^2 is: $0 \rightarrow 4 \rightarrow 3 \rightarrow 1 \rightarrow 2 \rightarrow 0$.

Observe that $\overline{w} = 0.25\widehat{w}^1 + 0.75\widehat{w}^2$ is a convex combination of \widehat{w}^1 and \widehat{w}^2 . Hence, $\begin{pmatrix} \overline{w} \\ \overline{y} \end{pmatrix}$ would be associated with $\tau(\widehat{w}^1)$ and $\tau(\widehat{w}^2)$ and their convex combination in any optimization problem over \overline{Q}_1 in which the objective function is a linear function of w .

Observe also that the associations above for $\begin{pmatrix} \overline{w} \\ \overline{y} \end{pmatrix}$ analogously hold for every $\begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1$. In other words, every $\begin{pmatrix} w \\ y \end{pmatrix} \in \overline{Q}_1$ would be associated to a number $\eta \in \mathbb{N}$ of TSP tours $\tau_1(w), \dots, \tau_\eta(w)$ and their convex combination in any optimization problem over \overline{Q}_1 in which the objective function is a linear function of w . Hence, if costs can be attached to w through a linear function in such a way that the total cost incurred at an extreme point of \mathcal{A}_n , \widehat{w}^i ($i \in \{1, \dots, m!\}$), is equal to the total cost of the travels involved in the associated

TSP tour, $\tau(\widehat{w}^i)$, then the TSP optimization problem would be solved as an LP over \overline{Q} . We believe it is not possible to “construct” such a cost association using w directly. However, as we will illustrate in the next section (Theorem 22), it is possible to “construct” such a cost association indirectly, using auxiliary variables which are based on w only. A *sufficient* condition for the model which includes these auxiliary variables (whatever they may end up being) to be able to solve the TSP optimization problem is that it (the model) be an EF of \mathcal{A}_n . Observe that, in other words, whether such an EF of \overline{Q}_1 projects to “the TSP polytope” (\mathcal{P}_n) or not is irrelevant to its being able to solve the TSP optimization problem.

3.2 General results

In this section, we will formally develop our thesis that a finding that there exists no polynomial-sized integral extended formulation of “the TSP polytope,” \mathcal{P}_n , does not imply that it is “impossible” for a polynomial-sized linear program to solve the TSP optimization problem. The general “flow” of our argumentation is as follows:

(a) The extreme points of the LAP polytope, \mathcal{A}_n , are in one-to-one correspondence with the TSP tours.

(b) *Provided* the descriptive variables for \mathcal{P}_n , y , and the constraints involving them in *Problem LP₂* can be discarded from *Problem LP₂* (hence reducing *Problem LP₂* to *Problem LP₀*) without changing the optimization problem at hand, *Problem LP₂* is an equivalent optimization problem to *Problem LP₀*. This equivalence is true regardless of whether *Polytope Q₁* projects to *Polytope P_n* or not.

(In the sense above, the y -variables and the constraints which involve them in *Problem LP₂* would be “redundant” for *Problem LP₂* (and equivalently for *Problem LP₀*), and they could be thought of as only serving “bookkeeping” purposes in *Problem LP₂*. Also, a sufficient condition for this *redundancy* is that Q_2 be an EF of Q_0 .)

(c) The combination of (a) and (b) above implies that *Problem LP₀* would solve the TSP optimization problem, *provided* the following conditions are satisfied:

(i) The extreme points of *Polytope Q₀* are in one-to-one correspondence with the extreme points of *Polytope A_n*;

(ii) The vector \tilde{c} is such that it “attaches” to any given extreme point of Q_0 , the cost of the TSP tour associated with that extreme point (i.e., for each $\widehat{x} \in \text{Ext}(Q_0)$, $\tilde{c}^t \widehat{x}$ is equal to the cost of the TSP tour associated with \widehat{x}).

(d) Hence, it is true that *Problem LP₀* would solve the TSP optimization problem, regardless of whether *Polytope Q₁* projects to “the TSP polytope,” \mathcal{P}_n , or not, *provided* the conditions specified in (b) and (c) above are satisfied.

(e) Hence, *provided* the conditions specified in (b) and (c) above are satisfied, “the TSP polytope” \mathcal{P}_n (and hence, the *barriers*) can be side-stepped via *Problem LP₀* in the process of solving the TSP optimization problem.

These ideas will now be developed formally. We start with the equivalence between *Problem LAP* and *Problem LP₁* illustrated in section 3.1, although this is not essential to our argumentation. We will henceforth assume the following extension relationships.

Assumption 15 We will henceforth assume that:

$$1. \pi_w(Q_0) = \left\{ w \in \mathbb{R}^{m^2} : \left(\exists x \in \mathbb{R}^k : \begin{pmatrix} w \\ x \end{pmatrix} \in Q_0 \right) \right\} = \mathcal{A}_n.$$

(Q_0 is an extended formulation (EF) of \mathcal{A}_n .)

$$2. \pi_w(Q_1) = \left\{ w \in \mathbb{R}^{m^2} : \left(\exists y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ y \end{pmatrix} \in Q_1 \right) \right\} = \mathcal{A}_n.$$

(Q_1 is an EF of \mathcal{A}_n .)

$$3. \pi_w(Q_2) = \left\{ w \in \mathbb{R}^{m^2} : \left(\exists \begin{pmatrix} x \\ y \end{pmatrix} \in \mathbb{R}^{k+\ell} : \begin{pmatrix} w \\ x \\ y \end{pmatrix} \in Q_2 \right) \right\} = \mathcal{A}_n.$$

(Q_2 is an EF of \mathcal{A}_n .)

$$4. \pi_{(w,x)}(Q_2) = \left\{ \begin{pmatrix} w \\ x \end{pmatrix} \in \mathbb{R}^{m^2+k} : \left(\exists y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ x \\ y \end{pmatrix} \in Q_2 \right) \right\} = Q_0.$$

(Q_2 is an EF of Q_0 .)

Also, the following notations will simplify the exposition.

Notation 16 :

$$1. \forall w \in \mathcal{A}_n, F_0(w) := \left\{ x \in \mathbb{R}^k : \begin{pmatrix} w \\ x \end{pmatrix} \in Q_0 \right\}.$$

$$2. \forall w \in \mathcal{A}_n, F_1(w) := \left\{ y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ y \end{pmatrix} \in Q_1 \right\}.$$

$$3. \forall w \in \mathcal{A}_n, \forall x \in \mathbb{R}^k : \begin{pmatrix} w \\ x \end{pmatrix} \in Q_0, F_2(w, x) := \left\{ y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ x \\ y \end{pmatrix} \in Q_2 \right\}.$$

The next theorem shows the equivalence between *Problem LAP* and *Problem LP₁*, a special case of which was illustrated in section 3.1.

Theorem 17 $w^* \in \mathcal{A}_n$ is optimal for *Problem LAP* if and only if $\begin{pmatrix} w^* \\ y^* \end{pmatrix}$ is optimal for *Problem LP₁* for each $y^* \in F_1(w^*)$.

Proof.

(\implies): Let w^* be an optimal solution for *Problem LAP*. In other words, we have:

$$\forall w^\dagger \in \mathcal{A}_n, c^t w^\dagger \geq c^t w^*. \quad (30)$$

Now, let $y^* \in F_1(w^*)$. Assume $\begin{pmatrix} w^* \\ y^* \end{pmatrix}$ is not optimal for Q_1 . Then, there must exist a feasible solution to *Problem LP₁* which has a lower value of the objective of *Problem LP₁* than that of $\begin{pmatrix} w^* \\ y^* \end{pmatrix}$. In other words, we must have:

$$\exists \begin{pmatrix} w^\dagger \\ y^\dagger \end{pmatrix} \in Q_1 : (c^t w^\dagger + \mathbf{0}^t \cdot y^\dagger) = c^t w^\dagger < (c^t w^* + \mathbf{0}^t \cdot y^*) = c^t w^*. \quad (31)$$

By Assumption 15.2,

$$\begin{pmatrix} w^\dagger \\ y^\dagger \end{pmatrix} \in Q_1 \implies w^\dagger \in \mathcal{A}_n. \quad (32)$$

Hence, (31) contradicts (30). Hence, we must have:

$$w^* \text{ optimal for } \textit{Problem LAP} \implies \begin{pmatrix} w^* \\ y^* \end{pmatrix} \text{ optimal for } LP_1 \text{ for each } y^* \in F_1(w^*).$$

(\impliedby): Assume $\begin{pmatrix} w^* \\ y^* \end{pmatrix} \in Q_1$ is optimal for *Problem LP₁* for each $y^* \in F_1(w^*)$. In other words, assume the following is true:

$$\forall \begin{pmatrix} w^\dagger \\ y^\dagger \end{pmatrix} \in Q_1, (c^t w^\dagger + \mathbf{0}^t \cdot y^\dagger) = c^t w^\dagger \geq (c^t w^* + \mathbf{0}^t \cdot y^*) = c^t w^*. \quad (33)$$

Now, assume w^* is not optimal for *Problem LP₁*. Then, we must have:

$$\forall y^* \in F_1(w^*), \exists w^\dagger \in \mathcal{A}_n : c^t w^\dagger < c^t w^* = (c^t w^* + \mathbf{0}^t \cdot y^*). \quad (34)$$

By Assumption 15.2,

$$\begin{pmatrix} w^* \\ y^* \end{pmatrix} \in Q_1 \implies w^* \in \mathcal{A}_n. \quad (35)$$

Hence, statement (34) contradicts statement (33). Hence, we must have:

$$\begin{aligned} & \forall w^* \in \mathcal{A}_n, \forall y^* \in F_1(w^*), \begin{pmatrix} w^* \\ y^* \end{pmatrix} \text{ optimal for } \textit{Problem LP}_1 \implies \\ & w^* \text{ optimal for } \textit{Problem LAP}. \end{aligned} \quad (36)$$

■

Similar to the relation of *Problem LP₁* to *Problem LAP*, *Problem LP₂* is equivalent to *Problem LP₀*. This is the object of the next theorem.

Theorem 18 $\begin{pmatrix} w^* \\ x^* \end{pmatrix} \in Q_0$ is optimal for *Problem LP₀* if and only if $\begin{pmatrix} w^* \\ x^* \\ y^* \end{pmatrix}$ is optimal for *Problem LP₂* for each $y^* \in F_2(w^*, x^*)$.

Proof. The proof steps are similar to those of Theorem 17.

(\implies): Let $\begin{pmatrix} w^* \\ x^* \end{pmatrix} \in Q_0$ be an optimal solution for *Problem LP₀*. In other words, let the following be true:

$$\forall \begin{pmatrix} w^\dagger \\ x^\dagger \end{pmatrix} \in Q_0, \quad (\mathbf{0}^t \cdot w^\dagger + \tilde{c}^t x^\dagger) = \tilde{c}^t x^t \geq (\mathbf{0}^t \cdot w^* + \tilde{c}^t x^*) = \tilde{c}^t x^*. \quad (37)$$

Let $y^* \in F_2(w^*, x^*)$. Assume $\begin{pmatrix} w^* \\ x^* \\ y^* \end{pmatrix}$ is not optimal for Q_2 . In other words, assume the following is true:

$$\begin{aligned} \exists \begin{pmatrix} w^\dagger \\ x^\dagger \\ y^\dagger \end{pmatrix} \in Q_2 : (\mathbf{0}^t \cdot w^\dagger + \tilde{c}^t x^\dagger + \mathbf{0}^t \cdot y^\dagger) &= \tilde{c}^t x^t \\ &< (\mathbf{0}^t \cdot w^* + \tilde{c}^t x^* + \mathbf{0}^t \cdot y^*) = \tilde{c}^t x^*. \end{aligned} \quad (38)$$

By Assumption 15.2,

$$\begin{pmatrix} w^\dagger \\ x^\dagger \\ y^\dagger \end{pmatrix} \in Q_2 \implies \begin{pmatrix} w^\dagger \\ x^\dagger \end{pmatrix} \in Q_0. \quad (39)$$

Hence, (38) contradicts (37). Hence, we must have:

$$\begin{aligned} \begin{pmatrix} w^* \\ x^* \end{pmatrix} \text{ optimal for } \textit{Problem LP}_0 &\implies \\ \begin{pmatrix} w^* \\ x^* \\ y^* \end{pmatrix} \text{ optimal for } \textit{Problem LP}_2 &\text{ for each } y^* \in F_2(w^*, x^*). \end{aligned} \quad (40)$$

(\impliedby): Assume $\begin{pmatrix} w^* \\ x^* \\ y^* \end{pmatrix} \in Q_2$ is optimal for *Problem LP₂* for each $y^* \in F_2(w^*, x^*)$. In other words, assume we have:

$$\begin{aligned} \forall y^* \in F_2(w^*, x^*), \forall \begin{pmatrix} w^\dagger \\ x^\dagger \\ y^\dagger \end{pmatrix} \in Q_2, \quad (\mathbf{0}^t \cdot w^\dagger + \tilde{c}^t x^\dagger + \mathbf{0}^t \cdot y^\dagger) &= \tilde{c}^t x^t \\ &\geq (\mathbf{0}^t \cdot w^* + \tilde{c}^t x^* + \mathbf{0}^t \cdot y^*) = \tilde{c}^t x^*. \end{aligned} \quad (41)$$

Now, assume $\begin{pmatrix} w^* \\ x^* \end{pmatrix}$ is not optimal for *Problem LP₀*. Then, we must have:

$$\exists \begin{pmatrix} w^\dagger \\ x^\dagger \end{pmatrix} \in Q_0 : (\mathbf{0}^t \cdot w^\dagger + \tilde{c}^t x^\dagger) = \tilde{c}^t x^\dagger < (\mathbf{0}^t \cdot w^* + \tilde{c}^t x^*) = \tilde{c}^t x^*. \quad (42)$$

By Assumption 15.2,

$$\begin{pmatrix} w^\dagger \\ x^\dagger \\ y^\dagger \end{pmatrix} \in Q_2 \implies \begin{pmatrix} w^\dagger \\ x^\dagger \end{pmatrix} \in Q_0. \quad (43)$$

Hence, statement (42) contradicts statement (41). Hence, we must have:

$$\begin{aligned} & \begin{pmatrix} w^* \\ x^* \\ y^* \end{pmatrix} \text{ optimal for } \textit{Problem LP}_2 \text{ for each } y^* \in F_2(w^*, x^*) \implies \\ & \begin{pmatrix} w^* \\ x^* \end{pmatrix} \text{ optimal for } \textit{Problem LP}_0. \end{aligned} \quad (44)$$

■

Our central thesis will now be formally stated.

Theorem 19 Assume that the extreme points of *Polytope* Q_0 are in one-to-one correspondence with the extreme points of the LAP polytope, \mathcal{A}_n . Let $\hat{w} \in \text{Ext}(\mathcal{A}_n)$ be the extreme point of \mathcal{A}_n associated with $\hat{x} \in \mathbb{R}^k$ which is such that $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} \in \text{Ext}(Q_0)$. Let $\tau(\hat{w})$ be the TSP tour corresponding to \hat{w} . Assume \tilde{c} is such that $\tilde{c}^t \hat{x}$ is equal to the cost of the travels involved in $\tau(\hat{w})$. Then, *Problem LP₀* correctly solves the TSP optimization problem, irrespective of Q_0 projecting to “the TSP polytope” (\mathcal{P}_n) or not.

Proof. The (premised) one-to-one correspondence between $\text{Ext}(\mathcal{A}_n)$ and $\text{Ext}(Q_0)$ and the one-to-one correspondence between $\text{Ext}(\mathcal{A}_n)$ and TSP tours imply (by transitivity):

$$\text{There exists a one-to-one correspondence between } \text{Ext}(Q_0) \text{ and TSP tours.} \quad (45)$$

Also, for each $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} \in \text{Ext}(Q_0)$, $\hat{w} \in \text{Ext}(\mathcal{A}_n)$, and the unique TSP tour, $\tau(\hat{w})$, associated with \hat{w} must also be associated with $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix}$. Hence, by premise, $\begin{pmatrix} \mathbf{0} \\ \tilde{c} \end{pmatrix}^t \cdot \begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} = \tilde{c}^t \hat{x}$ correctly accounts the total cost of the travels involved in $\tau(\hat{w})$, for each $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} \in \text{Ext}(Q_0)$. The theorem follows directly from the combination of this and Theorem 18. ■

Corollary 20 A finding that “the TSP polytope” does not have a polynomial-sized EF does not imply that there does not exist a polynomial-sized LP which can solve the TSP optimization problem.

Remark 21 If *Polytope* Q_0 is integral in addition to its extreme points being in one-to-one correspondence with the extreme points of *Polytope* \mathcal{A}_n , then there exist a polynomially bounded number of x -variables for which several alternate ways for “constructing” costs \tilde{c} so that TSP tour costs are correctly captured. This is shown in the next theorem. In order not to detract from the focus of this paper, we will leave the (difficult) issue of finding extension constraints of Q_0 (i.e., finding \mathbf{A} , \mathbf{B} , and \mathbf{b}) so that Q_0 is an exact EF of \mathcal{A}_n as an open topic for research.

Theorem 22 For each triplet of nodes of the TSPAG, $([i, p], [j, r], [k, s])$, define a variable $x_{[i,p][j,r][k,s]}$ and $\tilde{c} \in \mathbb{R}^{m^6}$ based on the TSP travel costs, d , as follows:

$$\tilde{c}_{[i,p][j,r][k,s]} := \begin{cases} d_{0i} + d_{ij} + d_{jk} & \text{if } (p = 1; r = 2; s = 3); \\ d_{jk} + d_{k0} & \text{if } (p = 1; r = m - 1; s = m); \\ d_{jk} & \text{if } (p = 1; 3 \leq r \leq m - 2; s = r + 1); \\ 0 & \text{Otherwise.} \end{cases}$$

Assume the following are valid for Q_0 :

- (i) $x_{[i,p][j,r][k,s]} > 0 \implies p < r < s$ and $i \neq j \neq k$;
- (ii) $x_{[i,p][j,r][k,s]} \leq \min(w_{[i,p]}, w_{[j,r]}, w_{[k,s]})$;
- (iii) $w_{[i,p]} + w_{[j,r]} + w_{[k,s]} - x_{[i,p][j,r][k,s]} \leq 2$.

Then, for each $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} \in \text{Ext}(Q_0)$,

$$\mathcal{V}(w, x) := \begin{pmatrix} \mathbf{0} \\ \tilde{c} \end{pmatrix}^t \cdot \begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} = \sum_{([i,p],[j,r],[k,s]) \in N^3} \tilde{c}_{[i,p][j,r][k,s]} \hat{x}_{[i,p][j,r][k,s]}$$

correctly accounts the total cost of the travels involved in the TSP tour associated with \hat{w} and $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix}$.

Proof. Let $\begin{pmatrix} \hat{w} \\ \hat{x} \end{pmatrix} \in \text{Ext}(Q_0)$. Then, by premise, $\hat{w} \in \text{Ext}(\mathcal{A}_n)$. Hence, letting the positive entries of \hat{w} be $\hat{w}_{u_1,1}, \hat{w}_{u_2,2}, \dots, \hat{w}_{u_m,m}$, we must have:

$$\begin{aligned} & \forall (p, q) \in S^2 : p \neq q, u_p \neq u_q; \text{ and} \\ & \forall [v, q] \in N, \hat{w}_{vq} = \begin{cases} 1 & \text{if } v = u_q \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (46)$$

Hence, the (unique) TSP tour corresponding to \widehat{w} can be stated as $\tau(\widehat{w}) := 0 \rightarrow u_1^k \rightarrow \dots \rightarrow u_m^k \rightarrow 0$ (as discussed earlier in section 2.2.2). The total cost of the travels involved in $\tau(\widehat{w})$ is:

$$TCost = d_{0,u_1} + d_{u_m,0} + \sum_{q=1}^{m-1} d_{u_q,u_{q+1}}. \quad (47)$$

To account the total cost involved in \widehat{x} , observe that statement (46) and conditions (i) – (iii) imply:

$$\begin{aligned} & \forall ([v_p, p], [v_r, r], [v_s, s]) \in N^3, \\ & \widehat{x}_{[v_p, p][v_r, r][v_s, s]} = \begin{cases} 1 & \text{if } v_p = u_p, v_r = u_r, v_s = u_s, p < r < s; \\ 0 & \text{otherwise.} \end{cases} \end{aligned} \quad (48)$$

Hence, focusing on the components of \widehat{x} and using (48), the costs incurred by \widehat{x} are as follows:

Component, $\widehat{x}_{[u_p, p][u_r, r][u_s, s]}$	Cost, $\widetilde{c}_{[u_p, p][u_r, r][u_s, s]}$
$p = 1; r = 2; s = 3$	$d_{0,u_1} + d_{u_1,u_2^k} + d_{u_2,u_3}$
$p = 1; r = m - 1; s = m$	$d_{u_{m-1},u_m} + d_{u_m,0}$
$p = 1; r = 3; s = 4$	d_{u_3,u_4}
\vdots	\vdots
$p = 1; r = m - 2; s = m - 1$	$d_{u_{m-2},u_{m-1}}$
Total cost attached to $\begin{pmatrix} \widehat{w} \\ \widehat{x} \end{pmatrix} =$	$d_{0,u_1} + d_{u_m,0} + \sum_{q=1}^{m-1} d_{u_q,u_{q+1}}$

Comparing the results above, we observe that the total cost associated with $\begin{pmatrix} \widehat{w} \\ \widehat{x} \end{pmatrix}$ is indeed equal to that of the TSP tour represented by \widehat{w} , $\tau(\widehat{w})$. ■

Hence, in particular, if Q_0 can be described by a polynomial-sized set of constraints in the space of the x -variables specified in Theorem 22 in such a way that it is integral and has extreme points which are in one-to-one correspondence with those of the LAP polytope \mathcal{A}_n , then the TSP optimization problem can be solved as a polynomial sized LP without recourse to “the TSP polytope” \mathcal{P}_n , and thereby side-stepping the existing *barriers*.

4 Conclusions

We have provided conditions under which the TSP optimization problem may be solvable without recourse to “the TSP polytope.” The possibility of these conditions is not ruled out by any existing result, and hence subject to this, we have shown that a finding that

“the TSP polytope” does not have a polynomial-sized *exact* extended formulation (EF) does not imply that it is “impossible” for a linear program (polynomial-sized or not) to solve the TSP optimization problem. Our demonstrations are based on a time-dependent/Assignment Problem representation of TSP tours. We have shown that provided an EF (polynomial-sized or not) of the Linear Assignment Problem (LAP) polytope is *exact* and that TSP tour costs can be accurately accounted through linear functions of its descriptive variables, that EF would solve the TSP optimization problem, irrespective of whether a further EF of it in the space of the city-to-city (*travel leg*) variables projects to “the TSP polytope” or not. We have also proposed variables which are auxiliary to the natural descriptive variables of the LAP polytope and in terms of which TSP tour costs can be correctly captured using linear functions. Also, because the combinatorial configurations of many of the other NP-Complete problems can be modeled as a LAP with “side-constraints,” our developments are directly applicable to many of the other NP-Complete problems.

References

- [1] Aprile, M. (2022). Extended formulations for matroid polytopes through randomized protocols. *Operations Research Letters*, 50:2, pp. 145-149.
- [2] Bazaraa, M.S., J.J. Jarvis and H.D. Sherali (2010). *Linear Programming and Network Flows*. Wiley, New York, NY.
- [3] Burkard, R., M. Dell’Amico, and S. Martello (2009). *Assignment Problems*. SIAM (Philadelphia).
- [4] Conforti, M., G. Cornuéjols, and G. Zambelli (2014). *Integer Programming (Graduate Texts in Mathematics, 271)*. Springer, New York.
- [5] Dantzig, G., R. Fulkerson, S. Johnson (1954). Solution of a Large-Scale Traveling-Salesman Problem. *Journal of the Operations Research Society of America* 2:4, pp. 393–410.
- [6] Faenza, Y., G. Oriolo, G. Stauffer (2021). Separation routine and extended formulations for the stable set problem in claw-free graphs. *Mathematical Programming*, 188:1, pp. 53-84.
- [7] Fiorini, S., S. Massar, S. Pokutta, H.R. Tiwary, and R. de Wolf (2015). Exponential Lower Bounds for Polytopes in Combinatorial Optimization. *Journal of the ACM* 62:2, Article No. 17.
- [8] Garey, M.R. and D.S. Johnson (1979). *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman and Company, New York, NY.
- [9] Karmarkar, N. (1984). A new polynomial-time algorithm for linear programming. *Combinatorica* 4, pp. 373–395.
- [10] Khachiyan, L.G. (1979). A Polynomial Algorithm in Linear Programming. *Doklady Akademii Nauk SSSR* 244, pp. 1093–1096. (English translation: *Soviet Mathematics Doklady* 20:1, pp. 191–194.)
- [11] Kwan, M., L. Saueremann, Y. Zhao (2022). Extension complexity of low-dimensional polytopes. *Transactions of the American Mathematical Society*, 375:6, pp. 4209-4250.
- [12] Lawler, E.L., J.K. Lenstra, A.H.G. Rinnooy Kan, and D.B. Shmoys (1985), eds. *The Traveling Salesman Problem: A Guided Tour of Combinatorial Optimization* (Wiley, New York).
- [13] Yannakakis, M. (1991). Expressing combinatorial optimization problems by linear programming. *Journal of Computer and System Sciences* 43:3, pp. 441-466.

Appendix: Tables of notations

Notation	Meaning
n	Number of TSP cities.
$m := n - 1$	Numbers of <i>levels</i> and <i>stages</i> respectively, of the TSPAG
$C := \{0, \dots, m\}$	Set of cities
$\forall (i, j) \in C^2, d_{ij}$	Cost of traveling from city i to city j
$\ell := n \times m$	Number of arcs in the <i>TSP graph</i>
\mathcal{P}_n	“ <i>the</i> TSP polytope” defined over n cities
$L := \{1, \dots, m\}$	Index set for the <i>levels</i> of the TSPAG
$S := \{1, \dots, m\}$	Index set for the <i>stages</i> of the TSPAG
$N := \{(l, s) \in (L, S)\}$	Set of nodes of the TSPAG
\mathcal{A}_n	The LAP polytope defined over the TSPAG associated with a n -city TSP
$(A)^t$	Transpose of matrix A
$Ext(B)$	Set of extreme points of polyhedron B
$\pi_x(B)$	Projection of polyhedron B on the space of vector x
\mathbb{R}	Set of real numbers
$\mathbb{R}_{\geq 0}$	Set of nonnegative real numbers
\mathbb{N}	Set of natural numbers

Table A.1: List of general notations

Notation	Meaning
$\forall w \in \mathcal{A}_n, F_0(w) := \left\{ x \in \mathbb{R}^k : \begin{pmatrix} w \\ x \end{pmatrix} \in Q_0 \right\}$	Set of x -vectors such that $\begin{pmatrix} w \\ x \end{pmatrix}$ is feasible for Q_0
$\forall w \in \mathcal{A}_n, F_1(w) := \left\{ y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ y \end{pmatrix} \in Q_1 \right\}$	Set of y -vectors such that $\begin{pmatrix} w \\ y \end{pmatrix}$ is feasible for Q_1
$\forall w \in \mathcal{A}_n, \forall x \in \mathbb{R}^k : \begin{pmatrix} w \\ y \end{pmatrix} \in Q_0,$ $F_2(w, x) := \left\{ y \in \mathbb{R}^\ell : \begin{pmatrix} w \\ x \\ y \end{pmatrix} \in Q_2 \right\}$	Set of y -vectors such that $\begin{pmatrix} w \\ x \\ y \end{pmatrix}$ is feasible for Q_2

Table A.2: Feasible sets definitions