"The important thing is not to stop questioning", including the symmetries on which is based the Standard Model

Daniel Sternheimer

To the memory of Moshe Flato and of Noriko Sakurai

Abstract. New fundamental physical theories can, so far a posteriori, be seen as emerging from existing ones via some kind of deformation. That is the basis for Flato's "deformation philosophy", of which the main paradigms are the physics revolutions from the beginning of the twentieth century, quantum mechanics (via deformation quantization) and special relativity. On the basis of these facts we describe two main directions by which symmetries of hadrons (strongly interacting elementary particles) may "emerge" by deforming in some sense (including quantization) the Anti de Sitter symmetry (AdS), itself a deformation of the Poincaré group of special relativity. The ultimate goal is to base on fundamental principles the dynamics of strong interactions, which originated half a century ago from empirically guessed "internal" symmetries. After a rapid presentation of the physical (hadrons) and mathematical (deformation theory) contexts, we review a possible explanation of photons as composites of AdS singletons (in a way compatible with QED) and of leptons as similar composites (massified by 5 Higgs, extending the electroweak model to 3 generations). Then we present a "model generating" multifaceted framework in which AdS would be deformed and quantized (possibly at root of unity and/or in manner not yet mathematically developed with noncommutative "parameters"). That would give (using deformations) a space-time origin to the "internal" symmetries of elementary particles, on which their dynamics were based, and either question, or give a conceptually solid base to, the Standard Model, in line with Einstein's quotation: "The important thing is not to stop questioning. Curiosity has its own reason for existing."

Mathematics Subject Classification (2010). Primary 81R50; Secondary 53D55, 17B37, 53Z05, 81S10.

Keywords. Symmetries of hadrons, models, Anti de Sitter, deformation theory, deformation quantization, singletons, quantum groups at root of unity, "quantum deformations".

This text is published in Geometric Methods in Physics, XXXII Workshop 2013 in Białowieża, Trends in Mathematics, 7-37, Springer (2014).

1. Introduction: the deformation philosophy and the present proposal

1.1. Why deformations?

However seductive the idea may be, the notion of "Theory of Everything" is to me unrealistic. In physics, knowingly or not, one makes approximations in order to have as manageable a theory (or model) as possible. That happens in particular when the aim is to describe the reality known at the time, even if one suspects that a more elaborate reality is yet to be discovered. The question is how to discover that reality. We claim, on the basis of past experience, that one should not extrapolate but rather "deform."

Indeed physical theories have their domain of applicability defined, e.g., by the relevant distances, velocities, energies, etc. involved. But the passages from one domain (of distances, etc.) to another do not happen in an uncontrolled way: experimental phenomena appear that cause a paradox and contradict accepted theories, in line with the famous quote by Fermi [24]: There are two possible outcomes: if the result confirms the hypothesis, then you've made a measurement. If the result is contrary to the hypothesis, then you've made a discovery.

Eventually a new fundamental constant enters, causing the formalism to be modified: the attached structures (symmetries, observables, states, etc.) *deform* the initial structure to a new structure which in the limit, when the new parameter goes to zero, "contracts" to the previous formalism. The problem is that (at least until there is no other way out) the physics community is gregarious. Singing "It ain't necessarily so" (which I am doing in this paper) is not well received.

A first example of the "deformation" phenomenon can be traced back to Antiquity, when it was gradually realized that the earth is not flat. [Yet nowadays some still dispute the fact!] In mathematics the first instances of deformations can be traced to the nineteenth century with Riemann surface theory, though the main developments happened a century later, in particular with the seminal analytic geometry works of Kodaira and Spencer [56] (and their lesser known interpretation by Grothendieck [45], where one can see in watermark his "EGA" that started a couple of years later). These deep geometric works were in some sense "linearized" in the theory of deformations of algebras by Gerstenhaber [44].

The realization that deformations are fundamental in the development of physics happened a couple of years later in France, when it was noticed that the Galilean invariance $(SO(3) \cdot \mathbb{R}^3 \cdot \mathbb{R}^4)$ of Newtonian mechanics is deformed, in the Gerstenhaber sense [44], to the Poincaré group of special relativity $(SO(3,1) \cdot \mathbb{R}^4)$. In spite of the fact that the composition law of symbols of pseudodifferential operators, essential in the Atiyah–Singer index theorem developed at that time (to the exposition of which I took part in Paris in the Séminaire Cartan–Schwartz 1963/64), was in effect a deformation of their abelian product, it took another ten years or so to develop the tools which enabled us to make explicit, rigorous and convincing, what was in the back of the mind of many: quantum mechanics is a deformation of classical mechanics. That developed into what became known as *deformation quantization* and its manifold avatars and more generally into the realization that quantization is deformation. This stumbling block being removed, the paramount importance of deformations in theoretical physics became clear [26], giving rise to what I call "Flato's deformation philosophy".

This paper being aimed at both physicists and mathematicians and dealing with so many topics, we may look overly schematic (even trivial) in many parts for readers coming from one or the other community. More details can be found in relatively recent reviews ([21, 76, 77] by myself, and many more by others) and references quoted therein. The hope is that both communities will get the flavor of (and maybe contribute to) the framework for models proposed here. It is based on developments I have witnessed since the early 1960s and in which Flato and I took part, sometimes in a controversial way. In numerous discussions I had with scientists around these ideas, especially in the past two years, I was surprised to notice that many had often only a vague idea of a number of the topics involved, not going beyond the views given in textbooks and/or educated popularizations. I am probably one of the very few who can (and dare) deal with all the topics involved in that unconventional manner. Part of the mathematical aspect is virgin territory and in any case requires an approach (which can be called "mathematical engineering") dealing more with specific examples than with very abstract developments. On the other hand the basic physical approach is unconventional, and some of the physical issues and models questioned here have for years been taught as facts in courses and presented as such in the literature.

1.2. A brief overlook of the paper

Towards the end of the nineteenth century many believed that, in particular with Newtonian mechanics (and gravitation) and electromagnetism, physics was well understood. Yet the best was to come. In the first half of last century appeared relativity and quantum mechanics, which we now can interpret as deformations. On the fundamental side the second half of last century was dominated by the interactions between elementary particles, classified (in increasing order of strength) as gravitational, weak, electromagnetic and strong. Quantum electrodynamics (QED), developed in the 1940s, explained electromagnetic interactions with an extremely high level of accuracy (even if the theory is not yet fully mathematically rigorous). In the 1970s it was combined with weak interactions in the electroweak model, which required the Higgs boson that was (most likely) now discovered in CERN.

After an outlook of the physical and mathematical context we shall indicate how, using AdS symmetry (a deformation of Poincaré) we can explain the photon (the basis of QED) as composite of two "singletons", massless particles in a 2+1 space-time (themselves composites of two harmonic oscillators). Then an extension of the electroweak model to the presently known 3 generations of leptons could explain how, in AdS, these can also be composites of singletons, massified by 5 Higgs.

Daniel Sternheimer

It is therefore tempting to try and obtain the symmetries of hadrons, on which their dynamics has been built, by deforming further AdS. That cannot be done in the category of Lie groups but can, e.g., in that of Hopf algebras (quantum groups). It turns out that these, at root of unity (often called "restricted quantum groups") are finite dimensional vector spaces, and have finite dimensional UIRs (unitary irreducible representations), an important feature of the presently used simple unitary symmetries. There are of course many other problems to address, which cannot be ignored, but if that direction produces a model which could fit experimental data, a revolution in our understanding of physics might follow.

That could be too much to hope for and more general deformations might be needed, in particular (also at roots of unity) multiparameter (e.g., with parameters in the group algebra of $\mathbb{Z}/n\mathbb{Z}$, denoted in the following by $\mathbb{Z}_{(n)}$), or a novel theory of deformations, not yet developed mathematically, with non-commutative "deformation parameter" (especially quaternions or belonging to the group algebra of \mathbb{S}_n , the permutation group of *n* elements, e.g., n = 3).

Both are largely virgin mathematical territory, and if successful we might have to "go back to the drawing board," for the theory and for the interpretation of many raw experimental data. That is a challenge worthy of the future generations, which in any case should give nontrivial mathematics.

2. A very schematic glimpse on the context: hadrons and their symmetries

In the fifties the number of known elementary particles increased so dramatically that Fermi quipped one day [24]: Young man, if I could remember the names of these particles, I would have been a botanist.

Clearly, already then, the theoretical need was felt, to bring some order into that fast increasing [8] flurry of particles. Two (related) natural ideas appeared: To apply in particle physics "spectroscopy" methods that were successful in molecular spectroscopy, in particular group theory [83]. And to try and treat some particles as "more elementary", considering others as composite.

A seldom mentioned caveat: In molecular spectroscopy, e.g., when a crystalline structure breaks rotational symmetry, which (for trigonal and tetragonal crystals) was the subject of Flato's M.Sc. Thesis [55] under Racah (defended in 1960 and still frontier when its main part was published in 1965 as his French "second thesis"), we know the forces, and their symmetries give the spectra (energy levels). In particle physics things occurred in reverse order: one guessed symmetries from the observed spectra, interpreted experimental data on that basis and developed dynamics compatible with them.

In the beginning, in order to explain the similar behavior of proton p and neutron n under strong interactions, a quantum number (isospin) was

introduced in the 1930s, related to a SU(2) symmetry. In the 1950s new particles were discovered in cosmic rays, that behaved "strangely" (e.g., they lived much longer than expected). So a new quantum number (strangeness) was introduced, which would be conserved by strong and electromagnetic interactions, but not by weak interactions. One of these is the baryon Λ . In 1956 Shoichi Sakata [71], extending an earlier proposal by Fermi and Yang (involving only protons and neutrons) came with the "Sakata model" according to which p, n and Λ are "more elementary" and the other particles are composites of these 3 and their antiparticles. This conceptually appealing model (maybe not as "sexy" as Yoko Sakata, a top model who was not born then) had a strong impact [66], in spite of the fact that a number of the experimental predictions it gave turned out to be wrong.

In the beginning of 1961 an idea (that was in the making before) appeared: since we have 2 quantum numbers (isospin and strangeness, we would now say "two generations") conserved in strong interactions, we should try a rank 2 compact Lie group to "put into nice boxes" the many particles we had. In particular three papers were written then: An elaborate paper [7] in which, "since it is as yet too early to establish a definite symmetry of the strong interactions," all 3 groups (types $A_2, B_2 = C_2$ and G_2), and more, were systematically studied. And two [43,65], in which only the simplest (SU(3),type A_2) was proposed. The known "octets" of 8 baryons of spin $\frac{1}{2}$ and of the 8 scalar (spin 0) and vector (spin 1) mesons fitted nicely in the 8-dimensional adjoint representation. (Hence the name "the eightfold way" coined by Gell-Mann, an allusion to the "Noble Eightfold Path of Buddhism".) In 1962 Lev Okun proposed "hadrons" as a common name for strongly interacting particles, the half-integer spin (fermions) baryons, usually heavier, and the integer spin (bosons) mesons. The 9 then known baryons of spin $\frac{3}{2}$ (4 Δ +3 Σ^* +2 Ξ^*) were associated with the 10 dimensional representation: the missing one (Ω^{-}) in the "decuplet" was discovered in 1964 with roughly the properties predicted by Gell-Mann in 1962. Big success! (Even if anyone can guess that after 4,3,2 comes 1...)

But what to do with the basic (3-dimensional) representations of SU(3), which can give (by tensor product and reduction into irreducible components) all other representations? In 1964 Murray Gell-Mann, and independently George Zweig, suggested that they could be associated with 3 entities (the same number as in the Sakata model) and their antiparticles. Zweig proposed to call them "aces" but Gell-Mann, with his feeling for a popular name, called them "quarks", a nonsense word which he imagined and shortly afterward found was used by James Joyce in "Finnegans Wake:"

> Three quarks for Muster Mark! Sure he has not got much of a bark And sure any he has it's all beside the mark.

Now, how could such "confined" quarks, which would have spin $\frac{1}{2}$ (not to mention fractional charge), coexist in a hadron, something forbidden by the Pauli exclusion principle? That same year O.W. Greenberg (and Y. Nambu)

proposed to give them different "colors", now labeled blue, green, and red. Eventually, since the 1970s, that gave rise to QCD (quantum chromodynamics) in parallel with QED but with nonabelian "gauge group" SU(3) instead of the abelian group U(1) in QED. In order to keep them together "gluons" were introduced, which carried the strong force. From that time on, the development of particle physics followed essentially a ballistic trajectory, and eventually its theory became more and more phenomenology-oriented – with the caveat that many raw experimental data are interpreted within the prevalent models.

In 1964 quarks came in 3 "flavors" (up, down, and strange) but the same year a number of people, in particular Sheldon Glashow, proposed a fourth flavor (named charm) for a variety of reasons, which became gradually more convincing until in 1974 a "charmed" meson J/Ψ was discovered, completing the 2 generations of quarks, in parallel with the 2 generations of leptons $(e \text{ and } \mu)$ and their associated neutrinos. The number of supposed quark flavors grew to the current six in 1973, when Makoto Kobayashi and Toshihide Maskawa noted that an experimental observation (CP violation) could be explained if there were another pair of quarks, eventually named bottom and top by Haim Harrari, and "observed" (with much heavier mass 1 than expected for the top) at Fermilab in 1977 and 1995 (resp.). In parallel, in 1974-1977, the existence of a heavier lepton τ was experimentally found, and its neutrino discovered in 2000. Kobayashi and Maskawa shared the 2008 Nobel prize in physics with Yoichiro Nambu who, already in 1960, described the mechanism of spontaneous symmetry breaking in particle physics. They were also awarded in 1985 the first J.J. Sakurai prize for Theoretical Particle Physics established, after JJ's premature death in 1982, with the American Physical Society (by his widow Noriko Sakurai, who in 2008 became my wife [25]); Nambu had received the J.J. Sakurai prize in 1994.

So now we have 3 generations of leptons and 3 of quarks (in 6 flavors and 3 colors). SU(3) is back in, with a different meaning than originally. Eventually the electroweak model was incorporated and elaborate dynamics built on that basis of empirical origin, and everything seems to fit.

In a series of recent papers (see [13] and references therein) Alain Connes and coworkers showed that "noncommutative geometry provides a promising framework for unification of all fundamental interactions including gravity." In the last paper, assuming that "space-time is a noncommutative space formed as a product of a continuous four dimensional manifold times a finite space" he develops a quite personal attempt to predict the Standard Model (possibly with 4 colors).

But what if the Standard Model was a colossus with clay feet (as in the interpretation by prophet Daniel of Nebuchadnezzar's dream: Book of Daniel, Chapter 2, verses 31–36)? What if it were "all beside the mark"??

 $^{^1{\}rm The}$ quark masses are not measurements, but parameters used in theoretical models and compatible with raw experimental data.

3. The mathematical context: Deformation theory and quantization

In this section, for the sake of self-completeness, we shall give a very brief summary of what can be found with more details in a number of books, papers and reviews (in particular [21,77]). Since quantization is a main paradigm for our "deformation philosophy", the idea is to give readers who would not know these already, some rudiments of deformation theory, of how quantum mechanics and field theory can be realized as a deformation of their classical counterparts, and of applications to symmetries (in particular the quantum group "avatar"). Educated readers or those who do not care too much about mathematical details may (at least for the time being ...) only browse through this Section. Note however that deformation quantization (as it is now known), introduced in the "founding papers" [6], is more than a mere reformulation of usual quantum mechanics; in particular it goes beyond canonical quantization (on $\mathbb{R}^{2\ell}$) and applies to general phase spaces (symplectic or Poisson manifolds).

3.1. The Gerstenhaber theory of deformations of algebras

A concise formulation of a Gerstenhaber deformation (over the field $\mathbb{K}[[\nu]]$ of formal series in a parameter ν with coefficients in a field \mathbb{K}) of an algebra (associative, Lie, bialgebra, etc.) over \mathbb{K} is [10, 44]:

Definition 1. A deformation of an algebra A over \mathbb{K} is an algebra \tilde{A} over $\mathbb{K}[[\nu]]$ such that $\tilde{A}/\nu\tilde{A} \approx A$. Two deformations \tilde{A} and \tilde{A}' are said equivalent if they are isomorphic over $\mathbb{K}[[\nu]]$ and \tilde{A} is said trivial if it is isomorphic to the original algebra A considered by base field extension as a $\mathbb{K}[[\nu]]$ -algebra.

For associative (resp. Lie) algebras, the above definition tells us that there exists a new product * (resp. bracket $[\cdot, \cdot]$) such that the new (deformed) algebra is again associative (resp. Lie). Denoting the original composition laws by ordinary product (resp. Lie bracket $\{\cdot, \cdot\}$) this means, for $u_1, u_2 \in A$ (we can extend this to $A[[\nu]]$ by $\mathbb{K}[[\nu]]$ -linearity), that we have the formal series expansions:

$$u_1 * u_2 = u_1 u_2 + \sum_{r=1}^{\infty} \nu^r C_r(u_1, u_2)$$
(1)

$$[u_1, u_2] = \{u_1, u_2\} + \sum_{r=1}^{\infty} \nu^r B_r(u_1, u_2)$$
(2)

where the bilinear maps $(A \times A \to A)$ C_r and (skew-symmetric) B_r are what are called 2-cochains in the respective cohomologies (Hochschild and Chevalley–Eilenberg), satisfying (resp.) $(u_1 * u_2) * u_3 = u_1 * (u_2 * u_3) \in \tilde{A}$ and $\mathcal{S}[[u_1, u_2], u_3] = 0$, for $u_1, u_2, u_3 \in A$, \mathcal{S} denoting summation over cyclic permutations, the leading term (resp. C_1 or B_1) being necessarily a 2-cocycle (the coefficient of ν in the preceding conditions may be taken as a definition of that term). For a (topological) bialgebra (an associative algebra A where we have in addition a coproduct $\Delta : A \longrightarrow A \otimes A$ and the obvious compatibility relations), denoting by \otimes_{ν} the tensor product of $\mathbb{K}[[\nu]]$ -modules, we can identify $\tilde{A} \otimes_{\nu} \tilde{A}$ with $(A \otimes A)[[\nu]]$, where $\hat{\otimes}$ denotes the algebraic tensor product completed with respect to some topology (e.g., projective for Fréchet nuclear topology on A). Then we have also a deformed coproduct $\tilde{\Delta} =$ $\Delta + \sum_{r=1}^{\infty} \nu^r D_r, D_r \in \mathcal{L}(A, A \otimes A)$ satisfying $\tilde{\Delta}(u_1 * u_2) = \tilde{\Delta}(u_1) * \tilde{\Delta}(u_2)$, where \mathcal{L} denotes the space of linear maps. In this context appropriate cohomologies can be introduced. There are natural additional requirements for Hopf algebras (see, e.g., [11]).

3.2. Deformation quantization

The above abstract definition should become less abstract when applied to an algebra N(W) of (differentiable) functions on a symplectic or Poisson manifold W, in particular those over phase space $\mathbb{R}^{2\ell}$ (with coordinates $p, q \in$ \mathbb{R}^{ℓ}) endowed with the Poisson bracket P of two functions u_1 and u_2 , defined on a Poisson manifold W as $P(u_1, u_2) = i(\Lambda)(du_1 \wedge du_2)$ (where *i* denotes the interior product, here of the 2-form $du_1 \wedge du_2$ with the 2-tensor Λ defining the Poisson structure on W, which in the case of a symplectic manifold is everywhere nonzero with for inverse a closed nondegenerate 2-form ω). For $W = \mathbb{R}^{2\ell}$, P can be written by setting r = 1 in the formula for the r^{th} power $(r \geq 1)$ of the bidifferential operator P (we sum over repeated indices):

$$P^{r}(u_{1}, u_{2}) = \Lambda^{i_{1}j_{1}} \dots \Lambda^{i_{r}j_{r}}(\partial_{i_{1}\dots i_{r}}u_{1})(\partial_{j_{1}\dots j_{r}}u_{2})$$
(3)

with $i_k, j_k = 1, \ldots, 2\ell$, $k = 1, \ldots, r$ and $(\Lambda^{i_k j_k}) = \binom{0-I}{I0}$. We can write deformations of the usual product of functions (deformations driven by the Poisson bracket) and of the Poisson bracket as what are now called the Moyal ("star") product and bracket, resp.

$$u_1 *_M u_2 = \exp(\nu P)(u_1, u_2) = u_1 u_2 + \sum_{r=1}^{\infty} \frac{\nu^r}{r!} P^r(u_1, u_2).$$
(4)

$$M(u_1, u_2) = \nu^{-1} \sinh(\nu P)(u_1, u_2) = P(u_1, u_2) + \sum_{r=1}^{\infty} \frac{\nu^{2r}}{(2r+1)!} P^{2r+1}(u_1, u_2).$$
(5)

These correspond (resp.) to the product and commutator of operators in the "canonical" quantization on $\mathbb{R}^{2\ell}$ of a function H(q, p) with inverse Fourier transform $\tilde{H}(\xi, \eta)$, given by (that formula was found by Hermann Weyl [80] as early as 1927 when the weight is $\varpi = 1$):

$$H \mapsto \hat{H} = \Omega_{\varpi}(H) = \int_{\mathbb{R}^{2\ell}} \tilde{H}(\xi, \eta) \exp(i(\hat{p}.\xi + \hat{q}.\eta)/\hbar) \varpi(\xi, \eta) d^{\ell} \xi d^{\ell} \eta \qquad (6)$$

which maps the classical function H into an operator on $L^2(\mathbb{R}^{2\ell})$, the "kernel" $\exp(i(\hat{p}.\xi + \hat{q}.\eta)/\hbar)$ being the corresponding unitary operator in the (projectively unique) representation of the Heisenberg group with generators \hat{p}_{α} and \hat{q}_{β} ($\alpha, \beta = 1, \ldots \ell$) satisfying the canonical commutation relations $[\hat{p}_{\alpha}, \hat{q}_{\beta}] = i\hbar \delta_{\alpha,\beta} I$. An inverse formula to that of the Weyl quantization formula was found in 1932 by Eugene Wigner [81] and maps an operator into what mathematicians call its symbol by a kind of trace formula: Ω_1 defines an isomorphism of Hilbert spaces between $L^2(\mathbb{R}^{2\ell})$ and Hilbert–Schmidt operators on $L^2(\mathbb{R}^{\ell})$ with inverse given by

$$u = (2\pi\hbar)^{-\ell} \operatorname{Tr}[\Omega_1(u) \exp((\xi \cdot \hat{p} + \eta \cdot \hat{q})/i\hbar)].$$
(7)

It is important to remember that "star products" exist as deformations (the skew-symmetric part of the leading term being the Poisson bracket P) of the ordinary product of functions in N(W) for any W [21], including when there are no Weyl or Wigner maps and no obvious Hilbert space treatment of quantization. They can also be defined for algebraic varieties, "manifolds with singularities", and (with some care) infinite-dimensional manifolds.

We refer, e.g., to [6, 21, 77] and especially references therein for more developments on deformation quantization and its many avatars. These include the notion of covariance of star products and the "star representations" (without operators) it permits. They include also quantum groups, which appeared in Leningrad around 1980 for entirely different reasons [23] but, especially after the seminal works of Drinfel'd [22] (who coined the name) and of Jimbo [53] (for quantized enveloping algebras) in the early 1980s, can be viewed as deformations of (topological) Hopf algebras (see, e.g., [10, 11, 75]).

In this connection it is worth remembering a prophetic general statement by Dirac [19], which applies to many situations in physics: Two points of view may be mathematically equivalent, and you may think

Two points of view may be mathematically equivalent, and you may think for that reason if you understand one of them you need not bother about the other and can neglect it. But it may be that one point of view may suggest a future development which another point does not suggest, and although in their present state the two points of view are equivalent they may lead to different possibilities for the future. Therefore, I think that we cannot afford to neglect any possible point of view for looking at Quantum Mechanics and in particular its relation to Classical Mechanics.

What Dirac had then in mind is certainly the quantization of constrained systems which he developed shortly afterward and by now can be viewed as a special case of deformation quantization. But the principle applies to many contexts and is even a most fruitful strategy to extend a framework beyond its initial context. A wonderful example is given by noncommutative geometry [14], now a frontier domain of mathematics with a wide variety of developments ranging from number theory to various areas of physics.

In order to show that important and concrete problems in physics can be treated in an *autonomous* manner using deformation quantization, without the need to introduce a Hilbert space (which for most physicists is still considered as a requirement of quantum theories) we treated in [6] a number of important problems, first and foremost the harmonic oscillator (the basic paradigm in many approaches), but also angular momentum, the hydrogen atom, and in general the definition of spectrum inside deformation quantization, without needing a Hilbert space. Not so many further applications have been developed since but the approach should eventually prove fruitful (even necessary) in many domains in which quantum phenomena play a role (including quantum computing). True, in many concrete examples we need (at least implicitely) "auxiliary conditions" to limit the possibly excessive freedom coming, in particular for spectra, from the absence of that Procrustean bed, the Hilbert space. But there also, too much freedom is better than not enough.

3.3. Further important deformations (and contractions)

3.3.1. An instance of multiparameter deformation quantization. A natural question is whether "the buck stops there" i.e., whether, like for Gerstenhaber deformations of simple Lie groups or algebras, the structure obtained is rigid, or whether some further deformations are possible. An answer to that question, looking for further deformations of N(W) with another parameter β (in addition to $\nu = i\frac{\hbar}{2}$), was given in [5] and applied to statistical mechanics and the so-called KMS states (with parameter $\beta = 1/kT$, T denoting the absolute temperature). It turns out that there is some intertwining which is not an equivalence of deformations: As a ν -deformation, the two-parameter "star product" is driven by a "conformal Poisson bracket" with conformal factor of the form $\exp(-\frac{1}{2}\beta H)$ for some Hamiltonian H.

3.3.2. Brief survey of a few aspects of quantum groups. The literature on quantum groups (and Hopf algebras) is so vast, diversified (and growing) that we shall refer the interested reader to his choice among the textbooks and papers dealing with the many aspects of that notion, often quite algebraic. A two-pages primer can be found in [61].

Roughly speaking quantum groups can often be considered [22] as deformations (in the sense of Definition 1) of an algebra of functions on a Poisson-Lie group (a Lie group G equipped with a Poisson bracket compatible with the group multiplication, e.g., a semi-simple Lie group), or the "dual aspect" [22,53] of a deformation of (some closure of) its enveloping algebra $\mathcal{U}(\mathfrak{g})$ equipped with its natural Hopf algebra structure, which is how the whole thing started in Leningrad around 1980. The first example was $\mathcal{U}_t(\mathfrak{sl}(2))$, an algebra with generators e, f, h as for $\mathfrak{sl}(2)$ but with "deformed" commutation relations that can be written somewhat formally (the deformation parameter t = 0 for $\mathfrak{sl}(2)$):

$$[h, e] = 2e, \ [h, f] = -2f, \ [e, f] = \sinh(th) / \sinh t \tag{8}$$

or more traditionally, as an algebra with generators E, F, K, K^{-1} (one often writes $K^{\pm 1} = q^{\pm H}$) and relations

$$KK^{-1} = K^{-1}K = 1, \ KEK^{-1} = q^2E, \ KFK^{-1} = q^{-2}F,$$
 (9)

$$EF - FE = \frac{K - K^{-1}}{q - q^{-1}},$$
(10)

For simplicity we shall not write here the expressions for the coproduct, counit and antipode, needed to show the Hopf algebra structure. For higher rank simple Lie algebras one has in addition trilinear relations (the deformed Serre relations), which complicate matters. All these can be found in the literature.

Note that the algebraic dual of a Hopf algebra is a also Hopf algebra only when these are finite-dimensional vector spaces, which is quite restrictive a requirement. In particular the Hopf algebras considered in quantum groups (except at root of unity), e.g., those of differentiable functions over a Poisson-Lie group, are (finitely generated) infinite dimensional vector spaces; but one can [10, 11] define on these spaces natural topologies (e.g., Fréchet nuclear) which in particular express the duality between them and "quantized enveloping algebras". Remember that for any connected Lie group Gwith Lie algebra \mathfrak{g} the elements of the enveloping algebra $\mathcal{U}(\mathfrak{g})$ can be considered as differential operators over G, i.e., as distributions with support at any point in G (e.g., the identity $e \in G$), which lie in the topological dual of the space of differentiable functions with compact support. That exhibits a "hidden group structure" [10] in Drinfeld's quantum groups, which [22] are not groups (and not always quantum ...)

3.3.3. About quantum groups at root of 1. As was noticed around 1990, in particular by Lusztig [59, 60] (see also, e.g., [17, 70]) the situation changes drastically when the deformation parameter is a root of unity. Then the Hopf algebras $\mathcal{U}_q(\mathfrak{g})$ become finite dimensional.

The case of $\mathcal{U}_q(\mathfrak{sl}(2))$ is well understood. For q a 2*p*-th root of unity, the finite-dimensionality of the algebra comes from the fact that, in addition to (9) and (10), one has the relations

$$K^{2p} = 1, \ E^p = 0, \ F^p = 0.$$
 (11)

In particular (see, e.g., [85]) all finite-dimensional indecomposable representations have been determined, as well as the indecomposable decomposition of tensor products [57]. The higher rank case is still largely virgin mathematical territory. It seems that one either needs a new approach, or to restrict oneself to particular cases, or both.

3.3.4. Multiparameter quantum groups. More generally it is natural to try and look from the start at deformations with several scalar parameters. That question seems to have been tackled for the first time, in the context of quantum groups, by Manin et al. [18,62], who called "nonstandard" these multiparameter deformations, and Reshetikhin [69], and later by Frønsdal [39,41]. But the notion does not seem to have drawn the attention it deserves, certainly not much in comparison with the many works on more traditional aspects of quantum groups. And more sophisticated questions such as what happens (outside the generic case), e.g., "at roots of unity" (whether the same root for all parameters or not) do not seem to have ever been considered for multiparameter quantum groups.

3.3.5. Nonscalar deformation "parameter". Other deformations, more general than those of Gerstenhaber type, were considered by Pinczon [68] and his student Nadaud [63, 64], in which the "parameter" acts on the algebra

(on the left, on the right, or both) instead of being a scalar. For instance one can have [68], for $\tilde{a} = \sum_n a_n \lambda^n$, $a_n \in A$, a left multiplication by λ of the form $\lambda \cdot \tilde{a} = \sum_n \sigma(a_n)\lambda^{n+1}$ where σ is an endomorphism of A. A similar deformation theory can be done in this case, with appropriate cohomologies, which gives new and interesting results.

In particular [68], while the Weyl algebra W_1 (generated by the Heisenberg Lie algebra \mathfrak{h}_1) is known to be Gerstenhaber-rigid, it can be nontrivially deformed in such a supersymmetric deformation theory to the supersymmetry enveloping algebra $\mathcal{U}(\mathfrak{osp}(1|2))$. Shortly thereafter [64], on the polynomial algebra $\mathbb{C}[x, y]$ in 2 variables, Moyal-like products of a new type were discovered; a more general situation was studied, where the relevant Hochschild cohomology is still valued in the algebra but with "twists" on both sides for the action of the deformation parameter on the algebra.

3.3.6. Contractions. Curiously, it is the (less precisely defined) inverse notion of *contraction* of symmetries that was first introduced in mathematical physics [51, 72]. Contractions, "limits of Lie algebras" as they were called in the first examples, can be viewed as an inverse of deformations – but not necessarily of Gerstenhaber-type deformations. We shall not expand on that "inverse" notion (see [78] for a more elaborate study) but give its flavor since it makes it easier to grasp the deformations of symmetries which are important in our presentation. A (finite dimensional) Lie algebra \mathfrak{g} can be described in a given basis L_i (i = 1, ..., n) by its structure constants $C_{i,j}^k$. The equations governing the skew-symmetry of the Lie bracket and the Jacobi identity ensure that the set of all structure constants lies on an algebraic variety in that n^3 dimensional space [58]. A contraction is obtained, e.g., when one makes a simple basis change of the form $L'_i = \varepsilon L_i$ on some of the basis elements, and lets $\varepsilon \to 0$. Take for example n = 3 and restrict to the 3-dimensional subspace of the algebraic variety of 3-dimensional Lie algebras with commutation relations $[L_1, L_2] = c_3 L_3$ and cyclic permutations. The semi-simple algebras $\mathfrak{so}(3)$ and $\mathfrak{so}(2,1)$ are obtained in the open set $c_1c_2c_3 \neq 0$. A contraction gives the Euclidean algebras, where one c_i is 0. The "coordinate axes" (two of the c_i 's are 0) give the Heisenberg algebra \mathfrak{h}_1 and the origin is the Abelian Lie algebra. That is of course a partial picture (e.g., solvable algebras are missing) but it is characteristic.

The above mentioned passage from the Poincaré Lie algebra to the Galilean is a higher dimensional version of such contractions of Lie algebras (multiply the "Lorentz boosts" generators M_{0j} by ε). A similar "trick" on the AdS₄ Lie algebra $\mathfrak{so}(3,2)$ gives the Poincaré Lie algebra. A traditional basis for the Poincaré Lie algebra is $M_{\mu\nu}$ for the Lorentz Lie algebra $\mathfrak{so}(3,1)$ and P_{μ} for the space-time translations (momentum generators), with $\mu, \nu \in \{0, 1, 2, 3\}$. The commutation relations for the conformal Lie algebra $\mathfrak{so}(4,2)$ in the basis $M_{\mu\nu}$ with $\mu, \nu \in \{0, 1, 2, 3, 5, 6\}$ can be written (see, e.g., [2]) $[M_{\mu\nu}, M_{\mu'\nu'}] = \eta_{\nu\mu'}M_{\mu\nu'} + \eta_{\mu\nu'}M_{\nu\mu'} - \eta_{\mu\mu'}M_{\nu\nu'} - \eta_{\nu\nu'}M_{\mu\mu'}$ with diagonal metric tensor $\eta_{\mu\mu}$ (equal, e.g., to +1 for $\mu = 0, 5$ and -1 for $\mu = 1, 2, 3, 6$). One can identify a Poincaré subalgebra by setting, e.g., $P_{\mu} = M_{\mu5} + M_{\mu6}$. If

one omits $\mu = 6$ one obtains the AdS₄ Lie algebra $\mathfrak{so}(3, 2)$ where the role of the Poincaré space translations is taken over by M_{j5} $(j \in \{1, 2, 3\})$ and that of the time translations by M_{05} . How to realize a contraction from AdS₄ to Poincaré for the massless representations used in Section 5 is described in [2]. Since it may be easier for physicists to grasp the notion of contraction, we mentioned it here and explicited the examples we use in familiar notations (possibly old fashioned, but with bases on \mathbb{R}).

4. On the connection between internal and external symmetries

We shall not extend our desire to be as self contained as possible to describing in full detail the Poincaré group and its UIRs (unitary irreducible representations), known since on the instigation of Dirac (his "famous brother-in-law" as he liked to call him) Wigner [82] published the paper that started the study of UIRs of non compact Lie groups. Those associated with free particles are usually denoted by D(m, s), where $m \ge 0$ is the mass and s the spin of the particle (for m > 0) or its helicity (for m = 0), associated with the "squared mass Casimir operator" (in the center of the Poincaré enveloping algebra) $P_{\mu}P^{\mu}$ and with the inducing representation of the "little group" (SO(3) and $SO(2) \cdot \mathbb{R}^2$, resp.). [The letter D, coming from the German "Darstellung", is often used to denote representations.]

In the early 1960s a natural question appeared: Is there any connection between the "internal symmetry" used in the classification of interacting elementary particles (tentatively SU(3) at the time), and the Poincaré symmetry whose UIRs are associated with free particles? The question is not innocent since 3 octets of different spins were associated with the same representation of SU(3) (the 8-dimensional adjoint representation). And the various families within an octet (the same applies to the decuplet) exhibit a mass spectrum. If there is a connection, one ought to describe a mechanism permitting all that. Of course an important issue is how to mathematically formulate the question.

4.1. No-go theorems, objections, counter-examples and generalizations

4.1.1. A Lie algebra no-go and counterexamples. In the "particle spectroscopy" spirit of the time, it was natural to look for a Lie algebra containing both symmetries (internal and external). In 1965, a year after quarks and color were proposed, appeared a "no-go theorem" as physicists like to call such results, due to L. O'Raifeartaigh [67]. It boiled down to the fact that, since the momentum generators P_{μ} are nilpotent in the Poincaré Lie algebra, they are nilpotent in any simple Lie algebra containing it, which forbids a discrete mass spectrum. Hence in order to have a mass spectrum the connection must be a direct sum. Almost everybody was happy, except that two trouble makers in France said: "It ain't necessarily so". Thanks to

Isidor Rabi (then president of the APS, who remembered Moshe Flato as a most brilliant student who often asked difficult questions during the course he gave for a quarter at the Hebrew University, invited by Giulio Racah) our objection was published shortly afterward [33] in the provocative form desired by Moshe. It was followed by counterexamples [34, 36]. The problem with the "proof" in [67] is that it implicitly assumed the existence of a common invariant domain of differentiable vectors for the whole Lie algebra, something which Wigner was careful to state as an assumption in [82] and was proved later for Banach Lie group representations by (in Wigner's own words) "a Swedish gentleman" [42]. Eventually the statement of [67] could be proved within the context of UIRs of finite dimensional Lie groups [54] and was further refined by several authors (especially L. O'Raifeartaigh). However we showed in [36] that a mass spectrum is possible when assuming only the Poincaré part to be integrable to a UIR, and there is no a priori reason why the additional observables should close to a finite dimensional group UIR. We gave also counterexamples [37] with a natural infinite-dimensional group and even showed [35] that it is possible to obtain any desired mass spectrum in the framework of finitely-generated infinite-dimensional associative algebras and unitary groups.

Like with many physical "theorems", a main issue is to decide what assumptions and what heuristic developments can be considered as "natural". While some flexibility can be accepted in proving positive results, no-go theorems should be taken with many grains of salt.

4.1.2. Subsequent developments and relative importance of the question. As we indicated in [36], such considerations apply also to a more sophisticated no-go theorem [12], formulated in the context of symmetries of the *S*-matrix, which could be applied also to infinite-dimensional groups. This very nice piece of work is still considered by most physicists (especially those who learned it at university) as definitely proving the direct sum connection (under hypotheses easily forgotten, some of which are even hidden in the apparently natural notations).

In retrospect one can say that a main impact of the latter result came through an attempt to get a supersymmetric extension [46], which showed one might get around the no-go in the supersymmetry context. That gave a big push to the latter. Incidentally it is generally considered that the "super-Poincaré group" of Wess and Zumino [79] is practically the first instance of supersymmetry. That is not quite correct. In particular already in 1967 (in CRAS) we introduced what we called "a Poincaré-like group", semi-direct product of the Lorentz group and \mathbb{R}^8 consisting of both vector and spinor translations, but (for fear of Pauli) we did not dare introduce anticommutators for spinorial translations together with commutators for space-time translations, so we remained in the Lie algebra framework. However one can find in [31] a physical application of that group in which the spinorial translations are multiplied by an operator F anticommuting with itself. That was in effect the first realization of the super-Poincaré group. Both Wess and Zumino told me some years ago that they were unaware of the fact and it seems that (except for Frønsdal) not many noticed it either.

5. Singleton physics

5.1. Singletons as "square roots" of massless particles

The contraction of AdS to Poincaré (in both structure and representations) is (cf. [2]) one of the justifications for calling "massless" some minimal weight UIRs of $Sp(\mathbb{R}^4)$ (the double covering of SO(3,2)). These are denoted by $D(E_0, s)$, the parameters being the lowest values of the energy and spin (resp.) for the compact subgroup $SO(2) \times SO(3)$. These irreducible representations are unitary provided $E_0 \geq s + 1$ for $s \geq 1$ and $E_0 \geq s + \frac{1}{2}$ for s = 0and $s = \frac{1}{2}$. The massless representations of SO(3,2) are thus defined (for $s \geq \frac{1}{2}$) as D(s + 1, s) and (for helicity zero) $D(1, 0) \oplus D(2, 0)$. At the limit of unitarity (when going down in the values of E_0 for fixed s) the Harish Chandra module $D(E_0, s)$ becomes indecomposable and the physical UIR appears as a quotient, a hall-mark of gauge theories. For $s \geq 1$ we get in the limit an indecomposable representation $D(s+1, s) \rightsquigarrow D(s+2, s-1)$, where \leadsto ("leaking into") is a shorthand notation [28] for what mathematicians would write as a short exact sequence of (infinite-dimensional) modules.

A complete classification of the UIRs of the (covering of) SO(p, 2) can be found, e.g., in [1]. For p = 3 the classification had been completed when Dirac [20] introduced the most degenerate "singleton" representations. The latter are irreducible and massless on a subgroup, the Poincaré subgroup of a 2 + 1 dimensional space-time, of which AdS is the conformal group. That is why (on the pattern of Dirac's "bra" and "ket") we call these representations $Di = D(1, \frac{1}{2})$ and $Rac = D(\frac{1}{2}, 0)$ for (resp.) the spinorial and scalar representations. The singleton representations have a fundamental property:

$$(Di \oplus Rac) \otimes (Di \oplus Rac) = (D(1,0) \oplus D(2,0)) \oplus 2 \bigoplus_{s=\frac{1}{2}}^{\infty} D(s+1,s).$$
(12)

The representations appearing in the decomposition are what we call massless representations of the AdS group, for a variety of good reasons [2]. For s = 0a split occurs because time is compact in AdS and one does not distinguish between positive and negative helicity, which is also the reason for the factor 2 in front of the sum. An extension to the conformal group SO(4, 2), which is operatorially unique for massless representations of the Poincaré group and (once a helicity sign is chosen) for those of SO(3, 2), or a contraction of the latter to the Poincaré group, restores the distinction between both helicity signs and provide other good reasons for calling massless representations of AdS those in the right hand side of (12).

Thus, in contradistinction with flat space, in AdS_4 , massless states are "composed" of two singletons. The flat space limit of a singleton is a vacuum

and, even in AdS_4 , the singletons are very poor in states: their (E, j) diagram has a single trajectory (hence the name given to them by Dirac), and is not a lattice like, e.g., for massless particles in AdS. In normal units a singleton with angular momentum j has energy $E = (j + \frac{1}{2})\rho$, where ρ is the curvature of the AdS₄ universe. This means that only a laboratory of cosmic dimensions can detect a j large enough for E to be measurable: one can say that the singletons are "naturally confined".

Like the $\operatorname{AdS}_n/\operatorname{CFT}_{n-1}$ correspondence, the symmetry part of which states essentially that SO(n-1,2) is the conformal group for n-1-dimensional space-time, singletons exist in any space-time of dimension $n \geq 3$ [3], n = 4being somewhat special. For n = 3 the analogue of (12) writes $(HO) \otimes (HO) =$ $Di \oplus Rac$ where (HO) denotes the harmonic oscillator representation of the metaplectic group (double covering of $SL(2, \mathbb{R})$, itself a double covering of SO(2, 1)) which is the sum of the discrete series representation $D(\frac{3}{4})$ and of the complementary series representation $D(\frac{1}{4})$. One thus has a kind of "dimensional reduction" by which ultimately massless particles can be considered as arising from the interaction of harmonic oscillators. I leave it to the reader to derive consequences from that fact, and maybe study connections with the challenging suggestions of Gerard 't Hooft (see, e.g., [49, 50] and references therein) dealing with "quantum determinism" and based in particular on cellular automata.

Remark 5.1.1. Phase space realization. Quadratic polynomials in 2ℓ real variables p_{α} and q_{β} ($\alpha, \beta \in \{1, \ldots, \ell\}$), satisfying the Heisenberg canonical commutation relations (CCR) $[p_{\alpha}, q_{\beta}] = \delta_{\alpha\beta}I$, generate a realization of the symplectic Lie algebra $\mathfrak{sp}(\mathbb{R}^{2\ell})$. Together with the linear polynomials (under anticommutators) they close to an irreducible realization of $\mathfrak{osp}(1|2\ell)$, the corresponding superalgebra ($\mathbb{Z}_{(2)}$ -graded). $[Di \oplus Rac$ and (HO) are special cases of the phenomenon for $\ell = 2, 1$ (resp.).] That fact allowed us in [6] to write a power series expansion in t of what we call the "star exponential" $\operatorname{Exp} * (tH/i\hbar)$ (corresponding in Weyl quantization to the unitary evolution operator) of the harmonic oscillator Hamiltonian $H = \frac{1}{2}(p^2 + q^2)$, while a theorem of Harish Chandra states that the character of a UIR (which in our formalism is the integral over phase space of the star-exponential) always has a singularity at the origin: the singularities for the two components in (HO) cancel at the origin, a true miracle which puzzles many specialists of Lie group representation theory!

5.2. AdS_4 dynamics

Until now we were concerned mainly with what can be called the "kinematical aspect" of the question, i.e., symmetries. However at some point one cannot avoid looking at the dynamics involved. In particular covariant field equations and Lagrangians will have to be studied. And indeed many papers were written, especially in the 1980s and 1990s by Flato, Frønsdal and coworkers, developing various aspects of singleton physics. These include BRST symmetry, conformal aspects and related indecomposable representations (in particular of the Gupta–Bleuler type), etc. Our purpose here is to build on these and on the "deformation philosophy" and not to give an extensive account of all these works, references to many of which can be found in Flato's last paper [29]. In the next two subsections we shall give a brief account of the two papers that are the most important from the point of view developed here, composite QED [28] and what can be called an electroweak model extended to 3 generations of leptons [40].

5.2.1. The Flato–Frønsdal "singletonic QED". Dynamics require in particular the consideration of field equations, initially at the first quantized level, in particular the analogue of the Klein–Gordon equation in AdS_4 for the *Rac.* There, as can be expected of massless (in 1+2 space) representations, gauges appear, and the physical states of the singletons are determined by the value of their fields on the cone at infinity of AdS_4 (see below; we have here a phenomenon of holography [48], in this case an AdS_4/CFT_3 correspondence).

We thus have to deal with indecomposable representations, triple extensions of UIR, as in the Gupta–Bleuler (GB) theory, and their tensor products. [It is also desirable to take into account conformal covariance at these GBtriplets level, which in addition permits distinguishing between positive and negative helicities (in AdS₄, the time variable being compact, the massless representations of SO(2,3) of helicity s > 0 contract (resp. extend in a unique way) to massless representations of helicity $\pm s$ of the Poincaré (resp. conformal) group.] The situation gets therefore much more involved, quite different from the flat space limit, which makes the theory even more interesting.

In order to test the procedure it is necessary to make sure that it is compatible with conventional Quantum Electrodynamics (QED), the best understood quantum field theory, at least at the physical level of rigor.

One is therefore led to see whether QED is compatible with a massless photon composed of two scalar singletons. For reasons explained, e.g., in [29] and references quoted therein, we consider for the *Rac*, the dipole equation $(\Box - \frac{5}{4}\rho)^2\phi = 0$ with the boundary conditions $r^{1/2}\phi < \infty$ as $r \to \infty$, which carries the indecomposable representation $D(\frac{1}{2},0) \rightsquigarrow D(\frac{5}{2},0)$. A remarkable fact is that this theory is a *topological field theory*; that is [27], the physical solutions manifest themselves only by their boundary values at $r \to \infty$: $\lim r^{1/2}\phi$ defines a field on the 3-dimensional boundary at infinity. There, on the boundary, gauge invariant interactions are possible and make a 3-dimensional conformal field theory (CFT).

However, if massless fields (in four dimensions) are singleton composites, then singletons must come to life as 4-dimensional objects, and this requires the introduction of unconventional statistics (neither Bose–Einstein nor Fermi–Dirac). The requirement that the bilinears have the properties of ordinary (massless) bosons or fermions tells us that the statistics of singletons must be of another sort. The basic idea is [28] that we can decompose the *Rac* field operator as $\phi(x) = \sum_{-\infty}^{\infty} \phi^j(x)a_j$ in terms of positive energy creation operators $a^{*j} = a_{-j}$ and annihilation operators a_j (with j > 0) without so far making any assumptions about their commutation relations. The choice of commutation relations comes later, when requiring that photons, considered

as 2-*Rac* fields, be Bose–Einstein quanta, i.e., their creation and annihilation operators satisfy the usual canonical commutation relations (CCR). The singletons are then subject to unconventional statistics (which is perfectly admissible since they are naturally confined), the total algebra being an interesting infinite-dimensional Lie algebra of a new type, a kind of "square root" of the CCR. An appropriate Fock space can then be built. Based on these principles, a (conformally covariant) composite QED theory was constructed [28], with all the good features of the usual theory—however years after QED was developed by Schwinger, Feynman, Tomonaga and Dyson.

Remark 5.2.1.1. Classical Electrodynamics as a covariant nonlinear PDE approach to coupled Maxwell–Dirac equations. Only relatively recently was classical electrodynamics (on 4 dimensional flat space-time) rigorously understood. By this we mean the proof of asymptotic completeness and global existence for the coupled Maxwell–Dirac equations, and a study of the infrared problem. That was done [32] with the third aspect of our trilogy (complementing deformation quantization and singleton physics), based on a theory of nonlinear group representations, plus a lot of hard analysis using spaces of initial data suggested by the linear group representations. The deformation quantization of that classical electrodynamics (e.g., on an infinite dimensional phase space of initial conditions) remains to be done.

5.2.2. Composite leptons, Frønsdal's extended electroweak model. A natural step, after QED, is to introduce compositeness in electroweak theory. Along the lines described above, that would require finding a kind of "square root of an infinite-dimensional superalgebra," with both CAR (canonical anticommutation relations) and CCR included: The creation and annihilation operators for the naturally confined Di or Rac need not satisfy CAR or CCR; they can be subject to unusual statistics, provided that the two-singleton states satisfy Fermi–Dirac or Bose–Einstein statistics depending on their nature. We would then have a (possibly \mathbb{Z} -)graded algebra where only the two-singleton states creation and annihilation operators satisfy CCR or CAR. That has yet to be done. Some steps in that direction have been initiated but the mathematical problems are formidable, even more so since now the three generations of leptons have to be considered.

But here a more pragmatic approach can be envisaged [40], triggered by experimental data showing oscillations between various generations of neutrinos. The latter can thus no more be considered as massless. This is not as surprising as it seems from the AdS point of view, because one of the attributes of masslessness is the presence of gauges. These are group theoretically associated with the limit of unitarity in the representations diagram, and the neutrino is above that limit in AdS: the Di is at the limit. Thus, all nine leptons can be treated on an equal footing.

It is then natural [40] to arrange them in a square table (L_{β}^{A}) , the rows being the 3 generations of leptons, each of which carry the Glashow representation of the 'weak group' $S_{W} = SU(2) \otimes U(1)$ and to consider the 9 leptons L_{β}^{A} (ν_{e}, e_{L}, e_{R} and similarly for the two other generations μ and τ) as composites, $L_{\beta}^{A} = R^{A}D_{\beta}$ $(A = N, L, R; \beta = \epsilon, \mu, \tau)$. We assume that the R^{A} s are Racs and carry the Glashow representation of S_{W} , while the D_{β} s are Dis, insensitive to S_{W} but transforming as a Glashow triplet under a 'flavor group' S_{F} isomorphic to S_{W} . To be more economical we also assume that the two U(1)s are identified, the same hypercharge group acting on both Dis and Racs. As explained in [40], the leptons are initially massless (as Di-Rac composites) and massified by (in effect, five) Higgs fields $K_{AB}^{\alpha\beta}$ that (like in the electroweak model) have a Yukawa coupling to the leptons. The model predicts, in parallel to the W^{\pm} and Z bosons, two new bosons C^{\pm} and C^{3} (hard to detect due to the large mass differences between the 3 generations of leptons) and explains the neutrino masses. It is qualitatively promising but the presence of too many free parameters limits its quantitative predictive power.

One could be tempted to add to the picture a deformation induced by the strong force and 18 quarks, which (with the 9 leptons) could be written in a cube and also considered composite (of maybe three constituents when the strong force is introduced). That might make this "composite Standard Model" more predictive. But introducing the hadrons brings in a significant quantitative change that should require a qualitative change, e.g., some further deformation of the AdS symmetry.

6. Hadrons and quantized Anti de Sitter

6.1. A beginning of a new picture

Instead of a "totalitarian" approach aiming towards a "theory of everything," at least inasmuch as elementary particles are concerned, we shall adopt an approach which is both more pragmatic and based on fundamental principles. Since symmetries were the starting point from which what is now the Standard Model emerged, and since free particles are governed by UIRs of the Poincaré group (the symmetry of special relativity), we shall start from the latter and proceed using our "deformation philosophy" as a guideline.

6.1.1. Photons, Leptons and Hadrons. As we have seen, assuming that in the "microworld", i.e., at some scale (to be made more precise eventually) the universe is endowed with a small negative curvature, the Poincaré group is deformed to AdS and the massless photon states can be dynamically (in a manner compatible with QED) considered as a 2-Rac state [28]. Then, extending the electroweak theory to the empirically discovered 3 generations, the 9 leptons can be considered, using the AdS deformation of the Poincaré group, as initially massless Di-Rac states, massified by 5 Higgs bosons [40].

The "tough cookie" is then how to explain hadrons and strong interactions. The deformation philosophy suggests to try and deform AdS, which is not possible as a group but can be done as a Hopf algebra, to a "quantum group", qAdS. In the "generic case" the obtained representation theory will not be very different from the AdS case. That is essentially due to the "Drinfeld twist" which intertwines between AdS and qAdS, even if that is not an equivalence of deformations (it is a kind of "outer automorphism"). But at root of unity the situation becomes drastically different: the Hopf algebra becomes finite-dimensional (which is not the case of the generic qAdS) and there are only a finite number of irreducible representations.

6.1.2. Remarks on the mathematical context. The fact that only a few irreducible representations may be relevant is both an encouraging feature and a restrictive one, even more so since for quantum groups at root of 1, the theory of tensor products of such representations (needed in order to consider these as describing interacting particles), which is "nice" in the generic case, is not straightforward. In particular the tensor products are usually indecomposable, extensions of direct sums of irreducibles defined by some cocycles (which could however be related to the "width" of the observed resonances). The phenomenon appears already in rank one (quantized $\mathfrak{sl}(2)$) at root of unity) [57], where the category of representations is not braided and the tensor products $R \otimes S$ and $S \otimes R$ of two representations can sometimes be different. The general theory for higher ranks seems hopeless. But for physical applications we do not need a general theory, possibly on the contrary: If only a few representations behave "nicely", and it turns out that Nature selects these, so much the better. Even for the Poincaré group only half of the UIRs are physically relevant, those of positive mass, and of zero mass and discrete helicity.] The case of rank 2 (in particular qAdS) seems more within reach, though the mathematics is highly non trivial. Some works are in progress in that direction, in particular by Jun Murakami who recently [16] studied quantum 6*j*-symbols for $SL(2,\mathbb{C})$. But a lot remains to be done in order to clarify the mathematical background.

6.2. Quantized AdS (in particular at some root of 1) and generalizations

6.2.1. Some ideas, problems and results around qAdS representations. Over 20 years ago appeared a concise and interesting paper [30], written, without sacrificing rigor, using a language (e.g., Bose creation and annihilation operators, supersymmetry, Fock space) that can appeal to physicists (but maybe less to mathematicians ...). In that paper, on the basis of a short panorama of singleton and massless representations of $\mathfrak{so}(3, 2)$, their supersymmetric extensions and (for the massless) imbedding in the conformal Lie algebra $\mathfrak{u}(2, 2)$, the authors dealt with q-deformations of that picture, especially q-singletons, q-massless representations, and the imbedding therein of q-deformations of $\mathfrak{sl}(2)$. They studied both the case of generic q and the case, to write explicitly, in the case of $\mathcal{U}_q(\mathfrak{so}(3, 2))$, defining relations similar to those of (9) and (10) (including now the "q-Serre relations") and to express in a more physical language the fact (discovered a few years before by Lusztig [60]) that one gets then finite-dimensional unitary representations.

Incidentally the "unitarization" of irreducible representations can be important for possible physical applications. For quantum groups at root of unity that has been studied, in particular for AdS, in [30, 73] and for many series of simple noncompact Lie algebras in [74], where in addition it is shown that the unitary highest weight modules of the classical case are recovered in the limit $q \to 1$.

Guided by the "deformation philosophy" we are thus led to look at what happens when AdS, the deformation of the Poincaré group when we assume a (tiny) negative curvature in some regions, is further deformed to the quantum group qAdS. The idea is that "internal symmetries" might arise as such deformations, which seems especially appealing at root of unity because we have then finite-dimensional unitary representations.

Now if we want to try and assign multiplets of particles to (irreducible) representations of qAdS at root of 1, a first step is to know what are the dimensions of these representations. These dimensions have been found (by Jun Murakami, work in progress) for sixth root of 1, to be: 1; 4,5; 10,14,16; 35,40; 81. We chose here p = 3 in the $2p^{\text{th}}$ root of 1, physically because there are 3 generations, and mathematically because for a variety of reasons one must take $p \geq 3$, so that is the first case.

The first nontrivial representations are, as can be expected in the case of a Lie algebra of type $B_2 \equiv C_2$, of dimensions 4 and 5 (that was 3 in the case of $\mathfrak{su}(3)$, hence quarks). Thus if we want to mimic what has been done for unitary symmetries, we might have to replace, e.g., the basic octet by two "quartets", unless a doubling of the dimension can be justified by some mathematical or physical reasons (maybe looking at a corresponding supersymmetry).

Note that while all (compact) simple groups of rank 2 were studied in detail in [7] from the point of view of strong interaction symmetries and eventually type A_2 emerged, and while the (finite-dimensional) representation theory of generic quantum groups is similar to the classical case, the restricted quantum groups are so different that the B_2 type cannot be excluded a priori. A notable difference is that the restricted quantum groups have only a finite number of finite-dimensional representations, which might be an advantage.

But the knowledge of the dimensions is only the beginning of the beginning. In particular we need to study the tensor products of the representations we want to use, which helps to describe what happens when two particles interact strongly and eventually produce other particles. In the case of roots of unity these tensor products typically give rise to indecomposable representations, essentially extensions of irreducibles given by some cocycles (somewhat like in the Gupta-Bleuler formalism for the electromagnetic field). The fact might here be related to the widths of the resonances produced, but that is so far only a conjecture.

The general study of such tensor products (beyond the rank 1 case, where it is already complicated) is nontrivial mathematically. As a "warm up exercise" for rank 2, it may be worth to start with the A_2 case, i.e., $\mathcal{U}_q(\mathfrak{sl}(3))$.

If we want to have for strongly interacting particles a picture similar to what has been done so far with unitary symmetries, we could first want to assign particle multiplets with some low-dimensional representations of qAdS. But since now we have a connection (via deformations) between the free (Poincaré) and the strongly interacting symmetry (qAdS), which conceptually is an advantage, we should imagine a mechanism explaining why to assign some spins (traditionally associated with the Poincaré group) to such multiplets, and how can we have a mass spectrum inside the multiplet.

Assuming we solve these (hard) "mathematical and physical homeworks", the physical task ahead of us is even more formidable: Re-examine critically half a century of particle physics, first from the phenomenological and experimental points of view on the basis of the new symmetries. "Going back to the drawing board," we should then re-examine the present phenomenology in the new framework, including interpretations of raw experimental data. These were so far made in the context of the standard model and the quarks hypothesis, starting from nucleons and a few other particles and explaining inductively the observations in accelerators and cosmic rays within that framework.

Note that, as we have seen in Section 5, a main success of the present theory (the electroweak model) is preserved by deforming Poincaré to AdS. The speculations in this subsection are natural extensions of that in order to try and describe strong interactions using our deformation philosophy.

Inasmuch as we would "simply" replace the internal symmetries by some qAdS at root of 1, we should also, from the theoretical point of view, reexamine the various aspects (e.g., QCD) of the dynamics that was built on the possibly "clay feet" of simple unitary symmetries. That is a colossal task, but not as much as it may seem because we are not starting from scratch. A lot of the sophisticated notions introduced and theoretical advances made in the past decades might be adapted to our "deformed" view of symmetries. That could include many parts of the string framework.

Even more so since in the same spirit, it is possible that more sophisticated (and largely unexplored) mathematics would require less drastic a departure from the present puzzle, the pieces of which fit so well (so far). We shall explain that in the following.

6.2.2. Generalizations: Multiparameter, superizations and affiniza-

tions. An essential part in the representation theory of the traditional internal symmetries like SU(n) (we take usually n = 3) boils down to questions of number theory, around the Weyl group (\mathbb{S}_n for SU(n)) and the center of the group ($\mathbb{Z}_{(n)}$ in that case). One is therefore led to study what can be said of quantum deformations of $\mathcal{U}(\mathfrak{so}(3,2))$ when we take for deformation parameter an element of the group algebra of the center, $\mathbb{CZ}_{(n)}$. That is, if we want to remain in the context already studied (see the next section for a more daring suggestion) of multiparameter deformations. While the generic case is relatively well understood (cf. Section 3.3.4), the case of root of 1 seems not to have been considered. It is not even clear whether one could (or should) take the same root of 1 for all the generators of the center. The "warping effect" of the roots of 1, which manifests itself already in (11), could play tricks. The same procedure can be applied to quantizing the superalgebra $\mathfrak{osp}(1|4)$ obtained [30,38] from the realization of $\mathfrak{so}(3,2)$) as quadratic homogeneous polynomials in 4 real variables $(p_1, p_2; q_1, q_2)$ by adding the linear terms (endowed with anticommutators).

In view of possibly incorporating dynamics into the symmetry picture, and in the spirit of the string framework of "blowing up" points, often a cause of singularities, into, e.g., strings, one may then want to consider loop algebras (maps from a closed string S^1 to the symmetry in question, e.g., $\mathfrak{so}(3,2)$), and their quantization. In the same vein one may want to consider a number of infinite-dimensional algebras (Kac–Moody, Virasoro, etc.) built on that pattern. That is a very active topic (cf. e.g., [47,52]) in which many results are available. But the kinds of specific examples we would need here have not been much studied and the root of unity case even less (mildly speaking). And then one may want to say something about the very hard question of maps from something more general than S^1 (e.g., a K_3 surface or a Calabi–Yau complex 3-fold) into some groups or algebras, and the possible quantization of such structures: these are totally virgin territory.

All these generalizations, even in the specific cases we would need here for possible applications, are at least valid mathematical questions.

6.2.3. "Quantum deformations" (with noncommutative "parameter"). The "best of both worlds" might however result from a challenging idea. Internal symmetries could emerge from deforming those of space-time in a more general sense that would include the use made of unitary symmetries like SU(n). A fringe benefit might even be to give a conceptually beautiful explanation to the fact that we observe 3 generations.

The idea would be to "quantize" the Gerstenhaber definition (1) of deformations of algebras, not simply by considering a parameter that acts on the algebra (as indicated in Section 3.3.5) but by trying to develop a similar theory with a noncommutative "parameter". That has not been done and is far from obvious. It is not even clear what (if any) cohomology would be needed.

In particular one may think of deformations with a quaternionic deformation parameter. The field \mathbb{H} of quaternions is the only number field extending that of complex numbers \mathbb{C} , but it is nonabelian. So we could speak of such a theory as a "quantum deformation" since one often calls "quantum" mathematical notions that extend existing ones by "plugging in" noncommutativity. As is well known, elements of \mathbb{H} can be written in the form a+bi+cj+dk with $a, b.c.d \in \mathbb{R}, i^2 = j^2 = k^2 = -1$ and i, j, k anticommuting (like Pauli matrices). Interchanging the roles of i, j, k would give a symmetry (\mathbb{S}_3 or SO(3)) which might explain why we have 3 (and only 3) generations. And deforming $\mathcal{U}(\mathfrak{so}(3,2))$ (the choice of that real form could be important) using such a (so far, hypothetical) quantum deformation theory, especially in some sense at root of unity, might give rise to internal symmetries for which the role of the Weyl group S_3 of SU(3) (and possibly of SU(3)) would permit to re-derive, this time on a fundamental basis, all what has been done with the representations of SU(3). We would still have to explain, e.g., spin assignments to multiplets, and much more. But now we would have a frame for that, and a subtle nontrivial connection between internal and external symmetries could be developed, with all its implications (especially concerning the dynamics involved), with a relatively modest adaptation of the present empirical models.

A variant of that would be to develop a theory of deformations parametrized by the group algebra of S_n . Except for the fact that we would have to assume, e.g., n = 3 and not "explain" why we have only 3 generations, the general idea would be similar, so we shall not repeat the above speculations.

More generally, a mathematical study of such "quantum deformations" should be of independent interest, even if that uses very abstract tools that may not make it directly applicable to the physical problems we started from.

6.2.4. Remark: Quantized AdS space and related cosmology. One should not only try and develop a theory of the fundamental constituents of the matter we know (even if it constitutes only about 4% of the universe) but also explain how that small part is created, and if possible how comes that we see so little antimatter. To this end also, quantizing AdS space-time might help.

Of course, in line with recent observational cosmology, our universe is probably, "in the large", asymptotically de Sitter, with positive curvature and invariance group SO(4,1), the other simple group deformation of the Poincaré group which however, unlike AdS, does not give room to a positive energy operator. At our scale, for most practical purposes, we can treat it as Minkowskian (flat). Focussing "deeper" we would then discover that it can be considered as Anti de Sitter. There one can explain photons and leptons as composites of singletons that live in AdS space-time. It is thus natural to try and quantize that AdS space-time. And in fact, in [9], we showed how to build such "quantized hyperbolic spheres", i.e., noncommutative spectral triples à la Connes, but in a Lorentzian context, which induce in (an open orbit in) AdS space-time a pseudo-Riemannian deformation triple similar (except for the compactness of the resolvent) to the triples developed for quantized spheres by Connes et al. (see, e.g., [15]). Such a "quantized AdS space" has a horizon which permits to consider it as a black hole (similar to the BTZ black holes [4], which exist for all AdS_n when $n \geq 3$). [A kind of groupoid structure might be needed if one wants to treat all 3 regions.]

For q an even root of unity, since the corresponding quantum AdS group has finite dimensional UIRs, such a quantized AdS black hole could be considered as "q-compact" in a sense to be made precise. As we mention in [9,76], in some regions of our universe, our Minkowski space-time could be, at very small distances, both deformed to anti de Sitter and quantized, to qAdS. These regions would appear as black holes which might be found at the edge of our expanding universe, a kind of "stem cells" of the initial singularity dispersed at the Big Bang. From these (that is so far mere speculation) might emerge matter, possibly first some kind of singletons that couple and become massified by interaction with, e.g., dark matter and/or dark energy. Such a scheme could be responsible, at very large distances, for the observed positive cosmological constant – and might bring us a bit closer to quantizing gravity, the Holy Grail of modern physics, whether or not that is a relevant question (even if very recent and well publicized observations of gravitational waves might indicate that quantizing gravity is needed).

7. Epilogue and a tentative "road map"

After such a long overview involving a fireworks of fundamental mathematical and physical notions, many of which need to be developed, a natural question (which most physicists will probably ask after going through the abstract) is: why argue with success?

After all, the Standard Model is considered (so far) as the ultimate description of particle physics. We were looking for a key to knowledge under a lantern (bei der Laterne), found one which turned out to open a door nearby (not the one sought initially but never mind) through which a blue angel M led us to a beautiful avenue with many ramifications. But what if M was Fata Morgana and that avenue eventually turns out to be a dead end? We would then need a powerful flashlight (deformations maybe) to find in a dark corner a strange key with which, with much effort, a hidden door can be opened and lead us to an avenue where part of our questions can be answered.

Still, physicists tend to have a rather positivist attitude and, in most areas of physics, one takes for granted some experimental facts without trying to explain them on the basis of fundamental principles. Why would particle physics be different? A first answer is given by Einstein: "Curiosity has its own reason for existing." Theoretical particle physics is certainly an area in which very fundamental questions can be asked, and answered, if needed with the help of sophisticated mathematics to be developed.

So, at the risk of being considered simple minded, I am asking the question of why the symmetries on which is based the standard model are what they are in the model, and not only what are they and how do they work. In other words, the question is: Is it necessarily so??

Another reason (more pragmatic) is that often in mathematics, questions originating in Nature tend to be more seminal than others imagined "out of the blue". That is, if I may say so, an experimental fact. It is therefore worth developing the new mathematical tools we have indicated, and those that their development will suggest. If that solves physical problems (those intended or others), so much the better. In any case that should give nontrivial mathematics. We express the general approach as follows: **Conjecture 2.** THE DEFORMATION CONJECTURE. Internal symmetries of elementary particles emerge from their relativistic counterparts by some form of deformation (possibly generalized, including quantization).

In particular we would like to realize a Quantum Deformation Dream:

- The above mentioned "Quantum Deformations" can be defined, then permit to define "QQgroups," including a "restricted" version thereof (at roots of unity) which would be finite-dimensional algebras, and the tensor products of their representations can be studied.
- Such a procedure can be applied to $\mathcal{U}(\mathfrak{so}(3,2))$ (and $\mathcal{U}(\mathfrak{so}(1,2))$ as a toy model), if needed along with a supersymmetric extension and maybe some kind of "affinization", and serve as a starting point for a well based theory of strong interactions.

Of course one cannot rule out that symmetries, one of the bases of "the unreasonable effectiveness of mathematics in the natural sciences" [84], turn out to have little role in particle physics. I strongly doubt it.

In the list of problems we have encountered, the "simplest" on the mathematical side seems to be what happens with tensor products of representations of qAdS (or of other rank 2 quantum groups) at root of unity. In a similar spirit it would be possibly more interesting to study representations of multiparameter deformations of AdS, including their tensor products.

After some mathematical and theoretical progress we should then try and see how that knowledge can be used to interpret the known strong interactions, step by step, starting with the earliest known particles. That requires both theoretical and phenomenological studies, and possibly also renewed experiments to check the theoretical results.

A more appealing approach is to try and develop a "quantum deformations" theory, in particular with quaternions. Then, if and when we have such a theory, would come the question to apply it to AdS both in mathematics and in possible theoretical physics applications.

Ultimately one would have to study in details the phenomenological implications of these developments. If they differ somewhat from the present interpretation of the raw experimental data, we would need to revise that interpretation and possibly to re-do some experiments.

Note that a "fringe benefit" of any such revision is that it can be done using the present experimental tools, if needed (as far as more ancient data are concerned) with the refinement of modern technology. The civil society is not likely to give us significantly more powerful accelerators.

All these are problems worthy of attack. It can be expected that they will prove their worth by hitting back. Starting from a primary question, I have asked many more, combined that with many notions and results developed during half a century of research in physical mathematics, and indicated avenues along which some of these might be answered. At 75 I leave it to the next generations to enter that promised land and tackle the many problems that will follow.

Acknowledgment

This panorama benefited from 35 years of active collaboration with Moshe Flato (and coworkers), from close to \infty discussions with many scientists (too numerous to be listed individually) and from the hospitality of Japanese colleagues during my so far over 10 years of relatively generous retirement from France, after 42 years of (sometimes bumpy) career in CNRS. I want also to thank the Editors of this volume for their friendly patience.

References

- E. Angelopoulos, Sur les représentations unitaires irréductibles de SO₀(p, 2), C. R. Acad. Sci. Paris Sér. I Math. **292** (1981), 469–471.
- [2] E. Angelopoulos, M. Flato, C. Fronsdal, and D. Sternheimer, Massless particles, conformal group, and de Sitter universe, Phys. Rev. D (3) 23 (1981), no. 6, 1278–1289.
- [3] E. Angelopoulos and M. Laoues, Singletons on AdS_n. Conférence Moshé Flato 1999, Vol. II (Dijon), 3–23, Math. Phys. Stud. 22, Kluwer Acad. Publ., Dordrecht, 2000.
- [4] M. Bañados, M. Henneaux, C. Teitelboim, and J. Zanelli, Geometry of the 2+1 black hole, Phys. Rev. D (3) 48 (1993), 1506–1525.
- [5] H. Basart, M. Flato, A. Lichnerowicz and D. Sternheimer, Deformation theory applied to quantization and statistical mechanics, Lett. Math. Phys. 8 (1984), 483–494.
- [6] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz and D. Sternheimer, Deformation theory and quantization I. Deformations of symplectic structures, and II. Physical applications Ann. Physics 111 (1978), 61–110 and 111–151.
- [7] R.E. Behrends, J. Dreitlein, C. Fronsdal, B.W. Lee, Simple groups and strong interaction symmetries. Rev. Modern Phys. 34 (1962) 1–40.
- [8] J. Beringer et al. (Particle Data Group), 2013 Review of Particle Physics, Phys. Rev. D86, 010001 (2012) and 2013 partial update for the 2014 edition, available at http://pdg.lbl.gov/
- [9] P. Bieliavsky, L. Claessens, D. Sternheimer and Y. Voglaire, Quantized anti de Sitter spaces and non-formal deformation quantizations of symplectic symmetric spaces, Poisson geometry in mathematics and physics, 1–24, Contemp. Math., 450, Amer. Math. Soc., Providence, RI, 2008.
- [10] P. Bonneau, M. Flato, M. Gerstenhaber and G. Pinczon, The hidden group structure of quantum groups: strong duality, rigidity and preferred deformations, Comm. Math. Phys. 161 (1994), 125–156.
- P. Bonneau, M. Gerstenhaber, A. Giaquinto and D. Sternheimer, Quantum groups and deformation quantization: explicit approaches and implicit aspects, J. Math. Phys. 45(10) (2004), 3703–3741.
- [12] S. Coleman and J. Mandula, All Possible Symmetries of the S Matrix, Phys. Rev. 159 (1967), 1251–1256.
- [13] A. Chamseddine, A. Connes and W. van Suijlekom, Beyond the Spectral Standard Model: Emergence of Pati-Salam Unification, JHEP 1311 (2013) 132.
- [14] A. Connes, Noncommutative Geometry, Academic Press, San Diego 1994.

- [15] A. Connes and M. Dubois-Violette, Moduli space and structure of noncommutative 3-spheres, Lett. Math. Phys. 66 (2003), 99–121.
- [16] F. Costantino and J. Murakami, On the SL(2,C) quantum 6j-symbols and their relation to the hyperbolic volume. Quantum Topol. 4 (2013), 303–351, and J. Murakami, Private communications (2013, 2014).
- [17] C. De Concini and V. Kac, Representations of quantum groups at roots of 1 pp. 471–506 in Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989), Progr. Math., 92, Birkhäuser Boston 1990.
- [18] E.E. Demidov, Yu.I. Manin, E.E. Mukhin, D.V. Zhdanovich, Nonstandard quantum deformations of GL(n) and constant solutions of the Yang-Baxter equation. Common trends in mathematics and quantum field theories (Kyoto, 1990). Progr. Theoret. Phys. Suppl. No. 102 (1990), 203–218 (1991).
- [19] P.A.M. Dirac, *The relation of Classical to Quantum mechanics*, Proc. Second Canadian Math. Congress, Vancouver, 1949, pp. 10–31. University of Toronto Press 1951.
- [20] P.A.M. Dirac, A remarkable representation of the 3+2 de Sitter group, J. Math. Phys. 4 (1963), 901–909.
- [21] G. Dito and D. Sternheimer, Deformation quantization: genesis, developments and metamorphoses, pp. 9-54 in: Deformation quantization (Strasbourg 2001), IRMA Lect. Math. Theor. Phys., 1, Walter de Gruyter, Berlin 2002 (math.QA/0201168)
- [22] V. Drinfel'd, Proceedings Quantum Groups, of $_{\mathrm{the}}$ International Congress of Mathematicians, Vol. 1, 2(Berkeley, Calif., 1986),798 - 820, Amer. Math. Soc., Providence, RI, 1987. Available as: http://www.mathunion.org/ICM/ICM1986.1/Main/icm1986.1.0798.0820.ocr.pdf
- [23] L.D. Faddeev, N.Yu. Reshetikhin and L.A. Takhtajan, Quantization of Lie groups and Lie algebras, pp. 129–139 in: Algebraic analysis, Vol. I, Academic Press, Boston 1988.
- [24] Quoted, e.g., in Helge Kragh, Quantum Generations (1999), 321. See http://todayinsci.com/QuotationsCategories/B_Cat/Biography-Quotations.htm See also http://www.goodreads.com/author/quotes/107005.Enrico_Fermi
- [25] Laura Fermi, Atoms in the Family: My Life with Enrico Fermi, University of Chicago Press (1954). [My uncle Richard Sternheimer, then MD at the University of Chicago, had been the physician of the physicists working for the Manhattan Project and remained the Fermi family doctor...]
- [26] M. Flato, Deformation view of physical theories, Czechoslovak J. Phys. B32(4) (1982), 472–475.
- [27] M. Flato and C. Fronsdal, Quantum field theory of singletons. The Rac, J. Math. Phys. 22 (1981), 1100–1105.
- [28] M. Flato and C. Fronsdal, Composite Electrodynamics, J. Geom. Phys. 5 (1988), 37–61. [See also: M. Flato, C. Fronsdal and D. Sternheimer, Singletons as a basis for composite conformal quantum electrodynamics, pp. 65–76, in Quantum Theories and Geometry, Mathematical Physics Studies 10, Kluwer Acad. Publ., Dordrecht 1988.]
- [29] M. Flato, C. Frønsdal and D. Sternheimer, Singletons, physics in AdS universe and oscillations of composite neutrinos, Lett. Math. Phys. 48 (1999), 109–119.

- [30] M. Flato, L.K. Hadjiivanov and I.T. Todorov, Quantum Deformations of Singletons and of Free Zero-Mass Fields, Foundations of Physics 23 (1993), 571– 586.
- [31] M. Flato and P. Hillion, Poincaré-like group associated with neutrino physics, and some applications, Phys. Rev. D(3) 1 (1970), 1667–1673.
- [32] M. Flato, J.C.H. Simon and E. Taflin, Asymptotic completeness, global existence and the infrared problem for the Maxwell-Dirac equations, Mem. Amer. Math. Soc. 127 (1997), no. 606, x+311 pp.
- [33] M. Flato and D. Sternheimer, Remarks on the connection between external and internal symmetries, Phys. Rev. Letters 15 (1965), 934–936.
- [34] M. Flato and D. Sternheimer, Local representations and mass-spectrum, Phys. Rev. Letters 16 (1966), 1185–1186.
- [35] M. Flato and D. Sternheimer, Nonuniqueness of Mass-Formulas from Infinite-Dimensional Groups and Algebras, Phys. Rev. Letters 19 (1967), 254–255.
- [36] M. Flato and D. Sternheimer, Poincaré partially integrable local representations and mass-spectrum, Comm. Math. Phys. 12 (1969), 296–303.
- [37] M. Flato and D. Sternheimer, On an infinite-dimensional group, Comm. Math. Phys. 14 (1969), 5–12.
- [38] C. Frønsdal, Dirac supermultiplet, Phys. Rev. D 26 (1982), 1988–1995.
- [39] C. Frønsdal, Generalization and exact deformations of quantum groups, Publ. Res. Inst. Math. Sci. 33 (1997), 91–149.
- [40] C. Frønsdal, Singletons and neutrinos, Lett. Math. Phys. 52 (2000), 51–59, Conference Moshé Flato 1999 (Dijon) (hep-th/9911241).
- [41] C. Frønsdal and A. Galindo, Deformations of multiparameter quantum gl(N), Lett. Math. Phys. 34 (1995), 25–36.
- [42] L. Gårding, Note on continuous representations of Lie groups, Proc. Nat. Acad. Sci. U.S.A. 33 (1947), 331–332.
- [43] M. Gell-Mann, Symmetries of baryons and mesons. Phys. Rev. (2) 125 (1962), 1067–1084. [California Institute of Technology Synchrotron Laboratory Report CTSL-20 (1961)].
- [44] M. Gerstenhaber, On the deformation of rings and algebras, Ann. Math. (2) 79 (1964), 59–103; and (IV), *ibid.* 99 (1974), 257–276.
- [45] A. Grothendieck, Techniques de construction en géométrie analytique, in: Familles d'Espaces complexes et Fondements de la Géométrie analytique, Séminaire H. Cartan 13 (1960/61), École Norm. Sup. Paris (1962).
- [46] R. Haag, J. Lopuszański and M. Sohnius, All possible generators of supersymmetries of the S-matrix, Nuclear Phys. B 88 (1975), 257–274.
- [47] D. Hernandez and M. Jimbo, Asymptotic representations and Drinfeld rational fractions, Compos. Math. 148 (2012), 1593–1623.
- [48] G. 't Hooft, Dimensional reduction in quantum gravity, pp. 284–296 in Salamfestschrift: a collection of talks (A. Ali, J. Ellis and S. Randjbar-Daemi eds.), World Scientific 1993 (gr-qc/9310026).
- [49] G. 't Hooft, The Fate of the Quantum, arXiv:1308.1007 [quant-ph].
- [50] G. 't Hooft, Hamiltonian formalism for integer-valued variables and integer time steps and a possible application in quantum physics, arXiv:1312.1229 [math-ph].

- [51] E. Inönü and E.P. Wigner, On the contraction of groups and their representations, Proc. Nat. Acad. Sci. U. S. A. 39 (1953), 510–524.
- [52] R-Q. Jian and M. Rosso, Braided cofree Hopf algebras and quantum multi-brace algebras, J. Reine Angew. Math. 667 (2012), 193–220.
- [53] M. Jimbo, A q-difference algebra of U(g) and the Yang-Baxter equation, Lett. Math. Phys. 10 (1985), 63–69.
- [54] R. Jost, Eine Bemerkung zu einem "Letter" von L. O'Raifeartaigh und einer Entgegnung von M. Flato und D. Sternheimer, Helv. Phys. Acta 39 (1966), 369–375.
- [55] M. Kibler, The Master thesis of Moshé Flato, Conférence Moshé Flato 1999, Vol. II (Dijon), 177–184, Math. Phys. Stud., 22, Kluwer Acad. Publ., Dordrecht, 2000.
- [56] K. Kodaira and D.C. Spencer, On deformations of complex analytic structures, Ann. Math. 67 (1958), 328–466.
- [57] H. Kondo and Y. Saito, Indecomposable decomposition of tensor products of modules over the restricted quantum universal enveloping algebra associated to sl₂, J. Algebra **330** (2011), 103–129.
- [58] M. Lévy-Nahas, Deformations and contractions of Lie algebras, J. Math. Phys. 8 (1967), 1211–1222.
- [59] G. Lusztig, Finite-dimensional Hopf algebras arising from quantized universal enveloping algebra, J. Amer. Math. Soc. 3 (1990), 257–296.
- [60] G. Lusztig, Quantum groups at roots of 1, Geom. Dedicata **35** (1990), 89–113.
- [61] S. Majid, What is ... a Quantum Group, Notices Amer. Math. Soc. 53 (2006), 30-31. http://www.ams.org/notices/200601/what-is.pdf
- [62] Yu.I. Manin, Topics in noncommutative geometry. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 1991.
- [63] F. Nadaud, Generalised deformations, Koszul resolutions, Moyal Products, Rev. Math. Phys. 10 (1998), 685–704.
 & Thèse, Dijon (January 2000).
- [64] F. Nadaud, Generalized Deformations and Hochschild Cohomology, Lett. Math. Phys. 58 (2001), 41–55.
- [65] Y. Ne'eman, Derivation of strong interactions from a gauge invariance. Nuclear Phys. 26 (1961), 222–229.
- [66] L.B. Okun, The impact of the Sakata model, Prog. Theor. Phys. Suppl. 167 (2007), 163–174.
- [67] L. O'Raifeartaigh, Mass differences and Lie algebras of finite order, Phys. Rev. Lett. 14 (1965), 575–577.
- [68] G. Pinczon, Non commutative deformation theory, Lett. Math. Phys. 41 (1997), 101–117.
- [69] N. Reshetikhin, Multiparameter quantum groups and twisted quasitriangular Hopf algebras, Lett. Math. Phys. 20 (1990), no. 4, 331–335.
- [70] N. Reshetikhin and V.G. Turaev, Invariants of 3-manifolds via link polynomials and quantum groups, Invent. Math. 103 (1991), 547–597.
- [71] S. Sakata, On a composite model for the new particles, Prog. Theor. Phys. (Letter) 16 (1956), 686–688.

- [72] I.E. Segal, A class of operator algebras which are determined by groups, Duke Math. J. 18 (1951), 221–265.
- [73] H. Steinacker, Finite-dimensional unitary representations of quantum anti-de Sitter groups at roots of unity. Comm. Math. Phys. 192 (1998), 687–706.
- [74] H. Steinacker, Unitary representations of noncompact quantum groups at roots of unity, Rev. Math. Phys. 13 (2001), 1035–1054.
- [75] D. Sternheimer, Quantization is deformation, pp. 331–352 in Noncommutative geometry and representation theory in mathematical physics, Contemp. Math. 391, Amer. Math. Soc., Providence, RI, 2005.
- [76] D. Sternheimer, The geometry of space-time and its deformations from a physical perspective, in From geometry to quantum mechanics, 287–301, Progr. Math. 252, Birkhäuser Boston, 2007.
- [77] D. Sternheimer, A very short presentation of deformation quantization, some of its developments in the past two decades, and conjectural perspectives. Travaux mathématiques. Volume XX, 205-228, Fac. Sci. Technol. Commun. Univ. Luxemb., Luxembourg, 2012. http://www.uni.lu/content/download/56018/661547/file/sternheimer.pdf
- [78] E. Weimar-Woods, Contractions, generalized Inönü-Wigner contractions and deformations of finite-dimensional Lie algebras, Rev. Math. Phys. 12 (2000), 1505–1529.
- [79] J. Wess and B. Zumino, Supergauge transformations in four dimensions, Nuclear Physics B 70 (1974), 39–50.
- [80] H. Weyl, The theory of groups and quantum mechanics, Dover, New-York 1931, edited translation of Gruppentheorie und Quantenmechanik, Hirzel Verlag, Leipzig 1928. Quantenmechanik und Gruppentheorie, Z. Physik 46 (1927), 1–46.
- [81] E.P. Wigner, Quantum corrections for thermodynamic equilibrium, Phys. Rev.
 - **40** (1932), 749–759.
- [82] E. Wigner, On unitary representations of the inhomogeneous Lorentz group, Ann. of Math. (2) 40 (1939), 149–204.
- [83] E. Wigner, Group theory: And its application to the quantum mechanics of atomic spectra. Expanded and improved ed. Translated from the German by J. J. Griffin. Pure and Applied Physics. Vol. 5, Academic Press, New York-London 1959.
- [84] E.P. Wigner, The unreasonable effectiveness of mathematics in the natural sciences, Comm. Pure Appl. Math. 13 (1960), 1–14.
- [85] Jie Xiao, Finite-dimensional representations of $U_t(sl(2))$ at roots of unity, Canad. J. Math. **49** (1997), 772–787.

Daniel Sternheimer

Department of Mathematics, Rikkyo University,

3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501, Japan,

& Institut de Mathématiques de Bourgogne, Université de Bourgogne BP 47870, F-21078 Dijon Cedex, France.

e-mail: Daniel.Sternheimer@u-bourgogne.fr