

# THE ALGEBRAIC AND ANALYTIC COMPACTIFICATIONS OF THE HITCHIN MODULI SPACE

SIQI HE, RAFFAELLA MAZZEO, XUESEN NA, AND RICHARD A. WENTWORTH

ABSTRACT. Following the work of Mazzeo-Swoboda-Weiss-Witt [30] and Mochizuki [33], there is a map  $\Xi$  between the algebraic compactification of the Dolbeault moduli space of  $\mathrm{SL}(2, \mathbb{C})$  Higgs bundles on a smooth projective curve coming from the  $\mathbb{C}^*$  action, and the analytic compactification of Hitchin's moduli space of solutions to the  $\mathrm{SU}(2)$  self-duality equations on a Riemann surface by adding solutions to the decoupled equations, known as "limiting configurations". This map extends the classical Kobayashi-Hitchin correspondence. The main result of this paper is that  $\Xi$  fails to be continuous at the boundary over a certain subset of the discriminant locus of the Hitchin fibration. This suggests the possibility of a third, refined compactification which dominates both.

## 1. INTRODUCTION

Let  $\Sigma$  be a closed Riemann surface of genus  $g \geq 2$ . The coarse Dolbeault moduli space of  $\mathrm{SL}(2, \mathbb{C})$  semistable Higgs bundles on  $\Sigma$ , denoted by  $\mathcal{M}_{\mathrm{Dol}}$ , and Hitchin's moduli space of solutions to the  $\mathrm{SU}(2)$  self-duality equations on  $\Sigma$ , denoted by  $\mathcal{M}_{\mathrm{Hit}}$ , have been extensively studied since their introduction over 35 years ago. The Kobayashi-Hitchin correspondence, proved in [21], gives a homeomorphism between these two moduli spaces:

$$\Xi : \mathcal{M}_{\mathrm{Dol}} \rightarrow \mathcal{M}_{\mathrm{Hit}} .$$

The space  $\mathcal{M}_{\mathrm{Dol}}$  is naturally a quasiprojective variety [35, 45], and similarly to monopole moduli spaces,  $\mathcal{M}_{\mathrm{Hit}}$  fails to be compact. Recently, there has been interest from several directions on natural compactifications of these two spaces. A key feature on the Dolbeault side is the existence of a  $\mathbb{C}^*$  action with the Białynicki-Birula property, and this may be used to define a completion of  $\mathcal{M}_{\mathrm{Dol}}$  as a projective variety [19, 7, 11]. The ideal points are identified with the  $\mathbb{C}^*$  orbits in the complement of the nilpotent cone of  $\mathcal{M}_{\mathrm{Dol}}$ . The Hitchin moduli space also admits a more recently introduced compactification,  $\overline{\mathcal{M}}_{\mathrm{Hit}}$ , based on the work of several authors (see [30, 33, 47]). The boundary of  $\overline{\mathcal{M}}_{\mathrm{Hit}}$  is given by gauge equivalence classes of *limiting configurations*. This compactification is relevant to many aspects of Hitchin's moduli space. For more details, we refer the reader to [10, 29, 14, 15, 36, 27, 3], and the references therein.

By the work of [30, 33], there is a natural extension

$$\overline{\Xi} : \overline{\mathcal{M}}_{\mathrm{Dol}} \rightarrow \overline{\mathcal{M}}_{\mathrm{Hit}}$$

of the Kobayashi-Hitchin correspondence to the two compactifications described above, and it is of interest to study the geometry of this map. This involves another key feature of Hitchin's moduli space; namely, spectral curves. Spectral curves and spectral data [22] play a central role in the realization of the Dolbeault moduli space as an algebraically complete integrable system  $\mathcal{H} : \mathcal{M}_{\mathrm{Dol}} \rightarrow \mathcal{B}$ . In the case of  $\mathrm{SL}(2, \mathbb{C})$ , the base  $\mathcal{B}$  is the space of holomorphic quadratic differentials on  $\Sigma$ . Given  $q \in H^0(K^2)$ , one obtains a (scheme theoretic) spectral curve  $S_q$ . This curve is reduced if  $q \neq 0$ , irreducible if  $q$  is not the square of an abelian differential, and smooth if  $q$  has simple zeros. We let  $\mathcal{B}^{\mathrm{reg}} \subset \mathcal{B}$  denote the open cone of quadratic differentials with simple zeros.

The ideal points of both compactifications  $\overline{\mathcal{M}}_{\text{Dol}}$  and  $\overline{\mathcal{M}}_{\text{Hit}}$  have associated nonzero quadratic differentials, and therefore spectral curves. We write  $\overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}}$  for the elements in  $\overline{\mathcal{M}}_{\text{Dol}}$  with smooth spectral curves, and  $\overline{\mathcal{M}}_{\text{Dol}}^{\text{sing}} = \overline{\mathcal{M}}_{\text{Dol}} \setminus \overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}}$  for those with singular spectral curves; similarly for  $\overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}$  and  $\overline{\mathcal{M}}_{\text{Hit}}^{\text{sing}}$ . We then have the following result.

**Theorem 1.1.** *The restriction of the compactified Kobayashi-Hitchin map  $\overline{\Xi} : \overline{\mathcal{M}}_{\text{Dol}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}$  to the locus with smooth associated spectral curves defines a homeomorphism  $\overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} \simeq \overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}$ . On the singular spectral curve locus, however,  $\overline{\Xi}^{\text{sing}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{sing}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{sing}}$  is neither surjective nor injective.*

It is convenient to analyze the behavior along rays in  $\mathcal{B}$ , where the spectral curve is simply rescaled. Let  $q \neq 0$  be a quadratic differential and  $\overline{\mathcal{M}}_{\text{Dol},q}, \overline{\mathcal{M}}_{\text{Hit},q}$  be the points with spectral curve  $S_q$ . The restriction of  $\overline{\Xi}$  gives us the map  $\overline{\Xi}_q : \overline{\mathcal{M}}_{\text{Dol},q} \rightarrow \overline{\mathcal{M}}_{\text{Hit},q}$ . We shall study the continuous behavior of  $\overline{\Xi}_{tq}$  for values  $tq$ ,  $t \in \mathcal{R}^+$ , as  $t \rightarrow +\infty$ . For convenience, we set  $\mathcal{M}_q := \overline{\mathcal{M}}_{\text{Dol},q} \cap \mathcal{M}_{\text{Dol}}$ . When  $q$  is irreducible, i.e. not a square, all elements in  $\mathcal{M}_q$  are stable. Via the Hitchin [23] and Beauville-Narasimhan-Ramanan (BNR) correspondence [1], this reduces the description of the fiber  $\mathcal{H}^{-1}(q)$  to the characterization of rank 1 torsion free sheaves on the integral curve  $S_q$ .

In [38], parameter spaces for rank 1 torsion free sheaves on algebraic curves with Gorenstein singularities were studied in the context of compactified Jacobians, and the crucial notion of a parabolic module was introduced. This was extensively investigated by Cook in [5, 4], partially following ideas of Bhosle [2]. For simple plane curve singularities of the type appearing in spectral curves, one makes use of the local classification of torsion free modules of Greuel-Knörrer [17]. These methods were applied to study the Hitchin fibration by Gothen-Oliveira in [16], see also [28] for recent study. In parallel, in [24], Horn defines a stratification of  $\mathcal{M}_q = \bigcup_D \mathcal{M}_{q,D}$  by certain effective divisors contained in the divisor of  $q$  (see Section 5.5, and also [26] for the more general situation). Using the results from these references, we have a reinterpretation of Mochizuki's construction [33]. This leads to the following result.

**Theorem 1.2.** *Let  $q \neq 0$  be an irreducible quadratic differential.*

- (1) *If  $q$  has only zeros of odd order, then  $\overline{\Xi}_q$  is continuous.*
- (2) *If  $q$  has at least one zero of even order, then for each  $D \neq 0$  there exists an integer  $n_D \geq 1$  so that for any Higgs bundle  $(\mathcal{F}, \psi) \in \mathcal{M}_{q,D}$ , there exist  $n_D$  sequences of Higgs bundles  $(\mathcal{E}_i^k, \varphi_i^k)$  with  $k = 1, \dots, n_D$  satisfying*
  - $\lim_{i \rightarrow \infty} (\mathcal{E}_i^k, \varphi_i^k) = (\mathcal{F}, \psi)$  for  $k = 1, \dots, n_D$ ,
  - and if we write

$$\eta^k := \lim_{i \rightarrow \infty} \overline{\Xi}_q(\mathcal{E}_i^k, \varphi_i^k) \quad , \quad \xi := \lim_{i \rightarrow \infty} \overline{\Xi}_q(\mathcal{F}, t_i \psi) \quad ,$$

*for some sequence  $t_i \in \mathbb{R}^+$ ,  $t_i \rightarrow +\infty$ , then  $\xi, \eta^1, \dots, \eta^{n_D}$  are  $n_D + 1$  different limiting configurations.*

When  $q$  is reducible, the description of Higgs bundles in  $\mathcal{M}_q$  becomes more complicated because, among other things, of the existence of strictly semistable objects. To understand this, we use the local descriptions of Gothen-Oliveira and Mochizuki (see [16, 33]). Our result, which focuses on the stable locus, is the following.

**Theorem 1.3.** *Suppose  $q$  is reducible. Let  $\overline{\mathcal{M}}_{\text{Dol},q}^{\text{st}}$  denote the stable locus of  $\overline{\mathcal{M}}_{\text{Dol},q}$ . If  $g \geq 3$ , then the restriction map  $\overline{\Xi}_q|_{\overline{\mathcal{M}}_{\text{Dol},q}^{\text{st}}}$  is discontinuous. However, if  $g = 2$ , the map  $\overline{\Xi}_q|_{\overline{\mathcal{M}}_{\text{Dol},q}^{\text{st}}}$  is continuous.*

We also note that there are also recent work of Mochizuki and Szabó [34] on the asymptotic behavior for family of Higgs bundles in higher rank case.

This paper is organized as follows: in Section 2, we provide an overview of Higgs bundles and BNR correspondence. In Section 3, we delve into the concepts of filtered bundles and their compactness properties. Section 4 defines the algebraic and analytic compactifications. Section 5 introduces parabolic modules and examines their connection to spectral curves. The main results for Hitchin fibers with irreducible singular spectral curves are established in Section 6. In Section 7, the results for the reducible case are proven. Finally, in Section 8, we construct the compactified Kobayashi-Hitchin map and prove the main results. The Appendix, based on the work of Greuel-Knörrer, calculates some invariants of rank 1 torsion free sheaves on the spectral curves we consider.

**Acknowledgements.** We extend our sincere gratitude to Takurō Mochizuki for his valuable insights and stimulating discussions during the BIRS conference in 2021. The authors also wish to express their gratitude to a great many people for their interest and helpful comments. Among them are Mark de Cataldo, Ron Donagi, Simon Donaldson, Laura Fredrickson, Johannes Horn, Laura Schaposnik, Shizhang Li, Jie Liu, Tony Pantev, Thomas Walpuski, Daxin Xu. S.H is supported by NSFC grant No.12288201. R.W.'s research is supported by NSF grants DMS-1906403 and DMS-2204346. The authors gratefully acknowledge the support and hospitality of the MSRI under NSF grant DMS-1928930 during the program “Analytic and Geometric Aspects of Gauge Theory”, Fall 2022.

## 2. BACKGROUND ON HIGGS BUNDLES

This section gives a very brief overview of the Dolbeault and Hitchin moduli spaces, spectral curve descriptions, and the nonabelian Hodge correspondence. For a more details on these topics, see [21, 23, 44, 48].

**2.1. Higgs bundles.** As in the Introduction, throughout this paper  $\Sigma$  will denote a closed Riemann surface of genus  $g \geq 2$  with structure sheaf  $\mathcal{O} = \mathcal{O}_\Sigma$  and canonical bundle  $K = K_\Sigma$ . Let  $E \rightarrow \Sigma$  be a complex vector bundle. A Higgs bundle consists of a pair  $(\mathcal{E}, \varphi)$ , where  $\mathcal{E}$  is a holomorphic bundle and  $\varphi \in H^0(\text{End}(\mathcal{E}) \otimes K)$ . If  $\text{rank}(\mathcal{E}) = 1$ , then a Higgs field is just an abelian differential  $\omega$ . The pair  $(\mathcal{E}, \varphi)$  is called an  $\text{SL}(2, \mathbb{C})$  Higgs bundle if  $\text{rank}(E) = 2$ ,  $\det(\mathcal{E})$  has a fixed isomorphism with the trivial bundle, and  $\text{Tr}(\varphi) = 0$ . In this paper we will focus mainly on  $\text{SL}(2, \mathbb{C})$  Higgs bundles, but the rank 1 case will also be important.

Let  $(\mathcal{E}, \varphi)$  be an  $\text{SL}(2, \mathbb{C})$  Higgs bundle. A (proper) Higgs subbundle of  $(\mathcal{E}, \varphi)$  is a holomorphic line bundle  $\mathcal{L} \subset \mathcal{E}$  that is  $\varphi$ -invariant, i.e.  $\varphi : \mathcal{L} \rightarrow \mathcal{L} \otimes K$ . In this case, the restriction  $\varphi_{\mathcal{L}} := \varphi|_{\mathcal{L}}$ , makes  $(\mathcal{L}, \varphi_{\mathcal{L}})$  a rank 1 Higgs bundle. Moreover,  $\varphi$  induces a Higgs bundle structure on the quotient  $\mathcal{E}/\mathcal{L}$ . We say  $(\mathcal{E}, \varphi)$  is stable (resp. semistable) if for all Higgs subbundles  $\mathcal{L}$ , then  $\deg \mathcal{L} < 0$  (resp.  $\deg \mathcal{L} \leq 0$ ). We say  $(\mathcal{E}, \varphi)$  is polystable if  $(\mathcal{E}, \varphi) \simeq (\mathcal{L}, \omega) \oplus (\mathcal{L}^{-1}, -\omega)$ , where  $\mathcal{L}$  is a degree zero holomorphic line bundle and  $\omega \in H^0(K)$ .

If  $(\mathcal{E}, \varphi)$  is strictly semistable, i.e. semistable but not stable, the Seshadri filtration [40] gives a unique Higgs subbundle  $0 \subset (\mathcal{L}, \omega) \subset (\mathcal{E}, \varphi)$  with  $\deg(\mathcal{L}) = \frac{1}{2} \deg(\mathcal{E}) = 0$ . Write  $(\mathcal{L}', \omega') := (\mathcal{E}, \varphi)/(\mathcal{L}, \omega)$ , then we have  $\omega' = -\omega$  and  $\mathcal{L}' = \mathcal{L}^{-1}$ . The associated graded bundle  $\text{Gr}(\mathcal{E}, \varphi) = (\mathcal{L}, \omega) \oplus (\mathcal{L}^{-1}, -\omega)$  of this filtration is a polystable  $\text{SL}(2, \mathbb{C})$  Higgs bundle. We say that  $(\mathcal{E}, \varphi)$  is S-equivalent to  $\text{Gr}(\mathcal{E}, \varphi)$ .

Holomorphic bundles  $\mathcal{E}$  with underlying  $C^\infty$  bundle  $E$  are in 1-1 correspondence with  $\bar{\partial}$ -operators  $\bar{\partial}_E : \Omega^0(E) \rightarrow \Omega^{0,1}(E)$ . We use the notation  $\mathcal{E} := (E, \bar{\partial}_E)$ . Let  $\mathcal{C}$  denote the space of pairs  $(\bar{\partial}_E, \varphi)$ ,  $\bar{\partial}_E \varphi = 0$ . Let  $\mathcal{C}^s$  and  $\mathcal{C}^{ss}$  denote the subspaces of  $\mathcal{C}$  where the Higgs bundles are stable (resp. semistable). The complex gauge transformation group  $\mathcal{G}_{\mathbb{C}} := \text{Aut}(E)$  has a right action on  $\mathcal{C}$  by defining for  $g \in \mathcal{G}_{\mathbb{C}}$ ,  $(\bar{\partial}_E, \varphi)g := (g^{-1} \circ \bar{\partial} \circ g, g^{-1} \circ \varphi \circ g)$ .

There is a quasiprojective scheme  $\mathcal{M}_{\text{Dol}}$  whose closed points are in 1-1 correspondence with polystable Higgs bundles constructed via (finite dimensional) Geometric Invariant Theory (see [35, 45]). In [12] it was shown that the infinite dimensional quotient  $\mathcal{C}^{\text{ss}} // \mathcal{G}_{\mathbb{C}}$ , where the double slash indicates that S-equivalent orbits are identified, admits the structure of a complex analytic space that is biholomorphic to the analytification  $\mathcal{M}_{\text{Dol}}^{\text{an}}$  of  $\mathcal{M}_{\text{Dol}}$ . Henceforth, we shall work in the complex analytic category, identify the algebro-geometric and gauge theoretic moduli spaces, and simply denote them both by  $\mathcal{M}_{\text{Dol}}$ . We note that the set of stable Higgs bundles modulo gauge transformations,  $\mathcal{M}_{\text{Dol}}^s := \mathcal{C}^s // \mathcal{G}_{\mathbb{C}}$ , is a geometric quotient and an open subset of  $\mathcal{M}_{\text{Dol}}$ .

Finally, notice that the a pair  $(\mathcal{E}, \varphi)$  is stable (resp. semistable) if and only if the same is true for  $(\mathcal{E}, \lambda\varphi)$ ,  $\lambda \in \mathbb{C}^*$ . Hence,  $\mathcal{M}_{\text{Dol}}$  admits an action of  $\mathbb{C}^*$  that preserves  $\mathcal{M}_{\text{Dol}}^s$ . While  $\mathcal{M}_{\text{Dol}}$  is only quasiprojective, the  $\mathbb{C}^*$  action satisfies the Białynicki-Birula property:

**Theorem 2.1** ([21, 44]). *For any  $[(\mathcal{E}, \varphi)] \in \mathcal{M}_{\text{Dol}}$ ,*

$$\lim_{\lambda \rightarrow 0} \lambda \cdot [(\mathcal{E}, \varphi)] = \lim_{\lambda \rightarrow 0} [(\mathcal{E}, \lambda\varphi)]$$

*exists in  $\mathcal{M}_{\text{Dol}}$ .*

**2.2. Spectral curves and the Hitchin fibration.** The Hitchin map is defined as

$$\mathcal{H} : \mathcal{M}_{\text{Dol}} \longrightarrow H^0(K^2), \quad [(\mathcal{E}, \varphi)] \mapsto \det(\varphi),$$

where  $H^0(K^2) =: \mathcal{B}$  is known as the Hitchin base. Hitchin [21, 23] showed that  $\mathcal{H}$  is a proper map and a fibration by abelian varieties over the open cone  $\mathcal{B}^{\text{reg}} \subset \mathcal{B}$  consisting of nonzero quadratic differentials with only simple zeros. The discriminant locus  $\mathcal{B}^{\text{sing}} := \mathcal{B} \setminus \mathcal{B}^{\text{reg}}$  consists of quadratic differentials that are either identically zero or have at least one zero with multiplicity. For  $q \in \mathcal{B}$ , let  $\mathcal{M}_q := \mathcal{H}^{-1}(q)$ . The “most singular fiber”  $\mathcal{M}_0$  is called the *nilpotent cone*.

Consider the total space  $\text{Tot}(K)$  of  $K$ , along with its projection to  $\Sigma$ , denoted as  $\pi : \text{Tot}(K) \rightarrow \Sigma$ . The pullback bundle  $\pi^*K$  has a tautological section, which we denote as  $\lambda \in H^0(\text{Tot}(K), \pi^*K)$ . Given any  $q \neq 0 \in H^0(K^2)$ , the *spectral curve*  $S_q$  associated with  $q$  is the zero scheme of the section  $\lambda^2 - \pi^*q \in H^0(\text{Tot}(K), \pi^*K)$ . This is a reduced, but possibly reducible, projective algebraic curve. The restriction of  $\pi$  to  $S_q$ , also denoted as  $\pi : S_q \rightarrow \Sigma$ , is a double covering branched along the zeros of  $q$ .

The spectral curve  $S_q$  is smooth if and only if  $q$  has only simple zeros. It is reducible if and only if  $q = -\omega \otimes \omega$  for some  $\omega \in H^0(K)$ . We shall refer to such quadratic differentials as *reducible*, and *irreducible* otherwise. There is a noteworthy observation regarding irreducible spectral curves.

**Proposition 2.2** (cf. [23]). *Let  $(\mathcal{E}, \varphi)$  be a Higgs bundle with  $q = \det(\varphi)$ , and suppose  $q$  is irreducible. Then  $(\mathcal{E}, \varphi)$  has no proper invariant subbundles. In particular,  $(\mathcal{E}, \varphi)$  is stable.*

*Proof.* Suppose  $\mathcal{L} \subset \mathcal{E}$  were  $\varphi$ -invariant, and let  $\varphi_{\mathcal{L}}$  be the restriction. Then

$$\det \varphi = -\frac{1}{2} \text{Tr}(\varphi^2) = -(\varphi_{\mathcal{L}})^2,$$

contradicting the assumption. □

Let us emphasize that being reducible is not the same as having only even zeros. To see this, suppose that  $\text{Div}(q) = 2\Lambda'$ . Then  $K \simeq \mathcal{O}(\Lambda') \otimes \mathcal{I}$ , where  $\mathcal{I}$  is a 2-torsion point in the Jacobian. The spectral curve  $S_q$  is reducible if and only if  $\mathcal{I}$  is trivial.

**2.3. Rank 1 torsion free sheaves and the BNR correspondence.** In this subsection, we provide some background on rank 1 torsion free sheaf theory over spectral curves in the context of the Hitchin and BNR correspondence developed in [23, 1].

Let  $S$  be a reduced and irreducible complex projective curve and  $\mathcal{O}_S$  its structure sheaf. The moduli space of invertible sheaves on  $S$  is denoted by  $\text{Pic}(S)$ , and  $\text{Pic}^d(S) \subset \text{Pic}(S)$  is the degree  $d$  component. If  $\mathcal{F}$  is a coherent analytic sheaf on  $S$ , we can define its cohomology groups  $H^i(X, \mathcal{F})$ . Since  $\dim S = 1$ ,  $H^i(X, \mathcal{F}) = 0$  for  $i \geq 2$ . The Euler characteristic is defined as  $\chi(\mathcal{F}) = \dim H^0(X, \mathcal{F}) - \dim H^1(X, \mathcal{F})$ . The degree of a torsion free sheaf  $\mathcal{F}$  is given by  $\deg(\mathcal{F}) = \chi(\mathcal{F}) - \text{rank}(\mathcal{F})\chi(\mathcal{O})$ . If  $\mathcal{F}$  is locally free, then  $\deg(\mathcal{F})$  coincides with the degree of the invertible sheaf  $\det(\mathcal{F})$ .

Let  $\overline{\text{Pic}}^d(S)$  be the moduli space of degree  $d$  rank 1 torsion free sheaves on  $S$ , and  $\overline{\text{Pic}}(S) = \prod_{d \in \mathbb{Z}} \overline{\text{Pic}}^d(S)$  [9]. Then  $\overline{\text{Pic}}^d(S)$  is an irreducible projective scheme containing  $\text{Pic}^d(S)$  as an open subscheme. When  $S$  is smooth, we have  $\overline{\text{Pic}}^d(S) = \text{Pic}^d(S)$ . The relationship to Higgs bundles is given by the following.

**Theorem 2.3.** [23, 1] *Let  $q \in H^0(K^2)$  be an irreducible quadratic differential with spectral curve  $S_q$ . There is a bijective correspondence between points in  $\overline{\text{Pic}}(S_q)$  and isomorphism classes of rank 2 Higgs pairs  $(\mathcal{E}, \varphi)$  with  $\text{Tr}(\varphi) = 0$  and  $\det(\varphi) = q$ . Explicitly: if  $\mathcal{L} \in \overline{\text{Pic}}(S_q)$ , then  $\mathcal{E} := \pi_*(\mathcal{L})$  is a rank 2 vector bundle and the homomorphism  $\pi_*\mathcal{L} \rightarrow \pi_*\mathcal{L} \otimes K \cong \pi_*(\mathcal{L} \otimes \pi^*K)$  given by the multiplication of the canonical section  $\lambda$  defines the Higgs field  $\varphi$ .*

This correspondence gives the very useful exact sequence

$$(1) \quad 0 \rightarrow \mathcal{L}(-\Delta) \rightarrow \pi^*\mathcal{E} \xrightarrow{\pi^*\varphi - \lambda} \pi^*\mathcal{E} \otimes \pi^*K \rightarrow \mathcal{L} \otimes \pi^*K \rightarrow 0,$$

where  $\mathcal{O}_S(\Delta) := K_S \otimes \pi^*K_\Sigma^{-1}$  is the ramification divisor. This will be used below.

Let  $q$  be a quadratic differential with only simple zeros, and define a divisor on  $S$  by  $\Lambda = \text{Div}(\lambda)$ . Then  $\Lambda$  is the ramification divisor of the map  $\pi : S \rightarrow \Sigma$ . By the Riemann-Hurwitz formula, the genus of  $S$  is  $g(S) = 4g - 3$ . Furthermore, for any  $\mathcal{L} \in \text{Pic}(S)$ , Riemann-Roch gives  $\deg(\pi_*\mathcal{L}) = \deg(\mathcal{L}) - (2g - 2)$ .

The  $\text{SL}(2, \mathbb{C})$  Higgs bundles are characterized by

$$(2) \quad \mathcal{T} := \{\mathcal{L} \in \text{Pic}^{2g-2}(S) \mid \det(\pi_*\mathcal{L}) = \mathcal{O}_\Sigma\}.$$

By the Hitchin-BNR correspondence (Theorem 2.3), the map  $\chi_{\text{BNR}} : \mathcal{T} \rightarrow \mathcal{M}_q$  is a bijection.

For the branched double cover  $\pi : S \rightarrow \Sigma$ , we have the norm map  $\text{Nm}_{S/\Sigma} : \text{Jac}(S) \rightarrow \text{Jac}(\Sigma)$ , where  $\text{Nm}_{S/\Sigma}(\mathcal{O}_S(D)) := \mathcal{O}_\Sigma(\pi(D))$ . The Prym variety is defined as

$$\text{Prym}(S/\Sigma) := \ker(\text{Nm}_{S/\Sigma}) = \{\mathcal{L} \in \text{Pic}(S) \mid \mathcal{L} \otimes \sigma^*\mathcal{L} = \mathcal{O}_S\},$$

where  $\sigma : S \rightarrow S$  is the involution. Also, we have  $\det(\pi_*\mathcal{L}) \cong \text{Nm}_{S/\Sigma}(\mathcal{L}) \otimes K^{-1}$ . Thus,  $\mathcal{T}$  can be expressed as

$$\mathcal{T} = \{\mathcal{L} \in \text{Pic}^{2g-2}(S) \mid \text{Nm}_{S/\Sigma}(\mathcal{L}) \cong K\}.$$

Hence,  $\mathcal{T}$  is a torsor over  $\text{Prym}(S, \Sigma)$ . Explicitly, by choosing  $\mathcal{L}_0 \in \mathcal{T}$ , we obtain an isomorphism  $\mathcal{T} \cong \text{Prym}(S, \Sigma)$  given by  $\mathcal{L} \rightarrow \mathcal{L} \otimes \mathcal{L}_0^{-1}$ .

Recall that we have defined  $\mathcal{M}_q := \mathcal{H}^{-1}(q)$ . To summarize, we have the following:

**Proposition 2.4.** *Let  $q$  be a quadratic differential with simple zeros. Then  $\mathcal{M}_q \cong \mathcal{T} \cong \text{Prym}(S, \Sigma)$ .*

If  $q \neq 0$  is irreducible but nongeneric, the spectral curve  $S$  is singular and irreducible. We may still define the set  $\overline{\mathcal{T}} \subset \overline{\text{Pic}}^{2g-2}(S)$  as follows:

$$\overline{\mathcal{T}} := \{\mathcal{L} \in \overline{\text{Pic}}^{2g-2}(S) \mid \det(\pi_*\mathcal{L}) \cong \mathcal{O}_\Sigma\},$$

We also set  $\mathcal{T} := \overline{\mathcal{T}} \cap \text{Pic}^{2g-2}$ . Then  $\overline{\mathcal{T}}$  is the natural compactification of  $\mathcal{T}$  induced by the inclusion  $\text{Pic}^{2g-2}(S) \subset \overline{\text{Pic}}(S)$ . The BNR correspondence, as stated in Theorem 2.3, implies that  $\chi_{\text{BNR}} : \overline{\mathcal{T}} \rightarrow \mathcal{M}_g$  is an isomorphism.

**2.4. The Hitchin moduli space and the nonabelian Hodge correspondence.** We now recall the well-known nonabelian Hodge theorem, which relates the space of flat  $\text{SL}(2, \mathbb{C})$  connections, Higgs bundles, and solutions to the Hitchin equations. This result was developed in the work of Hitchin [21], Simpson [42], Corlette [6], and Donaldson [8].

Let  $E$  be a trivial, smooth, rank 2 vector bundle over a Riemann surface  $\Sigma$ , and let  $H_0$  be a background Hermitian metric on  $E$ . We denote by  $\mathfrak{sl}(E)$  (resp.  $\mathfrak{su}(E)$ ) the bundle of traceless (resp. traceless skew-hermitian) endomorphisms of  $E$ . Let  $A$  be a unitary connection on  $E$  that induces the trivial connection on  $\det E$ , and let  $\phi \in \Omega^1(i\mathfrak{su}(E))$  be a Higgs field. The Hitchin equations for the pair  $(A, \phi)$  are given by:

$$(3) \quad F_A + \phi \wedge \phi = 0, \quad d_A \phi = d_A^* \phi = 0.$$

If we split the Higgs field into type:  $\phi = \varphi + \varphi^\dagger$ , with  $\varphi \in \Omega^{1,0}(\mathfrak{sl}(E))$ , then (3) is equivalent to:

$$(4) \quad F_A + [\varphi, \varphi^\dagger] = 0, \quad \bar{\partial}_A \varphi = 0.$$

Notice that  $(\bar{\partial}_A, \varphi)$  then defines an  $\text{SL}(2, \mathbb{C})$  Higgs bundle. The Hitchin moduli space, denoted by  $\mathcal{M}_{\text{Hit}}$ , is the moduli space of solutions to the Hitchin equation, given by

$$\mathcal{M}_{\text{Hit}} := \{(A, \phi) \mid (A, \phi) \text{ satisfies (3)}\} / \mathcal{G},$$

where  $\mathcal{G}$  is the gauge group of unitary automorphisms of  $E$ . Recall that a flat connection  $\mathcal{D}$  is called completely reducible if and only if it is a direct sum of irreducible flat connections. The Kobayashi-Hitchin correspondence can be summarized as follows:

**Theorem 2.5.** [21, 43, 6, 8] *A Higgs bundle  $(\mathcal{E}, \varphi)$  is polystable if and only if there exists a Hermitian metric  $H$  such that the corresponding Chern connection  $A$  and Higgs field  $\phi = \varphi + \varphi^\dagger$  solve the Hitchin equations (3). Moreover, the connection  $\mathcal{D}$  defined by  $\mathcal{D} = \nabla_A + \phi$  is a completely reducible flat connection, and it is irreducible if and only if  $(\mathcal{E}, \varphi)$  is stable.*

*Conversely, a flat connection  $\mathcal{D}$  is completely reducible if and only if there exists a Hermitian metric  $H$  on  $E$  such that when we express  $\mathcal{D} = \nabla_A + \varphi + \varphi^\dagger$ , we have  $\bar{\partial}_{\mathcal{E}} \varphi = 0$ . Moreover, the corresponding Higgs bundle  $(\mathcal{E}, \varphi)$  is polystable, and it is stable if and only if  $\mathcal{D}$  is irreducible.*

The nonabelian Hodge correspondence defines the following Kobayashi-Hitchin homeomorphism

$$\Xi : \mathcal{M}_{\text{Dol}} \rightarrow \mathcal{M}_{\text{Hit}},$$

which is a diffeomorphism to irreducible solutions of (3) when restricted to the stable locus.

Finally, we note that there is an action of  $S^1$  on  $\mathcal{M}_{\text{Hit}}$  defined by  $(A, \phi) \rightarrow (A, e^{i\theta} \cdot \phi)$ , where  $e^{i\theta} \cdot \phi = e^{i\theta} \varphi + e^{-i\theta} \varphi^\dagger$ . With respect to this and the  $S^1 \subset \mathbb{C}^*$  action on  $\mathcal{M}_{\text{Dol}}$ , the map  $\Xi$  is  $S^1$ -equivariant.

### 3. FILTERED BUNDLES AND COMPACTNESS

Filtered (or parabolic) bundles are described, for example, in [43]. They play a key role in the analytic compactification. This section provides a brief overview of filtered line bundles and demonstrates a compactness result.

**3.1. Filtered line bundles and nonabelian Hodge.** Let  $Z$  be a finite collection of distinct points on a closed Riemann surface  $\Sigma$ , and let  $\Sigma' = \Sigma \setminus Z$ . Viewing  $\Sigma$  as a projective algebraic curve, an algebraic line bundle  $L$  over  $\Sigma'$  is a line bundle defined by regular transition functions on Zariski open sets over  $\Sigma'$ . The sheaf of sections of  $L$  can be extended in infinitely many different ways over  $Z$  to obtain coherent (invertible) sheaves on  $\Sigma$ . The sections of  $L$  are then realized as meromorphic sections of an extension, regular on  $\Sigma'$ .

A *filtered line bundle*  $\mathcal{F}_*(L)$  is an algebraic line bundle  $L$  with a collection of coherent extensions  $L_\alpha$  across the punctures  $Z$  such that  $L_\alpha \subset L_\beta$  for  $\alpha \geq \beta$ , for a fixed sufficiently small  $\epsilon$   $L_{\alpha-\epsilon} = L_\alpha$ , and  $L_\alpha = L_{\alpha+1} \otimes \mathcal{O}_\Sigma(Z)$ . Let  $\text{Gr}_\alpha = L_{\alpha+\epsilon}/L_\alpha$  denote the quotient (torsion) sheaf. A value  $\alpha$  where  $\text{Gr}_\alpha \neq 0$  is called a jump. Since we are considering line bundles, there is exactly one jump  $\alpha_p \in [0, 1)$  for each  $p$  (i.e.  $p$  is in the support of  $\text{Gr}_{\alpha_p}$ ). The collection of jumps  $\{\alpha_p\}$  fully determines the filtered bundle structure. If we denote by  $\mathcal{L} := L_0$ , the degree of a filtered line bundle is defined as

$$\deg(\mathcal{F}_*(L)) = \deg(\mathcal{L}) + \sum_{p \in Z} \alpha_p .$$

Alternatively, a *weighted line bundle* is a pair  $(\mathcal{L}, \chi)$  where  $\mathcal{L} \rightarrow \Sigma$  is a holomorphic line bundle and  $\chi : Z \rightarrow \mathbb{R}$  is a weight function. The degree of a weighted bundle is defined as

$$\deg(\mathcal{L}, \chi) = \deg(\mathcal{L}) + \sum_{p \in Z} \chi_p .$$

The concepts of filtered line bundles and weighted line bundles are nearly equivalent. Namely, given a filtered line bundle  $\mathcal{F}_*(L)$ , we define  $\mathcal{L} := \mathcal{L}_0$  and  $\chi_p = \alpha_p$ . Conversely, given a weighted line bundle  $(\mathcal{L}, \chi)$ , let  $\alpha_p = \chi_p + n_p$ , where  $n_p \in \mathbb{Z}$  is the unique integer so that  $0 \leq \chi_p + n_p < 1$ . A filtered bundle  $\mathcal{F}_*(L)$  is then determined by setting  $L_0 = \mathcal{L}(-\sum_{p \in Z} n(\alpha_p)p)$  with jumps  $\alpha_p$ . Clearly,  $\deg(\mathcal{F}_*(L)) = \deg(\mathcal{L}, \chi)$ . We shall use the notation  $\mathcal{F}_*(\mathcal{L}, \chi)$  for the filtered bundle associated to a weighted bundle  $(\mathcal{L}, \chi)$  in this way.

Different weighted bundles can give rise to the same filtered bundle. The following is a fact that will be frequently used in this paper. If  $D = \sum_{x \in Z} d_x x$  is a divisor supported on  $Z$ , let

$$\chi_D(x) := \begin{cases} d_x , & x \in Z ; \\ 0 , & x \in \Sigma \setminus Z . \end{cases}$$

Then for any weighted bundle  $(\mathcal{L}, \chi)$  we have  $\mathcal{F}_*(\mathcal{L}(D), \chi - \chi_D) = \mathcal{F}_*(\mathcal{L}, \chi)$ .

Let  $(\mathcal{L}_1, \chi_1)$  and  $(\mathcal{L}_2, \chi_2)$  be two weighted line bundles. We define the tensor product

$$(\mathcal{L}_1, \chi_1) \otimes (\mathcal{L}_2, \chi_2) := (\mathcal{L}_1 \otimes \mathcal{L}_2, \chi_1 + \chi_2) .$$

Then the degree is additive on tensor products. For filtered bundles, we *define*

$$(5) \quad \mathcal{F}_*(\mathcal{L}_1, \chi_1) \otimes \mathcal{F}_*(\mathcal{L}_2, \chi_2) := \mathcal{F}_*(\mathcal{L}_1 \otimes \mathcal{L}_2, \chi_1 + \chi_2) .$$

The degree is again additive for the tensor product of filtered bundles. This agrees with the usual definition of tensor product for parabolic bundles.

### 3.2. Harmonic metrics for weighted line bundles.

**Proposition 3.1.** *Let  $(\mathcal{L}, \chi)$  be a degree 0 weighted bundle. Then there exists a Hermitian metric  $h$  such that:*

- (i) *the Chern connection  $A_h$  of  $(\mathcal{L}, h)$  is flat:  $F_{A_h} = 0$ ;*
- (ii) *for  $p \in Z$ , and  $(U_p, z)$  a holomorphic coordinate centered at  $p$ ,  $|z|^{-2\chi_p} h$  is a  $C^\infty$  Hermitian metric on  $\mathcal{L}|_{U_p}$ ;*
- (iii)  *$h$  is uniquely determined up to a multiplication by a nonzero constant.*

*Proof.* We first choose a background Hermitian metric  $h_0$  such that  $|z|^{-2\chi_p} h_0$  defines a  $\mathcal{C}^\infty$  Hermitian metric defined on  $U_p$ . Let  $A_{h_0}$  be the Chern connection and  $F_{A_0}$  be the curvature. We have  $\frac{\sqrt{-1}}{2\pi} \int_\Sigma F_{A_0} = \deg(\mathcal{L}, \chi) = 0$ . Therefore, there exists a  $\mathcal{C}^\infty$  function  $\rho$  such that  $\Delta\rho + \frac{\sqrt{-1}}{2\pi} \Lambda F_{A_0} = 0$ . We define  $h = h_0 e^\rho$ . For the corresponding Chern connection  $A_h$ , we have  $F_{A_h} = 0$ , which implies (i). (ii) follows from the property of  $h_0$  and  $\rho$  being a smooth function on  $\Sigma$ . As  $\rho$  is well-defined up to a constant,  $h$  is well-defined up to a constant, which implies (iii).  $\square$

The metric obtained above is called the *harmonic metric*. For a weighted bundle  $(\mathcal{L}, \chi)$ , the holomorphic bundle  $\mathcal{L}$  and the harmonic metric  $h$  define a filtration as follows. For  $\epsilon > 0$  sufficiently small, let

$$L_\alpha := \{s \in \mathcal{L}(*Z) \mid |s|_h \leq Cr^{\alpha-\epsilon} \text{ for some } C\}.$$

Here,  $r$  denotes the distance to  $Z$  in any smooth conformal metric on  $\Sigma$ . It is straightforward to check that this defines a filtered bundle that matches  $\mathcal{F}_*(\mathcal{L}, \chi)$  under the correspondence in the previous section.

Even though the harmonic metric is only well-defined up to a constant, the Chern connection  $A = (\mathcal{L}, h)$  is independent of this choice. The  $(1, 0)$  part of  $A$ , denoted  $\nabla_h$ , then defines logarithmic connections  $\nabla_h : L_\alpha \rightarrow L_\alpha \otimes K(Z)$ .

**3.3. Convergence of weighted line bundles.** In this subsection, we will consider the convergence of weighted line bundles and the main result we prove here might be a direct corollary of [34, Theorem 1.8]. For the reader's convenience, we present a short proof here for our situation.

Let  $(\Sigma_0, g_0)$  be a metrized Riemann surface (i.e. a Riemann surface  $\Sigma_0$  with conformal metric  $g_0$ ) with a holomorphic structure  $J_0$ . Consider a neighborhood  $U_1$  of  $J_0$  in the moduli space of holomorphic structures and a neighborhood  $U_2$  of  $g_0$  in the space of smooth metrics. We denote the product of these neighborhoods by  $U = U_1 \times U_2$ . We can define the fiber bundle  $\text{Pic}_U \rightarrow U$ , where each fiber is the Picard group defined by the holomorphic structure. Let  $(\Sigma_t = (C, J_t), g_t)$  be a family of metrized Riemann surfaces that converge smoothly to  $(\Sigma_0, g_0)$  as  $t \rightarrow 0$ . Let  $Z_t \subset \Sigma_t$  be a collection of a finite number of points that converge to  $Z_0$  in suitable symmetric products of  $C$ . For each  $p \in Z_0$ , we can write  $Z_t = \cup_{p \in Z_0} Z_{t,p}$  such that all points in  $Z_{t,p}$  converge to  $p$ . We define the convergence of weighted line bundles as follows:

**Definition 3.2.** *A family of weighted line bundles  $(\mathcal{L}_t, \chi_t)$  over  $\Sigma_t$  with weights  $\chi_t : Z_t \rightarrow \mathbb{R}$  converges to  $(\mathcal{L}_0, \chi_0)$  if*

- (i)  $\mathcal{L}_t$  converges to  $\mathcal{L}_0$  in  $\text{Pic}_U$ ,
- (ii) for all  $p \in Z_0$  and  $t$  sufficiently small,  $\sum_{q \in Z_{t,p}} \chi_t(q) = \chi_0(p)$ .

A sequence of filtered bundles  $\mathcal{F}_*(\mathcal{L}_t)$  converges to  $\mathcal{F}_*(\mathcal{L}_0)$  if the corresponding weighted bundles converge. The following theorem provides insight into the compactness of a sequence of weighted line bundles:

**Theorem 3.3.** *Let  $(\mathcal{L}_t, \chi_t)$  be a sequence of weighted line bundles over  $\Sigma_t \setminus Z_t$ , where  $\deg(\mathcal{L}_t, \chi_t) = 0$ , and let  $h_t$  be the corresponding harmonic metrics. If  $Z_t$  converges to  $Z_0$ , we write  $Z_t = \cup_{p \in Z_0} Z_{t,p}$ . Then there exists a weighted line bundle  $(\mathcal{L}_0, \chi_0)$  over  $Z_0$  with a harmonic metric  $h_0$  such that:*

- (i) After being rescaled by  $c_t > 0$ ,  $c_t h_t$  converges to  $h_0$  over  $\Sigma_0 \setminus Z_0$  in the  $\mathcal{C}_{\text{loc}}^\infty$  sense.
- (ii) Let  $\nabla_t$  be the unitary connection of  $h_t$ . Then on  $\Sigma_0 \setminus Z_0$ ,  $\lim_{t \rightarrow 0} \nabla_t = \nabla_0$  in  $\mathcal{C}_{\text{loc}}^\infty$ .



*Proof.* By the assumptions on weights,  $\deg(\mathcal{L}_t)$  is a fixed,  $t$ -independent constant. Let  $\gamma_t = (J_t, g_t)$  be a path in  $U$ . Then  $\text{Pic}_U|_{\gamma_t}$  is compact, and there exists an  $\mathcal{L}_0 \in \text{Pic}(\Sigma_0)$  such that  $\mathcal{L}_t$  converges to  $\mathcal{L}_0$ . For  $p \in Z_0$ , define  $\chi_0(p) = \sum_{q \in Z_{t,p}} \chi_t(q)$ , and thus obtain a weighted line bundle  $(\mathcal{L}_0, \chi_0)$ . We can choose a family of approximate harmonic metrics  $h_t^{\text{app}}$ , such that  $|z|^{-2\chi_p} h_t^{\text{app}}$  extends to a smooth metric in a neighborhood of  $p$  and  $h_t^{\text{app}}$  converges to  $h_0^{\text{app}}$  in  $\mathcal{C}_{\text{loc}}^\infty(\Sigma_0 \setminus Z_0)$ . Moreover, we write  $h_t = h_t^{\text{app}} e^{s_t}$ . After a suitable rescale of  $h_t$ , we can assume  $\|s_t\|_{L^2} = 1$ . Let  $\rho_t := \Delta_t h_t^{\text{app}}$  be the curvature defined by the metric  $h_t^{\text{app}}$ . Then  $s_t$  satisfies the equation  $\Delta_t s_t = \rho_t$  over  $\Sigma$ . As  $\rho_t$  converges to  $\rho_0 \in \mathcal{C}_{\text{loc}}^\infty(\Sigma \setminus Z_0)$ , and  $g_t$  is a family with bounded geometry, we obtain the estimate

$$\|s_t\|_{\mathcal{C}^{k+2,\alpha}(\Sigma)} \leq C_{k,\alpha} (\|\rho_t\|_{\mathcal{C}^{k,\alpha}(\Sigma)} + 1) ,$$

where  $C_{k,\alpha}$  is a  $t$ -independent constant. Therefore, passing to a subsequence,  $s_t$  converges to  $s_0$  in  $\mathcal{C}^\infty(\Sigma)$ , which implies (i). The assertion (ii) follows from (i).  $\square$

#### 4. THE ALGEBRAIC AND ANALYTIC COMPACTIFICATIONS

In this section, we introduce compactifications of the Dolbeault and Hitchin moduli spaces.

**4.1. The algebraic compactification of the Dolbeault moduli space.** In this subsection, we present the algebraic method for compactifying the Dolbeault moduli space. This technique is based on the  $\mathbb{C}^*$  action on  $\mathcal{M}_{\text{Dol}}$ , and was introduced in [41, 39, 19, 7, 27]. The gauge theoretic approach can be found in [11].

**Theorem 4.1** ([41, Thm. 11.2],[7]). *Let  $V$  be an algebraic variety with  $\mathbb{C}^*$  action. Suppose*

- (i) *the fixed point set of the  $\mathbb{C}^*$  action is proper,*
- (ii) *for every  $t \in \mathbb{C}^*$ ,  $v \in V$ , the limit  $\lim_{t \rightarrow 0} t \cdot v$  exists.*

*Then the space  $U := \{v \in V \mid \lim_{t \rightarrow \infty} t \cdot v \text{ does not exist}\}$  is open in  $V$ , and the quotient  $U/\mathbb{C}^*$  is separated and proper.*

We apply this to the Dolbeault moduli space. The first step is to study the possible isotropy subgroups.

**Lemma 4.2.** [19, Thm. 6.2] *Let  $\xi = [(\mathcal{E}, \varphi)]$  be a Higgs bundle equivalence class with  $\mathcal{H}(\xi) \neq 0$ . Then the stabilizer  $\Gamma_\xi$  of  $\xi$  for the  $\mathbb{C}^*$  action is either trivial or  $\mathbb{Z}/2$ . The latter case holds if and only if  $(\mathcal{E}, \varphi)$  and  $(\mathcal{E}, -\varphi)$  are complex gauge equivalent.*

*Proof.* For  $t \in \Gamma_\xi$ ,  $\mathcal{H}(t \cdot \xi) = t^2 \mathcal{H}(\xi)$ , hence  $t^2 = 1$  if  $\mathcal{H}(\xi) \neq 0$ .  $\square$

The fixed points of the  $\mathbb{Z}/2$  action correspond to real representations under the nonabelian Hodge correspondence [21, Sec. 10]. Hence by this Lemma, the space  $(\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))/\mathbb{C}^*$  has an orbifold structure.

By the properness of the Hitchin map  $\mathcal{H}$ , see Theorem 2.1, it follows that  $\lim_{t \rightarrow \infty} t \cdot \xi$  exists if and only if  $\mathcal{H}(\xi) = 0$ . Now define

$$(6) \quad \overline{\mathcal{M}}_{\text{Dol}} = \left\{ (\mathcal{M}_{\text{Dol}} \times \mathbb{C}^*) \coprod (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0)) \right\} / \mathbb{C}^* .$$

The analytic topology on the disjoint union is generated by open sets  $U \times W_1$  and  $V \times (W_2 \cap \mathbb{C}^*) \amalg V \cap (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))$ , where  $U, V \subset \mathcal{M}_{\text{Dol}}$ ,  $W_1, W_2 \subset \mathbb{C}$  are open, with  $0 \notin W_1$ ,  $0 \in W_2$ . The topology on  $\overline{\mathcal{M}}_{\text{Dol}}$  is then the quotient topology, and it is straightforward to see that with this topology, it is compact.

Since  $(\mathcal{M}_{\text{Dol}} \times \mathbb{C}^*)/\mathbb{C}^* = \mathcal{M}_{\text{Dol}}$ , there is a natural inclusion

$$\iota : \mathcal{M}_{\text{Dol}} \rightarrow \overline{\mathcal{M}}_{\text{Dol}}, \quad \iota(\xi) = [(\xi, 1)] ,$$

where brackets denote the equivalent class under the  $\mathbb{C}^*$  action. The boundary of  $\overline{\mathcal{M}}_{\text{Dol}}$  is

$$\partial\overline{\mathcal{M}}_{\text{Dol}} = \overline{\mathcal{M}}_{\text{Dol}} \setminus \iota(\mathcal{M}_{\text{Dol}}) = (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))/\mathbb{C}^* .$$

There is a *boundary map*

$$\iota_{\partial} : \mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0) \longrightarrow \partial\overline{\mathcal{M}}_{\text{Dol}}, \quad \xi \mapsto [(\xi, 0)] ,$$

which is invariant under the  $\mathbb{C}^*$  action, i.e.,  $\iota_{\partial}(\lambda\xi) = \iota_{\partial}(\xi)$  for  $\lambda \in \mathbb{C}^*$ .

The  $\mathbb{C}^*$  action on  $\mathcal{M}_{\text{Dol}}$  covers the (square of the) action on  $\mathcal{B}$ . Hence, it is natural to compactify  $\mathcal{B}$  by projectivizing:

$$\overline{\mathcal{B}} := \mathbb{P}(H^0(K^2) \oplus \mathbb{C}) .$$

The inclusion is given, as usual, by

$$\iota_0 : \mathcal{B} \rightarrow \overline{\mathcal{B}}, \quad \iota_0(q) = [q \times \{1\}] ,$$

where  $q \times \{1\} \in H^0(K^2) \oplus \mathbb{C}$ . We also define  $\partial\overline{\mathcal{B}} = \overline{\mathcal{B}} \setminus \iota_0(\mathcal{B}) \simeq \mathbb{P}(H^0(K^2))$ , with boundary inclusion map

$$\iota_{0,\partial} : \mathcal{B} \setminus \{0\} \rightarrow \partial\overline{\mathcal{B}}, \quad \iota_{0,\partial}(q) = [q \times \{0\}] .$$

The Hitchin map  $\mathcal{H} : \mathcal{M}_{\text{Dol}} \rightarrow \mathcal{B}$  extends to  $\overline{\mathcal{H}} : \overline{\mathcal{M}}_{\text{Dol}} \rightarrow \overline{\mathcal{B}}$ , where  $\overline{\mathcal{H}}|_{\mathcal{M}_{\text{Dol}}} := \iota_0 \circ \mathcal{H}$  and for every  $[(\mathcal{E}, \varphi)]/\mathbb{C}^* \in \partial\overline{\mathcal{M}}_{\text{Dol}}$ ,

$$\overline{\mathcal{H}}([( \mathcal{E}, \varphi )]/\mathbb{C}^*) := [(0, \mathcal{H}(\varphi))] \subset \overline{\mathcal{B}} .$$

This is well defined since  $\det(\varphi) \neq 0$  if  $[(\mathcal{E}, \varphi)]/\mathbb{C}^* \in \partial\overline{\mathcal{M}}_{\text{Dol}}$ , and

$$\begin{array}{ccc} \mathcal{M}_{\text{Dol}} & \xrightarrow{\iota} & \overline{\mathcal{M}}_{\text{Dol}} \\ \downarrow \mathcal{H} & & \downarrow \overline{\mathcal{H}} \\ \mathcal{B} & \xrightarrow{\iota_0} & \overline{\mathcal{B}} \end{array}$$

commutes.

There is a good algebraic structure on this algebraic compactification:

**Theorem 4.3.** [41, 39, 19, 7, 11] *The compactified space  $\overline{\mathcal{M}}_{\text{Dol}}$  is a normal projective variety, and  $\partial\overline{\mathcal{M}}_{\text{Dol}}$  is a Cartier divisor of  $\overline{\mathcal{M}}_{\text{Dol}}$ .*

The following characterization of sequential convergence is useful.

**Proposition 4.4.** *Let  $[(\mathcal{E}_i, \varphi_i)] \in \mathcal{M}_{\text{Dol}}$  be a sequence of Higgs bundles, and write  $q_i = \det(\varphi_i)$  and  $r_i = \|q_i\|_{L^2}^{\frac{1}{2}}$ . Suppose  $\limsup r_i \rightarrow \infty$ ; then up to subsequence:*

- (i) *there exists a Higgs bundle  $[(\widehat{\mathcal{E}}_{\infty}, \widehat{\varphi}_{\infty})]$  with  $\tilde{q}_{\infty} = \det(\widehat{\varphi}_{\infty})$  and  $\|\widehat{q}_{\infty}\|_{L^2} = 1$  such that  $\lim_{i \rightarrow \infty} [(\mathcal{E}_i, r_i^{-1}\varphi_i)] = [(\widehat{\mathcal{E}}_{\infty}, \widehat{\varphi}_{\infty})]$  in  $\mathcal{M}_{\text{Dol}}$  and  $\lim_{i \rightarrow \infty} r_i^{-1}q_i = \widehat{q}_{\infty}$  in  $H^0(K^2)$ ;*
- (ii)

$$\begin{aligned} \lim_{i \rightarrow \infty} \iota[(\mathcal{E}_i, \varphi_i)] &= \iota_{\partial}[(\widehat{\mathcal{E}}_{\infty}, \widehat{\varphi}_{\infty})] && \text{on } \overline{\mathcal{M}}_{\text{Dol}}, \\ \lim_{i \rightarrow \infty} \iota_0(q_i) &= \iota_{0,\partial}(\widehat{q}_{\infty}) && \text{on } \overline{\mathcal{B}}. \end{aligned}$$

*Proof.* The first point follows since the Hitchin map  $\mathcal{H}$  is proper and  $\mathcal{H}(r_i^{-1}\varphi_i)$  is bounded. The second follows directly from the definition.  $\square$

**4.2. The analytic compactification of the Hitchin moduli space.** We next describe the compactification of the Hitchin moduli space developed in [29, 33, 46].

4.2.1. *Decoupled Hitchin equations.* We begin by defining the decoupled Hitchin equations. Recall the notation from Section 2.4. Let  $Z$  be a finite set of distinct points in  $\Sigma$ . For a smooth unitary connection  $A$  on  $E|_{\Sigma \setminus Z}$  and smooth  $\phi = \varphi + \varphi^\dagger \in \Omega^1(\text{isu}(E))|_{\Sigma \setminus Z}$ , the *decoupled Hitchin equations* on  $\Sigma \setminus Z$  are:

$$(7) \quad F_A = 0, \quad [\varphi, \varphi^\dagger] = 0, \quad \bar{\partial}_A \varphi = 0.$$

Solutions to (7) may be quite singular near  $Z$ , so we make the following restriction:

**Definition 4.5.** *A solution  $(A, \phi)$  to the decoupled Hitchin equations over  $\Sigma \setminus Z$  is called admissible if  $\phi \neq 0$ , and  $|\phi|$  extends to a continuous function on  $\Sigma$  with  $|\phi|^{-1}(0) = Z$ .*

By a *limiting configuration* we always mean an admissible solution to the decoupled Hitchin equations. Clearly  $Z$  is determined by  $(A, \phi)$ . Admissibility guarantees that  $\det(\varphi)$  extends to a holomorphic quadratic differential  $q = \det(\varphi)$  on  $\Sigma$  with  $Z = q^{-1}(0)$  as a set. Hence, the spectral curve  $S_q$  is well-defined. We emphasize that  $Z$  may vary for different admissible solutions, but one always has that  $\#Z \leq 4g - 4$ .

The equivalence relation on limiting configurations is that  $(A_1, \phi_1) \sim (A_2, \phi_2)$  if  $Z_1 = Z_2$  and  $(A_1, \phi_1)g = (A_2, \phi_2)$  for a smooth unitary gauge transformation  $g$  on  $\Sigma \setminus Z_1$ . The moduli space of the decoupled Hitchin equations is then

$$\mathcal{M}_{\text{Hit}}^{\text{Lim}} = \{\text{admissible solutions to (7)}\} / \sim.$$

We denote by  $\mathcal{M}_{\text{Hit}, q}^{\text{Lim}}$  the elements in  $\mathcal{M}_{\text{Hit}}^{\text{Lim}}$  with a fixed quadratic differential  $q$ , so the equivalence relation is now induced by the action of the unitary gauge group over  $\Sigma \setminus Z$ ,  $Z = q^{-1}(0)$ .

There is a natural  $\mathbb{C}^*$  action on the moduli space  $\mathcal{M}_{\text{Hit}}^{\text{Lim}}$ : given  $(A, \phi = \varphi + \varphi^\dagger) \in \mathcal{M}_{\text{Hit}}^{\text{Lim}}$  and  $t \in \mathbb{C}^*$ , we set  $t \cdot [(A, \phi)] = [(A, t\varphi + \bar{t}\varphi^\dagger)]$ , which is also a solution to (7).

4.2.2. *Compactification of the Hitchin moduli space.* The following compactness result is due to Taubes [47] and Mochizuki [33] (see also [20]).

**Proposition 4.6.** *Let  $(A_i, \varphi_i)$  be a sequence of solutions to (3), with  $q_i = \det(\varphi_i) \in H^0(K^2)$ . Then*

- (i) *if  $\limsup \|q_i\|_{L^2(\Sigma)} < \infty$ , then there is a subsequence (also denoted  $\{i\}$ ), a smooth solution  $(A_\infty, \phi_\infty)$  to (3), and a sequence  $g_i$  of smooth unitary gauge transformations on  $\Sigma$ , such that  $(A_i, \varphi_i)g_i$  converges smoothly to  $(A_\infty, \phi_\infty)$  on  $\Sigma$ ;*
- (ii) *if  $\lim \|q_i\|_{L^2(\Sigma)} = \infty$ , then there is a subsequence (also denoted  $\{i\}$ ), and  $q_\infty \in H^0(K^2)$  so that*

$$\frac{q_i}{\|q_i\|_{L^2}} \longrightarrow q_\infty$$

*over  $\Sigma$ , and an admissible solution  $(A_\infty, \phi_\infty = \varphi_\infty + \varphi_\infty^\dagger)$  to (7), with  $Z_\infty := q_\infty^{-1}(0)$ , and smooth unitary gauge transformations  $g_i$  on  $\Sigma \setminus Z_\infty$ , such that over any open set  $\Omega \Subset \Sigma \setminus Z_\infty$ ,  $(A_i)g_i \rightarrow A_\infty$ , and*

$$\frac{g_i^{-1} \phi_i g_i}{\|\phi_i\|_{L^2}} \longrightarrow \phi_\infty$$

*smoothly on  $\Omega$ .*

The norm on  $H^0(\Sigma, K^2)$  can be chosen arbitrarily since it is a finite dimensional space.

There is also a compactness result for sequences of solutions in  $\mathcal{M}_{\text{Hit}}^{\text{Lim}}$ .

**Proposition 4.7.** *Let  $[(A_i, \phi_i = \varphi_i + \varphi_i^\dagger)] \in \mathcal{M}_{\text{Hit}}^{\text{Lim}}$  be a sequence of admissible solutions to (7), and let  $q_i = \det(\varphi_i)$  be the corresponding quadratic differentials. Then after passing to a subsequence, there are  $t_i \in \mathbb{C}^*$ , a limiting configuration  $(A_\infty, \phi_\infty = \varphi_\infty + \varphi_\infty^\dagger)$  with*

quadratic differential  $q_\infty = \det(\varphi_\infty) \neq 0$ , and a sequence  $g_i$  of smooth gauge transformations on  $\Sigma \setminus Z_\infty$ , such that:

- (i)  $t_i^2 q_i$  converges smoothly to  $q_\infty$ ,
- (ii) over any open set  $\Omega \Subset X \setminus Z_\infty$ ,  $(A_i, t_i \cdot \phi_i) g_i$  converges smoothly to  $(A_\infty, \phi_\infty)$ .

*Proof.* Write  $q_i = \det(\varphi_i) \in H^0(K^2)$ . Adjusting by  $t_i$  if necessary, we may assume  $q_i$  converges to  $q_\infty$  over  $\Sigma$ . Also, since  $F_{A_i} = 0$  over  $\Sigma \setminus Z_i$  and  $Z_i$  converges to  $Z_\infty$ , we can apply both Uhlenbeck compactness and the classical bootstrapping method to obtain  $A_\infty$  such that up to gauge  $A_i$  converges smoothly to  $A_\infty$  over  $\Sigma \setminus Z_\infty$ . Finally, the convergence of  $\varphi_i$  follows by the bound on  $q_i$ 's.  $\square$

4.2.3. *The topology on the compactified space.* We now carefully define the topology on the space  $\mathcal{M}_{\text{Hit}} \cup \mathcal{M}_{\text{Hit}}^{\text{Lim}}/\mathbb{C}^*$ . Choose a metric in the conformal class on  $\Sigma$ . Let  $W^{k,2}$  denote the Sobolev spaces on  $\Sigma$  of distributional sections with at least  $k$  derivatives in  $L^2$ . For a finite set of points  $Z \subset \Sigma$  (or indeed any closed subset),

$$W_{\text{loc}}^{k,2}(\Sigma \setminus Z) := \{f \mid f \in W^{k,2}(K), K \subset \Sigma \setminus Z, K \text{ compact}\}.$$

These definitions extend easily to the space of connections and  $\Omega^1(\text{isu}(E))$  for a Hermitian vector bundle  $(E, H_0)$  over  $\Sigma$  with a fixed smooth background connection.

Let  $\omega_n$  be a nested collection of open sets with  $\omega_n \subset \overline{\omega_n} \subset \omega_{n+1}$ , with  $\bigcup_n \omega_n = \Sigma \setminus Z$ . We then define the seminorms  $\|f\|_n := \|f\|_{W^{k,2}(\omega_n)}$ ; in terms of these,  $W_{\text{loc}}^{k,2}(\Sigma \setminus Z)$  a Fréchet space.

For any  $q \in H^0(K^2)$ , set  $Z_q := q^{-1}(0)$ , and consider the moduli space

$$\mathbb{M}_q = \{(A, \phi) \in \mathcal{M}_q \cap W^{k,2}(\Sigma)\} \cup \{(A, \phi) \in \mathcal{M}_{\text{Hit},q}^{\text{Lim}} \cap W_{\text{loc}}^{k,2}(\Sigma \setminus Z_q)\}/\mathbb{C}^*.$$

By classical bootstrapping of the gauge-theoretic elliptic equations,  $\mathbb{M}_q$  is independent of  $k \geq 2$ .

Next define  $\mathbb{M} := \bigcup_{q \in H^0(K^2)} \mathbb{M}_q$ . Its topology is generated by two types of open sets. For interior points  $\xi = [(A, \phi)] \in \mathcal{M}_{\text{Hit}} \subset \mathbb{M}$  we use the open sets

$$V_{\xi,\epsilon} := \{[(A', \phi')] \in \mathcal{M}_{\text{Hit}} \mid \|A' - A\|_{W^{k,2}(\Sigma)} + \|\phi' - \phi\|_{W^{k,2}(\Sigma)} < \epsilon\}$$

from the topology of  $\mathcal{M}_{\text{Hit}}$ . For any boundary point  $\xi_0 \in \mathcal{M}_{\text{Hit}}^{\text{Lim}}/\mathbb{C}^*$ , choose a representative  $(A_0, e^{i\theta} \phi_0)$  for some constant  $\theta$ . Let  $q = \det(\phi_0)$ , and fix any open set  $\omega \Subset \Sigma \setminus Z_q$ . Then, setting  $\mathcal{M}_{\text{Hit}}^* = \mathcal{M}_{\text{Hit}} \setminus \mathcal{H}^{-1}(0)$ ,

$$U_{\xi_0,\omega,\epsilon} := \{(A, \phi) \in \mathcal{M}_{\text{Hit}}^* \mid \|A - A_0\|_{W^{k,2}(\omega)} + \inf_{\theta \in S^1} \|\phi\|_{L^2}^{-\frac{1}{2}} \phi - e^{i\theta} \phi_0\|_{W^{k,2}(\omega)} < \epsilon, \|\phi\|_{L^2} > \epsilon\}$$

$$\bigcup \{(A, \phi) \in \mathcal{M}_{\text{Hit}}^{\text{Lim}} \mid \|A - A_0\|_{W^{k,2}(\omega)} + \|\phi - \phi_0\|_{W^{k,2}(\omega)} < \epsilon\}$$

defines an open set around  $\xi_0$ . The sets  $U_{\xi_0,\omega,\epsilon}$  and  $V_{\xi,\epsilon}$  generate the topology on  $\mathbb{M}$ .

**Theorem 4.8.** *The space  $\mathbb{M}$  is a Hausdorff and compact.*

*Proof.* The Hausdorff property follows from the definition of the topology. By Propositions 4.6 and 4.7,  $\mathbb{M}$  is sequentially compact. Moreover, using this explicit base for the topology  $\mathbb{M}$  is first countable, and hence compact.  $\square$

We may now define the compactification of the Hitchin moduli space as the closure  $\overline{\mathcal{M}}_{\text{Hit}} \subset \mathcal{M}_{\text{Hit}} \cup \mathcal{M}_{\text{Hit}}^{\text{Lim}}/\mathbb{C}^*$ ; we write  $\partial \overline{\mathcal{M}}_{\text{Hit}}$  for the boundary of the closure and  $\overline{\mathcal{M}}_{\text{Hit},q} := \overline{\mathcal{M}}_{\text{Hit}} \cap \mathbb{M}_q$  for the subset of elements with a fixed quadratic differential.

The following result is described in [30, 36, 31].

**Theorem 4.9.** [31, Prop. 3.3] *If  $q$  has only simple zeros, then  $\overline{\mathcal{M}}_{\text{Hit},q} = \mathbb{M}_q$ .*

In other words, the compactification of any slice where  $q$  does not lie in the discriminant locus is ‘the obvious one’.

## 5. PARABOLIC MODULES AND STRATIFICATION OF BNR DATA

In this section, we review the notion of a parabolic module, as described in [38, 5, 4, 16]. This concept leads to a partial normalization of the generalized Jacobian and Prym varieties of the spectral curve.

**5.1. Normalization of the spectral curve.** Let  $q \neq 0$  be a quadratic differential with an irreducible, singular spectral curve  $S = S_q$ . The zeros of  $q$  define a divisor  $\text{Div}(q) = \sum_{i=1}^{r_1} m_i p_i + \sum_{j=1}^{r_2} n_j p'_j$ , where the  $m_i$  and  $n_j$  are even and odd integers, respectively, and hence  $r_1$  and  $r_2$  are the numbers of even and odd zeros, respectively, counted without multiplicity. Write  $Z_{\text{even}} = \{p_1, \dots, p_{r_1}\}$ ,  $Z_{\text{odd}} = \{p'_1, \dots, p'_{r_2}\}$ , and  $Z = Z_{\text{even}} \cup Z_{\text{odd}}$ , so  $\#Z = r = r_1 + r_2$ .

The map  $\pi : S \rightarrow \Sigma$  is a double covering branched along  $Z$ ; hence, we may view  $p_i$  and  $p'_i$  as points in  $S$ . For  $x \in S$ , let  $\mathcal{O}_x$  be the local ring,  $\mathcal{O}_x^*$  its group of units, and  $R_x$  the completion. We say that  $S$  has an  $A_n$  singularity at  $x$  if  $R_x \cong \mathbb{C}[[r, s]]/(r^2 - s^{n+1})$ , where  $n \geq 1$ . If  $S$  has an  $A_1$  singularity at  $x$ , we call it a *nodal* singularity, and if  $S$  has an  $A_2$  singularity at  $x$ , we call it a *cusp* singularity.

Let  $p : \tilde{S} \rightarrow S$  be the normalization of  $S$ , and let  $\tilde{\pi} := \pi \circ p$ :

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{p} & S \\ & \searrow \tilde{\pi} & \downarrow \pi \\ & & \Sigma \end{array}$$

For even zeros  $p_i$  we write  $p^{-1}(p_i) = \{\tilde{p}_i^+, \tilde{p}_i^-\}$ , and for odd zeros  $p'_i$  we write  $p^{-1}(p'_i) = \tilde{p}'_i$ . Since  $\pi : S \rightarrow \Sigma$  is a branched double cover, the involution  $\sigma$  on  $S$  extends to an involution of  $\tilde{S}$  which we also denote by  $\sigma$ . Note that  $\sigma(\tilde{p}'_i) = \tilde{p}'_i$  while  $\sigma(\tilde{p}_i^\pm) = \tilde{p}_i^\mp$ .

The ramification divisor  $\Lambda' = \frac{1}{2} \sum_{i=1}^{r_1} m_i p_i + \sum_{j=1}^{r_2} (n_j - 1) p'_j$ , is a divisor on  $S$ , and there is an exact sequence:

$$(8) \quad 0 \longrightarrow \mathcal{O}_S \longrightarrow p_* \mathcal{O}_{\tilde{S}} \longrightarrow \mathcal{O}_{\Lambda'} \longrightarrow 0.$$

The genus of  $\tilde{S}$  is  $g(\tilde{S}) = 4g - 3 - \deg(\Lambda') = 2g - 1 + r_2/2$ .

**5.2. Jacobian under the pull-back of the normalization.** We now recall some facts about the Jacobian under the pull-back of the normalization, cf. [16]. Let  $x \in Z \subset S$  be a singular point, i.e. either  $x \in Z_{\text{even}}$  or  $x = q_j$  with  $n_j \geq 3$ . Let  $\tilde{\mathcal{O}}_x$  be the integral closure of  $\mathcal{O}_x$ . We take  $V := \prod_{x \in Z} \tilde{\mathcal{O}}_x^*/\mathcal{O}_x^*$ . Then we have the following well-known short exact sequence induced by the pull-back of the normalization.

$$(9) \quad 0 \longrightarrow V \longrightarrow \text{Jac}(S) \xrightarrow{p^*} \text{Jac}(\tilde{S}) \longrightarrow 0.$$

This will play an important role later on.

**5.2.1. Hitchin fiber.** Now we examine the locally free part  $\mathcal{T}$  of the Hitchin fiber under the pull-back. Here,  $\mathcal{T}$  is defined to be the set of  $L \in \text{Pic}^{2g-2}(S)$  such that  $\det(\pi_* L) = \mathcal{O}_\Sigma$ . For any  $L \in \text{Pic}(S)$ , from (8) we see that  $\det(\tilde{\pi}_* p^* L) \cong \det(\pi_* L) \otimes \mathcal{O}_\Sigma(\Lambda')$ . We define a new set,  $\tilde{\mathcal{T}}$ , as follows:

$$\tilde{\mathcal{T}} := \{\tilde{L} \in \text{Pic}^{2g-2}(\tilde{S}) \mid \det(\tilde{\pi}_* \tilde{L}) \cong \mathcal{O}(\Lambda')\}.$$

It follows that  $p^*$  maps  $\mathcal{T}$  to  $\tilde{\mathcal{T}}$ . Furthermore, if  $L_1, L_2 \in \text{Pic}(S)$  satisfy  $p^*L_1 \cong p^*L_2$ , then we have  $\pi_*L_1 \cong \pi_*L_2$ . This means that the fiber of  $p^* : \text{Jac}(S) \rightarrow \text{Jac}(\tilde{S})$  is the same as  $p^* : \mathcal{T} \rightarrow \tilde{\mathcal{T}}$ , resulting in the following exact sequence:

$$(10) \quad 0 \longrightarrow V \longrightarrow \mathcal{T} \xrightarrow{p^*} \tilde{\mathcal{T}} \longrightarrow 0 .$$

**5.3. Torsion free sheaves.** Now we present Cook's parametrization of rank 1 torsion free sheaves on curves with Gorenstein singularities (see [4, p. 40] and also [5, 38]). An explicit computation of the invariants used in this paper is provided in Appendix A. Let  $x \in Z$  be a singular point of  $S$ , and let  $L \rightarrow S$  be a rank 1 torsion free sheaf. We again let  $\mathcal{O}_x$  denote the local ring at  $x$ ,  $\tilde{\mathcal{O}}_x$  its integral closure,  $\mathcal{K}_x$  its fraction field, and  $\delta_x = \dim_{\mathbb{C}}(\tilde{\mathcal{O}}_x/\mathcal{O}_x)$ . According to [18, Lemma 1.1], there exists a fractional ideal  $I_x$  that is isomorphic to  $L_x$ , uniquely defined up to multiplication by a unit of  $\tilde{\mathcal{O}}_x$ , such that  $\mathcal{O}_x \subset I_x \subset \tilde{\mathcal{O}}_x$ . We define  $\ell_x := \dim_{\mathbb{C}}(I_x/\mathcal{O}_x)$  and  $b_x := \dim_{\mathbb{C}}(\text{Tor}(I_x \otimes_{\mathcal{O}_x} \tilde{\mathcal{O}}_x))$ . Then,  $\ell_x$  and  $b_x$  are invariants of  $L$ .

Let  $\mathcal{K}_x$  be the field of fractions of  $\mathcal{O}_x$ . The *conductor* of  $I_x \subset \tilde{\mathcal{O}}_x$  is defined to be

$$C(I_x) = \{u \in \mathcal{K}_x \mid u \cdot \tilde{\mathcal{O}}_x \subset I_x\} .$$

The singularity is characterized by the following dimensions:

$$C(\mathcal{O}_x) \subset C(I_x) \subset \underbrace{\mathcal{O}_x \subset I_x \subset \tilde{\mathcal{O}}_x}_{\delta_x - \ell_x} .$$

$\underbrace{\hspace{10em}}_{2\delta_x}$

For  $x = p_i \in Z_{\text{even}}$ , we have  $\delta_{p_i} = m_i/2$ , and there are two maximal ideals  $\mathfrak{m}_{\pm}$  in  $\tilde{\mathcal{O}}_x$  corresponding to the points  $\tilde{p}_i^{\pm}$ . We let  $(\tilde{\mathcal{O}}_{p_i}/C(I_{p_i}))_{\mathfrak{m}_{\pm}}$  be the localization by the ideals  $\mathfrak{m}_{\pm}$ , and define  $a_{\tilde{p}_i^{\pm}} := \dim_{\mathbb{C}}(\tilde{\mathcal{O}}_{p_i}/C(I_{p_i}))_{\mathfrak{m}_{\pm}}$ . Moreover, we have  $\dim_{\mathbb{C}}(\tilde{\mathcal{O}}_{p_i}/C(\mathcal{O}_{p_i}))_{\mathfrak{m}_{\pm}} = m_i/2 = \delta_{p_i}$ . By Appendix A,  $a_{\tilde{p}_i^{\pm}} = (m_i/2) - \ell_{p_i}$ , and therefore  $a_{\tilde{p}_i^+} + a_{\tilde{p}_i^-} = 2\delta_{p_i} - 2\ell_{p_i}$ , and also  $b_{p_i} = \ell_{p_i}$ . Define

$$V(L_{p_i}) = \{(c_i^+, c_i^-) \mid c_i^{\pm} \in \mathbb{Z}_{\geq 0}, c_i^+ + c_i^- = \ell_{p_i}\} .$$

For  $x = p'_i \in Z_{\text{odd}}$ , we have  $\delta_{p'_i} = (n_i - 1)/2$ , and the maximal ideal  $\mathfrak{m}$  of  $\tilde{\mathcal{O}}_x$  is unique. Define  $a_{\tilde{p}'_i} := \dim_{\mathbb{C}}(\tilde{\mathcal{O}}_{p'_i}/C(I_{p'_i}))_{\mathfrak{m}}$ . By Appendix A, we have  $a_{\tilde{p}'_i} = 2\delta_{p'_i} - 2\ell_{p'_i}$  and  $b_{p'_i} = \ell_{p'_i}$ . Moreover,  $\dim_{\mathbb{C}}(\tilde{\mathcal{O}}_{p'_i}/C(\mathcal{O}_{p'_i}))_{\mathfrak{m}} = n_i - 1 = 2\delta_{p'_i}$ . In this case we set  $V(L_{p'_i}) = \{\ell_{p'_i}\}$ .

Now consider modules compatible with  $L_x$ . Let  $\eta : \tilde{\mathcal{O}}_x \rightarrow \tilde{\mathcal{O}}_x/C(\mathcal{O}_x)$  be the quotient map. Define

$$S(L_x) := \{\mathcal{O}_x\text{-submodules } F_x \subset \tilde{\mathcal{O}}_x/C(\mathcal{O}_x) \mid \dim_{\mathbb{C}}(F_x) = \delta, \eta^{-1}(F_x) \cong L_x\} .$$

Hence, if  $J_x = \eta^{-1}(F_x)$  with  $F_x \in S(L_x)$ , there exists an ideal  $\mathfrak{t}_x$  such that  $J_x = \mathfrak{t}_x \cdot L_x$ . For  $x = p_i \in Z_{\text{even}}$ , we obtain two integers  $c_i^{\pm} = \dim_{\mathbb{C}}(\tilde{\mathcal{O}}_x/(\mathfrak{t}_x \cdot \tilde{\mathcal{O}}_x))_{\mathfrak{m}_{\pm}}$ . By [4, Lemma 6],  $(c_i^+, c_i^-) \in V(L_{p_i})$ , for  $x = p'_i \in Z_{\text{odd}}$ ,  $\dim_{\mathbb{C}}(\tilde{\mathcal{O}}_x/(\mathfrak{t}_x \cdot \tilde{\mathcal{O}}_x)) = \ell_{p'_i} \in V(L_{p'_i})$ , and these only depend on  $F_x$ . Hence, there is a well-defined map:

$$\kappa_x : S(L_x) \longrightarrow V(L_x) : \begin{cases} F_x \rightarrow (c_i^+, c_i^-) & \text{when } x = p_i, \\ F_x \rightarrow \ell_{p'_i} & \text{when } x = p'_i . \end{cases}$$

**Lemma 5.1.** [4, Lemma 6] *For  $x \in Z$ , the components of  $S(L_x)$  are parameterized by elements in  $V(L_x)$ .*

Set  $V(L) := \prod_{x \in Z} V(L_x)$  and  $S(L) := \prod_{x \in Z} S(L_x)$ . Write  $N(L) := |V(L)|$  for the number of points in  $V(L)$ . there is a map

$$\kappa := \prod_{x \in Z} \kappa_x : S(L) \longrightarrow V(L) .$$

For any  $\mathbf{c} \in V(L)$ , write  $\mathbf{c} = (c_1^\pm, \dots, c_{r_1}^\pm, \ell_{p'_1}, \dots, \ell_{p'_{r_2}})$ . Associate to this the divisor

$$D_{\mathbf{c}} = \sum_{i=1}^{r_1} (c_i^+ \tilde{p}_i^+ + c_i^- \tilde{p}_i^-) + \sum_{i=1}^{r_2} \ell_{p'_i} \tilde{p}'_i$$

on  $\tilde{S}$ . Composing  $\kappa$  with the map above, we define

$$(11) \quad \varkappa : S(L) \longrightarrow \text{Div}(\tilde{S}) : \prod_{x \in Z} F_x \mapsto \mathbf{c} \mapsto D_{\mathbf{c}} .$$

The following result is straightforward but important:

**Proposition 5.2.**  *$L$  is locally free if and only if  $\varkappa = 0$  on  $S(L)$ .*

*Proof.*  $L$  is locally free if and only if  $\ell_x = 0$  for  $x \in Z$ . The claim then follows directly from the definition of  $D_{\mathbf{c}}$ .  $\square$

**5.4. Parabolic modules.** In this subsection, we define the notion of a parabolic module, following [38, 5, 4].

First note that  $\dim_{\mathbb{C}}(\tilde{\mathcal{O}}_x/C(\mathcal{O}_x)) = 2\delta_x$ . Let  $\text{Gr}(\delta_x, \tilde{\mathcal{O}}_x/C(\mathcal{O}_x))$  be the Grassmannian of  $\delta_x$  dimensional subspaces of the vector space  $\tilde{\mathcal{O}}_x/C(\mathcal{O}_x)$ . Then  $\tilde{\mathcal{O}}_x^*$  acts on  $\text{Gr}(\delta_x, \tilde{\mathcal{O}}_x/C(\mathcal{O}_x))$  by multiplication, with fixed points corresponding to  $\delta_x$ -dimensional  $\mathcal{O}_x$  submodules of  $\tilde{\mathcal{O}}_x/C(\mathcal{O}_x)$ . We write  $\mathcal{P}(x)$  for the (reduced) variety of fixed points. This is a closed subvariety of  $\text{Gr}(\delta_x, \tilde{\mathcal{O}}_x/C(\mathcal{O}_x))$ .

Suppose  $x$  is an  $A_n$  singularity. For notation convenience, we write  $\mathcal{P}(A_n) := \mathcal{P}(x)$ . We have the following:

**Proposition 5.3.** [4, Prop. 2] *The following holds:*

- (i)  $\mathcal{P}(A_n)$  is connected and depends only on  $\delta_x$ . Also,  $\dim_{\mathbb{C}} \mathcal{P}(A_n) = n$ , and we have isomorphisms  $\mathcal{P}(A_{2n-1}) \cong \mathcal{P}(A_{2n})$ .
- (ii) If  $\mathcal{P}(A_0)$  is defined to be a point, then the inclusions  $\mathcal{P}(A_0) \subset \mathcal{P}(A_2) \subset \dots \subset \mathcal{P}(A_{2n})$  give a cell decomposition of  $\mathcal{P}(A_{2n})$ .
- (iii) The singular locus  $\text{Sing}(\mathcal{P}(A_{2n})) \cong \mathcal{P}(A_{2n-4})$ . In particular, it has codimension  $\geq 2$ . Moreover,  $\mathcal{P}(A_1) = \mathcal{P}(A_2) \cong \mathbb{C}P^1$ , and  $\mathcal{P}(A_4)$  is a quadric cone.

Define  $\mathcal{P}(S) = \prod_{x \in Z} \mathcal{P}(x)$ . This only depends on the curve singularity of  $S$ . Let  $J \in \text{Pic}(\tilde{S})$ . We have an isomorphism  $J_x \otimes \mathcal{O}_{\Lambda', x} \cong \tilde{\mathcal{O}}_x/C(\mathcal{O}_x)$  as  $\mathcal{O}_x$ -modules. More explicitly, as vector spaces,

$$J_{\tilde{p}_i^+}^{\oplus \frac{m_i}{2}} \oplus J_{\tilde{p}_i^-}^{\oplus \frac{m_i}{2}} \cong \tilde{\mathcal{O}}_{p_i}/C(\mathcal{O}_{p'_i}) , \quad J_{\tilde{p}'_i}^{\oplus (n_i-1)} \cong \tilde{\mathcal{O}}_{p'_i}/C(\mathcal{O}_{p'_i}) .$$

**Definition 5.4.** *A parabolic module  $\text{PMod}(\tilde{S})$  consists of pairs  $(J, v)$ , where  $J \in \text{Jac}(\tilde{S})$  and  $v = \prod_{x \in Z} v_x$ , with  $v_x \in J_x \otimes \mathcal{O}_{\Lambda', x}$ .*

By [4, p. 41],  $\text{PMod}(\tilde{S})$  has a natural algebraic structure. Let  $\text{pr} : \text{PMod}(\tilde{S}) \rightarrow \text{Jac}(\tilde{S})$  be the projection to the first component. Then  $\text{pr}$  defines a fibration of  $\text{PMod}(\tilde{S})$  with fiber  $\mathcal{P}(S)$ . Moreover, there is a finite morphism  $\tau : \text{PMod}(\tilde{S}) \rightarrow \overline{\text{Jac}}(S)$  defined by sending  $(J, v) \rightarrow L$ , where  $L$  is the kernel of the restriction map  $p_* J \rightarrow (J \otimes \mathcal{O}_{\Lambda})/v$ :

$$0 \longrightarrow L \longrightarrow p_* J \longrightarrow (J \otimes \mathcal{O}_{\Lambda})/v \longrightarrow 0 .$$

There is a diagram:

$$\begin{array}{ccc} \mathcal{P}(S) & \longrightarrow & \text{PMod}(\tilde{S}) \xrightarrow{\text{pr}} \text{Jac}(\tilde{S}) \\ & & \downarrow \tau \\ & & \overline{\text{Jac}}(S) \end{array} .$$

The map  $\tau$  may be regarded as the compactification of the pull-back normalization map  $p^*$  in (9).

**Theorem 5.5** ([4, Thm. 1]). *For the map  $\tau : \text{PMod}(\tilde{S}) \rightarrow \overline{\text{Jac}}(S)$  defined above,*

- (i)  $\tau$  is a finite morphism, where the fiber over  $L$  consists of  $N(L)$  points,
- (ii) The restriction  $\tau : \tau^{-1}\text{Jac}(S) \rightarrow \text{Jac}(S)$  is an isomorphism. Moreover, for  $L \in \text{Jac}(S)$ , we have  $\text{pr} \circ \tau^{-1}(L) = p^*(L)$ .
- (iii) Suppose  $\tau(J, v) = L$ . For  $x \in Z$ , we have  $v \in S(L)$ . Let  $D_v = \varkappa(v)$  be the divisor defined in (11). Then

$$0 \longrightarrow p^*L/\text{Tor}(p^*L) \longrightarrow J \longrightarrow \mathcal{O}_{D_v} \longrightarrow 0 .$$

In particular,  $p^*L/\text{Tor}(p^*L) = J(-D_v)$ .

Suppose all of the zeros of the quadratic differential  $q$  are odd. Then for  $L \in \overline{\text{Jac}}(S)$ ,  $N(L) = 1$ , and we can deduce the following.

**Corollary 5.6.** *If  $q^{-1}(0) = \{p'_1, \dots, p'_r\}$  and all zeroes have odd multiplicity, then  $\tau : \text{PMod}(\tilde{S}) \rightarrow \overline{\text{Jac}}(S)$  is a bijection. Moreover, for  $L \in \overline{\text{Jac}}(S)$  with  $\tau(J, v) = L$ , we have*

$$p^*L/\text{Tor}(p^*L) = J(-\sum \ell_{p'_i} \tilde{p}_i) .$$

We will now present an example of a parabolic module.

**Example 5.7** ([4, Ex. 2]). *Suppose  $q$  contains  $4g - 2$  simple zeros and one zero  $x$  of order 2. Then the spectral curve  $S$  has one nodal singularity at  $x$ . Denote  $p : \tilde{S} \rightarrow S$  the normalization, with  $p^{-1}(x) = \{\tilde{x}_+, \tilde{x}_-\}$ . Then  $\mathcal{P}(S) = \mathbb{C}P^1$ , and we obtain a fibration  $\mathbb{C}P^1 \rightarrow \text{PMod}(\tilde{S}) \rightarrow \text{Jac}(\tilde{S})$ . Let  $L \in \overline{\text{Jac}}(S)$ . If we write  $\tilde{L} := p^*L/\text{Tor}(p^*L)$ , then  $\tau^{-1}(L) = (\tilde{L} \otimes \mathcal{O}(\tilde{x}_+), v_+), (\tilde{L} \otimes \mathcal{O}(\tilde{x}_-), v_-)$ . We can define two sections:*

$$s_{\pm} : \overline{\text{Jac}}(S) \longrightarrow \text{PMod}(\tilde{S}), \quad s_{\pm} : L \longrightarrow (\tilde{L} \otimes \mathcal{O}(\tilde{x}_{\pm}), v_{\pm}) .$$

Then  $\overline{\text{Jac}}(S)$  is the quotient of  $\text{PMod}(\tilde{S})$  given by the identification  $\overline{\text{Jac}}(S) \cong \text{PMod}(\tilde{S})/(s_+ \sim \mathcal{O}(\tilde{x}_- - \tilde{x}_+)s_-)$ . In particular,  $\text{PMod}(\tilde{S})$  is not a fibration over  $\overline{\text{Jac}}(S)$ .

**Proposition 5.8.** *The singular set of  $\text{PMod}(\tilde{S})$  has codimension at least 2. Moreover, if the spectral curve  $S$  contains only cusp or nodal singularities, then  $\text{PMod}(\tilde{S})$  is smooth.*

*Proof.* As the singularities of  $\text{PMod}(\tilde{S})$  come from the space  $\mathcal{P}(S)$ , the claim follows from Proposition 5.3.  $\square$

Since we focus on  $\text{SL}(2, \mathbb{C})$  Higgs bundles, we must consider the parabolic module compactification of the fibration

$$0 \longrightarrow V \longrightarrow \mathcal{P} \xrightarrow{p^*} \text{Prym}(\tilde{S}/\Sigma) \longrightarrow 0 .$$



Setting,  $\widehat{\text{PMod}}(\tilde{S}) := \tau^{-1}(\overline{\mathcal{P}})$ , then there is a diagram from [16, p. 17]

$$(12) \quad \begin{array}{ccc} \mathcal{P}(S) & \longrightarrow & \widehat{\text{PMod}}(\tilde{S}) \xrightarrow{\text{pr}} \text{Prym}(\tilde{S}/\Sigma) \\ & & \downarrow \tau \\ & & \overline{\mathcal{P}} \end{array}$$

Theorem 5.5 proves that  $\text{pr} \circ \tau^{-1}|_{\mathcal{P}} = p^*$ .

**5.5. Stratifications of the BNR data.** Parabolic modules define a stratification of  $\overline{\mathcal{P}}$  and  $\overline{\mathcal{T}}$ . In the following,  $\pi : S \rightarrow \Sigma$  is a branched double cover,  $\sigma$  the associated involution on  $S$  and  $\sigma$  its extension to an involution on  $\tilde{S}$ .

For a rank 1 torsion free sheaf  $L \in \overline{\text{Pic}}(S)$ , consider the map

$$p_{\text{tf}}^* : \overline{\text{Pic}}(S) \longrightarrow \text{Pic}(\tilde{S}), \quad p_{\text{tf}}^*(L) := p^*L / \text{Tor}(p^*L)$$

i.e. the torsion free part of the pull-back to the normalization. By [37],  $p_{\text{tf}}^*(L) = p^*L$  at  $x \in \tilde{S}$  if and only if  $L$  is locally free at  $p(x) \in S$ .

**Definition 5.9** ([24]). *An effective divisor  $D \in \text{Div}(\tilde{S})$  is called a  $\sigma$ -divisor if*

- (i)  $D \leq \Lambda$  and  $\sigma^*D = D$ ,
- (ii) let  $x \in \text{Fix}(\sigma)$  and we write  $D|_x = d_x x$ , then  $d_x \equiv 0 \pmod{2}$ .

The  $\sigma$ -divisors play an important role in describing the singular Hitchin fibers.

**Proposition 5.10.** [24, 33] *Let  $L \in \overline{\mathcal{P}}$  and write  $\tilde{L} := p_{\text{tf}}^*L$ . Then we have  $\tilde{L} \otimes \sigma^*\tilde{L} = \mathcal{O}(\Lambda - D)$  for  $D$  an  $\sigma$ -divisor.*

The  $\sigma$ -divisors give a stratification of the BNR data  $\overline{\mathcal{T}}$  and  $\overline{\mathcal{P}}$ . For a  $\sigma$ -divisor  $D$ , define

$$(13) \quad \begin{aligned} \tilde{\mathcal{T}}_D &= \{J \in \text{Pic}(\tilde{S}) \mid J \otimes \sigma^*J = \mathcal{O}(\Lambda - D)\}; \\ \tilde{\mathcal{P}}_D &= \{J \in \text{Pic}(\tilde{S}) \mid J \otimes \sigma^*J = \mathcal{O}(-D)\}. \end{aligned}$$

Then by [24, Prop. 5.6],  $\tilde{\mathcal{T}}_D$  and  $\tilde{\mathcal{P}}_D$  are abelian torsors over  $\text{Prym}(\tilde{S}/\Sigma)$  with dimension  $g(\tilde{S}) - g(S) = g - 1 + \frac{1}{2}r_2$ . In addition, we define

$$(14) \quad \begin{aligned} \overline{\mathcal{T}}_D &= \{L \in \overline{\mathcal{T}} \mid p_{\text{tf}}^*L \in \tilde{\mathcal{T}}_D\}; \\ \overline{\mathcal{P}}_D &= \{L \in \overline{\mathcal{P}} \mid p_{\text{tf}}^*L \in \tilde{\mathcal{P}}_D\}. \end{aligned}$$

Then the partial order on divisors defines a stratification of  $\overline{\mathcal{T}}$  (resp.  $\overline{\mathcal{P}}$ ) by:  $\cup_{D' \leq D} \overline{\mathcal{T}}_{D'}$  (resp.  $\cup_{D' \leq D} \overline{\mathcal{P}}_{D'}$ ). The top strata are  $\overline{\mathcal{T}}_{D=0}$  (resp.  $\overline{\mathcal{P}}_{D=0}$ ), and these consist of the locally free sheaves. From the definition,  $\mathcal{T} = \overline{\mathcal{T}}_{D=0}$  and  $\mathcal{P} = \overline{\mathcal{P}}_{D=0}$ .

**Theorem 5.11.** [24, Thm. 6.2] (i) *Suppose  $q$  contains at least one zero of odd order. Then for each stratum indexed by  $\sigma$ -divisor  $D$ , if we write  $n_{ss}$  be the numbers of  $p$  such that  $D|_p = \Lambda|_p$ , then there are holomorphic fiber bundles*

$$(15) \quad \begin{aligned} (\mathbb{C}^*)^{k_1} \times \mathbb{C}^{k_2} &\longrightarrow \overline{\mathcal{T}}_D \xrightarrow{p_{\text{tf}}^*} \tilde{\mathcal{T}}_D; \\ (\mathbb{C}^*)^{k_1} \times \mathbb{C}^{k_2} &\longrightarrow \overline{\mathcal{P}}_D \xrightarrow{p_{\text{tf}}^*} \tilde{\mathcal{P}}_D, \end{aligned}$$

where  $k_1 = r_1 - n_{ss}$ ,  $k_2 = 2g - 2 - \frac{1}{2} \deg(D) - r_1 + n_{ss} - \frac{r_2}{2}$ , and  $r_1, r_2$  are the number of even and odd zeros. (ii) *Suppose  $q$  is irreducible but all zeros are of even order. Then there exists an analytic space  $\overline{\mathcal{T}}'_D$  and a double branched covering  $p : \overline{\mathcal{T}}'_D \rightarrow \overline{\mathcal{T}}_D$ , with  $\overline{\mathcal{T}}'_D$  a holomorphic fibration*

$$(\mathbb{C}^*)^{k_1} \times \mathbb{C}^{k_2} \longrightarrow \overline{\mathcal{T}}'_D \xrightarrow{p_{\text{tf}}^*} \tilde{\mathcal{T}}_D.$$

In particular,  $\dim(\overline{\mathcal{P}}_D) = \dim(\overline{\mathcal{T}}_D) = 3g - 3 - \frac{1}{2} \deg(D)$ .

As explained in [24], via the BNR correspondence the stratification above translates into a stratification of the Hitchin fiber. Let  $\mathcal{M}_q := \mathcal{H}^{-1}(q)$  be the Hitchin fiber of an irreducible quadratic differential  $q$ . Let  $\chi_{\text{BNR}} : \overline{\mathcal{T}} \rightarrow \mathcal{M}_q$  be the bijection in Theorem 2.3. Let  $D$  be a  $\sigma$ -divisor. Define  $\mathcal{M}_{q,D} := \chi_{\text{BNR}}(\overline{\mathcal{T}}_D)$ . Then the stratification of  $\overline{\mathcal{T}}$  induces a stratification on  $\mathcal{M}_q = \bigcup_D \mathcal{M}_{q,D}$ .

For each  $\sigma$ -divisor  $D$ , since  $\sigma^*D = D$ , we can write  $D' := \frac{1}{2}\tilde{\pi}(D)$ . Then  $D'$  is an effective divisor with  $\text{supp } D' \subset Z$ . Moreover, for  $x \in q^{-1}(0)$ ,  $D'_x \leq \frac{1}{2}[\text{ord}_x(q)]$ . Therefore,  $\mathcal{M}_q$  may be regarded as also being stratified by divisors  $D'$  defined over  $\Sigma$ .

**5.6. The structure of the parabolic module projection.** We now explain the relationship between the divisor  $D_v$  in Theorem 5.5 and the  $\sigma$ -divisor. Given  $L \in \overline{\mathcal{P}}$ , define

$$(16) \quad \begin{aligned} \mathcal{N}_L &:= \{(J, v) \in \widehat{\text{PMod}}(\tilde{S}) \mid \tau(J, v) = L\}; \\ \mathcal{D}_L &:= \{D_v \mid (J, v) \in \mathcal{N}_L\}, \end{aligned}$$

where  $\mathcal{N}_L$  is  $\tau^{-1}(L)$ , and  $\mathcal{D}_L$  is the collection of divisors  $D_v$  such that  $J(-D_v) = p_{\text{tf}}^*(L)$ . If  $L$  is locally free, then  $J = p^*L$ , and  $\mathcal{D}_L$  is empty. Moreover, if  $\tau(J, v) = \tau(J', v)$ , then  $J' = J(D_{v'} - D_v)$ .

The divisor  $D_v$  satisfies the following symmetry property:

**Proposition 5.12.** *Let  $D$  be a  $\sigma$ -invariant divisor and  $L \in \overline{\mathcal{P}}_D$ . For any  $D_v \in \mathcal{D}_L$ , we have  $D_v + \sigma^*D_v = D$ .*

*Proof.* Let  $\tau(J, v) = L$ . Then by Theorem 5.5, we have  $\tilde{L} = J(-D_v)$ , where  $\tilde{L} = p_{\text{tf}}^*(L)$ . As  $L \in \overline{\mathcal{P}}_D$  and  $J \in \text{Prym}(\tilde{S}/\Sigma)$ , we have  $\tilde{L} \otimes \sigma^*\tilde{L} = \mathcal{O}(-D)$  and  $J \otimes \sigma^*J = \mathcal{O}_{\tilde{S}}$ , which implies  $D_v + \sigma^*D_v = D$ .  $\square$

As a consequence, we have the following:

**Corollary 5.13.** *Suppose  $q$  has only zeroes of odd order. Then for  $L \in \overline{\mathcal{P}}_D$  and  $D_v \in \mathcal{D}_L$ , we have  $\sigma^*D_v = D_v$  and  $D_v = \frac{1}{2}D$ . In addition,  $\tau : \widehat{\text{PMod}}(\tilde{S}) \rightarrow \overline{\mathcal{P}}$  is a bijection.*

*Proof.* Since each zero has odd order  $\text{supp}(D_v) \subset \text{Fix}(\sigma)$ , which implies  $D_v = \sigma^*D_v$ . By Proposition 5.12, we must have  $D_v = \frac{1}{2}D$ .  $\square$

There are relationships between the integers appearing in the construction of the parabolic module:

**Lemma 5.14** ([17]). *Let  $D = \sum_{i=1}^{r_1} d_i(\tilde{p}_i^+ + \tilde{p}_i^-) + \sum_{i=1}^{r_2} d'_i \tilde{p}'_i$  be a  $\sigma$ -divisor, and let  $L \in \overline{\mathcal{P}}_D$ . Then we have*

- (i)  $\ell_{p_i} = d_i$  and  $\ell_{p'_i} = d'_i/2$ ;
- (ii)  $a_{\tilde{p}_i^+} = a_{\tilde{p}_i^-} = (m_i/2) - d_i$  and  $a_{\tilde{p}'_i} = n_i - 1 - d'_i$ .

*Proof.* Since  $L \in \overline{\mathcal{P}}_D$ , we have  $\dim \text{Tor}(p^*L_{p_i}) = d_i$  and  $\dim \text{Tor}(p^*L_{p'_i}) = d'_i/2$ . The claim then follows from Proposition A.1.  $\square$

The elements in  $\mathcal{D}_L$  can be explicitly computed.

**Proposition 5.15.** *Let  $D = \sum_{i=1}^{r_1} d_i(\tilde{p}_i^+ + \tilde{p}_i^-) + \sum_{i=1}^{r_2} d'_i \tilde{p}'_i$  be a  $\sigma$ -divisor, and let  $L \in \overline{\mathcal{P}}_D$ . Then  $N_L = \prod_{i=1}^{r_1} (d_i + 1)$ . In particular, the number  $N_L$  depends only on the  $\sigma$ -divisor  $D$ .*

*Proof.* By Lemma 5.14,  $V(L)$  can be rewritten as

$$V(L) = \{(c_1^\pm, \dots, c_{r_1}^\pm, c'_1 = l_{p'_1}, \dots, c'_{r_2} = l_{p'_{r_2}}) \mid c_i^+ + c_i^- = d_i, c_i^\pm \in \mathbb{Z}_{\geq 0}\},$$

which implies the claim. The condition  $D_v + \sigma^*D_v = D$  is automatically satisfied.  $\square$

If we define  $n_L$  to be the number of  $D_v \in \mathcal{D}_L$  such that  $\sigma^*D_v \neq D_v$ , then we have the following:

**Proposition 5.16.** (i)  $n_L$  is even;  
 (ii) if  $L \in \overline{\mathcal{P}}_D$  with

$$D = \sum_{i=1}^{r_1} d_i(\tilde{p}_i^+ + \tilde{p}_i^-) + \sum_{i=1}^{r_2} d'_i \tilde{p}'_i,$$

and if there exists  $i_0 \in \{1, \dots, r_1\}$  such that  $d_{i_0}$  is not even, then  $n_L = N_L$ ; otherwise,  $n_L = N_L - 1$ .

*Proof.* To prove (i), note that if  $\sigma^*D_v \neq D_v$ , then  $\sigma^*(\sigma^*D_v) \neq \sigma^*D_v$ , which means that  $n_L$  is even. For (ii), by Proposition 5.15,  $D_v = \sigma^*D_v$  for  $D_v \in \mathcal{D}_L$  if and only if  $c_i^+ = c_i^- = d_i/2$ . Therefore,  $n_L \neq N_L$  if and only if all  $d_i$  are even, which implies (ii).  $\square$

## 6. IRREDUCIBLE SINGULAR FIBERS AND THE MOCHIZUKI MAP

In this section, we provide a reinterpretation of the limiting configuration construction of a Higgs bundle on an irreducible fiber introduced by Mochizuki in [33] (see also [24]). We also investigate the relationship between limiting configurations and the stratification.

**6.1. Abelianization of a Higgs bundle.** Let  $q$  be a fixed irreducible quadratic differential with spectral curve  $S$ , with normalization  $p : \tilde{S} \rightarrow S$ . We define  $\tilde{K} := \tilde{\pi}^*K$  (but note that  $\tilde{K} \neq K_{\tilde{S}}$ ) and  $\tilde{q} := \tilde{\pi}^*q \in H^0(\tilde{K}^2)$ , where  $\tilde{\pi}$  is the double branched covering of  $\tilde{S} \rightarrow \Sigma$  associated to the branching set  $Z$  of  $q$ . Choose a square root  $\omega \in H^0(\tilde{K})$  such that  $\tilde{q} = -\omega \otimes \omega$ . Let  $\Lambda := \text{Div}(\omega)$  and  $\tilde{Z} := \text{supp}(\Lambda)$ . We can then write

$$\Lambda = \sum_{i=1}^{r_1} \frac{m_i}{2} (\tilde{p}_i^+ + \tilde{p}_i^-) + \sum_{j=1}^{r_2} n_j \tilde{p}'_j.$$

If  $\sigma : \tilde{S} \rightarrow \tilde{S}$  is the involution, then  $\sigma^*\omega = -\omega$ .

Let  $(\mathcal{E}, \varphi)$  be a Higgs bundle with  $\det \varphi = q$ . Consider the pullback  $(\tilde{\mathcal{E}}, \tilde{\varphi}) := (\tilde{\pi}^*\mathcal{E}, \tilde{\pi}^*\varphi)$ . We have  $\tilde{\varphi} \in H^0(\text{End}(\tilde{\mathcal{E}}) \otimes \tilde{K})$  and  $\tilde{q} = \det(\tilde{\varphi})$ . Since  $\tilde{q} = -\omega \otimes \omega$ ,  $\pm\omega$  are well-defined eigenvalues of  $\tilde{\varphi}$  over  $\tilde{S}$ . Let  $\tilde{\lambda}$  be the canonical section of  $\text{Tot}(\tilde{K})$ . The spectral curve for  $(\tilde{\mathcal{E}}, \tilde{\varphi})$  is defined by the equation

$$\tilde{S}' := \{\tilde{\lambda}^2 - \tilde{q} = 0\}.$$

The set  $\tilde{S}' = \text{Im}(\omega) \cup \text{Im}(-\omega) \subset \text{Tot}(\tilde{K})$  decomposes into two irreducible pieces.

Having fixed a choice of  $\omega$ , the eigenvalues of  $\tilde{\varphi}$  are globally well-defined, and we can define the line bundle  $\tilde{L}_+ \subset \tilde{\mathcal{E}}$  as  $\tilde{L}_+ := \ker(\tilde{\varphi} - \omega)$ . Since  $\sigma^*\omega = -\omega$ ,  $\tilde{L}_- = \sigma^*\tilde{L}_+ = \ker(\tilde{\varphi} + \omega)$ , and there is an isomorphism  $\tilde{\mathcal{E}}|_{\tilde{S} \setminus \tilde{Z}} \cong \tilde{L}_+ \oplus \tilde{L}_-|_{\tilde{S} \setminus \tilde{Z}}$ .

There is a local description of  $(\tilde{\mathcal{E}}, \tilde{\varphi})$ :

**Lemma 6.1.** [24, Lemma 5.1, Thm. 5.3], [33, Lemma 4.2] *Let  $x \in \tilde{Z}$  and write  $\Lambda|_x = m_x x$ . Let  $U$  be a holomorphic coordinate neighborhood of  $x$ . Then there exists a frame  $\mathbf{e} \in H^0(U, \tilde{K})$  such that, under a suitable trivialization of  $\mathcal{E}|_U \cong U \times \mathbb{C}^2$ , we can write*

$$(17) \quad \tilde{\varphi} = z^{d_x} \begin{pmatrix} 0 & 1 \\ z^{2m_x - 2d_x} & 0 \end{pmatrix} \otimes \mathbf{e}.$$

Moreover, if we define  $D := \sum_{x \in \tilde{Z}} d_x x$ , then  $D$  is a  $\sigma$ -divisor.

**Lemma 6.2.** [33, Sec. 4.1] *For the  $\tilde{L}_\pm$  defined above, we have  $\tilde{L}_+ \otimes \tilde{L}_- = \mathcal{O}(D - \Lambda)$ . Moreover, if we denote  $\tilde{L}_0 := \tilde{L}_+(\Lambda - D)$  and  $\tilde{L}_1 := \sigma^* \tilde{L}_0$ , then  $\tilde{L}_+ = \tilde{\mathcal{E}} \cap \tilde{L}_0$ ,  $\tilde{L}_- = \tilde{\mathcal{E}} \cap \tilde{L}_1$ , and we have the exact sequences*

$$\begin{aligned} 0 &\longrightarrow \tilde{L}_+ \longrightarrow \tilde{\mathcal{E}} \longrightarrow \tilde{L}_1 \longrightarrow 0 ; \\ 0 &\longrightarrow \tilde{L}_- \longrightarrow \tilde{\mathcal{E}} \longrightarrow \tilde{L}_0 \longrightarrow 0 . \end{aligned}$$

*Proof.* The inclusion of  $\tilde{L}_\pm \rightarrow \mathcal{E}$  defines an exact sequence of sheaves

$$0 \longrightarrow \mathcal{O}(\tilde{L}_+) \oplus \mathcal{O}(\tilde{L}_-) \longrightarrow \mathcal{O}(\mathcal{E}) \longrightarrow \mathcal{T} \longrightarrow 0 ,$$

where  $\mathcal{T}$  is a torsion sheaf with  $\text{supp } \mathcal{T} \subset \tilde{Z}$ . From the local description in (17), in the same trivialization,  $\tilde{L}_\pm$  are spanned by the bases  $s_\pm = \begin{pmatrix} 1 \\ \pm z^{m_x - d_x} \end{pmatrix}$ . Therefore, as  $\det(\mathcal{E}) = \mathcal{O}_\Sigma$ , we obtain  $\tilde{L}_+ \otimes \tilde{L}_- = \mathcal{O}(D - \Lambda)$ . Since  $s_+, s_-$  are linear independent away from  $z$ ,  $\tilde{\mathcal{E}}/\tilde{L}_+$  is locally generated by the section  $z^{d_x - m_x} s_-$ . Therefore,  $\tilde{\mathcal{E}}/\tilde{L}_+ \cong \tilde{L}_-(\Lambda - D) = \tilde{L}_1$ . Using the involution, we obtain the other exact sequence.  $\square$

Therefore, if  $\tilde{L} \otimes \sigma^* \tilde{L} = \mathcal{O}(D - \Lambda)$ , we have  $\tilde{L}_0 = \tilde{L}(\Lambda - D) \in \tilde{\mathcal{T}}_D$ . In summary, the construction above leads us to consider the composition of the following maps:

$$\begin{aligned} \delta : \mathcal{M}_q &\longrightarrow \text{Pic}(\tilde{S}) \xrightarrow{\otimes \mathcal{O}(\Lambda - D)} \tilde{\mathcal{T}}_D ; \\ (\mathcal{E}, \varphi) &\mapsto \tilde{L}_+ \mapsto \tilde{L}_+(\Lambda - D) , \end{aligned}$$

where the first map is given by taking the kernel of  $(\tilde{\pi}^* \varphi - \omega)|_{\tilde{\pi}^* \mathcal{E}}$ .

This construction is directly related to the torsion free pull-back. Recall that  $\chi_{\text{BNR}} : \overline{\mathcal{T}} \rightarrow \mathcal{M}_q$  is the BNR correspondence map, and  $p_{\text{tf}}^* : \overline{\text{Pic}}(S) \rightarrow \text{Pic}(\tilde{S})$  is the torsion free pull-back. Then we have

**Proposition 6.3.**  $\delta \circ \chi_{\text{BNR}} = p_{\text{tf}}^*$ . *In particular, if  $J \in \overline{\mathcal{T}}_D$ , then  $\delta \circ \chi_{\text{BNR}}(J) \in \tilde{\mathcal{T}}_D$ .*

*Proof.* Let  $J \in \overline{\mathcal{T}}$ , and write  $(\mathcal{E}, \varphi) = \chi_{\text{BNR}}(J)$ ,  $(\tilde{\mathcal{E}}, \tilde{\varphi}) := \tilde{\pi}^*(\mathcal{E}, \varphi)$ . From (1), we have the BNR exact sequence on  $S$ :

$$0 \longrightarrow J(-\Delta) \longrightarrow \pi^* \mathcal{E} \xrightarrow{\pi^* \varphi - \lambda} \pi^* \mathcal{E} \otimes \pi^* K \longrightarrow J \otimes \pi^* K \longrightarrow 0 .$$

As  $p^*$  is right exact, we obtain

$$\tilde{\mathcal{E}} \xrightarrow{\tilde{\varphi} - \tilde{\lambda}} \tilde{\mathcal{E}} \otimes \tilde{K} \longrightarrow p^* J \otimes \tilde{K} \longrightarrow 0 .$$

Since the spectral curve is  $\tilde{S}' = \text{Im}(\omega) \cup \text{Im}(-\omega)$ , we can consider the restriction to the component  $\text{Im}(\omega)$  and write  $\tilde{\lambda} = \omega$ ,  $\tilde{L}_\pm := \ker(\tilde{\varphi} \mp \omega)$ . We obtain an exact sequence

$$0 \longrightarrow \tilde{L}_+ \longrightarrow \tilde{\mathcal{E}} \xrightarrow{\tilde{\varphi} - \omega} \tilde{\mathcal{E}} \otimes \tilde{K} \longrightarrow p^* J \otimes \tilde{K} \longrightarrow 0 ,$$

which breaks into short exact sequences

$$\begin{aligned} 0 &\longrightarrow \tilde{L}_+ \longrightarrow \tilde{\mathcal{E}} \longrightarrow \text{Im}(\tilde{\varphi} - \omega) \longrightarrow 0 ; \\ 0 &\longrightarrow \text{Im}(\tilde{\varphi} - \omega) \longrightarrow \tilde{\mathcal{E}} \otimes \tilde{K} \longrightarrow p^* J \otimes \tilde{K} \longrightarrow 0 . \end{aligned}$$

Using the local trivialization in Lemma 6.1,  $\text{Im}(\tilde{\varphi} - \omega)$  is locally spanned by  $\begin{pmatrix} z^{d_x} \\ -z^{m_x} \end{pmatrix} \mathbf{e}$ . From Lemma 6.2, if we write  $\tilde{L}_0 := \tilde{L}_+(\Lambda - D)$  and  $\tilde{L}_1 := \sigma^* \tilde{L}_0$ , then

$$\delta \circ \chi_{\text{BNR}}(J) = \tilde{L}_+(\Lambda - D) .$$

Moreover, there is an isomorphism  $\text{Im}(\tilde{\varphi} - \omega) \cong \tilde{L}_1$ . Letting  $\tilde{L}'_1$  be the saturation of  $\tilde{L}_1$ , then we obtain the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \tilde{L}_1 & \longrightarrow & \tilde{\mathcal{E}} \otimes \tilde{K} & \longrightarrow & p^*J \otimes \tilde{K} \longrightarrow 0 \\ & & \downarrow i & & \downarrow \cong & & \downarrow \\ 0 & \longrightarrow & \tilde{L}'_1 & \longrightarrow & \tilde{\mathcal{E}} \otimes \tilde{K} & \longrightarrow & p_{\text{tf}}^*J \otimes \tilde{K} \longrightarrow 0 \end{array}$$

where  $i : \tilde{L}_1 \rightarrow \tilde{L}'_1$  is the natural inclusion. Moreover, in the same trivialization,  $\tilde{L}'_1$  is spanned by the section  $\begin{pmatrix} 1 \\ -z^{m_x-d_x} \end{pmatrix} \epsilon$ . Therefore,  $\tilde{L}'_1 \cong \tilde{L}_- \otimes \tilde{K}$  and from Lemma 6.2,  $p_{\text{tf}}^*J = \delta \circ \chi_{\text{BNR}}(J)$ .  $\square$

From [33, Sec. 4.1], if  $(\mathcal{E}, \varphi)$  is a Higgs bundle with  $(\mathcal{E}, \varphi) = \chi_{\text{BNR}}(L)$ , and  $\tilde{L}_0 = \delta \circ \chi_{\text{BNR}}(L)$ , then by Proposition 6.3,  $\tilde{L}_0 = p_{\text{tf}}^*(L)$ . We define a Higgs bundle  $(\tilde{\mathcal{E}}_0, \tilde{\varphi}_0)$  as follows

$$\tilde{\mathcal{E}}_0 = \tilde{L}_0 \oplus \sigma^* \tilde{L}_0, \quad \tilde{\varphi}_0 = \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix}.$$

Moreover,  $\tilde{\mathcal{E}}$  is an  $\mathcal{O}_{\tilde{S}}$  submodule of  $\tilde{\mathcal{E}}_0$  with a natural inclusion  $\iota : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}_0$  satisfying the following:

- (i) the induced morphism  $\tilde{\mathcal{E}} \rightarrow \tilde{L}_0, \tilde{\mathcal{E}} \rightarrow \sigma^* \tilde{L}_0$  is surjective,
- (ii) the restriction of  $\iota|_{\tilde{S} \setminus \tilde{Z}}$  is an isomorphism,
- (iii)  $\tilde{\varphi}_0 \circ \iota = \iota \circ \tilde{\varphi}$ .

We call  $(\tilde{\mathcal{E}}_0, \tilde{\varphi}_0)$  the *abelianization of the Higgs bundle*  $(\mathcal{E}, \varphi)$ .

**6.2. The construction of the algebraic Mochizuki map.** In this subsection, we define the algebraic Mochizuki map, as introduced in [33]. Recall that for any divisor  $D = \sum_{x \in Z} d_x x$ , there is a canonical weight function  $\chi_D(x) := d_x$  when  $x \in \text{supp } D$  and  $\chi_D(x) = 0$  when  $x \notin \text{supp } D$ . We also have the stratification  $\overline{\mathcal{T}} = \cup_D \overline{\mathcal{T}}_D$ , for  $\sigma$  divisors  $D$ . Let  $\mathcal{F}(\tilde{S})$  be the space of all degree zero filtered line bundles over  $\tilde{S}$ . The *algebraic Mochizuki map*  $\Theta^{\text{Moc}}$  is defined as

$$\Theta^{\text{Moc}} : \overline{\mathcal{T}} \longrightarrow \mathcal{F}(\tilde{S}), \quad L \mapsto \mathcal{F}_*(p_{\text{tf}}^*(L), \frac{1}{2}\chi_{D-\Lambda}).$$

**Example 6.4.** When  $q$  has only simple zeroes, this construction generalizes that of [30] (see also [13]). In the case of a quadratic differential with simple zeros, the spectral curve  $S$  is smooth, and every torsion free sheaf is locally free, so that  $\mathcal{T} = \overline{\mathcal{T}}$ . If  $Z = \{p_1, \dots, p_{4g-4}\}$  are the branch points of  $S$ , and  $\Lambda = \sum_{i=1}^{4g-4} p_i$ , then the weight function  $\frac{1}{2}\chi_{-\Lambda}$  assigns a weight of  $-\frac{1}{2}$  at each  $p_i$ . For  $L \in \mathcal{T}$ ,  $\Theta^{\text{Moc}}(L) = \mathcal{F}_*(L, \frac{1}{2}\chi_{-\Lambda})$ .

Below are some additional properties of  $\Theta^{\text{Moc}}$ .

**Proposition 6.5.**  $\Theta^{\text{Moc}}|_{\overline{\mathcal{T}}_D}$  is a continuous map.

*Proof.* This follows directly from the definition of  $\Theta^{\text{Moc}}$  and Theorem 3.3.  $\square$

From Theorem 5.11, we know that for a  $\sigma$ -divisor  $D$ , the preimage of the map  $p_{\text{tf}}^* : \overline{\mathcal{T}}_D \rightarrow \overline{\mathcal{T}}_D$  has dimension  $2g - 2 - \frac{1}{2} \deg(D) - r_2/2$ , where  $r_2$  is the number of odd zeros of  $q$ . Even for the top stratum  $D = 0$ ,  $p_{\text{tf}}^*$  is not injective if the spectral curve is not smooth. Let  $L_1, L_2 \in \overline{\mathcal{T}}_D$  with  $p_{\text{tf}}^*(L_1) = p_{\text{tf}}^*(L_2)$ . Then, based on the construction, we have  $\Theta^{\text{Moc}}(L_1) = \Theta^{\text{Moc}}(L_2)$ . In summary, we have the following result:

**Proposition 6.6.** *Let  $q \in H^0(K^2)$  be irreducible. Then  $\Theta^{\text{Moc}}$  is injective if and only if  $q$  has simple zeros.*

**6.3. Convergence of subsequences.** Fix a locally free  $L_0 \in \mathcal{T}$ . Using the isomorphism  $\psi_{L_0} : \overline{\mathcal{T}} \rightarrow \overline{\mathcal{P}}$  defined by  $\psi_{L_0}(L) = LL_0^{-1}$ , we can extend the Mochizuki map  $\Theta^{\text{Moc}}$  to  $\overline{\mathcal{P}}$ . For  $J \in \overline{\mathcal{P}}_D$ , we write  $\tilde{J} := p_{\text{tf}}^*(J)$  and choose the weight function  $\frac{1}{2}\chi_D$ . We then define:

$$\Theta_0^{\text{Moc}} : \overline{\mathcal{P}} \longrightarrow \mathcal{F}(\tilde{S}) , \quad J \mapsto \mathcal{F}_*(\tilde{J}, \frac{1}{2}\chi_D) .$$

**Proposition 6.7.** *The map  $\Theta_0^{\text{Moc}}$  satisfies the following properties:*

(i) *Let  $J \in \overline{\mathcal{P}}$  and  $L := L_0J$ , then*

$$\Theta_0^{\text{Moc}}(J) = \Theta^{\text{Moc}}(L) \otimes \Theta^{\text{Moc}}(L_0)^{-1} ,$$

*where  $\otimes$  is the tensor product for filtered line bundles (5).*

(ii) *Suppose  $L = \tau(I, v)$  with  $(\mathcal{I}, v) \in \widehat{\text{PMod}}(\tilde{S})$  and  $L \in \overline{\mathcal{P}}_D$ , then*

$$\Theta_0^{\text{Moc}} \circ \tau(I, v) = \mathcal{F}_*(I(-D_v), \frac{1}{2}\chi_{D_v + \sigma^*D_v}) ,$$

*where  $D_v$  is the corresponding divisor defined in Theorem 5.5.*

(iii) *If  $\sigma^*D_v = D_v$ , then  $\Theta_0^{\text{Moc}} \circ \tau(\mathcal{I}, v) = \mathcal{F}_*(\mathcal{I}, 0)$ , where 0 means to take parabolic weight 0 at any point.*

*Proof.* As  $L_0$  is locally free, we have  $p_{\text{tf}}^*J = (p^*L_0)^{-1} \otimes p_{\text{tf}}^*L$ . By definition,

$$\Theta_0^{\text{Moc}}(J) = \mathcal{F}_*(p_{\text{tf}}^*J, \frac{1}{2}\chi_D), \quad \Theta^{\text{Moc}}(L) = \mathcal{F}_*(p_{\text{tf}}^*L, \frac{1}{2}\chi_{D-\Lambda}), \quad \Theta^{\text{Moc}}(L_0) = \mathcal{F}_*(p_{\text{tf}}^*L_0, \frac{1}{2}\chi_{-\Lambda}),$$

which implies (i). For (ii), by Theorem 5.5,  $p_{\text{tf}}^*L = I(-D_v)$ , and from Proposition 5.12, we have  $D = D_v + \sigma^*D_v$ , which implies (ii). When  $\sigma^*D_v = D_v$ , we compute

$$\mathcal{F}_*(I(-D_v), \frac{1}{2}\chi_{D_v + \sigma^*D_v}) = \mathcal{F}_*(I(-D_v), \chi_{D_v}) = \mathcal{F}_*(I, 0),$$

which implies (iii).  $\square$

We now give a criterion for the continuity of the map  $\Theta^{\text{Moc}}$ . By this Proposition, it is sufficient to study the map  $\Theta_0^{\text{Moc}}$ . Recall that for  $L \in \overline{\mathcal{P}}$ , we have

$$\mathcal{N}_L := \{(J, v) \in \widehat{\text{PMod}}(\tilde{S}) \mid \tau(J, v) = L\}, \quad \mathcal{D}_L := \{D_v \mid (J, v) \in \mathcal{N}_L\},$$

and the number  $n_L$  is defined to be the number of divisors  $D_v \in \mathcal{D}_L$  such that  $\sigma^*D_v \neq D_v$ .

**Proposition 6.8.** *Let  $D$  be a  $\sigma$ -divisor,  $L \in \overline{\mathcal{P}}_D$ , and assume that  $\Theta_0^{\text{Moc}}$  is continuous at  $L$ . Then, for  $(J, v) \in \mathcal{N}_L$  and  $D_v \in \mathcal{D}_L$ , we have  $\sigma^*D_v = D_v$ , i.e.,  $n_L = 0$ .*

*Proof.* As the top stratum  $\mathcal{P}$  is dense in  $\overline{\mathcal{P}}$ , there exists a family of  $L_i \in \mathcal{P}$  such that  $\lim_{i \rightarrow \infty} L_i = L$ . Let  $(J_i, v_i) \in \widehat{\text{PMod}}(\tilde{S})$  be such that  $\tau(J_i, v_i) = L_i$ . Then,  $\lim_{i \rightarrow \infty} (J_i, v_i) = (J_\infty, v_\infty)$ , and  $\tau(J_\infty, v_\infty) = L$ . As  $L_i$  is locally free, we have  $D_{v_i} = 0$ . Moreover, by Theorem 5.5, we have  $p_{\text{tf}}^*L = J_\infty(-D_{v_\infty})$ , and from Proposition 5.12, we have  $D = D_{v_\infty} + \sigma^*D_{v_\infty}$ . By Proposition 6.7, we have

$$\Theta_0^{\text{Moc}}(L_i) = \Theta_0^{\text{Moc}} \circ \tau(J_i, v_i) = \mathcal{F}_*(J_i, 0) ,$$

and we compute

$$\lim_{i \rightarrow \infty} \Theta_0^{\text{Moc}}(L_i) = \mathcal{F}_*(J_\infty, 0) = \mathcal{F}_*(J_\infty(-D_{v_\infty}), \chi_{D_{v_\infty}}) .$$

Moreover, by Proposition 6.7, we have  $\Theta_0^{\text{Moc}}(L) = \mathcal{F}_*(J_\infty(-D_{v_\infty}), \frac{1}{2}(\chi_{D_{v_\infty}} + \chi_{\sigma^*D_{v_\infty}}))$ . Since  $\Theta_0^{\text{Moc}}$  is continuous on  $L$ , we have  $\lim_{i \rightarrow \infty} \Theta_0^{\text{Moc}}(L_i) = \Theta_0^{\text{Moc}}(L)$ , which implies that  $\chi_{D_{v_\infty}} = \chi_{\sigma^*D_{v_\infty}}$ .  $\square$

By Proposition 5.16,  $n_L > 0$  if and only if  $q$  has at least one zero of even order. Hence, the following is immediate.

**Corollary 6.9.** *Suppose  $q$  is irreducible and has a zero of even order, then  $\Theta_0^{\text{Moc}}$  is not continuous.*

When all zeros of  $q$  have odd order, however, we have the following.

**Proposition 6.10.** *If  $q$  is irreducible with all zeroes of odd order, then  $\Theta_0^{\text{Moc}}$  is continuous.*

*Proof.* Since all zeroes of  $q$  are odd, for any  $L \in \overline{\mathcal{P}}$ , we have  $n_L = 0$ . Let  $L_\infty \in \overline{\mathcal{P}}$  be fixed and let  $L_i \in \overline{\mathcal{P}}$  be any sequence such that  $\lim_{i \rightarrow \infty} L_i = L_\infty$ . Since  $\tau : \widehat{\text{PMod}}(\tilde{S}) \rightarrow \overline{\mathcal{P}}$  is bijective, we take  $(J_i, v_i) \in \widehat{\text{PMod}}(\tilde{S})$  with  $\tau(J_i, v_i) = L_i$ . Moreover, we assume  $\lim_{i \rightarrow \infty} (J_i, v_i) = (J_\infty, v_\infty)$  with  $\tau(J_\infty, v_\infty) = L_\infty$ . Since  $q$  only contains odd zeros, it follows that  $\text{supp } D_v \subset \text{Fix}(\sigma)$ . By Proposition 6.7, we have  $\Theta_0^{\text{Moc}}(L_i) = \mathcal{F}_*(J_i, 0)$ . Therefore, we have:

$$\lim_{i \rightarrow \infty} \Theta_0^{\text{Moc}}(L_i) = \lim_{i \rightarrow \infty} \mathcal{F}_*(J_i, 0) = \mathcal{F}_*(J_\infty, 0) = \Theta_0^{\text{Moc}}(L_\infty).$$

This concludes the proof.  $\square$

**Theorem 6.11.** *Suppose  $q$  is irreducible. For the map  $\Theta^{\text{Moc}} : \mathcal{M}_q \rightarrow \mathcal{F}(\tilde{S})$ , we have:*

- (i)  $\Theta^{\text{Moc}}$  is injective if and only if  $q$  only has only simple zeros;
- (ii) if  $q$  has only zeroes of odd order,  $\Theta^{\text{Moc}}$  is continuous;
- (iii) if  $q$  contains a zero of even order,  $\Theta^{\text{Moc}}$  is not continuous.

*Proof.* (i) follows from Proposition 6.6. (ii) follows from Proposition 6.10. (iii) follows from Corollary 6.9.  $\square$

**Proposition 6.12.** *Suppose  $n_L > 0$ . Then for  $k = 1, \dots, n_L$ , there exist sequences  $L_i^k$  with  $\lim_{i \rightarrow \infty} L_i^k = L$  such that if we denote  $\mathcal{F}_*^k := \lim_{i \rightarrow \infty} \Theta_0^{\text{Moc}}(L_i^k)$ ,  $\mathcal{F}_*^0 := \Theta_0^{\text{Moc}}(L)$ , then  $\mathcal{F}_*^{k_1} \neq \mathcal{F}_*^{k_2}$  for  $k_1 \neq k_2$ . Moreover, there exist  $\{D_1, \dots, D_{n_L}\} \subset \mathcal{D}_L$  such that  $\mathcal{F}_*^k = \mathcal{F}_*(p_{\text{tf}}^* L, \chi_{D_k})$ .*

*Proof.* By the definition of  $n_L$ , we can find  $(J^k, v^k)$  with  $\tau(J^k, v^k) = L$ . If we define  $D_k := D_{v^k}$ , then  $\sigma^* D_k \neq D_k$ . Moreover, by Theorem 5.5, we have  $p_{\text{tf}}^* L = J^k(-D_k)$ . As  $\tau^{-1}(\mathcal{P})$  is dense in  $\widehat{\text{PMod}}(\tilde{S})$ , for each  $(J^k, v^k)$ , we can find a sequence  $(J_i^k, v_i^k) \in \tau^{-1}(\mathcal{P})$  such that  $\lim_{i \rightarrow \infty} (J_i^k, v_i^k) = (J^k, v^k)$  and we define  $L_i^k := \tau(J_i^k, v_i^k)$ . Since  $L_i^k$  is locally free,  $D_{v_i^k} = 0$ , and thus  $\Theta_0^{\text{Moc}}(L_i^k) = \mathcal{F}_*(J_i^k, 0)$ . We compute

$$\lim_{i \rightarrow \infty} \Theta_0^{\text{Moc}}(L_i^k) = \mathcal{F}_*(J^k, 0) = \mathcal{F}_*(p_{\text{tf}}^* L, \chi_{D_k})$$

and  $\Theta_0^{\text{Moc}}(L) = \mathcal{F}_*(p_{\text{tf}}^* L, \frac{1}{2}\chi_D)$ . Based on our assumptions, we have  $D_{k_1} \neq D_{k_2}$  for  $k_1 \neq k_2$  and  $\sigma^* D_k \neq D_k$ , which implies that  $\chi_{D_{k_1}} \neq \chi_{D_{k_2}}$  for  $k_1 \neq k_2$  and  $\chi_{D_k} \neq \frac{1}{2}\chi_D$ .  $\square$

We now present a computation for the case of a simple nodal curve.

**Example 6.13.** *Let  $q$  be a quadratic differential with  $2g - 4$  simple zeros, and let  $x$  be an even zero of  $q$  of order two. Then  $S$  has a singular point, which we also denote by  $x$ . Let  $p : \tilde{S} \rightarrow S$  be the normalization map and  $p^{-1}(x) = \{x_1, x_2\}$ . Consider the  $\sigma$ -divisor  $D = x_1 + x_2$ , and let  $L \in \overline{\mathcal{P}}_D$ . Then  $n_L = 2$ , and we can write  $\mathcal{N}_L = (J_1, v_1), (J_2, v_2)$ , where  $D_{v_1} = x_1$  and  $D_{v_2} = x_2$ . Moreover, we have  $p_{\text{tf}}^* L = J_1 \otimes \mathcal{O}(-x_1) = J_2 \otimes \mathcal{O}(-x_2)$ . Let  $(\alpha, \beta)$  denote the parabolic weight that is equal to  $\alpha$  at  $x_1$ ,  $\beta$  at  $x_2$ , and  $\frac{1}{2}$  at all other zeros. Then the filtered bundles obtained in Proposition 6.12 are*

$$\mathcal{F}_*(p_{\text{tf}}^* L, (1, 0)) , \mathcal{F}_*(p_{\text{tf}}^* L, (0, 1)) , \mathcal{F}_*(p_{\text{tf}}^* L, (\frac{1}{2}, \frac{1}{2})) .$$

**6.4. Mochizuki's convergence theorem for irreducible fibers.** In this subsection, we recall Mochizuki's construction of the limiting configuration metric [33] and the convergence theorem.

**6.4.1. Limiting configuration metric.** Let  $q$  be an irreducible quadratic differential, and let  $(\mathcal{E}, \varphi) \in \mathcal{M}_q$  be a Higgs bundle with  $(\mathcal{E}, \varphi) = \chi_{\text{BNR}}(L)$ . We write  $\tilde{L}_0 = p_{\text{tf}}^* L$  and  $(\tilde{\mathcal{E}}, \tilde{\varphi}) := p^*(\mathcal{E}, \varphi)$ . Then the abelianization of  $(\mathcal{E}, \varphi)$ , which is a Higgs bundle over  $\tilde{S}$ , can be written as  $\tilde{\mathcal{E}}_0 = \tilde{L}_0 \oplus \sigma^* \tilde{L}_0$ ,  $\tilde{\varphi}_0 = \text{diag}(\omega, -\omega)$ . The natural inclusion  $\iota : (\tilde{\mathcal{E}}, \tilde{\varphi}) \rightarrow (\tilde{\mathcal{E}}_0, \tilde{\varphi}_0)$  is an isomorphism over  $\tilde{S} \setminus \tilde{Z}$ .

We define  $\tilde{h}_0$  to be the harmonic metric for the filtered bundle  $\Theta^{\text{Moc}}(L)$ ; this is well-defined up to a multiplicative constant. To fix this constant, assume that  $\sigma^* \tilde{h}_0 \otimes \tilde{h}_0 = 1$ . This gives a unique choice of  $\tilde{h}_0$ . We then define the metric  $\tilde{H}_0 = \text{diag}(\tilde{h}_0, \sigma^* \tilde{h}_0)$  on  $\tilde{\mathcal{E}}_0$ , with  $\det(\tilde{H}_0) = 1$ . For the resulting harmonic bundle  $(\tilde{\mathcal{E}}_0, \varphi_0, \tilde{H}_0)$ , we define  $\tilde{\nabla}_0$  to be the unitary connection determined by  $\tilde{H}_0$ . Since  $\tilde{H}_0$  is diagonal, it follows that  $F_{\tilde{\nabla}_0} = 0$ , and we have  $[\varphi_0, \varphi_0^{\dagger \tilde{H}_0}] = 0$ . Furthermore, as  $\iota$  is an isomorphism over  $\tilde{S} \setminus \tilde{Z}$ , the metric  $\tilde{H}_0$  also defines a metric on  $(\tilde{\mathcal{E}}, \tilde{\varphi})$  over  $\tilde{S} \setminus \tilde{Z}$ .

For any  $\tilde{x} \in \tilde{S} \setminus \tilde{Z}$  with  $x := p(\tilde{x})$ , we have isomorphisms

$$(\tilde{\mathcal{E}}_0, \tilde{\varphi}_0)|_{\sigma(\tilde{x})} \cong (\tilde{\mathcal{E}}_0, \tilde{\varphi}_0)|_{\tilde{x}} \cong (\tilde{\mathcal{E}}, \tilde{\varphi})|_{\tilde{x}} \cong (\mathcal{E}, \varphi)|_x.$$

Therefore,  $\tilde{H}_0$  induces a metric  $H^{\text{Lim}}$  on  $\Sigma \setminus Z$ , and we may consider  $H^{\text{Lim}}$  as the push-forward of  $\tilde{h}_0$ . In [25, Theorem 5.2], the push-forward metric can be explicitly written in local coordinates.

Over  $\Sigma \setminus Z$ , we write  $\nabla^{\text{Lim}}$  for the Chern connection of  $H^{\text{Lim}}$  and  $\varphi^{\dagger \text{Lim}}$  for the adjoint with respect to  $H^{\text{Lim}}$ . They satisfy the decoupled Hitchin equations over  $\Sigma \setminus Z$ . Thus from any Higgs bundle  $(\mathcal{E}, \varphi)$ , we obtain a limiting configuration  $(\nabla^{\text{Lim}}, \phi^{\text{Lim}} = \varphi + \varphi^{\dagger \text{Lim}}) \in \mathcal{M}_{\text{Hit}}^{\text{Lim}}$ . This leads us to define the *analytic Mochizuki map*  $\Upsilon^{\text{Moc}}$  as

$$(18) \quad \Upsilon^{\text{Moc}} : \mathcal{M}_q \longrightarrow \mathcal{M}_{\text{Hit}}^{\text{Lim}} : [(\mathcal{E}, \varphi)] \mapsto [(\nabla^{\text{Lim}}, \phi^{\text{Lim}})].$$

**6.4.2. The continuity of the limiting configurations.** We now introduce the main result of Mochizuki [33]. Fix  $(\mathcal{E}, \varphi) = \chi_{\text{BNR}}(L) \in \mathcal{M}_q$ . For any real parameter  $t$ ,  $(\mathcal{E}, t\varphi)$  is a stable Higgs bundle. By the Kobayashi-Hitchin correspondence, there exists a unique metric  $H_t$  solving the Hitchin equation. Denote by  $\nabla_t$  the unitary connection defined by  $H_t$  and write  $D_t = \nabla_t + t\phi_t$  for the full  $\text{SL}(2, \mathbb{C})$  flat connection. We then have:

**Theorem 6.14.** [33] *The family  $(\mathcal{E}, t\varphi)$  has a unique limiting configuration as limit for  $t \rightarrow \infty$ . Moreover, for any compact set  $K \subset \Sigma \setminus Z$ , let  $d := \min_{x \in K} |q|(x)$ . Then there exist  $t$ -independent constants  $C_k$  and  $C'_k$  such that*

$$|(\nabla_t, \phi_t) - \Upsilon^{\text{Moc}}(\mathcal{E}, \varphi)|_{C^k} \leq C_k e^{-C'_k t d}.$$

As the map  $\Upsilon^{\text{Moc}}$  is the composition of  $\Theta^{\text{Moc}} \circ \chi_{\text{BNR}}^{-1}$  with the nonabelian Hodge correspondence, the behavior of  $\Upsilon^{\text{Moc}}$  is the same as  $\Theta^{\text{Moc}}$ . By Theorem 6.11 and Proposition 6.12, we obtain:

**Theorem 6.15.** *Let  $q$  be an irreducible quadratic differential. The map  $\Upsilon^{\text{Moc}} : \mathcal{M}_q \rightarrow \mathcal{M}_{\text{Hit}}^{\text{Lim}}$  satisfies the following properties:*

- (i) *If all the zeroes of  $q$  are odd, then  $\Upsilon^{\text{Moc}}$  is continuous;*
- (ii) *If at least one zero of  $q$  is even, let  $\mathcal{M}_q = \bigcup \mathcal{M}_{q,D}$  be the stratification. For each  $(\mathcal{E}, \varphi) \in \mathcal{M}_{q,D}$ , there exists an integer  $n_D$  that only depends on  $D$ , and  $n_D$  sequences*



$\{(\mathcal{E}_i^k, \varphi_i^k)\}$  for  $k = 1, \dots, n_D$  such that  $\lim_{i \rightarrow \infty} (\mathcal{E}_i^k, \varphi_i^k) = (\mathcal{E}_\infty, \varphi_\infty)$  and for  $k_1 \neq k_2$ ,

$$\lim_{i \rightarrow \infty} \Upsilon^{\text{Moc}}(\mathcal{E}_i^{k_1}, \varphi_i^{k_1}) \neq \lim_{i \rightarrow \infty} \Upsilon^{\text{Moc}}(\mathcal{E}_i^{k_2}, \varphi_i^{k_2}) \neq \Upsilon^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty).$$

## 7. REDUCIBLE SINGULAR FIBER AND THE MOCHIZUKI MAP

We now investigate the properties of the Hitchin fiber associated with a reducible quadratic differential, as discussed in [16]. Additionally, we will provide an overview of Mochizuki's technique for constructing limiting configurations of Hitchin fibers for reducible quadratic differentials, as detailed in [33]. We also analyze the continuity of the Mochizuki map.

**7.1. Local description of a Higgs bundle.** In this subsection, we give a local description of Higgs bundles in the fiber of a reducible quadratic differential  $q$ . Write  $q = -\omega \otimes \omega$  with  $\omega \in H^0(K)$ ,  $\Lambda = \text{Div}(\omega)$ ,  $Z = \text{supp}(\Lambda)$ , and  $\mathcal{M}_q = \mathcal{H}^{-1}(q)$ . Compared to the irreducible case,  $\mathcal{M}_q$  contains strictly semistable Higgs bundles, so we let  $\mathcal{M}_q^{\text{st}}$  denote the stable locus. We should point out that there is an ambiguity up  $\pm$  in the choice of  $\omega$ , which actually plays an important role in the following.

7.1.1. *Local description.* Given a Higgs bundle  $(\mathcal{E}, \varphi)$  with  $\det(\varphi) = q$ , define line bundles

$$(19) \quad L_\pm := \ker(\varphi \pm \omega) .$$

Then the inclusion map  $L_\pm \rightarrow \mathcal{E}$  is injective. Similarly, we may also define an abelianization of  $(\mathcal{E}, \varphi)$  as  $(\mathcal{E}_0 = L_+ \oplus L_-, \varphi_0 = \text{diag}(\omega, -\omega))$ . We then have a natural inclusion  $\iota : \mathcal{E}_0 \rightarrow \mathcal{E}$ . Then  $\iota|_{\Sigma \setminus Z}$  is an isomorphism, and  $\varphi \circ \iota = \iota \circ \varphi_0$ .

It follows from [16, Prop. 7.10] that  $L_\pm$  are the only  $\varphi$ -invariant subbundles of  $\mathcal{E}$ . If we write  $d_\pm := \deg(L_\pm)$ , then  $(\mathcal{E}, \varphi)$  is stable (resp. semistable) if and only if  $d_\pm < 0$  (resp.  $\leq 0$ ). As  $\det(\mathcal{E}) = \mathcal{O}$ , the map  $\det(\iota) : L_+ \otimes L_- \rightarrow \mathcal{O}$  defines a divisor  $D = \text{Div}(\det(\iota))$  such that  $L_+ \otimes L_- = \mathcal{O}(-D)$ . Therefore, we obtain

$$d_+ + d_- + \deg D = 0 ,$$

and  $0 \leq D \leq \Lambda$ . The Higgs bundle  $(\mathcal{E}, \varphi)$  is semistable if and only if  $-\deg D \leq d_+ \leq 0$  and stable if the equalities do not hold. For the rest of this section, we always write  $D = \sum_{p \in Z} \ell_p p$ .

7.1.2. *Semistable settings.* As the fiber  $\mathcal{M}_q$  might contain strictly semistable Higgs bundles, we now explicitly enumerate all of the possible  $s$ -equivalent classes. When  $D = 0$ , then  $L_- = L_+^{-1}$  and  $\deg(L_+) = 0$ . The corresponding Higgs bundle is polystable and can be explicitly written as

$$\left( L \oplus L^{-1}, \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix} \right) ,$$

where  $L \in \text{Jac}(\Sigma)$ . When  $D \neq 0$ , suppose  $\deg(L_+) = -\deg(D)$ . Then  $L_- = L_+^{-1}(-D)$  and  $\deg(L_-) = 0$ . Under  $S$ -equivalence, the polystable Higgs bundle is

$$\left( L_+(D) \oplus L_+^{-1}(-D), \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix} \right) ,$$

where  $L_+ \in \text{Pic}^{-\deg(D)}(\Sigma)$ .

**7.2. Reducible spectral curves.** In this subsection, we introduce the algebraic data in [16] which describes the singular fiber with a reducible spectral curve. This plays a similar role to the parabolic modules. See [16, Sec. 7.1] for more details.

For any divisor  $D$ , and line bundle  $L$ , we define the space

$$H^0(D, L) = \bigoplus_{p \in \text{supp } D} \mathcal{O}(L)_p / \sim ,$$

where  $s_1 \sim s_2$  if and only if  $\text{ord}_p([s_1] - [s_2]) \geq D_p$ .

Let  $L \in \text{Pic}(\Sigma)$ , define the following subspaces of  $H^0(\Lambda, L^2 K)$ :

$$\mathcal{V}(D, L) := \{s \in H^0(\Lambda, L^2 K) \mid \text{ord}_p(s) = \Lambda_p - D_p, \text{ if } D_p > 0; s(p) = 0, \text{ if } D_p = 0\},$$

$$\mathcal{W}(D, L) = \{s \in H^0(\Lambda, L^2 K) \mid s|_{\text{supp}(\Lambda - D)} = 0\} .$$

One checks that  $\mathcal{W}(D, L) = \cup_{D' \leq D} \mathcal{V}(D', L)$ . Moreover, the space  $\mathcal{V}(D, L)$  is a linear subspace of  $H^0(\Lambda, L^2 K)$  with a hyperplane removed. In addition,  $\mathbb{C}^*$  acts on  $\mathcal{V}(D, L)$  by multiplication, and  $\dim(\mathcal{V}(D, L)/\mathbb{C}^*) = \deg(D) - 1$ .

We define the fibrations

$$p_m : \mathcal{V}(D, m) \longrightarrow \text{Pic}^m(\Sigma), \quad p_m : \mathcal{W}(D, m) \longrightarrow \text{Pic}^m(\Sigma)$$

such that for  $L \in \text{Pic}^m(\Sigma)$ , the fibers are  $\mathcal{V}(D, L)$  and  $\mathcal{W}(D, L)$ .

**7.2.1. Algebraic data from the extension.** The Higgs bundle  $(\mathcal{E}, \varphi)$  can be understood in terms of an extension of exact sequence. As  $\det(\mathcal{E}) = \mathcal{O}$ , we have the exact sequence

$$0 \longrightarrow L_+ \longrightarrow \mathcal{E} \longrightarrow L_+^{-1} \longrightarrow 0.$$

For each  $p \in Z$ , with  $U \subset \Sigma$  a neighborhood of  $p$ ,  $(\mathcal{E}, \varphi)$  can be written as the following splitting of  $\mathcal{C}^\infty$  bundles

$$\mathcal{E} = L_+ \oplus_{\mathcal{C}^\infty} L_+^{-1}, \quad \bar{\partial}_{\mathcal{E}} = \begin{pmatrix} \bar{\partial}_{L_+} & b \\ 0 & \bar{\partial}_{L_+^{-1}} \end{pmatrix}, \quad \varphi = \begin{pmatrix} \omega & c \\ 0 & -\omega \end{pmatrix}.$$

We would like to consider the restriction of  $\varphi + \omega \cdot \text{id}$  to  $\Lambda$ . As  $\omega|_\Lambda = 0$ ,  $\varphi + \omega \cdot \text{id}|_{L_+} = 0$  and the image of  $\varphi + \omega \cdot \text{id} \subset (L_+ \oplus 0) \otimes K \subset \mathcal{E} \otimes K$ . Therefore, the restriction of  $\varphi + \omega \cdot \text{id}$  to  $\Lambda$  defines a holomorphic map  $s : L_+^{-1}|_\Lambda \rightarrow L_+ K|_\Lambda$ , or equivalently a section  $s \in H^0(\Lambda, L_+^2 K)$ . Moreover, by [16, Lemma 7.12],  $\text{Div}(s) = \Lambda - D$ . Therefore, given any  $(\mathcal{E}, \varphi) \in \mathcal{M}_q$ , we obtain an  $L \in \text{Pic}^m(\Sigma)$  and an element in  $\mathcal{V}(D, L)$ . Moreover, the stability condition implies that  $0 \leq D \leq \Lambda$ , we have  $-\deg D \leq \deg L \leq 0$ .

**7.2.2. Inverse construction.** The inverse of the construction above also holds; for further details, see [16, Sec. 7] and [24, Sec. 5]. Given  $L \in \text{Pic}^m(\Sigma)$  and  $q \in \mathcal{V}(D, L)$ , we define a Higgs bundle via extensions as follows. From  $q, L$ , we have a short exact sequence of complexes of sheaves:

$$\begin{array}{ccccccc} & & C_1^* & & C_2^* & & C_3^* \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & L^2 & \xrightarrow{=} & L^2 & \xrightarrow{\text{pr}} & 0 \longrightarrow 0, \\ & & \downarrow \text{id} & & \downarrow c & & \downarrow 0 \\ 0 & \longrightarrow & L^2 & \xrightarrow{c} & L^2 K & \xrightarrow{\text{res}(\Lambda)} & L^2 K|_{\Lambda'} \longrightarrow 0 \end{array}$$

where, for a section  $s \in \Gamma(L^2)$ ,  $c(s) := \sqrt{-1}\omega s$ , and  $\text{res}(\Lambda)$  is the restriction map to the divisor  $\Lambda$ . The long exact sequence in hypercohomology implies that  $\text{res}(\Lambda)$  induces an isomorphism

$$\text{res}(\Lambda) : \mathbf{H}^1(C_2^*) \cong \mathbf{H}^1(C_3^*) = H^0(\Lambda, L^2K).$$

Moreover, we have  $\mathbf{H}^1(C_2^*) \cong H^1(\Sigma, L^2)$ , which parameterizes extensions

$$0 \longrightarrow L \longrightarrow \mathcal{E} \longrightarrow L^{-1} \longrightarrow 0.$$

From  $s \in H^0(\Lambda, L^2K)$  and  $\mathcal{E}$  above, we can find a section  $c \in \Gamma(L^2K)$ , and construct a Higgs bundle

$$(20) \quad E = L \oplus_{\mathcal{C}^\infty} L^{-1}, \quad \bar{\partial}_E = \begin{pmatrix} \bar{\partial}_L & b \\ 0 & \bar{\partial}_{L^{-1}} \end{pmatrix}, \quad \varphi = \begin{pmatrix} \omega & c \\ 0 & -\omega \end{pmatrix},$$

where  $\bar{\partial}c = 2b\omega$  for  $b \in \Omega^{0,1}(L^2)$  and  $c$  is an extension of  $q$ . For  $0 \leq D \leq \Lambda$  and  $-\deg D \leq m \leq 0$ , the construction above defines a map

$$\wp : \mathcal{V}(D, m) \longrightarrow \mathcal{M}_q, \quad s \in \mathcal{V}(D, L) \mapsto (\mathcal{E}, \varphi),$$

where  $(\mathcal{E}, \varphi)$  is the Higgs bundle constructed in (20). When  $D = 0$ ,  $\mathcal{V}(\Lambda, L) = \{0\}$  and the image of  $\wp : \mathcal{V}(\Lambda, 0) \rightarrow \mathcal{M}_q$  are the polystable Higgs bundles  $\mathcal{E} = L \oplus L^{-1}$ ,  $\varphi = \text{diag}(\omega, -\omega)$  such that  $L^2 \cong \mathcal{O}_\Sigma$ .

**Theorem 7.1.** [16, Thm. 7.7] *For  $0 \leq D \leq \Lambda$  and  $-\deg(D) \leq m_1 \leq 0$  and the map  $\wp : \mathcal{V}(D, m_1) \rightarrow \mathcal{M}_q$ , we have*

- (i) *for  $m_2 = -\deg(D) - m_1$ , we have  $\wp(\mathcal{V}(D, m_1)) = \wp(\mathcal{V}(D, m_2))$ ,*
- (ii) *for the  $\mathbb{C}^*$  action on  $\mathcal{V}(D, m_1)$  by multiplication, for  $\xi \in \mathcal{V}(D, m_1)$ ,  $\wp(\mathbb{C}^*\xi) = \wp(\xi)$ ,*
- (iii) *when  $m_1 \neq -\frac{1}{2}\deg(D)$ ,  $\wp : \mathcal{V}(D, m_1)/\mathbb{C}^* \rightarrow \mathcal{M}_q$  is an isomorphism onto its image,*
- (iv) *when  $m_1 = -\frac{1}{2}\deg(D)$ ,  $\wp : \mathcal{V}(D, m_1)/\mathbb{C}^* \rightarrow \mathcal{M}_q$  is a double branched covering, which branched along line bundles  $L \in \text{Pic}^{m_1}(\Sigma)$  such that  $L^2 \cong \mathcal{O}(-D)$ ,*
- (v) *when  $D = 0$ , then  $\wp : \mathcal{V}(\Lambda, 0) \rightarrow \mathcal{M}_q$  is a double branched covering, branched along  $L \in \text{Pic}^0(\Sigma)$  such that  $L^2 \cong \mathcal{O}$ .*

**Example 7.2.** *When  $g = 2$ , for  $q = -\omega \otimes \omega$ , we can write  $\Lambda = p_1 + p_2$  or  $\Lambda = 2p$ . In either case, the  $\mathcal{M}_q^{\text{st}} = \wp(\mathcal{V}(D, m))$  for  $-\deg(D) < m < 0$  and  $0 \leq D \leq \Lambda$ . Therefore,  $m = -1$ ,  $D = \Lambda$  and  $\wp(\mathcal{V}(\Lambda, -1)) = \mathcal{M}_q^{\text{st}}$ . Moreover, generically, the map  $\wp : (\mathcal{V}(\Lambda, -1))/\mathbb{C}^* \rightarrow \mathcal{M}_q^{\text{st}}$  is two-to-one.*

**7.3. The stratification of the singular fiber.** We now present two stratifications of  $\mathcal{M}_q$ . Recall that from any Higgs bundle  $(\mathcal{E}, \varphi)$  we obtain two line bundles  $L_\pm$  and a divisor  $D$ . There are two different stratifications: one given by the divisor  $D$  and the other by the degree of  $L_+$ .

**7.3.1. Divisor stratification.** We first discuss the stratification defined by the divisor. Indeed, using  $D$ , decompose into strata:  $\mathcal{M} = \bigcup_{0 \leq D \leq \Lambda} \mathcal{M}_D$ . As the definition of  $L_\pm$  depends on the choice of the square root, there is no natural map from  $\mathcal{M}_D$  to  $\text{Pic}(\Sigma)$ . Consider the following space:  $\mathbb{V}_D = \bigcup_{-\deg(D) \leq m \leq 0} \mathcal{V}(D, m)$ . This forms a fibration

$$\tau : \mathbb{V}_D \longrightarrow \bigcup_{-\deg(D) \leq m \leq 0} \text{Pic}^m(\Sigma).$$

Moreover, for  $L \in \text{Pic}^m(\Sigma)$ , we have  $\tau^{-1}(L) = \mathcal{V}(D, L)$  and  $\dim(\tau^{-1}(L)/\mathbb{C}^*) = \deg(D) - 1$ . By Theorem 7.1,  $\wp|_{\mathbb{V}_D} : \mathbb{V}_D \rightarrow \mathcal{M}_D$  is surjective. Moreover, since

$$\wp|_{\mathcal{V}(D, m)} = \wp|_{\mathcal{V}(D, -\deg(D) - m)}$$

generically,  $\wp|_{\mathbb{V}_D}$  is a two-to-one map.

In summary, we obtain the following map which characterizes the singular fiber.

$$\wp : \mathbb{V} = \bigcup_{0 \leq D \leq \Lambda} \mathbb{V}_D \rightarrow \mathcal{M}_q = \bigcup_{0 \leq D \leq \Lambda} \mathcal{M}_D.$$

The top stratum is given by  $D = \Lambda$ .

**7.3.2. Degree stratification.** We next introduce the stratification defined by degrees; this encodes how different divisor stratifications are pasted together.

For  $-(2g-2) \leq m \leq 0$  and  $L \in \text{Pic}^m(\Sigma)$ , define  $\mathbb{W}(L) := \bigcup_{\deg D \geq -m} \mathcal{V}(D, L)$ . This set is connected, based on the definition and [16, Lemma 7.14]. Moreover, if we define

$$\mathbb{W}_m := \bigcup_{-m \leq \deg D, 0 \leq D \leq \Lambda} \mathcal{V}(D, m), \quad \mathbb{W} := \bigcup_{-(2g-2) \leq m \leq 0} \mathbb{W}_m,$$

then we have  $\wp(\mathbb{W}) = \wp(\mathbb{V})$ .

We should also note that though  $\mathbb{W}_m \cap \mathbb{W}_n = \emptyset$  for any  $m \neq n$ ,  $\mathbb{W}$  is connected. As  $L_+, L_-$  are symmetric, by Theorem 7.1, we have  $\wp(\mathcal{V}(D, m)) = \wp(\mathcal{V}(D, -\deg(D) - m))$ , which implies that for any integer  $-(2g-2+m) \leq n \leq 0$ ,  $\wp\mathbb{W}_m \cap \wp\mathbb{W}_n \neq \emptyset$ .

We now give an example of the degree stratification when  $g = 2$ .

**Example 7.3.** Suppose  $\omega$  has only one zero with order 2. Then  $\Lambda = 2p$ , and all possible divisors are  $D_2 = 2p, D_1 = p, D_0 = 0$ . The degree stratification is

$$\begin{aligned} \mathbb{W}_{-2} &= \mathcal{V}(D_2, -2), \quad \mathbb{W}_{-1} = \mathcal{V}(D_2, -1) \cup \mathcal{V}(D_1, -1) \\ \mathbb{W}_0 &= \mathcal{V}(D_0, 0) \cup \mathcal{V}(D_1, 0) \cap \mathcal{V}(D_2, 0). \end{aligned}$$

The image of  $\wp(\mathcal{V}(D_2, -1))$  is stable,  $\wp(\mathcal{V}(D_0, 0))$  is poly-stable and  $\wp(\mathcal{W} \setminus (\mathcal{V}(D_2, -1) \cup \mathcal{V}(D_0, 0)))$  is semistable.

Moreover, we have  $\wp(\mathcal{V}(D_2, -2)) = \wp(\mathcal{V}(D_2, 0))$ ,  $\wp(\mathcal{V}(D_1, -1)) = \wp(\mathcal{V}(D_1, 0))$  and  $\wp|_{\mathcal{V}(D_2, -1)}$  is a branched covering. Moreover, we have  $\wp(\mathcal{V}(D_2, -1)) \cap \wp(\mathcal{V}(D_1, 0)) \neq 0$  and  $\wp(\mathcal{V}(D_2, -1)) \cap \wp(\mathcal{V}(D_0, 0)) = 0$ .

**7.4. Algebraic Mochizuki map.** Based on the study of the local rescaling properties of Higgs bundles, Mochizuki introduced a weight for each  $p \in Z$  in [33, Sec. 3]. To be more specific, let  $c$  be a real number. For each  $p \in Z$ , the weight we consider is given by  $\chi_p(c) = \min\{\ell_p, (m_p + 1)c + \ell_p/2\}$ .

By utilizing the global geometry of a Higgs bundle, we can uniquely determine the constant  $c$ . We aim to choose the sign of  $\omega$  such that  $d_+ \leq d_-$ .

**Lemma 7.4** ([33, Lemma 4.3]). *If  $(\mathcal{E}, \varphi)$  is stable, then there exists a unique constant  $c > 0$  such that*

$$d_+ + \sum_{p \in Z} \chi_p(c) = 0, \quad d_- + \sum_{p \in Z} (\ell_p - \chi_p(c)) = 0.$$

*Proof.* Since  $(\mathcal{E}, \varphi)$  is stable, we have  $-\sum \ell_p < d_{\pm} < 0$ . We define the function

$$(21) \quad f(c) = d_+ + \sum_p \chi_p(c),$$

which is strictly increasing. Moreover, for  $c$  sufficiently large,  $\chi_p(c) = \ell_p$ , and therefore  $f(c) = d_+ + \sum_p \ell_p = -d_- > 0$ . Additionally,  $f(0) = d_+ + \sum_p (\ell_p/2)$ . Since  $d_+ \leq d_-$  and  $d_+ + d_- + \sum_p \ell_p = 0$ , we obtain  $f(0) \leq 0$ . The monotonicity of  $f$  implies the existence of  $c_0$  such that  $f(c_0) = 0$ .  $\square$

From the construction, if  $d_+ \leq d_-$ , two weighted bundles  $(L_+, \chi_p(c_0))$  and  $(L_-, \ell_p - \chi_p(c_0))$  are obtained with weights  $\chi_p(c_0)$  and  $\ell_p - \chi_p(c_0)$  at each  $p \in Z$ , respectively. On the other hand, if  $d_+ \geq d_-$ , by symmetry, weighted bundles  $(L_+, \ell_p - \chi_p(c_0))$  and  $(L_-, \chi_p(c_0))$  are obtained. When  $(\mathcal{E}, \varphi)$  is strictly semistable, S-equivalent to  $(L, \omega) \oplus (L^{-1}, -\omega)$ , then we would like to consider the weighted bundles  $(L, 0) \oplus (L^{-1}, 0)$  with weight zero.

Next, we define the algebraic Mochizuki map. Let  $\mathcal{F}_\pm(\Sigma)$  be the space of rank 1 degree zero filtered bundles on  $\Sigma$ , and let  $\mathcal{F}_2(\Sigma) := \mathcal{F}_+(\Sigma) \oplus \mathcal{F}_-(\Sigma)$  be the direct sum. Fix a choice of  $\omega$ . Then from any Higgs bundle  $(\mathcal{E}, \varphi)$ , we obtain the subbundles  $L_\pm$  with degree  $d_\pm$  and define the algebraic Mochizuki map

$$\begin{aligned} \Theta^{\text{Moc}} : \mathcal{M}_q &\longrightarrow \mathcal{F}_2(\Sigma), \\ \Theta^{\text{Moc}}(\mathcal{E}, \varphi) &:= \begin{cases} \mathcal{F}_*(L_+, \chi_p(c_0)) \oplus \mathcal{F}_*(L_-, \ell_p - \chi_p(c_0)), & \text{if } d_+ \leq d_- \\ \mathcal{F}_*(L_+, \ell_p - \chi_p(c_0)) \oplus \mathcal{F}_*(L_-, \chi_p(c_0)), & \text{if } d_- \leq d_+ \end{cases}, \quad (\mathcal{E}, \varphi) \text{ stable}, \\ \Theta^{\text{Moc}}(\mathcal{E}, \varphi) &:= \mathcal{F}_*(L, 0) \oplus \mathcal{F}_*(L^{-1}, 0), \quad (\mathcal{E}, \varphi) \text{ semistable}. \end{aligned}$$

We list some properties of this map.

**Proposition 7.5.** *For  $\Theta^{\text{Moc}}$ , we have:*

- (i) *for each  $\mathcal{V}(D, m)$  with  $0 \leq D \leq \Lambda$ ,  $-\deg(D) \leq m \leq 0$ ,  $\Theta^{\text{Moc}}|_{\varphi(\mathcal{V}(D, m))}$  is continuous,*
- (ii) *for  $i=1, 2$  and  $s_i \in \mathbb{V}_D$  with  $(\mathcal{E}_i, \varphi_i) := \wp(s_i)$ , suppose  $\tau(s_1) = \tau(s_2)$ , then  $\Theta^{\text{Moc}}(\mathcal{E}_1, \varphi_1) = \Theta^{\text{Moc}}(\mathcal{E}_2, \varphi_2)$ . In particular,  $\Theta^{\text{Moc}}$  is not injective.*

*Proof.* The proof follows directly from the definition. □

**7.5. Exotic phenomena.** A Higgs bundle  $(\mathcal{E}, \varphi) \in \mathcal{M}_q$  is called “exotic” if the constant  $c$  in Lemma 7.4 satisfies  $c \neq 0$ . This new behavior only appears in the Hitchin fiber with reducible spectral curve. In this subsection, we aim to understand the exotic phenomenon of a Higgs bundle.

We first provide another expression for the weight in Lemma 7.4.

**Proposition 7.6.** *Suppose  $-\deg D < d_+ \leq -\frac{1}{2} \deg D$  and let  $Z_0 := \text{supp}(D)$ , then the constant  $c$  in Lemma 7.4 is given by*

$$c = \frac{d_+ + \frac{1}{2} \deg D}{\deg(\Lambda|_{Z_0}) + |Z_0|},$$

where  $\Lambda|_{Z_0}$  means the restriction of the divisor to  $Z_0$ , and  $|Z_0|$  means the number of points in  $Z_0$  without multiplicity.

*Proof.* By the definition of  $\chi_p(c)$ , if  $p \in Z_1$ ,  $\chi_p(c) = 0$  for any  $c \geq 0$ . Therefore, the choice of  $c$  is determined by the equation  $d_+ + \sum_{p \in Z_0} \min\{\ell_p, (m_p + 1)c + \frac{\ell_p}{2}\} = 0$ . We define the function  $F(c) := d_+ + \sum_{p \in Z_0} ((m_p + 1)c + \ell_p/2)$ , and let  $f(c)$  be the function defined in (21), then  $F(c) \geq f(c)$ ,  $F(0) = 0$  and for  $c$  sufficiently large,  $F(c) = f(c)$ .

We compute

$$\sum_{p \in Z_0} (m_p + 1)c = (\deg(\Lambda|_{Z_0}) + |Z_0|)c, \quad \sum_{p \in Z_0} (\ell_p/2) = \frac{1}{2} \deg D.$$

Then for

$$c_0 = \frac{d_+ + \frac{1}{2} \deg D}{\deg(\Lambda|_{Z_0}) + |Z_0|}$$

we have  $F(c_0) = 0$ . Therefore,  $f(c_0) = 0$ , which determines the choice of the constant  $c$  in Lemma 7.4. □

**Corollary 7.7.** *A Higgs bundle  $(\mathcal{E}, \varphi)$  is exotic if and only if its corresponding degrees  $d_{\pm}$  satisfy  $d_+ \neq d_-$ . Additionally, there are only a finite number of possible choices for  $c$  over  $\mathcal{M}_q$ .*

*Proof.* The result follows directly from the formulas in Proposition 7.6.  $\square$

We will compute some examples of possible weights in some special cases. First consider the generic case.

**Example 7.8.** *Suppose  $\omega$  has only simple zeros, therefore  $Z = \{p_1, \dots, p_{2g-2}\}$  and  $\Lambda = p_1 + \dots + p_{2g-2}$ . Given a stable Higgs bundle  $(\mathcal{E}, \varphi)$  with corresponding  $L_+, L_-, D, d_{\pm}$ , recall that from the stability condition, these degrees satisfy the followings:*

$$d_+ + d_- + \deg D = 0, \quad d_+ \leq d_-, \quad d_+ < 0, \quad d_- < 0, \quad -\deg D < d_+ \leq -\frac{1}{2} \deg D.$$

If we write  $Z_0 := \text{supp } D$ , then  $\chi_p = 0$  for  $p \in Z \setminus Z_0$  and  $\chi_p = 2c + 1/2$  for  $p \in Z_0$ . Moreover, we have  $d_+ + \deg(D)(2c + 1/2) = 0$ . Therefore,  $c = -\frac{d_+}{2\deg(D)} + \frac{1}{4}$ , and two weighted bundles we obtained are

$$\left(L_+, \left(-\frac{d_+}{\deg D} \Big|_{Z_0}, 0 \Big|_{Z \setminus Z_0}\right)\right), \quad \left(L_-, \left(1 + \frac{d_+}{\deg D} \Big|_{Z_0}, 1 \Big|_{Z \setminus Z_0}\right)\right).$$

Next, we consider the most nongeneric case.

**Example 7.9.** *Suppose  $\omega$  only contains one zero. Write  $\text{Div}(\omega) = (2g - 2)p$ . Then the possible divisors are  $D = \ell p$ , for  $0 \leq \ell \leq 2g - 2$ . Let  $L_+$  be a line bundle with  $d_+ := \deg(L_+)$  and  $-\ell < d_+ \leq \ell/2$ . Then the choice of  $c$  is determined by the equation*

$$d_+ + \min\{\ell, (2g - 1)c + \ell/2\} = 0.$$

As  $d_+ \neq -\ell$ , we must have  $c = -\frac{\ell}{2(2g-1)}$  and the corresponding weighted bundles are

$$(L_+, -d_+), \quad (L_+^{-1} \otimes \mathcal{O}(-\ell p), \ell - d_+).$$

Finally, we explicitly compute the limiting configurations when  $g = 2$  for the strictly semistable locus of the stratification in Example 7.3.

**Example 7.10.** *When  $g = 2$ , we consider  $\Lambda = 2p$  and  $D_i = ip$  for  $i = 0, 1, 2$ . Let  $p_m : \mathcal{V}(D, m) \rightarrow \text{Pic}^m(\Sigma)$  be the projection. Over  $\wp(\mathcal{V}(D_i, 0))$ , for  $L \in \text{Pic}^0(\Sigma)$ , the  $S$ -equivalence class for  $\wp(p_0^{-1}(L))$  is  $(L \oplus L^{-1}, \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix})$  and the corresponding weighted bundles are  $(L, 0) \oplus (L^{-1}, 0)$ . Over  $\wp(\mathcal{V}(D_2, -1))$ , for  $L \in \text{Pic}^{-1}(\Sigma)$ , let  $(\mathcal{E}, \varphi) = \wp(L, s \in \mathcal{V}(D_2, L))$ , then*

$$\Theta^{\text{Moc}}(\mathcal{E}, \varphi) = \mathcal{F}_*(L, 1) \oplus \mathcal{F}_*(L^{-1}(-D_2), 1) = \mathcal{F}_*(L(p), 0) \oplus \mathcal{F}_*(L^{-1}(p - D_2), 0).$$

**7.6. Discontinuous behavior.** In this subsection, we study the discontinuous behavior of  $\Theta^{\text{Moc}}$ . Consider a sequence of algebraic data  $(L_i, q_i) \in \mathbb{W}_m$ , where  $L_i \in \text{Pic}^m$  and  $q_i \in \mathcal{V}(D, L_i)$ . We assume that  $\lim_{i \rightarrow \infty} L_i = L_{\infty}$  in  $\text{Pic}^m$  and  $\lim_{i \rightarrow \infty} q_i = q_{\infty} \in \mathcal{V}(D_{\infty}, L)$ , for  $D_{\infty} \neq D$ . As the space  $\bigcup_{\deg D' \geq -m} \mathcal{V}(D', m)$  is connected, we can always find such a sequence.

We define  $L_+^i := L_i$  and  $L_-^i := L_i^{-1} \otimes \mathcal{O}(-D)$ , then by Lemma 7.4, the weight function we choose is independent of  $i$ , which we denote as  $\chi_{\pm}$ . In addition, we have

$$\lim_{i \rightarrow \infty} \Theta^{\text{Moc}} \circ \wp(L_i, q_i) = \mathcal{F}_*(L_{\infty}, \chi_+) \oplus \mathcal{F}_*(L_{\infty}^{-1}(-D), \chi_-).$$

For  $(L_\infty, q_\infty \in \mathcal{V}(D_\infty, L))$ , we write  $\chi_\pm^\infty$  be the corresponding weights, which only depends on  $D_\infty$  and  $m$ , then

$$\Theta^{\text{Moc}} \circ \wp(L_\infty, q_\infty) = \mathcal{F}_*(L_\infty, \chi_+^\infty) \oplus \mathcal{F}_*(L_\infty^{-1} \otimes \mathcal{O}(-D_\infty), \chi_-^\infty).$$

Therefore, we obtain

$$(22) \quad \begin{aligned} & \lim_{i \rightarrow \infty} \Theta^{\text{Moc}} \circ \wp(L_i, q_i) \\ &= \Theta^{\text{Moc}} \circ \wp(L_\infty, q_\infty) \otimes (\mathcal{F}_*(\mathcal{O}, \chi_+ - \chi_+^\infty) \oplus \mathcal{F}_*(\mathcal{O}(D_\infty - D), \chi_- - \chi_-^\infty)). \end{aligned}$$

**Proposition 7.11.** *When  $g \geq 3$ , there exists a sequence  $(\mathcal{E}_i, \varphi_i) \in \mathcal{M}_q$  of stable Higgs bundles with stable limit  $(\mathcal{E}_\infty, \varphi_\infty) = \lim_{i \rightarrow \infty} (\mathcal{E}_i, \varphi_i)$  such that*

$$\lim_{i \rightarrow \infty} \Theta^{\text{Moc}}(\mathcal{E}_i, \varphi_i) \neq \Theta^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty) .$$

*Proof.* We choose  $D = \Lambda$  and  $d_+ = -(g-1)$  with  $L_i = L \in \text{Pic}^{d_+}(\Sigma)$ , and study the degenerate behavior for a family  $q_i \in \mathcal{V}(\Lambda, L)$  which converges to  $q_\infty \in \mathcal{V}(D_\infty, L)$ . Here,  $D_\infty$  satisfies  $D_\infty \leq D$  and  $\deg(D_\infty) = \deg(D) - 1$ . As  $q_i$  lies in the top stratum, we can always find such a family. Take  $(\mathcal{E}_i, \varphi_i) = \wp(L_i, q_i)$  and  $(\mathcal{E}_\infty, \varphi_\infty) = \wp(L, q_\infty)$ . When  $g \geq 3$ , we have  $-\deg(D_\infty) < d_+ \leq -\frac{1}{2} \deg(D_\infty)$ , which implies  $(\mathcal{E}_\infty, \varphi_\infty)$  is a stable Higgs bundle.

Write  $D = \sum_p \ell_p$ . As  $(\mathcal{E}_i, \varphi_i)$  is nonexotic, the weights will be  $\chi_+(p) = \chi_-(p) = \ell_p/2$ . However, as  $\deg(D_\infty) \neq 2d_+$ ,  $(\mathcal{E}_\infty, \varphi_\infty)$  is exotic. By Proposition 7.6, if we write  $\chi_\pm^\infty(p)$  for the weight functions with corresponding constant  $c$ , then  $c > 0$ . Therefore, for  $p \neq p_0$ , we have  $\chi_+^\infty(p) = (m_p + 1)c + m_p/2 > m_p/2 = \chi_+(p)$ . By (22),  $\lim_{i \rightarrow \infty} \Theta^{\text{Moc}}(\mathcal{E}_i, \varphi_i) \neq \Theta^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty)$ .  $\square$

When  $g = 2$ , the stratification is simpler, and we have the following.

**Proposition 7.12.** *When  $g = 2$ , the following holds:*

- (i) *Suppose  $\Lambda = p_1 + p_2$  for  $p_1 \neq p_2$ , then  $\Theta^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is continuous. Moreover, there exists a sequence of stable Higgs bundles  $(\mathcal{E}_i, \varphi_i) \in \mathcal{M}_q$  where the limit  $(\mathcal{E}_\infty, \varphi_\infty) = \lim_{i \rightarrow \infty} (\mathcal{E}_i, \varphi_i)$  is semistable, and  $\gamma(0)$  is also semistable and furthermore*

$$\lim_{i \rightarrow \infty} \Theta^{\text{Moc}}(\mathcal{E}_i, \varphi_i) \neq \Theta^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty) .$$

- (ii) *Suppose  $\Lambda = 2p$ . Then  $\Theta^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is continuous.*

*Proof.* For (i), suppose  $\Lambda = p_1 + p_2$ , then by Example 7.2, we have  $\mathcal{M}_q^{\text{st}} = \wp(\mathcal{V}(\Lambda, -1))$ . By Proposition 7.5,  $\Theta_q^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is continuous.

However, for semistable elements, other strata must be taken into consideration. Take  $L \in \text{Pic}^{-1}(\Sigma)$  and  $q_i \in \mathcal{V}(\Lambda, L)$  such that  $q_i$  convergence to  $q_\infty \in \mathcal{V}(p_1, L)$ . We define  $(\mathcal{E}_i, \varphi_i) = \wp(L, q_i)$  and  $(\mathcal{E}_\infty, \varphi_\infty) = \wp(L, q_\infty)$ . For each  $i$ ,

$$\Theta^{\text{Moc}}(\mathcal{E}_i, \varphi_i) = \mathcal{F}_*(L, (\frac{1}{2}, \frac{1}{2})) \oplus \mathcal{F}_*(L^{-1}(-\Lambda), (\frac{1}{2}, \frac{1}{2})).$$

Moreover, we have

$$\Theta^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty) = \mathcal{F}_*(L(D), (0, 0)) \oplus \mathcal{F}_*(L^{-1}(-D), (0, 0)) \neq \lim_{i \rightarrow \infty} \Theta^{\text{Moc}}(\mathcal{E}_i, \varphi_i).$$

For (ii), by Example 7.3,  $\wp(\mathcal{V}(D_2, -1)) = \mathcal{M}_q^{\text{st}}$  and by Proposition 7.5,  $\Theta_q^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is continuous. We now consider the behavior of the filtered bundle when crossing the divisors.  $\square$

**7.7. The analytic Mochizuki map and limiting configurations.** In this subsection, we construct the analytic Mochizuki map for the Hitchin fiber with a reducible spectral curve. We also introduce the convergence theorem of Mochizuki as stated in [33] and examine the discontinuous behavior of the analytic Mochizuki map.

For  $(\mathcal{E}, \varphi) \in \mathcal{M}_q$ , we can express the abelianization as  $(\mathcal{E}_0, \varphi_0) = (L_+ \oplus L_-, \begin{pmatrix} \omega & 0 \\ 0 & -\omega \end{pmatrix})$ , thus  $\Theta^{\text{Moc}}(\mathcal{E}, \varphi) = \mathcal{F}_*(L_+, \chi_+) \oplus \mathcal{L}_-(L_-, \chi_-) \in \mathcal{F}_2(\Sigma)$ . Via the nonabelian Hodge correspondence for filtered bundles, we obtain two Hermitian metrics  $h_{\pm}^{\text{Lim}}$  with corresponding Chern connections  $A_{h_{\pm}^{\text{Lim}}}$ . These metrics satisfy the following proposition.

**Proposition 7.13.** [32, Lemma 4.4] *The metrics  $h_{\pm}^{\text{Lim}}$  over  $L_{\pm}$  satisfy*

- i)  $F_{A_{h_{\pm}^{\text{Lim}}}} = 0$  and  $h_+^{\text{Lim}} h_-^{\text{Lim}} = 1$ ,
- ii) for every  $p \in \Sigma$ , there exists an open neighborhood  $(U, z)$  with  $P = \{z = 0\}$  such that  $|z|^{-2\chi_p(\text{co})} h_+^{\text{Lim}}$  and  $|z|^{2\chi_p(\text{co})+2l_P} h_-^{\text{Lim}}$  extends smoothly to  $L_{\pm}|_U$ .

Now,  $H^{\text{Lim}} := h_+^{\text{Lim}} \oplus h_-^{\text{Lim}}$  is a metric on  $\mathcal{E}_0$  which induces a metric on  $(\mathcal{E}, \varphi)|_{\Sigma \setminus Z}$  because  $(\mathcal{E}, \varphi)|_{\Sigma \setminus Z} \cong (\mathcal{E}_0, \varphi_0)|_{\Sigma \setminus Z}$ . Let  $(A^{\text{Lim}}, \phi^{\text{Lim}})$  be the Chern connection defined by  $(\mathcal{E}, \varphi, H^{\text{Lim}})$  over  $\Sigma \setminus Z$ . Then  $(A^{\text{Lim}}, \phi^{\text{Lim}})$  is a limiting configuration that satisfies the decoupled Hitchin equations (7). The analytic Mochizuki map  $\Upsilon^{\text{Moc}}$  is defined as:

$$(23) \quad \Upsilon^{\text{Moc}} : \mathcal{M}_q \longrightarrow \mathcal{M}_{\text{Hit}}^{\text{Lim}}, \quad \Upsilon^{\text{Moc}}(\mathcal{E}, \varphi) = (A^{\text{Lim}}, \phi^{\text{Lim}}).$$

Note that  $H^{\text{Lim}}$  is not unique: for any constant  $c$ , the metric  $ch_+^{\text{Lim}} \oplus c^{-1}h_-^{\text{Lim}}$  defines the same Chern connection as  $H^{\text{Lim}}$ . In any case, the map  $\Upsilon^{\text{Moc}}$  is well-defined.

Suppose  $(\mathcal{E}, \varphi)$  is an S-equivalence class of a semistable Higgs bundle. Let  $H_t$  be the harmonic metric for  $(\mathcal{E}, t\varphi)$ . For each constant  $C > 0$ , define  $\mu_C$  to be the automorphism of  $L_+ \oplus L_-$  given by  $\mu_C = C \text{id}_{L_+} \oplus C^{-1} \text{id}_{L_-}$ . As  $\mathcal{E} \cong L_+ \oplus L_-$  on  $\Sigma \setminus Z$ ,  $\mu_C^* H_t$  can be regarded as a metric on  $\mathcal{E}|_{\Sigma \setminus Z}$ . Take any point  $x \in \Sigma \setminus Z$  and a frame  $e_x$  of  $L_+|_x$ , and define:

$$C(x, t) := \left( \frac{h_{L_+}^{\text{Lim}}(e_x, e_x)}{H_t(e_x, e_x)} \right)^{1/2}.$$

Writing  $\nabla_t + t\phi_t$  as the corresponding flat connection of  $(\mathcal{E}, t\varphi)$  under the nonabelian Hodge correspondence, then

**Theorem 7.14.** [33] *On any compact subset  $K$  of  $\Sigma \setminus Z$ ,  $\mu_{C(x,t)}^* H_t$  converges smoothly to  $H^{\text{Lim}}$ . Additionally, there exist  $t$ -independent constants  $C_k$  and  $C'_k$  such that*

$$|(\nabla_t, \phi_t) - \Upsilon^{\text{Moc}}(\mathcal{E}, \varphi)|_{C^k} \leq C_k e^{-C'_k d}.$$

Propositions 7.11 and 7.12 now give

**Theorem 7.15.** *When  $g \geq 3$ ,  $\Upsilon^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is discontinuous, and when  $g = 2$ ,  $\Upsilon^{\text{Moc}}|_{\mathcal{M}_q^{\text{st}}}$  is continuous.*

## 8. THE COMPACTIFIED KOBAYASHI-HITCHIN MAP

In this section, we define a compactified version of the Kobayashi-Hitchin map and prove the main theorem of our paper. The original Kobayashi-Hitchin map  $\Xi$  is a homeomorphism between the Dolbeault moduli space  $\mathcal{M}_{\text{Dol}}$  and the Hitchin moduli space  $\mathcal{M}_{\text{Hit}}$ . We wish to extend this to a map  $\overline{\Xi}$  from the compactified Dolbeault moduli space  $\overline{\mathcal{M}}_{\text{Dol}}$  to the compactification  $\overline{\mathcal{M}}_{\text{Hit}} \subset \mathcal{M}_{\text{Hit}} \cup \mathcal{M}_{\text{Hit}}^{\text{Lim}}$  of the Hitchin moduli space, and to study the properties of this extended map.



**8.1. The compactified Kobayashi-Hitchin map.** We first summarize the results obtained above. By the construction in Section 4, there is an identification  $\partial\overline{\mathcal{M}}_{\text{Dol}} \cong (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))/\mathbb{C}^*$ . Moreover, through (18) and (23), we have constructed the analytic Mochizuki map  $\Upsilon^{\text{Moc}} : \mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0) \rightarrow \mathcal{M}_{\text{Hit}}^{\text{Lim}}$ . Writing  $(A^{\text{Lim}}, \phi^{\text{Lim}} = \varphi + \varphi^{\dagger\text{Lim}}) = \Upsilon^{\text{Moc}}(\mathcal{E}, \varphi)$ , then for  $w \in \mathbb{C}^*$ , we have

$$\Upsilon^{\text{Moc}}(\mathcal{E}, w\varphi) = (A^{\text{Lim}}, \phi^{\text{Lim}} = w\varphi + \bar{w}\varphi^{\dagger\text{Lim}}).$$

Hence  $\Upsilon^{\text{Moc}}$  descends to a map  $\partial\overline{\Xi}$  between  $\mathbb{C}^*$  orbits.

$$\partial\overline{\Xi} : \partial\overline{\mathcal{M}}_{\text{Dol}} = (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))/\mathbb{C}^* \longrightarrow \mathcal{M}_{\text{Hit}}^{\text{Lim}}/\mathbb{C}^*,$$

Together with the initial Kobayashi-Hitchin map  $\Xi : \mathcal{M}_{\text{Dol}} \rightarrow \mathcal{M}_{\text{Hit}}$ , we obtain

$$\overline{\Xi} : \overline{\mathcal{M}}_{\text{Dol}} = \mathcal{M}_{\text{Dol}} \cup \partial\overline{\mathcal{M}}_{\text{Dol}} \longrightarrow \mathcal{M}_{\text{Hit}} \cup \mathcal{M}_{\text{Hit}}^{\text{Lim}}/\mathbb{C}^*.$$

Theorems 6.14 and 7.14 show that for a Higgs bundle  $(\mathcal{E}, \varphi) \in \mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0)$  and real  $t$ ,  $\lim_{t \rightarrow \infty} \Xi(\mathcal{E}, t\varphi) = \partial\overline{\Xi}[(\mathcal{E}, \varphi)/\mathbb{C}^*]$ . Thus the image of  $\overline{\Xi}$  lies in  $\overline{\mathcal{M}}_{\text{Hit}}$ , the closure of  $\mathcal{M}_{\text{Hit}}$  in  $\mathcal{M}_{\text{Hit}} \cup \mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0)$ .

There are natural compactified Hitchin maps  $\overline{\mathcal{H}}_{\text{Dol}} : \overline{\mathcal{M}}_{\text{Dol}} \rightarrow \overline{\mathcal{B}}$  and  $\overline{\mathcal{H}}_{\text{Hit}} : \overline{\mathcal{M}}_{\text{Hit}} \rightarrow \overline{\mathcal{B}}$  such that  $\overline{\mathcal{H}}_{\text{Hit}} \circ \overline{\Xi} = \overline{\mathcal{H}}_{\text{Dol}}$ .

In summary, there are commutative diagrams

$$\begin{array}{ccc} \mathcal{M}_{\text{Dol}} & \xrightarrow{\Xi} & \mathcal{M}_{\text{Hit}} \\ \downarrow & & \downarrow \\ \overline{\mathcal{M}}_{\text{Dol}} & \xrightarrow{\overline{\Xi}} & \overline{\mathcal{M}}_{\text{Hit}} \end{array}, \quad \begin{array}{ccc} \overline{\mathcal{M}}_{\text{Dol}} & \xrightarrow{\overline{\Xi}} & \overline{\mathcal{M}}_{\text{Hit}} \\ & \searrow \overline{\mathcal{H}}_{\text{Dol}} & \downarrow \overline{\mathcal{H}}_{\text{Hit}} \\ & & \overline{\mathcal{B}} \end{array}.$$

We now turn to the analysis of some properties of the compactified Kobayashi-Hitchin map. Define

$$\overline{\mathcal{B}}^{\text{reg}} = \{[(q, w)] \in \overline{\mathcal{B}} \mid 0 \neq q \text{ has simple zeros}\};$$

this represents the compactified space of quadratic differentials with simple zeros. Let  $\overline{\mathcal{B}}^{\text{sing}} = \overline{\mathcal{B}} \setminus \overline{\mathcal{B}}^{\text{reg}}$ . Additionally, define the open sets  $\overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} = \overline{\mathcal{H}}_{\text{Dol}}^{-1}(\overline{\mathcal{B}}^{\text{reg}})$  and  $\overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}} = \overline{\mathcal{H}}_{\text{Hit}}^{-1}(\overline{\mathcal{B}}^{\text{reg}})$  as the collections of elements with regular spectral curves. Now set  $\overline{\mathcal{M}}_{\text{Dol}}^{\text{sing}} = \overline{\mathcal{H}}_{\text{Dol}}^{-1}(\overline{\mathcal{B}}^{\text{sing}})$  and  $\overline{\mathcal{M}}_{\text{Hit}}^{\text{sing}} = \overline{\mathcal{H}}_{\text{Hit}}^{-1}(\overline{\mathcal{B}}^{\text{sing}})$  to be the sets of singular fibers. We can write  $\overline{\Xi} = \overline{\Xi}^{\text{reg}} \cup \overline{\Xi}^{\text{sing}}$ , where

$$\overline{\Xi}^{\text{reg}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} \longrightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}, \quad \overline{\Xi}^{\text{sing}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{sing}} \longrightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{sing}}.$$

**Proposition 8.1.** *The map  $\overline{\Xi}^{\text{reg}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}$  is bijective, whereas  $\overline{\Xi}^{\text{sing}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{sing}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{sing}}$  is neither surjective nor injective.*

*Proof.* The bijectivity of  $\overline{\Xi}^{\text{reg}}$  is established by Theorem 4.9. The nonsurjectivity and non-injectivity of  $\overline{\Xi}^{\text{sing}}$  follow from Theorem 6.15 and Theorem 7.15.  $\square$

**8.2. Continuity properties of the compactified Kobayashi-Hitchin map.** In this subsection, we prove that the continuity of the compactified Kobayashi-Hitchin map is fully determined by the continuity of the analytic Mochizuki map.

Let  $(\mathcal{E}_i, t_i\varphi_i)$  be a sequence of Higgs bundles with real numbers  $t_i \rightarrow +\infty$ ,  $\det(\varphi_i) = q_i$ ,  $Z_i = q_i^{-1}(0)$ , and  $\|q_i\|_{L^2} = 1$ . We denote  $\xi_i = [(\mathcal{E}_i, t_i\varphi_i)] \in \mathcal{M}_{\text{Dol}}$ . By the compactness of  $\overline{\mathcal{M}}_{\text{Dol}}$ , after passing to a subsequence, we may assume there is  $\xi_\infty \in \partial\overline{\mathcal{M}}_{\text{Dol}}$  such that  $\lim_{i \rightarrow \infty} \xi_i = \xi_\infty$ . Since  $\partial\overline{\mathcal{M}}_{\text{Dol}} \cong (\mathcal{M}_{\text{Dol}} \setminus \mathcal{H}^{-1}(0))/\mathbb{C}^*$ , we can select a representative  $(\mathcal{E}_\infty, \varphi_\infty)$  of  $\xi_\infty$ . By Lemma 4.4, we have that  $(\mathcal{E}_i, \varphi_i)$  converges to  $(\mathcal{E}_\infty, \varphi_\infty)$  in  $\mathcal{M}_{\text{Dol}}$ , and  $q_i$  converges to  $q_\infty$ . We write  $Z_\infty = q_\infty^{-1}(0)$ .

By Proposition 4.6,  $\lim_{i \rightarrow \infty} \overline{\Xi}(\mathcal{E}_i, t_i \varphi_i)$  exists. The following result establishes the continuity of this map with respect to the analytic Mochizuki map  $\Upsilon^{\text{Moc}}$ .

**Proposition 8.2.** *Under the previous convention,  $\lim_{i \rightarrow \infty} \overline{\Xi}(\xi_i) = \overline{\Xi}(\xi_\infty)$  if and only if  $\lim_{i \rightarrow \infty} \Upsilon^{\text{Moc}}(\mathcal{E}_i, \varphi_i) = \Upsilon^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty)$ . In other words,  $\overline{\Xi}$  is continuous at  $\xi_\infty$  if and only if  $\Upsilon^{\text{Moc}}$  is continuous at  $\xi_\infty$ .*

*Proof.* Set  $\overline{\Xi}(\mathcal{E}_i, t_i \varphi_i) = \Xi(\mathcal{E}_i, t_i \varphi_i) = A_i + t_i \phi_i$ ,  $\Upsilon^{\text{Moc}}(\mathcal{E}_i, \varphi_i) = (A_i^{\text{Lim}}, \phi_i^{\text{Lim}})$ , and  $\Upsilon^{\text{Moc}}(\mathcal{E}_\infty, \varphi_\infty) = (A_\infty^{\text{Lim}}, \phi_\infty^{\text{Lim}})$ . By Proposition 4.6, there exists a limiting configuration  $(A_\infty, \phi_\infty) := \lim_{i \rightarrow \infty} (A_i, \phi_i)$  over  $\Sigma \setminus Z_\infty$ . Let  $K$  be any compact set in  $\Sigma \setminus Z_\infty$ . Note that by the convergence assumption, there exists  $i_0$  such that  $Z_i \cap K = \emptyset$  for all  $i \geq i_0$ . Moreover, from Theorems 6.14 and 7.14, for  $d_i = \min_K |q_i|$ , there exist  $t$ -independent constants  $C, C' > 0$  such that (up to gauge transformations)

$$|(A_i, \phi_i) - (A_i^{\text{Lim}}, \phi_i^{\text{Lim}})|_{\mathcal{C}^k(K)} \leq C e^{-C' t_i d_i}.$$

The convergence is uniform and exponential for fixed  $K$ . Therefore, over  $K$ , the size of  $|(A_\infty, \phi_\infty) - (A_\infty^{\text{Lim}}, \phi_\infty^{\text{Lim}})|_{\mathcal{C}^k(K)}$  is the same as the size of  $|(A_i^{\text{Lim}}, \phi_i^{\text{Lim}}) - (A_\infty^{\text{Lim}}, \phi_\infty^{\text{Lim}})|_{\mathcal{C}^k(K)}$ . This proves the Proposition.  $\square$

8.2.1. *Continuity along rays.* We now investigate the behavior of the compactified Kobayashi-Hitchin map restricted to a singular fiber. Specifically, fix  $0 \neq q \in H^0(K^2)$ , and denote by  $[q]$  the  $\mathbb{C}^*$ -orbit of  $q \times 1$  in the compactified Hitchin base  $\overline{\mathcal{B}}$ . Define  $\overline{\mathcal{M}}_{\text{Dol}, q} := \overline{\mathcal{H}}_{\text{Dol}}^{-1}([q])$ ,  $\overline{\mathcal{M}}_{\text{Hit}, q} := \overline{\mathcal{H}}_{\text{Hit}}^{-1}([q])$ . Then the restriction of  $\overline{\Xi}$  on  $\overline{\mathcal{M}}_{\text{Dol}, q}$  defines a map  $\overline{\Xi}_q : \overline{\mathcal{M}}_{\text{Dol}, q} \rightarrow \overline{\mathcal{M}}_{\text{Hit}, q}$ .

**Theorem 8.3.** *Let  $q$  be an irreducible quadratic differential.*

- (i) *If  $q$  contains only zeroes of odd order, then  $\overline{\Xi}_q$  is continuous.*
- (ii) *If  $q$  contains a zero of even order, let  $\mathcal{M}_q = \cup_D \mathcal{M}_{q, D}$  be the stratification defined earlier. Then for each  $D \neq 0$ , there exists an integer  $n_D > 0$  such that for any Higgs bundle  $(\mathcal{F}, \psi) \in \mathcal{M}_{q, D}$ , there exist  $n_D$  sequences of Higgs bundles  $(\mathcal{E}_i^k, \varphi_i^k)$  with  $k = 1, \dots, n_D$  such that*
  - (a)  $\lim_{i \rightarrow \infty} (\mathcal{E}_i^k, \varphi_i^k) = (\mathcal{F}, \psi)$  for  $k = 1, \dots, n_D$ ,
  - (b) *if  $\lim_{i \rightarrow \infty} t_i = \infty$ , and we write*

$$\eta^k := \lim_{i \rightarrow \infty} \overline{\Xi}_q(\mathcal{E}_i^k, \varphi_i^k), \quad \xi := \lim_{i \rightarrow \infty} \overline{\Xi}_q(\mathcal{F}, t_i \psi),$$

*then  $\xi, \eta^1, \dots, \eta^{n_D}$  are  $n_D + 1$  different limiting configurations.*

*Proof.* This follows from Theorem 6.15 and Proposition 8.2.  $\square$

**Theorem 8.4.** *Suppose  $q$  is reducible, and let  $\overline{\mathcal{M}}_{\text{Dol}, q}^{\text{st}}$  be the stable locus of  $\overline{\mathcal{M}}_{\text{Dol}, q}$ . Then the restriction map  $\Theta_q^{\text{Moc}}|_{\overline{\mathcal{M}}_{\text{Dol}, q}^{\text{st}}} : \overline{\mathcal{M}}_{\text{Dol}, q}^{\text{st}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}, q}$  is discontinuous when  $g \geq 3$  and continuous when  $g = 2$ .*

*Proof.* This follows from Propositions 7.11 and 8.2.  $\square$

8.2.2. *Varying fiber.* With the conventions above, suppose  $(\mathcal{E}_i, \varphi_i)$  converges to  $(\mathcal{E}_\infty, \varphi_\infty)$  with  $q_\infty$  having only simple zeros, and  $\xi_i = (\mathcal{E}_i, t_i \varphi_i)$  converges to  $\xi_\infty$  on  $\overline{\mathcal{M}}_{\text{Dol}}$ . Since the condition of having only simple zeros is open, the  $q_i$  also have simple zeros for  $i$  sufficiently large.

**Proposition 8.5** (cf. [36, Thm. 2.12]). *Suppose  $q_\infty$  has only simple zeros. Then,  $\lim_{i \rightarrow \infty} \overline{\Xi}(\xi_i) = \overline{\Xi}(\xi_\infty)$ . In particular, the map  $\overline{\Xi}^{\text{reg}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}$  is continuous.*

*Proof.* Let  $S_i$  denote the spectral curve of  $(\mathcal{E}_i, \varphi_i)$  with branching locus  $Z_i$ . Also, let  $L_i := \chi_{BNR}^{-1}(\mathcal{E}_i, \varphi_i)$  be the eigenline bundles. By the construction in Section 6, we have  $\Upsilon^{\text{Moc}}(\xi_i) = \mathcal{F}_*(L_i, \chi_i)$ , where  $\chi_i = -\frac{1}{2}\chi_{Z_i}$ . Our assumption implies that  $\mathcal{F}_*(L_i, \chi_i)$  converges to  $\mathcal{F}_*(L_\infty, \chi_\infty)$  in the sense of Definition 3.2. Thus, by Theorem 3.3, we obtain the convergence of the limiting configurations:  $\lim_{i \rightarrow \infty} \Upsilon^{\text{Moc}}(\xi_i) = \Upsilon^{\text{Moc}}(\xi_\infty)$ . The claim follows from Proposition 8.2.  $\square$

**Theorem 8.6.** *The map  $\overline{\Xi}^{\text{reg}} : \overline{\mathcal{M}}_{\text{Dol}}^{\text{reg}} \rightarrow \overline{\mathcal{M}}_{\text{Hit}}^{\text{reg}}$  is a homeomorphism.*

*Proof.* By Theorem 4.9,  $\overline{\Xi}^{\text{reg}}$  is a bijection. Moreover, by Proposition 8.5,  $\overline{\Xi}^{\text{reg}}$  is continuous. Finally, that  $(\overline{\Xi}^{\text{reg}})^{-1}$  is continuous follows directly from the construction in [31].  $\square$

## APPENDIX A. CLASSIFICATION OF RANK 1 TORSION MODULES FOR $A_n$ SINGULARITIES

In this appendix, we review the classification result for rank 1 torsion free modules at  $A_n$  singularities, as given in [17]. We compute the integer invariants defined in Subsection 5.3.

Let  $S$  be the spectral curve of an  $\text{SL}(2, \mathbb{C})$  Higgs bundle, and  $p$  a singular point with local defining equation for  $x$  given by  $r^2 - s^{n+1} = 0$ ; this is an  $A_n$  singularity. Let  $p : \tilde{S} \rightarrow S$  be the normalization, where  $p^{-1}(x) = \{\tilde{x}_+, \tilde{x}_-\}$  if  $n$  is odd and  $p^{-1}(x) = \tilde{x}$  if  $n$  is even. We use  $\mathcal{O}_p$  to denote the local ring,  $R$  to denote the completion,  $K$  to denote the field of fractions, and  $\tilde{R}$  to denote the normalization.

**A.1.  $A_{2n}$  singularity.** The local equation is  $r^2 - s^{2n+1} = 0$ . The normalization induces a map between coordinate rings, and we can write

$$\psi : \mathbb{C}[r, s]/(r^2 - s^{2n+1}) \longrightarrow \mathbb{C}[t], \quad \psi(f(r, s)) = f(t^{2n}, t^2),$$

where  $\tilde{R} = \mathbb{C}[[t]]$  and  $R = \mathbb{C}[[t^2, t^{2n+1}]] \subset \tilde{R}$ . According to [17, Anh. (1.1)], any rank 1 torsion free  $R$ -module can be written as

$$M_k = R + R \cdot t^k \subset \tilde{R}, \quad k = 1, 3, \dots, 2n+1.$$

Here,  $M_k$  is a fractional ideal that satisfies  $R \subset M_k \subset \tilde{R}$ , with  $M_1 = \tilde{R}$  and  $M_{2n+1} = R$ .

We may express any  $f \in M_k$  as  $f = \sum_{i=0}^{\frac{k-1}{2}} f_{2i} t^{2i} + \sum_{i \geq k} f_i t^i$ , where  $f_i \in \mathbb{C}$ .

We are interested in the integers  $\ell_x := \dim_{\mathbb{C}}(M_k/R)$ ,  $a_{\tilde{x}} := \dim_{\mathbb{C}}(\tilde{R}/C(M_k))$  and  $b_x = \dim_{\mathbb{C}}(\text{Tor}(M_k \otimes_R \tilde{R}))$ . Thus, as a  $\mathbb{C}$ -vector space,  $M_k/R$  is generated by  $t^k, t^{k+2}, \dots, t^{2n-1}$ , implying that  $\ell_x = \frac{2n+1-k}{2}$ .

The conductor of  $M_k$  is given by  $C(M_k) = \{u \in K \mid u \cdot \tilde{R} \subset M_k\}$ . By the expression of  $M_k$  and a straightforward computation, we have  $C(M_k) = (t^{k-1})$ , where  $(t^{k-1})$  is the ideal in  $\tilde{R}$  generated by  $t^{k-1}$ . Thus,  $1, t, \dots, t^{k-2}$  will form a basis for  $\tilde{R}/C(M_k)$ , and we have  $a_{\tilde{x}} = k - 1$ . Therefore, we have  $a_{\tilde{x}} = 2n - 2\ell_x$ .

For  $i = 0, 1, \dots, \frac{2n-1-k}{2}$ , we define  $s_i = t^{k+2i} \otimes_R 1 - 1 \otimes t^{k+2i} \in M_k \otimes_R \tilde{R}$ . As  $k$  is odd,  $t^{2n+1-k-2i} \in R$  and  $t^{2n+1-k-2i} s_i = t^{2n+1} \otimes_R 1 - 1 \otimes t^{2n+1} = 0$ , where the last equality is because  $t^{2n+1} \in R$ . Moreover,  $\{s_1, \dots, s_{\frac{2n-1-k}{2}}\}$  form a basis of  $\text{Tor}(M_k \otimes_R \tilde{R})$ , thus  $b_x = \frac{2n+1-k}{2} = \ell_x$ .

**A.2.  $A_{2n-1}$  singularity.** The local equation is  $r^2 - s^{2n} = 0$ . The normalization induces a map between the coordinate rings:

$$\psi : \mathbb{C}[r, s]/(r^2 - s^{2n}) \longrightarrow \mathbb{C}[t] \oplus \mathbb{C}[t], \quad \psi(f(r, s)) = (f(t^n, t), f(-t^n, t)),$$

where  $\tilde{R} = \mathbb{C}[[t]] \oplus \mathbb{C}[[t]]$  and  $R = \mathbb{C}[[t, t], (t^n, -t^n)] \cong \mathbb{C}[[t, t], (t^n, 0)]$ . By [17, Anh. (2.1)], any rank 1 torsion free  $R$ -module can be written as:

$$M_k = R + R \cdot (t^k, 0) \subset \tilde{R}, \quad k = 0, 1, \dots, n.$$

Then,  $M_k$  is also a fractional ideal with  $R \subset M_k \subset \tilde{R}$ . Moreover,  $M_n = R$ , and  $M_0 = \tilde{R}$ .

As  $p^{-1}(x) = \{\tilde{x}_+, \tilde{x}_-\}$ ,  $\tilde{R}$  contains two maximal ideals,  $\mathfrak{m}_+ = ((t, 1))$ ,  $\mathfrak{m}_- = ((1, t))$ . For  $f \in M_k$ , we can express  $f$  as:

$$f = \sum_{i=0}^{k-1} f_{ii}(t^i, t^i) + \sum_{l \geq 0} f_{l0}(t^{k+l}, 0) + f_{0l}(0, t^{k+l}),$$

where  $f_{ij} \in \mathbb{C}$ . Therefore,  $\ell_x = \dim_{\mathbb{C}}(M_k/R) = n - k$ . Moreover, using the expression, we can compute the conductor  $C(M_k) = ((t^k, 1)) \cdot ((1, t^k))$ , which implies  $a_{\tilde{x}_{\pm}} = k$ . Similarly, for  $i = k, \dots, n-1$ , we define  $s_i = (t^i, 0) \otimes_R (1, 1) - (1, 1) \otimes_R (t^i, 0)$ , then  $(t, t)^{n-i} \cdot s_i = 0$  and  $\{s_k, \dots, s_{n-1}\}$  will be a basis for  $\text{Tor}(M_k \otimes_R \tilde{R})$  and  $b_x = \ell_x$ .

In summary, we have the following:

**Proposition A.1.** *For the integers defined above, we have:*

- (i) *for the  $A_{2n}$  singularity, we have  $a_{\tilde{x}} = 2n - 2\ell_x$  and  $b_x = \ell_x$ ,*
- (ii) *for the  $A_{2n-1}$  singularity, we have  $a_{\tilde{x}_{\pm}} = n - \ell_x$  and  $b_x = \ell_x$ .*

## REFERENCES

- [1] Arnaud Beauville, M. S. Narasimhan, and S. Ramanan. Spectral curves and the generalised theta divisor. *J. Reine Angew. Math.*, 398:169–179, 1989.
- [2] Usha Bhosle. Generalised parabolic bundles and applications to torsionfree sheaves on nodal curves. *Ark. Mat.*, 30(2):187–215, 1992.
- [3] Gao Chen and Nianzi Li. Asymptotic geometry of the moduli space of rank two irregular higgs bundles over the projective line. *arXiv preprint arXiv:2206.11883*, 2022.
- [4] P. R. Cook. Compactified Jacobians and curves with simple singularities. In *Algebraic geometry (Catania, 1993/Barcelona, 1994)*, volume 200 of *Lecture Notes in Pure and Appl. Math.*, pages 37–47. Dekker, New York, 1998.
- [5] Phillip Richard Cook. *Local and global aspects of the module theory of singular curves*. PhD thesis, University of Liverpool, 1993.
- [6] Kevin Corlette. Flat  $G$ -bundles with canonical metrics. *J. Differential Geom.*, 28(3):361–382, 1988.
- [7] Mark Andrea A. de Cataldo. Projective compactification of Dolbeault moduli spaces. *Int. Math. Res. Not. IMRN*, (5):3543–3570, 2021.
- [8] S. K. Donaldson. Twisted harmonic maps and the self-duality equations. *Proc. London Math. Soc. (3)*, 55(1):127–131, 1987.
- [9] Cyril D’Souza. Compactification of generalised Jacobians. *Proc. Indian Acad. Sci. Sect. A Math. Sci.*, 88(5):419–457, 1979.
- [10] David Dumas and Andrew Neitzke. Asymptotics of Hitchin’s metric on the Hitchin section. *Comm. Math. Phys.*, 367(1):127–150, 2019.
- [11] Yue Fan. An analytic approach to the quasi-projectivity of the moduli space of Higgs bundles. *Adv. Math.*, 406:Paper No. 108506, 33, 2022.
- [12] Yue Fan. Construction of the moduli space of Higgs bundles using analytic methods. *Mathematical Research Letters*, 29(4):1011–1048, 2022.
- [13] Laura Fredrickson. Generic ends of the moduli space of  $\text{SL}(n, \mathbb{C})$ -higgs bundles. *arXiv preprint arXiv:1810.01556*, 2018.
- [14] Laura Fredrickson. Exponential decay for the asymptotic geometry of the Hitchin metric. *Comm. Math. Phys.*, 375(2):1393–1426, 2020.
- [15] Laura Fredrickson, Rafe Mazzeo, Jan Swoboda, and Hartmut Weiss. Asymptotic geometry of the moduli space of parabolic  $\text{SL}(2, \mathbb{C})$ -higgs bundles. *J. Lond. Math. Soc. (2)*, 106(2):590–661, 2022.
- [16] Peter B. Gothen and André G. Oliveira. The singular fiber of the Hitchin map. *Int. Math. Res. Not. IMRN*, (5):1079–1121, 2013.
- [17] G.-M. Greuel and H. Knörrer. Einfache Kurvensingularitäten und torsionsfreie Moduln. *Math. Ann.*, 270(3):417–425, 1985.
- [18] Gert-Martin Greuel and Gerhard Pfister. Moduli spaces for torsion free modules on curve singularities. I. *J. Algebraic Geom.*, 2(1):81–135, 1993.
- [19] Tamás Hausel. Compactification of moduli of Higgs bundles. *J. Reine Angew. Math.*, 503:169–192, 1998.

- [20] Siqi He. The behavior of sequences of solutions to the Hitchin-Simpson equations. *arXiv preprint arXiv:2002.08109*, 2020.
- [21] N. J. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc. (3)*, 55(1):59–126, 1987.
- [22] N. J. Hitchin. Lie groups and Teichmüller space. *Topology*, 31(3):449–473, 1992.
- [23] Nigel Hitchin. Stable bundles and integrable systems. *Duke Math. J.*, 54(1):91–114, 1987.
- [24] Johannes Horn. Semi-abelian spectral data for singular fibres of the  $SL(2, \mathbb{C})$ -Hitchin system. *Int. Math. Res. Not. IMRN*, (5):3860–3917, 2022.
- [25] Johannes Horn.  $sl(2)$ -type singular fibres of the symplectic and odd orthogonal Hitchin system. *J. Topol.*, 15(1):1–38, 2022.
- [26] Johannes Horn and Xuesen Na. The geometry of singular Hitchin fibers via Hecke modifications and abelianisation. *in preparation*.
- [27] Ludmil Katzarkov, Alexander Noll, Pranav Pandit, and Carlos Simpson. Harmonic maps to buildings and singular perturbation theory. *Communications in Mathematical Physics*, 336:853–903, 2015.
- [28] Georgios Kydonakis, Hao Sun, and Lutian Zhao. Monodromy of rank 2 parabolic hitchin systems. *Journal of Geometry and Physics*, 171:104411, 2022.
- [29] Rafe Mazzeo, Jan Swoboda, Hartmut Weiß, and Frederik Witt. Limiting configurations for solutions of Hitchin’s equation. *Séminaire de théorie spectrale et géométrie*, 31:91–116, 2012-2014.
- [30] Rafe Mazzeo, Jan Swoboda, Hartmut Weiss, and Frederik Witt. Ends of the moduli space of Higgs bundles. *Duke Math. J.*, 165(12):2227–2271, 2016.
- [31] Rafe Mazzeo, Jan Swoboda, Hartmut Weiss, and Frederik Witt. Asymptotic geometry of the Hitchin metric. *Comm. Math. Phys.*, 367(1):151–191, 2019.
- [32] Takuro Mochizuki. Asymptotic behaviour of tame harmonic bundles and an application to pure twistor  $D$ -modules. I. *Mem. Amer. Math. Soc.*, 185(869):xii+324, 2007.
- [33] Takuro Mochizuki. Asymptotic behaviour of certain families of harmonic bundles on Riemann surfaces. *J. Topol.*, 9(4):1021–1073, 2016.
- [34] Takuro Mochizuki and Szilárd Szabó. Asymptotic behaviour of large-scale solutions of hitchin’s equations in higher rank. *arXiv preprint arXiv:2303.04913*, 2023.
- [35] Nitin Nitsure. Moduli space of semistable pairs on a curve. *Proc. London Math. Soc. (3)*, 62(2):275–300, 1991.
- [36] Andreas Ott, Jan Swoboda, Richard Wentworth, and Michael Wolf. Higgs bundles, harmonic maps, and pleated surfaces. *arXiv:2004.06071*, 2020.
- [37] Joshua H. Rabinowitz. On monoidal transformations of coherent sheaves. *Proc. Amer. Math. Soc.*, 74(2):389–390, 1979.
- [38] C. J. Rego. The compactified Jacobian. *Ann. Sci. École Norm. Sup. (4)*, 13(2):211–223, 1980.
- [39] Alexander Schmitt. Projective moduli for Hitchin pairs. *Internat. J. Math.*, 9(1):107–118, 1998.
- [40] C. S. Seshadri. Space of unitary vector bundles on a compact Riemann surface. *Ann. of Math. (2)*, 85:303–336, 1967.
- [41] Carlos Simpson. The Hodge filtration on nonabelian cohomology. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 217–281. Amer. Math. Soc., Providence, RI, 1997.
- [42] Carlos T. Simpson. Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization. *J. Amer. Math. Soc.*, 1(4):867–918, 1988.
- [43] Carlos T. Simpson. Harmonic bundles on noncompact curves. *J. Amer. Math. Soc.*, 3(3):713–770, 1990.
- [44] Carlos T. Simpson. Higgs bundles and local systems. *Inst. Hautes Études Sci. Publ. Math.*, (75):5–95, 1992.
- [45] Carlos T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. I. *Inst. Hautes Études Sci. Publ. Math.*, (79):47–129, 1994.
- [46] Clifford Henry Taubes. Compactness theorems for  $SL(2; \mathbb{C})$  generalizations of the 4-dimensional anti-self dual equations. *arXiv preprint arXiv:1307.6447*, 2013.
- [47] Clifford Henry Taubes.  $PSL(2; \mathbb{C})$  connections on 3-manifolds with  $L^2$  bounds on curvature. *Camb. J. Math.*, 1(2):239–397, 2013.
- [48] Richard A. Wentworth. Higgs bundles and local systems on Riemann surfaces. In *Geometry and quantization of moduli spaces*, Adv. Courses Math. CRM Barcelona, pages 165–219. Birkhäuser/Springer, Cham, 2016.

MORNINGSIDE CENTER OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING, 100190 CHINA  
*Email address:* `sqhe@amss.ac.cn`

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA, 94305  
*Email address:* `rmazzeo@stanford.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, CHAMPAIGN, IL 61820, USA  
*Email address:* `xna@illinois.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD 20742, USA  
*Email address:* `raw@umd.edu`