

# Spin-orbit coupled Hubbard skyrmions

Ryo Makuta\* and Chisa Hotta†

Department of Basic Science, University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan

(Dated: April 18, 2023)

When the material phases exhibit topological quantum numbers, they host defects protected by the nontrivial topology. Magnetic skyrmions are such “quantized” objects and although many of them are metals they had been most likely treated in theories as purely classical spin states. Here, we show that the electrons described by the Hubbard model with strong spin-orbit couplings can exhibit various nano- or flake-size skyrmions in their ground state. Importantly, the conducting electrons forming Fermi pockets themselves carry textured magnetic moments in both real and momentum space and on that top, possess Chern numbers in their energy bands. This quantum and conducting skyrmion is related to small skyrmions observed in atomic-layered compounds. We clarify how the effective magnetic interactions and magnetic anisotropies are tied to the spin-orbit coupling, and how they influence the stability of skyrmions beyond the phenomenology.

## I. INTRODUCTION

Skyrmions are topologically protected classical defects first discovered in a nonlinear field theory for mesons in high energy physics<sup>1</sup>. These studies are followed afterward by the intense discussions in the 1990s, regarding the charged excitations of quantum Hall ferromagnets as skyrmions<sup>2,3</sup>. The latest analog appears in magnetic materials<sup>4</sup> particularly in a family of non-centrosymmetric B20-type transition-metal silicides and germanides<sup>5–10</sup>, which have been experimentally established as platforms of skyrmions that extend from nanometers to micrometer scale. In these chiral ferromagnets with broken inversion symmetry, the classical descriptions based on field theories have successfully explained the phenomenological features of their swirling magnetic structures nowadays called Bloch-type<sup>11</sup>; the underlying helical magnetic structures running in different directions are combined and thermodynamically converted to skyrmions at a very particular range of finite temperature and magnetic-field strength. Later, another series called Néel-type skyrmions was found in the cycloidal magnets in noncentrosymmetric polar crystals<sup>12–15</sup>. Meanwhile, some other classes of skyrmions in centrosymmetric materials<sup>16,17</sup> show that the frustrated classical magnetic interactions<sup>18</sup> or RKKY interactions<sup>19–21</sup> can also bear very similar textures to Bloch-type ones.

These “classical skyrmions” are so far believed to be well understood by the classical-spin based Ginzburg-Landau theories or models<sup>11,22,23</sup>, which seemingly suggests it be buried in oblivion that most of the skyrmionic materials discovered so far are metals. Importantly, for skyrmions found in laboratories, the electrons itinerate as charge carriers which themselves carry spatially textured magnetic moments. This means that although the ordered moments of long wavelength may fit the classical description, their underlying origins should be far from being purely classical. Previously, the magnetic interactions discussed in classical theories were ferromagnetic Heisenberg and Dzyaloshinskii-Moriya (DM) interaction as well as magnetic anisotropies which are treated

as handweaving free parameters allowed from the crystal symmetry point of view; it remains unsettled how they stem from both spin-orbit coupling (SOC) and inversion-symmetry breaking.

The current challenge is to truly understand the SOC as the direct origin of skyrmions visibly formed by *conducting* electrons. It is found that the SOC Hubbard model can afford metallic skyrmions that carry both charges and spin moments while maintaining its topological feature as Chern numbers on top of their magnetic quantum winding numbers. Quite importantly, the observed skyrmions are different from the Bloch or Néel type sizable skyrmions previously reported at finite temperatures. They are either nano- or flake-size and appear in the ground state for a wide range of Coulomb interactions when the SOC is substantially large. It turns out that the DM interactions and the types of spin anisotropies for these insulating magnets are not at all independent parameters, being strictly governed by the SOC. Our conclusions urge the reconsideration of the models and conclusions drawn from classical theories.

## II. MODEL AND METHOD

We consider the Hubbard model at half-filling with Rashba SOC on the triangular lattice whose Hamiltonian is given as

$$\hat{\mathcal{H}} = - \sum_{\langle i,j \rangle} c_i^\dagger (t + i\lambda(\mathbf{n}_{ij} \cdot \boldsymbol{\sigma})) c_j + \text{h.c.} + U \sum_j \hat{n}_{j\uparrow} \hat{n}_{j\downarrow} - \sum_j c_j^\dagger (\mathbf{B} \cdot \boldsymbol{\sigma}) c_j, \quad (1)$$

where  $\langle i, j \rangle$  runs over all pairs of next nearest neighboring sites,  $\mathbf{c}_i^\dagger = (c_{i\uparrow}^\dagger, c_{i\downarrow}^\dagger)$  is the creation operator of up and down spin electrons,  $n_i$  is an electron number operator, and  $U$  and  $\mathbf{B}$  are on-site interaction and an external

magnetic field, respectively. The spin-dependent hopping integral  $\lambda$  originates from the SOC<sup>24,25</sup>, and when combined with the  $t$ -term we obtain the form,

$$t + i\lambda(\mathbf{n}_{ij} \cdot \boldsymbol{\sigma}) = t_{\text{eff}} e^{i(\theta/2)\mathbf{n}_{ij} \cdot \boldsymbol{\sigma}}, \quad (2)$$

using the SU(2) gauge field,  $U_{ij} = e^{i(\theta/2)\mathbf{n}_{ij} \cdot \boldsymbol{\sigma}}$ , where  $\boldsymbol{\sigma} = (\sigma^x, \sigma^y, \sigma^z)$  is the Pauli matrix,  $t_{\text{eff}} = \sqrt{t^2 + \lambda^2}$  and  $\theta = 2\arctan(\lambda/t)$ . The gauge field rotates the spin orientation by  $\theta$  about  $\mathbf{n}_{ij}$ -axis when the electron hops to the nearest neighbor sites as shown in Fig. 1(a).

The direction of the rotation axis  $\mathbf{n}_{ij}$  is determined by the crystal symmetry of the materials. For example, in the B20-type chiral magnets with  $P2_13$  or  $P4_132$  space groups,  $\mathbf{n}_{ij}$  should be parallel to the bond directions and favor helical structures. Whereas, for polar magnets of  $C_{nv}$  group,  $\mathbf{n}_{ij}$  should be perpendicular to bonds and host cycloidal magnetic structures. In the Hubbard model, the orientation of  $\mathbf{n}_{ij}$  turns out to be an important factor determining not only the kind of skyrmions but its stability. In our calculation, we consider  $C_{3v}$  type  $\mathbf{n}_{ij}$  placed inside the 2D plane (see Fig. 1(a)) and  $\mathbf{B}$  perpendicular to the plane unless otherwise noted.

The reason why Eq.(1) remained unexplored despite the importance of studying electronic models for skyrmions is purely because of technical difficulty. Standard numerical solvers that can be applied to Eq.(1) use the finite-size cluster that often misfits with the periods of incommensurate orders. In such cases, the stability of the true ground state is underestimated or overlooked without knowing its periods *a priori*. This artifact hindered the unbiased search of skyrmions in Hubbard models. Recently, however, a theoretical treatment called sine-square-deformed mean-field theory (SSDMF) has been developed<sup>26</sup> and was successfully applied to the same Hamiltonian as Eq.(1) on a square lattice<sup>27</sup>, whose phase diagram turned out to host a variety of magnetic structures including large-scale incommensurate spin-density-wave states, spiral, stripe, and vortex phases, which seriously compete with each other. There, the reliability about unbiasedness and quality of the SSDMF results are confirmed through the comparison with four different analytical and numerical methods, some of which take full account of the correlation effects. Indeed, SSDMF is not simply a lowest approximation method; it is shown to take account of the correlation effects beyond the mean-field approximation<sup>26</sup> possible due to the real-space renormalization effect caused by SSD<sup>28</sup>. We thus apply the SSDMF to Eq.(1) for the cluster shown in Fig. 1(a) and successfully disclose the basic properties of the ground state (for details see Methods).

We briefly outline the SSDMF theory<sup>26</sup>. We prepare a finite-size hexagonal-like cluster and multiply an envelope function to Eq.(1) that has a sine-squared functional form gradually decreasing from the maximum at the center ( $\mathbf{r}_i = 0$ ) toward the cluster edges (see Fig. 1(a)) given as  $f_{\text{SSD}}(\mathbf{r}) = (1 + \cos(\pi|\mathbf{r}|/R))/2$  with a radius  $R = \max_i |\mathbf{r}_i| + 1/2$ . The cluster size adopted are  $N = 90$  in

Fig. 1(a) and  $N = 198$  to confirm the convergence of the finite size effect.

The site-dependent mean fields  $\langle n_i \rangle$  and  $\langle \mathbf{S}_i \rangle$  are introduced and are determined self consistently. Our mean-field Hamiltonian with SSD is given by  $\mathcal{H}_{\text{MF}} = \mathcal{H}_0 + \mathcal{H}_{U,\text{MF}}$  where

$$\mathcal{H}_{U,\text{MF}} = U \sum_i f(\mathbf{r}_i) \left( \frac{\langle n_i \rangle}{2} n_i - 2 \langle \mathbf{S}_i \rangle \cdot \mathbf{S}_i \right) + E_c, \quad (3)$$

$$E_c = -U \sum_i f(\mathbf{r}_i) \left( \frac{\langle n_i \rangle^2}{4} - \langle \mathbf{S}_i \rangle^2 \right). \quad (4)$$

The obtained solutions are site-dependent when given the deformed Fourier transform as  $\langle \mathbf{S}_q \rangle = \sum_i f_{\text{SSD}}(\mathbf{r}_i) \langle \mathbf{S}_i \rangle e^{i\mathbf{q}\mathbf{r}_i} / \sum_i f_{\text{SSD}}(\mathbf{r}_i)$ , they yield an unbiased period that almost exactly reproduces that of the much larger cluster with the periodic boundary condition that safely accommodates that period. Previously, for example, the period of  $(q, q)$  ( $q = (1 - 1/\sqrt{10})\pi$ ) is found to be exactly detected using the  $8 \times 8$  cluster<sup>26</sup>. We perform the inverse Fourier transformation on  $\tilde{S}_i^\mu(\mathbf{k})$  and  $\tilde{n}_i(\mathbf{k})$  with normal periodic boundary conditions of  $N = 192$  clusters to obtain spin and particle densities on a uniform periodic lattice.

### III. RESULTS

#### A. Phase diagram

We first show the phase diagrams on the plane of  $\theta$  and  $B$  for  $U/t_{\text{eff}} = 8$  and 5 in Figs. 1(b) and 1(c). The  $\theta = 0$  case is the standard Hubbard model at half-filling; when  $U/t_{\text{eff}} = 8$ , the system is in a paramagnetic phase at  $B = 0$ , which is transformed at  $B/t_{\text{eff}} \geq 0.2$  to the state with all the triangles having two-up and one-down spin orientations, which is known as UUD phase in quantum magnetism. This result is consistent with the corresponding previous theories on the half-filled Hubbard model<sup>29</sup> showing the spin liquid Mott insulator at  $U/t \sim 8 - 10$  (while it is beyond our description) and  $120^\circ$  Néel ordering at  $U/t \gtrsim 10^{30-32}$ . At  $\theta = \pi$  with  $t = 0$  and  $\lambda = 1$ , an almost fully polarized ferromagnet appears, whose origin will be discussed shortly. In the shaded area of the ferromagnetic phase, the metastable solution reminiscent of the Néel-skyrmion appears which is discussed in Appendix A.

At intermediate  $\theta$ ,  $t$  and  $\lambda$  compete and generate various phases. The most prominent feature is the emergence of two small-size skyrmion phases at around  $0.2 \lesssim \theta/\pi \lesssim 0.5$ , which we call nano- and flake-skyrmions based on the similarity with the previously found skyrmions in quantum spin models<sup>33,34</sup>; their real space spin distributions and spin structural factors in the reciprocal space are shown in Figs. 1(d) and 1(e). They have site-centered




FIG. 1. (a) Schematic illustration of the rotation axis  $\mathbf{n}_{ij}$  included in the SU(2) gauge field  $U_{ij}$  when electrons hop from site  $-j$  to site  $-i$ .  $N = 90$  finite size cluster with SSD boundary conditions mainly used in the present calculation and the envelope function  $f_{\text{SSD}}(\mathbf{r})$  (see Methods). (b,c) Ground state phase diagram of Eq.(1) at  $U/t_{\text{eff}} = 8$  and 5. The two small panels give the density plots of the quantum winding number  $W$  and the metal-insulator phase boundaries for the two diagrams. (d,e,f) Realspace spin configuration of the SSDMF solutions for the nano-, flake- and three-sublattice AF-skyrmions. The lower two panels are the spin structural factors  $\langle S_{\mathbf{q}} \rangle$  in momentum space for the in-plane  $xy$ (left) and  $z$ -components(right panel).

and bond-centered characters with 7-site and 27-site per unit, respectively. The spin structural factors have peaks positioned at  $\mathbf{q} = (0.955\pi, 0)$  and  $(0.955\pi, 0.064\pi)$ , respectively. In the lower field region, the AF-skyrmion based on three sublattice structures of a 27-site unit is observed; the structural factors show peaks at different points from the nano- and flake-skyrmions. All of them are characterized as triple- $q$  states.

For classical skyrmions, a winding number is used to characterize the slowly varying continuous spin structures, which takes either an integer or half-integer numbers when the gradients of spins are integrated over the skyrmion unit. In quantum systems, the spin moments fluctuate and squeeze on discrete lattice sites and do not show global continuous structures. The replacement of the winding number is given by the skyrmion number  $W$  introduced in a Ref.[34] as the total solid angle of trian-

gular spins as

$$W = \frac{1}{4\pi} \sum_{\langle ijk \rangle} \text{atan} 2 \left( \frac{\langle \mathbf{S}_i \cdot \mathbf{S}_j \times \mathbf{S}_k \rangle}{S^3}, 1 + \frac{\langle \mathbf{S}_i \cdot \mathbf{S}_j \rangle + \langle \mathbf{S}_j \cdot \mathbf{S}_k \rangle + \langle \mathbf{S}_k \cdot \mathbf{S}_i \rangle}{S^2} \right), \quad (5)$$

where the summation is taken over the skyrmion unit for triangles  $\langle ijk \rangle$  with indices aligned counterclockwise. The value of  $W$  is not quantized and is relatively small<sup>35</sup>. However, it can overall capture the nano- and flake-skyr phases as shown in the insets of Fig. 1.

For other phases at  $\theta/\pi \sim 0.5$  with increasing the magnetic field, the system hosts stripes and 120°- or umbrella-like three-sublattice phases both denoted as “3sub AF”. The details of these magnetic structures are given in Appendix A.




FIG. 2. Metallic flake-skyrmion ground state obtained by SS-DMF at  $U/t_{\text{eff}} = 5$ ,  $B/t_{\text{eff}} = 0.6$  and  $\theta = 0.3\pi$ . (a) Spin configuration inside the 27-site unit. (b) Energy band structure consisting of 54 bands with Chern numbers shown on the r.h.s. Inset shows the magnified bands near the Fermi surface. (c) Fermi surfaces of valence (hole) and conduction (electron) bands. The bright/dark regions are those below/above the Fermi level. (d) Spin textures in momentum space ( $z$ -component as density plots and  $xy$ -components as arrows) carried by the valence and conduction bands. The Fermi pockets are given as the guide to the eye.

## B. Metallic skyrmions

The phase diagrams have insulating and metallic regions whose boundaries are indicated by the broken lines in the small panels of Fig. 1(b,c). In our framework, the flake-skyrmion at  $U/t_{\text{eff}} = 5$  can naturally appear as metals.

In the metallic phases, the magnetic moments shrink, and it becomes much more difficult to have a rigid regular magnetic structure compared to the insulating case. In our calculation particularly at  $U/t_{\text{eff}} = 5$ , the UUD, skyrlike and stripe-like phases surrounding the flake-skyrmion phase often show various numerical solutions with domains or irregular winding structures of spins. Contrastingly, the flake-skyrmion metal is rigid (see Fig. 2(a)).

Let us briefly show how we classify the states to be metallic or insulating. Once we obtain the spatial peri-

ods of the spin structures by the dominant peak positions  $S_\gamma(\mathbf{q})$ , we can construct a system with periodic boundary conditions of the size of that period. Based on the corresponding set of mean fields that work as one-body potentials, the energy band structure and the Fermi surface are obtained, which are shown in Figs. 2(b) and 2(c) for flake-skyrmion. The two bands cross the Fermi level that gives two hole pockets and electron pockets in different regions of the Brillouin zone. Since the valence and conduction bands do not contact, we can calculate the Chern numbers as

$$C_n = \frac{1}{2\pi} \sum_k \mathcal{I}m \ln \Delta U_l(k), \quad \Delta U_l(k) = \prod_{\text{plaquette}} \langle u_k | \nabla_k | u_k \rangle, \quad (6)$$

using the fidelity of Bloch wave function  $\Delta U_k$  along the discretized plaquette of the reciprocal space at  $k^36$ . The results are shown on the r.h.s. of the energy bands. The conducting and valence bands have  $C_n = 2$  and  $-1$ , respectively. This will give the quantized Hall conductivity that originates from the U(1) gauge generated from the texture of skyrmions. For comparison, we also show the energy bands of the insulating AF-skyrmion and 3sub-cluster phases with their Chern numbers in Appendix A.

Due to SOC, the spins are no longer good quantum numbers. Therefore, the eigenstates carry spin moments pointing in various directions; we show in Fig. 2(d) the spin textures on valence and conduction bands of the flake-skyrmion. Since the spin moments point almost opposite on different sides of the pockets, one may expect the spin current to be observed if this state is realized.

## C. Spin models at strong coupling limit

In our calculation, the large Néel-type skyrmion previously found in experiments and classical spin models of the same symmetry class are not observed even for parameters far outside the presented phase diagrams. To understand why some skyrmions appear for certain parameters and others do not, establishing the connection of Hubbard models with classical spin models is useful, since the skyrmions in the latter types of models are intensively studied.

The effective Hamiltonian of the strong coupling region  $U/t_{\text{eff}} \gg 1$  of Eq.(1) at  $B = 0$  can be obtained by the perturbation about  $t_{\text{eff}}/U$ . The second-order Hamiltonian includes the Heisenberg interaction, DM interaction, and the Kaplan-Shekhtman-Aharony-Entin-Wohlman (KSAEW) term as

$$\mathcal{H}^{(2)} = \sum_{\langle i,j \rangle} (J \mathbf{S}_i \cdot \mathbf{S}_j + \mathbf{D}_{ij} \cdot \mathbf{S}_i \times \mathbf{S}_j + K (\mathbf{n}_{ij} \cdot \mathbf{S}_i) (\mathbf{n}_{ij} \cdot \mathbf{S}_j)) \quad (7)$$

where  $J = A \cos \theta$ ,  $\mathbf{D}_{ij} = \mathbf{n}_{ij} A \sin \theta$ ,  $A = 4t_{\text{eff}}^2/U$ , and  $K = \sqrt{J^2 + D^2} - J$ . Importantly, the DM vector  $\mathbf{D}_{ij}$  and spin anisotropy vector both point toward the direction of the rotation axis of the SU(2) gauge,  $\mathbf{n}_{ij}$ , in Eq.(1).




FIG. 3. Phase diagram of the classical spin model including the Heiseberg interaction  $J = A \cos \theta$ , DM interaction  $D = A \sin \theta$  and spin anisotropy term  $K = \sqrt{J^2 + D^2} - J$ , taking  $t_{\text{eff}} = 1$  and  $U = 10$  i.e.,  $A = 4t_{\text{eff}}^2/U = 0.4$ . Panel (a) corresponds to the strong coupling limit of our SOC Hubbard model given in Eq.(7) with in-plane  $\mathbf{D}$  and  $\mathbf{n}$ , (b) is the one obtained by taking  $K = 0$ , and (c) show the case where  $\mathbf{n}_{ij}$  in Eq.(7) is pointing perpendicular to the  $xy$ -plane which gives  $\mathbf{D}$  and  $\mathbf{n} \perp xy$ -plane. In the lower panel of (a), the variation of  $J/A = \cos \theta$ ,  $D/A = \tan \theta$ ,  $K/A = (1 - \cos \theta)$  and the detailed descriptions of  $\mathbf{D}$  and  $\mathbf{n}$  for the three cases are summarized. The direction of vector  $\mathbf{n}_{ij}$  for the DM and anisotropy terms are bond and direction ( $i \rightarrow j$ ) dependent.

Here,  $\mathbf{S}_j$  in  $\mathcal{H}^{(2)}$  is a quantum spin-1/2. Whereas taking its large- $S$  limit and regarding Eq.(7) as classical Hamiltonian of classical vector spins still works to understand the magnetic nature of the original Hamiltonian; it is shown in the square lattice SOC Hubbard model that the magnetic structure at finite  $U$  and agrees well with that of the classical ones<sup>27</sup>. Figure 3(a) shows the classical phase diagram of  $\mathcal{H}^{(2)}$  where we set the units of the interactions as constant,  $A = 4t_{\text{eff}}^2/U = 0.4$  corresponding to  $t_{\text{eff}} = 1$  and  $U = 10$ . The nano-/flake-skyr phase appears at around  $\theta \sim 0.5\pi$  and  $B \lesssim 0.8$ , and the regions  $\theta \lesssim 0.4\pi$  and  $0.6\pi \lesssim \theta$  are dominated by the three-sublattice (UUD or  $120^\circ$ ) phase and polarized ferromagnetic phase, respectively. Here, we prepared all representative states that appear in the Hubbard phase diagram as candidates and compared their energies;  $120^\circ$ , UUD, nano-, flake-(19 site unit) and AF-skyrmions and large-skyrmion (Néel as well as other types) with an energetically optimized periodicity. For the nano-/flake phases, the two compete while nano-skyrmion is relatively stable. The large Néel-skyrmion state was energetically subtle and did not appear as the lowest energy state. This diagram is fully consistent with the SSDMF results for

Eq.(1).

The variation of  $J$ ,  $|\mathbf{D}_{ij}|$ ,  $K$  as functions of  $\theta$  shown in the lower panel of Fig. 3(a) helps to understand the energetics. At  $\theta < 0.5\pi$ , the Heisenberg exchange is antiferromagnetic ( $J > 0$ ) which becomes ferromagnetic ( $J < 0$ ) at  $\theta > 0.5\pi$ , having maximum amplitudes at  $\theta = 0$  and  $\pi$ . The DM interaction takes the maximum at  $\theta = 0.5\pi$ , while the  $K$  term is always positive and monotonously increases; this term dislikes the adjacent spins to point in a similar direction, particularly in parallel to  $\mathbf{n}_{ij}$ . Since Fig. 3(a) has  $\mathbf{n}_{ij}$  pointing in-plane, the large DM interaction at  $\theta \sim 0.5\pi$  induces the typical cycloidal variation of moments along the bond directions, while comparably large  $K$  favors the spins to vary quite rapidly in space, both cooperatively stabilize the nano/flake-skyrmion phase when  $B$  is not too large.

The large Néel skyrmion, which may appear if present at around  $\theta/\pi = 0.6 - 0.9$  is not favored here, because it suffers the substantial loss of KSAEW energy compared to the fully polarized state. The reason why the previous classical model could easily host large skyrmions was that the anisotropy interactions (different from KSAEW) and  $\mathbf{D}_{ij}$  are dealt independently as free parameters, which is not allowed in Hubbard models. To confirm this scenario, we calculated the phase diagram by artificially setting  $K = 0$  as shown in Fig. 3(b). At a small magnetic field next to the ferromagnetic phase the large Néel skyrmion is recovered. For further reference, we show in Appendix B the above-mentioned cases like  $\mathbf{D} \parallel xy$  and  $\mathbf{n}_{ij} \perp xy$  which host Néel skyrmion, while they are unphysical in considering the SOC Hubbard model as their origin.

Nonetheless, since the large Néel skyrmion is relevant not only to classical models but to materials, it is natural to expect another channel to realize it. We thus restart from the model Eq.(1) with  $\mathbf{n}_{ij}$  pointing perpendicular to the two-dimensional  $xy$ -plane for all bonds. This modification does not change the form of Eq.(7), and its classical phase diagram is shown in Fig. 3(b); the nano/flake-skyrmions no longer exist while the large-skyrmion appears at large  $\theta$  region instead of the polarized ferromagnets. The large skyrmion here equivalently favors Néel, Bloch, or their mixtures because  $\mathbf{n}_{ij} \cdot \mathbf{S}_i \times \mathbf{S}_j$  depends solely on the relative angles of classical spins projected onto the  $xy$  plane, which is identical for the two skyrmions. This situation is realized in  $\text{Mn}_{1.4}\text{Pt}_{0.9}\text{Pd}_{0.1}\text{Sn}$  where the mixture of Néel and Bloch structures is observed<sup>37,38</sup>.

At the fourth order of perturbation, the ring exchange interaction including SOC appears. We have examined the amplitude of these terms in Appendix C and found that they remain one order smaller than the other magnetic interactions. While these higher-order fluctuations will influence the amplitude of spins, their effect is small since the phase diagrams we saw in Figs. 1(b) and 1(c) are similar, although the degree of fluctuation is by orders of magnitude larger for the latter.

TABLE I. Details of noncentrosymmetric materials hosting skyrmions based on helical (he) and cycloidal (cy) magnets; space group, metallic/insulating (M/I), and the size of the observed skyrmions. Those belonging to B20-group ( $P2_13$  or  $P4_132$ ) are helical/conical magnets that form Bloch skyrmions, and the corresponding DM interactions are  $\mathbf{D}_{ij} \parallel$  bonds. Here, EuPtSi is classified in a different column since it is an RKKY system where  $S = 7/2$  localized magnetic moments are coupled to conduction electrons. The  $C_{nv}$ ,  $R_{3m}$  and  $P4_{cc}$  systems are cycloidal magnets that form Néel skyrmions, and the corresponding DM interactions are  $\mathbf{D}_{ij} \perp$  bonds which are pointing mainly in-plane. The latter groups correspond to our model.

	material	symm.	M/I	[nm]	ref
	MnSi	$P2_13$	M	18	[5]
	FeGe	$P2_13$	M	70	[8]
	MnSi $_{1-x}$ Ge $_x$	$P2_13$	M	60	[39,40]
	MnGe	$P2_13$	M	3	[41]
	Mn $_{1-x}$ Fe $_x$ Ge	$P2_13$	M	3-6	[42]
(he)	Cu $_2$ OSe $_3$	$P2_13$	I	62	[9,43]
	Fe $_{1-x}$ Co $_x$ Si	$P2_13$	I	30-230	[7]
	Co $_{10-x/2}$ Zn $_{10-x/2}$ Mn $_x$	$P2_13$	M	9	[44]
	Co $_{8-x/2}$ Fe $_x$ Zn $_x$ Mn $_4$	$P4_132$	M	120	[10,45]
	FeCo $_{0.5}$ Rh $_{0.5}$ Mo $_3$ N	$P4_132$	M	110	[46]
(he)	EuPtSi	$P2_13$	M	1.8	[47,48]
(he/cy)	Mn $_{1.4}$ Pt $_{0.9}$ Pd $_{0.1}$ Sn	$D_{2d}$	M	150	[37,38]
	Fe/Ir	$C_{3v}$	M	1	[49]
	PdFe/Ir	$C_{3v}$	M	3	[50]
	Ir/Co/Pt	$C_{3v}$	M	30-90	[51]
(cy)	CBST/BST	$C_{3v}$	M	–	[52]
	SrIrO $_3$ /BaTiO $_3$	$C_{4v}$	M	–	[53]
	SrRuO $_3$ /BaTiO $_3$	$C_{4v}$	M	–	[54]
	GaV $_4$ S $_8$	$R_{3m}$	I	17	[12]
(cy)	GaV $_4$ Se $_8$	$R_{3m}$	I	19	[13,55]
	VOSe $_2$ O $_5$	$P4_{cc}$	I	140	[15]

#### IV. SUMMARY AND DISCUSSIONS

We have so far clarified that the metallic and nano- or flake-size skyrmions can be realized in an SOC Hubbard model at half-filling in a certain range of magnetic field perpendicular to the triangular lattice plane. Key differences from the former theories are listed as follows; Firstly, the conducting electrons forming Fermi pockets contribute to skyrmions. They may remind of a spin-density wave state but is different in that the spin momentum is not conserved on each band, namely, each  $k$ -point at each energy band carries magnetic moment pointing in different directions. Second, each band carries nonzero-integer Chern numbers that shall contribute to Hall conductivity. Finally, the major effective magnetic interactions well-defined in the Mott insulating state,  $J$ ,  $D$  and  $K$ , are interrelated and are fully controlled by the single parameter  $\theta = 2 \arctan(\lambda/t)$ , originating from the substantial SOC. Accordingly, the large skyrmions that were easily observed in idealized and phenomenological classical spin models are not found.

To make the discussions relevant to laboratory exper-

iments, we present in Table I the representative non-centrosymmetric materials hosting skyrmions. They are classified into separate columns by double-lines as chiral magnets ( $P2_13$ ,  $P4_132$ ), atomic layers (Fe/Ir) or heterostructures (SrIrO $_3$ /BaTiO $_3$  interfaces), and bulk layered materials.

The ones based on helical magnets (he) are known as B20-type structures with distorted cubic lattices. The Bloch-skyrmions found there are mostly as large as 10-200 nm, and are treated phenomenologically by the Ginzburg-Landau-based or lattice-based classical spin models; the major interactions are the ferromagnetic Heisenberg and DM interactions with  $\mathbf{D}_{ij}$  pointing parallel to the bonds along the  $\hat{x}$ ,  $\hat{y}$ ,  $\hat{z}$ -directions that favor helical or conical magnets. These phenomenologies successfully reproduced the experimental phase diagram, at finite temperatures within a certain field strength<sup>11,22,23</sup>.

However, in reality, most of these magnets are metallic; for example, MnSi is a typical itinerant-electron magnet with a concentration of moments being  $0.4\mu_B$  per Mn sites, which cannot be simply treated as a localized large- $S$  magnet. Further, the microscopic origin of ferromagnetic interactions in such a situation is not clarified. Whereas, EuPtSi having the same space group is different; the Eu ions have well-localized  $S = 7/2$  that interacts with conducting  $s$ -electrons<sup>47,48</sup>, which can be regarded as the RKKY systems. The theories built on classical Kondo-lattice types of models naturally reconcile with this system and exhibit phase diagram with Bloch skyrmions<sup>21,56</sup> which are more or less the same as the ones commonly observed in B20-magnets. Recently, the DFT calculations have shown that the mixed valence state of MnSi can be separated into a relatively localized spin part and the conducting part<sup>57</sup>, where they propose the Kondo-like model and show that the Kondo-coupling strength determines the magnetic concentration and the types of skyrmions realized. It may compromise the phenomenology and the RKKY model for this primary but specific class of skyrmions.

The layered cycloidal magnets (cy) hosting Néel skyrmions have a different character. The major DM interactions allowed for these space groups have  $\mathbf{D}_{ij}$  perpendicular to bonds and are mostly pointing in-plane, which corresponds to the types of SOC we applied. Indeed, the SOC of these materials is substantially large<sup>58</sup>; the atomic SOC for  $3d$  (Fe,Co),  $4d$  (Ru) and  $5d$  (Ir,Pt) ions are 0.04–0.08 eV, 0.1–0.2 eV, and 0.3–0.6 eV, respectively, and when combined with the crystal field or the electric potentials coming from strong two-dimensionality, the electronic states on the clusters or ions have strong anisotropy and mixed spin-angular momentum. The electronic Hamiltonian in the presence of such effect is represented at the simplest by our Eq.(1)<sup>24,59</sup>. The value of  $\lambda/t$  or  $\theta$  is determined by the strength of SOC and the details of the crystal structures and can take substantially large values<sup>25</sup>.

We thus consider that these materials can be the platform of the skyrmions observed in our model. Their

skyrmions overall have a relatively smaller size than those of the B20-chiral magnets, which agrees with our results. In particular, the flake-size skyrmions in our model are the only one that can explain the metallic electron skyrmions observed in  $C_{nv}$  crystals. Here, the Mott insulating phases can be partially related to the previously studied spin models that include a certain amount of quantum fluctuation; the nano-like skyrmion is found in a semi-classical spin- $S$  model<sup>35</sup> that aimed to explain the 1 nm sized skyrmions found in Fe/Ir layers<sup>49</sup>. The flake-skyrmion can be compared with those observed in the fully quantum spin-1/2 system<sup>34</sup>. However, their magnetic interactions do not conform to the ones naturally obtained from Hubbard models (see Appendix B).

We stress that although previous theories assume the ferromagnetic Heisenberg interactions, it is elusive for electronic models without SOC. The superexchange interactions are so far the only natural realization for ferromagnetic exchange, which shall be the case applied to  $\text{GaV}_4\text{X}_8$  or  $\text{VOSe}_2\text{O}_5$  where the V-ions on  $\text{V}_4\text{X}_4$  or  $\text{VO}_5$  clusters carry spin-1/2 and interact through ligands<sup>12,13</sup>.

In the Rashba-type SOC Hubbard model on the square lattice before this study, the double- $q$  spiral phase with noncoplanar spin structures are found at  $0.45 \leq \theta/\pi \leq 0.65$  and  $U/t_{\text{eff}} \gtrsim 4^{27}$ , which is identified as an antiferromagnetic skyrmion lattice phase, and shall be relevant to  $\text{VOSe}_2\text{O}_5$ <sup>15</sup> and  $\text{SrXO}_3/\text{BaTiO}_3$  heterostructures<sup>53,54</sup>. The future perspective will be to apply a similar analysis to Hubbard models or other correlated electron models with different fillings or lattices, which may disclose the landscape of antisymmetric SOC Mott insulators and related SOC metals that host a variety of spin structures based on spin-split bands.

## V. ACKNOWLEDGEMENT

The work is supported by JSPS KAKENHI Grants No. JP21K03440 from the Ministry of Education, Science, Sports and Culture of Japan and from a Grant-in-Aid for Transformative Research Areas "The Natural Laws of Extreme Universe—A New Paradigm for Spacetime and Matter from Quantum Information" (KAKENHI Grant No. 21H05191) from JSPS of Japan.

### Appendix A: Several magnetic ground states in the phase diagrams

We present in Fig. 4 the spin configuration of each phase in Figs. 1(b) and 1(c) not referred to in detail in the main text. At  $U/t_{\text{eff}} = 8$  the 3 sub-cluster and 3 sub-AF (120°-like) magnetic states appear at large  $B$  whose moments are relatively large (Figs. 4(a,b)). The UUD state at  $\theta = 0$  given in Fig. 4(c) is the typical phase observed in quantum Heisenberg or Ising models on a triangular lattice in the magnetic field, often forming plateaus. The stripe phase emerges when the system

approaches the ferromagnetic phase at a small  $B$  which is shown in Fig. 4(d).

The states at  $U/t_{\text{eff}} = 5$  are mostly metallic, so the magnetic moments are much smaller than in the above cases. The flake-skyr phase (Fig. 4(e)) is very stable, which is the one discussed mainly in the main text. The 3-sublattice-cluster phase in Fig. 4(f) has a similar structure with nano-skyr while difficult to judge its periodicity, and so as the stripy-domains (Fig. 4(g)) that have similarity with Néel-type skyrmion.

The one found in the smaller  $\theta$  region (shaded) of the ferromagnetic phase in Fig. 1(b,c) in the main text has spin structure shown in Fig. 4(h), and is classified as a polarized ferromagnet, although there is a small but finite modulation of spin moments similar to Néel-type skyr. We have carefully examined the energy differences of this SSDMF solution with those of the fully polarized ferromagnet by choosing the proper cluster with periodic boundary condition and recalculating the solution for the uniform (non-SSD) Hamiltonian, finding that the fully polarized ferromagnet slightly overwhelms the Néel-like skyrmions. The parameter range where this kind of solution appears is shaded in the phase diagram. We expect that the Néel-like skyrmions may appear at smaller  $\theta$  part of this region at finite temperature as in the previously reported theories.

We next show the energy band structures of insulating AF-skyrmion and UUD 3 sub-like states in Fig. SreffS-ins. They are realized in the  $U/t_{\text{eff}} = 8$  phase diagram Fig. 1(b) at  $(B/t_{\text{eff}}, \theta/\pi) = (0.1, 0.5)$  and  $(0.45, 0.15)$ , respectively. The energy dispersion is suppressed compared to Fig. 3(b) and a large charge gap of  $\sim t_{\text{eff}}$  opens. The Chern numbers are shown on the r.h.s. of the energy bands. Since the real-space magnetic textures (AF-skyrmion is shown in Fig. 1(f) and 3 sub-like state similar to 3 sub-cluster in Fig. 4(a)) do not seem to differ too much from the metallic flake-skyrmion shown in Fig. 3(a) (main text), they have a nonzero winding number, while Chern numbers near the Fermi level are both zero for the latter and the valence band is 1 for the former.

### Appendix B: Comparison with several classical spin models

To understand how the types of magnetic interactions used in the previous studies influence the ground state phase diagram, we consider the analogues of second order Hamiltonian  $\mathcal{H}^{(2)}$  in Eq.(5) in the main text, taking  $\mathbf{S}_i$  as classical vector of size-1. In the main text, the results for Eq.(5) ( $\mathcal{H}^{(2)}$ ) with  $\mathbf{n}$  placed in the  $xy$ -plane perpendicular to the  $ij$ -bonds is shown in Fig. 3(a) where the parameters  $J$ ,  $D$  and  $K$  are the functions of  $\theta$ . This result was compared with its analogs naturally derived by setting  $K = 0$  in Fig. 3(b) or by taking  $\mathbf{n} \parallel z$ -axis in Fig. 3(c).

However, the types of interactions or anisotropies adopted in previous studies are different; in Ref.[35] they




FIG. 4. Real space spin textures of several phases found in Figs. 1(b) and 1(c) in the main text. (a)-(d)  $U/t_{\text{eff}} = 8$  states: 3 sub-cluster ( $B, \theta/\pi = (0.8, 0.2)$ ), 3 sub-AF ( $0.8, 0.4$ ), UUD ( $0.5, 0$ ), and stripe ( $0.1, 0.55$ ). (e)-(h)  $U/t_{\text{eff}} = 5$  states: flake-skyr ( $B, \theta/\pi = (0.2, 0.35)$ ), 3 sub-cluster ( $1, 0.3$ ), stripy-domains ( $0.2, 0.65$ ), and Néel-like (ferro) ( $0.7, 0.55$ ). The colors of the circles are the density plots of  $\langle S_i^z \rangle$  and arrows denote the in-plane element as vectors ( $\langle S_i^x \rangle, \langle S_i^y \rangle$ ).

took  $\mathbf{D} \perp$  bond in the  $xy$ -plane as in our case but considered easy-axis anisotropy which is valid only for  $S > 1$  as  $-K \sum_i (S_i^z)^2$ . To see this effect we calculate in Fig. 6(a) the classical ground state phase diagram by setting  $K$  as constant, while varying  $D$  and  $J$  as functions of  $\theta$ . The constant values,  $K = (4/U)(K_m/\sqrt{D_m^2 + 1})$  with  $K_m = 0.518$  and  $D_m = 2.16$ , are set to directly compare with the case of Ref.[35]. Their SkX1 phase has a similar structure to our nano/flake-skyrmion and is consistent with the stable skyr region in the phase diagram.

Regarding Ref.[60], they used  $\mathbf{D} \perp$  bond in the  $xy$ -plane but for spin anisotropies they set  $\mathbf{n} \parallel z$  (corresponding to the XXZ-model, taking the spin anisotropic exchange interaction), which does not conform to the




FIG. 5. Energy bands of insulating states at  $U/t_{\text{eff}} = 8$  with (a) AF-skyrmions ( $B/t_{\text{eff}}, \theta/\pi = (0.1, 0.5)$ ) shown in Fig. 1(f) in the main text and (b) UUD 3 sub-like state at  $(0.45, 0.15)$ . Chern numbers of bands are shown on the r.h.s. In (a) it has a nonzero value at the valence band.

symmetries provided by the SOC in the Hubbard model. The phase diagram obtained by varying  $J, D, K$  in the same manner as our model but taking  $\mathbf{n} \parallel z$  for the anisotropy term to follow Ref.[60] is shown in Fig. 6(b). The ones taking  $K = \sqrt{J^2 + D^2}/4$  as constant about  $\theta$  is also shown in Fig. 6(c). Broken lines are  $\theta = 0.66\pi$  and  $0.85\pi$  studied in Ref.[60] as  $J = -2D$  and  $J = -0.5D$ , respectively. If we interpret our nano/flake-skyr as their  $\text{SKX}_{3a}$ , the results of ours and theirs are slightly inconsistent in that the nano/flake-skyr appear only in  $J = -2D$  but not in  $J = -0.5D$ , while they argue that both cases have  $\text{SKX}_{3a}$ . In that context, our phase diagram is natural since the  $\theta = 0.85\pi$ , where  $|J| > |D|$  holds, favors larger size skyrmion.

Although these works stress the quantum mechanical magnetic fluctuation as being key to realize small skyrmions, the systematic comparison given here suggests that the relationships between magnetic interactions governed by SOC seems to play a more important role.

### Appendix C: Effect of ring exchange in fourth-order perturbation

In Fig. 3 in the main text, we examined  $\mathcal{H}^{(2)}$  including interactions originating from the second-order perturbation terms. Since the role of higher order terms is being discussed recently<sup>19</sup>, we additionally consider the fourth-order spin-dependent ring exchange term. By considering






FIG. 6. Ground state phase diagram of the classical spin model with different types of anisotropies. The interactions  $J = \cos\theta, D = \sin\theta$  are set as the same as Fig. 3(main text), with  $B$  given in unit of  $A = 4t_{\text{eff}}^2/U = 0.4$ . For panel (a) we take  $-K(S_i^z)^2$  with constant  $K$  given as  $(4/U)(K_m/\sqrt{D_m^2} + 1)$  with  $K_m \equiv K/J = 0.518$  and  $D_m = D/J = 2.16$  following Ref.[35]. In panels (b) and (c) we considered the XXZ type anisotropies by setting  $\mathbf{n} \parallel z$  and taking  $K = 1 - \cos\theta$  for (b) and  $K = \sqrt{J^2 + D^2}/4$  being constant for (c), to compare with Ref.[60] whose  $J = -2D$  and  $J = -0.5D$  cases are shown in broken lines  $\theta = 0.66\pi$  and  $0.85\pi$ , respectively.

the SOC hopping term  $\lambda$ , the ring exchange term is given in the form,

$$\mathcal{H}_{\text{ring}}^{(4)} = \sum_{abcd} \sum_{j=0}^4 \hat{R}_j[\mathbf{S}_a, \mathbf{S}_b, \mathbf{S}_c, \mathbf{S}_d] \cos^j \frac{\theta}{2} \sin^{4-j} \frac{\theta}{2}, \quad (\text{C1})$$

where  $\hat{R}_j$  is a four-body operator about the four spins  $a, b, c, d$  on a unit plaquette consisting of two triangles, and its form depends on  $j$ . At  $\theta = 0$  this term reduces to the normal ring exchange term of the Hubbard model at half-filling<sup>61</sup>, which takes the form,

$$J_4 \sum_{abcd} ((s_a \cdot s_b)(s_c \cdot s_d) + (s_a \cdot s_d)(s_b \cdot s_c) - (s_a \cdot s_c)(s_b \cdot s_d)). \quad (\text{C2})$$

This conventional ring exchange term is known to give a substantial influence on the magnetism particularly when  $U/t_{\text{eff}}$  becomes small near the metal-insulator phase boundary.

At  $\theta \neq 0$ , the coupling constant is given as  $J_4(\theta) = 80t^4 \cos(2\theta)/U^3$ . To see how much contribution this term gives compared to the second-order terms, we show in Fig. 7 the contour plot of  $J_4(\theta)/\sqrt{D^2 + J^2}$  as a function of  $\theta$  and  $U$  where we set  $t_{\text{eff}} = 1$ . At around




FIG. 7. Density (contour) plot of  $J_4/\sqrt{D^2 + J^2}$  on the plane of  $\theta$  and  $U$ , where we set  $t_{\text{eff}} = 1$  and compared the ratio of the major fourth-order ring exchange interaction and the 2nd order magnetic interaction  $\sqrt{D^2 + J^2} = A = 4t_{\text{eff}}^2/U$ .

$U = 8, |J_4|/\sqrt{D^2 + J^2} \lesssim 0.2$ , showing that the effect of ring exchange interaction is small for the insulating phase. However, when  $U = 5$  the contribution from the fourth-order terms are similar to the second order terms, while since the system is in the metallic region at  $\theta \lesssim 0.6$ , these magnetic interactions no longer make sense. We may interpret the enhancement of  $J_4$  as having an itinerancy in the magnetism, which increases the quantum fluctuation effect. Then, based on this analysis, the comparison between the phase diagrams in Figs. 1(b) and 1(c) (main text) can be made; although the magnetic phases become less distinct for the smaller  $U$ , the basic features of magnetism do not differ much. Therefore, we can conclude that the fourth-order terms may play only a secondary role for the stabilization of the skyrmion phases.

\* makuta-r@g.ecc.u-tokyo.ac.jp

† chisa@phys.c.u-tokyo.ac.jp

<sup>1</sup> Skyrme, T. H. A non-linear field theory. *Proc. R. Soc. Lond. A* **260**, 127—138 (1961).

<sup>2</sup> Sondhi, S. L., Karlhede, A., Kivelson, S. A. & Rezayi, E. H.

Skyrmions and the crossover from the integer to fractional quantum Hall effect at small Zeeman energies. *Phys. Rev. B* **47**, 16419–16426 (1993).

<sup>3</sup> Tycko, R., Barrett, S. E., Dabbagh, G., Pfeiffer, L. N. & West, K. W. Electronic states in gallium arsenide quantum

- wells probed by optically pumped NMR. *Science* **268**, 1460–1463 (1995).
- <sup>4</sup> Bogdanov, A. Thermodynamically stable vortexes in magnetically ordered crystals-mixed state of magnetics. *Zh. Eksp. Teor. Fiz.* **95**, 178–182 (1989).
  - <sup>5</sup> Mühlbauer, S. *et al.* Skyrmion Lattice in a Chiral Magnet. *Science* **323**, 915–919 (2009).
  - <sup>6</sup> Neubauer, A. *et al.* Topological Hall Effect in the A Phase of MnSi. *Phys. Rev. Lett.* **102**, 186602 (2009).
  - <sup>7</sup> Yu, X. Z. *et al.* Real-space observation of a two-dimensional skyrmion crystal. *Nature* **465**, 901–904 (2010).
  - <sup>8</sup> Yu, X. Z. *et al.* Near room-temperature formation of a skyrmion crystal in thin-films of the helimagnet FeGe. *Nature Materials* **10**, 106–109 (2011).
  - <sup>9</sup> Seki, S., Yu, X. Z., Ishiwata, S. & Tokura, Y. Observation of Skyrmions in a Multiferroic Material. *Science* **336**, 198–201 (2012).
  - <sup>10</sup> Tokunaga, Y. *et al.* A new class of chiral materials hosting magnetic skyrmions beyond room temperature. *Nature Communications* **6**, 7638 (2015).
  - <sup>11</sup> Nagaosa, N. & Tokura, Y. Topological properties and dynamics of magnetic skyrmions. *Nature Nanotechnology* **8**, 899–911 (2013).
  - <sup>12</sup> Kézsmárki, I. *et al.* Néel-type skyrmion lattice with confined orientation in the polar magnetic semiconductor GaV<sub>4</sub>S<sub>8</sub>. *Nature Materials* **14**, 1116–1122 (2015).
  - <sup>13</sup> Fujima, Y., Abe, N., Tokunaga, Y. & Arima, T. Thermodynamically stable skyrmion lattice at low temperatures in a bulk crystal of lacunar spinel GaV<sub>4</sub>Se<sub>8</sub>. *Phys. Rev. B* **95**, 180410 (2017).
  - <sup>14</sup> Butykai, Á. *et al.* Characteristics of ferroelectric-ferroelastic domains in Néel-type skyrmion host GaV<sub>4</sub>S<sub>8</sub>. *Scientific Reports* **7**, 44663 (2017).
  - <sup>15</sup> Kurumaji, T. *et al.* Néel-Type Skyrmion Lattice in the Tetragonal Polar Magnet VOSe<sub>2</sub>O<sub>5</sub>. *Phys. Rev. Lett.* **119**, 237201 (2017).
  - <sup>16</sup> Kurumaji, T. *et al.* Skyrmion lattice with a giant topological Hall effect in a frustrated triangular-lattice magnet. *Science* **365**, 914–918 (2019).
  - <sup>17</sup> Hirschberger, M. *et al.* Skyrmion phase and competing magnetic orders on a breathing kagomé lattice. *Nature Commun.* **10**, 5831 (2019).
  - <sup>18</sup> Okubo, T., Chung, S. & Kawamura, H. Multiple- $q$  States and the Skyrmion Lattice of the Triangular-Lattice Heisenberg Antiferromagnet under Magnetic Fields. *Phys. Rev. Lett.* **108**, 017206 (2012).
  - <sup>19</sup> Hayami, S., Ozawa, R. & Motome, Y. Effective bilinear-biquadratic model for noncoplanar ordering in itinerant magnets. *Phys. Rev. B* **95**, 224424 (2017).
  - <sup>20</sup> Wang, Z., Su, Y., Lin, S.-Z. & Batista, C. D. Skyrmion Crystal from RKKY Interaction Mediated by 2D Electron Gas. *Phys. Rev. Lett.* **124**, 207201 (2020).
  - <sup>21</sup> Mitsumoto, K. & Kawamura, H. Replica symmetry breaking in the RKKY skyrmion-crystal system. *Phys. Rev. B* **104**, 184432 (2021).
  - <sup>22</sup> Yi, S. D., Onoda, S., Nagaosa, N. & Han, J. H. Skyrmions and anomalous Hall effect in a Dzyaloshinskii-Moriya spiral magnet. *Phys. Rev. B* **80**, 054416 (2009).
  - <sup>23</sup> Buhrandt, S. & Fritz, L. Skyrmion lattice phase in three-dimensional chiral magnets from Monte Carlo simulations. *Phys. Rev. B* **88**, 195137 (2013).
  - <sup>24</sup> Witczak-Krempa, W., Chen, G., Kim, Y. B. & Balents, L. Correlated Quantum Phenomena in the Strong Spin-Orbit Regime. *Annual Review of Condensed Matter Physics* **5**, 57–82 (2014).
  - <sup>25</sup> Nakai, H. & Hotta, C. Perfect flat band with chirality and charge ordering out of strong spin-orbit interaction. *Nature Communications* **13** (2022).
  - <sup>26</sup> Kawano, M. & Hotta, C. Sine-square deformed mean-field theory. *Phys. Rev. Res.* **4**, L012033 (2022).
  - <sup>27</sup> Kawano, M. & Hotta, C. Phase diagram of the square-lattice Hubbard model with Rashba-type antisymmetric spin-orbit coupling. *Phys. Rev. B* **107**, 045123 (2023).
  - <sup>28</sup> Hotta, C., Nishimoto, S. & Shibata, N. Grand canonical finite size numerical approaches in one and two dimensions: Real space energy renormalization and edge state generation. *Phys. Rev. B* **87**, 115128 (2013).
  - <sup>29</sup> In standard mean-field approximation the magnetic insulators are overstabilized so that the UUD state shall appear at  $B = 0$  in both diagrams. However, SSDMF safely captures the paramagnetic ground state observed in previous studies taking account of the correlation effects at their maximum, indicating that it includes the higher-order correlation effect from the idea of energy renormalization done by SSD<sup>28</sup>. Indeed the 120°-based states appear in our phase diagram at the comparable  $U/t \sim 10$  as Ref.[32].
  - <sup>30</sup> Morita, H., Watanabe, S. & Imada, M. Nonmagnetic insulating states near the Mott transitions on lattice with geometrical frustration and implications for  $\kappa$ -ET<sub>2</sub>Cu<sub>2</sub>(CN)<sub>3</sub>. *J. Phys. Soc. Jpn.* **71**, 2109 (2002).
  - <sup>31</sup> Yoshioka, T., Koga, A. & Kawakami, N. Quantum Phase Transitions in the Hubbard Model on a Triangular Lattice. *Phys. Rev. Lett.* **103**, 036401 (2009).
  - <sup>32</sup> Shirakawa, T., Tohyama, T., Kokalj, J., Sota, S. & Yunoki, S. Ground-state phase diagram of the triangular lattice Hubbard model by the density-matrix renormalization group method. *Phys. Rev. B* **96**, 205130 (2017).
  - <sup>33</sup> Sotnikov, O. M. *et al.* Probing the topology of the quantum analog of a classical skyrmion. *Phys. Rev. B* **103**, L060404 (2021).
  - <sup>34</sup> Lohani, V., Hickey, C., Masell, J. & Rosch, A. Quantum Skyrmions in Frustrated Ferromagnets. *Phys. Rev. X* **9**, 041063 (2019).
  - <sup>35</sup> Mæland, K. & Sudbø, A. Quantum topological phase transitions in skyrmion crystals. *Phys. Rev. Res.* **4**, L032025 (2022).
  - <sup>36</sup> Fukui, T., Hatsugai, Y. & Suzuki, H. Chern Numbers in Discretized Brillouin Zone: Efficient Method of Computing (Spin) Hall Conductances. *J. Phys. Soc. Jpn.* **74**, 1674–1677 (2005).
  - <sup>37</sup> Nayak, A. K. *et al.* Magnetic antiskyrmions above room temperature in tetragonal Heusler materials. *Nature* **548**, 561–566 (2017).
  - <sup>38</sup> Peng, L. *et al.* Controlled transformation of skyrmions and antiskyrmions in a non-centrosymmetric magnet. *Nature Nanotechnology* **15**, 181–186 (2020).
  - <sup>39</sup> Shibata, K. *et al.* Towards control of the size and helicity of skyrmions in helimagnetic alloys by spin-orbit coupling. *Nature Nanotechnology* **8**, 723–728 (2013).
  - <sup>40</sup> Grigoriev, S. V. *et al.* Chiral Properties of Structure and Magnetism in Mn<sub>1-x</sub>Fe<sub>x</sub>Ge Compounds: When the Left and the Right are Fighting, Who Wins? *Phys. Rev. Lett.* **110**, 207201 (2013).
  - <sup>41</sup> Kanazawa, N. *et al.* Large Topological Hall Effect in a Short-Period Helimagnet MnGe. *Phys. Rev. Lett.* **106**, 156603 (2011).

- <sup>42</sup> Tanigaki, T. *et al.* Real-Space Observation of Short-Period Cubic Lattice of Skyrmions in MnGe. *Nano Letters* **15**, 5438–5442 (2015).
- <sup>43</sup> Adams, T. *et al.* Long-Wavelength Helimagnetic Order and Skyrmion Lattice Phase in Cu<sub>2</sub>OSeO<sub>3</sub>. *Phys. Rev. Lett.* **108**, 237204 (2012).
- <sup>44</sup> Fujishiro, Y. *et al.* Topological transitions among skyrmion- and hedgehog-lattice states in cubic chiral magnets. *Nature Communications* **10**, 1059 (2019).
- <sup>45</sup> Karube, K. *et al.* Controlling the helicity of magnetic skyrmions in a  $\beta$ -Mn-type high-temperature chiral magnet. *Phys. Rev. B* **98**, 155120 (2018).
- <sup>46</sup> Li, W. *et al.* Emergence of skyrmions from rich parent phases in the molybdenum nitrides. *Phys. Rev. B* **93**, 060409 (2016).
- <sup>47</sup> Kaneko, K. *et al.* Unique Helical Magnetic Order and Field-Induced Phase in Trillium Lattice Antiferromagnet EuPtSi. *Journal of the Physical Society of Japan* **88**, 013702 (2019).
- <sup>48</sup> Kakihana, M. *et al.* Giant Hall Resistivity and Magnetoresistance in Cubic Chiral Antiferromagnet EuPtSi. *Journal of the Physical Society of Japan* **87**, 023701 (2018).
- <sup>49</sup> Heinze, S. *et al.* Spontaneous atomic-scale magnetic skyrmion lattice in two dimensions. *Nature Physics* **7**, 713–718 (2011).
- <sup>50</sup> Romming, N. *et al.* Writing and Deleting Single Magnetic Skyrmions. *Science* **341**, 636–639 (2013).
- <sup>51</sup> Moreau-Luchaire, C. *et al.* Additive interfacial chiral interaction in multilayers for stabilization of small individual skyrmions at room temperature. *Nature Nanotechnology* **11**, 444–448 (2016).
- <sup>52</sup> Yasuda, K. *et al.* Geometric Hall effects in topological insulator heterostructures. *Nature Physics* **12**, 555–559 (2016).
- <sup>53</sup> Matsuno, J. *et al.* Interface-driven topological Hall effect in SrRuO<sub>3</sub>-SrIrO<sub>3</sub> bilayer. *Science Advances* **2** (2016).
- <sup>54</sup> Wang, L. *et al.* Ferroelectrically tunable magnetic skyrmions in ultrathin oxide heterostructures. *Nature Materials* **17**, 1087–1094 (2018).
- <sup>55</sup> Bordács, S. *et al.* Equilibrium Skyrmion Lattice Ground State in a Polar Easy-plane Magnet. *Scientific Reports* **7**, 7584 (2017).
- <sup>56</sup> Ozawa, R., Hayami, S. & Motome, Y. Zero-Field Skyrmions with a High Topological Number in Itinerant Magnets. *Phys. Rev. Lett.* **118**, 147205 (2017).
- <sup>57</sup> Choi, H., Tai, Y. Y. & Zhu, J. X. Spin-fermion model for skyrmions in MnGe derived from strong correlations. *Phys. Rev. B* **99**, 134437 (2019).
- <sup>58</sup> Koelling, D. D. & Harmon, B. A technique for relativistic spin-polarised calculations. *Journal of Physics C: Solid State Physics* **10**, 3107–3114 (1977).
- <sup>59</sup> Kurita, M., Yamaji, Y. & Imada, M. Topological Insulators from Spontaneous Symmetry Breaking Induced by Electron Correlation on Pyrochlore Lattices. *J. Phys. Soc. Jpn.* **80**, 044708 (2011).
- <sup>60</sup> Haller, A., Groenendijk, S., Habibi, A., Michels, A. & Schmidt, T. L. Quantum skyrmion lattices in Heisenberg ferromagnets. *Phys. Rev. Res.* **4**, 043113 (2022).
- <sup>61</sup> Takahashi, M. Half-filled Hubbard model at low temperature. *J. Phys. C* **10**, 1289–7301 (1977).