

ON THE COHOMOLOGY OF THE CLASSIFYING SPACES OF $SO(n)$ -GAUGE GROUPS OVER S^2

YUKI MINOWA

ABSTRACT. Let $\mathcal{G}_\alpha(X, G)$ be the G -gauge group over a space X corresponding to a map $\alpha: X \rightarrow BG$. We compute the integral cohomology of $B\mathcal{G}_1(S^2, SO(n))$ for $n = 3, 4$. We also show that the homology of $B\mathcal{G}_1(S^2, SO(n))$ is torsion free if and only if $n \leq 4$. As an application, we classify the homotopy types of $SO(n)$ -gauge groups over a Riemann surface for $n \leq 4$.

1. INTRODUCTION

Let G be a topological group, and let P be a principal G -bundle over a base space X . Then automorphisms of P are, by definition, G -equivariant self-maps of P covering the identity map of X . The *gauge group* of P is defined as the topological group of automorphisms of P . Let $\mathcal{G}_\alpha(X, G)$ denote the gauge group of the principal G -bundle over X corresponding to a map $\alpha: X \rightarrow BG$.

Gauge groups have been studied in several contexts of geometry, topology, and physics. Here, we briefly recall the homotopy theory of gauge groups. Given G and X , we have a family of principal G -bundles over X , and so we get a family of their gauge groups. There are many previous studies about the classification of the homotopy types of gauge groups in this family. As shown by Gottlieb [6], there is a natural homotopy equivalence

$$(1.1) \quad B\mathcal{G}_\alpha(X, G) \simeq \text{Map}(X, BG; \alpha)$$

where $\text{Map}(X, Y; f)$ denotes the path-component of the space of maps from X to Y containing a map $f: X \rightarrow Y$ (see also [1]). Then the classification of the homotopy types of gauge groups is closely related to the classical problem in algebraic topology, the classification of components of a mapping space [20]. There are some general results on the classification of the homotopy types of gauge groups [10, 12, 14, 22, 25]. Remarkably, Crabb and Sutherland [4] showed that there are only finitely many homotopy types of $\mathcal{G}_\alpha(X, G)$ as α ranges over all maps $X \rightarrow BG$ whenever G is a compact connected Lie group and X is a connected finite complex, while there are possibly infinitely many principal G -bundles over X .

The classification of the homotopy types of gauge groups is generalized to the classification of the A_n -types: Tsutaya [27] showed that there are only finitely many A_n -types for $n < \infty$ whenever G is a compact connected Lie group and X is a finite complex. On the other hand, Kishimoto and Tsutaya [17] showed that there are infinitely many A_∞ -types, i.e. the homotopy types of the classifying spaces, of gauge groups whenever G is a compact connected simple Lie group and $X = S^d$ with $\pi_{d-1}(G) \otimes \mathbb{Q} \neq 0$. In addition, there are many results on the classification for specific G and X [5, 7, 8, 9, 11, 13, 16, 18, 23, 24].

Although there are many results on the classification of the homotopy types of gauge groups, not much is known about their (co)homology. In particular, there are only a few results determining the (co)homology of the classifying spaces of gauge groups. Takeda [21] computed the integral cohomology ring of $B\mathcal{G}_k(S^2, U(n))$. Kishimoto and Theriault [15] proved that, for a compact connected simple Lie group

G of type (n_1, \dots, n_l) and a prime p larger than $n_l + 1$, there is an isomorphism

$$H_*(B\mathcal{G}_k(S^4, G); \mathbb{Z}/p) \cong H_*(BG; \mathbb{Z}/p) \otimes H_*(\Omega^3 G/3; \mathbb{Z}/p)$$

as \mathbb{Z}/p -vector spaces and so the mod p homology of $B\mathcal{G}_k(S^4, G)$ does not depend on the classifying map $k \in \mathbb{Z} \cong \pi_3(G)$. Partial results on the mod p (co)homology of $B\mathcal{G}_\alpha(X, G)$ for specific G and X are given by Choi [3], Masbaum [19] and Tsukuda [26]. As of today, Tsukuda's result [26] is the only nontrivial result on the integral or mod p (co)homology of $B\mathcal{G}_\alpha(S^d, G)$ for a Lie group G that has p -torsion in its homology.

In this paper, we study the cohomology of $B\mathcal{G}_k(S^2, SO(n))$ for $n \geq 3$, where $k = 0, 1 \in \mathbb{Z}/2 \cong \pi_1(SO(n))$. We will always assume $n \geq 3$ as long as we consider $SO(n)$. First, we will apply the result of Takeda [21] on the integral cohomology of $B\mathcal{G}_k(S^2, U(n))$ for determining the integral cohomology of $B\mathcal{G}_1(S^2, SO(n))$ for $n \leq 4$, where the result of Takeda includes a minor mistake and we will correct it. To state the result, we set notation. Let A be a graded abelian group of finite type. The *rational closure* of a subgroup B of A is defined by

$$B_{\mathcal{R}} = \{a \in A \mid na \in B \text{ for some } 0 \neq n \in \mathbb{Z}\}.$$

We also define a polynomial $s_n \in \mathbb{Z}[x_1, \dots, x_n]$ inductively by

$$(1.2) \quad s_0 = 1, \quad s_1 = x_1 \quad \text{and} \quad s_n = (-1)nx_n + \sum_{i=1}^{n-1} (-1)^{n+i+1} x_{n-i} s_i.$$

So if x_i is the i -th elementary symmetric function in t_1, \dots, t_n , then for $k \leq n$, we have

$$s_k = t_1^k + \dots + t_n^k.$$

Now we are ready to state the first result.

Theorem 1.1. (1) *There is an isomorphism*

$$H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z}) \cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / ((h_2, h_3, \dots)_{\mathcal{R}} + (x_1))$$

$$\text{where } |c_i| = |x_i| = 2i \text{ and } h_i = s_i - s_{i-1}c_1 + s_{i-2}c_2.$$

(2) *There is an isomorphism*

$$H^*(B\mathcal{G}_1(S^2, SO(4)); \mathbb{Z}) \cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] \otimes \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / I$$

$$\text{where } |c_i| = |x_i| = 2i, \quad h_i = s_i - s_{i-1}c_1 + s_{i-2}c_2 \text{ and}$$

$$I = (h_2 \otimes 1, 1 \otimes h_2, h_3 \otimes 1, 1 \otimes h_3, \dots)_{\mathcal{R}} + (c_1 \otimes 1 - 1 \otimes c_1, x_1 \otimes 1).$$

Next, as an application of Theorem 1.1, we will generalize the result of Tsukuda [26], in which he showed that the homology of the classifying space $B\mathcal{G}_1(S^2, SO(3))$ is torsion free.

Theorem 1.2. *The homology of $B\mathcal{G}_1(S^2, SO(n))$ is torsion free if and only if $n \leq 4$.*

Finally, as another application of Theorem 1.1, we will compute $\pi_1(\mathcal{G}_k(S^2, SO(n)))$ for $n \leq 4$. Then by combining with Theriault's homotopy decomposition of a gauge group over a Riemann surface [22], we can determine $\pi_1(\mathcal{G}_k(\Sigma_g, SO(n)))$ for $n \leq 4$ as follows, where Σ_g denotes a Riemann surface of genus g . Note that the homotopy set $[\Sigma_g, BSO(n)]$ is isomorphic to $\mathbb{Z}/2$.

Theorem 1.3. *For $n \leq 4$, there are isomorphisms*

$$\pi_1(\mathcal{G}_k(\Sigma_g, SO(n))) \cong \begin{cases} \mathbb{Z}^{n-2} \oplus \mathbb{Z}/2 & (k = 0) \\ \mathbb{Z}^{n-2} & (k = 1). \end{cases}$$

Then we get the following immediate corollary.

Corollary 1.4. *For $n \leq 4$, $\mathcal{G}_k(\Sigma_g, SO(n)) \simeq \mathcal{G}_l(\Sigma_g, SO(n))$ if and only if $k = l$.*

This paper is structured as follows. Section 2 computes the integral cohomology of $B\mathcal{G}_k(S^2, Spin^c(n))$ for $n = 3, 4$ by correcting the result of Takeda [21] on the integral cohomology of $B\mathcal{G}_k(S^2, U(n))$. Section 3 computes the integral cohomology ring of $B\mathcal{G}_1(S^2, SO(n))$ for $n = 3, 4$ and proves Theorem 1.1. Section 4 computes the mod 2 cohomology of $B\mathcal{G}_1(S^2, SO(n))$ for $n \geq 5$ and gives a proof of Theorem 1.2. Finally, Section 5 proves Theorem 1.3 as an application of the proof of Theorem 1.1.

2. COHOMOLOGY OF $B\mathcal{G}_k(S^2, Spin^c(n))$ FOR $n = 3, 4$

This section computes the integral cohomology of $B\mathcal{G}_k(S^2, Spin^c(n))$ for $n = 3, 4$ by correcting the result of Takeda [21] on the integral cohomology of $B\mathcal{G}_k(S^2, U(n))$. Recall that $Spin^c(n)$ is defined as the nontrivial extension of $SO(n)$ by S^1 :

$$1 \rightarrow S^1 \xrightarrow{\alpha} Spin^c(n) \xrightarrow{\pi} SO(n) \rightarrow 1.$$

This short exact sequence of Lie groups is natural with respect to the inclusions $i'_n: Spin^c(n) \rightarrow Spin^c(n+1)$ and $i_n: SO(n) \rightarrow SO(n+1)$. Then, in particular, there are exceptional isomorphisms:

$$Spin^c(3) \cong U(2) \quad \text{and} \quad Spin^c(4) \cong \{(A, B) \in U(2) \times U(2) \mid \det A = \det B\}$$

where $\alpha: S^1 \rightarrow Spin^c(3)$ and $i'_3: Spin^c(3) \rightarrow Spin^c(4)$ are given by

$$\alpha(e^{i\theta}) = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{i\theta} \end{pmatrix}, \quad i'_3(A) = (A, A).$$

So we can get the cohomology of $B\mathcal{G}_k(S^2, Spin^c(n))$ for $n = 3, 4$ from that of $B\mathcal{G}_k(S^2, U(2))$.

Takeda [21] described the integral cohomology of $B\mathcal{G}_k(S^2, U(n))$, but it includes a mistake. Then we correct it. Bott [2] studied the homology of the loop space of a Lie group, and in particular, we have

Proposition 2.1. *There is an isomorphism*

$$H^*(\Omega_0 U(n); \mathbb{Z}) \cong \mathbb{Z}[x_1, x_2, \dots] / (s_n, s_{n+1}, \dots)_{\mathcal{R}}, \quad |x_i| = 2i$$

where s_n is as in (1.2).

By (1.1), we have $B\mathcal{G}_k(S^2, U(n)) \simeq \text{Map}(S^2, BU(n); k)$. Then there is a homotopy fibration

$$(2.1) \quad \Omega_0 U(n) \xrightarrow{\iota} B\mathcal{G}_k(S^2, U(n)) \xrightarrow{\epsilon} BU(n)$$

given by the evaluation at the basepoint of S^2 . Recall that the cohomology of $BU(n)$ is given by

$$H^*(BU(n); \mathbb{Z}) \cong \mathbb{Z}[c_1, \dots, c_n], \quad |c_i| = 2i$$

where c_i is the universal i -th Chern class. Then

$$H^{\text{odd}}(BU(n); \mathbb{Z}) = 0 \quad \text{and} \quad H^{\text{odd}}(\Omega_0 U(n); \mathbb{Z}) = 0$$

by Proposition 2.1. Thus the integral cohomology Serre spectral sequence for the above homotopy fibration collapses at the E_2 -term and we get

Lemma 2.2. *The integral cohomology of $B\mathcal{G}_k(S^2, U(n))$ is a free abelian group and generated by $c_i = e^*(c_i)$ for $i = 1, \dots, n$ and x_j for $j \geq 1$ as an algebra, where $\iota^*(x_j) = x_j$ in Proposition 2.1.*

Let $c_0 = 1$ and $c_i = 0$ for $i > n$. We define

$$h_i = kc_i + \sum_{j=1}^i (-1)^j s_j c_{i-j} \quad \text{for } i \geq 1.$$

Takeda [21] showed that $h_i = 0$ in $H^*(B\mathcal{G}_k(S^2, U(n)); \mathbb{Z})$ for $i \geq n$ by constructing the element $x_j \in H^{2j}(B\mathcal{G}_k(S^2, U(n)); \mathbb{Z})$ explicitly. Thus by Lemma 2.2, there is a surjection

$$\mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots) \rightarrow H^*(B\mathcal{G}_k(S^2, U(n)); \mathbb{Z}).$$

Takeda [21] claimed that the surjection above is an isomorphism, which is false. To see this, let $n = 2$ and $k = 1$. Then

$$h_4 - c_1 h_3 - (c_1 x_1 + x_1^2) h_2 = 2y$$

where

$$y = c_1^2 x_2 + c_1 x_1 x_2 - x_1^2 x_2 - c_2 x_2 + x_2^2 + 2x_1 x_3 - 2x_4.$$

However, since the coefficient of x_4 in $f \in (h_2, h_3, \dots)$ is a multiple of 4, we have $y \notin (h_2, h_3, \dots)$. Thus the image of y in $\mathbb{Z}[c_1, c_2, x_1, x_2, \dots]/(h_2, h_3, \dots)$ is of order 2, while by Lemma 2.2, $H^*(B\mathcal{G}_1(S^2, U(2)); \mathbb{Z})$ is a free abelian group.

Note that, for a free abelian group A and its subgroup B , $B_{\mathcal{R}}$ is the minimal direct summand of A containing B . Then $\mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots)_{\mathcal{R}}$ is the maximal free abelian subgroup of $\mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots)$. Thus by Lemma 2.2, we have a surjection

$$(2.2) \quad \mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots)_{\mathcal{R}} \rightarrow H^*(B\mathcal{G}_k(S^2, U(n)); \mathbb{Z}).$$

Theorem 2.3. *There is an isomorphism*

$$H^*(B\mathcal{G}_k(S^2, U(n)); \mathbb{Z}) \cong \mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots)_{\mathcal{R}}.$$

Proof. Since the rational cohomology Serre spectral sequence for the homotopy fibration (2.1) collapses at the E_2 -term as above, it follows from Proposition 2.1 that the Poincaré series of the rational cohomology of $B\mathcal{G}_k(S^2, U(n))$ is given by

$$\prod_{i=1}^n \frac{1}{1-t^{2i}} \prod_{i=1}^{n-1} \frac{1}{1-t^{2i}}.$$

On the other hand, one can easily see that h_n, h_{n+1}, \dots is a regular sequence in $\mathbb{Q}[c_1, \dots, c_n, x_1, x_2, \dots]$ and so $(\mathbb{Z}[c_1, \dots, c_n, x_1, x_2, \dots]/(h_n, h_{n+1}, \dots)_{\mathcal{R}}) \otimes \mathbb{Q}$ has the same Poincaré series as $B\mathcal{G}_k(S^2, U(n))$. Thus the map (2.2) is injective, hence an isomorphism. \square

Corollary 2.4. *There is an isomorphism*

$$\begin{aligned} H^*(B\mathcal{G}_k(S^2, Spin^c(3)); \mathbb{Z}) &\cong H^*(B\mathcal{G}_k(S^2, U(2)); \mathbb{Z}) \\ &\cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots]/(h_2, h_3, \dots)_{\mathcal{R}}. \end{aligned}$$

Next, we compute the integral cohomology of $B\mathcal{G}_k(S^2, Spin^c(4))$. Recall that there is a short exact sequence of Lie groups

$$1 \rightarrow Spin^c(4) \xrightarrow{\alpha} U(2) \times U(2) \xrightarrow{\pi} S^1 \rightarrow 1$$

where α is the inclusion and $\pi(A, B) = \det(AB^{-1})$. We denote the composite map $S^2 \xrightarrow{k} BSpin^c(4) \xrightarrow{\alpha} BU(2) \times BU(2) \xrightarrow{p} BU(2)$ by \underline{k} , where p is the first projection. Since the induced map $\Omega\alpha: \Omega_0 Spin^c(4) \rightarrow \Omega_0(U(2) \times U(2))$ is a homotopy equivalence, we have

Lemma 2.5. *There is a homotopy commutative diagram*

$$\begin{array}{ccccc}
\Omega_0 Spin^c(4) & \xrightarrow{\cong} & \Omega_0(U(2) \times U(2)) & \longrightarrow & * \\
\downarrow \iota & & \downarrow \iota & & \downarrow \\
B\mathcal{G}_k(S^2, Spin^c(4)) & \xrightarrow{\alpha_*} & B\mathcal{G}_{(\underline{k}, \underline{k})}(S^2, U(2) \times U(2)) & \xrightarrow{\pi_*} & B\mathcal{G}_0(S^2, S^1) \\
\downarrow e & & \downarrow e & & \downarrow \wr \\
BSpin^c(4) & \xrightarrow{\alpha} & BU(2) \times BU(2) & \xrightarrow{\pi} & BS^1
\end{array}$$

where all columns and rows are homotopy fibrations.

Lemma 2.6. $H^*(BSpin^c(4); \mathbb{Z})$ is a free abelian group and there is an isomorphism

$$H^*(BSpin^c(4); \mathbb{Z}) \cong \mathbb{Z}[c_1, c_2] \otimes \mathbb{Z}[c_1, c_2] / (c_1 \otimes 1 - 1 \otimes c_1).$$

Proof. First, we consider $H^*(BSpin^c(4); \mathbb{Z})$. Consider the short exact sequence of Lie groups

$$1 \rightarrow SU(2) \times SU(2) \hookrightarrow Spin^c(4) \rightarrow S^1 \rightarrow 1.$$

Then there is a homotopy fibration

$$BSU(2) \times BSU(2) \rightarrow BSpin^c(4) \rightarrow BS^1.$$

Since

$$H^{\text{odd}}(BSU(2) \times BSU(2); \mathbb{Z}) = 0 \quad \text{and} \quad H^{\text{odd}}(BS^1; \mathbb{Z}) = 0,$$

the integral cohomology Serre spectral sequence for the homotopy fibration collapses at the E_2 -term by degree reasons. Thus we get $H^{\text{odd}}(BSpin^c(4); \mathbb{Z}) = 0$. Since both $H^*(BSU(2) \times BSU(2); \mathbb{Z})$ and $H^*(BS^1; \mathbb{Z})$ are free abelian groups, $H^*(BSpin^c(4); \mathbb{Z})$ is also a free abelian group. Next, we consider the homotopy fibration of the bottom row of the diagram in Lemma 2.5

$$BSpin^c(4) \xrightarrow{\alpha} BU(2) \times BU(2) \xrightarrow{\pi} BS^1.$$

Recall that $H^{\text{odd}}(BSpin^c(4); \mathbb{Z}) = 0$. By degree reasons, the integral cohomology Serre spectral sequence for the homotopy fibration collapses at the E_2 -term. Since $\pi^*(c_1) = c_1 \otimes 1 - 1 \otimes c_1$, we get the isomorphism in the statement. \square

Proposition 2.7. $H^*(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z})$ is a free abelian group and there is an isomorphism

$$H^*(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z}) \cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] \otimes \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / I$$

where

$$I = (h_2 \otimes 1, 1 \otimes h_2, h_3 \otimes 1, 1 \otimes h_3, \dots)_{\mathbb{R}} + (c_1 \otimes 1 - 1 \otimes c_1).$$

Proof. Recall that

$$H^{\text{odd}}(\Omega_0 Spin^c(4); \mathbb{Z}) \cong H^{\text{odd}}(\Omega S^3 \times \Omega S^3; \mathbb{Z}) = 0.$$

By Lemma 2.6, $H^{\text{odd}}(BSpin^c(4); \mathbb{Z}) = 0$. Then we get

$$H^{\text{odd}}(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z}) = 0,$$

since the integral cohomology Serre spectral sequence for the homotopy fibration of the left column of the diagram in Lemma 2.5

$$\Omega_0 Spin^c(4) \xrightarrow{\iota} B\mathcal{G}_k(S^2, Spin^c(4)) \xrightarrow{e} BSpin^c(4)$$

collapses at the E_2 -term by degree reasons. Since both $H^*(\Omega_0 Spin^c(4); \mathbb{Z})$ and $H^*(BSpin^c(4); \mathbb{Z})$ are free abelian groups, $H^*(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z})$ is also a free

abelian group. Then, since $H^{\text{odd}}(BS^1; \mathbb{Z}) = 0$, the integral cohomology Serre spectral sequence for the homotopy fibration of the middle row of the diagram in Lemma 2.5

$$B\mathcal{G}_k(S^2, Spin^c(4)) \xrightarrow{\alpha_*} B\mathcal{G}_{(\underline{k}, \underline{k})}(S^2, U(2) \times U(2)) \xrightarrow{\pi_*} BS^1$$

collapses at the E_2 -term by degree reasons. Furthermore, we have

$$(\pi_*)^*(c_1) = (\pi \circ e)^*(c_1) = e^* \circ \pi^*(c_1) = e^*(c_1 \otimes 1 - 1 \otimes c_1) = c_1 \otimes 1 - 1 \otimes c_1.$$

Thus we get

$$\begin{aligned} & H^*(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z}) \\ & \cong H^*(B\mathcal{G}_{(\underline{k}, \underline{k})}(S^2, U(2) \times U(2)); \mathbb{Z}) / ((\pi_*)^*(c_1)) \\ & \cong H^*(B\mathcal{G}_k(S^2, U(2)); \mathbb{Z}) \otimes H^*(B\mathcal{G}_k(S^2, U(2)); \mathbb{Z}) / (c_1 \otimes 1 - 1 \otimes c_1) \\ & \cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] \otimes \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / I \end{aligned}$$

and the proof is finished. \square

3. COHOMOLOGY OF $B\mathcal{G}_1(S^2, SO(n))$ FOR $n = 3, 4$

This section computes the integral cohomology ring of $B\mathcal{G}_1(S^2, SO(n))$ for $n = 3, 4$ and proves Theorem 1.1. Recall that there is an exact sequence of Lie groups

$$1 \rightarrow S^1 \xrightarrow{\alpha} Spin^c(n) \xrightarrow{\pi} SO(n) \rightarrow 1.$$

For an integer k , let \bar{k} be its mod 2 reduction. Since the induced map $\Omega\pi: \Omega_0 Spin^c(n) \rightarrow \Omega_0 SO(n)$ is a homotopy equivalence, we have

Lemma 3.1. *There is a homotopy commutative diagram*

$$\begin{array}{ccccc} * & \longrightarrow & \Omega_0 Spin^c(n) & \xrightarrow{\cong} & \Omega_0 SO(n) \\ \downarrow & & \downarrow \iota & & \downarrow \iota \\ B\mathcal{G}_0(S^2, S^1) & \xrightarrow{\alpha_*} & B\mathcal{G}_k(S^2, Spin^c(n)) & \xrightarrow{\pi_*} & B\mathcal{G}_{\bar{k}}(S^2, SO(n)) \\ \wr & & \downarrow e & & \downarrow e \\ BS^1 & \xrightarrow{\alpha} & BSpin^c(n) & \xrightarrow{\pi} & BSO(n) \end{array}$$

where all columns and rows are homotopy fibrations.

First, we compute the $n = 3$ case. By Lemma 3.1, there is a homotopy fibration

$$BS^1 \xrightarrow{\alpha_*} B\mathcal{G}_k(S^2, Spin^c(3)) \xrightarrow{\pi_*} B\mathcal{G}_{\bar{k}}(S^2, SO(3)).$$

Let $c_i, x_i \in H^{2i}(B\mathcal{G}_k(S^2, Spin^c(3)); \mathbb{Z})$ be the generators as in Corollary 2.4 and $c_1 \in H^2(BS^1; \mathbb{Z})$ be the first universal Chern class. Recall that, in the short exact sequence of Lie groups

$$1 \rightarrow S^1 \xrightarrow{\alpha} Spin^c(3) \cong U(2) \xrightarrow{\pi} SO(3) \rightarrow 1,$$

$\alpha: S^1 \rightarrow U(2)$ is given by $\alpha(e^{i\theta}) = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{i\theta} \end{pmatrix}$. Then we can take the generators $c_i \in H^{2i}(BSpin^c(3); \mathbb{Z})$ to satisfy $\alpha^*(c_1) = 2c_1$ and $\alpha^*(c_2) = c_1^2$. By applying Lemma 2.2, the following statement holds.

Lemma 3.2. $(\alpha_*)^*(c_1) = 2c_1$ and $(\alpha_*)^*(c_2) = c_1^2$.

Proposition 3.3. $(\alpha_*)^*(x_1) = kc_1$.

Proof. Since c_1 generates $H^2(BS^1; \mathbb{Z})$, there exists an integer l such that

$$(\alpha_*)^*(x_1) = (k - l)c_1.$$

By Corollary 2.4, we have

$$\begin{aligned} kc_2 - s_1c_1 + s_2 &= 0 \\ s_{n-2}c_2 - s_{n-1}c_1 + s_n &= 0 \quad \text{for } n \geq 3 \end{aligned}$$

in $H^{2i}(B\mathcal{G}_k(S^2, Spin^c(3)); \mathbb{Z})$. By Lemma 3.2, we have

$$(\alpha_*)^*(c_1) = 2c_1 \quad \text{and} \quad (\alpha_*)^*(c_2) = c_1^2.$$

Then we get

$$(\alpha_*)^*(s_n) = (k - nl)c_1^n$$

by induction. Recall the Girard-Newton formula

$$s_n = (-1)^n n \sum_{\substack{r_1+2r_2+\dots+nr_n=n \\ r_1 \geq 0, \dots, r_n \geq 0}} \frac{(r_1 + r_2 + \dots + r_n - 1)!}{r_1! r_2! \dots r_n!} \prod_{i=1}^n (-x_i)^{r_i}.$$

Then $s_p - x_1^p$ is divisible by p for p prime. On the other hand,

$$(\alpha_*)^*(s_p - x_1^p) = ((k - pl) - (k - l)^p)c_1^p = (((k - l) - (k - l)^p) - (p - 1)l)c_1^p$$

is divisible by p if and only if l is divisible by p . Then l is divisible by all primes and so we get $l = 0$. Thus

$$(\alpha_*)^*(x_1) = kc_1$$

and the proof is finished. \square

By Proposition 3.3, we have $(\alpha_*)^*(x_1) = c_1$ for $x_1 \in H^2(B\mathcal{G}_1(S^2, Spin^c(3)); \mathbb{Z})$. Now we consider the homotopy fibration

$$BS^1 \xrightarrow{\alpha_*} B\mathcal{G}_1(S^2, Spin^c(3)) \xrightarrow{\pi_*} B\mathcal{G}_1(S^2, SO(3))$$

and prove the first half of Theorem 1.1.

Proof of Theorem 1.1 (1). By the Leray-Hirsch theorem, we get an isomorphism of $H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z})$ -modules

$$\Phi: H^*(BS^1; \mathbb{Z}) \otimes H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z}) \rightarrow H^*(B\mathcal{G}_1(S^2, Spin^c(3)); \mathbb{Z})$$

defined by

$$\Phi\left(\sum c_1^m \otimes b\right) := \sum x_1^m (\pi_*)^*(b)$$

for $b \in H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z})$. Actually, Φ is also a ring isomorphism, since

$$\begin{aligned} \Phi(c_1^{m+m'} \otimes (bb')) &= x_1^{m+m'} (\pi_*)^*(bb') \\ &= (x_1^m (\pi_*)^*(b))(x_1^{m'} (\pi_*)^*(b')) \\ &= \Phi(c_1^m \otimes b)\Phi(c_1^{m'} \otimes b') \end{aligned}$$

for $m, m' \geq 0$ and $b, b' \in H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z})$. Since $\Phi(c_1 \otimes 1) = x_1$, we have

$$H^*(B\mathcal{G}_1(S^2, SO(3)); \mathbb{Z}) \cong (\mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / (h_2, h_3, \dots)_{\mathbb{R}}) / (x_1)$$

and the proof for $n = 3$ is done. \square

Next, we compute the $n = 4$ case. Since the diagram in Lemma 3.1 is natural with respect to the inclusions $i'_n: Spin^c(n) \rightarrow Spin^c(n+1)$ and $i_n: SO(n) \rightarrow SO(n+1)$, we have a homotopy commutative diagram

$$\begin{array}{ccccc} BS^1 & \xrightarrow{\alpha_*} & B\mathcal{G}_1(S^2, Spin^c(3)) & \xrightarrow{\pi_*} & B\mathcal{G}_1(S^2, SO(3)) \\ \parallel & & \downarrow (i'_3)_* & & \downarrow (i_3)_* \\ BS^1 & \xrightarrow{\alpha_*} & B\mathcal{G}_1(S^2, Spin^c(4)) & \xrightarrow{\pi_*} & B\mathcal{G}_1(S^2, SO(4)) \end{array}$$

where all rows are homotopy fibrations.

Proof of Theorem 1.1 (2). By Proposition 2.7, we have an isomorphism

$$H^*(B\mathcal{G}_k(S^2, Spin^c(4)); \mathbb{Z}) \cong \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] \otimes \mathbb{Z}[c_1, c_2, x_1, x_2, \dots] / I$$

where

$$I = (h_2 \otimes 1, 1 \otimes h_2, h_3 \otimes 1, 1 \otimes h_3, \dots)_{\mathbb{R}} + (c_1 \otimes 1 - 1 \otimes c_1).$$

Since the composite $Spin^c(3) \xrightarrow{i'_3} Spin^c(4) \hookrightarrow U(2) \times U(2)$ is the diagonal map, we have

$$((i'_3)_*)^*(c_i \otimes 1) = c_i \quad \text{and} \quad ((i'_3)_*)^*(x_i \otimes 1) = x_i.$$

Thus by Proposition 3.3, we get

$$(\alpha_*)^*(x_1 \otimes 1) = (\alpha_*)^* \circ ((i'_3)_*)^*(x_1 \otimes 1) = (\alpha_*)^*(x_1) = c_1.$$

Then by arguing as in the proof of Theorem 1.1 (1), we get

$$H^*(B\mathcal{G}_1(S^2, SO(4)); \mathbb{Z}) \cong H^*(B\mathcal{G}_1(S^2, Spin^c(4)); \mathbb{Z}) / (x_1 \otimes 1),$$

and the proof for $n = 4$ is complete. \square

Corollary 3.4. *The homology of $B\mathcal{G}_1(S^2, SO(n))$ is torsion free for $n \leq 4$.*

Proof. For $n \leq 4$, there is a module isomorphism

$$\Phi: H^*(BS^1; \mathbb{Z}) \otimes H^*(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}) \rightarrow H^*(B\mathcal{G}_1(S^2, Spin^c(n)); \mathbb{Z})$$

as seen in the proof of Theorem 1.1. Then $H^*(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z})$ is a direct summand of $H^*(B\mathcal{G}_1(S^2, Spin^c(n)); \mathbb{Z})$ which is a free abelian group for $n \leq 4$. Thus $H^*(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z})$ is also a free abelian group, completing the proof. \square

4. TORSION IN THE COHOMOLOGY OF $B\mathcal{G}_1(S^2, SO(n))$

This section gives a proof of Theorem 1.2. We consider the mod 2 cohomology of $B\mathcal{G}_1(S^2, SO(n))$. Let $i_n: SO(n) \rightarrow SO(n+1)$ be the inclusion.

Lemma 4.1. *There are isomorphisms*

$$H^2(\Omega_0 SO(n); \mathbb{Z}/2) \cong \begin{cases} \langle y_1 \rangle & (n = 3) \\ \langle z_1, y_1 \rangle & (n = 4) \\ \langle z_1 \rangle & (n \geq 5) \end{cases}$$

such that $(\Omega i_3)^*(y_1) = y_1$, $(\Omega i_3)^*(z_1) = 0$ and $(\Omega i_n)^*(z_1) = z_1$ for $n \geq 4$.

Proof. Since there is a natural homotopy equivalence $\Omega_0 SO(n) \simeq \Omega Spin(n)$, we prove the statement for $\Omega Spin(n)$. Let $j_n: Spin(n) \rightarrow Spin(n+1)$ be the inclusion. There is a commutative diagram

$$\begin{array}{ccccc} Spin(3) & \xrightarrow{j_3} & Spin(4) & \xrightarrow{j_4} & Spin(5) \\ \parallel & & \parallel & & \parallel \\ Sp(1) & \xrightarrow{\Delta} & Sp(1) \times Sp(1) & \xrightarrow{\rho} & Sp(2) \end{array}$$

where Δ is the diagonal map and ρ is the diagonal inclusion. Then there is a induced homotopy commutative diagram of the corresponding classifying spaces. Let $q \in H^4(BSp(n); \mathbb{Z}/2)$ be the mod 2 reduction of the first symplectic Pontryagin class. Then

$$\rho^*(q) = q \times 1 + 1 \times q \quad \text{and} \quad \Delta^*(q \times 1) = q = \Delta^*(1 \times q).$$

We define $y_1, z_1 \in H^2(\Omega Spin(n); \mathbb{Z}/2)$ by

$$\begin{aligned} y_1 &:= \sigma^2(q) & (n=3) \\ z_1 &:= \sigma^2(q \times 1 + 1 \times q), \quad y_1 := \sigma^2(q \times 1) & (n=4) \\ z_1 &:= \sigma^2(q) & (n \geq 5), \end{aligned}$$

where $\sigma^2: H^4(BSpin(n); \mathbb{Z}/2) \rightarrow H^2(\Omega Spin(n); \mathbb{Z}/2)$ denotes the cohomology double suspension. Since $BSpin(n)$ is 3-connected, σ^2 is an isomorphism. Moreover, we have $(\Omega j_n) \circ \sigma^2 = \sigma^2 \circ j_n$. Thus we get $(\Omega j_3)^*(y_1) = y_1$ and $(\Omega j_3)^*(z_1) = 0$. Since Ωj_n is a $(n-2)$ -equivalence, we also get $(\Omega i_n)^*(z_1) = z_1$ for $n \geq 4$, completing the proof. \square

Let $\{E_r(n), d_r\}$ denote the mod 2 cohomology Serre spectral sequence for the homotopy fibration

$$\Omega_0 SO(n) \xrightarrow{\iota} B\mathcal{G}_1(S^2, SO(n)) \xrightarrow{e} BSO(n).$$

Since $BSO(n)$ is simply-connected, there is an isomorphism

$$E_2(n) \cong H^*(BSO(n); \mathbb{Z}/2) \otimes H^*(\Omega_0 SO(n); \mathbb{Z}/2).$$

Recall that the mod 2 cohomology of $BSO(n)$ is given by

$$H^*(BSO(n); \mathbb{Z}/2) \cong \mathbb{Z}/2[w_2, w_3, \dots, w_n]$$

where $w_i \in H^i(BSO(n); \mathbb{Z}/2)$ is the universal i -th Stiefel-Whitney class. We also recall the classical result of Bott [2].

Lemma 4.2. *Let G be a connected compact Lie group. Then the integral homology of $\Omega_0 G$ is a free abelian group such that $H_{\text{odd}}(\Omega_0 G; \mathbb{Z}) = 0$.*

In particular, we have

$$H^{\text{odd}}(\Omega_0 SO(n); \mathbb{Z}/2) = 0.$$

Hence $E_2(n) = E_3(n)$ by degree reasons.

Lemma 4.3. *For $n \geq 5$, there is an isomorphism*

$$H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}/2) \cong (\mathbb{Z}/2)^2.$$

Proof. We have

$$E_3^{0,2}(n) = E_2^{0,2}(n) = \langle 1 \otimes z_1 \rangle, \quad E_3^{3,0}(n) = E_2^{3,0}(n) = \langle w_3 \otimes 1 \rangle.$$

Let $i_n^*: E_3(n+1) \rightarrow E_3(n)$ be the induced homomorphism. Then we have

$$i_n^*(d_3(1 \otimes z_1)) = d_3(1 \otimes (\Omega i_n)^*(z_1)) \quad \text{in} \quad E_3^{3,0}(n)$$

for $n \geq 3$. Since $i_3^*: E_3^{3,0}(n+1) \rightarrow E_3^{3,0}(n)$ is an isomorphism for $n \geq 3$ and

$$d_3(1 \otimes (\Omega i_3)^*(z_1)) = d_3(1 \otimes 0) = 0 \quad \text{in} \quad E_3^{3,0}(3)$$

by Lemma 4.1, we get

$$d_3(1 \otimes z_1) = 0 \quad \text{in} \quad E_3^{3,0}(n).$$

Thus $d_3: E_3^{0,2}(n) \rightarrow E_3^{3,0}(n)$ is the zero map and so $1 \otimes z_1$ is a permanent cycle. Since $w_2 \otimes 1 \in E_3^{2,0}(n) \cong \mathbb{Z}/2$ is also a permanent cycle by degree reasons, we get

$$\begin{aligned} H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}/2) &\cong E_\infty^{2,0}(n) \oplus E_\infty^{0,2}(n) \\ &\cong E_3^{2,0}(n) \oplus E_3^{0,2}(n) = \langle w_2 \otimes 1, 1 \otimes z_1 \rangle \end{aligned}$$

and the proof is finished. \square

Lemma 4.4. *For $n \geq 5$, there is an isomorphism*

$$H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}) \cong \mathbb{Z}.$$

Proof. Consider the integral cohomology Serre spectral sequence for the homotopy fibration

$$\Omega_0 SO(n) \xrightarrow{L} B\mathcal{G}_1(S^2, SO(n)) \xrightarrow{c} BSO(n).$$

Then its E_2 -term is isomorphic to $H^*(BSO(n); \mathbb{Z}) \otimes H^*(\Omega_0 SO(n); \mathbb{Z})$. Since

$$H^i(BSO(n); \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & (i = 0) \\ 0 & (i = 1, 2) \\ \mathbb{Z}/2 & (i = 3), \end{cases} \quad H^i(\Omega_0 SO(n); \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & (i = 0, 2) \\ 0 & (i = 1) \end{cases}$$

for $n \geq 5$, we have

$$H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}) \cong E_\infty^{0,2} \cong E_3^{0,2} \cong \mathbb{Z}$$

by degree reasons. Thus we get the isomorphism in the statement. \square

Proof of Theorem 1.2. By Lemma 4.3 we have

$$\begin{aligned} (\mathbb{Z}/2)^2 &\cong H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}/2) \\ &\cong (H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}) \otimes \mathbb{Z}/2) \oplus \text{Tor}(H^3(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}), \mathbb{Z}/2) \end{aligned}$$

for $n \geq 5$. Since $H^2(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}) \cong \mathbb{Z}$ by Lemma 4.4, we get

$$\text{Tor}(H^3(B\mathcal{G}_1(S^2, SO(n)); \mathbb{Z}), \mathbb{Z}/2) \cong \mathbb{Z}/2.$$

Thus by the universal coefficient theorem, the integral homology of $B\mathcal{G}_1(S^2, SO(n))$ has 2-torsion for $n \geq 5$. Therefore by Corollary 3.4, the proof is finished. \square

We have so far considered only 2-torsion in the homology of $B\mathcal{G}_1(S^2, SO(n))$. Now we consider p -torsion for p odd prime.

Proposition 4.5. *For p odd prime, $H^{\text{odd}}(B\mathcal{G}_k(S^2, SO(n)); \mathbb{Z}/p) = 0$.*

Proof. Consider the mod p cohomology Serre spectral sequence for the homotopy fibration

$$\Omega_0 SO(n) \xrightarrow{L} B\mathcal{G}_1(S^2, SO(n)) \xrightarrow{c} BSO(n).$$

Then its E_2 -term is isomorphic to $H^*(BSO(n); \mathbb{Z}/p) \otimes H^*(\Omega_0 SO(n); \mathbb{Z}/p)$. By Lemma 4.2 we have $H^{\text{odd}}(\Omega_0 SO(n); \mathbb{Z}/p) = 0$. On the other hand, it is well known that $H^{\text{odd}}(BSO(n); \mathbb{Z}/p) = 0$. Then by degree reasons, the spectral sequence collapses at the E_2 -term and the proof is done. \square

Therefore by applying the universal coefficient theorem, the following statement holds.

Corollary 4.6. *For p odd prime, the homology of $B\mathcal{G}_k(S^2, SO(n))$ has no p -torsion.*

5. HOMOTOPY TYPES OF $\mathcal{G}_k(\Sigma_g, SO(n))$

This section proves Theorem 1.3. We recall the result of Theriault [22].

Proposition 5.1. *There is a homotopy decomposition*

$$\mathcal{G}_k(\Sigma_g, U(n)) \simeq (\Omega U(n))^{2g} \times \mathcal{G}_k(S^2, U(n)).$$

One can easily see that the proof in [22] is also applied to $\mathcal{G}_k(\Sigma_g, SO(n))$. Thus we get a homotopy decomposition

$$(5.1) \quad \mathcal{G}_k(\Sigma_g, SO(n)) \simeq (\mathbb{Z}/2)^{2g} \times (\Omega Spin(n))^{2g} \times \mathcal{G}_k(S^2, SO(n)).$$

Proof of Theorem 1.3. Recall that $\Omega Spin(n)$ is simply-connected. By (5.1), we have

$$\begin{aligned} \pi_1(\mathcal{G}_k(\Sigma_g, SO(n))) &\cong \pi_1((\mathbb{Z}/2)^{2g}) \times \pi_1((\Omega Spin(n))^{2g}) \times \pi_1(\mathcal{G}_k(S^2, SO(n))) \\ &\cong \pi_1(\mathcal{G}_k(S^2, SO(n))). \end{aligned}$$

Since $\mathcal{G}_k(S^2, SO(n))$ is connected,

$$\pi_1(\mathcal{G}_k(S^2, SO(n))) \cong \pi_2(B\mathcal{G}_k(S^2, SO(n))) \cong H_2(B\mathcal{G}_k(S^2, SO(n)); \mathbb{Z})$$

by the Hurewicz theorem. Thus it is sufficient to prove

$$H_2(B\mathcal{G}_k(S^2, SO(n)); \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^{n-2} \oplus \mathbb{Z}/2 & (k=0) \\ \mathbb{Z}^{n-2} & (k=1). \end{cases}$$

The $k=1$ case follows from Theorem 1.1 and the universal coefficient theorem. Let $\{E'_r(n), d'_r\}$ denote the integral cohomology Serre spectral sequence for the homotopy fibration of the middle row of the diagram in Lemma 3.1

$$BS^1 \xrightarrow{\alpha_*} B\mathcal{G}_0(S^2, Spin^c(n)) \xrightarrow{\pi_*} B\mathcal{G}_0(S^2, SO(n)).$$

Since $BSO(n)$ is simply-connected, there is an isomorphism

$$E'_2(n) \cong H^*(B\mathcal{G}_0(S^2, SO(n)); \mathbb{Z}) \otimes H^*(BS^1; \mathbb{Z})$$

Then, since $H^{\text{odd}}(BS^1; \mathbb{Z}) = 0$, $E'_2(n) = E'_3(n)$ by degree reasons. Let $c_i, x_i \in H^{2i}(B\mathcal{G}_0(S^2, Spin^c(3)); \mathbb{Z})$ be the generators as in Corollary 2.4 and $c_1 \in H^2(BS^1; \mathbb{Z})$ be the first Chern class. For $n=3$, we have

$$(\alpha_*)^*(x_1) = 0$$

by Proposition 3.3. Since $(\alpha_*)^*(c_1) = 2c_1$ by Lemma 3.2, there is an element $x \in H^2(B\mathcal{G}_0(S^2, SO(3)); \mathbb{Z})$ such that

$$(\pi_*)^*(x) = x_1$$

and there is an element $w \in H^3(B\mathcal{G}_0(S^2, SO(3)); \mathbb{Z})$ of order 2 such that

$$d_3(1 \otimes c_1) = w \otimes 1.$$

By degree reasons, we have

$$H^i(B\mathcal{G}_0(S^2, SO(3)); \mathbb{Z}) \cong \begin{cases} \langle x \rangle \cong \mathbb{Z} & (i=2) \\ \langle w \rangle \cong \mathbb{Z}/2 & (i=3). \end{cases}$$

Thus by the universal coefficient theorem, we get

$$H_2(B\mathcal{G}_0(S^2, SO(3)); \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}/2.$$

For the $n=4$ case, we get

$$H^i(B\mathcal{G}_0(S^2, SO(4)); \mathbb{Z}) \cong \begin{cases} \mathbb{Z}^2 & (i=2) \\ \mathbb{Z}/2 & (i=3) \end{cases}$$

by arguing as in the $n = 3$ case. Thus

$$H_2(B\mathcal{G}_0(S^2, SO(4)); \mathbb{Z}) \cong \mathbb{Z}^2 \oplus \mathbb{Z}/2$$

and the proof is complete. \square

REFERENCES

- [1] M. F. Atiyah and R. Bott, The Yang–Mills equations over Riemann surfaces, *Philos. Trans. R. Soc. London, Ser. A* **308** (1983), 523–615. [1](#)
- [2] R. Bott, The space of loops on a Lie group, *Michigan Math. J.* **5** (1958), no. 1, 35–61. [3](#), [9](#)
- [3] Y. Choi, Homology of the classifying space of $Sp(n)$ Gauge groups, *Isr. J. Math.* **151** (2006), 167–177. [2](#)
- [4] M.C. Crabb and W.A. Sutherland, Counting homotopy types of gauge groups, *Proc. London Math. Soc.* **81** (2000), no. 3, 747–768. [1](#)
- [5] T. Cutler, The homotopy types of $U(n)$ -gauge groups over S^4 and $\mathbb{C}P^2$, *Homol. Homotopy Appl.* **20** (2018), no. 1, 5–36. [1](#)
- [6] D. H. Gottlieb, Applications of bundle map theory, *Trans. Am. Math. Soc.* **171** (1972), 23–50. [1](#)
- [7] H. Hamanaka and A. Kono, Unstable K^1 -group and homotopy type of certain gauge groups, *Proc. R. Soc. Edinb. A: Math.* **136** (2006), no. 1, 149–155. [1](#)
- [8] H. Hamanaka and A. Kono, Homotopy type of gauge groups of $SU(3)$ -bundles over S^6 , *Topol. Appl.* **154** (2007), no. 7, 1377–1380. [1](#)
- [9] S. Hasui, D. Kishimoto, A. Kono, T. Sato, The homotopy types of $PU(3)$ - and $PSp(2)$ -gauge groups, *Algebraic Geom. Topol.* **16** (2016), no. 3, 1813–1825. [1](#)
- [10] M. Kameko, D. Kishimoto, M. Takeda, Homotopy types of gauge groups over Riemann surfaces, to appear. [1](#)
- [11] Y. Kamiyama, D. Kishimoto, A. Kono and S. Tsukuda, Samelson products of $SO(3)$ and applications. *Glasgow Math. J.* **49** (2007), no. 2, 405–409. [1](#)
- [12] D. Kishimoto and A. Kono, Splitting of gauge groups, *Trans. Am. Math. Soc.* **362** (2010), no. 12, 6715–6731. [1](#)
- [13] D. Kishimoto and A. Kono, On the homotopy types of $Sp(n)$ gauge groups, *Algebraic Geom. Topol.* **19** (2019), no. 1, 491–502. [1](#)
- [14] D. Kishimoto, A. Kono and M. Tsutaya, On p -local homotopy types of gauge groups, *Proc. R. Soc. Edinb. A: Math.* **144** (2014), no. 1, 149–160. [1](#)
- [15] D. Kishimoto and S. Theriault, The mod- p homology of the classifying spaces of certain gauge groups, *Proc. R. Soc. Edinb. A: Math.* (2022), 1–13. [1](#)
- [16] D. Kishimoto, S. Theriault and M. Tsutaya, The homotopy types of G_2 -gauge groups, *Topol. Appl.* **228** (2017), 92–107. [1](#)
- [17] D. Kishimoto and M. Tsutaya, Infiniteness of A_∞ -types of gauge groups, *J. Topol.* **9** (2016), 181–191. [1](#)
- [18] A. Kono, A note on the homotopy type of certain gauge groups, *Proc. R. Soc. Edinb. A: Math.* **117** (1991), no. 3–4, 295–297. [1](#)
- [19] G. Masbaum, On the cohomology of the classifying space of the gauge group over some 4-complexes, *Bull. Soc. Math. Fr.* **119** (1991), no. 1, 1–31. [2](#)
- [20] S. B. Smith, The homotopy theory of function spaces: a survey, *Homotopy theory of function spaces and related topics* (American Mathematical Society, Providence RI, 2010), 3–39. [1](#)
- [21] M. Takeda, Cohomology of the classifying spaces of $U(n)$ -gauge groups over the 2-sphere, *Homol. Homotopy Appl.* **23** (2021), no. 1, 17–24. [1](#), [2](#), [3](#), [4](#)
- [22] S. Theriault, Odd primary homotopy decompositions of gauge groups, *Algebraic Geom. Topol.* **10** (2010), 535–564. [1](#), [2](#), [11](#)
- [23] S. Theriault, The homotopy types of $Sp(2)$ -gauge groups, *Kyoto J. Math.* **50** (2010), no. 3, 591–605. [1](#)
- [24] S. Theriault, The homotopy types of $SU(5)$ -gauge groups, *Osaka J. Math.* **52** (2015), no. 1, 15–29. [1](#)
- [25] S. Theriault, Homotopy decompositions of the classifying spaces of pointed gauge groups, *Pac. J. Math.* **300** (2019), no. 1, 215–231. [1](#)
- [26] S. Tsukuda, On the cohomology of the classifying space of a certain gauge group, *Proc. R. Soc. Edinb. A: Math.* **127** (1997), no. 2, 407–409. [2](#)
- [27] M. Tsutaya, Finiteness of A_n -equivalence types of gauge groups, *J. London Math. Soc.* **85** (2012), 142–164. [1](#)