

Three q -Painlevé equations with affine Weyl group symmetry of type $E_7^{(1)}$

By

Hidehito Nagao¹ and Yasuhiko Yamada²
(Akashi College¹ and Kobe University², Japan)

Abstract. Three q -Painlevé type equations are derived through degenerations of the q -Painlevé equation with affine Weyl group symmetry of type $q-E_8^{(1)}$. The three q -Painlevé type equations are associated with different realizations of the same symmetry/surface type $q-E_7^{(1)}/q-A_1^{(1)}$. We give three representations of affine Weyl group actions of type $q-E_7^{(1)}$, and relations among them. The three $q-E_7^{(1)}$ equations are also derived from the three representations respectively.

Key Words and Phrases. q -Painlevé equation, affine Weyl group symmetry of type $E_7^{(1)}$, rational surface of type $A_1^{(1)}$, singular point configuration.

2000 Mathematics Subject Classification Numbers. 14H70, 34M55, 37K20, 39A13.

1 Introduction

In view of the Weyl group symmetry, the discrete Painlevé equations with higher symmetries such as $E_8^{(1)}$, $E_7^{(1)}$, etc. are very complicated in general. Therefore we need some ideas to obtain their explicit forms (see for examples [1, 7, 8]). Namely, those equations admit simple expression only when we choose both the representations and the evolution directions properly.

Conversely, for a given explicit equation, one needs some efforts to understand it from geometric or symmetric points of view. In this paper, we study such problems in case of three $q-E_7^{(1)}$ equations arising from certain degenerations of the simple form of the $q-E_8^{(1)}$ equation [2, 10].

The purpose of this paper is to work out the following studies.

- To derive the three $q-E_7^{(1)}$ equations as degenerations of the $q-E_8^{(1)}$ equation.
- To understand the obtained three $q-E_7^{(1)}$ equations in view of the Weyl group symmetry.
- To clarify the relations among the three $q-E_7^{(1)}$ equations using the Weyl group symmetry.

This paper is organized as follows. In Section 2, we derive the three $q-E_7^{(1)}$ equations by using the degenerations of the $q-E_8^{(1)}$ equation. In Section 3, we give three representations of the affine Weyl group actions for the three $q-E_7^{(1)}$ equations without using the affine Weyl group symmetry, and relations among the three representations. In Section 4, we derive the three equations from the three representations. In Appendix A, we show a

relation between the $q-E_8^{(1)}$ equation given in [2, 10] and ORG form [7]. In Appendix B, we investigate a correspondence between two of the three $q-E_7^{(1)}$ equations. In Appendix C, we give three translations on the root variables corresponding to the three representations.

2 Three $q-E_7^{(1)}$ equations with different realizations of surface type $q-A_1^{(1)}$

In this section we recall the $q-E_8^{(1)}$ equation and will derive three $q-E_7^{(1)}$ equations by taking suitable degeneration limits.

Throughout this section, we use the following notations. Parameters $q, \kappa_1, \kappa_2, v_1, \dots, v_8 \in \mathbb{C}^\times$ are complex parameters with a constraint

$$\kappa_1^2 \kappa_2^2 = q \prod_{i=1}^8 v_i, \quad (2.1)$$

and f, g be dependent variables. Then we consider a time evolution T given as

$$T : (v_1, \dots, v_8, \kappa_1, \kappa_2, f, g) \mapsto (v_1, \dots, v_8, \frac{\kappa_1}{q}, q\kappa_2, \bar{f}, \bar{g}), \quad (2.2)$$

where $\bar{X} = T(X)$ and $\underline{X} = T^{-1}(X)$.

2.1 $q-E_8^{(1)}$ equation

The $q-E_8^{(1)}$ equation^{1 2} is given by (2.2) as the following simple form [2, 10]:

$$\begin{aligned} \frac{\{g - G(\frac{\kappa_1}{z})\}\{\underline{g} - \underline{G}(\frac{\kappa_1}{z})\}}{\{g - G(z)\}\{\underline{g} - \underline{G}(z)\}} &= \frac{U(\frac{\kappa_1}{z})}{U(z)}, \quad \text{for } f = F(z), \\ \frac{\{f - F(\frac{\kappa_2}{z})\}\{\bar{f} - \bar{F}(\frac{\kappa_2}{z})\}}{\{f - F(z)\}\{\bar{f} - \bar{F}(z)\}} &= \frac{U(\frac{\kappa_2}{z})}{U(z)}, \quad \text{for } g = G(z), \end{aligned} \quad (2.3)$$

where

$$F(z) = z + \frac{\kappa_1}{z}, \quad G(z) = z + \frac{\kappa_2}{z}, \quad U(z) = z^{-4} \prod_{i=1}^8 (z - v_i). \quad (2.4)$$

In next subsections, we will consider three degenerations of the $q-E_8^{(1)}$ equation (2.3).

2.2 Degeneration 1

Replacing parameters κ_i, v_j, g by

$$\kappa_i \rightarrow \varepsilon \kappa_i \quad (i = 1, 2), \quad v_j \rightarrow \varepsilon v_j \quad (i = 1, \dots, 4), \quad g \rightarrow \frac{1}{g}, \quad (2.6)$$

¹The $q-E_8^{(1)}$ was firstly presented in [7] and the birational form is more complicated than one given in [10]. See Appendix A for more details.

²See [2, 9, 10] for the Lax equation, and [2, 10] for hypergeometric special solutions of (2.3).

The eight singular points are

$$(f, g) = (F(v_i), G(v_i))_{i=1, \dots, 8} \quad (2.5)$$

on a curve of bidegree $(2, 2)$ with a node. This configuration is the surface type $q-A_0^{(1)}$ in Figure 1.




Figure 1: Surface type $q-A_0^{(1)}$

and taking a degeneration limit $\varepsilon \rightarrow 0$, we obtain the well-known $q-E_7^{(1)}$ equation³ [1]:

$$\begin{aligned} \frac{(fg - \frac{\kappa_1}{\kappa_2})(fg - \frac{q\kappa_1}{\kappa_2})}{(fg - 1)(\bar{f}g - 1)} &= \frac{\prod_{i=1}^4 (f - \frac{\kappa_1}{v_i})}{\prod_{i=5}^8 (f - v_i)}, \\ \frac{(fg - \frac{\kappa_1}{\kappa_2})(\bar{f}g - \frac{\kappa_1}{q\kappa_2})}{(fg - 1)(\bar{f}g - 1)} &= \frac{\prod_{i=1}^4 (g - \frac{v_i}{\kappa_2})}{\prod_{i=5}^8 (g - \frac{1}{v_i})}. \end{aligned} \quad (2.7)$$

The eight singular points are

$$(f, g) = \left(\frac{\kappa_1}{v_i}, \frac{v_i}{\kappa_2} \right)_{i=1}^4, \left(v_i, \frac{1}{v_i} \right)_{i=5}^8, \quad (2.8)$$

on the product of two curves of bidegree $(1, 1)$ and $(1, 1)$. This configuration for surface type $q-A_1^{(1)}$ is given in Figure 2. We call it configuration type 1.




Figure 2: Configuration type 1

2.3 Degeneration 2

Similarly, doing a replacement of parameters

$$\kappa_1 \rightarrow \varepsilon\kappa_1, \quad v_i \rightarrow \varepsilon v_i \quad (i = 1, 2), \quad (2.9)$$

and taking a degeneration limit $\varepsilon \rightarrow 0$, one gets the $q-E_7^{(1)}$ equation⁴ :

$$\begin{aligned} \{g - G(f)\}\{\bar{g} - \bar{G}(f)\} &= \frac{\prod_{i=3}^8 (f - v_i)}{f^2 \prod_{i=1}^2 (f - \frac{\kappa_1}{v_i})}, \\ \frac{(f - \frac{\kappa_2}{z_s})(\bar{f} - \frac{\kappa_2}{z_s})}{(f - z_s)(\bar{f} - z_s)} &= \frac{z_s^2 \prod_{i=3}^8 (\frac{\kappa_2}{z_s} - v_i)}{(\frac{\kappa_2}{z_s})^2 \prod_{i=3}^8 (z_s - v_i)}, \quad \text{for } g = G(z_s). \end{aligned} \quad (2.10)$$

³See [2, 3] for the Lax equation and hypergeometric special solutions of (2.7).

⁴In [4] the Lax equation and hypergeometric special solutions of (2.10) are given, and a relation between the Lax equations for (2.7) and (2.10) is also shown.

This has been regarded in [4, 5, 6] as a variation of the q - $E_7^{(1)}$ equation (2.7). The eight singular points are

$$(f, g) = (v_i, G(v_i))_{i=3}^8, \left(\frac{\kappa_1}{v_i}, \infty \right)_{i=1,2}, \quad (2.11)$$

on the product of a curve and line of bidegree (2, 1) and (0, 1). This configuration for the surface type q - $A_1^{(1)}$ is given in Figure 3. We call it configuration type 2.




Figure 3: Configuration type 2

2.4 Degeneration 3

Doing a replacement of parameters

$$v_1 \rightarrow \varepsilon v_1, \quad v_2 \rightarrow \varepsilon^{-1} v_2, \quad (2.12)$$

and applying a degeneration limit $\varepsilon \rightarrow 0$, we have the following novel evolution equation:

$$\begin{aligned} \frac{\{g - G(\frac{\kappa_1}{z})\}\{\underline{g} - \underline{G}(\frac{\kappa_1}{z})\}}{\{g - G(z)\}\{\underline{g} - \underline{G}(z)\}} &= \frac{U(\frac{\kappa_1}{z})}{U(z)}, \quad \text{for } f = F(z), \\ \frac{\{f - F(\frac{\kappa_2}{z})\}\{\bar{f} - \bar{F}(\frac{\kappa_2}{z})\}}{\{f - F(z)\}\{\bar{f} - \bar{F}(z)\}} &= \frac{U(\frac{\kappa_2}{z})}{U(z)}, \quad \text{for } g = G(z), \end{aligned} \quad (2.13)$$

where $F(z)$ and $G(z)$ are as in (2.4) but only $U(z)$ (2.4) is changed to $U(z) = z^{-3} \prod_{i=3}^8 (z - v_i)$. This equation (2.13) is a kind of the q - $E_7^{(1)}$ equation. The eight singular points are

$$(f, g) = (F(v_i), G(v_i))_{i=3}^8, \left(\frac{1}{\varepsilon}, \frac{1 - v_1 v_2 / \kappa_1}{(1 - v_1 v_2 / \kappa_2) \varepsilon} \right)_2, \quad (2.14)$$




Figure 4: Configuration type 3

on a curve of bidegree (2, 2) with a node. Here, the last point is a double point at (∞, ∞) with the limit $\lim_{\varepsilon \rightarrow 0} \frac{g}{f} = \frac{1 - v_1 v_2 / \kappa_1}{1 - v_1 v_2 / \kappa_2}$. This configuration for the surface type q - $A_1^{(1)}$ is given in Figure 4, and it is not considered so far. We call it configuration type 3.

Remark 2.4.1 *The equations (2.7) and (2.13) are equivalent with each other (see Appendix B).*

The actions on dependent variables x, y are

$$\begin{aligned}
s_0^{[1]} : x &\rightarrow \frac{1}{y}, \quad y \rightarrow \frac{1}{x}, \\
s_4^{[1]} : x &\rightarrow \frac{e_4(h_1 - h_2)x + h_2(e_4e_5 - h_1)xy - h_1(e_4e_5 - h_2)}{e_4(e_4e_5 - h_2)xy - e_4(e_4e_5 - h_1) - e_4e_5(h_1 - h_2)y}, \\
\pi^{[1]} : x &\rightarrow \frac{h_1}{x}, \quad y \rightarrow \frac{1}{h_2y}.
\end{aligned} \tag{3.5}$$

The actions (3.4) and (3.5) is equivalent to the actions given in [2, Section 8.4.2] up to trivial changes of parameters and variables.

3.2 Representation 2 of affine Weyl group actions

We consider $s_i^{[2]}$ and $\pi^{[2]}$ defined by

$$s_i^{[2]} = \varphi_{12}s_i^{[1]}\varphi_{12}^{-1}, \quad \pi^{[2]} = \varphi_{12}\pi^{[1]}\varphi_{12}^{-1}, \tag{3.6}$$

where the action of φ_{12} is given by

$$\begin{aligned}
\varphi_{12} : x &\rightarrow \frac{h_2 \{ (e_3 + e_4)x - e_3e_4 + h_2 - xy \}}{(e_3 + e_4)h_2 + (e_3e_4 - h_2)x - e_3e_4y}, \quad y \rightarrow \frac{1}{x}, \\
e_3 &\rightarrow \frac{h_1}{e_3}, \quad e_4 \rightarrow \frac{h_1}{e_4}, \quad h_1 \rightarrow \frac{h_1h_2}{e_3e_4}, \quad h_2 \rightarrow h_1.
\end{aligned} \tag{3.7}$$

Then one can obtain the actions

$$\begin{aligned}
s_0^{[2]} &= \mu_{34}, \quad s_1^{[2]} = r_{12}, \quad s_2^{[2]} = \nu_{23}, \quad s_3^{[2]} = r_{34}, \\
s_4^{[2]} &= r_{45}, \quad s_5^{[2]} = r_{56}, \quad s_6^{[2]} = r_{67}, \quad s_7^{[2]} = r_{78}, \\
\pi^{[2]} : h_2 &\leftrightarrow \frac{h_1^2h_2}{e_3e_4e_5e_6}, \quad e_1 \leftrightarrow e_8, \quad e_2 \leftrightarrow e_7, \quad e_i \leftrightarrow \frac{e_{\sigma(i)}}{h_1}, \quad \sigma = \begin{pmatrix} 3456 \\ 6543 \end{pmatrix},
\end{aligned} \tag{3.8}$$

and

$$\begin{aligned}
s_0^{[2]} : x &\rightarrow \frac{h_2(xy - a_1x + a_2 - h_2)}{(h_2 - a_2)x + a_2y - a_1h_2}, \\
s_2^{[2]} : y &\rightarrow \frac{e_2e_3xy - e_3(e_2e_3 - h_1)x - e_3h_1y - h_2(e_2e_3 - h_1)}{e_2e_3(x - e_3)}, \\
\pi^{[2]} : x &\rightarrow \frac{h_1}{x}, \quad y \rightarrow \frac{h_1 \{ x^2yh_2 - h_2b_1x^2 - (b_4 - h_2b_2 + h_2^2)x + b_4y - h_2b_3 \}}{b_4(xy - x^2 - h_2)},
\end{aligned} \tag{3.9}$$

where a_i (resp. b_i) is given by $\prod_{i=3}^4(1 + te_i) = \sum_{i=0}^2 a_it^i$ (resp. $\prod_{i=3}^6(1 + te_i) = \sum_{i=0}^4 b_it^i$).

3.3 Representation 3 of affine Weyl group actions

We consider $s_i^{[3]}$ and $\pi^{[3]}$ defined by

$$s_i^{[3]} = \varphi_{23} s_i^{[2]} \varphi_{23}^{-1}, \quad \pi^{[3]} = \varphi_{23} \pi^{[2]} \varphi_{23}^{-1}, \quad (3.10)$$

where the action of φ_{23} is given by

$$\begin{aligned} \varphi_{23} : x &\rightarrow \frac{e_3(h_1 - h_2) + h_2x - h_1y}{e_3x - e_3y - h_1 + h_2}, \quad y \rightarrow x, \\ e_2 &\rightarrow \frac{h_1}{e_2}, \quad e_3 \rightarrow \frac{h_1}{e_3}, \quad h_1 \rightarrow \frac{h_1h_2}{e_2e_3}, \quad h_2 \rightarrow h_1. \end{aligned} \quad (3.11)$$

Then one can get the actions

$$\begin{aligned} s_0^{[3]} &= r_{34}, \quad s_1^{[3]} = \nu_{12}, \quad s_2^{[3]} = c, \quad s_3^{[3]} = \nu_{34}, \\ s_4^{[3]} &= r_{45}, \quad s_5^{[3]} = r_{56}, \quad s_6^{[3]} = r_{67}, \quad s_7^{[3]} = r_{78}, \\ \pi^{[3]} : h_1 &\rightarrow \frac{e_2^2 e_3 e_4 e_5 e_6}{h_1^2 h_2^2}, \quad h_2 \rightarrow \frac{e_2^2 e_3 e_4 e_5 e_7}{h_1^2 h_2^2}, \quad e_1 \leftrightarrow e_8, \quad e_2 \rightarrow \frac{h_1^2 h_2^2}{e_2^2 e_3 e_4 e_5 e_6 e_7}, \\ e_3 &\rightarrow \frac{h_1 h_2}{e_2 e_4 e_5}, \quad e_4 \rightarrow \frac{h_1 h_2}{e_2 e_3 e_5}, \quad e_5 \rightarrow \frac{h_1 h_2}{e_2 e_3 e_4}, \quad e_6 \rightarrow \frac{h_2}{e_2}, \quad e_7 \rightarrow \frac{h_1}{e_2}, \end{aligned} \quad (3.12)$$

and

$$\begin{aligned} s_1^{[3]} : y &\rightarrow \frac{-h_2(e_1 e_2 - h_1)x + h_1(e_1 e_2 - h_2)y}{e_1 e_2 h_-}, \quad s_2^{[3]} : x \leftrightarrow y, \\ s_3^{[3]} : y &\rightarrow \frac{a_2 h_- x y + a_1 h_2 (a_2 - h_1)x - a_1 h_1 (a_2 - h_2)y + (a_2 - h_1)(a_2 - h_2)h_-}{a_2 \{(a_2 - h_2)x - (a_2 - h_1)y - a_1 h_-\}}, \\ \pi^{[3]} : x &\rightarrow \Phi, \quad y \rightarrow \Phi|_{e_6 \rightarrow e_7}. \end{aligned} \quad (3.13)$$

Here

$$\Phi = \frac{h_1 h_2 \{(b_4(x - y) - h_- b_3)(x - y) + (h_2 x - h_1 y + h_- b_1)(h_2 x - h_1 y) + b_2(h_-)^2\}}{e_2 b_4 \{h_2 x^2 + h_1 y^2 - h_+ x y + (h_-)^2\}}, \quad (3.14)$$

and h_{\pm} mean $h_1 \pm h_2$ respectively, and a_i, b_i are given below (3.9).

4 Three q - $E_7^{(1)}$ equations from affine Weyl group actions

In this section, the q - $E_7^{(1)}$ equations (2.7), (2.10), (2.13) will be derived from the three representations 1, 2, 3 of affine Weyl group actions in Section 3.

4.1 The q - $E_7^{(1)}$ equation from representation 1

Let us define the following composition of the Weyl group elements s_0, \dots, s_7 as

$$T_1 = (s_{04567345623451234})^2, \quad (4.1)$$

where $s_{i_0 i_1 \dots i_n} := s_{i_0} s_{i_1} \cdots s_{i_n}$. Then the action $T_1^{[1]}$ of T_1 in representation 1 is expressed as

$$T_1^{[1]} = (s_{04567345623451234}^{[1]})^2, \quad (4.2)$$

and we have

Proposition 4.1.1 *The action $T_1^{[1]}$ (4.2) gives the following equation*

$$\begin{aligned} \frac{T_1^{(1)-1}(x)y - 1}{T_1^{(1)-1}(x)y - \frac{qh_2}{h_1}xy - 1} \frac{xy - \frac{h_1}{h_2}}{xy - 1} &= \frac{\prod_{i=1}^4 (y - \frac{e_i}{h_2})}{\prod_{i=5}^8 (y - \frac{1}{e_i})}, \\ \frac{xT_1^{(1)}(y) - 1}{\frac{qh_1}{h_2}xT_1^{(1)}(y) - 1} \frac{\frac{h_2}{h_1}xy - 1}{xy - 1} &= \frac{\prod_{i=1}^4 (\frac{e_i x}{h_1} - 1)}{\prod_{i=5}^8 (\frac{x}{e_i} - 1)}. \end{aligned} \quad (4.3)$$

Here the equation (4.3) is equivalent to the q - $E_7^{(1)}$ equation (2.7), and it has the singular point configuration (i.e. configuration type 1 for surface type q - $A_1^{(1)}$ in Figure 2)

$$(x, y) = \left(\frac{h_1}{e_i}, \frac{e_i}{h_2} \right)_{i=1, \dots, 4}, \left(e_i, \frac{1}{e_i} \right)_{i=5, \dots, 8}. \quad (4.4)$$

Proof. Firstly, from the actions $s_i^{[1]}$ (3.4) and (3.5), the action $T_1^{[1]}$ (4.2) can be rewritten into the equation (4.3).

Next, under the transformation

$$v_i = \frac{e_i}{\lambda}, \quad \kappa_1 = \frac{h_2}{\lambda^2}, \quad \kappa_2 = \frac{h_1}{\lambda^2}, \quad f = \frac{1}{\lambda y}, \quad g = \frac{\lambda}{x}, \quad (4.5)$$

where $T_1^{[1]-1}(h_1) = \frac{qh_2^2}{h_1}$, $T_1^{[1]}(h_2) = \frac{qh_1^2}{h_2}$, $T_1^{[1]}(\lambda) = \frac{\lambda T_1^{[1]}(h_2)}{h_1}$, the equation (4.3) is equivalent to the q - $E_7^{(1)}$ equation (2.7), and it has the configuration (4.4). \square

4.2 The q - $E_7^{(1)}$ equation from representation 2

Let us define the following composition of the Weyl group elements s_0, \dots, s_7, π as

$$T_2 = \pi s_{56543765402345123406543765430}, \quad (4.6)$$

where $s_{i_0 i_1 \dots i_n} := s_{i_0} s_{i_1} \cdots s_{i_n}$. Then the action $T_2^{[2]}$ of T_2 in representation 2 is expressed as

$$T_2^{[2]} = \pi^{[2]} s_{56543765402345123406543765430}^{[2]}, \quad (4.7)$$

and we get

Proposition 4.2.1 *The action $T_2^{[2]}$ (4.7) gives the following equation*

$$\begin{aligned} \left(x - \frac{h_2}{qh_1} T_2^{[2]-1}(y) + \frac{h_2}{qx} \right) \left(x - y + \frac{h_2}{x} \right) &= \frac{\prod_{i=3}^8 (x - e_i)}{x^2 \prod_{i=1,2} (x - \frac{h_1}{e_i})}, \\ \frac{\{T_2^{[2]}(x) - \frac{T_2^{[2]}(h_1)}{z}\} (x - \frac{h_2}{z})}{\{T_2^{[2]}(x) - T_2^{[2]}(\frac{h_2}{h_1} \frac{z}{q})\} (x - z)} &= \frac{z^2 \prod_{i=3}^8 (\frac{h_2}{z} - e_i)}{(\frac{h_2}{z})^2 \prod_{i=3}^8 (z - e_i)} \quad \text{for } y = z + \frac{h_2}{z}. \end{aligned} \quad (4.8)$$

Here the equation (4.8) is equivalent to the q - $E_7^{(1)}$ equation (2.10), and it has the singular point configuration (i.e. configuration type 2 for surface type q - $A_1^{(1)}$ in Figure 3)

$$(x, y) = \left(e_i, e_i + \frac{h_2}{e_i} \right)_{i=3, \dots, 8}, \left(\frac{h_1}{e_i}, \infty \right)_{i=1, 2}. \quad (4.9)$$

Proof. Firstly, from the actions $s_i^{[2]}$, $\pi^{[2]}$ (3.8), (3.9), the action $T_2^{[2]}$ (4.7) can be rewritten into the equation (4.8).

Next, under the transformation

$$v_i = \frac{e_i}{\lambda}, \quad u = \frac{z}{\lambda}, \quad \kappa_i = \frac{h_i}{\lambda^2}, \quad f = \frac{x}{\lambda}, \quad g = \frac{y}{\lambda}, \quad (4.10)$$

where $T_2^{[2]}(h_1) = \frac{qe_1e_2h_2}{h_1}$, $T_2^{[2]^{-1}}(h_2) = \frac{qh_1^2}{h_2}$, $T_2^{[2]}(\lambda) = \frac{\lambda T_2^{[2]}(h_1)}{h_2}$, the equations (4.8) is equivalent to the equation (2.10), and it has the configuration (4.9). \square

4.3 The q - $E_7^{(1)}$ equation from representation 3

Let us define the following composition of the Weyl group elements s_0, \dots, s_7 as

$$T_3 = (s_{26756456734567104567345604534})^2, \quad (4.11)$$

where $s_{i_0i_1\dots i_n} := s_{i_0}s_{i_1}\cdots s_{i_n}$. Then the action $T_3^{[3]}$ of T_3 in representation 3 is expressed as

$$T_3^{[3]} = (s_{26756456734567104567345604534}^{[3]})^2, \quad (4.12)$$

and we obtain

Proposition 4.3.1 *The action $T_3^{[3]}$ (4.12) gives the following equation*

$$\begin{aligned} \frac{\{T_3^{[3]^{-1}}(y) - (z + \frac{T_3^{[3]^{-1}}(h_2)}{z})\}\{y - (\frac{h_1}{z} + \frac{h_2z}{h_1})\}}{\{T_3^{[3]^{-1}}(y) - (\frac{h_1}{z} + \frac{T_3^{[3]^{-1}}(h_2)z}{h_1})\}\{y - (z + \frac{h_2}{z})\}} &= \frac{U(\frac{h_1}{z})}{U(z)} \quad \text{for } x = z + \frac{h_1}{z}, \\ \frac{\{T_3^{[3]}(x) - (z + \frac{T_3^{[3]}(h_1)}{z})\}\{x - (\frac{h_2}{z} + \frac{h_1z}{h_2})\}}{\{T_3^{[3]}(x) - (\frac{h_2}{z} + \frac{T_3^{[3]}(h_1)z}{h_2})\}\{x - (z + \frac{h_1}{z})\}} &= \frac{U(\frac{h_2}{z})}{U(z)} \quad \text{for } y = z + \frac{h_2}{z}. \end{aligned} \quad (4.13)$$

Here the equation (4.13) is equivalent to the q - $E_7^{(1)}$ equation (2.13), and it has the singular point configuration (i.e. configuration type 3 for surface type q - $A_1^{(1)}$ in Figure 4)

$$(x, y) = \left(e_i + \frac{h_1}{e_i}, e_i + \frac{h_2}{e_i} \right)_{i=3, \dots, 8}, \left(\frac{1}{\varepsilon}, \frac{1 - e_1e_2/h_1}{(1 - e_1e_2/h_2)\varepsilon} \right)_2. \quad (4.14)$$

Proof. Firstly, from the actions $s_i^{[3]}$ (3.12) and (3.13), the action $T_3^{[3]}$ (4.12) can be rewritten into the equation (4.13).

Next, under the transformation

$$v_i = \frac{e_i}{\lambda}, \quad u = \frac{z}{\lambda}, \quad \kappa_i = \frac{h_i}{\lambda^2}, \quad f = \frac{x}{\lambda}, \quad g = \frac{y}{\lambda}, \quad (4.15)$$

where $T_3^{[3]}(h_1) = \frac{qh_2^2}{h_1}$, $T_3^{[3]^{-1}}(h_2) = \frac{qh_1^2}{h_2}$, $T_3^{[3]}(\lambda) = \frac{\lambda T_3^{[3]}(h_1)}{h_2}$, the equation (4.13) is equivalent to the equation (2.13), and it has the configuration (4.14). \square

Proposition 4.3.2 *Two actions T_1 (4.1) and T_3 (4.11) are transformed with each other under the following relation*

$$T_3 = \psi^{-1}T_1\psi, \quad (4.16)$$

where $\psi = s_{403423}$.

Proof. Thanks to the fundamental relations (3.1), one can obtain the relation $T_3^{-1}\psi^{-1}T_1\psi = 1$. \square

Remark 4.3.3 *It seems that the action T_2 (4.6) can not be transformed into the actions T_1 (4.1) and T_3 (4.11) since the action T_2 contains the Dynkin diagram automorphism π .*

5 Conclusion

In this paper, we obtained the following main results.

- We derived the three $q-E_7^{(1)}$ equations (2.7), (2.10), (2.13) by applying the degeneration through confluence of the $q-E_8^{(1)}$ equation in [2, 10]. The $q-E_7^{(1)}$ equations (2.7), (2.10), (2.13) have different realizations (2.8), (2.11), (2.14) (i.e. configuration types 1, 2, 3 for surface type $q-A_1^{(1)}$ in Figures 2, 3, 4) in viewpoint of realizations of eight singular points configuration on the curve of bidgree (2, 2) in $\mathbb{P}^1 \times \mathbb{P}^1$.
- We gave the three representations 1, 2, 3 of affine Weyl group actions with symmetry type $q-E_7^{(1)}$, and relations (3.6), (3.10) among the three representations 1, 2, 3. The $q-E_7^{(1)}$ equations (4.3), (4.8), (4.13) were derived from the actions (4.2), (4.7), (4.12) of the three representations 1, 2, 3 respectively.

A Relation between $q-E_8^{(1)}$ equation (2.3) and ORG from

We give a correspondence between the $q-E_8^{(1)}$ equation given in [2, 10] and ORG form [7]. The q -Painlevé equation with affine Weyl group symmetry of type $E_8^{(1)}$:

$$\begin{aligned} \frac{(\bar{f} - g)(f - g) - (\frac{\kappa_1}{q} - \kappa_2)(\kappa_1 - \kappa_2)\frac{1}{\kappa_2}}{(\frac{\bar{f}q}{\kappa_1} - \frac{g}{\kappa_2})(\frac{f}{\kappa_1} - \frac{g}{\kappa_2}) - (\frac{q}{\kappa_1} - \frac{1}{\kappa_2})(\frac{1}{\kappa_1} - \frac{1}{\kappa_2})\kappa_2} &= \frac{\kappa_1^2 A(\kappa_2, g)}{q B(\kappa_2, g)}, \\ \frac{(f - g)(f - g) - (\kappa_1 - \kappa_2)(\kappa_1 - \frac{\kappa_2}{q})\frac{1}{\kappa_1}}{(\frac{f}{\kappa_1} - \frac{g}{\kappa_2})(\frac{f}{\kappa_1} - \frac{gq}{\kappa_2}) - (\frac{1}{\kappa_1} - \frac{1}{\kappa_2})(\frac{1}{\kappa_1} - \frac{q}{\kappa_2})\kappa_2} &= \frac{\kappa_2^2 A(\kappa_1, f)}{q B(\kappa_1, f)}, \end{aligned} \quad (A.1)$$

were derived by Ohta-Rammani-Grammaticos in [7]. Here $A(h, x)$ and $B(h, x)$ are polynomials in x of degree 4 given by

$$\begin{aligned}
A(h, x) &= \left(\frac{m_0}{h^2} - \frac{m_2}{h} + m_4 - hm_6 + h^2m_8\right) + \left(\frac{m_1}{h^2} - m_5 + 2hm_7\right)x \\
&\quad + \left(-\frac{m_0}{h^3} + m_6 - 3hm_8\right)x^2 - m_7x^3 + m_8x^4, \\
B(h, x) &= \left(\frac{m_0}{h^2} - \frac{m_2}{h^2} + m_4 - hm_6 + h^2m_8\right) + \left(\frac{2m_1}{h^2} - \frac{m_3}{h} + hm_7\right)x \\
&\quad + \left(-\frac{3m_0}{h^3} + \frac{m_2}{h} - hm_8\right)x^2 - \frac{m_1}{h^3}x^3 + \frac{m_0}{h^4}x^4, \\
U(z) &= \frac{1}{z^4} \prod_{i=1}^8 (z - u_i) = \frac{1}{z^4} \sum_{i=0}^8 (-1)^i m_i z^i.
\end{aligned} \tag{A.2}$$

It is in [10] shown that the equation (2.3) is equivalent to the equation (A.1) under the relation

$$A\left(h, z + \frac{h}{z}\right) = \frac{zU(z) - \frac{h}{z}U\left(\frac{h}{z}\right)}{z - \frac{h}{z}}, \quad B\left(h, z + \frac{h}{z}\right) = \frac{zU\left(\frac{h}{z}\right) - \frac{h}{z}U(z)}{z - \frac{h}{z}}. \tag{A.3}$$

B Direct derivation of the relation between q - $E_7^{(1)}$ equations (2.7) and (2.13)

The relation between two kinds of q - $E_7^{(1)}$ equation (2.7) and (2.13) was obtained in Proposition 4.3.2. Here we give its another direct derivation without using the affine Weyl group symmetry. Replacing dependent variables (f, g) and parameters $(\kappa_1, \kappa_2, v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8)$ by $(x, \frac{1}{y})$ and $(\kappa_1, \kappa_2, \frac{\kappa_1\kappa_2}{qv_6v_7v_8}, v_3, v_4, v_5, \frac{\kappa_1\kappa_2}{v_3v_4v_5}, v_6, v_7, v_8)$, then the q - $E_7^{(1)}$ equation (2.7) is rewritten as

$$\frac{x - \frac{\kappa_1}{\kappa_2}y}{x - y} \frac{x - \frac{\kappa_1 q}{\kappa_2}y}{x - y} = \frac{x - \frac{qv_6v_7v_8}{\kappa_2}}{x - \frac{\kappa_1\kappa_2}{v_3v_4v_5}} \frac{\prod_{i=3}^5 (x - \frac{\kappa_1}{v_i})}{\prod_{i=6}^8 (x - v_i)}, \tag{B.1}$$

$$\frac{y - \frac{\kappa_2}{\kappa_1}x}{y - x} \frac{y - \frac{\kappa_2 q}{\kappa_1}\bar{x}}{y - \bar{x}} = \frac{y - \frac{qv_6v_7v_8}{\kappa_1}}{y - \frac{\kappa_1\kappa_2}{v_3v_4v_5}} \frac{\prod_{i=3}^5 (y - \frac{\kappa_2}{v_i})}{\prod_{i=6}^8 (y - v_i)}. \tag{B.2}$$

Here the replacement satisfies the constraint (2.1). Replacing the variable z by u , the q - $E_7^{(1)}$ equation (2.13) is rewritten as

$$\frac{\frac{\kappa_1}{u} + \frac{\kappa_2}{\kappa_1}u - g}{u + \frac{\kappa_2}{u} - g} \frac{\frac{\kappa_1}{u} + \frac{\kappa_2}{q\kappa_1}u - \underline{g}}{u + \frac{\kappa_2}{qu} - \underline{g}} = \frac{u^6}{\kappa_1^3} \prod_{i=3}^8 \frac{\kappa_1 - v_i}{u - v_i} \quad \text{for } f = u + \frac{\kappa_1}{u}, \tag{B.3}$$

$$\frac{\frac{\kappa_2}{u} + \frac{\kappa_1}{\kappa_2}u - f}{u + \frac{\kappa_1}{u} - f} \frac{\frac{\kappa_2}{u} + \frac{\kappa_1}{q\kappa_2}u - \bar{f}}{u + \frac{\kappa_1}{qu} - \bar{f}} = \frac{u^6}{\kappa_2^3} \prod_{i=3}^8 \frac{\kappa_2 - v_i}{u - v_i} \quad \text{for } g = u + \frac{\kappa_2}{u}. \tag{B.4}$$

In order to relate (B.1), (B.2) and (B.3), (B.4), we put

$$x + \frac{\kappa_1}{x} - f = \frac{(x - y) \prod_{i=3}^5 (x - \frac{\kappa_1}{v_i})}{x(x - \frac{\kappa_1\kappa_2}{v_3v_4v_5})(x - \frac{\kappa_1}{\kappa_2}y)}, \tag{B.5}$$

$$y + \frac{\kappa_2}{y} - g = \frac{(y-x) \prod_{i=3}^5 (y - \frac{\kappa_2}{v_i})}{y(y - \frac{\kappa_1 \kappa_2}{v_3 v_4 v_5})(y - \frac{\kappa_2}{\kappa_1} x)}. \quad (\text{B.6})$$

Then we get the following two Lemmas,

Lemma B.0.1 *The relations (B.5) and (B.6) give the birational transformation: $(f, g) \leftrightarrow (x, y)$.*

Proof. Obviously, the relations (B.5) and (B.6) are the rational transform $(x, y) \rightarrow (f, g)$. Conversely, using $\prod_{i=3}^5 (u - v_i) = u^3 - \mu_1 u^2 + \mu_2 u - \mu_3$, the relations (B.5) and (B.6) are written as

$$x = \kappa_1 \frac{f^2 \kappa_2 - f g \kappa_1 + f(\mu_3 - \kappa_2 \mu_1) + g(\kappa_1 \mu_1 - \mu_3) + (\kappa_1 - \kappa_2)(\kappa_1 - \mu_2)}{f^2 \mu_3 - f g \mu_3 + f \kappa_1 (\kappa_2 - \mu_2) + g \kappa_1 (\mu_2 - \kappa_1) + (\kappa_1 - \kappa_2)(\kappa_1 \mu_1 - \mu_3)}, \quad (\text{B.7})$$

$$y = \kappa_2 \frac{g^2 \kappa_1 - f g \kappa_2 + g(\mu_3 - \kappa_1 \mu_1) + f(\kappa_2 \mu_1 - \mu_3) + (\kappa_2 - \kappa_1)(\kappa_2 - \mu_2)}{g^2 \mu_3 - f g \mu_3 + g \kappa_2 (\kappa_1 - \mu_2) + f \kappa_2 (\mu_2 - \kappa_2) + (\kappa_2 - \kappa_1)(\kappa_2 \mu_1 - \mu_3)}, \quad (\text{B.8})$$

and they are the rational transform $(f, g) \rightarrow (x, y)$. \square

Lemma B.0.2 *The birational transformations (B.5), (B.6) (or equivalently (B.7), (B.8)) can be written as*

$$\frac{x - \frac{\kappa_1}{u}}{x - u} \frac{u + \frac{\kappa_2}{u} - g}{\frac{\kappa_1}{u} + \frac{\kappa_2}{\kappa_1} u - g} = \frac{\kappa_1^2}{u^4} \prod_{i=3}^5 \frac{u - v_i}{\frac{\kappa_1}{u} - v_i} \quad \text{for } f = u + \frac{\kappa_1}{u}, \quad (\text{B.9})$$

$$\frac{y - \frac{\kappa_2}{u}}{y - u} \frac{u + \frac{\kappa_1}{u} - f}{\frac{\kappa_2}{u} + \frac{\kappa_1}{\kappa_2} u - f} = \frac{\kappa_2^2}{u^4} \prod_{i=3}^5 \frac{u - v_i}{\frac{\kappa_2}{u} - v_i} \quad \text{for } g = u + \frac{\kappa_2}{u}. \quad (\text{B.10})$$

Proof. Under a condition $f = u + \frac{\kappa_1}{u}$, eliminating y from the relations (B.5) and (B.6) (i.e. (B.5) and (B.8)), we have the relation (B.9). Under a condition $g = u + \frac{\kappa_2}{u}$, eliminating x from the relations (B.5) and (B.6) (i.e. (B.6) and (B.7)), we have the relation (B.10). \square

Thanks to Lemmas B.0.1 and B.0.2, we have

Proposition B.0.3 *The equations (B.1), (B.2) and the equations (B.3), (B.4) are equivalent with each other under the relations⁵ (B.5) and (B.6)*

Proof. Combining (B.1) and (B.5), we have

$$x + \frac{\kappa_1}{x} - f = \frac{(x - \frac{q\kappa_1}{\kappa_2} y) \prod_{i=6}^8 (x - v_i)}{x(x - y)(x - \frac{qv_6 v_7 v_8}{\kappa_2})}. \quad (\text{B.11})$$

On the other hand, using (B.2) and (B.6), one has

$$y + \frac{\kappa_2}{y} - g = \frac{(y - \frac{q\kappa_2}{\kappa_1} \bar{x}) \prod_{i=6}^8 (y - v_i)}{y(y - \bar{x})(y - \frac{qv_6 v_7 v_8}{\kappa_1})}. \quad (\text{B.12})$$

⁵The relations (B.5) and (B.6) corresponds to the relation (4.16) in viewpoint of affine Weyl group symmetry.

Substituting (B.11) into y in (B.12), we get

$$\frac{u + \frac{\kappa_2}{qu} - \underline{g}}{\frac{\kappa_1}{u} + \frac{\kappa_2}{q\kappa_1}u - \underline{g}} = \frac{\kappa_1}{u^2} \frac{x - \frac{\kappa_1}{u}}{x - u} \prod_{i=6}^8 \frac{u - v_i}{\frac{\kappa_1}{u} - v_i} \quad \text{for } f = u + \frac{\kappa_1}{u}. \quad (\text{B.13})$$

Therefore the equation (B.3) is derived from the relations (B.9) and (B.13).

Similarly, substituting (B.12) into \bar{x} in (B.11), we obtain

$$\frac{u + \frac{\kappa_1}{qu} - \bar{f}}{\frac{\kappa_2}{u} + \frac{\kappa_1}{q\kappa_2}u - \bar{f}} = \frac{\kappa_2}{u^2} \frac{y - \frac{\kappa_2}{u}}{y - u} \prod_{i=6}^8 \frac{u - v_i}{\frac{\kappa_2}{u} - v_i} \quad \text{for } g = u + \frac{\kappa_2}{u}. \quad (\text{B.14})$$

Therefore the equation (B.4) is derived from the relations (B.10) and (B.14). \square

C Three translations on the root variables

For the Dynkin diagram

$$\begin{array}{ccccccc} & & & \circ \alpha_0 & & & \\ & & & | & & & \\ \circ \alpha_1 & - & \circ \alpha_2 & - & \circ \alpha_3 & - & \circ \alpha_4 & - & \circ \alpha_5 & - & \circ \alpha_6 & - & \circ \alpha_7 \end{array} \quad \delta_0 \iff \delta_1, \quad (\text{C.1})$$

we consider a pair of root bases $\{\alpha_i\}$, $\{\delta_i\}$ as symmetry/surface type $q\text{-}E_7^{(1)}/q\text{-}A_1^{(1)}$. Then we give three representations of root bases $\{\alpha_i^{[m]}\}$, $\{\delta_i^{[m]}\}$ ($m = 1, 2, 3$) as symmetry/surface type $E_7^{(1)}/A_1^{(1)}$ associated with the point configurations (4.4), (4.9), (4.14). We also show how the translations $T_1^{[1]}$ (4.2), $T_2^{[2]}$ (4.7), $T_3^{[3]}$ (4.12) act on the root variables $\alpha_i^{[1]}$, $\alpha_i^{[2]}$, $\alpha_i^{[3]}$ explicitly.

C.1 Case of representation 1

The root bases for representation 1 are

$$\begin{aligned} E_7^{(1)} : \quad & \alpha_0^{[1]} = H_1 - H_2, \quad \alpha_1^{[1]} = E_2 - E_1, \quad \alpha_2^{[1]} = E_3 - E_2, \quad \alpha_3^{[1]} = E_4 - E_3, \\ & \alpha_4^{[1]} = H_2 - E_4 - E_5, \quad \alpha_5^{[1]} = E_5 - E_6, \quad \alpha_6^{[1]} = E_6 - E_7, \quad \alpha_7^{[1]} = E_7 - E_8, \\ A_1^{(1)} : \quad & \delta_0^{[1]} = H_1 + H_2 - E_1 - E_2 - E_3 - E_4, \quad \delta_1^{[1]} = H_1 + H_2 - E_5 - E_6 - E_7 - E_8. \end{aligned} \quad (\text{C.2})$$

This realization for symmetry/surface type $E_7^{(1)}/A_1^{(1)}$ are associated with the actions $s_i^{[1]}$, $\pi^{[1]}$ /configuration (4.4) (i.e. configuration type 1 in Figure 2).

Proposition C.1.1 *The action $T_1^{[1]}$ (4.2) on the root variables $\alpha_i^{[1]}$ in (C.2) are given by the translation*

$$T_1^{[1]}(\alpha_0^{[1]}) = q^2 \alpha_0^{[1]}, \quad T_1^{[1]}(\alpha_4^{[1]}) = \frac{\alpha_4^{[1]}}{q}, \quad T_1^{[1]}(\alpha_i^{[1]}) = \alpha_i^{[1]} \quad (i = 1, 2, 3, 5, 6, 7), \quad (\text{C.3})$$

Proof. Reading the relation (C.2) multiplicatively and applying (3.4), we obtain the desired result (C.3). \square

C.2 Case of representation 2

The root bases for representation 2 are

$$\begin{aligned}
E_7^{(1)} : \quad & \alpha_0^{[2]} = H_2 - E_3 - E_4, \quad \alpha_1^{[2]} = E_2 - E_1, \quad \alpha_2^{[2]} = H_1 - E_2 - E_3, \quad \alpha_3 = E_3 - E_4, \\
& \alpha_4^{[2]} = E_4 - E_5, \quad \alpha_5^{[2]} = E_5 - E_6, \quad \alpha_6^{[2]} = E_6 - E_7, \quad \alpha_7^{[2]} = E_7 - E_8, \\
A_1^{(1)} : \quad & \delta_0^{[2]} = 2H_1 + H_2 - E_3 - E_4 - E_5 - E_6 - E_7 - E_8, \quad \delta_1^{[2]} = H_2 - E_1 - E_2.
\end{aligned} \tag{C.4}$$

This realization for symmetry/surface type $E_7^{(1)}/A_1^{(1)}$ are associated with the actions $s_i^{[2]}, \pi^{[2]}$ /configuration (4.9) (i.e. configuration type 2 in Figure 3).

Proposition C.2.1 *The action $T_2^{[2]}$ (4.7) on the root variables $\alpha_i^{[2]}$ in (C.4) are given by the translation*

$$T_2^{[2]}(\alpha_0^{[2]}) = \frac{\alpha_0^{[2]}}{q}, \quad T_2^{[2]}(\alpha_2^{[2]}) = q\alpha_2^{[2]}, \quad T_2^{[2]}(\alpha_i^{[2]}) = \alpha_i^{[2]} \quad (i = 1, 3, 4, 5, 6, 7). \tag{C.5}$$

Proof. Reading the relation (C.4) multiplicatively and applying (3.8), we obtain the desired result (C.5). \square

C.3 Case of representation 3

The root bases for representation 3 are

$$\begin{aligned}
E_7^{(1)} : \quad & \alpha_0^{[3]} = E_3 - E_4, \quad \alpha_1^{[3]} = H_1 - E_1 - E_2, \quad \alpha_2^{[3]} = H_2 - H_1, \quad \alpha_3^{[3]} = H_1 - E_3 - E_4, \\
& \alpha_4^{[3]} = E_4 - E_5, \quad \alpha_5^{[3]} = E_5 - E_6, \quad \alpha_6^{[3]} = E_6 - E_7, \quad \alpha_7^{[3]} = E_7 - E_8, \\
A_1^{(1)} : \quad & \delta_0^{[3]} = 2H_1 + 2H_2 - 2E_2 - E_3 - E_4 - E_5 - E_6 - E_7 - E_8, \quad \delta_1^{[3]} = E_2 - E_1.
\end{aligned} \tag{C.6}$$

This realization for symmetry/surface type $E_7^{(1)}/A_1^{(1)}$ are associated with the actions $s_i^{[3]}, \pi^{[3]}$ /configuration (4.14) (i.e. configuration type 3 in Figure 4).

Proposition C.3.1 *The action $T_3^{[3]}$ (4.12) on the root variables $\alpha_i^{[3]}$ in (C.6) are given by the translation*

$$T_3^{[3]}(\alpha_i^{[3]}) = \frac{\alpha_i^{[3]}}{q}, \quad (i=1,3) \quad T_3^{[3]}(\alpha_2^{[3]}) = q^2\alpha_2^{[3]}, \quad T_3^{[3]}(\alpha_i^{[3]}) = \alpha_i^{[3]} \quad (i = 0, 2, 4, 5, 6, 7). \tag{C.7}$$

Proof. Reading the relation (C.6) multiplicatively and applying (3.12), we obtain the desired result (C.7). \square

Acknowledgment. The authors shall be thankful to Professor Tetsu Masuda, Masatoshi Noumi, Hidetaka Sakai, for valuable discussions. This work is partially supported by JSPS KAKENHI (22H01116), JSPS KAKENHI (19K14579) and JSPS KAKENHI (23K03173).

References

- [1] Grammaticos B., Ramani A.: On a novel q -discrete analogue of the Painlevé VI equation. *Phys. Lett. A.* **257** (1999), 288–292.
- [2] Kajiwara K., Noumi M., Yamada Y.: *Geometric aspects of Painlevé equations*. *J. Phys. A: Math. Theor.* **50** (2017), 073001–073164 (Topical Review).
- [3] Nagao H.: *The Padé interpolation method applied to q -Painlevé equations*. *Lett. Math. Phys.* **105** (2015), Issue 4, 503–521.
- [4] Nagao H.: *A variation of the q -Painlevé system with affine Weyl group symmetry of type $E_7^{(1)}$* . *SIGMA.* **13** (2017), 92–109.
- [5] Nagao H., Yamada Y.: *Study of q -Garnier system by Padé method*. *Funkcialaj Ekvacioj.* **61** (2018), 109–133.
- [6] Nagao H., Yamada Y.: *Variations of the q -Garnier system*. *J. Phys. A: Math. Theor.* **51** (2018), 135204–135222.
- [7] Ohta Y., Ramani A., Grammaticos B.: *An affine Weyl group approach to the eight-parameter discrete Painlevé equation*. *J. Phys.* **A34** (2001), 10523–10532.
- [8] Sakai H.: *Rational surfaces with affine root systems and geometry of the Painlevé equations*. *Commun. Math. Phys.* **220** (2001), 165–221.
- [9] Yamada Y.: *Lax formalism for q -Painlevé equations with affine Weyl group symmetry of type $E_n^{(1)}$* . *IMRN.* **17** (2011), 3823–3838.
- [10] Yamada Y.: *A simple expression for discrete Painlevé equations*. *RIMS Kokyuroku Bessatsu.* **B47** (2014), 087–095.