

# Fate of dissipative hierarchy of timescales in the presence of unitary dynamics

Nick D. Hartmann, Jimin L. Li, and David J. Luitz  
*Institute of Physics, University of Bonn, Nußallee 12, 53115 Bonn*

The generic behavior of purely dissipative open quantum many-body systems with local dissipation processes can be investigated using random matrix theory, revealing a hierarchy of decay timescales of observables organized by their complexity as shown in [Wang *et al.*, *Phys. Rev. Lett.* **124**, 100604 (2020)]. This hierarchy is reflected in distinct eigenvalue clusters of the Lindbladian. Here, we analyze how this spectrum evolves when unitary dynamics is present, both for the case of strongly and weakly dissipative dynamics. In the strongly dissipative case, the unitary dynamics can be treated perturbatively and it turns out that the locality of the Hamiltonian determines how susceptible the spectrum is to such a perturbation. For the physically most relevant case of (dissipative) two-body interactions, we find that the correction in the first order of the perturbation vanishes, leading to the relative robustness of the spectral features. For weak dissipation, the spectrum flows into clusters with well-separated eigenmodes, which we identify to be the local symmetries of the Hamiltonian.

## I. INTRODUCTION

The eigenvalues of unitary quantum many-body systems typically exhibit level repulsion. It was realized early on that while the precise location of all eigenvalues can only be analytically described in rare, integrable systems, their statistical properties in the vast majority of systems are well captured by random matrix theory [1–8]. This random matrix description provides a powerful framework, correctly predicting the distribution of gaps between neighboring eigenvalues, and further providing insight on the corresponding structure of eigenvectors and their link to thermalization [9–14].

It is exceedingly difficult to fully isolate a real quantum system from its environment, which is a source of dissipation and incoherent dynamics. Systems in contact with their environment are therefore described in the framework of open quantum systems. In the simplest class of Markovian open quantum systems, the dynamics of the density matrix is determined by the Lindblad master equation [15] and the generator of the dynamics is a Lindbladian superoperator, rather than just the Hamiltonian. The spectrum of the Lindbladian is generally complex with nonpositive real parts, such that at long times a state from the manifold corresponding to the zero eigenvalue is reached.

Analogously to the approach for unitary quantum systems, random matrix descriptions were developed to identify universal features of Markovian open systems. These theories are based on an ensemble of random matrices which are at the same time valid Lindbladians and hence generate completely positive trace preserving maps. It was found that non-local, purely dissipative Liouvillians exhibit a lemon-shaped support on the complex plane [16–20], which is similar to the spindle-shaped spectral support found in classical random Master equations [21].

With a non-vanishing Hamiltonian part, the spectral support is deformed to an elliptic shape [16], and various spectral transitions in the weak and strong dissipation

limits [22, 23] and transitions of the steady-state properties [17, 24] were identified.

To bridge the gap between these purely random matrix theories and physical systems, the locality of (dissipative) interactions in a complexity theoretic sense was included. In particular, the most physically relevant case of two body interactions leads to a random matrix ensemble with an additional striking structure: Instead of one lemon-shaped eigenvalue cluster, well-separated clusters of eigenvalues were identified, which correspond to observables of different locality and relax to the stationary on a hierarchy of timescales [25]. This phenomenological prediction was recently verified in experiments on a noisy qubit platform [26] and extended to include the case of strongly varying dissipation strengths, which introduce additional, metastable, structure [27].

Both, the presence of a Hamiltonian in an otherwise unstructured Lindbladian and locality in a purely dissipative system strongly deform the spectrum. In this paper, we investigate the interplay of these competing tendencies and consider a random open quantum system with nontrivial, local, unitary evolution and local dissipation channels.

## II. MODEL

For Markovian open quantum systems, the Liouvillian superoperator  $\mathcal{L}$  generates the dynamics of density matrices and the Lindblad master equation reads

$$\frac{d\rho}{dt} = \mathcal{L}(\rho) = \alpha\mathcal{L}_U(\rho) + \mathcal{L}_D(\rho) \quad (1)$$

$$= -i\alpha[H, \rho] + \sum_{n,m=1}^{N_L} K_{nm} \left[ L_n \rho L_m^\dagger - \frac{1}{2} \{L_m^\dagger L_n, \rho\} \right], \quad (2)$$

where  $H$  is the Hamiltonian yielding unitary evolution by  $\mathcal{L}_U$  and  $\{L_n\}_{n=1,\dots,N_L}$  is the set of  $N_L$  jump operators that represent the dissipation channels coupled by

the positive semidefinite Kossakowski matrix  $K$  [28], and  $1/\alpha \geq 0$  parametrizes the relative strength of dissipation.

For a generic modeling of dissipative quantum many-body spin systems, we consider  $\ell$  spins  $S = 1/2$  with a Hilbert space dimension  $N = 2^\ell$  and a Liouville operator space dimension  $N^2 = 4^\ell$ . The locality of dissipative interactions is introduced by choosing the jump operators to be normalized Pauli strings

$$L_{\vec{\mu}} = \frac{1}{\sqrt{N}} \sigma_{\mu_1} \otimes \sigma_{\mu_2} \otimes \cdots \otimes \sigma_{\mu_\ell}, \quad \mu_i \in \{0, x, y, z\}, \quad (3)$$

of maximal weight  $k_{\vec{\mu}} = \sum_{i=1}^{\ell} (1 - \delta_{\mu_i, 0}) \leq k_{\max}$ , i.e. there are at most  $k_{\max}$  non-identity operators in the string. We note that Pauli strings with weight  $k$  can be interpreted as  $k$ -body interactions. These operators are traceless  $\text{Tr}(L_n) = 0$  and satisfy orthonormality  $\text{Tr}(L_n^\dagger L_m) = \delta_{n,m}$  [29].

We consider one and two-body dissipation channels ( $k_{\max} = 2$ ), so the total number of traceless Lindblad operators is  $N_L = 3\ell + 9\ell(\ell - 1)/2$  and we choose the Kossakowski matrix  $K$  to be a random matrix. To ensure that  $K$  is positive semidefinite, we first sample a diagonal matrix  $D$  with i.i.d. non-negative entries from a uniform distribution. The matrix  $D$  is normalized by  $\text{Tr}D = N$ , and then rotated into a random unitary basis by a unitary transformation  $U$  sampled from the Haar measure to yield  $K = U^\dagger D U$ . The mean of the off-diagonal elements vanishes,  $\text{mean}(K_{ij}) = 0$  with a standard deviation  $\text{std}(K_{ij}) = d/\sqrt{6N_L}$ , where  $d$  is the mean of the diagonal elements  $\text{mean}(K_{ii}) = N/N_L = d$ . Consequently,  $K$  is diagonally dominant and can therefore be separated into a large diagonal term and a small off-diagonal term, which is the basis for the perturbation theory proposed in Ref. [25].

For the unitary dynamics we focus on 2-body interactions, in the form of all possible Pauli strings of weight  $k = 2$  [30], and start from a generic Hamiltonian with all-to-all interactions [31]

$$H = \sum_{s,s'=\{x,y,z\}} \sum_{i<j}^{\ell} J_{i,j}^{s,s'} \sigma_i^s \sigma_j^{s'}, \quad (4)$$

where the random real coupling constants  $J_{i,j}^{s,s'}$  follow independent Gaussian distributions centered at 0. The notation  $\sigma_i^s = I \otimes \sigma^s \otimes I$  is the weight one Pauli string acting on site  $i$ . We normalize the Hamiltonian by  $\text{Tr}(H^2) = N$  and with the following standard deviation

$$\sigma = \text{std}(J_{i,j}^{s,s'}) = \sqrt{2/(9\ell(\ell - 1))}. \quad (5)$$

Before we provide our analytic results in full detail, it is worth outlining the strategy of our analysis. Our starting point is the purely dissipative case  $\alpha = 0$ , for which the spectrum of the Liouvillian  $\mathcal{L}$  splits into well-separated eigenvalue clusters. We consider a three term decomposition of  $\mathcal{L}$  into  $\mathcal{L}_D^0 + \mathcal{L}_D^1 + \alpha\mathcal{L}_U$  and the key insight is that  $\mathcal{L}_D^0$  is diagonal in the basis of Pauli strings.

For small  $\alpha$ , we can treat the unitary part  $\alpha\mathcal{L}_U$  as a perturbation and analyze the deformation of the spectrum. In the last step, we include the off-diagonal part of the dissipator  $\mathcal{L}_D^1$ , which leads to further corrections in the spectrum but does not change the physical structure of the eigenmodes significantly.

### III. SPECTRUM

Figure 1 shows the eigenvalues of the Liouvillian  $\alpha\mathcal{L}_U + \mathcal{L}_D$  from one random realization of  $J$  and  $K$  for different relative strengths  $\alpha$  of the unitary dynamics. The spectrum is complex with a non-positive real part such that the continuous dynamical map is completely positive and trace preserving (CPTP) [32]. For  $\alpha = 0$ , the dynamics is purely dissipative and eigenvalues are organized in well-separated clusters [25]. This structure can be unraveled by the following arguments: the diagonally dominant nature of  $K$  allows us to split the dissipator into two terms

$$\mathcal{L}_D = \mathcal{L}_D^0 + \mathcal{L}_D^1, \quad (6)$$

where  $\mathcal{L}_D^0$  contains the diagonal part with Kossakowski matrix  $K_0 = d\mathbf{1}$  and  $\mathcal{L}_D^1$  the off-diagonal perturbation  $K' = K - d\mathbf{1}$  correspondingly. The first term is diagonal in the Pauli string basis, and the eigenvalues are determined only by the weight  $k$  of the eigenvectors [25], which leads to highly degenerate eigenvalues  $\lambda_0(k)$ . Degenerate perturbation theory with  $\mathcal{L}_D^1$  in each degenerate subspace lifts this degeneracy and eigenvalue clusters emerge, centered around  $\lambda_0(k)$ .

Once the unitary component is added ( $\alpha > 0$ ), we observe an attraction of the eigenvalue clusters along the real axes and the expansion of cluster height in the imaginary direction [33, 34]. We observe a linear scaling of the imaginary cluster extent with  $\alpha$  upon adding a local Hamiltonian, which also holds for a non-local Hamiltonian [16]. Gradually increasing the unitary strength  $\alpha$  leads to clusters merging when their boundaries touch. Furthermore, the complex spacing ratio of the merged cluster equals the Ginibre ensemble, which confirms its level repulsion nature [35], see Fig. 2.

The spectral influence of the unitary term also appears in the support of the eigenmodes. For  $\alpha = 0$ , the support of eigenmodes is well approximated by the degenerate subspaces of  $\mathcal{L}_D^0$ , which are spanned by all Pauli strings with fixed weight  $k$ . The degeneracy of these eigenvalues is lifted by  $\mathcal{L}_D^1$ , leading to eigenvalue clusters with eigenmodes composed of fixed weight Pauli strings.

For  $\alpha > 0$ , these clusters tend to merge and the eigenmodes become smeared out over Pauli strings of different weights. We demonstrate this change of the support of eigenmodes by considering the overlap between every left eigenmode  $\rho_i$  of the Liouvillian with an operator  $\hat{O}^{(2)}$  given by a random superposition of Pauli strings




FIG. 1. Eigenvalue spectrum of one realization of the random Liouvillian for a system size  $\ell = 6$ , with weight-2  $\mathcal{L}_U$  and up-to-weight-2 ( $k \leq 2$ )  $\mathcal{L}_D$  at eight different relative strengths  $\alpha$  of the unitary component. Note that in the bottom panels we rescale the imaginary part of the eigenvalues by a factor of  $1/\alpha$  to put these panels on the same scale. The color indicates the overlap  $\text{Tr}(\rho_i \hat{O}^{(2)})$  of the eigenmodes  $\rho_i$  with a random weight-2 operator  $\hat{O}^{(2)}$ , indicating the two-body operator content of the eigenmodes. The gray lines below the spectrum give the analytic predictions for the cluster centers using  $\mathcal{L}_D^0$  [25]. (a) Purely dissipative case ( $\alpha = 0$ ), the spectrum consists of separated eigenvalue clusters that are supported by weight  $k$  Pauli strings. (b)–(g) Increasing the contribution of unitary dynamics, the clusters merge into each other and show an expansion in the imaginary direction. The red lines above the spectrum indicate the prediction of the shifted cluster centers  $\lambda_0(k) + \langle \overline{\lambda}_{2i}(k) \rangle \alpha^2$  by perturbation theory, and the red dots indicate the real part of the mean of the still separated clusters. (h) For strong unitary dynamics, the eigenvalues are concentrated around  $-1$  with a separated eigenvalue near the eigenvalue  $\lambda_0(k=2)$  of  $\mathcal{L}_D^0$  (gray line marked with 2).

of fixed weight  $k = 2$ . This is shown in the colorbar of Fig. 1. For  $\alpha = 0$ , the eigenvalue cluster labeled 2 has a strong weight-2 operator support. Upon increasing  $\alpha$ , the weight-2 nature gets diluted as Pauli strings from different degenerate subspaces start contributing.

In the large  $\alpha$  limit, all clusters merge with the real part concentrating around eigenvalue  $-1$  except a persistent decay mode constructed by Pauli strings of weight 2. This outlier, labeled by the gray bar in Fig. 1(h), is a unique feature of unitary locality and is absent in random ensembles [17].

### A. Strong dissipation: Perturbation Theory

In this section, we provide a quantitative understanding of the observed cluster attraction by treating the uni-

tary term  $\alpha\mathcal{L}_U$  as a perturbation for small  $\alpha$  to an otherwise purely dissipative Liouvillian  $\mathcal{L}_D$ . For  $\alpha = 0$ , we start from  $\mathcal{L}_D^0$  in Eq. (6), which is diagonal in the Pauli string basis due to the CPTP condition of the Liouvillian superoperator. Furthermore, the diagonal elements of  $\mathcal{L}_D^0$  in the Pauli string basis are identical for strings of the same weight, see Fig. 3(a). In Ref. [25], it was worked out that the diagonal dominant structure of  $\mathcal{L}_D$  in the Pauli string basis leads to the separation of eigenvalue clusters for the low weights Pauli strings, which is robust in the thermodynamical limit. This suggests that discarding  $\mathcal{L}_D^1$  well approximates  $\mathcal{L} = \mathcal{L}_D + \alpha\mathcal{L}_U$  qualitatively. In the rest of this section, we derive analytic results of  $\mathcal{L}_D^0 + \alpha\mathcal{L}_U$  to understand the spectral transition.

We begin by treating the unitary generator  $\mathcal{L}_U$  for a 2-local Hamiltonian as a perturbation and representing




FIG. 2. Probability density of the complex spacing ratio  $r = |\lambda_0 - \lambda_{nn}|/|\lambda_0 - \lambda_{nnn}|$  for spectra as shown in Fig. 1 for  $\ell = 6$  from 100 realizations. Due to the symmetry with respect to the real axis, we only consider eigenvalues  $\lambda$  with  $\text{Re}(\lambda) > 0$ . The complex spacing ratio for all four unitary strengths shows clear level repulsion and agrees with the Ginibre ensemble. The purely unitary case is also considered for comparison ( $p(r) \approx 1$ ).

it as a matrix  $\mathcal{L}_U$  in the orthonormal Pauli string basis, which is the eigenbasis of  $\mathcal{L}_D^0$ . The matrix elements are given by

$$\mathcal{L}_{x,y}^U = \text{Tr}(S_x^k \mathcal{L}_U [S_y^{k'}]) = i \text{Tr}([S_x^k, S_y^{k'}] H). \quad (7)$$

Eq. (7) encodes how the unitary term non-trivially couples different weight sectors of the Liouvillian  $\mathcal{L}_D$ , and hence potentially mixes previously well-separated eigenvalue clusters. Therefore, it is crucial to understand the conditions for nonzero matrix elements between sectors of different weights  $k$ .

First, we note that the commutator  $[S_x^k, S_y^{k'}]$  is also a Pauli string and evidently must yield a term of the Hamiltonian (any weight-2 Pauli string in this work) to give a non-zero trace and hence a non-vanishing matrix element  $\mathcal{L}_{x,y}^U$ . We find that  $\text{Tr}([S_x^k, S_y^{k'}] H) \neq 0$  is only possible if  $|k - k'| = 1$ , i.e., the weight of two strings differs by one. To see this, we consider two Pauli strings with all possible ranges of  $|k - k'|$ .

For any two Pauli strings  $S_x^k$  and  $S_y^{k'}$  with  $|k - k'| \geq 2$ , there are at least two sites of  $S_x^k$  containing non-identity Pauli strings where  $S_y^{k'}$  is identity. The commutator of the remaining sites cannot yield an identity Pauli string since this would contradict  $\text{Tr}([A, B]) = \text{Tr}(AB) - \text{Tr}(BA) = 0$ . Consequently, the commutator  $[S_x^k, S_y^{k'}]$  either vanishes or has a weight higher than the 2-local Hamiltonian, hence yielding a zero element  $\mathcal{L}_{x,y}^U$ .

For  $k = k'$ , we tripartite two strings as the following: 1.  $n_1$  different non-identity Pauli matrices on the same sites, 2. a staggered arrangement of non-identity Pauli matrices and identities on  $n_2$  sites, and 3. the same Pauli matrices on the remaining sites with  $n_3$  non-identity Pauli

matrices:

$$\begin{aligned} S_x^k &= A \otimes \sigma_i \otimes \cdots \otimes \sigma_j \otimes I \otimes \cdots \otimes I \otimes \sigma_\alpha \otimes \cdots \otimes \sigma_\gamma \\ S_y^{k'} &= A \otimes \underbrace{I \otimes \cdots \otimes I}_{n_2 \text{ sites}} \otimes \sigma_k \otimes \cdots \otimes \sigma_l \otimes \underbrace{\sigma_\beta \otimes \cdots \otimes \sigma_\delta}_{n_1 \text{ sites}} \\ A &= I \otimes \cdots \otimes I \otimes \underbrace{\sigma_n \otimes \cdots \otimes \sigma_m}_{n_3 \text{ sites}}. \end{aligned} \quad (8)$$

Counting all non-identity Pauli matrices  $2n_1 + n_2 + 2n_3 = 2k$  in both strings shows that  $n_2$  is even. Evaluating the commutator of string pairs with non-zero  $n_2$  gives either 0 or a weight higher than two, requiring a higher locality Hamiltonian to produce non-zero matrix elements. For  $n_2 = 0$ , the commutator yields either 0 or a string with odd weight, which is orthogonal to the Hamiltonian that contains only terms of weight two.

Therefore, a 2-local Hamiltonian couples a weight  $k$  sector to the  $k \pm 1$  sectors of the Pauli string basis, i.e.  $\mathcal{L}_{x,y}^U$  is block-subdiagonal as shown in Fig. 3(c). The non-zero matrix elements are given by  $\mathcal{L}_{x,y}^U = 2J_{i,j}^{s,s'}$ , where  $(s, s', i < j)$  encode the Pauli string of weight 2 obtained by  $[S_x^k, S_y^{k'}] = 2\sigma_i^s \sigma_j^{s'}$ . Using combinatorics, we observe that the numbers of non-zero matrix elements are equal for different rows within the same weight block; for a fixed  $S_x^k$  and weight  $k' = k \pm 1$ , it is

$$\begin{aligned} h(k, k' = k + 1) &= 6k(\ell - k), \\ h(k, k' = k - 1) &= 2k(k - 1). \end{aligned} \quad (9)$$

Focusing on  $\mathcal{L}_D^0 + \mathcal{L}_U$ , we obtain analytic predictions for the centers of eigenvalue clusters, and the role of  $\mathcal{L}_U$  is to couple Pauli strings with different weights  $k$  and  $k \pm 1$ . In the last step, we include  $\mathcal{L}_D^1$ , introducing further renormalization of the resulting eigenvalues. This results in the scattering of the full spectrum around the analytic predictions for the centers of eigenvalue clusters.

To obtain a quantitative understanding, we expand the eigenvalues up to the second order in the strength  $\alpha$  of the perturbation  $\lambda_i = \lambda_{0i} + \lambda_{1i}\alpha + \lambda_{2i}\alpha^2 + \mathcal{O}(\alpha^3)$ . For a degenerate  $\lambda_{0i}$ , the first-order correction is given by diagonalizing  $\mathcal{L}^U$  in the respective degenerate sub-block. Using the fact that  $\mathcal{L}^{U\dagger} = -\mathcal{L}^U$ , the first-order perturbation is purely imaginary. We note that while the eigenvalues of  $\mathcal{L}_D^0$  are determined by the weight of the Pauli string eigenmodes, two subspaces of different weights can have the same eigenvalue.

However, as a consequence of vanishing diagonal blocks in the block-structure of  $\mathcal{L}^U$ , we do not expect a linear scaling of the cluster height for clusters supported by only a single weight, as observed in Fig. 4. This also holds for clusters supported by a degenerate subspace that contains weights differing by more than one. The exceptions are the weights  $k_\pm = 3\ell/4 \pm 1/2$  for  $\ell \equiv 2 \pmod{4}$  sectors, which are consecutive and share the same eigenvalue. In this case, we expect a strong response of the spectrum to the perturbation  $\mathcal{L}_U$ .




FIG. 3. Single realization of the supermatrix elements of  $\mathcal{L} = \mathcal{L}_D^0 + \mathcal{L}_D^1 + \mathcal{L}_U$  in the Pauli String basis, where the shade indicates the absolute value of the matrix elements. (a) and (b) show the diagonal dominant nature of  $\mathcal{L}_D$  with eigenvalues that only depend in first order ( $\mathcal{L}_D^0$ ) only on the Pauli weight. (c) shows a block-subdiagonal structure defined by Eq. (7). By combinatorics (cf. Eq. (9)) it is possible to count the number of non-vanishing elements in the rows (one row indicated by arrows). The lower right blue frame shows exemplary how the matrix  $\mathcal{L}_{k_-, k_+}^{U(2)}$  (cf. Eq. (10)) can look like for adjacent weights  $k_-, k_+$  (here 2, 3).

We diagonalize  $\mathcal{L}^U$  in the union of the  $k_+$  and  $k_-$ -local operator subspaces,

$$\mathcal{L}_{k_-, k_+}^{U(2)} = \begin{pmatrix} 0 & \mathbf{V} \\ -\mathbf{V}^T & 0 \end{pmatrix} \quad (10)$$

where  $\mathbf{V}$  is a real matrix of matrix dimension  $n_{k_-} \times n_{k_+}$ , and  $n_k := \binom{\ell}{k} 3^k$  is the number of Pauli strings with weight  $k$ .

To quantify the imaginary perturbation of the clusters, we calculate the mean of the squared modulus of the eigenvalues  $|\lambda|^2$ . For a square matrix  $\mathbf{M}_{N \times N}$ , this is related to the matrix elements via  $|\lambda|^2 = \frac{1}{N} \text{Tr}(\mathbf{M}^\dagger \mathbf{M}) = \frac{1}{N} \sum_{i,j} |\mathbf{M}_{ij}|^2$ . One obtains an analytic prediction by further Gaussian averaging this formula. However, it is not possible to observe the weight- $k_\pm$  cluster independently since its separation to the other clusters is already broken by  $\mathcal{L}_D^1$ . Still, as shown in Fig. 1, the eigenvalue cluster with the largest negative real part shows a clear linear perturbation along the imaginary axis, since the weight-4 and 5 Pauli strings belong to the same degenerate subspace.

In the weak dissipation limit, the spread of the eigenvalues is only determined by the Hamiltonian. Since the eigenvalues of  $\mathcal{L}_U$  are given by  $\lambda_U = i(E_n - E_m)$ , where  $E_i$  are the eigenvalues of  $H$ , we can calculate the standard deviation  $\text{std}(\text{Im}(\lambda_U)) = \sqrt{2}$  using the normalization of the Hamiltonian  $\text{Tr}(H^2) = N$ . The numerical results shown in Fig. 4 agree with the scaling of the imaginary cluster extent with  $\alpha\sqrt{2}$ .

Next, we move to the second-order perturbation and calculate the mean of the perturbed cluster to see the

attraction phenomenon. Overall, we find the mean of the second-order perturbation in one  $k$ -local subspace is given by

$$\overline{\lambda_{2i}}(k) = \frac{1}{n_k} \sum_i \sum_{m \neq k, j} \frac{\langle x_{0i}^{(k)} | \mathcal{L}_U | x_{0j}^{(m)} \rangle \langle x_{0j}^{(m)} | \mathcal{L}_U | x_{0i}^{(k)} \rangle}{\lambda_0(m) - \lambda_0(k)} \quad (11)$$

where the sum does not depend on the particular basis  $\{|x_{0i}^{(k)}\rangle\}$  of a degenerate subspace containing weight  $k$  Pauli strings. In the Pauli string basis, this sum reads

$$\overline{\lambda_{2i}}(k) = \frac{1}{n_k} \sum_i \sum_{m \neq k, j} \frac{(\mathcal{L}_{ij}^{(km)})^2}{\lambda_0(m) - \lambda_0(k)}, \quad (12)$$

where  $\mathcal{L}_{ij}^{(km)}$  are the matrix elements in the block of the matrix representation given by a  $k$ -local and a  $m$ -local Pauli string. Recall that  $\mathcal{L}_{x,y}^U$  are Gaussian distributed with standard deviation  $2\sigma$  and that there are  $h(k, k-1) + h(k, k+1)$  non-vanishing entries in each row in the weight- $k$ -block. This indicates that the mean of the perturbed cluster is given by Gaussian averaging of Eq. (12)

$$\langle \overline{\lambda_{2i}}(k) \rangle = \frac{8}{9\ell(\ell-1)} \sum_{m \in \{k \pm 1\}} \frac{h(k, m)}{\lambda_0(k) - \lambda_0(m)}, \quad (13)$$

and reduces to  $\langle \overline{\lambda_{2i}}(k) \rangle \rightarrow 1/\ell^2$  in the thermodynamic limit. As indicated in Fig. 4, the analytic prediction of the mean agrees well with exact diagonalization results.




FIG. 4. Numerical calculation of the influence of unitary dynamics for different relative dissipation strength  $1/\alpha$  for one realization of  $\mathcal{L}_U$  and  $\mathcal{L}_D$ . (a) dots: numerical calculation of the shift of the cluster centers  $\bar{\lambda}(k, \alpha) - \bar{\lambda}(k, 0)$ , lines: analytic predictions given in Eq. (13). The deviation of the numerical results from the analytic results is expected due to  $\mathcal{L}_D^1$ . The shift of the cluster centers is clearly second order in  $\alpha$ . (b) shows the standard deviation  $\text{std}(\text{Im}(\lambda))$  of the eigenvalues from the real axis for different relative strengths of the dissipation  $\alpha$ . A constant offset is expected for  $\alpha = 0$  due to  $\mathcal{L}_D^1$ . We observe no significant scaling for clusters supported by eigenmodes of a single weight (see data points for  $k \in \{1, 2\}$ ), whereas the clusters that contain adjacent localities with the same analytic center ( $k = 4, 5$ ) scale approximately linear with  $\alpha$ . This supports the analytic results for the first-order perturbation.

### B. Weak dissipation

For a dominant unitary part, see Fig. 1(h), we observe a concentrated eigenvalue cluster around  $-1$  [16], but also a single persistent 2-local Pauli string supported decay mode. Since the Pauli strings span the full  $4^\ell$  dimensional Liouville operator space, one can have a decay mode equal to the Hamiltonian, which yields a trivial commutator  $\mathcal{L}_U(H) = -i[H, H] = 0$ . The weight-2-only nature of Hamiltonian makes this persistent decay mode share the same locality and lives in the 2-local subspace of  $\mathcal{L}_D^0$ . Consequently, it is an eigenvector of  $\mathcal{L}_D^0$  and  $\mathcal{L}_D^0 + \alpha\mathcal{L}_U$  with eigenvalue  $\lambda_0(2)$ . Taking  $\mathcal{L}_D^1$  as a small perturbation approximates the persistent decay mode  $\rho \approx H$  of the full Liouvillian  $\mathcal{L}$  with eigenvalue  $\lambda \approx \lambda_0(2)$ .

## IV. APPLICATION: HEISENBERG CHAIN

As a concrete example, we consider the XXX antiferromagnetic Heisenberg Hamiltonian

$$H = J \sum_{i=1}^{\ell} (\sigma_x^i \otimes \sigma_x^{i+1} + \sigma_y^i \otimes \sigma_y^{i+1} + \sigma_z^i \otimes \sigma_z^{i+1}), \quad (14)$$




FIG. 5. Spectrum of the Lindbladian for the 1D XXX Heisenberg chain with up-to-2-local random dissipation from a single realization. The upper panels demonstrate the attractions of eigenvalue clusters. The lower panel indicates the persistent clusters in the opposite limit. The left horizontal lines in the weak dissipation limit (bottom right) show the spectrum of the purely unitary Liouvillian  $\mathcal{L}_U$ . The vertical lines with numbers show the expected positions of persistent eigenmodes of the respective weights.

where  $J = 1/\sqrt{3\ell} > 0$  describes the coupling of two neighboring spins and implies the normalization  $\text{Tr}(H^2) = N$ . We further use periodic boundary conditions (PBC) with random 1- and 2-local dissipation. Figure 5 shows the Lindbladian spectrum of such a 1D random Heisenberg chain for different strengths of the unitary component. When the unitary strength  $\alpha$  is weak, the clusters show a similar attraction and merging behavior to the totally random 2-local Hamiltonian.

In the weakly dissipative case (large  $\alpha$ ), we observe the eigenvalues ordered into lines with the same imaginary part due to the discrete spectrum nature of the Heisenberg Hamiltonian and remaining well-separated clusters. The emergence of the extra persistent eigenmodes is a manifestation of the symmetry properties of the Hamiltonian. The XXX spin chain has a  $\text{SU}(2)$  symmetry algebra, which means the total spin operators

$$\rho_\alpha^{(1)} = \sum_{i=1}^{\ell} \sigma_\alpha^i \quad \alpha \in \{x, y, z\}$$

satisfy  $[H, \rho_\alpha^{(1)}] = 0$ . We note that  $\rho_\alpha^{(1)}$  are sums of Pauli strings of weight 1. This local symmetry of the XXX chain implies the existence of 3 persistent eigenmodes of weight 1 with an eigenvalue that is independent of the unitary component of  $\mathcal{L}$ .

Furthermore, we also find the following 7 eigenmodes

of weight 2 that commute with the Hamiltonian

$$\rho_1^{(2)} = H, \quad \rho_\alpha^{(2)} = \sum_{i \neq j} \sigma_\alpha^i \otimes \sigma_\alpha^j, \quad \alpha \in \{x, y, z\},$$

$$\rho_{\alpha\beta}^{(2)} = \sum_{i \neq j} \left( \sigma_\alpha^i \otimes \sigma_\alpha^j - \sigma_\beta^i \otimes \sigma_\beta^j \right), \quad \alpha \neq \beta \in \{x, y, z\},$$

to make up 10 persistent eigenmodes in total. Other conserved operators of higher weight may be observed as separated eigenvalues at the respective positions but flow to the cluster with a real part  $-1$  in the presented case.




FIG. 6. Single realization of the supermatrix elements of a 2-local Heisenberg  $\mathcal{L}_U$  ( $\ell = 4$ ) in the Pauli String basis. The XXX chain only consists of Pauli strings of weight 2, hence it only couples the adjacent weight sectors of  $\mathcal{L}_D^0$ , as predicted by Eq. (7).

The terms of the XXX chain are a subset of the totally random 2-local  $\mathcal{L}_U$ . Thus, the same perturbative treatment holds and leads to a sparse unitary matrix Eq.

(7). Figure 6 shows the realization of the supermatrix elements of a 2-local Heisenberg  $\mathcal{L}_U$  in the Pauli string basis. The weight 2 nature of the XXX chain only couples the Pauli string of adjacent weights.

## V. CONCLUSION

In this work, we present a perturbative study on how local Hamiltonians modify the hierarchy of relaxation timescales in local strongly dissipative Liouvillians. The modified timescales are determined by the weight of the Hamiltonian terms in the Pauli string basis via coupling different weight sectors of the Liouvillian. For an even-local (sum of weight  $k$  strings,  $k$  even) unitary, the separation of the eigenvalue clusters, which physically correspond to the hierarchy of relaxation, is stable up to second-order in the strength  $\alpha$  of the unitary contribution; the imaginary part of the eigenvalues depend linearly on  $\alpha$  as a first-order perturbation effect, but the real parts (responsible for the separation of eigenvalue clusters) change only in second order in  $\alpha$ . For a random 2-local Hamiltonian, we find the second-order perturbation changes of the average eigenvalues in each eigenvalue cluster to behave like  $(\alpha/\ell)^2$  in the thermodynamic limit. Our perturbation analysis is applicable to the non-random XXX antiferromagnetic Heisenberg chain, where we find one persistent eigenmode in the weak dissipation limit, which we identify as the Hamiltonian. In addition to this Hamiltonian mode, we find further persistent modes in the XXX chain given by its  $SU(2)$  operators.

## ACKNOWLEDGMENTS

This project was supported by the Deutsche Forschungsgemeinschaft (DFG) through the cluster of excellence ML4Q (EXC 2004, project-id 390534769). We further acknowledge support from the QuantERA II Programme that has received funding from the European Union's Horizon 2020 research innovation programme (GA 101017733), and from the Deutsche Forschungsgemeinschaft through the project DQUANT (project-id 499347025).

- 
- [1] E. P. Wigner, Random Matrices in Physics, *SIAM Rev.* **9**, 1 (1967).
  - [2] F. J. Dyson, Statistical Theory of the Energy Levels of Complex Systems. I, *J. Math. Phys.* **3**, 140 (1962).
  - [3] F. J. Dyson, Statistical Theory of the Energy Levels of Complex Systems. II, *J. Math. Phys.* **3**, 157 (1962).
  - [4] F. J. Dyson, Statistical Theory of the Energy Levels of Complex Systems. III, *J. Math. Phys.* **3**, 166 (1962).
  - [5] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
  - [6] C. W. J. Beenakker, Random-matrix theory of quantum transport, *Rev. Mod. Phys.* **69**, 731 (1997).
  - [7] J. Ginibre, Statistical Ensembles of Complex, Quaternion, and Real Matrices, *J. Math. Phys.* **6**, 440 (1965).
  - [8] O. Bohigas, M. J. Giannoni, and C. Schmit, Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws, *Phys. Rev. Lett.* **52**, 1 (1984).
  - [9] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
  - [10] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems,

- Nature* **452**, 854 (2008).
- [11] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From Quantum Chaos and Eigenstate Thermalization to Statistical Mechanics and Thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [12] J. M. Deutsch, Eigenstate thermalization hypothesis, *Rep. Prog. Phys.* **81**, 082001 (2018).
- [13] D. J. Luitz and Y. Bar Lev, The ergodic side of the many-body localization transition, *Ann. Phys.* **529**, 1600350 (2017).
- [14] D. A. Abanin and Z. Papić, Recent progress in many-body localization, *Ann. Phys.* **529**, 1700169 (2017).
- [15] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of N-level systems, *J. Math. Phys.* **17**, 821 (1976).
- [16] S. Denisov, T. Lapyeva, W. Tarnowski, D. Chruściński, and K. Życzkowski, Universal Spectra of Random Lindblad Operators, *Phys. Rev. Lett.* **123**, 140403 (2019).
- [17] L. Sá, P. Ribeiro, and T. Prosen, Spectral and steady-state properties of random Liouvillians, *J. Phys. A Math. Theor.* **53**, 305303 (2020).
- [18] W. Tarnowski, I. Yusipov, T. Lapyeva, S. Denisov, D. Chruściński, and K. Życzkowski, Random generators of Markovian evolution: A quantum-classical transition by superdecoherence, *Phys. Rev. E* **104**, 034118 (2021).
- [19] T. Can, Random Lindblad Dynamics, [arXiv:1902.01442](https://arxiv.org/abs/1902.01442) (2019).
- [20] S. Lange and C. Timm, Random-matrix theory for the Lindblad master equation, *Chaos* **31**, 023101 (2021).
- [21] C. Timm, Random transition-rate matrices for the master equation, *Phys. Rev. E* **80**, 021140 (2009).
- [22] T. Can, V. Oganesyan, D. Orgad, and S. Gopalakrishnan, Spectral gaps and mid-gap states in random quantum master equations, [arXiv:1902.01414](https://arxiv.org/abs/1902.01414) (2019).
- [23] T. Haga, M. Nakagawa, R. Hamazaki, and M. Ueda, Quasiparticles of Decoherence Processes in Open Quantum Many-Body Systems: Incoherentons, [arXiv:2211.14991](https://arxiv.org/abs/2211.14991) (2022).
- [24] J. Costa, P. Ribeiro, A. de Luca, T. Prosen, and L. Sá, Spectral and steady-state properties of fermionic random quadratic Liouvillians, [arxiv:2210.07959](https://arxiv.org/abs/2210.07959) (2022).
- [25] K. Wang, F. Piazza, and D. J. Luitz, Hierarchy of Relaxation Timescales in Local Random Liouvillians, *Phys. Rev. Lett.* **124**, 100604 (2020).
- [26] O. E. Sommer, F. Piazza, and D. J. Luitz, Many-body hierarchy of dissipative timescales in a quantum computer, *Phys. Rev. Research* **3**, 023190 (2021).
- [27] J. L. Li, D. C. Rose, J. P. Garrahan, and D. J. Luitz, Random matrix theory for quantum and classical metastability in local Liouvillians, *Phys. Rev. B* **105**, L180201 (2022).
- [28] H. Breuer, P. Breuer, F. Petruccione, and S. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2002).
- [29] A. Y. Kitaev, A. H. Shen, and M. N. Vyalyi, *Classical and Quantum Computation* (Boston, MA, USA, 2002).
- [30] S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum Heisenberg magnet, *Phys. Rev. Lett.* **70**, 3339 (1993).
- [31] Note that the effect of introducing spatial locality was studied in Ref. [25], leading to further spectral structures which we do not consider here.
- [32] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys* **48** (1976).
- [33] Y.-N. Zhou, L. Mao, and H. Zhai, Renyi Entropy Dynamics and Lindblad Spectrum for Open Quantum System, *Phys. Rev. Research* **3**, 043060 (2021).
- [34] V. Popkov and C. Presilla, Full spectrum of the Liouvillian of open dissipative quantum systems in the Zeno limit, *Phys. Rev. Lett.* **126**, 190402 (2021).
- [35] L. Sá, P. Ribeiro, and T. Prosen, Complex spacing ratios: A signature of dissipative quantum chaos, *Phys. Rev. X* **10**, 021019 (2020).