# Density Elicitation with applications in Probabilistic Loops\*

Andrey Kofnov<sup>1</sup>, Ezio Bartocci<sup>2</sup>, and Efstathia Bura<sup>1</sup>

<sup>1</sup> Applied Statistics, Faculty of Mathematics and Geoinformation, TU Wien <sup>2</sup> Faculty of Informatics, TU Wien

Abstract. Probabilistic loops can be employed to implement and to model different processes ranging from software to cyber-physical systems. One main challenge is how to automatically estimate the distribution of the underlying continuous random variables symbolically and without sampling. We develop an approach, which we call K-series estimation, to approximate statically the joint and marginal distributions of a vector of random variables updated in a probabilistic non-nested loop with polynomial and non-polynomial assignments. Our approach is a general estimation method for an unknown probability density function with bounded support. It naturally complements algorithms for automatic derivation of moments in probabilistic loops such as [2,24]. Its only requirement is a finite number of moments of the unknown density. We show that Gram-Charlier (GC) series, a widely used estimation method, is a special case of K-series when the normal probability density function is used as reference distribution. We provide also a formulation suitable for estimating both univariate and multivariate distributions. We demonstrate the feasibility of our approach using multiple examples from the literature.

**Keywords:** Probabilistic programs · Prob-solvable loops · Polynomial Chaos Expansion · Probability density function · Density recovery

### 1 Introduction

Probabilistic programming languages (PPL) [1] are imperative/functional programming languages providing first-class programming constructs to take random decisions and to draw samples from discrete and continuous distributions. PPL are very popular in many different applications including computer security/privacy protocols [8], distributed consensus algorithms [15], randomized algorithms [29], generative machine learning models [13] and scenario-based testing [11] of cyber-physical systems operating in uncertain environments.

One of the main challenges in probabilistic programming analysis is how to automatically estimate the probabilistic density function of the program random

<sup>&</sup>lt;sup>\*</sup> The research in this paper has been funded by the Vienna Science and Technology Fund (WWTF) [10.47379/ICT19018], the TU Wien Doctoral College (SecInt), and the FWF research project P 30690-N35.

variables without directly sampling the probabilistic program [12]. This becomes even more challenging in the presence of loops with potentially infinite execution. Recently, this problem has started to be studied for a class of probabilistic programs called *prob-solvable* loops [18,24]. The structure of a prob-solvable loop [2] consists of a set of initialization statements followed by a non-nested loop body where the variables are updated according to polynomial assignments and/or by drawing samples from statistical distributions determined by their moments. For this class of probabilistic loops, it is possible to compute automatically [2] moments of any order of the program random variables as closed-form expressions in the number of iterations using symbolic summation and polynomial algebra. [20] extends this framework to approximate moments of any order for probabilistic loops with non-polynomial assignments. This is achieved by taking advantage of the theory of polynomial chaos expansion [43] in order to approximate non-linear functional updates as sums of orthogonal polynomials. Such a model is given in the left panel of Fig. 1, which encodes in the form of a probabilistic loop with non-polynomial (trigonometric) assignments, the motion of a differential-drive mobile robot in the presence of external disturbances from [17]. The position of the robot on the 2D plane is reflected in the X, Y coordinates, and its orientation in  $\theta$ . The speed of the left and right wheels are constant and already implemented into the equations. External disturbances are modeled as  $\Omega_l \sim \text{Uniform}(-0.1, 0.1), \ \Omega_r \sim \text{Beta}(1, 3)$ . The initialization of the location variables X, Y is also random (Uniform (-0.1, 0.1)) and the angle  $\theta$  is initialized as Gaussian with mean 0 and variance 0.1.

There are several methods in statistics to infer the distribution of a random sample. In absence of additional information, nonparametric density estimation is used (e.g., histogram, k-NN, kernel density estimation, etc. [34]). In this paper, we consider the problem of deriving the distribution of random variables when we know a finite number of their moments. Such random quantities are generated, for example, in probabilistic loops, as that of Fig. 1, where the moments of the random location variables X and Y are computable at each iteration using the approach in [20]. Alternatively, one can compute the sample moments of a large number of draws of the random variable(s) resulting from repeated execution of the program at specific iterations.

In this paper we develop a method, which we call *K*-series estimation, to recover probability density functions (pdfs) with bounded support from the finite set of the moments,  $m_{i_1,i_2,...,i_k} = \mathbb{E} \left( X_1^{i_1} X_2^{i_2} \cdot \ldots \cdot X_k^{i_k} \right), 0 \leq i_j \leq d_j, d_j \in \mathbb{N}, j =$  $1, \ldots, k$ , of a random vector  $\mathbf{X} = (X_1, \ldots, X_k)^T$  using orthogonal polynomials. The results of our approach on deriving the joint distribution of the location coordinates of the robot in Fig. 1 are depicted in the top two right and bottom panels. Assuming the distribution of (X, Y) is characterizable by its moments, we derive the joint and marginal distributions of the location variables X and Y from a finite set of moments at iteration t = 25. We use the first 6 moments for the marginals and the first 48 moments for the joint distribution, obtained via simulation ( $10^6$  replications of the program in top left panel of Fig. 1). The middle and right top panels plot the marginal pdfs of X and Y, respectively.



**Fig. 1:** Probabilistic loop with non-polynomial assignment for the Differential-Drive Mobile Robot [17] (top left), the approximations of the marginal distributions with K-series (top right) and the approximation of the joint distribution with K-series (bottom left) and comparison of estimated with true histogram (bottom right).

The histograms are created based on  $10^6$  draws from the true marginals. Our estimates are in dashed blue and agree almost perfectly with the true pdfs. The left bottom panel of Fig. 1 plots the estimated joint pdf of (X, Y). The right panel draws comparative frequency bar plots of  $10^6$  true and estimated values of the bivariate random variable (X, Y) over 2-dimensional grids of the support of the bivariate distribution. Our estimate (red bars) is almost the same as the true joint pdf (blue bars) over the grid.

The features of our method are (a) its ease of computation and application, (b) its speed and (c) its ability to recover multivariate distributions. Regarding (c), [7] derived a multivariate generalized Gram–Charlier expansion that expresses an unknown multivariate pdf in terms of vector cumulants and vector derivatives of a reference pdf that involves computation of vector Hermite polynomials, which is computationally challenging.

Paper organization In Section 2 we discuss the related work, while in Section 3 we introduce the notion of univariate and multivariate K-series demonstrating how they generalize other methods. In Section 4 we show how to combine K-series with other methods that compute the statistical higher-order moments as symbolic expression in the number of the loop iterations. Section 5 provides the experimental evaluation of our approach, while in Section 6 we draw our conclusion.

### 2 Related Work

The problem of estimating statically the probability distribution of random variables in probabilistic loops has been addressed only recently in [18,24]. The work in [18] considers the class of Prob-solvable loops [2] only with polynomial assignments that generate a single random variable. [18] estimate univariate distributions with Maximum Entropy (ME) [4,23] and Gram-Charlier (GC) expansion [14]. The first approach (ME) maximizes the Shannon information entropy subject to a finite set of moments provided as input. ME cannot be expressed symbolically in terms of the moments in the number of the loop iterations. GC expansion estimates the unknown probability density function in terms of its cumulants that can be computed from its moments and admits a closed-form expression in the number of the loop iterations, so that the resulting pdf is a symbolic expression in the number of the loop iteration. [24] extends the class of Prob-solvable loops [2] by transforming probabilistic non-nested loops with *if-statements* over discrete variables with finite value domain or with conditions over continuous random variables that does not depend on the previous iteration into polynomial assignments. [24] apply the GC expansion to estimate the distribution of the random variables generated in this new class of programs. GC uses the normal as reference distribution, which limits its accuracy as an estimator of non bell-shaped distributions. Our method, K-series, does not suffer from such limitation as any distribution can be used as reference. In effect, we show that GC is a special case of K-series when a normal reference is used.

[40,10,27] are the closest approaches we found in the literature to ours. The first is based on the moments of both the unknown target and the reference pdf. K-series, on the other hand, uses only the moments of the unknown pdf in the construction of its estimate, removing the need for an additional tuning parameter; that is, the number of moments of the reference. Also, [40] developed no theory on the statistical properties of the proposed estimator nor did they provide any connection with the Gram-Charlier or other series estimators. Our approach is general in that it allows for any reference distribution, whose effect is incorporated in the construction of the orthogonal polynomials via the Gram-Schmidt orthogonalization procedure and can be tailored to improve the accuracy of the estimation. [10,27] use the same estimation procedure and conditions as ours. The important difference is the lack development of statistical properties of the estimator. [10] focuses on ascertaining that the fundamental assumption (our Assumption 2) under which the method is valid. We prove the convergence of the K-series estimator to the true pdf in  $L_1$  and  $L_2$ . A separate contribution is the recovery of the unknown target univariate and multivariate distributions of probabilistic systems extending from robotics to macroeconomics and biology.

Our approach is also applicable to the framework of [20], where non-polynomial  $L_2$  functions are represented as a finite series of polynomials using Polynomial Chaos Expansion. [20] renders probabilistic loops with general continuous functional assignments compatible with the automatic tool of [2] for exact moment computation. Furthermore, in contrast to GC, K-series can estimate the pdf of

a *vector* of random variables; that is, multivariate distributions. Thus, K-series can derive the pdf(s) for a wide class of probabilistic programs with general functional assignments that generate multiple random variables.

### 3 K-series

We call *K*-series estimation the method we develop herein to recover the joint and marginal distributions of a vector of random variables given a finite number of their moments. Our proposal generalizes Gram-Charlier (GC) series to estimate an unknown probability density function (pdf) with bounded support. Both methods require a known reference distribution in order to derive the unknown continuous pdf. The usage of a normal reference pdf is instrumental in GC series as it dictates the choice of Hermite polynomials. Our approach allows using any continuous pdf provided its support covers the support of the target pdf we want to estimate. A similar, yet complex method, was proposed in [40] as a computational tool. We present the univariate and its multivariate extension separately.

#### 3.1 Univariate K-series

Let X be a continuous random variable, supported on an arbitrary interval  $\Omega \subseteq \mathbb{R}$ . Let the cumulative distribution function (cdf)  $F_X(x)$  of X be continuously differentiable on  $\Omega$  and the corresponding pdf  $f(x) = dF_X(x)/dx$  be non-negative upper bounded everywhere on  $\Omega$  with countable zeros. Let  $M = \{m_1, m_2, \ldots, m_n, \ldots\}$  be the set of all moments of the random variable X and suppose only the first n are known. We denote this finite subset of M by  $M_n = \{m_1, \ldots, m_n\}, n \in \mathbb{N}$  and the vector with elements the moments in  $M_n$  by  $\mathbf{m}_n = (1, m_1, \ldots, m_n)^T$ . Boldface symbols will denote vectors and matrices throughout the paper.

**Definition 1.** A probability density function is said to be exponentially integrable, if there exists a positive a > 0 such that  $\int_{\mathbb{R}} \exp\{a|x|\}f(x)dx < \infty$  (see

### *[9,28])*.

A pdf supported on an infinite set is uniquely identifiable by its moments if and only if it is exponentially integrable (see [9]). Let  $\phi(x)$  be an arbitrary continuous pdf with support  $\Theta$ , where  $\Omega \subseteq \Theta$ , that is positive everywhere on  $\Theta$ . We require that either  $\Theta$  is infinite and that  $\phi(x)$  is uniquely identifiable by its moments, or  $\Theta$  is finite (bounded). Let  $H = \{h_0(x), h_1(x), \dots, h_n(x)\}, h_0(x) \equiv 1$  be a sequence of orthonormal polynomials on  $\Theta$  with respect to  $\phi(x)$ ; i.e.,

$$\langle h_i, h_j \rangle_{\phi} = \int_{\Theta} h_i(x) h_j(x) \phi(x) dx = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}.$$
 (1)

**Definition 2.** A function l(x) is said to belong to  $L_p(\Sigma, \rho)$  if  $\int_{\Sigma} |l|^p(x)\rho(x)dx < \infty$  (see [32]).

Throughout, at least one of the following two assumptions is assumed to hold.

**Assumption 1** The support  $\Omega$  of the pdf of X is a bounded set.

**Assumption 2** The ratio  $f(x)/\phi(x)$  is in  $L_1(\Omega, f)$ .

We define  $\widetilde{f}(x)$  on  $\Theta$  to be

$$\widetilde{f}(x) = \begin{cases} f(x), & x \in \Omega, \\ 0, & x \in \Theta \setminus \Omega. \end{cases}$$
(2)

Since *H* is an orthonormal system on  $\Theta$  with respect to pdf  $\phi$ , any function in  $L_2(\Theta, \phi)$  can be expanded into a Fourier series (see, e.g., [21] or [31]) along the *H* basis elements. Under Assumption 1 or 2,  $g(x) = \tilde{f}(x)/\phi(x)$  satisfies

$$\int_{\Theta} g^2(x)\phi(x)dx = \int_{\Theta \setminus \Omega} \frac{\widetilde{f}(x)}{\phi(x)}\widetilde{f}(x)dx + \int_{\Omega} \frac{f(x)}{\phi(x)}dF_X(x) < \infty,$$
(3)

so that  $g(x) \in L_2(\Theta, \phi)$ . In consequence, g has a Fourier series representation

$$g(x) = \sum_{i=0}^{\infty} \alpha_i h_i(x), \tag{4}$$

with

$$\begin{aligned} \alpha_i &= \langle g, h_i \rangle_{\phi} = \int\limits_{\Theta} g(x) h_i(x) \phi(x) dx = \int\limits_{\Theta} \frac{\widetilde{f}(x)}{\phi(x)} h_i(x) \phi(x) dx \\ &= \int\limits_{\Theta} \widetilde{f}(x) h_i(x) dx = \int\limits_{\Omega} f(x) h_i(x) dx + \int\limits_{\Theta/\Omega} \widetilde{f}(x) h_i(x) dx = \langle 1, h_i \rangle_f \,. \end{aligned}$$

From (2) and (4), an estimator of f is

$$\hat{f}(x) = \phi(x) \sum_{i=0}^{n} \langle 1, h_i \rangle_f h_i(x).$$
(5)

Each polynomial  $h_i(x)$  is a sum of monomials,  $h_i(x) = \sum_{j=0}^{i} a_{ij} x^j$ , i = 0, ..., n. Since the first *n* moments of f(x) are known, we can compute inner products by

$$\langle 1, h_i \rangle_f = \sum_{j=0}^i a_{ij} \langle 1, x^j \rangle_f = \sum_{j=0}^i a_{ij} m_j.$$
 (6)

where  $m_j$  is the *j*th raw moment of X for j = 0, ..., i, i = 0, ..., n.

**Definition 3.** The series based estimator (5) of the pdf f of X is called a K-series estimator with reference  $\phi$ , or just K-series.

Let  $\mathbf{A} = \{a_{ij}\}_{i,j=0}^{n}$  be a lower triangular matrix with entries the coefficients of the *ordered* vector of polynomials  $h_i(x)$ ,  $\mathbf{h}_n(x) = (h_0(x), \ldots, h_n(x))^T$  from H. Then, (5) can also be computed as

$$\hat{f}(x) = \phi(x) \left(\mathbf{A} \cdot \mathbf{m}_n\right)^T \cdot \mathbf{h}_n(x).$$
(7)

The only assumption the *K*-series estimator requires is that the unknown target pdf f has bounded support and the support of the reference  $\phi$  is large enough to cover it. The only constraint for the choice of the reference distribution is that it is continuous with support larger than that of the target pdf. Any such pdf can serve as reference and thus polynomials  $h_i$  of any order in (5) can be computed using the Gram-Schmidt orthogonalization procedure.

#### 3.2 Examples of K-series estimation

We illustrate the K-series estimation approach with two examples: for the first (Truncated Exponential), we let the target pdf be truncated exponential with known parameters and support and derive its first two moments and its K-series estimate. In the second (Irwin-Hall Distribution), we express the distribution generating algorithm as a prob-solvable loop, compute its *exact* moments using the POLAR tool [24] and its K-series estimate.

**Truncated Exponential.** Suppose  $X \sim Trunc Exp(1, [0, 1])$  and that its first two moments are known. That is, let us suppose, that on the known support  $\Omega = [0, 1]$  we need to recover pdf using the first 2 moments from the set  $M_2 = \{m_1 = 0.418023, m_2 = 0.254070\}$ . We let the reference distribution be the uniform with the same support as the target unknown distribution; i.e.,  $\phi(x) = 1$ for  $x \in [0, 1]$ . Since  $\phi$  is uniform, we use the shifted and scaled Legendre polynomials as the orthonormal basis in the series (see [43]). That is,

$$h_0 = 1, \quad h_1 = \sqrt{3}(2x - 1), \quad h_2 = \sqrt{5}(6x^2 - 6x + 1).$$
 (8)

In order to compute the unknown pdf estimator in (5), we need to compute the  $\alpha_i$  coefficients in (4). By (6), this requires the substitution of  $x^i$  with the corresponding moment  $m_i$  in  $M_2$  for i = 1, 2. Doing so yields  $\alpha_0 = 1, \alpha_1 = \sqrt{3}(2 \cdot 0.418023 - 1) = -0.283976, \alpha_2 = \sqrt{5}(6 \cdot 0.25407 - 6 \cdot 0.418023 + 1) = 0.036407$ . The K-series estimator is

$$\hat{f}(x) = 1 - 0.283976 \cdot h_1(x) + 0.036407 \cdot h_2(x).$$

It almost overlaps the true truncated exponential pdf in panel (a) of Figure 2.

The Irwin-Hall Distribution. Irwin–Hall is the probability distribution of a sum of independent uniform random variables on the unit interval (uniform



**Fig. 2:** K-series approximation of the truncated exponential distribution (panel (a)) and the Irwin-Hall distribution (panel (b))

sum distribution). That is,  $X \sim \text{Irwin-Hall}(t)$  if  $X = \sum_{i=1}^{t} U_i$ , for  $U_i$  independent and identically distributed (i.i.d.) as Uniform(0,1). This distribution, parameterized by the number of its summands, is encodable as the Prob-solvable loop in Listing 1.1.

Listing 1.1: Irwin-Hall

 $\begin{array}{l} x:=0\\ while(True):\\ u_x:=Uniform(0,1)\\ x:=x+u_x\\ end \end{array}$ 

At each iteration t, the support of x is (0, t). The K-series estimator requires knowledge of a finite number of moments. Since the Irwin-Hall distribution is equivalent to a Prob-solvable loop, its exact n first moments can be computed with the algorithm in [2]:

$$M(t) = \left\{ \frac{t}{2}, \frac{t(3t+1)}{12}, \frac{t^2(t+1)}{8}, \frac{t(15t^3+30t^2+5t-2)}{240}, \frac{t^2(3t^3+10t^2+5t-2)}{96}, \frac{t(63t^5+315t^4+315t^3-91t^2-42t+16)}{4032}, \dots \right\}$$

The first 6 moments of Irwin-Hall(3) are  $M_6(3) = \left\{\frac{3}{2}, \frac{5}{2}, \frac{9}{2}, \frac{43}{5}, \frac{69}{4}, \frac{3025}{84}\right\}$ . We use the Uniform[0,3] as reference and construct the K-series estimator of the pdf of x at iteration t = 3 with the 6 first moments and the first 7 shifted and scaled Legendre polynomials. The true pdf and its K-series estimate are plotted in panel (b) of Figure 2, where we can see their close agreement.

### 3.3 K-series as a generalization of other estimation methods

The K-series density estimator generalizes the widely used Gram-Charlier (GC) density estimator. The GC series approximates the pdf f of a random variable X in terms of its cumulants and a known reference distribution  $\phi$  (see, e.g., [6,19]). The GC (type-)A estimate of the pdf f of X is given by

$$f_{\rm GC}(x) = \phi(x) \sum_{n=0}^{\infty} (-1)^n c_n H e_n(x) , \qquad (9)$$

where  $c_n = (-1)^n \int_{-\infty}^{\infty} f(t) He_n(t) dt/n!$ ,  $\phi$  is the standard normal pdf, and

$$He_n(x) = n! \sum_{k=0}^{[n/2]} \frac{(-1)^k x^{n-2k}}{k! (n-2k)! 2^k}.$$

Proposition 1 shows that the GC series A estimator in (9) is a special case of the K-series estimator.

**Proposition 1.** Suppose the reference  $pdf \phi$  is normal with mean and variance corresponding to the first and second moments of the target pdf f. Then, the K-series estimator (5) equals Gram-Charlier estimator (9).

Proposition 1 is easy to obtain using the standard normal as reference pdf and replacing the polynomials  $h_i$  in (5) by  $He_i/\sqrt{i!}$ .

[25] used the method of moments estimation to develop an algorithm for an n-order polynomial approximation of a pdf, based on its consecutive n first moments. They referred to the estimation algorithm as *Method of Moments (MM)*. MM estimates parameters of a target distribution f by equating sample moments with the corresponding moments of the distribution. The approximation is carried out on the interval where they wish to maximize accuracy. In practice, this is the same as assuming finite or bounded support. [25] showed in practice that MM beats the GC expansion for several distributions, such as the Weibull on a positive finite support.

The MM algorithm starts by choosing an interval [a, b] that is thought to contain most of the mass of the target unknown distribution. Using a finite set of moments  $\{m_1, \ldots, m_n\}$  and  $m_0 = 1$ , MM constructs a polynomial estimator  $\hat{f}(x)$  by solving a linear system of equations,

$$m_{i} = \int_{a}^{b} x^{i} \hat{f}(x) dx, \ i = 0, \dots, n,$$
(10)

which yields the coefficients  $p_i$  of the series representation  $\hat{f}_{MM}(x) = \sum_{i=0}^n p_i x^i$ .

Let  $\mathbf{m}_n = (1, m_1, \dots, m_n)^T$ ,  $\mathbf{p}_n = (p_0, p_1, \dots, p_n)^T$ , and  $\mathbf{x}_n = (1, x, \dots, x^n)^T$ . The linear system (10) can be expressed in matrix form as

$$\mathbf{m}_n = \mathbf{M}_{ab} \cdot \mathbf{p}_n,\tag{11}$$

where  $\mathbf{M}_{ab}$  is the matrix with elements the integrals of powers of x over the interval [a, b],

$$\mathbf{M}_{ab} = \begin{pmatrix} b-a & \frac{b^2-a^2}{2} & \dots & \frac{b^{n+1}-a^{n+1}}{n+1} \\ \frac{b^2-a^2}{2} & \frac{b^3-a^3}{3} & \dots & \frac{b^{n+2}-a^{n+2}}{n+2} \\ \vdots & \ddots & \vdots \\ \frac{b^{n+1}-a^{n+1}}{n+1} & \frac{b^{n+2}-a^{n+2}}{n+2} & \dots & \frac{b^{2n+1}-a^{2n+1}}{2n+1} \end{pmatrix}$$
(12)

Theorem 1 shows that the Method of Moments of [25] is a special case of the K-series estimator.

**Theorem 1.** Suppose the reference  $pdf \phi$  is the uniform with the same support as the target pdf f. Then, the MM estimator coincides with the K-series estimator (5).

*Proof.* Suppose f is supported on (a, b). Then,  $\phi(x) = \phi = 1/(b-a)$ . Let  $\{l_j(x) = \sum_{i=0}^{j} \lambda_{ji} x^i\}_{j=0}^n$  be the set of the first n shifted scaled Legendre polynomials that are orthonormal on [a, b], so that  $\mathbf{\Lambda} = (\lambda_{ji})_{j,i=0}^n$  is a lower triangular matrix.

Every polynomial of degree n can be expressed as a weighted sum of polynomials of degree up to n. In such a case, we can represent the MM estimator  $\hat{f}_{\text{MM}} = \sum_{i=0}^{n} p_i x^i$  as a weighted sum of Legendre polynomials  $1, l_1(x), \ldots, l_n(x)$  with weight coefficients  $\phi \cdot a_j, j = 0, \ldots, n$ . Then,

$$\hat{f}_{\rm MM}(x) = \sum_{i=0}^{n} p_i x^i = \phi \sum_{j=0}^{n} a_j l_j(x) = \phi \sum_{j=0}^{n} a_j \sum_{i=0}^{j} \lambda_{ji} x^i,$$

or, equivalently,

 $\mathbf{p}_n^T \mathbf{x}_n = \phi \cdot \mathbf{a}_n^T \mathbf{l}_n = \phi \cdot \mathbf{a}_n^T \mathbf{\Lambda} \mathbf{x}_n,$ where  $\mathbf{l}_n = (1, l_1(x), \dots, l_n(x))^T$  and  $\mathbf{a}_n = (1, a_1, \dots, a_n)^T$ . Then,

$$\mathbf{p}_n = \phi \cdot \boldsymbol{\Lambda}^T \mathbf{a}_n$$

Now, (11) implies

$$\mathbf{m}_n = \mathbf{M}_{ab} \cdot \mathbf{p}_n = \phi \cdot \mathbf{M}_{ab} \cdot \mathbf{\Lambda}^T \mathbf{a}_n$$

In the matrix form of the K-series estimator (7),  $\mathbf{\Lambda} \cdot \mathbf{m}_n = \mathbf{a}_n$ . It suffices to show that

$$\phi \cdot \mathbf{M}_{ab} \cdot \boldsymbol{\Lambda}^T = \boldsymbol{\Lambda}^{-1}. \tag{13}$$

The matrix  $\phi \cdot \mathbf{M}_{ab}$  contains the moments of the uniform distribution. Therefore,  $\boldsymbol{A}$  is a matrix with entries the coefficients of orthonormal polynomials and the left lower triangular factor of the Cholesky decomposition of the moment matrix  $\phi \cdot \mathbf{M}_{ab}$ . Thus, (13) follows from [37, Prop. 2(i)].

**Theorem 2.** Let  $\phi(x)$  be continuous, positive everywhere on  $\Theta: \Omega \subseteq \Theta$  and either (a)  $\Theta$  is infinite and  $\phi(x)$  is uniquely identifiable by its moments, or (b)  $\Theta$  is finite (bounded). Assume Assumption 1 or 2 holds. Then, the K-series estimator (5) converges to the true pdf (2),  $\tilde{f}(x)$ , in  $L_1(\Theta, 1)$ . Moreover, if  $\phi(x)$ is a uniform pdf, it converges in  $L_2(\Theta, 1)$ . Proof.

$$\left\|\frac{\tilde{f}(x)}{\phi(x)} - \sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right\|_{\phi}^{2} = \int_{\Theta} \left[\frac{\tilde{f}(x)}{\phi(x)} - \sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right]^{2} \phi(x)dx$$

$$= \int_{\Theta} \left[\tilde{f}(x) - \phi(x)\sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right]^{2} \frac{1}{\phi(x)}dx$$

$$= \int_{\Theta} \phi(x)dx \int_{\Theta} \left[\tilde{f}(x) - \phi(x)\sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right]^{2} \frac{1}{\phi(x)}dx$$

$$= ||\sqrt{\phi(x)}||_{1}^{2} \cdot \left\|\left(\tilde{f}(x) - \phi(x)\sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right) \frac{1}{\sqrt{\phi(x)}}\right\|_{1}^{2}$$

$$\geq \left(\int_{\Theta} \left|\tilde{f}(x) - \phi(x)\sum_{i=0}^{n} \alpha_{i}h_{i}(x)\right|dx\right)^{2}, \quad (14)$$

where the last inequality is due to Cauchy-Schwarz inequality. The function  $\tilde{f}(x)$  in (2) is a density. In the case where  $\Theta$  is bounded,  $\phi(x)$  is uniquely identifiable by its moments. When  $\Theta$  is infinite,  $\phi(x)$  is exponentially integrable by the assumption (a) of the theorem. Hence, for all  $n \geq 1$ ,  $\phi(x) \left| \sum_{i=0}^{n} \alpha_i h_i(x) \right|$  is integrable.

Now, since the truncated series  $\sum_{i=0}^{n} \alpha_i h_i(x)$  in (4) converges to  $g(x) = \tilde{f}(x)/\phi(x)$ in  $L_2(\Theta, \phi)$ , as  $n \to \infty$ , (14) obtains that the K-series estimator (5) converges to the extended true pdf  $\tilde{f}(x)$  in  $L_1(\Theta, 1)$ .

Next, suppose  $\phi(x)$  is the pdf of the uniform distribution, so that  $\Theta$  is bounded, and  $\phi(x) = c$ . Then,

$$\left\|\frac{\widetilde{f}}{\phi} - \sum_{i=0}^{n} \alpha_{i} h_{i}(x)\right\|_{\phi}^{2} = \int_{\Theta} \left[\frac{\widetilde{f}(x)}{\phi(x)} - \sum_{i=0}^{n} \alpha_{i} h_{i}(x)\right]^{2} \phi(x) dx$$
$$= \frac{1}{c} \int_{\Theta} \left[\widetilde{f}(x) - c \sum_{i=0}^{n} \alpha_{i} h_{i}(x)\right]^{2} dx$$

Hence,  $\tilde{f}(x) = c \cdot g(x)$  is in  $L_2(\Theta, 1)$ , and  $\int_{\Theta} c^2 \left[\sum_{i=0}^n \alpha_i h_i(x)\right]^2 dx$  is an integral of a polynomial over a bounded interval, so that the K-series estimator (5) converges to the true pdf  $\tilde{f}(x)$  in  $L_2(\Theta, 1)$ .

#### 3.4 Multivariate K-series

K-series density estimation is easily generalisable to multivariate distributions by considering the product of independent univariate distributions as the reference

11

pdf. The coefficients of the corresponding multivariate orthogonal polynomials recover the multivariate dependence structure via their joint moments.

Let  $\mathbf{X} = (X_1, \ldots, X_k)^T$  be a vector of continuous random variables with joint non-negative pdf  $f(\mathbf{x})$ , upper bounded and supported on  $\Omega$  with countable zeros. Suppose that there exists a k-dimensional compact cube  $\mathcal{Q}$ , such that  $\Omega \subseteq \mathcal{Q}$ . We assume that a finite number of moments, not necessarily an equal number for all, is known for each  $X_i$ ,  $i = 1, \ldots, k$ , and all cross-product moments are also known. That is, we assume the set  $M_{d_1,\ldots,d_k}$  is known.

$$M_{d_1,\dots,d_k} = \left\{ m_{i_1,\dots,i_k} = \mathbb{E}\left( X_1^{i_1}\dots X_k^{i_k} \right) : i_j = 0,\dots,d_j, d_j \in \mathbb{N}, j = 1,\dots,k \right\}$$
(15)

Let  $\mathbf{Z} = (Z_1, \ldots, Z_k)^T$  be a vector of continuous independent random variables and  $\tilde{\phi}(\mathbf{z}) = \prod_{i=0}^k \phi_i(z_i)$  be its pdf with support  $\Theta$ , where  $\Omega \subseteq \Theta$ , that is positive everywhere on  $\Theta$ . We require that either  $\Theta$  is unbounded and that  $\tilde{\phi}(\mathbf{z})$  is uniquely identifiable by its moments, or  $\Theta$  is bounded (see [28]). Let

$$\tilde{h}_{i_1,\dots,i_k}(\mathbf{z}) = \prod_{j=1}^k h_{i_j}^j(z_j),$$
(16)

where  $h_{i_j}^j(z_j)$  is a polynomial of degree  $i_j$  that belongs to the set of orthogonal polynomials with respect to  $\phi(z_j)$ ,  $i_j = 0, \ldots, d_j$ ,  $j = 1, \ldots, k$ , that are calculated with the Gram-Schmidt orthogonalization procedure<sup>3</sup>. The set  $H = \{\tilde{h}_{i_1,\ldots,i_k}(\mathbf{z}), i_j = 0, \ldots, d_j, d_j \in \mathbb{N}, j = 1, \ldots, k\}$  contains the k-variate orthonormal polynomials on  $\Theta$  with respect to  $\tilde{\phi}(\mathbf{z})$ .

As in the univariate case, we require Assumption 1 hold and we let

$$\widetilde{f}(\mathbf{z}) = \begin{cases} f(\mathbf{z}), & \mathbf{z} \in \Omega, \\ 0, & \mathbf{z} \in \Theta \setminus \Omega. \end{cases}$$
(17)

Then,  $\tilde{f}(\mathbf{z})/\tilde{\phi}(\mathbf{z}) = g(\mathbf{z})$  which is approximated by

$$\hat{g}(\mathbf{z}) = \sum_{\substack{i_j \in \{0, \dots, d_j\}, \\ j=1, \dots, k}} \alpha(i_1, \dots, i_k) \tilde{h}_{i_1, \dots, i_k}(\mathbf{z}) = \sum_{\substack{i_j \in \{0, \dots, d_j\}, \\ j=1, \dots, k}} \alpha(i_1, \dots, i_k) \prod_{j=1}^{k} h_{i_j}^j(z_j),$$

<sup>&</sup>lt;sup>3</sup> Generalized PCE typically entails using orthogonal basis polynomials specific to the distribution of the basic variables, according to the Askey scheme of [43,42]. We opted for the most general procedure that can be used for any basic continuous variable distribution.

where the Fourier coefficients  $\alpha(i_1, \ldots, i_k)$  are calculated as follows,

$$\alpha(i_{1},\ldots,i_{k}) = \left\langle g,\tilde{h}_{i_{1},\ldots,i_{k}}\right\rangle_{\widetilde{\phi}} = \int_{\Theta} g(\mathbf{z})\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})\widetilde{\phi}(\mathbf{z})d\mathbf{z}$$
$$= \int_{\Theta} \frac{\widetilde{f}(\mathbf{z})}{\widetilde{\phi}(\mathbf{z})}\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})\widetilde{\phi}(\mathbf{z})d\mathbf{z} = \int_{\Theta} \widetilde{f}(\mathbf{z})\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})d\mathbf{z}$$
$$= \int_{\Omega} f(\mathbf{z})\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})d\mathbf{z} + \int_{\Theta/\Omega} \widetilde{f}(\mathbf{z})\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})d\mathbf{z}$$
$$= \left\langle 1,\tilde{h}_{i_{1},\ldots,i_{k}}(\mathbf{z})\right\rangle_{f}.$$
(18)

Since for all  $i_j = 0, \ldots, d_j, j = 1, \ldots, k, h_{i_j}^j(z_j) = \sum_{l=0}^{i_j} a_{i_j l}^j z_j^l$ , their product is

$$\tilde{h}_{i_1,\dots,i_k}(\mathbf{z}) = \prod_{j=1}^k h_{i_j}^j(z_j) = \prod_{j=1}^k \sum_{l=0}^{i_j} a_{i_j l}^j z_j^l = \sum_{\substack{l_j \in \{0,\dots,i_j\}, \\ j=1,\dots,k}} z_1^{l_1} \cdots z_k^{l_k} \prod_{j=1}^k a_{i_j l_j}^j.$$

Assuming all first cross-moments of  $f(\mathbf{z})$ ,  $m_{l_1,\ldots,l_k} = \mathbb{E}_f\left(z_1^{l_1}\cdots z_k^{l_k}\right)$  are known, we can compute (18) as

$$\left\langle 1, \tilde{h}_{i_1,\dots,i_k}(\mathbf{z}) \right\rangle_f = \sum_{\substack{l_j \in \{0,\dots,i_j\}, \\ j=1,\dots,k}} \left\langle 1, z_1^{l_1} \cdots z_k^{l_k} \right\rangle_f \prod_{j=1}^k a_{i_j l_j}^j = \sum_{\substack{l_j \in \{0,\dots,i_j\}, \\ j=1,\dots,k}} m_{l_1,\dots,l_k} \prod_{j=1}^k a_{i_j l_j}^j \right\rangle$$
(19)

The multivariate K-series estimator of f is

$$\hat{f}(\mathbf{x}) = \widetilde{\phi}(\mathbf{x}) \sum_{\substack{i_j \in \{0, \dots, d_j\}, \\ j=1, \dots, k}} \left\langle 1, \widetilde{h}_{i_1, \dots, i_k}(\mathbf{z}) \right\rangle_{\mathbf{f}} \widetilde{h}_{i_1, \dots, i_k}(\mathbf{z}).$$
(20)

The next example illustrates the algorithm.

**Truncated Bivariate Normal.** Suppose we want to recover the joint pdf of two random variables X and Y on a set  $\Omega = [-2, 2] \times [-4, 5]$  using their first eight cross-moments,

$$\left( m_{x^{j}y^{i}} = \mathbb{E}(X^{j}Y^{i}) \right)_{i,j=0,\dots,2} = \begin{pmatrix} m_{x^{0}y^{0}} m_{x^{1}y^{0}} m_{x^{2}y^{0}} \\ m_{x^{0}y^{1}} m_{x^{1}y^{1}} m_{x^{2}y^{1}} \\ m_{x^{0}y^{2}} m_{x^{1}y^{2}} m_{x^{2}y^{2}} \end{pmatrix} = \begin{pmatrix} 1.00000 \ 0.71721 \ 1.13054 \\ 1.99556 \ 1.43124 \ 2.25606 \\ 4.96894 \ 3.56379 \ 5.61757 \end{pmatrix} .$$

$$(21)$$

We choose the reference marginal pdfs be both truncated normal  $\phi_x(z_x)$  and  $\phi_y(z_y)$  with  $Z_x \sim Trunc \mathcal{N}(m_x, m_{x^2} - m_x^2, [-2, 2]) = Trunc \mathcal{N}(0.71721, 0.61614,$ 

13

[-2, 2]), and  $Z_y \sim Trunc \mathcal{N}(m_y, m_{y^2} - m_y^2, [-4, 5]) = Trunc \mathcal{N}(1.99556, 0.98667, [-4, 5])$ , respectively.

We construct sets of univariate orthonormal polynomials using, for example, the Gram-Schmidt orthogonalization procedure, and obtain

 $\begin{array}{c} h_0^x(z_x)=1, & h_0^y(z_y)=1, \\ h_1^x(z_x)=1.42119z_x-0.89705, & h_1^y(z_y)=1.01307z_y-2.01751, \\ h_2^x(z_x)=1.58907z_x^2-1.63885z_x-0.38542, & h_2^y(z_y)=0.74083z_y^2-2.92557z_y+2.16624 \end{array}$ 

Hence, starting from a reference joint pdf that is the product of the pdfs of the independent random variables  $Z_x$  and  $Z_y$ ,  $\tilde{\phi}(z_x, z_y) = \phi_x(z_x)\phi_y(z_y)$ , the multivariate orthogonal polynomials are simply all the pairwise products of univariate polynomials:

$$\begin{split} \dot{h}_{0,0}(z_x, z_y) &= 1 \\ \tilde{h}_{0,1}(z_x, z_y) &= 1.01307 z_y - 2.01751 \\ \tilde{h}_{0,2}(z_x, z_y) &= 0.74083 z_y^2 - 2.92557 z_y + 2.16624 \\ \tilde{h}_{1,0}(z_x, z_y) &= 1.42119 z_x - 0.89705 \\ \tilde{h}_{1,1}(z_x, z_y) &= 1.43976 z_x z_y - 2.86727 z_x - 0.90877 z_y + 1.80981 \\ \tilde{h}_{1,2}(z_x, z_y) &= 1.05286 z_x z_y^2 - 4.15779 z_x z_y + 3.07864 z_x - 0.66456 z_y^2 + 2.62438 z_y - 1.94323 \\ \tilde{h}_{2,0}(z_x, z_y) &= 1.58907 z_x^2 - 1.63885 z_x - 0.38542 \\ \tilde{h}_{2,1}(z_x, z_y) &= 1.60984 z_x^2 z_y - 3.20596 z_x^2 - 1.66027 z_x z_y + 3.30634 z_x - 0.39046 z_y + 0.77759 \\ \tilde{h}_{2,2}(z_x, z_y) &= 1.17723 z_x^2 z_y^2 - 4.64894 z_x^2 z_y + 3.44231 z_x^2 - 1.21411 z_x z_y^2 + 4.79457 z_x z_y \\ &\quad - 3.55014 z_x - 0.28553 z_y^2 + 1.12757 z_y - 0.83491 \end{split}$$

In order to compute the coefficients  $\alpha(i_1, i_2)$  of the PCE (20) along the reference pdf  $\tilde{\phi}(z_x, z_y)$  for each polynomial  $\tilde{h}_{i_1, i_2}(z_x, z_y)$ , according to (19), we need to substitute every monomial factor  $z_x^j z_y^i$  by the corresponding moment  $m_{x^j y^i}$  from (21) in each polynomial. For example, the coefficient of  $\tilde{h}_{1,1}(z_x, z_y)$  is  $1.43976m_{xy} - 2.86727m_x - 0.90877m_y + 1.80981 = 1.43976 \cdot 1.43124 - 2.86727 \cdot 0.71721 - 0.90877 \cdot 1.99556 + 1.80981 = 0.00051$ . The resulting estimator is

$$\begin{aligned} \hat{f}(z_x, z_y) &= \phi_1(z_x)\phi_2(z_y) \sum_{i_1, i_2=(0,0)}^{(2,2)} \alpha(i_1, i_2)\tilde{h}_{i_1, i_2}(z_x, z_y) \\ &= \phi_1(z_x)\phi_2(z_y) \times \left[ 1 + 0.00415 \cdot \tilde{h}_{0,1}(z_x, z_y) + 0.00924 \cdot \tilde{h}_{0,2}(z_x, z_y) \right. \\ &+ 0.12224 \cdot \tilde{h}_{1,0}(z_x, z_y) + 0.00051 \cdot \tilde{h}_{1,1}(z_x, z_y) + 0.00113 \cdot \tilde{h}_{1,2}(z_x, z_y) \\ &+ 0.23568 \cdot \tilde{h}_{2,0}(z_x, z_y) + 0.00098 \cdot \tilde{h}_{2,1}(z_x, z_y) + 0.00218 \cdot \tilde{h}_{2,2}(z_x, z_y) \right] \end{aligned}$$

The estimated bivariate density is plotted in Fig. 3 (a). In panel (b), we plot the frequencies of X and Y under the true f(x, y) (blue bars) and its K-series (red



**Fig. 3:** K-series estimates of the truncated bivariate normal distribution f(x, y) = Trunc Normal((1, 2), (1, 1), -0.3, [-2, 2], [-4, 5]).

bars) pdf estimate over a 2D grid comprising of eight parallelograms, where we can see their close agreement.

A similar approach of estimating the likelihood ratio of target and reference pdfs had appeared in [10,27], without any statistical guarantees of its accuracy. Our K-series estimation method was independently developed and we prove its convergence to the true pdf, at least in  $L_1$ . We also show that the Method of Moments of [25] is a special case of K-series.

### 4 Symbolic K-series representation along iterations

The K-series estimator can be expressed as a quantitative invariant in the sense that its formula is a function of loop iterations. In the univariate case, the K-series estimator (5) of the unknown pdf of the random variable X is  $\hat{f}(x) = \phi(x) \sum_{i=0}^{n} \left( \sum_{j=0}^{i} a_{ij} m_j \right) h_i(x)$ , where  $m_j = \mathbb{E}(X^j)$ . The estimator is a function of the moments of X, which in turn, vary along iterations in a probabilistic loop. That is, the K-series estimator can be equivalently expressed as

$$\hat{f}_t(x) = \phi(x) \sum_{i=0}^n \left( \sum_{j=0}^i a_{ij} m_j(t) \right) h_i(x),$$
(22)

where  $m_j(t) = \mathbb{E}(X^j(t))$  is the moment of the random variable X at iteration t. Formula (22) is the symbolic representation of the K-series pdf estimator as a function of iteration number.

Similarly, the multivariate K-series estimator (20) can be written as

$$\hat{\mathbf{f}}(\mathbf{x}) = \widetilde{\phi}(\mathbf{x}) \sum_{\substack{i_j \in \{0, \dots, d_j\}, \\ j=1, \dots, k}} \left( \sum_{\substack{l_j \in \{0, \dots, i_j\}, \\ j=1, \dots, k}} m_{l_1, \dots, l_k}(t) \prod_{j=1}^k a_{i_j l_j}^j \right) \widetilde{h}_{i_1, \dots, i_k}(\mathbf{z})$$
(23)

where  $m_{l_1,\ldots,l_k}(t) = \mathbb{E}\left(X_1^{l_1}(t) \cdot \ldots \cdot X_k^{l_k}(t)\right)$  at iteration t, since the moments of the random vector depend on the iteration in a probabilistic loop. We illustrate the representation (22) with an example in Appendix 4.

### 5 Experiments

We carried out K-series estimation of the distributions of the random variables generated in the execution of the probabilistic loops in Figs. 4, 9 and 13. Panels A and A1 in Fig. 4 encode the same turning vehicle model [33,20] with the difference that the variance of the basic random variables  $\psi$  and  $w_2$  in A is about 3 times larger than in A1. The effect of this on the joint and marginal distributions of X, Y can be seen in Figs. 7 and 8. In the case of larger variance (A, Fig. 7), the reference is truncated normal on  $(-18, 18) \times (-20, 20)$  with mean the sample mean and variance the sample variance of the marginal distributions of X and Y, respectively. For the smaller variance case (A1, Fig. 8), the reference pdf is truncated normal on  $(1, 18) \times (-15, 15)$  with mean the sample mean and variance 4 for X and the sample variance of the Y distribution. While the support of X is not important, the accuracy of the estimation depends on the variance for X. The K-series estimator is a sum of weighted orthonormal polynomials whose Gram-Schmidt orthogonalization with respect to the reference distribution involves the variance of the generated variables in the denominator. Thus, when the variance is very small, the fraction explodes and the estimation becomes highly numerically unstable. This can be managed by increasing the variance of the reference, as done in Fig. 8.

In Fig. 4, Panel B encodes the Taylor rule [39,20], C the 2D robotic arm [5], D the rimless wheel walker [35], and E the Vasicek [41,18] model. The Taylor rule (B), rimless wheel (D) and Vasicek model (E) generate a single random variable at each iteration. We plot the histograms from sampling the probabilistic loop programs (blue) and the overlaid pdf K-series estimates of these models in Fig. 5. The 2D robotic arm model generates a bivariate random variable. We plot the marginal pdf K-series estimates in the right panels, the joint pdf approximation in the bottom left panel, and the comparison of the true (blue bars) with our estimate (red bars) over twelve 2D parallelogram grid in the top left panel of Fig. 6. The moments of the true distribution were computed with the method in [20] for the Taylor rule, and in [2] for the rimless wheel, Vasicek and 2D robotic arm models. We used the following reference functions: truncated normal on (-30, 30) for the Taylor rule, truncated normal on (0, 10) for rimless wheel, normal distribution for the Vasicek model, and truncated normal on  $(260, 280) \times (525, 540)$  for the 2D robotic arm model. For all univariate and the bivariate models, our K-series estimator exhibits excellent estimation accuracy.

We explore the robustness of our method to violations of the assumption of continuity of random variables in Fig. 9 and Fig. 13 where we estimate the distributions of random variables generated in Prob-solvable loops with discrete random components. Panel A in Fig. 9 encodes the *Stuttering P model* in [2] and panel B the piece-wise deterministic process, or *PDP model*, modeling gene circuits that can be used to estimate the bivariate distribution of protein x and the mRNA levels y in a gene [16]. For the *Stuttering P* model, we used a truncated normal distribution on (0, 50) with true mean and variance as the reference pdf. For the *PDP* model, we used the truncated normal on (100, 1800) for X and uniform on (8, 80) for Y as reference pdfs to estimate marginal pdfs of X and Y and joint pdf (X, Y). The parameters of the truncated normal distribution are the exact mean and variance of marginal pdf of the corresponding variable computed using [2]. We also estimated the pdfs of the random variables in 1D and 2D random walks and report the results in Appendix 3.

We carry out formal statistical tests of goodness-of-fit of our estimates and the true (sampling) pdfs in Table 5 in Appendix 2.

### 6 Conclusion

We presented K-series, a density estimation method based on a series of a basis of orthogonal polynomials that target the unknown density via the choice of the reference probability density function. K-series is a general density estimation method that only requires a finite number of moments of the unknown pdf and includes existing series based density estimation, such as Gram-Charlier, as special cases.

K-series is a natural complement to automated tools for exact moment computation in probabilistic loops, such as Polar [24] and Mora [3], that can handle polynomial assignments and *if-then-else* conditions with certain restrictions [24]. For non-polynomial assignments, [20] provides approximations of exact moments of the target pdf. However, the use of approximations increases the error accumulation in K-series estimation (or any estimation). As future work, we plan to explore the Fourier series representation of functions in sines and cosines in conjunction with [17], which obtains exact moments for sine and cosine assignments, to reduce the estimation error for fixed loop iterations. Finally, we plan to implement a tool to automate the entire procedure.



Fig. 4: Probabilistic loops: (A) Turning vehicle model [33,20], (A1) Small variance Turning vehicle model [33,20], (B) Taylor rule [39,20], (C) 2D Robotic Arm [5], (D) Rimless Wheel Walker [36], (E) Vasicek model (truncated version) [41,18].



**Fig. 5:** K-series estimates of pdf for variable **A**) *i* at iterations t = 20 in Fig. 4 (B): Taylor rule model, **B**) *x* at iteration t = 2000 in Fig. 4 (D): rimless wheel model, **C**) *r* at iteration t = 100 in Fig. 4 (E): Vasicek model.



**Fig. 6:** Robotic arm model Fig. 4 (C): K-series estimates of the marginal pdfs of X and Y (right upper and lower panels), the joint (lower left panel) and comparison bar plot (upper left panel) at iteration t = 100.



**Fig. 7:** Turning vehicle model Fig. 4 (A): K-series estimates of the marginal pdfs of X and Y (right upper and lower panels), the joint (lower left panel) and comparison bar plot (upper left panel) at iteration t = 20.



Fig. 8: Small variance Turning vehicle model Fig. 4 (A1): K-series estimates of the marginal pdfs of X and Y (right upper and lower panels), the joint (lower left panel) and comparison bar plot (upper left panel) at iteration t = 20.



**Fig. 9:** K-series estimates of the pdf of variable S in Stuttering P model [2] (A) at iteration t = 10, marginal pdfs for variables X, Y and joint distribution for variables (X, Y) in PDP model [16] (B) at the iterations t = 100.

### Appendix 1. Effect of Reference Distribution

We study the effect of the choice of the reference distribution in K-series on estimation accuracy. We consider reference distributions with the same support as the target unknown pdf f, with bounded support that contains the support of f and with infinite support in absence of any knowledge about the possible values of the target distribution.

Target pdf $f$	Reference pdf $\phi$
Trunc Exp $(\lambda = 2/3, [0, 4])$	
	Uniform(0,4)
	Trunc Normal( $\mathbb{E}(f), \mathbb{Var}(f), [0, 4]$ )
	Uniform(-2, 6)
	Trunc Normal( $\mathbb{E}(f)$ , $\mathbb{Var}(f)$ , $[-2, 6]$ )
	$Normal(\mathbb{E}(f), \mathbb{Var}(f))$
Trunc Gamma( $\alpha = 2, \beta = 0.5, [0, 5]$ )	$= (f_{j})_{j} + (f_{j})_{j}$
	$\phi \sim Uniform(0,5)$
	Trunc Normal( $\mathbb{E}(f)$ , $\mathbb{Var}(f)$ , $[0, 5]$ )
	Uniform(-2,7)
	Trunc Normal( $\mathbb{E}(f)$ , $\mathbb{Var}(f)$ , $[-2, 7]$ )
	$Normal(\mathbb{E}(f), \mathbb{Var}(f))$
Continuous Bernoulli( $\pi = 0.3$ )	$= (f_{j})_{j} + (f_{j})_{j}$
0 (	$\phi \sim Uniform(0,1)$
	Trunc Normal( $\mathbb{E}(f), \mathbb{Var}(f), [0, 1]$ )
	Uniform(-2,3)
	Trunc Normal( $\mathbb{E}(f)$ , $\mathbb{Var}(f)$ , $[-2, 3]$ )
	$Normal(\mathbb{E}(f), \mathbb{Var}(f))$
Trunc Normal(1.5, 5, 76, [-6, 6])	$= (f_{j})_{j} + (f_{j})_{j}$
(, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,	$\phi \sim Uniform(-6,6)$
	Trunc Normal( $\mathbb{E}(f)$ , $\mathbb{Var}(f)$ , $[-6, 6]$ )
	Uniform(-8,8)
	Trunc Normal( $\mathbb{E}(f), \mathbb{Var}(f), [-8, 8]$ )
	$Normal(\mathbb{E}(f), \mathbb{V}ar(f))$
	( (0 ) / (0 ) /

Table 1: Target and reference distributions.

Table 1 lists the combinations of target and reference distributions we consider in our experiments. We plot the true target pdfs (red) and the K-series estimates for different number of moments using reference pdfs with the same support as the target in Figure 10. Our method does not suffer from the numerical instability associated with closeness to zero. In most cases, the uniform reference pdf works better on exact support.

In Figure 11, we plot the true four pdfs in Table 1 and their K-series estimates using different number of moments and the uniform reference supported on an interval that contains the support of the target pdf. Specifically, the reference

pdf is supported on the interval that extends by 2 units the true support in either side. The estimation improves significantly as the number of moments increases. The left panels of Figure 12 plot the true pdfs and their K-series estimates using different number of moments and a truncated normal reference supported on the interval that extends by 2 units the true support in both ends. The right panels of Figure 12 plot the true pdfs and their K-series estimates using different number of moments and a normal reference pdf supported on the entire real line.

Visual inspection of these plots indicates that the estimation is better if the support of all reference pdfs is close to the support of the target pdf. The uniform reference distribution results in accurate estimates provided its support is close to the support of the true pdf. On the other hand, both truncated and regular normal reference pdfs lead to accurate K-series estimates the closer the target pdf is to a normal. Moreover, the truncated normal distribution tends to work better on a support wider than the true in comparison with the uniform.

Formal assessment of the estimation accuracy is carried out with Kolmogorov-Smirnov tests. Tables 2, 3 and 4 report the values of the Kolmogorov-Smirnov test statistic comparing the K-series estimates with the true pdfs and whether the null of their equality is rejected for different numbers of moments and reference distributions. The sample size for both the estimated and true distribution is 1000. The critical values are 0.0607 and 0.0479 for significance levels 0.05 and 0.2, respectively.

Target pdf $f$	M	Uniform (Same support)	Trunc Normal (Same support)
$\overline{\text{Trunc Gamma}(\alpha = 2, \beta = 0.5, [0, 5])}$			
	2	0.0172 🖌 !	0.0188 🖌 !
	3	0.0031 🖌	0.0093 🗸
	5	< 1e - 4	0.0033 🗸
	8	$< 1e - 4 \checkmark !$	0.0002 🗸 !
Trunc Normal $(1.5, 5.76, [-6, 6])$			0.0011.11
	2	0.0617 ×	0.0011
	4	0.0122	< 1e - 4
$O_{a} = \frac{1}{2} \left( - 0.2 \right)$	1	0.0002 🔽 !	< 1e - 4 V
Continuous Bernoulli( $\pi = 0.3$ )	9		0.0104
	<u>చ</u>	< 1e - 4	0.0124
	D O	< 1e - 4	
Trung $Fr() = 2/2 [0, 4]$	0	< 1e - 4	< 1e - 4
$\operatorname{Hunc}\operatorname{Exp}(\lambda=2/3,[0,4])$	2	0.0082	0 0010 🗸 🛔
		0.0082	0.0212
	4	10 - 1	0.0023
	0	< 1e - 4 🚩 !	0.0003 🚩 1

 $\checkmark$  Null hypothesis is not rejected at significance level 0.05.

✓! Null hypothesis is not rejected at significance level 0.2.

× Null hypothesis is rejected at significance level 0.05.

 Table 2: Kolmogorov-Smirnov distances and significance test results for reference distribu 

 tions on the same support as the true pdf.



Fig. 10: K-series estimates of the truncated exponential pdf, the truncated gamma pdf, the truncated normal pdf and the continuous Bernoulli with uniform reference (Method of Moments [25]), left panels) and truncated normal (right panels) on exact support.

23



Fig. 11: Approximations of the truncated exponential pdf, the truncated gamma pdf, the truncated normal pdf and the continuous Bernoulli using K-series with uniform reference on the extended support.

Target pdf $f$	M	Uniform (Extended support)
$\overline{\text{Trunc Gamma}(\alpha = 2, \beta = 0.5, [0, 5])}$		
	4	0.0213 🖌 !
	8	0.0186 🖌 !
	10	0.0152 🖌 !
Trunc Normal $(1.5, 5.76, [-6, 6])$		
	5	0.0099 🖌 !
	7	0.0061 🖌 !
	10	0.0048 🖌 !
Continuous Bernoulli( $\pi = 0.3$ )		
	5	0.2285 🗡
	10	0.0939 🗡
	17	0.0579 🗸
Trunc $\operatorname{Exp}(\lambda = 2/3, [0, 4])$		
	6	0.1099 🗡
	10	0.0713 🗡
	15	0.0546 🖌

Null hypothesis is not rejected at significance level 0.05.

✓! Null hypothesis is not rejected at significance level 0.2.

× Null hypothesis is rejected at significance level 0.05.

 
 Table 3: Kolmogorov-Smirnov distances and significance test results for the uniform reference distribution on extended support.

Target pdf $f$	M	Trunc Normal (Extended support)	Normal (real line)
Trunc Gamma( $\alpha = 2, \beta = 0.5, [0, 5]$ )			
	6	0.0172 🖌 !	0.0202 🖌 !
	8	0.0158 🖌	0.0169
	$1\check{0}$	0.0132 🖌 !	0.0033 🗸 !
Trunc Normal $(1.5, 5.76, [-6, 6])$			
	2	0.0171 🖌 !	0.0182 🖌 !
	$\overline{5}$	0.0071 🖌	0.0095 🖌
	10	0.0044 🖌 !	0.0066 🖌 !
Continuous Bernoulli( $\pi = 0.3$ )			
	5	0.0516 🖌	0.0527 🖌
	Ř	0.0374 🖌	0.0387 🖌 !
	12	0.0340 🖌 !	0.0352 🖌 !
Trunc Exp $(\lambda = 2/3, [0, 4])$			
$\mathbf{F}(\mathbf{r}, \mathbf{r}, \mathbf{r}) = \mathbf{r}(\mathbf{r}, \mathbf{r})$	6	0.0667 🗡	0.0757 🗡
	10	0.0558 🖌	0.0617 🗡
	15	0.0391 🖌 !	0.0524 🖌

Null hypothesis is not rejected at significance level 0.05.
 Null hypothesis is not rejected at significance level 0.2.
 Null hypothesis is rejected at significance level 0.05.

Table 4: Kolmogorov-Smirnov distances and significance test results for truncated normal on extended support and normal reference distributions.



Fig. 12: Approximations of the truncated exponential pdf, the truncated gamma pdf, the truncated normal pdf and the continuous Bernoulli using K-series with truncated normal reference on the extended support (left) and normal reference on the whole real line (Gram-Charlier, right).

## Appendix 2. Kolmogorov-Smirnov and Energy Tests for Equality of Distributions

The Kolmogorov-Smirnov (K-S) test [26] compares two cumulative distribution functions (cdfs). We compute the cdf  $\hat{F}_{\rm KS}$  of the estimated pdf  $\hat{f}_{\rm KS}$ . We also compute the (empirical) cdf  $F_{Sample}$  of the data resulting from sampling the probabilistic program variables. The 2-sample Kolmogorov-Smirnov (K-S) test statistic for testing equality of the population (true) cdfs is

$$D_{\rm KS} = \max\left(|F_{\rm KS}(x) - F_{Sample}(x)|\right),\tag{24}$$

where  $N_1$  and  $N_2$  are the sample sizes from the K-series and empirical cdf, respectively. We reject the equality of the two distributions if

$$D_{\rm KS} > c(\alpha) \sqrt{\frac{N_1 + N_2}{N_1 \cdot N_2}} = \sqrt{-\frac{1}{2} \ln \frac{\alpha}{2}} \sqrt{\frac{N_1 + N_2}{N_1 \cdot N_2}}$$

at significance level  $\alpha$ .

The two-sample E-statistic for testing for equality of multivariate distributions proposed by [38] is the *energy* distance  $e(S_1, S_2)$ , which is defined by

$$e(S_1, S_2) = \frac{N_1 N_2}{N_1 + N_2} \left( 2D_{12} - D_{11} - D_{22} \right),$$

for two samples  $S_1, S_2$  of respective sizes  $N_1, N_2$ , where

$$D_{ij} = \frac{1}{N_i N_j} \sum_{p=1}^{N_i} \sum_{q=1}^{N_j} ||\mathbf{X}_{ip} - \mathbf{X}_{jq}||, \ i, j = 1, 2,$$

 $|| \cdot ||$  denotes the Euclidean norm, and  $\mathbf{X}_{1p}$  denotes the *p*-th and  $\mathbf{X}_{2q}$  the *q*-th (vector-valued) observations in the first and second sample, respectively. The test is implemented by nonparametric bootstrap, an approximate permutation test in the R-package energy [30].

We used the Kolmorov-Smirnov test to compare univariate distributions and the *energy* [38] test for multivariate distributions. We draw 1000 observations from the sampling ("true") and estimated distributions. The critical values are 0.0607 and 0.0479 for significance levels 0.05 and 0.2, respectively. Except for very few instances, when a small number of moments is used in the K-series estimation, our estimate is statistically the same as the true distribution. We also test the agreement of the K-series with the GC estimates. When the true distribution is similar to normal, K-series is statistically indistinguishable from

Gram-Charlier. But when the true distribution is not close to normal, K-series provides a far more accurate estimate than Gram-Charlier.

Problem	Var	M	KS Distance	KS Distance (GC)	Energy test (p-value)
Differential-Drive Robot					
	X	6	0.00069 🖌 !	0.00072 🖌 !	
	Y	6	0.00059 🖌 !	0.00059 🖌 !	
	(X,Y)	48			0.4700
PDP					
	X	2	0.00664 🖌 !	0.00680 🖌 !	
	Y	6	0.00033 🖌 !	0.05190 🖌	
	(X, Y)	48			0.4250
Turning vehicle				0.00100.41	
	X	8	0.00807 🖌 !	0.02109 🗸 !	
	Y	8	0.00494 🖌 !	0.01030 🖌 !	
	(X, Y)	80			0.4150
Turning vehicle					
(small variance)	X	8	0 02614 🖌 !	0 11054 <b>X</b>	
	Y	8	0.00070	0.00169 🖌 !	
	(X, Y)	80	0.00010	0.00100 •	0.5000
Tordon mula madal	,				
Taylor Tule model	i	6	0.00037 🖌 !	0.00037 🖌 !	
2D Robotic Arm					
	X	2	0.00037 🖌 !	0.00037 🖌 !	
	Y	2	0.00048 🖌 !	0.00048 🖌 !	
	(X,Y)	8			0.9650
Rimless Wheel Walker					
	X	2	0.00180 🖌 !	0.00180 🖌 !	
Vasicek model	r	2	0.00074 🖌 !	0.00074 🖌 !	
1D Random Walk			0.00012.0.0		
	x	2	0.03834 🖌 !	0.03834 🖌 !	
2D Random Walk					
	X	2	0.02743 🖌 !	0.02743 🗸 !	
	Y	2	0.02714 🖌 !	0.02714 🖌 !	
	(X, Y)	8			0.4902
Stuttering P					
-	s	2	0.00351 🖌 !	0.00354 🖌 !	

Null hypothesis is not rejected at significance level 0.05.
Null hypothesis is not rejected at significance level 0.2.

× Null hypothesis is rejected at significance level 0.05.

Table 5: Kolmogorov-Smirnov distances for univariate distributions and testing for equality of multivariate distributions.

### Appendix 3. 1D and 2D Random Walk

Panel A in Fig. 13 describes the 1D Random Walk [22], and panel B the 2D Random Walk [22]. For the 1D Random walk we used a truncated normal distribution on (-98, 102) as reference. For the 2D Random walk, we used two independent truncated normal distributions on  $(-100, 100) \times (-100, 100)$  with true means and variances of corresponding marginal pdfs obtained with the algorithm in [2]. The K-series estimator exhibits excellent performance for both 1D and 2D random walks, as can be seen in Fig. 13.



**Fig. 13:** K-series estimates of the pdf of X in 1D Random Walk (A) [22] at iteration t = 100, marginal pdfs for variables X, Y and joint distribution of (X, Y) in 2D Random Walk (B) [22] at the iterations t = 100.

### Appendix 4. Symbolic K-series representation example

Consider the probabilistic loop in Fig 14(A): the target random variable r is modeled as the minimum of random variables x and y. Variable y is uniformly distributed on (0, 20), while x follows a mean-reverting process and is affected by the stochastic shock  $\theta \sim \text{Uniform}(-8, 8)$  at each iteration. We can now use the approach from [2] to estimate moments for arbitrary iterations, and use them to receive the symbolic expression for the pdf of r for the corresponding iteration. Since min $(\cdot, \cdot)$  is a non-polynomial function, we apply [20] to represent min $(\cdot, \cdot)$ as a PCE. The transformation is illustrated in Fig. 14.



**Fig. 14:** (A) Probabilistic loop with non-polynomial assignment, (B) Transformation of the program A using Polynomial Chaos Expansion [20], by replacing the function  $\min(\cdot, \cdot)$  with the polynomial G(x, y).

Once the transformation is carried out, the program in Fig 14 (B) can be handled using the algorithm in [2]. The equations estimating first four moments for each iteration are in the left panel of Fig. 15. We choose the uniform distribution on (0, 20) as the reference pdf. We compute the shifted and scaled Legendre polynomials, substitute the moment equations as functions of t as in (22). In a similar manner, we can derive the symbolic expression of the pdf estimate for any arbitrary iteration t. The right panel of Fig 15 plots the pdf estimate of the random variable r at iteration t = 30.

$$\hat{f}_{30}(r) = 5.165866e - 7*r^4 + 2.561246e - 5*r^3 - 0.001472*r^2 + 0.012320*r + 0.055246e - 5*r^3 - 0.001472*r^2 + 0.0014$$



Fig. 15: Left panel: First four moments expressed symbolically in the number of iterations. Right panel: Comparison bar plot between the histogram of the sampling pdf and the symbolic K-series estimation at t = 30.

### References

- Barthe, G., Katoen, J.P., Silva, A.: Foundations of Probabilistic Programming. Cambridge University Press (2020)
- Bartocci, E., Kovács, L., Stankovic, M.: Automatic generation of moment-based invariants for prob-solvable loops. In: Proc. of ATVA 2019: the 17th International Symposium on Automated Technology for Verification and Analysis. LNCS, vol. 11781. Springer (2019). https://doi.org/10.1007/978-3-030-31784-3
- Bartocci, E., Kovács, L., Stankovic, M.: Mora automatic generation of momentbased invariants. In: Proc. of TACAS 2020: the 26th International Conference on Tools and Algorithms for the Construction and Analysis of Systems. LNCS, vol. 12078, pp. 492–498. Springer (2020). https://doi.org/10.1007/978-3-030-45190-5
- Biswas, P., Bhattacharya, A.K.: Function Reconstruction as a Classical Moment Problem: A Maximum Entropy Approach. Journal of Physics A: Mathematical and Theoretical 43(405003), 1–19 (2010)
- Bouissou, O., Goubault, E., Putot, S., Chakarov, A., Sankaranarayanan, S.: Uncertainty propagation using probabilistic affine forms and concentration of measure inequalities. In: International Conference on Tools and Algorithms for the Construction and Analysis of Systems. pp. 225–243. Springer (2016)
- Cramér, H.: Mathematical Methods of Statistics. Princeton Univ. Press, Princeton, NJ (1957), http://www.jstor.org/stable/j.ctt1bpm9r4
- Dharmani, Bhaveshkumar C: Multivariate generalized gram-charlier series in vector notations. Journal of Mathematical Chemistry 56, 1631–1655 (2018). https://doi.org/10.1007/s10910-018-0878-5, https://doi.org/10.1007/ s10910-018-0878-5
- Dwork, C.: Differential privacy. In: Proc. of ICALP 2006: the 33rd International Colloquium on Automata, Languages and Programming. LNCS, vol. 4052, pp. 1–12. Springer (2006). https://doi.org/10.1007/11787006
- Ernst, Oliver G., Mugler, Antje, Starkloff, Hans-Jörg, Ullmann, Elisabeth: On the convergence of generalized polynomial chaos expansions. ESAIM: M2AN 46(2), 317–339 (2012). https://doi.org/10.1051/m2an/2011045, https://doi.org/10. 1051/m2an/2011045
- Filipović, D., Mayerhofer, E., Schneider, P.: Density approximations for multivariate affine jump-diffusion processes. Journal of Econometrics 176(2), 93-111 (2013). https://doi.org/https://doi.org/10.1016/j.jeconom.2012.12.003, https:// www.sciencedirect.com/science/article/pii/S0304407613000596
- Fremont, D.J., Dreossi, T., Ghosh, S., Yue, X., Sangiovanni-Vincentelli, A.L., Seshia, S.A.: Scenic: a language for scenario specification and scene generation. In: Proc. of PLDI 2019: the 40th ACM SIGPLAN Conference on Programming Language Design and Implementation. pp. 63–78. ACM (2019). https://doi.org/10.1145/3314221
- Gehr, T., Misailovic, S., Vechev, M.T.: PSI: exact symbolic inference for probabilistic programs. In: Proc. of CAV 2016: the 28th International Conference on Computer Aided Verification. LNCS, vol. 9779, pp. 62–83. Springer (2016). https://doi.org/10.1007/978-3-319-41528-4\_4
- Ghahramani, Z.: Probabilistic machine learning and artificial intelligence. Nature 521(7553), 452–459 (2015). https://doi.org/10.1038/nature14541, https://doi.org/10.1038/nature14541
- 14. Hald, A.: The Early History of the Cumulants and the Gram-Charlier Series. International Statistical Review **68**(2), 137–153 (2000)

- Herman, T.: Probabilistic self-stabilization. Inf. Process. Lett. 35(2), 63–67 (1990). https://doi.org/10.1016/0020-0190(90)90107-9
- Innocentini, G.C.P., Hodgkinson, A., Radulescu, O.: Time dependent stochastic mrna and protein synthesis in piecewise-deterministic models of gene networks. Frontiers in Physics 6 (2018). https://doi.org/10.3389/fphy.2018.00046, https:// www.frontiersin.org/articles/10.3389/fphy.2018.00046
- Jasour, A., Wang, A., Williams, B.C.: Moment-based exact uncertainty propagation through nonlinear stochastic autonomous systems. CoRR abs/2101.12490 (2021), https://arxiv.org/abs/2101.12490
- Karimi, A., Moosbrugger, M., Stankovič, M., Kovács, L., Bartocci, E., Bura, E.: Distribution estimation for probabilistic loops. In: Ábrahám, E., Paolieri, M. (eds.) Quantitative Evaluation of Systems. pp. 26–42. Springer International Publishing, Cham (2022)
- Kendall, M., Stuart, A.: The Advanced Theory of Statistics. Volume 1: Distribution Theory. Macmillan, New York, NY (1977)
- Kofnov, A., Moosbrugger, M., Stankovič, M., Bartocci, E., Bura, E.: Moment-based invariants for probabilistic loops with non-polynomial assignments. In: Ábrahám, E., Paolieri, M. (eds.) Quantitative Evaluation of Systems. pp. 3–25. Springer International Publishing, Cham (2022)
- Kolmogorov, A., Fomin, S.: Elements of the Theory of Functions and Functional Analysis. Nauka, Moscow, 4 edn. (1976)
- 22. Kura, S., Urabe, N., Hasuo, I.: Tail probabilities for randomized program runtimes via martingales for higher moments. In: Vojnar, T., Zhang, L. (eds.) Tools and Algorithms for the Construction and Analysis of Systems. pp. 135–153. Springer International Publishing, Cham (2019)
- Lebaz, N., Cockx, A., Spérandio, M., Morchain, J.: Reconstruction of a Distribution From a Finite Number of Its Moments: A Comparative Study in the Case of Depolymerization Process. Computers & Chemical Engineering 84, 326–337 (2016)
- Moosbrugger, M., Stankovič, M., Bartocci, E., Kovács, L.: This is the moment for probabilistic loops. Proc. ACM Program. Lang. 6(OOPSLA2) (oct 2022). https://doi.org/10.1145/3563341, https://doi.org/10.1145/3563341
- Munkhammar, J., Mattsson, L., Rydén, J.: Polynomial probability distribution estimation using the method of moments. PLOS ONE 12(4), 1-14 (04 2017). https://doi.org/10.1371/journal.pone.0174573, https://doi.org/10.1371/journal.pone.0174573
- Myles Hollander, Douglas A. Wolfe, E.C.: Nonparametric Statistical Methods. John Wiley & Sons, New York-London-Sydney, 3 edn. (2013)
- 27. Provost, S.B., Tae Ha, H.: On the inversion of certain moment matrices. Linear Algebra and its Applications 430(10), 2650–2658 (2009). https://doi.org/https://doi.org/10.1016/j.laa.2008.10.024, https://www.sciencedirect.com/science/article/pii/S0024379508005223, special Issue devoted to selected papers presented at the 16th International Workshop on Matrices and Statistics (IWMS-2007)
- 28. Rahman, S.: A polynomial chaos expansion in dependent random variables. Journal of Mathematical Analysis and Applications 464(1), 749-775 (2018). https://doi.org/https://doi.org/10.1016/j.jmaa.2018.04.032, https://www.sciencedirect.com/science/article/pii/S0022247X18303305
- Rajeev Motwani, P.R.: Randomized Algorithms. Cambridge University Press (1995)

- 34 Kofnov et al.
- Rizzo, M., Szekely, G.: energy: E-Statistics: Multivariate Inference via the Energy of Data (2022), https://CRAN.R-project.org/package=energy, r package version 1.7-11
- Rudin, W.: Principles of mathematical analysis. McGraw-Hill New York, 3d ed. edn. (1976), http://www.loc.gov/catdir/toc/mh031/75017903.html
- Rudin, W.: Real and Complex Analysis. McGraw-Hill Science/Engineering/Math (May 1986)
- 33. Sankaranarayanan, S., Chou, Y., Goubault, E., Putot, S.: Reasoning about uncertainties in discrete-time dynamical systems using polynomial forms. In: Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M.F., Lin, H. (eds.) Advances in Neural Information Processing Systems. vol. 33, pp. 17502-17513. Curran Associates, Inc. (2020), https://proceedings.neurips.cc/paper/2020/ file/ca886eb9edb61a42256192745c72cd79-Paper.pdf
- Silverman, B.: Density Estimation for Statistics and Data Analysis. Routledge, 1 edn. (1998). https://doi.org/https://doi.org/10.1201/9781315140919
- Steinhardt, J., Tedrake, R.: Finite-time regional verification of stochastic non-linear systems. The International Journal of Robotics Research 31(7), 901–923 (2012)
- 36. Steinhardt, J., Tedrake, R.: Finite-time regional verification of stochastic non-linear systems. Int. J. Robotics Res. **31**(7), 901–923 (2012). https://doi.org/10.1177/0278364912444146
- P.J.: Α 37. Szabłowski, few remarks on orthogonal polynomials. Applied Mathematics and Computation **252**, 215 - 228(2015).https://doi.org/https://doi.org/10.1016/j.amc.2014.11.112, https://www. sciencedirect.com/science/article/pii/S0096300314016452
- Székely, G.J., Rizzo, M.L.: Testing for equal distributions in high dimension. InterStat, vol. 5 (November 2004)
- 39. Taylor, J.B.: Discretion versus policy rules in practice. Carnegie-Rochester Conference Series on Public Policy 39(1), 195-214 (December 1993), https://ideas.repec.org/a/eee/crcspp/v39y1993ip195-214.html
- Tekel, J., Cohen, L.: Constructing and estimating probability distributions from moments. In: Sadjadi, F.A., Mahalanobis, A. (eds.) Automatic Target Recognition XXII. vol. 8391, p. 83910E. International Society for Optics and Photonics, SPIE (2012). https://doi.org/10.1117/12.919443, https://doi.org/10.1117/12. 919443
- 41. Vasicek, O.: An equilibrium characterization of the term struc-Journal of Financial Economics 5(2),177 - 188(1977).ture. https://doi.org/https://doi.org/10.1016/0304-405X(77)90016-2, https: //www.sciencedirect.com/science/article/pii/0304405X77900162
- 42. Xiu, D.: Numerical Methods for Stochastic Computations: A Spectral Method Approach. Princeton University Press (2010), http://www.jstor.org/stable/j.ctv7h0skv
- Xiu, D., Karniadakis, G.: The Wiener-Askey polynomial chaos for stochastic differential equations. SIAM J. Sci. Comput. 24(2), 619–644 (Feb 2002). https://doi.org/10.1137/S1064827501387826