

The q -Onsager algebra and the quantum torus

Owen Goff

Abstract

The q -Onsager algebra, denoted O_q , is defined by two generators W_0, W_1 and two relations called the q -Dolan-Grady relations. Recently, Terwilliger introduced some elements of O_q , said to be alternating. These elements are denoted

$$\{W_{-k}\}_{k=0}^{\infty}, \quad \{W_{k+1}\}_{k=0}^{\infty}, \quad \{G_{k+1}\}_{k=0}^{\infty}, \quad \{\tilde{G}_{k+1}\}_{k=0}^{\infty}.$$

The alternating elements of O_q are defined recursively. By construction, they are polynomials in W_0 and W_1 . It is currently unknown how to express these polynomials in closed form.

In this paper, we consider an algebra T_q , called the quantum torus. We present a basis for T_q and define an algebra homomorphism $p : O_q \mapsto T_q$. In our main result, we express the p -images of the alternating elements of O_q in the basis for T_q . These expressions are in a closed form that we find attractive.

Keywords. q -Onsager algebra; q -Dolan-Grady relations; quantum torus; alternating central extension.

2020 Mathematics Subject Classification. Primary: 17B37. Secondary: 05E10, 16T20.

1 Introduction

The q -Onsager algebra, which we denote O_q , first appeared in the topic of algebraic graph theory [12, Lemma 5.4]. Applications of O_q in algebraic combinatorics appear in many papers, including [10, 11, 13]. The algebra O_q has recently been applied to statistical mechanics [1–3, 5, 7] and quantum groups [6, Section 2.2]. The algebra O_q is defined by generators W_0, W_1 and two relations called the q -Dolan-Grady relations.

We mentioned that O_q appears in statistical mechanics; let us expand on that. In [5, eqn. 4], Pascal Baseilhac and Kozo Koizumi described an algebra similar to O_q while researching operators on boundary integrable systems with hidden

symmetries. Following [16], we denote this algebra by \mathcal{O}_q . This algebra is defined by generators and relations. The generators are

$$\{\mathcal{W}_{-k}\}_{k=0}^{\infty}, \quad \{\mathcal{W}_{k+1}\}_{k=0}^{\infty}, \quad \{\mathcal{G}_{k+1}\}_{k=0}^{\infty}, \quad \{\tilde{\mathcal{G}}_{k+1}\}_{k=0}^{\infty}. \quad (1)$$

The relations are given in [7, Definition 3.1], as well as Definition 3.5 below. The name and notation for \mathcal{O}_q have evolved over time; it is currently known as the alternating central extension of the q -Onsager algebra, as in [15]. The elements displayed in equation (1) are called the alternating generators of \mathcal{O}_q ; see [15] and [16].

Motivated by [5], Paul Terwilliger showed in [15] that there exists an algebra isomorphism from \mathcal{O}_q to the tensor product of O_q and the polynomial algebra over countably many commuting variables. We denote this isomorphism by φ , and we denote the variables by $\{z_i\}_{i=1}^{\infty}$. For $i \geq 1$, let $\mathcal{Z}_i \in \mathcal{O}_q$ denote the φ -preimage of $1 \otimes z_i$. By [16, Definition 11.2], there exists an algebra homomorphism $\gamma : \mathcal{O}_q \mapsto O_q$ that sends $\mathcal{W}_0 \mapsto W_0$, $\mathcal{W}_1 \mapsto W_1$, and $\mathcal{Z}_i \mapsto 0$ for $i \geq 1$.

The γ -images of the alternating generators of \mathcal{O}_q are called the alternating elements of O_q . By construction, each alternating element of O_q is a polynomial in W_0 and W_1 . Recursive formulas for these polynomials are known; see [15, Lemma 8.22]. However, closed forms for the polynomials are not currently known.

In this paper, we discuss an algebra T_q , called the quantum torus [9]. The algebra T_q is defined by the generators x, y, x^{-1}, y^{-1} and the relations

$$xx^{-1} = 1 = x^{-1}x, \quad yy^{-1} = 1 = y^{-1}y, \quad xy = q^2yx.$$

It is known that the elements $\{x^i y^j \mid i, j \in \mathbb{Z}\}$ form a basis for T_q ; see [9, p. 3].

In this paper, we introduce the algebra homomorphism $p : \mathcal{O}_q \rightarrow T_q$ that sends $\mathcal{W}_0 \mapsto x + x^{-1}$ and $\mathcal{W}_1 \mapsto y + y^{-1}$. We consider the p -images of the alternating elements of \mathcal{O}_q . Our main objective is to express these p -images in closed form in the above basis. To reach our objective, we make use of generating functions. We give two versions of our main result: one expressed using generating functions and one without them. The main results of this paper are Theorem 8.1 and Theorem 9.1.

The paper is organized as follows. In Sections 3–5, we give definitions and results concerning \mathcal{O}_q , O_q , and T_q . Section 6 covers generating functions, and in Section 7 we provide notation and formulas. In Section 8 we state and prove our main result in terms of generating functions. In Section 9 we present our main result without using generating functions.

2 Notations

In this section, we introduce some notation that will be used throughout the paper.

- Let K be a field of characteristic 0.
- All algebras in this paper are associative, unital, and over K .
- For u, v in any algebra, define $[u, v] = uv - vu$. We call $[u, v]$ the *commutator* of u and v . For nonzero $r \in K$, define $[u, v]_r = ruv - r^{-1}vu$. We call $[u, v]_r$ the *r -commutator* of u and v .
- Fix nonzero $q \in K$.
- Let \mathbb{N} denote the set of natural numbers $\{0, 1, 2, \dots\}$. Let \mathbb{Z} denote the set of integers. Let \mathbb{C} denote the set of complex numbers.
- For $n \in \mathbb{N}$, define $[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}$. We are using the notation from [14].
- By an *automorphism* of an algebra \mathcal{A} , we mean an algebra isomorphism from \mathcal{A} to itself.
- The *opposite algebra* of an algebra \mathcal{A} , denoted \mathcal{A}^{op} , is the algebra on the vector space \mathcal{A} such that for $a, b \in \mathcal{A}^{op}$, ab (in \mathcal{A}^{op}) = ba (in \mathcal{A}). Note that if \mathcal{A} is commutative, then $\mathcal{A} = \mathcal{A}^{op}$.
- By an *antiautomorphism* of an algebra \mathcal{A} , we mean an algebra isomorphism from \mathcal{A} to \mathcal{A}^{op} . In other words, an antiautomorphism of \mathcal{A} is a K -linear bijection $\alpha : \mathcal{A} \rightarrow \mathcal{A}$ that reverses the order of multiplication: $\alpha(ab) = \alpha(b)\alpha(a)$ for all $a, b \in \mathcal{A}$.

3 The q -Onsager algebra

In this section, we consider the q -Onsager algebra and its alternating central extension.

Definition 3.1. Let O_q denote the algebra defined by generators W_0, W_1 and relations

$$[W_0, [W_0, [W_0, W_1]_q]_{q^{-1}}] = -(q^2 - q^{-2})^2 [W_0, W_1], \quad (2)$$

$$[W_1, [W_1, [W_1, W_0]_q]_{q^{-1}}] = -(q^2 - q^{-2})^2 [W_1, W_0]. \quad (3)$$

We call O_q the *q -Onsager algebra*.

Equations (2) and (3) are called the *q -Dolan-Grady relations*. We define an automorphism and an antiautomorphism of O_q .

Lemma 3.2. (See [16, Lemma 3.3].) *There exists an automorphism σ of O_q that sends*

$$W_0 \mapsto W_1, \quad W_1 \mapsto W_0.$$

Lemma 3.3. (See [16, Lemma 3.4].) *There exists an antiautomorphism \dagger of O_q that sends*

$$W_0 \mapsto W_0, \quad W_1 \mapsto W_1.$$

Lemma 3.4. *With reference to Lemmas 3.2 and 3.3, we have $\sigma \circ \dagger = \dagger \circ \sigma$ in O_q . That is to say, this diagram commutes:*

$$\begin{array}{ccc} O_q & \xrightarrow{\dagger} & O_q \\ \sigma \downarrow & & \downarrow \sigma \\ O_q & \xrightarrow{\dagger} & O_q \end{array}$$

Proof. Chase W_0 and W_1 around the diagram. □

We recall the alternating central extension of O_q .

Definition 3.5. (See [7, Definition 3.1]) Define the algebra \mathcal{O}_q by the generators

$$\{\mathcal{W}_{-k}\}_{k=0}^\infty, \quad \{\mathcal{W}_{k+1}\}_{k=0}^\infty, \quad \{\mathcal{G}_{k+1}\}_{k=0}^\infty, \quad \{\tilde{\mathcal{G}}_{k+1}\}_{k=0}^\infty, \quad (4)$$

and the following relations. For $k, \ell \in \mathbb{N}$,

$$[\mathcal{W}_0, \mathcal{W}_{k+1}] = [\mathcal{W}_{-k}, \mathcal{W}_1] = (\tilde{\mathcal{G}}_{k+1} - \mathcal{G}_{k+1})/(q + q^{-1}), \quad (5)$$

$$[\mathcal{W}_0, \mathcal{G}_{k+1}]_q = [\tilde{\mathcal{G}}_{k+1}, \mathcal{W}_0]_q = \rho\mathcal{W}_{-k-1} - \rho\mathcal{W}_{k+1}, \quad (6)$$

$$[\mathcal{G}_{k+1}, \mathcal{W}_1]_q = [\mathcal{W}_1, \tilde{\mathcal{G}}_{k+1}]_q = \rho\mathcal{W}_{k+2} - \rho\mathcal{W}_{-k}, \quad (7)$$

$$[\mathcal{W}_{-k}, \mathcal{W}_{-\ell}] = 0, \quad (8)$$

$$[\mathcal{W}_{k+1}, \mathcal{W}_{\ell+1}] = 0, \quad (9)$$

$$[\mathcal{W}_{-k}, \mathcal{W}_{\ell+1}] + [\mathcal{W}_{k+1}, \mathcal{W}_{-\ell}] = 0, \quad (10)$$

$$[\mathcal{W}_{-k}, \mathcal{G}_{\ell+1}] + [\mathcal{G}_{k+1}, \mathcal{W}_{-\ell}] = 0, \quad (11)$$

$$[\mathcal{W}_{-k}, \tilde{\mathcal{G}}_{\ell+1}] + [\tilde{\mathcal{G}}_{k+1}, \mathcal{W}_{-\ell}] = 0, \quad (12)$$

$$[\mathcal{W}_{k+1}, \mathcal{G}_{\ell+1}] + [\mathcal{G}_{k+1}, \mathcal{W}_{\ell+1}] = 0, \quad (13)$$

$$[\mathcal{W}_{k+1}, \tilde{\mathcal{G}}_{\ell+1}] + [\tilde{\mathcal{G}}_{k+1}, \mathcal{W}_{\ell+1}] = 0, \quad (14)$$

$$[\mathcal{G}_{k+1}, \mathcal{G}_{\ell+1}] = 0, \quad (15)$$

$$[\tilde{\mathcal{G}}_{k+1}, \tilde{\mathcal{G}}_{\ell+1}] = 0, \quad (16)$$

$$[\tilde{\mathcal{G}}_{k+1}, \mathcal{G}_{\ell+1}] + [\mathcal{G}_{k+1}, \tilde{\mathcal{G}}_{\ell+1}] = 0. \quad (17)$$

In these equations, $\rho = -(q^2 - q^{-2})^2$.

The algebra \mathcal{O}_q is called the *alternating central extension* of O_q . The elements in (4) are called the *alternating generators* of \mathcal{O}_q .

For notational convenience, we define

$$\mathcal{G}_0 = -(q - q^{-1})(q + q^{-1})^2, \quad \tilde{\mathcal{G}}_0 = -(q - q^{-1})(q + q^{-1})^2.$$

For the remainder of this section, we explain how O_q and \mathcal{O}_q are related.

Proposition 3.6. (See [3, Section 3].) *In \mathcal{O}_q , we have*

$$\begin{aligned} [\mathcal{W}_0, [\mathcal{W}_0, [\mathcal{W}_0, \mathcal{W}_1]_q]_{q^{-1}}] &= -(q^2 - q^{-2})^2 [\mathcal{W}_0, \mathcal{W}_1], \\ [\mathcal{W}_1, [\mathcal{W}_1, [\mathcal{W}_1, \mathcal{W}_0]_q]_{q^{-1}}] &= -(q^2 - q^{-2})^2 [\mathcal{W}_1, \mathcal{W}_0]. \end{aligned}$$

We have defined an automorphism and an antiautomorphism of \mathcal{O}_q . We define analogous maps for \mathcal{O}_q .

Lemma 3.7. (See [15, Lemma 3.1].) *There exists an automorphism σ of \mathcal{O}_q that sends*

$$\mathcal{W}_{-k} \mapsto \mathcal{W}_{k+1}, \quad \mathcal{W}_{k+1} \mapsto \mathcal{W}_{-k}, \quad \mathcal{G}_{k+1} \mapsto \tilde{\mathcal{G}}_{k+1}, \quad \tilde{\mathcal{G}}_{k+1} \mapsto \mathcal{G}_{k+1}$$

for all $k \in \mathbb{N}$.

Lemma 3.8. (See [15, Lemma 3.2].) *There exists an antiautomorphism \dagger of \mathcal{O}_q that sends*

$$\mathcal{W}_{-k} \mapsto \mathcal{W}_{-k}, \quad \mathcal{W}_{k+1} \mapsto \mathcal{W}_{k+1}, \quad \mathcal{G}_{k+1} \mapsto \tilde{\mathcal{G}}_{k+1}, \quad \tilde{\mathcal{G}}_{k+1} \mapsto \mathcal{G}_{k+1}$$

for all $k \in \mathbb{N}$.

Lemma 3.9. *This diagram commutes:*

$$\begin{array}{ccc} \mathcal{O}_q & \xrightarrow{\dagger} & \mathcal{O}_q \\ \sigma \downarrow & & \downarrow \sigma \\ \mathcal{O}_q & \xrightarrow{\dagger} & \mathcal{O}_q \end{array}$$

Proof. Chase the alternating generators of \mathcal{O}_q around the diagram. □

We bring in some notation. Let $\{z_i\}_{i=1}^\infty$ be mutually commuting indeterminates. Let $K[z_1, z_2, \dots]$ denote the algebra of polynomials in $\{z_i\}_{i=1}^\infty$ that have coefficients in K .

Proposition 3.10. *There exists an algebra isomorphism*

$$\phi : \mathcal{O}_q \otimes K[z_1, z_2, \dots] \mapsto \mathcal{O}_q$$

such that

$$\phi(W_0 \otimes 1) = \mathcal{W}_0, \quad \phi(W_1 \otimes 1) = \mathcal{W}_1,$$

and these diagrams commute:

$$\begin{array}{ccc} \mathcal{O}_q \otimes K[z_1, z_2, \dots] & \xrightarrow{\phi} & \mathcal{O}_q \\ \sigma \otimes id \downarrow & & \downarrow \sigma \\ \mathcal{O}_q \otimes K[z_1, z_2, \dots] & \xrightarrow{\phi} & \mathcal{O}_q \end{array} \quad \begin{array}{ccc} \mathcal{O}_q \otimes K[z_1, z_2, \dots] & \xrightarrow{\phi} & \mathcal{O}_q \\ \dagger \otimes id \downarrow & & \downarrow \dagger \\ \mathcal{O}_q \otimes K[z_1, z_2, \dots] & \xrightarrow{\phi} & \mathcal{O}_q \end{array}$$

Proof. The map ϕ is the isomorphism defined in [16, Proposition 8.8]. The diagrams commute by [16, Proposition 8.9]. \square

Definition 3.11. For $i \geq 1$, define $\mathcal{Z}_i = \phi(1 \otimes z_i)$. For notational convenience, define $\mathcal{Z}_0 = (q + q^{-1})^2$.

Lemma 3.12. *The elements*

$$\mathcal{W}_0, \quad \mathcal{W}_1, \quad \{\mathcal{Z}_i\}_{i=1}^{\infty}$$

generate \mathcal{O}_q . Moreover, the elements $\{\mathcal{Z}_i\}_{i=1}^{\infty}$ are central in \mathcal{O}_q .

Proof. By Proposition 3.10 and since $1 \otimes z_i$ is central in $O_q \otimes K[z_1, z_2, \dots]$ for $i \geq 1$. \square

Definition 3.13. Define an algebra homomorphism $\epsilon : K[z_1, z_2, \dots] \mapsto K$ that sends $z_i \mapsto 0$ for $i \geq 1$.

Definition 3.14. (See [16, Definition 11.2].) Let $\gamma : \mathcal{O}_q \mapsto O_q$ denote the composition

$$\gamma : \mathcal{O}_q \xrightarrow{\phi^{-1}} O_q \otimes K[z_1, z_2, \dots] \xrightarrow{id \otimes \epsilon} O_q \otimes K \xrightarrow{x \otimes 1 \mapsto x} O_q,$$

where ϕ is from Proposition 3.10.

Note that γ is an algebra homomorphism.

Lemma 3.15. *The map γ sends*

$$\mathcal{W}_0 \mapsto W_0, \quad \mathcal{W}_1 \mapsto W_1, \quad \mathcal{Z}_i \mapsto 0, \quad i \geq 1.$$

Proof. Routine consequence of Definition 3.14. \square

Definition 3.16. (See [16, Definition 11.5].) For $k \in \mathbb{N}$, define

$$W_{-k} = \gamma(\mathcal{W}_{-k}), \quad W_{k+1} = \gamma(\mathcal{W}_{k+1}), \quad G_{k+1} = \gamma(\mathcal{G}_{k+1}), \quad \tilde{G}_{k+1} = \gamma(\tilde{\mathcal{G}}_{k+1}).$$

We call these elements the *alternating elements* of O_q . For notational convenience, we define

$$G_0 = - (q - q^{-1}) (q + q^{-1})^2, \quad \tilde{G}_0 = - (q - q^{-1}) (q + q^{-1})^2.$$

Lemma 3.17. *These diagrams commute:*

$$\begin{array}{ccc} \mathcal{O}_q & \xrightarrow{\sigma} & \mathcal{O}_q & & \mathcal{O}_q & \xrightarrow{\dagger} & \mathcal{O}_q \\ \gamma \downarrow & & \downarrow \gamma & & \gamma \downarrow & & \downarrow \gamma \\ O_q & \xrightarrow{\sigma} & O_q & & O_q & \xrightarrow{\dagger} & O_q \end{array}$$

Proof. Chase \mathcal{W}_0 , \mathcal{W}_1 , and \mathcal{Z}_i around each diagram. \square

Corollary 3.18. *Pick $k \in \mathbb{N}$. The automorphism σ of O_q sends*

$$W_{-k} \mapsto W_{k+1}, \quad W_{k+1} \mapsto W_{-k}, \quad G_{k+1} \mapsto \tilde{G}_{k+1}, \quad \tilde{G}_{k+1} \mapsto G_{k+1}.$$

The antiautomorphism \dagger of O_q sends

$$W_{-k} \mapsto W_{-k}, \quad W_{k+1} \mapsto W_{k+1}, \quad G_{k+1} \mapsto \tilde{G}_{k+1}, \quad \tilde{G}_{k+1} \mapsto G_{k+1}.$$

Proof. Chase \mathcal{W}_{-k} , \mathcal{W}_{k+1} , \mathcal{G}_{k+1} , and $\tilde{\mathcal{G}}_{k+1}$ around the diagrams in Lemma 3.17. \square

4 The quantum torus

In this section we consider an algebra called the quantum torus, denoted T_q . We review some properties of T_q and display a basis for the vector space T_q . Then we introduce an algebra homomorphism $p : O_q \mapsto T_q$ and consider the p -images of the alternating elements of O_q .

Definition 4.1. (See [9].) Define the algebra T_q by generators

$$x, y, x^{-1}, y^{-1}$$

and relations

$$xx^{-1} = 1 = x^{-1}x, \quad yy^{-1} = 1 = y^{-1}y, \quad xy = q^2yx. \quad (18)$$

The algebra T_q is called the *quantum torus*.

Lemma 4.2. *The following relations hold in T_q :*

$$\begin{aligned} xy &= q^2yx, \\ x^{-1}y &= q^{-2}yx^{-1}, \\ x^{-1}y^{-1} &= q^2y^{-1}x^{-1}, \\ xy^{-1} &= q^{-2}y^{-1}x. \end{aligned}$$

Proof. Routine consequence of the relations in (18). □

We display a basis for the vector space T_q .

Lemma 4.3. (See [9, p. 3].) *The vector space T_q has a basis consisting of $\{x^a y^b \mid a, b \in \mathbb{Z}\}$.*

Definition 4.4. For the algebra T_q , define

$$w_0 = x + x^{-1}, \quad w_1 = y + y^{-1}. \quad (19)$$

We describe how w_0 and w_1 are related.

Lemma 4.5. *In the algebra T_q , we have*

$$[w_0, [w_0, w_1]_q]_{q^{-1}} = -(q^2 - q^{-2})^2 w_1, \quad (20)$$

$$[w_1, [w_1, w_0]_q]_{q^{-1}} = -(q^2 - q^{-2})^2 w_0. \quad (21)$$

Proof. Write (20) and (21) in terms of x and y using (19) and simplify the result using Lemma 4.2. □

Remark. A variation of Definition 4.4 and Lemma 4.5 can be found in [4, eqn. (30) and above eqn. (32)].

Corollary 4.6. *In the algebra T_q , we have*

$$[w_0, [w_0, [w_0, w_1]_q]_{q^{-1}}] = -(q^2 - q^{-2})^2 [w_0, w_1], \quad (22)$$

$$[w_1, [w_1, [w_1, w_0]_q]_{q^{-1}}] = -(q^2 - q^{-2})^2 [w_1, w_0]. \quad (23)$$

Proof. To obtain equation (22), take the commutator of w_0 with both sides of equation (20). To obtain equation (23), take the commutator of w_1 with both sides of equation (21). \square

Proposition 4.7. *There exists an algebra homomorphism $p : O_q \mapsto T_q$ that sends $W_0 \mapsto w_0$ and $W_1 \mapsto w_1$.*

Proof. The relations in Corollary 4.6 are the q -Dolan-Grady relations. \square

Definition 4.8. For $k \in \mathbb{N}$, we define the following elements of T_q :

$$w_{-k} = p(W_{-k}), \quad w_{k+1} = p(W_{k+1}), \quad g_{k+1} = p(G_{k+1}), \quad \tilde{g}_{k+1} = p(\tilde{G}_{k+1}). \quad (24)$$

The elements in (24) are called the *alternating elements* of T_q .

For notational convenience, we define

$$g_0 = -(q - q^{-1})(q + q^{-1})^2, \quad \tilde{g}_0 = -(q - q^{-1})(q + q^{-1})^2.$$

Since O_q is generated by W_0 and W_1 , the alternating elements of O_q are polynomials in W_0 and W_1 . However, only recursive formulas for these polynomials are known. The situation is different for T_q . We will express the alternating elements of T_q in the basis for T_q given in Lemma 4.3. This will provide attractive closed forms for the alternating elements of T_q .

5 An automorphism and antiautomorphism of T_q

In this section we define an automorphism and an antiautomorphism of T_q .

Lemma 5.1. *There exists an automorphism τ of T_q that sends*

$$x \mapsto y^{-1}, \quad y \mapsto x, \quad x^{-1} \mapsto y, \quad y^{-1} \mapsto x^{-1}.$$

Proof. Define

$$x_\tau = y^{-1}, \quad y_\tau = x, \quad (x^{-1})_\tau = y, \quad (y^{-1})_\tau = x^{-1}.$$

One routinely checks that x_τ , y_τ , $(x^{-1})_\tau$, and $(y^{-1})_\tau$ satisfy the defining relations in (18).

Consequently, there is an algebra homomorphism $\tau : T_q \rightarrow T_q$ that sends

$$x \mapsto x_\tau, \quad y \mapsto y_\tau, \quad x^{-1} \mapsto (x^{-1})_\tau, \quad y^{-1} \mapsto (y^{-1})_\tau.$$

Note that τ^4 fixes x , y , x^{-1} , and y^{-1} . Hence, τ^4 is the identity map, so the map τ is invertible. Thus τ is an automorphism of T_q . \square

Lemma 5.2. *There exists an antiautomorphism \ddagger of T_q that sends*

$$x \mapsto x^{-1}, \quad y \mapsto y, \quad x^{-1} \mapsto x, \quad y^{-1} \mapsto y^{-1}.$$

Proof. Define

$$x_{\ddagger} = x^{-1}, \quad y_{\ddagger} = y, \quad (x^{-1})_{\ddagger} = x, \quad (y^{-1})_{\ddagger} = y^{-1}.$$

The opposite algebra T_q^{op} is explained in Section 2. By this explanation and (18), the defining relations of T_q^{op} are

$$xx^{-1} = 1 = x^{-1}x, \quad yy^{-1} = 1 = y^{-1}y, \quad yx = q^2xy.$$

These relations are satisfied by $x_{\ddagger}, y_{\ddagger}, x_{\ddagger}^{-1}, y_{\ddagger}^{-1}$.

Therefore, there is an algebra homomorphism $\ddagger : T_q \rightarrow T_q^{\text{op}}$ that sends

$$x \mapsto x_{\ddagger}, \quad y \mapsto y_{\ddagger}, \quad x^{-1} \mapsto (x^{-1})_{\ddagger}, \quad y^{-1} \mapsto (y^{-1})_{\ddagger}.$$

Note that \ddagger^2 fixes x, y, x^{-1} , and y^{-1} , so \ddagger^2 is the identity map of T_q . It follows that \ddagger is a bijection and hence an antiautomorphism of T_q . □

Lemma 5.3. *These diagrams commute:*

$$\begin{array}{ccc} O_q & \xrightarrow{\sigma} & O_q \\ p \downarrow & & \downarrow p \\ T_q & \xrightarrow{\tau} & T_q \end{array} \qquad \begin{array}{ccc} O_q & \xrightarrow{\ddagger} & O_q \\ p \downarrow & & \downarrow p \\ T_q & \xrightarrow{\ddagger} & T_q \end{array}$$

Proof. Chase W_0 and W_1 around each diagram. □

6 Some generating functions

Our goal for the paper is to write the alternating elements of T_q in the basis from Lemma 4.3. We do so using generating functions. In this section we define these generating functions and discuss their properties.

Definition 6.1. (See [16, Definition 7.1].) For the algebra \mathcal{O}_q , define the generating functions

$$\begin{aligned} \mathcal{W}^-(t) &= \sum_{i=0}^{\infty} \mathcal{W}_{-i} t^i, & \mathcal{W}^+(t) &= \sum_{i=0}^{\infty} \mathcal{W}_{i+1} t^i, \\ \mathcal{G}(t) &= \sum_{i=0}^{\infty} \mathcal{G}_i t^i, & \tilde{\mathcal{G}}(t) &= \sum_{i=0}^{\infty} \tilde{\mathcal{G}}_i t^i. \end{aligned} \tag{25}$$

Definition 6.2. (See [16, Definition 12.1].) For the algebra O_q , define the generating functions

$$\begin{aligned} W^-(t) &= \sum_{i=0}^{\infty} W_{-i}t^i, & W^+(t) &= \sum_{i=0}^{\infty} W_{i+1}t^i, \\ G(t) &= \sum_{i=0}^{\infty} G_i t^i, & \tilde{G}(t) &= \sum_{i=0}^{\infty} \tilde{G}_i t^i. \end{aligned} \tag{26}$$

Definition 6.3. For the algebra T_q , define the generating functions

$$\begin{aligned} w^-(t) &= \sum_{i=0}^{\infty} w_{-i}t^i, & w^+(t) &= \sum_{i=0}^{\infty} w_{i+1}t^i, \\ g(t) &= \sum_{i=0}^{\infty} g_i t^i, & \tilde{g}(t) &= \sum_{i=0}^{\infty} \tilde{g}_i t^i. \end{aligned} \tag{27}$$

Recall the algebra homomorphism γ from Definition 3.14.

Lemma 6.4. *The map γ sends the generating functions in (25) to the generating functions in (26) as follows.*

<i>argument in O_q</i>	$\mathcal{W}^-(t)$	$\mathcal{W}^+(t)$	$\mathcal{G}(t)$	$\tilde{\mathcal{G}}(t)$
<i>image in O_q</i>	$W^-(t)$	$W^+(t)$	$G(t)$	$\tilde{G}(t)$

Proof. By Definition 3.16. □

Recall the algebra homomorphism p from Proposition 4.7.

Lemma 6.5. *The map p sends the generating functions in (26) to the generating functions in (27) as follows.*

<i>argument in O_q</i>	$W^-(t)$	$W^+(t)$	$G(t)$	$\tilde{G}(t)$
<i>image in T_q</i>	$w^-(t)$	$w^+(t)$	$g(t)$	$\tilde{g}(t)$

Proof. By Definition 4.8. □

Next, we express the relations (5)–(17) in terms of generating functions.

Lemma 6.6. (See [16, Lemma 7.2].) *The following relations hold in the algebra*

\mathcal{O}_q .

$$[\mathcal{W}_0, \mathcal{W}^+(t)] = [\mathcal{W}^-(t), \mathcal{W}_1] = t^{-1}(\tilde{\mathcal{G}}(t) - \mathcal{G}(t))/(q + q^{-1}), \quad (28)$$

$$[\mathcal{W}_0, \mathcal{G}(t)]_q = [\tilde{\mathcal{G}}(t), \mathcal{W}_0]_q = \rho \mathcal{W}^-(t) - \rho t \mathcal{W}^+(t), \quad (29)$$

$$[\mathcal{G}(t), \mathcal{W}_1]_q = [\mathcal{W}_1, \tilde{\mathcal{G}}(t)]_q = \rho \mathcal{W}^+(t) - \rho t \mathcal{W}^-(t), \quad (30)$$

$$[\mathcal{W}^-(s), \mathcal{W}^-(t)] = 0, \quad [\mathcal{W}^+(s), \mathcal{W}^+(t)] = 0, \quad (31)$$

$$[\mathcal{W}^-(s), \mathcal{W}^+(t)] + [\mathcal{W}^+(s), \mathcal{W}^-(t)] = 0, \quad (32)$$

$$s[\mathcal{W}^-(s), \mathcal{G}(t)] + t[\mathcal{G}(s), \mathcal{W}^-(t)] = 0, \quad (33)$$

$$s[\mathcal{W}^-(s), \tilde{\mathcal{G}}(t)] + t[\tilde{\mathcal{G}}(s), \mathcal{W}^-(t)] = 0, \quad (34)$$

$$s[\mathcal{W}^+(s), \mathcal{G}(t)] + t[\mathcal{G}(s), \mathcal{W}^+(t)] = 0, \quad (35)$$

$$s[\mathcal{W}^+(s), \tilde{\mathcal{G}}(t)] + t[\tilde{\mathcal{G}}(s), \mathcal{W}^+(t)] = 0, \quad (36)$$

$$[\mathcal{G}(s), \mathcal{G}(t)] = 0, \quad [\tilde{\mathcal{G}}(s), \tilde{\mathcal{G}}(t)] = 0, \quad (37)$$

$$[\tilde{\mathcal{G}}(s), \mathcal{G}(t)] + [\mathcal{G}(s), \tilde{\mathcal{G}}(t)] = 0. \quad (38)$$

In these equations, $\rho = -(q^2 - q^{-2})^2$. We turn our attention to \mathcal{O}_q .

Lemma 6.7. *The following equations hold in the algebra \mathcal{O}_q .*

$$[W_0, W^+(t)] = [W^-(t), W_1] = t^{-1}(\tilde{G}(t) - G(t))/(q + q^{-1}), \quad (39)$$

$$[W_0, G(t)]_q = [\tilde{G}(t), W_0]_q = \rho W^-(t) - \rho t W^+(t), \quad (40)$$

$$[G(t), W_1]_q = [W_1, \tilde{G}(t)]_q = \rho W^+(t) - \rho t W^-(t), \quad (41)$$

$$[W^-(s), W^-(t)] = 0, \quad [W^+(s), W^+(t)] = 0, \quad (42)$$

$$[W^-(s), W^+(t)] + [W^+(s), W^-(t)] = 0, \quad (43)$$

$$s[W^-(s), G(t)] + t[G(s), W^-(t)] = 0, \quad (44)$$

$$s[W^-(s), \tilde{G}(t)] + t[\tilde{G}(s), W^-(t)] = 0, \quad (45)$$

$$s[W^+(s), G(t)] + t[G(s), W^+(t)] = 0, \quad (46)$$

$$s[W^+(s), \tilde{G}(t)] + t[\tilde{G}(s), W^+(t)] = 0, \quad (47)$$

$$[G(s), G(t)] = 0, \quad [\tilde{G}(s), \tilde{G}(t)] = 0, \quad (48)$$

$$[\tilde{G}(s), G(t)] + [G(s), \tilde{G}(t)] = 0. \quad (49)$$

Proof. Apply the map γ to relations (28)–(38). □

We now turn our attention to T_q .

Lemma 6.8. *The following equations hold in the algebra T_q .*

$$[w_0, w^+(t)] = [w^-(t), w_1] = t^{-1}(\tilde{g}(t) - g(t))/(q + q^{-1}), \quad (50)$$

$$[w_0, g(t)]_q = [\tilde{g}(t), w_0]_q = \rho w^-(t) - \rho t w^+(t), \quad (51)$$

$$[g(t), w_1]_q = [w_1, \tilde{g}(t)]_q = \rho w^+(t) - \rho t w^-(t), \quad (52)$$

$$[w^-(s), w^-(t)] = 0, \quad [w^+(s), w^+(t)] = 0, \quad (53)$$

$$[w^-(s), w^+(t)] + [w^+(s), w^-(t)] = 0, \quad (54)$$

$$s[w^-(s), g(t)] + t[g(s), w^-(t)] = 0, \quad (55)$$

$$s[w^-(s), \tilde{g}(t)] + t[\tilde{g}(s), w^-(t)] = 0, \quad (56)$$

$$s[w^+(s), g(t)] + t[g(s), w^+(t)] = 0, \quad (57)$$

$$s[w^+(s), \tilde{g}(t)] + t[\tilde{g}(s), w^+(t)] = 0, \quad (58)$$

$$[g(s), g(t)] = 0, \quad [\tilde{g}(s), \tilde{g}(t)] = 0, \quad (59)$$

$$[\tilde{g}(s), g(t)] + [g(s), \tilde{g}(t)] = 0. \quad (60)$$

Proof. Apply the map p to relations (39)–(49). □

7 The generating function $\omega(t)$

In this section, we introduce a function that will become useful later on.

Definition 7.1. Define the generating function

$$\omega(t) = \sum_{i=0}^{\infty} \binom{2i}{i} \left(\frac{t}{q + q^{-1}} \right)^{2i}.$$

Note that $\omega(0) = 1$.

Remark: The function $\omega(t)$ is the power series expansion of

$$\left(1 - \frac{4t^2}{(q + q^{-1})^2} \right)^{-\frac{1}{2}}.$$

However, we will not make use of this fact.

Define

$$T = \frac{q + q^{-1}}{qt + q^{-1}t^{-1}}, \quad S = \frac{q + q^{-1}}{qt^{-1} + q^{-1}t}. \quad (61)$$

Following [15], we express T and S as power series:

$$T = (q + q^{-1}) \sum_{\ell=0}^{\infty} (-1)^\ell q^{2\ell+1} t^{2\ell+1}, \quad S = (q + q^{-1}) \sum_{\ell=0}^{\infty} (-1)^\ell q^{-(2\ell+1)} t^{2\ell+1}. \quad (62)$$

Lemma 7.2. *The following identities hold:*

$$\omega(T) = \frac{1 + q^2 t^2}{1 - q^2 t^2}, \quad \omega(S) = \frac{1 + q^{-2} t^2}{1 - q^{-2} t^2}. \quad (63)$$

Proof. We first verify the equation on the left in (63). From (61) and Definition 7.1, we obtain

$$\omega(T) = \sum_{i=0}^{\infty} \binom{2i}{i} \left(\frac{1}{qt + q^{-1}t^{-1}} \right)^{2i}.$$

Recall a special case of the Newton binomial theorem [8, Theorem 5.5.1], which states that for any integer a ,

$$(1 + z)^a = \sum_{j=0}^{\infty} \binom{a}{j} z^j,$$

where we define

$$\binom{a}{j} = \frac{a(a-1)\cdots(a-j+1)}{j!}.$$

One readily checks that

$$\binom{-a}{j} = (-1)^j \binom{a+j-1}{j}.$$

We express $(1 + q^2t^2)^{-1}\omega(T)$ as a power series in t . We have

$$\begin{aligned} (1 + q^2t^2)^{-1}\omega(T) &= \sum_{i=0}^{\infty} \binom{2i}{i} \frac{(qt)^{2i}}{(1 + q^2t^2)^{2i+1}} \\ &= \sum_{i=0}^{\infty} \binom{2i}{i} q^{2i} t^{2i} \sum_{j=0}^{\infty} (-1)^j \binom{2i+j}{j} q^{2j} t^{2j} \\ &= \sum_{n=0}^{\infty} \sum_{i=0}^n (-1)^{n-i} \binom{2i}{i} q^{2n} t^{2n} \binom{n+i}{n-i} \\ &= \sum_{n=0}^{\infty} c_n q^{2n} t^{2n} \end{aligned}$$

where

$$c_n = \sum_{i=0}^n (-1)^{n-i} \binom{2i}{i} \binom{n+i}{n-i}. \quad (64)$$

We claim that $c_n = 1$ for all $n \geq 0$. Let n be given and consider the expression

$$\psi(x) = (1 - x)^{-n-1} (1 - x)^n. \quad (65)$$

We will compute the coefficient of x^n in two ways. First, notice that

$$\psi(x) = (1 - x)^{-n-1} (1 - x)^n = (1 - x)^{-1} = \sum_{i=0}^{\infty} x^i$$

so the coefficient of x^n in $\psi(x)$ is equal to 1.

Second, we expand (65) using the Newton binomial theorem. We have

$$\psi(x) = (1-x)^{-n-1}(1-x)^n = \sum_{s=0}^{\infty} x^s \sum_{i=0}^s (-1)^{s+i} \binom{n+i}{i} \binom{n}{s-i}. \quad (66)$$

In the double sum on the right of (66), the coefficient of x^n is

$$\sum_{i=0}^n (-1)^{n+i} \binom{n+i}{i} \binom{n}{n-i} = \sum_{i=0}^n (-1)^{n-i} \binom{2i}{i} \binom{n+i}{n-i}. \quad (67)$$

The right side of (67) matches the right side of (64), and therefore is equal to c_n . Hence, we have shown that $c_n = 1$, and the claim is proven. By the claim, we have

$$(1+q^2t^2)^{-1}\omega(T) = \sum_{n=0}^{\infty} q^{2n}t^{2n} = \frac{1}{1-q^2t^2}.$$

We have verified the equation on the left in (63). The equation on the right in (63) is similarly verified. \square

8 The first main result

Recall the generating functions from Definition 6.3. In this section, we express these generating functions in the basis for T_q given in Lemma 4.3. We now state our first main result.

Theorem 8.1. *In the algebra T_q ,*

$$w^-(t) = \omega(t) (x + x^{-1}), \quad (68)$$

$$w^+(t) = \omega(t) (y + y^{-1}), \quad (69)$$

$$g(t) = (q^2 - q^{-2}) \omega(t) (t(qx^{-1}y + qxy^{-1}) - q - q^{-1}), \quad (70)$$

$$\tilde{g}(t) = (q^2 - q^{-2}) \omega(t) (t(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - q - q^{-1}). \quad (71)$$

Theorem 8.1 will be proven at the end of this section.

Definition 8.2. In the algebra \mathcal{O}_q , define

$$\mathcal{Z}(t) = \sum_{i=0}^{\infty} \mathcal{Z}_i (q + q^{-1})^i t^i,$$

where \mathcal{Z}_i is from Definition 3.11.

We present two lemmas about $\mathcal{Z}(t)$. Recall S, T from (61) and the homomorphism γ from Definition 3.14.

Lemma 8.3. *The map γ sends $\mathcal{Z}(t) \mapsto (q + q^{-1})^2$.*

Proof. By Lemma 3.15, $\gamma(\mathcal{Z}_i) = 0$ for $i \geq 1$. Recall from Definition 3.11 that $\mathcal{Z}_0 = (q + q^{-1})^2$. The result follows. \square

Lemma 8.4. (See [15, Definition 8.4].) *In the algebra \mathcal{O}_q ,*

$$\begin{aligned} \mathcal{Z}(t) = & t^{-1}STW^-(S)W^+(T) + tSTW^+(S)W^-(T) - q^2STW^-(S)W^-(T) \\ & - q^{-2}STW^+(S)W^+(T) + (q^2 - q^{-2})^{-2}\mathcal{G}(S)\tilde{\mathcal{G}}(T). \end{aligned} \quad (72)$$

Proof. Let $\mathcal{Z}^c(t)$ denote the expression on the right side of equation (72). We expand S and T using the power series in (62). These expansions each have zero constant term, so $t^{-1}ST$ can be written as a power series in t . Hence $\mathcal{Z}^c(t)$ is a power series, which we express as

$$\mathcal{Z}^c(t) = \sum_{i=0}^{\infty} \mathcal{Z}_i^c (q + q^{-1})^i t^i.$$

For $i \geq 1$, the map ϕ sends $1 \otimes z_i \mapsto \mathcal{Z}_i^c$ by [15, Lemma 9.3], so $\mathcal{Z}_i^c = \mathcal{Z}_i$. By [15, Lemma 8.18], we have $\mathcal{Z}_0^c = (q + q^{-1})^2 = \mathcal{Z}_0$. By these observations, $\mathcal{Z}^c(t) = \mathcal{Z}(t)$ and the result follows. \square

Corollary 8.5. *In the algebra \mathcal{O}_q ,*

$$\begin{aligned} (q + q^{-1})^2 = & t^{-1}STW^-(S)W^+(T) + tSTW^+(S)W^-(T) - q^2STW^-(S)W^-(T) \\ & - q^{-2}STW^+(S)W^+(T) + (q^2 - q^{-2})^{-2}G(S)\tilde{G}(T). \end{aligned} \quad (73)$$

Proof. Routine consequence of Lemmas 8.3 and 8.4. \square

Corollary 8.6. *In the algebra T_q ,*

$$\begin{aligned} (q + q^{-1})^2 = & t^{-1}STw^-(S)w^+(T) + tSTw^+(S)w^-(T) - q^2STw^-(S)w^-(T) \\ & - q^{-2}STw^+(S)w^+(T) + (q^2 - q^{-2})^{-2}g(S)\tilde{g}(T). \end{aligned} \quad (74)$$

Proof. Apply p to both sides of equation (73). \square

Proof of Theorem 8.1. By [15, Lemma 8.22], there is a unique set of generating functions in \mathcal{O}_q ,

$$\mathcal{W}_c^-(t), \quad \mathcal{W}_c^+(t), \quad \mathcal{G}_c(t), \quad \tilde{\mathcal{G}}_c(t) \quad (75)$$

that satisfy equations (28)–(38), equation (72), and the conditions

$$\mathcal{W}_c^-(0) = \mathcal{W}_0, \quad \mathcal{W}_c^+(0) = \mathcal{W}_1. \quad (76)$$

Applying $\rho \circ \gamma$ to (75) and (76), we see that there is a unique set of generating functions in T_q ,

$$w_c^-(t), \quad w_c^+(t), \quad g_c(t), \quad \tilde{g}_c(t),$$

that satisfy equations (50)–(60), equation (74), and the conditions

$$w_c^-(0) = w_0, \quad w_c^+(0) = w_1. \quad (77)$$

We are going to display this unique solution. Going forward, we define

$$w_c^-(t), \quad w_c^+(t), \quad g_c(t), \quad \tilde{g}_c(t)$$

to be the right-hand sides of equations (68)–(71), in that order.

Since $\omega(0) = 1$, these candidates satisfy (77). Therefore, it suffices to show that these candidates satisfy equations (74) and (50)–(60).

We begin with (74). We claim that in the algebra T_q ,

$$(q + q^{-1})^2 = t^{-1}STw_c^-(S)w_c^+(T) + tSTw_c^+(S)w_c^-(T) - q^2STw_c^-(S)w_c^-(T) - q^{-2}STw_c^+(S)w_c^+(T) + (q^2 - q^{-2})^{-2}g_c(S)\tilde{g}_c(T). \quad (78)$$

To show this, we evaluate equations (68)–(71) using (61) and (63) to obtain the following equations:

$$w_c^-(S) = \frac{1 + q^{-2}t^2}{1 - q^{-2}t^2} (x + x^{-1}), \quad (79)$$

$$w_c^+(S) = \frac{1 + q^{-2}t^2}{1 - q^{-2}t^2} (y + y^{-1}), \quad (80)$$

$$w_c^-(T) = \frac{1 + q^2t^2}{1 - q^2t^2} (x + x^{-1}), \quad (81)$$

$$w_c^+(T) = \frac{1 + q^2t^2}{1 - q^2t^2} (y + y^{-1}), \quad (82)$$

$$g_c(S) = \frac{(q + q^{-1})^2(q - q^{-1})(1 + q^{-2}t^2)}{(1 - q^{-2}t^2)(qt^{-1} + q^{-1}t)} (qxy^{-1} + qx^{-1}y - qt^{-1} - q^{-1}t), \quad (83)$$

$$\tilde{g}_c(T) = \frac{(q + q^{-1})^2(q - q^{-1})(1 + q^2t^2)}{(1 - q^2t^2)(qt + q^{-1}t^{-1})} (q^{-1}xy + q^{-1}x^{-1}y^{-1} - qt - q^{-1}t^{-1}). \quad (84)$$

To verify (78), we evaluate the expression on its right using equations (79)–(84). We have now verified (78), and with it (74). The equations (50)–(60) can be verified in a straightforward manner. The reader is welcome to see Appendix A for more details.

We have now verified that the candidate functions satisfy (50)–(60) and (74), so the proof is complete. \square

9 The second main result

In this section we use the formulas in Theorem 8.1 to express the alternating elements of T_q in the basis for T_q given in Lemma 4.3.

Throughout this section, j is a natural number. The following is our second main result.

Theorem 9.1. *In the table below, we express the alternating elements of T_q in the basis for T_q from Lemma 4.3. For $k \in \mathbb{N}$,*

	Case $k = 2j$	Case $k = 2j + 1$
w_{k+1}	$\binom{2j}{j} (q + q^{-1})^{-2j} (y + y^{-1})$	0
w_{-k}	$\binom{2j}{j} (q + q^{-1})^{-2j} (x + x^{-1})$	0
g_k	$-\binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{2-2j}$	$\binom{2j}{j} q (q - q^{-1}) (q + q^{-1})^{1-2j} (xy^{-1} + x^{-1}y)$
\tilde{g}_k	$-\binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{2-2j}$	$\binom{2j}{j} q^{-1} (q - q^{-1}) (q + q^{-1})^{1-2j} (xy + x^{-1}y^{-1})$

Proof. We begin with \tilde{g}_k . Expanding $\omega(t)$ in the right side of (71), we have

$$\tilde{g}(t) = (q^2 - q^{-2}) (t(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - (q + q^{-1})) \sum_{i=0}^{\infty} \binom{2i}{i} \frac{t^{2i}}{(q + q^{-1})^{2i}}. \quad (85)$$

We examine the coefficient of t^k in (85) and obtain the following equations:

$$\tilde{g}_k = -\binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{2-2j} \quad \text{for } k = 2j, \quad (86)$$

$$\tilde{g}_k = \binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{1-2j} (q^{-1}xy + q^{-1}x^{-1}y^{-1}) \quad \text{for } k = 2j + 1. \quad (87)$$

Expanding $\omega(t)$ in the right side of (70), we have

$$g(t) = (q^2 - q^{-2}) (t(qxy^{-1} + qx^{-1}y) - (q + q^{-1})) \sum_{i=0}^{\infty} \binom{2i}{i} \frac{t^{2i}}{(q + q^{-1})^{2i}}. \quad (88)$$

We examine the coefficient of t^k in (88) and obtain the following equations:

$$g_k = -\binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{2-2j} \quad \text{for } k = 2j, \quad (89)$$

$$g_k = \binom{2j}{j} (q - q^{-1}) (q + q^{-1})^{1-2j} (qxy^{-1} + qx^{-1}y) \quad \text{for } k = 2j + 1. \quad (90)$$

Expanding $\omega(t)$ in the right side of (69), we have

$$w^+(t) = (y + y^{-1}) \sum_{i=0}^{\infty} \binom{2i}{i} \frac{t^{2i}}{(q + q^{-1})^{2i}}. \quad (91)$$

We examine the coefficient of t^k in (91) and obtain the following equations:

$$w_{k+1} = \binom{2j}{j} (q + q^{-1})^{-2j} (y + y^{-1}) \quad \text{for } k = 2j, \quad (92)$$

$$w_{k+1} = 0 \quad \text{for } k = 2j + 1. \quad (93)$$

Expanding $\omega(t)$ in the right side of (68), we have

$$w^-(t) = (x + x^{-1}) \sum_{i=0}^{\infty} \binom{2i}{i} \frac{t^{2i}}{(q + q^{-1})^{2i}}. \quad (94)$$

We examine the coefficient of t^k in (94) and obtain the following equations:

$$w_{-k} = \binom{2j}{j} (q + q^{-1})^{-2j} (x + x^{-1}) \quad \text{for } k = 2j, \quad (95)$$

$$w_{-k} = 0 \quad \text{for } k = 2j + 1. \quad (96)$$

□

We end this section with a comment. From the table given in Theorem 9.1, we see that each alternating element of T_q is a linear combination of the following nine elements:

$$1, \quad x, \quad y, \quad x^{-1}, \quad y^{-1}, \quad xy, \quad xy^{-1}, \quad x^{-1}y, \quad x^{-1}y^{-1}.$$

The above elements are linearly independent. Therefore, the alternating elements of T_q are contained in a 9-dimensional subspace of T_q .

Appendix A: Details from the proof of Theorem 8.1

This appendix clarifies some details regarding the proof of Theorem 8.1. We seek to show that the generating functions in (68)–(71) satisfy relations (50)–(60).

To simplify our arguments, we use the fact that the automorphism τ of T_q sends

$$w^+(t) \mapsto w^-(t), \quad w^-(t) \mapsto w^+(t), \quad g(t) \mapsto \tilde{g}(t), \quad \tilde{g}(t) \mapsto g(t).$$

We consider (50)–(60) individually.

- We show that $[w_0, w^+(t)] = [w^-(t), w_1]$.

By Definition 4.4, we have

$$\begin{aligned} [w_0, w^+(t)] &= [w_0, \omega(t)(y + y^{-1})] = \omega(t)[x + x^{-1}, y + y^{-1}] \\ &= [\omega(t)(x + x^{-1}), w_1] = [w^-(t), w_1]. \end{aligned}$$

- We show that $[w^-(t), w_1] = \frac{\tilde{g}(t) - g(t)}{t(q + q^{-1})}$.

We have

$$\begin{aligned} t^{-1} \frac{\tilde{g}(t) - g(t)}{q + q^{-1}} &= t^{-1} \frac{(q^2 - q^{-2})t\omega(t)(q^{-1}xy + q^{-1}x^{-1}y^{-1} - qxy^{-1} - qx^{-1}y)}{q + q^{-1}} \\ &= (q - q^{-1})\omega(t)(q^{-1}xy + q^{-1}x^{-1}y^{-1} - qxy^{-1} - qx^{-1}y) \\ &= \omega(t)[w_0, w_1] \\ &= [w^-(t), w_1]. \end{aligned}$$

- We show that $[w_0, g(t)]_q = \rho w^-(t) - \rho t w^+(t)$.

We have

$$\begin{aligned} [w_0, g(t)]_q &= qw_0g(t) - q^{-1}g(t)w_0 \\ &= (q^2 - q^{-2})\omega(t)(q(x + x^{-1})(tqxy^{-1} + tqx^{-1}y - q - q^{-1}) \\ &\quad - q^{-1}(tqxy^{-1} + tqx^{-1}y - q - q^{-1})(x + x^{-1})) \\ &= (q^2 - q^{-2})\omega(t)((tq^2y + tq^2x^{-2}y + tq^2y^{-1} + tq^2x^2y^{-1} \\ &\quad - q^2x - q^2x^{-1} - x - x^{-1}) - (tq^{-2}y + tq^{-2}y^{-1} \\ &\quad + tq^2x^{-2}y + tq^2x^2y^{-1} - x - x^{-1} - q^{-2}x - q^{-2}x^{-1})) \\ &= (q^2 - q^{-2})\omega(t)(q^2 - q^{-2})(ty + ty^{-1} - x - x^{-1}) \\ &= -(q^2 - q^{-2})^2(w^-(t) - tw^+(t)) \\ &= \rho w^-(t) - \rho t w^+(t). \end{aligned}$$

- We show that $[w_1, \tilde{g}(t)]_q = \rho w^+(t) - \rho t w^-(t)$.

Apply the τ automorphism to the above equation.

- We show that $[\tilde{g}(t), w_0]_q = \rho w^-(t) - \rho t w^+(t)$.

We have

$$\begin{aligned}
[\tilde{g}(t), w_0]_q &= q\tilde{g}(t)w_0 - q^{-1}w_0\tilde{g}(t) \\
&= (q^2 - q^{-2})\omega(t) (q(tq^{-1}xy + tq^{-1}x^{-1}y^{-1} - q - q^{-1})(x + x^{-1}) \\
&\quad - q^{-1}(x + x^{-1})(tq^{-1}xy + tq^{-1}x^{-1}y^{-1} - q - q^{-1})) \\
&= (q^2 - q^{-2})\omega(t) ((tq^2y + tq^{-2}x^2y + tq^2y^{-1} + tq^{-2}x^{-2}y^{-1} \\
&\quad - q^2x - q^2x^{-1} - x - x^{-1}) - (tq^{-2}y + tq^{-2}y^{-1} \\
&\quad + tq^{-2}x^{-2}y^{-1} + tq^{-2}x^2y - x - x^{-1} - q^{-2}x - q^{-2}x^{-1})) \\
&= (q^2 - q^{-2})\omega(t) (q^2 - q^{-2}) (ty + ty^{-1} - x - x^{-1}) \\
&= - (q^2 - q^{-2})^2 (w^-(t) - tw^+(t)) \\
&= \rho w^-(t) - \rho t w^+(t).
\end{aligned}$$

- We show that $[g(t), w_1]_q = \rho w^+(t) - \rho t w^-(t)$.

Apply the τ automorphism to the above equation.

- We show that $[w^-(s), w^-(t)] = 0$.

We have

$$[w^-(s), w^-(t)] = \omega(s)\omega(t)[x + x^{-1}, x + x^{-1}] = 0.$$

- We show that $[w^+(s), w^+(t)] = 0$.

We have

$$[w^+(s), w^+(t)] = \omega(s)\omega(t)[y + y^{-1}, y + y^{-1}] = 0.$$

- We show that $[w^-(s), w^+(t)] + [w^+(s), w^-(t)] = 0$.

We have

$$\begin{aligned}
[w^-(s), w^+(t)] &= \omega(s)\omega(t)[x + x^{-1}, y + y^{-1}] = -\omega(s)\omega(t)[y + y^{-1}, x + x^{-1}] \\
&= -[w^+(s), w^-(t)].
\end{aligned}$$

- We show that $s[w^-(s), g(t)] + t[g(s), w^-(t)] = 0$.

We have

$$\begin{aligned}
s[w^-(s), g(t)] &= (q^2 - q^{-2})s\omega(s)\omega(t) [x + x^{-1}, t(qxy^{-1} + qx^{-1}y) - q - q^{-1}] \\
&= (q^2 - q^{-2})st\omega(s)\omega(t)[x + x^{-1}, qxy^{-1} + qx^{-1}y] \\
&= -(q^2 - q^{-2})t\omega(t)\omega(s) [s(qxy^{-1} + qx^{-1}y) - q - q^{-1}, x + x^{-1}] \\
&= -t[g(s), w^-(t)].
\end{aligned}$$

- We show that $s[w^+(s), \tilde{g}(t)] + t[\tilde{g}(s), w^+(t)] = 0$.

Apply τ to each side of the previous equation.

- We show that $s[w^-(s), \tilde{g}(t)] + t[\tilde{g}(s), w^-(t)] = 0$.

We have

$$\begin{aligned}
s[w^-(s), \tilde{g}(t)] &= (q^2 - q^{-2})s\omega(s)\omega(t) [x + x^{-1}, t(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - q - q^{-1}] \\
&= (q^2 - q^{-2})st\omega(s)\omega(t)[x + x^{-1}, q^{-1}xy + q^{-1}x^{-1}y^{-1}] \\
&= -(q^2 - q^{-2})t\omega(t)\omega(s)[s(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - q - q^{-1}, x + x^{-1}] \\
&= -t[\tilde{g}(s), w^-(t)].
\end{aligned}$$

- We show that $s[w^+(s), g(t)] + t[g(s), w^+(t)] = 0$. Apply τ to each side of the previous equation.
- We show that $[g(s), g(t)] = 0$.

We have

$$\begin{aligned}
[g(s), g(t)] &= (q^2 - q^{-2})^2\omega(s)\omega(t) [s(qxy^{-1} + qx^{-1}y) - q - q^{-1}, \\
&\quad t(qxy^{-1} + qx^{-1}y) - q - q^{-1}] \\
&= (q^2 - q^{-2})^2st\omega(s)\omega(t)[qxy^{-1} + qx^{-1}y, qxy^{-1} + qx^{-1}y] \\
&= 0.
\end{aligned}$$

- We show that $[\tilde{g}(s), \tilde{g}(t)] = 0$.
- Apply τ to each side of the previous equation.
- We show that $[\tilde{g}(s), g(t)] + [g(s), \tilde{g}(t)] = 0$.

We have

$$\begin{aligned}
[\tilde{g}(s), g(t)] &= (q^2 - q^{-2})^2\omega(s)\omega(t) [s(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - q - q^{-1}, \\
&\quad t(qxy^{-1} + qx^{-1}y) - q - q^{-1}] \\
&= (q^2 - q^{-2})^2st\omega(s)\omega(t)[q^{-1}xy + q^{-1}x^{-1}y^{-1}, qxy^{-1} + qx^{-1}y] \\
&= -(q^2 - q^{-2})^2\omega(s)\omega(t) [s(qxy^{-1} + qx^{-1}y) - q - q^{-1}, \\
&\quad t(q^{-1}xy + q^{-1}x^{-1}y^{-1}) - q - q^{-1}] \\
&= -[g(s), \tilde{g}(t)].
\end{aligned}$$

We have now verified the equations (50)–(60), so the argument is complete.

Acknowledgement

The author is presently a graduate student at the University of Wisconsin–Madison. He is deeply grateful to his research advisor, Professor Paul Terwilliger, for many of the resources used and for insightful comments to improve the clarity and flow of this paper.

References

- [1] P. Baseilhac. Deformed Dolan-Grady relations in quantum integrable models. *Nuclear Phys. B* **709** (2005), 491–521, [arXiv:hep-th/0404149](#).
- [2] P. Baseilhac, S. Belliard. Generalized q -Onsager algebras and boundary affine Toda field theories. *Lett. Math. Phys.* **93** (2010), 213–228, [arXiv:0906.1215](#).
- [3] P. Baseilhac, S. Belliard. An attractive basis for the q -Onsager algebra, [arXiv:1704.02950](#).
- [4] P. Baseilhac, K. Koizumi. Sine-Gordon quantum field theory on the half-line with quantum boundary degrees of freedom. *Nuclear Phys. B* **649** (2003), 491–510, [arXiv:hep-th/0208005](#).
- [5] P. Baseilhac, K. Koizumi. A new (in)finite dimensional algebra for quantum integrable models. *Nuclear Phys. B* **720** (2005), 325–347, [arXiv:math-ph/0503036](#).
- [6] P. Baseilhac, S. Kolb. Braid group action and root vectors for the q -Onsager algebra. *Transform. Groups* **25** (2020), 363–389, [arXiv:1706.08747v3](#).
- [7] P. Baseilhac, K. Shigechi. A new current algebra and the reflection equation. *Lett. Math. Phys.* **92** (2010), 47–65, [arXiv:0906.1482](#).
- [8] R. Brualdi. *Introductory Combinatorics*, Fifth Edition, Pearson/Prentice Hall, Upper Saddle River, 2009.
- [9] A. Gupta. Modules over quantum Laurent polynomials. *J. Aust. Math. Soc.* **91** (2011), 323–341.
- [10] T. Ito. TD-pairs and the q -Onsager algebra. *Sugaku Expositions* **32** (2019), 205–232.
- [11] T. Ito, K. Tanabe, P. Terwilliger. Some algebra related to P - and Q -polynomial association schemes. *Codes and association schemes* (Piscataway, NJ, 1999), 167–192, *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.* **56**, Amer. Math. Soc., Providence, RI, 2001.
- [12] P. Terwilliger. The subconstituent algebra of an association scheme (Part III) *J. Algebraic Combin.* **2** (1993), 177–210.
- [13] P. Terwilliger. Two relations that generalize the q -Serre relations and the Dolan-Grady relations. *Physics and Combinatorics 1999 (Nagoya)*, 379–398, World Scientific Publishing, River Edge, NJ, 2001, [arXiv:math.QA/0307016](#).
- [14] P. Terwilliger. A conjecture concerning the q -Onsager algebra. *Nuclear Phys. B* **966** (2021) 115391, 26 pp., [arXiv:2101.09860](#).

- [15] P. Terwilliger. The alternating central extension of the q -Onsager algebra. *Comm. Math. Phys.* **387** (2021), 1771–1819, [arXiv:2103.03028v1](#).
- [16] P. Terwilliger. The q -Onsager algebra and its alternating central extension. *Nuclear Phys. B* **975** (2022), Paper No. 115662, 37 pp., [arXiv:2106.14041v1](#).

Owen Goff
Department of Mathematics
University of Wisconsin
480 Lincoln Drive
Madison, WI 53706-1388 USA
Email: ogoff@wisc.edu