

# Consistent Linear Orders for Supermajority Rules\*

Yasunori Okumura<sup>†</sup>

October 13, 2023

## Abstract

We consider linear orders of finite alternatives that are constructed by aggregating the preferences of individuals. We focus on a linear order that is consistent with the collective preference relation, which is constructed by one of the supermajority rules and modified using two procedures if there exist some cycles. One modification procedure uses the transitive closure, and the other uses the Suzumura consistent closure. We derive two sets of linear orders that are consistent with the (modified) collective preference relations formed by *any* of the supermajority rules and show that these sets are generally not empty. These sets of linear orders are closely related to those obtained through the ranked pairs method and the Schulze method. Finally, we consider two social choice correspondences whose output is one of the sets introduced above, and show that the correspondences satisfy the four properties: the extended Condorcet principle, the Pareto principle, the independence of clones, and the reversal symmetry.

**JEL Classification Numbers: D71; D72**

**Keywords: Supermajority rule, Voting, Schulze method; Ranked pairs method; Linear order; Transitive closure; Suzumura consistent closure**

---

\*A part of results of this manuscript is based on discussions with Minoru Kitahara. I would like to express my appreciation to him. This work was supported by JSPS KAKENHI Grant Numbers 20K01675 and 22K01402.

<sup>†</sup>Address: 2-1-6, Etchujima, Koto-ku, Tokyo, 135-8533 Japan. Phone:+81-3-5245-7300. Fax:+81-3-5245-7300. E-mail: okuyasu@gs.econ.keio.ac.jp

# 1 Introduction

We consider linear orders of finite alternatives that are obtained by aggregating the preferences of a number of individuals. Although Arrow's (1963) impossibility theorem applies such a linear order, there are several methods (without independence of irrelevant alternatives) introduced in previous studies to derive linear orders from the preferences of individuals such as the Borda method<sup>1</sup> (Borda 1781) and the Kemeny-Young method (Kemeny (1958), Young and Levenglick (1978), Young (1988)), the Tideman's ranked pairs method (Tideman (1987, 2006)) and the Schulze method (Schulze (2011, 2018)). A linear order is useful not only for choosing one alternative but also for choosing  $n(\geq 2)$  alternative(s) from  $n + 1$  or more, and for selecting substitutes from the chosen ones.

We focus on sets of the linear orders of alternatives derived as follows. First, for a profile of preferences of individuals and each of the supermajority rules, collective (or social) preference relations by aggregating the preferences of individuals are derived. Second, if the collective preference relations obtained by a supermajority rule have some cycles à la Condorcet (1784), then we modify it by using one of two procedures. Third, for each supermajority rules, a set of linear orders that are consistent with the profile of the (modified) collective preference relations is derived. Finally, we derive the intersection of the sets of consistent linear orders derived by all supermajority rules, which is shown to be nonempty with unrestricted domain of the individual preferences. In summary, we derive the set of linear orders that are consistent with the profiles of collective preferences constructed by *any* of the supermajority rules, although the profiles are modified if they have some cycles.

We explain our approach in more detail and its advantage. To have the collective preference relations, we use the supermajority rules, where a level of the rule is given by  $\alpha \in [1/2, 1)$ . In the supermajority rule with  $\alpha \in [1/2, 1)$ , an alternative  $a$  is (collectively) pairwise preferred to alternative  $b$  if and only if more than  $100 \times \alpha\%$  of individuals prefer  $a$  to  $b$ . Note that if  $\alpha = 1/2$ , then this is the simple majority rule; and if  $\alpha$  is almost 1, then this is the Pareto rule. Let  $R_\alpha$  be the collective preference relation for given  $\alpha \in [1/2, 1)$ .

We say that a linear order is consistent with a preference relation if for any two alternatives  $a$  and  $b$  such that  $a$  is pairwise preferred to  $b$  but not

---

<sup>1</sup>Pattanaik (2002) calls that the method to have an order by using the Borda count the Borda ranking rule. Here, we simply call it the Borda method.

vice versa,  $a$  has a higher rank than  $b$ . If  $R_\alpha$  has some cycles, then there is no consistent linear order of  $R_\alpha$ . Moreover, Greenberg (1979) shows that  $R_\alpha$  may have some cycles unless  $\alpha$  is almost 1 (the Pareto rule). Thus, we modify  $R_\alpha$  using two procedures: the transitive closure procedure by Bordes (1976) and the Suzumura-consistent closure procedure.<sup>2</sup> Denoting the modified  $R_\alpha$  by  $T(R_\alpha)$  and  $S(R_\alpha)$  through the former and latter procedures respectively, we derive sets of consistent linear orders for  $T(R_\alpha)$  and  $S(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ . Subsequently, we define the  $T$ -order set (resp. the  $S$ -order set) as the intersection of sets of consistent linear orders for  $T(R_\alpha)$  (resp.  $S(R_\alpha)$ ) for all  $\alpha \in [1/2, 1)$ . We show that neither the  $T$ -order set nor the  $S$ -order set is empty for any profile of individual preference relations.

An advantage of our approach is that it is not necessary to choose proper levels of the supermajority rule, because we focus on the intersection of the sets of consistent linear orders of  $T(R_\alpha)$  or  $S(R_\alpha)$  for *all*  $\alpha \in [1/2, 1)$ . To the contrary, using a single  $\alpha$  would cause us to overlook a significant amount of information. When we focus on  $R_\alpha$ ,  $(a, b) \in R_\alpha$  (resp.  $(a, b) \notin R_\alpha$ ) even if just a little bit more than (resp. bit less than or equal to)  $100 \times \alpha\%$  of individuals prefer  $a$  to  $b$ . That is, the differences in proportions are hardly captured in  $R_\alpha$  for one  $\alpha$ . On the other hand, our approach can capture the differences in proportions, because we consider  $R_\alpha$  for *all*  $\alpha \in [1/2, 1)$ .

Moreover, since there are both advantages and disadvantages to adopt a high (or low)  $\alpha$  as introduced later, it is difficult to determine proper  $\alpha$ . On the one hand, we consider a low  $\alpha$ . Related to a low  $\alpha$ , Arrow (1977) present the social choice method using  $T(R_{1/2})$  as one of the promising methods.<sup>3</sup> However, a collective preference relation  $R_\alpha$  with a low  $\alpha$  tends to have many cycles. Thus, if  $\alpha$  is low, then significant modifications to the collective preference relation may be necessary. Then, as warned by Arrow (1977), we may treat too many alternatives as indifferent with the modifications. Suzumura (2012) highlights a more critical drawback of this method; that is, in the case where  $\alpha$  is sufficiently low, even if every individual prefers an alternative to another, the latter may have a higher rank than that of the former in a consistent linear order of  $T(R_\alpha)$  or  $S(R_\alpha)$ .

On the other hand, although cycles cannot completely be eliminated unless  $\alpha$  is almost 1, they become rare event if  $\alpha$  is high. See for example Caplin

---

<sup>2</sup>According to the interpretation of Caplin and Nalebuff (1988), Condorcet also considers the transitive closure procedure to immune to the cycles.

Bossert et al. (2005) firstly define the Suzumura-consistency.

<sup>3</sup>Arrow (1977) mentions that this method is proposed by Donald Campbell and Gorges Bordes. In fact, Bordes (1976) introduce the characteristics of this method.

and Nalebuff (1988) and Balasko and Crès (1997) on the fact.<sup>4</sup> Therefore, if  $\alpha$  is high, then we rarely need to modify the collective preferences. However, as shown by Tovey (1997), the scarceness of cycles in the sense of Balasko and Crès (1997) is a result of the incompleteness of the collective preference relation.<sup>5</sup> Thus, we may also treat too many alternatives as indifferent if  $\alpha$  is large. Moreover, the Condorcet loser, which is the alternative that pairwise loses any other with the simple majority rule, may have the highest rank in a consistent linear order of  $T(R_\alpha)$  or  $S(R_\alpha)$  with a sufficiently large  $\alpha$ .

To understand these points, we consider the following example. Suppose that there are four alternatives  $a$ ,  $b$ ,  $c$  and  $d$ . The ratio of individuals who prefer  $a$  to  $b$  is 0.7, that of those who prefer  $b$  to  $c$  is 0.66 and that of those who prefer  $c$  to  $a$  is 0.64. Moreover, the ratios of individuals who prefer  $d$  to each of the others is 0.6 and therefore  $d$  is the Condorcet loser. Note that in this case, the modified collective preferences with the two procedures are the same for all  $\alpha \in [1/2, 1)$ . Then, the characteristics of the consistent linear order of the modified collective preference relation with each  $\alpha \in [1/2, 1)$  is given in the following table. If  $0.64 \leq \alpha < 1$ , then there is no cycle but  $d$  can have the highest position in some consistent linear orders. Therefore, adopting a high  $\alpha$  may increase consistent linear orders and some of them is such that the Condorcet loser has the highest ranking.

$\alpha$	Characteristics of consistent linear orders
$[0.5, 0.6)$	$d$ has the lowest rank
$[0.6, 0.64)$	Any
$[0.64, 0.66)$	the ranks of $a$ and $b$ are respectively higher than those of $b$ and $c$
$[0.66, 0.7)$	the rank of $a$ is higher than $b$
$[0.7, 1)$	Any

This example implies that it is hard to decide a proper  $\alpha \in [1/2, 1)$  for deriving a “good” consistent linear order. However, in our approach, there

---

<sup>4</sup>A result of Nakamura (1979) also claims this fact, because the Nakamura number, which is the largest number of alternatives ensuring no cycle (plus one), is increasing in the number of individuals times  $\alpha$ .

Further, there is a vast literature on the acyclic domain, which is the set of preferences that guarantee nonexistence of cycles. On this literature, Black (1958) is the pioneer and Greenberg (1979), Slutsky (1979), Coughlin (1981), Caplin and Nalebuff (1988), Saari (2014) and Gjorgjiev and Xeferis (2015), and Crès and Ünver (2017) consider the acyclic domains with supermajority rules. Their results imply that the domain would be wider when  $\alpha$  rises.

<sup>5</sup>Thus, Tovey (1997) states that “An interesting research problem would be how to modify the model to retain the scarceness of cycles but permit more complete aggregate preferences.” In this study, we try to solve this problem.

is no need to decide it at all. In fact, in the example, the linear order in the order of  $a$ ,  $b$ ,  $c$  and  $d$  is the unique linear order that is in the  $T$ -order set, which is in this case equivalent to the  $S$ -order set.

Our results are especially related to two voting methods: the ranked pairs method (Tideman (1987, 2006)) and the Schulze method (Schulze (2011, 2018)), which are two of the influential methods. Their common feature is that they firstly derive a binary relation from the preferences of individuals. As summarized by Tideman (2006), these two methods share many important properties. Moreover, the Schulze method is used in several real world elections (Schulze (2018)). First, we show that any linear order obtained by the ranked pairs method is in the  $S$ -order set. Second, we show that the set of linear orders that can be obtained by the Schulze method is equivalent to the  $T$ -order set. This result provides new description and interpretation of the Schulze method. Since the  $T$ -order set is a subset of the  $S$ -order set, any linear order obtained by the Schulze method is also in the  $S$ -order set. These results imply both the ranked pairs method and the Schulze method are in the class of methods that choose one from the  $S$ -order set. Consequently, this clarifies that the relationship between these methods, despite being not well-known, is interconnected.

Finally, we consider two social welfare correspondences whose outputs are the  $S$ -order set and the  $T$ -order set, respectively. We show that the correspondence satisfies four properties: the extended Condorcet criterion, the Pareto principle, the independence of clones and the reversal symmetry, which are originally introduced by Truchon (1998), Arrow (1963), Tideman (1987) and Saari (1993), respectively.

## 2 Model

Let  $A$  be the finite set of alternatives where  $|A| \geq 3$ . Let  $R \subseteq A \times A$  be a binary relation on  $A$ . In this study, we focus only on a irreflexive binary relation; that is,  $(a, a) \notin R$  for all  $a \in A$ . The asymmetric part of  $R$  denoted by  $P(R)$  is a binary relation on  $A$  satisfying  $(a, b) \in P(R)$  if and only if  $(a, b) \in R$  and  $(b, a) \notin R$ . As mentioned particularly in the subsequent section, a binary relation on  $A$  denoted by  $R$  represents not only the preference of an individual but a collective (or social) preference that is obtained by aggregating the preferences of a number of individuals via each of the supermajority rules.

We define the following properties of binary relations. A binary relation  $R$  on a set  $A$  is

**complete** if for all  $a, b \in A$  such that  $a \neq b$ ,  $(a, b) \in R$  or  $(b, a) \in R$ ,

**negatively transitive** if  $[(a, b) \notin R \text{ and } (b, c) \notin R]$  implies  $(a, c) \notin R$ , for all  $a, b, c \in A$ ,

**transitive** if  $[(a, b) \in R \text{ and } (b, c) \in R]$  implies  $(a, c) \in R$ , for all  $a, b, c \in A$ ,

**Suzumura consistent** if for all  $K \in \mathbb{N} \setminus \{1\}$  and for all  $a_0, a_1, \dots, a_K \in A$ ,  $(a_{k-1}, a_k) \in R$  for all  $k \in \{1, \dots, K\}$  implies  $(a_K, a_0) \notin P(R)$ ,

**P-acyclic** if for all  $K \in \mathbb{N} \setminus \{1\}$  and for all  $a_0, a_1, \dots, a_K \in A$ ,  $(a_{k-1}, a_k) \in P(R)$  for all  $k \in \{1, \dots, K\}$  implies  $(a_K, a_0) \notin P(R)$ .

**asymmetric** if for all  $a, b \in A$ ,  $(a, b) \in R$  implies  $(b, a) \notin R$ .

The following facts are due to Suzumura (1983) and Bossert and Suzumura (2010).

- Lemma 1**
1. *If  $R$  is transitive, then it is Suzumura consistent.*
  2. *If  $R$  is Suzumura consistent, then it is P-acyclic.*
  3. *If  $R$  is asymmetric, then it is P-acyclic if and only if it is Suzumura consistent.*

A finite sequence of alternatives  $(a_1, a_2, \dots, a_m)$  where  $a_i \in A$  for all  $i = 1, \dots, m$  ( $\geq 2$ ) is called a **path for  $R$  from  $a$  to  $b$**  if  $a_1 = a$ ,  $a_m = b$ ,  $(a_l, a_{l+1}) \in R$  for all  $l = 1, 2, \dots, m - 1$ , and  $a_1, a_2, \dots, a_m$  are distinct, where  $m$  is the length of the path.<sup>6</sup> Let  $\mathcal{B}([a, b], R)$  be the set all paths for  $R$  from  $a$  to  $b$ . In words, if there is a path for  $R$  from  $a$  to  $b$  whose length is 2; that is, if  $(a, b) \in R$ , then we say that  $a$  directly beats  $b$  for  $R$ . If  $\mathcal{B}([a, b], R) \neq \emptyset$  but the length of any path from  $a$  to  $b$  is 3 or more, then we say that  $a$  indirectly beats  $b$ , for  $R$ . If there are two paths for  $R$  from  $a$  to  $b$  and that from  $b$  to  $a$ , then there is a **cycle** for  $R$  (containing  $a$  and  $b$ ). A cycle for  $P(R)$  is called a **P-cycle** for  $R$ . Trivially, if  $R$  is P-acyclic, then there is no P-cycle for  $R$ .

Let a **partial order** be asymmetric and transitive binary relation. Let a **weak order** be a partial order that satisfies negative transitivity. Moreover, let a **linear order** be a weak order that satisfies completeness. If  $R$  is a linear order where  $(a_1, a_2), (a_2, a_3), \dots, (a_{|A|-1}, a_{|A|}) \in R$  and  $a_1, a_2, \dots, a_{|A|}$  are distinct, then we simply write

$$R : a_1, a_2, \dots, a_{|A|}.$$

---

<sup>6</sup>In several previous studies, a path is referred to as a “beatpath”.

**Definition 1** For a binary relation  $R$  on  $A$ , let  $\mathcal{C}(R)$  be the set of linear orders which are **consistent with**  $R$  if any  $\hat{R} \in \mathcal{C}(R)$  is a linear order on  $A$  and  $(a, b) \in P(R)$  implies  $(a, b) \in \hat{R}$ .<sup>7</sup>

That is, for any  $\hat{R} \in \mathcal{C}(R)$ , if  $(a, b) \in P(R)$ , then  $(a, b) \in \hat{R}$  and since  $\hat{R}$  is a linear order,  $(a, b) \in P(\hat{R})$ . On the other hand, if either  $[(a, b) \in R$  and  $(b, a) \in R]$  or  $[(a, b) \notin R$  and  $(b, a) \notin R]$  holds, either  $(a, b) \in \hat{R}$  or  $(b, a) \in \hat{R}$  is allowed to hold.

We provide a class of algorithms to construct a linear order  $\hat{R} \in \mathcal{C}(R)$  for a P-acyclic  $R$ .

First, for  $A' \subseteq A$  a binary relation  $R$ ,  $M(A', R)$  be a maximal set; that is,

$$M(A', R) = \{a \in A' \mid (b, a) \notin P(R) \text{ for all } b \in A' \setminus \{a\}\}.$$

We consider the following algorithms.

**Step 1:** Let  $a_1 \in M(A, R)$ .

**Step  $t = 2, \dots, |A|$ :** Let  $a_t \in M(A \setminus \{a_1, \dots, a_{t-1}\}, R)$ .

Finally, we have a linear order  $\hat{R} : a_1, \dots, a_{|A|}$ .

Since  $M(A', R)$  may not be singleton for some  $A' \subseteq A$ , this is a class of algorithms that we call the **sequential maximal ordering** (hereafter **SMO**) class. If  $R$  is P-acyclic, then each algorithm in the SMO class results in a consistent linear order with  $R$ , because as shown by Bossert and Suzumura (2010),  $R$  is P-acyclic and thus the maximal set is not empty for any step.

**Proposition 1**  $\mathcal{C}(R) \neq \emptyset$  if and only if  $R$  is P-acyclic. Moreover, a linear order is consistent with  $R$  if and only if it is obtained by an algorithm within the SMO class with  $R$ .

**Proof.** The only-if-part of the first result is trivial. Moreover, the if-part of the first result is trivial from the second one. Thus, we only show the second result. We use the following result, which is due to Bossert and Suzumura (2010, Theorem 2.4).

**Remark 1** If  $R$  is P-acyclic, then  $M(A', \succ_s)$  is non-empty for any finite  $A' \subseteq A$ .

---

<sup>7</sup>In the terminology of set theory,  $\hat{R} \in \mathcal{C}(R)$  if and only if  $\hat{R}$  is an (linear order) extension of  $P(R)$ ; that is,  $\hat{R}$  is a super set of  $P(R)$  and a linear order. Thus, if  $R$  is asymmetric,  $\hat{R} \in \mathcal{C}(R)$  if and only if  $\hat{R}$  is an extension of  $R$ .

By Remark 1, if  $R$  is P-acyclic, the maximal set is not empty for any step and thus each algorithm in the SMO class is well-defined.

First, suppose that  $\hat{R} : a_1, a_2, \dots, a_{|A|}$  is a linear order obtained by the SMO algorithm class. Then, for all  $t$  and  $t'$  such that  $1 \leq t < t' \leq |A|$ ,  $(a_t, a_{t'}) \notin P(R)$ . Therefore,  $\hat{R} \in \mathcal{C}(R)$ . Second, suppose  $\hat{R} (: a_1, a_2, \dots, a_{|A|}) \in \mathcal{C}(R)$ . Suppose not; that is,  $\hat{R}$  is not obtained by any algorithm within the SMO class with  $R$ . Then, we can let  $t$  be the smallest integer such that  $a_t \notin M(A \setminus \{a_1, \dots, a_{t-1}\}, R)$ ; that is, there is  $t' > t$  such that  $(a_{t'}, a_t) \in P(R)$ , which contradicts  $\hat{R} \in \mathcal{C}(R)$ . **Q.E.D.**

Next, we focus on the intersection of the sets of consistent linear orders for several binary relations.

**Lemma 2** *Let  $R_1, \dots, R_J$  be binary relations. Then,*

$$\bigcap_{j \in \{1, \dots, J\}} \mathcal{C}(R_j) \subseteq \mathcal{C} \left( \bigcup_{j \in \{1, \dots, J\}} P(R_j) \right). \quad (1)$$

**Proof.** If the left-hand side of (1) is empty, then this is trivial. Thus, we assume that it is not empty and let  $\hat{R}$  be an element in it. Suppose not; that is,  $\hat{R}$  not an element in the right-hand of (1). Then, there are  $a, b \in A$  such that  $(a, b) \in \hat{R}$  but  $(b, a) \in \bigcup_{j \in \{1, \dots, J\}} P(R_j)$ . This implies that  $(b, a) \in P(R_j)$ ; that is,  $(b, a) \in R_j$  and  $(a, b) \notin R_j$  for some  $j = 1, \dots, J$ . However, these imply  $\hat{R} \notin \mathcal{C}(R_j)$ , which is a contradiction. **Q.E.D.**

If some conditions are satisfied, then (1) holds with equality; that is, we have the following result, which is the most important technical result of this study.

**Theorem 1** *If  $R_1 \subseteq R_2 \subseteq \dots \subseteq R_J$  and  $R_j$  is P-acyclic for all  $j \in \{1, \dots, J\}$ , then*

$$\bigcap_{j \in \{1, \dots, J\}} \mathcal{C}(R_j) = \mathcal{C} \left( \bigcup_{j \in \{1, \dots, J\}} P(R_j) \right).$$

**Proof.** By Lemma 2, it is sufficient to show

$$\bigcap_{j \in \{1, \dots, J\}} \mathcal{C}(R_j) \supseteq \mathcal{C} \left( \bigcup_{j \in \{1, \dots, J\}} P(R_j) \right). \quad (2)$$



If the right-hand side of (2) is empty, (2) is trivial. Thus, suppose that  $\hat{R}$  is an element of the right-hand side of (2). Then, toward a contradiction, suppose that  $\hat{R}$  is not an element of the left-hand side of (2). By the P-acyclicity of  $R_j$  for all  $j \in \{1, \dots, J\}$  and Proposition 1,  $\mathcal{C}(R_j) \neq \emptyset$  for all  $j \in \{1, \dots, J\}$ . Therefore, there are  $a, b \in A$  and  $j' \in \{1, \dots, J\}$  such that  $(a, b) \in \hat{R}$ ,  $(b, a) \in R_{j'}$  and  $(a, b) \notin R_{j'}$ . Since  $\hat{R}$  is asymmetric,  $(b, a) \notin \hat{R}$ . Since  $(b, a) \in P(R_{j'})$  and  $(a, b) \in \hat{R}$ , for some  $j'' \in \{1, \dots, J\}$ ,  $(a, b) \in P(R_{j''})$ ; that is,  $(a, b) \in R_{j''}$  and  $(b, a) \notin R_{j''}$ . If  $R_{j''} \subseteq R_{j'}$ , then  $(a, b) \in R_{j''}$  implies  $(a, b) \in R_{j'}$ , which is a contradiction. Thus,  $R_{j'} \subseteq R_{j''}$ , but  $(b, a) \in R_{j'}$  implies  $(b, a) \in R_{j''}$ , which is a contradiction. Therefore, we have Theorem 1. **Q.E.D.**

Note that even if the conditions of Theorem 1 are satisfied, the right-hand side of (2) may be empty. For example, suppose

$$R_1 = \{(a, b), (b, c)\}, \text{ and } R_2 = R_1 \cup \{(c, b), (c, a)\}.$$

Then,  $P(R_1) \cup P(R_2) = \{(a, b), (b, c), (c, a)\}$  has a P-cycle. Therefore,  $R_1 \subseteq R_2$  and they are P-acyclic, but (2) is empty. In our model in the subsequent sections, (2) is not empty.

We consider the following example to understand Theorem 1.

### Example 1

Let  $A = \{a, b, c, d\}$ ,  $R_1 = \emptyset$ ,  $R_2 = \{(a, b)\}$ ,  $R_3 = \{(a, b), (b, c)\}$ ,  $R_4 = R_3 \cup \{(b, a), (a, c), (c, a), (c, b)\}$ ,  $R_5 = R_4 \cup \{(a, d), (b, d), (c, d)\}$ . Then,  $R_j$  is P-acyclic for all  $i \in \{1, \dots, 5\}$  and  $R_1 \subseteq R_2 \subseteq \dots \subseteq R_5$ . We have

$$\bigcup_{i \in \{1, \dots, I\}} P(R_j) = \{(a, b), (b, c), (a, d), (b, d), (c, d)\},$$

$$\mathcal{C}\left(\bigcup_{j \in \{1, \dots, J\}} P(R_j)\right) = \{R\}, \text{ where } R: a, b, c, d,$$

and  $R$  is a consistent linear order for  $R_j$  for all  $j \in \{1, \dots, 5\}$ . Moreover,  $R \in \mathcal{C}(R_j)$  for all  $j \in \{1, \dots, 5\}$ . Therefore, Theorem 1 is satisfied in this example.

Since  $R$  may not be P-acyclic, by Proposition 1, there may be no linear order that is consistent with  $R$ . Thus, we modify  $R$  to have its consistent linear order, even if  $R$  has some P-cycle.

First, a binary relation  $T(R)$  is a **transitive closure** of  $R$  if  $T(R)$  is the smallest transitive binary relation containing  $R$ . This modification is introduced by Bordes (1976) to escape from Condorcet cycles. By Bossert and Suzumura (2010, Theorem 2.1),  $T(R)$  is calculated as follows. For a binary relation  $R$ ,  $(a, b) \in T(R)$  if and only if  $\mathcal{B}([a, b], R) \neq \emptyset$ . Since  $T(R)$  is transitive and Lemma 1 holds,  $T(R)$  is also P-acyclic. Note that as is shown by Deb (1977),  $M(A, T(R))$  is equivalent to the Schwartz set for an asymmetric  $R$ .

Second, a binary relation  $S(R)$  is a **Suzumura-consistent closure** of  $R$  if  $S(R)$  is the smallest Suzumura-consistent binary relation containing  $R$ . Bossert and Suzumura (2010, Theorem 2.8) show that for a binary relation  $R$ ,

$$S(R) = R \cup \{(a, b) \in A \times A \mid (a, b) \in T(R) \text{ and } (b, a) \in R\}.$$

Since  $S(R)$  is Suzumura consistent,  $S(R)$  is also P-acyclic.

We immediately have for any  $R$ ,  $R \subseteq S(R) \subseteq T(R)$  and thus  $S(R)$  is a smaller modification of  $R$  comparing to  $T(R)$ . By Lemma 1 and Proposition 1, we can have a consistent linear order by using  $S(R)$ . Moreover, we have the following results.

**Lemma 3**  $P(S(R)) \subseteq P(T(R))$  and  $P(S(R)) \subseteq P(R)$ .

**Proof.** Let  $(a, b) \in P(S(R))$ ; that is,  $(a, b) \in S(R)$  and  $(b, a) \notin S(R)$ . Since  $S(R) \subseteq T(R)$ ,  $(a, b) \in T(R)$ .

First, we show  $P(S(R)) \subseteq P(T(R))$ . Suppose not; that is,  $(a, b) \notin P(T(R))$ . Since  $(a, b) \in T(R)$ ,  $(b, a) \in T(R)$ . Then,  $(b, a) \in T(R)$  and  $(b, a) \notin S(R)$  imply  $(a, b) \notin R$ . Then,  $(a, b) \in T(R)$  and  $R \subseteq T(R)$  are not compatible. Hence  $P(S(R)) \subseteq P(T(R))$ .

Second, we show  $P(S(R)) \subseteq P(R)$ . Since  $(b, a) \notin S(R)$ ,  $(b, a) \notin R$ . Moreover, since  $(b, a) \notin R$  and  $(a, b) \in S(R)$ ,  $(a, b) \in R$ . Therefore,  $(a, b) \in P(R)$ . **Q.E.D.**

The following result is due to Suzumura (1983) and Bossert and Suzumura (2010).

**Lemma 4** If  $R \subseteq R'$ , then  $T(R) \subseteq T(R')$  and  $S(R) \subseteq S(R')$ .

We have the following result.

**Lemma 5** For any  $R$ ,

$$\emptyset \neq \mathcal{C}(T(R)) \subseteq \mathcal{C}(S(R)).$$

**Proof.** By Lemma 1,  $T(R)$  is P-acyclic. By Proposition 1,  $\emptyset \neq \mathcal{C}(T(R))$ . Thus, we can let  $\hat{R} \in \mathcal{C}(T(R))$ . For any  $(a, b) \in P(T(R))$ ,  $(a, b) \in \hat{R}$ . Then, since  $P(S(R)) \subseteq P(T(R))$  (by Lemma 4), for any  $(a, b) \in P(S(R))$ ,  $(a, b) \in \hat{R}$ . Therefore,  $\hat{R} \in \mathcal{C}(S(R))$ . **Q.E.D.**

We provide a sufficient condition where the three sets are equivalent.

**Lemma 6** *If  $R$  is asymmetric and P-acyclic, then*

$$\mathcal{C}(T(R)) = \mathcal{C}(S(R)) = \mathcal{C}(R). \quad (3)$$

**Proof.** First, by Lemma 1, if  $R$  is asymmetric and P-acyclic, then  $R$  is Suzumura-consistent and thus  $S(R) = R$ . We have  $\mathcal{C}(S(R)) = \mathcal{C}(R)$ .

By Lemma 5, showing  $\mathcal{C}(S(R)) \subseteq \mathcal{C}(T(R))$  is sufficient. Suppose not; that is, there is  $\hat{R} \in \mathcal{C}(S(R)) \setminus \mathcal{C}(T(R))$ . Then, there is  $(a, b) \in \hat{R}$  such that  $(b, a) \in P(T(R)) \setminus P(S(R))$ . Then,  $(a, b) \notin T(R)$  implies  $(a, b) \notin S(R)$ . Therefore,  $(b, a) \in T(R)$ ; that is,  $(a_1, a_2, \dots, a_m) \in \mathcal{B}([b, a], R)$ . If  $(a_l, a_k) \in T(R)$  for some  $m \geq l > k \geq 1$ , then there is a P-cycle, because  $R$  is asymmetric. Thus,  $(a_l, a_k) \notin T(R)$  and moreover  $(a_l, a_k) \notin S(R)$  for all  $m \geq l > k \geq 1$ . Therefore,  $(a_l, a_{l+1}) \in P(S(R))$  for all  $l = 1, 2, \dots, m-1$ . Therefore,  $(a_l, a_{l+1}) \in \hat{R}$  for all  $l = 1, 2, \dots, m-1$ . Since  $\hat{R}$  is transitive,  $(b, a) \in \hat{R}$  contradicting  $(a, b) \in \hat{R}$  and  $\hat{R}$  is asymmetric. **Q.E.D.**

However, the three sets are not equivalent in general, we consider the following example.

### Example 2

Let  $R = \{(a, b), (b, c), (c, a), (d, a)\}$ . Then,  $R$  is not P-acyclic and, by Proposition 1,  $\mathcal{C}(R) = \emptyset$ . In this case,

$$S(R) = R \cup \{(b, a), (c, b), (a, c)\}$$

Therefore, any linear order  $\hat{R}$  such that  $(d, a) \in \hat{R}$  is in  $\mathcal{C}(S(R))$ . On the other hand,

$$T(R) = S(R) \cup \{(d, b), (d, c)\}$$

A linear order is in  $\mathcal{C}(T(R))$  if and only if  $d$  is the highest position in the linear order. Therefore, for example,  $\hat{R} : c, b, d, a$  is in  $\mathcal{C}(S(R))$  but not in  $\mathcal{C}(T(R))$ . Hence  $\mathcal{C}(T(R)) \subsetneq \mathcal{C}(S(R))$  in this example.

**Corollary 1** *Suppose that  $R$  is asymmetric. Then,  $\mathcal{C}(R) \neq \emptyset$  if and only if (3) holds.*

**Proof.** First, by Proposition 1 and Lemma 6, if  $\mathcal{C}(R) \neq \emptyset$ , then  $R$  is P-acyclic and thus (3) holds.

Second, by Lemma 5, neither  $\mathcal{C}(T(R))$  nor  $\mathcal{C}(S(R))$  is empty. Therefore, if  $\mathcal{C}(R) \neq \emptyset$ , then (3) does not hold. **Q.E.D.**

In the subsequent sections, in the case where  $\mathcal{C}(R) \neq \emptyset$ , we use  $T(R)$  or  $S(R)$  instead of  $R$ . Corollary 1 implies that in the other case, (3) holds. Thus, we focus on  $\mathcal{C}(R)$  as long as it is nonempty.

### 3 Collective preference relations

Let  $I$  be the set of finite individuals (or voters) where  $|I| \geq 3$ . Each individual  $i \in I$  has a binary relation  $R^i$  over  $A$  where  $(a, b) \in R^i$  (resp.  $(a, b) \in P(R^i)$ ) for  $a, b \in A$  implies that individual  $i$  weakly (resp. strictly) prefers  $a$  to  $b$ .

We introduce supermajority rules for constructing collective preference relations by aggregating the preferences of individuals. Let

$$N[a, b] = |\{i \in I \mid (a, b) \in P(R^i)\}|,$$

which is the number of individuals who strictly prefer  $a$  to  $b$ . Moreover, let  $f : \mathbb{Z}_+ \times \mathbb{Z}_+ \rightarrow [0, 1]$  be such that for any  $n \geq 0$  and  $m \geq 0$  satisfying  $n + m \leq |I|$ ,  $f(n + 1, m) > f(n, m)$ ,  $f(n, m + 1) < f(n, m)$ , and  $f(n, m) = 1/2$  if  $n = m$ . For example,  $f(n, m)$  can be

$$\begin{aligned} f(n, m) &= \frac{n}{n + m} \text{ if } n + m > 0 \\ &= \frac{1}{2} \text{ if } n + m = 0, \text{ or} \end{aligned} \tag{4}$$

$$f(n, m) = \frac{|I| + (n - m)}{2|I|}. \tag{5}$$

For notational simplicity, we let  $\Phi[a, b] = f(N[a, b], N[b, a])$ .<sup>8</sup> We let  $R_\alpha \subseteq A \times A$  be a collective preference relation for  $\alpha \in [1/2, 1)$  that satisfies  $(a, b) \in R_\alpha$  if and only if  $\Phi[a, b] > \alpha$ . Hereafter, we fix  $f$ . In the terminology

---

<sup>8</sup>If  $R^i$  is a linear order for all  $i \in I$ , then  $\Phi[\cdot, \cdot]$  defined by (4) is equivalent to that defined by (5), because, in that case,  $N[a, b] + N[b, a] = |I|$ .

of Schulze (2011),  $\Phi[a, b]$  represents the degree of the strength of the link between  $a$  and  $b$ . Moreover, if (4) and (5), then the strength of a link is respectively measured by ratio and margin, respectively.

Since many previous studies focus only on  $R_{1/2}$ , we introduce the following terms with regards to  $R_{1/2}$ . An alternative  $a$  is said to be the **Condorcet winner** if  $(a, b) \in R_{1/2}$  for all  $b \in A \setminus \{a\}$ . On the other hand,  $a$  is said to be the **Condorcet loser** if  $(b, a) \in R_{1/2}$  for all  $b \in A \setminus \{a\}$ .

We consider an example to understand the construction of the binary relations.

### Example 3

There are  $A = \{a, b, c, d\}$  and  $I = \{1, \dots, 9\}$ . All individuals have a linear preference order on  $A$  and

$$\begin{aligned} R^i &: d, a, b, c, \text{ for all } i = 1, 2, 3, \\ R^{i'} &: b, c, a, d, \text{ for all } i' = 4, 5, 6, \\ R^{i''} &: c, a, b, d, \text{ for all } i'' = 7, 8, \\ R^9 &: d, c, a, b. \end{aligned}$$

Since this is a linear order,  $N[x, y] = 9 - N[y, x]$  for all  $x, y \in A$  such that  $x \neq y$ . Moreover,  $N[a, b] = N[b, c] = N[c, a] = 6$  and  $N[x, d] = 5$  for all  $x \in A \setminus \{d\}$ . Then,  $d$  is the Condorcet loser. When we adopt (4) or (5),  $\Phi[a, b] = \Phi[b, c] = \Phi[c, a] = 2/3$  and  $\Phi[x, d] = 5/9$  for all  $x \in A \setminus \{d\}$ . Then,

$$\begin{aligned} R_\alpha &= \{(a, b), (b, c), (c, a), (a, d), (b, d), (c, d)\} \text{ for all } \alpha \in [1/2, 5/9), \\ R_{\alpha'} &= \{(a, b), (b, c), (c, a)\} \text{ for all } \alpha' \in [5/9, 2/3), \\ R_{\alpha''} &= \emptyset \text{ for all } \alpha'' \in [2/3, 1). \end{aligned}$$

We have the following result on  $R_\alpha$ .

- Lemma 7**
1. For all  $\alpha \in [1/2, 1)$ ,  $R_\alpha$  is asymmetric.
  2. For all  $\alpha, \beta \in [1/2, 1)$  such that  $\alpha \geq \beta$ ,  $R_\alpha \subseteq R_\beta$ .
  3. For all  $\alpha, \beta \in [1/2, 1)$  such that  $\alpha \geq \beta$ , if  $R_\beta$  is  $P$ -acyclic, then  $R_\alpha$  is also  $P$ -acyclic, and moreover, if  $R_\alpha$  is not  $P$ -acyclic, then  $R_\beta$  is also not  $P$ -acyclic.
  4.  $(a, b) \in R_{1/2}$  if and only if  $N[a, b] > N[b, a]$

**Proof.** Fix  $\alpha, \beta$  such that  $1/2 \leq \beta \leq \alpha < 1$ . We show the first result. Suppose  $(a, b) \in R_\alpha$ . Then,  $\Phi[a, b] = f(N[a, b], N[b, a]) > \alpha \geq 1/2$ . Since  $f(N[b, a], N[b, a]) = 1/2$  and  $f(n, N[b, a]) \leq f(N[b, a], N[b, a])$  for all  $n \leq N[b, a]$ , we have  $N[a, b] > N[b, a]$ . Then,  $\Phi[b, a] = f(N[b, a], N[a, b]) \leq f(N[b, a], N[b, a])$  and  $f(N[b, a], N[a, b]) \leq 1/2 \leq \alpha$ . Hence,  $(a, b) \notin R_\alpha$ . Therefore,  $R_\alpha$  is asymmetric.

We show the second result. If  $(a, b) \in R_\alpha$ , then  $\Phi[a, b] > \alpha \geq \beta$  and thus  $(a, b) \in R_\beta$ .

We show the third result. First, a subset of a P-acyclic binary relation is also P-acyclic. Therefore, by the third result, we have the latter part of the third result. Second, suppose  $R_\alpha$  has a P-cycle. By the first and second results,  $R_\beta$  is asymmetric and  $R_\alpha \subseteq R_\beta$ . Therefore,  $R_\beta$  also has the P-cycle and thus we have the former part of the third result.

We show the fourth result. First, suppose  $N[a, b] > N[b, a]$ . Then,

$$\Phi[a, b] = f(N[a, b], N[b, a]) > f(N[b, a], N[b, a]) = 1/2.$$

Second, suppose  $N[a, b] \leq N[b, a]$ . Then,

$$\Phi[a, b] = f(N[a, b], N[b, a]) \leq f(N[b, a], N[b, a]) = 1/2.$$

Therefore, we have the fourth result. **Q.E.D.**

## 4 Consistent Linear Orders

In this section, we arbitrarily fix a profile of preference relations of all individuals  $(R^i)_{i \in I}$  and do not explicitly consider it excepting an example. Instead of  $(R^i)_{i \in I}$ , we focus on  $R_\alpha$  constructed from it for all  $\alpha \in [1/2, 1)$ .

### 4.1 P-acyclic rules

First, we focus only on P-acyclic  $R_\alpha$ . Let  $\mathcal{P}_{AC} = \{R_\alpha \mid R_\alpha \text{ is P-acyclic}\}$ . Note that  $R_\alpha \in \mathcal{P}_{AC}$  if  $\alpha$  is sufficiently large (the Pareto rule). Thus, we can let

$$\alpha^* = \min_{\alpha \in [1/2, 1)} \{\alpha \mid R_\alpha \in \mathcal{P}_{AC}\}.$$

We have the following result.

#### Proposition 2

$$\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha) = \mathcal{C}(R_{\alpha^*}) \neq \emptyset.$$

**Proof.** First, let  $1 > \alpha \geq \beta \geq 0$ . By the third result of Lemma 7, if  $R_\beta$  is P-acyclic, then  $R_\alpha$  is also P-acyclic, and if  $R_\alpha$  has a P-cycle, then  $R_\beta$  also has a P-cycle. Therefore,  $R_\alpha$  is P-acyclic if and only if  $\alpha \geq \alpha^*$ . We have  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha) = \bigcap_{\alpha \geq \alpha^*} \mathcal{C}(R_\alpha)$ . Since  $R_\alpha$  is asymmetric,  $P(R_\alpha) = R_\alpha$ . Moreover, since  $R_{\alpha^*} \subseteq R_\alpha$  for all  $\alpha \in [\alpha^*, 1)$ ,  $\bigcup_{\alpha \geq \alpha^*} P(R_\alpha) = \bigcup_{\alpha \geq \alpha^*} R_\alpha = R_{\alpha^*}$ . Thus, by Theorem 1, we have  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha) = \mathcal{C}(R_{\alpha^*})$ . Finally, by Proposition 1,  $\mathcal{C}(R_{\alpha^*}) \neq \emptyset$ . **Q.E.D.**

Considering P-acyclic  $R_\alpha$  only is in a similar spirit to the Simpson-Kramer min-max method introduced in the last section.<sup>9</sup> There is a similar disadvantage to that of the method; that is, the set  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha)$  may include some improper linear orders. To show this fact, we reconsider Example 3. Then,  $\alpha^* = 2/3$  and  $R_\alpha = \emptyset$  for all  $\alpha \geq \alpha^* = 2/3$ . Therefore, any linear order is in  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha)$ . Since  $d$  is the Condorcet loser, this includes a linear order such that the Condorcet loser as the first best alternative. This is because, in this approach, to escape from a cycle, we ignore the fact that nine-of-five (less than two-of-three) individuals disprefer  $d$  to any other alternatives. Hence,  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha)$  may include some improper linear orders. Thus, next, we consider a smaller set of linear orders that does not include such improper linear orders.

## 4.2 Suzumura-consistent Closure

Next, we consider a procedure to use the Suzumura-consistent closure of  $R$ . As mentioned above, even if  $R$  is P-acyclic,  $S(R)$  is not. Therefore, we can consider a linear order that is consistent with  $S(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ . First, we consider a linear order that is consistent with  $S(R_{1/2})$ . Then, there may include some improper linear orders. To show this fact, we consider an example introduced by Suzumura (2012).

### Example 4

There are  $A = \{a, a', b, c\}$  and  $I = \{1, 2, 3\}$ . All individuals have a linear order preference on  $A$  and

$$R^1 : a, a', c, b, \quad R^2 : b, c, a, a', \quad R^3 : c, a, a', b.$$

---

<sup>9</sup>However, for any  $\hat{R}(: a_1, \dots, a_{|I|}) \in \bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha)$ ,  $a_1$  may not be the Simpson-Kramer min-max winner. The Simpson-Kramer min-max method is formally introduced in the last section.

Then,

$$R_{1/2} = \{(a, a'), (a, b), (a', b), (c, a), (c, a'), (b, c)\},$$

$T(R_{1/2}) = S(R_{1/2}) = \emptyset$ . Then,  $R \in \mathcal{C}(S(R_{1/2}))$  such that  $R : a', c, b, a$ , which may be improper linear order, because all individuals prefer  $a$  to  $a'$ .

Therefore, we consider a smaller set of linear orders called the  **$S$ -order set** denoted by  $\mathcal{S}$ ; that is,

$$\mathcal{S} = \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(S(R_\alpha)), \quad (6)$$

which includes only linear orders that are consistent with  $S(R_\alpha)$  for *all*  $\alpha \in [1/2, 1)$ .

**Theorem 2**

$$\mathcal{S} = \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \right) \subseteq \bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha) \quad (7)$$

**Proof.** By Lemmata 4 and 7,  $S(R_\alpha) \subseteq S(R_{1/2})$  for all  $\alpha \in [1/2, 1)$ . Since  $R_{1/2}$  is finite, we can let be finite binary relations  $R(1), R(2), \dots, R(I)$  such that for each  $\alpha \in [1/2, 1)$ ,  $S(R_\alpha) = R(i)$  for some  $i = 1, \dots, I$ , and  $R(1) \subseteq R(2) \subseteq \dots \subseteq R(I)$ , because of Lemmata 4 and 7. Moreover, by Lemma 1, since  $S(R_\alpha)$  is P-acyclic for all  $\alpha \in [1/2, 1)$ ,  $R(i)$  is P-acyclic for all  $i = 1, \dots, I$ . By Theorem 1, we have the equality of (7).

Since  $R_{\alpha^*}$  is P-acyclic,  $\mathcal{C}(R_{\alpha^*}) \neq \emptyset$ . Then, by Corollary 1,  $S(R_{\alpha^*}) = R_{\alpha^*}$ . Since  $R_{\alpha^*}$  is asymmetric,  $P(S(R_{\alpha^*})) = R_{\alpha^*}$ . Moreover, by Proposition 2,  $\bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha) = \mathcal{C}(R_{\alpha^*})$ . Now, if  $\hat{R} \in \mathcal{S}$ , then  $(a, b) \in P(S(R_{\alpha^*}))$  implies  $(a, b) \in \hat{R}$ . Thus,  $\hat{R} \in \mathcal{C}(R_{\alpha^*}) = \bigcap_{\alpha: R_\alpha \in \mathcal{P}_{AC}} \mathcal{C}(R_\alpha)$ . **Q.E.D.**

Next, to show the nonemptyness of (6), we consider the **ranked pairs method**, which is a rule to construct a linear order from  $(R^i)_{i \in I}$  and introduced by Tideman (1987).

Let  $\pi : A \times A \rightarrow \{1, 2, \dots, |A|^2\}$  be such that  $\pi(a, b) \neq \pi(c, d)$  for any  $(a, b) \neq (c, d)$  and, if  $\Phi[a, b] > \Phi[c, d]$ , then  $\pi(a, b) < \pi(c, d)$ . Let  $\Pi$  be a set of all possible  $\pi$ .

**Step 0** Fix  $\pi \in \Pi$ .



**Step**  $z$  ( $= 1, \dots, |A|^2$ ) Let  $(a, b) \in A \times A$  be such that  $\pi(a, b) = z$ . If  $a = b$  or  $\mathcal{B}([b, a], R_{z-1}^\pi) \neq \emptyset$ , then let  $R_z^\pi = R_{z-1}^\pi$ . Otherwise, then let  $R_z^\pi = R_{z-1}^\pi \cup \{(a, b)\}$ .

For the result of this algorithm, we let  $R_{|A|^2}^\pi = R^\pi$ .

**Proposition 3** For any  $\pi \in \Pi$ ,

$$R^\pi \in \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \right) = \mathcal{S}. \quad (8)$$

**Proof.** The asymmetry of  $R^\pi$  is trivial. First, we show the completeness of  $R^\pi$ . Suppose not; that is,  $(a, b) \notin R^\pi$  and  $(b, a) \notin R^\pi$ . Then,  $\mathcal{B}([b, a], R_{\pi(a,b)-1}^\pi) \neq \emptyset$  and  $\mathcal{B}([a, b], R_{\pi(b,a)-1}^\pi) \neq \emptyset$ .

In Step 1, there is no path from  $a$  to  $b$  or from  $b$  to  $a$  for  $R_1^\pi = \emptyset$ . On the other hand, since  $(a, b) \notin R^\pi$  and  $(b, a) \notin R^\pi$ , there are at least two paths from  $a$  to  $b$  and from  $b$  to  $a$  for  $R_{|A|^2}^\pi = R^\pi$  in Step  $|A|^2$ . Thus, they are made in the process. Let  $(a_1, a_2, \dots, a_m)$  be a path for  $R^\pi$  such that it is firstly made among the paths from  $a$  to  $b$  and from  $b$  to  $a$  in the process. Without loss of generality, suppose  $(a_1, a_2, \dots, a_m)$  is from  $a$  to  $b$ . Then, consider **Step**  $z$  such that a path  $(b_1, b_2, \dots, b_{m'})$  from  $b$  to  $a$  for  $R^\pi$  is newly made. Then, we can let  $\pi(b_l, b_{l+1}) = z$  for some  $l$ ; that is,  $R_z^\pi = R_{z-1}^\pi \cup \{(b_l, b_{l+1})\}$  and  $(b_k, b_{k+1}) \in R_{z-1}^\pi$  for all  $k \neq l$ . Then, since  $a_1 = b_{m'} = a$  and  $b_1 = a_m = b$ ,

$$(b_{l+1}, b_{l+2}, \dots, b_{m'} (= a_1), a_2, \dots, a_m (= b_1), \dots, b_l)$$

is a path for  $R_{z-1}^\pi$  from  $b_{l+1}$  to  $b_l$ ; that is,  $\mathcal{B}([b_{l+1}, b_l], R_{z-1}^\pi) \neq \emptyset$ . However, this contradicts  $R_z^\pi = R_{z-1}^\pi \cup \{(b_l, b_{l+1})\}$ . Therefore,  $R^\pi$  is complete.

Second, we show transitivity of  $R^\pi$ . Suppose  $(a, b), (b, c) \in R^\pi$ . Then, showing  $(c, a) \notin R^\pi$  is sufficient, because  $R^\pi$  is complete. First, suppose  $\pi(c, a) > \max\{\pi(a, b), \pi(b, c)\}$ . Then, since  $(a, b, c) \in \mathcal{B}([a, c], R_{\pi(c,a)-1}^\pi) \neq \emptyset$ ,  $(c, a) \notin R^\pi$ . Second, suppose  $\pi(b, c) > \max\{\pi(a, b), \pi(c, a)\}$ . Then, since  $\mathcal{B}([c, b], R_{\pi(b,c)-1}^\pi) = \emptyset$  and  $(a, b) \in R_{\pi(b,c)-1}^\pi$ ,  $(c, a, b) \notin \mathcal{B}([c, b], R_{\pi(b,c)-1}^\pi)$ . Thus,  $(c, a) \notin R_{\pi(b,c)-1}^\pi$  and  $(c, a) \notin R^\pi$ . Third, suppose  $\pi(a, b) > \max\{\pi(b, c), \pi(c, a)\}$ . Then, since  $\mathcal{B}([b, a], R_{\pi(a,b)-1}^\pi) = \emptyset$  and  $(b, c) \in R_{\pi(a,b)-1}^\pi$ ,  $(c, a) \notin R_{\pi(a,b)-1}^\pi$  and hence  $(c, a) \notin R^\pi$ . Therefore,  $R^\pi$  is transitive.

Finally, we show (8). Since we have already shown that  $R^\pi$  is a linear order, showing  $\bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \subseteq R^\pi$  for any  $\pi \in \Pi$  is sufficient. Let

$(a, b) \in P(S(R_\alpha))$  for some  $\alpha \in [1/2, 1)$ . By Lemma 3,  $(a, b) \in P(R_\alpha)$  and thus  $(a, b) \in R_\alpha$  and  $(b, a) \notin R_\alpha$ . Suppose not; that is,  $(a, b) \notin R^\pi$  for some  $\pi \in \Pi$ . Let  $z = \pi(a, b)$ . Then, we can let  $(a_1, a_2, \dots, a_m) \in \mathcal{B}([b, a], R_{z-1}^\pi) \neq \emptyset$ , where  $a_1 = b$  and  $a_m = a$ . By the construction,  $(a_l, a_{l+1}) \in R_{z-1}^\pi$  for all  $l = 1, \dots, m-1$  implies  $\pi(a_l, a_{l+1}) > \pi(a, b)$  and thus  $\Phi[a_l, a_{l+1}] \geq \Phi[a, b]$ . Since  $(a, b) \in R_\alpha$ ,  $\Phi[a_l, a_{l+1}] \geq \Phi[a, b] > \alpha$ . Thus,  $(a_l, a_{l+1}) \in R_\alpha$  for all  $l = 1, \dots, m-1$ . Hence  $(a_1, a_2, \dots, a_m) \in \mathcal{B}([b, a], R_\alpha)$ . Then, since  $(a, b) \in R_\alpha$ ,  $(b, a) \in S(R_\alpha)$  but this contradicts  $(a, b) \in P(S(R_\alpha))$ . **Q.E.D.**

This result implies that any linear order obtained by the ranked pairs method is consistent with the Suzumura-consistent closures of all supermajority binary relations. We immediately have the following result.

**Corollary 2**  $\mathcal{S} \neq \emptyset$ .

### 4.3 Transitive Closure

Next, we consider a method to use the transitive closure of  $R$ . We consider  $\bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$  and show that it is equivalent to the binary relation constructed via the Schulze method, which is introduced below.

For  $(a_1, a_2, \dots, a_m) \in \mathcal{B}[(a, b), R_{1/2}]$ , let the **strength** of  $(a_1, a_2, \dots, a_m)$  be

$$ST((a_1, \dots, a_m)) = \min_{j=1, \dots, m-1} [\Phi[a_j, a_{j+1}]].$$

Furthermore, let

$$\begin{aligned} B[a, b] &= \max_{p \in \mathcal{B}[(a, b), R_{1/2}]} ST(p) \text{ if } \mathcal{B}[(a, b), R_{1/2}] \neq \emptyset \\ &= 0 \text{ if } \mathcal{B}[(a, b), R_{1/2}] = \emptyset, \end{aligned}$$

which represents the strength of the strongest path from  $a$  to  $b$ . Let  $R^{Sc}$  be a binary relation obtained by the **Schulze method**; that is,  $(a, b) \in R^{Sc}$  if and only if  $B[a, b] > B[b, a]$ . Note that, as shown by Schulze (2011),  $R^{Sc}$  is a partial order. Therefore, by Lemma 1 and Proposition 1,  $\mathcal{C}(R^{Sc}) \neq \emptyset$ .

We have the following result.

**Proposition 4**

$$R^{Sc} = \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)).$$

**Proof.** First, we show  $R^{Sc} \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ . To show this, we firstly show the following result.

**Claim 1** Suppose  $B[b, a] \in [1/2, 1)$ . First,  $\mathcal{B}[(b, a), R_{B[b, a]}] = \emptyset$ . Second, if  $B[a, b] > B[b, a]$ , then  $\mathcal{B}[(a, b), R_{B[b, a]}] \neq \emptyset$ .

**Proof of Claim 1.** Suppose  $B[b, a] \in [1/2, 1)$ . First, let  $(a_1, a_2, \dots, a_m)$  be an arbitrary sequence of distinct alternatives such that  $a_1 = b$  and  $a_m = a$ . By the definition of  $B[b, a]$ ,  $B[b, a] \geq \Phi[a_j, a_{j+1}]$  for some  $j = 1, \dots, m-1$ . Therefore,  $(a_1, a_2, \dots, a_m) \notin \mathcal{B}[(b, a), R_{B[b, a]}]$  and thus  $\mathcal{B}[(b, a), R_{B[b, a]}] = \emptyset$ .

Second, suppose  $B[a, b] > B[b, a]$ . By the definition of  $B[a, b]$ , there is a sequence of distinct alternatives  $(a_1, a_2, \dots, a_m)$  such that  $a_1 = a$ ,  $a_m = b$  and  $\Phi[a_j, a_{j+1}] \geq B[a, b] > B[b, a]$  for all  $j = 1, \dots, m-1$ . Then,  $(a_1, a_2, \dots, a_m) \in \mathcal{B}[(a, b), R_{B[b, a]}]$ . Thus,  $\mathcal{B}[(a, b), R_{B[b, a]}] \neq \emptyset$ . **Q.E.D.**

Now, we show  $R^{Sc} \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ . Let  $(a, b) \in R^{Sc}$ ; that is,  $B[a, b] > B[b, a]$ . First, suppose  $B[b, a] < 1/2$ . Then, by the definition,  $B[b, a] = 0$  and  $\mathcal{B}[(b, a), R_{1/2}] = \emptyset$ . Moreover, by the second result of Lemma 7,  $\mathcal{B}[(b, a), R_\alpha] = \emptyset$  for all  $\alpha \in [1/2, 1)$ . Thus,  $(b, a) \notin T(R_\alpha)$  for all  $\alpha \in [1/2, 1)$  and  $(b, a) \notin \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ . Next, we show  $(a, b) \in T(R_{1/2})$ . Since  $\mathcal{B}[(b, a), R_{1/2}] = \emptyset$ ,  $(b, a) \notin R_{1/2}$ . By the fourth result of Lemma 7,  $N[a, b] > N[b, a]$  and thus,  $(a, b) \in R_{1/2} \subseteq T(R_{1/2})$ . Thus, in this case,

$$(a, b) \in P(T(R_{1/2})) \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)).$$

Second, suppose  $B[b, a] \in [1/2, 1)$ . By Claim 1,  $\mathcal{B}[(b, a), R_{B[b, a]}] = \emptyset$  and  $\mathcal{B}[(a, b), R_{B[b, a]}] \neq \emptyset$ . That is,  $(a, b) \in T(R_{B[b, a]})$  and  $(b, a) \notin T(R_{B[b, a]})$ . Since  $B[b, a] \in [1/2, 1)$ , we have

$$(a, b) \in P(T(R_{B[b, a]})) \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)).$$

Therefore,  $R^{Sc} \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ .

Next, we show  $R^{Sc} \supseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ . Fix some  $\alpha \in [1/2, 1)$  and let  $(a, b) \in P(T(R_\alpha))$ ; that is,  $(a, b) \in T(R_\alpha)$  and  $(b, a) \notin T(R_\alpha)$ . Toward a contradiction, suppose  $(a, b) \notin R^{Sc}$ . Then,  $B[a, b] \leq B[b, a]$ . First, suppose  $B[a, b] < B[b, a]$ . Then  $(b, a) \in R^{Sc}$ . Since  $R^{Sc} \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ ,  $(b, a) \in P(T(R_{\alpha'}))$ ; that is,  $(b, a) \in T(R_{\alpha'})$  and  $(a, b) \notin T(R_{\alpha'})$ , for some  $\alpha' \in [1/2, 1)$ . First, suppose  $\alpha' < \alpha$ . Then, by Lemma 4 and the second

result of Lemma 7,  $T(R_\alpha) \subseteq T(R_{\alpha'})$ , which contradicts  $(a, b) \in T(R_\alpha)$  and  $(a, b) \notin T(R_{\alpha'})$ . Second, suppose  $\alpha' \geq \alpha$ . Then,  $T(R_{\alpha'}) \subseteq T(R_\alpha)$ , which contradicts  $(b, a) \notin T(R_\alpha)$  and  $(b, a) \in T(R_{\alpha'})$ . Therefore,  $(a, b) \in R^{Sc}$  and  $R^{Sc} \supseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ .

Second, suppose  $B[a, b] = B[b, a]$ . Then, since  $(a, b) \in T(R_\alpha)$ ,  $\mathcal{B}[(a, b), R_\alpha] \neq \emptyset$ . By Claim 1,  $\alpha < B[a, b] = B[b, a]$ . Then, by the definition of  $B[b, a]$ , there is a sequence of alternatives  $(a_1, a_2, \dots, a_m)$  such that  $a_1, a_2, \dots, a_m$  are distinct,  $a_1 = b$  and  $a_m = a$ , where  $\Phi[a_j, a_{j+1}] \geq B[b, a] > \alpha$  for all  $j = 1, \dots, m-1$ . Then,  $(a_1, a_2, \dots, a_m) \in \mathcal{B}[(b, a), R_\alpha]$  that contradicts  $(b, a) \notin T(R_\alpha)$ . **Q.E.D.**

Let

$$\mathcal{T} = \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(T(R_\alpha))$$

be the ***T*-order set**. We have the following result.

**Theorem 3**

$$\mathcal{T} = \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) \right) = \mathcal{C}(R^{Sc}).$$

**Proof.** First, we can show

$$\mathcal{T} = \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) \right)$$

by following the proof of the corresponding part of Theorem 2. Moreover, by Proposition 4,

$$\mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) \right) = \mathcal{C}(R^{Sc}).$$

**Q.E.D.**

This result can be rewritten as  $R \in \mathcal{C}(R^{Sc})$  if and only if  $R \in \mathcal{C}(T(R_\alpha))$  for all  $\alpha \in [1/2, 1)$ . Thus, any linear order that is consistent with the binary relation constructed by the Schulze method is also consistent with the transitive closure of the binary relation constructed by any supermajority

rules. Moreover, any linear order that is consistent with the transitive closures of all supermajority binary relations is also consistent with the binary relation constructed by Schulze method. Therefore, the Schulze method is interpreted as that obtains a linear order that is consistent with the binary relations derived by all supermajority rules and modified via the transitive closure procedure when they have some cycles.

By Theorem 2, we immediately have the following results. First, as shown by Schulze (2011),  $R^{Sc}$  is transitive and thus we have the following result.

**Corollary 3**  $\mathcal{T} \neq \emptyset$ .

Second, by Lemma 3,

$$\bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \subseteq \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)).$$

Therefore, by Theorems 1 and 2, we have the following result.

**Corollary 4**  $\mathcal{T} \subseteq \mathcal{S}$ .

By Proposition 4 and Corollary 4, the linear orders that are derived by the Schulze method and the ranked pairs method are in  $\mathcal{S}$ . Thus, the both methods belongs to the class of methods that choose one from  $\mathcal{S}$ . However, any linear orders that are derived by the ranked pairs method may not be in  $\mathcal{T}$ . Furthermore, there may be some linear orders in  $\mathcal{S}$  that are derived by neither of them. We show these facts by providing an example in the Appendix.

## 5 Properties of social choice correspondence

In this section, we consider a social choice correspondence  $F_A : \mathcal{W}^{|I|} \rightarrow 2^{\mathcal{L}}$ , where  $\mathcal{W}$  and  $\mathcal{L}$  are the sets of all weak orders and all linear orders, respectively. Let  $\mathbf{R} = (R^i)_{i \in I} \in \mathcal{W}^{|I|}$  represent a profile of the preference of individuals. We focus only on  $F_A(\mathbf{R})$  is either  $\mathcal{S}$  or  $\mathcal{T}$  for  $\mathbf{R}$ . Thus, let  $F_A^S(\mathbf{R}) = \mathcal{S}$  and  $F_A^T(\mathbf{R}) = \mathcal{T}$  for all  $\mathbf{R} \in \mathcal{W}^{|I|}$ .<sup>10</sup> Note that  $F_A^T(\mathbf{R}) \subseteq F_A^S(\mathbf{R})$  for all  $\mathbf{R} \in \mathcal{W}^{|I|}$ , because of Corollary 4. We show that these social choice correspondences satisfy four properties.

---

<sup>10</sup>The correspondences are also dependent on  $f(n, m)$ . Here, we fix  $f(n, m)$  such as (4) or (5).

## 5.1 Extended Condorcet criterion

We introduce the property introduced by Truchon (1998) and called the extended Condorcet criterion.

Let  $\mathcal{A}(\mathbf{R})$  be the set of all partitions of  $A$  for  $\mathbf{R} \in \mathcal{W}^{|I|}$  such that for all

$$\mathbf{A} = \{A_1, \dots, A_X\} \in \mathcal{A}(\mathbf{R}),$$

all  $A_x, A_{x'} \in \mathbf{A}$  with  $x < x'$ , all  $a \in A_x$  and all  $a' \in A_{x'}$ ,  $N[a, a'] > N[a', a]$ . A linear order  $R'$  is said to satisfy the **extended Condorcet criterion** for  $\mathbf{R}$  if for all  $\mathbf{A} \in \mathcal{A}(\mathbf{R})$ , all  $A_x, A_{x'} \in \mathbf{A}$  with  $x < x'$ , all  $a \in A_x$  and all  $a' \in A_{x'}$ ,  $(a, a') \in \hat{R}$ .

Truchon (1998) shows that any linear order that is derived by the Kemeny-Young method also satisfies the extended Condorcet criterion for any  $\mathbf{R} \in \mathcal{W}^{|I|}$ .

**Proposition 5** *All elements in  $F_A^S(\mathbf{R})$  satisfy the extended Condorcet criterion for  $\mathbf{R}$ .*

**Proof.** Fix  $\mathbf{A} = \{A_1, \dots, A_X\} \in \mathcal{A}(\mathbf{R})$ . Let  $R' \in F_A^S(\mathbf{R})$ . Suppose not; that is,  $R'$  does not satisfy the extended Condorcet criterion. Then, there exists  $(a', a) \in R'$  such that  $a \in A_x$  and  $a' \in A_{x'}$  with  $x < x'$ . Note that  $N[a, a'] > N[a', a]$ . By the fourth result of Lemma 7,  $N[a, a'] > N[a', a]$  implies  $(a, a') \in R_{1/2}$  and thus  $(a, a') \in S(R_{1/2})$ . If  $(a, a') \in P(S(R_{1/2}))$ , then  $(a', a) \notin R'$ . We have  $(a, a') \notin P(S(R_{1/2}))$  and thus  $(a', a) \notin S(R_{1/2})$ . Since  $R_\alpha \subseteq S(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ ,  $(a', a) \in S(R_{1/2}) \setminus R_{1/2}$ . Therefore,  $(a', a) \in T(R_{1/2})$ ; that is, there is a path  $(a_1, a_2, \dots, a_m) \in \mathcal{B}[(a', a), R_{1/2}]$ , where  $a_1 = a'$ , and  $a_m = a$  and  $(a_l, a_{l+1}) \in R_{1/2}$  for all  $l = 1, 2, \dots, m-1$ . By the fourth result of Lemma 7,  $(a_{m-1}, a_m) \in R_{1/2}$  implies  $N[a_{m-1}, a_m] > N[a_m, a_{m-1}]$ . Hence we can let  $a_{m-1} \in A_{x_1}$  and  $a_m \in A_{x_0}$  for  $x_1 > x_0$ . Likewise,  $a_{m-z} \in A_{x_z}$  for  $z = 1, \dots, m-1$  and  $x_{m-1} < \dots < x_2 < x_1 < x_0$ . Since  $a_m = a$  and  $a_1 = a'$ ,  $x_0 = x$  and  $x_{m-1} = x'$ , but this contradicts  $x < x'$ . **Q.E.D.**

Now, we say that a social choice correspondence  $F_A$  satisfies the **extended Condorcet criterion** if for all  $\mathbf{R} \in \mathcal{W}^{|I|}$ , each element in  $F_A(\mathbf{R})$  satisfies the extended Condorcet criterion for  $\mathbf{R}$ .

The following results are immediate from Proposition 5 and  $F_A^T(\mathbf{R}) \subseteq F_A^S(\mathbf{R})$  for all  $\mathbf{R} \in \mathcal{W}^{|I|}$ .

**Corollary 5** *Both  $F_A^S$  and  $F_A^T$  satisfy the extended Condorcet criterion.*

**Corollary 6** *For any  $\mathbf{R} \in \mathcal{W}^{|I|}$  and  $\hat{R} (: a_1, \dots, a_{|I|}) \in F_A^S(\mathbf{R}) (\supseteq F_A^T(\mathbf{R}))$ ,*

1. if  $a$  is the Condorcet for winner  $\mathbf{R}$ , then  $a = a_1$ ,
2. if  $b$  is the Condorcet loser for  $\mathbf{R}$ , then  $b = a_{|I|}$ .

## 5.2 Pareto principle

Next, we consider the Pareto principle. A linear order  $\hat{R}$  is said to satisfy the **Pareto principle** for  $\mathbf{R}$  if for any  $a, b \in A$  such that  $(b, a) \notin P(R^i)$  for all  $i \in I$  and  $(a, b) \in P(R^i)$  for some  $i \in I$ ,  $(a, b) \in \hat{R}$ .<sup>11</sup> A social choice correspondence  $F_A$  satisfies the **Pareto principle** if  $\hat{R}$  satisfies the **Pareto principle** for  $\mathbf{R}$  for all  $\hat{R} \in F_A(\mathbf{R})$  and all  $\mathbf{R} \in \mathcal{W}^{|I|}$ .

First, we show the following result.

**Lemma 8** *If  $\alpha \in [1/2, 1)$  is sufficiently large, then  $R_\alpha$  is P-acyclic.*

**Proof.** We fix  $\alpha \in [f(|I| - 1, 1), 1)$ . Then,  $(a, b) \in R_\alpha$  implies  $f(N[a, b], N[b, a]) > f(|I| - 1, 1)$ . Therefore,  $N[a, b] = |I|$  and/or  $N[b, a] = 0$ . If  $N[a, b] = |I|$ ; that is, if all individuals prefer  $a$  to  $b$ , then  $N[b, a] = 0$ . Therefore,  $(a, b) \in R_\alpha$  implies  $N[b, a] = 0$ ; that is, no individual prefers  $b$  to  $a$ . Moreover, since  $f(0, 0) = 1/2$ ,  $N[a, b] \geq 1$ ; that is, at least one individual prefers  $a$  to  $b$ .

Now, we show that  $R_\alpha$  is P-acyclic. Suppose that there is  $(a_0, a_1, \dots, a_m) \in \mathcal{B}[(k, j), P(R_\alpha)]$ . We show  $(a_K, a_0) \notin P(R_\alpha)$ . Since  $(a_0, a_1) \in R_\alpha$ , there is at least one individual  $i$  such that  $(a_0, a_1) \in R^i$ . Moreover, since  $N[b, a] = 0$ ,  $(a_l, a_{l-1}) \notin P(R^i)$  for all  $l = 1, \dots, m$ . We show  $(a_0, a_2) \in R^i$ . If  $(a_1, a_2) \in R^i$ , then  $(a_0, a_2) \in R^i$  because  $R^i$  is transitive. Suppose  $(a_1, a_2) \notin R^i$ . Then,  $(a_2, a_1) \notin P(R^i)$  implies  $(a_2, a_1) \notin R^i$ . Moreover, toward a contradiction, we assume  $(a_0, a_2) \notin R^i$ . Since  $R^i$  is negatively transitive,  $(a_0, a_2) \notin R^i$  and  $(a_2, a_1) \notin R^i$  imply  $(a_0, a_1) \notin R^i$ , which is a contradiction. Therefore,  $(a_0, a_2) \in R^i$ . Likewise, we can show  $(a_0, a_K) \in R^i$  and thus  $(a_K, a_0) \notin R_\alpha$ . Therefore,  $R_\alpha$  is P-acyclic. **Q.E.D.**

We consider the two cases. First, suppose that each individual has a linear order preference.

**Proposition 6** *Suppose that  $R^i$  is a linear order for all  $i \in I$ . For any  $\hat{R} \in \mathcal{C}(S(R_\alpha))$  for sufficiently large  $\alpha \in [1/2, 1)$ ,  $\hat{R}$  satisfies the Pareto principle.*

<sup>11</sup>There are several kinds of the Pareto principle. Our definition is in the spirit of (i) of the ‘‘strong Pareto’’ defined by Bossert and Suzumura (2010, P.129). Schulze (2018) also considers the Pareto principle, but his definitions are different from ours.

**Proof.** We fix  $\alpha \in [f(|I| - 1, 1), 1)$ . Suppose that  $(b, a) \notin P(R^i)$  for all  $i \in I$  and  $(a, b) \in P(R^i)$  for some  $i \in I$ ,  $(a, b) \in \hat{R}$ . If  $R^i$  is a linear order, then  $(b, a) \notin P(R^i)$  implies that  $(a, b) \in P(R^i)$ . Therefore,  $N[b, a] = 0$  and  $N[a, b] = |I|$ . Then,  $\Phi[a, b] = f(|I|, 0) = 1$  and  $\Phi[b, a] = f(0, |I|) < 1/2$ . Thus,  $(a, b) \in R_\alpha$  and  $(b, a) \notin R_\alpha$ . By Lemmata 1 and 8,  $R_\alpha$  is Suzumura-consistent and thus  $S(R_\alpha) = R_\alpha$ . Then,  $(a, b) \in P(S(R_\alpha)) = P(R_\alpha)$  and hence  $(a, b) \in \hat{R}$ . **Q.E.D.**

Second, suppose that each individual has a weak order preference. In this case, some additional assumption is necessary.

**Proposition 7** *Suppose that  $R^i$  is a weak order for all  $i \in I$ , and  $f(n, 0) = 1 > f(m, 1)$  for any  $n > 0$  and any  $m \geq 0$ . For any  $\hat{R} \in \mathcal{C}(S(R_\alpha))$  for sufficiently large  $\alpha \in [1/2, 1)$ ,  $\hat{R}$  satisfies the Pareto principle.*

**Proof.** We fix  $\alpha \in [f(|I| - 1, 1), 1)$ . Then, we have  $f(m, n) \leq \alpha$  and  $f(n, 0) = 1 > \alpha$  for all  $m \in [0, |I| - 1]$  and all  $n > 0$ . Then,  $(a, b) \in R_\alpha$  if and only if  $(b, a) \notin R^i$  for all  $i \in I$  and  $(a, b) \in R^i$  for some  $i \in I$ .

By Lemmata 1, 8 and 9,  $R_\alpha$  is Suzumura-consistent and thus  $S(R_\alpha) = R_\alpha$ . By Proposition 1, we can let  $\hat{R} \in \mathcal{C}(R_\alpha) = \mathcal{C}(S(R_\alpha))$ .

Now, suppose that there are  $a, b \in A$  such that  $(b, a) \notin P(R^i)$  for all  $i \in I$  and  $(a, b) \in P(R^i)$  for some  $i \in I$ . Then, since  $N[b, a] = 0$  and  $N[a, b] > 0$ ,  $(a, b) \in P(R_\alpha)$ . Hence  $(a, b) \in \hat{R}$ . **Q.E.D.**

Note that  $f(n, 0) = 1 > f(m, 1)$  is satisfied if (4) holds, but it is not satisfied if (5) holds. We provide an example with (5) (without  $f(n, 0) = 1 > f(m, 1)$ ) such that some linear orders on  $A$  in both  $\mathcal{T}$  and  $\mathcal{S}$  does not satisfy the Pareto principle.

### Example 5

We modify Example 4. There are  $A = \{a, a', b, c\}$  and  $I = \{1, 2, 3\}$ . All individuals have a weak order preference on  $A$  and

$$R^1 : [a, a'], c, b, \quad R^2 : b, c, [a, a'], \quad R^3 : c, a, a', b,$$

meaning that  $a$  and  $a'$  are indifferent alternatives for individuals 1 and 2. Note that  $(a', a) \notin P(R^i)$  for all  $i \in I$  and  $(a, a') \in P(R^3)$ . Thus, for any linear order  $\hat{R}$  that satisfies the Pareto principle,  $(a, a') \in \hat{R}$ .



Suppose (5) holds. Then, for  $\alpha \in [1/2, 2/3]$  and  $\alpha' \in (2/3, 1)$ ,

$$\begin{aligned} R_\alpha &= \{(a, a'), (a, b), (a', b), (c, a), (c, a'), (b, c)\}, \\ P(T(R_\alpha)) &= P(S(R_\alpha)) = \emptyset, \\ R_{\alpha'} &= T(R_{\alpha'}) = S(R_{\alpha'}) = \emptyset, \end{aligned}$$

Any linear orders on  $A$  are in both  $\mathcal{T}$  and  $\mathcal{S}$ . Thus, some linear orders on  $A$  in both  $\mathcal{T}$  and  $\mathcal{S}$  such as  $\hat{R} : a', c, b, a$  do not satisfy the Pareto principle.

On the other hand, suppose (4) holds. Then, for  $\alpha \in [1/2, 2/3]$  and  $\alpha' \in (2/3, 1)$ ,

$$\begin{aligned} R_\alpha &= \{(a, a'), (a, b), (a', b), (c, a), (c, a'), (b, c)\}, \\ P(T(R_\alpha)) &= P(S(R_\alpha)) = \emptyset, \\ R_{\alpha'} &= T(R_{\alpha'}) = S(R_{\alpha'}) = \{(a, a')\}, \end{aligned}$$

Thus, in this case, any linear order  $\hat{R} \in \mathcal{S} (= \mathcal{T})$  satisfies  $(a, a') \in \hat{R}$ .

We have the following result.

**Corollary 7** *If  $f(n, 0) = 1 > f(m, 1)$  for any  $n > 0$  and any  $m \geq 0$ , then both  $F_A^S$  and  $F_A^T$  satisfy the Pareto principle.*

### 5.3 Independence of clones

We consider the next property called the **independence of clones**, which is in the same spirit with that introduced by Tideman (1987).

We let  $b' \in A'(b)$  be a **clone** of  $b \in A$ . For notational convenience, let  $b \in A'(b)$ . Let  $R_{cl}^i$  be a binary relation representing the preference of  $i$  over  $A_{cl} = A \cup A'(b) \setminus \{b\}$  such that  $R^i \subseteq R_{cl}^i$ , for all  $a \in A$  and all  $b' \in A'(b)$ ,  $(a, b) \in R^i$  if and only if  $(a, b') \in R_{cl}^i$ , and  $(b, a) \in R^i$  if and only if  $(b', a) \in R_{cl}^i$ .

Let  $R_{\alpha, cl}$  be the binary relation constructed from  $\mathbf{R}_{cl} = (R_{cl}^i)_{i \in I}$  such that  $(d, e) \in R_{\alpha, cl}$  if and only if

$$\Phi_{cl}[d, e] = f(N_{cl}[d, e], N_{cl}[e, d]) > \alpha,$$

where

$$N_{cl}[d, e] = |\{i \in I \mid (d, e) \in P(R_{cl}^i)\}|.$$

Then, for all  $a \in A$  and all  $b' \in A'(b)$ ,  $\Phi[a, b] = \Phi_{cl}[a, b']$  and  $\Phi[b, a] = \Phi_{cl}[b', a]$  and therefore,  $(a, b) \in R_\alpha$  if and only if  $(a, b') \in R_{\alpha, cl}$ , and  $(b, a) \in R_\alpha$  if and only if  $(b, a') \in R_{\alpha, cl}$ .

**Lemma 9** For all  $a \in A$  and all  $b' \in A'(b)$ ,  $(a, b) \in T(R_\alpha)$  if and only if  $(a, b') \in T(R_{\alpha,cl})$ , and  $(b, a) \in T(R_\alpha)$  if and only if  $(b, a') \in T(R_{\alpha,cl})$  and moreover,  $(a, b) \in S(R_\alpha)$  if and only if  $(a, b') \in S(R_{\alpha,cl})$ , and  $(b, a) \in S(R_\alpha)$  if and only if  $(b', a) \in S(R_{\alpha,cl})$ .

**Proof.** We show  $(a, b) \in T(R_\alpha)$  if and only if  $(a, b') \in T(R_{\alpha,cl})$ . Since the only-if part is trivial, we show the if-part. Suppose  $(a, b') \in T(R_{\alpha,cl})$ . Then, there is a path  $(a'_1, a'_2, \dots, a'_m)$  from  $a'_1 = a$  to  $a'_m = b'$  such that  $m \geq 2$ ,  $(a'_l, a'_{l+1}) \in R_{\alpha,cl}$  for all  $l = 1, 2, \dots, m-1$ . We let  $a'_l \in A'(b)$  where  $l$  is the smallest among  $\{a'_1, a'_2, \dots, a'_m\} \cap A'(b)$ . Then, there is a path  $(a'_1, a'_2, \dots, a'_{l-1}, b)$  from  $a$  to  $b$  and thus  $(a, b) \in T(R_\alpha)$ . Since the proof is similar, we omit to prove  $(b, a) \in T(R_\alpha)$  if and only if  $(b, a') \in T(R_{\alpha,cl})$ .

Since the proofs of the remaining parts are also trivial, we omit them.

**Q.E.D.**

We say that a linear order  $\hat{R}$  on  $A$  is a **shrink** of a linear order  $\hat{R}_{cl}$  on  $A \cup A'(b) \setminus \{b\}$  if (i) there is  $b' \in A'(b)$  such that for all  $a \in A \setminus \{b\}$  satisfying  $(a, b') \in \hat{R}_{cl}$ ,  $(a, b) \in \hat{R}$  and (ii)

$$\hat{R} = \hat{R}_{cl} \cap ((A \setminus \{b\}) \times (A \setminus \{b\})).$$

In words, to construct a shrink, we first replace the position of an alternative  $A'(b)$  to  $b$  in  $\hat{R}_{cl}$  and second eliminate  $A \setminus \{b\}$  from  $\hat{R}_{cl}$ .

For example, let  $A = \{a, b, c, d\}$  and  $A'(b) = \{b, b', b''\}$ ,  $\hat{R}_{cl} : a, b, c, b', d, b''$ . Then, there are three shrinks of  $\hat{R}_{cl}$

$$\hat{R} : a, b, c, d, \quad \hat{R}' : a, c, b, d \quad \text{and} \quad \hat{R}'' : a, c, d, b$$

because  $b'$  and  $b''$  are the clones of  $b$ .

We say that  $F_A(\mathbf{R})$  satisfies the **independence of clones** if  $\hat{R} \in F_A(\mathbf{R})$  if and only if it is a shrink of some  $\hat{R}_{cl} \in F_{A_{cl}}(\mathbf{R}_{cl})$ .

First, we show the following result.

**Proposition 8** 1.  $\hat{R} \in \mathcal{C}(T(R_\alpha))$  if and only if it is a shrink of  $\hat{R}_{cl} \in \mathcal{C}(T(R_{\alpha,cl}))$ .

2.  $\hat{R} \in \mathcal{C}(S(R_\alpha))$  if and only if it is a shrink of  $\hat{R}_{cl} \in \mathcal{C}(S(R_{\alpha,cl}))$ .

**Proof.** First, let  $\hat{R}$  be a shrink of  $\hat{R}_{cl} \in \mathcal{C}(T(R_{\alpha,cl}))$  where  $\hat{R}_{cl} : a_1, a_2, \dots, a_{|I|+|A'(b)|-1}$ . We show  $\hat{R} \in \mathcal{C}(T(R_\alpha))$ . Let  $a_l \in A'(b)$  such that for all  $a \in A \setminus \{b\}$  satisfying  $(a, a_l) \in \hat{R}_{cl}$ ,  $(a, b) \in \hat{R}$ . Then, By Lemma

9, for any  $a_k, a_{k'} \in A \setminus \{b\}$  such that  $1 \leq k < k' \leq |I| + |A'(b)| - 1$ ,  $(a_{k'}, a_k) \notin P(T(R_{\alpha,cl}))$  implies  $(a_k, a_{k'}) \notin P(T(R_{\alpha,cl}))$ . Further, suppose  $k < l < k'$ . Then,  $(a_l, a_k) \notin P(T(R_{\alpha,cl}))$  and  $(a_{k'}, a_l) \notin P(T(R_{\alpha,cl}))$  imply  $(b, a_k) \notin P(T(R_{\alpha,cl}))$  and  $(a_{k'}, b) \notin P(T(R_{\alpha,cl}))$ , respectively, because  $a_l$  is a clone of  $b$ .

Second, let  $\hat{R} \in \mathcal{C}(T(R_\alpha))$  be  $\hat{R} : a_1, a_2, \dots, a_{|I|}$  and  $a_l = b$ . We show that  $\hat{R}$  is a shrink of some  $\hat{R}_{cl} \in \mathcal{C}(T(R_{\alpha,cl}))$ . For  $R_{\alpha,cl}$ , we derive a consistent linear order of  $T(R_{\alpha,cl})$  where the set of alternatives is only  $A'(b)$ . Since  $T(R_{\alpha,cl})$  is acyclic, we can derive such a linear order. Let  $R' : b_1, \dots, b_{|A'(b)|}$  be one of such linear orders on  $A'(b)$ .

We consider  $\hat{R}_{cl} : a'_1, a'_2, \dots, a'_{|I|+|A'(b)|-1}$  such that (i)  $a'_k = a_k$  for all  $k = 1, \dots, l-1$ , (ii)  $a'_{l+z-1} = b_z$  for  $z = 1, \dots, |A'(b)|$  and (iii)  $a_k = a'_{k+|A'(b)|-1}$  for  $k = l+1, \dots, |I|$ . Then, by Lemma 9,  $\hat{R}_{cl} \in \mathcal{C}(T(R_{\alpha,cl}))$ .

Since we can similarly show the second result, we omit the proof. **Q.E.D.**

By Proposition 8, we immediately have the following result.

**Corollary 8** *Both  $F_A^S$  and  $F_A^T$  satisfy the independence of clones.*

Note that this corollary does *not* imply that any social choice function that chooses a linear order in the  $S$ -order set satisfies the independence of clones. In fact, Zavist and Tideman (1989) show that the ranked pairs method without any restriction on  $\Pi$  does not satisfy the independence of clones.<sup>12</sup>

## 5.4 Reversal symmetry

In this subsection, we consider the reversal symmetry, which is originally introduced by Saari (1994).

Fix  $a, b \in A$ . Let  $R_{a \leftrightarrow b}^i$  such that  $(b, a) \in R_{a \leftrightarrow b}^i$  if  $(a, b) \in R^i$  and  $(a, b) \in R_{a \leftrightarrow b}^i$  if  $(b, a) \in R^i$ . Moreover, let  $R_{\alpha, a \leftrightarrow b}$  be the binary relation constructed from a preference profile  $\mathbf{R}_{a \leftrightarrow b} = (R_{a \leftrightarrow b}^i)_{i \in I}$  such that  $(d, e) \in R_{\alpha, a \leftrightarrow b}$  if and only if

$$\Phi_{a \leftrightarrow b}[d, e] = f(N_{a \leftrightarrow b}[d, e], N_{a \leftrightarrow b}[e, d]) > \alpha,$$

where

$$N_{a \leftrightarrow b}[d, e] = |\{i \in I \mid (d, e) \in P(R_{a \leftrightarrow b}^i)\}|.$$

---

<sup>12</sup>On the other hand, they also show that if  $\Pi$  is restricted to be impartial, then this method satisfies the clone independence.

Then,  $(a, x) \in T(R_\alpha) \Leftrightarrow (b, x) \in T(R_{\alpha, a \leftrightarrow b})$ ,  $(x, a) \in T(R_\alpha) \Leftrightarrow (x, b) \in T(R_{\alpha, a \leftrightarrow b})$ , and  $(x, b) \in T(R_\alpha) \Leftrightarrow (x, a) \in T(R_{\alpha, a \leftrightarrow b})$ , for all  $x \in A \setminus \{a, b\}$ . Moreover,  $(a, b) \in T(R_\alpha) \Leftrightarrow (b, a) \in T(R_{\alpha, a \leftrightarrow b})$ ,  $(b, a) \in T(R_\alpha) \Leftrightarrow (a, b) \in T(R_{\alpha, a \leftrightarrow b})$  and  $(x, y) \in T(R_\alpha) \Leftrightarrow (x, y) \in T(R_{\alpha, a \leftrightarrow b})$  for all  $x, y \in A \setminus \{a, b\}$ . The same outcomes are obtained even if  $T$  is replaced by  $S$ ; that is,  $(a, x) \in S(R_\alpha) \Leftrightarrow (b, x) \in S(R_{\alpha, a \leftrightarrow b})$ ,  $\dots$ .

We say that  $F_A$  satisfies the **reversal symmetry** if  $\hat{R} \in F_A(\mathbf{R})$  if and only if  $\hat{R}_{a \leftrightarrow b} \in F_A(\mathbf{R}_{a \leftrightarrow b})$ , where  $\hat{R}$  and  $\hat{R}_{a \leftrightarrow b}$  such that  $(a, b) \in \hat{R}$  if and only if  $(b, a) \in \hat{R}_{a \leftrightarrow b}$  and  $\hat{R} \setminus \{(a, b)\} = \hat{R}_{a \leftrightarrow b} \setminus \{(b, a)\}$ .

First, we show the following result.

**Proposition 9** *Let  $\hat{R}$  and  $\hat{R}_{a \leftrightarrow b}$  be linear orders on  $A$  such that  $(a, b) \in \hat{R}$  if and only if  $(b, a) \in \hat{R}_{a \leftrightarrow b}$  and  $\hat{R} \setminus \{(a, b)\} = \hat{R}_{a \leftrightarrow b} \setminus \{(b, a)\}$ . First,  $\hat{R} \in \mathcal{C}(T(R_\alpha))$  if and only if  $\hat{R} \in \mathcal{C}(T(R_{\alpha, a \leftrightarrow b}))$  and, second,  $\hat{R} \in \mathcal{C}(S(R_\alpha))$  if and only if  $\hat{R} \in \mathcal{C}(S(R_{\alpha, a \leftrightarrow b}))$ .*

Since this proposition is trivial from the facts above, we omit to provide here.

By Proposition 9, we immediately have the following result.

**Corollary 9** *Both  $F_A^S$  and  $F_A^T$  satisfy the reversal symmetry.*

## 6 Concluding Remarks

Our method to have a linear order is related to two influential methods; the Tideman's ranked pairs method and the Schulze method. They are types of the Condorcet methods, which attain the linear order with the same characteristics as those in Corollary 6. Although the Condorcet methods receive a lot of attention in the studies of voting, they are not widely used in the real world. A main reason of this fact may be that it is hard to make general voters understand the methods such as the ranked pairs method and the Schulze method.<sup>13</sup> Our results provide interpretations of these two methods

---

<sup>13</sup>For example, Poundstone (2008, Ch.13) mentions that "Maybe the biggest hitch with Condorcet voting is the simplicity issue. Condorcet supporters talk up the populist appeal of the John Wayne, last-man-standing premise. Soccer moms and NASCAR dads will nod their heads to that. The thing is, its impossible to explain Condorcet voting any further without talking about cycles and "beat-paths" and "cloneproof Schwartz sequential dropping." This is where the androids mask falls off and its all wires and microchips inside. Non-Ph.D.s run screaming for the exits. So far, Condorcet voting has tended to appeal to the kind of people who can write Javascript code for it."

and they may contribute to explain them well, because the explanation with the supermajority rules is relatively not be hard to understand.

Finally, we consider several remaining issues. First, we provide some remarks when our approach is used to choose only one winner. Second, we introduce a computationally efficient method to derive the  $S$ -order set.

## 6.1 Single winner election

In this subsection, we focus on methods to have only one winner in election. First, we consider the **Simpson-Kramer min-max method**, which is in a similar spirit to that considered in 4.1. We try to refine the set of winners of this method, which is calculated as

$$W^{SK} = \arg \min_{a \in A} \left[ \max_{b \in A} [\Phi [b, a]] \right].$$

Then, we have the following result.

**Remark 2**

$$W^{SK} = \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) \neq \emptyset} M(A, R_\alpha). \quad (9)$$

**Proof.** We use the following result.

**Claim 2** *If  $a \in M(A, R_\alpha)$  for  $\alpha \in [1/2, 1)$ ,  $a \in M(A, R_{\alpha'})$  for  $\alpha' \geq \alpha$ .*

First, we show

$$W^{SK} \subseteq \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) \neq \emptyset} M(A, R_\alpha).$$

Let  $a$  be in the left-hand side of (9) and  $b \in A$  is such that  $\Phi [b, a] \geq \Phi [c, a]$  for any  $c \in A \setminus \{a\}$ . Then,  $a \in M(A, R_{\Phi [b, a]})$ . Moreover, by Claim 1,  $a \in M(A, R_\alpha)$  for all  $\alpha \geq \Phi [b, a]$ . On the other hand,  $M(A, R_\alpha) = \emptyset$  for all  $\alpha < \Phi [b, a]$  because, for all  $c \in A$ , there is  $d \in A$  such that  $\Phi [d, c] > \alpha$ . Therefore,  $a$  also belongs to the right-hand side of (9)

Second, we

$$W^{SK} \supseteq \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) \neq \emptyset} M(A, R_\alpha).$$

Let  $a$  be in the right-hand side of (9) and

$$\alpha^{**} = \min \{ \alpha \in [1/2, 1) \mid M(A, R_\alpha) \neq \emptyset \}.$$

By Claim 1,  $a \in M(A, R_{\alpha^*})$ . Since  $M(A, R_\alpha) = \emptyset$  for  $\alpha < \alpha^{**}$ , for all  $c \in A$ , there is  $d$  such that  $\Phi[d, c] \geq \alpha^{**}$ . On the other hand, since  $a \in M(A, R_{\alpha^*}) \setminus M(A, R_\alpha)$  for any  $\alpha < \alpha^{**}$ ,  $\alpha^{**} \geq \Phi[c, a]$  for all  $c \in A$  and there is  $b \in A$  such that  $\alpha^{**} = \Phi[b, a]$ . Therefore,  $a$  also belongs to the left-hand side of (9). **Q.E.D.**

In words, the intersection of nonempty maximal sets for  $\alpha \in [1/2, 1)$  is equivalent to the set of winners of the Simpson-Kramer min-max method. However, any element of the intersection may be a Condorcet loser. To show this, we reexamine Example 3. In this example

$$\begin{aligned} M(A, R_\alpha) &= \emptyset \text{ if } \alpha \in [1/2, 5/9) \\ &= \{d\} \text{ if } \alpha \in [5/9, 2/3) \\ &= A \text{ if } \alpha \in [2/3, 1). \end{aligned} \tag{10}$$

Therefore,

$$W^{SK} = \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) \neq \emptyset} M(A, R_\alpha) = \{d\}.$$

As is mentioned above,  $d$  is the Condorcet loser but the unique winner of the Simpson-Kramer min-max method. Since  $d$  is the the unique winner, contrary to the case of 4.1, we cannot refine  $W^{SK}$  by using a similar method to that introduced in this study. To be more precise, we have the following result.

**Remark 3**

$$\left( \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) \neq \emptyset} M(A, R_\alpha) \right) \cap \left( \bigcap_{\alpha \in [1/2, 1): M(A, R_\alpha) = \emptyset} M(A, S(R_\alpha)) \right) \tag{11}$$

*may be empty.*

We use  $R_\alpha$  for all  $\alpha \in [1/2, 1)$  but modify it when the maximal sets for  $R_\alpha$  is empty. In some cases, we cannot choose any winner. We reexamine Example 3 to show this fact. If  $\alpha \in [1/2, 5/9)$ ,  $M(A, R_\alpha) = \emptyset$  and  $M(A, S(R_\alpha)) = \{a, b, c\}$ . By (10), in this example, (11) is empty. This implies that we cannot refine the Simpson-Kramer min-max winner by this approach.

Next, we reconsider the Schulze method and the ranked pairs method, which are introduced as methods determining the single winner. The set

of winners of Schulze method is  $W^{Sc} = M(A, R^{Sc})$  and that of the ranked pairs method is

$$W^{RP} = \{a \in A \mid R^\pi(1) = a \text{ for some } \pi \in \Pi\}.$$

We have the following results.

**Corollary 10**

$$W^{Sc} = \bigcap_{\alpha \in [1/2, 1)} M(A, T(R_\alpha)) = M\left(A, \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))\right).$$

This implies that the set of winners of Schulze method is equivalent to the intersection of the maximal sets for  $T(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ . Since  $R_\alpha$  is asymmetric for all  $\alpha \in [1/2, 1)$  (the first result of Lemma 7),  $M(A, T(R_\alpha))$  is equivalent to the Schwartz set for  $R_\alpha$ . Therefore,  $W^{Sc}$  is the intersection of the Schwartz sets for  $R_\alpha$  for all  $\alpha \in [1/2, 1)$ .

**Corollary 11**

$$W^{Sc} \cup W^{RP} \subseteq \bigcap_{\alpha \in [1/2, 1)} M(A, S(R_\alpha)) = M\left(A, \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))\right). \quad (12)$$

This implies that the sets of winners of the Schulze method and the ranked pairs method are subsets of the intersection of the maximal sets for  $S(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ .

## 6.2 Method to Derive Linear Orders

We provide methods to derive the  $S$ -order set and the  $T$ -order set, respectively. First, by Lemma 8, the construction method of  $\bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$  is the same as the Schulze rule. Therefore, we can use the method introduced by Schulze (2011). Here, we provide a method to construct  $\bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))$ .

**Proposition 10**  $(a, b) \in \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))$  if and only if  $\Phi[a, b] > \max\{1/2, B[b, a]\}$ .

**Proof.** First, we show the if-part. Suppose  $\Phi[a, b] > 1/2$  and  $\Phi[a, b] > B[b, a]$ . Then, for any  $\alpha \in [1/2, \Phi[a, b]]$ ,  $(a, b) \in R_\alpha \subseteq S(R_\alpha)$ . First, we show that  $(b, a) \notin S(R_\alpha)$  for some  $\alpha \in [1/2, \Phi[a, b]]$ . Since  $S(R_\alpha) \subseteq R_\alpha$ ,  $(b, a) \notin R_\alpha$  for any  $\alpha \in [1/2, \Phi[a, b]]$ . Moreover, by the second result of Lemma 7,  $(b, a) \notin R_\alpha$  for all  $\alpha \in [1/2, 1)$ . If  $B[b, a] < 1/2$ , then  $(b, a) \notin$

$T(R_\alpha)$  and therefore,  $(b, a) \notin S(R_\alpha)$  for all  $\alpha \in [1/2, 1)$ . Thus, we assume  $B[b, a] \geq 1/2$ . In this case,  $(b, a) \in T(R_\alpha)$  if and only if  $\alpha \in [1/2, B[b, a]]$ . Therefore, for  $\alpha \in (B[b, a], \Phi[a, b]]$ ,  $(b, a) \notin T(R_\alpha)$ . Since  $S(R_\alpha) \subseteq T(R_\alpha)$ ,  $(b, a) \notin S(R_\alpha)$ , for  $\alpha \in (B[b, a], \Phi[a, b]]$ . Therefore, since  $(a, b) \in S(R_\alpha)$  for any  $\alpha \in [1/2, \Phi[a, b]]$ ,  $(a, b) \in P(S(R_\alpha))$  for  $\alpha \in (B[b, a], \Phi[a, b]]$ .

Second, we show the only-if-part. Suppose  $(a, b) \in \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))$ . Then, we can fix  $\alpha \in [1/2, 1)$  such that  $(a, b) \in S(R_\alpha)$  and  $(b, a) \notin S(R_\alpha)$ . Then, by Lemma 3,  $(a, b) \in P(S(R_\alpha))$  implies  $(a, b) \in P(R_\alpha)$ ; that is,  $(a, b) \in R_\alpha$  and  $(b, a) \notin R_\alpha$ . Thus,  $\Phi[a, b] > 1/2$ . Next, since  $S(R_\alpha) \subseteq T(R_\alpha)$  and  $(b, a) \notin S(R_\alpha)$ ,  $(b, a) \notin T(R_\alpha)$ . Then,  $\alpha$  must be more than  $B[b, a]$  and less than or equal to  $\Phi[a, b]$ . Hence  $\Phi[a, b] > B[b, a]$ . **Q.E.D.**

Since we can have  $B[a, b]$  for all  $a, b \in A$  by using Dijkstra's algorithm, we can computationally efficiently construct  $\bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))$ .

By using  $\bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha))$  and  $\bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha))$ , we can derive the  $S$ -order set and the  $T$ -order set by using the class of algorithms in the previous section, because of Theorems 1 and 2.

For asymmetric  $R$ , let

$$\mathcal{A}(R) = \{R' \mid R' \subseteq R, R' \text{ is } P\text{-acyclic}\}$$

For asymmetric  $R$ , we let  $R'$  be a **maximum acyclic relation** if  $R' \in \mathcal{A}(R)$  and there is no  $R'' \in \mathcal{A}(R)$  such that  $R' \subseteq R''$ .

If  $R'$  is a maximum acyclic relation for  $R$ , then  $\mathcal{T}(R) \subseteq R'$ .

$$\bigcap_{\alpha \in [1/2, 1)} \mathcal{A}(R_\alpha) \neq \emptyset \text{ and } \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) \subseteq \bigcap_{\alpha \in [1/2, 1)} \mathcal{A}(R_\alpha).$$

Let  $C \subseteq R_\alpha$  be a  $P$ -cycle of  $R_\alpha$ . That is,  $(a_1^C, a_2^C, \dots, a_m^C)$  be alternatives containing  $C$  where  $a_1, a_2, \dots, a_M$  are distinct, where  $(a_m^C, a_{m+1}^C) \in R_{1/2}$  for all  $m = 1, \dots, M$  and  $a_{M+1}^C = a_1^C$ . Let

$$\Phi[a, b] = f(N[a, b], N[b, a])$$

For  $C = (a_1^C, a_2^C, \dots, a_m^C) \in \mathcal{B}[(a, b), R_{1/2}]$ , let

$$W_C = \{(a_j^C, a_{j+1}^C) \mid \Phi[a_j^C, a_{j+1}^C] \leq \Phi[a_m^C, a_{m+1}^C] \text{ for all } m = 1, \dots, M\}$$

That it,  $W_C$  is the set of weakest links of  $C$ .

be the set of  $\Phi[a_j, a_{j+1}]$

$$ST(C) = \min_{j=1, \dots, m} [\Phi[a_j, a_{j+1}]].$$



## References

- Arrow, K.J. (1963) *Social Choice and Individual Values*. Yale University Press, New Haven, CT.
- Arrow, K.J. (1977) Current Developments in the Theory of Social Choice, *Social Research* 44, 607-622.
- Balasko, Y., Crès H., (1997) The probability of Condorcet cycles and super majority rules, *Journal of Economic Theory* 75, 237-270.
- Black, D. (1958) *The Theory of Committees and Elections*. Cambridge University Press, London.
- Borda, J.-C. (1781) *Memoires sue les Elections au Scrutin*, *Histoires de l'Academie Royale des Sciences*. English translation by A. de Grazia. *Mathematical Derivation of an Election System*, *Isis* 44 (1953).
- Bordes, G. (1976) Consistency, Rationality, and Collective Choice, *Review of Economic Studies* 43, 451-457.
- Bossert, W., Sprumont Y., Suzumura, K. (2005) Consistent Rationalizability, *Economica* 72, 85–200.
- Bossert, W., Suzumura, K. (2010) *Consistency, Choice, and Rationality*, Harvard University Press.
- Caplin, A., Nalebuff, B. (1988) On 64% majority rule, *Econometrica* 56, 784-814.
- Coughlin, P.J. (1981) Necessary and Sufficient Conditions for  $\delta$ -relative Majority Voting Equilibria. *Econometrica* 49(5), 1223–1224
- Condorcet, M. (1785) *Essai Sur L'Application de L'Analyse a la Probabilite Des Decisions Rendues a la Pluralite Des Voix*. Paris.
- Crès H., Ünver, M. U. (2017) Toward a 50%-majority Equilibrium When Voters are Symmetrically Distributed, *Mathematical Social Sciences* 90, 145-149.
- Deb, R. (1977) On Schwartz's Rule. *Journal of Economic Theory* 16, 103-110.
- Gjorgjiev, R., Xefteris, D. (2015) Transitive Supermajority Rule Relations, *Economics Theory Bulletin* 3, 299–312.

- Green-Armytage, J., Tideman, N. (2020) Selecting the Runoff-pair, *Public Choice* 182, 119–137
- Greenberg, J. (1979) Consistent majority rule over compact sets of alternatives, *Econometrica* 47, 627-663.
- Nakamura, K. (1979) The voters in a simple game with ordinal preferences, *International Journal of Game Theory* 8, 55–61.
- Pattanaik, P.K. (2002) Positional Rules of Collective Decision-making, in *Handbook of Social Choice and Welfare*. Vol. 1, Edited by Arrow, K.J. Sen A.K., and Suzumura K., North-Holland. 361-394.
- Saari, D.G. (1994) *Geometry of Voting*, Springer Nature.
- Schwartz, T. (1972) Rationality and the Myth of the Maximum. *Nous* 6, 97-117.
- Schulze, M. (2011) A New Monotonic, Clone-independent, Reversal Symmetric, and Condorcet-consistent Single-winner Election Method, *Social Choice and Welfare* 36(2), 267–303.
- Slutsky, S. (1979) Equilibrium under  $\alpha$ -majority voting, *Econometrica* 47(5), 1113–1125.
- Suzumura, K. (1983) *Rational Choice, Collective Decisions and Social Welfare*, New York: Cambridge University Press.
- Suzumura, K. (2012) *Introduction to Social Choice Theory* (in Japanese), Tokyo: Toyo Keizai Inc.
- Tideman, T. N. (1987) Independence of Clones as a Criterion for Voting Rules, *Social Choice and Welfare* 4(3), 185–206.
- Tideman, T. N. (2006) *Collective Decisions and Voting: The Potential for Public Choice*, Ashgate Publishing.
- Tovey, C. A. (1997) Probabilities of Preferences and Cycles with Super Majority Rules, *Journal of Economic Theory* 75, 271-279.
- Truchon, M. (1998) An Extension of the Concordet Criterion and Kemeny Orders, *Mimeo*
- Young, H. P. (1988) Condorcet's theory of voting, *American Political Science Review* 82(4), 1231–1244.

Zavist, T.M., Tideman, T.N. (1989) Complete Independence of Clones in the Ranked Pairs Rule, *Social Choice and Welfare* 6, 167–173.

## Appendix: Example

In the examples so far,  $S(R_\alpha) = T(R_\alpha)$  for any  $\alpha \in [1/2, 1)$  and thus

$$\bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(S(R_\alpha)) = \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(T(R_\alpha))$$

are satisfied. Here, we provide an example where the equality above does not hold. Moreover, in the example, the linear orders obtained by the Schulze method and the ranked pairs method differ. Moreover, we discuss some other important voting methods by using the example.

Let  $A = \{a, b, c, d, e\}$  and  $I = \{1, \dots, 47\}$ . All individuals have a linear order on  $A$  and

- $R^i : a, c, b, e, d$  for  $i = 1, \dots, 5,$
- $: a, d, e, c, b$  for  $i = 6, \dots, 10,$
- $: c, a, b, e, d$  for  $i = 11, 12, 13$
- $: c, a, e, b, d$  for  $i = 14, \dots, 20,$
- $: b, c, d, a, e$  for  $i = 21$
- $: b, e, d, a, c$  for  $i = 22, \dots, 29,$
- $: c, b, a, d, e$  for  $i = 30, 31,$
- $: e, b, a, d, c$  for  $i = 32, \dots, 39,$
- $: d, c, b, a, e$  for  $i = 40$
- $: d, c, e, b, a$  for  $i = 41, \dots, 44,$
- $: d, e, c, b, a$  for  $i = 45, 46, 47.$

Then, the following table summarizes the voting results, where the number of a cell represents that of individuals who prefers the row alternative to the column one.

	a	b	c	d	e
a		20	26	30	24
b	27		17	34	20
c	21	30		18	23
d	17	13	29		16
e	23	27	24	31	

Then  $R_{1/2}$  is represented as **Figure 1**; that is, there is an arrow from  $j$  to  $k$  exists if  $(j, k) \in R_{1/2}$  or equivalently if 24 or more individuals prefer  $j$  to  $k$ , where the number written nearby the arrow from  $j$  to  $k$  represents the number of individuals who prefer  $j$  to  $k$ .

Therefore,  $R_\alpha$ ,  $S(R_\alpha)$  and  $T(R_\alpha)$  are as follows. First, for  $\alpha \in [1/2, 24/47)$ ,

$$\begin{aligned} R_\alpha &= \left\{ (a, c), (a, d), (a, e), (b, a), (b, d), \right. \\ &\quad \left. (c, b), (c, e), (d, c), (e, b), (e, d) \right\}, \\ S(R_\alpha) &= \{(a, d), (b, d)\} = T(R_\alpha). \end{aligned}$$

Second, for  $\alpha \in [24/47, 26/47)$ ,

$$\begin{aligned} R_\alpha &= R_{1/2} \setminus \{(a, e), (c, e)\}, \\ P(S(R_\alpha)) &= \{(e, b), (e, d)\}, \\ P(T(R_\alpha)) &= P(S(R_\alpha)) \cup \{(e, a), (e, c)\}. \end{aligned}$$

Third, for  $\alpha \in [26/47, 27/47)$ ,

$$\begin{aligned} R_\alpha &= R_{24/47} \setminus \{(a, c)\}, \\ P(S(R_\alpha)) &= \{(e, b), (e, d)\}, \\ P(T(R_\alpha)) &= P(S(R_\alpha)) \cup \{(e, a), (e, c)\}. \end{aligned}$$

Fourth, for  $\alpha \in [27/47, 29/47)$ ,

$$\begin{aligned} R_\alpha &= R_{26/47} \setminus \{(b, a), (e, b)\} \\ P(S(R_\alpha)) &= \{(a, d), (e, d)\}, \\ P(T(R_\alpha)) &= P(S(R_\alpha)) \cup \{(a, b), (a, c), (e, c)\}. \end{aligned}$$

Fifth, for  $\alpha \in [29/47, 30/47)$ ,

$$\begin{aligned} R_\alpha &= R_{27/47} \setminus \{(c, d)\}, \\ P(S(R_\alpha)) &= R_\alpha = \{(a, d), (b, d), (c, b), (e, d)\} \\ P(T(R_\alpha)) &= P(S(R_\alpha)) \cup \{(c, d)\}. \end{aligned}$$

Sixth, for  $\alpha \in [30/47, 31/47)$ ,

$$\begin{aligned} R_\alpha &= R_{29/47} \setminus \{(a, d), (c, b)\}, \\ P(S(R_\alpha)) &= R_\alpha = \{(b, d), (e, d)\} = P(T(R_\alpha)). \end{aligned}$$

Seventh, for  $\alpha \in [31/47, 34/47)$ ,

$$\begin{aligned} R_\alpha &= R_{30/47} \setminus \{(e, d)\}, \\ P(S(R_\alpha)) &= R_\alpha = \{(b, d)\} = P(T(R_\alpha)). \end{aligned}$$

Eighth, for  $\alpha \in [34/47, 1)$ ,

$$R_\alpha = S(R_\alpha) = \emptyset, \quad \alpha \in [34/47, 1).$$

Among them, we explain the case where  $\alpha \in [27/47, 29/47)$ . In this case,  $R_\alpha$  is represented as Figure 2; that is, a thick solid arrow from  $j$  to  $k$  exists if  $(j, k) \in R_\alpha$  or equivalently if 29 or more individuals prefer  $j$  to  $k$ . Since there is a P-cycle  $(b, d), (d, c), (c, b) \in R_\alpha, (d, b), (c, d), (b, c) \in S(R_\alpha)$ , which are represented by the thin solid arrows in Figure 2. Moreover, since  $(a, d), (d, c) \in R_\alpha, (a, c) \in T(R_\alpha) \setminus S(R_\alpha)$ . Similarly,  $(a, b), (a, c), (e, b), (e, c) \in T(R_\alpha) \setminus S(R_\alpha)$ , which are represented by the dot lines in Figure 1.

Thus,

$$\begin{aligned} \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) &= \{(a, d), (b, d), (c, b), (e, b), (e, d)\}, \\ \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) &= \left( \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \right) \cup \\ &\quad \{(a, b), (a, c), (c, d), (e, a), (e, c)\}. \end{aligned}$$

Then, we have

$$\begin{aligned} \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(S(R_\alpha)) &= \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(S(R_\alpha)) \right) = \{R_1^*, \dots, R_8^*\}, \\ \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(T(R_\alpha)) &= \mathcal{C} \left( \bigcup_{\alpha \in [1/2, 1)} P(T(R_\alpha)) \right) = \{R_6^*\}, \end{aligned}$$

where

$$\begin{aligned} R_1^* &: a, c, e, b, d, & R_2^* &: a, e, c, b, d, \\ R_3^* &: c, a, e, b, d, & R_4^* &: c, e, a, b, d, \\ R_5^* &: c, e, b, a, d, & R_6^* &: e, a, c, b, d, \\ R_7^* &: e, c, a, b, d, & R_8^* &: e, c, b, a, d. \end{aligned}$$

Note that  $R_6^* = R^{Sc}$  and  $R_5^* = R^\pi$  for any  $\pi \in \Pi$ .

Next, we focus on other important voting methods to determine a collective linear order. First, we consider the Kemeny-Young method. By this method, in the example above, we have linear order  $R^{KY} : e, b, a, d, c$ . In this case,

$$R^{KY} \notin \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(S(R_\alpha)),$$

because if  $\alpha \in [29/47, 30/47)$ , then  $(c, b) \in P(S(R_\alpha)) = R_\alpha$ .

Second, we consider the Borda method.<sup>14</sup> By this method, in the example above, we have linear order  $R^B : e, a, b, c, d$ . Then, we also have

$$R^B \notin \bigcap_{\alpha \in [1/2, 1)} \mathcal{C}(S(R_\alpha)),$$

because if  $\alpha \in [29/47, 30/47)$ , then  $(c, b) \in P(S(R_\alpha)) = R_\alpha$ .

---

<sup>14</sup>Since there is no Condorcet cycle in  $R_{1/2}$ , the result of the Black method is the same as that of the Borda rule.