

Two Applications of Topological Fixed Point Theory

Chaitanya Ambi*

April 20, 2023

Abstract

We give a new proof of Cartan's fixed point theorem using topological fixed point theory. For an odd dimensional, simply connected and complete manifold having non-positive curvature, we further prove that every isometry with finite order not only fixes a point, but also maps a *pencil* of geodesics to itself.

We apply similar techniques to the automorphism group of a smoothly bounded domain in the complex plane having connectivity three or more. We show that any prime dividing the order of this finite group must divide a certain integer depending only on the connectivity of the domain.

Keywords: Topological Fixed Point Theory, Automorphism Groups

MSC[2020]: 57S05, 57S25

1 Introduction.

The geometric properties of a complete Riemannian manifold M with non-positive curvature are intimately linked with its fundamental group. Specifically, the structure of the isometry group $Isom(M)$ has been a field of intense investigation. It is not easy to ascertain whether a geometric property is, in fact, topological. As an attempt in this direction, we present a

*Chennai Mathematical Institute, H1, SIPCOT IT Park, Siruseri, Kelambakkam 603103, India.

Email: chaitanya.ambi@gmail.com

new proof of the following version of Cartan's fixed point theorem (see [6], Chap. 6, Thm. 23, p. 164). A stronger version of the same for precompact subgroups of $Isom(M)$ is known (see [2], Thm. 1.4.6, p. 21).

Theorem 1.1 (E. Cartan). *Consider a complete, simply connected Riemannian manifold M of non-positive curvature. If $\sigma \in Isom(M)$ is an element of finite order, then*

$$Fix(\sigma) \neq \emptyset.$$

The above theorem can be proved by considering the centre of mass of the orbit of σ . In this article, we present a proof using topological fixed point theory. Our approach also reveals the following fact which is interesting by its own. By a *pencil* of geodesics, we shall mean a (non-intersecting) family of geodesics any two of which lie within a finite Hausdorff distance of one another.

Corollary 1.1. *In addition to the hypotheses of Cartan's Theorem 1.1, assume that $\dim M$ is odd. Then there exists a pencil Γ of geodesics such that*

$$\sigma(\Gamma) = \Gamma.$$

Next, certain domains in the complex space \mathbb{C}^n also become complete manifolds with non-positive curvature when equipped with the Bergmann metric. In this vein, we consider the group of conformal automorphisms of a bounded planar domain (with smooth boundary) having finite connectivity (see [5], Def. 4.2.2, p. 85). It is well-known that such a group is finite whenever the connectivity exceeds two (see [5], Thm. 12.2.3, p. 263). We prove the following result about such domains (where $Aut(\cdot)$ denotes the group of conformal automorphisms of the respective domain).

Theorem 1.2. *Let $\Omega_k \subset \mathbb{C}$ be any bounded planar domain having connectivity $k + 1$ (with $k \geq 2$) whose boundary consists of finitely many smooth Jordan curves. If p is a prime integer dividing the order of $Aut(\Omega_k)$, then we have*

$$k^3 \equiv k \pmod{p}.$$

As far as the author knows, it is an open problem to characterise the finite groups which may occur as the automorphism groups of planar domains. In

particular, by [4], we have the following sharp upper bound for $k \geq 2$.

$$\begin{aligned} \#Aut(\Omega_k) &\leq 2k && \text{for } k \neq 4, 6, 8, 12, 20, \\ & && \#Aut(\Omega_4) \leq 12, \\ & && \#Aut(\Omega_6), \#Aut(\Omega_8) \leq 24, \\ & && \#Aut(\Omega_{12}), \#Aut(\Omega_{20}) \leq 60. \end{aligned}$$

Theorem 1.2 shows that there are further restrictions on the possible order of an element in $Aut(\Omega_k)$. Hence, the subgroups of $Aut(\Omega_k)$ are constrained by the value of k . For instance, the cyclic group of order 7 cannot occur as a subgroup of $Aut(\Omega_k)$ for any $k < 6$ (otherwise, we would have $7|(k^3 - k)$). Note that the upper bound mentioned above does not exclude this possibility as $\#Aut(\Omega_5) \leq 10$.

2 Proof of Theorem 1.1.

By the Cartan-Hadamard Theorem (see [6], Chap. 6, Thm. 22, p. 162), M is diffeomorphic to \mathbb{R}^n , where $n = \dim M$. We compactify M by adding its ideal boundary $M(\infty)$ to obtain the *Gromov compactification*

$$\bar{M} := M \cup M(\infty).$$

We refer the reader to [1], Chap, 3, pp. 22-32 for the details. In fact, \bar{M} is homeomorphic to the closed unit ball $\mathbb{B}^n \subset \mathbb{R}^n$.

Next, let $\sigma \in Isom(M)$ have a finite order. It is known that σ induces a homeomorphism $\bar{\sigma}$ of $\bar{M} \approx \mathbb{B}^n$.

We shall argue inductively to reduce the dimension n . By Brouwer's fixed point theorem, $\bar{\sigma}$ has a fixed point $x \in \bar{M}$. If $x \in M(\infty)$, we observe that (see [1], Chap. 3, p. 30) $\bar{\sigma}$ maps a *horosphere* centred at x to itself. This horosphere in \bar{M} is diffeomorphic to \mathbb{B}^{n-1} . So, we may repeat the whole argument to keep reducing the dimension. At each stage, the restriction of $\bar{\sigma}$ fixes a point lying either in M or on $M(\infty)$.

If the latter case were to occur until the dimension reaches two, the argument would yield a geodesic $\gamma \subset M$ satisfying

$$\sigma(\gamma) = \gamma.$$

But now, γ is *isometric* with \mathbb{R} (equipped with the standard Euclidean metric). Any isometry of the latter either fixes a point or has infinite order. As σ is of finite order, we get a fixed point as asserted. This completes the argument.

2.1 Proof of Corollary 1.1.

As in the proof of Theorem 1.1, the extended map $\bar{\sigma}$ restricts to a homeomorphism of $M(\infty)$. Since we want to find a family of undirected geodesics (and not geodesic rays), we may well disregard the directions to obtain the quotient of $M(\infty)$ by $\mathbb{Z}/2\mathbb{Z}$ action. As $M(\infty) \approx \mathbb{S}^{n-1}$, the quotient space is homeomorphic to the real projective space $\mathbb{R}P^{n-1}$. This space also has an action induced by that of $\bar{\sigma}$.

Assume the contrary that the above action induced by $\bar{\sigma}$ has no fixed points. Let $m \in \mathbb{N}$ be the smallest power of $\bar{\sigma}$ whose Lefschetz number is nonzero. It is clear that m cannot exceed the order of $\bar{\sigma}$ because $(n-1)$ is even and so

$$\lambda(\text{Id}_{\mathbb{R}P^{n-1}}) = \chi(\mathbb{R}P^{n-1}) = 1.$$

If $m \neq 1$, choose any prime p which divides m . Set $\bar{\tau} := \bar{\sigma}^{m/p}$. For the action induced by $\bar{\tau}$ on the quotient space mentioned above, we must have

$$0 = \lambda(\bar{\tau}) \equiv \lambda(\bar{\tau}^p) \pmod{p}.$$

Here, we have used the \pmod{p} Lefschetz Theorem (see [3], Chap. 9, Thm. 3.4, p. 231). Thus, we arrive at the contradiction that

$$0 \equiv 1 = \chi(\mathbb{R}P^{n-1}) \pmod{p}.$$

Hence, the action induced by $\bar{\sigma}$ itself fixes some point. This fixed point corresponds precisely to a pencil Γ of geodesics as asserted.

3 Proof of Theorem 1.2.

Proof.

Select an arbitrary element $\sigma \in \text{Aut}(\Omega_k)$ having a prime order p . By [5] (see Thm. 1.4.3, p. 22), we have

$$\# \text{Fix}(\sigma) \leq 2. \tag{1}$$

Since $\Omega := \Omega_k \setminus \text{Fix}(\sigma)$ is not compact, we shall work with *generalised* Lefschetz numbers (as in [3], Def. 3.1, p. 421). Note that Ω is a *Lefschetz space*

(by [3], Chap. 5, Thm. 4.1, p. 423).

Now, the restriction of σ to Ω is an automorphism without fixed points. Therefore, the generalised Lefschetz number $\lambda(\sigma)$ of σ must vanish. As Ω has the homotopy type of a wedge of $(k + \#Fix(\sigma))$ circles, we obtain (using [3], Chap. 9, Prop. 4.7. p. 233)

$$\chi(\Omega) = 1 - k - \#Fix(\sigma) \equiv 0 \pmod{p}.$$

In view of Eq. (1), this implies that p must divide either k or one of $k \pm 1$. Thus, the assertion of the Theorem follows. \square

Acknowledgement. *The author would like to thank Chennai Mathematical Institute for support by a post-doctoral fellowship.*

References

- [1] Werner Ballmann, Mikhael Gromov, and Viktor Schroeder. *Manifolds of nonpositive curvature*, volume 61 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1985.
- [2] Patrick B. Eberlein. *Geometry of nonpositively curved manifolds*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1996.
- [3] Andrzej Granas and James Dugundji. *Fixed point theory*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [4] Maurice Heins. On the number of 1-1 directly conformal maps which a multiply-connected plane region of finite connectivity $p(> 2)$ admits onto itself. *Bull. Amer. Math. Soc.*, 52:454–457, 1946.
- [5] Steven G. Krantz. *Geometric function theory*. Cornerstones. Birkhäuser Boston, Inc., Boston, MA, 2006. Explorations in complex analysis.
- [6] Peter Petersen. *Riemannian geometry*, volume 171 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2006.