Studying continuous, time-varying, and/or complex exposures using longitudinal modified treatment policies

Katherine L. Hoffman^{*1}, Diego Salazar-Barreto², Nicholas T. Williams³, Kara E. Rudolph^{†3}, and Iván Díaz^{†4}

¹Division of Biostatistics, Department of Population Health Science, Weill Cornell Medicine, New York, NY, USA.

²School of Industrial Engineering, University of Los Andes, Bogotá, Colombia.

³Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA.

⁴Division of Biostatistics, Department of Population Health, New York University Grossman School of Medicine, New York University, New York, NY, USA.

May 15, 2024

Abstract

This tutorial discusses methodology for causal inference using longitudinal modified treatment policies. This method facilitates the mathematical formalization, identification, and estimation of many novel parameters, and mathematically generalizes many commonly used parameters, such as the average treatment effect. Longitudinal modified treatment policies apply to a wide variety of exposures, including binary, multivariate, and continuous, and can accommodate time-varying treatments and confounders, competing risks, loss-to-follow-up, as well as survival, binary, or continuous outcomes. Longitudinal modified treatment policies can be seen as an extension of static and dynamic interventions to involve the natural value of treatment, and, like dynamic interventions, can be used to define alternative estimands with a positivity assumption that is more likely to be satisfied than estimands corresponding to static interventions. This tutorial aims to illustrate several practical uses of the longitudinal modified treatment policy methodology, including describing different estimation strategies and their corresponding advantages and disadvantages. We provide numerous examples of types of research questions which can be answered using longitudinal modified treatment policies. We go into more depth with one of these examples—specifically, estimating the effect of delaying intubation on critically ill COVID-19 patients' mortality. We demonstrate the use of the open-source R package *lmtp* to estimate the effects, and we provide code on https://github.com/kathoffman/Imtp-tutorial.

^{*}Corresponding author:

kh3233@cumc.columbia.edu

⁷²² W. 168th St, NY, NY 10032

[†]Authors contributed equally.

Introduction

Consider the research question, "does early initiation of invasive mechanical ventilation (intubation) for coronavirus 2019 (COVID-19) patients increase mortality rates?". This query is the subject of active critical care research^{1–11}, but it is difficult to translate into a meaningful estimand and estimation procedure for multiple reasons. First, intubation can take place throughout a patient's course of hospitalization, meaning it is a time-varying exposure, or treatment, of interest. Timevarying exposures require specific methodology from the causal inference literature to properly accomodate time-dependent confounding^{12,13}. Second, respiratory support is a multilevel exposure, minimally consisting of categories such as "no oxygen support," "non-invasive respiratory support," and "invasive mechanical ventilation." The majority of the causal inference literature focuses on dichotomous exposures and interventions, e.g. treat everyone vs. treat no one, whereas methodology which can incorporate multilevel or continuous exposures is limited. Finally, a key assumption for causal inference is positivity, which roughly states that the intervention considered must occur with positive probability within all strata of confounders. For time-varying exposures, this positive probability must hold at each time point¹⁴.

Longitudinal modified treatment policies may offer a solution to the aforementioned challenges. This method provides an approach to translating complex research questions into a broad range of causal estimands, identifying such quantities, and estimating them. In brief, a modified treatment policy allows a hypothetical intervention to depend on the *natural value of treatment*, i.e. the value that treatment would take at time t if an intervention was discontinued right before time t^{15} . A modified treatment policy for a time-varying, or longitudinal, exposure, is a longitudinal modified treatment policy. The methodology for longitudinal modified treatment policies not only mathematically generalizes modified treatment policies, but can also be seen as an extension of more commonly known interventions such as *static* and *dynamic* interventions. The ?methodology thus allows for a wide range of interventions^{15–22} including those on binary, categorical, continuous, and multiple exposures. Longitudinal modified treatment policies can also accommodate many types of outcomes including binary, continuous, or time-to-event outcomes with possible competing risks and informative right-censoring. Finally, longitudinal modified treatment policies can address violations of the positivity assumption because they allow researchers to define, identify, and estimate alternative estimands for which positivity holds by design.

In this tutorial, we provide a guide to understanding and applying the longitudinal modified treatment policy methodology. We begin by reviewing static and dynamic interventions to see how longitudinal modified treatment policies fit into the broader causal inference literature, and to understand how they mathematically generalize static and dynamic interventions, as well as some other interventions which depend on the natural value of treatment. We then discuss specifics of the longitudinal modified treatment policy methodology, including defining interventions and identifying estimands. We highlight several estimation procedures and provide numerous examples of research questions that can be addressed with longitudinal modified treatment policies. Lastly, we demonstrate an analysis on a real-world longitudinal observational data set. One way to operationalize the initially proposed question of the effect of intubation timing policies on mortality is to consider early versus late intubation, relative to the actual time patients were intubated. We illustrate one way of defining that query using a longitudinal modified treatment policy and provide detailed descriptions of the study design and analytical methods, as well as code and synthetic data to aid future researchers.

Notation and general setup

We begin with a sample of independent and identically distributed observations $Z_1, ..., Z_n$ drawn from a distribution P. This P represents a longitudinal process and may contain any number of time points, but for simplicity we will describe a distribution with only two time points, $t \in \{0, 1\}$. For each unit in the study, we observe a set of random variables $Z = (L_0, A_0, L_1, A_1, Y)$. At the first time point, baseline covariates L_0 affect the baseline exposure, A_0 . At the second time point, we observe covariates L_1 and exposure A_1 , which are themselves affected by L_0 and A_0 , and have the potential to change from their respective baseline values (time-varying). An outcome Y is measured at the end of a defined follow up period. Each endogenous variable L_0, A_0, L_1, A_1 , and Y has a corresponding exogenous variable U, representing the unmeasured, external factors affecting each measured process. We may use the following simplified directed acyclic graph²³ shown in Figure 1 to denote the set-up.



Figure 1: Simplified directed acyclic graph representing a two-time point data structure.

We will use H_t as a shorthand notation for the history of data measured up to right before A_t . For example, $H_0 = L_0$, and $H_1 = (L_0, A_0, L_1)$. We conceptualize causal interventions, or treatment policies, in terms of hypothetical interventions on nodes of the DAG²⁴. First, consider a user-given function $d_0(a_0, h_0, \epsilon_0)$ which maps a treatment value a_0 , a history h_0 , and a randomizer ϵ_0 into a potential treatment value. The intervention at time t = 0 is defined by removing node A_0 from the DAG and replacing it with $A_0^d = d_0(A_0, H_0, \epsilon_0)$. This assignment generates counterfactual data $H_1(A_0^d)$ and $A_1(A_0^d)$, where we use notation $X(A_0^d)$ to denote the counterfactual value of X that would have been observed had treatment at time 0, A_0 been assigned according to d. $H_1(A_0^d)$ is referred to as the counterfactual history and $A_1(A_0^d)$ is referred to as the *natural value* of treatment^{15,16,21}, i.e., the value that treatment would have taken if the intervention is performed at time t = 0 but discontinued thereafter. At time t = 1, the intervention is likewise defined by a function $d_1(a_1, h_1, \epsilon_1)$. However, at t = 1 (and all subsequent times if there are more than two time points), the function must be applied applied to both the natural value of treatment *and* the counterfactual history. That is, at time t = 1, the intervention is defined by removing node A_1 from the DAG and replacing it with $A_1^d = d_1(A_1(A_0^d), H_1(A_0^d), \epsilon_1)$.

We refer to these longitudinal interventions, and the subsequent methods to identify and estimate effects under such interventions, as longitudinal modified treatment policies. We now give examples of how the functions d_t may be defined, explain how they mathematically generalize static and dynamic interventions, and discuss novel and useful interventions that may be defined using this setup.

A review of static and dynamic interventions

The function $d_t(a_t, h_t, \epsilon_t)$ that defines the intervention or treatment policy can be categorized by the inputs it non-trivially depends on. An intervention which depends on no inputs, meaning it is applied as a constant across all study units, is a static intervention. For this tutorial, we refer to an intervention which depends only on a study unit's past covariates as a dynamic intervention. Finally, we call an intervention which depends on a study unit's natural value of treatment (and possibly past covariates, and possibly a randomizer ϵ_t) a modified treatment policy. We summarize these hierarchical categories in Table 1.

Static interventions

In a static intervention, $d_t(a_t, h_t, \epsilon_t)$ is a constant value, so it does not actually vary with a_t, h_t , nor ϵ_t .

Example 1 (Average treatment effect). For the two time point example, one might examine the counterfactual outcomes in a hypothetical world in which all units are treated at both time points $(d_t = 1 \text{ for } t \in \{0, 1\})$, and contrast them to a hypothetical world in which no units are treated at either time point $(d_t = 0 \text{ for } t \in \{0, 1\})$, giving rise to the well-known average treatment effect.

Dynamic interventions

In a dynamic intervention, the function d_t assigns a treatment value according to a unit's covariate history h_t , but does not vary with a_t nor ϵ_t . This is often used in observational studies when study units need to meet an indication of interest for a treatment or policy to reasonably begin, for example, a severity of illness indicator or socioeconomic threshold.

Example 2 (Corticosteroids for COVID-19 hospitalized patients). One dynamic treatment regime application is to study the effect of initiating a corticosteroids regimen for COVID-19 patients²⁵. For example, we might estimate mortality under a hypothetical policy where corticosteroids are administered for six days if and when a COVID-19 patient first meets a severity of illness criteria (i.e. low levels of blood oxygen). In notation,

$$\mathsf{d}_t(h_t) = \begin{cases} 1 & \text{if } l_s^* = 1 \text{ for any } s \in \{t - 5, \dots, t\} \\ 0 & \text{otherwise,} \end{cases}$$

where L_t^* is a variable in H_s that denotes the first instance of low levels of blood oxygen.

Modified treatment policies

In contrast to static and dynamic interventions, in a modified treatment policy the intervention function $d_t(a_t, h_t, \epsilon_t)$ non-trivially depends on the natural value of treatment a_t , and perhaps on h_t and/or ϵ_t .

Example 3 (Threshold intervention). One example of a modified treatment policy is a threshold function, where all natural exposure values which fall outside of a certain boundary are intervened upon to meet a constant value. This type of intervention could be used to assess the effect of lifestyle interventions, for example, intervening on individuals' average number of drinks per week and estimating the risk of coronary heart disease¹⁷. If we categorize drinks per week as 1 = "none," 2 = "1-5," 3 = "6-10," 4 = "11-15," and 5 = ">25", and we intervene to lower all individuals in the highest two drinks-per-week categories to "6-10," we can consider that intervention in notation as,

$$\mathsf{d}_t(a_t) = \begin{cases} a_t & \text{if } a_t < 4\\ 3 & \text{otherwise.} \end{cases}$$

Example 4 (Smoking cessation policy). Another example is a hypothetical policy in which half of all current smokers quit smoking forever¹⁶. This intervention is motivated by the infeasibility of studying a world in which all current smokers quit smoking forever, since genetics, environment, and many other factors will always create some portion of current smokers who will never quit. Letting A_t denote a random variable denoting smoking and ϵ_t a random draw from a uniform distribution in (0, 1), this intervention may be represented in notation as

$$\mathsf{d}_t(a_t, \epsilon_t) = \begin{cases} 0 & \text{ if } \epsilon_t < 0.5 \text{ and } a_t = 1\\ a_t & \text{ otherwise,} \end{cases}$$

Example 5 (Multiplicative or additive shift functions). Shift functions assign treatment by modifying the natural value of the exposure by some constant δ . This intervention can be additive onto the exposure value, such as estimating the effect of a hypothetical intervention to reduce lung cancer resection surgeries lasting longer than 60 minutes by 15 minutes²⁰.

$$\mathsf{d}_t(a_t) = \begin{cases} a_t & \text{if } a_t \le 60\\ a_t - 15 & \text{otherwise.} \end{cases}$$

This shift function could also change the exposure on a multiplicative scale. For example, we may be interested in studying the effect of an intervention which doubles the number of street lights for roads with less than 10 lights per mile on nighttime automobile accidents.

$$\mathsf{d}_t(a_t) = \begin{cases} a_t & \text{if } a_t \ge 10\\ 2a_t & \text{otherwise.} \end{cases}$$

We provide additional examples of interesting interventions in the eAppendix. Readers should also note that some interventions we refer to as modified treatment policies²⁰ in this tutorial are called interventions which depend on the natural value of treatment,¹⁵ and are related to methods for stochastic treatment regimes which shift the natural population- or individual-level values of treatment^{19,26} discussed in other literature.

Causal estimands and identifying parameter

Once an intervention is specified, the counterfactual outcomes of observations under a specific d are denoted as $Y(\bar{A}^{d}_{\tau})$, where \bar{A} indicates the history of measurements of A for all time points, i.e. $\bar{A} = (A_1, \ldots, A_{\tau})$. Causal effects are defined as a distribution of contrasts of $Y(\bar{A}^{d}_{\tau})$ under different interventions, d' and d*. In this tutorial, we focus on $E[Y(\bar{A}^{d'}_{\tau}) - Y(\bar{A}^{d^*}_{\tau})]$ as our causal estimand of interest. The functions d' and d* may be any type of intervention, including "no intervention."

The next step in a formal causal inference analysis is to write the counterfactual expectation $E[Y(\bar{A}_{\tau}^{d'})]$ as a formula that depends only on the observed data distribution—i.e., an identifying formula. This will generally require assumptions, some of which are untestable with the data available. The mathematically rigorous form of the assumptions is given elsewhere^{21,22}, but we state them here in simple terms:

Assumption 1 (Positivity or common support¹⁵). If it is possible to find an observation with history h_t with an exposure of a_t , then it is also possible to find an observation with history h_t with an exposure $d(a_t, h_t, \epsilon_t)$.

Assumption 2 (Strong sequential randomization²²). This assumption states that all common causes of the intervention variable A_t and $(U_{L,t+1}, U_{A,t+1})$ are measured and recorded in H_t . This is generally satisfied if H_t contains all common causes of A_t and $(L_{t+1}, A_{t+1}, \ldots, L_{\tau}, A_{\tau}, Y)$, where τ is the last time point in the study.

Assumption 3 (Weak sequential randomization^{21,22}). This assumption states that all common causes of the intervention variable A_t and $(U_{L,t+1})$ are measured and recorded in H_t^{17} , and is generally satisfied if H_t contains all common causes of A_t and $(L_{t+1}, \ldots, L_{\tau}, Y)$.

Identification of longitudinal modified treatment policy estimands requires the strong version of sequential randomization. Interventions that do not depend on the natural value of treatment, such as static and dynamic interventions, require positivity and the weak version of sequential randomization.

Positivity

Violations to positivity can be *structural*, meaning there are certain characteristics of an individual or unit that will never yield receipt of the treatment assignment under the intervention. This type of positivity violation will not improve even with an infinite sample size. Violations to positivity can also be *practical*, meaning due to random chance or small datasets, there are certain covariate combinations with zero or near-zero predicted probabilities of treatment. For time-varying treatments, this positive probability must be maintained at each time point¹⁴. Practical positivity violations can increase the finite bias and variance of estimates and severely threaten the validity of casual inference analyses when not addressed¹⁴. By design, non-static interventions (e.g. dynamic treatment rules, modified treatment policies) may help define estimands with plausible positivity, since the function d can be modified to affect the exposure of only observations which are not subject to positivity violations, either structurally or practically.

This can be seen in the interventions described above, for instance, the additive shift in Example 5. We can conceptualize a world in which a continuous exposure is instead observed at

some fixed value higher or lower than it was factually observed for every unit in the study; for example, if surgery times were 15 minutes shorter for all lung resection biopsies. However, this type of uniform hypothetical modification is destined for structural positivity violations, because at the lowest end of the observed exposure range, there will by definition be no support for the intervened exposure level $d(a_t)$ (much less conditional on the observation's history h_t). This can be avoided by constraining the range of a_t affected by the hypothetical intervention, so that no $d(a_t)$ values are produced outside the observed range of A. The intervention function can also be modified to accommodate any other remaining structural or practical positivity violations. For example, clinical knowledge may inform us that a treatment of interest will never be administered after a certain amount of time since a disease diagnosis has passed, so the hypothetical intervention would restrict the values of t in which the intervention can occur. Alternatively, if there is not enough support in the data for individuals of a certain covariate strata or at a naturally observed exposure level to receive the intervention, a different estimand can be defined.

Identification formula

Under Assumptions 1 and 2, or 1 and 3, the estimand is identified by the generalized g-formula²⁷. A re-expression of this generalized g-formula^{22,28} involves recursively defining the expected outcome under the intervention, conditional on the observation's observed exposure and history, beginning at the final time point, and proceeding until the earliest time point. We illustrate the g-formula for two time points below:

- 1. Start with the conditional expectation of the outcome Y given $A_1 = a_1$ and $H_1 = h_1$. Let this function be denoted $Q_1(a_1, h_1)$.
- 2. Evaluate the above conditional expectation of Y if A_1 were changed to $d_1(A_1, H_1)$, which results in a pseudo outcome $\tilde{Y}_1 = Q_1(A_1^d, H_1)$.
- 3. Let the true expectation of \tilde{Y}_1 conditional on $A_0 = a_0$ and $H_0 = h_0$ be denoted $Q_0(a_0, h_0)$.
- 4. Evaluate the above expectation of \tilde{Y}_1 if A_0 were changed to $d_0(A_0, H_0)$, which results in $\tilde{Y}_0 = Q_0(A_0^d, H_0)$.
- 5. Under the identifying assumptions, we have $\mathsf{E}[Y(\bar{A}^{\mathsf{d}}_{\tau})] = \mathsf{E}[\tilde{Y}_{0}]$.

Estimation

Once a causal estimand is defined and identified, the researcher's task is to estimate the statistical quantity, e.g. $E[\tilde{Y}_0]$. We now discuss several estimators, both parametric and non-parametric, and provide their algorithm steps in pseudo-R code in the eAppendix.

Parametric estimation

The simplest option for estimation is to fit parametric outcome regressions for each step of the g-formula identification result. This "plug-in" esimator is often referred to as the parametric g-formula or g-computation. Another option is to use an estimator which relies on the exposure

mechanism, for example, the Inverse Probability Weighting (IPW) estimator. IPW estimation involves reweighting the observed outcome by a quantity which represents the likelihood the intervention was received, conditional on covariates.

Obtaining point estimations with the g-computation and IPW algorithms is computationally straightforward. If the exposure regression for IPW or outcome regression for g-computation are estimated using pre-specified parametric statistical models, standard errors for the estimate can be computed using bootstrapping or the Delta method. However, in causal models with large numbers of covariates and/or complex mathematical relations between confounders, exposures, and outcomes, parametric models are hard to pre-specify, and they are unlikely to consistently estimate the regressions. If the regression for the outcome (for g-computation) or treatment (for IPW) are misspecified, the final estimates will be biased.

One way to alleviate model misspecification is to use flexible approaches which incorporate model selection (e.g. LASSO, splines, boosting, random forests, ensembles thereof, etc.) to estimate the exposure or outcome regressions. Unfortunately, there is generally not statistical theory to support the standard errors of the g-computation or IPW estimators with such data-adaptive regressions. Standard inferential tools such as the bootstrap will fail because these estimators generally do not have an asymptotically normal distribution after using data-adaptive regressions²⁹. Thus, other methods are needed to accommodate both model selection and flexible regression techniques while still allowing for statistical inference.

Non-parametric estimation

Here we will advocate for the use of general, non-parametric estimators for longitudinal modified treatment policies²². These estimators use both an exposure and outcome regression, and allow the use of machine learning to estimate the regressions while still obtaining valid statistical uncertainty quantification on the final estimates. These estimators also remain consistent under inconsistent estimation of at most one of the nuisance parameters.

The two non-parametric estimators encoded in the R package $lmtp^{30,31}$ are Targeted Maximum Likelihood Estimation (TMLE)^{22,29,32} and Sequentially Doubly Robust estimation^{22,33,34}. A third non-parametric estimator, iterative TMLE (iTMLE), is not encoded in the R package but could be adapted from Luedtke et al.³³. TMLE is a doubly robust estimator for a time-varying treatment in the sense that it is consistent as long as all outcome regressions for times t > s are consistently estimated, and all treatment mechanisms for times $t \le s$ are consistent, for some time s. In contrast, sequentially double robust and iTMLE are sequentially doubly robust in that they are consistent if for all times t, either the outcome or the treatment mechanism are consistently estimated^{22,33}. Since TMLE and iTMLE are substitution estimators, they are guaranteed to produce estimates which remain within the observed outcome range. Sequentially double robust and iTMLE produce estimators with more robustness than TMLE.

Table 2 compares the statistical properties of various estimators. We provide practical guidance for choosing between estimation techniques in the Appendix. Of note, for the statistical properties of these estimators to hold, certain technical requirements must be met. These requirements are detailed in the Appendix. All of the aforementioned examples meet these requirements.

Longitudinal modified treatment policies in practice

In the worked example, we estimate the effect of delaying intubation on mortality. Other examples of longitudinal modified treatment policies applications in similar populations include studying the effect of a delay in intubation on an outcome of acute kidney injury, where death is a competing risk³⁵, or studying an intervention on a continuous measure of hypoxia in acute respiratory distress patients²². The methodology for for longitudinal modified treatment policies also accommodates interventions involving multiple treatments, such as delaying intubation by one day *and* increasing fluid intake.

There are multiple other examples of researchers using longitudinal modified treatment policies to answer real-world problems. Nugent and Balzer³⁶ used a longitudinal modified treatment policy to study the effects of mobility on COVID-19 case rates. Specifically, they studied the effect of a longitudinal modified shift on the observed mobility distribution on the number of newly reported cases per 100,000 residents two weeks ahead. A similar analysis could be done to look at interventions on masking or vaccination policies. Mobility, masking, and vaccination are important examples of when static or dynamic policy estimands may be unappealing because of the geographical, political, and cultural variation that exists even within relatively small regions. This applies to environmental exposures and health policies as well. For instance, Rudolph et al.³⁷ used longitudinal modified treatment policies to estimate the effects of Naxolone access laws on opioid overdose rates.

Other examples of longitudinal modified treatment policy applications include Jafarzadeh et al. ³⁸'s study of the effect of an intervention on knee pain scores over time on an outcome of knee replacement surgery. Huling et al. ³⁹ investigated the effects of interventions to public health nursing on the behaviors of clients in the Colorado Nurse Support Program. Mehta et al. ⁴⁰ researched whether an increase in primary care physicians has an effect on post-operative outcomes in patients undergoing elective total joint replacement. Cooray et al. ⁴¹ studied hypothetical interventions on tooth retention and social participation among older adults in Japan. Ikeda et al. ⁴² estimated effects of incremental changes in body mass index percentage on back pain. Tables 3 and 4 show additional applications of longitudinal modified treatment policy applications, as well as examples of possible static and dynamic treatment rules answering similar research questions.

Illustrative example

Motivation

In the following application, we study a clinical question, "what is the effect of delaying intubation of invasive mechanical ventilation on mortality for hospitalized COVID-19 patients in New York City's first COVID-19 wave?" Studying the effect of intubation timing is particularly ill-suited for static interventions because there is no scenario in which intubation at a certain study time could be uniformly applied across all critically ill patients. A dynamic intervention, which could help to evaluate a world in which patients are intubated when they meet a certain severity threshold, e.g. an oxygen saturation breakpoint, is less clinically relevant because intubation guidance may vary considerably between providers dependent on training, hospital policies, and ventilator availabil-ity^{43,44}. For this reason, a longitudinal modified treatment policy that varies the natural time of

intubation by a minimal amount of time may be a more realistic hypothetical intervention to study when considering the mechanistic effect of a delay-in-intubation strategy on mortality.

Measures and analysis

The population of interest is adult patients hospitalized and diagnosed with COVID-19 during Spring 2020. The cohort contains 3,059 patients who were admitted to NewYork-Presbyterian's Cornell, Queens, and Lower Manhattan locations between March 1-May 15, 2020^{35,45,46}. This research was approved with a waiver of informed consent by the Weill Cornell Medicine Institutional Review Board (IRB) 20-04021909.

The exposure of interest is the maximum level of daily supplemental oxygen support. This can take three categories, 0: no supplemental oxygen, 1: non-invasive supplemental oxygen support, and 2: invasive mechanical ventilation. The intervention of interest describes a hypothetical world in which patients who require invasive mechanical ventilation have their intubation delayed by one day, and instead receive non-invasive mechanical ventilation supplemental oxygen support on the observed day of intubation.

$$\mathsf{d}_t(a_t, h_t) = \begin{cases} 1 & \text{if } a_t = 2 \text{ and } a_s \leq 1 \text{ for all } s < t, \\ a_t & \text{otherwise} \end{cases}$$

Since patients in this observational data are subject to loss to follow up via discharge or transfer, the intervention additionally includes observing patients through all 14 days from hospitalization.

The primary outcome is time to death within 14 days from hospitalization. Hospital discharge or transfer to an external hospital system is considered an informative loss to follow-up. The causal estimand of interest is the difference in 14-day mortality rates between a hypothetical world in which there was a one-day delay in intubation and no loss to follow-up, and a hypothetical world in which there was no loss to follow-up and no delay in intubation.

Since the intervention is a longitudinal modified treatment policy, we require positivity and strong sequential randomization to identify our parameter of interest. The common causes we assume to satisfy the latter requirement include 37 baseline confounders and 14 time-varying confounders per time point.

The R package *lmtp*^{30,31} was used to obtain estimates of the difference in 14-day mortality rates under the two proposed interventions using sequentially double robust estimation. A superlearner^{47–49} library of various candidate learners was used to estimate the intervention and outcome estimation. We demonstrate code on synthetic data at https://github.com/kathoffman/lmtptutorial and provide several additional details in the Appendix.

Results

The estimated 14-day mortality incidence under no intervention on intubation was 0.211 (95% CI 0.193-0.229). The same incidence under a hypothetical longitudinal modified treatment policy in which intubation were delayed by 1 day was 0.219 (0.202-0.236). The estimated incidences across all time points are shown in Figure 2.

Figure 2: Panel A: Estimated incidence of mortality between a longitudinal modified treatment policy of delaying intubation (blue) and no intervention (red). Panel B: Estimated incidence difference in mortality if the longitudinal modified treatment policy were implemented during Spring 2020. In both panels, 95% simultaneous confidence bands⁵⁰ cover the sets of point estimates.



Discussion

The longitudinal modified treatment policy methodology allows researchers to define, identify, and estimate scientifically relevant estimands, including those that involve challenges such as loss-to-follow-up, survival analysis, missing exposures, competing risks, and interventions that include multiple exposures and/or continuous exposures. This method is particularly useful for designing estimands with a higher chance of meeting the positivity assumption. In addition, the existing packages that implement doubly and sequentially robust estimators enable researchers to take advantage of statistical learning algorithms to estimate the intervention and outcome mechanisms, thereby increasing the likelihood of estimator consistency.

While the longitudinal modified treatment policy methodology expands the applied researcher's toolbox to accommodate longitudinal interventions which non-trivially depend on the natural value of treatment, there are considerations and limitations to its implementation in realworld applications. First, longitudinal modified treatment policies (as well as static and dynamic interventions in a longitudinal setting) require discretizing time over intervals. Depending on the data collection process, this may cause issues in temporality or loss of data granularity. This discretization may also cause issues in small sample sizes if there are very few outcomes within a certain time point and the applied researcher hopes to estimate the outcome regressions using any statistical learning algorithm which segments the data for training/testing. Second, although this method may be used to formulate alternative estimands that satisfy the respective positivity assumption, the alternative estimand must be scientifically relevant. In addition, practical positivity violations may still occur, and possible solutions, such as truncating the density ratios at a certain threshold, are arbitrary and the potential for bias is unclear. Third, some particular estimator applications may be computationally intensive. Despite these limitations, we hope the explanation of longitudinal modified treatment policies, illustrative example, and corresponding Github repository are a useful toolset for researchers hoping to implement this method in their applied work.

Tables

Table 1: An overview of common intervention types in the causal inference literature and whether they are estimable using the *lmtp* R package.

Intervention and definition	Estimable with <i>lmtp</i> ?
Static: all units receive the same treatment assignment	Yes ^a
Dynamic: A unit's treatment assignment is determined according to their covariate history ^b	Yes ^a
Modified: A unit's treatment assignment is determined according to their natural exposure value, and possibly their covariate history ^{b,c}	See assumptions ^d
	C

^a The software should only be used with discrete exposures. The software will output a result with a continuous exposure, but this estimator will not have good statistical properties.

^b The covariate history could include prior exposure values under the intervention function.

^c A modified treatment policy may also depend on a randomizer.

^d Assumptions require the intervention function d does not depend on the distribution P, and, one of: (1) the exposure is discrete or (2) the exposure is continuous but d satisfies piecewise smooth invertibility. See Technical Requirements in Appendix for more details.

Table 2: Comparison of statistical properties for five possible estimators for longitudinal modified treatment policies: g-computation (G-COMP), inverse probability weighting (IPW), targeted maximum likelihood estimation (TMLE), sequentially doubly robust estimation (SDR), and iterative TMLE (iTMLE).

Statistical Property	G-COMP	IPW	TMLE	SDR	iTMLE ^a
Uses outcome regression	\checkmark		\checkmark	\checkmark	✓
Uses treatment regression		\checkmark	\checkmark	\checkmark	✓
Doubly robust ^b			\checkmark	\checkmark	\checkmark
Sequentially doubly robust ^c				\checkmark	✓
Valid inference ^d using parametric regressions (i.e. generalized linear models)	✓	~	~	√	√
Valid inference ^d using data-adaptive regressions (i.e. machine learning)			~	~	✓
Guaranteed to stay within observed outcome range	\checkmark		\checkmark		\checkmark

^a The iTMLE estimator is not currently available within the R package lmtp.

^b The estimator is consistent as long as all outcome regressions for times t > s are consistently estimated, and all treatment mechanisms for times $t \le s$ are consistent, for some time s.

^c The estimator is consistent if, for every time point, either the outcome regression or the treatment mechanism is consistently estimated.

^d Includes standard errors, confidence intervals, and p-values.

Table 3: Examples of static, dynamic, and modified interventions for (1) binary and (2) continuous point-in-time exposures. These can be expanded to any categorical exposure. We denote a random variable drawn from Bernoulli distribution with probability 0.5 with ϵ unless otherwise noted.

Exposure	Intervention	"What if"	Shift Notation
Point-in-time Binary, e.g. vaping (a = 1)	Static	no one vapes	$d(a,h,\epsilon)=0$
	Dynamic	only those working non-standard work hours $(l^* = 1)$ vape	$d(a,h,\epsilon) = \begin{cases} 1 & \text{ if } l^* = 1 \\ 0 & \text{ otherwise} \end{cases}$
	Modified	a random half of current vapers stop vaping	$d(a,h,\epsilon) = \begin{cases} \epsilon & \text{ if } a = 1\\ a & \text{ otherwise} \end{cases}$
Point-in-time Continuous, e.g. exposure to pollution as measured by the Air Quality Index (AQI) scale	Static	all counties are exposed to an AQI of 10	$d(a,h,\epsilon)=10$
	Dynamic	all urban $(l^* = 1)$ counties are exposed to an AQI of 40 and all rural $(l^* = 0)$ counties are exposed to an AQI of 20	$d(a,h,\epsilon) = \begin{cases} 40 & \text{if } l^* = 1\\ 20 & \text{otherwise} \end{cases}$
	Modified	all counties with an AQI higher than 20 are exposed to an AQI 10% lower than what they were naturally exposed to	$d(a,h,\epsilon) = \begin{cases} a \times 0.9 & \text{if } a > 20 \\ a & \text{otherwise} \end{cases}$

Table 4: Examples of static, dynamic, and modified interventions for a binary time-varying exposure. We index study time at t = 0 and denote a random variable drawn from Bernoulli distribution with probability 0.5 with ϵ_t .

Exposure	Intervention	"What if"	Shift Notation	
Time-varying Binary, e.g. corticosteroids	Static	all patients receive corticosteroids for the first 6 days of hospitalization	$d_t(a_t, h_t, \epsilon_t) = \begin{cases} 1 & \text{ if } t \leq 5\\ 0 & \text{ otherwise} \end{cases}$	
receipt $(a_t = 1)$	Dynamic	patients receive corticosteroids for 6 days once they become hypoxic $(l_t^* = 1)$	$d_t(a_t, h_t, \epsilon_t) = \begin{cases} 1 & \text{if } l_s^* = 1 \text{ for } s \in \{t - 5, \dots, t \\ 0 & \text{otherwise} \end{cases}$	ţ}
	Modified	patients' receipt of corticosteroids is delayed by 1 day	$d_t(a_t,h_t,\epsilon_t) = \begin{cases} 0 & \text{ if } a_t = 1 \text{ and } a_{t-1} = 0 \\ a_t & \text{ otherwise} \end{cases}$	

eAppendix

Additional Intervention Examples

Dynamic interventions

Example 6. [Dynamic antiretroviral initiation for HIV patients] One of the first uses of dynamic interventions was in the context of HIV, where investigators were interested in the effect of initiating antiretroviral therapy for a person with HIV if their CD4 count falls below a threshold, e.g. 200 cells/ μ l⁵¹. This can be described mathematically as

$$\mathsf{d}_t(h_t) = \begin{cases} 1 & \text{if } l_t^* < 200 \\ 0 & \text{otherwise,} \end{cases}$$

where L_t^* is a variable in H_t that denotes CD4 T-cell count.

Stochastic interventions

An additional category of interventions can be defined as functions which vary with some usergiven randomizer ϵ_t (and possibly an individual unit's prior history h_t) but not a_t . We will call these stochastic interventions. There are many real world examples of stochastic interventions, such as the lottery treatments of military drafts⁵², twin births⁵³, and college roommate assignments⁵⁴. Here we will focus on two stochastic interventions as they relate to the natural value of treatment. Of note, we do not consider regimes where ϵ has a distribution that depends on P.

Example 7 (Incremental propensity score interventions based on the odds ratio). A recently developed type of stochastic interventions is Kennedy⁵⁵'s incremental propensity score interventions (IPSI), which shifts a unit's probability of receiving treatment conditional on their according to some constant δ . First, a shifted propensity score is defined:

$$\pi_t^{\mathsf{d}}(h_t) = \frac{\delta \pi_t(h_t)}{\delta \pi_t(h_t) + 1 - \pi_t(h_t)}$$

Then, a random variable ϵ_t is drawn from a uniform distribution in (0,1) and the intervention is defined as:

$$\mathsf{d}_t(h_t, \epsilon_t, \pi_t) = I(\epsilon_t < \pi_t^{\mathsf{d}}(h_t)),$$

This IPSI is said to be based on an odds ratio because the odds ratio of $\pi_t^d(h_t)$ to $\pi_t(h_t)$ is equal to δ . Naimi et al.⁵⁶ used an IPSI to study the causal relationship between vegetable density consumption and the risk of preeclampsia among pregnant women. They studied whether preeclampsia would increase if women's propensity of eating a minimum amount of vegetables increased by an odds ratio of δ , for example, $\delta = 1.5$ would mean a woman with a propensity of 0.35 increases to 0.45.

Modified Treatment Policies

Example 8 (Incremental propensity score based on the risk ratio). Another type of IPSI begins with a slightly different set-up than Kennedy⁵⁵. Instead of relying on the treatment mechanism,

there is a random draw ϵ_t from a Uniform (0,1) distribution. If the draw is less than some δ , then the treatment assignment is the natural value of treatment. If not, it is some constant value, for example 0.

$$\mathsf{d}_t(a_t, \epsilon_t) = \begin{cases} a_t \text{ if } \epsilon_t < \delta, \\ 0 \text{ otherwise} \end{cases}$$

This intervention is said to be based on the risk ratio because $\delta = \pi_t^{d}(h_t)/\pi_t(h_t)$, where $\pi_t^{d}(h_t)$ is the post-intervention propensity score and $\pi_t(h_t)$ is the propensity score in the observed data generating mechanism. This intervention is motivated by Wen et al.⁵⁷, who propose an intervention where treatment is a random draw from the shifted distribution and results in the same identification formula. Wen et al.⁵⁷ argue the risk ratio interpretation may be more intuitive for collaborators, and they demonstrate their intervention in an application studying the effect of PrEP usage increases on sexually transmitted infection rates.

Estimation Algorithms

G-Computation Algorithm

The G-Computation algorithm steps and pseudo-code for a simple example with two time points is shown below.

1. Fit a generalized linear model (GLM) for Y conditional on $A_1 = a_1$ and $H_1 = h_1$. Call this $\hat{Q}_1(a_1, h_1)$.

 $Q1_hat \leftarrow glm(outcome = Y, predictors = \{H_1, A_1\})$

2. Modify the data set used in step (1) so that the values in the column for A_1 are changed to A_1^d . Obtain the predictions for the model \hat{Q}_1 using this modified data set. These are pseudo-outcomes \tilde{Y}_1 .

pseudo_Y1 \leftarrow predict(fit = Q1_hat, new data = { $H_1, A_1 = A_1^d$ })

3. Fit a generalized linear model (GLM) for \tilde{Y}_1 conditional on $A_0 = a_0$ and $H_0 = h_0$. Call this $\hat{Q}_0(a_0, h_0)$.

 $QO_hat \leftarrow glm(outcome = pseudo_Y1, predictors = \{H_0, A_0\})$

4. Modify the data set used in step (3) so that the values in the column for A_0 are changed to A_0^d . Obtain the predictions for the model \hat{Q}_0 using this modified data set. These are pseudo-outcomes \tilde{Y}_0 .

pseudo_Y0 \leftarrow predict(fit = Q1_hat, new data = { $H_0, A_0 = A_0^d$ })

5. Average \tilde{Y}_0 , i.e. compute $\hat{\mathsf{E}}[\tilde{Y}_0]$.

 $estimate \leftarrow mean(pseudo_Y0)$

Inverse Probability Weighting Algorithm

The IPW estimator relies on a density ratio $r_t = g_t^d(a_t \mid h_t)/g_t(a_t \mid h_t)$, where $g_t^d(a_t \mid h_t)$ is the density of the intervened exposure, and $g_t(a_t \mid h_t)$ is the density of the naturally observed exposure. Practically, r_t can be computed using a clever classification trick proposed in Qin⁵⁸, Cheng and Chu⁵⁹ and utilized in Díaz et al.²². The IPW algorithm with pseudo-code for a simple example with two time points is as follows.

1. Duplicate each row of the data set so that there are 2n rows. The first row of a duplicated pair should contain observed values A_1 , and the second row should be modified so that $A_1 = A_1^d$. A new column Λ_1 should be created that is 0 if $A_1 = A_1$ and 1 if $A_1 = A_1^d$.

```
\begin{split} & \texttt{data} \leftarrow \texttt{matrix}(\{H_1, A_1, \Lambda_1 = 0\}\}) \\ & \texttt{data\_copy} \leftarrow \texttt{matrix}(\{H_1, A_1 = A_1^\texttt{d}, \Lambda_1 = 1\}) \\ & \texttt{dr\_data} \leftarrow \texttt{bind\_rows}(\texttt{data, data\_copy}) \end{split}
```

- 2. Fit a logistic regression for Λ_1 conditional on $A_1 = a_1$ and $H_1 = h_1$. The odds ratio at a given a_1 and h_1 is the estimate of the density ratio at time 1, \hat{r}_1 .
 - $r1_fit \leftarrow glm(outcome = \Lambda_1, predictors = \{H_1, A_1\}, data = dr_data)$
 - $r1_hat \leftarrow predict(fit = r1_fit, prediction = odds ratio, new data = data)$
- 3. Create a new column in the duplicated data set which contains values of A_0 in the first row of a duplicated pair. The second row in a pair should be modified so that $A_0 = A_0^d$. A new column Λ_0 should be created that is 0 if $A_0 = A_0$ and 1 if $A_0 = A_0^d$.



Figure 3: Illustration of the data used for density ratio estimation with the classification trick.

4. Fit a logistic regression for Λ_0 conditional on $A_0 = a_0$ and $H_0 = h_0$. The odds ratio at a given a_0 and h_0 is the estimate of the density ratio at time 0, \hat{r}_0 .

 $r0_fit \leftarrow glm(outcome = \Lambda_0, predictors = \{H_0, A_0\}, data = dr_data)$

 $r0_hat \leftarrow predict(fit = r0_fit, prediction = odds ratio, new data = data)$

5. Multiply \hat{r}_1 and \hat{r}_0 together. This is the cumulative density ratio. Multiply the cumulative density ratio by Y. Compute the average. This is $\hat{E}[\tilde{Y}_0]$.

 $\texttt{estimate} \gets \texttt{mean}(\texttt{r1_hat} \ \texttt{*} \ \texttt{r0_hat} \ \texttt{*} \ Y)$

Targeted Maximum Likelihood Estimation (TMLE) Algorithm

- 1. Perform the same steps as for the IPW algorithm to produce estimates of \hat{r}_1 and \hat{r}_0 .
- 2. Fit a generalized linear model (GLM) for Y conditional on $A_1 = a_1$ and $H_1 = h_1$. Call this $\hat{Q}_1(a_1, h_1)$.

 $Q1_hat \leftarrow glm(outcome = Y, predictors = \{H_1, A_1\})$

3. Modify the data set used in step (2) so that the values in the column for A_1 are changed to A_1^d . Obtain predictions for the model \hat{Q}_1 using this modified data set and when $A_0 = a_0$ and $H_0 = h_0$.

 $\texttt{Y1d} \gets \texttt{predict}(\texttt{fit} = \texttt{Q1_hat}, \texttt{new data} = \{H_1, A_1 = A_1^{\texttt{d}}\})$

 $\texttt{Y1} \leftarrow \texttt{predict}(\texttt{fit} = \texttt{Q1_hat}, \texttt{new data} = \{H_1, A_1\})$

4. Fit a logistic regression model for Y with an offset equal to the logit transformation of the expected value of Y under model $\hat{Q}_1(a_1, h_1)$ and weights \hat{r}_1 . Call this $\tilde{Q}_1(a_1, h_1)$.

 $Q1_tilde \leftarrow glm(outcome = Y, offset = qlogis(Y1), weights = r1_hat)$

- 5. Update the predictions Y1d using $\tilde{Q}_1(a_1, h_1)$. These are pseudo-outcomes \tilde{Y}_1 . pseudo_Y1 \leftarrow plogis(qlogis(Y1d) + coef(Q1_tilde))
- 6. Repeat steps 2 to 5 replacing Y, A_1 , H_1 , and $\hat{r_1}$ with \tilde{Y}_1 , A_0 , H_0 , and $\hat{r_0} \times \hat{r_1}$ to obtain \tilde{Y}_0 .
- 7. Compute $E[\tilde{Y}_0]$. estimate \leftarrow mean(pseudo_Y0)

Sequentially Doubly Robust (SDR) Estimation Algorithm

- 1. Perform steps 1 to 3 as for the TMLE algorithm.
- 2. Using Y, \hat{r}_1 , Y1, and Y1d apply the SDR transformation from Díaz et al.²² to obtain the pseudo-outcome \tilde{Y}_1 .

 $\texttt{pseudo}_{\texttt{Y1}} \leftarrow \texttt{r1}_{\texttt{hat}} \ \texttt{*} (Y \texttt{-} \texttt{Y1}) \texttt{+} \ \texttt{Y1d}$

3. Fit a generalized linear model (GLM) for \tilde{Y}_1 conditional on $A_0 = a_0$ and $H_0 = h_0$. Call this $\hat{Q}_0(a_0, h_0)$.

 $\texttt{QO_hat} \gets \texttt{glm}(\texttt{outcome} = \texttt{pseudo}_\texttt{Y1}, \texttt{predictors} = \{H_0, A_0\})$

4. Modify the data set used in step (3) so that the values in the column for A_0 are changed to a_0^d . Obtain predictions for the model \hat{Q}_0 using this modified data set and when $A_0 = a_0$ and $H_0 = h_0$.

YOd ← predict(fit = Q0_hat, new data = $\{H_0, A_0 = a_0^d\}$) YO ← predict(fit = Q0_hat, new data = $\{H_0, A_0\}$)

- $10 \leftarrow \text{predict}(\text{In} = \text{QO}_{\text{In}}, \text{new data} = \{11_0, 11_0\}$
- 5. Using pseudo_Y1, Y0d, Y0, r1_hat, and r0_hat obtain an estimate of the un-centered efficient influence function, $\hat{\phi}_1$.

```
\begin{array}{l} \texttt{m\_d} \leftarrow \texttt{matrix}(\texttt{Y1d}, Y) \\ \texttt{m} \leftarrow \texttt{matrix}(\texttt{Y0}, \texttt{Y1}) \\ \texttt{r} \leftarrow \texttt{matrix}(\texttt{r0\_hat} \ * \ \texttt{r1\_hat}, \texttt{r1\_hat}) \\ \texttt{phi\_hat} \leftarrow \texttt{rowSums}(\texttt{r} \ * (\texttt{m\_d} \ - \ \texttt{m})) + \ \texttt{Y0d} \end{array}
```

6. Compute $\mathsf{E}[\hat{\phi}_1]$

 $\texttt{estimate} \gets \texttt{mean}(\texttt{phi_hat})$

Technical requirements for theoretical guarantees of estimators

The \sqrt{n} -consistency and asymptotic normality of the estimators proposed in Díaz et al.²², and therefore the correctness of the confidence intervals output by the R software $lmtp^{30,31}$, relies on two important technical requirements. First, the function d must not depend on the distribution of the data. Second, the exposure must be discrete or, if it is continuous, $d(\cdot, h_t, \epsilon_t)$ (as a function of a_t) must be piecewise smooth invertible^{20,22}. If these requirements are violated, important theoretical properties of the estimators such as \sqrt{n} -consistent will fail. This means that uncertainty quantification measures such as confidence intervals, p-values, and standard errors outputted by the package will be incorrect.

If the first requirement is violated, it may be possible to develop \sqrt{n} -consistent estimators. For example, Kennedy⁵⁵ developed non-parametric \sqrt{n} -consistent estimators for the IPSI shown in Example 7, in which d depends on the treatment mechanism. However, the IPSI in Example 8, which does not depend on the data distribution, can be estimated with the proposed estimators. Here it is important to note that the the estimators for the odds ratio IPSI of Kennedy⁵⁵ are not doubly robust (and such estimators possibly cannot be constructed, since d depends on the propensity score), whereas the estimators of the risk-ratio IPSI will be doubly robust. The threshold function as it is in Example 3 meets these technical requirements because the exposure is discrete, however, if the exposure were instead continuous, it would not meet the second requirement because a threshold d is not piecewise smooth invertible. However, the modified treatment policies in Examples 4 and 5 meet both requirements and are estimable using the proposed estimators.

Practical guidance and considerations

In practice we recommend using TMLE if the user is not concerned about model misspecification, because the estimates are guaranteed to remain within the observed outcome bounds. However, if model misspecification is a concern, SDR is significantly more robust, although there is a potential for estimates which fall outside the outcome bounds. An effective approach against model misspecification is to use the superlearner algorithm for estimating the exposure and outcome mechanisms. Superlearning combines the predictions of multiple pre-specified statistical learning models via weighting to produce final estimates which are proven to perform as well as possible in large sample sizes⁴⁹.

To assess positivity violations, we recommend the researcher to visually inspect or examine summary statistics of the density ratios. Extremely high values of density ratios (the quantification of "high" being context-dependent) are akin to propensity score values at or near zero, generally indicate positivity violations, and may create unstable estimates. If violations are detected, the researcher may be interested in modifying the portion of the population receiving the intervention, or in cases of continuous exposures, making the intervened exposure level closer to the observed exposure level. Interventions should be designed to avoid both structural and practical positivity violations present in the data, the latter of which can potentially be caught in the exploratory data analysis stage. One option if the researcher cannot find a solution by study design to eliminate positivity violations is to truncate the density ratios as a certain threshold, e.g. the 98th quantile. This is akin to truncating a propensity score, and may have potential for biases⁶⁰.

Analysts using longitudinal modified treatment policies for time-varying or survival outcomes will also note that discrete outcome intervals are required. The researcher must find a balance in choosing intervals that are both scientifically relevant (e.g. week-long intervals for a critically ill patient population would be too long) and in which there are still a number of outcomes which occur within the chosen interval. If the outcome is rare and the sample size is small, or if the sample size is very large and computationally intensive, the researcher may need to adjust the number of cross-fitting and/or superlearning cross-validation folds.

Additional Application Details

Motivation

The motivation for studying a delay in intubation policy for COVID-19 patients is further detailed in Díaz et al. ³⁵'s illustrative application section.

Methods

Baseline confounders include age, sex, race, ethnicity, body mass index (BMI), comorbidities (cerebral vascular event, hypertension, diabetes mellitus, cirrhosis, chronic obstructive pulmonary disease, active cancer, asthma, interstitial lung disease, chronic kidney disease, immunosuppression, HIV-infection, and home oxygen use), and hospital admission location. Time-dependent confounders include vital signs (heart rate, pulse oximetry percentage, respiratory rate, temperature, systolic/diastolic blood pressure), laboratory results (blood urea nitrogen (BUN)- creatinine ratio, creatinine, neutrophils, lymphocytes, platelets, bilirubin, blood glucose, D-dimers, C-reactive protein, activated partial thromboplastin time, prothrombin time, arterial partial pressures of oxygen

and carbon dioxide), and concurrent pharmaceutical treatments. Concurrent treatments included vasopressors, diuretics, Angiotensin-converting enzyme (ACE) and Angiotensin receptor blockers (ARBs), hydroxychloroquine, and tocilizumab.

In the case of multiple time-varying confounders measured in one day, the clinically worst value was used. Although data was relatively complete due to manual abstraction efforts, there were instances of patients missing laboratory results at one or multiple time points. A combination of last observation carried forward and an indicator for missing values was used to handle this informative missingness within the treatment and outcome regression models⁶¹.

The Superlearner ensemble algorithm utilized 5-fold cross-validation and candidate libraries included generalized linear models, multivariate adaptive regression splines⁶², random forests⁶³, and extreme gradient boosted trees⁶⁴ An assumption was made that the time-varying covariates from the previous two days (i.e. lag of 2 days) was sufficient to capture the mechanisms to reflect the collection of laboratory results at minimum 48-hour intervals. A 5-fold cross-fitting component was implemented for the final estimator to prevent variation in a certain sample split from biasing the final estimate⁶⁵. All analyses were conducted in R Version 4.1.2 with the packages tidyverse⁶⁶ for data cleaning and plotting, and Imtp³⁰ and SuperLearner⁶⁷ for estimation.

Discussion

There are limitations specific to our illustrative example. While the estimated effect of a less aggressive intubation strategy can help to understand an underlying biological or mechanistic process, it may not provide clinical guidance if the treatment strategy changes over time. The estimates themselves depend on the natural value of treatment, and this is dependent on the state of clinical practice during the study time frame (Spring 2020). In addition, we cannot rule out unmeasured confounding in the exposure, outcome, and loss to follow-up mechanisms. We also cannot be sure whether the informative right censoring is correctly specified, since patients were lost to follow-up for different reasons (e.g. discharge to home vs. assisted living).

Further reading

Researchers intending to learn more about longitudinal modified treatment policies may find the original longitudinal modified treatment policy methods research by Díaz et al.²² useful, as well the extension to competing risks (with a similar application as this tutorial) in Díaz et al.³⁵. The R software *lmtp* is explained in Williams and Díaz³¹.

The longitudinal modified treatment policy methodology is based upon decades of work in epidemiology, biostatistics, and related fields. Interested readers should see related methods research and applications such as Young et al.¹⁵, Robins et al.¹⁶, Taubman et al.¹⁷, Díaz and van der Laan¹⁹, Haneuse and Rotnitzky²⁰, Richardson and Robins²¹.

References

- [1] Jamuna K. Krishnan, Mangala Rajan, Benjamin R. Baer, Katherine L. Hoffman, Mark N. Alshak, Kerri I. Aronson, Parag Goyal, Chiomah Ezeomah, Shanna S. Hill, Fernando J. Martinez, Meredith L. Turetz, Martin T. Wells, Monika M. Safford, and Edward J. Schenck. Assessing mortality differences across acute respiratory failure management strategies in covid-19. *Journal of Critical Care*, 70:154045, 2022. ISSN 0883-9441. doi: https://doi.org/10.1016/j.jcrc.2022.154045. URL https://www.sciencedirect.com/science/article/pii/S0883944122000740.
- [2] Jaime B Hyman, Evan S Leibner, Pranai Tandon, Natalia N Egorova, Adel Bassily-Marcus, Roopa Kohli-Seth, Varun Arvind, Helena L Chang, Hung-Mo Lin, and Matthew A Levin. Timing of intubation and in-hospital mortality in patients with coronavirus disease 2019. *Critical care explorations*, 2(10), 2020.
- [3] Yong Hoon Lee, Keum-Ju Choi, Sun Ha Choi, Shin Yup Lee, Kyung Chan Kim, Eun Jin Kim, and Jaehee Lee. Clinical significance of timing of intubation in critically ill patients with covid-19: a multi-center retrospective study. *Journal of clinical medicine*, 9(9):2847, 2020.
- [4] Benjamin McKay, Matthew Meyers, Leah Rivard, Holly Stankewicz, Jill C Stoltzfus, and Guhan Rammohan. Comparison of early and late intubation in covid-19 and its effect on mortality. *International Journal of Environmental Research and Public Health*, 19(5):3075, 2022.
- [5] Denio A Ridjab, Ignatius Ivan, Fanny Budiman, and Dafsah A Juzar. Outcome in early vs late intubation among covid-19 patients with acute respiratory distress syndrome: an updated systematic review and meta-analysis. *Scientific reports*, 12(1):21588, 2022.
- [6] Eleni Papoutsi, Vassilis G Giannakoulis, Eleni Xourgia, Christina Routsi, Anastasia Kotanidou, and Ilias I Siempos. Effect of timing of intubation on clinical outcomes of critically ill patients with covid-19: a systematic review and meta-analysis of non-randomized cohort studies. *Critical Care*, 25:1–9, 2021.
- [7] Alfonso C Hernandez-Romieu, Max W Adelman, Maxwell A Hockstein, Chad J Robichaux, Johnathan A Edwards, Jane C Fazio, James M Blum, Craig S Jabaley, Mark Caridi-Scheible, Greg S Martin, et al. Timing of intubation and mortality among critically ill coronavirus disease 2019 patients: a single-center cohort study. *Critical care medicine*, 2020.
- [8] Atul Matta, Siddique Chaudhary, Kevin Bryan Lo, Robert DeJoy III, Fahad Gul, Ricardo Torres, Neal Chaisson, and Gabriel Patarroyo-Aponte. Timing of intubation and its implications on outcomes in critically ill patients with coronavirus disease 2019 infection. *Critical care explorations*, 2(10), 2020.
- [9] Aloknath Pandya, Navjot Ariyana Kaur, Daniel Sacher, Oisin O'Corragain, Daniel Salerno, Parag Desai, Sameep Sehgal, Matthew Gordon, Rohit Gupta, Nathaniel Marchetti, et al. Ventilatory mechanics in early vs late intubation in a cohort of coronavirus disease 2019 patients with ards: a single center's experience. *Chest*, 159(2):653–656, 2021.

- [10] Ricard Mellado-Artigas, Bruno L Ferreyro, Federico Angriman, María Hernández-Sanz, Egoitz Arruti, Antoni Torres, Jesús Villar, Laurent Brochard, and Carlos Ferrando. Highflow nasal oxygen in patients with covid-19-associated acute respiratory failure. *Critical Care*, 25(1):1–10, 2021.
- [11] Avni A Bavishi, Ruben J Mylvaganam, Rishi Agarwal, Ryan J Avery, Michael J Cuttica, NU COVID Investigators, et al. Timing of intubation in coronavirus disease 2019: a study of ventilator mechanics, imaging, findings, and outcomes. *Critical Care Explorations*, 3(5): e0415, 2021.
- [12] Mohammad Ali Mansournia, Mahyar Etminan, Goodarz Danaei, Jay S Kaufman, and Gary Collins. Handling time varying confounding in observational research. *bmj*, 359, 2017.
- [13] MA Hernán and JM Robins. *Causal Inference: What If.* Boca Raton: Chapman & Hall, 2023.
- [14] Maya L Petersen, Kristin E Porter, Susan Gruber, Yue Wang, and Mark J van der Laan. Diagnosing and responding to violations in the positivity assumption. *Statistical methods in medical research*, 21(1):31–54, 2012.
- [15] Jessica G. Young, Miguel A. Hernán, and James M. Robins. Identification, estimation and approximation of risk under interventions that depend on the natural value of treatment using observational data. *Epidemiologic Methods*, 3(1):1–19, 2014. doi: doi:10.1515/ em-2012-0001. URL https://doi.org/10.1515/em-2012-0001.
- [16] James M Robins, Miguel A Hernán, and Uwe Siebert. Effects of multiple interventions. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*, 1:2191–2230, 2004.
- [17] Sarah L Taubman, James M Robins, Murray A Mittleman, and Miguel A Hernán. Intervening on risk factors for coronary heart disease: an application of the parametric g-formula. *International journal of epidemiology*, 38(6):1599–1611, 2009.
- [18] Ilya Shpitser and Judea Pearl. Effects of treatment on the treated: Identification and generalization. *arXiv preprint arXiv:1205.2615*, 2012.
- [19] Iván Díaz and Mark van der Laan. Population intervention causal effects based on stochastic interventions. *Biometrics*, 68(2):541–549, 2012.
- [20] Sebastian Haneuse and Andrea Rotnitzky. Estimation of the effect of interventions that modify the received treatment. *Statistics in medicine*, 32(30):5260–5277, 2013.
- [21] Thomas S Richardson and James M Robins. Single world intervention graphs (swigs): A unification of the counterfactual and graphical approaches to causality. *Center for the Statistics and the Social Sciences, University of Washington Series. Working Paper*, 128(30):65–78, 2013.

- [22] Iván Díaz, Nicholas Williams, Katherine L. Hoffman, and Edward J. Schenck. Nonparametric Causal Effects Based on Longitudinal Modified Treatment Policies. *Journal of the American Statistical Association*, pages 1–16, September 2021. ISSN 0162-1459, 1537-274X. doi: 10.1080/01621459.2021.1955691. URL https://www.tandfonline.com/doi/full/10. 1080/01621459.2021.1955691.
- [23] Judea Pearl. Graphs, causality, and structural equation models. *Sociological Methods & Research*, 27(2):226–284, 1998.
- [24] Judea Pearl, Madelyn Glymour, and Nicholas P Jewell. Causal inference in statistics: A primer. 2016. *Google Ascholar there is no corresponding record for this reference*, 2016.
- [25] Katherine L Hoffman, Edward J Schenck, Michael J Satlin, William Whalen, Di Pan, Nicholas Williams, and Iván Díaz. Comparison of a target trial emulation framework vs cox regression to estimate the association of corticosteroids with covid-19 mortality. JAMA Network Open, 5(10):e2234425–e2234425, 2022.
- [26] Mark J van der Laan and Sherri Rose. *Targeted learning in data science*, chapter 14, pages 219–232. Springer, 2018.
- [27] James M Robins. A new approach to causal inference in mortality studies with sustained exposure periods - application to control of the healthy worker survivor effect. *Mathematical Modelling*, 7:1393–1512, 1986.
- [28] Heejung Bang and James M Robins. Doubly robust estimation in missing data and causal inference models. *Biometrics*, 61(4):962–973, 2005.
- [29] Mark J van der Laan, Sherri Rose, et al. *Targeted learning: causal inference for observational and experimental data*, volume 10. Springer, 2011.
- [30] Nicholas T Williams and Iván Díaz. Imtp: Non-parametric Causal Effects of Feasible Interventions Based on Modified Treatment Policies, 2021. URL https://github.com/ nt-williams/lmtp. R package version 1.3.1.
- [31] Nicholas Williams and Iván Díaz. lmtp: An r package for estimating the causal effects of modified treatment policies. *Observational Studies*, 9(2):103–122, 2023.
- [32] Mark J van der Laan and Susan Gruber. Targeted minimum loss based estimation of causal effects of multiple time point interventions. *The international journal of biostatistics*, 8(1), 2012.
- [33] Alexander R Luedtke, Oleg Sofrygin, Mark J van der Laan, and Marco Carone. Sequential double robustness in right-censored longitudinal models. arXiv preprint arXiv:1705.02459, 2017.
- [34] Andrea Rotnitzky, James Robins, and Lucia Babino. On the multiply robust estimation of the mean of the g-functional. *arXiv preprint arXiv:1705.08582*, 2017.

- [35] Iván Díaz, Katherine L Hoffman, and Nima S Hejazi. Causal survival analysis under competing risks using longitudinal modified treatment policies. *arXiv preprint arXiv:2202.03513*, 2022.
- [36] Joshua R Nugent and Laura B Balzer. Evaluating shifts in mobility and covid-19 case rates in us counties: A demonstration of modified treatment policies for causal inference with continuous exposures. *arXiv preprint arXiv:2110.12529*, 2021.
- [37] Kara E Rudolph, Catherine Gimbrone, Ellicott C Matthay, Iván Díaz, Corey S Davis, Katherine Keyes, and Magdalena Cerdá. When effects cannot be estimated: redefining estimands to understand the effects of naloxone access laws. *Epidemiology*, 33(5):689–698, 2022.
- [38] S Reza Jafarzadeh, Tuhina Neogi, Daniel K White, and David T Felson. The relationship of pain reduction with prevention of knee replacement under dynamic intervention strategies. *Arthritis & Rheumatology*, 74(10):1668–1675, 2022.
- [39] Jared D Huling, Robin R Austin, Sheng-Chieh Lu, Mary M Doran, Vicki J Swarr, and Karen A Monsen. Public health nurse tailored home visiting and parenting behavior for families at risk for referral to child welfare services, colorado: 2018–2019. *American Journal of Public Health*, 112(S3):S306–S313, 2022.
- [40] Bella Mehta, Collin Brantner, Nicholas Williams, Jackie Szymonifka, Iris Navarro-Millan, Lisa A Mandl, Anne R Bass, Linda A Russell, Michael L Parks, Mark P Figgie, et al. Primary care provider density and elective total joint replacement outcomes. *Arthroplasty today*, 10: 73–78, 2021.
- [41] U. Cooray, G. Tsakos, A. Heilmann, R.G. Watt, K. Takeuchi, K. Kondo, K. Osaka, and J. Aida. Impact of teeth on social participation: Modified treatment policy approach. *Journal of Dental Research*, 102(8):887–894, 2023. doi: 10.1177/00220345231164106. URL https://doi.org/10.1177/00220345231164106. PMID: 37085984.
- [42] Takaaki Ikeda, Upul Cooray, Yuta Suzuki, Anna Kinugawa, Masayasu Murakami, and Ken Osaka. Changes in Body Mass Index on the Risk of Back Pain: Estimating the Impacts of Weight Gain and Loss. *The Journals of Gerontology: Series A*, 78(6):973–979, 09 2022. ISSN 1758-535X. doi: 10.1093/gerona/glac184. URL https://doi.org/10.1093/gerona/glac184.
- [43] Martin J Tobin, Franco Laghi, and Amal Jubran. Caution about early intubation and mechanical ventilation in covid-19. *Annals of intensive care*, 10(1):1–3, 2020.
- [44] Gavin D Perkins, Keith Couper, Bronwen Connolly, J Kenneth Baillie, Judy M Bradley, Paul Dark, Anthony De Soyza, Ellen Gorman, Alasdair Gray, Louisa Hamilton, et al. Recovery-respiratory support: respiratory strategies for patients with suspected or proven covid-19 respiratory failure; continuous positive airway pressure, high-flow nasal oxygen, and standard care: a structured summary of a study protocol for a randomised controlled trial. *Trials*, 21: 1–3, 2020.

- [45] Parag Goyal, Justin J Choi, Laura C Pinheiro, Edward J Schenck, Ruijun Chen, Assem Jabri, Michael J Satlin, Thomas R Campion Jr, Musarrat Nahid, Joanna B Ringel, et al. Clinical characteristics of covid-19 in new york city. *New England Journal of Medicine*, 382(24): 2372–2374, 2020.
- [46] Edward J Schenck, Katherine L Hoffman, Marika Cusick, Joseph Kabariti, Evan T Sholle, and Thomas R Campion Jr. Critical care database for advanced research (cedar): An automated method to support intensive care units with electronic health record data. *Journal of Biomedical Informatics*, 118:103789, 2021.
- [47] David H Wolpert. Stacked generalization. Neural Networks, 5(2):241–259, 1992.
- [48] Leo Breiman. Stacked regressions. *Machine Learning*, 24(1):49–64, 1996.
- [49] Mark J van der Laan, Eric C Polley, and Alan E Hubbard. Super learner. Statistical Applications in Genetics & Molecular Biology, 6(25):Article 25, 2007.
- [50] Ted Westling, Mark J van der Laan, and Marco Carone. Correcting an estimator of a multivariate monotone function with isotonic regression. *Electronic journal of statistics*, 14(2): 3032, 2020.
- [51] Miguel A Hernán, Emilie Lanoy, Dominique Costagliola, and James M Robins. Comparison of dynamic treatment regimes via inverse probability weighting. *Basic & clinical pharmacology & toxicology*, 98(3):237–242, 2006.
- [52] Joshua D Angrist. Lifetime earnings and the vietnam era draft lottery: evidence from social security administrative records. *The american economic review*, pages 313–336, 1990.
- [53] Matt McGue, Merete Osler, and Kaare Christensen. Causal inference and observational research: The utility of twins. *Perspectives on psychological science*, 5(5):546–556, 2010.
- [54] Bruce Sacerdote. Peer effects with random assignment: Results for dartmouth roommates. *The Quarterly journal of economics*, 116(2):681–704, 2001.
- [55] Edward H Kennedy. Nonparametric causal effects based on incremental propensity score interventions. *Journal of the American Statistical Association*, 114(526):645–656, 2019.
- [56] Ashley I Naimi, Jacqueline E Rudolph, Edward H Kennedy, Abigail Cartus, Sharon I Kirkpatrick, David M Haas, Hyagriv Simhan, and Lisa M Bodnar. Incremental propensity score effects for time-fixed exposures. *Epidemiology (Cambridge, Mass.)*, 32(2):202, 2021.
- [57] Lan Wen, Julia L Marcus, and Jessica G Young. Intervention treatment distributions that depend on the observed treatment process and model double robustness in causal survival analysis. *Statistical Methods in Medical Research*, page 09622802221146311, 2023.
- [58] Jing Qin. Inferences for case-control and semiparametric two-sample density ratio models. *Biometrika*, 85(3):619–630, 1998.
- [59] Kuang Fu Cheng and Chih-Kang Chu. Semiparametric density estimation under a twosample density ratio model. *Bernoulli*, 10(4):583–604, 2004.

- [60] Maxime Léger, Arthur Chatton, Florent Le Borgne, Romain Pirracchio, Sigismond Lasocki, and Yohann Foucher. Causal inference in case of near-violation of positivity: comparison of methods. *Biometrical Journal*, 64(8):1389–1403, 2022.
- [61] A Burton and DG Altman. Missing covariate data within cancer prognostic studies: a review of current reporting and proposed guidelines. *British journal of cancer*, 91(1):4–8, 2004.
- [62] Stephen Milborrow. earth: Multivariate Adaptive Regression Splines, 2021. URL https: //CRAN.R-project.org/package=earth. R package version 5.3.1.
- [63] Marvin N. Wright and Andreas Ziegler. ranger: A fast implementation of random forests for high dimensional data in C++ and R. *Journal of Statistical Software*, 77(1):1–17, 2017. doi: 10.18637/jss.v077.i01.
- [64] Tianqi Chen, Tong He, Michael Benesty, Vadim Khotilovich, Yuan Tang, Hyunsu Cho, Kailong Chen, Rory Mitchell, Ignacio Cano, Tianyi Zhou, Mu Li, Junyuan Xie, Min Lin, Yifeng Geng, and Yutian Li. *xgboost: Extreme Gradient Boosting*, 2021. URL https: //CRAN.R-project.org/package=xgboost. R package version 1.5.0.2.
- [65] Paul N Zivich and Alexander Breskin. Machine learning for causal inference: on the use of cross-fit estimators. *Epidemiology (Cambridge, Mass.)*, 32(3):393, 2021.
- [66] Hadley Wickham, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino Mc-Gowan, Romain François, Garrett Grolemund, Alex Hayes, Lionel Henry, Jim Hester, Max Kuhn, Thomas Lin Pedersen, Evan Miller, Stephan Milton Bache, Kirill Müller, Jeroen Ooms, David Robinson, Dana Paige Seidel, Vitalie Spinu, Kohske Takahashi, Davis Vaughan, Claus Wilke, Kara Woo, and Hiroaki Yutani. Welcome to the tidyverse. *Journal of Open Source Software*, 4(43):1686, 2019. doi: 10.21105/joss.01686.
- [67] Eric Polley, Erin LeDell, Chris Kennedy, and Mark van der Laan. SuperLearner: Super Learner Prediction, 2021. URL https://CRAN.R-project.org/package= SuperLearner. R package version 2.0-28.