

A short proof of the surjectivity of the period map on K3 manifolds

Hongyi Liu

Abstract

In this note, we give a simple proof of the Todorov's surjectivity result on the period map of K3 surfaces in a differential geometric setting. Our proof makes use of collapsing geometry of hyperkähler 4-manifolds developed by Sun-Zhang in [4], and does not rely on the solution to the Calabi conjecture.

On an oriented smooth 4-manifold, a hyperkähler metric g is a Riemannian metric with holonomy contained in $SU(2)$. This is equivalent to saying that the bundle of self-dual forms is flat and trivial, which implies that there is a triple $\omega = (\omega_1, \omega_2, \omega_3)$ of closed 2-forms satisfying $\omega_\alpha \wedge \omega_\beta = 2\delta_{\alpha\beta} \text{dVol}_g$. Such a triple is often referred to as a hyperkähler triple.

Let \mathcal{K} be the K3 manifold, which is by definition the unique oriented smooth 4-manifold underlying a complex K3 surface. It is simply connected and the intersection form on $\Lambda \equiv H^2(\mathcal{K}; \mathbb{Z})$ has signature $(3, 19)$. Denote by \mathcal{N} the set of all hyperkähler metrics on \mathcal{K} with diameter 1. Given $g \in \mathcal{N}$, the space \mathbb{H}_g^+ of self-dual harmonic 2-forms with respect to g is a 3-dimensional subspace in $\Lambda_{\mathbb{R}} \equiv H^2(\mathcal{K}; \mathbb{R})$ which is positive definite with respect to the intersection form. Indeed, any choice of a hyperkähler triple gives rise to a basis of \mathbb{H}_g^+ , and they are up to a constant $O(3)$ rotation.

Define the positive Grassmannian Gr^+ to be the space of all 3-dimensional positive definite subspaces of $\Lambda_{\mathbb{R}}$. It is an open subset in the standard Grassmannian $Gr(3, \Lambda_{\mathbb{R}})$. We define the period map

$$\mathcal{P} : \mathcal{N} \rightarrow Gr^+; g \mapsto \mathbb{H}_g^+.$$

The diffeomorphism group $\mathbf{Diff}(\mathcal{K})$ acts on \mathcal{N} by $\varphi.g = \varphi^*g$, which induces a homomorphism $\Phi : \mathbf{Diff}(\mathcal{K}) \rightarrow \text{Aut}(\Lambda)$, where $\Gamma := \text{Aut}(\Lambda)$ is the automorphism group of the lattice Λ preserving the intersection form. There is a natural action of Γ on Gr^+ , hence \mathcal{P} induces a map

$$\underline{\mathcal{P}} : \mathcal{M} = \mathcal{N}/\mathbf{Diff}(\mathcal{K}) \rightarrow \mathcal{D} \equiv Gr^+/\Gamma. \quad (1)$$

The left-hand side is the set of isometry classes of hyperkähler metrics on \mathcal{K} . It is endowed with a natural Cheeger-Gromov topology. A sequence $[g_j]$ converges to $[g_\infty]$ if there are $\varphi_j \in \mathbf{Diff}(\mathcal{K})$ such that $\varphi_j^*g_j$ converges smoothly to g_∞ . Since hyperkähler metrics are Ricci-flat, it follows from the Cheeger-Colding theory that this topology coincides with the Gromov-Hausdorff topology. We also endow the period domain \mathcal{D} with the quotient topology. One can check that $\underline{\mathcal{P}}$ is continuous, and the Γ action on Gr^+ is properly discontinuous.

For any homology class $\delta \in H_2(\mathcal{K}; \mathbb{Z}) \cong H^2(\mathcal{K}; \mathbb{Z}) = \Lambda$ with $\delta.\delta = -2$, we define ξ^\perp to be subspace in Gr^+ consisting of hyperkähler metrics g such that $\int_\delta \xi = 0$ for all $\xi \in \mathbb{H}_g^+$. We denote

$$Gr^{+, \circ} = Gr^+ \setminus \bigcup_{\delta \in \Lambda, \delta.\delta = -2} \delta^\perp$$

and $\mathcal{D}^\circ = Gr^{+, \circ}/\Gamma$.

Theorem 1. *The image of $\underline{\mathcal{P}}$ is \mathcal{D}° .*

Acknowledgements. The author thanks Song Sun for suggesting the problem and pointing out that the results in [4] may help in this problem. The author is supported by Simons Collaboration Grant on Special Holonomy in Geometry, Analysis, and Physics (488633, S.S.).

We prove Theorem 1 in a few steps.

Step 1. We show that the image of \mathcal{P} is contained in $Gr^{+, \circ}$. In particular, the image of $\underline{\mathcal{P}}$ is contained in \mathcal{D}° .

The proof that we know uses some complex geometry. Suppose there is a $g \in \mathcal{N}$ and $0 \neq \delta \in H_2(\mathcal{K}; \mathbb{Z})$ such that $\delta.\delta \geq -2$ and $\int_\delta \xi = 0$ for all $\xi \in \mathbb{H}_g^+$. Choose a hyperkähler triple $\omega = (\omega_1, \omega_2, \omega_3)$ for g . Then with respect to a compatible complex structure J we see ω_1 is a Kähler form, and $\Omega = \omega_2 + \sqrt{-1}\omega_3$ is a holomorphic volume form. By the Hodge decomposition it follows that δ is a $(1, 1)$ class with respect to J . So

$\delta = c_1(L)$ for some non-trivial holomorphic line bundle L . By the Hirzebruch-Riemann-Roch theorem it follows that $h^0(\mathcal{K}, L) + h^0(\mathcal{K}, L^{-1}) = h^1(\mathcal{K}, L) + 2 + \frac{1}{2}\delta^2 > 0$. Without loss of generality we assume L has a non-zero holomorphic section S . Its zero set is a complex curve dual to δ . It follows that $\int_{\delta} \omega > 0$. Contradiction.

Step 2: We show that \mathcal{P} is an open map. In particular, the image of $\underline{\mathcal{P}}$ is open.

This follows from the standard deformation theory. We outline the argument, see [4] for details. Suppose $g \in \mathcal{N}$. We fix a hyperkähler triple ω associated to g . In the following, we will often identify an element in $\Omega_+^2 \otimes \mathbb{R}^3$ (i.e., a triple of self-dual 2-forms) with a 3×3 matrix-valued function in $\Omega^0 \otimes \mathbb{R}^{3 \times 3}$: $\eta \in \Omega_+^2 \otimes \mathbb{R}^3$ corresponds to $A = (A_{\alpha\beta}) \in \Omega^0 \otimes \mathbb{R}^{3 \times 3}$ if $\eta_{\alpha} = \sum_{\beta=1}^3 A_{\alpha\beta} \omega_{\beta}$, or concisely $\eta = A \cdot \omega$. We claim that for any fixed small triple of anti-self-dual harmonic 2-forms $\mathbf{h}^- \in \mathbb{H}_g^- \otimes \mathbb{R}^3$, $\omega' := \omega + \mathbf{h}^- + \mathbf{h}^+ + dd^*(\mathbf{f} \cdot \omega)$ defines a hyperkähler triple for some small $(\mathbf{h}^+, \mathbf{f}) \in \mathbb{H}_g^+ \otimes \mathbb{R}^3 \oplus C^{2,\gamma}(\Omega_+^2 \otimes \mathbb{R}^3)$.

Denote $\mathcal{S}_0(\mathbb{R}^3)$ the set of trace-free 3×3 symmetric matrices, and \mathfrak{F} the inverse of the map $\mathcal{S}_0(\mathbb{R}^3) \rightarrow \mathcal{S}_0(\mathbb{R}^3)$, $A \mapsto \text{tf}(A + A^T + AA^T)$ near 0, where $\text{tf}(B) = B - \frac{1}{3}\text{Tr}(B)$. Denote $\mathfrak{A} \subset C^{2,\gamma}(\Omega_+^2 \otimes \mathbb{R}^3)$, $\mathfrak{B} \subset C^{\gamma}(\Omega_+^2 \otimes \mathbb{R}^3)$, $\mathfrak{C} \subset \mathbb{H}_g^+ \otimes \mathbb{R}^3$ denote the subspace consisting of trace-free symmetric matrices, respectively. For $\mathbf{u} = (\mathbf{h}^+, \mathbf{f}) \in \mathfrak{C} \oplus \mathfrak{A}$, define $\mathcal{F} : \mathfrak{C} \oplus \mathfrak{A} \rightarrow \mathfrak{B}$ by $\mathcal{F}(\mathbf{u}) := \mathbf{h}^+ + d^+ d^*(\mathbf{f} \cdot \omega) - \mathfrak{F}(-\text{tf}(S_{\mathbf{h}^- + d^+ d^*(\mathbf{f} \cdot \omega)}))$, where $S_{\theta^-} = (\theta_{\alpha}^- \wedge \theta_{\beta}^- / 2d\text{Vol}_g)$. The condition ω' being hyperkähler is equivalent to the equation $\mathcal{F}(\mathbf{u}) = 0$.

To solve the equation, we write $\mathcal{F}(\mathbf{u}) = \mathcal{L}(\mathbf{u}) + \mathcal{N}(\mathbf{u})$, where $\mathcal{L}(\mathbf{u}) = \mathbf{h}^+ + d^+ d^*(\mathbf{f} \cdot \omega) = \mathbf{h}^+ + (\Delta_g \mathbf{f}) \cdot \omega$, $\mathcal{N}(\mathbf{u}) = -\mathfrak{F}(-\text{tf}(S_{\mathbf{h}^- + d^+ d^*(\mathbf{f} \cdot \omega)}))$. Then by standard elliptic theory that \mathcal{L} is a bounded linear map which is surjective with a bounded right inverse, and $\|\mathcal{N}(\mathbf{u}) - \mathcal{N}(\mathbf{v})\| \leq C(\|\mathbf{h}^-\| + \|\mathbf{u}\| + \|\mathbf{v}\|)(\|\mathbf{u} - \mathbf{v}\|)$. Then the implicit function theorem implies that there exists a $\delta > 0$ such that for any $\|\mathbf{h}^-\| < \delta$, $\mathcal{F}(\mathbf{u}) = 0$ has a solution \mathbf{u} with $\|\mathbf{u}\| < C(\delta)$, which finishes the proof of the claim. Now the map $\Psi : \mathbb{H}_g^- \otimes \mathbb{R}^3 \rightarrow Gr^+$, $\mathbf{h}^- \mapsto \text{span}\{\omega + \mathbf{h}^-\}$ defines a homeomorphism from a neighborhood of 0 to a neighborhood of $\mathcal{P}(g)$, and $\text{span}\{\omega + \mathbf{h}^-\} = \text{span}\{\omega + \mathbf{h}^- + \mathbf{h}^+ + dd^*(\mathbf{f} \cdot \omega)\}$ as elements in Gr^+ , it follows that the image of a neighborhood of g under \mathcal{P} contains a neighborhood of $\mathcal{P}(g)$, hence \mathcal{P} is an open map.

Step 3. We show that $\underline{\mathcal{P}} : \mathcal{M} \rightarrow \mathcal{D}^{\circ}$ is a proper map.

Suppose otherwise, then we can find a sequence of hyperkähler metrics $g_j \in \mathcal{N}$ which do not converge smoothly modulo $\mathbf{Diff}(\mathcal{K})$, but there exist $\gamma_j \in \Gamma$ such that $\gamma_j \cdot \mathcal{P}(g_j)$ converges to a positive 3-dimensional subspace P_{∞} in $Gr^{+, \circ}$. Choose a hyperkähler triple $\omega_j = (\omega_{j,1}, \omega_{j,2}, \omega_{j,3})$ for g_j . Denote $v_j^4 = 2\text{Vol}(g_j)$. We define the renormalized triple $\tilde{\omega}_j = v_j^{-2} \omega_j$, then $\int_{\mathcal{K}} \tilde{\omega}_{j,\alpha} \wedge \tilde{\omega}_{j,\beta} = \delta_{\alpha\beta}$.

Now we fix an norm $\|\cdot\|$ on $\Lambda_{\mathbb{R}}$. By abusing notation we also denote by $\|\cdot\|$ the standard norm on \mathbb{R}^3 , or the induced norm on $\Lambda_{\mathbb{R}} \otimes \mathbb{R}^3$.

Lemma 2. $\|\gamma_j \cdot [\tilde{\omega}_j]\|$ is uniformly bounded.

Proof. Otherwise, by passing to a subsequence and $O(3)$ rotations we may assume only the first component of $\gamma_j \cdot [\tilde{\omega}_j]$ is non-zero and $\|\gamma_j \cdot [\tilde{\omega}_j]\| = \lambda_j \rightarrow \infty$. Denote $\zeta_j = \lambda_j^{-1} \gamma_j \cdot [\tilde{\omega}_j]$, then passing to a subsequence we may assume ζ_j converges to an element ζ_{∞} in $\Lambda_{\mathbb{R}}$ with $\|\zeta_{\infty}\| = 1$ and with $\zeta_{\infty} \cup \zeta_{\infty} = 0$. But the line spanned by ζ_{∞} is contained in P_{∞} which is positive definite with respect to the intersection form. Contradiction. \square

Given the Lemma, by passing to a subsequence we may assume $\gamma_j \cdot [\tilde{\omega}_j]$ converges to a limit η_{∞} in $\Lambda_{\mathbb{R}} \otimes \mathbb{R}^3$. Notice $[\eta_{\infty, \alpha}] \cup [\eta_{\infty, \beta}] = \delta_{\alpha\beta}$, so η_{∞} forms a basis for P_{∞} .

Proposition 3. *Passing to a subsequence, for j large, there exists a non-zero homology class $C_j \in H_2(\mathcal{K}, \mathbb{Z})$ satisfying $C_j \cdot C_j \in \{0, -2\}$, and $\|\int_{C_j} \tilde{\omega}_j\| \rightarrow 0$.*

Proof. Passing to a subsequence we may assume (\mathcal{K}, g_j) converges to a Gromov-Hausdorff limit X_{∞} , which is a compact metric space.

If $v_j \geq \epsilon > 0$ for all j , then it follows from the classical results ([1, 2, 5]) that X_{∞} is a hyperkähler orbifold. Let $p_j \in \mathcal{K}$ be such that $\lambda_j := \max_{\mathcal{K}} |Rm(g_j)|$ is achieved at p_j . By assumption $\lambda_j \rightarrow \infty$. Then passing to a subsequence we can take a pointed Gromov-Hausdorff limit of $(\mathcal{K}, p_j, \lambda_j^{1/2} g_j)$ to get a complete ALE hyperkähler 4-manifold Z . By Kronheimer's classification [3] we know Z must contain a homology class C_{∞} with $C_{\infty} \cdot C_{\infty} = -2$. This gives rise to a sequence of -2 class C_j in \mathcal{K} such that $\|\int_{C_j} \omega_j\| \rightarrow 0$, so $\|\int_{C_j} \tilde{\omega}_j\| \rightarrow 0$ as well. Notice here we only need the topological classification in Kronheimer's result.

If $v_j \rightarrow 0$, then the conclusion follows from the results of Sun-Zhang [4]. The point is that away from finitely many points, the hyperkähler triple ω_j (up to $O(3)$ rotations) has almost local nilpotent symmetry, and it can be perturbed to a new hyperkähler triple ω'_j such that ω'_j has local nilpotent symmetry and $\omega_j - \omega'_j$ is exact. In particular, ω'_j has explicit expression and the integration of ω'_j over a cycle is the same as the integration of

ω_j over that cycle. We divide into 3 cases. The first two cases only use the analysis over the regular region in [4].

- $\dim X_\infty = 2$. In this case ω'_j is locally T^2 -invariant. Take C_j to be the class of a T^2 fiber. Then we get $\int_{C_j} \omega_{j,1} = \int_{C_j} \omega_{j,2} = 0$ and $\int_{C_j} \omega_{j,3} \sim v_j^4$. We get $\|\int_{C_j} \tilde{\omega}_j\| \sim v_j^2 \rightarrow 0$.
- $\dim X_\infty = 1$. Locally there are two cases, either there is a T^3 symmetry or a Heisenberg symmetry. In the former case the metric corresponding to ω'_j is locally a flat product $S^1(r_{j,1}) \times S^1(r_{j,2}) \times S^1(r_{j,3}) \times I$, where I is an open interval and we assume $r_{j,1} \leq r_{j,2} \leq r_{j,3}$, then we take C_j to be the homology class of $S^1(r_{j,1}) \times S^1(r_{j,2})$, then $\int_{C_j} \omega_{j,1} = \int_{C_j} \omega_{j,2} = 0$ and $0 < \int_{C_j} \omega_{j,3} = r_{j,1}r_{j,2}$ whereas $v_j^4 \sim r_{j,1}r_{j,2}r_{j,3}$. So $\|\int_{C_j} \tilde{\omega}_j\| \sim r_{j,1}^{1/2}r_{j,2}^{1/2}r_{j,3}^{-1/2} \rightarrow 0$. In the latter case the metric corresponding to ω'_j is locally given by the Gibbons-Hawking ansatz applied to a nonconstant linear function on $T^2 \times I$. Take C_j to be the homology class of the 2-torus given by the total space of the corresponding circle bundle over a circle S^1 in $T^2 \times I$, then one can arrange that $\int_{C_j} \omega_{j,1} = \int_{C_j} \omega_{j,2} = 0$ and $0 < \int_{C_j} \omega_{j,3} \sim r_{j,1}r_{j,2}$, where $r_{j,1}$ is the size of the S^1 fiber in the Gibbons-Hawking construction, and $r_{j,2}$ is the size of the flat T^2 base. Notice the volume $v_j^4 \sim r_{j,1}r_{j,2}^2$. So $\|\int_{C_j} \tilde{\omega}_j\| \sim r_{j,1}^{1/2} \rightarrow 0$.
- $\dim X_\infty = 3$. Here we need some global result from [4]. It is proved there that X_∞ must be a flat orbifold T^3/\mathbb{Z}_2 . Then ω'_j is given by Gibbons-Hawking construction on the complement of a small neighborhood of the orbifold points. We can take C_j to be the total space of the circle bundle over a closed geodesic $S^1 \subset T^3/\mathbb{Z}_2$. Then we may arrange that $\int_{C_j} \omega_{j,1} = \int_{C_j} \omega_{j,2} = 0$ while $0 < \int_{C_j} \omega_{j,3} = r_j \sim v_j^4$. So $\|\int_{C_j} \tilde{\omega}_j\| \sim r_j^{1/2} \rightarrow 0$.

□

Now we derive a contradiction. Denote $C'_j = \gamma_j^{-1}C_j$. Since C'_j is integral and non-zero, we know $\|C'_j\|$ has a uniform positive lower bound. By passing to a subsequence we may assume $\|C'_j\|^{-1}C'_j$ converges to a limit $C'_\infty \in \Lambda_\mathbb{R}$ with $\|C'_\infty\| = 1$. First suppose $C_j.C_j = 0$, then $C'_\infty.C'_\infty = 0$ and

$$\int_{C'_\infty} \eta_\infty = \lim_{j \rightarrow \infty} \|C'_j\|^{-1} \int_{C'_j} \eta_\infty = \lim_{j \rightarrow \infty} \|C'_j\|^{-1} \int_{C'_j} \gamma_j \cdot [\tilde{\omega}_j] = \lim_{j \rightarrow \infty} \|C'_j\|^{-1} \int_{C'_j} \tilde{\omega}_j = 0 \quad (2)$$

This contradicts the fact that the intersection form on $\Lambda_\mathbb{R}$ has signature $(3, 19)$. Now suppose $C_j.C_j = -2$. If $\|C'_j\|$ is unbounded, then $C'_\infty.C'_\infty = 0$, and we have $\int_{C'_\infty} \eta_\infty = 0$ as in (2), hence we obtain a contradiction in the same way. If $\|C'_j\|$ is bounded, by passing to a further subsequence we may assume C'_j converges to a limit $C''_\infty \in \Lambda$ with $C''_\infty.C''_\infty = -2$ and $\int_{C''_\infty} \eta_\infty = 0$. This contradicts to $P_\infty \in Gr^{+, \circ}$ and finishes the proof of Step 3.

Finally, $\underline{P} : \mathcal{M} \rightarrow \mathcal{D}^\circ$ being proper and \mathcal{D}° being locally compact, Hausdorff imply $\underline{P} : \mathcal{M} \rightarrow \mathcal{D}^\circ$ is a closed map. Together with the image of \underline{P} being open, \mathcal{D}° being connected, we conclude $\underline{P} : \mathcal{M} \rightarrow \mathcal{D}^\circ$ is surjective.

References

- [1] Michael T Anderson. Ricci curvature bounds and Einstein metrics on compact manifolds. *Journal of the American Mathematical Society*, pages 455–490, 1989.
- [2] Shigetoshi Bando, Atsushi Kasue, and Hiraku Nakajima. On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth. *Inventiones mathematicae*, 97(2):313–349, 1989.
- [3] Peter B Kronheimer. A Torelli-type theorem for gravitational instantons. *Journal of differential geometry*, 29(3):685–697, 1989.
- [4] Song Sun and Ruobing Zhang. Collapsing geometry of hyperkaehler 4-manifolds and applications. *arXiv preprint arXiv:2108.12991*, 2021.
- [5] Gang Tian. On Calabi’s conjecture for complex surfaces with positive first Chern class. *Inventiones mathematicae*, 101(1):101–172, 1990.