Classical 6j-symbols for finite dimensional representation of the algebra \mathfrak{gl}_3

D.V. Artamonov

Lomonosov Moscow Stater Unversity artamonov@econ.msu.ru

April 20, 2023

Abstract

In the paper an explicit formula for an arbitrary 6j-symbol for finitedimensional irreducible representations of the algebra \mathfrak{gl}_3 is derived. A 6j-symbol is written as a result of substitution of ± 1 into a series of hypergeometric type, which is similar to a Γ -series, which is a simplest example of a multivariate series of hypergeometric type. The selection rulers for a 6j-symbol are derived.

1 Introduction

1.1 The Racah coefficients

The Racah coefficients for a Lie algebra g are defined as follows. Consider irreducible representations V^1 , V^2 , V^3 of a given algebra and consider their tensor product $V^1 \otimes V^2 \otimes V^3$. Brackets in this work can be arranged in two ways:

$$(V^1 \otimes V^2) \otimes V^3$$
 or $V^1 \otimes (V^2 \otimes V^3)$.

One can split $V^1 \otimes V^2 \otimes V^3$ into a sum of irreducible representations in two ways.

1. The first way. First split into irreducible $V^1 \otimes V^2$:

$$V^1 \otimes V^2 = \bigoplus_U Mult_U^{V^1, V^2} \otimes U, \tag{1}$$

where U is an irreducible representation and $Mult_U^{V^1,V^2}$ is a multiplicity space. This is a vector space which is not supplied with an action g. Take a tensor product of (2) and V^3 from the right, one obtains

$$(V^1 \otimes V^2) \otimes V^3 = \bigoplus_{U,W} Mult_U^{V^1,V^2} \otimes Mult_W^{U,V^3} \otimes W$$
(2)

2. The second way. First split into irreducible $V^2 \otimes V^3$:

$$V^2 \otimes V^3 = \bigoplus_U Mult_H^{V^2, V^3} \otimes H, \tag{3}$$

and then

$$V^{1} \otimes (V^{2} \otimes V^{3}) = \bigoplus_{U,W} Mult_{H}^{V^{2},V^{3}} \otimes Mult_{W}^{V^{1},H} \otimes W$$

$$\tag{4}$$

There exists an isomorphism $\Phi: (V^1 \otimes V^2) \otimes V^3 \to V^1 \otimes (V^2 \otimes V^3)$, which gives a mapping

$$\Phi: \bigoplus_{U,W} Mult_U^{V^1,V^2} \otimes Mult_W^{U,V^3} \to \bigoplus_{U,W} Mult_H^{V^2,V^3} \otimes Mult_W^{V^1,H}$$
(5)

Definition 1. The Racah mapping is an induced by Φ mapping

$$W \begin{cases} V^1 & V^2 & U \\ V^3 & W & H \end{cases} : Mult_U^{V^1, V^2} \otimes Mult_W^{U, V^3} \to Mult_H^{V^2, V^3} \otimes Mult_W^{V^1, H}$$
(6)

When one fixes the basis, there appear matrix elements of this mapping. They are called *the Racah coefficients*. Let s be an index listing the basis vectors in the multiplicity space/ Then one has a notation for the Racah coefficients

$$W \begin{cases} V^{1} & V^{2} & U \\ V^{3} & W & H \end{cases}_{s_{3},s_{4}}^{s_{1},s_{2}},$$
(7)

where s_1, s_2, s_3, s_4 are indices of base vectors in $Mult_U^{V^1, V^2}, Mult_W^{U, V^3}, Mult_W^{V^2, V^3}, Mult_W^{V^1, H}$.

From this definition, the role of these coefficients from the point of view of representation theory is clear. Let's consider a category of finite-dimensional representations and let us pass on to its Grothendieck ring. Equivalently, this means passing from the category of finite-dimensional representations to the ring of their characters. Then some information about the category is lost. For example, one looses the information contained in Racah coefficients [1]. In some cases from the Grothendieck ring and Racah coefficients, the category of representations can be restored.

The Racah coefficients have applications in quantum mechanics. They were introduced by Racah in [3]. In the case of $g = \mathfrak{sl}_2$, these coefficients are discussed in all textbooks on quantum mechanics, for example [3], [4], and textbooks on

the theory of angular momentum, for example [5], [6]. In the case of finitedimensional irreducible representations of \mathfrak{sl}_2 , there are explicit formulas for the Racah coefficients in which they are expressed as the values of the hypergeometric function [7]. At the same time, the problem of calculating of the Racah coefficients for infinite-dimensional representations of \mathfrak{sl}_2 is still relevant [8], [9].

The Racah coefficients for \mathfrak{sl}_3 are also important in applications [10], [11]. The general formula for a 6j-symbol for \mathfrak{sl}_3 is unknown (see introduction to the dissertation [12]). Explicit formulas were obtained for some classes of representations [13], [14], [15], [16]. In papers [17], [18] there were also calculated some Racah coefficients for \mathfrak{sl}_3 . These Racah coefficients play an important technical role in calculation of some Clebsch-Gordan coefficients for \mathfrak{gl}_3 .

It should also be noted that even more attention is currently being paid to quantum Racah coefficients, that is, the Racah coefficients for quantum Lie algebras. It is not possible to give a complete overview of this activity, we mention only recent works [19], [20], where the Racah coefficients for algebras other than $U_q(\mathfrak{sl}_2)$ are considered.

1.2 6j-symbols

These coefficients were introduced even earlier than the Racah coefficients by Wigner ¹. To define them, we first introduce 3j-symbols. Let V^1 , V^2 , V^3 be representations of g, $\{v_{\alpha}^1\}$, $\{v_{\beta}^2\}$, $\{v_{\gamma}^3\}$ are the bases in these representations.

Definition 2. A 3*j*-symbol is a collection of coefficients

$$\begin{pmatrix} V^1 & V^2 & V^3 \\ v_{\alpha}^1 & v_{\beta}^2 & v_{\gamma}^3 \end{pmatrix}^f \in \mathbb{C}$$

such that

$$\sum_{\alpha,\beta,\gamma} \begin{pmatrix} V^1 & V^2 & V^3 \\ v_{\alpha}^1 & v_{\beta}^2 & v_{\gamma}^3 \end{pmatrix}^f v_{\alpha}^1 \otimes v_{\beta}^2 \otimes v_{\gamma}^3 = f \in V^1 \otimes V^2 \otimes V^3,$$

where f is a semi-invariant for the action of \mathfrak{gl}_3 . That is f is an eigenvector for the action of the Cartan subalgebra and it vanishes under the action of root elements.

One can easily see a relation between 3j-symbols and Clebsch-Gordan coefficients. To write it, one takes a splitting (2). Let us introduce a basis e_s in the multiplicity space and put $U^s := e_s \otimes U$, then (2) one can be write the following:

¹these coefficients are not called the Wigner coefficients, since the term "Wigner coefficients" is used for the coefficients that decompose a tensor product of two representations into irreducible ones, they are also Clebsch-Gordan coefficients

 $V^1\otimes V^2=\bigoplus_{s,U}U^s.$ Choose the bases $\{v^1_\alpha\},\,\{v^2_\beta\},\,\{u^s_\gamma\}$ in these spaces. Then one can write

$$u_{\gamma}^{s} = \sum_{\alpha,\beta} D_{V^{1},V^{2};\alpha,\beta}^{U,\gamma,s} v_{\alpha}^{1} \otimes v_{\beta}^{2}.$$
(8)

The coefficient $D_{V^1,V^2;\alpha,\beta}^{U,\gamma,s} \in \mathbb{C}$ is called the Clebsch-Gordan coefficient, it is related to the 3*j*-symbol by the rule

$$D^{U,\gamma,s}_{V^1,V^2;\alpha,\beta} = \begin{pmatrix} V^1 & V^2 & \bar{U} \\ v^1_{\alpha} & v^2_{\beta} & \bar{u}_{\gamma} \end{pmatrix}^s,$$

where \bar{U} and \bar{u}_{γ} denote the contragradient representation and the dual basis in it. The index s of a 3*j*-symbol carries the following information. The space of 3*j*-symbols with given internal indexes is isomorphic to the multiplicity space. Therefore, by fixing the basis in the multiplicity space, one fixes a basic 3*j*symbols.

Definition 3. A 6*j*-symbol is a paring of 3*j*-symbols by the ruler:

$$\begin{cases}
 V^{1} \quad V^{2} \quad U \\
 V^{3} \quad W \quad H
 \end{cases}^{s_{1},s_{2}} := \sum_{\alpha_{1},\dots,\alpha_{6}} \left(\begin{matrix} \bar{V}^{1} \quad \bar{V}^{2} \quad U \\
 \bar{v}_{\alpha_{1}}^{1} \quad \bar{v}_{\alpha_{2}}^{2} \quad u_{\alpha_{4}} \end{matrix} \right)^{s_{1}} \cdot \left(\begin{matrix} \bar{U} \quad \bar{V}^{3} \quad W \\
 \bar{u}_{\alpha_{4}} \quad \bar{v}_{\alpha_{3}}^{3} \quad w_{\alpha_{5}} \end{matrix} \right)^{s_{2}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} \quad \bar{H} \\
 V^{2}_{\alpha_{2}} \quad v^{3}_{\alpha_{3}} \quad \bar{h}_{\alpha_{6}} \end{matrix} \right)^{s_{3}} \cdot \left(\begin{matrix} V^{1} \quad H \quad \bar{W} \\
 v^{1}_{\alpha_{1}} \quad h_{\alpha_{6}} \quad \bar{w}_{\alpha_{5}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{1}_{\alpha_{3}} \quad \bar{h}_{\alpha_{6}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{1} \quad H \quad \bar{W} \\
 v^{1}_{\alpha_{1}} \quad h_{\alpha_{6}} \quad \bar{w}_{\alpha_{5}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} \quad \bar{h}_{\alpha_{6}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{1} \quad H \quad \bar{W} \\
 v^{1}_{\alpha_{1}} \quad h_{\alpha_{6}} \quad \bar{w}_{\alpha_{5}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} \quad \bar{h}_{\alpha_{6}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{1}_{\alpha_{1}} & h_{\alpha_{6}} \quad \bar{w}_{\alpha_{5}} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{2}_{\alpha_{2}} & v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \\
 v^{3}_{\alpha_{3}} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} \quad V^{3} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} \quad V^{3} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} \quad V^{3} & \bar{V} \end{matrix} \right)^{s_{4}} \cdot \left(\begin{matrix} V^{2} \quad V^{3} \quad V^{3} & \bar{V} \end{matrix} \right)^{s_{4}}$$

This expression should be understood as follows: the Lie algebra g acts on the 3j-symbols by acting on lower indices. We form a semiinvariant of 4-x 3jsymbols by pairing indexes so that in two 3j-symbols only one pair of indexes paired.

Let's now write out a relationship between the Racah coefficients and 6*j*-symbols. In the expression below, we use the fact that there is a duality between the spaces $Mult_U^{V^1,V^2}$ and $Mult_{\overline{U}}^{\overline{V}^1,\overline{V}^2}$. Thus if *s* is an index of a basic vector in $Mult_{\overline{U}}^{\overline{V}^1,\overline{V}^2}$, then \overline{s} is the index of the dual basic vector in $Mult_{\overline{U}}^{\overline{V}^1,\overline{V}^2}$.

$$W \begin{cases} V^1 & V^2 & U \\ V^3 & W & H \end{cases}_{s_3, s_4}^{\bar{s}_1, \bar{s}_2} = \begin{cases} V^1 & V^2 & U \\ V^3 & W & H \end{cases}_{s_3, s_4}^{s_1, s_2}$$

Below we deal with 6j-symbols.

1.3 The results of the paper

In this paper, we obtain a simple explicit formula for an arbitrary 6j symbol for finite-dimensional irreducible representations of \mathfrak{gl}_3 . This can be done by using

the following ideas.

First, A-GKZ realization of a representation of \mathfrak{gl}_3 is used. This realization is described in detail in [21]. The space of a representation is realized as a subspace in the space of polynomials in the variables A_X , $X \subset \{1, 2, 3\}$, antisymmetric in X, but not obeying other relations. The representation space is described as the space of polynomial solutions of a partial differential equation called the antisymmetrized Gelfand-Kapranov-Zelevinsky equation (A-GKZ for short). In this realization, it is possible to write explicitly functions in variables A_X corresponding to the Gelfand-Tsetlin basis vectors (see the original work [22], as well as [21]). It is important that in this model there is an explicitly written scalar product.

Secondly, an explicitly construction of the multiplicity index s for the Clebsch-Gordan coefficients and 3j-symbols from [23] is used.

Thirdly, an explicit and simple formula for an arbitrary 3j-symbol obtained in [21] is use ².

These ideas allow us to find an explicit simple formula for an arbitrary 6*j*-symbol for finite-dimensional irreducible representations in terms of the values of a hypergeometric type function.

The structure of this paper is the following. The introductory Section 2 describes A-GKZ and functional realizations of representations of \mathfrak{gl}_3 . A solution of the multiplicity problem in the decomposition of a the tensor product of \mathfrak{gl}_3 into the sum of irreducibles is given. A a formula for a 3j-symbol using scalar products is presented.

In the main Section 3 an arbitrary 6j-symbol for the algebra \mathfrak{gl}_3 is calculated. First of all, in the Section 3.1 an implicit formula is given via a scalar product (Lemma 1). Then in the Section 3.2 the selection rules for the 6j-symbol are proved (Theorem 3). Finally, in the section 3.3 an explicit formula is derived for the 6j-symbol (Theorem 4).

2 The basic notions

2.1 A-hypergeometric functions

A detailed information about a Γ -series can be found in [24].

Let $B \subset \mathbb{Z}^N$ be a lattice, let $\mu \in \mathbb{Z}^N$ be a fixed vector. Define a hypergeometric Γ -series in the variables $z_1, ..., z_N$ by the formula

$$\mathcal{F}_{\mu}(z,B) = \sum_{b\in B} \frac{z^{b+\mu}}{\Gamma(b+\mu+1)},\tag{10}$$

 $^{^{2}}$ in [23], an explicit, but much more cumbersome solution of the same problem is obtained.

where $z = (z_1, ..., z_N)$. We use a multi-index notation in the numerator and denominator

$$z^{b+\mu} := \prod_{i=1}^{N} z_i^{b_i+\mu_i}, \ \Gamma(b+\mu+1) := \prod_{i=1}^{N} \Gamma(b_i+\mu_i+1).$$

B Remark 1. A more compact formula for (10) is the following: $\mathcal{F}_{\mu}(z, B) = \sum_{x \in \mu + B} \frac{z^x}{\Gamma(x+1)}$.

Note that if at least one of the components of the vector $b + \mu$ is a negative integer, then the corresponding term in (10) turns to zero. Due to this, there is be only a finite number of terms in the series considered in the paper. For simplicity, we write factorials instead of Γ -functions.

A Γ -series satisfies the Gelfand-Kapranov-Zelevinsky system (the GKZ system for short). Let us write it explicitly in the case $z = (z_3, z_1, z_2, z_{1,3}, z_{2,3}, z_{1,2})$, $B = \mathbb{Z} < v = e_1 - e_2 - e_{1,3} + e_{2,3} >$:

$$\begin{pmatrix} \frac{\partial^2}{\partial z_1 \partial z_{2,3}} - \frac{\partial^2}{\partial z_2 \partial z_{1,3}} \end{pmatrix} \mathcal{F}_{\mu}(z, B) = 0,$$

$$(z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2}) \mathcal{F}_{\mu}(z, B) = (\mu_1 + \mu_2) \mathcal{F}_{\mu}(z, B),$$

$$(z_1 \frac{\partial}{\partial z_1} + z_{1,3} \frac{\partial}{\partial z_{1,3}}) \mathcal{F}_{\mu}(z, B) = (\mu_1 + \mu_{1,3}) \mathcal{F}_{\mu}(z, B),$$

$$(z_1 \frac{\partial}{\partial z_1} - z_{2,3} \frac{\partial}{\partial z_{2,3}}) \mathcal{F}_{\mu}(z, B) = (\mu_1 - \mu_{2,3}) \mathcal{F}_{\mu}(z, B),$$

$$z_3 \frac{\partial}{\partial z_3} \mathcal{F}_{\mu}(z, B) = \mu_3 \mathcal{F}_{\mu}(z, B),$$

$$z_{1,2} \frac{\partial}{\partial z_{1,2}} \mathcal{F}_{\mu}(z, B) = \mu_{1,2} \mathcal{F}_{\mu}(z, B).$$

$$(11)$$

2.2 A-GKZ realization of a representation of \mathfrak{gl}_3

Throughout the rest of the work, we talk about finite-dimensional irreducible representations.

Details can be found in [21]. Consider the variables A_X , where $X \subset \{1, 2, 3\}$ is a proper subset, antisymmetric in X, but not obeying other relations. On these variables there is an action \mathfrak{gl}_3 , defined by the rule

$$E_{i,j}A_X = \begin{cases} A_{X|_{j \mapsto i}}, & j \in X \\ 0 \text{ otherwise} \end{cases}$$
(12)

Here $X \mid_{j \mapsto i}$ denotes a substitution of j by i.

Consider an equation called the antisymmetrized GKZ equation (the A-GKZ equation):

$$\left(\frac{\partial^2}{\partial A_1 A_{2,3}} - \frac{\partial^2}{\partial A_2 A_{1,3}} + \frac{\partial^2}{\partial A_3 A_{1,2}}\right)F = 0 \tag{13}$$

B Remark 2. The term A-GKZ is explained as follows. Consider the system (11), leave only the first equation and "antisymmetrize" it by adding a third term. The resulting equation is the equation A-GKZ.

One has.

Theorem 1 ([21]). The space of polynomial solutions (13) is invariant under the action of \mathfrak{gl}_3 . As a representation, the space of polynomial solutions is a direct sum with multiplicity 1 of all finite-dimensional irreducible representations with $m_3 = 0$. The space of an irreducible representation with the highest weight $[m_1, m_2, 0]$ has the highest vector

$$A_1^{m_1-m_2}A_{1,2}^{m_2}$$

and consists of all polynomial solutions whose homogeneous degree in variables A_X , |X| = 1 is equal to $m_1 - m_2$, and homogeneous degree in variables A_X , |X| = 2 is equal to m_2 .

Let us write a base in the space of polynomial solutions. Consider the space \mathbb{C}^6 with coordinates $A_1, A_2, A_3, A_{1,2}, A_{1,3}, A_{2,3}$ and define the vectors

$$v = e_1 - e_2 - e_{1,3} + e_{2,3}, \quad r = e_3 + e_{1,2} - e_1 - e_{2,3},$$
 (14)

Now consider a Γ -series associated with a lattice $B = \mathbb{Z} < v >$ and a vector $\mu \in \mathbb{Z}^6$:

$$\mathcal{F}_{\mu}(A,B) := \sum_{t \in \mathbb{Z}} \frac{A^{\mu+tv}}{(\mu+tv)!},\tag{15}$$

where we used the multi-index notations

$$A^{\mu+tv} = \prod_{X} A_{X}^{\mu_{X}+tv_{X}}, \quad (mu+tv)! = \prod_{X} (\mu_{X}+tv_{X})!.$$

Then (see calculations in [23], [25] or [21]) the basis in the space of polynomial solutions consists of non-zero functions of type

$$F_{\mu}(A) := \sum_{s \in \mathbb{Z}_{\geq 0}} q_s^{\mu} \zeta_A^s \mathcal{F}_{\mu-sr}(A), \tag{16}$$

where

$$t_{0} = 1, \quad t_{s}^{\mu} = \frac{1}{s(s+1) + s(\mu_{1} + \mu_{2} + \mu_{1,3} + \mu_{2,3})}, \text{ in the case } s > 0$$

$$q_{\mu}^{s} = \frac{t_{s}^{\mu}}{\sum_{s' \in \mathbb{Z}_{\geq 0}} t_{s'}^{\mu}}$$

$$r = e_{3} + e_{1,2} - e_{1} - e_{2,3}, \quad \zeta_{A} = A_{1}A_{2,3} - A_{2}A_{1,3}.$$
(17)

Note that actually $F_{\mu}(A)$ depends not on the vector μ , but on the shifted lattice $\Pi = \mu + \mathbb{Z} < v >$.

Moreover, when this function is nonzero, then it is nothing else but the Gelfand-Tsetlin basic vector. Namely, the Gelfand-Cetlin diagram

$$(m_{p,q}) = \begin{pmatrix} m_{1,3} & m_{2,3} & m_{3,3} \\ & m_{1,2} & m_{2,2} \\ & & m_{1,1} & \end{pmatrix}$$

corresponds to such a function $F_{\mu}(A)$, that the shifted lattice $\Pi = \mu + \mathbb{Z} < v >$ is given by the equations

$$\delta \in \Pi \Leftrightarrow \begin{cases} \delta_1 + \delta_2 + \delta_3 + \delta_{1,2} + \delta_{1,3} + \delta_{2,3} = m_{1,3}, \\ \delta_{1,2} + \delta_{1,3} + \delta_{2,3} = m_{2,3}, \\ \delta_1 + \delta_2 + \delta_{1,2} + \delta_{1,3} + \delta_{2,3} = m_{1,2}, \\ \delta_{1,2} = m_{2,2}, \\ \delta_1 + \delta_{1,2} + \delta_{1,3} = m_{1,1} \end{cases}$$
(18)

Moreover, this is a one-to-one correspondence between the Gelfand-Tsetlin diagrams and nonzero functions $F_{\mu}(A)$.

It is important that in the A-GKZ realization there is an explicitly formula for an invariant scalar product

$$\langle f(A), g(A) \rangle = f(\frac{\partial}{\partial A})g(A) \mid_{A=0}$$
 (19)

Also note that if the representation of V is realized in the space of polynomials in the variables A_X : $V = \{h(A)\}$, then the contragradient representation is realized in the space of polynomials in the operators $\frac{\partial}{\partial A_X}$. The action of \mathfrak{gl}_3 on the differential operators is generated by an the action on functions of A_X . In this case $\bar{V} = \{h(\frac{\partial}{\partial A_X}) : h(A) \in V\}$. The pairing is given by a formula similar to (19):

$$\{h_1(A), h_2(\frac{\partial}{\partial A_X})\} >= h_2(\frac{\partial}{\partial A_X})h_1(A) \mid_{A=0}.$$
 (20)

Since the base $F_{\mu}(A)$ is the Gelfand-Tsetlin base, which is orthogonal, then the dual o $F_{\mu}(A)$ base is the base $\frac{1}{|F_{\mu}|^2}F_{\mu}(\frac{\partial}{\partial A})$.

2.3 The functional realization

We need also another realization, details about it can be found in [26]. Functions on the group GL_3 form a representation of GL_3 . On a function f(g), $g \in GL_3$, an element of the group $X \in GL_3$ acts by right shifts according to the rule

$$(Xf)(g) = f(gX). \tag{21}$$

Passing to an infinitesimal version of this action, one gets that the space of all functions on GL_3 is a representation of \mathfrak{gl}_3 .

Any finite-dimensional irreducible representation can be embedded into the space of function on \mathfrak{gl}_3 . More precise, if $[m_1, m_2, m_3]$ is the highest weight, then in the space of all functions there is a highest vector with such a weight, which is explicitly written as follows.

Let a_i^j , i, j = 1, 2, 3 be a function of a matrix element on the group GL_3 . Here j is the row index and i is the column index. Also put

$$a_{i_1,\dots,i_k} := det(a_i^j)_{i=i_1,\dots,i_k}^{j=1,\dots,k},$$
(22)

where one takes a determinant of the submatrix in the matrix (a_i^j) formed by rows indexed by 1, ..., k and columns $i_1, ..., i_k$. The operator $E_{i,j}$ acts on a determinant by acting onto column indexes, according to a formula similar to (12)

$$E_{i,j}a_X = \begin{cases} a_{X|_{j \mapsto i}}, j \in X\\ 0 \text{ otherwise.} \end{cases}$$
(23)

There is a mapping from A-GKZ realization to a functional one, consisting in replacing

$$A_X \mapsto a_x. \tag{24}$$

The fundamental difference between this realization and the A-GKZ realization is that the determinants of a_x obey the Plucker relation

$$a_1 a_{2,3} - a_2 a_{1,3} + a_3 a_{1,2} = 0, (25)$$

and the variables A_X are independent.

On the one hand, this simplifies the description of the representation space. So, there is a Theorem (cf. with the Theorem 1).

Theorem 2. The space of an irreducible representation with the highest weight $[m_1, m_2, 0]$ and the highest vector

$$a_1^{m_1-m_2}a_{1,2}^{m_2}$$

consists of all polynomials whose homogeneous degree in variables a_X , |X| = 1is equal to $m_1 - m_2$, and the homogeneous degree in variables a_X , |X| = 2 is equal to m_2 . But on the other hand, many calculations become more complicated. So the formula for the invariant scalar product (19) is incorrect when A_X is mechanically replaced by a_X .

Nevertheless, this following is true. When substituting (24), the function $F_{\mu}(A)$ passes into $\mathcal{F}_{\mu}(a, B)$. So $\mathcal{F}_{\mu}(a, B)$ is a Gelfand-Tsetlin basic vector in a functional realization.

2.4 A solution of the multiplicity problem for 3*j*-symbols

Let us construct explicitly the multiplicity index of s for 3j-symbols. The triple tensor product can be realized in the function space on $GL_3 \times GL_3 \times GL_3$. Functions of matrix elements on these factors GL_3 are denoted as a_i^j , b_i^j , c_i^j . Similar letters will denote the determinants of matrices composed of these matrix elements.

Let us construct explicitly a semiinvariant vectors f. It is easy to understand that semiinvariants are the following expressions

$$f = \frac{(abc)^{\tau_1} (aac)^{\tau_2} (acc)^{\tau_3} (aab)^{\tau_4} (abb)^{\tau_5} (bbc)^{\tau_6} (bcc)^{\tau_7} (aabbcc)^{\tau_8}}{\tau_1 ! \tau_2 ! \tau_3 ! \tau_4 ! \tau_5 ! \tau_6 ! \tau_7 ! \tau_8 !},$$
(26)

where

$$(abc) = det \begin{pmatrix} a_1^1 & a_2^1 & a_3^1 \\ b_1^1 & b_2^1 & b_3^1 \\ c_1^1 & c_2^1 & c_3^1 \end{pmatrix}, \quad (aac) = det \begin{pmatrix} a_1^1 & a_2^1 & a_3^1 \\ a_1^2 & a_2^2 & a_3^2 \\ c_1^1 & c_2^1 & c_3^1 \end{pmatrix}$$
 Pë.C,.Pr. (27)

Here

$$(aabbcc) := (\tilde{a}\tilde{b}\tilde{c}), \ \tilde{a}_1^1 := a_{2,3}, \ \tilde{a}_2^1 := -a_{1,3}, \ \tilde{a}_3^1 := a_{1,2},$$

 $\tilde{b}_i^1, \, \tilde{c}_i^1$ are defined analogously.

In this case, f belongs to the tensor product $V^1 \otimes V^2 \otimes V^3$ of representations with highest weights $[m_1, m_2, 0]$, $[m'_1, m'_2, 0]$, $[M_1, M_2, 0]$ if and only if the following conditions hold

$$m_{1} = \tau_{1} + \tau_{2} + \tau_{3} + \tau_{4} + \tau_{5} + \tau_{8}, \quad m_{2} = \tau_{2} + \tau_{4} + \tau_{8},$$

$$m_{1}' = \tau_{1} + \tau_{4} + \tau_{5} + \tau_{6} + \tau_{7} + \tau_{8}, \quad m_{2}' = \tau_{5} + \tau_{6} + \tau_{8},$$

$$M_{1} = \tau_{1} + \tau_{2} + \tau_{3} + \tau_{6} + \tau_{7} + \tau_{8}, \quad M_{2} = \tau_{3} + \tau_{7} + \tau_{8}$$
(28)

Proposition 1 (see Proposition 2 in [23]). In the space of 3*j*-symbols with the same internal indexes, there is a set of generators consisting of 3*j*-symbols indexed by semiinvariant functions f of the form (26), consistent with the higher weights in the upper row of the 3*j*-symbol according to the rule (28). To get a basis, it is necessary to leave f such that either $\tau_1 = 0$ or $\tau_8 = 0$.

2.5 A formulas for a 3*j*-symbol

In representations, we take bases of the type $F_{\mu}(A)$, but in order to coordinate the notation with (9), we use the index α_i to enumerate the basis vectors. Thus, as $v_{\alpha_1}^1$ we take $F_{\alpha_1}(A^1)$, as $v_{\alpha_2}^2$ we take $F_{\alpha_2}(A^2)$, etc.

Also, as a multiplicity index for the 3j-symbol, we use a function defines a semiinvariant.

The calculations in the present paper are based on the formula for a 3jsymbols in the basis F_{α} obtained in [21]. Let G be a semiinvariant in $V^1 \otimes V^2 \otimes$ V^3 in the A-GKZ realization and $G(A, B, C) = \sum_{\alpha'_1, \alpha'_2, \alpha'_3} c_{\alpha'_1, \alpha'_2, \alpha'_3} F_{\alpha'_1}(A) \otimes$ $F_{\alpha'_2}(B) \otimes F_{\alpha'_3}(C)$, where

$$c_{\alpha_1,\alpha_2,\alpha_3} = \begin{pmatrix} V^1 & V^2 & V^3 \\ F_{\alpha_1} & F_{\alpha_2} & F_{\alpha_3} \end{pmatrix}^G$$

The vectors F_{α} are orthogonal, since they are vectors of the Gelfand-Tsetlin basis, so

$$c_{\alpha_1,\alpha_2,\alpha_3} = \frac{\langle G(A, B, C), F_{\alpha_1}(A)F_{\alpha_2}(B)F_{\alpha_3}(C) \rangle}{|F_{\alpha_1}(A)|^2 |F_{\alpha_2}(B)|^2 |F_{\alpha_3}(C)|^2}$$

Also note the following. Let the semiinvariant G in the A-GKZ realization correspond to a semiinvariant f in the functional realization given by the function (26). Then

$$G(A, B, C) = f(A, B, C) + r(A, B, C),$$

where r(A, B, C) belongs to the ideal of relations between the determinants a_X , b_X , c_X . This ideal is generated by the relations (25), as well as similar relations for b_X , c_X . Then

$$< r(A, B, C), F_{\alpha_1}(A)F_{\alpha_2}(B)F_{\alpha_3}(C) >= 0,$$

since $F_{\alpha_1}(A)$, $F_{\alpha_2}(B)$, $F_{\alpha_3}(C)$ are the solution of A-GKZ equations (13), and the scalar product is given by (19). So, a 3*j*-symbol is calculated in the following manner

$$\begin{pmatrix} V^1 & V^2 & V^3 \\ F_{\alpha_1} & F_{\alpha_2} & F_{\alpha_3} \end{pmatrix}^f = \frac{\langle f(A, B, C), F_{\alpha_1}(A)F_{\alpha_2}(B)F_{\alpha_3}(C) \rangle}{|F_{\alpha_1}(A)|^2 |F_{\alpha_2}(B)|^2 |F_{\alpha_3}(C)|^2}$$
(29)

3 6*j*-symbols

Let's start calculating the 6j-symbol (9). First, we give an expression for it in terms of scalar products, and then we give an explicit formula through the value of a hypergeometric type function, with ± 1 substituted as arguments.

3.1An expression though scalar products

Let's write an expression for the 6j symbol (9). Let the multiplicity indices s_1, \ldots, s_4 in (9) correspond to the functions f_1, \ldots, f_4 .

We need to calculate 3j-symbols for the contragradient representation and the dual basis. Let use a realization of a contragradient representation described at the end of the section 2.3. The a basis dual to $F_{\alpha_i}(A^i)$ is the basis of $\frac{1}{|F_{\alpha_i}|^2}F_{\alpha_i}(\frac{\partial}{\partial A^i}).$ Note that a 3*j*-symbol of the form

$$\begin{pmatrix} \bar{V}^1 & \bar{V}^2 & U \\ F_{\alpha_1}(\frac{\partial}{\partial A^1}) & F_{\alpha_2}(\frac{\partial}{\partial A^2}) & F_{\alpha_4}(A^4) \end{pmatrix}^f$$

can be calculated as follows:

$$\begin{pmatrix} \bar{V}^1 & \bar{V}^2 & U \\ F_{\alpha_1}(\frac{\partial}{\partial A^1}) & F_{\alpha_2}(\frac{\partial}{\partial A^2}) & F_{\alpha_4}(A^4) \end{pmatrix}^f = \frac{\langle f(\frac{\partial}{\partial A^1}, \frac{\partial}{\partial A^2}, A^4), F_{\alpha_1}(\frac{\partial}{\partial A^1})F_{\alpha_2}(\frac{\partial}{\partial A^2})F_{\alpha_4}(A^4) \rangle}{|F_{\alpha_1}(\frac{\partial}{\partial A^1})|^2 |F_{\alpha_2}(\frac{\partial}{\partial A^2})|^2 |F_{\alpha_4}(A^4)|^2}$$
(30)

The scalar product in case when there is a function not of a variable, but of a differentiation operator is calculated using a formula similar to (19). One has

$$< f(\frac{\partial}{\partial A^{1}}, \frac{\partial}{\partial A^{2}}, A^{4}), F_{\alpha_{1}}(\frac{\partial}{\partial A^{1}})F_{\alpha_{2}}(\frac{\partial}{\partial A^{2}})F_{\alpha_{4}}(A^{4}) > = < f(A^{1}, A^{2}, A^{4}), F_{\alpha_{1}}(A^{1})F_{\alpha_{2}}(\partial A^{2})F_{\alpha_{4}}(A^{4}) > = |F_{\alpha_{1}}(A^{1})|^{2} = |F_{\alpha_{1}}(\frac{\partial}{\partial A^{1}})|^{2}, \dots$$

$$(31)$$

Bases $F_{\alpha_1}(A^1)$ and $F_{\alpha_1}(\frac{\partial}{\partial A^1})$ etc. are not dual, the basis dual to $F_{\alpha_1}(A^1)$ is $\frac{1}{|F_{\alpha_1}|^2}F_{\alpha_1}(\frac{\partial}{\partial A^1})$. So the 6*j*-symbol is expressed in terms of the considered 3j-symbols (30) as follows

$$\begin{cases} V^{1} \quad V^{2} \quad U \\ V^{3} \quad W \quad H \end{cases} \stackrel{f_{1},f_{2}}{\stackrel{f_{2}}{=}} := \sum_{\alpha_{1},\dots,\alpha_{6}} \begin{pmatrix} \bar{V}^{1} & \bar{V}^{2} & U \\ F_{\alpha_{1}}(\frac{\partial}{\partial A^{1}}) & F_{\alpha_{2}}(\frac{\partial}{\partial A^{2}}) & F_{\alpha_{4}}(A^{4}) \end{pmatrix}^{f_{1}} \cdot \\ \cdot \begin{pmatrix} \bar{U} & \bar{V}^{3} & W \\ F_{\alpha_{4}}(\frac{\partial}{\partial A^{4}}) & F_{\alpha_{3}}(\frac{\partial}{\partial A^{3}}) & F_{\alpha_{4}}(A^{5}) \end{pmatrix}^{f_{2}} \cdot \\ \cdot \begin{pmatrix} V^{2} & V^{3} & \bar{H} \\ F_{\alpha_{2}}(A^{2}) & F_{\alpha_{3}}(A^{3}) & F_{\alpha_{6}}(\frac{\partial}{\partial A^{6}}) \end{pmatrix}^{f_{3}} \cdot \begin{pmatrix} V^{1} & H & \bar{W} \\ F_{\alpha_{1}}(A^{1}) & F_{\alpha_{6}}(A^{6}) & F_{\alpha_{5}}(\frac{\partial}{\partial A^{5}}) \end{pmatrix}^{f_{4}} \cdot |F_{\alpha_{1}}|^{2} \cdot \dots \cdot |F_{\alpha_{6}}|^{2} \end{cases}$$
(32)

Take the expressions (30) and substitute them in (32). Consider (31). At the same time the expression $|F_{\alpha_i}|^2$ occurring at the end of (9) are written as $F_{\alpha_i}(\frac{\partial}{\partial A^i})F_{\alpha_i}(A^i)\mid_{A=0} \mathbf{B}$

One obtains B

c

ΒB

$$\begin{cases} V^1 & V^2 & U \\ V^3 & W & H \end{cases}_{f_3, f_4}^{f_1, f_2} = \sum_{\alpha_1, \dots, \alpha_6} \frac{\langle f_1, F_{\alpha_1} F_{\alpha_2} F_{\alpha_4} \rangle}{|F_{\alpha_1}|^2 |F_{\alpha_2}|^2 |F_{\alpha_4}|^2} F_{\alpha_1}(\frac{\partial}{\partial A^1}) F_{\alpha_2}(\frac{\partial}{\partial A^2}) F_{\alpha^4}(A^4) \cdot \frac{\langle f_2, F_{\alpha_4} F_{\alpha_3} F_{\alpha_5} \rangle}{|F_{\alpha_4}|^2 |F_{\alpha_3}|^2 |F_{\alpha_5}|^2} F_{\alpha_4}(\frac{\partial}{\partial A^4}) F_{\alpha_3}(\frac{\partial}{\partial A^3}) F_{\alpha_5}(A^5) \dots |_{A_1 = \dots = A_6 = 0} \end{cases}$$

Now write

$$f_1(\frac{\partial}{\partial A^1}, \frac{\partial}{\partial A^2}, A^4) = \sum \frac{\langle f_1, F_{\alpha_1} F_{\alpha_2} F_{\alpha_4} \rangle}{|F_{\alpha_1}|^2 |F_{\alpha_2}|^2 |F_{\alpha_4}|^2} F_{\alpha_1}(\frac{\partial}{\partial A^1}) F_{\alpha_2}(\frac{\partial}{\partial A^2}) F_{\alpha^4}(A^4),$$

and analogous expressions for B B $f_2(\frac{\partial}{\partial A^4}, \frac{\partial}{\partial A^5}, A^5)$, $f_3(\frac{\partial}{\partial A^2}, A^3, \frac{\partial}{\partial A^6})$, $f_4(A^1, A^6, \frac{\partial}{\partial A^5})$. Using that B $\{F_{\alpha_i}(A^i), F_{\alpha'_i}(\frac{\partial}{\partial A^i})\} = |F_{\alpha_i}|^2$ if $\alpha_i = \alpha'_i$ and 0 otherwise one

gets

LEmma 1.

$$\begin{cases} V^{1} \quad V^{2} \quad U \\ V^{3} \quad W \quad H \end{cases}_{f_{3}, f_{4}}^{f_{1}, f_{2}} = f_{1}(\frac{\partial}{\partial A^{1}}, \frac{\partial}{\partial A^{2}}, A^{4})f_{2}(\frac{\partial}{\partial A^{4}}, \frac{\partial}{\partial A^{3}}, A^{5}) \cdot \\ \cdot f_{3}(\frac{\partial}{\partial A^{2}}, A^{3}, \frac{\partial}{\partial A^{6}})f_{4}(A^{1}, A^{6}, \frac{\partial}{\partial A^{5}}) \cdot |_{A_{1}=\ldots=A_{6}=0}, \end{cases}$$
(33)

where it is assumed that this expression is evaluated as follows. At first it is represented as the sum of monomials of variables and differential operators. We multiply variables and differential operators as if they commute. Then, in each monomial, the differential operators act on the product of the variables occurring in the monomial. Finally, zeros are substituted instead of all the variables.

3.2The selection rulers

Let's find the conditions necessary for the 6j symbol to be nonzero. To do this, we consider supports of various functions.

Definition 4. A support of a function f of variables $Z = \{z_1, ..., z_N\}$ is defined as follows. Decompose the function into a power series, take the set of exponents $\delta \in \mathbb{Z}^N$ of monomials $Z^{\delta} := z_1^{\delta_1} \cdot \ldots \cdot z_N^{\delta_N}$, occurring in the decomposition of f. The obtained set of exponents is called the support of the function.

The support is denoted as supp f.

In [21] the function f of type (26) is considered as a function of ther variables

$$Z = \{ [c_1a_{2,3}], [c_2a_{1,3}], [c_3a_{1,2}], [a_1c_{2,3}], [a_2c_{1,3}], [a_3c_{1,2}], [c_1b_{2,3}], [c_2b_{1,3}], [c_3b_{1,2}], [b_1c_{2,3}], [b_2c_{1,3}], [b_3c_{1,2}], [b_1c_{2,3}], [b_2c_{1,3}], [b_3c_{1,2}], [a_1b_{2,3}], [b_2c_{1,3}], [b_3c_{1,2}], [a_1b_{2,3}], [a_2b_{1,3}], [a_3b_{1,2}], [a_1b_2c_3], [a_2b_3c_1], [a_3b_1c_2], -[a_2b_1c_3], [a_1b_3c_2], [a_3b_2c_1], [a_{2,3}b_{1,3}c_{1,2}], [a_{1,3}b_{1,2}c_{2,3}], [a_{1,2}b_{2,3}c_{1,3}], -[a_{1,3}b_{2,3}c_{1,2}], [a_{2,3}b_{1,2}c_{1,3}], [a_{1,2}b_{1,3}c_{2,3}] \}$$

$$(34)$$

Thus the summands in the determinants (cas), (ac), ..., (aabbcc) are identified with these variables.

In [21] it is shown that the support of the function f of type (26) has the form of an intersection of a shifted lattice and a positive octant

$$suppf = \left(\bigcap_{i} (\delta_i \ge 0)\right) \cap (\kappa + \mathcal{B}) \subset \mathbb{Z}^{30},$$

where 30 is the number of variables Z. The lattice $\mathcal{B} \subset \mathbb{Z}^{30}$ is generated by the vectors

$$p_{1} = e_{[c_{1}a_{2,3}]} - e_{[c_{2}a_{1,3}]}, \quad p_{2} = e_{[c_{1}a_{2,3}]} - e_{[c_{3}a_{1,2}]}, \quad p_{3} = e_{[a_{1}c_{2,3}]} - e_{[a_{2}c_{1,3}]}, \\ p_{4} = e_{[a_{1}c_{2,3}]} - e_{[a_{3}c_{1,2}]}, \quad p_{5} = e_{[c_{1}b_{2,3}]} - e_{[c_{2}b_{1,3}]}, \quad p_{6} = e_{[c_{1}b_{2,3}]} - e_{[c_{3}b_{1,2}]}, \\ p_{7} = e_{[b_{1}c_{2,3}]} - e_{[b_{2}c_{1,3}]}, \quad p_{8} = e_{[b_{1}c_{2,3}]} - e_{[b_{3}c_{1,2}]}, \quad p_{9} = e_{[a_{1}b_{2,3}]} - e_{[a_{2}b_{1,3}]}, \\ p_{10} = e_{[a_{1}b_{2,3}]} - e_{[a_{3}b_{1,2}]}, \quad p_{11} = e_{[b_{1}a_{2,3}]} - e_{[b_{2}a_{1,3}]}, \quad p_{12} = e_{[b_{1}a_{2,3}]} - e_{[b_{3}a_{1,2}]}, \\ p_{13} = e_{[a_{1}b_{2}c_{3}]} - e_{[a_{2}b_{3}c_{1}]}, \quad p_{14} = e_{[a_{1}b_{2}c_{3}]} - e_{[a_{3}b_{1}c_{2}]}, \quad p_{15} = e_{[a_{1}b_{2}c_{3}]} - e_{[a_{2}b_{1}c_{3}]} \\ p_{16} = e_{[a_{1}b_{2}c_{3}]} - e_{[a_{1}b_{3}c_{2}]}, \quad p_{17} = e_{[a_{1}b_{2}c_{3}]} - e_{[a_{3}b_{2}c_{1}]}, \quad p_{18} = e_{[a_{2,3}b_{1,3}c_{1,2}]} - e_{[a_{1,3}b_{1,2}c_{2,3}]}, \\ p_{19} = e_{[a_{2,3}b_{1,3}c_{1,2}]} - e_{[a_{1,2}b_{2,3}c_{1,3}]}, \quad p_{20} = e_{[a_{2,3}b_{1,3}c_{1,2}]} - e_{[a_{1,3}b_{2,3}c_{1,2}]}, \\ p_{21} = e_{[a_{2,3}b_{1,3}c_{1,2}]} - e_{[a_{1,2}b_{1,3}c_{2,3}]}, \quad p_{22} = e_{[a_{2,3}b_{1,3}c_{1,2}]} - e_{[a_{2,3}b_{1,2}c_{1,3}]}, \end{cases}$$

and κ is the vector

$$\kappa = (\tau_2, 0, 0, \tau_3, 0, 0, \tau_6, 0, 0, \tau_7, 0, 0, \tau_4, 0, 0, \tau_5, 0, 0, \tau_1, 0, 0, 0, 0, 0, \tau_8, 0, 0, 0, 0)$$

Moreover, f is Γ series in variables Z, taken with the sign ± 1 , constructed from the lattice \mathcal{B} and the shift vector κ . One takes a variable with the - sign if it occurs in a determinant with the minus sign.

The function f can also be understood in a more familiar way as a function of A_X , B_X , C_X . The transition from variables Z to variables A_X , B_X , C_X is carried out using an obvious substitution

$$[c_1 a_{2,3}] \mapsto C_1 A_{2,3}, \dots \tag{35}$$

There appears mappings pr_a, pr_b, pr_c from the space of exponents for variables Z to the space of exponents for variables A_X, B_X, C_X , respectively:

$$pr_a, pr_b, pr_c : \mathbb{Z}^{30} \to \mathbb{Z}^6.$$

Here \mathbb{Z}^{30} is a lattice of exponents of monomials in variables Z, and \mathbb{Z}^6 is a lattice of exponents of monomials in variables A_X (or B_X , or C_X). In \mathbb{Z}^{30} one has a basis indexed by the variables Z, and \mathbb{Z}^6 has a basis indexed by proper subsets of $X \subset \{1, 2, 3\}$.

Now we can formulate the condition necessary for the 6j-symbol to be nonzero. Consider the expression (33). Let

$$supp f_i = \left(\bigcap_i (\delta_i \ge 0)\right) \cap (\kappa_i + \mathcal{B}_i) \subset \mathbb{Z}^{30}, i = 1, ..., 4$$

put

$$H = supp f_1 \oplus supp f_2 \oplus supp f_3 \oplus supp f_4 \subset \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30}$$
(36)

For each i = 1, ..., 4 there is a mapping $pr_a^i, pr_b^i, pr_c^i : \mathbb{Z}^{30} \to \mathbb{Z}^6$. One can take their direct sum and get a mapping $pr_a^i \oplus pr_b^i \oplus pr_c^i : \mathbb{Z}^{30} \to (\mathbb{Z}^6)^{\oplus 3}$. Consider the mapping

$$pr := \bigoplus_{I=1}^{4} pr_a^i \oplus pr_b^i \oplus pr_c^i : \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \to \bigoplus_{i=1}^{4} (\mathbb{Z}^6)^{\oplus 3}.$$
(37)

In the image of $\bigoplus_{i=1}^{4} (\mathbb{Z}^6)^{\oplus 3}$ one can introduce a basis $e_X^{A^j,i}$, indexed as follows. The lower index is a proper subset of $X \subset \{1, 2, 3\}$. Thus, these indexes list the basis vectors in the selected \mathbb{Z}^6 . There are two upper indexes. The first corresponds to the name of one of the variables (or differential operator), which is substituted in f_i in (??). Thus, fixing this index determines the choice of one of the three terms in $(\mathbb{Z}^6)^{\oplus 3}$. The second upper index is the index i, specifying the term $(\mathbb{Z}^6)^{\oplus 3}$ in $\bigoplus_{i=1}^4 (\mathbb{Z}^6)^{\oplus 3}$.

Thus in the image of $(\mathbb{Z}^6)^{\oplus 3}$, corresponding to f_1 , one has the basis

In the first summand
$$\mathbb{Z}^6$$
: $e_1^{A^1,1}, e_2^{A^1,1}, e_3^{A^1,1}e_{1,2}^{A^1,1}, e_{1,3}^{A^1,1}, e_{2,3}^{A^1,1}$.
In the second summand \mathbb{Z}^6 : $e_1^{A^2,1}, e_2^{A^2,1}, e_3^{A^2,1}e_{1,2}^{A^2,1}, e_{1,3}^{A^2,1}, e_{2,3}^{A^2,1}$.
In the third summand \mathbb{Z}^6 : $e_1^{A^4,1}, e_2^{A^4,1}, e_3^{A^4,1}e_{1,2}^{A^4,1}, e_{1,3}^{A^4,1}, e_{2,3}^{A^1,1}$.

In the image of $(\mathbb{Z}^6)^{\oplus 3}$, corresponding to f_2 , one has the basis

In the first summand \mathbb{Z}^6 : $e_1^{A^4,2}, e_2^{A^4,2}, e_3^{A^4,2}e_{1,2}^{A^4,2}, e_{1,3}^{A^4,2}, e_{2,3}^{A^4,2}$. In the second summand \mathbb{Z}^6 : $e_1^{A^3,2}, e_2^{A^3,2}, e_3^{A^3,2}e_{1,2}^{A^3,2}, e_{1,3}^{A^3,2}, e_{2,3}^{A^3,2}$. In the third summand \mathbb{Z}^6 : $e_1^{A^5,2}, e_2^{A^5,2}, e_3^{A^4,2}e_{1,2}^{A^5,2}, e_{1,3}^{A^5,2}, e_{2,3}^{A^5,2}$.

In the image of $(\mathbb{Z}^6)^{\oplus 3}$, corresponding to f_3 , one has the basis

In the first summand \mathbb{Z}^6 : $e_1^{A^2,3}, e_2^{A^2,3}, e_3^{A^2,3}, e_{1,2}^{A^2,3}, e_{1,3}^{A^2,3}, e_{2,3}^{A^2,3}$. In the second summand \mathbb{Z}^6 : $e_1^{A^3,3}, e_2^{A^3,3}, e_3^{A^3,3}, e_{1,2}^{A^3,3}, e_{1,3}^{A^3,3}, e_{2,3}^{A^3,3}$. In the third summand \mathbb{Z}^6 : $e_1^{A^6,3}, e_2^{A^6,3}, e_3^{A^6,3}, e_{1,2}^{A^6,3}, e_{1,3}^{A^6,3}, e_{2,3}^{A^6,3}$.

In the image of $(\mathbb{Z}^6)^{\oplus 3}$, corresponding to f_4 , one has the basis

In the first summand \mathbb{Z}^6 : $e_1^{A^1,4}, e_2^{A^1,4}, e_3^{A^1,4}, e_{1,2}^{A^1,4}, e_{1,3}^{A^1,4}, e_{2,3}^{A^1,4}$. In the second summand \mathbb{Z}^6 : $e_1^{A^6,4}, e_2^{A^6,4}, e_3^{A^6,4}, e_{1,2}^{A^6,4}, e_{1,3}^{A^6,4}, e_{2,3}^{A^6,4}$. In the third summand \mathbb{Z}^6 : $e_1^{A^5,4}, e_2^{A^5,4}, e_3^{A^5,4}, e_{1,2}^{A^5,4}, e_{1,3}^{A^5,4}, e_{2,3}^{A^5,4}$.

Definition 5. Using the pairing in (33), let us define a sublattice $D \subset \bigoplus_{i=1}^{4} (\mathbb{Z}^6)^{\oplus 3}$ as a lattice generated vectors for all possible $X \subset \{1, 2, 3\}$, which are obtained as differences of vectors with the same X, the same variable A^j , but corresponding to different f_i .

The lattice D is generated by vector

$$\begin{split} & e_X^{A^1,1} - e_X^{A^1,4}, \ \ e_X^{A^2,1} - e_X^{A^2,3}, \ \ e_X^{A^3,2} - e_X^{A^3,4}, \\ & e_X^{A^4,1} - e_X^{A^4,2}, \ \ e_X^{A^5,2} - e_X^{A^5,4}, \ \ e_X^{A^6,3} - e_X^{A^6,4} \end{split}$$

Note the following. Let us be given a monomial which is obtained from the decomposition of (33). It gives a non-zero contribution if for each variable A_X^j , j = 1, ..., 6 the exponent of this variable coincides with the exponent of the differentiation by A_X^j . The fact of existence of such monomials can be reformulated in terms of supports. The result is the following. **Theorem 3.** In order for the 6*j*-symbol (33) to be non-zero, it is necessary that

$$H \cap pr^{-1}(D) \neq \emptyset.$$

where H is defined in (36), and D is defined in the definition of 5.

B Remark 3. This condition can also be formulated in terms of supports of functions f_i as functions of variables A^j . It can be seen that the support $\operatorname{supp}_A f_i$ (a support of a function of variables A^j) is an image of the support supp f_i (a support of a function of variables Z) undert the mapping $pr_a^i \oplus pr_b^i \oplus pr_c^i$, and therefore is also a shifted lattice. Then the necessary condition for 6j-symbol to be non-zero is formulated as follows: $\operatorname{supp}_A(f_1f_2f_3f_4) = pr(H)$ has a non-empty intersection with D. To calculate the value of a 6j-symbol, it is more convenient for us to deal with Z variables.

3.3 A formula for a 6*j*-symbol

The functions f_1, f_2, f_3, f_4 can be considered as functions of the variables Z_1, Z_2, Z_3, Z_4 , where Z_i is a set of variables obtained from the set (34) by replacing the symbols a, b, c by symbols of variables (or differential operators), substituted in f_i in (33). More precise,

For
$$f_1: a, b, c \mapsto A^1, A^2, A^4$$
, for $f_2: a, b, c \mapsto A^4, A^5, A^6$,
for $f_3: a, b, c \mapsto A^2, A^3, A^5$, for $f_4: a, b, c \mapsto A^1, A^6, A^5$ (38)

The computation of the expression (33) can be described as follows. First we consider $f_1...f_4$ as functions of the collection of variables Z_1, Z_2, Z_3, Z_4 . Let us write $f_1...f_4$ is the sums of the products of monomials in these variables. We leave only the terms whose support is in $H \cap pr^{-1}(D) \neq \emptyset$. Next, each of the variables from the set Z_1, Z_2, Z_3, Z_4 , is replaced by the ruler (35) (one takes into account the correspondence of (38)). Thus we pass to the variables A_X^j or $\frac{\partial}{\partial A_X^j}$ in according to the variable or the differential operator is involved in f_i in (33). Multiply them as if they were commuting. After that, in the resulting monomial in A_X^j and $\frac{\partial}{\partial t_X^j}$ one applies the differential operators to variables and substitutes zero instead of all the variables.

So, if one focuses on the variable A_1^1 that occurs in the monomial, then our actions look as follows. Such a symbol occurs in the notation of variables included in the sets Z_1 and Z_4 . We take the monomial resulting from the decomposition of $f_1...f_4$. Let its support belong to $H \cap pr^{-1}(D)$. Let's write it explicitly together with the coefficient at this monomial. It was noted above that $f_1,...,f_4$ are Γ series in the variables Z_1, Z_2, Z_3, Z_4 , so that the coefficient at the monomial is a product of the inverse quantities to the factorials of degrees:

$$\underbrace{\frac{[A_1^1 A_{2,3}^2]^{\beta_1}}{\beta_1!} \underbrace{[A_1^1 A_{2,3}^4]^{\beta_2}}_{\text{PeP} \cdot f_1} \dots \cdots \underbrace{\frac{[A_1^1 A_{2,3}^6]^{\gamma_1}}{\gamma_1!} \underbrace{[A_1^1 A_{2,3}^5]^{\gamma_2}}_{\text{PeP} \cdot f_4}}_{\text{PeP} \cdot f_4} \dots (39)$$

Next, one calculates the sum of the exponents of the variables, whose notation contains the symbol A_1^1 . For factors originating from f_1 , this sum is equal to $\beta_1 + \beta_2 + ...$, and for factors originating from f_4 , this sum is equal to $\gamma_1 + \gamma_2 + ...$ The fact that the supports belong to $H \cap pr^{-1}(D)$ implies that $\beta_1 + \beta_2 + ... = \gamma_1 + \gamma_2 + ...$ When passing to A_X^j or $\frac{\partial}{\partial A_X^j}$, one substitutes $\frac{\partial}{\partial A_1^1}$ into factors originating from f_1 , and one substitutes A_1^1 in factors originating from f_4 . After applying the differential operator to the variable A_1^1 and substituting zero instead of A_1^1 in (39), all symbols A_1^1 are actually deleted and a numerical multiplier $(\beta_1 + \beta_2 + ...)!$ is added in the top.

After performing similar actions with all the variables A_X^j , the monomial (39) turns into a numeric fraction. In its denominator occur the factorials (in the multi-index sense) of the degree of the monomial as a monomial of the variables $Z_1, ..., Z_4$ and at the top occur the factorials of the degree (again in the multi-index sense) of the monomial as a monomial in the variables A_X^j .

Let us proceed to the formulation of the main result of this paper. The set $H \cap pr^{-1}(D)$ is a shifted lattice. Therefore, for some vector \varkappa and some lattice $L \subset \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30}$, one can write

$$H \cap pr^{-1}(D) = \varkappa + L \subset \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \oplus \mathbb{Z}^{30} \tag{40}$$

There is a projection pr defined by the formula (37). One associates with the shifted lattice $\varkappa + L$ a series of hypergeometric type (in fact, being a finite sum) in the variables $Z = \{Z_1, ..., Z_4\}$, defined by a formula related to (10):

$$\mathcal{J}_{\gamma}(Z;L) = \sum_{x \in \varkappa + L} \frac{\Gamma(pr(x) + 1)Z^x}{\Gamma(x+1)}$$
(41)

Theorem 4. The 6*j*-symbol (33) is equal to $\mathcal{J}_{\gamma}(\pm 1; L)$, where 1 is substituted instead of those variables that are entering in the determinants in f_i the sign + and -1 is substituted instead of those variables that are entering in the determinants with the sign -.

The shifted lattice $\varkappa + L$ is defined in (40), where $H = \oplus H_i$, $H_i = supp f_i$; D is given in the Definition 5, and pr is defined in (37).

References

 Etingof P., Gelaki S., Nikshich D., Ostrik V., Tensor Categories, AMS: Mathematical Surveys and Monographs, Providence, V 205; 2015.

- [2] Racah, G. Theory of Complex Spectra II. Physical Review. 1942, v. 62, N 9–10, 438–462.
- [3] Wigner, E. P. Group Theory and its Application to the Quantum Mechanics of Atomic Spectra. New York : Academic Press, 1959
- [4] L. D. Landau; E. M. Lifshitz (1977). Quantum Mechanics: Non-Relativistic Theory. Vol. 3 (3rd ed.). Pergamon Press. 2nd ed. (1965)
- [5] D. Varshalovich, A. Moskalev, V. Khersonskii: Quantum theory of angular momentum. World Scientific, Singapur 1989.
- [6] L. C. Biedenharn, J. D. Louck: Angular Momentum in Quantum Physics, Addison-Wesley, 1981
- [7] N. Ja. Vilenkin, A. U. Klimyk, Representation of Lie Groups and Special Functions, Volume 1: Simplest Lie Groups, Special Functions and Integral Transforms, Springer Dordrecht, 1991.
- [8] S. E. Derkachov, G. A. Sarkissian, V. P. Spiridonov, Elliptic hypergeometric function and 6j-symbols for the SL(2, C) group, Theoret. and Math. Phys., 213:1 (2022), 1406–1422
- [9] S. E. Derkachev, A. V. Ivanov, Racah coefficients for the group SL(2, ℝ)", Questions of quantum field theory and statistical physics. Part 28, Zap. Nauchn. Sem. POMI, 509, POMI, St. Petersburg, 2021, 99–112
- [10] Rebbi C., Slansky R. Crossing Matrices for SU(2) and SU(3), Reviews of Modern Physics. 1970. v. 42, 1. p. 68.
- [11] Arnold P. Landau-Pomeranchuk-Migdal effect in sequential bremsstrahlung: From large-N QCD to N = 3 via the SU(N) analog of Wigner 6 j symbols, Physical Review D. 2019. v. 100, N 3. p. 034030.
- [12] Sleptsov, A dissertation.
- [13] Butler P., Wybourne B. Calculation of j and jm symbols for arbitrary compact groups. I. Methodology. International Journal of Quantum Chemistry. 1976. v. 10, N 4. p. 581–598.
- [14] Hecht K. A simple class of U(N) Racah coefficients and their application. Communications in Mathematical Physics. 1975. v. 41, N 2. p. 135–156.
- [15] Gustafson R. A Whipple, Transformation for Hypergeometric Series in U(N) and Multivariable Hypergeometric Orthogonal Polynomials. SIAM journal on mathematical analysis. 1987. v. 18, N2 . p. 495–530.

- [16] Wong M. On the multiplicity-free Wigner and Racah coefficients of U(n). Journal of Mathematical Physics. 1979. v. 20, N 12. p. 2391–2397.
- [17] Biedenharn L.C., Louck J. D., Canonical Unit Adjoint Tensor Operators in U(n), J. Math. Phys. v. 11, 2368 (1970);
- [18] Biedenharn L.C. , Louck J. D., Chacon E. , Ciftan M., On the structure of the canonical tensor operators in the unitary groups. I. An extension of the pattern calculus rules and the canonical splitting in U(3), J. Math. Phys. 1972, v. 13, p. 1957
- [19] Mironov A., Morozov A., Sleptsov A. On 6j-symbols for symmetric representations of $U_q(SU(n))$. JETP Letters. 2017. v. 106, N 10. p. 630–636.
- [20] Alekseev V., Morozov A., Sleptsov A. Multiplicity-free $U_q(SU(n))$ 6 j symbols: Relations, asymptotics, symmetries. Nuclear Physics B., 2020. v. 960. p. 115164.
- [21] D. V. Artamonov, Formulas for calculating the 3j-symbols of the representations of the Lie algebra gl_3 for the Gelfand-Tsetlin bases, Sibirsk. Mat. Zh., 63:4 (2022), 717–735
- [22] Biedenharn L.C., Baid G.E., On the representations of semisimple Lie Groups II, J. Math. Phys., 1963, v. 4, N 12, p. 1449–1466
- [23] Artamonov D. V., The Clebsh-Gordan coefficients for the algebra gl3 and hypergeometric functions, Algebra i Analiz, 33:1 (2021), 1–29
- [24] Gel'fand I. M., Graev M. I., Retakh V. S., General hypergeometric systems of equations and series of hypergeometric types Tok, Russian Math. Surveys, 47:4 (1992), 1–88
- [25] Artamonov D.V., Antisymmetrization of the Gelfand-Kapranov-Zelevinskij Systems B ,J Math Sci , v. 255, N 5, C Í. 535-542
- [26] Zhelobenko D.P, Compact Lie groups and the representations, Translations of Mathematical Monographs Volume: 40; 1973