

Φ_3^4 measures on compact Riemannian 3-manifolds

I. BAILLEUL and N. V. DANG and L. FERDINAND and T.D. TÔ

Abstract. We construct the Φ_3^4 measure on an arbitrary 3-dimensional compact Riemannian manifold without boundary as an invariant probability measure of a singular stochastic partial differential equation. Proving the nontriviality and the covariance under Riemannian isometries of that measure gives for the first time a non-perturbative, non-topological interacting Euclidean quantum field theory on curved spaces in dimension 3. This answers a longstanding open problem of constructive quantum field theory on curved 3 dimensional backgrounds. To control analytically several Feynman diagrams appearing in the construction of a number of random fields, we introduce a novel approach of renormalization using microlocal and harmonic analysis. This allows to obtain a renormalized equation which involves some universal constants independent of the manifold. We also define a new vectorial Cole-Hopf transform which allows to deal with the vectorial Φ_3^4 model where Φ is now a bundle valued random field. In a companion paper, we develop in a self-contained way all the tools from paradifferential and microlocal analysis that we use to build in our manifold setting a number of analytic and probabilistic objects.

Contents

1. Introduction	1
2. Long time well-posedness and a priori estimate	6
3. Scaling, regularity and microlocal extension	16
4. Induction on Feynman amplitudes	28
5. Random fields from renormalization	40
6. Invariant measure	63
7. Non-constant coupling function and vector bundle cases	68
A. Littlewood-Paley-Stein projectors	72

1 – Introduction

In the setting of the discrete d -dimensional torus $\Lambda_d = \left(\frac{\mathbb{Z}}{N\mathbb{Z}}\right)^d$, for $d \geq 2$, a field is represented by a real-valued function σ on Λ_d . Write $i \sim j$ when two points i and j are neighbours in Λ_d . One can assign an energy

$$(1.1) \quad S(\sigma) = \frac{1}{2} \sum_{i \sim j} |\sigma_i - \sigma_j|^2 + \frac{1}{2} \sum_i |\sigma_i|^2$$

to any field σ and define a Gibbs probability measure ν_{Λ_d} proportional to

$$e^{-S(\sigma)} \prod_{i \in \Lambda_d} d\sigma_i.$$

A random variable with values in the space $\mathcal{E}_d = \mathbb{R}^{\Lambda_d}$ of fields, with law ν_{Λ_d} , is called a discrete Gaussian free field. The continuum analogue of this random variable is the Gaussian free field on the torus \mathbb{T}^d , characterized by the fact that it is a random centered Gaussian field ζ with covariance $(1 - \Delta)^{-1}$ on the torus \mathbb{T}^d . This means that for any smooth real-valued test function $f \in C^\infty(\mathbb{T}^d)$ the random variable $\zeta(f)$ is Gaussian with zero mean and covariance $\langle f, (1 - \Delta)^{-1} f \rangle_{L^2(\mathbb{T}^d)}$. One can construct ζ as a random distribution that is almost surely of Besov-Hölder regularity $-\frac{d-2}{2} - \varepsilon$, for all $\varepsilon > 0$. Note the dependence of the regularity exponent on the dimension. Back to the discrete setting, we denote by $\Delta_{\mathcal{E}_d}$ the canonical Laplace operator on the field space $\mathcal{E}_d \simeq \mathbb{R}^{\Lambda_d}$ and by $\mathbf{1}$ the constant function on \mathcal{E}_d equal to 1. A dynamical picture of the Gibbs measure ν_{Λ_d} can be obtained from the genuine identity

$$\nabla_{\mathcal{E}_d} \mathbf{1}_{\mathcal{E}_d} = 0$$

by rewriting it under the form

$$\nabla_{\mathcal{E}_d} (e^{-S} \nabla_{\mathcal{E}_d} e^S) (e^{-S}) = 0.$$

The density e^{-S} appears here as an element of the kernel of the dual of the conjugated operator

$$\{\nabla_{\mathcal{E}_d} (e^{-S} \nabla_{\mathcal{E}_d} e^S)\}^* = \Delta_{\mathcal{E}_d} - (\nabla_{\mathcal{E}_d} S) \cdot \nabla_{\mathcal{E}_d}.$$

This allows a construction of ν_{Λ_d} as the invariant measure of the Markov process on \mathcal{E}_d with generator $\Delta_{\mathcal{E}_d} - (\nabla_{\mathcal{E}_d} S) \cdot \nabla_{\mathcal{E}_d}$ – provided this Markov process has indeed a unique invariant probability measure. The diffusion associated with the operator $\Delta_{\mathcal{E}_d} - (\nabla_{\mathcal{E}_d} S) \cdot \nabla_{\mathcal{E}_d}$ is the solution of the stochastic differential equation in \mathcal{E}_d

$$dz_t = \sqrt{2}dw_t - \nabla_{\mathcal{E}_d} S(z_t)dt,$$

for a Brownian motion w in \mathcal{E}_d .

The energy S in (1.1) contains a kinetic term $\sum_{i \sim j}$ and a potential term \sum_i . One can add to the potential term an additional bit of the form

$$Q(\sigma) = \sum_i Q(\sigma_i).$$

for a real-valued function Q , typically a real polynomial bounded from below for the so-called Ginzburg–Landau models. The corresponding Gibbs probability measure μ_{Λ_d} can then be seen as a perturbation of the discrete Gaussian free field probability measure ν_{Λ_d}

$$\mu_{\Lambda_d}(d\sigma) \sim e^{-Q(\sigma)}\nu_{\Lambda_d}(d\sigma).$$

The dynamics on \mathcal{E}_d associated with this measure is given as above by the stochastic differential equation

$$(1.2) \quad dz_t = \sqrt{2}dw_t - \nabla_{\mathcal{E}_d}(S + Q)(z_t)dt.$$

There is a problem for taking the formal continuum limit of these measures as the Gaussian free field measure is supported on a set of distributions of negative regularity, so an expression like $\int Q(\sigma)$ does not make sense for a nonlinear function Q of a distribution σ . It is the aim of the Euclidean quantum field theory of scalar fields to make sense of and construct such measures for some particular examples of potentials Q . The Φ_3^4 measure corresponds to $Q(a) = a^4/4$ in a 3-dimensional setting.

Quantum field theory was developed as a theory describing interactions of the elementary particles. It is arguably one of the most successful physical theory of the 20th century and has led to remarkable physical predictions with unprecedented numerical accuracy. However in spite of its success in theoretical physics a complete mathematical formulation and understanding of quantum field theory is still work in progress. A major difficulty in the subject comes from the divergences inherent to the formulation of the theory. In quantum field theory the perturbative calculation of any physical process involves a summation over an infinite number of virtual multi-particle states which is generically divergent, hence produces infinities. The divergences of perturbation theory in quantum field theory are directly linked to its short distance structure which is highly non-trivial because its description involves the infinity of multi-particle states. These divergences must be carefully subtracted in some organized way compatible with physical requirements such as locality, causality, unitarity. The methods developed to deal with these infinities were called *renormalization*. Constructive quantum field theory provides one of the mathematically rigorous approaches to quantum field theory. It was developed in the 70s with seminal contributions of Albeverio, Brydges, Feldman, Fröhlich, Gallavotti, Gawedzki, Glimm, Guerra, Jaffe, Kupiainen, Nelson, Rivasseau, Seiler, Sénéor, Spencer, Simon, Symanzik and Wightman to name but a few – see e.g. Glimm & Jaffe’s book [40] and the references inside for an account of the early achievements in this domain. One of the first successes of constructive quantum field theory was the construction of the so called $P(\phi)_2$ -model on \mathbb{R}^2 and of the ϕ_3^4 theory on \mathbb{R}^3 by Glimm & Jaffe [38, 39] in the 70s. The recent breakthroughs by Hairer [51] and Gubinelli, Perkowski & Imkeller [49] allowed several authors to recover the results of Glimm & Jaffe following the stochastic quantization program of Parisi & Wu [73], using only PDE and probabilistic techniques without the intricate combinatorial methods from the constructive school. However we would like to emphasize that the powerful cluster expansion techniques developed by the constructive school allowed, for both $P(\phi)_2, \phi_3^4$ theories, to show the Borel summability of the partition function in the couplings, to control the mass gap (exponential decay of the correlations), to study phase transitions (when there is no longer exponential decay of correlations), to investigate fine properties of the spectrum and scattering properties of these theories and finally, one of the deepest results of the constructive school was the stability proofs of Yang–Mills in 3 and 4 dimensions due to Balaban [12, 13]. Many of these

results are currently out of reach of the stochastic methods: We refer to the book of Rivasseau [77] for more information on the topic. The dynamical construction of the Φ_3^4 measure was first done in finite volume on the flat 3-dimensional torus by Mourrat & Weber [68], Hairer & Mattingly [52], Hairer & Schönbauer [53], Albeverio & Kusuoka [3], then in the infinite volume 3-dimensional Euclidean space by different authors – Albeverio & Kusuoka [4], Moinat & Weber [66], Gubinelli & Hofmanová [45, 46], Barashkov & Gubinelli [14, 15], and Duch [32, 33], using different methods. A crucial integrability property of the ϕ_3^4 measure was proved in Hairer & Steele’s work [54]. Gubinelli’s lecture notes [43, 44] provide a remarkable source of inspiration and information on the stochastic quantization approach to the construction of the Φ_3^4 measure.

Yet most results in constructive quantum field theory are proved in the geometric settings of either \mathbb{R}^3 or the flat torus \mathbb{T}^3 . On the other hand quantum field theory on curved spacetimes has been studied since the 70s. Recent breakthroughs by Brunetti & Fredenhagen [20], Hollands & Wald [55, 56] and Rejzner [76] on Lorentzian manifolds (see [27] for a detailed mathematical exposition of part of this approach) and Kopper & Müller [61] in a Riemannian setting, lead to complete proofs of *perturbative renormalization* on (pseudo)-Riemannian manifolds of dimension less than or equal to 4. We also mention the approach of Costello [24] that is designed to work on Riemannian manifolds, possibly with a boundary after Albert’s work [2]. However all these results are perturbative and construct quantum field theory objects as formal power series. They do not provide any probability measures, Hilbert spaces or operators in a straightforward way. Constructive quantum fields on manifolds have been earlier addressed only on *compact surfaces* in [74] by Pickrell and [30] by Dimock for the $P(\varphi)_2$ theories, in [62] by Lévy for the 2d Yang-Mills theory and in [48] by Guillarmou, Kupiainen, Rhodes & Vargas for the Liouville field theory on Riemann surfaces. In Lorentz signature, we would also like to mention the work [16] of Barata, Jaekel & Mund who managed to define the $P(\varphi)_2$ theory on 2-dimensional de Sitter space, extending previous work [36] of Figari, Høegh-Krohn & Nappi. From the PDE side, the constructions of Gibbs measures on Riemannian surfaces that we are aware of, come from [22] by Burq, Thomann & Tzvetkov for dynamical $P(\Phi)_2$ and from [72] by Oh, Robert, Tzvetkov & Wang for the dynamical Liouville model. It is therefore a longstanding open problem in both constructive quantum field theory and field theory on curved spaces to construct the ϕ_3^4 measure on an arbitrary closed, compact 3-dimensional Riemannian manifold. As emphasized by Witten in the recent work [80]

“If a theory exists perturbatively in curved spacetime, and non-perturbatively in flat spacetime, one would expect that it works non-perturbatively in curved spacetime. Unfortunately, not much is available in terms of rigorous theorems, except for special models like two-dimensional conformal field theories. That reflects the general mathematical difficulty of understanding quantum field theory rigorously. One would think that rigorous results for a superrenormalizable theory in curved spacetime might be relatively accessible, but such results are not available.”

Let us mention that both the 2d Yang-Mills and Liouville theories are *integrable* as mentioned above by Witten. Our work seems to give the first construction of a non-integrable, interacting quantum field theory on 3-manifolds.

Let (M, g) stand for a closed 3-dimensional Riemannian manifold. This work is dedicated to constructing the Φ_3^4 measure over M , formally the ill-defined functional integral measure

$$\frac{e^{-\frac{1}{2} \int_M (|\nabla u|^2 + u^2) - \frac{1}{4} \int_M u^4 [\mathcal{D}u]}}{\int e^{-\frac{1}{2} \int_M (|\nabla u|^2 + u^2) - \frac{1}{4} \int_M u^4 [\mathcal{D}u]}}$$

as an invariant probability measure of the dynamics

$$(1.3) \quad (\partial_t + P)u = -u^3 + \sqrt{2}\xi$$

where ξ stands for a spacetime white noise and $P \stackrel{\text{def}}{=} (1 - \Delta_g)$. The noise ξ plays in a continuum setting the role of the Brownian motion w in (1.2) in a discrete setting while the terms Pu and u^3 come from the gradient terms of the energy S and the quartic potential Q respectively. The construction of the Φ_3^4 measure as the hopefully unique invariant probability measure of this dynamics was first put forward by Parisi & Wu in a famous work of the early 80s; it comes under the name of *stochastic quantization*. Note that we define the Φ_3^4 measure as an invariant measure from Equation (1.3). This is a priori not equivalent to obtaining the measure as a scaling limit of lattice models, a highly non-trivial issue on manifolds since there is no canonical way of

discretising quantum field theories on manifolds. We do not try to relate here our construction of the Φ_3^4 measure with any such limiting procedure.

Equation (1.3) involves the fundamental problem of considering a nonlinear function of a distribution. Spacetime white noise on a 3-dimensional Riemannian manifold has indeed almost surely a parabolic¹ Besov-Hölder regularity $-5/2 - \varepsilon$, for all $\varepsilon > 0$, so one does not expect from a possible solution u to equation (1.3) that it has parabolic regularity better than $-1/2 - \varepsilon$, as a consequence of Schauder estimate. The term u^3 in (1.3) is thus ill-defined. This kind of problem in a stochastic partial differential equation (PDE) is characteristic of the class of singular stochastic PDEs, a field that was opened around 2014 by the groundbreaking works of M. Hairer on regularity structures [51] and Gubinelli, Imkeller & Perkowski [49] on paracontrolled calculus. The tools of regularity structures and paracontrolled calculus were used to run the stochastic quantization approach for constructing of the Φ_3^4 measure over a 3-dimensional torus and Euclidean space in a series of works. Local well-posedness of equation (1.3) was proved first by Hairer in [51] – see also Catellier & Chouk’s work [23] for a proof of that result with the tools of paracontrolled calculus. Mourrat & Weber proved in [68] an a priori estimate that gives the long time existence (and well-posedness) of the solution to (1.3) and the existence of an invariant probability measure. The uniqueness of this invariant probability measure comes from the works of Hairer & Mattingly [52] on the strong Markov property of transition semigroups associated to singular stochastic PDEs, and Hairer & Schönbauer [53] on the support of the laws of solutions to singular stochastic PDEs. See Hairer & Steele’s work [54] for more references.

None of the previous works are readily available in a manifold setting, and so far the only works on singular stochastic PDEs in a manifold setting can be divided in two groups:

- The works [7, 8, 9] of Bailleul & Bernicot, and the works [29], [70] and [11] of Dahlqvist, Diehl & Driver, Mouzard and Bailleul, Dang & Mouzard on the Anderson operator on a 2-dimensional Riemannian manifold. We refer to [71] for a lucid exposition of some of the above results.
- The works [22] by Burq, Thomann & Tzvetkov for dynamical $P(\Phi)_2$ and from [72] by Oh, Robert, Tzvetkov & Wang for the dynamical Liouville model, both works deal with interacting QFT measures on surfaces.

We would also like to mention the work of Hairer & Singh [42] which develops a generalisation of the original Theory of Regularity Structures which is able to treat SPDEs on manifolds with values in vector bundles in full generality. Their work gives *only short time existence* for the SPDE which overlaps with our own result. However, to build the QFT measure requires the long time existence and the coming down from infinity property for the solutions of the SPDE, which are not covered by [42].

The aim of the present work together with our companion work [10] is to develop in a self-contained way all the tools needed to run the analysis in a 3 dimensional closed Riemannian manifold. On the purely analytical side

- We follow Jagannath & Perkowski’s simple approach [57] of equation (1.3) to prove that this equation is locally well-posed. Their formulation of the problem avoids the use of regularity structures or paracontrolled calculus. We freely use several tools of pseudodifferential and paradifferential calculus on manifolds which are developed in detail in [10].
- We study the vectorial Φ_3^4 model, sometimes called the $O(N)$ model in the physics literature, where Φ is now E -valued, with (E, h) some Hermitian vector bundle over M . In this case the SPDE reads

$$(\partial_t + P)\Phi = -\Phi \langle \Phi, \Phi \rangle_h + \sqrt{2}\xi_E$$

where Δ is a generalized laplacian acting on E -valued sections, ξ_E is some E -valued white noise and the Φ_3^4 measure is still invariant under the Markovian dynamics. The crucial ingredient for this part is a novel vectorial Cole-Hopf transform that we introduce in definition 42.

¹Meaning the time variable has weight 2 whereas space variables have weight 1.

- We give a simple and short proof of an L^p “coming down from infinity” property satisfied by the solution to equation (1.3) using energy methods. The longtime existence of a unique solution to equation (1.3) follows as a consequence.
- As usual in the study of singular stochastic PDEs we need to feed the analytic machinery with a number of random distributions whose formal definitions involve some ill-defined products and whose actual definitions involve some probabilistic constructions based on regularization and renormalization. Our approach to the renormalization problem relies on microlocal analysis. It is a far reaching generalization of the Epstein-Glaser point of view where we benefit from the many improvements contained in [75, 20, 55, 56]. We reduce the problem of renormalization to an extension problem for distributions on a configuration space defined outside all the diagonals, for which we develop a general machinery. To control analytically the Feynman amplitudes appearing in the stochastic bounds we feed our renormalization machine with several microlocal estimates of distributional kernels which are done separately in [10].

We note that there is also a new approach to SPDE’s relying on the Epstein-Glaser renormalization in the works [31] by Dappiaggi, Drago, Rinaldi & Zambotti and [19] by Bonicelli, Dappiaggi & Rinaldi. However it seems that these authors work only at a perturbative level whereas our results are non-perturbative.

One remarkable feature of our approach is that we are able to renormalize Equation (1.3) using universal counterterms – they do not depend on the metric on M . We emphasize that fact in the following statement where $\varepsilon \in (0, 1/8)$ is a positive constant and $\xi_r \stackrel{\text{def}}{=} e^{-rP}\xi$ stands for a *space regularization* of ξ by the heat operator – so ξ_r is still white in time. Set

$$(1.4) \quad \begin{aligned} a_r &\stackrel{\text{def}}{=} \frac{r^{-1/2}}{4\sqrt{2}\pi^{3/2}}, \\ b_r &\stackrel{\text{def}}{=} \frac{|\log r|}{32\pi^2}. \end{aligned}$$

Theorem 1 – Pick $\phi \in C^{-1/2-\varepsilon}(M)$. The equation

$$(1.5) \quad (\partial_t - \Delta + 1)u_r = \sqrt{2}\xi_r - u_r^3 + 3(a_r - b_r)u_r$$

with initial condition ϕ has a unique solution over $[0, \infty) \times M$ in some appropriate function space. For any $0 < T < \infty$ this random variable converges in probability in $C([0, T], C^{-1/2-\varepsilon}(M))$ as $r > 0$ goes to 0 to a limit u .

The function u is what we define as the solution to Equation (1.3) and it turns out to be a Markov process. The a priori estimate encoded in the coming down from infinity property provides a compactness statement from which the existence of an invariant probability measure for the Markovian dynamics follows.

Theorem 2 – The dynamics of u is Markovian and its associated semigroup on $C^{-1/2-\varepsilon}(M)$ has an invariant non-Gaussian probability measure.

A Φ_3^4 measure over M is that invariant measure. Note that the constants a_r, b_r in (1.5) are universal in the sense that they do not depend on the Riemannian metric on M . They depend however on the regularization scheme we use, here the heat regularization in space which is *fully covariant* with respect to the Riemannian structure.

The study of Equation (1.3) on an arbitrary fixed time interval $[0, T]$ is the object of Section 2. We follow [57] which yields a robust formulation of Equation (1.3) which avoids the use of regularity structures or paracontrolled calculus. The local in time well-posedness of (1.3) is proved in Section 2.1. We get the longtime existence from the r -uniform L^p ‘coming down from infinity’ property satisfied by u_r , proved in Section 2.2. This property is proved in the formulation of Equation (1.3) as in [57]. The results of Section 2 show that u_r depends continuously on a finite family $\widehat{\xi}_r$ of multilinear functionals of ξ_r . The functional setting needed to prove their convergence in some appropriate space as r goes to 0 is detailed in Section 3. A crucial role is played here by a set of distributions with given wavefront sets and a certain scaling property with respect to some submanifolds. The notion of scaling field is introduced in Section 3.1 and the preceding

set of distributions is introduced in Section 3.2. We prove our main workhorse in Section 3.3, Theorem 13. It provides a numerical criterion for a distribution defined outside a submanifold, with some wavefront set bound and some scaling property with respect to that submanifold, to have a possibly unique extension to the whole manifold. We draw consequences of this general statement for the particular case of a configuration space in Section 4.1. The relevance of Theorem 13 to the convergence problem for $\widehat{\xi}_r$ stems from the fact that we can formulate the latter as a quantitative extension problem. This point of view is inspired by the Epstein-Glaser approach of renormalization. The convergence of $\widehat{\xi}_r$ in an appropriate space is the object of Section 5. At that point of the analysis of Equation (1.3) one can make sense of its $r = 0$ version as a Markovian Feller dynamics in the state space $C^{-1/2-\varepsilon}(M)$, for any fixed sufficiently small $\varepsilon > 0$. The r -uniform L^p coming down property from Section 2.2 is used to get the existence of an invariant probability measure for this dynamics. The non-triviality of this invariant probability measure is proved in Section 6.2. Last Section 7 is dedicated to the construction of a Φ_3^4 measure corresponding to a space-dependent coupling constant and also we discuss the bundle case. The detailed proofs of a number of basic tools from microlocal and harmonic analysis are given in our companion paper [10]. They are mainly put to work in our analysis of the convergence of the enhanced noise $\widehat{\xi}_r$ in Section 5.

Acknowledgements – We would like to thank C. Bellingeri, C. Brouder, N. Burq, C. Dappiaggi, P. Duch, C. Gérard, C. Guillarmou, F. Hélein, K. Lê, D. Manchon, P.T. Nam, S. Nonnenmacher, V. Rivasseau, G. Rivière, N. Tzvetkov, F. Vignes-Tourneret for interesting questions, remarks, comments on the present work when we were in some preliminary stage and also simply for expressing some interest and motivating us to pursue. Special thanks are due to Y. Bonthonneau, C. Brouder, J. Dereziński, M. Hairer and M. Wrochna for their very useful comments on a preliminary version of our work. Special thanks to A. Mouzard who found an important mistake in a previous version and for his strong support in the revision process. We also would like to thank M. Gubinelli, Y. Dabrowski, M. Lewin who suggested us that certain techniques from microlocal analysis used for QFT on curved spaces could be merged with constructive or SPDE methods in order to prove the present results. N.V.D. acknowledges the support of the Institut Universitaire de France. The authors would like to thank the ANR grant SMOOTH "ANR-22-CE40-0017", QFG "ANR-20-CE40-0018" and MARGE "ANR-CE40-0011-01" for support.

Notations – Given $0 < T < \infty$ and a Banach space B we write $C_T B$ for $C([0, T], B)$. The parabolic Besov-Hölder spaces on space time are denoted by $C^\gamma((a, b) \times M)$, the Besov spaces on the manifold M are denoted by $B_{p,q}^a(M)$, these spaces being defined in Definition 29. The cotangent space to M is denoted by T^*M , the conormal to a submanifold E of M is denoted by $N^*(E)$, and we denote by $dv(x)$ the volume form on M . Throughout this paper we are given a finite cover of M with some open charts $M = \cup_{i \in I} U_i$, where I is a fixed finite set. By localising on these charts, we define in Appendix A using the flat Laplacian some Littlewood-Paley blocks $(P_j^i, \widetilde{P}_j^i)_{j \geq -1}^{i \in I}$ on M – see (A.1), along with some paraproducts and resonant term $(\prec_i, \succ_i, \odot_i)_{i \in I}$ such that for any smooth functions $a, b \in \mathcal{D}(M)$ one has

2 – Long time well-posedness and a priori estimate

We prove the existence of a unique solution to equation (1.3) over any fixed time interval $[0, T]$, for an arbitrary initial condition in $C^{-1/2-\varepsilon}(M)$, for $\varepsilon > 0$ small enough. We adopt here the robust approach of [57]. They use a clever change of variable to reformulate the equation as a non-singular parabolic partial differential equation (2.4) with random coefficients of regularity no worse than $-1/2-\varepsilon$. This allows to solve the equation locally in time by a fixed point argument set in a classical functional space without resorting to regularity structures or paracontrolled calculus. Section 2.2 is dedicated to proving an L^p estimate on the solution to equation (2.4) that is independent of the initial condition. This plays a crucial role in proving the existence of an invariant probability measure for (1.3) by an argument using compactness.

2.1 Local in time well-posedness

This section is dedicated to proving the local well-posedness of a solution to equation (1.3), uniformly over $r > 0$. In [57], the authors noticed that a clever reformulation of the equation brings its study back to the study of a nonsingular stochastic PDE for which local in time well-posedness follows from an elementary fixed point argument.

Some distributions in the list (2.1) below involve an operator \odot , called *resonant operator*, that we introduce formally in Appendix A; its precise definition here does not matter other than the fact that it is well-defined and continuous from $B_{p_1, q_1}^{\alpha_1}(M) \times B_{p_2, q_2}^{\alpha_2}(M)$ into some Besov space if and only if $\alpha_1 + \alpha_2 > 0$, in which case it takes values in $B_{p, q}^{\alpha_1 + \alpha_2}(M)$, for some integrability exponents p, q whose precise value does not matter here. For $\Lambda_1 \in B_{p_1, q_1}^{\alpha_1}(M)$ and $\Lambda_2 \in B_{p_2, q_2}^{\alpha_2}(M)$, the product $\Lambda_1 \Lambda_2$ is well-defined if and only if $\Lambda_1 \odot \Lambda_2$ is well-defined.

2.1.1 The enhanced noise. We regularize ξ in space (only) using the heat kernel and set

$$\xi_r \stackrel{\text{def}}{=} e^{-rP}(\xi);$$

this is a white noise in time with values in a space of regular functions. The fact that ξ appears in an additive form in (1.3) does not make it necessary to regularize it in time. Regularising ξ only in space makes clear the Markovian character of the renormalized equation (1.5). Denote by \mathcal{L}^{-1} , respectively $\underline{\mathcal{L}}^{-1}$, the resolvent operator of $\partial_t + P$ with null initial condition at time 0, respectively at time $-\infty$. Explicitly, $\mathcal{L}^{-1}f(t, \cdot) = \int_0^t e^{-(t-s)P} f(s, \cdot) ds$ and $\underline{\mathcal{L}}^{-1}f(t) = \int_{-\infty}^t e^{-(t-s)P} f(s, \cdot) ds$ for any function f on $\mathbb{R} \times M$. The operator $\underline{\mathcal{L}}^{-1}$ provides stationary solutions. Recall from (1.4) the definitions of the constants a_r and b_r and set

$$\mathfrak{I}_r \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\xi_r)$$

and

$$\mathfrak{V}_r \stackrel{\text{def}}{=} (\mathfrak{I}_r)^2 - a_r, \quad \mathfrak{Y}_r \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\mathfrak{V}_r)$$

$$\mathfrak{W}_r \stackrel{\text{def}}{=} (\mathfrak{I}_r)^3 - 3a_r \mathfrak{I}_r, \quad \mathfrak{Y}_r^{\circ} \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\mathfrak{W}_r)$$

and

$$(2.1) \quad \widehat{\xi}_r \stackrel{\text{def}}{=} \left(\xi_r, \mathfrak{V}_r, \mathfrak{W}_r, \mathfrak{Y}_r^{\circ} \odot \mathfrak{I}_r, \mathfrak{Y}_r \odot \mathfrak{V}_r - \frac{b_r}{3}, |\nabla \mathfrak{Y}_r^{\circ}|^2 - \frac{b_r}{3}, \mathfrak{Y}_r^{\circ} \odot \mathfrak{V}_r - b_r \mathfrak{I}_r \right).$$

One has $\xi_r \in \mathcal{C}^{-5/2-\varepsilon}([0, T] \times M)$ and the restriction to any time interval $[0, T]$ of the other components of $\widehat{\xi}_r$ is seen as an element of the product space

$$(2.2) \quad C_T C^{-1-2\varepsilon}(M) \times \mathcal{C}^{-3/2-3\varepsilon}([0, T] \times M) \times \mathcal{C}^{-4\varepsilon}([0, T] \times M)^3 \times \mathcal{C}^{-1/2-5\varepsilon}([0, T] \times M).$$

Note that only \mathfrak{V}_r is an element of a space of the form $C_T C^\gamma(M)$, the other terms in $\widehat{\xi}_r$ are elements of a parabolic space of negative regularity, which is less precise than being an element in a space of the form $C_T C^\gamma(M)$ for a negative exponent γ . This is sufficient for our needs. The enhancement $\widehat{\xi}_r$ can be seen as a placeholder for a number of products that are not well-defined in the zero regularization limit. We will see in Section 5 that $\widehat{\xi}_r$ converges in all the $L^p(\Omega)$ spaces, $1 \leq p < \infty$, as $r > 0$ goes to 0, to a limit that does not depend on the mollification used to define ξ_r from ξ . Using the operator $\underline{\mathcal{L}}^{-1}$ rather than the operator \mathcal{L}^{-1} in the definitions of \mathfrak{Y}_r and \mathfrak{Y}_r° builds some random distributions that are stationary in time. This property will be useful in Section 6 to get a compactness statement on the family of laws of the solutions to (1.5).

2.1.2 Jagannath & Perkowski's formulation of equation (1.5). Set

$$v_{r, \text{ref}} \stackrel{\text{def}}{=} 3 \underline{\mathcal{L}}^{-1} \left(e^{3\mathfrak{Y}_r^{\circ}} \left\{ \mathfrak{Y}_r^{\circ} \mathfrak{V}_r - b_r (\mathfrak{I}_r + \mathfrak{Y}_r^{\circ}) \right\} \right).$$

This is an element of $C_T C^{1-\varepsilon}(M)$. The starting point for the analysis of the renormalized form (1.5) of equation (1.3) is that u_r is a solution to (1.5) if and only if

$$(2.3) \quad v_r \stackrel{\text{def}}{=} e^{3\mathfrak{Y}_r^{\circ}} \left(u_r - \mathfrak{I}_r + \mathfrak{Y}_r^{\circ} \right) - v_{r, \text{ref}}$$

is a solution of a particular equation of the form

$$(2.4) \quad (\partial_t + P)v_r = -6\nabla\mathfrak{Y}_r^\circ \cdot \nabla v_r - e^{-6\mathfrak{Y}_r^\circ} v_r^3 + Z_{2,r}v_r^2 + Z_{1,r}v_r + Z_{0,r},$$

where $Z_{2,r}, Z_{1,r}, Z_{0,r}$ are elements of $C_T C^{-1/2-\eta}(M)$, for all $\eta > 0$, that depend continuously on $\widehat{\xi}_r$ – see equation (2.4) in [57] (We deduce the regularity properties of the Z_i from the fact that \mathcal{L}^{-1} sends continuously $\mathcal{C}^\gamma([0, T] \times M)$ into $C_T C^{\gamma+2}(M)$ when $-2 < \gamma < 0$).

We now solve equation (2.4) with an arbitrary initial condition in $C^{-1/2-\varepsilon}(M)$ – [57] only considered the case of an initial condition that differs from $\mathfrak{I}_r(0)$ by an element of $C^{3/2-\varepsilon}(M)$. For that purpose, and for exponents $\alpha > 0, \beta \in \mathbb{R}$, we introduce the spaces $\langle \alpha, \beta \rangle$ made up of all functions $v \in C([0, T], C^\beta(M))$ such that

$$t^\alpha \|u(t)\|_{L^\infty} \xrightarrow[t \downarrow 0]{} 0$$

and

$$\|v\|_{\langle \alpha, \beta \rangle} \stackrel{\text{def}}{=} \max \left\{ \sup_{0 < t \leq T} t^\alpha \|v(t)\|_{C^\beta}, \sup_{0 \leq t \neq s \leq T} \frac{\|t^\alpha v(t) - s^\alpha v(s)\|_{L^\infty}}{|s - t|^{\beta/2}} \right\} < \infty.$$

(The use of such weighted spaces is suggested in [57]; we use here the same spaces as in Section 6 of Gubinelli & Perkowski's work [47].) The free propagation map

$$(\mathcal{F}a)(t) \stackrel{\text{def}}{=} e^{-tP} a$$

sends for instance $C^\beta(M)$ into $\langle \alpha, \beta + 2\alpha \rangle$, for all $\beta \in \{\mathbb{R} \setminus \mathbb{N}\}$ and $\alpha > 0$, and one has for all $0 \leq \delta < \min(\beta, 2\alpha)$

$$(2.5) \quad \|\mathcal{L}^{-1}(f)\|_{\langle \alpha - \delta/2, \beta - \delta \rangle} \lesssim \|f\|_{\langle \alpha, \beta - 2 \rangle}$$

This inequality allows to trade some explosion rate against some regularity. We also have for the same range of exponents and all $f \in \langle \alpha, \beta - 2 \rangle$

$$(2.6) \quad \|f\|_{\langle \alpha - \delta/2, \beta - \delta \rangle} \lesssim \|f\|_{\langle \alpha, \beta \rangle}.$$

These statements correspond in our setting to Lemma 6.6 and Lemma 6.8 in Gubinelli & Perkowski's work [47] – a proof is given in our companion work [10]. Note that since the different components of the enhanced noise are stationary they do not take value 0 at time 0. The initial condition for v_r is thus different from the initial condition for u_r . We keep the notation ϕ for the initial condition for u_r and write ϕ' for the initial condition for v_r . We will repeatedly use the estimate

$$(2.7) \quad \|fg\|_{C^{\alpha \wedge \beta}} \leq \|f\|_{C^\alpha} \|g\|_{C^\beta},$$

if $\alpha + \beta > 0$ which follows immediately from Proposition 43.

Proposition 3 – Pick $\varepsilon' = 4\varepsilon$ and set $\alpha_0 \stackrel{\text{def}}{=} 3/4 + (\varepsilon + \varepsilon')/2$. For any $\phi' \in C^{-1/2-\varepsilon}(M)$ there exists a positive time T^* such that for all $0 < T < T^*$ equation (2.4) has a unique solution

$$v_r \in C_T C^{-1/2-\varepsilon}(M) \cap \langle \alpha_0, 1 + \varepsilon' \rangle$$

with initial condition ϕ' . This solution depends continuously on $\widehat{\xi}_r$ and $\phi' \in C^{-1/2-\varepsilon}(M)$, and for any small positive λ these exist $T_\lambda \in (\lambda, T^*)$ such that $u \in C([\lambda, T_\lambda], C^{3/2-4\varepsilon}(M))$.

Proof – First, remark that $\lim_{t \downarrow 0} t^{\alpha_0} \mathcal{F}(\phi') = 0$, since $\|\mathcal{F}(\phi')\|_{L^\infty} \leq t^{-1/4-\varepsilon/2} \|\phi'\|_{-1/2-\varepsilon}$, so $\mathcal{F}(\phi') \in C_T C^{-1/2-\varepsilon}(M) \cap \langle \alpha_0, 1 + \varepsilon' \rangle$. We use a standard Picard iteration argument for the map

$$(2.8) \quad F(v) \stackrel{\text{def}}{=} \mathcal{F}(\phi') + \mathcal{L}^{-1} \left(-6\nabla\mathfrak{Y}_r^\circ \cdot \nabla v - e^{-6\mathfrak{Y}_r^\circ} v^3 + Z_{2,r}v^2 + Z_{1,r}v + Z_{0,r} \right).$$

Denote B_R the ball of radius $R = 4\|\phi'\|_{C^{-1/2-\varepsilon}}$ in $C_T C^{-1/2-\varepsilon}(M) \cap \langle \alpha_0, 1 + \varepsilon' \rangle$. Let $v_1, v_2 \in B_R$. Our first goal is to get a bound of the form

$$\|F(v_2) - F(v_1)\|_{\langle \alpha_0, 1 + \varepsilon' \rangle} \lesssim_{\widehat{\xi}_r} T^{\varepsilon/2} (R + R^2) \|v_2 - v_1\|_{\langle \alpha_0, 1 + \varepsilon' \rangle}.$$

meaning F is a contraction for the $(\alpha_0, 1 + \varepsilon')$ norm by choosing T small enough. We have

$$\begin{aligned} \|F(v_2) - F(v_1)\|_{(\alpha_0, 1 + \varepsilon')} &\leq \|\mathcal{L}^{-1}(6\nabla\mathcal{Y}_r \cdot \nabla(v_2 - v_1))\|_{(\alpha_0, 1 + \varepsilon')} + \|\mathcal{L}^{-1}(e^{-6\mathcal{Y}_r}(v_2^3 - v_1^3))\|_{(\alpha_0, 1 + \varepsilon')} \\ &\quad + \|\mathcal{L}^{-1}(Z_{2,r}(v_2^2 - v_1^2))\|_{(\alpha_0, 1 + \varepsilon')} + \|\mathcal{L}^{-1}(Z_{1,r}(v_2 - v_1))\|_{(\alpha_0, 1 + \varepsilon')}. \end{aligned}$$

Since $\nabla\mathcal{Y}_r \in C_T C^{-\eta}(M)$ for all $\eta > 0$, we first use the estimate (2.5) with $\delta = 0$ and (2.7) to get

$$\begin{aligned} \|\mathcal{L}^{-1}(\nabla\mathcal{Y}_r \cdot \nabla(v_2 - v_1))\|_{(\alpha_0, 1 + \varepsilon')} &\lesssim \sup_{0 < t \leq T} t^{\alpha_0} \|\nabla\mathcal{Y}_r \cdot \nabla(v_2 - v_1)\|_{C^{-\eta}} \\ &\lesssim_{\widehat{\xi}_r} \sup_{0 < t \leq T} t^{\alpha_0} \|\nabla(v_2 - v_1)\|_{C^{2\eta}} \\ &\lesssim T^{\alpha_0 - \alpha_1} \|v_2 - v_1\|_{(\alpha_1, 1 + 2\eta)}, \quad (\alpha_1 = 3/4 + (\varepsilon + 2\eta)/2) \\ &\lesssim T^{\varepsilon'/2 - \eta} \|v_2 - v_1\|_{(\alpha_1, 1 + 2\eta)}, \quad (\eta = \varepsilon'/4) \\ &\lesssim T^{\varepsilon'/4} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')}, \quad (\text{by (2.6) with } \delta = \varepsilon' - 2\eta). \end{aligned}$$

Now using again (2.5), (2.6), (2.7) and the fact that $\exp(-6\mathcal{Y}_r) \in C^{1-\eta}(M)$ for all $\eta > 0$, we have, for $\varepsilon' = 4\varepsilon$

$$\begin{aligned} \|\mathcal{L}^{-1}(\exp(-6\mathcal{Y}_r)(v_1^3 - v_2^3))\|_{(\alpha_0, 1 + \varepsilon')} &\lesssim_{\widehat{\xi}_r} \sup_{0 < t \leq T} t^{\alpha_0} \|(v_2 - v_1)(v_2^2 + v_1^2 + v_2 v_1)(t)\|_{C^\eta} \\ &\lesssim T^{\alpha_0 - 3\alpha'_0} \|v_2 - v_1\|_{(\alpha'_0, \eta)} (\|v_2\|_{(\alpha'_0, \eta)}^2 + \|v_1\|_{(\alpha'_0, \eta)}^2 + \|v_2\|_{(\alpha'_0, \eta)} \|v_1\|_{(\alpha'_0, \eta)}), \quad (\alpha'_0 = 1/4 + (\varepsilon + \eta)/2) \\ &\lesssim T^{(\varepsilon' - 2\varepsilon - 3\eta)/2} \|v_2 - v_1\|_{(\alpha'_0, \eta)} (\|v_2\|_{(\alpha'_0, \eta)}^2 + \|v_1\|_{(\alpha'_0, \eta)}^2 + \|v_2\|_{(\alpha'_0, \eta)} \|v_1\|_{(\alpha'_0, \eta)}) \\ &\lesssim T^{\varepsilon/2} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')} (\|v_2\|_{(\alpha_0, 1 + \varepsilon')} + \|v_1\|_{(\alpha_0, 1 + \varepsilon')} + \|v_2\|_{(\alpha_0, 1 + \varepsilon')} \|v_1\|_{(\alpha_0, 1 + \varepsilon')}), \end{aligned}$$

choosing $\eta = \varepsilon/3$ and using (2.6) in the last inequality. Next, with the same argument we have for $\varepsilon' = 4\varepsilon$, and setting $\alpha'_0 = \frac{1}{2} + (\varepsilon + \eta)/2$, $\alpha''_0 = \frac{1}{4} + (\varepsilon + \eta)/2$ in the third inequality

$$\begin{aligned} \|\mathcal{L}^{-1}(Z_{2,r}(v_2^2 - v_1^2))\|_{(\alpha_0, 1 + \varepsilon')} &\lesssim_{\widehat{\xi}_r} \sup_{0 < t \leq T} t^{\alpha_0} \|(v_2^2 - v_1^2)(t)\|_{C^{1/2+\eta}} \\ &\lesssim \sup_{0 < t \leq T} t^{\alpha_0} \|v_2(t) - v_1(t)\|_{C^{1/2+\eta}} (\|v_2(t)\|_\eta + \|v_1(t)\|_\eta) \\ &\leq T^{\alpha_0 - \alpha'_0 - \alpha''_0} \|v_2 - v_1\|_{(\alpha'_0, 1/2+\eta)} (\|v_2\|_{(\alpha''_0, \eta)} + \|v_1\|_{(\alpha''_0, \eta)}) \\ &\leq T^{(\varepsilon' - 2\varepsilon - 2\eta)/2} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')} (\|v_2\|_{(\alpha_0, 1 + \varepsilon')} + \|v_1\|_{(\alpha_0, 1 + \varepsilon')}) \\ &\leq T^{\varepsilon/2} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')} (\|v_2\|_{(\alpha_0, 1 + \varepsilon')} + \|v_1\|_{(\alpha_0, 1 + \varepsilon')}), \end{aligned}$$

using (2.6) in the fourth inequality and choosing $\eta = \varepsilon/2$ in the last inequality. Similarly, we get

$$\|\mathcal{L}^{-1}(Z_{1,r}(v_2 - v_1))\|_{(\alpha_0, 1 + \varepsilon')} \lesssim_{\widehat{\xi}_r} T^{\varepsilon/2} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')}.$$

Therefore

$$\|F(v_2) - F(v_1)\|_{(\alpha_0, 1 + \varepsilon')} \lesssim_{\widehat{\xi}_r} T^{\varepsilon/2} (R + R^2) \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')}.$$

The rest is to estimate $F(v_1) - F(v_2)$ in $C_T C^{-1/2-\varepsilon}$. Since $v_1, v_2 \in B_R$, for any $s \in [0, T]$, $i = 1, 2$ and $0 \leq \delta < 1 + \varepsilon'$, it follows from (2.6) that $\|v_i\|_{(\alpha_0 - \delta/2, 1 + \varepsilon' - \delta)} \leq R$, hence we have

$$(2.9) \quad \|v_i(s)\|_{C^{-1/2-\varepsilon}} \leq R \quad \text{and} \quad \|v_i\|_{C^{1+\varepsilon'-\delta}} \leq R s^{-(\alpha_0 - \delta/2)}.$$

Recall the following estimate which is used many times below: if $\|u(s)\|_{C^\beta} \lesssim s^{-\gamma}$, for $\gamma < 1$, then

$$(2.10) \quad \|\mathcal{L}^{-1}u\|_{C^\beta} \lesssim \int_0^T s^{-\gamma} \lesssim T^{1-\gamma}.$$

Using (2.9) and the fact that $\nabla\mathcal{Y}_r \in C_T C^{-\varepsilon'/4}(M)$, we have

$$\|\nabla\mathcal{Y}_r \cdot \nabla(v_2 - v_1)(s)\|_{C^{-1/2-\varepsilon}} \lesssim_{\widehat{\xi}_r} \|\nabla(v_2 - v_1)(s)\|_{C^{\varepsilon'}} \lesssim \|v_2 - v_1\|_{C^{1+\varepsilon'}} \lesssim s^{-\alpha_0} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')},$$

hence by (2.10)

$$\|\mathcal{L}^{-1}(\nabla\mathcal{Y}_r \cdot \nabla(v_2 - v_1))\|_{C^{-1/2-\varepsilon}} \lesssim_{\widehat{\xi}_r} T^{1-\alpha_0} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')} = T^{1/4 - \varepsilon/2 - \varepsilon'/2} \|v_2 - v_1\|_{(\alpha_0, 1 + \varepsilon')}.$$

Again, by (2.9) and the fact that $\mathfrak{Y}_r \in C_T C^{1-\eta}(M)$, and $\|uv\|_{C^\beta} \leq \|u\|_{L^\infty} \|v\|_{C^\beta} + \|v\|_{L^\infty} \|u\|_{C^\beta}$, for $\beta \in (0, 1)$,

$$\begin{aligned} \|(\exp(-6\mathfrak{Y}_r)(v_1^3 - v_2^3))(s)\|_{-1/2-\varepsilon} &\lesssim_{\widehat{\xi}_r} \|(v_2 - v_1)(v_2^2 + v_1^2 + v_2v_1)(t)\|_{C^{-1/2-\varepsilon}} \\ &\lesssim \|v_2 - v_1\|_{C^{-1/2-\varepsilon}} \|v_2^2 + v_1^2 + v_2v_1\|_{C^{1/2+2\varepsilon}} \\ &\lesssim \|v_2 - v_1\|_{C^{-1/2-\varepsilon}} \sum_{i,j=1}^2 \|v_i\|_{C^\varepsilon} \|v_j\|_{C^{1/2+2\varepsilon}} \\ &\lesssim \|v_2 - v_1\|_{C^{-1/2-\varepsilon}} s^{-3/4-5\varepsilon/2} R^2 \end{aligned}$$

where we use (2.9) for both $\|v_i\|_{C^\varepsilon}$ and $\|v_j\|_{C^{1/2+2\varepsilon}}$ in the last inequality. Hence by (2.10) we have

$$\|\mathcal{L}^{-1}((\exp(-6\mathfrak{Y}_r)(v_1^3 - v_2^3))\|_{-1/2-\varepsilon} \lesssim_{\widehat{\xi}_r} T^{1/4-5\varepsilon/2} R^2 \|v_2 - v_1\|_{C^{-1/2-\varepsilon}}.$$

Now, with the same argument and (2.6), for $\eta > \varepsilon$, we have

$$\begin{aligned} \|(Z_{2,r}(v_2^2 - v_1^2))(s)\|_{C^{-1/2-\varepsilon}} &\lesssim_{\widehat{\xi}_r} \|(v_2^2 - v_1^2)(s)\|_{C^{1/2+\eta}} \\ &\lesssim (\|v_1\|_{C^\eta} + \|v_2\|_{C^\eta}) \|v_1 - v_2\|_{1/2+\eta} \\ &\lesssim s^{-(\alpha_0 - (1+\varepsilon' - \eta)/2)} R s^{-1/2 - \varepsilon/2 - \eta/2} \|v_1 - v_2\|_{(\frac{1}{2} + \frac{\varepsilon+\eta}{2}, \frac{1}{2} + \eta)} \\ &\lesssim s^{-(3/4 + \varepsilon + \eta)} R \|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')}, \end{aligned}$$

hence, choosing $\eta = 2\varepsilon$ yields

$$\|\mathcal{L}^{-1}(Z_{2,r}(v_2^2 - v_1^2))\|_{-1/2-\varepsilon} \lesssim_{\widehat{\xi}_r} T^{1/4-3\varepsilon} R \|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')}.$$

Similarly, we get

$$\|\mathcal{L}^{-1}(Z_{1,r}(v_2 - v_1))\|_{C^{-1/2-\varepsilon}} \lesssim_{\widehat{\xi}_r} T^{1/2-3\varepsilon/2} \|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')}.$$

Therefore

$$\begin{aligned} \|F(v_2) - F(v_1)\|_{C^{-1/2-\varepsilon}} &\lesssim_{\widehat{\xi}_r} T^{1/4-5\varepsilon/2} R^2 \|v_2 - v_1\|_{C^{-1/2-\varepsilon}} \\ &\quad + (T^{1/4-\varepsilon/2-\varepsilon'/2} + T^{1/4-3\varepsilon} R + T^{1/2-3\varepsilon/2}) \|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')}. \end{aligned}$$

Combining the estimates above we infer that for $T > 0$ sufficiently small, depending on $\|\phi'\|_{C^{-1/2-\varepsilon}}$, $\widehat{\xi}_r$, the map F is a contraction on the ball of radius $4\|\phi'\|_{C^{-1/2-\varepsilon}}$ in $C_T C^{-1/2-\varepsilon}(M) \cap (\alpha_0, 1 + \varepsilon')$. The unique fixed point is our solution on $[0, T]$. Taking the supremum of all such T gives the maximal existence time T^* . Once we know that v takes values in $(\alpha_0, 1 + \varepsilon')$ we can restart the fixed point procedure from a positive time, with an initial condition that is now of Hölder regularity $(1 + \varepsilon')$. It is elementary to adapt the preceding estimates to see that now the solution will take values in $C^{3/2-4\varepsilon}(M)$.

For the continuous dependence on $\widehat{\xi}_r$ and the initial data, we define

$$F(\widehat{\xi}_r, \phi, v) \stackrel{\text{def}}{=} e^{-tP}(\phi) + \mathcal{L}^{-1}\left(-6\nabla\mathfrak{Y}_r(\widehat{\xi}_r) \cdot \nabla v - e^{-6\mathfrak{Y}_r(\widehat{\xi}_r)} v^3 + Z_{2,r}(\widehat{\xi}_r) v^2 + Z_{1,r}(\widehat{\xi}_r) v + Z_{0,r}(\widehat{\xi}_r)\right).$$

Let $K > 0$ be a uniform constant satisfying

$$(2.11) \quad \|e^{-tP}\phi\|_{C^{-1/2-\varepsilon}} \leq K\|\phi\|_{C^{-1/2-\varepsilon}} \quad \text{and} \quad \|e^{-tP}\phi\|_{(\alpha_0, 1 + \varepsilon')} \leq K\|\phi\|_{C^{-1/2-\varepsilon}}.$$

Take the ball B_R in $C^{-1/2-\varepsilon}(M)$. Since F depends linearly on $\widehat{\xi}_r$ and $\exp(-6\mathfrak{Y}_r)$, by the same arguments above, for any $\phi \in B_R$ and we can choose $T = T(R, \widehat{\xi}_r, \widehat{\xi}'_r)$ small enough such that $C(T) < 1/2$ and

$$\|F(\widehat{\xi}_r, \phi, v_1) - F(\widehat{\xi}'_r, \phi, v_2)\|_{C_T C^{-1/2-\varepsilon}} \leq C(T) (\|v_1 - v_2\|_{C_T C^{-1/2-\varepsilon}} + \|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')} + \|\widehat{\xi}_r - \widehat{\xi}'_r\|)$$

and

$$\|F(\widehat{\xi}_r, \phi, v_1) - F(\widehat{\xi}'_r, \phi, v_2)\|_{(\alpha_0, 1 + \varepsilon')} \leq C(T) (\|v_1 - v_2\|_{(\alpha_0, 1 + \varepsilon')} + \|\widehat{\xi}_r - \widehat{\xi}'_r\|).$$

Now for $\phi_1, \phi_2 \in B_R$ we have

$$\begin{aligned}
& \|v_r(\widehat{\xi}_r, \phi_1) - v_r(\widehat{\xi}_r, \phi_2)\|_{C_T C^{1/2-\varepsilon}} = \|F(\widehat{\xi}_r, \phi_1, v_r(\phi_1)) - F(\widehat{\xi}_r, \phi_2, v_r(\phi_2))\|_{C_T C^{1/2-\varepsilon}} \\
& \leq \|F(\widehat{\xi}_r, \phi_1, v_r(\phi_1)) - F(\widehat{\xi}_r, \phi_2, v_r(\phi_1))\|_{C_T C^{1/2-\varepsilon}} + \|F(\widehat{\xi}_r, \phi_2, v_r(\phi_1)) - F(\widehat{\xi}_r, \phi_2, v_r(\phi_2))\|_{C_T C^{1/2-\varepsilon}} \\
& \leq K\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}} + C(T)\left(\|v_r(\phi_1) - v_r(\phi_2)\|_{C_T C^{1/2-\varepsilon}} + \|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} + \|\widehat{\xi}_r - \widehat{\xi}_r'\|\right).
\end{aligned}$$

Similarly we have

$$\begin{aligned}
\|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} & = \|F(\widehat{\xi}_r, \phi_1, v_r(\phi_1)) - F(\widehat{\xi}_r, \phi_2, v_r(\phi_2))\|_{(\alpha_0, 1+\varepsilon')} \\
& \leq K\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}} + C(T)\left(\|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} + \|\widehat{\xi}_r - \widehat{\xi}_r'\|\right),
\end{aligned}$$

so

$$\begin{aligned}
\|v_r(\phi_1) - v_r(\phi_2)\|_{C_T C^{1/2-\varepsilon}} + \|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} & \\
& \leq 2K\|\phi_1 - \phi_2\|_{C_T C^{1/2-\varepsilon}} + C(T)\|v_r(\phi_1) - v_r(\phi_2)\|_{C_T C^{-1/2-\varepsilon}} \\
& \quad + 2C(T)\|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} + 2\|\widehat{\xi}_r - \widehat{\xi}_r'\|,
\end{aligned}$$

and we read on the estimate

$$\begin{aligned}
& \|v_r(\phi_1) - v_r(\phi_2)\|_{C_T C^{1/2-\varepsilon}} + \|v_r(\phi_1) - v_r(\phi_2)\|_{(\alpha_0, 1+\varepsilon')} \\
& \leq \frac{1}{1 - 2C(T)} \left(2K\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}} + 2C(T)\|\widehat{\xi}_r - \widehat{\xi}_r'\| \right)
\end{aligned}$$

the continuous dependence of the solution on $\widehat{\xi}$ and the initial data. \triangleright

We see from the proof that T^* depends only on $\widehat{\xi}_r$ and $\phi' \in C^{-1/2-\varepsilon}(M)$. The following additional piece of information will be useful when proving the coming down from infinity property by energy methods in the next section.

Lemma 4 – Let $0 < t_0 < t_1$. For $\beta = 3/2 - \varepsilon$ and any $\kappa \leq \beta/2$, then $t \mapsto v_r(t, x)$ is κ -Hölder continuous as a function from $[t_0, t_1]$ to L^∞ .

Proof – By the change of variable $t \mapsto t - t_0$, we can assume $t_0 = 0, t_1 = T > 0$ and $v_r \in C_T C^{3/2-\varepsilon}(M)$. We now show the Hölder regularity of v_r at time 0, the adaptation to arbitrary times is straightforward. We have

$$v_r(t, \cdot) = e^{-tP} v_r(0) + \int_0^t e^{-(t-s)P} \left(-6\nabla_Y^{\varphi_r} \cdot \nabla v_r - e^{-6\varphi_r} v_r^3 + Z_{2,r} v_r^2 + Z_{1,r} v_r + Z_{0,r} \right) (s) ds.$$

We first remark that

$$(2.12) \quad |v_r(t, \cdot) - v_r(0)| \leq |e^{-tP} v_r(0) - v_r(0)| + \left| \int_0^t e^{-(t-s)P} (f_r(s)) ds \right|,$$

where $f_r = -6\nabla_Y^{\varphi_r} \cdot \nabla v_r - e^{-6\varphi_r} v_r^3 + Z_{2,r} v_r^2 + Z_{1,r} v_r + Z_{0,r}$. It follows from the time regularity of the heat flow that $\|(1 - e^{-tP})h\|_{L^\infty} \lesssim t^{\frac{\beta}{2}} \|h\|_{C^\beta}$ for $0 \leq \beta \leq 2$, hence

$$(2.13) \quad \|e^{-tP} v_r(0) - v_r(0)\|_{L^\infty} \lesssim t^{\beta/2} \|v_r(0)\|_{C^\beta}.$$

Since $v_r \in C_T C^\beta$, with $\beta = 3/2 - \varepsilon$, $\nabla_Y^{\varphi_r} \in C_T C^{-\varepsilon}$, $e^{-6\varphi_r} \in C_T C^{1-\varepsilon}$, $\mathbf{Z}_r = \{Z_{0,r}, Z_{1,r}, Z_{2,r}\} \subset C_T C^{-\alpha}$ with $\alpha = 1/2 + \varepsilon'$, using $\|gh\|_{C^{\alpha' \wedge \beta'}} \lesssim \|g\|_{C^{\alpha'}} \|h\|_{C^{\beta'}}$ for $\alpha' + \beta' > 0$ we have

$$\|f_r(s, \cdot)\|_{C^{-\alpha}} \lesssim \mathbf{Z}_r \|v_r\|_{C_T C^\beta} \lesssim 1.$$

Then the estimate $\|e^{-tP} h\|_{C^{\eta+\gamma}} \lesssim t^{-\gamma/2} \|h\|_{C^\eta}$, for $\gamma \geq 0$ implies

$$\left\| \int_0^t e^{-(t-s)P} (f_r(s)) ds \right\|_{C^{\varepsilon'}} \lesssim \int_0^t (t-s)^{-(\alpha-\varepsilon')/2} \|f_r(s)\|_{C^{-\alpha}} ds \lesssim t^{1-\alpha/2-\varepsilon'/2}.$$

By choosing $\varepsilon' \geq \varepsilon/2$ we have $\kappa \stackrel{\text{def}}{=} 1 - \alpha/2 - \varepsilon'/2 \leq \beta/2$, therefore u is κ -Hölder from $[0, T]$ to L^∞ . \triangleright

As in Proposition 6.8 of Mourrat & Weber's work [67] it follows from this property that the function

$$t \in (0, T] \mapsto \|v_r(t)\|_{L^p}^p$$

satisfies the equation

$$(2.14) \quad \frac{1}{p} (\|v_r(t)\|_{L^p}^p - \|v_r(s)\|_{L^p}^p) = \int_s^t (v_r^{p-1}, \Delta v_r) - \int_s^t \left(\int_M e^{-6\mathcal{Y}_r^\circ(s_1)} v_r^{p+2}(s_1) \right) ds_1 \\ + \int_s^t \left(\frac{1}{p} (-6\nabla\mathcal{Y}_r^\circ, \nabla v_r^p) + (Z_2, v_r^{p+1}) + (Z_1, v_r^p) + (Z_0, v_r^{p-1}) \right).$$

2.2 Long time existence and coming down from infinity

We show in this section that the superlinear attractive drift $-\exp(-6\mathcal{Y}_r^\circ) v_r^3$ in equation (2.4) entails an a priori bound on the $L^p(M)$ norm of the solution away from the initial time that is independent of the initial condition. This bound entails the long time existence of the solution v_r to (2.4) and is the key to proving the existence of an invariant probability measure for the dynamics (1.3) via a compactness argument. This point will be developed in Section 6.

We rewrite equation (2.4) in the form

$$(2.15) \quad (\partial_t + P + B_r \nabla) v_r = -A_r v_r^3 + Z_{2,r} v_r^2 + Z_{1,r} v_r + Z_{0,r},$$

with

$$B_r \stackrel{\text{def}}{=} 6\nabla\mathcal{Y}_r^\circ \in C_T C^{-\eta}(M), \quad A_r \stackrel{\text{def}}{=} e^{-6\mathcal{Y}_r^\circ} \in C_T C^{1-\eta}(M),$$

and $Z_{i,r} \in C_T C^{-1/2-\eta}(M)$, for all $\eta > 0$.

Theorem 5 – *The solution $v_r(t) \in C^{3/2-\varepsilon}(M)$ exists for all times $t > 0$. Pick an even integer $p \geq 8$. There is a random variable $C(p, \widehat{\xi}_r|_{[0,t]})$ that depends only on the restriction to the interval $[0, t]$ of $\widehat{\xi}_r$ such that one has*

$$(2.16) \quad \|v_r(t)\|_{L^p(M)} \leq C(p, \widehat{\xi}_r|_{[0,t]}) \max \left\{ \frac{1}{\sqrt{t}}, 1 \right\}$$

for all $t > 0$, independently of the initial condition $\phi' \in C^{-1/2-\varepsilon}(M)$.

The upper bound in (2.16) is in particular independent of the initial condition in (2.15); this phenomenon is called *coming down from infinity*. We note for later use that keeping track of the implicit constants in the computations below gives an upper in (2.16) takes for $1 \leq t \leq 2$ the form

$$(2.17) \quad (1 + \|\widehat{\xi}_r\|)^\gamma \left(\exp(\gamma \|\mathcal{Y}_r^\circ\|_{L^\infty([0,2] \times M)}) + 1 \right)$$

for some positive constants $\gamma = \gamma(p)$, $\gamma' = \gamma'(p)$, up to a multiplicative constant. We denoted here by $\|\widehat{\xi}_r\|$ the norm of $\widehat{\xi}_r$ seen as an element of the product space where $\widehat{\xi}_r$ takes its values. We use a priori energy estimates to prove Theorem 5, following the strategy initiated by Mourrat & Weber in their proof of a similar result in [68], Theorem 7.1 therein. Gubinelli & Hofmanová also used energy estimates in their work [45] on the Φ_3^4 measure on \mathbb{R}^3 . See also the proof of Proposition 3.7 in the work [78] of Tsatsoulis & Weber for an implementation of that strategy in the 2-dimensional torus.

We will use in the remainder of this section the shorthand notation

$$B_p^\gamma(M) \stackrel{\text{def}}{=} B_{p,\infty}^\gamma(M)$$

for any $\gamma \in \mathbb{R}$ and $1 \leq p \leq \infty$. Set

$$F_r(t) \stackrel{\text{def}}{=} \|v_r(t)\|_{p+2}^p + \|v_r(t)\|_{B^{\frac{1+2\varepsilon}{p+2}}}^{\frac{p}{3}}.$$

We prove below that one has for all $0 < T_0 \leq s < t \leq T < T^* \wedge 1$

$$(2.18) \quad \|v_r(t)\|_{L^p}^p + \int_s^t F_r(s_1)^{\frac{p+2}{p}} ds_1 \lesssim_{\widehat{\xi}_r} 1 + F_r(s).$$

This inequality shows that, all $0 < T_0 \leq s < t \leq T$

$$(2.19) \quad \int_s^t F_r(s_1)^{\frac{p+2}{p}} ds_1 \lesssim_{\widehat{\xi}_r} 1 + F_r(s).$$

It then follows from a modified version of Mourrat & Weber's comparison test recalled in Proposition 44 of Appendix A that there is an integer $N \geq 1$ and sequence of times $T_0 = t_0 < t_1 < t_2 < \dots < t_N = T$ such that for all $n \in \{0, \dots, N-1\}$,

$$F_r(t_n) \lesssim_{\hat{\xi}_r} 1 + t_{n+1}^{-\frac{p}{2}},$$

for an implicit constant that does not depend on T_0, T . Pick $t \in [T_0, T]$. There exists $n \in \{0, \dots, N-1\}$ such that $t \in [t_n, t_{n+1}]$. Moreover, by (2.18) with $s = t_n$, we have

$$\|v_r(t)\|_{L^p}^p \lesssim_{\hat{\xi}_r} 1 + F_r(t_n) \lesssim_{\hat{\xi}_r} 1 + t_{n+1}^{-\frac{p}{2}} \lesssim_{\hat{\xi}_r} 1 + t^{-\frac{p}{2}}.$$

This bound holds for T_0 arbitrarily small and $T = 1$, and can be repeated on $[1, 2]$, etc, so that the uniform estimate (2.16) follows. Recall $p > 6$ so the space $L^p(M)$ is continuously embedded into the space $C^{-1/2-\varepsilon}(M)$. Given that T^* depends only on the restriction to $[0, T^*]$ of $\hat{\xi}_r$ and the initial condition $\phi' \in C^{-1/2-\varepsilon}(M)$ the uniform estimate (2.16) and the continuous injection of $L^p(M)$ into $C^{-1/2-\varepsilon}(M)$ imply we can extend the solution though $T^* \wedge 1$, hence $T^* > 1$. Then we can repeat the same argument on the interval $[1, 2]$ and so on to get the long time existence of v_r .

On a technical level, our proof of Theorem 5 will only use the fractional Leibniz rule from Proposition 45 and the elementary interpolation result from Proposition 46, both recalled in Appendix A. Last, recall Young inequality that gives the existence for any positive δ of a constant C_δ such that one has

$$ab \leq \delta a^{p'} + \delta^{-\frac{q'}{p'}} b^{q'},$$

for all positive a, b and exponent $1 < p' < \infty$ with conjugate exponent q' . The proof of (2.18) requires two intermediate results stated as lemmas.

Lemma 6 – *For every $0 < s < t \leq T$, we have*

$$(2.20) \quad \|v_r(t)\|_{L^p}^p + \int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \lesssim_{\hat{\xi}_r} 1 + \|v_r(s)\|_{L^{p+2}}^p + \int_s^t \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}} ds_1.$$

Proof – Pairing the equation with v^{p-1} with respect to the L^2 scalar product yields identity (2.14). As A is positive and bounded below and $(v^{p-1}, -\Delta v)$ is positive, since p is an even integer greater than 4, we obtain

$$(2.21) \quad \|v_r(t)\|_{L^p}^p + \int_s^t \|v_r\|_{L^{p+2}}^{p+2} \lesssim_{\hat{\xi}_r} \|v_r(s)\|_{L^p}^p + \int_s^t (B_r, \nabla v_r^p) + (Z_{2,r}, v_r^{p+1}) + (Z_{1,r}, v_r^p) + (Z_{0,r}, v_r^{p-1}).$$

where the implicit constant is $p \left(\exp(6\|\mathfrak{Y}_r\|_{L^\infty([0,2] \times M)}) + 1 \right)$.

We bound the different terms in the right hand side of (2.21). Recall that since B_r is an element of $B_{\infty, \infty}^{-\varepsilon'}(M)$ for all $\varepsilon' > 0$ it is an element of $B_{1, \infty}^{-\varepsilon}(M)$. By the fractional Leibniz rule from Proposition 45 and Young inequality we have for $|(B_r, \nabla v_r^p)|$, up to a $\hat{\xi}_r$ -dependent multiplicative constant, the upper bound

$$\|\nabla v_r^p\|_{B_1^\varepsilon} \lesssim \|v_r^p\|_{B_1^{1+\varepsilon}} \lesssim \|v_r^{p-1}\|_{L^{\frac{p+2}{p-1}}} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}} \lesssim \|v_r\|_{L^{p+2}}^{p-1} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}} \lesssim \delta \|v_r\|_{L^{p+2}}^{p+2} + \delta^{-\frac{p-1}{3}} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}},$$

where $\delta > 0$ is arbitrarily small. For the other terms, we have first

$$|(Z_{2,r}, v_r^{p+1})| \lesssim_{\hat{\xi}_r} \|v_r^{p+1}\|_{B_1^{\frac{1+\varepsilon}{2}}} \lesssim \|v_r^p\|_{L^{\frac{p+2}{p}}} \|v_r\|_{B_{\frac{p+2}{2}}^{\frac{1+\varepsilon}{2}}} \lesssim \|v_r\|_{L^{p+2}}^p \|v_r\|_{B_{\frac{p+2}{2}}^{\frac{1+\varepsilon}{2}}}.$$

Here we interpolate the last term to obtain

$$\|v_r\|_{B_{\frac{p+2}{2}}^{\frac{1+\varepsilon}{2}}} \lesssim \|v_r\|_{L^{p+2}}^{\frac{1}{2}} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{1}{2}},$$

and we deduce that

$$|(Z_{2,r}, v_r^{p+1})| \lesssim_{\hat{\xi}_r} \|v_r\|_{L^{p+2}}^{p+\frac{1}{2}} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{1}{2}} \leq \delta \|v_r\|_{L^{p+2}}^{p+2} + C_\delta \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}},$$

using Young inequality in the second inequality, here $C_\delta = \delta^{-\frac{2p+1}{6}}$.

Similar estimates hold for the $Z_{1,r}$ and $Z_{0,r}$ terms. We have

$$\begin{aligned} |(Z_{1,r}, v_r^p)| &\lesssim_{\widehat{\xi}_r} \|v_r^p\|_{B_1^{1+\varepsilon}} \lesssim \|v_r^{p-1}\|_{L^{\frac{p+2}{p-1}}} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}} \lesssim \|v_r\|_{L^{p+2}}^{p-1} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}} \\ &\lesssim \delta \|v_r\|_{L^{p+2}}^{p+2} + \delta^{-\frac{p-1}{3}} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}}. \end{aligned}$$

and

$$\begin{aligned} |(Z_{0,r}, v_r^{p-1})| &\lesssim_{\widehat{\xi}_r} \|v_r^{p-1}\|_{B_1^{1+\varepsilon}} \lesssim \|v_r^{p-2}\|_{L^{\frac{p+2}{p-2}}} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{4}}}} \lesssim \|v_r\|_{L^{p+2}}^{p-2} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}} \\ &\lesssim \delta \|v_r\|_{L^{p+2}}^{\frac{(p+2)(p-2)}{p-1}} + \delta^{-\frac{p-1}{3}} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}} \lesssim 1 + \delta^{\frac{p-1}{p-2}} \|v_r\|_{L^{p+2}}^{p+2} + \delta^{-\frac{p-1}{3}} \|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}}. \end{aligned}$$

One can then absorb the δ terms of these upper bounds in the corresponding L^{p+2} term in the left hand side of (2.21) to get the result by integrating in time on the interval (s, t) . Since we choose $\delta \lesssim (1 + \|\widehat{\xi}_r\|)^{-1}$, then $C_\delta, \delta^{-\frac{p-1}{3}} \gtrsim (1 + \|\widehat{\xi}_r\|)^\gamma$ for some $\gamma > 0$ depending on p . Combining with the implicit constant in (2.21) we obtain that the implicit constant in (2.20) is of form

$$(1 + \|\widehat{\xi}_r\|)^\gamma \left(\exp(\gamma \|\mathcal{Y}_r\|_{L^\infty([0,2] \times M)}) + 1 \right)$$

for some $\gamma, \gamma' > 0$ depending on p . The implicit constants in the next steps will be obtained in the same way. \triangleright

Lemma 7 – For $0 \leq s < t < T < T^* \wedge 1$ we have

$$(2.22) \quad \int_s^t \|v_r(s_1)\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}} ds_1 \lesssim_{\widehat{\xi}_r} 1 + F_r(s) + \int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1.$$

Note that we have a $B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}$ norm involved in (2.20):

$$\|v_r(t)\|_{L^p}^p + \int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \lesssim \|v_r(s)\|_{L^{p+2}}^p + \int_s^t \|v_r(s_1)\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}} ds_1$$

while we estimate in (2.22) a stronger $B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}$ norm. We postpone for a second the proof of Proposition 7 and explain now how we get the estimate (2.18)

$$\|v_r(t)\|_{L^p}^p + \int_s^t F_r(s_1)^{\frac{p+2}{p}} ds_1 \lesssim_{\widehat{\xi}_r} 1 + F_r(s)$$

from (2.20) and (2.22). Since

$$F_r(s_1)^{\frac{p+2}{p}} \lesssim \|v_r(s_1)\|_{L^{p+2}}^{p+2} + \|v_r(s_1)\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\frac{p+2}{3}}$$

we see as a consequence of (2.22) that one gets (2.18) if one proves that

$$(2.23) \quad \|v_r(t)\|_{L^p}^p + \int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \lesssim_{\widehat{\xi}_r} 1 + F_r(s).$$

We start from the inequality (2.20) and use the interpolation estimate

$$\|v_r\|_{B^{\frac{1+\varepsilon}{\frac{p+2}{3}}}} \lesssim \|v_r\|_{L^{\frac{p+2}{3}}}^{1-\theta_\varepsilon} \|v_r\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\theta_\varepsilon} \leq \delta \|v_r\|_{L^{\frac{p+2}{3}}}^3 + C_\delta \|v_r\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\sigma_\varepsilon} \leq \delta \|v_r\|_{L^{p+2}}^3 + C_\delta \|v_r\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\sigma_\varepsilon},$$

with

$$\theta_\varepsilon \stackrel{\text{def}}{=} \frac{1+\varepsilon}{1+2\varepsilon} < 1, \quad \sigma_\varepsilon \stackrel{\text{def}}{=} \frac{3\theta_\varepsilon}{2+\theta_\varepsilon} < 1.$$

We feed this upper bound inside (2.20); the contribution of the small factor involving the L^{p+2} norm of v_r can be absorbed in the corresponding term in the left hand side of (2.20), so we have

$$(2.24) \quad \|v_r(t)\|_{L^p}^p + \int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \lesssim \|v_r(s)\|_{L^{p+2}}^p + \int_s^t \|v_r(s_1)\|_{B^{\frac{1+2\varepsilon}{\frac{p+2}{3}}}}^{\sigma_\varepsilon \frac{p+2}{3}} ds_1.$$

We use Young inequality once more to bound

$$\|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\sigma_\varepsilon} \lesssim \delta \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} + C_\delta.$$

By choosing δ small enough we can absorb the L^{p+2} term that comes from (2.22) in the left hand side of (2.24) and use that $\|v_r(s)\|_{L^{p+2}}^p \leq F_r(s)$ to get (2.23) from (2.24).

Proof of Lemma 7 – We proceed in two steps.

Step 1. We first prove that one has

$$(2.25) \quad \|v_r(t)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} \lesssim_{\widehat{\xi}_r} \|e^{-(t-s)P} v_r(s)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} + \left(\int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \right)^{\frac{3}{p+2}} + \left(\int_s^t \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{p+2} ds_1 \right)^{\frac{3}{p+2}}.$$

We look at each term in the expression for $v_r(t) - e^{-(t-s)P} v_r(s)$

$$\int_s^t e^{-(t-s_1)P} \left(-A_r(s_1)v_r(s_1)^3 + B_r(s_1)\nabla v_r(s_1) + Z_{2,r}(s_1)v_r(s_1)^2 + Z_{1,r}(s_1)v_r(s_1) + Z_{0,r}(s_1) \right) ds_1.$$

One has

$$\begin{aligned} \left\| \int_s^t e^{-(t-s_1)P} (A_r(s_1)v_r(s_1)^3) ds_1 \right\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} &\lesssim \int_s^t \|e^{-(t-s_1)P} (A_r(s_1)v_r(s_1)^3)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} ds_1 \\ &\lesssim \int_s^t (t-s_1)^{-\frac{1+2\varepsilon}{2}} \|A_r(s_1)v_r(s_1)^3\|_{L^{\frac{p+2}{3}}} ds_1 \\ &\lesssim_{\widehat{\xi}_r} \int_s^t (t-s_1)^{-\frac{1+2\varepsilon}{2}} \|v_r(s_1)\|_{L^{p+2}}^3 ds_1 \\ &\lesssim \left(\int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} \right)^{\frac{3}{p+2}}, \end{aligned}$$

where we used Hölder inequality, the integrability in time of $(t-s_1)^{-\frac{(1+2\varepsilon)(p+2)}{2(p-1)}}$ and the fact that $s \leq s_1 \leq t < T < 1$. Similarly, we have

$$\begin{aligned} \left\| \int_s^t e^{-(t-s_1)P} (B_r(s_1)\nabla v_r(s_1)) ds_1 \right\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} &\lesssim \int_s^t \|e^{-(t-s_1)P} (B_r(s_1)\nabla v_r(s_1))\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} ds_1 \\ &\lesssim \int_s^t (t-s_1)^{-\frac{1+3\varepsilon}{2}} \|B_r(s_1)\nabla v_r(s_1)\|_{B_{\frac{p+2}{3}}^{-\varepsilon}} ds_1 \\ &\lesssim_{\widehat{\xi}_r} \int_s^t (t-s_1)^{-\frac{1+3\varepsilon}{2}} \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}} ds_1 \\ &\lesssim \left(\int_s^t \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{p+2} \right)^{\frac{3}{p+2}}. \end{aligned}$$

Next we have

$$\begin{aligned} \left\| \int_s^t e^{-(t-s_1)P} (Z_{2,r}(s_1)v_r(s_1)^2) ds_1 \right\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} &\lesssim \int_s^t \|e^{-(t-s_1)P} (Z_{2,r}(s_1)v_r(s_1)^2)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} ds_1 \\ &\lesssim \int_s^t (t-s_1)^{-\frac{1+2\varepsilon+\frac{1+\varepsilon}{2}}{2}} \|Z_{2,r}(s_1)v_r(s_1)^2\|_{B_{\frac{p+2}{3}}^{-\frac{1+\varepsilon}{2}}} ds_1. \end{aligned}$$

Using the interpolation result from Proposition 46 and Young inequality we have

$$\|Z_{2,r}v_r^2\|_{B_{\frac{p+2}{3}}^{-\frac{1+\varepsilon}{2}}} \lesssim_{\widehat{\xi}_r} \|v_r^2\|_{B_{\frac{p+2}{3}}^{\frac{1+\varepsilon}{2}}} \lesssim \|v_r\|_{L^{p+2}} \|v_r\|_{B_{\frac{p+2}{3}}^{\frac{1+\varepsilon}{2}}} \lesssim \|v_r\|_{L^{p+2}}^{\frac{3}{2}} \|v_r\|_{B_{\frac{p+2}{3}}^{\frac{1+\varepsilon}{2}}}^{\frac{1}{2}} \lesssim \|v_r\|_{L^{p+2}}^{p+2} + \|v_r\|_{B_{\frac{p+2}{3}}^{\frac{1+\varepsilon}{2}}}^{\frac{p+2}{3}},$$

and the desired estimate follows as in the previous terms. One proceeds in exactly the same way to prove similar estimates on the $Z_{1,r}$ and $Z_{0,r}$ terms. We leave the details to the interested reader.

Step 2. We first rewrite (2.25) at $t = s_1$ and rise this inequality to the power $\frac{p+2}{3}$. It yields the upper bound

$$\begin{aligned} \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} &\lesssim_{\widehat{\xi}_r} \|e^{-(s_1-s)P} v_r(s)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} + \int_s^{s_1} \|v_r\|_{L^{p+2}}^{p+2} + \int_s^{s_1} \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}} \\ &\lesssim_{\widehat{\xi}_r} \|e^{-(s_1-s)P} v_r(s)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} + \int_s^t \|v_r\|_{L^{p+2}}^{p+2} + \int_s^t \|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}}, \end{aligned}$$

which holds for $t \geq s_1$, using the fact that $\int_s^{s_1} \|v_r\|^\alpha$ is increasing in s_1 whatever the exponent α the norm on v_r . Integrating on $s_1 \in [s, t]$, we obtain

$$(2.26) \quad \begin{aligned} \left(\int_s^t \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} ds_1 \right)^{\frac{3}{p+2}} &\lesssim_{\widehat{\xi}_r} \left(\int_s^t \|e^{-(s_1-s)P} v_r(s)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p+2}{3}} ds_1 \right)^{\frac{3}{p+2}} \\ &\quad + (t-s)^{\frac{3}{p+2}} \left[\left(\int_s^t \|v_r(s_1)\|_{L^{p+2}}^{p+2} ds_1 \right)^{\frac{3}{p+2}} + \left(\int_s^t \|v_r(s_1)\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}}^{\frac{p+2}{3}} ds_1 \right)^{\frac{3}{p+2}} \right], \end{aligned}$$

as the function $[\dots]$ after $(t-s)^{\frac{3}{p+2}}$ in the right hand side is increasing. We bound the first term in (2.26) by $\|v_r(s)\|_{B_{\frac{p+2}{3}}^{(1+2\varepsilon)(1-\frac{3}{p+2})}}^{\frac{p+2}{3}}$ using the fact that the linear continuous map

$$e^{-(s_1-s)P} : B_{\frac{p+2}{3}}^{(1+2\varepsilon)(1-\frac{3}{p+2})}(M) \rightarrow B_{\frac{p+2}{3}}^{1+2\varepsilon}(M)$$

has a norm bounded above by $(s_1-s)^{-\frac{3(1+2\varepsilon)}{2(p+2)}} \leq (s_1-s)^{-1/2}$, a quantity that is integrable over the interval (s, t) . The interpolation estimate

$$\|v_r(s)\|_{B_{\frac{p+2}{3}}^{(1+2\varepsilon)(1-\frac{3}{p+2})}}^{\frac{p+2}{3}} \lesssim \|v_r(s)\|_{L^{\frac{p+2}{3}}} \|v(s)\|_{B_{\frac{p+2}{3}}^{\frac{p+2}{3}-1}}^{\frac{p+2}{3}-1} \lesssim \|v_r(s)\|_{L^{\frac{p+2}{3}}}^p + \|v_r(s)\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{p}{3}} = F_r(s),$$

gives $F_r(s)$ as a final upper bound for this term. Now, with v_r evaluated at time s_1 and $\theta_\varepsilon = \frac{1+\varepsilon}{1+2\varepsilon}$, we have

$$\|v_r\|_{B_{\frac{p+2}{3}}^{1+\varepsilon}} \lesssim \|v_r\|_{L^{\frac{p+2}{3}}}^{1-\theta_\varepsilon} \|v_r\|_{B_{\frac{p+2}{3}}^{\theta_\varepsilon}}^{\theta_\varepsilon} \lesssim \|v_r\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}^{\frac{3\theta_\varepsilon}{2+\theta_\varepsilon}} + \|v_r\|_{L^{p+2}}^3 \leq \delta \|v_r\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}} + C_\delta (1 + \|v_r\|_{L^{p+2}}^3),$$

for some $C_\delta > 0$ and δ small enough so that the term related to $\delta \|v_r\|_{B_{\frac{p+2}{3}}^{1+2\varepsilon}}$ can be absorbed by the left hand side of (2.26). This gives inequality (2.22). \triangleright

3 – Scaling fields, regularity and microlocal extension

We state in this section an extension result, Theorem 13, that provides conditions under which a distribution on a manifold defined outside a submanifold can be extended to the whole manifold. The quantification of this extension result involves the notion of scaling field that is introduced in Section 3.1. Such vector fields are also known as Euler vector fields. Some function spaces associated with a given scaling field are introduced in Section 3.2, they allow to measure the singularity of distributions when we scale along certain submanifolds of some given ambient space. They generalise the weakly homogeneous distributions introduced by Meyer in [65]. An extension Theorem 13 is proven in Section 3.3, given a distribution whose blow up is moderate along a given submanifold, it allows to extend canonically the distribution to the whole ambient space in the spirit of the definition of the principal value. This statement is put to work in the particular setting of a configuration space in Section 4.1 to give a useful extension result for a class of Feynman amplitudes – see Theorem 27. Moreover in Theorem 27, we will need to control the blow up of our Feynman amplitudes in two steps, first when all points collapse on the deepest diagonal, then when

all points collapse to a single given point – this difficulty comes from the absence of translation invariance. This requires a variant of the extension Theorem 13 stated in Theorem 14, in which we scale w.r.t a given submanifold first, and then w.r.t. a given point, in order to control the blow up of our distributions w.r.t. both scalings.

We work in Sections 3.1 to 3.3 in the setting of a smooth manifold \mathcal{X} where a smooth submanifold $\mathcal{Y} \subset \mathcal{X}$ is given.

3.1 Scaling fields

Definition – Let $\mathfrak{I}_{\mathcal{Y}}$ be the ideal of smooth real valued functions on \mathcal{X} that vanish on \mathcal{Y} . Set for $k \geq 1$:

$$\mathfrak{I}_{\mathcal{Y}}^k \stackrel{\text{def}}{=} \{f_1 \dots f_k; (f_1, \dots, f_k) \in \mathfrak{I}_{\mathcal{Y}} \times \dots \times \mathfrak{I}_{\mathcal{Y}}\}.$$

A vector field ρ defined on a neighbourhood of \mathcal{Y} is called an isotropic **scaling field** for $\mathcal{Y} \subset \mathcal{X}$ if for all $f \in \mathfrak{I}_{\mathcal{Y}}$

$$f \in \rho f + \mathfrak{I}_{\mathcal{Y}}^2.$$

This type of vector field is also called an *Euler vector field* in the literature. Denote by n the dimension of \mathcal{X} and by d the dimension of \mathcal{Y} . If ρ is a scaling field for $\mathcal{Y} \subset \mathcal{X}$ there exists a neighbourhood of \mathcal{Y} that is stable by the backward semiflow $(e^{-s\rho})_{s \geq 0}$ of ρ and every point $y \in \mathcal{Y}$ has a neighbourhood U_y in \mathcal{X} on which coordinates

$$h = (h_1, \dots, h_n) : U \rightarrow \mathbb{R}^n$$

are defined and such that

$$U_y \cap \mathcal{Y} = h^{-1}(\mathbb{R}^d)$$

with $\mathbb{R}^d \subset \mathbb{R}^n$, and

$$\rho = \sum_{i=d+1}^n h^i \partial_{h^i}.$$

A proof of existence of a stable neighbourhood can be found in Lemma 2.4 of [28] and the normal form theorem can be found in Proposition 2.5 of [28] – see also Lemma 2.1 in [63]. The example of the configuration space of ℓ points in \mathbb{R}^k will be particularly relevant for us. The scaling field ρ whose flow reads

$$e^{-t\rho}(x_1, \dots, x_\ell) = \left(x_1, e^{-t}(x_2 - x_1) + x_1, \dots, e^{-t}(x_\ell - x_1) + x_1 \right)$$

will move all points towards the deepest diagonal and its dynamics is tangent to all the larger diagonals. In the sequel, we will only work with product (sub)manifolds of the form

$$(3.1) \quad \begin{aligned} \mathcal{X} &= \mathbb{R}^p \times X, \\ \mathcal{Y} &= (\{0\} \times \mathbb{R}^q) \times Y \end{aligned}$$

with $(\{0\} \times \mathbb{R}^q) \subset \mathbb{R}^p$ and $Y \subset X$, and some non-isotropic scaling fields, of the form

$$(3.2) \quad \rho = \sum_{j=1}^q 2t_j \partial_{t_j} + \rho_Y$$

for the canonical coordinates $(t_j)_{1 \leq j \leq q}$ on \mathbb{R}^q and a scaling field ρ_Y for $Y \subset X$. This is an example of a **weighted vector field on a weighted manifold**. We give a formal definition in the case where our submanifold \mathcal{Y} is the transverse intersection $\mathcal{Y}_1 \cap \mathcal{Y}_2$ where $\mathcal{Y}_1 = (\{0\} \times \mathbb{R}^q) \times X$ and $\mathcal{Y}_2 = (\mathbb{R}^p \times Y)$. Then one has a description of the ideal $\mathfrak{I}_{\mathcal{Y}}$ as the product $\mathfrak{I}_{\mathcal{Y}_1} \mathfrak{I}_{\mathcal{Y}_2}$ so the ideal $\mathfrak{I}_{\mathcal{Y}}$ has a bifiltration

$$\mathfrak{I}_{\mathcal{Y}_1}^m \mathfrak{I}_{\mathcal{Y}_2}^n \subset \dots \subset \mathfrak{I}_{\mathcal{Y}_1} \mathfrak{I}_{\mathcal{Y}_2} = \mathfrak{I}_{\mathcal{Y}}.$$

Now assume we want to put a weight 2 to powers of $\mathfrak{I}_{\mathcal{Y}_1}$ and weight 1 to powers of $\mathfrak{I}_{\mathcal{Y}_2}$. We want to give an intrinsic characterization for vector fields of the form given by Equation (3.2).

Definition – A vector field ρ defined on a neighbourhood of \mathcal{Y} is called a *parabolic scaling field* for $\mathcal{Y} \subset \mathcal{X}$ if for all (m, n) and for all $f \in \mathfrak{I}_{\mathcal{Y}_1}^m \mathfrak{I}_{\mathcal{Y}_2}^n$ one has

$$\rho f - (2m + n)f \in \left(\mathfrak{I}_{\mathcal{Y}_1}^{m+1} \mathfrak{I}_{\mathcal{Y}_2}^n + \mathfrak{I}_{\mathcal{Y}_1}^m \mathfrak{I}_{\mathcal{Y}_2}^{n+1} \right).$$

In the sequel we will simply call ‘scaling fields’ some parabolic scaling fields as we will only work with such fields. The **weighted co-dimension** of \mathcal{Y} is defined here as

$$\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) \stackrel{\text{def}}{=} 2(p - q) + \dim(X) - \dim(Y).$$

3.2 Function spaces associated with scaling fields

We assume from now on that \mathcal{X} has a Riemannian structure and we denote by $K_r(x, y)$ its heat kernel. We also let $U \subset \mathcal{X}$ be an open set and Γ be a closed conic set in $T^*U \setminus \{0\}$.

Definition 8 – We denote by $\mathcal{D}'_\Gamma(U)$ the space of distributions on U whose wave front set is contained in Γ . This is a locally convex topological vector space endowed with a natural normal topology invented by Y. Dabrowski – see [25, p. 823] and [18] for results about why this topology is well-behaved with respect to some natural operations on distributions. The seminorms defining its topology are

- given a chart $\kappa : \Omega \subset U \mapsto \mathbb{R}^{\dim(\mathcal{X})}$, an integer $N \in \mathbb{N}$, $\chi \in C_c^\infty(\kappa(\Omega))$, and a cone $V \subset \mathbb{R}^{n^*}$ such that

$$\text{supp}(\chi) \times V \cap \kappa_*\Gamma = \emptyset, \quad \text{where} \quad \kappa_*\Gamma = \left\{ (\kappa(x); ({}^t d\kappa_x)^{-1}(\xi)); (x; \xi) \in \Gamma \right\},$$

we have the norm

$$\|\Lambda\|_{N, V, \chi, \kappa} = \sup_{\xi \in V} (1 + |\xi|)^N |(\widehat{\kappa_*\Lambda})\chi(\xi)|;$$

- the seminorms of the strong topology of distributions

$$\sup_{\chi \in \mathcal{B}} |\langle \Lambda, \chi \rangle|,$$

where \mathcal{B} is a bounded set of $C_c^\infty(\mathcal{X})$ which means that there is some compact K such that $\text{supp}(f) \subset K$ for all $f \in \mathcal{B}$, and for any differential operator Q one has $\sup_{\chi \in \mathcal{B}} \|Q\chi\|_{L^\infty(K)} < \infty$.

To be bounded in $\mathcal{D}'_\Gamma(U)$ will always mean that all the above seminorms are bounded.

The following elementary example will play an important role in the sequel. Assume $k \in \mathbb{N}$ is of the form $d_1 + d_2 + d_3$ with $d_i \in \mathbb{N} \setminus \{0\}$ and $\mathbb{R}^k = \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \times \mathbb{R}^{d_3}$, and denote here by ρ the linear vector field on \mathbb{R}^k whose restriction to \mathbb{R}^{d_1} is null, whose restriction to \mathbb{R}^{d_2} is the identity and whose restriction to \mathbb{R}^{d_3} is twice the identity. So for $z = (x, y, t) \in \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \times \mathbb{R}^{d_3}$ one has

$$\rho = y\partial_y + 2t\partial_t.$$

The third component $t \in \mathbb{R}^{d_3}$ has the meaning of a type of time coordinates. This vector field over \mathbb{R}^k will be our model scaling field in a parabolic setting.

Lemma 9 – The family of distribution

$$\delta(z' - e^{-\ell\rho}z) \quad (1 \leq \ell \leq +\infty)$$

on $\mathbb{R}^k \times \mathbb{R}^k$ is bounded in $\mathcal{D}'_{\Gamma_\rho}(\mathbb{R}^k \times \mathbb{R}^k)$, where

$$\Gamma_\rho = \bigcup_{1 \leq \ell \leq +\infty} \left\{ ((z, e^{-\ell\rho}z), (\lambda, e^{\ell\rho}\lambda)); (z, \lambda) \in T^*\mathbb{R}^k \right\} \subset T^*(\mathbb{R}^k \times \mathbb{R}^k).$$

This estimate can also be used to give an upper bound on the wave front set of the resolvent $(\rho + z)^{-1}$ which implies the radial type estimates for ρ . This is very similar in spirit to the radial estimates from the works of Melrose [64], Vasy [79] or Dyatlov & Zworski [34].

Proof – Note that the distributions $\delta(z' - e^{-\ell\rho}z) \in \mathcal{D}'(\mathbb{R}^k \times \mathbb{R}^k)$ are nothing but the Schwartz kernels of the transfer operator $\varphi \in C^\infty(\mathbb{R}^k) \mapsto e^{-\ell\rho^*}\varphi \in C^\infty(\mathbb{R}^k)$, so we will use the identification $[e^{-\ell\rho^*}] = \delta(z' - e^{-\ell\rho}z)$. The fact that this family of distributions is bounded (weak boundedness implies strong boundedness by uniform boundedness) automatically follows from the continuity of the pull-back of a distribution by a smooth family of diffeomorphisms and the strong convergence of $\delta(z' - e^{-\ell\rho}z)$ to $\delta(z' - (x, 0, 0))$ when ℓ goes to infinity, for $z = (x, y, t)$.

Fix an arbitrary compact subset $K \subset \mathbb{R}^{d_1+d_2+d_3}$ that is stable by the scaling maps

$$(x, y, \mathbf{t}) \mapsto (x, e^{-\ell}y, e^{-2\ell}\mathbf{t}) \quad (\ell \geq 0).$$

Then we shall restrict the Schwartz kernel $[e^{-\ell\rho}]$ to $K \times K$. It means we estimate this wave front set near the diagonal but for arbitrary large ℓ . Choose some test functions χ_1, χ_2 in $C_K^\infty(\mathbb{R}^{d_1+d_2+d_3})$, supported in K . In local coordinates we have

$$\begin{aligned} & \int_{\mathbb{R}^{d_1+d_2+d_3}} e^{i\xi_2 \cdot x + i\eta_2 \cdot y + i\tau_2 \cdot t} \chi_2(x, y, t) e^{-t\rho^*} (\chi_1 e^{i\xi_1 \cdot x + i\eta_1 \cdot y + i\tau_1 \cdot t}) dx dy dt \\ &= e^{-\ell(d_2+2)} \int_{\mathbb{R}^{d_1+d_2+d_3}} e^{i\xi_2 \cdot x + i\eta_2 \cdot y + i\tau_2 t} \chi_2(x, y, t) (\chi_1(x, e^{-\ell}y, e^{-2\ell}t) e^{i\xi_1 \cdot x + i e^{-\ell}\eta_1 \cdot y + e^{-2\ell}\tau_1 \cdot t}) dx dy dt \\ &= \widehat{\chi_\ell}(\xi_1 + \xi_2, \eta_2 + e^{-\ell}\eta_1, \tau_2 + e^{-2\ell}\tau_1) \end{aligned}$$

where

$$\chi_\ell(x, y, t) = \chi_2(x, y, t) \chi_1(x, e^{-\ell}y, e^{-2\ell}t)$$

is a bounded family of smooth compactly supported functions (this is crucial) when $\ell \in [0, +\infty)$. We then have for any $N \geq 1$ the upper bound

$$(3.3) \quad |\widehat{\chi_\ell}(\xi, \eta, \tau)| \leq C_N (1 + |\xi| + |\eta| + |\tau|)^{-N}$$

where the constant C_N does not depend on $\ell \in [0, +\infty)$. Hence in any closed conic set V which does not meet the subset

$$\Lambda = \left\{ (\xi, -\xi, \eta, -e^{-\ell}\eta, \tau, -e^{-2\ell}\tau) \in (\mathbb{R}^k \times \mathbb{R}^k)^*, \ell \geq 1 \right\},$$

there exists some $\varepsilon > 0$ such that for all $(\xi_1, \xi_2, \eta_1, \eta_2, \tau_1, \tau_2) \in V \subset (\mathbb{R}^k \times \mathbb{R}^k)^*$ we have for all $\ell \geq 1$ the inequality

$$\left| (\xi_1 + \xi_2, e^{-s}\eta_1 + \eta_2, e^{-2s}\tau_1 + \tau_2) \right| \geq \varepsilon (|\xi_1| + |\xi_2| + |\eta_1| + |\eta_2| + |\tau_1| + |\tau_2|).$$

This implies the following Fourier bound

$$\begin{aligned} & \left| \int_{\mathbb{R}^k} e^{i\xi_2 \cdot x + i\eta_2 \cdot y + i\tau_2 \cdot t} \chi_2(x, y, t) e^{-\ell\rho^*} (\chi_1 e^{i\xi_1 \cdot x + i\eta_1 \cdot y + i\tau_1 \cdot t}) dx dy dt \right| \\ & \leq C_N \left(1 + |\xi_1 + \xi_2| + |\eta_2 + e^{-s}\eta_1| + |\tau_2 + e^{-2s}\tau_1| \right)^{-N} \\ & \leq C_N \varepsilon^{-N} \left(1 + |\xi_1| + |\xi_2| + |\eta_1| + |\eta_2| + |\tau_1| + |\tau_2| \right)^{-N} \end{aligned}$$

for all $\ell \geq 1$ and $(\xi_1, \xi_2, \eta_1, \eta_2, \tau_1, \tau_2) \in V \subset (\mathbb{R}^k \times \mathbb{R}^k)^*$. The previous bound analyzes the wave front set of the family $\delta(z' - e^{-\ell\rho}(z))$ near $T^*(K \times K) \subset T^*(\mathbb{R}^k \times \mathbb{R}^k)$. Since K is arbitrary the family of distributions $\delta(z' - e^{-\ell\rho}(z))$ is bounded in $\mathcal{D}'_{\Gamma_\rho}(\mathbb{R}^k \times \mathbb{R}^k)$, with $\Gamma_\rho \subset T^*(\mathbb{R}^k \times \mathbb{R}^k)$ given by

$$\begin{aligned} \Gamma_\rho = & \underbrace{\left\{ (x, x, 0, 0, 0, 0; \xi, -\xi, 0, \eta_2, 0, \tau_2); (\xi, \eta_2, \tau_2) \neq (0, 0, 0) \right\}}_{\text{the radial set which is the conormal of the singular set of } \rho} \\ & \cup \left\{ (x, y, t, x, e^{-\ell}y, e^{-2\ell}t; \xi, \eta, \tau, -\xi, -e^{-\ell}\eta, -e^{-2\ell}\tau); \ell \geq 1, (\xi, \eta, \tau) \neq (0, 0, 0) \right\}. \end{aligned}$$

This concludes the proof. \triangleright

Set

$$(3.4) \quad \pi(z) \stackrel{\text{def}}{=} (x, 0, 0).$$

Lemma 9 is useful to give a description of the Taylor subtraction operation. The Taylor subtractors of order 0 and 1 read

$$\begin{aligned} R_0 : \varphi &\mapsto \varphi - \varphi \circ \pi, \\ R_1 : \varphi &\mapsto \varphi - \varphi(x, 0, 0) - y \cdot \partial_y \varphi(x, 0, 0) - t \partial_t \varphi(x, 0, 0), \end{aligned}$$

We call these operators R_0 and R_1 , with the letter R chosen for ‘remainder’. Denote generically by $[\Lambda]$ the Schwartz kernel of an operator Λ .

Proposition 10 – *The operators R_0, R_1 have Schwartz kernel*

$$\begin{aligned} [R_0] &= \int_0^\infty [\rho e^{-a\rho^*}] da, \\ [R_1] &= \int_0^\infty [(1 - e^a + e^a \rho) \rho e^{-a\rho^*}] da - [\rho] \end{aligned}$$

and

$$\begin{aligned} [e^{-\ell\rho^*} R_0] &= \int_0^\infty [\rho e^{-(\ell+a)\rho^*}] da, \\ [e^{-\ell\rho^*} R_1] &= \int_0^\infty [(1 - e^a + e^a \rho) \rho e^{-(\ell+a)\rho^*}] da - [e^{-\ell\rho^*} \rho]. \end{aligned}$$

and the families of distributions $([(e^{-\ell\rho})^* R_0])_{0 \leq \ell \leq +\infty}$ and $([(e^{-\ell\rho})^* R_1])_{0 \leq \ell \leq +\infty}$ are bounded in $\mathcal{D}'_{\Gamma_\rho}(\mathbb{R}^k \times \mathbb{R}^k)$.

Proof – We write a detailed proof for R_0 ; the proof for R_1 is very similar and left to the reader. For a test function χ with compact support on $\mathbb{R}^k \times \mathbb{R}^k$ write $\rho_1 \chi$ for the action of the vector field on the first component of χ . We have

$$\langle [\rho(e^{-\ell\rho})^*], \chi \rangle = - \int_{\mathbb{R}^k} (\rho_1 \chi)(z, e^{-\ell\rho}(z)) dz$$

and since $(\rho_1 \chi)(z, e^{-\ell\rho} z)$ vanishes along the singular set $\{y = 0, t = 0\}$ the integrand is of order $e^{-\ell}$, so the integral is converging. The wavefront bound follows from the wave front bound on the propagator $[(e^{-\ell\rho})^*]$ and the fact that the wave front of a distribution is stable under the action the vector field ρ . \triangleright

We come back to the general setting of an open subset $U \subset \mathcal{X}$ and assume we are given a closed conic set Γ in $T^*U \setminus \{0\}$. It is a classical fact that for $\alpha < 0$ the Besov space $C^\alpha(\mathcal{X}) = B_{\infty, \infty}^\alpha(\mathcal{X})$ can be characterized as the set of distributions Λ on \mathcal{X} such that

$$\sup_{x \in \mathcal{X}} \sup_{0 < r \leq 1} r^{-\frac{\alpha}{2}} |\langle \Lambda, K_r(x, \cdot) \rangle| < \infty.$$

A distribution $\Lambda \in \mathcal{D}'(U)$ is an element of $\mathcal{D}'_\Gamma(U)$ iff for all pseudodifferential operators Q with Schwarz kernel compactly supported in $U \times U$ and whose symbol vanishes on Γ one has for every compact subset C of \mathcal{X}

$$(3.5) \quad \sup_{x \in C} \sup_{0 < r \leq 1} |\langle \Lambda, Q K_r(x, \cdot) \rangle| < \infty.$$

One can describe an element of $C_{\text{loc}}^\alpha(U)$, with $\alpha < 0$, with wave front set in Γ in terms similar to (3.5) as the set of distributions $\Lambda \in C_{\text{loc}}^\alpha(U)$ iff for all pseudodifferential operators Q with Schwarz kernel compactly supported in $U \times U$ and whose symbol vanishes on Γ one has for every compact subset C of \mathcal{X}

$$\sup_{x \in C} \sup_{0 < r \leq 1} r^{-\frac{\alpha}{2}} |\langle \Lambda, (I + Q) K_r(x, \cdot) \rangle| < \infty.$$

The I element ensures that Λ is Hölder whereas the operator Q is here to test the smoothness of Λ outside of the wave front set.

We now come back to the parabolic setting $\mathcal{Y} \subset \mathcal{X}$ of (3.1) in Section 3.1 and denote by ρ a parabolic scaling field for this embedding. Let U stand for an open set of \mathcal{X} which is stable by the backward semiflow of ρ , in the sense that $e^{-\ell\rho}(U) \subset U$ for all $\ell \geq 0$.

Definition 11 – *Given some closed conic set $\Gamma \subset T^*U \setminus \{0\}$ we denote by $(e^{-\ell\rho})^* \Gamma$ the set defined as*

$$(e^{-\ell\rho})^* \Gamma \stackrel{\text{def}}{=} \left\{ (e^{\ell\rho} x; ({}^t \text{de}^{\ell\rho})_x^{-1}(\xi)); (x; \xi) \in \Gamma \right\}.$$

Then we assume that the lifted flow of $e^{-\ell\rho}$ leaves the conic set Γ stable i.e.

$$(e^{-\ell\rho})^* \Gamma \subset \Gamma$$

for all $\ell \geq 0$.

Having described the necessary geometric framework, we are ready to state the definition of the scaling spaces that will be used extensively in the present work.

Definition – For $\alpha < 0$ and $a \in \mathbb{R}$ we define the **scaling space** $\mathcal{S}_\Gamma^{a,\rho;\alpha}(U)$ of distributions $\Lambda \in \mathcal{D}'(U)$ with the following property. For all pseudodifferential operators Q with Schwarz kernel compactly supported in $U \times U$ and whose symbol vanishes on Γ , for each compact set $C \subset U$

$$\sup_{\ell \geq 1} \sup_{x \in C} \sup_{0 < r \leq 1} e^{a\ell} r^{-\frac{\alpha}{2}} \left| \langle (e^{-\ell\rho})^* \Lambda, (I + Q)K_r(x, \cdot) \rangle \right| < \infty.$$

We define $\mathcal{S}_\Gamma^a(U)$ as the union over α of all the spaces $\mathcal{S}_\Gamma^{a,\rho;\alpha}(U)$, for $a \in \mathbb{R}$ fixed and ρ a parabolic scaling field for the inclusion $\mathcal{Y} \subset \mathcal{X}$ whose backward semiflows leave Γ fixed. This implies in particular that $N^*(\mathcal{Y} \subset \mathcal{X}) \subset \Gamma$. The letter ‘ \mathcal{S} ’ is chosen for *scaling*. The exponent a retains the scaling property and Γ retains some information on the wavefront set. Note that the space $\mathcal{S}_\Gamma^a(U)$ is a priori larger than the space of conormal distributions with wavefront set in $N^*(\mathcal{Y} \subset U)$ since elements in $\mathcal{S}_\Gamma^a(U)$ might have some wavefront set contained in the cone Γ which is not necessarily included in $N^*(\mathcal{Y} \subset U)$. The notation $\mathcal{S}_\Gamma^a(U)$ does not emphasize the dependence of this space on the inclusion $\mathcal{Y} \subset \mathcal{X}$. This will always be clear for us from the context. An elementary example is given by the principal value of $1/|x|$ in \mathbb{R} , where $\mathcal{Y} = \{0\} \subset \mathbb{R}$ and it has scaling exponent $a = -1$ and wavefront set $\Gamma = T_0^*\mathbb{R}$. Note the fact that for all element $\Lambda \in \mathcal{S}_\Gamma^a(U)$ the family of scaled distributions $(e^{a\ell} e^{-\ell\rho^*} \Lambda)_{\ell \geq 0}$ is bounded in \mathcal{D}'_Γ . The next proposition gives an example of an element of some space $\mathcal{S}^{a,\rho;0}(U)$ for some scaling exponent a and some scaling field ρ . For $n \geq 2$, denote by \mathbf{d}_n the diagonal of M^n . Denote by ρ_n a scaling field on M^n for the inclusion $\mathbf{d}_n \subset M^n$ and define on $M^2 \times \mathbb{R}^2$ the parabolic scaling field

$$\rho = 2(t - s)\partial_s + \rho_2.$$

Denote also by

$$\pi_{\leq 2} : (x_1, \dots, x_n) \in M^n \rightarrow (x_1, x_2) \in M^2$$

the canonical projection on the first two components.

Lemma 12 – Let M be a closed manifold and $A_r(x, y)$ be a smooth kernel on $M^2 \setminus \mathbf{d}_2$ such that one can associate to any small enough open set U a coordinate system in which one has for all multiindices α, β

$$(3.6) \quad \left| \partial_{s,t}^\alpha \partial_{x,y}^\beta A_{|t-s|}(x, y) \right| \lesssim (\sqrt{t-s} + |y-x|)^{-a-2|\alpha|-|\beta|}.$$

Then the family

$$\left(e^{\ell a} (e^{-\ell\rho})^* \pi_{\leq 2}^* A_{|t-s|}(\cdot, \cdot) \right)_{\ell \geq 0}$$

is bounded in

$$\mathcal{D}'_{N^*(\{s=t\})} \left((M^n \times \mathbb{R}) \setminus (\pi_{\leq 2}^* \mathbf{d}_2 \cap \{s=t\}) \right),$$

that is

$$\pi_{\leq 2}^* A_{|t-s|}(\cdot, \cdot) \in \mathcal{S}_{N^*(\{s=t\})}^{a,\rho;0} \left((M^n \times \mathbb{R}) \setminus (\pi_{\leq 2}^* \mathbf{d}_2 \cap \{s=t\}) \right).$$

In the sequel we denote by \mathcal{K}^a the C^∞ -module of kernels $A_r(x, y)$ as above depending on two variables endowed with the weakest topology containing the $C^\infty([0, +\infty) \times (M^2 \setminus \mathbf{d}_2))$ topology and which makes all the seminorms defined by the estimates (3.6) continuous.

Proof – We first localize in a neighbourhood $U \times U$ of the diagonal since K is smooth off-diagonal. It is enough to prove the claim for $A_{|t-s|}(x, y)\chi_1(y)\chi_2(x)$ where $\chi_i \in C_c^\infty(U)$ and use a partition of unity to get the global result. In $U \times U$ we pull-back everything to the configuration space, which we write with a slight abuse of notations

$$\pi_{\leq 2}^*(A\chi_1\chi_2)(t, s, x_1, \dots, x_n) = A_{|t-s|}(x_1, x_2)\chi_1(x_1)\chi_2(x_2).$$

We already know that this kernel satisfies some bound of the form

$$\left| A(t, s, x_1, x_2)\chi_1(x_1)\chi_2(x_2) \right| \lesssim \left(\sqrt{|t-s|} + |x_1 - x_2| \right)^{-a}.$$

Somehow we would like to flow both sides of the inequality by the parabolic dynamics $(e^{-\ell\rho})^*$ and bound the term $e^{-\ell\rho^*} \left(\sqrt{|t-s|} + |x_1 - x_2| \right)^{-a}$ asymptotically when ℓ goes to $+\infty$. We use for

that purpose the Normal Form Theorem for the space part of the isotropic scaling fields

$$\rho_{[n]} \stackrel{\text{def}}{=} \sum_{k=2}^n h_k \cdot \partial_{h_k},$$

for some new coordinates $(h_k)_{k=2}^n$ that vanish at order 1 along the deep space diagonal \mathbf{d}_n . The fact that $x_1 - x_2$ vanishes at first order along \mathbf{d}_n implies by Taylor expansion at first order that

$$(3.7) \quad x_1 - x_2 = L(h) + \mathcal{O}(|h|^2)$$

where $L(h)$ is a linear function of $(h_k)_{k=2}^n$. One then has

$$(e^{-\ell\rho_{[n]}})^*(x_1 - x_2) = (e^{-\ell\rho_{[n]}})^*L(h) + \mathcal{O}(e^{-2\ell}|h|^2) = L(e^{-\ell}h) + \mathcal{O}(e^{-2\ell}|h|^2),$$

and an exponential lower bound of the form

$$e^{-\ell}|x_1 - x_2| \lesssim |(e^{-\ell\rho_{[n]}})^*(x_1 - x_2)|$$

which yields the desired bound

$$\left| \partial_t^\alpha \partial_x^\beta e^{-\ell\rho^*} \pi^*(A\chi_1\chi_2)(t, s, x_1, \dots, x_n) \right| \lesssim e^{\ell a} \left(\sqrt{|t-s|} + |x_1 - x_2| \right)^{-a-2|\alpha|-|\beta|}$$

and proves the claim. The above bound allows for instance to justify that the singularities when $x_1 \neq x_2$ are conormal along the equal time region $t = s$ since we are smooth on each half region $t \geq s$ and $s > t$. \triangleright

3.3 The canonical extension

3.3.1 The basic extension result. Let us use a unique notation $\mathbf{0}$ for the zero section of any vector bundle.

Definition – Let \mathcal{X} be a smooth manifold and $\mathcal{Y} \subset \mathcal{X}$. A closed conic set $\Gamma \subset T^*(\mathcal{X} \setminus \mathcal{Y}) \setminus \mathbf{0}$ is said to satisfy the **conormal landing condition** if its closure $\tilde{\Gamma}$ in $T^*(\mathcal{X}) \setminus \mathbf{0}$ satisfies $\tilde{\Gamma} \subset (\Gamma \cup N^*(\mathcal{Y}))$.

In what follows, we will use the terminology *smooth weighted manifolds* for smooth weighted manifolds of product type as introduced in subsection 3.1.

Theorem 13 – Let \mathcal{X} be a smooth weighted manifold and $\mathcal{Y} \subset \mathcal{X}$ and $\Gamma \subset T^*(\mathcal{X} \setminus \mathcal{Y}) \setminus \mathbf{0}$ be a closed conic set that satisfy the conormal landing condition. Assume we are given a family $(\Lambda_\varepsilon)_{0 < \varepsilon \leq 1}$ of distributions on \mathcal{X} that converge in $\mathcal{D}'(\mathcal{X} \setminus \mathcal{Y})$ as ε goes to 0 to an element $\Lambda \in \mathcal{S}'_\Gamma(\mathcal{X} \setminus \mathcal{Y})$.

(a) If

$$-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) < a$$

then Λ has a unique extension into a distribution over \mathcal{X} such that the convergence of Λ_ε to Λ occurs in $\mathcal{S}'_{\Gamma \cup N^*(\mathcal{Y})}(\mathcal{X})$.

(b) If

$$-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1 \leq a < -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$$

there exists a family $\Lambda_{\mathcal{Y}, \varepsilon}$ of distributions supported on \mathcal{Y} , with wavefront set in $N^*(\mathcal{Y})$ such that $\Lambda_\varepsilon - \Lambda_{\mathcal{Y}, \varepsilon}$ has a limit in $\mathcal{D}'(\mathcal{X})$ and the convergence occurs in $\mathcal{S}'_{\Gamma \cup N^*(\mathcal{Y})}(\mathcal{X})$ for all

$$a' < a.$$

The particular case where $\mathcal{Y} = \{p\}$ is a point will be used in Theorem 14 below. We loose any information on the wavefront set of the extension at p in that case, so the convergence of Λ_ε to Λ happens in $\mathcal{S}'_{\Gamma \cup T^*_p M}(\mathcal{X})$, for $b \in \{a, a'\}$, depending on the situation.

Proof – We follow the proof of similar results proved in an elliptic setting in [25] – see Theorem 1.10, Theorem 4.4 and Section 6 therein. We give here the main arguments to emphasize the differences with [25] that come from our parabolic setting. The main idea of the proof is to start from a continuous partition of unity which approximates the constant function 1 but vanishes near \mathcal{Y} . Then we multiply the distribution Θ by this partition of unity to product an approximation of Θ outside \mathcal{Y} and use the assumption on the weak homogeneity of Θ near \mathcal{Y} to conclude that the approximation genuinely converges to some given distribution Θ^+ which is the desired extension.

Let ρ be a scaling field for the inclusion $\mathcal{Y} \subset \mathcal{X}$ such that $\Lambda \in \mathcal{S}_{\Gamma \cup N^*(\mathcal{Y})}^{a, \rho; \alpha}(\mathcal{X})$ for some $\alpha \in \mathbb{R}$, and let χ be a smooth function equal to 1 in a neighborhood of \mathcal{Y} stable by the backward semiflow of ρ and such that χ vanishes outside some larger neighborhood.

(1) We first use the normal form theorem to reduce our problem to the model case of a distribution on \mathbb{R}^k , with $k = d_1 + d_2 + d_3$ with coordinates (x, y, t) , the scaling field $\rho = y\partial_y + 2t\partial_t$ is linear and globally defined, and the extension is done with respect to the linear subspace $\mathbb{R}^{d_1} \subset \mathbb{R}^k$. We work in that setting in the remainder of the proof. Let then $(\Theta_\varepsilon)_{0 < \varepsilon \leq 1}$ be a family of distributions on \mathbb{R}^k . We assume that the Θ_ε converge as ε goes to 0 to an element $\Theta \in \mathcal{S}'_{\Gamma}(\mathbb{R}^k \setminus \mathbb{R}^{d_1})$ where $a > -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$.

(2) Pick $0 < \ell_0$ and think of it as being large. We use the continuous partition of unity

$$\text{Id} - (e^{\ell_0 \rho})^* \chi = \text{Id} - \chi + \int_0^{\ell_0} (e^{\ell \rho})^* (-\rho \chi) d\ell$$

to define an extension of our distribution Θ . We set for convenience $\bar{\chi} \stackrel{\text{def}}{=} -\rho \chi$; its support does not meet \mathcal{Y} . We have for any test function $f \in \mathcal{D}(\mathbb{R}^k)$

$$\langle \Theta(1 - (e^{\ell_0 \rho})^* \chi), f \rangle = \langle \Theta(1 - \chi), f \rangle + \int_0^{\ell_0} e^{-\ell(d_2 + 2d_3)} \langle \bar{\chi}(e^{-\ell \rho})^* \Theta, (e^{-\ell \rho})^* f \rangle d\ell.$$

The exponential factor $e^{-\ell(d_2 + 2d_3)}$ comes from the Jacobian of the flow of $e^{-\ell \rho}$. Note that $d_2 + 2d_3 = \text{codim}_w(\mathcal{Y} \subset \mathcal{X})$. If $\Gamma \subset T^*\mathbb{R}^k \setminus \mathbf{0}$ stands for a closed conic set invariant by the lifted dynamics of $(e^{-\ell \rho})^*$ such that $\Gamma \cap T^*\mathbb{R}^{d_1} \subset N^*(\mathbb{R}^{d_1})$, our choice of scaling exponent a ensures that the family

$$(\Theta^{(\ell)})_{\ell \geq 0} \stackrel{\text{def}}{=} (e^{a\ell}(e^{-\ell \rho})^* \Theta)_{\ell \geq 0}$$

is bounded in $\mathcal{D}'_{\Gamma}(\mathbb{R}^k)$. One then has for the Schwartz kernels

$$[\Theta(1 - (e^{\ell_0 \rho})^* \chi)](z, z') = [\Theta(1 - \chi)](z, z') + \int_0^{\ell_0} e^{-\ell(a + d_2 + 2d_3)} (\bar{\chi} \Theta^{(\ell)})(z) \delta(z' - e^{-\ell \rho}(z)) d\ell.$$

We know from the hypocontinuity theorem on the Hörmander product of distributions [18, Thm 6.1 p. 219] that the family

$$(\bar{\chi} \Theta^{(\ell)})(z) \delta(z' - e^{-\ell \rho}(z))$$

with $\ell \geq \ell_1$ large enough, is bounded in $\mathcal{D}'_{\Gamma}(\mathbb{R}^k)$, where

$$\bar{\Gamma} \stackrel{\text{def}}{=} (\Gamma \times \mathbf{0}) \cup \Gamma_\rho \cup ((\Gamma \times \mathbf{0}) + \Gamma_\rho).$$

For the moment this means that the ℓ_0 -dependent family of distributions associated with the kernels

$$(\star) \stackrel{\text{def}}{=} \int_0^{\ell_0} e^{-\ell(a + d_2 + 2d_3)} (\bar{\chi} \Theta^{(\ell)})(z) \delta(z' - e^{-\ell \rho}(z)) d\ell$$

is bounded in $\mathcal{D}'_{\bar{\Gamma}}(\mathbb{R}^k \times \mathbb{R}^k)$ uniformly in $\ell_0 \geq \ell_1$. In particular the integral converges in $\mathcal{D}'_{\bar{\Gamma}}(\mathbb{R}^k \times \mathbb{R}^k)$ when ℓ_0 goes to $+\infty$. Now we interpret the integral over the variable z as a push-forward along the fibers of the linear projection

$$p_2 : (z, z') \mapsto z'.$$

The pushforward Theorem yields that $p_{2*}(\cdot)$ is bounded in $\mathcal{D}'_{p_{2*}\bar{\Gamma}}(\mathbb{R}^k)$ where

$$p_{2*}\bar{\Gamma} = (p_{2*}\Gamma_\rho) \cup p_{2*}(\Gamma \times \mathbf{0} + \Gamma_\rho)$$

and

$$p_{2*}\Gamma_\rho = \{(x, 0, 0), (0, \eta, \tau)\} \\ p_{2*}(\Gamma \times \mathbf{0} + \Gamma_\rho) = \left\{ \left(e^{-\ell \rho}(z), (e^{-\ell \rho})^*(\lambda) \right); (z, \lambda) \in \Gamma, 0 \leq \ell \leq +\infty \right\} \subset \Gamma \cup N^*(\mathcal{Y})$$

since the cone Γ is invariant by the lifted flow of $e^{-\ell \rho}$ provided $\ell < +\infty$ and the limit points of the form $\lim_{\ell \uparrow \infty} (e^{-\ell \rho}(z), (e^{-\ell \rho})^*(\lambda))$ for $(z, \lambda) \in \Gamma$ must belong to the conormal $N^*(\mathcal{Y})$ by the conormal landing condition on Γ . It is at this precise place that we are using the conormal landing condition assumption on Γ . The distributions $\Theta(1 - (e^{\ell \rho})^* \chi)$ are thus converging in $\mathcal{D}'_{\Gamma \cup N^*(\mathbb{R}^{d_1})}(\mathbb{R}^k)$

to

$$\langle \Theta^+, f \rangle = \langle \Theta(1 - \chi), f \rangle + \int_0^\infty e^{-\ell(d_2+2d_3)} \langle \bar{\chi}(e^{-\ell\rho})^* \Theta, f \circ e^{-\ell\rho} \rangle d\ell.$$

The uniqueness of the extension Θ^+ follows from the continuity of all the operations involved in above. To see the scaling property of the extension we note that the family

$$(\Theta^{(\ell)})_{\ell \geq 0} = (e^{a\ell}(e^{-\ell\rho})^* \Theta)_{\ell \geq 0}$$

is bounded in $\mathcal{D}'_\Gamma(\mathbb{R}^k)$, since $\Theta \in \mathcal{S}'_\Gamma(\mathbb{R}^k)$ means that the family $(e^{a\ell}(e^{-\ell\rho})^* \Theta)_{\ell \geq 0}$ is bounded in $\mathcal{D}'_\Gamma(\mathbb{R}^k)$ by definition of $\mathcal{D}'_\Gamma(\mathbb{R}^k)$ and $\mathcal{S}'_\Gamma(\mathbb{R}^k)$, and observe that

$$e^{\ell a}(e^{-\ell\rho})^* \Theta^+ = p_{2*} \left(\int_0^\infty e^{-\ell(a+d_2+2d_3)} ((\Theta^{\ell+a})' \bar{\chi} \otimes 1) [e^{-\ell\rho}] d\ell \right).$$

(3) In the borderline case our proof follows closely [25, Prop 4.9 p. 841] except we work in a parabolic setting. We proceed as above with $(e^{-\ell\rho})^*$ replaced by $(e^{-\ell\rho})^* R_0$ if $-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1 < a \leq -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$ and $(e^{-s\rho})^* R_1$ if $a = -\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1$. So our extension reads

$$\langle \Theta^+, f \rangle = \langle \Theta(1 - \chi), f \rangle + \int_0^\infty e^{-\ell(d_2+2d_3)} \langle \bar{\chi}(e^{-\ell\rho})^* \Theta, (R_i f) \circ e^{-\ell\rho} \rangle d\ell,$$

where $R_i f, i = 0, 1$ is obtained from f by Taylor subtraction. The integral converges absolutely since $[e^{-\ell\rho} R_i] = \mathcal{O}_{\mathcal{D}'_\Gamma}(e^{-\ell(1+i)})$ and the map

$$\Theta \mapsto \Theta^+$$

is continuous from $\mathcal{S}'_\Gamma(\mathcal{X} \setminus \mathcal{Y})$ to $\mathcal{S}'_{\Gamma \cup N^*(\mathcal{Y})}(\mathcal{X})$ for all $a' < a$ as we will see below when we check the weak homogeneity of the extension Θ^+ . This shows that when

$$-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1 < a \leq -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$$

one can take

$$\Lambda_{\mathbb{R}^{d_1}, \varepsilon}(f) = \int_0^\infty e^{-\ell(d_2+2d_3)} \langle \Theta_\varepsilon, \bar{\chi} \circ e^{-\ell\rho} \Pi(f) \rangle d\ell$$

and when

$$a = -\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1$$

one can take

$$\Lambda_{\mathbb{R}^{d_1}, \varepsilon}(f) = \int_0^\infty e^{-\ell(d_2+2d_3)} \langle \Theta_\varepsilon, \bar{\chi} \circ e^{-\ell\rho} \Pi(f) + t(\partial_t f)(\cdot, 0, 0) + y \cdot (\partial_y f)(\cdot, 0, 0) \rangle d\ell.$$

For simplicity, in the remainder of the proof we shall specialize to the case $-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1 < a \leq -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$. By the wave front set condition on Θ_ε one can always decompose $\Lambda_{\mathbb{R}^{d_1}, \varepsilon}$ under the product form

$$\Lambda_{\mathbb{R}^{d_1}, \varepsilon} = c_\varepsilon \Lambda_{\mathbb{R}^{d_1}}$$

where

$$\Lambda_{\mathbb{R}^{d_1}} = \Pi$$

is a distribution independent of ε , supported on \mathbb{R}^{d_1} , with wavefront set contained in $N^*(\mathbb{R}^{d_1})$, and the function c_ε is given by

$$c_\varepsilon(x) = \int_0^\infty e^{-\ell(d_2+2d_3)} \Theta_\varepsilon(x, y, t) \bar{\chi}(e^{-\ell\rho}(x, y, t)) d\ell dy dt.$$

To check the weak homogeneity bound write

$$e^{\ell' a} \langle e^{-\ell' \rho} \Theta^+, f \rangle = \int_0^\infty \langle \Theta^{(\ell)} \bar{\chi}, e^{-(\ell-\ell')\rho} \varphi - \Pi(\varphi) \rangle d\ell$$

and observe that the support of $e^{-(\ell-\ell')\rho} \varphi$ meets the support of $\bar{\chi}$ only if $\ell \geq C + \ell'$ for a constant C that depends only on the support of $\bar{\chi}$. So the integral can be split in

$$- \int_0^{C+\ell'} \langle \Theta^{(\ell)} \bar{\chi}, \Pi(\varphi) \rangle d\ell + \int_{C+\ell'}^\infty \langle \Theta^{(\ell)} \bar{\chi}, e^{-(\ell-\ell')\rho} \varphi - \Pi(\varphi) \rangle d\ell$$

A change of variable shows that the second term is uniformly bounded in ℓ' whereas the first term is bounded above by $(C + \ell') \|\varphi\|_{C^0}$. This concludes the proof that $e^{\ell' a} e^{-\ell' \rho} \bar{U} = \mathcal{O}_{\mathcal{D}'}(a)$. \triangleright

In the case (b), note that the extension is no longer unique. Any two extensions differ by some conormal distribution supported on \mathcal{Y} whose order is 0 if $-\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1 < a \leq -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$ and of order 1 if $a = -\text{codim}_w(\mathcal{Y} \subset \mathcal{X}) - 1$.

3.3.2 Two step canonical extensions. We need the following refinement of the canonical extension. It involves a two step extension procedure.

Theorem 14 – *Let $\mathcal{Y} \subset \mathcal{X}$ be a closed embedded submanifold in the ambient smooth manifold \mathcal{X} , $p \in \mathcal{Y}$ a point in \mathcal{Y} , a_1, a_2 some real numbers. Assume we are given a smooth function χ such that $\chi = 1$ near \mathcal{Y} and $\chi = 0$ outside some larger neighbourhood. Set*

$$\Omega \stackrel{\text{def}}{=} \left\{ m \in \mathcal{X} : \text{dist}(m, p) > 2\text{dist}(m, \mathcal{Y}) \right\}.$$

Let Θ be a distribution in $\mathcal{D}'(\mathcal{X} \setminus \mathcal{Y})$ such that there exists a scaling field ρ_1 with respect to $\{p\}$ and a scaling field ρ_2 with respect to \mathcal{Y} , such that both flows of ρ_1 and ρ_2 preserve the inclusion $p \in \mathcal{Y}$. Furthermore assume that

$$\begin{aligned} & - \text{for every test function } \varphi \in C_c^\infty(\Omega \setminus \mathcal{Y}), \\ & \quad e^{\ell_2 a_2 + \ell_1 a_1} \langle e^{-\ell_2 \rho_2^*} e^{-\ell_1 \rho_1^*} (\chi \Theta), \varphi \rangle \quad (\ell_1, \ell_2 \geq 0) \end{aligned}$$

is bounded;

$$\begin{aligned} & - \text{for every test function } \varphi \in C_c^\infty(\mathcal{X} \setminus \mathcal{Y}), \\ & \quad e^{\ell_1 a_1} \langle e^{-\ell_1 \rho_1^*} (\chi \Theta), \varphi \rangle \quad (\ell_1 \geq 0) \end{aligned}$$

is bounded.

If

$$\begin{aligned} a_1 &> -\text{dim}_w(\mathcal{X}), \\ a_2 &> -\text{codim}_w(\mathcal{Y} \subset \mathcal{X}), \end{aligned}$$

then there exists a canonical extension of Θ in $\mathcal{D}'(\mathcal{X})$.

Note that we no longer need any microlocal control in the above statements, only weak topology statements. Let us emphasize that our double scaling assumption only applies when our test function is in the domain $\Omega \setminus \mathcal{Y}$ whereas the second simple scaling assumption applies only when we scale toward the point p . In the Feynman amplitude context of Section 4.1 both scaling fields will be admissible in the sense of definition 16 hence the fact that the inclusion $p \in \mathcal{Y}$ is preserved by both dynamics will be immediate.

Proof – We reduce the proof to some normal form. We cover the whole of \mathcal{X} by some locally finite open cover, use charts and a subordinated partition of unity $\sum \chi_i = 1$. It suffices to prove the same claim for $\Theta \chi$ where χ supported in some open chart $\kappa : \mathcal{X} \supset U \rightarrow \mathbb{R}^k$ containing p , where $\kappa(\mathcal{Y} \cap U) \subset \mathbb{R}^{d_1+d_2} \times \{0, 0\}$ and $\kappa(p) = 0 \in \mathbb{R}^k$. So we are reduced to study the distribution $\kappa_* (\Theta \chi)$ which is compactly supported and satisfies the assumption of our Theorem on $\mathcal{X} = \mathbb{R}^k$ with $k = d_1 + d_2 + d_3 + d_4$ with coordinates (x, t, y, s) such that

$$\begin{aligned} \{p\} &= \{x = 0, t = 0, y = 0, s = 0\}, \quad \text{and} \quad \mathcal{Y} = \{y = 0, s = 0\} = \mathbb{R}^{d_1+d_2} \times \{0\}, \\ \Omega &= \left\{ \sqrt{|x|^2 + |t|^2 + |y|^2 + |s|^2} > 2\sqrt{|y|^2 + |s|^2} \right\}. \end{aligned}$$

Near p we have from the Linearization Theorem proved in [25, Prop 2.3] or [63] the identities

$$e^{-\ell E_1} = e^{-\ell \rho_1} \circ U_1(\ell), \quad \text{and} \quad e^{-\ell E_2} = e^{-\ell \rho_2} \circ U_2(\ell)$$

where E_1, E_2 are the linear scaling fields reading

$$\begin{aligned} E_1 &= 2t \cdot \partial_t + x \cdot \partial_x + 2s \cdot \partial_s + y \cdot \partial_y, \\ E_2 &= 2s \cdot \partial_s + y \cdot \partial_y, \end{aligned}$$

and $U_1(\ell)$ and $U_2(\ell)$ are two smooth germs of diffeomorphisms near p depending smoothly on $e^{-\ell}$ when $\ell \uparrow \infty$, and both U_1, U_2 have limits when $\ell \uparrow \infty$ that also are smooth germs of diffeomorphisms

near $p \in \kappa(U)$. Then, for every test function $\varphi \in C_c^\infty(\Omega \setminus \mathcal{Y})$, we have

$$\begin{aligned} \langle e^{-\ell E_1^*} e^{-r E_2^*}(\Theta\chi), \varphi \rangle &= \langle U_1(\ell) e^{-\ell \rho_1^*} e^{-r E_2^*}(\Theta\chi), \varphi \rangle \\ &= \langle e^{-\ell \rho_1^*} e^{-r E_2^*}(\Theta\chi), U_1(\ell)^{-1*} \varphi \rangle \\ &= \langle e^{-\ell \rho_1^*} U_2(r)^* e^{-r \rho_2^*}(\Theta\chi), U_1(\ell)^{-1*} \varphi \rangle \\ &= \langle (e^{-\ell \rho_1^*} U_2(r)^* e^{u \rho_1^*}) e^{-\ell \rho_1^*} e^{-r \rho_2^*}(\Theta\chi), U_1(\ell)^{-1*} \varphi \rangle \\ &= \langle e^{-\ell \rho_1^*} e^{-r \rho_2^*}(\Theta\chi), \Psi_{\ell,r}^{-1*} U_1(\ell)^{-1*} \varphi \rangle, \end{aligned}$$

where

$$\Psi_{\ell,r} \stackrel{\text{def}}{=} e^{-\ell \rho_1^*} U_2(r)^* e^{\ell \rho_1^*}.$$

At this stage we make the observation that the families $(\Psi_{u,r}^{-1*} U_1(u)^{-1*} \varphi)_{u \geq 0, r \geq 0}$ are bounded families of test functions since $(\Psi_{u,r})_{u \geq 0, r \geq 0}$ is a bounded family of smooth germs of diffeomorphisms. By the Banach-Steinhaus Theorem the family of distributions

$$(e^{\ell a_1 + r a_2} e^{-\ell \rho_1^*} e^{-r \rho_2^*} \Theta\chi)_{\ell, r \geq 0}$$

is weakly, hence strongly, bounded. From this we deduce that

$$e^{\ell a_1 + r a_2} \langle e^{-\ell \rho_1^*} e^{-r \rho_2^*} \Theta\chi, \Psi_{\ell,r}^{-1*} U_1(\ell)^{-1*} \varphi \rangle$$

is bounded for all test functions $\varphi \in C_c^\infty(\Omega \setminus \mathcal{Y})$. Therefore the family $e^{\ell a_1 + r a_2} e^{-\ell E_1^*} e^{-r E_2^*}(\Theta\chi)$ is also bounded in $\mathcal{D}'(\Omega \setminus \mathcal{Y})$. From the Linearization Theorem, without loss of generality, we may thus choose some linear scaling fields ρ_1 and ρ_2 with respect to $\{p\}$ and \mathcal{Y} , respectively, that read

$$\begin{aligned} \rho_1 &= 2t \cdot \partial_t + x \cdot \partial_x + 2s \cdot \partial_s + y \cdot \partial_y, \\ \rho_2 &= 2s \cdot \partial_s + y \cdot \partial_y. \end{aligned}$$

Since Θ is a distribution on $\mathcal{D}'(\Omega \setminus \mathcal{Y})$ with scaling degree a_2 with respect to \mathcal{Y} and since $a_2 > -\text{codim}_w(\mathcal{Y} \subset \mathcal{X})$, the extension theorem 13 implies that we can extend Θ to a distribution $\mathcal{R}\Theta$ on Ω , recall that $p \notin \Omega$. The first extension $\mathcal{R}\Theta$ for Θ extending from $\Omega \setminus \mathcal{Y}$ to the larger space Ω satisfies

$$\mathcal{R}\Theta = \Theta(1 - \chi) + \underbrace{\int_0^\infty (e^{r \rho_2^*} \psi) \Theta dr}_{\text{integral}}$$

for every χ such that $\chi = 1$ near \mathcal{Y} , $\chi = 0$ outside some larger neighbourhood (recall that here $\psi = -\rho_2 \chi$) where the integral underbraced converges in $\mathcal{D}'(\Omega)$ by the assumption on the scaling degree of Θ with respect to \mathcal{Y} . Indeed for any test function $\varphi \in C_c^\infty(\Omega)$ one has

$$\left\langle \int_0^\infty (e^{r \rho_2^*} \psi) \Theta dr, \varphi \right\rangle = \int_0^\infty e^{-r(d_3 + 2d_4 + a_2)} \langle (e^{r a_2} e^{-r \rho_2^*} \Theta), \psi e^{-r \rho_2^*} \varphi \rangle dr$$

where it is immediate that the integrand converges since $d_3 + 2d_4 + a_2 > 0$, the family $(\psi e^{-r \rho_2^*} \varphi)_{r \geq 0}$ forms a bounded family of test functions in $C_c^\infty(\Omega \setminus \mathcal{Y})$ and $(e^{r a_2} e^{-r \rho_2^*} \Theta)_{r \geq 0}$ is a bounded family of distributions in $\mathcal{D}'(\Omega \setminus \mathcal{Y})$. We also have from the identity for $T > 0$

$$1 - \chi + \int_0^T e^{r \rho_2^*} \psi dr = 1 - e^{T \rho_2^*} \chi,$$

and the absolute convergence of the integral $\int_0^\infty (e^{r \rho_2^*} \psi) \Theta dr$, that $\mathcal{R}\Theta$ can be defined as the limit

$$\mathcal{R}\Theta = \lim_{T \uparrow \infty} \Theta(1 - e^{T \rho_2^*} \chi).$$

Here we make the observation that for every isomorphism $f : \mathbb{R}^k \mapsto \mathbb{R}^k$ that stabilizes \mathcal{Y} and commutes with ρ_2 we have the identity

$$\mathcal{R}\Theta = \Theta(1 - f^* \chi) + \underbrace{\int_0^\infty (e^{r \rho_2^*} f^* \psi) \Theta dr}_{\text{integral}}.$$

Since $f^* \chi$ still satisfies the same technical assumptions as χ and $-\rho_2 f^* \chi = -f^* \rho_2 \chi = f^* \psi$, the integral underbraced also converges in $\mathcal{D}'(\Omega)$. The key observation is the continuous partition of

unity identity for $T > 0$

$$1 - \chi + \int_0^T e^{r\rho_2^*} \psi \, dr = 1 - e^{T\rho_2^*} \chi,$$

which implies

$$\begin{aligned} 1 - \chi + \int_0^T e^{r\rho_2^*} \psi \, dr - \left(1 - f^* \chi + \int_0^T e^{r\rho_2^*} f^* \psi \, dr \right) \\ = 1 - e^{T\rho_2^*} \chi - (1 - e^{T\rho_2^*} f^* \chi) = e^{T\rho_2^*} f^* \chi - e^{T\rho_2^*} \chi = \int_T^\infty e^{r\rho_2^*} (\psi - f^* \psi) \, dr, \end{aligned}$$

where the last equality holds true in $C^\infty(\Omega \setminus \mathcal{Y})$. Then, we can control the difference

$$\Theta(1 - e^{T\rho_2^*} \chi) - \Theta(1 - e^{T\rho_2^*} f^* \chi) = \Theta(e^{T\rho_2^*} (f^* \chi - \chi)) = \int_T^\infty e^{r\rho_2^*} (\psi - f^* \psi) \Theta \, dr.$$

Repeating the above estimate, for every test function $\varphi \in C_c^\infty(\Omega \setminus \mathcal{Y})$

$$\left| \left\langle \int_T^\infty e^{r\rho_2^*} (\psi - f^* \psi) \Theta \, dr, \varphi \right\rangle \right| \lesssim \int_T^\infty e^{-r(d_2 + 2d_3 + a_2)} \, dr \longrightarrow 0$$

as $T \uparrow \infty$.

The whole point is that we start from the information that Θ defined on $\Omega \setminus \mathcal{Y}$ is weakly homogeneous with respect to $\{p\}$ of degree a_1 and we need to make sure that the same property still holds true for the first extension $\mathcal{R}\Theta$. Therefore we need to check that the partial extension $\mathcal{R}\Theta \in \mathcal{D}'(\Omega)$ (recall $p \notin \Omega$) is still weakly homogeneous of degree a_1 when we scale with respect to $\{p\}$.

We scale the renormalized $\mathcal{R}\Theta$ using $e^{-t\rho_1}$ which gives

$$\begin{aligned} e^{\ell a_1} e^{-\ell \rho_1^*} \mathcal{R}\Theta &= e^{\ell a_1} e^{-\ell \rho_1^*} \left(\Theta(1 - f^* \chi) + \underbrace{\int_0^\infty (e^{r\rho_2^*} f^* \psi) \Theta \, dr}_{\text{underbraced term}} \right) \\ &= e^{\ell a_1} (e^{-\ell \rho_1^*} \Theta) (1 - \chi) + \underbrace{\int_0^\infty (e^{r\rho_2^*} \psi) (e^{\ell a_1} e^{-\ell \rho_1^*} \Theta) \, dr}_{\text{underbraced term}}, \end{aligned}$$

choosing $f = e^{\ell \rho_1^*}$ and using the fact the flows $e^{-r\rho_2}$, $e^{-\ell \rho_1}$ commute and the absolute convergence of the underbraced term. Then we make the second observation that the family $(e^{\ell a_1} e^{-\ell \rho_1^*} \Theta)_{\ell \geq 0}$ satisfies the assumptions of the extension Theorem 13 applied to the scaling with respect to $\{p\}$ **uniformly** in the parameter $\ell \geq 0$. Using the boundedness of the extension map \mathcal{R} , this concludes that $\mathcal{R}\Theta \in \mathcal{D}'(\Omega)$ is weakly homogeneous of degree a_1 with respect to $\{p\}$.

We can now conclude, choose a partition of unity $1 = \chi_\Omega + (1 - \chi_\Omega)$ where $\chi_\Omega \in C^\infty(\mathbb{R}^k \setminus \{0\})$ is scale invariant with respect to ρ_1 , $\chi_\Omega|_\Omega = 1$ near \mathcal{Y} and $\chi_\Omega = 0$ outside Ω , then note that the two families

$$(e^{\ell a_1} e^{-\ell \rho_1^*} \chi_\Omega \mathcal{R}\Theta)_{\ell \geq 0}$$

and

$$(e^{\ell a_1} e^{-\ell \rho_1^*} (1 - \chi_\Omega) \Theta)_{\ell \geq 0}$$

are bounded in $\mathcal{D}'(\mathbb{R}^k \setminus \{0\})$. Hence applying the extension Theorem 13 to the sum $\chi_\Omega \mathcal{R}\Theta + (1 - \chi_\Omega) \Theta$ yields that $\Theta = \chi_\Omega \mathcal{R}\Theta + (1 - \chi_\Omega) \Theta \in \mathcal{D}'(\mathbb{R}^k \setminus \{0\})$ has a canonical extension in $\mathcal{D}'(\mathbb{R}^k)$ which is the result we wanted. \triangleright

Examples On \mathbb{R}^2 , fix $p = (p_1, p_2) \in \mathbb{R}^2$ and let

$$\varphi(x_1, x_2) \stackrel{\text{def}}{=} |x_1 - x_2|^{a_2} (|x_1 - p_1| + |x_2 - p_2|)^{(a_1 - a_2)}.$$

It is weakly homogeneous of degree a_2 with respect to the diagonal $\{x_1 = x_2\}$ and of degree a_1 with respect to the point p . Theorem 14 therefore applies to φ provided $a_2 > -1$ and $a_1 > -2$.

3.3.3 Invariance properties of scaling spaces. We end this subsection by stating some invariance results which will be useful later, in the proofs of Theorems 27 and Theorem 14, when we need to make a particular choice of scaling field in the proof and then use the fact that the blow-up does

not depend on this choice. In the next proposition all the scaling fields are relative to the closed embedding $\mathcal{Y} \subset \mathcal{X}$.

Proposition 15 – Assume $a \in \mathbb{R}$, $\mathcal{Y} \subset \mathcal{X}$ and the distribution $\Lambda \in \mathcal{D}'(\mathcal{X})$, resp. $\mathcal{D}'(\mathcal{X} \setminus \mathcal{Y})$, is weakly homogeneous of degree a for some scaling field ρ , in the sense that the family

$$(e^{\ell a} e^{-\ell \rho^*}(\chi \Lambda))_{\ell \geq 0}$$

is bounded in $\mathcal{D}'(\mathcal{X})$, resp. $\mathcal{D}'(\mathcal{X} \setminus \mathcal{Y})$, for some scaling field ρ and some χ such that $\chi = 1$ near \mathcal{Y} and ρ is well-defined near $\text{supp}(\chi)$. Then Λ is weakly homogeneous of degree a for every scaling fields. More precisely, for any scaling field $\tilde{\rho}$ with respect to \mathcal{Y} , every function $\tilde{\chi}$ supported in the domain of $\tilde{\rho}$ such that $\tilde{\chi} = 1$ near \mathcal{Y} ,

$$(3.8) \quad \text{the family of distributions } (e^{t a} e^{-t \tilde{\rho}^*}(\tilde{\chi} \Lambda))_{t \geq 0} \text{ is bounded in } \mathcal{D}'(\mathcal{X}), \text{ resp. } \mathcal{D}'(\mathcal{X} \setminus \mathcal{Y}).$$

In the sequel, we shall denote by $\mathcal{S}^a(\mathcal{X})$ (resp. $\mathcal{S}^a(\mathcal{X} \setminus \mathcal{Y})$) the space of distributions such that (3.8) holds. Consequently, assume we are given an open subset $\mathcal{U} \subset \mathcal{X}$ such that \mathcal{U} is stable by ρ_1 and ρ_2 . Then every $\Lambda \in \mathcal{D}'(\mathcal{U})$ which is weakly homogeneous of degree a for ρ_1 is also weakly homogeneous of the same degree a for ρ_2 . In the sequel, we shall denote by $\mathcal{S}^a(\mathcal{U})$ the space of such distributions.

We note that the space $\mathcal{S}^a(\mathcal{X})$, resp. $\mathcal{S}^a(\mathcal{X} \setminus \mathcal{Y})$, is intrinsic, defined independently from the choice of scaling fields. These spaces do not contain microlocal informations hence the topologies of the spaces \mathcal{S}^a are weaker than the topologies of spaces \mathcal{S}_Γ^a .

Proof – The proof can be found in [25, Thm 3.3 p. 828] and relies on the linearization Theorem for scaling fields proved in [25] and also in the beautiful exposition of Meinrenken [63, Lemma 2.1 p. 226]. We sketch a proof for completeness. We localize near $x \in \mathcal{Y}$, we choose some chart $\kappa : U \ni x \mapsto \mathbb{R}^{d_1+d_2+d_3}$ in which $\kappa(\mathcal{Y} \cap U) = (\mathbb{R}^{d_1} \times \{0\}) \cap \kappa(U)$. Then once we push ρ by the linear chart κ we still get a scaling field that we abusively denote by ρ . In these coordinates the vector field ρ has the form $\rho = 2t \cdot \partial_t + y \cdot \partial_y + R(t, y, \partial_t) + H(t, y, \partial_y)$ where R and H vanish at order 2 when (t, y) go to $(0, 0) \in \mathbb{R}^{d_2+d_3}$. Choose χ supported in the open chart U , $\kappa_*(\Lambda \chi)$ is a distribution in $\mathbb{R}^{d_1+d_2+d_3}$ weakly homogeneous of degree a under scaling by ρ . With no loss of generality it suffices to prove the weak homogeneity of $\kappa_*(\Lambda \chi)$ when we scale with a different scaling field ρ_2 , the general claim can be deduced by localising plus gluing with a partition of unity without problem. Then the linearization proof [25, Prop 2.3 p. 826] tells us that we have an equation of the form

$$e^{-\ell \rho} \circ U(\ell) = e^{-\ell \rho_2}$$

where $U(\ell)$ is a family of diffeomorphism germs near $(x, 0, 0)$ which has a limit when $\ell \uparrow \infty$, the limit is still a diffeomorphism germ. So for every test function $\varphi \in C_c^\infty(\kappa(U))$ one has

$$e^{\ell a} \langle e^{-\ell \rho_2^*} \kappa_*(\Lambda \chi), \varphi \rangle = e^{\ell a} \langle U(\ell)^* e^{-\ell \rho^*} \kappa_*(\Lambda \chi), \varphi \rangle = e^{\ell a} \langle e^{-\ell \rho^*} \kappa_*(\Lambda \chi), U(\ell)^{-1*} \varphi \rangle,$$

where the last quantity on the right hand side is bounded when $\ell \geq 0$ since $(U(\ell)^{-1*} \varphi)_{\ell \geq 0}$ is a bounded family of test functions and $(e^{\ell a} e^{-\ell \rho^*} \kappa_*(\Lambda \chi))_{\ell \geq 0}$ is a bounded family of distributions by assumption. Then we just proved that $(e^{\ell a} e^{-\ell \rho_2^*} \kappa_*(\Lambda \chi))_{\ell \geq 0}$ is a weakly, hence strongly, bounded family of distributions which concludes the proof. \triangleright

4 – Induction on Feynman amplitudes

In the present section we describe a general induction to control analytically the Feynman amplitudes which appear when we study the regularities of the stochastic trees appearing in our equation. This is the content of the main Theorem 27. To state this Theorem precisely, we need to introduce several layers of formalism. We first describe correctly the configuration spaces on which we work which is the content of Subsection 4.1, then we give the definition of the Feynman graphs in subsection 4.2. There is a subtlety in the stochastic estimates, since we work in the non stationary setting (our manifold M is not a flat torus or \mathbb{S}^3 with the round metric) we need to control the size of the pointwise value of Littlewood–Paley blocks, we refer the reader to Lemma 30 and Definition 31 where we discuss precisely these matters. In terms of Feynman amplitudes, this forces us to introduce a certain distinguished propagator \mathcal{Q}_x^γ in which depends on some point

$x \in M$, and scale the whole amplitude when all points collapse to the given point $x \in M$, we control the size of the amplitude uniformly in $x \in M$. Mathematically, this requires the definition of some pointed scaling space in Subsection 4.2.1. Finally the section concludes with the inductive proof of Theorem 27 which involves a double scaling, where we need to control the growth of the amplitude when all points collide and also when all points collide on a given $x \in M$, uniformly in $x \in M$. This makes the proof much more involved than usual Weinberg convergence like Theorems in usual Quantum Field Theories. It is also because of these double scalings that we need the double extension Theorem 14. For several graphs that appear in our stochastic estimates which are enumerated in the first paragraph in Section 5.4, we need to feed the induction Theorem 27 with more informations, this uses subtle smoothing properties of the Schwartz kernel of the resonant product which is discussed in Subsection 5.4.1 and also some control of certain renormalized subamplitude in Subsection 5.4.2, we refer the reader to the relevant parts for more details but this can be read after one is used to the general inductive machinery.

4.1 Configuration space

The Feynman amplitudes which arise from the stochastic estimates involve the analysis of products of distributional kernels on space-time configuration spaces. In the present section we introduce the necessary general formalism for analysing these amplitudes.

4.1.1 Diagonals and scaling fields. In the sequel we write

$$\mathcal{M} \stackrel{\text{def}}{=} \mathbb{R} \times M$$

and work on

$$\mathbb{R}^p \times M^{p+q} \simeq \mathcal{M}^p \times M^q$$

for $p, q \geq 0$. Given a fixed $t \in \mathbb{R}$ we always view some elements of $\mathbb{R}^p \times M^{p+q}$ as some elements of \mathcal{M}^{p+q} using the mapping

$$(4.1) \quad \mathcal{M}^p \times M^q \simeq \mathcal{M}^p \times (\{t\} \times M)^q \subset \mathcal{M}^{p+q},$$

that is to say attributing the time t to the purely spatial points. The configuration space \mathcal{M}^n of $n = p + q$ points in \mathcal{M} will play a particular role in the sequel. Given a distinguished point $(t, x) \in \mathcal{M}$, writing $m_i = (t_i, x_i) \in \mathcal{M}$, for $I \subset \{1, \dots, n\}$ we denote by

$$(4.2) \quad \begin{aligned} \mathbf{d}_I &\stackrel{\text{def}}{=} \left\{ m = (m_1, \dots, m_n); x_i = x_j \text{ and } t_i = t_j = t \text{ for } i \neq j \text{ if } (i, j) \in I^2 \right\}, \\ &\text{when space-time points labelled by } I \text{ collide;} \\ \mathbf{d}_{I, (t, x)} &\stackrel{\text{def}}{=} \left\{ m = (m_1, \dots, m_n), m_i = (t, x) \text{ if } i \in I \right\}, \quad \text{the marked diagonal} \\ &\text{where all space-time points labelled by } I \text{ collide to } (t, x), \\ \mathcal{T}_I &\stackrel{\text{def}}{=} \left\{ m = (m_1, \dots, m_n), t_i = t_j = t \text{ for } i \neq j \text{ if } (i, j) \in I^2 \right\}, \\ &\text{when only time components labelled by } I \text{ coincide with } t \end{aligned}$$

the corresponding diagonals in the product spaces. Let us make several important observations. First the marked diagonal $\mathbf{d}_{[n], (t, x)}$ is in reality just **one point** $((t, x), \dots, (t, x))$ in the configuration space \mathcal{M}^n . Second one has the natural inclusion relations

$$\mathbf{d}_I \subset \mathcal{T}_I \text{ and } \mathbf{d}_I \subset \mathbf{d}_J \text{ and } \mathcal{T}_I \subset \mathcal{T}_J \text{ if } J \subset I.$$

We denote by

$$\mathbf{d}_n \stackrel{\text{def}}{=} \left\{ ((t, x_1), \dots, (t, x_1)) \in \mathcal{M}^n : x_1 \in M \right\},$$

and

$$\mathbf{d}_{n, (t, x)} \stackrel{\text{def}}{=} \left\{ ((t, x), \dots, (t, x)) \right\} \subset \mathcal{M}^n.$$

In the sequel, to take (4.1) into account, we will work on some submanifold $\mathcal{M}^p \times (\{t\} \times M)^q = \mathcal{T}_J \subset \mathcal{M}^n$ for some fixed $J \subset \{1, \dots, n\}$, $|J| = q$, where all time variables indexed by J coincide and equal t .

Examples We give two examples of stochastic estimates we will meet in the sequel so that the reader can see in what type of space-time domains we need to integrate our Feynman amplitudes.

In the sequel we will calculate $\mathbb{E}[\mathcal{Q}_x^\gamma(\varphi^\circ(t), \varphi^\circ(t))]$ which can be represented as a Feynman amplitude (underbraced below) tested against the constant function 1 that reads

$$\begin{aligned} & \mathbb{E}[\mathcal{Q}_x^\gamma(\varphi^\circ(t), \varphi^\circ(t))] \\ &= \int_{(-\infty, t]^2 \times M^4} \underbrace{G^{(3)}(s_a - s_b, x_a, x_b) \underline{\mathcal{L}}^{-1}(t - s_a, y_*, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_*, x_b) \mathcal{Q}_x^\gamma(y_*, z_*)}_{\text{Feynman amplitude}} \end{aligned}$$

for some function $G^{(3)}$. The integration is with respect to $dv(x_a, x_b, y_*, z_*) ds_a ds_b$. In this case we integrate a certain Feynman amplitude underbraced on some domain $\mathcal{T}_{\{c, d\}} \subset \mathcal{M}^4$ with two space-time variables (s_a, x_a) , (s_b, x_b) and two space variable (s_c, y_*) , (s_d, z_*) promoted to space-time variables by writing $s_c = s_d = t$. On the other hand, when we want to prove the continuity in time using a Kolmogorov type argument we will also consider

$$\begin{aligned} & \mathbb{E}[\mathcal{Q}_x^\gamma(\varphi^\circ(t_1), \varphi^\circ(t_2))] \\ &= \int_{(-\infty, t]^2 \times M^4} \underbrace{G^{(3)}(s_a - s_b, x_a, x_b) \underline{\mathcal{L}}^{-1}(t_1 - s_a, y_*, x_a) \underline{\mathcal{L}}^{-1}(t_2 - s_a, z_*, x_b) \mathcal{Q}_x^\gamma(y_*, z_*)}_{\text{Feynman amplitude}}. \end{aligned}$$

In this case we integrate over $\mathcal{T}_{\{c, d\}} \subset \mathcal{M}^4$ where the last two time variables are taken equal to some fixed times t_1, t_2 . With this example we see that we also need to fix the time variables.

We will also use a **particular class of scaling fields that will leave all \mathcal{T}_J and \mathbf{d}_I stable**, such class of scaling fields will be called **admissible**.

Definition 16 – (Admissible scaling fields) *Pick some open chart $U \subset M, \kappa : U \mapsto \mathbb{R}^d$ such that $\kappa(U) \subset \mathbb{R}^d$ is an open convex ball – this is always possible up to making things smaller.*

We define a scaling field $\rho_{[p]}$ in U from its flow given by for any $(x_1, \dots, x_p) \in \kappa(U)^p$ by

$$(x_1, \dots, x_p) \in U^p \mapsto \left(x_1, e^{-t}(x_2 - x_1) + x_1, \dots, e^{-t}(x_p - x_1) + x_1 \right) \in \kappa(U)^p;$$

it means $\rho_{[p]}$ reads $\sum_{j=2}^p (x_j - x_1) \partial_{x_j}$ in the above coordinate chart. We define the local scaling field on $(\mathbb{R} \times U)^p$, with local coordinates $(s_i, x_i)_{1 \leq i \leq p}$, setting

$$\rho = 2 \left(\sum_{j=2}^p (s_j - s_1) \partial_{s_j} \right) + \rho_{[p]} \stackrel{\text{def}}{=} 2\rho_{\text{times}} + \rho_{[p]}.$$

We obtain global scaling fields by gluing together the above local objects. Consider a cover $\cup_i U_i^p$ of some neighborhood of the space diagonal and choose $\chi \in C_c^\infty(\cup_i U_i^p)$ such that $\chi = 1$ near the space diagonal. For a subordinated partition of unity $\sum \chi_i = 1$ of $\text{supp}(\chi)$, we set $\rho = 2\rho_{\text{times}} + \chi \sum_i \chi_i \rho_i$ where each local scaling field $\rho_i \in C^\infty(T(U_i^p))$ is constructed in some charts as above.

The same construction also works when we scale towards some marked points, in all charts $\kappa : U \mapsto M$ containing x , we decide that we scale with the flow

$$(x_1, \dots, x_p) \in U^p \mapsto \left(e^{-t}(x_1 - x) + x, e^{-t}(x_2 - x) + x, \dots, e^{-t}(x_p - x) + x \right) \in \kappa(U)^p,$$

it means $\rho_{[p]}$ reads $\sum_{j=1}^p (x_j - x) \partial_{x_j}$ in the above coordinate chart. If the chart does not contain x , we decide our flow is trivial (the generator of the flow is the zero vector field) and $\rho_{[p]} = 0$ in such a chart. We define the local scaling field on $(\mathbb{R} \times U)^p$, with local coordinates $(s_i, x_i)_{1 \leq i \leq p}$, setting

$$\rho_{(t, x)} = 2 \left(\sum_{j=1}^p (s_j - t) \partial_{s_j} \right) + \rho_{[p]} \stackrel{\text{def}}{=} 2\rho_{\text{times}} + \rho_{[p]}.$$

We obtain global scaling fields by gluing together the above local objects. Consider a cover $\cup_i U_i^p$ of some neighborhood of the space diagonal and choose $\chi \in C_c^\infty(\cup_i U_i^p)$ such that $\chi = 1$ near the space diagonal. For a subordinated partition of unity $\sum \chi_i = 1$ of $\text{supp}(\chi)$, we set $\rho = 2\rho_{\text{times}} + \chi \sum_i \chi_i \rho_i$ where each local scaling field $\rho_i \in C^\infty(T(U_i^p))$ is constructed in charts as above.

We make the observation that in all the above situations the constructed vector fields generate some dynamics which preserve all the diagonals $\mathbf{d}_I, \mathcal{T}_I$ or all the marked diagonals $\mathbf{d}_{I, (t, x)}$ in the case of scaling fields $\rho_{(t, x)}$ on marked points **by construction** (they preserve these diagonals in the charts locally, hence globally by gluing). The above definition shows that *admissible* scaling

fields are abundant. Admissible scaling fields enjoy another remarkable property. The cotangent lift of $e^{-u\rho}$ stabilizes all the conormal bundles of all the partial diagonals.

Lemma 17 – For all $I \subset \{1, \dots, p\}$ and $\ell \geq 0$, for all admissible scaling fields ρ , one has

$$e^{-\ell\rho^*} (N^*(\mathbf{d}_I \subset \mathcal{M}^p)) \subset N^*(\mathbf{d}_I \subset \mathcal{M}^p).$$

Proof – If $v \in T_x \mathbf{d}_I$, $de_x^{\ell\rho}(v) \in T_{e^{\ell\rho}(x)} \mathbf{d}_I$ (since the flow stabilizes the diagonals) implies that if $\xi \in T_{e^{\ell\rho}(x)} \mathbf{d}_I^\perp$ then $\langle \xi, de_x^{\ell\rho}(v) \rangle = 0$ for all $v \in T_x \mathbf{d}_I$ hence ${}^t de_x^{\ell\rho}(v)(\xi) \in T_x \mathbf{d}_I^\perp = N_x^*(\mathbf{d}_I)$, which concludes our proof. \triangleright

We will typically be given a family $(\Lambda_\varepsilon)_{0 < \varepsilon \leq 1}$ of distributions on $\mathcal{T}_J \setminus (\cup_{I \subset \{1, \dots, p\}} \mathbf{d}_I)$ that converge to a limit as a distribution outside all the diagonals of \mathcal{T}_J . We will use Theorem 13 to extend it to the whole of \mathcal{T}_J by an inductive procedure under some scaling-type assumptions. The inductive structure of the extension procedure will come from the geometric form of *Popineau & Stora's lemma*, which we recall here. We associate to $I \subset \{1, \dots, p\}$, the open set:

$$\mathcal{O}_I \stackrel{\text{def}}{=} \left\{ m = (m_1, \dots, m_p) \in \mathcal{M}^p ; m_i \neq m_j \ \forall (i, j) \in I \times I^c \right\} \subset \mathcal{M}^p.$$

Lemma 18 – One has

$$\mathcal{M}^p \setminus \mathbf{d}_p = \bigcup_{I \subset \{1, \dots, p\}} \mathcal{O}_I$$

and there is an associated smooth partition of the unity, $1 = \sum_{I \subset \{1, \dots, p\}} \eta_I \in C^\infty(\mathcal{M}^p \setminus \mathbf{d}_p)$, with the family $(\eta_I \circ e^{-\ell\rho})_{\ell \geq 0}$ bounded in $C^\infty(\mathcal{M}^p \setminus \mathbf{d}_p)$ for every admissible scaling field ρ with respect to the deepest diagonal $\mathbf{d}_p \subset \mathcal{M}^p$.

The proof is simple and can be found in [26, Lemma 6.3]. The proof of the claim on the family $(\eta_I \circ e^{-\ell\rho})_{\ell \geq 0}$ can be found in [27, Lemma 6.3.1 p. 131]. Since $\mathcal{T}_J \subset \mathcal{M}^p$ for $J \subset \{1, \dots, p\}$, the above partition of unity induces naturally a partition of unity on $\mathcal{T}_J \setminus \mathbf{d}_p$ with the same properties.

4.1.2 Hörmander product of distributions. In the simplest cases the distributions Λ_ε will be given as some products of distributions, with each factor depending possibly only on a subset of the variables \mathcal{M}^p . The easiest case in which to make sense of such products relies on Hörmander's product theorem [18, Thm 6.1 p. 219] and gives the following statement.

Lemma 19 – If $\Lambda_1 \in \mathcal{D}'(\mathcal{M}^p)$ depends only on the first $1 \leq k < p$ components of \mathcal{M}^p and $\Lambda_2 \in \mathcal{D}'(\mathcal{M}^p)$ depends only on the last $p - (k - 1)$ components, so they have only one component in common, and

$$\begin{aligned} WF(\Lambda_1) &\subset \bigcup_{I \subset \{1, \dots, k\}} N^*(\mathbf{d}_I) \cup N^*(\mathcal{T}_I), \\ WF(\Lambda_2) &\subset \bigcup_{J \subset \{k, \dots, p\}} N^*(\mathbf{d}_J) \cup N^*(\mathcal{T}_J), \end{aligned}$$

then the product $\Lambda_1 \Lambda_2$ is well-defined in $\mathcal{D}'(\mathcal{M}^p)$ and

$$WF(\Lambda_1 \Lambda_2) \subset (WF(\Lambda_1) + WF(\Lambda_2)) \cup WF(\Lambda_1) \cup WF(\Lambda_2).$$

Proof – Denote by λ a generic element of $T^*\mathcal{M}$. If $(\lambda_1, \dots, \lambda_k, 0, \dots, 0)$ and $(0, \dots, 0, \mu_k, \mu_{k+1}, \dots, \mu_p)$ stand for some non-null elements of $T^*(\mathcal{M}^p)$ such that

$$\sum \lambda_i = 0, \quad \sum \mu_j = 0,$$

then the convex sum

$$(\lambda_1, \dots, \lambda_k, 0, \dots, 0) + (0, \dots, 0, \mu_k, \mu_{k+1}, \dots, \mu_p) = (\lambda_1, \dots, \lambda_k + \mu_k, \mu_{k+1}, \dots, \mu_p)$$

cannot vanish. This implies that $WF(\Lambda_1) + WF(\Lambda_2)$ does not meet the zero section $\{0\}$ and one can apply Hörmander's Theorem [18, Thm 6.1 p. 219] which yields the existence of the distributional product $\Lambda_1 \Lambda_2$ together with a bound on the wave front set $WF(\Lambda_1 \Lambda_2)$ of the product. \triangleright

We give another important consequence of Theorem 13 before talking about Feynman amplitudes.

Proposition 20 – Let $\mathcal{Y} \subset \mathcal{X}$ be a closed embedding and let ρ stand for a parabolic scaling field for the inclusion $\mathcal{Y} \subset \mathcal{X}$. Assume we are given some closed conic sets Γ_1, Γ_2 in $T^*\mathcal{X} \setminus \mathcal{Y}$ such that

$$\mathbf{0} \notin (\Gamma_1 + \Gamma_2).$$

and such that $e^{-\ell\rho^*}(\Gamma_i) \subset \Gamma_i, \forall \ell \geq 0$ and $i \in \{1, 2\}$. Assume also that we are given two distributions

$$\Lambda_1 \in \mathcal{S}_{\Gamma_1}^{s_1, \rho}(\mathcal{X} \setminus \mathcal{Y}) \cap \mathcal{D}'(\mathcal{X}), \quad \Lambda_2 \in \mathcal{S}_{\Gamma_2}^{s_2, \rho}(\mathcal{X}) \cap \mathcal{D}'(\mathcal{X})$$

so the product $\Lambda_1 \Lambda_2$ is well-defined on $\mathcal{U} \subset \mathcal{X} \setminus \mathcal{Y}$. If

$$s_1 + s_2 > -\text{codim}_w(\mathcal{Y})$$

then this product has a unique extension as an element of $\mathcal{S}_{\Gamma}^{s_1+s_2, \rho}(\mathcal{X})$ with

$$\Gamma = \Gamma_1 \cup \Gamma_2 \cup (\Gamma_1 + \Gamma_2) \cup N^*(\mathcal{Y} \subset \mathcal{X}).$$

The condition $\mathbf{0} \notin (\Gamma_1 + \Gamma_2)$ ensures that the distributional product $\Lambda_1 \Lambda_2$ is well-defined at least on $\mathcal{X} \setminus \mathcal{Y}$ and the stability of Γ_1, Γ_2 under the lifted cotangent flow ensures that the convex sum $(\Gamma_1 + \Gamma_2) \subset T^*\mathcal{X} \setminus \mathcal{Y}$ satisfies the conormal landing condition for the inclusion $\mathcal{Y} \subset \mathcal{X}$. The statement of the above proposition means that for any mollification $\Lambda_1^\varepsilon, \Lambda_2^\varepsilon$ of these distributions which converge in the respective functional spaces the product $\Lambda_1^\varepsilon \Lambda_2^\varepsilon$ is converging in $\mathcal{S}_{\Gamma}^{s_1+s_2, \rho}(\mathcal{M}^p)$ to a limit independent of the mollification.

4.2 Feynman graphs and Feynman amplitudes

We are now ready to define the Feynman graphs we are using in order to control the Hölder-Besov norms of the tree appearing in the construction of the Φ_3^4 measure. For an oriented finite graph (V, E) with vertex set V and edge set E we denote by $v(e)_-, v(e)_+$ its two vertices, according to its orientation.

Definition – A **Feynman graph for Φ_3^4** is an oriented finite graph $\mathcal{G}_{(t,x)} = (V, E)$ with p vertices in V and an edge set E with no two edges joining a given pair of vertices, along with

- a distinguished edge $e_{\text{ref}} \in E$,
- a subset J of V which indicates which times are set equal to the fixed $t \in \mathbb{R}$,
- for each vertex $v \in V$, a variable $z_v = (t_v, x_v) \in (\mathbb{R} \times M)$, with the restriction that for the two vertices $v(e_{\text{ref}})_-, v(e_{\text{ref}})_+$ attached to e_{ref} , one has $z_{v(e_{\text{ref}})_\pm} = (t, x_{v(e_{\text{ref}})_\pm})$ for some $x_{v(e_{\text{ref}})_\pm} \in M$.

We set

$$\tilde{E} \stackrel{\text{def}}{=} E \setminus \{e_{\text{ref}}\}.$$

We furthermore assume that the following facts hold.

- The set V of vertices can be partitioned as

$$V = V' \sqcup V_A$$

where V' is a disjoint union of singletons and V_A is a disjoint union of

$$n_{\mathcal{G}} \stackrel{\text{def}}{=} |A|$$

triples of vertices indexed by a finite set A , with each triple made up of a distinguished pair of vertices and another vertex. For $j \in [n_{\mathcal{G}}]$ such a triple in V_A reads $(v_*^j, (v_1^j, v_2^j))$ where (v_1^j, v_2^j) is the distinguished pair and v_*^j the remaining vertex.

- For every $j \in [n_{\mathcal{G}}]$ there is no edge in the graph relating v_1^j to v_2^j or one of these points to v_*^j .
- We are given for each edge $e \in E$ a kernel $K_e \in \mathcal{K}^{a_e}$ for some scaling exponent $a_e \in \mathbb{R}$. (The space \mathcal{K}^{a_e} was defined in Lemma 12). Moreover for the distinguished edge $e_{\text{ref}} \in E$ the corresponding kernel is given by $K_{e_{\text{ref}}} = \mathcal{Q}_x^\gamma$ and e_{ref} is the only edge in E whose kernel K_e is of the form \mathcal{Q}_y^γ for some $y \in M$.

See figure 4.2 below for an illustration. In the sequel we often omit the base point (t, x) , writing \mathcal{G} instead of $\mathcal{G}_{(t,x)}$. Since renormalization also involves the analysis of singularities of Feynman subgraphs we also need a notion of Feynman subgraphs adapted to our specific setting.

Definition – A **Feynman subgraph** $\mathcal{G}_1 = (V_1, E_1) \subset \mathcal{G}_{(t,x)} = (V, E)$ is the data of

- some subset V_1 of the vertices V of \mathcal{G} ,
- some subset E_1 of the edges E of \mathcal{G} such that V_1 can be partitioned as $V_1 = V'_1 \sqcup V'_A$ and $V'_A \subset V_A$ respects the partitioning of V_A , any triple $(v_*^j, (v_1^j, v_2^j))$, $j \in n_{\mathcal{G}_1}$ in V'_A corresponds to a triple in V_A ;

along with the conditions that every edge $e \in E_1$ has its bounding vertices $v(e)_-, v(e)_+$ in V_1 , and that the subgraph \mathcal{G}_1 does not necessarily contain the distinguished edge e_{ref} of \mathcal{G} . A (sub)graph \mathcal{G} is said to be irreducible if it cannot be disconnected by removing exactly one edge $e \in E$.

We also need to recall the notion of loops for the Feynman graphs we consider, this is given by the usual Euler formula.

Definition – Given a Feynman graph \mathcal{G} whose rule is defined above we define the **number of loops for our exotic Feynman graph \mathcal{G}** as

$$b_1(\mathcal{G}) \stackrel{\text{def}}{=} |E(\mathcal{G})| - |V'(\mathcal{G})| - n_{\mathcal{G}} + 1.$$

We are given some distributions

$$\begin{aligned} [\odot](\cdot, \cdot, \cdot) &\in \mathcal{S}_{N^*(\mathbf{d}_3)}^{-6}(M^3), \\ \mathcal{Q}_x^\gamma &\in \mathcal{S}_{T^*(x,x)M^2}^{-6-2\gamma}(M^2), \end{aligned}$$

where the scaling of $[\odot]$ is with respect to the deepest diagonal $\mathbf{d}_3 = \{(y, y, y) : y \in M\} \subset M^3$ and the scaling of \mathcal{Q}_x^γ is with respect to the point (x, x) . Here $[\odot]$ is a general notation for the kernel $[\odot_i]$ of the resonant product in some chart $i \in I$ which is defined by

$$[\odot_i](x, y, z) \stackrel{\text{def}}{=} \sum_{|k-\ell| \leq 1} P_k^i(x, y) \tilde{P}_\ell^i(x, z),$$

where P_k^i, \tilde{P}_ℓ^i are the Littlewood-Paley blocks introduced in Appendix A, and \mathcal{Q}_x^γ is the kernel probing the regularity of the trees, introduced below in Definition 31. The weak homogeneity exponent -6 for $[\odot_i]$ comes from the fact that $[\odot_i] \in \mathcal{D}'(M^3)$ is the Schwartz kernel of the resonant product and that our manifold M has dimension 3.

As a last piece of notation let us introduce a function space \mathcal{R}^γ adapted to the kernels varying with a parameter $x \in M$, such as the kernel \mathcal{Q}_x^γ . Note that we already know that given $x \in M$ the kernel $\mathcal{Q}_x^\gamma \in \mathcal{D}'(M \times M)$ is singular at (x, x) and smooth everywhere else. Also, the singular locus is moving with $x \in M$. Our goal is to define a correct functional space \mathcal{R}^γ that measures the singularities of \mathcal{Q}_x^γ uniformly in $x \in M$. This is done in the following definition.

Definition 21 – Fix $x_0 \in M$, along with a pair of closed neighbourhoods $U_{x_0} \Subset \Omega_{x_0} \subset M$ of x_0 , and define a conical neighbourhood

$$\mathcal{C}_{x_0} \stackrel{\text{def}}{=} \bigcup_{x \in \Omega_{x_0}} T_{(x,x)}^* M^2$$

of $T_{(x_0, x_0)}^* M^2$. Then we define \mathcal{R}^γ as the space of all families $(\mathcal{Q}_x)_{x \in U_{x_0}}$ of distributions

$$\mathcal{Q}_x \in \mathcal{D}'(M^2) \cap C^\infty(M^2 \setminus \{(x, x)\})$$

such that

$$\left(e^{-(6+2\gamma)\ell} e^{-\ell \rho_x} \mathcal{Q}_x \right)_{\ell \geq 0, x \in U_{x_0}}$$

is bounded in $\mathcal{D}'_{\mathcal{C}_{x_0}}(M^2)$ for every family of scaling fields ρ_x scaling with respect to (x, x) .

We can now define the Feynman amplitudes.

Definition 22 – We view $[\odot]$ and \mathcal{Q}_x^γ as some distributions on M^3 and M^2 by pull-back by the canonical projection from M^p to M^p . Denote by $\mathbf{d}_{V'}$ the diagonals of $(\mathbb{R} \times M)^p$, for $V' \subset V$. The amplitude $\mathcal{A}^{\mathcal{G}}$ associated with the graph $\mathcal{G}_{(t,x)}$ is the distribution on $\mathcal{T}_j \setminus \bigcup_{V' \subset V} \mathbf{d}_{V'}$ defined by the product

$$\mathcal{A}_{\mathcal{G}}(z_1, \dots, z_p) \stackrel{\text{def}}{=} \mathcal{Q}_x^\gamma(x_{v(e_{\text{ref}})_-}, x_{v(e_{\text{ref}})_+}) \prod_{e \in \tilde{E}} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{1 \leq j \leq n_{\mathcal{G}}} [\odot](x_{v_*^j}, x_{v_1^j}, x_{v_2^j})$$

with the second product corresponding to all of V_A , see Figure 4.2 for a fully detailed example. We talk of \mathcal{A}_G as the **Feynman amplitude associated with \mathcal{G}** .

The following fact is a direct consequence of Lemma 19. It allows two things in the analysis of Feynman amplitudes

- (a) If a Feynman graph is a tree, i.e. it contains no loops, then the corresponding Feynman amplitude \mathcal{A}_G is always well-defined as distribution on the corresponding configuration space.
- (b) We can reduce the analysis to **irreducible graphs** which contain at least one loop since joining two subgraphs by a bridge is always well-defined microlocally.

These two facts are detailed in the next two statements. The following Lemma states that Feynman trees are always well defined.

Lemma 23 – *If $\mathcal{G} = (V, E)$ is a tree, for every $e \in \tilde{E}$, each two point kernel K_e belongs to the module \mathcal{K}^{a_e} , $a_e \in \mathbb{R}$ endowed with the topology of Lemma 12, each three point kernel*

$$[\odot](\cdot, \cdot, \cdot) \in \mathcal{S}_{N^*(\mathbf{d}_3 \subset M^3)}^{-6}(M^3)$$

where the scaling is with respect to $\mathbf{d}_3 \subset M^3$ and the eventual marked edge e_{ref} is associated with the propagator \mathcal{Q}_x^γ which belongs to the topological space \mathcal{R}^γ from Definition 21. Then the multilinear map

$$\left([\odot], (K_e)_{e \in E}, \mathcal{Q}_x^\gamma\right) \in \mathcal{S}_{N^*(\mathbf{d}_3 \subset M^3)}^{-6}(M^3) \times \prod_{e \in E} \mathcal{K}^{a_e} \times \mathcal{R}^\gamma \mapsto \mathcal{A}_G \in \mathcal{D}'_\Gamma(\mathcal{T}_J)$$

where

$$\Gamma = \bigcup_{V' \subset V} N^*(\mathbf{d}_{V'}) \cup N^*(\mathcal{T}_{V'})$$

is continuous.

The simplest example which illustrates the above claim is the composition of pseudo-differential kernels. If we represent each kernel by an edge then composition can be interpreted in terms of Feynman rules as gluing the edges at one common vertex and this is always perfectly well defined, the diagonal singularities of the kernels do not matter. We now state a useful corollary of Lemma 23 which allows to restrict the analysis of Feynman amplitudes to *connected irreducible subgraphs*.

Corollary 24 – *Let \mathcal{A}_G be a Feynman amplitude which is obtained by joining two irreducible amplitudes $\mathcal{A}_{G_1}, \mathcal{A}_{G_2}(z_j, \mathbf{z}_2)$ by a propagator $K \in \mathcal{D}'(M^2)$ whose wave front set is in the conormal $N^*(\mathbf{d}_2 \subset M^2)$ to the diagonal*

$$\mathcal{A}_{G_1}(\mathbf{z}_1, z_i)K(z_i, z_j)\mathcal{A}_{G_2}(z_j, \mathbf{z}_2)$$

then if \mathcal{A}_{G_1} and $\mathcal{A}_{G_2}(z_j, \mathbf{z}_2)$ are some well defined distributions with wave front set in

$$(4.3) \quad \begin{aligned} WF(\mathcal{A}_{G_1}) &\subset \bigcup_{V'_1 \subset V_1} N^*(\mathbf{d}_{V'_1}) \cup N^*(\mathcal{T}_{V'_1}), \\ WF(\mathcal{A}_{G_2}) &\subset \bigcup_{V'_2 \subset V_2} N^*(\mathbf{d}_{V'_2}) \cup N^*(\mathcal{T}_{V'_2}), \end{aligned}$$

then the global amplitude \mathcal{A}_G is a well-defined distribution with wave front set included in

$$\bigcup_{V' \subset V} N^*(\mathbf{d}_{V'}) \cup N^*(\mathcal{T}_V).$$

The weak homogeneity of \mathcal{A}_G is then the sum of the weak homogeneities of the subamplitudes and of the kernel K .

Proof – Since the amplitude of a reducible graph reads

$$\mathcal{A}_{G_1}(\mathbf{z}_1, z_i)K(z_i, z_j)\mathcal{A}_{G_2}(z_j, \mathbf{z}_2)$$

for some collective variables $(\mathbf{z}_1, z_i), (z_j, \mathbf{z}_2)$ partitioning $\{z_v\}_{v \in V}$ and corresponding to a partition $V = V_1 \cup V_2$ of V , where $K(z_i, z_j)$ has wave front contained in $N^*(\mathbf{d}_2 \subset (\mathbb{R} \times M)^2) \cup N^*(\{t_i = t_j\})$ and (4.3) holds, one can apply Lemma 19 twice to

$$\mathcal{A}_{G_1}(\mathbf{z}_1, z_i)K(z_i, z_j)$$

then to

$$(\mathcal{A}_{\mathcal{G}_1}(\mathbf{z}_1, z_i)K(z_i, z_j))\mathcal{A}_{\mathcal{G}_2}(z_j, \mathbf{z}_2).$$

It shows that the product is well defined so the only difficulty is to treat the amplitudes $\mathcal{A}_{\mathcal{G}_1}$ and $\mathcal{A}_{\mathcal{G}_2}$. \triangleright




FIGURE 1. An example of Feynman graph $\mathcal{G}(= \mathcal{G}_{24})$ with $n_{\mathcal{G}} = 2$ along with the contributions to $\mathcal{A}_{\mathcal{G}}$ of its edges and vertices. Space-time vertices are pictured by red nodes, purely space vertices (with time set to the value t) are pictured by blue nodes, and black nodes represent the noises.

4.2.1 Pointed scaling spaces. We next define a parametrized version of the functional scaling spaces \mathcal{S}_I^a in configuration space which generalizes Definition 21. The elements in our new functional space depend on some space time point $z = (t, x) \in \mathcal{M}$. We will use these functional spaces when we will scale the whole Feynman amplitude $\mathcal{A}_{\mathcal{G}}$ with respect to the marked diagonal $(z, \dots, z) \subset \mathcal{M}^p$ and test that everything is uniform in $z \in \mathcal{M}$. We start by describing the geometrical setting. We are given:

- For every $z_0 = (t_0, x_0)$ an open neighborhood U_{z_0} of z_0 in \mathcal{M} so that $U_{z_0}^p$ is a neighborhood of $(z_0, \dots, z_0) \in \mathcal{M}^p$;
- a continuous family of scaling fields $\rho_z \in C^\infty(T(\mathcal{M}^p))$, $z \in U_{z_0}$ on \mathcal{M}^p such that ρ_z scales with respect to $\{(z, \dots, z)\} \subset U_{z_0}^p$ and $e^{-\ell\rho_z}U_{z_0}^p \subset U_{z_0}^p$ for every $z_0 \in U_{z_0}$, $\ell \geq 0$ and the flow of ρ_z preserves all the diagonals \mathbf{d}_I and marked diagonals $\mathbf{d}_{I,z}$.

We need to give an important example which shows that such geometric setting is non-empty and that one can always produce such setting.

Example In a local product chart $(a, b) \times U$, $U \subset M$, for every $z = (t, x) \in (a, b) \times U$ in the chart, the typical example of such a vector field reads

$$\rho_z = 2(t_1 - t)\partial_{t_1} + \dots + 2(t_p - t)\partial_{t_p} + (x_1 - x) \cdot \partial_{x_1} + \dots + (x_p - x) \cdot \partial_{x_p}.$$

Then we define the functional data. As above we define the functional spaces associated to distributions which are weakly homogeneous at a space-time diagonal $(z, \dots, z) \in \mathcal{M}^I$, in a way which is uniform in $z \in \mathcal{M}$. Here is an example that may help fixing the setting.

Example The function

$$(\mathbb{R}^3)^p \ni (x_1, \dots, x_p) \mapsto \left(|x_1 - x| \cdots |x_p - x| \right)^{-\frac{1}{2}}$$

is a function on the configuration space $(\mathbb{R}^3)^p$ with a singular locus $\{x_1 = \cdots = x_p = x\}$ which is moving with x .

Recall from (4.2) the definitions of the sets \mathbf{d}_I and \mathcal{T}_I .

Definition 25 – Fix $a \in \mathbb{R}$, $J \subset \{1, \dots, p\}$ and $z_0 = (t_0, x_0) \in \mathcal{M}$. Choose some neighbourhood U_{z_0} of z_0 . Then define $\mathcal{S}^a(\mathcal{T}_J; U_{z_0})$ as the set of families $(T_z)_{z \in U_{z_0}}$ of distributions $T_z \in \mathcal{D}'(\mathcal{T}_J \setminus \mathbf{d}_{p,z})$ such that

$$\left(e^{\ell a} e^{-\ell \rho_z * T_z} \right)_{\ell \geq 0, z \in U_{z_0}}$$

is bounded in $\mathcal{D}'(\mathcal{T}_J)$ for every ρ_z as above.

Implicitly, in the definition of \mathcal{T}_J there is a time variable t that we view as parameter where some time variables are taken to be equal to this time parameter t , t is not equal to t_0 but $(t, x) \in U_{t_0, x_0}$.

The next statement makes explicit a construction of such families; it is concerned with recentering.

Proposition 26 – Assume we are given a distribution $T \in \mathcal{S}_\Gamma^a(\mathcal{T}_J)$ with $\Gamma = N^*(\mathbf{d}_I \subset \mathcal{T}_J)$ for some $I \subset \{1, \dots, p\}$. Then for any admissible ρ_z scaling with respect to $\mathbf{d}_{V(\mathcal{G}),z}$ the family

$$\left(e^{\ell a} e^{-\ell \rho_z * T} \right)_{\ell \geq 0}$$

is bounded in $\mathcal{D}'(\mathcal{T}_J)$ uniformly in z .

Let $I \subset \{1, \dots, p\}$ and denote by $\pi : \mathcal{M}^p \mapsto \mathcal{M}^I$ the canonical projection map. Given $T \in \mathcal{S}_\Gamma^a(\mathcal{M}^I)$ where $\Gamma = \cup_{I' \subset I} N^* \mathbf{d}_{I'}$, and we scale with respect to \mathbf{d}_I . Then $\pi^* T \in \mathcal{S}_{\pi^* \Gamma}^a(\mathcal{M}^p)$ where we scale with respect to the deepest diagonal \mathbf{d}_p using admissible scaling fields.

Roughly speaking, in practice, it means you have a subamplitude $\mathcal{A}_{\mathcal{G}'}$ with a certain scaling degree with respect to its deepest diagonal $\mathbf{d}_{V(\mathcal{G}'})$, then since we would like to see how this subamplitude scales inside a bigger graph, we need to lift this to a bigger configuration space $\mathcal{M}^{V(\mathcal{G})}$, and we would like to examine the scaling degree with respect to the deepest diagonal $\mathbf{d}_{V(\mathcal{G})}$ of the larger graph or the marked diagonal $\mathbf{d}_{V(\mathcal{G}),z}$ of the larger graph. We need to ensure the scaling degree is unaffected under changes of the scaling dynamics.

Proof – The proof is very similar to the proof of [27, Lemma 6.4.5 p. 144]. By the invariance results under scalings from Proposition 15, the proof reduces to the comparison of linear scalings w.r.t. different points. So we are reduced to the following generic situation, we work on (\mathbb{R}^{1+d}) and we would like to compare the two linear scaling flows :

$$e^{-s\rho_{z_0}} : (t, x) \in \mathbb{R}^{1+d} \mapsto (e^{-2s}(t - t_0) + t_0, e^{-s}(x - x_0) + x_0) \in \mathbb{R}^{1+d}$$

which scales w.r.t. $z_0 = (t_0, x_0)$ and the second scaling

$$e^{-s\rho_{z_1}}(t, x) \in \mathbb{R}^{1+d} \mapsto (e^{-2s}(t - t_1) + t_1, e^{-s}(x - x_1) + x_1) \in \mathbb{R}^{1+d}$$

which scales w.r.t. a different point $z_1 = (t_1, x_1)$.

Then just observe the identity $e^{-s\rho_{z_1}} = (e^{-s\rho_{z_1}} e^{s\rho_{z_0}}) e^{-s\rho_{z_0}}$ where the composition $\Phi(s) \stackrel{\text{def}}{=} e^{-s\rho_{z_1}} e^{s\rho_{z_0}}$ has a smooth linear limit when $s \rightarrow +\infty$ as can be easily inspected by a direct calculation: $\Phi(s) : (t, x) \mapsto (t + (e^{-2s} - 1)(t_1 - t_0), x + (e^{-s} - 1)(x_1 - x_0))$. \triangleright

In spite of the fact that these spaces are rather ad hoc and not really intrinsic, they are sufficient to capture uniformity in $x \in M$ and therefore to control the size of the Feynman amplitudes uniformly in x .

4.2.2 The inductive Theorem for convergent amplitudes. Our next goal is to describe a recursive algorithm that controls the convergence as a distribution over the space \mathcal{T}_J . For a given graph \mathcal{G} with marked edges, given a fixed $J \subset V(\mathcal{G})$, we prove in the next statement that for every $x_0 \in M$ and every compact neighborhood U_{x_0} of x_0 , the following multilinear Feynman map is continuous

under suitable conditions on the weak homogeneity a_e of the two point kernels $(K_e)_{e \in E}$

$$(4.4) \quad \left([\odot], (K_e)_{e \in E}, \mathcal{Q}_x^\gamma \right) \in \mathcal{S}_{N^*(\mathbf{d}_3 \subset M^3)}^{-6}(M^3) \times \prod_{e \in E} \mathcal{K}^{a_e} \times \mathcal{R}^\gamma \longmapsto \mathcal{A}_{\mathcal{G},x} \in \mathcal{S}_\Gamma^{a_1}(\mathcal{T}_J) \cap \mathcal{S}^{a_2}(\mathcal{T}_J; U_{x_0})$$

where

$$\begin{aligned} \Gamma &= \bigcup_{V' \subset V(\mathcal{G})} \left(N^*(\mathbf{d}_{V'} \subset \mathcal{T}_J) \cup N^*(\mathcal{T}_{V'} \subset \mathcal{T}_J) \right) \\ a_1 &= -6n_{\mathcal{G}} - \sum_{e \in E \setminus \{e_{ref}\}} a_e \\ a_2 &= a_1 - 6 - 2\gamma. \end{aligned}$$

The space $\mathcal{S}^{a_2}(\mathcal{T}_J; U_{x_0})$ which appears in Definition 25 accounts for the fact that our estimates should be uniform in $x \in U_{x_0}$ when we scale with respect to the marked diagonal $\mathbf{d}_{V(\mathcal{G}),x}$. Recall also that we work on $\mathcal{T}_J \subset \mathcal{M}^{V(\mathcal{G})}$ for some $J \subset V(\mathcal{G})$ because we take into account that our amplitudes are integrated on regions where certain time variables coincide. If the graph \mathcal{G} has no distinguished edge e_{ref} , then we do not need to test the regularity in the space $\mathcal{S}^{a_2}(\mathcal{T}_J; U_{x_0})$ and the target functional space is just $\mathcal{S}_\Gamma^{a_1}(\mathcal{T}_J)$ for $a_1 = -6n_{\mathcal{G}} - \sum_{e \in E} a_e$.

We only consider below some subgraphs $\mathcal{G}' = (V', E')$ of \mathcal{G} which contain all the points of a given triple if ever they contain one of them. Recall all our analysis takes place in the submanifold \mathcal{T}_J of \mathcal{M}^p . With a slight abuse of notation we will also denote by $\mathbf{d}_{V'}$, the diagonal $\mathbf{d}_{V'} \cap \mathcal{T}_J$.

For any marked subgraph \mathcal{G}' we set

$$\begin{aligned} a_{\mathcal{G}',1} &\stackrel{\text{def}}{=} -6n_{\mathcal{G}'} - \sum_{e \in E' \setminus \{e_{ref}\}} a_e \\ a_{\mathcal{G}',2} &\stackrel{\text{def}}{=} a_{\mathcal{G}',1} - 6 - 2\gamma, \end{aligned}$$

while for any subgraph \mathcal{G}' with no marked/reference edge we set $a_{\mathcal{G}'} \stackrel{\text{def}}{=} -6n_{\mathcal{G}'} - \sum_{e \in E'} a_e$ and

$$\Gamma_{\mathcal{G}'} \stackrel{\text{def}}{=} \bigcup_{V'' \subset V'} \left\{ N^*(\mathbf{d}_{V''} \subset \mathcal{T}_J) \cup N^*(\mathcal{T}_{V''} \subset \mathcal{T}_J) \right\}.$$

A given Feynman graph \mathcal{G} will be said to *have a loop* if $b_1(\mathcal{G}) > 0$.

Theorem 27 – *The following holds.*

(a) *If every connected irreducible subgraph $\mathcal{G}' = (V', E')$ of \mathcal{G} with a loop satisfies*

$$(4.5) \quad \begin{aligned} a_{\mathcal{G}',1} + \text{codim}_w(\mathbf{d}_{V'}) &> 0, \\ a_{\mathcal{G}',2} + \text{codim}_w(\mathbf{d}_{V',(\mathbf{t},x)}) &> 0, \end{aligned}$$

then the Feynman map defined by Equation 4.4 is continuous.

(b) *If $\mathcal{G} = (V, E)$ is a graph that contains no distinguished edge q , every connected irreducible strict subgraph of \mathcal{G} with a loop satisfies Condition (4.5) and \mathcal{G} satisfies*

$$a_{\mathcal{G}} + \text{codim}_w(\mathbf{d}_V \subset \mathcal{T}_J) > -1,$$

then there exists a family $\Lambda_{\mathcal{Y},\varepsilon}$ of distributions supported on \mathbf{d}_V , with wavefront set in $N^(\mathbf{d}_V \subset \mathcal{T}_J)$ such that $\Lambda_\varepsilon - \Lambda_{\mathcal{Y},\varepsilon}$ has a limit in $\mathcal{D}'(\mathcal{T}_J)$ and the convergence occurs in $\mathcal{S}_{\Gamma_{\mathcal{G}} \cup N^*(\mathbf{d}_V)}^{a'}(\mathcal{T}_J)$ for all*

$$a' < -\text{codim}_w(\mathbf{d}_V \subset \mathcal{T}_J).$$

Before we give the proof of Theorem 27 an important Remark is in order.

Remark – We aim to point an important fact. The above result is general, but relies on the fact that the irreducible subgraphs are convergent. It turns out that we took care to implement the fact that for any of the Feynman graphs that we need to study, the most divergent subgraph is the graph itself. Note that this would not be the case if we were to replace all the cubic kernels $[\odot](x, y, z)$ by $\delta_x(y)\delta_x(z)$. Indeed while $[\odot]$ and the product of two δ s have the same weak homogeneity (-6) with respect to the deep diagonal $\{x = y = z\}$ they do not share the same microlocal properties. We will further elaborate on this point later in Remark 36. In particular

when we deal with the resonant products $\mathfrak{V} \odot \mathfrak{Y}$ and $\mathfrak{V} \odot \mathfrak{Y}^\circ$ in Section 5, this will imply that the subgraph containing the covariance $G^{(2)}(z_1, z_2)$ of \mathfrak{V} and the probe operator $\mathcal{Q}_x^\gamma(y_1, y_2)$ is very convergent, since these two operators are linked by $[\odot](y_1, z_1, w_1)$ and $[\odot](y_2, z_2, w_2)$ for w_i far from y_i, z_i . Conversely, replacing $[\odot](y_i, z_i, w_i)$ by $\delta_{y_i}(z_i)\delta_{y_i}(w_i)$, then this subgraph would be more divergent than the graph itself, which corresponds to the fact that while $\mathfrak{V} \odot \mathfrak{Y}$ and $\mathfrak{V} \odot \mathfrak{Y}^\circ$ are of regularities 0^- and $(-1/2)^-$, the products $\mathfrak{V} \times \mathfrak{Y}$ and $\mathfrak{V} \times \mathfrak{Y}^\circ$ are both of regularity $(-1)^-$.

Proof – Our proof proceeds by induction on subgraphs for the inclusion relation. For a subset $V_1 = V_1' \cup V_1'' \subset V(\mathcal{G})$ of vertices and $E' \subset E(\mathcal{G})$ of edges, we denote by $\mathcal{G}_{V_1, E'}$ the corresponding subgraph.

The initialization of the induction is immediate, the simplest subgraphs are just the propagators or the kernel of the resonant product which have their respective wave front sets on the conormals of the diagonals.

Given a graph \mathcal{G} constructed according to our rules, assume that all subgraphs \mathcal{G}' of \mathcal{G} have well-defined Feynman amplitudes $\mathcal{A}_{\mathcal{G}'}$ with wave front set in $\Gamma_{\mathcal{G}'}$, without loss of generality, we may assume that \mathcal{G} has p vertices which means that $V(\mathcal{G})$ is in bijection with $\{1, \dots, p\}$ hence $\mathcal{M}^{V(\mathcal{G})} \simeq \mathcal{M}^p$. For every $I \subsetneq V(\mathcal{G}) \simeq \{1, \dots, p\}$ we consider the open subset \mathcal{U}_I defined as follows

$$(4.6) \quad \mathcal{U}_I \stackrel{\text{def}}{=} \left\{ (m_1, \dots, m_p) \in \mathcal{M}^{V(\mathcal{G})}; i \in I, j \in V(\mathcal{G}) \setminus I, m_i \neq m_j \right\}.$$

By the Popineau-Stora covering Lemma 18 we have the covering $\mathcal{T}_J \setminus \mathbf{d}_{V(\mathcal{G})} = \cup_{I \subsetneq V(\mathcal{G})} \mathcal{U}_I$. The idea is to restrict $\mathcal{A}_{\mathcal{G}}|_{\mathcal{U}_I}$, then to factor the restricted amplitude as a product of subamplitudes which are well-defined by the inductive assumption and a product of smooth kernels.

For given a subset $I \subsetneq V(\mathcal{G})$ of vertices, we set

$$I' \stackrel{\text{def}}{=} V(\mathcal{G}) \setminus I, \quad (I, I') \stackrel{\text{def}}{=} (I \times I') \cup (I' \times I).$$

If K is another subset of $V(\mathcal{G})$, we write $K \not\subset I^{(\prime)}$ to say that K is neither a subset of I nor of I' . With this notation the amplitude $\mathcal{A}_{\mathcal{G}}$ factors as

$$\mathcal{A}_{\mathcal{G}} = \mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}} \left(\prod_{e: (v(e)_-, v(e)_+) \in (I, I')} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{j: \{v_1^j, v_2^j, v_3^j\} \not\subset I^{(\prime)}} [\odot](x_{v_1^j}, x_{v_2^j}, x_{v_3^j}) \right),$$

where the subgraph \mathcal{G}_I , resp. $\mathcal{G}_{I'}$, is some subgraph of \mathcal{G} with only vertices in I , resp. I' , and edges $e \in E$ such that both vertices $v(e)_-, v(e)_+ \in I$, resp. $v(e)_-, v(e)_+ \in I'$, bounding e belong to I , resp. I' . More importantly, the product on the resonant kernels indexed by integers j runs over the kernels $[\odot]$ whose vertices (v_1^j, v_2^j, v_3^j) are not entirely contained neither in I nor I' : some vertices amongst them are in I and some others are in I' . Note the fact that the product on resonant kernels might well be empty, for instance if all distinguished triples are contained either in I or I' .

Now observe that when we restrict the amplitude $\mathcal{A}_{\mathcal{G}}$ on the open subset \mathcal{U}_I , this yields

$$\mathcal{A}_{\mathcal{G}}|_{\mathcal{U}_I} = \underbrace{\mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}}}_{\text{exterior product}} \underbrace{\left(\prod_{e: (v(e)_-, v(e)_+) \in (I, I')} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{j: \{v_1^j, v_2^j, v_3^j\} \not\subset I^{(\prime)}} [\odot](x_{v_1^j}, x_{v_2^j}, x_{v_3^j}) \right)}_{\in C^\infty},$$

where the restriction to open subset \mathcal{U}_I of the product inside the parenthesis is smooth, and the product $\mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}}$ denotes an exterior tensor product of distributions which is always well-defined. We know by induction that the two amplitudes $\mathcal{A}_{\mathcal{G}_I}((z_j)_{j \in I})$ and $\mathcal{A}_{\mathcal{G}_{I'}}((z_j)_{j \in I'})$ are well-defined distributions with wave front sets

$$WF(\mathcal{A}_{\mathcal{G}_I}) \subset \Gamma_I \stackrel{\text{def}}{=} \bigcup_{K \subset I} N^*(\mathbf{d}_K \subset \mathcal{T}_J) \cup N^*(\mathcal{T}_K \subset \mathcal{T}_J),$$

and

$$WF(\mathcal{A}_{\mathcal{G}_{I'}}) \subset \Gamma_{I'} \stackrel{\text{def}}{=} \bigcup_{K' \subset I'} N^*(\mathbf{d}_{K'} \subset \mathcal{T}_J) \cup N^*(\mathcal{T}_{K'} \subset \mathcal{T}_J).$$

Analysing the graph \mathcal{G} , there are two situations, either \mathcal{G} contains some distinguished edge \mathcal{Q}_x^γ in which case we should study the scaling behaviour in two steps, first scale on the partial diagonals $\mathbf{d}_{V(\mathcal{G})}$ where all points labelled by $V(\mathcal{G})$ collide, then study the scaling on the marked diagonals $\mathbf{d}_{V(\mathcal{G}), (t,x)}$ (recall the marked diagonal is just one point in $\mathcal{M}^{|V(\mathcal{G})|}$) where all points labelled by $V(\mathcal{G})$ collide uniformly on (t,x) , we use the spaces of Definition 25 to control the uniformity in the parameter (t,x) . If the graph contains no distinguished edge, then we just scale in one step with respect to $\mathbf{d}_{V(\mathcal{G})}$.

Assume we are in the first situation where \mathcal{G} contains a distinguished edge, the second case is simpler to handle. Choosing any admissible scaling field ρ scaling with respect to the deep diagonal $\mathbf{d}_{V(\mathcal{G})}$ and using the stability of \mathcal{U}_I by such scaling field (a consequence of admissibility of ρ , we made an essential use of the fact that the flow by scaling fields ρ from Definition 16 preserves both time and space-time diagonals $\mathcal{T}_J, \mathbf{d}_I$ hence every $\mathcal{U}_I, I \subsetneq V(\mathcal{G})$). The weak homogeneity of each term in factor does not depend on the choice of scaling field with respect to $\mathbf{d}_{V(\mathcal{G})}$ by Proposition 15. Moreover, we also use the property that the symplectic lifts of these scaling fields preserve the conormals of all time and space-time diagonals $N^*(\mathcal{T}_I), N^*(\mathbf{d}_I)$ so that the cones $\Gamma_I, \Gamma_{I'}$ containing microsingularities of $\mathcal{A}_{\mathcal{G}_I}, \mathcal{A}_{\mathcal{G}_{I'}}$ are stable by the lifted flow: $e^{-\ell\rho^*}\Gamma_I \subset \Gamma_I, e^{-\ell\rho^*}\Gamma_{I'} \subset \Gamma_{I'}$ by Lemma 17. Using the fact that each element $\eta_I \in C^\infty(M^{d_{V(\mathcal{G})}} \setminus \mathbf{d}_{V(\mathcal{G})})$ of the partition of unity belongs to $\mathcal{S}_0^0(M^{V(\mathcal{G})} \setminus \mathbf{d}_{V(\mathcal{G})})$, it has scaling degree 0 in $C^\infty(M^{d_{V(\mathcal{G})}}) \setminus \mathbf{d}_{V(\mathcal{G})}$ in the sense that

$$(e^{-\ell\rho^*}\eta_I)_{\ell \geq 0}$$

is a bounded family in $C^\infty(M^{V(\mathcal{G})} \setminus \mathbf{d}_{V(\mathcal{G})})$. We can thus define the renormalized product

$$\eta_I \mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}} \left(\prod_{e: (v(e)_-, v(e)_+) \in (I, I')} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{j: \{v_1^j, v_2^j, v_*^j\} \not\subset I'} [\odot](x_{v_*^j}, x_{v_1^j}, x_{v_2^j}) \right)$$

on $\mathcal{T}_J \setminus \mathbf{d}_{V(\mathcal{G}), (t,x)}$ by applying Proposition 20 twice and we find that the product is well-defined since the weak homogeneity of the above product is $> -\text{codim}_w(\mathbf{d}_{V(\mathcal{G})})$ by assumption. We insist that when we scale with respect to $\mathbf{d}_{V(\mathcal{G})}$ the weak homogeneity of \mathcal{Q}_x^γ is 0 by Lemma 28. By summation over I we conclude that $\mathcal{A}_{\mathcal{G}}$ is well-defined on $\mathcal{T}_J \setminus \mathbf{d}_{V(\mathcal{G}), (t,x)}$.

To define an extension on the whole configuration space \mathcal{T}_J we need the two step extension from Theorem 14. We now need to apply Proposition 20 and now Theorem 14 a second time to the expression

$$\eta_I \mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}} \left(\prod_{e: (v(e)_-, v(e)_+) \in (I, I')} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{j: \{v_1^j, v_2^j, v_*^j\} \not\subset I'} [\odot](x_{v_*^j}, x_{v_1^j}, x_{v_2^j}) \right)$$

but for the scaling fields with respect to the distinguished diagonal $\mathbf{d}_{V(\mathcal{G}), (t,x)}$. *It is at this point that we are using our second inductive assumption on the weak homogeneity of all subgraphs containing the distinguished edge*, equivalently all subamplitudes containing the propagator \mathcal{Q}_x^γ . We make an induction on all marked subgraphs whose amplitude contains \mathcal{Q}_x^γ , the induction starts with \mathcal{Q}_x^γ itself. We use the fact that the scaling degree of \mathcal{Q}_x^γ with respect to $\mathbf{d}_{V(\mathcal{G})} \setminus \mathbf{d}_{V(\mathcal{G}), (t,x)}$ equals 0, while when scaling with respect to $\mathbf{d}_{V(\mathcal{G}), (t,x)}$, the scaling degree of \mathcal{Q}_x^γ is now given by $-6 - 2\gamma$. This last claim is proven in Lemma 28 below.

At this stage given a graph \mathcal{G} , assume by induction that all subgraphs \mathcal{G}' containing the distinguished edge have scaling degree with respect to the marked diagonal $> -\text{codim}_w(\mathbf{d}_{V(\mathcal{G}'), (t,x)})$. Then every product defined on $\mathcal{T}_J \setminus \mathbf{d}_{V(\mathcal{G})}$

$$\eta_I \mathcal{A}_{\mathcal{G}_I} \mathcal{A}_{\mathcal{G}_{I'}} \left(\prod_{e: (v(e)_-, v(e)_+) \in (I, I')} K_e(z_{v(e)_-}, z_{v(e)_+}) \prod_{j: \{v_1^j, v_2^j, v_*^j\} \not\subset I'} [\odot](x_{v_*^j}, x_{v_1^j}, x_{v_2^j}) \right)$$

will be weakly homogeneous of degree $a_{\mathcal{G}, 2} = \sum_e a_e - 6n_{\mathcal{G}} - 6 - 2\gamma$ when we scale with respect to the marked diagonal $\mathbf{d}_{V(\mathcal{G}), (t,x)}$ where we need to include the scaling degree of \mathcal{Q}_x^γ which equals $-6 - 2\gamma$. So each piece above satisfies the assumptions of Theorem 14 hence extends to \mathcal{T}_J .

The uniform estimates in x essentially follow from the stability assumptions of the cones by change of scaling in Proposition 26. \triangleright

Lemma 28 – Fix $d \geq 2$ and denote by $\pi_{\leq 2} : \mathcal{M}^d \rightarrow \mathcal{M}^2$ the canonical projection. For all values of $\gamma \in \mathbb{R}$ the kernel $\pi_{\leq 2}^* \mathcal{Q}_x^\gamma$ is a smooth germ near $\mathbf{d}_d \setminus \mathbf{d}_{d,x}$ in $\mathbb{R}^2 \times \mathcal{M}^d \setminus \mathbf{d}_{d,x}$. We define

$$\mathcal{O} \stackrel{\text{def}}{=} \left\{ m \in \mathbb{R}^2 \times \mathcal{M}^d : \text{dist}(m, \mathbf{d}_{d,x}) > 2\text{dist}(m, \mathbf{d}_d) \right\}.$$

Then $\pi_{\leq 2}^* \mathcal{Q}_x^\gamma$ is weakly homogeneous of degree 0 when scaling with respect to $\mathbf{d}_d \setminus \mathbf{d}_{d,x}$ in $\Omega \setminus \mathbf{d}_d$ in the following sense: For every compact $K \Subset \Omega$, for all scaling field ρ scaling with respect to \mathbf{d}_d , there exists $\ell_K > 0$ such that the family

$$\left(\{ e^{-\ell\rho} \pi_{\leq 2}^* \mathcal{Q}_x^\gamma \}|_K \right)_{\ell \geq \ell_K}$$

is bounded in $C^\infty(K)$.

Proof – For any element $m \in \mathcal{O}$ we know that the limit point $m_\infty = \lim_{t \uparrow \infty} e^{-\ell\rho}(m) \in \mathbf{d}_d \setminus \mathbf{d}_{d,x}$ has the form $m_\infty = (y, \dots, y, t, t)$ for $y \neq x$ and therefore $\pi_{\leq 2}^* \mathcal{Q}_x^\gamma(\cdot)$ is smooth near the limit point $m_\infty = (y, \dots, y, t, t)$ since $\pi_{\leq 2}^* \mathcal{Q}_x^\gamma(m_\infty) = \mathcal{Q}_x^\gamma(y, y)$ for $y \neq x$. Now by a continuity argument, we know that for any element $m \in \mathcal{O}$ there exists a neighbourhood $V_m \subset \Omega$ and $\ell_m > 0$, such that for all $\ell \geq \ell_m$ one has $e^{-\ell\rho}(V_m) \subset \Omega$ and $\text{dist}(\pi_{\leq 2}(e^{-\ell\rho}(V_m)), (x, x)) \geq \delta > 0$ hence $\pi_{\leq 2}^* \mathcal{Q}_x^\gamma|_{e^{-\ell\rho}(V_m)}$ is smooth uniformly in $\ell \geq \ell_m$ which yields the claim by compactness. \triangleright

5 – Random fields from renormalization

Recall the definition (2.1) of the enhancement $\widehat{\xi}_r$ of the regularized noise ξ_r

$$\widehat{\xi}_r \stackrel{\text{def}}{=} \left(\xi_r, \mathfrak{V}_r, \mathfrak{W}_r, \mathfrak{Y}_r \odot \mathfrak{I}_r, \mathfrak{Y}_r \odot \mathfrak{V}_r - \frac{b_r}{3}, |\nabla \mathfrak{Y}_r|^2 - \frac{b_r}{3}, \mathfrak{Y}_r \odot \mathfrak{V}_r - b_r \mathfrak{I}_r \right).$$

We use the index r to emphasize the regularization of the noise. We show in this section the convergence of $\widehat{\xi}_r$ to some limit enhanced noise in its natural space (2.2). The random fields are constructed by means of some covariance computations, which finally yield the existence of the stochastic objects in some Hölder-Besov spaces by a Kolmogorov type argument. In this section we need to extract a Littlewood-Paley block of our stochastic trees, then control the size of the Littlewood-Paley block evaluated at some point $x \in M$, uniformly in $x \in M$. We need some uniform control because of the loss of translation invariance, which is the main difficulty of this section and of SPDE on non-homogeneous Riemannian manifolds. We start the section by defining abstractly in Section 5.1 the framework of Kolmogorov estimates in our non stationary setting where we need pointwise bounds. This also requires the introduction of some distinguished propagator \mathcal{Q}_x^γ which depends on a base point $x \in M$. Then we put our definition in action by discussing the stochastic estimates of Wick monomials where things are simple. The reader who wants to get some familiarity with our approach is urged to first read how we handle the regularities of the simple Wick monomials $\mathfrak{I}, \mathfrak{V}, \mathfrak{W}$ in Lemma 33 and sections 5.2.2 and 5.2.3. A novelty of our approach for the Wick monomials is that we renormalize in a **locally covariant way** which is the only way to ensure some form of locality. We refer the reader to Section 5.2.1 for a more detailed discussion. The estimates on Wick monomials should also be useful for readers interested by simpler Quantum Field Theories on surfaces.

Then, starting from Section 5.3, we deal with more complicated graphs coming from the higher chaoses, as enumerated in the first paragraph in Section 5.4. While some of the amplitude arising from these computations can be directly handled using Theorem 27, for several graphs that appear in our stochastic estimates we need to feed the induction Theorem 27 with more informations, this uses subtle smoothing properties of the Schwartz kernel of the resonant product which is discussed in Section 5.4.1 and also some control of certain renormalized subamplitude in Section 5.4.2. We refer the reader to the relevant parts for more details but this can be read after one is used to the general inductive machinery. The amplitude containing a more singular subamplitude require a more careful analysis, taking place in Section 5.4. We also devote Section 5.5.3 to the exact calculation of the counterterms from the divergent subamplitude; this uses subtle arguments involving heat asymptotics, stationary phase and the borderline case of our extension Theorem 13 where we need a renormalization.

5.1 Kolmogorov type argument

We start by giving the definitions of the Besov spaces that we need for our analysis.

Definition 29 – In the sequel we fix I a finite set and $(U_i, \kappa_i)_{i \in I}$ a cover of M with $\kappa_i : U_i \rightarrow \mathbb{R}^3$, along with a quadratic partition of unity $(\chi_i)_{i \in I}$ on the closed manifold M subordinated to $(U_i)_{i \in I}$. For any $\alpha \in \mathbb{R}$ and $p, q \in [1, \infty]$ we define the Besov space $B_{pq}^\alpha(M)$ on the closed manifold M as the completion of $C^\infty(M)$ with the norm

$$(5.1) \quad \|f\|_{B_{pq}^\alpha(M)} \stackrel{\text{def}}{=} \max_{i \in I} \|\kappa_{i*}(\chi_i f)\|_{B_{pq}^\alpha(\mathbb{R}^3)} \stackrel{\text{def}}{=} \max_{i \in I} \left\| 2^{k\alpha} \|\Delta_k(\kappa_{i*}(\chi_i f))\|_{L^p(\mathbb{R}^3)} \right\|_{\ell_k^q},$$

where the operators Δ_k for $k \geq -1$ are Littlewood-Paley blocks on the flat space \mathbb{R}^3 .

We slightly rewrite this norm as follows. We fix a collection $(\psi_i)_{i \in I}$ of functions on \mathbb{R}^3 such that $\text{supp}(\psi_i) \subset \kappa_i(U_i)$ and $\psi_i|_{\kappa_i(\text{supp}(\chi_i))} = 1$; there is enough room, since the support of χ_i does not fill U_i . Then we prove in the companion paper [10] that the norm defined in (5.1) is equivalent to

$$(5.2) \quad \max_{i \in I} \left\| 2^{k\alpha} \|\psi_i \Delta_k(\kappa_{i*}(\chi_i f))\|_{L^p(\mathbb{R}^3)} \right\|_{\ell_k^q}.$$

We therefore define for $i \in I$ and $k \geq -1$ some Littlewood-Paley blocks on the manifold M by

$$(5.3) \quad P_k^i(f) \stackrel{\text{def}}{=} \kappa_i^*[\psi_i \Delta_k(\kappa_{i*}(\chi_i f))].$$

With this notation in hand $\|f\|_{B_{pq}^\alpha(M)}$ therefore equivalent to

$$(5.4) \quad \max_{i \in I} \left\| 2^{k\alpha} \|P_k^i(\cdot)\|_{L^p(M)} \right\|_{\ell_k^q}.$$

In the sequel, to lighten the notation, we often let $B_{p,q}^\alpha$ stand for $B_{pq}^\alpha(M)$ and define

$$C^\alpha \stackrel{\text{def}}{=} B_{\infty, \infty}^\alpha(M).$$

The equivalence of (5.1) with (5.2) is shown in the proof of [10, Lemma 2.10], where we prove that on \mathbb{R}^3 the commutator of a Littlewood-Paley block with multiplication by a smooth compactly supported function is a smoothing operator. In the sequel we denote by $\psi \circ |\cdot| \in \mathcal{S}(\mathbb{R})$ the Schwartz kernel of Δ_0 in \mathbb{R}^3 ; its Fourier transform η is supported in an annulus. Note that for any $k \geq 0$, the Schwartz kernel of Δ_k is given by

$$\Delta_k(x - y) = 2^{3k} \psi(2^k |x - y|).$$

We are now ready to state our Kolmogorov type lemma. It turns out that due to the lack of stationarity of the stochastic objects on manifolds (as opposed to the flat case, or the sphere), the inequalities stated in Lemma 30 below involve a point-wise bound requiring to be uniform in some base-point $x \in M$. The mechanics involved here is classical; we give the details for the reader's convenience.

Lemma 30 – Fix $T > 0$, $n \in \mathbb{N}$ and let u be a random distribution in $\mathcal{D}'([0, T] \times M)$ which belongs to the Wiener chaos of degree n .

– Assume one has

$$(5.5) \quad \mathbb{E}[(P_k^i u)(t, x)^2] \lesssim 2^{-2k\gamma}$$

uniformly in $x \in M$, $i \in I$ and $k \geq 0$ for some $\gamma \in \mathbb{R}$, for some time $0 \leq t \leq T$. Then for every $s < \gamma$ and $p > 1$ we have

$$\mathbb{E}[\|u(t)\|_{C^s}^p] \lesssim_p 1.$$

– If, in addition to (5.5), the distribution u further satisfies

$$(5.6) \quad \mathbb{E}[(P_k^i u)(t_1, x) - (P_k^i u)(t_2, x)]^2 \lesssim 2^{-2k\gamma} |t_1 - t_2|^{2\beta}$$

uniformly in $t_1, t_2, x \in [0, T] \times [0, T] \times M$ and $(i, k) \in I \times \mathbb{N}$ for some $\gamma \in \mathbb{R}, \beta > 0$ then for every $s < \gamma$ and $\theta < \beta$ we have

$$u \in C_T^\theta C^s.$$

Proof – We concentrate on the second item as the first item is proved by a similar reasoning. By Sobolev embedding by have

$$\begin{aligned} \mathbb{E} \left[\|u(t_1) - u(t_2)\|_{C^s}^p \right] &\lesssim \mathbb{E} \left[\|u(t_1) - u(t_2)\|_{B_{p,p}^{s+3/p}}^p \right] \\ &\lesssim \max_{i \in I} \sum_{k \geq 0} 2^{pk(s+3/p)} \int_M \mathbb{E} \left[|(P_k^i u)(t_1, x) - (P_k^i u)(t_2, x)|^p \right] dx. \end{aligned}$$

Then the Gaussian hypercontractivity estimate implies that

$$\begin{aligned} \mathbb{E} \left[\|u(t_1) - u(t_2)\|_{C^s}^p \right] &\lesssim \max_{i \in I} \sum_{k \geq 0} 2^{pk(s+3/p)} \int_M \mathbb{E} \left[\{(P_k^i u)(t_1, x) - (P_k^i u)(t_2, x)\}^2 \right]^{\frac{p}{2}} dx \\ &\lesssim \max_{i \in I} \sum_{k \geq 0} \sup_{x \in M} \left(2^{2k(s+3/p)} \mathbb{E} \left[\{(P_k^i u)(t_1, x) - (P_k^i u)(t_2, x)\}^2 \right] \right)^{\frac{p}{2}}. \end{aligned}$$

Therefore using the hypothesis (5.6) we end up with

$$\mathbb{E} \left[\|u(t_1) - u(t_2)\|_{C^s}^p \right] \lesssim \sum_{k \geq 0} 2^{pk(s+3/p-\gamma)} |t_1 - t_2|^{p\beta} \lesssim |t_1 - t_2|^{p\beta},$$

which holds provided we choose p large enough to have $s < \gamma - 3/p$. The proof now follows using a classical Kolmogorov argument in time. \triangleright

We denote by $P_k^i(x, y)$ the smooth Schwartz kernel of the operator P_k^i .

Definition 31 – We introduce a bilinear kernel depending on a point $x \in M$ and $\gamma \in \mathbb{R}$ allowing to probe the regularities of objects as follows

$$(5.7) \quad \mathcal{Q}_x^\gamma(y_1, y_2) \stackrel{\text{def}}{=} \sum_{i \in I} \sum_{k \geq -1} 2^{2k\gamma} P_k^i(x, y_1) P_k^i(x, y_2).$$

Note that if there exists $\gamma \in \mathbb{R}$ such that one has

$$\mathbb{E} \left[\mathcal{Q}_x^\gamma(u(t, \cdot), u(t, \cdot)) \right] = \mathbb{E} \left[\sum_{k \geq -1} 2^{2k\gamma} (P_k^i u)(t, x)^2 \right] = \sum_{k \geq -1} 2^{2k\gamma} \mathbb{E} \left[(P_k^i u)(t, x)^2 \right] < \infty$$

uniformly in $x \in M$, then we have

$$\mathbb{E} \left[(P_k^i u)(t, x)^2 \right] \lesssim 2^{-2k\gamma}$$

uniformly in $k \geq -1$.

The microlocal properties of the kernel \mathcal{Q}_x^γ are stated in the following lemma.

Lemma 32 – Fix $x \in M$ and $\gamma \in \mathbb{R}$. The series defining \mathcal{Q}_x^γ converges in $C^\infty(M^2 \setminus \{(x, x)\})$ and in the space $\mathcal{D}'_{\Gamma_x}(M \times M)$ of distributions with wave front set

$$\Gamma_x = T_{(x,x)}^* M^2.$$

Moreover it is weakly homogeneous of degree $-6 - 2\gamma$ with respect to the scaling towards $\mathcal{Y} = (x, x)$, that is to say we have $\mathcal{Q}_x^\gamma \in \mathcal{S}_{\Gamma_x}^{-6-2\gamma}$ with a control uniform in x .

Proof – For the convergence in $C^\infty(M^2 \setminus \{(x, x)\})$, in a local chart indexed by $i \in I$, the kernel of \mathcal{Q}_x^γ is given by

$$\sum_{k \geq 0} 2^{k(6+2\gamma)} \psi_i^2(\tilde{x}) (\kappa_{i*} \chi_i)^{\otimes 2}(y_1, y_2) \psi(2^k |y_1 - \tilde{x}|) \psi(2^k |y_2 - \tilde{x}|),$$

where $\psi \circ |\cdot|$ is a Schwartz function on \mathbb{R} with Fourier transform η supported on an annulus, and $\tilde{x} \stackrel{\text{def}}{=} \kappa_i(x)$. For $y_i \neq \tilde{x}$ we have for any $N \geq 0$ and any $\alpha \in \mathbb{N}^3$

$$|\partial_{y_i}^\alpha \psi(2^k |y_i - \tilde{x}|)| \lesssim 2^{k|\alpha|} (1 + 2^k |y_i - \tilde{x}|)^{-N} \lesssim 2^{k|\alpha| - Nk}.$$

Therefore choosing N large enough ensures the convergence of the series in C^∞ .

We now turn to the proof of the convergence of the series in the sense of distributions in $\mathcal{D}'_{\Gamma_x}(M \times M)$. We use different arguments depending on the sign of γ .

• Suppose that $\gamma < 0$. Then we have $\|2^{3k}\psi(2^k|\cdot - \tilde{x}|\|_{L^1} = \|\psi\|_{L^1}$ and the series (5.7) converges absolutely in L^1 since

$$\begin{aligned} \left\| \sum_{k \geq 0} 2^{k(6+2\gamma)} \psi_i^2(\tilde{x})(\kappa_{i*}\chi_i)^{\otimes 2}(y_1, y_2) \psi(2^k|y_1 - \tilde{x}|) \psi(2^k|y_2 - \tilde{x}|) \right\|_{L^1_{y_1, y_2}} \\ \lesssim \sum_{k \geq 0} 2^{k\gamma} \|\psi\|_{L^1}^2 \|\psi_i\|_{L^\infty}^2 \|\chi_i\|_{L^\infty}^2 \lesssim 1. \end{aligned}$$

• Suppose now that $\gamma \geq 0$. We must prove that the series converges in the distributional sense. For every test function $\varphi \in C_c^\infty(\mathbb{R}^3 \times \mathbb{R}^3)$ we must prove the convergence of the series

$$\begin{aligned} \sum_{k \geq 0} 2^{k(6+2\gamma)} \int_{(\mathbb{R}^3)^2} \varphi(y_1, y_2) \psi_i^2(\tilde{x})(\kappa_{i*}\chi_i)^{\otimes 2}(y_1, y_2) \psi(2^k|y_1 - \tilde{x}|) \psi(2^k|y_2 - \tilde{x}|) dy_1 dy_2 \\ = \sum_{k \geq 0} 2^{2k\gamma} \int_{(\mathbb{R}^3)^2} \varphi(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \\ \times \psi_i^2(\tilde{x})(\kappa_{i*}\chi_i)^{\otimes 2}(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) dy_1 dy_2. \end{aligned}$$

We rename by $\tilde{\varphi}$ the test function

$$\tilde{\varphi}(y_1, y_2) \stackrel{\text{def}}{=} \varphi(y_1, y_2)(\kappa_{i*}\chi_i)^{\otimes 2}(y_1, y_2).$$

The series therefore rewrites as

$$\sum_{k \geq 0} 2^{2k\gamma} \int_{(\mathbb{R}^3)^2} \tilde{\varphi}(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) \psi_i^2(\tilde{x}) dy_1 dy_2.$$

Note that the product $\psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) \psi_i^2(\tilde{x})$ is a Schwartz function in y_1, y_2 uniformly in \tilde{x} . The idea is to Taylor expand $\tilde{\varphi}$ about (\tilde{x}, \tilde{x}) at order N with integral remainder. This yields

$$\tilde{\varphi} = P_N + R_N, \quad R_N(y_1, y_2) = \mathcal{O}((|y_1 - \tilde{x}| \vee |y_2 - \tilde{x}|)^{N+1}),$$

where P_N is a polynomial function of (y_1, y_2) centered at (\tilde{x}, \tilde{x}) . The rescaled $\tilde{\varphi}$ reads

$$\begin{aligned} \tilde{\varphi}(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \\ = P_N(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) + R_N(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}), \end{aligned}$$

where $R_N = \mathcal{O}(2^{-k(N+1)})$ depends on the jets of order $N+1$ of $\tilde{\varphi}$. Injecting the above decomposition in our series yields

$$\sum_{k \geq 0} 2^{2k\gamma} \int_{(\mathbb{R}^3)^2} (P_N + R_N)(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) \psi_i^2(\tilde{x}) dy_1 dy_2.$$

Now we make use of the fact that we have

$$\int_{(\mathbb{R}^3)^2} P_N(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) \psi_i^2(\tilde{x}) dy_1 dy_2 = 0,$$

since the support of η is included in an annulus, which implies that its integral against all polynomials vanish. The series therefore simplifies as

$$\begin{aligned} \sum_{k \geq 0} 2^{2k\gamma} \int_{(\mathbb{R}^3)^2} \underbrace{R_N(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x})}_{\mathcal{O}(2^{-k(N+1)})} \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) \psi_i^2(\tilde{x}) dy_1 dy_2 \\ \lesssim \sum_{k=0}^{\infty} 2^{k(2\gamma-N-1)} \sum_{|\alpha|=N+1} \|\partial^\alpha \tilde{\varphi}\|_{L^1}, \end{aligned}$$

where the series on the right hand side converges as soon as $N+1 > 2\gamma$ and we control the convergence from the $N+1$ derivatives of $\tilde{\varphi}$ hence η . We just proved that \mathcal{Q}'_x converges as distribution of order $N+1$ for all $N+1 > 2\gamma$.

Finally we need to control the weak homogeneity when we scale towards (x, x) . Working in a chart $i \in I$ this reduces to estimate the weak homogeneity near (x, x) of the series

$$U \stackrel{\text{def}}{=} \sum_{k \geq 0} 2^{k(6+2\gamma)} \psi(2^k |y_1 - \tilde{x}|) \psi(2^k |y_2 - \tilde{x}|) \in \mathcal{S}'(\mathbb{R}^3 \times \mathbb{R}^3).$$

By duality it suffices to estimate the weak homogeneity at ∞ of its Fourier transform

$$\begin{aligned} \widehat{U}(\xi_1, \xi_2) &= \sum_{k \geq 0} 2^{k(6+2\gamma)} \mathcal{F}[\psi(2^k |\cdot - \tilde{x}|)](\xi_1) \mathcal{F}[\psi(2^k |\cdot - \tilde{x}|)](\xi_2) \\ &= \sum_{k \geq 0} 2^{2k\gamma} e^{i\tilde{x} \cdot (\xi_1 + \xi_2)} \eta(2^{-k} \xi_1) \eta(2^{-k} \xi_2). \end{aligned}$$

Here we can use the fact that $\eta(2^{-k} \xi)$ is zero unless $2^{k-1} \leq |\xi| \leq 2^{k+1}$, so that both ξ_1 and ξ_2 are localized at scale 2^j for some $j \geq 0$, which yields the bound

$$\left| \sum_{k \geq 0} 2^{2k\gamma} e^{i\tilde{x} \cdot (\xi_1 + \xi_2)} \eta(2^{-k} \xi_1) \eta(2^{-k} \xi_2) \right| = \mathcal{O}(|\xi_1|^{2\gamma} \vee |\xi_2|^{2\gamma})$$

uniform in \tilde{x} , hence the Fourier transform \widehat{U} is such that the family of tempered distributions $(\lambda^{-2\gamma} \widehat{U}(\lambda \cdot))_{\lambda \geq 1}$ is bounded. By Plancherel this entails that the family of distributions

$$(\lambda^{-2\gamma-6} U(\lambda \cdot + (\tilde{x}, \tilde{x})))_{\lambda \in (0,1]}$$

is bounded too, which concludes the proof. \triangleright

As an example we discuss below the regularity of the dynamical free field.

Lemma 33 – Fix $T > 0$ and $\varepsilon > 0$. Then, we have $\mathfrak{F} \in C_T \mathcal{C}^{-1/2-\varepsilon}$.

Proof – We start by performing the estimate at some fixed time $t > 0$. By stationary in time of \mathfrak{F} one has for any $x \in M$ the t -independent bound

$$\mathbb{E}[\mathcal{Q}_x^\gamma(\mathfrak{F}(t, \cdot), \mathfrak{F}(t, \cdot))] \lesssim \int_{M^2} \mathcal{Q}_x^\gamma(y_1, y_2) P^{-1}(y_1, y_2) dv(y_1) dv(y_2).$$

Next we use the representation (5.3) of the Schwartz kernel of P_k^i , which implies that pushing the estimate in the chart $\kappa_i(U_i)$ we have

$$\mathbb{E}[(P_k^i \mathfrak{F})(t, x)^2] \lesssim 2^{6k} \int_{(\mathbb{R}^3)^2} \psi_i(\tilde{x})^2 \psi(2^k |y_1 - \tilde{x}|) \psi(2^k |y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2}(y_1, y_2) \kappa_{i*}^{\otimes 2} P^{-1}(y_1, y_2) dy_1 dy_2$$

where we denote $\tilde{x} \stackrel{\text{def}}{=} \kappa_i(x)$. Also, denoting by $\kappa_{i*}^{\otimes 2} P^{-1} =: P_{i,i}^{-1}$ the pulled-back Green function, we obtain

$$\begin{aligned} \mathbb{E}[(P_k^i \mathfrak{F})(t, x)^2] &\lesssim \int_{(\mathbb{R}^3)^2} \psi_i(\tilde{x})^2 \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2} \left(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x} \right) \\ &\quad \times P_{i,i}^{-1} \left(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x} \right) dy_1 dy_2. \end{aligned}$$

Since \tilde{x} is chosen uniformly in a compact set, $\psi(|y_1 - \tilde{x}|), \psi(|y_2 - \tilde{x}|)$ are Schwartz in y_1, y_2 and $P_{i,i}^{-1} \left(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x} \right)$ is integrable in y_1, y_2 near $y_1 = y_2$, so the second integral is well-defined. Using the bound on the rescaled Green function

$$P_{i,i}^{-1} \left(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x} \right) \lesssim 2^k |y_1 - y_2|^{-1},$$

we can now conclude that we have

$$\mathbb{E}[(P_k^i \mathfrak{F})(t, x)^2] \lesssim 2^k \int_{(\mathbb{R}^3)^2} |\psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|)| |y_1 - y_2|^{-1} dy_1 dy_2 \lesssim 2^k.$$

We now show that \mathfrak{F} is indeed continuous in time. To do so, let us first introduce the notation

$$G_{i,i}^{(1)}(t_1 - t_2, x, y) \stackrel{\text{def}}{=} \kappa_{i*}^{\otimes 2} \left(\frac{e^{-|t_1 - t_2|P}}{P} \right) (x, y).$$

The quantity $\mathbb{E}[(P_k^{i\hat{\eta}}(t_1, x) - P_k^{i\hat{\eta}}(t_2, x))^2]$ is bounded above by

$$\left| \int_{(\mathbb{R}^3)^2} \psi_i(\tilde{x})^2 \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2} \left(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x} \right) \right. \\ \left. \times \left(P_{i,i}^{-1}(2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) - G_{i,i}^{(1)}(t_1 - t_2, 2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \right) \right|$$

for an integral with respect to $dy_1 dy_2$. Our goal is to bound the difference of kernels $P_{i,i}^{-1} - G_{i,i}^{(1)}(r, \bullet)$. To do so, we use the identity

$$\frac{1 - e^{-rP}}{P} = \int_0^r e^{-sP} ds$$

which expresses the difference as an integral of the heat operator. Then we can inject in the integral expression the bound on the heat kernel on the product chart

$$|e^{-sP}(y_1, y_2)| \lesssim s^{-3/2} e^{-s} e^{-C|y_1 - y_2|^2/s},$$

where $C > 0$. We obtain the estimate for $\beta \in (0, 1)$

$$\begin{aligned} \left| P_{i,i}^{-1}(y_1, y_2) - G_{i,i}^{(1)}(r, y_1, y_2) \right| &\lesssim \int_0^r s^{-3/2} e^{-C|y_1 - y_2|^2/s-s} ds \lesssim \int_{r^{-1}}^\infty e^{-Cs|y_1 - y_2|^2} e^{-\frac{1}{s}} s^{-\frac{1}{2}} ds \\ &\lesssim |y_1 - y_2|^{-1} \int_{r^{-1}|y_1 - y_2|^2}^\infty e^{-Cs - \frac{1}{s}} s^{-\frac{1}{2}} ds \lesssim |y_1 - y_2|^{-1} \left(e^{-\frac{|y_1 - y_2|^2}{r}} \right) \\ (5.8) \quad &\lesssim |y_1 - y_2|^{-1} \left(1 + \frac{|y_1 - y_2|^2}{r} \right)^{-N} \lesssim |y_1 - y_2|^{-1-\beta} r^{\frac{\beta}{2}}. \end{aligned}$$

Replacing in the difference estimate finally yields

$$\mathbb{E} \left[\left((P_k^{i\hat{\eta}}(t_1, x) - P_k^{i\hat{\eta}}(t_2, x))^2 \right) \right] \lesssim 2^{k(1+\beta)} |t_1 - t_2|^{\frac{\beta}{2}}.$$

▷

Together with Kolmogorov's classical regularity theorem what we said above justifies that $\hat{\eta}$ takes almost surely its values in $C_T^\eta C^{-1/2-\varepsilon}(M)$, for all small enough $\eta > 0$ and all $\varepsilon > 0$, and its norm in the corresponding space has moments of any finite order.

Remark – In the sequel we will not dwell on the proof of the estimate (5.6) for larger trees. However it is by now classical that when computing in charts the covariance of $(P_k^i u)(t_1, x) - (P_k^i u)(t_2, x)$ the amplitudes appearing can be reorganised by means of some telescoping sums in order to introduce a terms $P_{i,i}^{-1} - G_{i,i}^{(1)}(r, \bullet)$. In view of (5.8) this term always yields a good factor in time modulo a small loss in spacial regularity.

5.2 The Wick monomials

5.2.1 Locally covariant Wick renormalization. In the sequel we will systematically first estimate regularities using Wick renormalization. Then we shall compare the usual Wick renormalization which is not *locally covariant* with a *locally covariant* renormalization in which we only subtract *universal quantities* at the cost of producing objects which do not belong to homogeneous Wiener chaoses. Let us discuss the notion of *local covariance* with some simple example which also explains why the usual Wick renormalization fails to be locally covariant. A *function valued invariant of Riemannian structure* is a function c which assigns to each manifold M and Riemannian structure g on M some function $c[g]$ on M which is *locally covariant* in the following sense: given $f : M' \hookrightarrow M$ a diffeomorphism onto an open submanifold of M , and a Riemannian structure g on M , then c must satisfy the equation $f^* c[g] = c[f^* g]$. Moreover, we require that c depends smoothly on the Riemannian metric g . In fact, one can prove such locally covariant function c should depend at every point on finite jets of the metric and c has an invariance property under changes of coordinates [35, section 8 p. 76]. This definition is partly motivated by [37, 2.4 p. 160] and [5, p. 282] on local index theory and also by the notion of local covariance arising in algebraic Quantum Field Theory [76, 60, 21]. In the usual Wick renormalization for some massive Gaussian free field ϕ on some Riemannian 3-manifold (M, g) with covariance P^{-1} , one first mollifies ϕ

via heat regularization. This yields a random smooth function $\phi_r \stackrel{\text{def}}{=} e^{-rP}\phi$. To renormalize the square ϕ_r^2 , one subtracts to the square $(\phi_r)^2(x)$ of the mollified field at x , the counterterm $a_r(x) = e^{-2rP}P^{-1}(x, x)$ and it is well-known that the difference $(\phi_r)^2(\cdot) - a_r(\cdot)$ will converge as random distribution when $r \downarrow 0$. However, the counterterm $a_r(x) = e^{-2rP}P^{-1}(x, x)$ that we subtracted is **nonlocal** in the metric g at x , it depends on the global Riemannian geometry of (M, g) and not on finite jets of the metric g at x . Hence such a_r is not *locally covariant* in the above sense. Now we observe that the diagonal value $e^{-2rP}P^{-1}(x, x)$ has an asymptotic expansion of the form:

$$e^{-2rP}P^{-1}(x, x) \sim \frac{1}{4\pi^{\frac{3}{2}}r^{\frac{1}{2}}} + \mathcal{O}(1).$$

If instead of subtracting the diagonal value of $e^{-2rP}P^{-1}$ one subtracted its **singular part**: $(\phi_r)^2(\cdot) - \frac{1}{4\pi^{\frac{3}{2}}r^{\frac{1}{2}}}$, then one would still get a random distribution at the limit when $r \downarrow 0$ but the covariantly renormalized Wick square : ϕ^2 : would no longer have zero expectation. So one may wonder why is it so important to subtract only locally covariant quantities ? The answer lies in the deep notion of locality in quantum field theory. It is a folklore result in quantum fields on curved backgrounds that subtracting non locally covariant counterterms is incompatible with locality in the sense of Atiyah-Segal. Let us quote the beautiful discussion on the regularization of tadpoles and the relation with Atiyah-Segal gluing in [58, 1.2 p. 1852]: *"In various treatments of scalar theory, tadpole diagrams were set to zero (this corresponds to a particular renormalization scheme – in flat space, this is tantamount to normal ordering, ...). However, in our framework this prescription contradicts locality in Atiyah-Segal sense, ... One good solution is to prescribe to the tadpole diagrams the zeta-regularized diagonal value of the Green's function. We prove that assigning to a surface its zeta-regularized tadpole is compatible with locality, ... However there are other consistent prescriptions (for instance, the tadpole regularized via point-splitting and subtracting the singular term, ...). This turns out to be related to Wilson's idea of RG flow in the space of interaction potentials, ..."* We refer the interested reader to [58, Section 5 p. 1885] for further details on this central topic of quantum fields on curved backgrounds. So in the present paper, we follow a similar strategy as in the previous example and try to subtract only locally covariant quantities, in fact we shall see that we subtract universal quantities that do not even depend on the metric g .

5.2.2 The quadratic term. The argument used for the study of \mathfrak{I} , without regularization, works for the study of \mathfrak{V}_r , with regularization. Since ξ_r is regularized in space \mathfrak{I}_r is here a function on $[0, T] \times M$. Set

$$a_r(z) \stackrel{\text{def}}{=} \mathbb{E}[\mathfrak{I}_r^2(z)], \quad \overline{\mathfrak{V}}_r(z) \stackrel{\text{def}}{=} \mathfrak{I}_r^2(z) - a_r(z).$$

To treat the regularity of $\overline{\mathfrak{V}}$, we will follow a similar strategy as for \mathfrak{I} and try to control

$$\mathbb{E}[(P_k^i \overline{\mathfrak{V}})(t, x)^2]$$

when i becomes large and uniformly in $x \in M$. Denote by $G_r^{(2)}(t-s)$ the operator with kernel

$$\left(\frac{e^{-(2r+|t-s|)P}}{P}(x, y) \right)^2.$$

The operators $G_r^{(2)}(0)$ take values in $\Psi^{-1}(M)$ and $G_r^{(2)}(t)$ is of order $|t|^{\gamma/2}$ in $\Psi^{-1+\gamma}(M)$ since $e^{-|t|P}P^{-1}$ is of order $|t|^\gamma$ in $\Psi^{-2+2\gamma}(M)$. This holds uniformly in $r \in [0, 1]$. By definition, and using the Wick formula, we can compute the quantity $\mathbb{E}[P_k^i \overline{\mathfrak{V}}(t, x)^2]$, which is equal to

$$\begin{aligned} & \mathbb{E}[(P_k^i \overline{\mathfrak{V}})(t, x)^2] \\ &= \int_{M^2} G^{(2)}(t_1 - t_2, y_1, y_2) P_k^i(x, y_1) P_k^i(x, y_2) dv(y_1) dv(y_2) \\ &\lesssim \int_{(\mathbb{R}^3)^2} G_{i,i}^{(2)}(t_1 - t_2, y_1, y_2) \psi_i(\tilde{x})^2 \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2} (2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}) \\ &\lesssim 2^{2k} \int_{(\mathbb{R}^3)^2} |y_1 - y_2|^{-2} \psi_i(\tilde{x})^2 \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2} (2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x}), \end{aligned}$$

for some integrals over $(\mathbb{R}^3)^2$ with respect to $dy_1 dy_2$, and where again $\tilde{x} = \kappa_i(x)$ and $G_{i,i}^{(2)} = \kappa_{i*}^{\otimes 2} G^{(2)}$. We used the estimate

$$|G_{i,i}^{(2)}(t_1 - t_2, x_1, x_2)| \lesssim |x_1 - x_2|^{-2}$$

and the changes of variables

$$y_1 \mapsto 2^{-i}(y_1 - \tilde{x}) + \tilde{x}, \quad y_2 \mapsto 2^{-i}(y_2 - \tilde{x}) + \tilde{x}.$$

We can now conclude since the integral

$$\int_{(\mathbb{R}^3)^2} |y_1 - y_2|^{-2} \psi_i(\tilde{x})^2 \psi(|y_1 - \tilde{x}|) \psi(|y_2 - \tilde{x}|) (\kappa_{i*} \chi_i)^{\otimes 2} (2^{-k}(y_1 - \tilde{x}) + \tilde{x}, 2^{-k}(y_2 - \tilde{x}) + \tilde{x})$$

with respect to $dy_1 dy_2$ is bounded uniformly in $x \in \text{supp}(\psi_i)$ because $\psi(|y - \tilde{x}|)$ is a Schwartz function of y .

We conclude as above from Kolmogorov's regularity theorem that $\bar{\mathcal{V}}_r \in C_T^\eta C^{-1-\varepsilon}(M)$ for all $\eta > 0, \varepsilon > 0$, uniformly in $r \in (0, 1]$, and we further get from the r -uniform and continuity of the different quantities as functions of r the convergence of $\bar{\mathcal{V}}_r$ in $C_T^\eta C^{-1-\varepsilon}(M)$ to a limit which we denote by $\bar{\mathcal{V}}$.

In the present paper, we choose to define the **renormalization in a locally covariant way** with respect to the Riemannian metric g . Therefore, we shall subtract from $\hat{\mathcal{V}}_r^2$ only the **singular part** of the function $a_r(z) = \mathbb{E}[\hat{\mathcal{V}}_r^2(z)]$. We actually prove that this singular part a_r is actually some universal constant. The function $a_r(z)$ differs from the constant a_r , but for $z = (t, x)$ we have

$$a_r(z) = \frac{e^{-2rP}}{P}(x, x) = \int_{2r}^1 e^{-sP}(x, x) ds + b(x),$$

where b is a smooth function. The small time asymptotics of the heat kernel then tells us that $e^{-sP}(x, x) = \frac{1}{(4\pi s)^{\frac{3}{2}}} + \mathcal{O}(s^{-\frac{1}{2}})$, so that the function $a_r(\cdot) - a_r$ is indeed bounded, uniformly in $r \in (0, 1]$. To prove that the difference $a_r(\cdot) - a_r$ is smooth, we rely on the description of the heat kernel of [41, def 2.1 p. 6]. In local coordinates in some open subset U , the heat kernel can be represented as $s^{-\frac{3}{2}} A(a, \frac{x-y}{\sqrt{s}}, y)$ where $A \in C^\infty([0, +\infty)_{\frac{1}{2}} \times \mathbb{R}^3 \times U)$. Hence $a_r(x) - a_r = \int_{2r}^1 s^{-\frac{3}{2}} (A(\sqrt{s}, 0, x) - A(0, 0, x)) ds$ converges together with all its derivatives in x since $\partial_x^\beta s^{-\frac{3}{2}} (A(\sqrt{s}, 0, x) - A(0, 0, x)) = \mathcal{O}(s^{-\frac{1}{2}})$ for all multi-indices β . Hence the convergent integral defining $a_r(x) - a_r$ depends smoothly on $x \in U$.

The convergence of $\mathcal{V}_r \stackrel{\text{def}}{=} \hat{\mathcal{V}}_r^2 - a_r$ in $C_T^\eta C^{-1-\varepsilon}(M)$ to a limit which we denote by \mathcal{V} follows as a consequence. We note that as a random variable, \mathcal{V} is not a homogeneous element in the chaos of order 2 of the Gaussian noise since we did not subtract the full expectation. However it differs from a homogeneous element by a deterministic smooth function hence \mathcal{V} has moments of any order $1 \leq r < \infty$ that are equivalent to its second moment. In the sequel, we will always prove stochastic estimates for homogeneous elements in Wiener chaoses and then justify why the locally covariant renormalization, subtracting only universal local quantities, still yields a stochastic object with the same analytic properties.

5.2.3 The cubic term. For the stochastic term \mathfrak{V}° , we do not bound the regularity by hands anymore and use the microlocal machinery. Recall that our aim is to find the range of $\gamma \in \mathbb{R}$ so that

$$\mathbb{E}[\mathcal{Q}_x^\gamma(\mathfrak{V}^\circ(t, \cdot), \mathfrak{V}^\circ(t, \cdot))] < \infty$$

uniformly in $x \in M$. By application of Wick's Theorem we have

$$\begin{aligned} & \mathbb{E}[\mathcal{Q}_x^\gamma(\mathfrak{V}^\circ(t, \cdot), \mathfrak{V}^\circ(t, \cdot))] \\ &= \int_{(-\infty, t]^2 \times M^4} G^{(3)}(s_1 - s_2, z_1, z_2) e^{-|t-s_1|P}(y_1, z_1) e^{-|t-s_2|P}(y_2, z_2) \mathcal{Q}_x^\gamma(y_1, y_2), \end{aligned}$$

for an integral with respect to $dv(y_1, y_2, z_1, z_2) ds_1 ds_2$, where we denote by $G_r^{(3)}(t_1 - t_2)$ the operator with kernel

$$\left(\frac{e^{-(2r+|t_1-t_2|)P}}{P}(x, y) \right)^3.$$

This operator belongs to $\mathcal{S}_\Gamma^{-3}(\mathbb{R}^2 \times M^2)$ with

$$\Gamma = N^*\left(\{t = s\} \times \mathbf{d}_2 \subset \mathbb{R}^2 \times M^2\right) \cup N^*\left(\{t = s\} \subset \mathbb{R}^2 \times M^2\right).$$

By the general theory described above, the only irreducible subgraph of the Feynman graph is the graph itself, therefore the Feynman amplitude is well-defined on $(-\infty, t]^2 \times M^4 \setminus \Sigma$ where the singular locus Σ corresponds to the generalized diagonal

$$\underbrace{\{y_1 = y_2 = z_1 = z_2 = x\}}_{\text{codim}_w=12} \cap \underbrace{\{s_1 = s_2 = t\}}_{\text{codim}_w=4} \subset (-\infty, t]^2 \times M^4$$

which has weighted codimension 16. The sum of the weak homogeneities of the kernels in the Feynman amplitude is given by

$$-3 + 2(-3) - 6 - 2\gamma,$$

so that is finite for $\gamma < \frac{1}{2}$ by Theorem 27.

To conclude this section, let us now discuss the general strategy for larger random fields. First, a general object τ_m lying in the p -th inhomogeneous Gaussian chaos can be expanded onto the p first homogeneous chaoses, so that its covariance can be controlled by the covariances of the projections and subsequently by some amplitudes indexed by Feynman graphs, denoted \mathcal{A}_{mn} for $n = p, p-2$, etc. The next step is to localize the amplitudes in local charts using the partition of unity $1 = \sum_{i \in I} \chi_i$ subordinated to the cover $M = \cup_{i \in I} U_i$, and to control for every $i \in I$ the localized version \mathcal{A}_{mn}^i . The amplitudes \mathcal{A}_{mn} correspond to products of kernels, and the aim is to identify the range of γ for which they are well defined using item **(a)** of Theorem 27. However, two difficulties occur:

- (a) Because it contains a divergent subgraph, the term coming from the lowest chaos, \mathcal{A}_{m0} or \mathcal{A}_{m1} , can not be directly handled using item **(a)** of Theorem 27, and rather requires the use of item **(b)**: this amounts to subtracting a local counterterm.
- (b) The terms \mathcal{A}_{mn} coming from higher chaoses do not contain any divergent subgraph and could therefore be defined using item **(a)** of Theorem 27. However, due to the fact that \mathcal{A}_{mn} contains a subgraph including \mathcal{Q}_x^γ with worse weak homogeneity than the whole graph, this would deteriorate the value of γ . This last case require a special treatment leveraging the precise microlocal properties of $[\odot]$, which is performed in Section 5.4.

5.3 Diagrammatic notation for higher chaoses

To handle the quartic and quintic terms, we first need to introduce more notation. From Appendix A, we define a family $(P_k^i, \tilde{P}_k^i)_{k \in \mathbb{N}, i \in I}$ of generalized Littlewood-Paley-Stein projectors indexed by the frequency 2^k and the chart index $i \in I$. The localized resonant product \odot_i , where i is the chart index, is defined in the appendix as $u \odot_i v = \sum_{|k-\ell| \leq 1} P_k^i(u) \tilde{P}_\ell^i(v)$. The goal of the present subsection is to deal with the localized objects

$$\tau_1 = \mathfrak{Y}_r \odot_i \mathfrak{I}_r, \quad \bar{\tau}_2 = \mathfrak{Y}_r \odot_i \mathfrak{V}_r - \chi_i \frac{b_r}{3}, \quad \bar{\tau}_3 = \chi_i |\nabla \mathfrak{Y}_r|^2 - \chi_i \frac{b_r}{3}, \quad \bar{\tau}_4 = \mathfrak{Y}_r \odot_i \mathfrak{V}_r - \chi_i b_r \mathfrak{I}_r,$$

where $\chi_i \in C_c^\infty(U_i)$ is the cut-off function used to define \odot_i , and find for each of them the range of exponents γ for which our test criterion is verified.

Note that the trees are defined with the localized resonant term \odot_i made with these projectors and recall from Appendix A that these resonant terms are not commutative, in the sense that $A \odot_i B \neq B \odot_i A$. However, we do not have to worry about the definitions of τ_1 and $\bar{\tau}_2$, since the analytic properties of P_k^i and \tilde{P}_k^i are similar, which entails that for every A, B , the construction of the renormalized part of $B \odot_i A$ is totally equivalent to that of $A \odot_i B$ which we provide here.

Denote by $[\odot_i](x, y, z)$ the kernel of the localized resonant operator \odot_i on the chart $U_i \subset M$. Recall that we write

$$\underline{\mathcal{L}}^{-1}(t-s, x, y) = \mathbf{1}_{(-\infty, t]}(s) e^{-(t-s)P}(x, y)$$

and

$$\begin{aligned}
G_r^{(p)}(t-s, x, y) &= \left(\left\{ e^{-|t-s|P} P^{-1} \right\} (x, y) \right)^p, & 1 \leq p \leq 3, \\
[\odot_i](x, y, z) &= \sum_{|k-\ell| \leq 1} P_k^i(x, y) \tilde{P}_\ell^i(x, z), & i \in I, \\
\mathcal{Q}_x^\gamma(y, z) &= \sum_{i \in I} \sum_{k \geq -1} 2^{2k\gamma} P_k^i(x, y) P_k^i(x, z), & \gamma \in \mathbb{R}, x \in M.
\end{aligned}$$

These kernels are in some spaces of the form \mathcal{S}_Γ^a for different ambient spaces, scaling exponents a and wavefront sets as follows.

- The kernel $\underline{\mathcal{L}}^{-1}(t-s, x, y)$ has scaling exponent -3 and wavefront set $N^*(\{t=s\} \times \mathbf{d}_2 \subset \mathbb{R}^2 \times M^2)$.
- The kernel $G_r^{(p)}(t-s, x, y)$ have scaling exponent $-p$ and wavefront set $N^*(\{t=s\} \subset \mathbb{R}^2 \times M^2) \cup N^*(\{t=s\} \times \mathbf{d}_2 \subset \mathbb{R}^2 \times M^2)$.
- The kernel $[\odot_i](x, y, z)$ has scaling exponent -6 and wavefront set $N^*(\{x=y=z\} \subset M^3)$.
- The kernel $\mathcal{Q}_x^\gamma(y_1, y_2)$ has scaling exponent $-6 - 2\gamma$ and wavefront set $T_{(x,x)}^*(M^2)$.

Beware that in the case of \mathcal{Q}_x^γ , the singular locus depends on the marked point $x \in M$. In this list the kernels on $(\mathbb{R} \times M)^2$ satisfy a local diagonal bound of the form

$$|\partial_{\sqrt{t}, \sqrt{s}, y_1, y_2}^\alpha K| \lesssim \left(\sqrt{|t-s|} + |y_1 - y_2| \right)^{-a-|\alpha|}$$

for the corresponding scaling exponent a . We also often see the kernels $\underline{\mathcal{L}}^{-1}$, \mathcal{Q}_x^γ and $G_r^{(1)}$ as some time dependent space operators $X(t_1 - t_2)$ whose kernels are then given by $X(t_1 - t_2, y_1, y_2)$. The proof of the above microlocal bounds is done in detail in our companion work [10].

We use a pictorial representation of the τ_i in which the black dot \bullet represents a resonant operator and the noises are coloured circles. In a given graph noises of the same color are integrated outside all diagonals of the corresponding set of variables. In the present subsection, it will be convenient to first discuss stochastic estimates for Wick ordered elements which live in homogeneous Wiener chaoses, then explain why our locally covariant renormalization yields stochastic elements that differ from the Wick renormalized ones only up to higher regularity elements. The Wiener chaos decomposition of the τ_i is

$$\begin{aligned}
\tau_1 &= \text{[diagram]} = \text{[diagram]} + 3 \text{[diagram]}, & \tau_2 &= \text{[diagram]} = \text{[diagram]} + 2 \text{[diagram]} + 2 \text{[diagram]}, \\
\tau_3 &= \text{[diagram]} = \text{[diagram]} + 2 \text{[diagram]} + 2 \text{[diagram]}, & \tau_4 &= \text{[diagram]} = \text{[diagram]} + 6 \text{[diagram]} + 6 \text{[diagram]}.
\end{aligned}$$

where black edges stand for the kernel of $\underline{\mathcal{L}}^{-1}$ and green edges stand for the kernel of $\Delta \underline{\mathcal{L}}^{-1}$. The reason for the presence of the Laplacian operators in τ_3 will be made clear in Section 5.5.4.

We write for $m \in \{1, 2, 3\}$ $\tau_m = \tau_{m4} + \tau_{m2} + \tau_{m0}$ with τ_{mn} in the homogeneous chaos of degree $n \in \{0, 2, 4\}$ and accordingly $\tau_4 = \tau_{45} + \tau_{43} + \tau_{41}$. By orthogonality

$$\mathbb{E} \left[\mathcal{Q}_x^\gamma(\tau_m(t, \cdot), \tau_m(t, \cdot)) \right] = \sum_{n \in \mathbb{N}} \mathbb{E} \left[\mathcal{Q}_x^\gamma(\tau_{mn}(t, \cdot), \tau_{mn}(t, \cdot)) \right].$$

Remember that we are not interested in the τ_m themselves but in their renormalized versions generically written $\bar{\tau}_m = \tau_m - c_{m,r}$, with $c_{m,r}$ for ‘counterterm’, for which we still have the

orthogonality relation

$$\begin{aligned} & \mathbb{E} \left[\mathcal{Q}_x^\gamma(\bar{\tau}_m(t, \cdot), \bar{\tau}_m(t, \cdot)) \right] \\ &= \sum_{n \geq 2} \mathbb{E} \left[\mathcal{Q}_x^\gamma(\tau_{mn}(t, \cdot), \tau_{mn}(t, \cdot)) \right] + \mathbb{E} \left[\mathcal{Q}_x^\gamma((\tau_{mp}(t, \cdot) - c_{k,r}), (\tau_{mp}(t, \cdot) - c_{m,r})) \right], \end{aligned}$$

where $p = 0$ for $m \in \{1, 2, 3\}$ and $p = 1$ for $m = 4$. Also, note that each element τ_{mn} in the Wiener chaos can be written as $F_n(\cdot; \xi^{\otimes n} \cdot)$ where each $F_n : C^\infty(\mathbb{R} \times M)^{\otimes n} \rightarrow \mathcal{D}'(\mathbb{R} \times M)$ is a multilinear functional valued in distributions involving iterated integrals of the heat operator, products represented by the trees. The following elementary well-known result tells us that we only need to bound very symmetric Feynman diagrams to bound the preceding expectations.

Lemma 34 – *Let $F \in L^2(M^n)$ be a function of n variables on some compact Riemannian manifold (M, g) , ξ the white noise on (M, g) . Then:*

$$\mathbb{E} \left[\langle F, \cdot; \xi^{\otimes n} \cdot \rangle^2 \right] \leq \|F\|_{L^2(M^n)}^2.$$

Proof – We define the symmetrization operator S_n as $S_n(\varphi)(x_1, \dots, x_n) = \frac{1}{n!} \sum_{\sigma \in S_n} \varphi(x_{\sigma(1)}, \dots, x_{\sigma(n)})$. Note the important fact that S_n is self-adjoint on $L^2(M^n)$ hence S_n is the orthogonal projector on symmetric L^2 functions. By the Itô isometry property,

$$\mathbb{E} \left(\langle F, \cdot; \xi^{\otimes n} \cdot \rangle^2 \right) = \mathbb{E} \left(\langle S_n F, \cdot; \xi^{\otimes n} \cdot \rangle^2 \right) = \|S_n F\|_{L^2(M^n)}^2$$

hence to prove the Lemma it suffices to note that $\|S_n F\|_{L^2(M^n)} \leq \|F\|_{L^2(M^n)}$ which is obvious since S_n is an orthogonal projector. \triangleright

Each expectation can therefore be bounded by a quantity of the form

$$\int_{(\mathbb{R} \times M)^{n+2}} F_n(x_1; y_1, \dots, y_n) F_n(x_2; y_1, \dots, y_n) \mathcal{Q}_x^\gamma(x_1, x_2) dy dx,$$

which can be represented by some mirror symmetric Feynman diagram. Using a **purple edge** for the kernel of the operator \mathcal{Q}_x^γ , a dotted edge for $\underline{\mathcal{L}}^{-1}$, a black edge for $G_r^{(1)}$, and a **green edge** for $\Delta \underline{\mathcal{L}}^{-1}$, we have

$$\begin{aligned} \mathbb{E} \left[\mathcal{Q}_x^\gamma(\bar{\tau}_1(t, \cdot), \bar{\tau}_1(t, \cdot)) \right] &\lesssim \text{diagram 1} + \text{diagram 2} =: \mathcal{G}_{14} + \mathcal{G}_{12}, \\ \mathbb{E} \left[\mathcal{Q}_x^\gamma(\bar{\tau}_2(t, \cdot), \bar{\tau}_2(t, \cdot)) \right] &\lesssim \text{diagram 3} + \text{diagram 4} + \mathbb{E} \left[\mathcal{Q}_x^\gamma((\tau_{20}(t, \cdot) - \chi_i \frac{b_r}{3}), (\tau_{20}(t, \cdot) - \chi_i \frac{b_r}{3})) \right] \\ &=: \mathcal{G}_{24} + \mathcal{G}_{22} + \mathcal{G}_{20}, \\ \mathbb{E} \left[\mathcal{Q}_x^\gamma(\bar{\tau}_3(t, \cdot), \bar{\tau}_3(t, \cdot)) \right] &\lesssim \text{diagram 5} + \text{diagram 6} + \mathbb{E} \left[\mathcal{Q}_x^\gamma((\tau_{30}(t, \cdot) - \chi_i \frac{b_r}{3}), (\tau_{30}(t, \cdot) - \chi_i \frac{b_r}{3})) \right] \\ &=: \mathcal{G}_{34} + \mathcal{G}_{32} + \mathcal{G}_{30}, \\ \mathbb{E} \left[\mathcal{Q}_x^\gamma(\bar{\tau}_4(t, \cdot), \bar{\tau}_4(t, \cdot)) \right] &\lesssim \text{diagram 7} + \text{diagram 8} + \mathbb{E} \left[\mathcal{Q}_x^\gamma((\tau_{41}(t, \cdot) - \chi_i b_r \mathring{1}_r), (\tau_{41}(t, \cdot) - \chi_i b_r \mathring{1}_r)) \right] \\ &=: \mathcal{G}_{45} + \mathcal{G}_{43} + \mathcal{G}_{41}. \end{aligned}$$

We will use the notation \mathcal{G}_{mn} for positive quantity represented by the mirror graph associated with τ_{mn} and the notation \mathcal{A}_{mn} for the associated distribution on the corresponding configuration space ; we use Theorem 27 to prove their convergence. We use the word ‘*amplitude*’ to talk about any of the \mathcal{A}_{mn} . The terms involving τ_{20} and τ_{30} are treated differently from the mirror graphs. Recall that the vertex set is partitioned as $V = V' \sqcup V_A$. We always denote by x_a, x_b , etc. the space points associated with the vertices in V' , while in our setting V_A will always contain two triplets $((v_1^j, v_2^j), v_*^j)$ whose associated space points are denoted $((y_1, y_2), y_*)$ and $((z_1, z_2), z_*)$ for $n_{\mathcal{G}} = 2$.

Our aim is now to control all of the eleven amplitudes \mathcal{A}_{mn} using Theorem 27. It turns out that this naive approach will only work for \mathcal{A}_{12} and, after renormalization, \mathcal{A}_{20} and \mathcal{A}_{30} . All the remaining eight amplitudes need a special care for one of their subamplitudes. This analysis is

performed in the next section. Once the problematic subamplitudes are constructed, it will then suffice to plug this knowledge inside the recursion of the proof of Theorem 27, and this will yield the existence of \mathcal{A}_{mn} .

5.4 Modification of the inductive proof

As anticipated, while Theorem 27 turns out to be sufficient to handle some graphs and subgraphs of the graphs introduced in Section 5.3, yet most of the total amplitudes need a particular care in order to be controlled:

- (a) The amplitudes of the graphs \mathcal{G}_{14} , \mathcal{G}_{24} , and \mathcal{G}_{45} are given for $(m, n) \in \{(1, 4), (2, 4), (4, 5)\}$ by

$$\mathcal{A}_{mn} = [\odot](y_*, y_1, y_2) [\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(k)}(0, y_1, z_1) \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) G_r^{(\ell)}(s_a - s_b, x_a, x_b),$$

where $(k, \ell) = (1, 3)$ for \mathcal{G}_{14} , $(k, \ell) = (2, 2)$ for \mathcal{G}_{24} , and $(k, \ell) = (2, 3)$ for \mathcal{G}_{45} , and have a subgraph with amplitude reading

$$\mathcal{C}^{(k)}(y_*, y_1, y_2, z_*, z_1, z_2) \stackrel{\text{def}}{=} [\odot](y_*, y_1, y_2) [\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(k)}(0, y_1, z_1)$$

which is *less convergent* (in the sense of the power counting from Theorem 27) than the whole graph: since this is a one loop subgraph, naively applying the power counting criterion from Theorem 27) on the deepest diagonal $\{y_* = y_1 = y_2 = z_* = z_1 = z_2 = x\}$ would result in a potential loss in the regularity γ . The exact same phenomenon also occurs for the amplitude (5.16) of the graph \mathcal{G}_{34} whose subamplitude

$$\tilde{\mathcal{C}}^{(2)}(y_*, y_1, y_2, z_*, z_1, z_2, x_c, x_d, s_c, s_d) \stackrel{\text{def}}{=} [\odot](y_*, y_1, y_2) [\odot](z_*, z_1, z_2) \times \mathcal{Q}_x^\gamma(y_*, z_*) \Delta \underline{\mathcal{L}}^{-1}(t - s_c, y_1, x_c) \Delta \underline{\mathcal{L}}^{-1}(t - s_d, z_1, x_d) G_r^{(2)}(s_c - s_d, x_c, x_d)$$

is also *less convergent* than the whole amplitude, which equally forbids the use of a naive power counting argument.

- (b) The amplitudes of the graphs \mathcal{G}_{22} and \mathcal{G}_{43} are given for $(m, n) \in \{(2, 2), (4, 3)\}$ by

$$\mathcal{A}_{mn} = [\odot](y_*, y_1, y_2) [\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(1)}(0, y_1, z_1) \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) G_r^{(1)}(t - s_a, y_1, x_a) G_r^{(1)}(t - s_b, y_2, x_b) G_r^{(\ell)}(s_a - s_b, x_a, x_b),$$

where $\ell = 1$ for \mathcal{G}_{22} and $\ell = 2$ for \mathcal{G}_{43} , and have a subgraph with amplitude $\mathcal{C}^{(1)}$ which is *less convergent* than the whole graph (marginally for \mathcal{G}_{43}). Again the same phenomenon occurs for the amplitude (5.17) of the graph \mathcal{G}_{32} that has a subamplitude $\tilde{\mathcal{C}}^{(1)}$ defined as $\tilde{\mathcal{C}}^{(2)}$ above with $G_r^{(2)}$ replaced with $G_r^{(1)}$ which is *less convergent* than the whole amplitude.

- (c) The amplitude of the graph \mathcal{G}_{41} , which is given by

$$(5.9) \quad \mathcal{A}_{41} = \mathcal{Q}_x^\gamma(y_*, z_*) \mathcal{B}(y_*, y_1, y_2, x_a, s_a) \mathcal{B}(z_*, z_1, z_2, x_b, s_b) G_r^{(1)}(s_a - s_b, x_a, x_b),$$

where

$$(5.10) \quad \mathcal{B}(y_*, y_1, y_2, x_a, s_a) \stackrel{\text{def}}{=} [\odot](y_*, y_1, y_2) G_r^{(2)}(t - s_a, y_1, x_a) \underline{\mathcal{L}}^{-1}(t - s_a, y_1, x_a) - b_r \delta_{\mathbf{d}_4},$$

requires a renormalization to define the subamplitudes

$$\mathcal{B}(y_*, y_1, y_2, x_a, s_a) \text{ and } \mathcal{B}(z_*, z_1, z_2, x_b, s_b).$$

Fortunately, for the items (a) and (b), the presence of the kernel $[\odot]$ does prevent this loss in the value of γ from occurring, due to a subtle mechanism which is not taken into account by the recursion of Theorem 27. Indeed, while the kernel $[\odot](y_*, y_1, y_2)$ is divergent on the deep diagonal $\{y_* = y_1 = y_2\}$, it is *smooth* as long as one of the three points is distinct from the two others and applying any smoothing operator at one of its input variables, indexed by (y_1, y_2) , makes the *whole kernel smooth*. In particular, provided the kernel $[\odot]$ is tested against a smooth kernel in one of its two input variables (y_1, y_2) , then it is smooth in the two remaining variables – see Lemma 35. In terms of multilinear operators, this translates the important fact that

$$\text{for every } f \in \mathcal{D}'(M) \text{ and } g \in C^\infty(M), f \odot g \in C^\infty(M).$$

To take both this effect and the renormalization of the divergent subamplitudes of \mathcal{A}_{41} into account, we introduce in this section a minor modification of the induction in the proof of Theorem 27. More precisely, the strategy is to first construct the subamplitudes by hand, and then to inject our knowledge of the subamplitudes into the induction yielding Theorem 27. Indeed, if an amplitude \mathcal{A} contains the subamplitude \mathcal{B} , $\mathcal{C}^{(k)}$ or $\tilde{\mathcal{C}}^{(k)}$, then we will add the estimates on the subamplitude to the hypothesis of Theorem 27 when applying it to \mathcal{A} , which will yield control over \mathcal{A} avoiding the issue which would naively be caused by the subamplitude.

5.4.1 The smoothing effect from the kernel of the resonant product. This section is devoted to items (a) and (b) above, that is to say with the construction of the subamplitudes $\mathcal{C}^{(k)}$ and $\tilde{\mathcal{C}}^{(k)}$ for $k \in \{1, 2\}$. Note that by a naive application of Theorem 13, it would be possible to construct these subamplitudes for every $\gamma < -k/2$. However, to avoid losing some regularity, the aim is to actually construct them for any $\gamma \in \mathbb{R}$. This is possible thanks to the lemma below, which expresses the smoothing property of the kernel $[\odot]$. Informally, in the sequel, we denote the smoothing effect described in the following lemma as *shielding effect*.

Lemma 35 – Fix a chart label $i \in I$, and define a collection of scaling fields $(\rho_{i,x})_{x \in M}$ as follows: for $x \in U_i$, the scaling field $\rho_{i,x}$ is such that $\kappa_i(e^{-t\rho_{i,x}}z) = e^{-t}(\kappa_i(z) - \kappa_i(x)) + \kappa_i(x)$, while if $x \notin U_i$ we choose for $\rho_{i,x}$ any scaling field with respect to $x \in M \setminus U_i$. Also fix $\Xi_i \in C_c^\infty(M^3)$ such that $\Xi_i(y_*, y_1, y_2)$ has support in y_2 contained in the open chart U_i .

Then, the following holds: for any $x \in M$, $\varphi \in \mathcal{D}(M)$, and any scaling vector field $\rho_{i,x}$ scaling with respect to x chosen as above, the partial integration

$$I_u^i(y_*, y_1) \stackrel{\text{def}}{=} e^{-6u} \int_M (e^{-u\rho_{i,x^*}}[\odot]) (y_*, y_1, y_2) \varphi(y_2) \Xi_i(y_*, y_1, y_2) dv(y_2)$$

is an element of $C^\infty(M^2)$ bounded **uniformly** in $u \in [0, +\infty)$ and $x \in M$.

Proof – Recall that in the chart $i \in I$, we have

$$[\odot_i](y_*, y_1, y_2) = \sum_{|k-l| \leq 1} 2^{3k+3l} \frac{\psi_i(y_*) \tilde{\psi}_i(y_*)}{|g_i(y_1)| |g_i(y_2)|} \psi(2^k |y_1 - y_*|) \psi(2^l |y_2 - y_*|) \kappa_{i^*}^{\otimes 2}(\chi_i \otimes \tilde{\chi}_i)(y_1, y_2),$$

where $|g_i| \stackrel{\text{def}}{=} \sqrt{\det(\kappa_{i^*} g)}$, and that the family

$$e^{-6u} (e^{-u\rho_{i,x^*}}[\odot_i]), \quad u \in [0, +\infty), \quad x \in M,$$

is bounded in $\mathcal{D}'_{N^*(\mathfrak{d}_3)}(M^3)$. We first assume without loss of generality that $x \in U_i$ then we choose a **very specific scaling vector field** adapted to our chart structure as

$$e^{-u\rho_{i,x}}(y_*, y_1, y_2) \stackrel{\text{def}}{=} (e^{-u}(y_* - x) + x, e^{-u}(y_1 - x) + x, e^{-u}(y_2 - x) + x),$$

so the scaling is linear in the chart $\kappa_i : U_i \rightarrow \mathbb{R}^3$. Here and throughout the proof, with a slight abuse of notation, for any $a \in M$ we also use the notation a to denote $\kappa_i(a)$. With the same above of notation, we identify I_u^i with its localization on U_i , which thus reads

$$\begin{aligned} I_u^i(y_*, y_1) &= \sum_{|k-l| \leq 1} 2^{3k+3l} e^{-6u} \int_{\mathbb{R}^3} \psi_i(e^{-u}(y_* - x) + x) \tilde{\psi}_i(e^{-u}(y_* - x) + x) \\ &\quad \times \frac{\psi(2^k e^{-u} |y_1 - y_*|) \psi(2^l e^{-u} |y_2 - y_*|)}{|g_i|(e^{-u}(y_1 - y_*) + y_*) |g_i|(e^{-u}(y_2 - y_*) + y_*)} \\ &\quad \times \kappa_{i^*}^{\otimes 2}(\chi_i \otimes \tilde{\chi}_i)(e^{-u}(y_1 - x) + x, e^{-u}(y_2 - x) + x) \varphi(y_2) \Xi_i(y_*, y_1, y_2) dy_2, \end{aligned}$$

Our goal is to prove that the above function I_u^i is smooth in the variables (y_*, y_1) uniformly in the scale parameter $u \geq 0$ and in $x \in \mathbb{R}^3$. At this point in our proof, it is crucial that the scaling by $\rho_{i,x}$ is the same scaling as the one used to define the Littlewood-Paley blocks. We consider the series with $k = l$ in I_u^i , other series are treated similarly, which reads

$$\sum_{k=0}^{\infty} 2^{6k} e^{-6u} \psi(2^k e^{-u} |y_1 - y_*|) \psi(2^k e^{-u} |y_2 - y_*|) \in \mathcal{S}'(\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{R}^3),$$

and write $e^{-u} = 2^{-\ell}$. The above family rewrites

$$\sum_{k=0}^{\infty} 2^{6(k-\ell)} \psi(2^{k-\ell}|y_1 - y_*|) \psi(2^{k-\ell}|y_2 - y_*|) = \sum_{k=-\ell}^{\infty} 2^{6k} \psi(2^k|y_1 - y_*|) \psi(2^k|y_2 - y_*|),$$

the summation getting shifted. Moreover, introduce the family of smooth functions

$$F_{\ell,x}(y_*, y_1, y_2)$$

defined by the formula

$$\frac{\psi_i(2^{-\ell}(y_* - x) + x) \tilde{\psi}_i(2^{-\ell}(y_* - x) + x) \kappa_{i_*}^{\otimes 2}(\chi_i \otimes \tilde{\chi}_i)(2^{-\ell}(y_1 - x) + x, 2^{-\ell}(y_2 - x) + x) \varphi(y_2) \Xi_i(y_*, y_1, y_2)}{|g_i|(2^{-\ell}(y_1 - y_*) + y_*) |g_i|(2^{-\ell}(y_2 - y_*) + y_*)}.$$

They are bounded in C^∞ as functions of the variables y_*, y_1, y_2 uniformly in ℓ, x , and have compact support in the variable y_2 in some compact $K \Subset \mathbb{R}^3$. By Plancherel's theorem applied to the variable y_2 , we have

$$\begin{aligned} \sum_{k=-\ell}^{\infty} 2^{6k} \int_{\mathbb{R}^3} \psi(2^k|y_1 - y_*|) \psi(2^k|y_2 - y_*|) F_{\ell,x}(y_*, y_1, y_2) dy_2 \\ = \sum_{k=-\ell}^{\infty} 2^{3k} \int_{\mathbb{R}^3} e^{i\xi_2 \cdot y_*} \psi(2^k|y_1 - y_*|) \eta(2^{-k}\xi_2) \mathcal{F}_{y_2} F_{\ell,x}(y_*, y_1, \xi_2) d\xi_2, \end{aligned}$$

where we recall that η denotes the Fourier transform of $\psi \circ |\cdot|$. Using the fact that $F_{\ell,x}$ is smooth, which entails the bound $\partial_{y_*, y_1}^\alpha \mathcal{F}_{y_2} F_{\ell,x}(y_*, y_1, \xi_2) = \mathcal{O}(\langle \xi_2 \rangle^{-2N})$, we obtain

$$\sum_{k=1}^{\infty} 2^{3k} \int_{\mathbb{R}^3} e^{i\xi_2 \cdot y_*} \psi(2^k|y_1 - y_*|) \eta(2^{-k}\xi_2) \mathcal{F}_{y_2} F_{\ell,x}(y_*, y_1, \xi_2) d\xi_2 \lesssim \sum_{k=1}^{\infty} 2^{(3-N)k} \int_{\mathbb{R}^3} \langle \xi_2 \rangle^{-N} d\xi_2,$$

which choosing N large enough is absolutely convergent. Uniform in $x \in \mathbb{R}^3$ bounds on the derivatives can be obtained in a similar fashion. Regarding the sum over k negative we have

$$\begin{aligned} \left| \sum_{k=-\ell}^0 2^{3k} \int_{\mathbb{R}^3} e^{i\xi_2 \cdot y_*} \psi(2^k|y_1 - y_*|) \eta(2^{-k}\xi_2) \mathcal{F}_{y_2} F_{\ell,x}(y_*, y_1, \xi_2) d\xi_2 \right| \\ \lesssim \sum_{k=0}^{\ell} 2^{-3k} \int_{\mathbb{R}^3} \eta(2^k\xi_2) \langle \xi_2 \rangle^{-N} d\xi_2 \lesssim 1, \end{aligned}$$

uniformly in $\ell \geq 0$. Indeed $\sum_{k=0}^{\ell} 2^{-3k} \eta(2^k\xi_2) \leq \sum_{k=0}^{\ell} \eta(2^k\xi_2) \lesssim 1$ by definition the Littlewood-Paley blocks. This concludes the proof of the claim. \triangleright

Remark 36 – We note that this shielding phenomenon is second microlocal in nature and cannot be captured by classical (“conical”) wave front set analysis. In fact, it can be only captured semiclassically. Let us just explain things in the flat case. Indeed, the wave front set of both $[\odot](y_*, y_1, y_2)$ and $\delta_{y_*}(y_1) \delta_{y_*}(y_2)$ are contained in the conormal of the deepest diagonal

$$\{y_1 = y_2 = y_*; \eta_1 + \eta_2 + \eta_* = 0\}$$

but we see that the Fourier support of $[\odot]$ is smaller than the hyperplane $\{\eta_1 + \eta_2 + \eta_* = 0\}$, it is contained in the subset $\{\eta_1 + \eta_2 + \eta_* = 0, |\eta_1| \simeq |\eta_2|\}$ which is not a closed conical subset ($\{\eta_1 + \eta_2 + \eta_* = 0, \eta_1 \neq 0, \eta_2 \neq 0\}$ is not a closed conical subset) and is responsible for the shielding. The purpose of the above lemma is therefore to capture such phenomenon on manifolds using local charts.

With the above shielding lemma in hand we are ready to construct the subamplitudes $\mathcal{C}^{(k)}$ and $\tilde{\mathcal{C}}^{(k)}$. Note that since these amplitudes contain \mathcal{Q}_x^γ , we aim to construct them both on $\mathbf{d}_6 \setminus \mathbf{d}_{6,x}$ and $\mathbf{d}_{6,x}$, where it has a different weak homogeneity.

Lemma 37 – Fix $k \in \{1, 2\}$. For any $\gamma \in \mathbb{R}$, the distribution $\mathcal{C}^{(k)}$ is well-defined on M^6 . More precisely, we have

$$\mathcal{C}^{(k)} \in \mathcal{S}^{-18-k-2\gamma}(M^6; U_{x_0}) \cap \mathcal{S}_\Gamma^{-12-k}$$

uniformly in $x \in U_{x_0}$, where

$$\begin{aligned} \Gamma = & N^*(\mathbf{d}_6) \cup N^*\{y_* = y_1 = y_2\} \cup N^*\{z_* = z_1 = z_2\} \cup N^*\{y_1 = z_1\} \\ & \cup N^*\{y_* = y_1 = y_2 = z_1\} \cup N^*\{z_* = z_1 = z_2 = y_1\}. \end{aligned}$$

Moreover, a similar statement holds for $\tilde{\mathcal{C}}^{(k)}$, which is well-defined as a distribution on $\mathbb{R}^2 \times M^8$ and belongs to $\mathcal{S}^{-18-k-2\gamma}(\mathbb{R}^2 \times M^8; U_{x_0}) \cap \mathcal{S}_{\tilde{\Gamma}}^{-12-k}$ where $\tilde{\Gamma}$ is a union of conormals defined accordingly.

Remark – With this lemma in hand, we will be able to inject our knowledge on $\mathcal{C}^{(k)}$ and $\tilde{\mathcal{C}}^{(k)}$ inside the recurrence of the proof of Theorem 27, when applied to an amplitude \mathcal{A} containing these subamplitudes. Since they exist for every $\gamma \in \mathbb{R}$, they will therefore not contribute to the upper bound on γ , which will actually come from the scaling of the whole amplitude \mathcal{A} itself as desired.

Proof – We prove the statement for $\mathcal{C}^{(k)}$, the proof for $\tilde{\mathcal{C}}^{(k)}$ being a straightforward modification ((5.15) shows that $\Delta \underline{\mathcal{L}}^{-1}$ actually scales like a Dirac, so that its presence does not affect the argument). Recall that this subamplitude reads

$$\mathcal{C}^{(k)}(y_*, y_1, y_2, z_*, z_1, z_2) = [\odot](y_*, y_1, y_2)[\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(1)}(0, y_1, z_1).$$

We first claim that $\mathcal{C}^{(k)}$ is well-defined as a distribution on M^6 for all $\gamma \in \mathbb{R}$ (and not just $\gamma < -1/2$). Indeed, when testing it against some $\varphi \in \mathcal{D}(M^6)$, one can first perform the partial integration

$$\int_M [\odot](y_*, y_1, y_2) \varphi(y_*, y_1, y_2, z_*, z_1, z_2) dv(y_2).$$

An application of Lemma 35 shows that this expression is a smooth function on M^6 , and testing it against $[\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(1)}(0, y_1, z_1)$ proves the claim.

Regarding the scaling on Γ , observe that

$$\begin{aligned} & \underbrace{\int_M e^{-6u} (e^{-u\rho_i^*} [\odot_i]) (y_*, y_1, y_2) \varphi(y_*, y_1, y_2, z_*, z_1, z_2) dv(y_2)} \\ & \times e^{-(6+k)u} e^{-u\rho_i^*} \left([\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(k)}(0, y_1, z_1) \right) \end{aligned}$$

is bounded since $[\odot](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(k)}(0, y_1, z_1)$ is a tree and the underbraced term is bounded by Lemma 35. Note that we used the fact that \mathcal{Q}_x^γ is weakly homogeneous of degree 0, which is proven in Lemma 28. It follows that

$$e^{-(12+k)u} e^{-u\rho_i^*} \left([\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(k)}(0, y_1, z_1) \right)$$

is bounded in $\mathcal{D}'_\Gamma(M^6)$. Then, by Proposition 15, the same holds with any scaling field ρ with respect to \mathbf{d}_6 , which concludes after summation over $i \in I$ that $\mathcal{C}^{(k)} \in \mathcal{S}_\Gamma^{-12-k}$.

It remains to scale on the deep marked diagonal $\mathbf{d}_{6,x}$. This directly stems from Theorem 14, and from an easy modification of the above discussion, since the only propagator whose scaling degree changes when we scale with respect to \mathbf{d}_6 or $\mathbf{d}_{6,x}$ is \mathcal{Q}_x^γ , as discussed in Lemma 28. Here we need to rely on Proposition 26 which allows to keep the same scaling degrees for the new scaling field ρ_x . \triangleright

5.4.2 Renormalising the divergent subamplitude. This section is devoted to the construction of the subamplitude \mathcal{B} , which is performed in the following Lemma.

Lemma 38 – For any $\gamma < 0$, the distribution \mathcal{B} is well-defined on $M^4 \times \mathbb{R}$. More precisely, we have

$$\mathcal{B} \in \mathcal{S}_\Gamma^{-11},$$

where Γ is equal to

$$\begin{aligned} & N^*(\{s_a = t\}) \cup N^*(\{s_a = t, y_1 = x_a\}) \cup N^*(\{s_a = t, y_2 = x_a\}) \cup N^*(\{y_* = y_1 = y_2\}) \\ & \cup N^*(\{s_a = t, y_1 = y_2 = x_a\}) \cup N^*(\{s_a = t, y^* = y_1 = y_2\}) \cup N^*(\{s_a = t, y^* = y_1 = y_2 = x_a\}). \end{aligned}$$

Remark – Again, with this lemma in hand, we will be able to inject our knowledge on \mathcal{B} inside the recurrence of the proof of Theorem 27, when applied to an amplitude \mathcal{A}_{41} which contains two copies of this subamplitude.

Proof – Recall the definition (5.10) of \mathcal{B} and note that in (5.10), $\delta_{\mathbf{d}_4}$ is the unique distribution on $M^4 \times \mathbb{R}$ supported on $\mathbf{d}_4 = \{s_a = t, y_* = y_1 = y_2 = x_a\}$ and such that for all $\varphi \in C_c^\infty(M^4 \times \mathbb{R})$, one has

$$\begin{aligned} \langle \delta_{\mathbf{d}_4}, \varphi \rangle_{\mathbb{R} \times M^4} &\stackrel{\text{def}}{=} \int_{M^4 \times \mathbb{R}} \delta_{\mathbf{d}_4}(y_*, y_1, y_2, x_a, t, s_a) \varphi(y_*, y_1, y_2, x_a, t, s_a) dv_{M^4 \times \mathbb{R}} \\ &= \int_{y_* \in M} \varphi(y_*, y_*, y_*, y_*, t, t) dv_M(y), \end{aligned}$$

where both $dv_{M^4 \times \mathbb{R}}$ and dv_M are the natural volume forms on the respective Riemannian manifolds, so that $\delta_{\mathbf{d}_4}$ only depends on the choice of Riemannian volume form on M . We denote by $\tilde{\mathcal{B}}$ the unrenormalized amplitude of \mathcal{B} which reads

$$\tilde{\mathcal{B}}(y_*, y_1, y_2, x_a, s_a) \stackrel{\text{def}}{=} [\odot](y_*, y_1, y_2) G_r^{(2)}(t - s_a, y_1, x_a) \underline{\mathcal{L}}^{-1}(t - s_a, y_1, x_a),$$

and verifies $\mathcal{B} = \tilde{\mathcal{B}} - b_r \delta_{\mathbf{d}_4}$. Moreover, to simplify the notation, we write y_a rather than x_a , and therefore focus on $\tilde{\mathcal{B}}(y_*, y_1, y_2, y_a, s_a)$. Finally, throughout the proof, given $y_* \in M$ we use the short-hand notation $z = (y_*, y_*, y_*, y_*, t) \in M^4 \times \mathbb{R}$.

The definition of \mathcal{B} is a consequence of item **(b)** in Theorem 27, and of the discussion performed in Section 5.5.3. Indeed, naively, we know that $\tilde{\mathcal{B}}$ belongs to $\mathcal{S}_\Gamma^{-11}(M^4 \times \mathbb{R}^2 \setminus \mathbf{d}_4)$. But the weighted codimension of \mathbf{d}_4 also equals 11, so we are in case **(b)** of Theorem 13 and our extension requires a renormalization as in Hadamard's finite parts. We thus aim to show that one can renormalize $\tilde{\mathcal{B}}$ by subtracting some explicit distributional counterterm $\chi_i c_r$ proportional to $\delta_{\mathbf{d}_4}$. A priori $c_r = c_r(y_*, t)$ is a function on $M \times \mathbb{R}$, but we also want to show that one can actually take c_r constant equal to b_r .

The first step in our proof is to implement in real conditions the abstract extension Theorem 13 with counterterms, also controlling the wave front of the extension. The second step is to explicitly compute the abstract counterterm c_r whose existence is given by Theorem 13 in terms of trace densities of some operators, this computation is similar to the one which is done for the quartic term $\bar{\tau}_2$ in section 5.5.3.

Let us now study the renormalization problem for $\tilde{\mathcal{B}}(y_*, y_1, y_2, y_a, s_a)$ locally in $U_i^4 \times \mathbb{R}$ for $i \in I$, since the diagonal $\mathbf{d}_4 = \{y_* = \dots = y_a, s_a = t\}$ can be covered by such sets, so that we can recover the global extension just from working with the local extensions. Fix $\chi_i : M^4 \rightarrow \mathbb{R}_{\geq 0}$ with support in U_i^4 , identically equal to 1 in a neighbourhood of $\{y_* = \dots = x_a\} \cap U_i^4$. Here and from now on, abusing notation, use the same notation to denote both elements of U_i^4 and their coordinates. As in the extension Theorem 13, we use the parabolic scaling defined by the scaling field $\rho = 2(t-s)\partial_s + \sum_{j \in \{1,2,a\}} (y_j - y_*) \cdot \partial_{y_j}$ whose semiflow $(e^{-t\rho})_{t \geq 0}$ leaves U_i^4 stable. Then we have the continuous partition of unity formula $\chi_i = \int_0^\infty e^{u\rho} w_i du$ where we set $w_i \stackrel{\text{def}}{=} -\rho \chi_i \in C^\infty(U_i^4)$ which vanishes near $\{y_* = \dots = x_a\} \cap U_i^4$. Moreover, we denote for $k \in \mathbb{N}_{\geq 1}$ by

$$\text{dens}_k(y_1, \dots, y_k) \stackrel{\text{def}}{=} \prod_{j=1}^k \sqrt{\det(g)_{y_j}}$$

the density of the Riemannian volume on U_i^k endowed with the product metric with respect to the Lebesgue measure $\prod_{j=1}^k dy_j$.

Fix a test function $\varphi \in C_c^\infty(U_i^4 \times \mathbb{R})$. We decompose the pairing of $\tilde{\mathcal{B}}$ with φ as

$$\begin{aligned}
\langle \tilde{\mathcal{B}}, \varphi \rangle &= \langle \tilde{\mathcal{B}}(1 - \chi_i), \varphi \rangle \\
&+ \int_0^\infty \int_{-\infty}^t \int_{U_i^4} \left(w_i e^{-u\rho^*} \tilde{\mathcal{B}} e^{-u\rho^*} (\Pi_z(\varphi \text{dens}_4)) \right) (y_*, y_1, y_2, y_a, s_a) e^{-11u} \prod_{j \in \{*, 1, 2, a\}} dy_j ds_a du \\
&+ \int_0^\infty \int_{-\infty}^t \int_{U_i^4} (\tilde{\mathcal{B}} e^{u\rho^*} w_i) (y_*, y_1, y_2, y_a, s_a) (\varphi \text{dens}_4)(z) \prod_{j \in \{*, 1, 2, a\}} dy_j ds_a du,
\end{aligned}$$

where we used the notation $\Pi_z(\varphi \text{dens}_4) \stackrel{\text{def}}{=} \varphi \text{dens}_4 - (\varphi \text{dens}_4)(z)$ to denote the recentering at the point $z = (y_*, y_*, y_*, y_*, t)$, and the factor e^{-11u} comes from the Jacobian of the semi-flow generated by ρ . Moreover, observe that $e^{-u\rho^*} \tilde{\mathcal{B}} = \mathcal{O}_{\mathcal{D}'}(e^{11u})$ and that $\Pi_z(\varphi \text{dens}_4)$ vanishes on $\mathbf{d}_4 \cap (U_i^4 \times \mathbb{R})$, which implies that

$$e^{-u\rho^*} (\Pi_z(\varphi \text{dens}_4)) = \mathcal{O}(e^{-u}).$$

The integral over u in the second term of the r.h.s. is therefore absolutely convergent (in fact there is a subtlety related to compactness, but it is immediate to check that the product $\psi_i e^{-u\rho^*} (\Pi_z(\varphi \text{dens}_4))$ forms a bounded family of test functions). We have thus proved that the first two terms of the r.h.s. have well-defined limits as $r \downarrow 0$. It remains to identify the counterterm as the divergent part of the third term, which we denote by $\mathfrak{D}(\cdot)$. Note that all the quantities $f = f(r)$ that we compute and which depend on the cut-off r are contained in some algebra $\mathcal{O}_{>0}$ of functions of the cut-off r which are smooth functions of $r \in (0, 1]$ and have log-polyhomogeneous expansions as $r \downarrow 0$, $f(r) = ar^{-1/2} + b \log r^{-1} + c + o(1)$ where $(a, b, c) \in \mathbb{R}^3$. We define the divergent part of any element $f \in \mathcal{O}_{>0}$ as $\mathfrak{D}(f) \stackrel{\text{def}}{=} ar^{-1/2} + b \log r^{-1}$.

With this definition in hand, the divergent part of the third term verifies

$$\begin{aligned}
\mathfrak{D} \left(\int_0^\infty \int_{-\infty}^t \int_{U_i^4} (\tilde{\mathcal{B}} e^{u\rho^*} w_i) (y_*, y_1, y_2, y_a, s_a) (\varphi \text{dens}_4)(z) \prod_{j \in \{*, 1, 2, a\}} dy_j ds_a du \right) \\
= \int_{U_i} \mathfrak{D} \left(\int_{-\infty}^t \int_{U_i^3} (\tilde{\mathcal{B}} \chi_i) (y_*, y_1, y_2, y_a, s_a) \text{dens}_3(y_*, y_*, y_*) \prod_{j \in \{1, 2, a\}} dy_j ds_a \right) \varphi(z) dv(y_*) \\
= \langle c_r \delta_{\mathbf{d}_4}, \varphi \rangle,
\end{aligned}$$

which implies by definition of $\delta_{\mathbf{d}_4}$ that the counterterm reads

$$\begin{aligned}
c_r(y_*, t) &= \mathfrak{D} \left(\int_{-\infty}^t \int_{U_i^3} (\tilde{\mathcal{B}} \chi_i) (y_*, y_1, y_2, y_a, s_a) \text{dens}_3(y_*, y_*, y_*) \prod_{j \in \{1, 2, a\}} dy_j ds_a \right) \\
(5.11) \quad &= \mathfrak{D} \left(\int_{-\infty}^t \int_{U_i^3} (\tilde{\mathcal{B}} \chi_i) (y_*, y_1, y_2, y_a, s_a) \text{dens}_3(y_1, y_2, y_a) \prod_{j \in \{1, 2, a\}} dy_j ds_a \right).
\end{aligned}$$

To go from the first to the second line of this last equality, again we used the fact that

$$\text{dens}_3(y_1, y_2, y_a) - \text{dens}_3(y_*, y_*, y_*)$$

vanishes on $\{y_* = \dots = y_a\}$, so that it does not contribute to the singular part of the above integral, which gives us the freedom to change the evaluation point of the volume elements. The reader might also wonder why we could interchange the extraction of singular parts with the integration: this is due to the fact that the integrand admits an asymptotic expansion. To conclude about the expression of $c_r(y_*, t)$, since in (5.11) we reconstructed the volume form $\prod_{j \in \{1, 2, a\}} dv(y_j) = \text{dens}_3(y_1, y_2, y_a) \prod_{j \in \{1, 2, a\}} dy_j$, it remains to plug the expression of $\tilde{\mathcal{B}}$, and we obtain

$$c_r(y_*, t) = \mathfrak{D} \left(\int_{-\infty}^t \sum_{|k-\ell| \leq 1} P_k^i(y_*, \cdot) \circ \tilde{\chi}_i \circ \left(G_r^{(2)}(t-s) \circ e^{-(t-s)P} \right) \circ \tilde{\chi}_i \circ \tilde{P}_\ell^i(\cdot, y_*) ds \right),$$

where the P_k^i, \tilde{P}_ℓ^i are the Littlewood-Paley-Stein projectors, the functions $\tilde{\chi}_i \in C_c^\infty(U_i)$ are arbitrary test functions such that $\tilde{\chi}_i = 1$ near y_* and where we used the explicit definition of $[\circ_i]$ in terms of the Littlewood-Paley-Stein projectors. This is indeed the desired expression (5.12),

the composition $\left(\int_{-\infty}^t \underline{\mathcal{L}}^{-1}(t-s) \circ G^{(1)}(t-s) ds\right)$ belongs to Ψ_P^{-3} , hence it belongs to $\Psi^{-4}(M)$ uniformly in t and is trace class on M . The series of pseudodifferential operators

$$\begin{aligned} & \sum_{|k-\ell| \leq 1} \int_{-\infty}^t \left(P_k^i \circ e^{-(t-s)P} \circ G^{(1)}(t-s) \circ \tilde{P}_\ell^i \right) (y_*, y_*) ds = \\ & \sum_{|k-\ell| \leq 1} \int_{-\infty}^t \left\{ \left(P_k^i \circ \tilde{P}_\ell^i \circ e^{-(t-s)P} \circ G^{(1)}(t-s) \right) + \left(P_k^i \circ [e^{-(t-s)P} \circ G^{(1)}(t-s), \tilde{P}_\ell^i] \right) \right\} (y_*, y_*) \end{aligned}$$

will also converge in $\Psi^{-4}(M)$ since the series $\sum_{|k-\ell| \leq 1} P_k^i \circ \tilde{P}_\ell^i$ converges in $\Psi_{0,1}^0(M)$ and the commutator $[e^{-(t-s)P} \circ G^{(1)}(t-s), \tilde{P}_\ell^i]$ is bounded in $\Psi^{-5}(M)$ uniformly in (ℓ, i) since the sequence $(\tilde{P}_\ell^i)_{\ell,i}$ is bounded in $\Psi^0(M)$. Therefore the second series involving the commutator term converges in $\Psi^{-5}(M)$. We use for that purpose a commutator identity that says that for every pseudodifferential operator $A \in \Psi^m(M)$, the series


$$\sum_{|k-\ell| \leq 1} \left(P_k^i A \tilde{P}_\ell^i - A P_k^i \tilde{P}_\ell^i \right)$$

converges as a pseudodifferential operator of order $m-1$ that we prove in our companion work. Finally this implies that the term

$$\int [\odot_i](x_*, x_1, x_2) \underline{\mathcal{L}}^{-1}((t, x_2), (s, x_a)) G_r^{(1)}((t, x_1), (s, x_a)) ds dx_1 dx_2 dx_a$$

is a well-defined smooth function.

5.5.2 Elementary bounds for τ_2 . Again, \mathcal{A}_{24} and \mathcal{A}_{22} need the special treatment from Section 5.4 since they respectively have the subamplitudes $\mathcal{C}^{(2)}$ and $\mathcal{C}^{(1)}$: Lemma 37 states that these subamplitudes are indeed well-defined for every $\gamma \in \mathbb{R}$.


To complete the argument, we check for the reader's convenience the weak homogeneities of the amplitudes of all the other subgraphs. The amplitude \mathcal{A}_{24} of $\mathcal{G}_{24} =$  is

$$\begin{aligned} & [\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(2)}(0, y_1, z_1) \\ & \quad \times \underline{\mathcal{L}}^{-1}(t-s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t-s_b, z_2, x_b) G_r^{(2)}(s_a-s_b, x_a, x_b). \end{aligned}$$

Here again this graph has four closed, irreducible, connected subgraphs and we calculate the weak homogeneity of \mathcal{A}_{24} itself by summing the weak homogeneity of all analytical objects, the kernels $G_r^{(2)}, [\odot_i], \underline{\mathcal{L}}^{-1}, \mathcal{Q}^\gamma$, appearing in the amplitude

$$\begin{aligned} \sum \text{weak homogeneities} &= 2(-2) + 2(-6) + 2(-3) - 6 - 2\gamma = -28 - 2\gamma \\ &> -\text{codim}_w(\{s_a = s_b = t, y_* = \dots = x_b\}) = -4 - 24 = -28, \end{aligned}$$

hence $\gamma < 0$. Repeating this verification for all subgraphs yields the result that $\mathcal{G}_{24} < \infty$ for all $\gamma < 0$.

The verification for the graph $\mathcal{G}_{22} =$  of amplitude

$$\begin{aligned} \mathcal{A}_{22} &= [\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(1)}(t-s_a, y_1, x_a) G_r^{(1)}(t-s_b, z_1, x_b) \\ & \quad \times \underline{\mathcal{L}}^{-1}(t-s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t-s_b, z_2, x_b) G_r^{(1)}(s_a-s_b, x_a, x_b) \end{aligned}$$

is similar, for instance the subamplitude

$$[\odot_i](y_*, y_1, y_2) G_r^{(1)}(t-s_a, y_1, x_a)$$

is weakly homogeneous of degree $-6-1 = -7 > \text{codim}_w(\{s_a = t, y_* = y_1 = y_2 = x_a\}) = -2-9 = -11$ and yields the same range of regularity exponent. In a similar way as what we did for τ_1 , the locally covariant renormalized τ_2 will differ from the Wick renormalized τ_2 by homogeneous terms in Wiener chaoses of order 2 and 0 which have the form $P_1 \overset{\circ}{\mathcal{Y}} + P_2 \mathcal{V} + f_3$ where P_1, P_2 are smoothing operators and f_3 is a smooth function. We used the fact that both $\overset{\circ}{\mathcal{Y}}$ and \mathcal{V} differ from their Wick

renormalized version by a smooth function and also that for any smooth function $f \in C^\infty(M)$, the multiplication operator by the localized resonant product $u \in \mathcal{D}'(M) \mapsto f \odot_i u \in C^\infty(M)$ is smoothing.

5.5.3 The divergent part of τ_{20} . Recall that we denote by $[\odot_i]$ the kernel of \odot_i . An immediate calculation yields for $z = (t, y_*)$

$$(5.12) \quad \begin{aligned} \tau_{20}(z) &= \mathbb{E}[\tau_2(z)] = 2 \int_{-\infty}^t \int_{M^3} [\odot_i](y_*, y_1, y_2) \mathcal{L}^{-1}(t - s_a, y_2, x_a) G_r^{(2)}(t - s_a, y_1, x_a) dv(y_1, y_2, x_a) ds_a \\ &= 2 \sum_{|k-\ell| \leq 1} \int_{-\infty}^t P_k^i(y_*, \cdot) \circ G_r^{(2)}(s - t) \circ e^{-(t-s)P} \circ \tilde{P}_\ell^i(\cdot, y_*) ds, \end{aligned}$$

with the shorthand notation $dv(y_1, y_2, x_a) \stackrel{\text{def}}{=} dv(y_1)dv(y_2)dv(x_a)$ for the Riemannian volume form on M^3 , with dv the corresponding volume form on M .

§1. Preliminary analysis. Recall that

$$G_r^{(2)}(t - s, x, y) = \left(\left\{ e^{-(|t-s|+2r)P} P^{-1} \right\} (x, y) \right)^2.$$

We reformulate the integral

$$(5.13) \quad \int_{-\infty}^t G_r^{(2)}(t - s) \circ e^{-(t-s)P} ds$$

as the composition of two operators in the parabolic calculus $(\Psi_P^\alpha)_{\alpha \in \mathbb{R}}$ defined in our companion work [10]. This calculus extends the heat calculus $(\Psi_H^\alpha)_{\alpha \in \mathbb{R}}$ as defined in Grieser's note [41] in order to include the kernels $G_r^{(p)}$, $p \in \{1, 2, 3\}$ needed to define φ_3^4 amplitudes. Since $G_r^{(2)} \in \Psi_P^{-\frac{3}{2}}$ uniformly in $r > 0$ and $e^{-tP} \in \Psi_H^{-1}$, we prove in [10] that the composition $\int_{-\infty}^t G_r^{(2)}(s, t) \circ e^{-(t-s)P} ds$ defines an element in $\Psi_P^{-\frac{5}{2}+\gamma}$ for any $\gamma > 0$, uniformly in $r > 0$. Hence it is a pseudodifferential operator depending continuously on t of order $-3 + 2\gamma$, by the comparison Theorem viewing parabolic operators as parameter dependent pseudodifferential operators in [10]. It follows that the t -indexed family of operators (5.13) is bounded in $\Psi^{-3+2\gamma}(M)$ and fails to be trace class when the regularization parameter r tends to 0. Our goal in the sequel of this section is to extract the singular part of this operator.

First we need to disentangle this operator in (5.12) from the Littlewood-Paley-Stein projectors P_k^i, \tilde{P}_ℓ^i . We use for that purpose a commutator identity that says that for every pseudodifferential operator $A \in \Psi^m(M)$, the series

$$\sum_{|k-\ell| \leq 1} \left(P_k^i A \tilde{P}_\ell^i - A P_k^i \tilde{P}_\ell^i \right)$$

converges as a pseudodifferential operator of order $m - 1$. The operator

$$(5.14) \quad \sum_{|k-\ell| \leq 1} \left(P_k^i G_r^{(2)}(t - s) e^{-(t-s)P} \tilde{P}_\ell^i - G_r^{(2)}(t - s) e^{-(t-s)P} P_k^i \tilde{P}_\ell^i \right)$$

is in particular in $\Psi^{-4+2\gamma}(M)$ uniformly in $r \in [0, 1]$, for any $0 < \gamma < 1/2$, so it is trace class.

Denote by δ_{y_*} is the unique distribution depending on the Riemannian volume form $dv(y)$ such that

$$\int_M \delta_{y_*}(z) f(z) dv(z) = f(y_*),$$

for all bounded measurable functions f . With this notation one has

$$\tau_{20}(z) = 2 \int_{-\infty}^t \sum_{|k-\ell| \leq 1} \left\langle P_k^i(\delta_{y_*}), G_r^{(2)}(t - s) e^{-(t-s)P} \tilde{P}_\ell^i(\delta_{y_*}) \right\rangle ds$$

and we see from the preceding regularity result for the commutator (5.14) that the singular part $\mathfrak{D}\tau_{20}(z)$ of $\tau_{20}(z)$ coincides with the singular part of

$$\left\langle \left(\left\{ \int_{-\infty}^t G_r^{(2)}(t - s) e^{-(t-s)P} ds \right\} \delta_{y_*} \right) \odot_i \delta_{y_*}, 1 \right\rangle.$$

Given that the localized paraproduct of any two distributions is always well-defined, and that (cf Appendix A)

$$\chi_i(uv) = u \odot_i v + u \prec_i v + u \succ_i v,$$

this localized singular part coincides with the singular part of

$$\chi_i(y_*) \left\{ \int_{-\infty}^t G_r^{(2)}(t-s) e^{-(t-s)P} ds \right\} (y_*, y_*).$$

§2. *Explicit computation of the divergent part.* We proceed in two steps by localising first in space and time and then by using the heat kernel asymptotics.

§2.1. *Space and time localisation.* Fix $z = (t, y_*)$. We start by localising in space near y_* . Let χ stand for a smooth indicator function of a neighbourhood U_{y_*} of y_* in M – without loss of generality the domain of a chart. Since the quantity

$$\int_{-\infty}^t \int_M G_r^{(2)}(t-s)(y_*, y) (1 - \chi(y e^{-(t-s)P}(y, y_*))) dv(y) ds$$

has a well-defined limit when $r \downarrow 0$, we concentrate on

$$\begin{aligned} (\star) &\stackrel{\text{def}}{=} \int_{-\infty}^t \int_{U_{y_*}} G_r^{(2)}(t-s)(y_*, y) e^{-(t-s)P}(y, y_*) \chi(y) dv(y) ds \\ &= \int_0^\infty \left(\int_{[a+2r, \infty)^2} \int_{U_{y_*}} e^{-s_1 P}(y_*, y) e^{-s_2 P}(y_*, y) e^{-aP}(y_*, y) \chi(y) dv(y) ds_1 ds_2 \right) da. \end{aligned}$$

§2.2. *Use the heat kernel asymptotics.* We have near y_*

$$e^{-sP}(y_*, y) = \frac{1}{(4\pi s)^{\frac{3}{2}}} e^{-\frac{|y_* - y|_g^2(y_*)}{4s}} e^{-s} + R(s, y_*, \frac{y_* - y}{\sqrt{s}}) =: K^0(s, y_*, y) + R\left(s, y_*, \frac{y_* - y}{\sqrt{s}}\right)$$

for a remainder term $R \in \Psi_H^{-2}(M)$ in the heat calculus as defined in [41], that is $R(s, y_*, \frac{y_* - y}{\sqrt{s}})$ has the same estimates as s times the heat kernel itself. It follows from that fact that replacing any of the heat kernels $e^{-s_1 P}, e^{-s_2 P}, e^{-aP}$ by a remainder term R gives a contribution to (\star) that remains uniformly bounded for $r \in [0, 1]$. We can therefore keep in our computations only the leading term of the heat expansion. Integrating first with respect to y the stationary phase in U_{y_*} gives the asymptotics

$$\begin{aligned} &\int_{U_{y_*}} K^0(s_1, y_*, y) K^0(s_2, y_*, y) \chi(y) K^0(a, y_*, y) dv(y) \\ &= \frac{e^{-(s_1 + s_2 + a)} (4\pi)^{-\frac{9}{2}}}{(s_1 s_2 a)^{\frac{3}{2}}} \int_{U_{y_*}} e^{-\frac{|y_* - y|_g^2(y_*)}{4}} (s_1^{-1} + s_2^{-1} + a^{-1}) \chi(y) \sqrt{\det(g)_y} dy \\ &= \frac{e^{-(s_1 + s_2 + a)} \chi(y_*) \sqrt{\det(g)_{y_*}} (4\pi)^{-3}}{(s_1^{-1} + s_2^{-1} + a^{-1})^{\frac{3}{2}} (s_1 s_2 a)^{\frac{3}{2}} \sqrt{\det \text{Hess}(|\cdot - y_*|_g^2(y_*))}(y_*)}} \\ &\quad + \mathcal{O}\left((s_1 s_2 a)^{-\frac{3}{2}} (s_1^{-1} + s_2^{-1} + a^{-1})^{-\frac{5}{2}}\right) \\ &= \frac{e^{-(s_1 + s_2 + a)} (4\pi)^{-3}}{(s_1^{-1} + s_2^{-1} + a^{-1})^{\frac{3}{2}} (s_1 s_2 a)^{\frac{3}{2}}} + \mathcal{O}\left((s_1 s_2 a)^{-\frac{3}{2}} (s_1^{-1} + s_2^{-1} + a^{-1})^{-\frac{5}{2}}\right), \end{aligned}$$

with an error term $\mathcal{O}((s_1 s_2 a)^{-\frac{3}{2}} (s_1^{-1} + s_2^{-1} + a^{-1})^{-\frac{5}{2}})$ bounded uniformly in the y_* variable. Note that to go from the second to the last line we used $\chi(y_*) = 1$, along with the fact that the determinant of the Hessian equals the determinant of g evaluated at y_* .

Here, note that the divergence of the integral in a only takes place when $a \downarrow 0$, so that in order to seek the divergent part, one can restrict the integration over a to $[0, 1]$. Moreover, the exponential decay $e^{-(s_1 + s_2 + a)}$ will not help, so that we can discard it when computing the divergent part. In the end, we are therefore left with computing the singular part of

$$\begin{aligned}
\frac{1}{(4\pi)^3} \int_0^1 \int_{[a+2r, \infty)^2} \frac{ds_1 ds_2 da}{(s_2 a + s_1 a + s_1 s_2)^{\frac{3}{2}}} &= \frac{1}{(4\pi)^3} \int_0^1 \frac{da}{a+2r} \int_{[1, \infty)^2} \frac{d\alpha d\beta}{(\alpha + \beta + \alpha\beta - \frac{2r(\alpha+\beta)}{a+2r})^{\frac{3}{2}}} \\
&= -\frac{1}{(4\pi)^3} \int_{[1, \infty)^2} \frac{d\alpha d\beta}{(\alpha + \beta + \alpha\beta)^{\frac{3}{2}}} \log r + \mathcal{O}(1) \\
&= -\frac{1}{3} \times \frac{\log r}{32\pi^2} + \mathcal{O}(1).
\end{aligned}$$

On the first line, we did the two changes of variables $s_1 = \alpha(a+2r)$ and $s_2 = \beta(a+2r)$, while on the second line the integral over α, β can be computed explicitly. This finally tells us that the diverging part of τ_{20} is exactly given by $\chi_i \frac{\log r}{3}$ for every $i \in I$. To conclude, note that what is left of τ_{20} after the subtraction of the divergent part defines a smooth function of y_* .

5.5.4 Bounds on $\bar{\tau}_3$. We first briefly discuss the counterterm for $\bar{\tau}_3$ and we will do the stochastic estimates in a second step. Note that for any test function φ we have from an integration by parts

$$\int_M \chi_i |\nabla \Upsilon_r^\varphi|^2 \varphi = - \int_M (\Delta \Upsilon_r^\varphi) \Upsilon_r \chi_i \varphi - \int_M \langle \nabla \Upsilon_r^\varphi, \nabla(\chi_i \varphi) \rangle \Upsilon_r^\varphi.$$

The singular part of $|\nabla \Upsilon_r^\varphi|^2$ as a random distribution is thus the same as the singular part of $(\Delta \Upsilon_r^\varphi) \Upsilon_r^\varphi$. The latter is also the singular part of

$$\begin{aligned}
\mathbb{E} \left[(P \Upsilon_r^\varphi) \Upsilon_r^\varphi \right] &= \int_{(-\infty, t]^2} \text{tr}_{L^2} \left(P e^{-(t-s_1)P} \circ G_r^{(2)}(s_1 - s_2) \circ e^{-(t-s_2)P} \right) ds_1 ds_2 \\
&= \int_{(-\infty, t]^2} \text{tr}_{L^2} \left(P e^{-(t-s_1)P} \circ G_r^{(2)}(s_1 - s_2) \circ e^{-(t-s_2)P} \right) ds_1 ds_2 \\
&= \int_{(-\infty, t]^2} \text{tr}_{L^2} \left(G_r^{(2)}(s_1 - s_2) \circ P e^{-(2t-s_1-s_2)P} \right) ds_1 ds_2
\end{aligned}$$

so changing variables for $s'_1 = t - s_1$ and $s'_2 = s_1 - s_2$ gives

$$\begin{aligned}
\int_{(-\infty, t]^2} \text{tr}_{L^2} \left(G_r^{(2)}(s'_2) \circ P e^{-(2s'_1+s'_2)P} \right) ds'_1 ds'_2 &= \int_{(-\infty, t]^2} \text{tr}_{L^2} \left(G_r^{(2)}(s'_2) \circ \frac{d}{ds'_1} e^{-(2s'_1+s'_2)P} \right) ds'_1 ds'_2 \\
&= \int_{-\infty}^t \text{tr}_{L^2} \left(e^{-s'_2 P} \circ G_r^{(2)}(s'_2) \right) ds'_2,
\end{aligned}$$

where we use time permutation symmetry of $G_r^{(2)}$ and cyclicity of the L^2 -trace. This quantity is equal to the divergent part of τ_{20} . We next discuss the regularity of $\bar{\tau}_3$. First, we need to isolate the resonant part in the scalar product. Since we work in the manifold setting, note that we need to define carefully the resonant scalar product of two vector fields in $C^\infty(TM)$. For $s_1, s_2 \in C^\infty(TM)^2$, using the notations and conventions of the Appendix A, we define $\langle s_1 \odot_i s_2 \rangle_{TM}$ as:

$$\langle s_1 \odot_i s_2 \rangle_{TM} \stackrel{\text{def}}{=} \kappa_i^* ((\kappa_{i*} g)^{\mu\nu} \kappa_{i*} \psi_i (\kappa_{i*}(\chi_i s_1)_\mu \odot \kappa_{i*}(\tilde{\chi}_i s_2)_\nu))$$

where $\kappa_i, \psi_i, \chi_i, \tilde{\chi}_i$ come from our definition of resonant product, $(\kappa_{i*} g)$ is the metric g induced by the charts $\kappa_i : U_i \mapsto \kappa_i(U_i) \subset \mathbb{R}^d$. Similarly we have

$$\langle s_1 \prec_i s_2 \rangle_{TM} \stackrel{\text{def}}{=} \kappa_i^* ((\kappa_{i*} g)^{\mu\nu} \kappa_{i*} \psi_i (\kappa_{i*}(\chi_i s_1)_\mu \prec \kappa_{i*}(\tilde{\chi}_i s_2)_\nu))$$

and we recover the usual decomposition

$$\langle s_1, s_2 \rangle_{TM} = \sum_i \langle s_1 \odot_i s_2 \rangle_{TM} + \langle s_1 \prec_i s_2 \rangle_{TM} + \langle s_1 \succ_i s_2 \rangle_{TM}$$

for the scalar product on sections of TM . We need to prove that such resonant scalar product satisfies some approximate integration by parts identity and we are careful since the Laplacian is no longer translation invariant. For every $Y \in C^\infty(M)$ a calculation relying on the definitions of both resonant product and resonant scalar product:

$$\langle \nabla Y \odot_i \nabla Y \rangle_{TM} = \chi_1(Y \odot_1 Y) + (PY) \odot_2 Y + Y \odot_i \Delta Y$$

where χ_1 is a smooth function, P is a differential operator of order 1 on M with smooth coefficients, \odot_1, \odot_2 are localized resonant type products which might differ from the original \odot_i in the choice of

smooth cut-off functions involved in the definition but have the exact same analytical properties from Proposition 43 and the last term involves the localized resonant product of Y with ΔY . From the point of view of regularities, for $Y = \mathring{Y}_r \in \mathcal{C}^{1-2\varepsilon}([0, T] \times M)$, $\chi_1(\mathring{Y}_r \odot_1 \mathring{Y}_r) \in C_T \mathcal{C}^{2-4\varepsilon}(M)$ and $(P\mathring{Y}_r) \odot_2 \mathring{Y}_r \in C_T \mathcal{C}^{1-4\varepsilon}(M)$ and finally we are reduced to the study of $\mathring{Y}_r \odot_i \Delta \mathring{Y}_r$. This is now really very similar to what we did for the graph τ_2 except there is an extra propagator $\Delta \underline{\mathcal{L}}^{-1}$ in all the Feynman amplitudes. However microlocal estimates from our companion work [10] actually show that

$$(5.15) \quad \Delta \underline{\mathcal{L}}^{-1} \in \mathcal{S}_{N^*(\{t=s\} \times \mathbf{d}_2 \subset \mathbb{R}^2 \times M^2)}^{-5}(\mathbb{R}^2 \times M^2).$$

Now we repeat the stochastic estimates on $\mathbb{R}^4 \times M^{10}$ taking this new kernel and its weak homogeneity into account. For instance, the amplitude controlling the homogeneous chaos of orders 2 and 4 in the chaos decomposition of $\mathring{Y}_r \odot_i \Delta \mathring{Y}_r$ now read

$$(5.16) \quad \mathcal{A}_{34} = [\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) \\ G_r^{(2)}(s_a - s_b, x_a, x_b) \Delta \underline{\mathcal{L}}^{-1}(t - s_c, y_1, x_c) \Delta \underline{\mathcal{L}}^{-1}(t - s_d, z_1, x_d) G_r^{(2)}(s_c - s_d, x_c, x_d),$$

$$(5.17) \quad \mathcal{A}_{32} = [\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) \\ G_r^{(1)}(s_a - s_b, x_a, x_b) \Delta \underline{\mathcal{L}}^{-1}(t - s_c, y_1, x_c) \Delta \underline{\mathcal{L}}^{-1}(t - s_d, z_1, x_d) G_r^{(1)}(s_c - s_d, x_c, x_d) \\ G_r^{(1)}(s_a - s_c, x_a, x_c) G_r^{(1)}(s_b - s_d, x_b, x_d).$$


These two amplitudes respectively have the subamplitudes $\tilde{\mathcal{C}}^{(2)}$ and $\tilde{\mathcal{C}}^{(1)}$, which are dealt with in Section 5.4 (see Lemma 37). Here again, it remains to calculate the weak homogeneities of the amplitudes and all the other subamplitudes, by summing the weak homogeneity of all analytical objects, the kernels $G_r^{(2)}$, $[\odot_i]$, $\underline{\mathcal{L}}^{-1}$, $\Delta \underline{\mathcal{L}}^{-1}$, \mathcal{Q}_x^γ . For \mathcal{A}_{34} , this yields

$$\begin{aligned} \sum \text{weak homogeneities} &= 2(-2) + 2(-6) + 2(-3) - 6 - 2\gamma + 2(-5) = -38 - 2\gamma \\ &> -\text{codim}_w(\{s_a = s_b = s_c = s_d = t, y_* = \dots = x_d = x\}) \\ &= -8 - 30 = -38, \end{aligned}$$


hence we obtain $\gamma < 0$. Repeating this verification for all subgraphs and for all amplitudes controlling the term homogeneous of order 2 in the chaos decomposition yields the result that $\mathring{Y}_r \odot_i \Delta \mathring{Y}_r \in \mathcal{C}^{-4\varepsilon}([0, T] \times M)$ almost surely.

The locally covariant renormalisation for $\bar{\tau}_3$ differs from the Wick one by $f_1 \cdot \nabla^\circ \mathring{Y} + f_2$ where f_1 is a smooth vector and f_2 a smooth function which has higher regularity than $|\nabla^\circ \mathring{Y}|^2$.

5.5.5 The quintic term. The first two graphs respectively have the subgraphs with amplitude $\mathcal{C}^{(2)}$ and $\mathcal{C}^{(1)}$, which are dealt with in Section 5.4 (see Lemma 37). To complete the argument, one extracts all the other closed, connected irreducible graphs and finds the range of parameter γ so that they satisfy the criterion of Theorem 27. For the reader's convenience, recall that the

amplitude \mathcal{A}_{45} of $\mathcal{G}_{45} = \text{

$$[\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(2)}(0, y_1, z_1) \\ \times \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) G_r^{(3)}(s_a - s_b, x_a, x_b),$$$

and the amplitude \mathcal{A}_{43} of $\mathcal{G}_{43} = \text{

$$\mathcal{A}_{43} = [\odot_i](y_*, y_1, y_2) [\odot_i](z_*, z_1, z_2) \mathcal{Q}_x^\gamma(y_*, z_*) G_r^{(1)}(0, y_1, z_1) G_r^{(1)}(t - s_a, y_1, x_a) G_r^{(1)}(t - s_b, z_1, x_b) \\ \times \underline{\mathcal{L}}^{-1}(t - s_a, y_2, x_a) \underline{\mathcal{L}}^{-1}(t - s_b, z_2, x_b) G_r^{(2)}(s_a - s_b, x_a, x_b).$$$

The reader can check that the scaling degrees of the subgraphs yield the range $\gamma < -\frac{1}{2}$.

It remains to handle the last contribution, which verifies

$$\mathbb{E}[\mathcal{Q}_x^\gamma((\tau_{41}(t, \cdot) - \chi_i b_r \mathring{I}_r), (\tau_{41}(t, \cdot) - \chi_i b_r \mathring{I}_r))] = \langle \mathcal{A}_{41}, 1 \rangle,$$

where \mathcal{A}_{41} is given by (5.9). The amplitude \mathcal{A}_{41} is indeed well-defined as a distribution of $M^8 \times \mathbb{R}^2$ by Theorem 27 for $\gamma < -1/2$, taking the renormalized subamplitude \mathcal{B} (that is constructed in Lemma 38) into account in the induction over all subgraphs of \mathcal{A}_{41} .

We conclude our discussion by pointing out the difference between the locally covariant and Wick renormalisations. The locally covariant renormalisation of $\overset{\circ}{\Psi} \odot_i \mathcal{V}$ differs from the Wick renormalisation by $(\underline{\mathcal{L}}^{-1}(f^\circ)) \odot_i \mathcal{V} + P(\overset{\circ}{\Psi})$ where f is a smooth function and P a smoothing operator and therefore $(\underline{\mathcal{L}}^{-1}(f^\circ)) \odot_i \mathcal{V} \in \mathcal{C}^{\frac{1}{2}-\varepsilon}$, $\forall \varepsilon > 0$ which is absorbed in the $\mathcal{C}^{-\frac{1}{2}-0}$ regularity of $\overset{\circ}{\Psi} \odot_i \mathcal{V}$.

The results of this section justify the convergence in $L^2(\Omega)$, hence in any $L^p(\Omega)$ with $p < \infty$, of $\widehat{\xi}_r$ to a limit random variable $\widehat{\xi}$ in its natural space. The convergence *in probability* of $v_r \in (\alpha_0, 1 + \varepsilon')$ to a limit v in that space follows as a consequence of the pathwise continuity of v_r with respect to $\widehat{\xi}_r$ obtained from the fixed point construction of v_r . Formula (2.3) relating v_r to u_r shows the convergence *in probability* of $u_r \in C_T C^{-1/2-\varepsilon}(M)$ to a limit u in that space.

6 – Invariant measure

We prove in Section 6.1 that the dynamics generated by equation (1.3) is Markovian and that its semigroup has the Feller property. The existence of an invariant measure is obtained from a compactness argument building on the L^p coming down from infinity property of Theorem 5. We prove in Section 6.2 that the invariant probability measure is non-Gaussian.

6.1 A Markovian dynamics

Denote by \mathcal{F}_t the usual augmentation of the σ -algebra generated by the random variables $\xi(f)$, for functions $f \in L^2(\mathbb{R} \times M)$ that are null on $[t, \infty) \times M$. Since ξ_r is white in time the dynamics

$$(\partial_t + P)u_r = -u_r^3 + 3(a_r - b_r)u_r + \sqrt{2}\xi_r$$

generates an $(\mathcal{F}_t)_{t \geq 0}$ -Markov process. For looking at the restriction to a finite time interval $[0, T]$ of this process it is convenient to extend functions on $[0, T]$ into functions on $[0, +\infty)$ that are constant on $[T, +\infty)$. For $t \in \mathbb{R}$ and any $(s, x) \in \mathbb{R} \times M$ set

$$\tau_t(s, x) \stackrel{\text{def}}{=} (s - t, x).$$

Denote by $\theta_s : \Omega \rightarrow \Omega$, $s \geq 0$ a family of measurable shifts on (Ω, \mathcal{F}) such that one has

$$(\xi \circ \theta_s, f) = (\xi, f \circ \tau_s)$$

for all s and all L^2 test functions f . The Markov property for

$$u_r : \Omega \times [0, T] \times C^{-1/2-\varepsilon}(M) \rightarrow C^{-1/2-\varepsilon}(M)$$

writes

$$(6.1) \quad \mathbb{E}[F(u_r(s + \cdot, \phi)) \mathbf{1}_E] = \mathbb{E}[F(u_r \circ \theta_s(\cdot, u_r(s, \phi))) \mathbf{1}_E],$$

for any bounded measurable cylindrical functional F on $C((0, T], C^{-1/2-\varepsilon}(M))$ and all events $E \in \mathcal{F}_s$, with $0 \leq s \leq T$ arbitrary. We need the following quantitative stability result to pass to the zero r limit in (6.1).

Lemma 39 – *Fix some positive times $t_1 < \dots < t_k$. There exists two positive constants γ, γ' such that the restriction of the functions*

$$\phi \in C^{-1/2-\varepsilon}(M) \mapsto u_r(t_i, \phi) \in C^{-1/2-\varepsilon}(M), \quad (1 \leq i \leq k)$$

to any centered ball of $C^{-1/2-\varepsilon}(M)$ with radius $R > 0$ is Lipschitz continuous, with Lipschitz constant bounded above by an explicit function of R and $\widehat{\xi}_r$.

Proof – This result is obtained from the exact same statement for the functions $v_r(t_i, \cdot)$. The relation

$$u_r(t, \phi_1) - u_r(t, \phi_2) = e^{-3\overset{\circ}{\Psi}_r(t)}(v_r(t, \phi_1') - v_r(t, \phi_2')),$$

with

$$\phi_i = \mathfrak{I}_r(0) - \mathfrak{Y}_r^\circ(0) + e^{-3\mathfrak{Y}_r^\circ(0)}(\phi'_i + v_{\text{ref}}(0))$$

allows to transport the locally Lipschitz character of v_r to u_r . It suffices to prove the statement with $k = 1$ and $t_1 = 1$; we prove in that case that $\phi \in C^{-1/2-\varepsilon}(M) \mapsto v_r(1, \phi) \in C^{-1/2-\varepsilon}(M)$ is locally Lipschitz. Define

$$(6.2) \quad F(\phi, v) \stackrel{\text{def}}{=} e^{-tP}(\phi) + \mathcal{L}^{-1}\left(-6\nabla\mathfrak{Y}_r^\circ \cdot \nabla v - e^{-6\mathfrak{Y}_r^\circ}v^3 + Z_{2,r}v^2 + Z_{1,r}v + Z_{0,r}\right).$$

Let $K > 0$ be a uniform constant satisfying

$$(6.3) \quad \|e^{-tP}\phi\|_{C^{-1/2-\varepsilon}} \leq K\|\phi\|_{C^{-1/2-\varepsilon}} \quad \text{and} \quad \|e^{-tP}\phi\|_{(|\alpha_0, 1+\varepsilon'|)} \leq K\|\phi\|_{C^{-1/2-\varepsilon}}.$$

Take the ball B_R in $C^{-1/2-\varepsilon}(M)$. It follows from the proof of Proposition 3 that for any $\phi \in B_R$, there exists $T = T(\widehat{\xi}_r|_{[0,2]}, R)$ and a constant $C(T) < 1/2$ only depending on T such that

$$\|F(\phi, v_1) - F(\phi, v_2)\|_{C_T C^{-1/2-\varepsilon}} \leq C(T)(\|v_1 - v_2\|_{C_T C^{-1/2-\varepsilon}} + \|v_1 - v_2\|_{(|\alpha_0, 1+\varepsilon'|)})$$

and

$$\|F(\phi, v_1) - F(\phi, v_2)\|_{(|\alpha_0, 1+\varepsilon'|)} \leq C(T)\|v_1 - v_2\|_{(|\alpha_0, 1+\varepsilon'|)}.$$

Now by the same argument in the proof of Proposition 3, we infer that for $\phi_1, \phi_2 \in B_R$,

$$\begin{aligned} & \|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{C_T C^{-1/2-\varepsilon}} \\ & \leq K\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}} + C(T)(\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{C_T C^{-1/2-\varepsilon}} + \|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{(|\alpha_0, 1+\varepsilon'|)}) \end{aligned}$$

and

$$\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{(|\alpha_0, 1+\varepsilon'|)} \leq K\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}} + C(T)\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{(|\alpha_0, 1+\varepsilon'|)}.$$

This implies that

$$\begin{aligned} & \|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{C_T C^{-1/2-\varepsilon}} + \|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{(|\alpha_0, 1+\varepsilon'|)} \\ & \leq 2K\|\phi_1 - \phi_2\|_{C_T C^{-1/2-\varepsilon}} + C(T)\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{C_T C^{-1/2-\varepsilon}} \\ & \quad + 2C(T)\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{(|\alpha_0, 1+\varepsilon'|)}, \end{aligned}$$

hence

$$\|v_r(\cdot, \phi_1) - v_r(\cdot, \phi_2)\|_{C_T C^{-1/2-\varepsilon}} \leq \frac{2K}{1-C}\|\phi_1 - \phi_2\|_{C^{-1/2-\varepsilon}}.$$

By the L^p a priori estimate of Theorem 5, if $T < 1$ then

$$\|v_r(t, \phi)\|_{C^{-1/2-\varepsilon}} \leq R' \stackrel{\text{def}}{=} \frac{C(\widehat{\xi}_r|_{[0,2]})}{T^{1/2}}, \quad (T/2 \leq t \leq 1).$$

Then as above we can get a short time $T' = T'(\widehat{\xi}_r|_{[0,2]}, R')$ such that the map $F(\cdot, \cdot)$ is contracting with a constant $C(T') < 1/2$ for any initial condition in $B_{R'} \subset C^{-1/2-\varepsilon}(M)$. Since $v_r(t) \in B_{R'}$ for all $t \in [T/2, 1]$ we can divide the interval $[T/2, 1]$ into subintervals $[t_j, t_j + T']$ and repeat our process above to get

$$\|v_r(t, \phi_1) - v_r(t, \phi_2)\|_{C^{-1/2-\varepsilon}} \leq \frac{2K}{1-C(T')} \|v_r(t_j, \phi_1) - v_r(t_j, \phi_2)\|_{C^{-1/2-\varepsilon}},$$

on each small interval $[t_j, t_j + T']$. Combining all yields that $\phi \mapsto v_r(1, \phi)$ is locally Lipschitz. \triangleright

Proposition 40 – *The dynamics of u is Markovian and its associated semigroup $(\mathcal{P}_t)_{t \geq 0}$ on $C^{-1/2-\varepsilon}(M)$ has the Feller property.*

Proof – Given any $\eta > 0$ it follows from Lemma 39 there is an $R(\eta) > 0$ such that outside an event of probability η the random variables $(u_r \circ \theta_s(\cdot, \cdot))_{0 < r \leq 1}$ are (r -uniformly) uniformly continuous in their second argument on the centered ball of $C^{-1/2-\varepsilon}(M)$ of radius $R(\eta)$ and $u_r(s, \phi)$ is converging to $u(s, \phi)$ with $|u(s, \phi)|_{C^{-1/2-\varepsilon}} \leq R(\eta)$. The process $u_r \circ \theta_s(\cdot, u_r(s, \phi))$ is thus converging in probability to $u \circ \theta_s(\cdot, u(s, \phi))$, so one can get the Markov property of the limit process u by passing to the zero r limit in (6.1) along a subsequence r_k where the convergence of u_{r_k} is almost sure, using dominated convergence.

The Feller property of the semigroup $(\mathcal{P}_t)_{t \geq 0}$, that is the fact that it sends the space of continuous functions on $C^{-1/2-\varepsilon}(M)$ into itself, is a direct consequence of the pathwise continuous dependence

of the solution u to (1.3) with respect to the initial condition ϕ and dominated convergence in the expression $(\mathcal{P}_t f)(\phi) = \mathbb{E}[f(u(t, \phi))]$. \triangleright

Proposition 41 – *The semigroup $(\mathcal{P}_t)_{t \geq 0}$ has an invariant probability measure.*

Proof – Recall we turned equation (1.5) on u_r into equation (2.4) on v_r , with abstract form (2.15). Coming back to

$$(6.4) \quad u = \mathfrak{I} - \mathfrak{Y}^\circ + e^{-3\mathfrak{Y}^\circ}(v + v_{\text{ref}}),$$

seen as an element of $C_T C^{-1/2-\varepsilon}(M)$, one can write for any fixed time

$$\begin{aligned} & \{ \|u(t)\|_{C^{-1/2-\varepsilon}} > 3m \} \\ & \subset \left\{ \|\mathfrak{I}(t)\|_{C^{-1/2-\varepsilon}} > m \right\} \cup \left\{ \|\mathfrak{Y}^\circ(t)\|_{C^{-1/2-\varepsilon}} > m \right\} \cup \left\{ \|e^{-3\mathfrak{Y}^\circ}(v + v_{\text{ref}})\|_{C^{-1/2-\varepsilon}} > m \right\}, \end{aligned}$$

with

$$\mathbb{P}(\|\mathfrak{I}(t)\|_{C^{-1/2-\varepsilon}} > m) + \mathbb{P}(\|\mathfrak{Y}^\circ(t)\|_{C^{-1/2-\varepsilon}} > m) = o_m(1)$$

uniformly in $t \geq 0$ by stationarity. We also have

$$\begin{aligned} & \mathbb{P}\left(\|e^{-3\mathfrak{Y}^\circ(t)}(v + v_{\text{ref}})(t)\|_{C^{-1/2-\varepsilon}} > m\right) \\ & \leq \mathbb{P}\left(\|e^{-3\mathfrak{Y}^\circ(t)}\|_{C^{-1/2-\varepsilon}} \geq c\right) + \mathbb{P}\left(\|(v + v_{\text{ref}})(t)\|_{C^{-1/2-\varepsilon}} > \frac{m}{c}\right) \\ & \leq o_c(1) + \mathbb{P}\left(\|v(t)\|_{C^{-1/2-\varepsilon}} > \frac{m}{2c}\right) + \mathbb{P}\left(\|v_{\text{ref}}(t)\|_{C^{-1/2-\varepsilon}} > \frac{m}{2c}\right) \\ & \leq o_c(1) + o_{m/c}(1). \end{aligned}$$

The $o_m(1)$ function does not depend on t by stationarity. In the last step we used the ϕ -independent estimate (2.17) quantifying the upper bound (2.16) in the coming down from infinity property together with the stationarity of v_{ref} . This gives the t -uniform and ϕ -independent estimate

$$(6.5) \quad \mathbb{P}(\|u(t)\|_{C^{-1/2-\varepsilon}} > 3m) = o_m(1).$$

We have been cautious to construct an enhanced noise whose law is stationary in time. This property together with the independence of the estimate (6.5) with respect to the initial condition allows then to propagate (6.5) uniformly in time by restarting fictively the dynamics every integer time while keeping an upper bound $o_m(1)$ that does not depend on the interval considered. The family of laws $\mathcal{L}(u_r(t, \phi))$ of $u_r(t, \phi)$ is thus tight in $C^{-1/2-2\varepsilon}(M)$, independently of the regularization parameter $r \in [0, 1]$ and the initial condition $\phi \in C^{-1/2-\varepsilon}(M)$, uniformly in $t \geq 1$. It follows that for any $\phi \in C^{-1/2-2\varepsilon}(M)$ the probability measures on $C^{-1/2-\varepsilon}(M)$

$$\frac{1}{T-1} \int_1^T \delta_{\mathcal{L}(u(t, \phi))} dt, \quad (T \geq 2)$$

have a weak limit along a subsequence of times T tending to infinity. The Feller property of the semigroup generated by (1.3) ensures that this weak limit is an invariant probability measure of the dynamics. \triangleright

6.2 Non-triviality of the ϕ_3^4 measure

Some care is needed when working with Jagannath & Perkowski's representation

$$u_r = \mathfrak{I}_r - \mathfrak{Y}_r^\circ + e^{-3\mathfrak{Y}_r^\circ}(v_r + v_{r, \text{ref}})$$

of u_r when it comes to taking the expectation of some quantities. This is related to the fact that the random variable \mathfrak{Y} being a quadratic polynomial of a Gaussian noise the random variable $\exp(-3\mathfrak{Y}^\circ)$ may not be integrable. This a priori makes tricky to say anything about the integrability of $u_r(t)$ from its description in terms of $v_r(t)$. To circumvent this problem we follow Jagannath & Perkowski' suggestion to trade $\exp(-\mathfrak{Y}^\circ)$ for $\exp(-P_{\geq n} \mathfrak{Y}^\circ)$ in their change of unknown (2.3). The operator

$$P_{\geq n} = \sum_{i \in I} \sum_{|k| \geq n} P_k^i$$

removes a number of initial terms of a Littlewood-Paley expansion. One can thus *choose n random* so that

$$\|P_{\geq n}^{\mathfrak{Y}}\|_{C_T C^{1-\eta}} \leq 1.$$

Set

$$f_n \stackrel{\text{def}}{=} \sum_{i \in I} \sum_{|k| \leq n-1} P_k^i(\mathfrak{Y}) \in C_T C^\infty(M).$$

This change of unknown adds a term $f_n v$ into the equation for v , which only changes Z_1 for a new Z_1 that is still an element of $C_T C^{-1/2-\varepsilon}(M)$ and is a polynomial of the noise. As $P_k^i P_\ell^j = 0$ for $|k - \ell|$ greater than a fixed constant we have

$$\|f_n\|_{C_T C^{1-2\varepsilon}} \lesssim \|\mathfrak{Y}\|_{C_T C^{1-2\varepsilon}}$$

independently of our definition of the random integer n . So the new Z_1 has finite moments of any order. We get from the estimates (2.16) and (2.17) quantifying of the coming down, with $\exp(-\mathfrak{Y})$ now replaced by $\exp(-P_{\geq n}^{\mathfrak{Y}})$, and the formula (2.3) relating u and v the fact that

$$(6.6) \quad u(1) = \mathfrak{Y}(1) + \mathfrak{Y}^{\circ\circ}(1) + (\star)$$

for an element $(\star) \in C^{1-\eta}(M)$ whose norm belongs to all the $L^q(\Omega)$ spaces, $1 \leq q < \infty$, uniformly with respect to the initial condition ϕ of u . We now see clearly that $u(1) \in C^{-1/2-\varepsilon}(M)$ belongs to all the $L^q(\Omega)$ spaces, $1 \leq q < \infty$, *uniformly with respect to the initial condition ϕ .* We assume in the remainder of this section that we work with this version of Jagannath & Perkowski's equation.

The mechanics of the proof that the Φ_3^4 measure is non-Gaussian is well-known. We write it here for completeness and follow for that purpose the lecture notes [44] of Gubinelli – Section 6.4 therein, after Gubinelli & Hofmanová's work [46]. Assume ϕ is random, with law the invariant measure of the dynamics, so $u(1)$ itself has the same law. Consider the heat regularisation $e^{r\Delta}u(1)$ of our solution u at time 1. In this subsection, for simplicity, we shall assume that we used true Wick ordering for the renormalisation which simplifies the discussion and allows to use true orthogonality properties of the Wiener chaos decomposition. Our argument is of semiclassical nature, we will use the small r asymptotic behaviour of heat kernels to justify nontriviality – so r somehow plays the role of a semiclassical parameter. If the Φ_3^4 measure were Gaussian the random variable $e^{-rP}(u(1))$ would also be Gaussian uniformly when $r > 0$ goes to 0. So its truncated four point function

$$\begin{aligned} C_4^r &= C_4 \left(e^{-rP}(u(1)), e^{-rP}(u(1)), e^{-rP}(u(1)), e^{-rP}(u(1)) \right) \\ &\stackrel{\text{def}}{=} \mathbb{E} [e^{-rP}(u(1))^4] - 3 \mathbb{E} [e^{-rP}(u(1))^2]^2 \end{aligned}$$

would be identically null uniformly in $r \in (0, 1]$. The sufficient integrability of the different elements of the decomposition

$$u(1) = \mathfrak{Y}(1) - \mathfrak{Y}^{\circ\circ}(1) + e^{-3P_{\geq n}^{\mathfrak{Y}}(1)} \mathfrak{Y}^{\circ\circ}(1) (v(1) + v_{\text{ref}}(1)),$$

allows to plug it inside the formula for the fourth order cumulant and use Wick's Theorem to get

$$\begin{aligned} &C_4 \left(e^{-rP}(u(1)), e^{-rP}(u(1)), e^{-rP}(u(1)), e^{-rP}(u(1)) \right) \\ &= 24 \int_{-\infty}^t G_r^{(3)}(t-s) \circ e^{-(t-s+r)P}(x, x) ds + 216 \int_{(-\infty, t]^2} \int_{y_1, y_2 \in U^2} G_0^{(2)}(s_1 - s_2, y_1, y_2) \\ &\quad \times e^{-(r+t-s_1)P}(y_1, x) e^{-(r+t-s_2)P}(y_2, x) G_r^{(1)}(t-s_1, y_1, x) G_r^{(1)}(t-s_2, y_2, x) dy_{12} ds_{12} \\ &\quad + \mathbb{E} \left[(e^{-rP} \mathfrak{Y}) P \left(e^{-rP} \mathfrak{Y}^{\circ\circ}(1), e^{-rP} e^{-3P_{\geq n}^{\mathfrak{Y}}(1)} \mathfrak{Y}^{\circ\circ}(1) (v(1) + v_{\text{ref}}(1)) \right) \right] \end{aligned}$$

where P is some polynomial functional in its stochastic arguments. We have many cancellations in the above expression since Gaussian cumulants only retain connected Feynman graphs and we also use orthogonality of homogeneous Wiener chaoses of different degrees. The remainder has the corresponding decay

$$\mathbb{E} \left[\mathfrak{Y}(1) P \left(e^{-rP} \mathfrak{Y}^{\circ\circ}(1), e^{-rP} e^{-3P_{\geq n}^{\mathfrak{Y}}(1)} \mathfrak{Y}^{\circ\circ}(1) (v(1) + v_{\text{ref}}(1)) \right) \right] = \mathcal{O}(r^{-\frac{1}{4}})$$

since we just need to recall that the remainder only involves the terms,

$$e^{-rP}\mathcal{Y}(1) = \mathcal{O}(r^{-\frac{1}{4}}), \quad e^{-rP}\mathcal{Y}^{\text{qo}}(1) = \mathcal{O}(1), \quad e^{-rP}e^{-3P_{\geq n}}\mathcal{Y}^{\text{qo}}(1)(v(1) + v_{\text{ref}}(1)) = \mathcal{O}(1)$$

since they are Hölder regular in $\mathcal{C}^{\frac{1}{2}-0}$ and \mathcal{C}^{1-0} respectively. (We used the fact that we can probe the space Hölder regularity by testing against heat kernels: $\sup_{\varepsilon \in (0,1]} \varepsilon^{-\frac{\delta}{2}} \|e^{-\varepsilon P}u\|_{L^\infty(M)} \lesssim \|u\|_{C^\delta(M)}$ and also we made an implicit use of Besov embeddings, $\forall \delta > 0, \|\cdot\|_{C^{s-\frac{d}{p}-\delta}} \lesssim \|\cdot\|_{B_{p,p}^s}$ together with hypercontractive estimates which allows us to consider expectations of Hölder norms.)

Let us study in detail the asymptotics of the first term on the right hand side of the equation for C_4^r which has a Feynman integral interpretation. For every $x \in U$, choose some cut-off function $\chi \in C_c^\infty(U)$ which equals 1 near x that we use to localize the asymptotics as in the calculation of counterterms, then we can extract the small r leading asymptotics as

$$\begin{aligned} & \int_{-\infty}^t G_r^{(3)}(t-s) \circ e^{-(t-s+r)P}(x, x) ds \\ & \simeq \int_0^\infty \int_{y \in U} \chi(y) \left(\int_{[a+r, +\infty)^3} \prod_{i=1}^3 e^{-s_i P}(x, y) ds_i \right) e^{-(a+r)P}(y, x) da. \end{aligned}$$

We compute the integral with respect to y first; this reads

$$\begin{aligned} & \int \left(\prod_{i=1}^3 K^0(s_i, x, y) \right) K^0(a+r, x, y) \chi(y) dy \\ & \simeq \frac{e^{-(s_1+s_2+s_3+a+r)} (4\pi)^{-6}}{(s_1 s_2 s_3 (a+r))^{\frac{3}{2}}} \int_U e^{-\frac{|x-y|g(x)}{4} (s_1^{-1}+s_2^{-1}+s_3^{-1}+(a+r)^{-1})} \chi(y) \det(g)_y^{\frac{1}{2}} dy \\ & \simeq \frac{(4\pi)^{-\frac{9}{2}} e^{-(s_1+s_2+s_3+a+r)}}{(s_1 s_2 s_3 (a+r))^{\frac{3}{2}} (s_1^{-1} + s_2^{-1} + s_3^{-1} + (a+r)^{-1})^{\frac{3}{2}}} \\ & \quad + \mathcal{O} \left((s_1 s_2 s_3 (a+r))^{-\frac{3}{2}} (s_1^{-1} + s_2^{-1} + s_3^{-1} + (a+r)^{-1})^{-\frac{5}{2}} \right) \end{aligned}$$

where we only keep the leading terms in the heat asymptotic expansion and use a stationary phase estimate. It is possible, as we did for the counterterms, to show that the integral with respect to a, s_1, s_2, s_3 of the $\mathcal{O}(\dots)$ term gives subleading asymptotics compared to the leading term. We are reduced after a change of variables to the asymptotics of the following integral

$$\int_0^1 (a+r)^{-\frac{3}{2}} \left(\int_{[1, +\infty)^3} (a_2 a_3 + a_1 a_3 + a_1 a_2 + a_1 a_2 a_3)^{-\frac{3}{2}} da_{123} \right) da \simeq cr^{-\frac{1}{2}}$$

for some non-null constant c . The next term in the formula for C_4^r is

$$\begin{aligned} & \int_{(-\infty, t]^2} \int_{y_1, y_2 \in U^2} G_0^{(2)}(s_1 - s_2, y_1, y_2) e^{-(r+t-s_1)P}(y_1, x) e^{-(r+t-s_2)P}(y_2, x) \\ & \quad \times G_r^{(1)}(t - s_1, y_1, x) G_r^{(1)}(t - s_2, y_2, x) dy_1 dy_2 ds_{12} \end{aligned}$$

which is bounded by a constant multiple of the integral over $(-\infty, t]^2 \times U^2$ of

$$\left(\sqrt{|s_1 - s_2|} + |y_1 - y_2| \right)^{-2} \left(\sqrt{|r+t-s_1|} + |y_1 - x| \right)^{-4} \left(\sqrt{|r+t-s_2|} + |y_2 - x| \right)^{-4}.$$

Making first the change of variables $s_i \mapsto r^2(s_i - t) + s_i$, $y_i \mapsto r(y_i - x) + x$ and then using polar coordinates gives $\mathcal{O}(|\log r|)$ as an upper bound for that integral, therefore the cumulant C_4^r blows-up like $24cr^{-\frac{1}{2}}$ when $r > 0$ goes to 0. This shows that C_4^r does not vanish asymptotically and that the Φ_3^4 measure is non-Gaussian.

For a Φ_3^4 measure obtained as above as weak limit of Birkhoff averages, the covariance property under Riemannian isometries is clear from its construction and the fact that the renormalisation constants a_r, b_r do not depend on which Riemannian metric is used: given a field ϕ on (M, g) whose law is a Φ_3^4 measure, let $f : M' \mapsto M$ be a smooth diffeomorphism, then the pulled-back field $f^*\phi$ on (M', f^*g) will have the law of a Φ_3^4 measure of the SPDE (1.3) for the metric f^*g .

Such measure gives for the first time a non-perturbative, non-topological interacting quantum field theory on 3-dimensional curved Riemannian spaces. We prove in [6] that the semigroup on $C^{-1/2-\varepsilon}(M)$ generated by the dynamics (1.3) has a unique invariant probability measure. This uniqueness result yields a stronger notion of covariance.

7 – Non-constant coupling function and vector bundle cases

7.1 Non-constant coupling function

Denote by $\mu_{\text{GFF}}(d\phi)$ the Gaussian free field measure on M . We construct in this section a Φ_3^4 measure on M with formal description

$$(7.1) \quad \nu(d\phi) \propto e^{-\int_M \lambda(x)\phi^4(x)dx} \mu_{\text{GFF}}(d\phi)$$

corresponding to a space dependent coupling constant $\lambda(x)$, say a C^∞ function of x such that $\lambda > 0$ on M . We considered so far the case $\lambda = 1$. We show that the results proved to analyse this special case allow to deal with the general case. We construct the measure (7.1) as an invariant measure of a Markovian evolution in $C^{-1/2-\varepsilon}(M)$. The goal of this section is to prove that the **counterterms are local functionals** in the coupling function $\lambda \in C^\infty(M, \mathbb{R}_{>0})$ which is a deep feature of renormalisation: *The divergent counterterms you need to subtract at some point x depends only on finite jets of the Lagrangian functional density at the same point x .* We refer to the work [1] of Abdesselam which also deals with space dependent couplings for some hierarchical model in 3d.

Let us make a detailed calculation to determine the fine structure of the divergences that arise in the equation as well as to establish the locality of the counterterms. We use the same regularisation ξ_r as above. Start from $\mathcal{L}u = \xi_r - \lambda u^3$, set a first decomposition $u = \mathfrak{I}_r + Z$, hence $\mathcal{L}(\mathfrak{I}_r + Z) = \xi_r - (\mathfrak{I}_r + Z)^3$ therefore

$$\begin{aligned} \mathcal{L}Z &= -\lambda \left(\mathfrak{I}_r^3 + 3Z\mathfrak{I}_r^2 + 3Z^2\mathfrak{I}_r + Z^3 \right) = -\lambda \left(\mathfrak{V}_r + 3a_r\mathfrak{I}_r + 3Z\mathfrak{V}_r + 3Za_r + 3Z^2\mathfrak{I}_r + Z^3 \right) \\ &= -\lambda \left(\mathfrak{V}_r + 3Z\mathfrak{V}_r + 3Z^2\mathfrak{I}_r + Z^3 \right) - 3\lambda a_r u \end{aligned}$$

where we compare the covariant Wick renormalized powers \mathfrak{V}_r with the nonrenormalized powers and the counterterms are shown in **red** to clearly see the difference.

This motivates to define a new regularized equation for u_r as

$$\mathcal{L}u = \xi_r - \lambda u^3 + 3\lambda a_r u$$

where adding the counterterm in **red** has the effect of Wick renormalising the trees appearing on the r.h.s of the equation for Z . Therefore, in what follows, all trees that appear are covariantly Wick renormalized. We also define the following new stochastic trees decorated by the subscript $\cdot\lambda$:

$$\mathfrak{Y}_{r,\lambda}^{\circ\circ} \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\lambda \mathfrak{V}_r)$$

and

$$\mathfrak{Y}_{r,\lambda}^{\circ} \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\lambda \mathfrak{V}_r),$$

this means they have an extra coupling function λ inserted at the vertex to avoid confusion with all previous trees introduced earlier where the coupling function was set to $\lambda = 1$.

Now we introduce a further decomposition writing u as $u = \mathfrak{I}_r - \underbrace{\mathfrak{Y}_{r,\lambda}^{\circ\circ}}_{=Z} + R$ where we used a

subscript $\cdot\lambda$ to denote the fact that the nonlinear term u^3 has become λu^3 which affects the trees in the equation. Then the next remainder R satisfies the new equation:

$$\mathcal{L}R = -3\lambda \mathfrak{V}_r \left(R - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right) - 3\lambda \mathfrak{I}_r \left(R - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^2 - \lambda \left(R - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^3$$

with Wick renormalized r.h.s. Let us anticipate a bit on what follows and try to guess what problematic terms we will encounter next. We spot two problematic terms in the equation for R , first we expect that R has regularity $1-$ and \mathfrak{V}_r has regularity $-1-$ hence the product $R\mathfrak{V}_r$ is

ill-defined with the borderline regularity. Furthermore the product $3\lambda\mathcal{V}_r\overset{\circ}{\mathcal{Y}}_{r,\lambda}$ is clearly ill-defined and requires a renormalisation. We will discuss later how to deal with this term.

As in the work of Jagannath–Perkowski, we introduce a Cole–Hopf transform $v_r \stackrel{\text{def}}{=} e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}} R$ to kill the borderline ill-defined product $R\mathcal{V}_r$. A long and tedious calculation yields the following equation for v_r :

$$\begin{aligned} \mathcal{L}v_r = & 9|\nabla\overset{\circ}{\mathcal{Y}}_{r,\lambda}|^2v_r - 3v_r\overset{\circ}{\mathcal{Y}}_{r,\lambda} - 6\nabla(\overset{\circ}{\mathcal{Y}}_{r,\lambda})\nabla v_r + 3\lambda e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}}\mathcal{V}_r\overset{\circ}{\mathcal{Y}}_{r,\lambda} \\ & - 3\lambda e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}} \left(e^{-3\overset{\circ}{\mathcal{Y}}_{r,\lambda}}v_r - \overset{\circ}{\mathcal{Y}}_{r,\lambda} \right)^2 \mathfrak{I}_r - \lambda e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}} \left(e^{-3\overset{\circ}{\mathcal{Y}}_{r,\lambda}}v_r - \overset{\circ}{\mathcal{Y}}_{r,\lambda} \right)^3. \end{aligned}$$

Let us again try to identify the origin of new divergences in this new equation. It may arise from the product $e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}} \left(\lambda\mathcal{V}_r\overset{\circ}{\mathcal{Y}}_{r,\lambda} \right)$ where we need to renormalize the quintic term in parenthesis. Once it is renormalized it will be of regularity $-1 - 0$ and we still need to discuss how we can make sense of the product with $e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}}$ which is of regularity $1-$. As usual, by the paraproduct decomposition, it is the resonant term $\lambda\mathcal{V}_r \odot \overset{\circ}{\mathcal{Y}}_{r,\lambda}$ which requires a renormalisation which is very similar to what we did for the quintic term τ_4 except for the insertion of the coupling function λ at the vertices. Let us indicate what changes must be made on the treatment of τ_{41} in order to renormalize this term with the coupling function λ . The divergent subamplitude \mathcal{A}_r now reads

$$\mathcal{A}_r(t, s, x_*, x_1, x_2, x_a) \stackrel{\text{def}}{=} [\odot](x_*, x_1, x_2)\underline{\mathcal{L}}^{-1}((t, x_1), (s, x_a))G_r^{(2)}((t, x_2), (s, x_a))\lambda(x_2)\lambda(x_a)$$

where the function λ appearing in the subamplitude \mathcal{A}_r is taken at two different points x_2 and x_a . Now if one repeats the proof of Lemma 38 with the function λ inserted at the right places, one ends up with a counterterm of the form

$$c_r(t_1, x_*) = \mathcal{S} \int_{-\infty}^{t_1} \sum_{0 \leq |k-\ell| \leq 1, i} P_k^i(x_*, \cdot) \circ \tilde{\chi} \lambda \circ \left(G_r^{(2)}(t-s) \circ e^{-(t-s)P} \right) \circ \lambda \tilde{\chi} \circ \tilde{P}_\ell^i(\cdot, x_*) ds$$

where the P_k^i, \tilde{P}_ℓ^i are the Littlewood–Paley–Stein projectors, the functions $\tilde{\chi} \in C_c^\infty(U)$ are arbitrary test functions such that $\tilde{\chi} = 1$ near x_* and where we used the explicit definition of $[\odot]$ in terms of the Littlewood–Paley–Stein projectors. Observe that changing the position of λ at two places in the above composition of operators inserts two commutator terms and since the commutators lower the pseudodifferential order $[\Psi^{m_1}, \Psi^{m_2}] \in \Psi^{m_1+m_2-1}$, we get

$$\begin{aligned} c_r(t, x_*) &= \mathcal{S} \int_{-\infty}^t \sum_{0 \leq |k-\ell| \leq 1, i} \lambda^2(x_*) P_k^i(x_*, \cdot) \circ \tilde{\chi} \circ \left(G_r^{(2)}(t-s) \circ e^{-(t-s)P} \right) \circ \tilde{\chi} \circ \tilde{P}_\ell^i(\cdot, x_*) ds \\ &+ \text{Trace density of some element in } \Psi^{-4}(M) \\ &= \mathcal{S} \int_{-\infty}^t \sum_{0 \leq |k-\ell| \leq 1, i} \lambda^2(x_*) P_k^i(x_*, \cdot) \circ \tilde{\chi} \circ \left(G_r^{(2)}(t-s) \circ e^{-(t-s)P} \right) \circ \tilde{\chi} \circ \tilde{P}_\ell^i(\cdot, x_*) ds \end{aligned}$$

since elements in $\Psi^{-4}(M)$ are trace class. Therefore as we saw in Section 5.5.5, we find that the term

$$\lambda\mathcal{V}_r \odot \overset{\circ}{\mathcal{Y}}_{r,\lambda} - \lambda^2 b_r \mathfrak{I}_r$$

converges as $r \downarrow 0$ in $\mathcal{C}^{-1/2-5\varepsilon}([0, T] \times M)$ in $L^2(\Omega)$. There is no problem for multiplying it with

$$\exp(3\overset{\circ}{\mathcal{Y}}_{r,\lambda}) \in \mathcal{C}^{1-2\varepsilon}([0, T] \times M)$$

and send r to 0. For this reason, in the ill-defined product $\underbrace{e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}}}_{\text{well-defined}} \left(\underbrace{\lambda\mathcal{V}_r\overset{\circ}{\mathcal{Y}}_{r,\lambda} - \lambda^2 b_r \mathfrak{I}_r}_{\text{ill-defined}} \right)$ where the term in parenthesis is well-defined at the limit $r \downarrow 0$ but the product of the two terms underbraced is ill-defined at the limit $r \downarrow 0$.

In our companion paper [10], we show that using two renormalisations yields the existence of

$$e^{3\overset{\circ}{\mathcal{Y}}_{r,\lambda}} \left(\lambda\mathcal{V}_r\overset{\circ}{\mathcal{Y}}_{r,\lambda} - \lambda^2 b_r \mathfrak{I}_r - \lambda^2 b_r \overset{\circ}{\mathcal{Y}}_{r,\lambda} \right)$$

as a random variable valued in $\mathcal{C}^{-4\varepsilon}([0, T] \times M) + C_T \mathcal{C}^{\frac{1}{2}-7\varepsilon} + C_T \mathcal{C}^{-1-2\varepsilon}(M)$ when $r \downarrow 0$. Now we define a new element ϕ_r by the equation:

$$v_r = Y + \phi_r = \underbrace{\underline{\mathcal{L}}^{-1} \left(3e^{3\mathfrak{Y}_{r,\lambda}} \left(\lambda \mathfrak{V}_r \mathfrak{Y}_{r,\lambda}^{\circ\circ} - \lambda^2 b_r \left(\mathfrak{I}_r + \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right) \right) \right)}_{=Y} + \phi_r$$

where $Y \in \mathcal{C}^{1-\varepsilon}([0, T] \times M)$, $\forall \varepsilon > 0$ as $r \downarrow 0$.

We rewrite the equation for v_r in terms of both (Y, ϕ_r) , the goal of introducing Y is that it is well-defined thanks to the double renormalisations we just performed and this equation makes appear new divergent terms:

$$\begin{aligned} \mathcal{L}\phi_r = & 9|\nabla \mathfrak{Y}_{r,\lambda}^{\circ\circ}|^2(Y + \phi_r) - 3(Y + \phi_r)\mathfrak{Y}_{r,\lambda}^{\circ\circ} - 6\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})\nabla(Y + \phi_r) + 3\lambda^2 b_r e^{3\mathfrak{Y}_{r,\lambda}} \left(\mathfrak{I}_r + \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right) \\ & - 3\lambda e^{3\mathfrak{Y}_{r,\lambda}} \left(e^{-3\mathfrak{Y}_{r,\lambda}}(Y + \phi_r) - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^2 \mathfrak{I}_r - \lambda e^{3\mathfrak{Y}_{r,\lambda}} \left(e^{-3\mathfrak{Y}_{r,\lambda}}(Y + \phi_r) - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^3. \end{aligned}$$

The first divergent term in purple on the r.h.s comes from the counterterm in the definition of Y . The next terms which are ill-posed read $9|\nabla \mathfrak{Y}_{r,\lambda}^{\circ\circ}|^2$ and $\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})\nabla Y$ since Y is a priori in $C_T \mathcal{C}^{1-2\varepsilon}(M)$ and $\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ}) \in C_T \mathcal{C}^{-2\varepsilon}(M)$ hence the scalar product will be ill-defined. In the companion work [10], we also show how to extract the singular term from this scalar product. More precisely:

$$\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})\nabla Y = |\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})|^2 3e^{3\mathfrak{Y}_{r,\lambda}} \mathfrak{Y}_{r,\lambda}^{\circ\circ} + \text{well-defined.}$$

We isolated the singular term at the limit $r \downarrow 0$ as $|\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})|^2$. It remains to explain how to extract the counterterm of $|\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})|^2$ while taking into account the presence of the coupling function λ . This was done at the end of Section 5.5.4. Similarly, the singular part of $|\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})|^2$ is the same as the singular part of $(\Delta \mathfrak{Y}_{r,\lambda}^{\circ\circ})\mathfrak{Y}_{r,\lambda}^{\circ\circ}$ whose divergent part now reads

$$\mathcal{S} \left(\int_{-\infty}^t tr_{L^2} \left(\lambda \circ e^{-s'_2 P} \circ \lambda \circ G_r^{(2)}(s'_2) ds'_2 \right) \right)$$

where the λ in the above expression is viewed as multiplication operator. Note that the difference $\int_{-\infty}^t tr_{L^2} \left((\lambda \circ e^{-s'_2 P} \circ \lambda - \lambda^2 \circ e^{-s'_2 P}) \circ G_r^{(2)}(s'_2) ds'_2 \right)$ is regular when $r \downarrow 0$ simply by the fact that $\lambda(x)\lambda(y) - \lambda^2(x)$ vanishes near the diagonal $x = y$ and by simple power counting argument as we did when we studied amplitudes. Finally, the singular part

$$\mathcal{S} \left(\int_{-\infty}^t tr_{L^2} \left(\lambda^2 \circ e^{-s'_2 P} \circ G_r^{(2)}(s'_2) ds'_2 \right) \right) = \lambda^2(x) \frac{b_r}{3}$$

as we did already calculate for quartic graphs. This means that

$$\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})\nabla Y - \lambda^2(x) b_r e^{3\mathfrak{Y}_{r,\lambda}} \mathfrak{Y}_{r,\lambda}^{\circ\circ}$$

has a well-defined limit when $r \downarrow 0$.

In the same way the limit $9|\nabla \mathfrak{Y}_{r,\lambda}^{\circ\circ}|^2 - 9\lambda^2 \frac{b_r}{3}$ has a well-defined limit. Let us add and subtract the divergent terms in the equation for ϕ_r to single out the divergences which are again represented in *red*, this now reads

$$\begin{aligned} \mathcal{L}\phi_r = & (9|\nabla \mathfrak{Y}_{r,\lambda}^{\circ\circ}|^2 - 3\lambda^2 b_r)(Y + \phi_r) - 3(Y + \phi_r)\mathfrak{Y}_{r,\lambda}^{\circ\circ} - 6\nabla(\mathfrak{Y}_{r,\lambda}^{\circ\circ})\nabla(Y + \phi_r) \\ & + 6\lambda^2(x) b_r e^{3\mathfrak{Y}_{r,\lambda}} \mathfrak{Y}_{r,\lambda}^{\circ\circ} + 3\lambda^2 b_r (Y + \phi_r) - 6\lambda^2 b_r e^{3\mathfrak{Y}_{r,\lambda}} \mathfrak{Y}_{r,\lambda}^{\circ\circ} + 3\lambda^2 b_r e^{3\mathfrak{Y}_{r,\lambda}} \left(\mathfrak{I}_r + \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right) \\ & - 3\lambda e^{3\mathfrak{Y}_{r,\lambda}} \left(e^{-3\mathfrak{Y}_{r,\lambda}}(Y + \phi_r) - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^2 \mathfrak{I}_r - \lambda e^{3\mathfrak{Y}_{r,\lambda}} \left(e^{-3\mathfrak{Y}_{r,\lambda}}(Y + \phi_r) - \mathfrak{Y}_{r,\lambda}^{\circ\circ} \right)^3. \end{aligned}$$

Now note that the sum of all the terms in red equals $3\lambda^2 b_r e^{3\mathfrak{Y}_{r,\lambda}} u$, so introduce the function

$$(7.2) \quad v_{r,\text{ref},\lambda} \stackrel{\text{def}}{=} 3\underline{\mathcal{L}}^{-1} \left(\lambda e^{3\mathfrak{Y}_{r,\lambda}} \left\{ \mathfrak{Y}_{r,\lambda}^{\circ\circ} \mathfrak{V}_r - b_r (\mathfrak{I}_r + \lambda \mathfrak{Y}_{r,\lambda}^{\circ\circ}) \right\} \right).$$

This is an element of $C_T C^{1-\varepsilon}(M)$ that converges in any of these spaces as r goes to 0 in $L^2(\Omega)$. Setting

$$v_r \stackrel{\text{def}}{=} e^{3\mathfrak{Y}_r} \left(u_r - \mathfrak{I}_r + \lambda \mathfrak{Y}_r^{\circ\circ} \right) - v_{r,\text{ref},\lambda}$$

we see that u_r is a solution to the renormalized equation

$$(7.3) \quad \mathcal{L}u = \xi_r - \lambda u^3 + (3\lambda a_r - 3\lambda^2 b_r) u$$

has a well-defined solution in the limit $r \downarrow 0$, the solution enjoys all the properties we proved in case λ is constant. if and only if v_r is a solution to an equation of the form

$$\mathcal{L}v_r = b_{r,\lambda} \nabla v_r - a_{r,\lambda} v_r^3 + Z_{2,r,\lambda} v_r^2 + Z_{1,r,\lambda} v_r + Z_{0,r,\lambda}$$

where $b_{r,\lambda} \in C_T C^{-\varepsilon}(M)$, $a_{r,\lambda} \in C_T C^{1-2\varepsilon}(M)$ and $Z_{i,r,\lambda} \in C_T C^{-1/2-\varepsilon}(M)$ all converge in their spaces as $r > 0$ goes to 0 in $L^2(\Omega)$, for all $\varepsilon > 0$. This analysis of equation (7.3) puts us in a position to go in the present setting over all the different steps that we have done above to construct the Φ_3^4 measure when $\lambda = 1$ and provides a construction of the Φ_3^4 measure in a setting where the coupling constant is space dependent.

7.2 The Φ_3^4 vectorial model in the bundle case

We conclude this section by showing that our result also holds when ϕ is taken to be the section of a vector bundle. This corresponds to the construction of the vectorial ϕ_3^4 measure (which is sometimes called the $O(N)$ -model in the physics literature). We summarize here what changes need to be done in the bundle case, while more technical details are postponed in our companion paper. First, we consider a Hermitian vector bundle $E \mapsto M$, smooth (resp. C^α , distributional) sections of E are denoted by $\Gamma^\infty(M, E) = C^\infty(E)$ (resp $\Gamma^\alpha(M, E) = C^\alpha(E)$, $\Gamma^{-\infty}(M, E) = \mathcal{D}'(E)$) and we are given some generalized Laplacian Δ_g which means Δ_g is a symmetric differential operator acting on $C^\infty(E)$ such that its principal symbol is positive definite, symmetric, diagonal, it reads $g_{\mu\nu}(x) \xi^\mu \xi^\nu \otimes Id_{End(E_x)}$ as a function on $C^\infty(T^*M, End(E))$ where $g_{\mu\nu}$ is the induced Riemannian cometric on T^*M . We furthermore assume that $-\langle \varphi, \Delta_g \varphi \rangle_{L^2(E)} \geq 0$ for all $\varphi \in C^\infty(E)$. The above in particular implies that $-\Delta_g$ is a nonnegative, elliptic second order operator. The corresponding heat operator now reads $\mathcal{L} = \partial_t + 1 - \Delta_g$. It is well-known from elliptic theory that $P := 1 - \Delta_g$ has self-adjoint extension as $P : H^2(E) \mapsto L^2(E)$, P has compact self-adjoint resolvent with discrete real spectrum in $\mathbb{R}_{\geq 0}$ and the eigenfunctions of P form an L^2 -basis of the space $L^2(E)$ of L^2 sections of E . In this case, we can define some E -valued white noise as $\xi = \sum_{\lambda \in \sigma(P)} c_\lambda e_\lambda$ where the sum runs over the eigenvalues of P , e_λ are the eigensections of P and $c_\lambda \sim \mathcal{N}(0, 1)$ are i.i.d gaussian random variables. The E -valued GFF reads $P^{-\frac{1}{2}} \xi$. The goal is to make sense of the Gibbs measure

$$\nu(d\phi) \propto e^{-\int_M \lambda \langle \varphi, \varphi \rangle_E} \mu_{GFF}(d\phi)$$

where $\langle \cdot, \cdot \rangle_E$ denotes the Hermitian scalar product of E , the interaction term now reads $(\langle \varphi, \varphi \rangle_E)^2$ in the vectorial case and $\lambda \in C^\infty(M, \mathbb{R}_{>0})$ is the coupling function.

The corresponding vectorial ϕ_3^4 SPDE reads:

$$(7.4) \quad \mathcal{L}u = -\lambda u \langle u, u \rangle + (\text{rk}(E) + 2)(\lambda a_r - \lambda^2 b_r) u + \xi_r$$

where u is an E -valued random distribution over space time $\mathbb{R} \times M$, $\xi_r = e^{-rP} \xi$ and a_r, b_r are the *exact same constants than in the scalar case*. All E -valued Besov (resp Hölder, Sobolev) distributions are defined almost exactly like in the scalar case using local charts on M and local trivialisations of $E \mapsto M$, we denote them by $\mathcal{B}_{p,q}^s(E)$ (resp $C^s(E), H^s(E)$). Because the analytical properties of the heat kernel $(e^{-tP})_{t \geq 0}$ acting on sections of E are exactly the same as in the scalar case, both inverses \mathcal{L}^{-1} and $\underline{\mathcal{L}}^{-1}$ are well-defined with the exact same definitions and they have the exact same analytical properties as in the scalar case. The symbol \mathfrak{I}_r still denotes $\underline{\mathcal{L}}^{-1} \xi_r$. Because of the classical results on the asymptotic expansion of the heat kernel in the bundle case [17], the key idea is that the singularities are valued in diagonal elements in $C^\infty(End(E))$, we immediately find that the covariant Wick renormalisation for the cubic power reads $\mathfrak{V}_r \stackrel{\text{def}}{=} \langle \mathfrak{I}_r, \mathfrak{I}_r \rangle \mathfrak{I}_r - (\text{rk}(E) + 2) a_r \mathfrak{I}_r$ for the *same universal constant a_r* as in the scalar case and $\text{rk}(E)$ is the rank of the vector bundle E . Beware that the cubic vertex has a new meaning, it is a Hermitian scalar product in the fibers of E times an element of a fiber of E since u^3 has become $\langle u, u \rangle_E u$. The new stochastic tree now

reads

$$\mathring{\Psi}_{r,\lambda}^{\circ} \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\lambda \mathring{\mathcal{V}}_r) \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}(\lambda(\langle \mathring{\mathfrak{I}}_r, \mathring{\mathfrak{I}}_r \rangle \mathring{\mathfrak{I}}_r - (\text{rk}(E) + 2)a_r \mathring{\mathfrak{I}}_r)).$$

Then we introduce a similar decomposition as: $u = \mathring{\mathfrak{I}}_r - \mathring{\Psi}_{r,\lambda}^{\circ} + R$. Writing the equation satisfied by R , we see that the new term in the bundle case we need to eliminate is the borderline ill-defined product $-\lambda \langle \mathring{\mathfrak{I}}_r, \mathring{\mathfrak{I}}_r \rangle R - 2\lambda \langle R, \mathring{\mathfrak{I}}_r \rangle \mathring{\mathfrak{I}}_r$. We define some random endomorphism $\mathring{\mathcal{V}}_r$ acting on smooth sections $C^\infty(E)$ as

$$\mathring{\mathcal{V}}_r : T \in C^\infty(E) \mapsto \langle \mathring{\mathfrak{I}}_r, \mathring{\mathfrak{I}}_r \rangle T + \langle T, \mathring{\mathfrak{I}}_r \rangle \mathring{\mathfrak{I}}_r - (\text{rk}(E) + 2)a_r T \in \mathcal{D}'(E).$$

Observe that with this definition, it holds $3\mathring{\mathcal{V}}_r = \mathring{\mathcal{V}}_r(\mathring{\mathfrak{I}}_r) - 2(\text{rk}(E) + 2)\mathring{\mathfrak{I}}_r$, which is consistent with the fact that $\mathring{\mathcal{V}}_r$ is the renormalized version of $3\mathring{\mathfrak{I}}_r^2$. The bundle morphism $\mathring{\mathcal{V}}_r$ is *local* since it is $C^\infty(M)$ -linear hence it can be identified canonically with some random element in $\mathcal{D}'(M, \text{End}(E))$. Therefore, we define a new *vectorial Cole–Hopf transform* needed in the bundle case in terms of the above random endomorphism $\mathring{\mathcal{V}}_r$ as follows:

Definition 42 – *In the above notation, we define the random bundle map:*

$$(7.5) \quad R \stackrel{\text{def}}{=} e^{-\underline{\mathcal{L}}^{-1}\lambda \mathring{\mathcal{V}}_r}(v_r)$$

where similar stochastic estimates in the bundle case allow to prove that $\underline{\mathcal{L}}^{-1}\lambda \mathring{\mathcal{V}}_r$ is a.s. in $\mathcal{C}^{1-\varepsilon}(M, \text{End}(E))$ for all $\varepsilon > 0$.

Accordingly, one also defines the stochastic object

$$v_{r,\text{ref}} \stackrel{\text{def}}{=} \underline{\mathcal{L}}^{-1}\left(e^{\underline{\mathcal{L}}^{-1}\lambda \mathring{\mathcal{V}}_r} \left\{ \mathring{\mathcal{V}}_r(\mathring{\Psi}_{r,\lambda}^{\circ}) - (\text{rk}(E) + 2)b_r(\mathring{\mathfrak{I}}_r + \mathring{\Psi}_{r,\lambda}^{\circ}) \right\}\right),$$

that obeys the same estimate in the bundle case and is a.s. in $C_T \mathcal{C}^{1-\varepsilon}$ for all $\varepsilon > 0$. Similarly, define

$$\begin{aligned} \tau_1 : T \in C^\infty(E) &\mapsto \langle \mathring{\mathfrak{I}}_r \odot \mathring{\Psi}_{r,\lambda}^{\circ} \rangle T + \langle T, (\mathring{\mathfrak{I}}_r \odot \mathring{\Psi}_{r,\lambda}^{\circ}) \rangle + \langle T, (\mathring{\Psi}_{r,\lambda}^{\circ} \odot \mathring{\mathfrak{I}}_r) \rangle, \\ \tau_2 : T \in C^\infty(E) &\mapsto \mathring{\mathcal{V}}_r \odot (\underline{\mathcal{L}}^{-1}(\lambda \mathring{\mathcal{V}}_r(T))) - (\text{rk}(E) + 2)b_r T, \\ \tau_3 : T \in C^\infty(E) &\mapsto \nabla \underline{\mathcal{L}}^{-1}(\lambda \mathring{\mathcal{V}}_r) \odot (\nabla \underline{\mathcal{L}}^{-1}(\lambda \mathring{\mathcal{V}}_r(T))) - (\text{rk}(E) + 2)b_r T, \\ \tau_4 &\stackrel{\text{def}}{=} \mathring{\mathcal{V}}_r \odot (\mathring{\Psi}_{r,\lambda}^{\circ}) - (\text{rk}(E) + 2)b_r \mathring{\mathfrak{I}}_r. \end{aligned}$$

τ_1 is local and belongs to $C_T \mathcal{C}^{0-}(M, \text{End}(E))$, while τ_2 and τ_3 are not local, and only belong to $C_T \mathcal{L}(C^\infty(E), \mathcal{C}^{0-}(E))$. Finally, it holds $\tau_4 \in C_T \mathcal{C}^{-1/2-}(E)$. Contrary to the second Wick power, we do not need τ_2 and τ_3 to be local, since we do not aim to rise them to some exponent or take their exponential, and always evaluate them at some $T \in \mathcal{C}^\alpha(E)$. The proofs that these objects are correctly renormalized, and that $v_{r,\text{ref}}$ obeys the same estimates can be found in our companion paper. Regarding the global existence estimates, the L^p estimates can be obtained exactly in the same way in the vector bundle case, taking care of pairing the equation with $\langle v, v \rangle^{(p-2)/2} v$ instead of v^{p-1} , and to perform this pairing both in E and in $L^2(M)$.

A – Littlewood-Paley-Stein projectors

There are several ways of defining some Littlewood-Paley type projectors on function spaces over a manifold – see [59, 7, 8, 50, 70] for a sample. We choose here an intermediate road and use the classical Littlewood-Paley projectors over \mathbb{R}^d to define a number of operators on functions spaces over M using local charts. This allows to import at low cost some known regularity properties of the corresponding objects from the flat to the curved setting. We denote as usual by $B_{p,q}^\gamma(M)$ the Besov spaces over M and by $C^\gamma(M)$ the Besov-Hölder space $B_{\infty,\infty}^\gamma(M)$, with associated norm denoted by $\|\cdot\|_{C^\gamma}$.

Let then denote by

$$a' \prec b' \stackrel{\text{def}}{=} \sum_{-1 \leq j < k-1} (\Delta_j a')(\Delta_k b')$$

the paraproduct of some distributions a' and b' on \mathbb{R}^d , and write

$$a' \odot b' \stackrel{\text{def}}{=} \sum_{|j-k| \leq 1} (\Delta_j a')(\Delta_k b')$$

for the resonance of a' and b' whenever the latter is defined. Let $(U_i, \kappa_i)_i$ denote a finite open cover of M by some charts, with κ_i a smooth diffeomorphism between $U_i \subset M$ and $\kappa_i(U_i) \subset \mathbb{R}^d$. Let $(\chi_i)_i$ be a partition of unity subordinated to $(U_i)_i$, so $\sum_i \chi_i = 1$, with $\chi_i \in C_c^\infty(U_i)$. Choose also for every index i a function $\tilde{\chi}_i \in C_c^\infty(U_i)$ such that $\tilde{\chi}_i$ equals 1 on the support of χ_i and some function $\psi_i \in C_c^\infty(\kappa_i(U_i))$ which equals 1 on the support of $\kappa_{i*}(\tilde{\chi}_i)$. Given some smooth functions a, b on M we have the decomposition

$$\begin{aligned} ab &= \sum_{i \in I} (a\chi_i)(b\tilde{\chi}_i) = \sum_{i \in I} \kappa_i^* [\kappa_{i*}(a\chi_i)] \kappa_i^* [\kappa_{i*}(b\tilde{\chi}_i)] \\ &= \sum_{i \in I} \kappa_i^* [(\kappa_{i*}(a\chi_i)(\kappa_{i*}(b\tilde{\chi}_i))] = \sum_{i \in I} \kappa_i^* [\psi_i \kappa_{i*}(a\chi_i) \kappa_{i*}(b\tilde{\chi}_i)] \\ &= \sum_{i \in I} \kappa_i^* [\psi_i (\kappa_{i*}(a\chi_i) \prec \kappa_{i*}(b\tilde{\chi}_i))] + \sum_{i \in I} \kappa_i^* [\psi_i (\kappa_{i*}(a\chi_i) \odot \kappa_{i*}(b\tilde{\chi}_i))] \\ &\quad + \sum_{i \in I} \kappa_i^* [\psi_i (\kappa_{i*}(a\chi_i) \succ \kappa_{i*}(b\tilde{\chi}_i))]. \end{aligned}$$

Actually for arbitrary $\chi_i, \tilde{\chi}_i \in C_c^\infty(U_i)^2$ such that $\tilde{\chi}_i = 1$ on the support of χ_i , we set the generalized Littlewood-Paley-Stein projectors

$$(A.1) \quad P_k^i(a) \stackrel{\text{def}}{=} \kappa_i^* [\psi_i \Delta_k (\kappa_{i*}(\chi_i a))], \text{ and } \tilde{P}_k^i(a) \stackrel{\text{def}}{=} \kappa_i^* [\tilde{\psi}_i \Delta_k (\kappa_{i*}(\tilde{\chi}_i a))],$$

where $\tilde{\psi}_i \in C_c^\infty(\kappa_i(U_i))$ equals 1 on the support of ψ_i . We do not necessarily require that $\sum_{i \in I} \tilde{\chi}_i = 1$.

On the manifold M , recall $i \in I$ denotes a chart index, we define generalized chart localized operations as:

$$a \prec_i b \stackrel{\text{def}}{=} \sum_{-1 \leq j < k - 1} P_j^i a \tilde{P}_k^i b, \quad a \succ_i b \stackrel{\text{def}}{=} \sum_{i \in I} \sum_{-1 \leq k < j - 1} P_j^i a \tilde{P}_k^i b, \quad \text{and } a \odot_i b \stackrel{\text{def}}{=} \sum_{i \in I} \sum_{|j-k| \leq 1} P_j^i a \tilde{P}_k^i b.$$

In particular when $\sum_{i \in I} \chi_i = 1$, the above operations decompose the product ab on M as:

$$ab = \sum_{i \in I} (a \prec_i b + a \odot_i b + a \succ_i b).$$

Note the important fact that the definition of the resonant product and paraproducts are asymmetrical, therefore they are noncommutative meaning that $a \prec b \neq b \succ a$, however all the regularity properties are similar as in the flat case. We collect in the next two statements some regularity properties of these operators and refer the reader to our companion work [10] for their proofs.

Proposition 43 – *One has the following continuity estimates. For every chart index i ,*

- For p, p_1, p_2, q, q_1, q_2 in $[1, +\infty]$ with $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$ and $\frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{q}$

(a) For $\gamma_2 \in \mathbb{R}$

$$\|a \prec_i b\|_{B_{p,q_2}^{\gamma_2}} \lesssim \|a\|_{L^{p_1}} \|b\|_{B_{p_2,q_2}^{\gamma_2}},$$

(b) For $\gamma_1 < 0$ and $\gamma_2 \in \mathbb{R}$

$$\|a \prec_i b\|_{B_{p,q}^{\gamma_1+\gamma_2}} \lesssim \|a\|_{B_{p_1,q_1}^{\gamma_1}} \|b\|_{B_{p_2,q_2}^{\gamma_2}}$$

(c) For any $\gamma_1, \gamma_2 \in \mathbb{R}$ with $\gamma_1 + \gamma_2 > 0$ one has

$$\|a \odot_i b\|_{B_{p,q}^{\gamma_1+\gamma_2}} \lesssim \|a\|_{B_{p_1,q_1}^{\gamma_1}} \|b\|_{B_{p_2,q_2}^{\gamma_2}}.$$

We recall from Lemma 7.2 of Mourrat & Weber's work [68] the following comparison test that we used in our proof of Theorem 5.

Proposition 44 – *Let a continuous function $F : [0, T] \rightarrow [0, +\infty)$ that satisfies the inequality*

$$(A.2) \quad \int_s^t F(s_1)^\lambda ds_1 \leq c(F(s) + 1)$$

for all $0 \leq s \leq t \leq T$, for some exponent $\lambda > 1$ and some positive constant c . Then there is a sequence of times $t_0 = 0 < t_1 < \dots < t_N = T$ such that one has

$$F(t_n) \leq 1 + 2^{\frac{\lambda}{\lambda-1}} \left(\frac{c}{1 - 2^{-(\lambda-1)}} \right)^{\frac{1}{\lambda-1}} t_{n+1}^{-\frac{1}{\lambda-1}},$$

for all $0 \leq n \leq N - 1$.

Proof – We include a proof following closely Mourrat-Weber’s comparison test in a slightly different setting compared to them since we have $F(s) + 1$ rather than $F(s)$ on the right hand side of the inequality (A.2). We first define $t_0 = 0$, then given some time t_n , consider $t_{n+1}^* = t_n + c2^\lambda(1 + F(t_n))^{1-\lambda}$, if $t_{n+1}^* \geq T$ we stop the algorithm and set $N = n + 1$, $t_{n+1} = T$ and verify that the conclusion of the statement holds. Otherwise, choosing t_{n+1} such that $F(t_{n+1}) = \inf_{t_n < s < t_{n+1}^*} F(s)$ yields a bound of the form $F(t_{n+1}) \leq \frac{1+F(t_n)}{2}$. By iteration, this yields a bound of the form $F(t_{n+1}) \leq \frac{F(t_0)-1}{2^{n+1}} + 1$. Note that for n large enough, since $\lambda > 1$

$$t_{n+1}^* - t_n = c2^\lambda(1 + F(t_n))^{1-\lambda} \geq c2^\lambda \left(2 + \frac{F(t_0) - 1}{2^n} \right)^{1-\lambda} \geq c2^\lambda(2 + 1/3)^{1-\lambda} \geq 2c,$$

hence the algorithm must terminate for n large enough after finite number of iterations. Now we need to check the conclusion $t_n = \sum_{k=0}^{n-1} (t_{k+1} - t_k) \leq c2^\lambda \sum_i (1 + F(t_i))^{1-\lambda}$. Note that since $F(t_i) \geq (F(t_n) - 1)2^{n-i} + 1$ then $(1 + F(t_i))^{1-\lambda} \leq ((F(t_n) - 1)2^{n-i} + 2)^{1-\lambda}$, so we have

$$\begin{aligned} t_{n+1} &\leq c2^\lambda \sum_{i=0}^n ((F(t_n) - 1)2^{n-i} + 2)^{1-\lambda} \leq c2^\lambda (F(t_n) - 1)^{1-\lambda} \sum_{i=0}^n 2^{(n-i)(1-\lambda)} \\ &\leq c2^\lambda (F(t_n) - 1)^{1-\lambda} \frac{1}{1 - 2^{1-\lambda}}, \end{aligned}$$

which yields the estimate from the statement. \triangleright

Last we recall the fractional Leibniz rule and an elementary interpolation result used in the proof of the coming down property in Section 2.2, we prove these results in [10].

Proposition 45 – Let $\alpha > 0$, $r \in \mathbb{N}$ and $p, p_1, p_2, q \in [1, \infty]$ such that

$$\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}.$$

Then

$$\|u^{r+1}\|_{B_{p,q}^\alpha} \lesssim \|u^r\|_{L^{p_1}} \|u\|_{B_{p_2,q}^\alpha}.$$

Proposition 46 – Let $\alpha_1, \alpha_2 \in \mathbb{R}$ and $p_1, p_2, q_1, q_2 \in [1, \infty]$ and $\theta \in [0, 1]$. Define $\alpha = \theta\alpha_1 + (1-\theta)\alpha_2$, and $p, q \in [1, \infty]$ by

$$\frac{1}{p} = \frac{\theta}{p_1} + \frac{1-\theta}{p_2} \quad \text{and} \quad \frac{1}{q} = \frac{\theta}{q_1} + \frac{1-\theta}{q_2}.$$

Then

$$\|u\|_{B_{p,q}^\alpha} \lesssim \|u\|_{B_{p_1,q_1}^{\alpha_1}}^\theta \|u\|_{B_{p_2,q_2}^{\alpha_2}}^{1-\theta}.$$

References

- [1] A. Abdesselam, *Towards three-dimensional conformal probability*. p-Adic Numbers, Ultrametric Analysis and Applications 10 (2018): 233-252.
- [2] B.I. Albert, *Heat kernel renormalisation on manifolds with boundary*. arXiv:1609.02220, (2016).
- [3] S. Albeverio and S. Kusuoka, *The invariant measure and the flow associated to the Φ_3^4 -quantum field model*. Ann. Sc. Norm. Super. Pisa. Cl. Sci. **5**(20):1359–1427, (2020).
- [4] S. Albeverio and S. Kusuoka, *Construction of a non-Gaussian and rotation-invariant Φ_3^4 -measure and associated flow on \mathbb{R}^3 through stochastic quantisation*. arXiv:2102.08040, (2021).
- [5] Atiyah, M., R. Bott, and V. K. Patodi. *On the heat equation and the index theorem*. Inventiones Mathematicae 19.4 (1973): 279-330.
- [6] I. Bailleul, *Uniqueness of the Φ_3^4 measures on closed Riemannian 3-manifolds*. arXiv:2306.07616 (2023).
- [7] I. Bailleul and F. Bernicot, *Heat semigroup and singular PDEs*. J. Funct. Anal., **270**:3344–3452, (2016).
- [8] I. Bailleul and F. Bernicot and D. Frey, *Spacetime para-products for paracontrolled calculus, 3d-PAM and multiplicative Burgers equations*. Ann. Sc. Éc. Norm. Sup., **51**:1399–1457, 2018.

- [9] I. Bailleul and F. Bernicot, *High order paracontrolled calculus*. Forum Math., Sigma, **7**(e-44):1–93, (2019).
- [10] I. Bailleul and V.N. Dang and L. Ferdinand and T. D. Tô, *Global harmonic analysis for Φ_3^4 on closed Riemannian manifolds*. arXiv:2306.07757, (2023)
- [11] I. Bailleul and V.N. Dang and A. Mouzard, *Analysis of the Anderson operator*. arXiv:2201.04705, (2022).
- [12] Balaban, Tadeusz. *Ultraviolet stability of three-dimensional lattice pure gauge field theories*. Communications in mathematical physics 102.2 (1985): 255-275.
- [13] Balaban, Tadeusz. *Renormalization group approach to lattice gauge field theories: I. Generation of effective actions in a small field approximation and a coupling constant renormalization in four dimensions*. Communications in mathematical physics 109 (1987): 249-301.
- [14] N. Barashkov and M. Gubinelli, *A variational method for Φ_3^4* . Duke Math. J., **169**(17): 3339–3415, (2020).
- [15] N. Barashkov and M. Gubinelli, *The Φ_3^4 measure via Girsanov theorem*. Elec. J. Probab., **26**:1–29, (2021).
- [16] Barata, João CA, Christian D. Jäkel, and Jens Mund. *Interacting quantum fields on de Sitter Space*. arXiv preprint arXiv:1607.02265 (2016).
- [17] Berline, Nicole, Ezra Getzler, and Michele Vergne. *Heat kernels and Dirac operators*. Springer, 2003.
- [18] Ch. Brouder and N.V. Dang and F. H elein, Continuity of the fundamental operations on distributions having a specified wave front set (with a counterexample by Semyon Alesker). *Studia Mathematica*, **232**(3):201–226, (2016).
- [19] Bonicelli, Alberto, Claudio Dappiaggi, and Paolo Rinaldi. *An Algebraic and Microlocal Approach to the Stochastic Nonlinear Schrödinger Equation*. Annales Henri Poincaré. 2023.
- [20] R. Brunetti and K. Fredenhagen, *Microlocal Analysis and Interacting Quantum Field Theories: Renormalization on Physical Backgrounds*. Comm. Math. Phys., **208**: 623-661, (2000).
- [21] Brunetti, Romeo, Klaus Fredenhagen, and Rainer Verch. *The generally covariant locality principle—a new paradigm for local quantum field theory*. Communications in Mathematical Physics 237 (2003): 31-68.
- [22] Burq, Nicolas, Laurent Thomann, and Nikolay Tzvetkov. *Remarks on the Gibbs measures for nonlinear dispersive equations*. Annales de la Faculté des sciences de Toulouse: Mathématiques. Vol. 27. No. 3. 2018.
- [23] R. Catellier and K. Chouk, *Paracontrolled distributions and the 3-dimensional stochastic quantization equation*. Ann. Probab., **46**(5):2621–2679, (2018).
- [24] K. Costello, *Renormalization and effective field theory*. Vol. 170. American Mathematical Society, 2011.
- [25] N.V. Dang, *The extension of distributions on manifolds, a microlocal approach*. Annales Henri Poincaré, **17**(4), (2016).
- [26] N.V. Dang and E. Herscovich, *Renormalization of quantum field theory on Riemannian manifolds*. Reviews in Math. Phys., **31**(6):1950017, (2019).
- [27] N.V. Dang, *Renormalization of quantum field theory on curved space-times, a causal approach*. arXiv preprint arXiv:1312.5674, (2013).
- [28] N.V. Dang and M. Wrochna, *Dynamical residues of Lorentzian spectral zeta functions*. arXiv preprint arXiv:2108.07529, to appear in J.  Ec. Polytechnique, (2022+).
- [29] Dahlqvist, Antoine, Joscha Diehl, and Bruce K. Driver. *The parabolic Anderson model on Riemann surfaces*. Probability Theory and Related Fields 174.1-2 (2019): 369-444.
- [30] Dimock, J. *Markov quantum fields on a manifold*. Reviews in Mathematical Physics 16.02 (2004): 243-255.
- [31] Dappiaggi, C., Drago, N., Rinaldi, P., and Zambotti, L. (2022). *A microlocal approach to renormalization in stochastic PDEs*. Communications in Contemporary Mathematics, 24(07), 2150075.
- [32] P. Duch, *Flow equation approach to singular stochastic PDEs*. arXiv preprint arXiv:2109.11380 (2021)
- [33] P. Duch, *Renormalization of singular elliptic stochastic PDEs using flow equation*. arXiv preprint arXiv:2201.05031 (2022)
- [34] S. Dyatlov and M. Zworski. *Dynamical zeta functions for Anosov flows via microlocal analysis*. Ann. Sc.  Ec. Norm. Sup., **49**:543–577, (2016).
- [35] Fefferman, Charles, and C. Robin Graham. *The ambient metric (AM-178)*. Princeton University Press, 2012.
- [36] Figari, Rodolfo, Raphael H oegh-Krohn, and Chiara R. Nappi. *Interacting relativistic boson fields in the de Sitter universe with two space-time dimensions*. Communications in Mathematical Physics 44 (1975): 265-278.
- [37] Gilkey, Peter B. *Invariance theory: the heat equation and the Atiyah-Singer index theorem*. Vol. 16. CRC press, 2018.
- [38] J. Glimm, *Boson fields with the Φ^4 interaction in three dimensions*. Comm. Math. Phys. **10**(1):1–47, (1968).
- [39] J. Glimm and A. Jaffe. *Positivity of the ϕ_3^4 Hamiltonian*. Fortschritte der Physik. Progress of Physics, 21:327–376, 1973.
- [40] J. Glimm and A. Jaffe, *Quantum Physics – A Functional Integral Point of View*. Springer, (1987).
- [41] D. Grieser, *Notes on heat kernel asymptotics*. <http://www.staff.unioldenburg.de/daniel.grieser/wwwlehre/Schriebe/heat.pdf>, (2004).
- [42] M. Hairer and H. Singh. *Regularity Structures on Manifolds and Vector bundles*. arXiv:2308.05049, (2023).
- [43] M. Gubinelli, *Lectures on Stochastic Quantization of ϕ_3^4 – Part I: From QM to Equantum field theory*. https://www.iam.uni-bonn.de/fileadmin/user_upload/gubinelli/sq-lectures-milan/sq-notes-1.pdf
- [44] M. Gubinelli, *Lectures on Stochastic Quantization of ϕ_3^4 – Part II: Stochastic quantization*. https://www.iam.uni-bonn.de/fileadmin/user_upload/gubinelli/sq-lectures-milan/sq-notes-2.pdf
- [45] M. Gubinelli and M. Hofmanova, *Global solutions to elliptic and parabolic Φ^4 models in Euclidean space*. Comm. Math. Phys., **368**(3):1201–1266, (2019).
- [46] M. Gubinelli and M. Hofmanova, *A PDE construction of the Euclidean Φ_3^4 quantum field theory*. Comm. Math. Phys., **384**(1):1–75, (2021).

- [47] M. Gubinelli and N. Perkowski, *KPZ reloaded*. *Comm. Math. Phys.*, **349**:165–269, (2017).
- [48] Guillarmou, C., Kupiainen, A., Rhodes, R., & Vargas, V. (2021). *Segal’s axioms and bootstrap for Liouville Theory*. arXiv preprint arXiv:2112.14859.
- [49] M. Gubinelli and P. Imkeller and N. Perkoswki, *Paracontrolled distributions and singular PDEs*. *Forum Math. Pi*, **3**-e6:1–75, (2015).
- [50] C. Guillarmou and Th. de Poyferré *A paradifferential approach for hyperbolic dynamical systems and applications*. *Tunisian J. Math.*, **4**:673–718, (2022).
- [51] M. Hairer, *A theory of regularity structures*. *Invent. Math.*, **198**(2):269–504, (2014).
- [52] M. Hairer and J. Mattingly, *The strong Feller property for singular stochastic PDEs*. *Ann. Institut H. Poincaré B*, **54**(3):1314–1340, (2018).
- [53] M. Hairer and P. Schönbauer, *The support of singular stochastic PDEs*. *Forum Math. Pi*, **10**(e1):1–127, (2022).
- [54] M. Hairer and R. Steele, *The Φ_3^4 measure has sub-Gaussian tails*. *J. Stat. Phys.*, **186**, (2022).
- [55] S. Hollands and R. M. Wald, *Local Wick Polynomials and Time Ordered Products of Quantum Fields in Curved Spacetime*. *Comm. Math. Phys.*, **223**:289–326, (2001).
- [56] S. Hollands and R. M. Wald, *Existence of local covariant time ordered products of quantum fields in curved spacetime*. *Comm. Math. Phys.*, **231**:309–345, (2002).
- [57] A. Jagannath and N. Perkowski, *A simple construction of the dynamical ϕ_3^4 model*. *Trans. Amer. Math. Soc.*, **376**:1507–1522, (2023).
- [58] Kandel, Santosh, Pavel Mnev, and Konstantin Wernli. *Two-dimensional perturbative scalar QFT and Atiyah–Segal gluing*. *Advances in Theoretical and Mathematical Physics* 25.7 (2021): 1847–1952.
- [59] S. Klainerman and I. Rodnianski, *A geometric approach to the Littlewood–Paley theory*. *Geom. Funct. Anal.*, **16**(1):126–163, (2006).
- [60] Khavkine, Igor, and Valter Moretti. *Analytic dependence is an unnecessary requirement in renormalization of locally covariant QFT*. *Communications in Mathematical Physics* 344.2 (2016): 581–620.
- [61] Ch. Kopper and V.F. Müller, *Renormalization Proof for Massive Theory on Riemannian Manifolds*. *Comm. Math. Phys.* **275**(2):331–372, (2007).
- [62] Lévy, Thierry. *Yang–Mills measure on compact surfaces*. American Mathematical Soc., 2003.
- [63] E. Meinrenken, *Euler-like vector fields, normal forms, and isotropic embeddings*. *Indagationes Mathematicae*, **32**(1): 224–245, (2021).
- [64] R. B. Melrose. *Spectral and scattering theory for the Laplacian on asymptotically Euclidian spaces*. Marcel Dekker, 1994.
- [65] Meyer, Yves. *Wavelets, vibrations and scalings*. No. 9. American Mathematical Soc., 1998.
- [66] A. Moinat and H. Weber, *Space-time localization for the dynamic Φ_3^4 model*. *Comm. Pure Appl. Math.*, **73**(12):2519–2555, (2020).
- [67] J.C. Mourrat and H. Weber, *Global well-posedness of the dynamic Φ^4 model in the plane*. *Ann. Probab.*, **45**(4):2398–2476, (2017).
- [68] J.C. Mourrat and H. Weber, *The dynamics of Φ_3^4 model comes down from infinity*. *Comm. Math. Phys.*, **356**(3):673–753, (2017).
- [69] J.C. Mourrat and H. Weber and W. Xu, *Construction of Φ_3^4 Diagrams for Pedestrians*. Meeting on Particle Systems and PDE’s. Springer, Cham, (2015).
- [70] A. Mouzard, *The Anderson Hamiltonian on a two-dimensional Riemannian manifold*. arXiv:2009.03549, to appear in *Ann. Institut H. Poincaré*, (2023).
- [71] A. Mouzard, *A paracontrolled calculus for singular stochastic PDEs on manifold : to infinity and beyond*. Phd thesis
- [72] Oh, T., Robert, T., Tzvetkov, N., and Wang, Y. (2020). *Stochastic quantization of Liouville conformal field theory*. arXiv preprint arXiv:2004.04194.
- [73] Parisi, G., Wu, Y.S. *Perturbation theory without gauge fixing*. *Scientia Sinica. Zhongguo Kexue* 24(4), 483–496 (1981)
- [74] Pickrell, Doug. *$P(\phi)_2$ quantum field theories and Segal’s axioms*. *Communications in Mathematical Physics* 280.2 (2008): 403–425.
- [75] G. Popineau and R. Stora; *A pedagogical remark on the main theorem of perturbative renormalization theory*. *Nuclear Physics B*, **912**:70–78, (2016).
- [76] Rejzner, Kasia. *Perturbative algebraic quantum field theory*. *Math. Phys. Stud.*, Springer (2016).
- [77] Rivasseau, Vincent. *From perturbative to constructive renormalization*. Vol. 46. Princeton University Press, 2014.
- [78] P. Tsatsoulis and H. Weber, *Spectral Gap for the Stochastic Quantization Equation on the 2-dimensional Torus*. *Ann. Inst. H. Poincaré Probab. Statist.*, **54**(3):1204–1249, (2018).
- [79] A. Vasy. *Microlocal analysis of asymptotically hyperbolic and Kerr-de Sitter spaces*. *Inventiones Math.*, 194:381–513, 2013. With an appendix by S. Dyatlov
- [80] E. Witten, *Why Does Quantum Field Theory In Curved Spacetime Make Sense? And What Happens To The Algebra of Observables In The Thermodynamic Limit? Dialogues Between Physics and Mathematics: CN Yang at 100*. Cham: Springer International Publishing, 241–284, (2022).

• I. Bailleul – Univ Brest, CNRS UMR 6205, Laboratoire de Mathématiques de Bretagne Atlantique, France. *E-mail*: ismael.bailleul@univ-brest.fr

- N.V. Dang – Sorbonne Université and Université Paris Cité, CNRS, IMJ-PRG, F-75005 Paris, France.
Institut Universitaire de France, Paris, France. *E-mail:* dang@imj-prg.fr
- L. Ferdinand – Laboratoire de Physique des 2 infinis Irène Joliot-Curie, UMR 9012, Université Paris-Saclay, Orsay, France. *E-mail:* lferdinand@ijclab.in2p3.fr
- T.D. Tô – Sorbonne Université and Université Paris Cité, CNRS, IMJ-PRG, F-75005 Paris, France. *E-mail:* tat-dat.to@imj-prg.fr