A new copula regression model for hierarchical data.

Talagbé Gabin Akpo^{a,b}, Louis-Paul Rivest^{a,*}

^aDépartement de Mathematiques et de Statistique, Université Laval, 2325, rue de l'Université, GIV 0A6, Québec, Canada. ^bInstitut National de la Recherche Scientifique, Centre Armand-Frappier Santé Biotechnologie, 531 boulevard des Prairies, H7V 1B7, Laval, Canada

Abstract

This paper proposes multivariate copula models for hierarchical data. They account for two types of correlation: one is between variables measured on the same unit and the other is a correlation between units in the same cluster. This model is used to carry out copula regression for hierarchical data that gives cluster specific prediction curves. In the simple case where a cluster contains two units and where two variables are measured on each one, the new model is constructed within a D-vine. Then we focus on situations where two variables are measured on the units of a cluster of arbitrary size. The proposed copula density has an explicit form; it is expressed in terms of three copula families. We study the properties of the model; compare it to the linear mixed model and end with special cases. When the three copula families and the marginal distributions are normal, the model is equivalent to a normal linear mixed model with random, cluster specific, intercepts. The method to select the three copula families and to estimate their parameters are proposed. We perform a Monte Carlo study of the parameter estimators. A data set on the marks of students in several school is used to implement the proposed model and to compare its performance to standard normal mixed linear models. *Keywords:* Exchangeability, Heterogeneity, Normal linear mixed models, Vine copula.

1. Introduction

A simple bivariate copula regression predicts dependent variable y using independent variable x. It proceeds by selecting a bivariate copula summarizing the relationship between x and y. The copula regression predictor is then constructed using characteristics of the conditional distribution of y given x derived from the selected copula, see [17], [14] and [5] exemplify this method while [18] derive its asymptotic properties and [1] investigates predictions errors. This method is easily extended a multivariate explanatory variable x.

The goal of this work is to generalize the basic copula regression model to hierarchical data: *x* and *y* are observed on units that are in clusters and one would like to include a cluster effect in the copula regression predictions. The classical regression model for hierarchical data is a normal linear mixed model, with cluster specific random slopes and intercepts, see [2], [25], [16] and [10].

^{*}Corresponding author. Email address : Louis-paul.rivest@mat.ulaval.ca

Preprint submitted to Elsevier

The exchangeable copula families have been proposed to model the residual dependency within cluster in this context, see [20] and [11]; [23] provide a survival data application of this approach.

A model for the joint distribution of all the *x* and the *y* variables in a cluster is first constructed. The conditions that the model must fulfill in order to yield suitable predictions are given in Section 2. It is required to meet an exchangeability assumption: permuting the units in a cluster does not change the joint distribution of the variables. It also relies on a partial conditional independence assumption that insures that the prediction of *y* for a unit does not depend on the *x* values for the other units in the cluster. The proposed model is then constructed within a D-vine in the simple case of a cluster containing two units. The general model, for clusters of arbitrary size, is introduced in Section 3. Afterwards, we study its properties by showing that, in particular cases, conditional versions are equivalent to the models of [2] and of [20]. We use the proposed copula to do cluster specific predictions. The copula model is then implemented in a data set of [10] and compared with standard normal mixed linear models.

2. Model construction : Exchangeability and conditional independence

This section considers that $d \ge 2$ variables are measured on all the units in a cluster; subscript *j* represents a unit, j = 1, ..., n, where *n* is the size of the cluster. The dependent variable for unit *j* is Y_j while X_j is the corresponding vector of d - 1 explanatory variables and $Z_j = (X_j^{\top}, Y_j)^{\top}$, is the vector of the *d* variables measured on unit *j*. Let $F_{d,1:n}(z_1, ..., z_n)$ be the joint cumulative distribution function (cdf) of the *nd* variables measured in the cluster, where $z_j = (x_j^{\top}, y_j)^{\top}$. The model is constructed using copulas; it is therefore of interest to define $F(x) = (F_1(x_1), ..., F_{d-1}(x_{d-1}))$ and G(y) as the marginal distributions of respectively X_j and Y_j which are the same for all the units. We let $U_j = F(X_j)$ and $V_j = G(Y_j)$ be random variables with uniform margins and $c_{d,1:n} \{(u_1, v_1), ..., (u_n, v_n)\}$ be the copula density for the joint distribution of $\{(U_j^{\top}, V_j)^{\top} : j = 1, ..., n\}$. We now give some conditions for the family of joint distributions $F_{d,1:n}(z_1, ..., z_n) : n = 2, 3, ...$ to give useful regression models.

2.1. Exchangeability

We consider the family of cumulative distribution functions defined by

$$\mathcal{F}_d = \left\{ F_{d,1:n}(z_1, \dots, z_n) : z_j \in \mathbb{R}^d, \ j = 1, \dots, n, \ n = 1, 2 \dots \right\}.$$

The familly \mathcal{F}_d is said to be *d*-exchangeable if, for all $n \ge 2$, $F_{d,1:n} \in \mathcal{F}_d$ satisfies the following conditions

i) **Permutation invariance :** For all permutations $\{\pi(1), \ldots, \pi(n)\}$ of $\{1, 2, \ldots, n\}$,

$$F_{d,1:n}(z_1,\ldots,z_n) = F_{d,1:n}\{z_{\pi(1)},\ldots,z_{\pi(n)}\}.$$
(1)

ii) **Closure on marginalization:** For any $r \le n$,

$$F_{d,1:r}(z_1,...,z_r) = F_{d,1:n}(z_1,...,z_r,\infty,...,\infty).$$
 (2)

This definition is similar to the classical definition of univariate exchangeability that is given in [15]. A simple example of *d*-exchangeability is a multivariate one way ANOVA model with random effects, $Z_j = A + E_j$, where A and E_j : j = 1, ..., n are $d \times 1$ independent random vectors and $\{E_j : j = 1, ..., n\}$ have the same distribution. The next proposition gives the form of the correlation matrix for a *d*-exchangeable random vector.

Proposition 1. Let $\{\mathbf{Z}_1, \dots, \mathbf{Z}_n\}$ be a set of $nd \times 1$ random vectors, verifying the definition of *d*-exchangeability given by (1) and (2). Then, the $nd \times nd$ correlation matrix of Pearson, Spearman, and Kendall's between these *n* vectors have the form

$$I_{n} \otimes \Sigma_{w} + J_{n} \otimes \Sigma_{b} = \begin{pmatrix} \Sigma_{w} + \Sigma_{b} & \Sigma_{b} & \dots & \Sigma_{b} \\ \Sigma_{b} & \Sigma_{w} + \Sigma_{b} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \Sigma_{b} \\ \Sigma_{b} & \dots & \Sigma_{b} & \Sigma_{w} + \Sigma_{b} \end{pmatrix},$$
(3)

where J_n is and $n \times n$ matrix of ones and \otimes denotes the Kroenecker product. Moreover, for Pearson and Spearman correlations, the matrices Σ_w and Σ_b are positive semi definite.

2.2. Partial conditional independence

This section proposes an assumption concerning the dependency within each cluster.

Definition 1. The random variables Y_j and $\{X_k : k \neq j\}$ are assumed to be independent, given X_j . This is a partial conditional independence assumption that can be written as

$$Y_j \perp \{X_k : k \neq j\} | X_j, \ j = 1, \dots, n.$$
 (4)

This condition is weaker than the conditional independence assumption underlying the standard regression model that can be formulated as

$$Y_{j} \perp \{(X_{k}, Y_{k}) : k \neq j\} | X_{j}, j = 1, \dots, n.$$

We now implement the exchangeability and the independence assumption within a *D*-vine for the joint distribution of two units within a cluster when d = 2, that is when $\mathbf{Z} = (X, Y)^{\mathsf{T}}$.

2.3. A D-vine construction of the model when d = n = 2

This section constructs a copula density function for the random variables in a cluster containing two units and verifying the properties of partial conditional independence and 2-exchangeability. The 4 random variables are $U_1 = F(X_1)$, $V_1 = G(Y_1)$, $U_2 = F(X_2)$ and $V_2 = G(Y_2)$. The four trees for the proposed *D*-vine are given in Figure 1.



Figure 1: Graphical representation of the proposed D-vine.

The decomposition involves 6 bivariate copulas $C_{U_1V_1}$, $C_{U_1U_2}$, $C_{U_2V_2}$, $C_{V_1U_2;U_1}$, $C_{U_1V_2;U_2}$ and $C_{V_1V_2;U_1U_2}$. The density of the multivariate copula corresponding to the trees given in Figure 1 can be derived from Joe [13, pp. 108]. It is given by

$$c\{(u_{1}, v_{1}), (u_{2}, v_{2})\} = c_{U_{1}V_{1}}(u_{1}, v_{1})c_{U_{2}V_{2}}(u_{2}, v_{2})c_{U_{1}U_{2}}(u_{1}, u_{2})c_{V_{1}U_{2};U_{1}}\left\{C_{V_{1}|U_{1}}(v_{1}|u_{1}), C_{U_{2}|U_{1}}(u_{2}|u_{1})\right\}$$

$$\times c_{U_{1}V_{2};U_{2}}\left\{C_{U_{1}|U_{2}}(u_{1}|u_{2}), C_{V_{2}|U_{2}}(v_{2}|u_{2})\right\}c_{V_{1}V_{2};U_{1}U_{2}}\left\{C_{V_{1}|U_{1}U_{2}}(v_{1}|u_{1}, u_{2}), C_{V_{2}|U_{1}U_{2}}(v_{2}|u_{1}, u_{2})\right\},$$
(5)

where the conditional distribution functions $C_{V|U}$ and $C_{W|UV}$ are defined by

$$C_{V|U}(v|u) = \frac{C_{UV}(u,v)}{\partial u}, \quad C_{W|UV}(w|u,v) = \frac{\partial C_{WV;U} \{ C_{V|U}(v|u), C_{W|U}(w|u) \}}{\partial C_{V|U}(v|u)}.$$
(6)

This vine is assumed to fulfill the simplifying assumption: the copulas associated to the second and third trees do not depend on the conditioning variables. Discussions of this vine decomposition can be found in [19], [8], and [7]. We would like the density (5) to fulfill the conditions for 2-exchangeability, see (1) and (2), and for partial conditional independence, see (4).

The density of the copula is 2-exchangeable if $c \{(u_1, v_1), (u_2, v_2)\} = c \{(u_2, v_2), (u_1, v_1)\}$. This requires a unique copula density for the dependency between U and V, that is $c_{U_1V_1}(u, v) = c_{U_2V_2}(u, v)$, for $u, v \in (0, 1)$. Also, the copula for the dependency between U_1 and U_2 needs to be symmetric, that is $c_{U_1U_2}(u, v) = c_{U_1U_2}(v, u)$ and $c_{V_1V_2;U_1U_2}(u, v) = c_{V_1V_2;U_1U_2}(v, u)$. This definition also entails restrictions on the conditional copula densities $c_{V_1U_2|U_1}$ and $c_{U_1V_2|U_2}$. However for (5) to fulfill the partial conditional independence condition (4) these two copulas need to be equal to the independence copula. This leads to following copula density

$$c\{(u_1, v_1), (u_2, v_2)\} = c_{U_1 U_2}(u_1, u_2) \times \left\{ c_{U_1 V_1}(u_1, v_1) c_{U_1 V_1}(u_2, v_2) \right\} c_{V_1 V_2; U_1 U_2} \{ C_{V_1 | U_1}(v_1 | u_1), C_{V_1 | U_1}(v_2 | u_2) \}.$$

$$(7)$$

It involves an arbitrary copula density c_{UV} for the relationship between U and V and two symmetric copula densities, $c_{U_1U_2}$ and $c_{V_1V_2;U_1U_2}$.

It is interesting to consider the special case where the three copulas in (7) are normal. Let ρ_1 represent the correlation between U_1 and U_2 , ρ_2 that between U and V and ρ_3 be the residual correlation. The joint copula density (7) for (U_1, U_2, V_1, V_2) is normal, see Joe [13, pp. 119]. Its correlation matrix is given by

In (8), the result that $\mathbb{E}\left\{\Phi^{-1}(U_1)\Phi^{-1}(V_2)\right\} = \rho_1\rho_2$ is obtained by conditioning on U_2 while the variance-covariance matrix of $\left\{\Phi^{-1}(V_1), \Phi^{-1}(V_2)\right\}$ knowing (U_1, U_2) is used to find that $\rho_3 = \left\{\mathbb{E}\left\{\Phi^{-1}(V_1)\Phi^{-1}(V_2)\right\} - \rho_1\rho_2^2\right\}/(1-\rho_2^2)$.

Given that (U_1, U_2) are distributed according to a normal copula with correlation ρ_1 , one obtains the normal copula with correlation matrix (8) for the joint distribution of (U_1, U_2, V_1, V_2) if V_1 and V_2 are defined by

$$\Phi^{-1}(V_j) = \rho_2 \Phi^{-1}(U_j) + \sqrt{\rho_3} (1 - \rho_2^2)^{1/2} A + (1 - \rho_2^2)^{1/2} (1 - \rho_3)^{1/2} E_j, \quad j = 1, 2,$$
(9)

where A, E_1, E_2 are independent with a N(0, 1) distribution. Model (9) is similar to the linear mixed model of [2]. The term $\rho_2 \Phi^{-1}(U_j)$ represents the contribution of the explanatory variables to the regression; it is fixed if one conditions on (U_1, U_2) . The cluster specific random intercept, $\sqrt{\rho_3}(1-\rho_2^2)^{1/2}A$, is independent of the experimental error, $(1-\rho_2^2)^{1/2}(1-\rho_3)^{1/2}E_j$. One can easily show that the correlation matrix of $\{\Phi^{-1}(U_1), \Phi^{-1}(U_2), \Phi^{-1}(V_1), \Phi^{-1}(V_2)\}$ entering in (9) is given by (8). Note also that (9) is easily generalized to clusters of size n > 2. This is also true of the general model (7). This leads to the general 2-exchangeable model that is proposed in the next section.

3. A multivariate 2-exchangeable copula model

This section proposes a method to construct densities for the joint distribution of $Z_j = (X_j, Y_j)^T$, for j = 1, ..., n that meets constraints of 2-exchangeability and of partial conditional independence presented in Section 2. The joint cdf for $\{Z_j : j = 1, ..., n\}$ is denoted $F_{2,1:n}$, where the first index, 2, refers to the dimension of Z while the index 1 : n means that it concerns n units labelled from 1 to n. This joint distribution involves marginal distributions F(x) and G(y) for X and Y and a copula density $c_{2,1:n} \{(u_1, v_1), ..., (u_n, v_n)\}$. This density depends on $c^{(2)}$, a copula density for the relationship between X and Y, and two families of 1-exchangeable copula densities, $\{c_{1,1:n}^{(1)}(u_1, ..., u_n) : n \ge 2\}$ and $\{c_{1,1:n}^{(3)}(u_1, ..., u_n) : n \ge 2\}$ for the dependency between the X variables and the copula regression residuals within a cluster respectively. These two families are assumed to fulfill conditions (1) and (2) for d = 1.

The general form for the 2-exchangeable copula density is

$$c_{2,1:n}\left\{(u_1, v_1), \dots, (u_n, v_n)\right\} = c_{1,1:n}^{(1)}(u_1, \dots, u_n) \times \prod_{j=1}^n \left\{c^{(2)}(u_j, v_j)\right\} \times c_{1,1:n}^{(3)}\left\{C_{2|1}(v_1|u_1), \dots, C_{2|1}(v_n|u_n)\right\},$$
(10)

where $u_j, v_j \in [0, 1], j = 1, ..., n$ and the conditional distribution $C_{2|1}$ is deduced from equation (6) with C_{UV} replaced by $C^{(2)}$. For n = 2, equation (10) reduces to the *D*-vine copula density (7)

To prove that (10) meets the conditions (1) and (2) for 2-exchangeability and (4) for conditional independence we integrate the proposed joint density for $\{(U_1, V_1), \dots, (U_n, V_n)\}$ in (10) for $v_n \in (0, 1)$. To carry this out, it is convenient to change variable, $w_n = C_{2|1}(v_n|u_n)$. The jacobian is $dw_n = c^{(2)}(u_u, v_n)dv_n$. Using the closure on marginalization property of copula family $C_{1,1:n}^{(3)}$, the integral is equal to

$$c_{1,1:n}^{(1)}(u_1,\ldots,u_n) \times \prod_{j=1}^{n-1} c^{(2)}(u_j,v_j) \times c_{1,1:(n-1)}^{(3)} \{ C_{2|1}(v_1|u_1),\ldots,C_{2|1}(v_{n-1}|u_{n-1}) \}$$
(11)

The variable u_n only appears in $c_{1,1,n}^{(1)}(u_1, \ldots, u_n)$. Using the closure on marginalization property of copula family $C_{1,1,n}^{(1)}$, the integral on u_n gives the density (10) for the (n - 1) pairs $(U_1, V_1), \ldots, (U_{n-1}, V_{n-1})$. Thus (10) defines a proper copula density that meets requirements (1) and (2) for 2-exchangeability. To prove the partial conditional independence assumption (4) one integrates (11) for $v_{n-1}, v_{n-2}, \ldots, v_2 \in (0, 1)$. This is easily carried by changing variables, $w_j = C_{2|1}(v_j|u_j), j = 2, \ldots, n - 1$. The joint density of (V_1, U_1, \ldots, U_n) is given by $c_{1,1,n}^{(1)}(u_1, \ldots, u_n)c^{(2)}(u_1, v_1)$; thus, given U_1, V_1 and (U_2, \ldots, U_n) are independent and (4) holds.

The derivations in the previous paragraph have highlighted a key property of the proposed model. If the density of $\{(U_1, V_1), \dots, (U_n, V_n)\}$ is (10) then the two vectors (U_1, \dots, U_n) and $\{W_1 = C_{2|1}(V_1|U_1), \dots, W_n = C_{2|1}(V_n|U_n)\}$ are independent with densities respectively given by $c_{1,1,n}^{(1)}(u_1, \dots, u_n)$ and $c_{1,1,n}^{(3)}(w_1, \dots, w_n)$. This is summarized in the following proposition.

Proposition 2. Let $(U_1, V_1), \ldots, (U_n, V_n)$, be a set of n random vectors, whose joint density is given by (10). If $W_j = C_{2|1}(V_j|U_j)$, $j = 1, \ldots, n$, where the distribution function $C_{2|1}$ comes from (6), then the random vectors (U_1, \ldots, U_n) and (W_1, \ldots, W_n) are independent with respective densities $c_{1,1;n}^{(1)}$ and $c_{1,1;n}^{(3)}$.

The result of Proposition 2 suggests the following algorithm to simulate a random vector with a density given of (10):

- Step 1 : Simulate (U_1, \ldots, U_n) according to the exchangeable copula $C_{1,1:n}^{(1)}(u_1, \ldots, u_n)$;
- Step 2 : Simulate (W_1, \ldots, W_n) according to the exchangeable copula $C_{1,1:n}^{(3)}(w_1, \ldots, w_n)$;
- Step 3 : Solve, the equations in V_j defined by $W_j = C_{2|1}(V_j|U_j)$, j = 1, ..., n, where $C_{2|1}$ comes from the equation (6). The the joint density of $(U_1, V_1, ..., U_n, V_n)$ is given by (10).

This algorithm differs from the proposal of Czado [6, pp. 136] to simulate from a D-vine, that goes through the vine sequentially.

It is interesting to construct the joint density of $\{(X_1, Y_1), \dots, (X_n, Y_n)\}$ from (10). It involves the joint marginal density of (X_1, \dots, X_n) ,

$$g_{1,1:n}^{(1)}(x_1,\ldots,x_n) = c_{1,1:n}^{(1)} \{F(x_1),\ldots,F(x_n)\} \prod_{j=1}^n f(x_j)$$

The conditional densities of Y_j given X_j , for j = 1, ..., n,

$$\prod_{j=1}^{n} \left[g(y_j) c^{(2)} \left\{ G(y_j), F(x_j) \right\} \right],$$

and a term for the residual dependency within clusters:

$$c_{1,1:n}^{(3)}\left[C_{2|1}\left\{G(y_1)|F(x_1)\right\},\ldots,C_{2|1}\left\{G(y_n)|F(x_n)\right\}\right].$$

This highlights a step wise construction of the 2-exchangeable model: first comes the specification of the marginal distribution for X, then that for the conditional distribution of Y given X and one finally adds a component for the residual dependency within a cluster.

We now assume that the copula $C^{(2)}$ is normal with correlation ρ_2 . The conditional distribution in (10) and its inverse are given by

$$C_{2|1}(v|u) = \Phi\left\{\frac{\Phi^{-1}(v) - \rho_2 \Phi^{-1}(u)}{\sqrt{1 - \rho_2^2}}\right\} \text{ and } C_{2|1}^{-1}(t|u) = \Phi\left\{\Phi^{-1}(t)\sqrt{1 - \rho_2^2} + \rho_2 \Phi^{-1}(u)\right\},$$

for $u, v, t \in [0, 1]$, see [3] for similar results. Using Proposition 2, the dependent variable Y_j for a unit in a cluster of size n with explanatory variable x_j can be expressed as $Y_j = G^{-1}[C_{2|1}^{-1}\{W_j|F(x_j)\}]$, where W_j is the entry of a vector with distribution $C_{1,1:n}^{(3)}$. This gives the following linear model :

$$\Phi^{-1}\{G(Y_j)\} = \rho_2 \Phi^{-1}\{F(x_j)\} + \sqrt{1 - \rho_2^2} \Phi^{-1}(W_j) \cdot$$

If, in addition, the marginal distributions of X and Y are normal: $F(y) = \Phi\{(x - \mu_1)/\sigma_1\}$ and $G(y) = \Phi\{(y - \mu_2)/\sigma_2\}$ where $\mu_1, \mu_2, \sigma_1^2, \sigma_2^2$ are respectively the marginal means and variances, then the model becomes:

$$Y_{j} = \beta_{0} + \beta_{1} x_{j} + \sigma_{e} \Phi^{-1}(W_{j}), \qquad (12)$$

where $\beta_0 = \mu_2 + \beta_1 \mu_1$, $\beta_1 = \rho_2 \sigma_2 / \sigma_1$, and $\sigma_e^2 = \sigma_2^2 (1 - \rho_2^2)$. In (12) the conditional marginal distribution of Y_j is normal.

The joint distribution depends on the copula $C_{1,1:n}^{(3)}$. These models are investigated in [20]. If this copula is normal and exchangeable, with correlation ρ_3 , then (12) reduces to the normal mixed model of [2]. Finally note that the conditional density in (12) for (Y_1, \ldots, Y_n) given $(X_1 = x_1, \ldots, X_n = x_n)$ is given by

$$f(y_1, \dots, y_n | x_1, \dots, x_n) = \frac{1}{(2\pi)^{n/2} \sigma_e^n} \exp\left\{-\frac{1}{2\sigma_e^2} \sum_{j=1}^n (y_j - \beta_0 - \beta_1 x_j)^2\right\} c_{1,1:n}^{(3)} \left[\Phi\left\{(y_1 - \beta_0 - \beta_1 x_1)/\sigma_e\right\}, \dots, \Phi\left\{(y_n - \beta_0 - \beta_1 x_n)/\sigma_e\right\}\right] \cdot \frac{1}{(2\pi)^{n/2} \sigma_e^n} \exp\left\{-\frac{1}{2\sigma_e^2} \sum_{j=1}^n (y_j - \beta_0 - \beta_1 x_j)^2\right\} c_{1,1:n}^{(3)} \left[\Phi\left\{(y_1 - \beta_0 - \beta_1 x_1)/\sigma_e\right\}, \dots, \Phi\left\{(y_n - \beta_0 - \beta_1 x_n)/\sigma_e\right\}\right] \cdot \frac{1}{(2\pi)^{n/2} \sigma_e^n} \exp\left\{-\frac{1}{2\sigma_e^2} \sum_{j=1}^n (y_j - \beta_0 - \beta_1 x_j)^2\right\} c_{1,1:n}^{(3)} \left[\Phi\left\{(y_1 - \beta_0 - \beta_1 x_1)/\sigma_e\right\}, \dots, \Phi\left\{(y_n - \beta_0 - \beta_1 x_n)/\sigma_e\right\}\right] \cdot \frac{1}{(2\pi)^{n/2} \sigma_e^n} \exp\left\{-\frac{1}{2\sigma_e^2} \sum_{j=1}^n (y_j - \beta_0 - \beta_1 x_j)^2\right\} c_{1,1:n}^{(3)} \left[\Phi\left\{(y_1 - \beta_0 - \beta_1 x_1)/\sigma_e\right\}, \dots, \Phi\left\{(y_n - \beta_0 - \beta_1 x_n)/\sigma_e\right\}\right\}$$

Thus model (12) can easily be fitted by maximum likelihood.

3.1. Predictions with the 2-exchangeable copula model

Suppose that (n - 1) units $\{(x_1, y_1), \dots, (x_{n-1}, y_{n-1})\}$ have been observed in a cluster. This section investigates the conditional distribution of Y_n given x_n in that cluster. A closed form expression for the conditional expectation of Y_n given x_n and $\{(x_1, y_1), \dots, (x_{n-1}, y_{n-1})\}$ is derived. Illustrations of the prediction curves for various specifications of the copula $C^{(2)}$ are presented.

The conditional density, $g_P(y_n)$ of Y_n given x_n and $\{(x_1, y_1), \dots, (x_{n-1}, y_{n-1})\}$ is expressed in terms of $w_j = C_{2|1} \{G(y_j)|F(x_j)\}$, $j = 1, \dots, n-1$. It is given by the ratio of (10) over (11), this yields

$$g_P(y_n) = g(y_n)c^{(2)} \{F(x_n), G(y_n)\} \frac{c_{1,1:n}^{(3)} [w_1, \dots, w_{n-1}, C_{2|1} \{G(y_n) | F(x_n)\}]}{c_{1,1:(n-1)}^{(3)} (w_1, \dots, w_{n-1})}.$$
(13)

Observe that the conditional density of $W = C_{2|1} \{G(Y_n)|F(x_n)\}$, given $\{(x_1, y_1), \dots, (x_{n-1}, y_{n-1})\}$, is simply $c_{1,1:n}^{(3)}(w_1, \dots, w_{n-1}, w)/c_{1,1:(n-1)}^{(3)}(w_1, \dots, w_{n-1})$. This is the conditional distribution of $(W_n|W_1, \dots, W_{n-1})$, when (W_1, \dots, W_n) is distributed according to copula $C_{1,1:n}^{(3)}$. Thus one can easily simulate from (13). Starting from the equation (13) and making a change of variable $w = C_{2|1} \{G(y_n)|F(x_n)\}$, we easily obtain the result.

The best predictor of the unknown Y_n is its conditional expectation, given $\{x_n, (x_1, y_1), \dots, (x_{n-1}, y_{n-1})\}$, where $w_j = \left[C_{2|1}\left\{G(y_j)|F(x_j)\right\}\right]$ it can be expressed as

$$\mathbb{E}_{P}(Y_{n}) = \int_{0}^{1} G^{-1} \left[C_{2|1}^{-1} \{ w | F(x_{n}) \} \right] \frac{c_{1,1:n}^{(3)}(w_{1}, \dots, w_{n-1}, w)}{c_{1,1:(n-1)}^{(3)}(w_{1}, \dots, w_{n-1})} dw$$
(14)

If the copula $C_{1,1:n}^{(3)}$ is the independence copula this reduces to a standard, unconditional copula regression for $C^{(2)}$, see [14], [5] or [18]. Taking the expectation of $\mathbb{E}_P(Y_n)$ with respect to the distribution of $\{(X_1, Y_1), \ldots, (X_{n-1}, Y_{n-1})\}$ also gives the unconditional copula regression curve for $C^{(2)}$. Thus the proposed model gives regression curves that vary between clusters and their expectation is equal to the marginal copula regression based on $C^{(2)}$.

We suppose that the copula $C_{1,1,n}^{(3)}$ is exchangeable normal with correlation ρ_3 . Thus if $W_j = C_{2|1} \{G(Y_j)|F(x_j)\}$, j = 1, ..., n, then the cumulative distribution function of random vector $\{\Phi^{-1}(W_1), ..., \Phi^{-1}(W_n)\}$ is a multivariate normal distribution with an exchangeable correlation matrix $\Sigma(n, \rho_3)$ whose entries are 1 on the diagonal and ρ_3 off the diagonal. Using standard properties of the multivariate normal distribution, the conditional distribution of $\Phi^{-1}(W_n)$ knowing $\{\Phi^{-1}(W_1), ..., \Phi^{-1}(W_{n-1})\}$ is univariate normal with mean μ_0 and variance σ_0^2 defined by

$$\mu_0 = \frac{(n-1)\rho_3 \sum_{j=1}^{n-1} \Phi^{-1}(W_j)/(n-1)}{1 + (n-2)\rho_3}, \quad \sigma_0^2 = \frac{(1-\rho_3) \{1 + (n-1)\rho_3\}}{1 + (n-2)\rho_3}, \tag{15}$$

see for instance, [20]. Thus the conditional distribution of random variable $\Phi^{-1}(W_n) = \Phi^{-1}[C_{2|1}\{G(Y_n)|F(x_n)\}$ is a $N(\mu_0, \sigma_0^2)$. The final form of (14) when $C_{1,1:n}^{(3)}$ is a normal copula is therefore given by

$$\mathbb{E}_{P}(Y_{n}) = \int_{\mathbb{R}} G^{-1} \left[C_{2|1}^{-1} \left\{ \Phi(\mu_{0} + \sigma_{0}z) | F(x_{n}) \right\} \right] \phi(z) dz, \tag{16}$$

where $\phi(z) = \exp(-z^2/2)/\sqrt{2\pi}$. The conditional expectation of the equation (16) is easily evaluated using the Gauss-Hermite quadrature method, see [22]. Note also that the density function (13) can be expressed as:

$$g_{P}(y_{n}) = g(y_{n})\frac{c^{(2)}\{F(x_{n}), G(y_{n})\}}{\sigma_{0}} \times \exp\left\{-\frac{1}{2\sigma_{0}^{2}}\left(\Phi^{-1}\left[C_{2|1}\{G(y_{n})|F(x_{n})\}\right] - \mu_{0}\right)^{2} + \frac{1}{2}\Phi^{-1}\left[C_{2|1}\{G(y_{n})|F(x_{n})\}\right]^{2}\right\}.$$
 (17)

The corresponding quantile function of g_p has a simple form, namely

$$G_P^{-1}(w) = G^{-1}\left(C_{2|1}^{-1}\left[\Phi\{\mu_0 + \sigma_0\Phi^{-1}(w)\}|F(x_n)\cdot\right]\right),$$

Thus the median and various quantiles of the conditional distribution of Y_n are easily evaluated.

When *n* is large, $\mu_0 \approx \sum_{j=1}^{n-1} \Phi^{-1}(W_j)/(n-1)$ varies between clusters according to a $N(0,\rho_3)$ distribution. This defines the range of possible cluster specific regression curves. Using equation (16), we construct the prediction curves for three copulas $C^{(2)}$ when the margins *F* and *G* are the standard normal distribution. The curves are presented in Figure 2. They correspond to Kendall correlation coefficient τ_2 equal to (0.3, 0.6, 0.9) for each copula $C^{(2)}$. Three copula families $C^{(2)}$ are used: Normal, Frank and Clayton copulas. The dependence parameters are deduced from τ_2 following Joe [13, pp. 58, 166 and 168] and using the function *iTau* of the **R**-package copula, see Ivan Kojadinovic and Jun Yan [12]. We also fix the size n = 21 and the Kendall tau of $C_{1,1:n}^{(3)}$ is $\tau_3 = 0.1$ corresponding to $\rho_3 = 0.16$. Prediction curves are plotted for the quantiles $q \in \{1/10, 5/10, 9/10\}$ of $\sum_{j=1}^{n-1} \Phi^{-1}(W_j)/(n-1)$ whose approximate distribution is a $N(0, \rho_3)$.



Figure 2: Curves of $\mathbb{E}(Y|x = x_n, (x_1, y_1), \dots, (x_{n-1}, y_{n-1}))$ with a normal copula (first line), Clayton copula (second line), and Frank copula (third line) for quantiles q = 1/10 (dotted line), q = 5/10 (full traint) and q = 9/10 (dotdash line) of the between cluster effects.

The graphs exemplify the impact of the dependency in $C^{(2)}$ on the prediction curves. Indeed, when the copula $C^{(2)}$ is a normal copula, we obtain, in Figure 2, parallel regression lines corresponding to the mixed linear regression model of [2]. For Clayton's copula, the regression curves are close for small values of *x*; this might be related to the lower tail dependency of that copula family. When the correlation τ_2 increases, the regression curves tend to a unique straight lines. This agrees with the conditional form of the model given in equations (9) and (12) where the random part of the model tends to 0 as ρ_2 increases to 1. In Figure 2 we used a relatively small value for the error dependence parameter in $C_{1,1:n}^{(3)}$; such a small residual dependency is found in many applications such as the analysis presented in Section 5.

4. Copula selection and parameter estimation for a 2-exchangeable model

The data set to analyze is $\{(X_{ij}, Y_{ij}) : i = 1, ..., m; j = 1, ..., n_i\}$, where index *i* is for clusters and *j* is for units within clusters. We assume that the joint distribution of the $2n_i$ variables in cluster *i* is determined by two marginal distributions $F(x|\alpha)$, $G(y|\beta)$ and by copula density (10) that involves a bivariate copula density, $c^{(2)}(u, v; \delta_2)$ for the marginal relationship between X and

Y and two 1-exchangeable copula families with densities, $c_{1,1;n}^{(1)}(u_1, \ldots, u_n; \delta_1)$ and $c_{1,1;n}^{(3)}(w_1, \ldots, w_n; \delta_3)$. The goal of this section is to identify parametric families for the five components of this model and to estimate their parameters that are put in vector $\theta = (\alpha, \beta, \delta_1, \delta_2, \delta_3)$. Following the model building procedure presented in chapter 5 of Joe [13] and chapters 7 and 8 of Czado [6]. The model components are selected sequentially, using techniques that are presented in this section.

4.1. Determination of the model components

The determination of the marginal distributions is done independently for the two margins. Competing models are compared on the basis of AIC criteria calculated as if the units were independent. Preliminary, or IFM, estimators see Joe [13, section 5.5], $(\tilde{\alpha}, \tilde{\beta})$ are obtained by maximizing

$$\mathcal{L}_F = \sum_{i=1}^m \sum_{j=1}^{n_i} \log\left\{f(x_{ij}|\alpha)\right\} \text{ and } \mathcal{L}_G = \sum_{i=1}^m \sum_{j=1}^{n_i} \log\left\{g(y_{ij}|\beta)\right\}$$

The next step is to calculate the pseudo observations, defined by

$$\tilde{u}_{ij} = F(x_{ij}|\tilde{\alpha}), \quad \tilde{v}_{ij} = G(y_{ij}|\tilde{\beta})$$
(18)

The $N = \sum n_i$ pairs $(\tilde{u}_{ij}, \tilde{v}_{ij})$ are then used to select a copula $C^{(2)}$ for the (X, Y) dependency. Note that the bivariate empirical distribution function of the pseudo-observations $(\tilde{u}_{ij}, \tilde{v}_{ij})$ is a consistent estimator of $C^{(2)}$ when the number of clusters m goes to infinity provided that the cluster sizes are bounded, $n_i < n_{max}$ for each i. The within cluster dependency impacts the variances of the estimators however this is overlooked at this stage and standard methods, proposed in Joe [13] and Czado [6], are used to select a copula family $C^{(2)}$ for the bivariate sample. Competing models are compared on the basis of their AIC and an IFM estimator $\tilde{\delta}_2$ of the parameter of the selected copula family is obtained by maximizing

$$\mathcal{L}_2 = \sum_{i=1}^m \sum_{j=1}^{n_i} \log \left[c^{(2)}(\tilde{u}_{ij}, \tilde{v}_{ij}; \delta_2) \right]$$

To assess the within cluster dependency associated with copula families $C_{1,1:n}^{(1)}$ and $C_{1,1:n}^{(3)}$, we use the unit level version of the exchangeable Kendall's tau introduced in Romdhani et al. [21]. It is evaluated using the proportion of concordant pairs $\{(x_{ij}, x_{i\ell}), (x_{kr}, x_{ks})\}$ among the $\sum_{i>k} n_i(n_i - 1)n_k(n_k - 1)$ possible pairs of ordered observations coming from different clusters. Graphical methods to select a family of exchangeable copulas are proposed in Rivest et al. [20]. Models can also be compared on the basis of their AIC, and an IFM estimator of δ_1 is obtained by maximizing

$$\mathcal{L}_{1} = \sum_{i=1}^{m} \log \left\{ c_{1,1:n_{i}}^{(1)} (\tilde{u}_{i1}, \ldots, \tilde{u}_{in_{i}}; \delta_{1}) \right\}.$$

The selection of the copula family $C_{1,1:n}^{(3)}$ is based on the pseudo observations $\tilde{w}_{ij} = C_{2|1}^{(2)}(\tilde{v}_{ij}|\tilde{u}_{ij};\tilde{\delta}_2)$. The exchangeable Kendall's tau evaluated on $\{\tilde{w}_{ij}: i = 1, ..., m; j = 1, ..., n_i\}$ can be used to assess the within cluster dependency of the residuals. IFM

estimator $\tilde{\delta}_3$ is obtained by maximizing

$$\mathcal{L}_{3} = \sum_{i=1}^{m} \log \left\{ c_{1,1:n_{i}}^{(3)}(\tilde{w}_{i1},\ldots,\tilde{w}_{in_{i}};\delta_{3}) \right\}$$

As note in Joe [13, section 5.5] the five estimating equations associated with the IFM estimation of θ can be combined in a multivariate estimating equation that yield the IFM estimator $\tilde{\theta}$. The standard asymptotic theory, presented in Tsiatis [24, pp. 30], applies. It shows that the joint asymptotic distribution of $\tilde{\theta} - \theta$, as the number of clusters *m* goes to infinity, is a centered multivariate normal distribution with a sandwich covariance matrix. As expected, this sandwich variance estimator accounts for the within cluster dependence.

4.2. Maximum likelihood estimation of the parameters

Once parametric families for the five model components have been identified, optimal estimators of the parameters are obtained by maximum likelihood. This section discusses the properties of the maximum likelihood estimator for the parameter vector θ .

The log-likelihood for θ is equal to

$$\begin{aligned} \mathcal{L}(\theta) &= \sum_{i=1}^{m} \sum_{j=1}^{n_{i}} \log\left\{f(x_{ij}|\alpha)\right\} + \sum_{i=1}^{m} \sum_{j=1}^{n_{i}} \log\left[c^{(2)}\left\{F(x_{ij}|\alpha), G(y_{ij}|\beta); \delta_{2}\right\}\right] \\ &+ \sum_{i=1}^{m} \sum_{j=1}^{n_{i}} \log\left\{g(y_{ij}|\beta)\right\} + \sum_{i=1}^{m} \log\left[c^{(1)}_{1,1:n_{i}}\left\{F(x_{i1}|\alpha), \dots, F(x_{in_{i}}|\alpha); \delta_{1}\right\}\right] \\ &+ \sum_{i=1}^{m} \log\left[c^{(3)}_{1,1:n_{i}}\left[C_{2|1}\left\{G(y_{i1}|\beta)|F(x_{i1}|\alpha)\right\}, \dots, C_{2|1}\left\{G(y_{in_{i}}|\beta)|F(x_{in_{i}}|\alpha)\right\}; \delta_{3}\right]\right] \end{aligned}$$

This function is easily maximized once parametric families for the two margins in the model and the three copulas are selected. This yields $\hat{\theta}$ the maximum likelihood estimator of the parameter vector. This maximization is carried out using an optimiser such the R-function *optim* or *nlminb*. Minus the hessian of $\mathcal{L}(\theta)$, evaluated at $\hat{\theta}$, is the observed Fisher information for the model. Its inverse is the asymptotic covariance matrix of $\hat{\theta} - \theta$. It can be used to calculate standard error estimates for all the parameters that have been estimated. At this stage, likelihood ratio tests comparing nested candidate parametric families for $C^{(2)}$ can also be carried out to validate the IFM model selection step that ignored the within cluster dependency.

When the margins F(x), G(y) and the copula $C^{(2)}$ are normal, the likelihood can be split into a marginal likelihood for the parameters of F(x) and $C_{1,1:n}^{(1)}$ times a conditional likelihood for the regression parameters ($\beta_0, \beta_1, \sigma_e^2$) and the parameter for $C_{1,1:n}^{(3)}$, see equation (12). The standard normal mixed linear model of [2] falls into that category as its parameters are estimated using a conditional likelihood. In general the parameters are intertwined in a complicated way and their estimation relies the log-likelihood $\mathcal{L}(\theta)$.

4.3. A Monte Carlo investigation of the sampling properties of IFM and ML estimators

For a 2-exchangeable copula model with parameters θ , $\hat{\theta}$ and $\tilde{\theta}$ are the maximum likelihood and the IFM estimators. This section investigates their sampling properties when *m* is finite. This is done using Monte-Carlo simulations where the expectation and the variance of an estimator $\hat{\psi}$ are approximated by

$$\mathbb{E}_{B}(\hat{\psi}) = \frac{1}{B} \sum_{b=1}^{B} \hat{\psi}_{b}, \quad \mathbb{V}_{B}(\hat{\psi}) = \frac{1}{B-1} \sum_{i=1}^{B} \left\{ \hat{\psi}_{b} - \mathbb{E}_{B}(\hat{\psi}) \right\}^{2},$$

where *b* indexes the estimates obtained in the B = 1000 Monte-Carlo simulations. The expectations and the variances of IFM estimators $\tilde{\psi}$ are evaluated in a similar way.

Throughout this Monte Carlo study of the copula 2-exchangeable model, the margins *F* and *G* are normal distributions with mean $\mu_1 = \mu_2 = 0$ and variance $\sigma_1^2 = \sigma_2^2 = 1$. The exchangeable copulas $C_{1,1:n}^{(1)}$ and $C_{1,1:n}^{(3)}$ belong to the normal family with correlation $\rho_1 = 0.31$ and $\rho_3 = 0.16$ respectively, corresponding to Kendall's tau of 0.2 and 0.1. Such small levels of within cluster association are often found in applications. In the study correlations are parameterized in terms of their logit,

$$\eta = \log\left(\frac{\rho}{1-\rho}\right). \tag{19}$$

Three copulas $C^{(2)}$ are investigated: the normal copula, the Clayton copula and a two-parameter Khoudraji copula [9] that features an asymmetric relationship between U and V. It is defined by

$$C^{(2)}(u, v; \rho, \kappa) = u^{1-\kappa} C_{\rho}(u^{\kappa}, v), \quad u, v \in [0, 1],$$
(20)

where C_{ρ} is a normal copula with correlation $\rho \in (0, 1)$ and $\kappa \in (0, 1)$ is the asymmetry parameter. In the simulations κ is parameterized in terms of its logit, see (19). Two values for the Kendall's tau τ_2 of copula $C^{(2)}$, 0.4 and 0.6, are considered. Two sample sizes, m = 10, 50, are investigated; their corresponding cluster sizes are n = 30 and n = 18 respectively. Tables 1 to 2 present the expectations and the variances of the estimators of the copula parameters.

$ au_2(\eta_2)$	т	Method	$\eta_1 = -0.80$	η_2	$\eta_3 = -1.69$
0.4(0.35)	10	MV	-0.96(1.90)	0.35(0.37)	-1.81(2.29)
	10	IFM	-1.00(2.20)	0.34(0.62)	-1.89(2.41)
	50	MV	-0.85(0.58)	0.35(0.12)	-1.74(0.75)
	50	IFM	-0.86(0.63)	0.35(0.16)	-1.76(0.80)
0.6(1.44)	10	MV	-0.90(1.97)	1.45(0.31)	-1.83(2.24)
	10	IFM	-0.95(2.30)	1.44(0.41)	-1.93(2.50)
	50	MV	-0.84(0.53)	1.44(0.09)	-1.73(0.70)
	50	IFM	-0.85(0.60)	1.44(0.11)	-1.75(0.76)

Table 1: Expectations of the estimators and their with variances multiplied by 10, in parenthesis, when $C^{(2)}$ is a normal copula.

Table 2: Expectations of the estimators and their with variances multiplied by 10, in parenthesis, when $C^{(2)}$ is Clayton copula.

$ au_2(\delta_2)$	т	Method	$\eta_1 = -0.80$	δ_2	$\eta_3 = -1.69$
0.4(1.33)	10	MV	-0.88(1.63)	1.35(0.66)	-1.82(2.36)
	10	IFM	-0.98(2.18)	1.33(0.84)	-1.91(2.68)
	50	MV	-0.83(0.46)	1.33(0.15)	-1.73(0.67)
	50	IFM	-0.85(0.60)	1.32(0.19)	-1.75(0.73)
0.6(3)	10	MV	-0.88(1.48)	3.02(2.39)	-1.80(2.48)
	10	IFM	-0.98(2.11)	2.96(3.13)	-1.91(2.56)
	50	MV	-0.82(0.45)	3.01(0.61)	-1.73(0.71)
	50	IFM	-0.86(0.59)	2.99(0.77)	-1.75(0.77)

$ au_2$	т	Method	$\eta_1 = -0.80$	$\eta_{ ho} = 0.75$	$\eta_{\kappa} = 1.52$	$\eta_3 = -1.69$
0.4	10	MV	-0.90(1.83)	0.80(0.55)	1.51(2.42)	-1.85(2.13)
	10	IFM	-0.95(2.08)	0.75(0.65)	1.49(3.25)	-1.89(2.49)
	50	MV	-0.82(0.51)	0.77(0.12)	1.51(0.98)	-1.73(0.63)
	50	IFM	-0.83(0.58)	0.76(0.18)	1.51(1.44)	-1.74(0.79)
$ au_2$	т	Method	$\eta_1 = -0.80$	$\eta_{ ho} = 1.45$	$\eta_{\kappa} = 3.48$	$\eta_3 = -1.69$
0.6	10	MV	-0.86(1.77)	1.49(0.36)	3.39(5.30)	-1.81(2.20)
	10	IFM	-0.94(2.09)	1.45(0.38)	3.62(6.34)	-1.88(2.74)
		MV	-0.81(0.49)	1.47(0.11)	3.48(2.14)	-1.77(0.73)
	50					

Table 3: Expectations of the estimators and their with variances multiplied by 10, in parenthesis, when $C^{(2)}$ is Khoudraji copula.

In Tables 1 to 3, all estimators have negligible biases. The discussion focuses on variances. As expected the strength of the U-V association in $C^{(2)}$ does not impact the precision of the two estimators for η_1 . The loss of precision for the IFM estimators is larger for the parameters of copula $C^{(2)}$ than for the parameters of the other 2 copula families. The efficiency of the maximum likelihood estimator is larger at m = 10 than at m = 50. The Supplementary Material provides additional simulations for unequal sample sizes $\{n_i\}$ within clusters. Unequal sample sizes are associated to a small loss of precision for all estimators. Overall the two estimation methods give similar results; this supports the proposal of Section 4 to use IFM estimators to select the components of the proposed copula model. The detailed simulation results, including a presentation of the sampling properties of the estimators for (μ_k, σ_k) for k = 1, 2, are presented in the Supplementary Material.

5. Modeling math grades with a 2-exchangeable copula model

This section revisits a data set discussed in [10]. It concerns math grades in fourth and seventh year measured on $N = \sum n_i =$ 728 students in m = 48 primary schools. The cluster sample sizes n_i vary between 4 and 40. The fourth year mark (X) and the seventh year mark (Y) vary between 0 and 40. We map them to the (0,1) interval using the transform $M \rightarrow (M + 1/2)/41$. To break ties in the grades a small random perturbation was added to each one.



Figure 3: Scatter plot of *Y* versus *X* with a smooth.

One objective of the analysis presented here is to construct predictive models for Y given X whose support is (0, 1). Another objective is to investigate whether the 2-exchangeable copula model can capture the non-linearity seen in Figure 3 that gives a scatter plot of the N = 728 data points and a smooth. The goal is to contrast an analysis carried out with copulas to the one reported in [10] that is based on standard normal linear mixed models.

5.1. Selection of the marginal distributions for X and Y

The candidates families for *F* and *G* are the beta (denoted \mathcal{B}) and the generalized beta (denoted $\mathcal{GB3}$) distributions, see [4]. Indeed, by the construction of histogram of the distribution, we arrive at an asymmetric law. If *X* has a $\mathcal{B}(\alpha, \beta)$ distribution then $Y = X/\{\lambda + (1 - \lambda)X\}$ has for $\lambda \in (0, 1)$ a $\mathcal{GB3}(\alpha, \beta, \lambda)$ distribution whose density is given by

$$\frac{\lambda^{\alpha}\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)}\frac{y^{\alpha-1}(1-y)^{\beta-1}}{\{1-(1-\lambda)y\}^{\alpha+\beta}}, \quad 0 \le y \le 1.$$

Table 4 compares the fit of these two distributions to the two margins. As stated in Section 4.1, this preliminary analysis does not account for the within cluster dependency. Thus the *pse*, for pseudo standard error, ignores the within classroom dependency.

Variable	Model	$ ilde{ heta}$	pse	AIC
X	${\mathcal B}$	(4.280,2.332)	(0.222,0.115)	-542.88
	GB3	(4.280,2.330,1)	(0.222,0.115,NA)	-540.80
Y	${\mathcal B}$	5.24,1.79	(0.28,0.08)	-789.57
	GB3	(2.616,2.319,0.29)	(0.303,0.241,0.069)	-826.96

Table 4: Fit of the \mathcal{B} and of the \mathcal{GB} 3 distributions to the two margins.

The best fitting models are respectively the beta end the generalized beta for X and Y.

5.2. Selection of the bivariate copula $C^{(2)}$

The fist step is to calculate the pseudo observations \tilde{u}_{ij} et \tilde{v}_{ij} for $j = 1, ..., n_i$ and i = 1, ..., m defined in (18). The Kendall's tau is 0.49 (*pse* = 0.03) so there is a relatively strong association between the two variables. Following [13, chap. 1], Kendall's tau is calculated for the sub-samples in the 4 quadrants of the unit square. This reveals a stronger association for large grades than for smaller ones. Also a 0.1 difference between the Kendall's tau for the upper left and lower quadrant suggests that some of the asymmetry seen in the Figure 3 is left once the margins' effect has been factored out.

Several copulas were fitted to this bivariate sample using the functions BiCopEst in the R package VineCopula and fitCopula in copula. To capture the asymmetry in the data we used to Khoudraji device, see (20) to create asymmetric alternatives. The best fitting copula in Table 5 is the survival Khoudraji normal copula given by

$$C^{(2)}(u, v | \rho_2, \kappa_1, \kappa_2) = u + v - 1 + (1 - u)^{1 - \kappa_1} (1 - v)^{1 - \kappa_2} C_{\rho_2} \{ (1 - u)^{\kappa_1}, (1 - v)^{\kappa_2} \},\$$

where C_{ρ_2} is the bivariate normal copula with correlation ρ_2 . The density of a copula in this three parameter family can be evaluated using functions of copula. The conditional distribution, $w = \partial C^{(2)}(u, v|\rho_2, \kappa_1, \kappa_2)/\partial u$ has the following explicit form.

$$w = 1 - (1 - \kappa_1) (1 - u)^{-\kappa_1} (1 - v)^{1 - \kappa_2} C_{\rho_2} \{ (1 - u)^{\kappa_1}, (1 - v)^{\kappa_2} \} - \kappa_1 (1 - v)^{1 - \kappa_2} \Phi \left[\frac{\Phi^{-1} \{ (1 - v)^{\kappa_2} \} - \rho_2 \Phi^{-1} \{ (1 - u)^{\kappa_1} \}}{\sqrt{1 - \rho_2^2}} \right] \cdot (21)$$

In Table 5, Survival Khoudraji-Normal2 refers to a two parameter version of this copula obtained by setting $\kappa_2 = 1$.

Copula <i>C</i> ⁽²⁾	$ ilde{ heta}$	pse	AIC
Normal	0.682	0.016	-454.32
Survival-Gumbel	1.892	0.057	-459.5
Survival Khoudraji-Normal	(0.791, 0.822, 0.960)	(0.023, 0.043, 0.029)	-474.10
Survival Khoudraji-Normal2	(0.762, 0.837)	(0.021, 0.045)	-468.64

Table 5: Paramater estimates, their pseudo standard errors *pse* and the AIC for several copulas for the (u, v) relationship.

5.3. Selection of the exchangeable copula families $C_{1,1:n}^{(1)}$ and $C_{1,1:n}^{(3)}$

The exchangeable Kendall's tau for X and Y are respectively 0.046 (se = .017) and 0.095 (se = 0.035) showing a stronger school effect in the seventh year. The fit of several families of copulas for the joint distribution of { \tilde{u}_{ij} } and { \tilde{w}_{ij} }, evaluated using (21), are compared by maximizing the pseudo log-likelihoods \mathcal{L}_1 and \mathcal{L}_3 . The results are reported in Table 6. The normal family is the best choice for both $C_{1,1:n}^{(1)}$ and $C_{1,1:n}^{(3)}$.

Table 6: Parameter estimates, their pseudo standard errors *pse*, and the AIC for three families for $C_{1,1:n}^{(1)}$ and $C_{1,1:n}^{(3)}$.

		$C_{1,1:n}^{(1)}$			$C_{1,1:n}^{(3)}$	
Family	$ ilde{ heta}$	pse	AIC	$ ilde{ heta}$	pse	AIC
Frank	0.251	0.138	-5.69	0.935	0.179	-69.76
Gumbel	1.040	0.022	-4.64	1.109	0.024	-66017
Normal	0.064	0.025	-12.76	0.167	0.035	-80.03

To complete the analysis we carry out a full maximum likelihood estimation of the 10 parameters for the components of the proposed model. The parameter estimate for κ_2 is very close to 1. We first carry out a likelihood ratio test for H_0 : $\kappa_2 = 1$ using the full likelihood. This gives $\chi^2_{1,obs} = 8.2$ for a p-value of 0.4%. Thus the full model, with 10 parameters, is definitive ; the estimates and standard errors (*se*) for the parameters are reported in Table 7.

Component	$\hat{ heta}$	se
$F\left(\mathcal{B} ight)$	$(\hat{\alpha}_1, \hat{\beta}_1) = (4.271, 2.359)$	(0.235, 0.124)
$G(\mathcal{GB}3)$	$(\hat{\alpha}_2,\hat{\beta}_2,\hat{\lambda})=(2.457,2.470,0.248)$	(0.245, 0.254, 0.052)
$C_{1,1:n}^{(1)}$ (Normal)	$\hat{\rho}_1 = 0.063$	0.026
$C^{(2)}$ (Survival Khoudraji-Normal)	$(\hat{\rho},\hat{\kappa}_1,\hat{\kappa}_2)=(0.795,0.822,0.959)$	(0.024, 0.046, 0.029)
$C_{1,1:n}^{(3)}$ (Normal)	$\hat{\rho}_3 = 0.161$	0.040

Table 7: Maximum likelihood estimators of the parameters for the full model.

The fit of the final model summarized in Table 7 is illustrated using two schools, numbered 1, with $n_1 = 19$, and 30, with $n_{30} = 31$. Their respective values of μ_0 , see (15), are -0.464 and 0.810; this means that, for the same math4 mark, the math7 grade in School 30 are higher than in School 1. This can be seen in Figure 4 that gives the two regression curves, constructed using (16), that use the Gauss-Hermite quadrature method to approximate the normal integrals for each *x*-value.



Figure 4: Scatter plots and regression curves for Schools 1 (+ character and full line) and 30 (× character and dashed line).

Conditional residuals for the fitted copula models can be defined as $y_{ij} - \hat{y}_{ij}$, where \hat{y}_{ij} is evaluated as the predicted value at x_{ij} , using (16), where the values of μ_0 and σ_0 , see (15), are those for school *i*. The conditional residuals of the copula model (*CM*) can be compared to those of mixed linear models. The first one, *ML*1, has a random school intercept while the second one, *ML*2, has possibly dependent random slope and intercept. The mean squared errors and inter quartile ranges (IQR) of the residuals for *CM*, *ML*1, and *ML*2, are (0.0107, 0.0112, 0.0101) and (0.1082, 0.1186, 0.1098) respectively. Thus, in agreement with the analysis

reported in Table 5, the fit of ML1 is poor. The fits of CM and ML2 are very similar as the latter captures the between school change in slope than can be seen in Figure 4. CM has a larger residual MSE; however it has a smaller residual IQR and it gives a smaller absolute residual than ML2 for 53% of the data points.



Figure 5: Densities for the predicted Y values for X = 0.4 (continuous lines) and X = 0.9 (dashed lines) for Schools 1, in black, and 30, in red.

The key advantage of the 2-exchangeable copula model is that it allows prediction intervals for *Y* to depend on both, the School and the *X*-value. This is illustrated in Figure 5 that gives the predictive densities for mark *Y*, given by formula(17), for X = 0.4 and X = 0.9 in Schools 1 and 30. The variability of *Y* is larger at X = 0.4 and in School 1. This can be seen by looking at the corresponding 95% prediction intervals for *Y* that are given in Table 8.

School	X = 0.4	X = 0.9
1	(0.23, 0.89)	(0.66,0.97)
30	(0.44,0.97)	(0.86,0.99)

Table 8: 95% prediction intervals for X = 0.4 and X = 0.9 in Schools 1 and 30.

6. Conclusion

The 2-exchangeable copula model proposed in this work provides flexible methods to predict variable Y knowning X in a hierarchical data set. A key feature of the proposed methodology highlighted in Section 4 is the flexibility of the predictive densities for Y given X. Its shape and its support can depend on the known X value for Y and on the cluster. An outstanding problem is whether the proposed model can be generalized to two or more continuous explanatory variables. The key to such a generalization is the availability of flexible (d - 1)-exchangeable families $C_{d-1,1:n}^{(1)}$ for the joint distribution of the explanatory variables for all the unit in a cluster. An elliptical copula, with a correlation matrix given by (3) could be used. This specifies partially the copula $C^{(2)}$ for the joint distribution of the X and the Y variables on a unit. A vine decomposition could possibly be used to complete the specification of $C^{(2)}$. These problems will be the object of future investigations.

Supplementary Material

The supplementary material contains the proof of Proposition and additional results of the Monte Carlo simulation study.

Acknowledgments and Miscellaneous

We would like to thank Étienne Marceau for his critical reading of a previous version of this work.

Funding

The support of the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.

Disclosure Statement

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Data availability

The data is available upon request.

References

- [1] E. F. Acar, P. Azimaee, M. E. Hoque, Predictive assessment of copula models, Canadian Journal of Statistics, 47 (2019) 8–26.
- [2] G. E. Battese, R. M. Harter, W. A. Fuller, An error-components model for prediction of county crop areas using survey and satellite data, *Journal of the American Statistical Association*, 83 (1988) 28–36.
- [3] C. Bernard, C. Czado, Conditional quantiles and tail dependence, Journal of Multivariate Analysis, 138 (2015) 104–126.
- [4] W. M. Cockriel, J. B. McDonald, Two multivariate generalized beta families, *Communications in Statistics Theory and Methods*, 47 (2018) 5688–5701.

- [5] G. J. Crane, J. Van der Hoek, Conditional expectation formulae for copulas, *Australian and New Zealand Journal of Statistics* 50 (2008) 53–67.
- [6] C. Czado, Analyzing dependent data with vine copulas : a practical guide with R, 2019.
- [7] C. Czado, T. Nagler, Vine copula based modeling, Annual Review of Statistics and Its Application, 9 (2022) 453-477.
- [8] J. Dissmann, E. C. Brechmann, C. Czado, D. Kurowicka, Selecting and estimating regular vine copulae and application to financial returns, *Computational Statistics and Data Analysis*, 59 (2013) 52–69.
- [9] C. Genest, K. Ghoudi, L.-P. Rivest, Discussion of "understanding relationships using copulas, by E. Frees and E. Valdez", North American Actuarial Journal, 3 (1998) 543–552.
- [10] H. Goldstein, Multilevel Statistical Models, Wiley, 2nd edition, 2011.
- [11] K. Grover, E. F. Acar, M. Torabi, Copula-based predictions in small area estimation, Canadian Journal of Statistics, 48 (2020) 685-711.
- [12] Ivan Kojadinovic, Jun Yan, Modeling multivariate distributions with continuous margins using the copula R package, *Journal of Statistical Software*, 34 (2010) 1–20.
- [13] H. Joe, Dependence modelling with copulas, Chapman and Hall, 2014.
- [14] P. Kumar, M. Shoukri, Copula based prediction models: an application to an aortic regurgitation study, *BMC Medical Research Methodology*, 7 (2007) 1–9.
- [15] J. F. Mai, M. Scherer, Simulating copulas: stochastic models, sampling algorithms, and applications, Quantitative Finance, 2012.
- [16] C. E. McCulloch, S. R. Searle, Generalized, linear, and mixed models, John Wiley et Sons, 2001.
- [17] R. B. Nelsen, A introduction to copulas, Springer, 2006.
- [18] H. Noh, A. E. Ghouch, T. Bouezmarni, Copula-based regression estimation and inference, *Journal of the American Statistical Association*, 108 (2013) 676–688.
- [19] A. Panagiotelis, C. Czado, H. Joe, Pair copula constructions for multivariate discrete data, *Journal of the American Statistical Association*, 107 (2012) 1063–1072.
- [20] L. Rivest, F. Verret, S. Baillargeon, Unit level small area estimation with copulas, The Canadian Journal of Statistics/ La revue Canadienne de statistique, 44 (2016) 397–415.
- [21] H. Romdhani, L. Lakhal-Chaieb, L.-P. Rivest, An exchangeable kendall's tau for clustered data, *Canadian Journal of Statistics*, 42 (2014) 384–403.
- [22] L. A. Stefanski, D. D. Boos, The calculus of m-estimation, The American Statistician, 56 (2002) 29-38.
- [23] C.-L. Su, J. G. Neslehova, W. Wang, Modelling hierarchical clustered censored data with the hierarchical kendall copula, *Canadian Journal of Statistics*, 47 (2019) 182–203.
- [24] A. A. Tsiatis, Semiparametric theory and missing data, Springer, 2006.
- [25] G. Verbeke, G. Molenberghs, Linear mixed models for longitudinal data, Springer, 2000.