

SPARSE SAMPLING AND DILATION OPERATIONS ON A GIBBS WEIGHTED TREE, AND MULTIFRACTAL FORMALISM

JULIEN BARRAL AND STÉPHANE SEURET

ABSTRACT. In this article, starting from a Gibbs capacity, we build a new random capacity by applying two simple operators, the first one introducing some redundancy and the second one performing a random sampling. Depending on the values of the two parameters ruling the redundancy and the sampling, the new capacity has very different multifractal behaviors. In particular, the multifractal spectrum of the capacity may contain two to four phase transitions, and the multifractal formalism may hold only on a strict subset (sometimes, reduced to a single point) of the spectrum's domain.

1. INTRODUCTION

Multifractal analysis of capacities defined on \mathbb{R}^d ($d \geq 1$) is a natural generalisation of multifractal analysis of Radon measures. Our purpose in this article is to perform the multifractal analysis of a family of random capacities parametrised by two indices ϱ and η , ϱ being a redundancy index and η a lacunarity index. These new families of multifractal objects are easy to build, and have a versatile multifractal structure, exhibiting many and diverse phase transitions.

Let us now be more specific. We consider the set $\text{Cap}([0, 1]^d)$ of capacities on $[0, 1]^d$, i.e. non-decreasing non-negative mappings $\mu : \mathcal{B}([0, 1]^d) \rightarrow \mathbb{R}^+$, where $\mathcal{B}([0, 1]^d)$ stands for the Borel subsets of $[0, 1]^d$. One way to produce such capacities is, starting from a sequence $c = (c_I)_{I \in \mathcal{D}}$ of complex numbers indexed by the set \mathcal{D} of dyadic subcubes of $[0, 1]^d$, to impose some hierarchical structure to c as follows.

Definition 1.1. *For every $c = (c_I)_{I \in \mathcal{D}}$, the capacity $M(c) \in \text{Cap}([0, 1]^d)$ is defined by*

$$M(c) : E \in \mathcal{B}([0, 1]^d) \mapsto \sup\{|c_J| : J \in \mathcal{D}, J \subset E\}.$$

For instance, when $c = (c_I)_{I \in \mathcal{D}}$ is the collection of wavelet coefficients of a Hölder continuous function $f : [0, 1] \rightarrow \mathbb{R}$ (see Section 2.4 for details), the sequence $(M(c)(3I))_{I \in \mathcal{D}}$ where $3I$ is the union of the cube I and all of its neighboring cubes of same generation, coincides with the so-called wavelet leaders of f . Then, performing the multifractal analysis of the function f amounts to perform the multifractal analysis of the capacity $(M(c)(3I))_{I \in \mathcal{D}}$. Actually, the model of multifractal capacities considered in this paper, of which the model studied by the authors in [5] is a particular case, was motivated by the lacunary wavelet series introduced by Jaffard in [13], see Section 2.4.

In [5], we defined the following random sparse sampling on capacities. A probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is fixed for the rest of the paper. For $j \in \mathbb{N}$, denote by \mathcal{D}_j the set of dyadic cubes of generation j included in $[0, 1]^d$. We denote by $g(I)$ is the generation of a dyadic cube $I \subset [0, 1]^d$, i.e. $g(I) = j$ whenever $I \in \mathcal{D}_j$. By definition, $\mathcal{D} = \bigcup_{j \geq 0} \mathcal{D}_j$.

Definition 1.2. *Let $\eta \in (0, 1)$ and let $(p_I)_{I \in \mathcal{D}}$ be a family of independent random variables such that each p_I is a Bernoulli variable of parameter $2^{-g(I)d(1-\eta)}$. For every $\mu \in \text{Cap}([0, 1]^d)$, one introduces the sequence $(S_\eta \mu)$ defined as*

$$(S_\eta \mu) = (S_\eta \mu(I))_{I \in \mathcal{D}} = (p_I \mu(I))_{I \in \mathcal{D}}.$$

Then, the random capacity $M_\eta(\mu)$ is defined as

$$M_\eta(\mu) := M(S_\eta\mu).$$

To get $M_\eta(\mu)$, one first applies a random sparse sampling to the sequence $(S_\eta\mu)$. Observe that as $j \rightarrow \infty$, this sampling operation keeps asymptotically about $2^{jd\eta}$ among the 2^{jd} values $\mu(I)$ for $I \in \mathcal{D}_j$, and it puts the other values to 0. Then, from this sparse sequence one gets $M_\eta(\mu)$ by re-imposing a hierarchical structure via the operator M defined above.

When $\mu(E) = \lambda_\gamma(E) := (\mathcal{L}^d(E))^\gamma$, where \mathcal{L}^d stands for the Lebesgue measure and $\gamma > 0$, the capacity $M_\eta(\lambda_\gamma)$ is very similar to the wavelet leaders associated to the lacunary wavelet series considered by Jaffard in [13]. While λ_γ is essentially the simplest monofractal object (in a sense to be precised below) fully supported over $[0, 1]^d$, the capacity $M_\eta(\lambda_\gamma)$ is a rich multifractal object. In particular, it obeys the multifractal formalism in the following sense.

The topological support of $\mu \in \text{Cap}([0, 1]^d)$ is defined as

$$\text{supp}(\mu) = \{x \in [0, 1]^d : \mu(B(x, r)) > 0, \forall r > 0\}.$$

If $\text{supp}(\mu) \neq \emptyset$, set

$$(1) \quad \tau_\mu : q \in \mathbb{R} \mapsto \liminf_{j \rightarrow \infty} \tau_{\mu, j}(q), \quad \text{where } \tau_{\mu, j}(q) := \frac{-1}{j} \log_2 \sum_{I \in \mathcal{D}_j, \mu(I) > 0} \mu(3I)^q.$$

An equivalent definition is

$$\tau_\mu(q) = \liminf_{r \rightarrow 0^+} \frac{\log \sup \sum_i \mu(B(x_i, r))^q}{\log(r)}$$

where the supremum is taken over all families of disjoint closed balls $\{B(x_i, r)\}_i$ of radii r with centers $x_i \in \text{supp}(\mu)$, and $\mu(B(x, r)) := \mu(B(x, r) \cap [0, 1]^d)$. This function is concave and non-decreasing, and it takes values in $\mathbb{R} \cup \{-\infty\}$.

The *singularity*, or *multifractal spectrum* of μ is then defined as

$$\sigma_\mu : H \mapsto \dim \underline{E}_\mu(H), \quad H \in \mathbb{R},$$

where

$$\underline{E}_\mu(H) = \left\{ x \in \text{supp}(\mu) : \liminf_{r \rightarrow 0^+} \frac{\log(\mu(B(x, r)))}{\log(r)} = H \right\},$$

and where \dim stands for the Hausdorff dimension, and by convention, $\dim E = -\infty$ if and only if E is empty. The singularity spectrum provides a fine geometric description of μ at small scales by giving the Hausdorff dimension of the iso-Hölder sets $\underline{E}_\mu(H)$. Note that since μ is a capacity, the set $\underline{E}_\mu(H)$ is empty when $H < 0$. It is always true (see [7, 19, 22, 16] for measures and [17] for capacities), that

$$(2) \quad \sigma_\mu(H) \leq \tau_\mu^*(H), \quad \forall H \in \mathbb{R},$$

where for any function $f : \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty\}$, the Legendre transform f^* of f is defined as

$$f^* : H \in \mathbb{R} \mapsto \inf\{Hq - \tau_\mu(q) : q \in \mathbb{R}\}.$$

The domain of such a function f , that is the set of points where it takes real values, will be denoted by $\text{dom}(f)$.

The multifractal formalism holds for μ at H when the inequality (2) turns out to be an equality. It holds for μ when (2) is an equality for all $H \in \mathbb{R}$.

A capacity μ is monofractal when τ_μ is linear or more generally if $\underline{E}_\mu(H_0) = \text{supp}(\mu)$ for some $H_0 \in \mathbb{R}_+$, and μ is called multifractal when there are at least two distinct non-negative exponents H and H' such that $\underline{E}_\mu(H) \neq \emptyset \neq \underline{E}_\mu(H')$.


Coming back to the capacity $\lambda_\gamma(E) = (\mathcal{L}^d(E))^\gamma$, for $\gamma > 0$, it is clear that

$$\sigma_{\lambda_\gamma}(H) = \begin{cases} d & \text{if } H = d\gamma \\ -\infty & \text{otherwise} \end{cases} \quad \text{and} \quad \tau_{\lambda_\gamma}(q) = d(\gamma q - 1), q \in \mathbb{R}.$$

Consequently, λ_γ is monofractal and it obeys the multifractal formalism. The situation is more involved for $M_\eta(\lambda_\gamma)$: it is shown in [5] (see also [13] for a slightly different model) that almost surely,

$$\sigma_{M_\eta(\lambda_\gamma)}(H) = \begin{cases} \frac{\eta}{\gamma}H & \text{if } H \in [d\gamma, d\gamma/\eta] \\ -\infty & \text{otherwise,} \end{cases} \quad \text{and} \quad \tau_{M_\eta(\lambda_\gamma)}(q) = \begin{cases} \eta^{-1}d(\gamma q - \eta) & \text{if } q < 1/\gamma \\ d(\gamma q - \eta) & \text{if } q \geq 1/\gamma \end{cases}$$

Thus, $M_\eta(\lambda_\gamma)$ is multifractal, and it also obeys the multifractal formalism.



Let us now introduce some redundancy in the constructions.

Definition 1.3. For $\varrho \in (0, 1]$, $j \in \mathbb{N}$ and $I \in \mathcal{D}_j$, let I^ϱ denote the unique dyadic cube in $\mathcal{D}_{[ej]}$ of sidelength $2^{-[ej]}$ containing I .

The ϱ -dilation operation on $\text{Cap}([0, 1]^d)$ is defined by

$$D_\varrho\mu = M(c_\varrho(\mu)), \quad \text{where } c_\varrho(\mu) = (\mu(I^\varrho))_{I \in \mathcal{D}}.$$

Note that D_1 is the identity map.

When $\varrho < 1$, some redundancy is introduced in $D_\varrho\mu$, since each $I \in \mathcal{D}_j$ shares the same cube I^ϱ with the $2^{j-[ej]}$ dyadic subcubes of I^ϱ of generation j . In addition, observe that especially if the measure μ is doubling, the measures $\mu(I^\varrho)$ and $\mu((I')^\varrho)$ for two neighbor cubes I and I' are very comparable.

Remark that the family $(\lambda_\gamma)_{\gamma > 0}$ possesses the rather exceptional property that $D_\varrho\lambda_\gamma$ still obeys the multifractal formalism. Indeed, $D_\varrho\lambda_\gamma \approx \lambda_{\varrho\gamma}$, where for two capacities μ and μ' , $\mu \approx \mu'$ means that there exists $C \geq 1$ such that $C^{-1}\mu(I) \leq \mu'(I) \leq C\mu(I)$ for all $I \in \mathcal{D}$.

If $\mu \in \text{Cap}([0, 1]^d)$ and $\text{supp}(\mu) \neq \emptyset$, one checks that

$$\sigma_{D_\varrho\mu}(H) = \sigma_\mu(H/\varrho), \forall H \in \mathbb{R}$$

and that if $\text{supp}(\mu) = [0, 1]^d$,

$$\tau_{D_\varrho\mu}(q) = \varrho\tau_\mu(q) + d(\varrho - 1), \forall q \in \mathbb{R},$$

whose Legendre transform equals $\varrho\tau_\mu^*(\cdot/\varrho) + d(1 - \varrho)$. In particular, if μ obeys the multifractal formalism, then $\sigma_{D_\varrho\mu} = \tau_\mu^*(\cdot/\varrho)$, so that $D_\varrho\mu$ obeys the multifractal formalism if and only if $\tau_\mu^*(\cdot/\varrho) = \tau_{D_\varrho\mu}^* = \varrho\tau_\mu^*(\cdot/\varrho) + d(1 - \varrho)$, i.e. τ_μ^* takes its values in $\{d, -\infty\}$ if $\varrho < 1$. Assuming that $\tau_\mu'(0)$ exists, and setting $\gamma = \tau_\mu'(0)/d$, this equality implies that necessarily $\tau_\mu = \tau_{\lambda_\gamma}$ and $\sigma_\mu = \sigma_{\lambda_\gamma}$. So, as soon as μ is multifractal and τ_μ is differentiable at 0, the capacity $D_\varrho\mu$ for $\varrho < 1$ cannot obey the multifractal formalism when μ does.

Let us introduce a notation (that we already used to define $\lambda_\gamma = (\mathcal{L}_d)^\gamma$): for any $\gamma > 0$ and $\mu \in \text{Cap}([0, 1]^d)$, the capacity μ^γ is defined by

$$(3) \quad \mu^\gamma : E \in \mathcal{B}([0, 1]^d) \mapsto \mu(E)^\gamma.$$

Using the previous observations, one sees that for any $\mu \in \text{Cap}([0, 1]^d)$, $\gamma > 0$ and $\varrho \in (0, 1]$, that $(D_\varrho(\mu))^\gamma = D_\varrho(\mu^\gamma)$, and

$$\sigma_{D_\varrho(\mu^\gamma)} = \sigma_\mu(\cdot/\gamma\varrho), \quad \tau_{D_\varrho\mu}(q) = \varrho\tau_\mu(\gamma\cdot) + d(\varrho - 1), \quad \text{and} \quad \sigma_{D_\varrho(\mu)} = \sigma_{(\mu^\varrho)}.$$

It is clear that μ^γ satisfies the multifractal formalism as soon as μ does. Also, if $\mu \in \text{Cap}([0, 1]^d)$ satisfies the three properties (i) μ obeys the multifractal formalism, (ii) μ is multifractal, and (iii) τ_μ is differentiable at 0, then the capacities $D_\varrho(\mu^{1/\varrho})$, $0 < \varrho < 1$, share the same singularity spectrum as μ but none of them obeys the multifractal formalism.

In this article, we are interested in combining D_ϱ and M_η .

Definition 1.4. For every $\mu \in \text{Cap}([0, 1]^d)$, for every $\eta \in (0, 1]$ and $0 < \varrho \leq 1/\eta$, the capacity $M_{\varrho, \eta}(\mu)$ is defined by

$$M_{\varrho, \eta}(\mu) = M_\eta(D_{\varrho\eta}\mu) = M(S_\eta(D_{\varrho\eta}\mu)).$$

The two operators are somehow complementary: on one side, $D_{\varrho\eta}$ creates redundancy since $\varrho\eta$ is always less than 1, while on the other side M_η introduces lacunarity.

Our goal is to study the multifractal nature of the elements of $\{M_{\varrho, \eta}(\mu) : 0 < \varrho \leq 1/\eta\}$ when μ is a *Gibbs capacity*, defined as follows.

Let $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$ be \mathbb{Z}^d -invariant Hölder continuous potential. It is standard that the sequence of Radon measures

$$\nu_n(dx) = \frac{\exp(S_n\varphi(x))}{\int_{[0, 1]^d} \exp(S_n\varphi(t)) \mathcal{L}^d(dt)} \mathcal{L}^d(dx), \quad \text{where } S_n\varphi(x) = \sum_{k=0}^{n-1} \varphi(2^k x),$$

converges vaguely to a measure fully supported on \mathbb{R}^d (see [6]). Denote by ν the restriction of this *Gibbs measure* to $[0, 1]^d$.

Definition 1.5. A *Gibbs capacity* μ is a capacity $\mu \in \text{Cap}([0, 1]^d)$ such that $\mu \approx \nu^\gamma$, where $\gamma > 0$ and ν is such a *Gibbs measure* on $[0, 1]^d$ as above.

Classical results on Gibbs measures (see [23, 8, 19]) give :

- One has

$$\tau_\nu(q) = (qP(\varphi) - P(q\varphi))/\log(2), \quad q \in \mathbb{R}$$

where $P(q\varphi) = \lim_{n \rightarrow +\infty} \frac{1}{n} \log \int_{[0, 1]^d} 2^n \exp(S_n(q\varphi)(x)) \mathcal{L}^d(dx) \in \mathbb{R}$ is the topological pressure of $q\varphi$. In particular $\tau_\nu = \lim_{j \rightarrow \infty} \tau_{\nu, j}$.

- τ_ν is analytic, concave increasing and $\lim_{q \rightarrow +\infty} \tau_\nu(q) = +\infty$.
- ν satisfies the multifractal formalism: $\sigma_\nu = \tau_\nu^*$.

Hence, the same properties are true for any Gibbs capacity $\mu \approx \nu^\gamma$ with $\gamma > 0$.

The family of Gibbs capacities is obviously invariant under the operator $\mu \mapsto \mu^\gamma$. It is known that a Gibbs capacity $\mu \approx \nu^\gamma$ is multifractal if and only if the Gibbs measure ν is not equivalent to the Lebesgue measure, that is when φ is not cohomologous to a constant on $\mathbb{R}^d/\mathbb{Z}^d$ endowed with the dynamics $\times 2$ (see [6]).

The situation studied in [5]¹ corresponds to the extreme case $\varrho = 1/\eta$, that is $M_{\varrho, \eta}(\mu) = M(S_\eta(D_1\mu)) = M_\eta(\mu)$. There, we proved that $M_\eta(\mu)$ still satisfies the multifractal formalism. In the present paper, we complete the previous results for a Gibbs capacity μ as follows:

¹In [5] we considered capacities of the form $\mu(I) = K\nu(I)^\alpha 2^{-\beta j}$ for $I \in \mathcal{D}_j$, with $K > 0$, $\alpha \geq 0$, $\beta \geq 0$ and $\alpha + \beta > 0$, which is essentially equivalent to assuming $\mu \approx \nu^\gamma$ upon modifying φ .

- when $\varrho \in [1, 1/\eta)$, $M_{\varrho,\eta}(\mu)$ obeys the multifractal formalism (although $D_{\varrho\eta}(\mu)$ does not when μ is multifractal) - see Theorem 2.5 ;
- when $0 < \varrho < 1$ and μ is multifractal, $M_{\varrho,\eta}$ obeys the multifractal formalism only on a strict subset of $\text{dom}(\sigma_{M_{\varrho,\eta}})$ - see Theorems 2.1 and 2.4.

As a consequence of our study, even if all the capacities $D_{\varrho}(\mu^{1/\varrho})$, for $0 < \varrho < 1$, share the same singularity spectrum (by the above observations), the capacities $M_{\eta}(D_{\varrho}(\mu^{1/\varrho}))$, $0 < \varrho < 1$, have distinct singularity spectra, and obey the multifractal formalism if and only if $\varrho \in [\eta, 1]$.

In this article, we will focus on the cases $\varrho = 1$ and $0 < \varrho < 1$, since these are the situations where new interesting phenomena occur. We only give a statement for $1 < \varrho \leq 1/\eta$ since the result and its proof can be obtained by adapting the proofs of the case $\varrho = 1/\eta$ considered in [5].

Before stating precisely our results, it is worth saying a word about the properties of the L^q and singularity spectra of $M_{\varrho,\eta}(\mu)$ when μ is Gibbs. While τ_{μ} is analytic, it will be shown that for any $\varrho \in (0, 1/\eta]$, $\tau_{M_{\varrho,\eta}(\mu)}$ always presents a first order phase transition (i.e. a point of non differentiability), and in some cases even a second order phase transition (point of differentiability but not twice differentiability). Moreover, $\tau_{M_{\varrho,\eta}(\mu)}$ is analytic outside its phase transition(s) point(s). Also, while σ_{μ} is analytic over the interior of its domain when $\varrho \in (0, 1/\eta]$, this is not the case for $\sigma_{M_{\varrho,\eta}(\mu)}$. Indeed, $\sigma_{M_{\varrho,\eta}(\mu)}$ is still concave but may contain three to five consecutive analytic parts, and can be non differentiable at some point.

2. STATEMENT OF THE MAIN RESULTS

A multifractal Gibbs capacity μ is fixed, as well as $\eta \in (0, 1)$. Set $H_{\min} = \tau'_{\mu}(+\infty)$ and $H_{\max} = \tau'_{\mu}(-\infty)$. We start with the case $\varrho = 1$, which is both the simplest one and a transition between the case $\varrho \in (1, 1/\eta]$, for which the multifractal behavior of $M_{\varrho,\eta}(\mu)$ is similar to that in the case $\varrho = 1/\eta$ studied in [5] and the multifractal formalism holds, and the case $\varrho \in (0, 1)$, which is more versatile and for which the multifractal formalism holds only partially.

2.1. The case $\varrho = 1$. This situation corresponds to the case where there is an almost compensation between the redundancy process and the random sampling. Indeed, with this choice of parameters, every $\mu(I)$ with $I \in \mathcal{D}_j$ is replaced by $\mu(I^{\eta})$, but only $\sim 2^{j d \eta}$ of them are kept. Since the surviving cubes are chosen uniformly in $[0, 1]^d$, intuitively the surviving cubes I after sampling are such that the union of the corresponding cubes I^{η} cover $[0, 1]^d$ (this is not true, but not far from it).

Theorem 2.1. *Let μ be a multifractal Gibbs capacity, and $\eta \in (0, 1)$. Let q_1 be the unique solution of the equation $\tau_{\mu}(q) = 0$, and let $H_1 = \tau'_{\mu}(q_1)$. Equivalently, $q_1 = \sigma'_{\mu}(H_1)$.*

With probability 1, $M_{1,\eta}(\mu)$ satisfies the multifractal formalism with

$$(4) \quad \sigma_{M_{1,\eta}(\mu)}(H) = \begin{cases} \eta \sigma_{\mu}(H/\eta) & \text{if } \eta H_{\min} \leq H < \eta H_1, \\ \frac{\sigma_{\mu}(H_1)}{H_1} H & \text{if } \eta H_1 \leq H < H_1, \\ \sigma_{\mu}(H) & \text{if } H_1 \leq H < H_{\max}. \end{cases}$$

and $\sigma_{M_{1,\eta}(\mu)}(H) = -\infty$ otherwise. Also,

$$(5) \quad \tau_{M_{1,\eta}(\mu)}(q) = \sigma_{M_{1,\eta}(\mu)}^*(q) = \begin{cases} \tau_{\mu}(q) & \text{if } q < q_1, \\ \eta \tau_{\mu}(q) & \text{if } q \geq q_1. \end{cases}$$

The continuity of $\sigma_{M_{1,\eta}(\mu)}$ can be checked directly on the formula. The intermediate linear phase is the unique interpolation between $H \mapsto \eta \sigma_{\mu}(H/\eta)$ and $H \mapsto \sigma_{\mu}(H)$ which makes the resulting spectrum concave.

2.2. **The case $0 < \varrho < 1$.** This situation, in which there is a lot of redundancy in the values $\mu(I^\varrho)$, is quite versatile and involves some extra parameters. The existence of the parameters introduced below, as well as their justification and their properties, will be studied in Section 5.

- Let θ_ϱ be the mapping defined as

$$\theta_\varrho : H \in [H_{\min}, \tau'_\mu(0)] \mapsto \frac{\varrho H \sigma_\mu(H)}{d(1-\varrho) + \varrho \sigma_\mu(H)}.$$

It increases continuously from $\frac{\varrho H_{\min} \sigma_\mu(H_{\min})}{d(1-\varrho) + \varrho \sigma_\mu(H_{\min})}$ to $\varrho \tau'_\mu(0)$.

- Define q_ϱ as the unique $q \in \mathbb{R}$ such that $\tau_\mu(q) = \frac{d(1-\varrho)}{\varrho}$ and set $H_\varrho = \tau'_\mu(q_\varrho)$.

Let us now define a mapping, that will be the multifractal spectrum of $\mathbf{M}_{\varrho,\eta}(\mu)$.

Definition 2.2. (1) Suppose that $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho}$. Then set

$$(6) \quad D_{\mu,\varrho,\eta}(H) = \begin{cases} \eta(d(1-\varrho) + \varrho \sigma_\mu(H/\varrho\eta)) & \text{if } \varrho\eta H_{\min} \leq H < \varrho\eta H_\varrho, \\ \frac{\sigma_\mu(H_\varrho)}{\theta_\varrho(H_\varrho)} H & \text{if } \varrho\eta H_\varrho \leq H < \theta_\varrho(H_\varrho), \\ \sigma_\mu(\theta_\varrho^{-1}(H)) & \text{if } \theta_\varrho(H_\varrho) \leq H < \varrho\tau'_\mu(0), \\ \sigma_\mu(H/\varrho) & \text{if } \varrho\tau'_\mu(0) \leq H \leq \varrho H_{\max}, \end{cases}$$

and $D_{\mu,\varrho,\eta}(H) = -\infty$ otherwise.

(2) Suppose that $\sigma_\mu(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho}$. Let $H_{\varrho,\eta} \in [H_{\min}, \tau'_\mu(0)]$ be the unique solution to

$$\sigma_\mu(H_{\varrho,\eta}) = \frac{d(1-\varrho)}{1/\eta-\varrho}.$$

This can be equivalently rewritten as $\varrho\eta H_{\varrho,\eta} = \theta_\varrho(H_{\varrho,\eta})$

(a) If $H_{\varrho,\eta} < H_\varrho$, then set

$$(7) \quad D_{\mu,\varrho,\eta}(H) = \begin{cases} \sigma_\mu(H/\varrho\eta) & \text{if } \varrho\eta H_{\min} \leq H < \varrho\eta H_{\varrho,\eta}, \\ \eta(d(1-\varrho) + \varrho \sigma_\mu(H/\varrho\eta)) & \text{if } \varrho\eta H_{\varrho,\eta} \leq H < \varrho\eta H_\varrho, \\ \frac{\sigma_\mu(H_\varrho)}{\theta_\varrho(H_\varrho)} H & \text{if } \varrho\eta H_\varrho \leq H < \theta_\varrho(H_\varrho), \\ \sigma_\mu(\theta_\varrho^{-1}(H)) & \text{if } \theta_\varrho(H_\varrho) \leq H < \varrho\tau'_\mu(0), \\ \sigma_\mu(H/\varrho) & \text{if } \varrho\tau'_\mu(0) \leq H \leq \varrho H_{\max}, \end{cases}$$

and $D_{\mu,\varrho,\eta}(H) = -\infty$ otherwise.

(b) If $H_{\varrho,\eta} \geq H_\varrho$, then set

$$(8) \quad D_{\mu,\varrho,\eta}(H) = \begin{cases} \sigma_\mu(H/\varrho\eta) & \text{if } \varrho\eta H_{\min} \leq H < \varrho\eta H_{\varrho,\eta}, \\ \sigma_\mu(\theta_\varrho^{-1}(H)) & \text{if } \varrho\eta H_{\varrho,\eta} \leq H < \varrho\tau'_\mu(0), \\ \sigma_\mu(H/\varrho) & \text{if } \varrho\tau'_\mu(0) \leq H \leq \varrho H_{\max}, \end{cases}$$

and $D_{\mu,\varrho,\eta}(H) = -\infty$ otherwise.

Observe that when $H_{\varrho,\eta} = H_\varrho$, then $\varrho\eta H_{\varrho,\eta} = \varrho\eta H_\varrho = \theta_\varrho(H_\varrho)$, and the second and third intervals in (7) merge to a single point. These intervals do not matter in (8) when $H_{\varrho,\eta} \geq H_\varrho$.

The formulas are not easy to handle with, but they are driven by the model. In particular, in each case there are different phases. The continuity of $D_{\mu,\varrho,\eta}$ can be checked on the formulas, and its other properties (differentiability, concavity) are studied in Section 5 and gathered in the next proposition. One can already notice that in (6) and (7) there is a linear part in the spectrum.




FIGURE 1. **Left:** Spectrum of $M_{\varrho, \eta}(\mu)$ for a given spectrum of μ when $\varrho = 0.8$, $\eta = 0.8$, case **2a**: 5 phases. **Right:** Spectrum when $\varrho = 0.7$, $\eta = 0.4$, case **2b**: 3 phases. Original spectrum in dashed blue, Legendre spectrum in dashed black.

Proposition 2.3. *For all $\varrho \in (0, 1)$, the function $D_{\mu, \varrho, \eta}$ is concave, continuous, and analytic on every open interval given by the different formulas in Definition 2.2. Moreover, $D_{\mu, \varrho, \eta}$ is differentiable, except at $\varrho\eta H_{\varrho, \eta}$ in cases (2a) and (2b) of Definition 2.2.*

Our main theorem is the following.

Theorem 2.4. *Let μ be a multifractal Gibbs capacity on $[0, 1]^d$, and $\eta \in (0, 1)$. For all $\varrho \in (0, 1)$, with probability 1:*

- (1) $\sigma_{M_{\varrho, \eta}(\mu)} = D_{\mu, \varrho, \eta}$.
- (2) One has

$$\tau_{M_{\varrho, \eta}(\mu)}(q) = \begin{cases} d(\varrho - 1) + \varrho\tau_{\mu}(q) & \text{if } q < q_{\varrho}, \\ \eta(d(\varrho - 1) + \varrho\tau_{\mu}(q)) & \text{if } q \geq q_{\varrho}. \end{cases}$$

- (3) Set

$$I_{\varrho, \eta} = \begin{cases} [\varrho\eta H_{\min}, \theta_{\varrho}(H_{\varrho})] \cup \{\varrho\tau'_{\mu}(0)\} & \text{if } \sigma_{\mu}(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho} \\ [\varrho\eta H_{\varrho, \eta}, \theta_{\varrho}(H_{\varrho})] \cup \{\varrho\tau'_{\mu}(0)\} & \text{if } \sigma_{\mu}(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho} \text{ and } H_{\varrho, \eta} < H_{\varrho}, \\ \{\varrho\eta H_{\varrho, \eta}\} \cup \{\varrho\tau'_{\mu}(0)\} & \text{if } \sigma_{\mu}(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho} \text{ and } H_{\varrho, \eta} = H_{\varrho} \\ \{\varrho\tau'_{\mu}(0)\} & \text{if } \sigma_{\mu}(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho} \text{ and } H_{\varrho, \eta} > H_{\varrho}. \end{cases}$$

The multifractal formalism holds for $M_{\varrho, \eta}(\mu)$ over $I_{\varrho, \eta}$ and it fails over $\text{dom}(\sigma_{M_{\varrho, \eta}(\mu)}) \setminus I_{\varrho, \eta}$.

So, we are able to compute in each case the multifractal spectrum and the L^q -spectrum, but these two quantities do not form a Legendre pair (i.e. they are not Legendre transform one of each other). A very careful analysis must be done to find a sharper upper bound for $\sigma_{M_{\varrho, \eta}(\mu)}$ than the one provided by the multifractal formalism.

One may wonder if alternatively the multifractal formalism developed by Olsen in [19] does hold for $M_{\varrho, \eta}(\mu)$. However this formalism is tailored to study the Hausdorff dimension of the level sets of the local dimension, while (for all ρ), these sets $E_{M_{\varrho, \eta}(\mu)}(H)$ (see Definition 3.1) are empty for H in a non-trivial subinterval of $\text{dom}(\sigma_{M_{\varrho, \eta}(\mu)})$.

2.3. The case $1 < \varrho \leq 1/\eta$. In this case, there is less redundancy. For $\varrho \in (1, 1/\eta]$, let us introduce some parameters:

- Let q_{ϱ} be the unique solution of $\varrho - 1 + \varrho\tau_{\mu}(q) = 0$, and $H_{\varrho} = \tau'_{\mu}(q_{\varrho})$.
- Let $\tilde{H}_{\varrho} = \min\{H \geq 0 : \sigma_{\mu}(H) \geq d(1 - 1/\varrho)\}$.




FIGURE 2. **Left:** Spectrum of $M_{1,\eta}(\mu)$ for a given spectrum of μ when $\eta = 0.7$: 3 phases. **Right:** Spectrum of $M_{\varrho,\eta}(\mu)$ when $\varrho = 0.55$, $\eta = 0.5$, case 1: 4 phases. Original spectrum in dashed blue, Legendre spectrum in dashed black (for $\rho < 1$ only).

- Finally, $\widehat{H}_\varrho = -\frac{\tau_\mu(q_\varrho)}{q_\varrho}$.

Theorem 2.5. *Let μ be a multifractal Gibbs capacity, and $\eta \in (0, 1)$. For all $\varrho \in (1, 1/\eta]$, with probability 1, $M_{\varrho,\eta}$ satisfies the multifractal formalism with*

$$\sigma_{M_{\varrho,\eta}(\mu)}(H) = \begin{cases} \eta(1 - \varrho + \varrho\sigma_\mu(H/\varrho\eta)) & \text{if } \varrho\eta\widehat{H}_\varrho \leq H < \varrho\eta H_\varrho, \\ q_\varrho H & \text{if } \varrho\eta H_\varrho \leq H < H_\varrho + \widehat{H}_\varrho, \\ \sigma_\mu(H - \widehat{H}_\varrho) & \text{if } H_\varrho + \widehat{H}_\varrho \leq H \leq \tau'_\mu(\infty) + \widehat{H}_\varrho \end{cases}$$

and $\sigma_{M_{\varrho,\eta}(\mu)}(H) = -\infty$ otherwise. In particular,

$$\tau_{M_{\varrho,\eta}(\mu)}(q) = \sigma_{M_{\varrho,\eta}(\mu)}^*(q) = \begin{cases} \tau_\mu(q) - \frac{\tau_\mu(q_\varrho)}{q_\varrho} \cdot q & \text{if } q \leq q_\varrho, \\ \eta(\varrho - 1 + \varrho\tau_\mu(q)) & \text{if } q_\varrho < q < \widetilde{q}_\varrho, \\ \eta\varrho\tau'_\mu(\widetilde{q}_\varrho) \cdot q & \text{if } \widetilde{q}_\varrho < +\infty \text{ and } q \geq \widetilde{q}_\varrho. \end{cases}$$

There are two phase transitions in $\sigma_{M_{\varrho,\eta}(\mu)}$, and one or two in $\tau_{M_{\varrho,\eta}(\mu)}$.

This situation for $\varrho \in (1, 1/\eta)$ is very similar to the case $\varrho = 1/\eta$ studied in [5], and with some work, one can adapt easily the arguments of [5]. Note that in [5], when $\varrho = 1/\eta$, we define H_ϱ equivalently as the unique real number such that the tangent to the graph of σ_μ at the point $(H_\varrho, \sigma_\mu(H_\varrho))$ passes through $(0, d(1 - 1/\varrho))$, and \widehat{H}_ϱ as $-\frac{\tau_\mu(\sigma'_\mu(H_\varrho))}{\sigma'_\mu(H_\varrho)}$. These equivalent formulations do hold when $\varrho \in (1, 1/\eta)$.

We choose not to give the proof of Theorem 2.5, and to focus on Theorems 2.1 and 2.4, since these are two situations where new phenomena appear, in particular where the multifractal formalism does not hold any more.

Let also note that the graph of the multifractal spectrum of $M_{\varrho,\eta}(\mu)$ depends continuously (in the Hausdorff distance metric) on $\eta \in (0, 1)$ and $\varrho \in (0, 1/\eta]$.

2.4. Lacunary wavelet series. Let us quickly explain the connexion with lacunary wavelet series. In [13], Jaffard studied real random processes called lacunary wavelet series, essentially equivalent

to the following model when $d = 1$: for $\gamma > 0$ and $\eta < 1$, let

$$L_{\gamma,\eta}(x) = \sum_{j \geq 1} \sum_{k=0}^{2^j-1} 2^{-j\gamma} X_{j,k} \psi(2^j x - k), \quad x \in \mathbb{R},$$

where ψ is a wavelet (see [18] for precise definitions), and the $X_{j,k}$ are independent random variables such that each $X_{j,k}$ is a Bernoulli variable with parameter $2^{-j\eta}$. Recalling Definition 1.1, it turns out [15, 14] that analyzing the pointwise regularity and the multifractal properties of μ is equivalent to performing the multifractal analysis the capacity $M(c_w)$, where the sequence c_w is the wavelet coefficients of $L_{\gamma,\eta}$: $c_w = (2^{-j\gamma} X_{j,k})_{j \geq 1, k \in \{0, \dots, 2^j-1\}}$.

Hence our model can be seen as a generalisation of the $L_{\gamma,\eta}$ process, where for $I \in \mathcal{D}_j$ the coefficient $2^{-j\gamma}$ (which can be interpreted as $(\mathcal{L}^1(I))^\gamma$ or $\mathcal{L}^1(I^\gamma)$) is replaced by $\nu(I)^\gamma$ for a Gibbs measure μ . In other words, instead of working from the uniform Lebesgue measure, we start with a multifractal measure and obtain a random process with a much richer structure.

Let us mention [11] where some variations of Jaffard's model are considered to construct random functions supported on Cantor sets, and whose restrictions to their support have a piecewise linear non-decreasing multifractal spectrum and do not obey the multifractal formalism. Other interesting multifractal random wavelets series are considered in [1, 3, 10].

2.5. Final remark on τ_{M_ϱ} and the large deviations spectra of M_ϱ .

Definition 2.6. Let $\mu \in \text{Cap}([0, 1]^d)$ with $\text{supp}(\mu) \neq \emptyset$. For $H \in \mathbb{R}$, $j \in \mathbb{N}$ and $\varepsilon > 0$, let

$$\mathcal{E}_\mu(j, H \pm \varepsilon) = \left\{ w \in \Sigma_j : \frac{\log_2 \mu(I_w)}{-j} \in [H - \varepsilon, H + \varepsilon] \right\}.$$

Then, the lower and upper large deviations spectra of μ are respectively defined as

$$\begin{aligned} \underline{f}_\mu(H) &= \lim_{\varepsilon \rightarrow 0} \liminf_{j \rightarrow +\infty} \frac{\log_2 \#\mathcal{E}_\mu(j, H \pm \varepsilon)}{j} \\ \text{and } \bar{f}_\mu(H) &= \lim_{\varepsilon \rightarrow 0} \limsup_{j \rightarrow +\infty} \frac{\log_2 \#\mathcal{E}_\mu(j, H \pm \varepsilon)}{j}. \end{aligned}$$

Let now μ be a multifractal Gibbs capacity. The equality $\sigma_\mu(H) = \tau_\mu^*(H)$ can be strengthened into $\dim E_\mu(H) = \tau_\mu^*(H)$ for all $H \in \mathbb{R}$ [8, 20] (see Definition 3.1 for the definition of $E_\mu(H)$); together with the standard inequalities $\dim E_\mu(H) \leq \underline{f}_\mu(H) \leq \bar{f}_\mu(H) \leq \tau_\mu^*(H)$ (valid for any capacity [19, 16, 17]), this yields $\underline{f}_\mu = \bar{f}_\mu = \tau_\mu^*$ [21, 19, 12]. Alternatively, this comes from the facts that τ_μ is differentiable and $\tau_\mu = \lim_{j \rightarrow \infty} \tau_{\mu,j}$, and the Gartner-Ellis theorem [9]. It is not hard to deduce from the studies conducted in this paper and its companion [5] that for all $\varrho \in (0, 1/\eta)$, with probability 1, one has $\tau_{M_{\varrho,\eta}(\mu)} = \lim_{j \rightarrow \infty} \tau_{M_{\varrho,\eta}(\mu),j}$ and $\underline{f}_{M_{\varrho,\eta}(\mu)} = \bar{f}_{M_{\varrho,\eta}(\mu)} = \tau_{M_{\varrho,\eta}(\mu)}^*$.

2.6. Organisation of the article. In Section 3 we recall additional properties associated with the multifractal formalism of Gibbs capacities and measures, as well as some properties of the random sampling process that were already established in [5]. In Section 4, the case $\rho = 1$ is treated (Theorem 2.1). Section 5 contains the proof of the redundant case $\rho \in (0, 1)$, Theorem 2.4.

3. MORE ON THE MULTIFRACTAL FORMALISM, GIBBS CAPACITIES, AND THE RANDOM SAMPLING

3.1. Some notations. For every $j \geq 1$, let $\Sigma_j = (\{0, 1\}^d)^j$. Let $\Sigma^* = \bigcup_{j \geq 0} \Sigma_j$ and $\Sigma = (\{0, 1\}^d)^\mathbb{N}$ be respectively the sets of finite and infinite words over the alphabet $\{0, 1\}^d$. By convention $(\{0, 1\}^d)^0$ contains the empty word denoted by ϵ . The set Σ^* acts on $\Sigma^* \cup \Sigma$ by concatenation, that

is if $u \in \Sigma^*$ and $v \in \Sigma^* \cup \Sigma$, the concatenation $u \cdot v$ of u and v is obtained by adding the prefix u to the word v . If $u \in \Sigma^*$, it defines the cylinder $[u]$ consisting of all infinite words with common prefix u . Let $\mathcal{C} = \{[u] : u \in \Sigma^*\}$. The length of an element u of $\Sigma^* \cup \Sigma$, that is its number of letters, will be denoted by $|u|$. If $0 \leq j \leq |u|$, $u|_j$ stands for the prefix of u of length j . The set Σ is endowed with the standard ultrametric distance $d : (x, y) \mapsto 2^{-|x \wedge y|}$, where $x \wedge y$ is the longest common prefix of x and y .

The coding map $\pi : t = (t_n^{(1)}, \dots, t_n^{(d)})_{n \geq 1} \in \Sigma \mapsto \left(\sum_{n=1}^{\infty} t_n^{(i)} 2^{-n} \right)_{1 \leq i \leq d} \in [0, 1]^d$ maps each cylinder $[w] \in \mathcal{C}$ onto the closed nontrivial dyadic cube

$$I_w = \prod_{i=1}^d \left[\sum_{k=1}^{|w|} w_k^{(i)} 2^{-k}, 2^{-|w|} + \sum_{k=1}^{|w|} w_n^{(i)} 2^{-k} \right].$$

We define

$$(9) \quad x_w = \left(\sum_{k=1}^j w_k^{(i)} 2^{-k} \right)_{1 \leq i \leq d}.$$

If $x = (x^{(1)}, x^{(2)}, \dots, x^{(d)}) \in [0, 1]^d$ has no dyadic component, then x is encoded by a unique $w = (w^{(1)}, w^{(2)}, \dots, w^{(d)}) \in \Sigma$, and $I_j(x)$ stands for $I_{w|_j}$. When $x^{(i)}$ is dyadic, $w^{(i)}$ is chosen as the largest element of $\{0, 1\}^{\mathbb{N}^+}$ in lexicographical order which encodes $x^{(i)}$. In both cases, $w|_j$ is also denoted $x|_j$.

3.2. Additional properties associated with multifractal formalism and Gibbs capacities.

Definition 3.1. Let $\mu \in \text{Cap}([0, 1]^d)$ with full support. For $x \in [0, 1]^d$, the lower and upper local dimensions of μ at x are respectively defined as

$$\underline{\dim}(\mu, x) = \liminf_{r \rightarrow 0^+} \frac{\log \mu(B(x, r))}{\log r} \quad \text{and} \quad \overline{\dim}(\mu, x) = \limsup_{r \rightarrow 0^+} \frac{\log \mu(B(x, r))}{\log r}.$$

When $\underline{\dim}(\mu, x) = \overline{\dim}(\mu, x)$, their common value is denoted by $\dim(\mu, x)$.

For $H \in \mathbb{R}$, set

$$\begin{aligned} \underline{E}_\mu(H) &= \left\{ x \in [0, 1]^d : \underline{\dim}(\mu, x) = H \right\}, \\ \overline{E}_\mu(H) &= \left\{ x \in [0, 1]^d : \overline{\dim}(\mu, x) = H \right\}, \\ E_\mu(H) &= \underline{E}_\mu(H) \cap \overline{E}_\mu(H). \end{aligned}$$

Observe that for every $x \in [0, 1]^d$,

$$(10) \quad \underline{\dim}(\mu, x) = \liminf_{j \rightarrow +\infty} \frac{\log \mu \left(\bigcup_{I \in \mathcal{N}(I_j(x))} I \right)}{-j} = \liminf_{j \rightarrow +\infty} \frac{\log \max \left(\mu(I) : I \in \mathcal{N}(I_j(x)) \right)}{-j},$$

where $I_j(x)$ stands for the unique dyadic cube of generation j that contains x , and where the notation $\mathcal{N}(I)$ for $I \in \mathcal{D}_j$ stands for the family of dyadic cubes neighbor to I of same generation:

$$(11) \quad \text{for } I \in \mathcal{D}_j, \quad \mathcal{N}(I) = \{I' \in \mathcal{D}_j : \bar{I} \cap \overline{I'} \neq \emptyset\}$$

contains $I_j(x)$ and its $3^d - 1$ neighboring dyadic cubes at generation j . By abuse of notations, we will sometimes write $w \in \mathcal{N}(I)$ for $I \in \mathcal{D}_j$ to describe all the dyadic cubes $I_w \in \mathcal{D}_j$ belonging to $\mathcal{N}(I)$.

Definition 3.2. For any fully supported capacity $\mu \in \text{Cap}([0, 1]^d)$, define the level sets

$$\begin{aligned} \underline{E}_\mu^{\leq}(H) &= \{x \in [0, 1]^d : \underline{\dim}(\mu, x) \leq H\} \\ \text{and } \overline{E}_\mu^{\geq}(H) &= \{x \in [0, 1]^d : \overline{\dim}(\mu, x) \geq H\}. \end{aligned}$$

The multifractal formalism also claims that

$$(12) \quad \dim \underline{E}_\mu^{\leq}(H) \leq \tau_\mu^*(H) \text{ if } H \leq \tau_\mu'(0^-),$$

$$(13) \quad \text{and } \dim \overline{E}_\mu^{\geq}(H) \leq \tau_\mu^*(H) \text{ if } H \geq \tau_\mu'(0^+).$$

Let now μ be a multifractal Gibbs capacity μ . One always has:

- τ_μ is a limit (not only a liminf), i.e.

$$(14) \quad \tau_\mu(q) = \lim_{j \rightarrow \infty} \tau_{\mu, j}(q), \quad \text{for every } q \in \mathbb{R}.$$

- τ_μ is strictly concave, analytic (due to the properties recalled in the previous section), and σ_μ is strictly concave, and real analytic over (H_{\min}, H_{\max}) .
- μ is quasi-multiplicative : There exists a constant $C \geq 1$ such that for every finite words $w, w' \in \Sigma^*$,

$$(15) \quad C^{-1} \mu(I_w) \mu(I_{w'}) \leq \mu(I_{ww'}) \leq C \mu(I_w) \mu(I_{w'}).$$

- There are two exponents $0 < \alpha_m \leq \alpha_M < +\infty$ such that for every finite word $w \in \Sigma^*$,

$$(16) \quad 2^{-|w|\alpha_M} \leq \mu(I_w) \leq 2^{-|w|\alpha_m}.$$

- μ is doubling.

These properties will be used repeatedly in the following.

3.3. Basic properties of the distribution of the surviving vertices. All these properties are proved in [5]. If $w \in \Sigma^*$, then p_{I_w} is simply denoted p_w .

Definition 3.3. If $w \in \Sigma^*$ and $p_w = 1$, say that w is a surviving vertex (or a survivor).

For every $j \geq 1$, denote by $\mathcal{S}_j(\eta)$ the (random) set of surviving vertices in Σ_j :

$$\mathcal{S}_j(\eta) := \{w \in \Sigma_j : p_w = 1\}.$$

Recall that x_w , defined by (9), is the dyadic point corresponding to the projection of the finite word $w \in \Sigma_j$ to $[0, 1]^d$. The first question concerns the distribution of the points x_w , for $w \in \mathcal{S}_j(\eta)$.

Definition 3.4. For every $j \geq 1$, and every finite word $W \in \Sigma^*$, one sets

$$\mathcal{S}_j(\eta, W) = \{w \in \mathcal{S}_j(\eta) : I_w \subset I_W\}.$$

The set $\mathcal{S}_j(\eta, W)$ describes the surviving vertices at generation j included in I_W . Obviously, for every $J \leq j$,

$$\mathcal{S}_j(\eta) = \bigcup_{W \in \Sigma_J} \mathcal{S}_j(\eta, W).$$

Lemma 3.5. [5] There exists a positive sequence $(\varepsilon_j)_{j \geq 1}$ converging to 0 such that, with probability 1, for every j large enough, for every $W \in \Sigma_{\lfloor j(\eta - \varepsilon_j) \rfloor}$, one has $\mathcal{S}_j(\eta, W) \neq \emptyset$.

So, every cylinder of generation $\lfloor j(\eta - \varepsilon_j) \rfloor$ contains a surviving vertex w of generation j .

The sequence $(\varepsilon_j)_{j \geq 1}$ is now fixed.

Lemma 3.5 has the following consequence: Almost surely, the set of points belonging to an infinite number of balls of the form $B(x_w, 2^{-\lfloor |w|(\eta - \varepsilon_{|w|}) \rfloor})$ with $p_w = 1$ is exactly the whole cube $[0, 1]^d$, i.e.

$$(17) \quad [0, 1]^d = \limsup_{j \rightarrow +\infty} \bigcup_{w \in \mathcal{S}_j(\eta)} B(x_w, 2^{-\lfloor |w|(\eta - \varepsilon_{|w|}) \rfloor}).$$

Next we obtain an upper bound for the cardinality of $\mathcal{S}_j(\eta, W)$ when $W \in \Sigma_{\lfloor \eta j \rfloor}$.

Lemma 3.6. *With probability one, for every large j and every $W \in \Sigma_{\lfloor \eta j \rfloor}$, $\#\mathcal{S}_j(\eta, W) \leq j$.*

As a conclusion, one keeps in mind the intuition that every cylinder $W \in \Sigma_{\lfloor \eta j \rfloor}$ contains at least one, but not much more than one surviving vertex $w \in \mathcal{S}_j(\eta)$.

4. PROOF OF THEOREM 2.1

Recalling Definitions 1.4, one has

$$M_{1,\eta}(\mu) = M_\eta(D_\eta \mu) = M(S_\eta(D_\eta \mu)).$$

By Definitions 1.1 and 1.2, each $M_{1,\eta}(\mu)(I)$, for $I \in \mathcal{D}_j$, can be written as

$$M_{1,\eta}(\mu)(I) = \sup_{I' \subset I, p_{I'} = 1} \mu((I')^\eta).$$

For sake of simplicity, $M_{1,\eta}(\mu)$ is denoted by M_1 in this section.

4.1. Upper and lower bound for the weights of M_1 on dyadic cubes. By some abuse of notations, we sometimes identify a dyadic cube $I \subset [0, 1]^d$ of generation j with the corresponding word $w \in \Sigma_j$.

Lemma 4.1. *With probability 1, there exists a positive sequence $(\tilde{\varepsilon}_j)_{j \geq 1}$ such that for every $w \in \Sigma^*$,*

$$(18) \quad \mu(I_w) 2^{-|w| \tilde{\varepsilon}_{|w|}} \leq M_1(I_w) \leq \mu(I_w^\eta).$$

Proof. Fix $w \in \Sigma_j$. In order to compute $M_1(I_w)$, one needs to find the largest value $\mu((I')^\eta)$ for $I' \subset I_w$ and $p_{I'} = 1$.

The upper bound in (18) simply follows from the hierarchical structure of the initial Gibbs capacity μ . Indeed, the greatest value that may appear in the computation of $M_1(I_w)$ is $\mu(I_w^\eta)$.

Further, we estimate the lower bound. The covering lemma 3.5 ensures that when $j = |w|$ is large enough, there is at least one surviving vertex $w' \in \mathcal{S}_{\lfloor j/(\eta - \varepsilon_j) \rfloor}(\eta)$ such that $I_{w'} \subset I_w$. Hence

$$M_1(I_w) \geq \mu(I_{w'}^\eta).$$

The cube $I_{w'}^\eta$ corresponds to a cube $I_{\tilde{w}}$, where \tilde{w} is the prefix of the word w' of length $\lfloor \eta |w'| \rfloor = \lfloor \eta \lfloor j/(\eta - \varepsilon_j) \rfloor \rfloor$, which satisfies for large j (i.e. for ε_j small)

$$j(1 + \varepsilon_j/\eta) - \eta - 1 \leq \lfloor \eta |w'| \rfloor \leq j(1 + 2\varepsilon_j/\eta).$$

Since $I_{w'} \subset I_w$, w is a prefix of the word corresponding to the cube $I_{w'}^\eta$, so we can write using the quasi-Bernoulli property (15) of μ that

$$\frac{1}{C} \mu(I_w) \mu(J) \leq \mu(I_{w'}^\eta) \leq C \mu(I_w) \mu(J)$$

for some dyadic cube J of generation between $j\varepsilon_j/\eta - \eta - 1$ and $j2\varepsilon_j/\eta$.

Using the uniform bounds (16) for μ to bound $\mu(J)$, recalling $j = |w|$, we have

$$\frac{1}{C} \mu(I_w) 2^{-(|w| \varepsilon_{|w|} \eta - \eta - 1) \alpha_M} \leq \mu(I_{w'}^\eta) \leq C \mu(I_w) 2^{-(|w| \varepsilon_{|w|} \eta - \eta - 1) \alpha_m}.$$

This can be rewritten, for some positive sequence $(\tilde{\varepsilon}_j)_{j \geq 1}$ converging to zero, as

$$(19) \quad 2^{-|w|\tilde{\varepsilon}_{|w|}} \leq \frac{\mu(I_{w'})^\eta}{\mu(I_w)} \leq 2^{|w|\tilde{\varepsilon}_{|w|}},$$

hence the result. \square

4.2. Local dimensions of M_1 , μ , and approximation rates. The value of the local dimensions of M_1 at a point $x \in [0, 1]^d$ depends on how close to the "surviving" wavelet coefficients x are. Hence, we introduce an approximation rate and some following approximating sets.

4.2.1. Approximation rate and local dimensions of M_1 . Let us denote, for every $w \in \Sigma^*$,

$$(20) \quad \ell_w = 2^{-\lfloor |w|(\eta - \varepsilon_{|w|}) \rfloor}.$$

Recall that by (17), almost surely, one has the covering property

$$[0, 1]^d = \limsup_{j \rightarrow +\infty} \bigcup_{k \in \mathcal{S}_j(\eta)} B(x_w, \ell_w).$$

Definition 4.2. For every $\delta \geq 1$, we define

$$\mathcal{A}_\delta = \limsup_{j \rightarrow +\infty} \bigcup_{k \in \mathcal{S}_j(\eta)} B(x_w, (\ell_w)^\delta).$$

The set \mathcal{A}_δ contains those points x which are approximated at a rate $\delta \geq 1$ by those dyadic numbers x_w such that $p_w = 1$. Obviously, $\mathcal{A}_1 = [0, 1]^d$ almost surely.

Definition 4.3. For every $x \in [0, 1]^d$, the approximation rate of x is the real number

$$\delta_x = \sup\{\delta \geq 1 : x \in \mathcal{A}_\delta\}.$$

A remarkable property is that, at the points x which are badly approximated by the surviving dyadic numbers, the local dimension of M_1 is easily expressed in terms of the local dimension of μ .

Proposition 4.4. Almost surely, for every $x \in [0, 1]^d$, if $\delta_x = 1$, then $\underline{\dim}(M_1, x) = \underline{\dim}(\mu, x)$.

Proof. Let $x \in [0, 1]^d$ be such that $\delta_x = 1$. Recall that $I_j(x) \in \mathcal{D}_j$ is the unique dyadic cube of generation j that contains x . Our goal is to estimate $M_1(B(x, 2^{-j}))$. Observe that by (10), it is equivalent to estimate

$$(21) \quad \tilde{M}_1(I_j(x)) = \max\left(M_1(I) : I \in \mathcal{N}(I_j(x))\right),$$

where $\mathcal{N}(I)$ was defined in (11).

Since $\delta_x = 1$, for every $\varepsilon > 0$, x does not belong to $\mathcal{A}_{1+\varepsilon}$. In other words, there exists an integer $J_{\varepsilon, x}$ such that for any $j \geq J_{\varepsilon, x}$, x does not belong to $\bigcup_{w \in \mathcal{S}_j(\eta)} B(x_w, (\ell_w)^{1+\varepsilon})$.

Let us use this to estimate from above the value of $\tilde{M}_1(I_w)$, where $w \in \Sigma_j$ is the unique word such that $I_j(x) = I_w$. To do so, it is enough to consider the dyadic cubes $I_{w'}$ such that:

- $|w'| \geq |w| = j$,
- $w' \in \mathcal{S}_{j'}(\eta)$,
- $x_{w'} \in \bigcup_{I \in \mathcal{N}(I_j(x))} I$.

As noticed in the proof of Lemma 4.1, it is enough to consider the words w' such that $j \leq |w'| \leq j/(\eta - \varepsilon_j)$. Moreover, for j large enough, it is not possible that $w' \in \mathcal{S}_{j'}(\eta)$ satisfies $j' \leq j/(\eta + \varepsilon)$ and $|x_{w'} - x| \leq 2 \cdot 2^{-j}$. This would imply that

$$|x_{w'} - x| \leq 2 \cdot 2^{-j} \leq 2 \cdot 2^{-j'(\eta + \varepsilon)} < (\ell_{w'})^{1+\varepsilon},$$

contradicting $\delta_x = 1$. Hence, the value of $M_1(I_w)$ is reached for some word w' of length satisfying $j/(\eta + \varepsilon) \leq j' \leq j/(\eta - \varepsilon_j)$. Using the same kind of estimates as in Lemma 4.1 and the doubling property of μ , the corresponding value $\mu(I_{w'})$ can be bounded by

$$(22) \quad \mu(I_w)2^{-j\tilde{\varepsilon}_j} \leq \mu(I_{w'}) \leq \mu(I_w)2^{j\tilde{\varepsilon}_j},$$

for some other sequence $\tilde{\varepsilon}_j$ independent of x (we keep the same notation $(\tilde{\varepsilon}_j)$ for the new sequence, to keep them simple). Hence $\tilde{M}_1(I_w)$ satisfies

$$(23) \quad \max(\mu(I) : I \in \mathcal{N}(I_j(x))) 2^{-j\tilde{\varepsilon}_j} \leq \tilde{M}_1(I_w) \leq \max(\mu(I) : I \in \mathcal{N}(I_j(x))) 2^{j\tilde{\varepsilon}_j}.$$

Taking the liminf in (23), we conclude the proof of Proposition 4.4. \square

4.2.2. Some approximating sets; heterogeneous ubiquity. In order to better understand the relation between the local behavior of μ and the approximation rate, we introduce some approximating sets.

Definition 4.5. Let $H \geq 0$, $\delta \geq 1$ and let $\tilde{\xi} := (\xi_j)_{j \geq 1}$ be a positive sequence. A word $w \in \Sigma^*$ is said to satisfy property $\mathcal{P}(H, \tilde{\xi})$ whenever

$$(24) \quad |I_w^\eta|^{H+\xi_{|w|}} \leq \mu(I_w^\eta) \leq |I_w^\eta|^{H-\xi_{|w|}}.$$

Set

$$\mathcal{A}_{H, \tilde{\xi}, \delta} = \bigcap_{J \geq 1} \bigcup_{j \geq J} \bigcup_{w \in \mathcal{S}_j(\eta): w \text{ satisfies } \mathcal{P}(H, \tilde{\xi})} B(x_w, (\ell_w)^\delta),$$

The set $\mathcal{A}_{H, \tilde{\xi}, \delta}$ contains those points x which are approximated at a rate δ by dyadic points x_w satisfying the property that the measure μ has a controlled local Hölder behavior around x_w .

Proposition 4.6. If $\tilde{\xi}$ converges to 0 and $x \in \mathcal{A}_{H, \tilde{\xi}, \delta}$, then

$$\eta H \leq \underline{\dim}(M_1, x) \leq \frac{H}{\min(\delta, 1/\eta)}.$$

Proof. The left inequality is true for all x 's, and follows simply from (18).

Let $x \in \mathcal{A}_{H, \tilde{\xi}, \delta}$. There exists an infinite sequence of words $(w_n)_{n \geq 1}$ such that

- $|x - x_{w_n}| \leq (\ell_{w_n})^\delta = 2^{-\delta \lfloor |w_n|(\eta - \varepsilon_{|w_n|}) \rfloor}$,
- $p_{w_n} = 1$,
- $2^{-\lfloor \eta |w_n| \rfloor (H + \xi_{|w_n|})} \leq \mu(I_{w_n}^\eta) \leq 2 \cdot 2^{-\lfloor \eta |w_n| \rfloor (H - \xi_{|w_n|})}$.

The last property follows from (24).

Set $\tilde{\delta} = \min(\delta, 1/\eta)$. For every $n \geq 1$, let $J_n = \lfloor \tilde{\delta} \lfloor |w_n|(\eta - \varepsilon_{|w_n|}) \rfloor \rfloor$. We are going to estimate $\tilde{M}_1(I_{J_n}(x))$ from below. To do so, as noticed in the proofs of Lemma 4.4 and Proposition 4.1, it is enough to consider the words w' such that $J_n \leq |w'| \leq J_n/(\eta - \varepsilon_{J_n})$.

In particular, by construction the dyadic cube I_{w_n} satisfies that $I_{w_n} \subset \bigcup_{I \in \mathcal{N}(I_{J_n}(x))} I$ and

$$J_n \leq \tilde{\delta} |w_n|(\eta - \varepsilon_{|w_n|}) \leq |w_n| \leq \frac{J_n}{\tilde{\delta}(\eta - \varepsilon_{|w_n|})} \leq \frac{J_n}{\eta - \varepsilon_{J_n}},$$

since $1 \leq \tilde{\delta} \leq 1/\eta$. So,

$$\tilde{M}_1(I_{J_n}(x)) \geq \mu(I_{w_n}^\eta) \geq 2^{-\lfloor \eta |w_n| \rfloor (H + \xi_{|w_n|})} \geq C 2^{-(H + \xi_{|w_n|}) \eta \frac{J_n}{\tilde{\delta}(\eta - \varepsilon_{|w_n|})}} \geq C 2^{-J_n \cdot \frac{(H + \xi_{|w_n|})}{\tilde{\delta}} \cdot \frac{1}{1 - \varepsilon_{|w_n|}/\eta}}.$$

Consequently,

$$\liminf_{n \rightarrow +\infty} \frac{\log \tilde{M}_1(I_{J_n}(x))}{\log 2^{-J_n}} \leq \lim_{n \rightarrow +\infty} \frac{(H + \xi_{|w_n|})}{\tilde{\delta}} \cdot \frac{1}{1 - \varepsilon_{|w_n|}/\eta} = \frac{H}{\tilde{\delta}}.$$

□

A remarkable property is that the Hausdorff dimension of the sets $\mathcal{A}_{H, \tilde{\xi}, \delta}$ can be bounded from below using the notion of *heterogeneous ubiquity* developed in [2, 4]. Next theorem corresponds to Theorem 2.7 in [4], or in a weaker form to Theorem 1.2 in [2], in the case $\varrho = 1$.

Theorem 4.7. *Let μ be a Gibbs capacity as in Definition 1.5. With probability 1, since (17) holds true, for all $\delta \geq 1$ and $H \geq 0$ such that $\sigma_\mu(H) > 0$, there exists a sequence $\tilde{\xi}$ decreasing to zero and a Borel probability measure $m_{H, \delta}$ such that:*

- $m_{H, \delta}(\mathcal{A}_{H, \tilde{\xi}, \delta}) > 0$,
- for any set E of Hausdorff dimension less than $\sigma_\mu(H)/\delta$, one has $m_{H, \delta}(E) = 0$.

In other words, the lower Hausdorff dimension of $m_{H, \delta}$ is larger than or equal to $\sigma_\mu(H)/\delta$. In particular, $\dim \mathcal{A}_{H, \tilde{\xi}, \delta} \geq \sigma_\mu(H)/\delta$.

Remark 4.8. (1) Actually the theorems of [2, 4] are proved when the condition (24) is replaced by

$$(25) \quad (\ell_w)^{H + \xi_{|w|}} \leq \mu(B(x_w, \ell_w)) \leq (\ell_w)^{H - \xi_{|w|}}.$$

One easily checks that in the proofs of [2, 4], (25) can be replaced by (24). Indeed, recalling (20), the doubling property of the Gibbs capacity μ implies that $\mu(B(x_w, \ell_w))$ and $\mu(I_w^\eta)$ are comparable, since both $B(x_w, \ell_w)$ and I_w^η contain x_w and have comparable diameters.

(2) Theorem 2.7 of [4] deals with $H \in (H_{\min}, H_{\max})$, but a simple adaptation of the proof makes it possible to extend the result to $H \in \{H_{\min}, H_{\max}\}$.

4.3. Upper bound for the spectrum of M_1 . To find an upper bound for σ_{M_1} , we use the inequality (2). Recall that q_1 is the unique solution of the equation $\tau_\mu(q) = 0$.

Proposition 4.9. *Almost surely, for every $q \in \mathbb{R}$, we have*

$$(26) \quad \tau_{M_1}(q) \geq T(q) := \begin{cases} \tau_\mu(q) & \text{if } q < q_1, \\ \eta \tau_\mu(q) & \text{if } q \geq q_1. \end{cases}$$

Proof. Let $j \geq 1$. Recalling (1) and (21), one has to estimate the sum $\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q$.

- **Case $q < 0$:** using the lower bound in (18), one has

$$\begin{aligned} \sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q &\leq \sum_{w \in \Sigma_j} \max(M_1(I) : I \in \mathcal{N}(I_j(x)))^q \\ &\leq 3^{d+1} \sum_{w \in \Sigma_j} \mu(I_w)^q 2^{-q|w| \tilde{\varepsilon}_{|w|}} \leq 3^{d+1} \cdot 2^{-j(\tau_{\mu, j}(q) + q \tilde{\varepsilon}_j)}. \end{aligned}$$

Since $\lim_{j \rightarrow +\infty} \tilde{\varepsilon}_j = 0$, letting j go to infinity yields $\tau_{M_1}(q) \geq \tau_\mu(q)$.

- **Case $q > 0$:** the upper bound in (18) is actually too crude. Using the covering property 3.5, one knows that for every word $w \in \Sigma_j$, the value of $\tilde{M}_1(I_w)$ is reached for one word w' whose length is between j and $j/(\eta - \varepsilon_j)$. Conversely, each word $w' \in \mathcal{S}_{j'}(\eta)$ with $j \leq j' \leq j/(\eta - \varepsilon_j)$ may contribute to at most 3^d values of $\tilde{M}_1(I_w)$ for $|w| = j$. Hence, one has

$$\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta - \varepsilon_j) \rfloor} \sum_{w \in \mathcal{S}_{j'}(\eta)} \mu(I_w^\eta)^q \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta - \varepsilon_j) \rfloor} \sum_{w \in \Sigma_{j'}} p_w \mu(I_w^\eta)^q$$

Recalling that when $w' \in \Sigma_{j'}$, $p_{w'}$ is a Bernoulli variable with parameter $2^{-dj'(1-\eta)}$, one gets

$$\mathbb{E}\left(\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q\right) \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \Sigma_{j'}} 2^{-dj'(1-\eta)} \mu(I_w^\eta)^q.$$

Now, observe that each cube $I_{w'}$, for $w' \in \Sigma_{\lfloor \eta j' \rfloor}$ and $j \leq j' \leq j/(\eta - \varepsilon_j)$, contains $2^{d(j' - \lfloor j' \eta \rfloor)}$ dyadic cubes of generation $2^{j'}$. In other words, each $w' \in \Sigma_{\lfloor \eta j' \rfloor}$ with $j \leq j' \leq j/(\eta - \varepsilon_j)$ appears $2^{d(j' - \lfloor j' \eta \rfloor)}$ times in the previous double sum as a cube of the form I_w^η with $w \in \Sigma_{j'}$. Thus,

$$(27) \quad \mathbb{E}\left(\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q\right) \leq C \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \Sigma_{\lfloor j' \eta \rfloor}} \mu(I_w)^q \leq C \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} 2^{-\lfloor j' \eta \rfloor \tau_{\mu, \lfloor j' \eta \rfloor}(q)},$$

for some constant C depending only on the dimension d . The behavior of the last sum depends on the value of $\tau_\mu(q)$.

• **Subcase $q > 0$, $\tau_\mu(q) < 0$:** this coincides with the fact that $q < q_1$. Fix $s < \tau_\mu(q)$ and $\varepsilon = (\tau_\mu(q) - s)/3$. Recalling (14), for j' sufficiently large, $|\tau_{\mu, \lfloor j' \eta \rfloor}(q) - \tau_\mu(q)| \leq \varepsilon$. Then, since $\tau_\mu(q) < 0$, one has

$$\mathbb{E}\left(\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q\right) \leq C \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} 2^{-\lfloor j' \eta \rfloor (\tau_\mu(q) - \varepsilon)} \leq C 2^{-j\eta/(\eta-\varepsilon_j)(\tau_\mu(q) - \varepsilon)}.$$

Thus,

$$\mathbb{E}\left(\sum_{j \geq 1} 2^{js} \sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q\right) \leq C \sum_{j \geq 1} 2^{j(s - \eta/(\eta-\varepsilon_j)(\tau_\mu(q) - \varepsilon))} < \infty,$$

since when j becomes large, $s - \eta/(\eta - \varepsilon_j)(\tau_\mu(q) - \varepsilon) \leq -\varepsilon$. Hence, almost surely,

$$\sum_{j \geq 1} 2^{js} \sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q = \sum_{j \geq 1} 2^{js} 2^{-j\tau_{M_1, j}(q)} < \infty.$$

Consequently, $\tau_{M_1}(q) \geq s$, and letting s tend to $\tau_\mu(q)$ one gets $\tau_{M_1}(q) \geq \tau_\mu(q)$ almost surely.

• **Subcase $q > 0$, $\tau_\mu(q) > 0$:** this corresponds to the values $q > q_1$. The arguments above can be adapted here. Fix $s < \tau_\mu(q)$, $\varepsilon = (\tau_\mu(q) - s)/3$ and j' large. Since $\tau_\mu(q) > 0$, (27) gives in this case (since it involves a sum of a geometric sequence)

$$\mathbb{E}\left(\sum_{w \in \Sigma_j} \tilde{M}_1(I_w)^q\right) \leq C 2^{-j\eta(\tau_\mu(q) - \varepsilon)}.$$

Letting j tend to infinity, and then ε to zero, we obtain $\tau_{M_1}(q) \geq \eta\tau_\mu(q)$, almost surely.

• **Subcase $q > 0$, $\tau_\mu(q) = 0$:** In this case, a simple calculation gives $\tau_{M_1}(q) \geq \tau_\mu(q)$.

Finally, the previous lower bounds hold for each q almost surely, but τ_{M_1} and τ_μ being concave functions, they hold almost surely for all q . \square

Corollary 4.10. *With probability 1, one has $\sigma_{M_1} \leq \tau_{M_1}^* \leq T^*$.*

Proof. This is a direct consequence of the multifractal formalism (2), i.e. $\sigma_{M_1} \leq \tau_{M_1}^*$, together with the inequality $\tau_{M_1} \geq T$ established in the previous Proposition 4.9, so that $\tau_{M_1}^* \leq T^*$. \square

Simple calculations show that the Legendre transform of T coincides with the formula (4), i.e.

$$(28) \quad T^*(H) = \begin{cases} \eta\sigma_\mu(H/\eta) & \text{if } \eta H_{\min} \leq H < \eta H_1 \\ q_1 H & \text{if } \eta H_1 \leq H < H_1 \\ \sigma_\mu(H) & \text{if } H_1 \leq H < H_{\max} \end{cases}$$

and $T^*(H) = -\infty$ otherwise. It remains to prove that $\sigma_{M_1}(\mu) \geq T^*$.

4.4. Lower bound for the singularity spectrum of M_1 . We distinguish the three regions given by formula (4) that we claim to hold for σ_{M_1} . Recall that q_1 is the unique solution of the equation $\tau_\mu(q) = 0$, $H_1 = \tau'_\mu(q_1)$, and that this implies that $q_1 = \sigma'_\mu(H_1)$.

Observe first that by (18), necessarily $\underline{\dim}(M_1, x) \geq \eta H_{\min}$ for every $x \in [0, 1]^d$.

4.4.1. Case $\eta H_{\min} \leq H \leq \eta H_1$. Observe that we are in the increasing part of the spectrum. Heuristically, the value of the Hausdorff dimension of $\underline{E}_{M_1}(H) = \{x : \underline{\dim}(M_1, x) = H\}$ is obtained for the points x having an approximation rate equal to $1/\eta$ (recall Definition 4.3).

Let $\eta H_{\min} \leq H \leq \eta H_1$ and set $\tilde{H} = H/\eta$. If $\tilde{H} = H_{\min}$ and $\sigma_\mu(H_{\min}) = 0$, since it is easy to construct a decreasing sequence $\tilde{\xi}$ such that $\mathcal{A}_{H_{\min}, 1/\eta, \tilde{\xi}} \neq \emptyset$, and for all $x \in \mathcal{A}_{H_{\min}, 1/\eta, \tilde{\xi}}$ one has both $\underline{\dim}(M_1, x) \geq \eta H_{\min}$ by the above observation and $\underline{\dim}(M_1, x) \leq \eta H_{\min} \leq H_{\min}$ by Proposition 4.6, we get $\underline{E}_{M_1}(\eta H_{\min}) \neq \emptyset$ hence $\dim \sigma_{M_1}(\eta H_{\min}) \geq 0 = \sigma_\mu(H_{\min}) = T^*(\eta H_{\min})$.

If $\tilde{H} \neq H_{\min}$ or $\sigma_\mu(H_{\min}) > 0$, by Theorem 4.7 applied with \tilde{H} and $\delta = 1/\eta$, there exists a sequence $\tilde{\xi}$ and a measure $m_{\tilde{H}, 1/\eta}$ with the properties:

- $m_{\tilde{H}, 1/\eta}(\mathcal{A}_{\tilde{H}, 1/\eta, \tilde{\xi}}) > 0$,
- $m_{\tilde{H}, 1/\eta}(E) = 0$ for any Borel set E such that $\dim E < \sigma_\mu(\tilde{H})/(1/\eta) = \eta\sigma_\mu(\tilde{H})$.

Applying Proposition 4.6, every point $x \in \mathcal{A}_{\tilde{H}, 1/\eta, \tilde{\xi}}$ satisfies

$$\underline{\dim}(M_1, x) \leq \frac{\tilde{H}}{\min(1/\eta, 1/\eta)} = \eta\tilde{H} = H.$$

We deduce (recall Definition 3.2) that $m_{\tilde{H}, 1/\eta}(\underline{E}_{M_1}^{\leq}(H)) > 0$.

Further, note that the inequality $\tau_{M_1} \geq T$ implies that $\tau'_{M_1}(0^+) \geq T'(0)$. Consequently, since $H \leq T'(0) = \tau'_\mu(0)$, one deduces from (12) that for every $n \geq 1$,

$$\dim \underline{E}_{M_1}^{\leq}(H - 1/n) \leq (\tau_{M_1})^*(H - 1/n) \leq \eta\sigma_\mu\left(\frac{H - 1/n}{\eta}\right) = \eta\sigma_\mu\left(\tilde{H} - \frac{1}{n\eta}\right) < \eta\sigma_\mu(\tilde{H}).$$

Hence $m_{\tilde{H}, 1/\eta}(\underline{E}_{M_1}^{\leq}(H - 1/n)) = 0$. Finally, since

$$(29) \quad \underline{E}_{M_1}^{\leq}(H) = \underline{E}_{M_1}(H) \cup \bigcup_{n \geq 1} \underline{E}_{M_1}^{\leq}(H - 1/n),$$

one obtains $m_{\tilde{H}, 1/\eta}(\underline{E}_{M_1}(H)) > 0$, which implies the expected lower bound $\sigma_{M_1}(H) \geq \eta\sigma_\mu(\tilde{H})$.

4.4.2. Case $\eta H_1 < h \leq H_1$. The proof is similar to that used for $\eta H_{\min} \leq H \leq \eta H_1$, except that here, the value of the Hausdorff dimension of $\underline{E}_{M_1}(H)$ is obtained for points x which have an approximation rate between 1 and $1/\eta$.

Let $\eta H_1 < H \leq H_1$ and $\delta = \frac{H_1}{H} \in [1, 1/\eta)$. One can apply Theorem 4.7 with H_1 and δ . We reproduce exactly the same arguments as in the previous subsection. There exists a sequence $\tilde{\xi}$ and a measure $m_{H_1, \delta}$ with the properties:

- $m_{H_1, \delta}(\mathcal{A}_{H_1, \delta, \tilde{\xi}}) > 0$,
- $m_{H_1, \delta}(E) = 0$ for any Borel set E such that

$$\dim E < \frac{\sigma_\mu(H_1)}{\delta} = \frac{\sigma_\mu(H_1)}{H_1} H = q_1 H,$$

where the last equality comes from the equalities $\tau_\mu(q_1) = 0$, $H_1 = \tau'_\mu(q_1)$ and $\sigma_\mu(\tau'_\mu(q)) = \tau_\mu^*(\tau'_\mu(q)) = q\tau'_\mu(q) - \tau_\mu(q)$ for all $q \in \mathbb{R}$.

Further, applying Proposition 4.6, every point $x \in \mathcal{A}_{H_1, \delta, \tilde{\xi}}$ satisfies

$$\underline{\dim}(\mathbf{M}_1, x) \leq \frac{H_1}{\min(\delta, 1/\eta)} = \frac{H_1}{\delta} = \frac{H_1}{H_1/H} = H.$$

Consequently, $m_{H_1, \delta}(\underline{E}_{\mathbf{M}_1}^{\leq}(H)) > 0$, and by the same argument as above,

$$\dim \underline{E}_{\mathbf{M}_1}^{\leq}(H - 1/n) \leq (\tau_{\mathbf{M}_1})^*(H - 1/n) \leq q_1(H - 1/n),$$

hence $m_{H_1, \delta}(\underline{E}_{\mathbf{M}_1}^{\leq}(H - 1/n)) = 0$. Finally, using (29), necessarily $m_{H_1, \delta}(\underline{E}_{\mathbf{M}_1}(H)) > 0$, hence $\sigma_{\mathbf{M}_1}(H) \geq q_1 H$, which again is the expected lower bound.

4.4.3. Case $H \geq H_1$. We will need the following proposition. Recall that a Borel probability measure ν on \mathbb{R}^d is said to be exact dimensional with dimension $D \geq 0$ if $\nu(E_\nu(D)) = 1$, and that in this case $\nu(E) = 0$ for all Borel sets such that $\dim E > D$.

Proposition 4.11. [5, Theorem 5(1)] *With probability 1, for all $H \in [H_{\min}, H_{\max}]$, there exists an exact dimensional Borel probability measure μ_H of dimension $\sigma_\mu(H)$ such that*

$$\mu_H \left(\{x \in [0, 1]^d : \dim(\mu, x) = H\} \cap \{x \in [0, 1]^d : \delta_x = 1\} \right) = \mu_H \left(\{x \in [0, 1]^d : \dim(\mu, x) = H\} \right).$$

Let us mention that if $\mu \approx \nu^\gamma$ where ν is a Gibbs measure associated with a Hölder potential φ , then for $H \in (H_{\min}, H_{\max})$, $H = \tau'_\mu(q)$ for some q and μ_H can be taken equal to the Gibbs measure associated with the potential $q\gamma\varphi$.

Now, if $H \geq H_1$, by Proposition 4.4, any $z \in \{x \in [0, 1]^d : \dim(\mu, x) = H\} \cap \{x \in [0, 1]^d : \delta_x = 1\}$ satisfies $\underline{\dim}(\mathbf{M}_1, z) = H$. Hence μ_H is supported on $\underline{E}_{\mathbf{M}_\varrho}(H)$, so $\sigma_{\mathbf{M}_1}(H) \geq \sigma_\mu(H)$, which is the expected lower bound.

5. PROOF OF THEOREM 2.4

Here, $\varrho \in (0, 1)$ and $\mathbf{M}_{\varrho, \eta}$ is simply denoted by \mathbf{M}_ϱ .

Since $\varrho < 1$, there is a lot of redundancy since the dilation parameter over-compensates the sampling.

Recalling Definition 1.4, each $\mathbf{M}_\varrho(I)$, for $I \in \mathcal{D}_j$, can be written as

$$\mathbf{M}_\varrho(I) = \sup_{I' \subset I, p_{I'}=1} \mu((I')^{\varrho\eta}),$$

and let us introduce by analogy with (21)

$$(30) \quad \widetilde{\mathbf{M}}_\varrho(I) = \max(\mathbf{M}_\varrho(I) : I \in \mathcal{N}(I_j(x))).$$

Computing $\underline{\dim}(\mathbf{M}_\varrho, x)$ amounts to estimating the behavior of $\widetilde{\mathbf{M}}_\varrho(I_j(x))$ when j tends to infinity.

5.1. **Some auxiliary notations and properties.** We omit the subscript μ in the following definitions, although the quantities depend on σ_μ .

Definition 5.1. For all $H \in [H_{\min}, \tau'_\mu(0)]$ and $\delta \geq 1$, define

$$(31) \quad \delta_\varrho(H) = \frac{d(1-\varrho) + \varrho\sigma_\mu(H)}{\sigma_\mu(H)}$$

$$(32) \quad \theta_\varrho(H) = \frac{\varrho H}{\delta_\varrho(H)}.$$

Observe that:

- The mapping $H \mapsto \delta_\varrho(H)$ decreases from $\frac{d(1-\varrho) + \varrho\sigma_\mu(H_{\min})}{\sigma_\mu(H_{\min})}$ to 1 over $[H_{\min}, \tau'_\mu(0)]$ (in particular, $\delta_\varrho(H) \geq 1$).
- The mapping $H \mapsto \theta_\varrho(H)$ increases continuously from $\frac{\varrho H_{\min}\sigma_\mu(H_{\min})}{d(1-\varrho) + \varrho\sigma_\mu(H_{\min})}$ to $\varrho\tau'_\mu(0)$.
- Since $\delta_\varrho(H) \geq 1$, $\theta_\varrho(H) \leq \varrho H$, with equality only if $\delta_\varrho(H) = 1$, i.e. $H = \tau'_\mu(0)$.

Additional parameters are needed (they were defined in the introduction, but we give more details now):

- Let q_ϱ be the unique q such that

$$(33) \quad \tau_\mu(q_\varrho) = \frac{d(1-\varrho)}{\varrho},$$

and set

$$H_\varrho = \tau'_\mu(q_\varrho).$$

Recall that this q_ϱ exists since τ_μ is an homeomorphism of \mathbb{R} . The mapping

$$(34) \quad H \in [H_{\min}, H_{\max}] \mapsto \frac{d(1-\varrho) + \varrho\sigma_\mu(H)}{H}$$

is increasing on $[H_{\min}, H_\varrho]$ and decreasing on $[H_\varrho, H_{\max}]$. Indeed, the derivative of this mapping vanishes when $\varrho H\sigma'_\mu(H) - d(1-\varrho) - \varrho\sigma_\mu(H) = 0$, i.e. $H\sigma'_\mu(H) - \sigma_\mu(H) = \frac{d(1-\varrho)}{\varrho}$. The fact that $\tau_\mu = \sigma_\mu^*$ allows to conclude. So the mapping reaches its maximum at H_ϱ .

- A direct calculation shows that

$$(35) \quad \frac{d(1-\varrho) + \varrho\sigma_\mu(H_\varrho)}{\varrho H_\varrho} = \frac{\sigma_\mu(H_\varrho)}{\theta_\varrho(H_\varrho)} = q_\varrho.$$

Indeed, the first equality comes from (31) and (32). The second one follows from (33) and the fact that, by Legendre transform, $\sigma_\mu(H_\varrho) = q_\varrho H_\varrho - \tau_\mu(q_\varrho)$.

Next, as explained in Definition 2.2, three cases must be distinguished depending on the behavior of the multifractal spectrum σ_μ of μ .

- **Case 1.** $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho}$: For every $H \in [H_{\min}, \tau'_\mu(0)]$, $\sigma_\mu(H) > \frac{d(1-\varrho)}{1/\eta-\varrho}$, and $\delta_\varrho(H) \leq \delta_\varrho(H_{\min}) = \frac{d(1-\varrho) + \varrho\sigma_\mu(H_{\min})}{\sigma_\mu(H_{\min})} < 1/\eta$. In particular, $\varrho\eta H_\varrho < \theta_\varrho(H_\varrho)$.

Notice that $\frac{d(1-\varrho)}{1/\eta-\varrho}$ is always smaller than d , and that it decreases from ηd to 0 as ϱ increases from 0 to 1.

As a conclusion,

$$(36) \quad \varrho\eta H_{\min} < \varrho\eta H_\varrho < \theta_\varrho(H_\varrho) < \varrho H_\varrho < \varrho\tau'_\mu(0) \leq \varrho H_{\max},$$

which explains the regions in formula (6).

- **Case 2.** $\sigma_\mu(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho}$: By continuity of σ_μ in its increasing part, there exists a (necessarily unique) $H = H_{\varrho,\eta} \in [H_{\min}, \tau'_\mu(0)]$ such that

$$(37) \quad \sigma_\mu(H_{\varrho,\eta}) = \frac{d(1-\varrho)}{1/\eta-\varrho}.$$

This is equivalently written as

$$\delta_\varrho(H_{\varrho,\eta}) = 1/\eta.$$

Note that $H_{\varrho,\eta} = H_\varrho$ if and only if $q_\varrho \tau'_\mu(q_\varrho) = d(1-\varrho)/(1-\varrho\eta)$.

- **Case 2.a.** $H_{\varrho,\eta} < H_\varrho$: here $\delta_\varrho(H_\varrho) < 1/\eta$, so $\varrho\eta H_{\varrho,\eta} < \varrho\eta H_\varrho < \theta_\varrho(H_\varrho)$, and

$$(38) \quad \varrho\eta H_{\min} < \varrho\eta H_{\varrho,\eta} < \varrho\eta H_\varrho < \theta_\varrho(H_\varrho) < \varrho H_\varrho < \varrho\tau'_\mu(0) \leq \varrho H_{\max},$$

which explains the regions in formula (7).

- **Case 2.b.** $H_{\varrho,\eta} \geq H_\varrho$: now $\delta_\varrho(H_\varrho) \geq 1/\eta$, so $\theta_\varrho(H_\varrho) \leq \varrho\eta H_\varrho \leq \varrho\eta H_{\varrho,\eta}$, and the proof will show that only the value of $\varrho\eta H_{\varrho,\eta}$ matters. Hence, the relevant regions in (8) are

$$(39) \quad \varrho\eta H_{\min} < \varrho\eta H_{\varrho,\eta} < \varrho\tau'_\mu(0) \leq \varrho H_{\max}.$$

One has the following simple rewriting of (31):

$$(40) \quad \frac{d(1-\varrho) + \varrho\sigma_\mu(\theta_\varrho^{-1}(H))}{\delta_\varrho(\theta_\varrho^{-1}(H))} = \sigma_\mu(\theta_\varrho^{-1}(H)).$$

All these formulas will be useful when looking for the optimal values of the parameters to compute the Hausdorff dimensions of the level sets $E_{M_\varrho}(H)$ in the corresponding intervals of values for H .

We conclude this section with the proof of Proposition 2.3.

Proof of Proposition 2.3. The regularity properties (continuity, analyticity on the region's interior) of $D_{\mu,\varrho,\eta}$ are easily checked. It is also immediate that among the intervals distinguished in the formulas (6), (7) and (8) providing $D_{\mu,\varrho,\eta}$, the only one over which the concavity of $D_{\mu,\varrho,\eta}$ is not clear is the interval where $\sigma_\mu(\theta_\varrho^{-1}(H))$. Once this concavity property is verified, the global concavity of $D_{\mu,\varrho,\eta}$ is immediate for the Cases 1 and 2.b., and it follows from the fact that $D'_{\mu,\varrho,\eta}(\varrho\eta H_{\varrho,\eta}^-) = \eta^{-1}D'_{\mu,\varrho,\eta}(\varrho\eta H_{\varrho,\eta}^+) > D'_{\mu,\varrho,\eta}(\varrho\eta H_{\varrho,\eta}^+)$ in Case 2.a.

To prove that $H \mapsto \sigma_\mu(\theta_\varrho^{-1}(H))$ is concave over $[H_\varrho, \varrho\tau'_\mu(0)]$ (and so over $[H_{\varrho,\eta}, \varrho\tau'_\mu(0)]$ in Case 2.b.), one rather shows that its reciprocal map $F = \theta_\varrho \circ \sigma_\mu^{-1}$ is convex over $[\sigma_\mu(H_\varrho), d]$. Indeed, one sees that $F(u) = \varrho \frac{u\sigma_\mu^{-1}(u)}{d(1-\varrho)+\varrho u}$, and a calculation shows that

$$F''(u) = \frac{\varrho}{(d(1-\varrho) + \varrho u)^3} (G(u) + 2d(1-\varrho)h(u)),$$

where, setting $g = \sigma_\mu^{-1}$, $h(u) = g'(u)(d(1-\varrho) + \varrho u) - \varrho g(u)$, and G is a polynomial function with positive coefficients in the variables $u, g(u), g'(u), g''(u)$, which are all non-negative over $[\sigma_\mu(H_\varrho), d]$ (using the concavity of σ_μ). Moreover, setting $H = g(u)$, one has $g'(u) = \sigma'_\mu(H)^{-1}$. So $h(u) \geq 0$ if and only if $d(1-\varrho) \geq \varrho(\sigma'_\mu(H)H - \sigma_\mu(H))$, that is $H = \tau'_\mu(q)$ with $\tau_\mu(q) \leq d(1-\varrho)/\varrho$, i.e. $H \geq H_\varrho$. Consequently, F'' is positive over $[\sigma_\mu(H_\varrho), d]$, hence the desired result. \square

5.2. **Rewriting the formula for $D_{\mu,\varrho,\eta}$.** Let us define the mapping (again, we omit the dependence on μ to keep the notations as light as possible)

$$(41) \quad m_\varrho(H, \delta) = \min \left(\frac{d(1-\varrho) + \varrho\sigma_\mu(H)}{\delta}, \sigma_\mu(H) \right), \quad H \in \mathbb{R}, \delta \geq 1.$$

The main proposition of this section, key to compute the multifractal spectrum of M_ϱ , is the following alternative formula for $D_{\mu,\varrho,\eta}$. This formula reflects the intuition that the fact that M_ϱ has at x a local dimension equal to H , results from a compromise between the local behavior of μ at x and the approximation rate of x by the surviving vertices.

Proposition 5.2. *If $H \geq 0$, set*

$$(42) \quad \tilde{D}_{\mu,\varrho,\eta}(H) = \max \left\{ m_\varrho(u, \delta) : 0 \leq \frac{\varrho u}{\delta} \leq H, 1 \leq \delta \leq 1/\eta \right\}.$$

One has $D_{\mu,\varrho,\eta} = \tilde{D}_{\mu,\varrho,\eta}$ over the interval $[0, \varrho\tau'_\mu(0)]$.

Proof. Observe that when $u < H_{\min}$, $\sigma_\mu(u) = -\infty$, so $m_\varrho(u, \delta) = -\infty$. So it is enough to consider $u \in [H_{\min}, H_{\max}]$.

Fact: When $\delta = \delta_\varrho(u)$ and $u \leq \varrho\tau'_\mu(0)$ then $\sigma_\mu(u) = \frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\delta}$. When $\delta < \delta_\varrho(u)$, the minimum in (41) with respect to u is reached by $\sigma_\mu(u)$, while when $\delta \geq \delta_\varrho(u)$ it is reached by $\frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\delta}$.

Case 1. $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta - \varrho}$: Let $H \in [0, \varrho\tau'_\mu(0)]$.

For all $u \in [H_{\min}, \tau'_\mu(0)]$, $\sigma_\mu(u) \geq \sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta - \varrho}$, so

$$\delta_\varrho(u) = \varrho + \frac{d(1-\varrho)}{\sigma_\mu(u)} < 1/\eta.$$

(i) When $1 \leq \delta \leq \delta_\varrho(u)$ is fixed, by the fact above, the minimum in (41) is reached at $\sigma_\mu(u)$. So, when looking for the maximum in (42), one tries to maximize $\sigma_\mu(u)$ for $H_{\min} \leq \frac{\varrho u}{\delta} \leq H$, $1 \leq \delta \leq \delta_\varrho(u)$. This maximum is reached for some u when $\delta = \delta_\varrho(u)$, i.e. one has to maximize $\sigma_\mu(u)$ for $H_{\min} \leq \frac{\varrho u}{\delta_\varrho(u)} \leq H$. The maximum is reached when $\frac{\varrho u}{\delta_\varrho(u)} = H$, i.e. when $u = \theta_\varrho^{-1}(H)$ (the reciprocal image of H by θ_ϱ), and its value is $\sigma_\mu(\theta_\varrho^{-1}(H))$.

(ii) Now, when $\delta_\varrho(u) \leq \delta \leq 1/\eta$, the minimum in (41) is reached at $\frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\delta}$. So one has to maximize $\frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\delta}$ for $H_{\min} \leq \frac{\varrho u}{\delta} \leq H$, $\delta_\varrho(u) \leq \delta \leq 1/\eta$. Given δ , this is maximal when $\varrho u/\delta = H$, and it equals $H \frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\varrho u}$. Since $\delta_\varrho(u) \leq \delta$, this implies that $\varrho u/\delta_\varrho(u) \geq H$, i.e. $u \geq \theta_\varrho^{-1}(H)$. In addition, since $\delta \leq 1/\eta$, the optimal u satisfies $u = H/(\varrho\eta)$. As a conclusion, the maximum in (42) in this range of δ 's can be written

$$\max \left\{ H \frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{\varrho u} : \theta_\varrho^{-1}(H) \leq u \leq H/\varrho\eta \right\}.$$

Recall that the mapping $u \mapsto \frac{d(1-\varrho) + \varrho\sigma_\mu(u)}{u}$ increases over $[H_{\min}, H_\varrho]$ and decreases over $[H_\varrho, \tau'_\mu(0)]$. Hence, the value of the above maximum gives birth to three phases, depending on the position of H_ϱ with respect to the interval $[\theta_\varrho^{-1}(H), H/\varrho\eta]$ (or by symmetry, on the position of H with respect to the interval $[\varrho\eta H_\varrho, \theta_\varrho(H_\varrho)]$).

More precisely:

- when $\varrho\eta H_{\min} \leq H \leq \varrho\eta H_\varrho$, the maximum is reached when $u = H/\varrho\eta$ (the corresponding δ is $1/\eta$), and it equals $\eta(d(1-\varrho) + \varrho\sigma_\mu(H/\varrho\eta))$.

- when $\varrho\eta H_\varrho \leq H \leq \theta_\varrho(H_\varrho)$, it is reached at $u = H_\varrho$ (with $\delta = \delta_\varrho(H_\varrho)H_\varrho/H$), and it equals $H \frac{d(1-\varrho)+\varrho\sigma_\mu(H_\varrho)}{\varrho H_\varrho} = H \frac{\sigma_\mu(\theta_\varrho(H_\varrho))}{\theta_\varrho(H_\varrho)}$ (here, we used (35)).
- when $\theta_\varrho(H_\varrho) \leq H \leq \varrho\tau'_\mu(0)$, it is reached at $u = \theta_\varrho^{-1}(H)$ (with $\delta = \delta_\varrho(H_\varrho)$), and it equals $H \frac{d(1-\varrho)+\varrho\sigma_\mu(\theta_\varrho^{-1}(H))}{\varrho\theta_\varrho^{-1}(H)} = \sigma_\mu(\theta_\varrho^{-1}(H))$, where the last equality comes from (32).

Putting the previous estimates together shows that $\tilde{D}_{\mu,\varrho,\eta}$ indeed coincides with $D_{\mu,\varrho,\eta}$, given by formula (6).

Case 2.a. $\sigma_\mu(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho}$ and $H_{\varrho,\eta} < H_\varrho$:

Recall the definition (37) of $H_{\varrho,\eta}$, as well as the series of inequalities (38).

(i) When $\varrho\eta H_{\min} \leq H \leq \varrho\eta H_{\varrho,\eta}$, all those $u \in [H_{\min}, \tau'_\mu(0)]$ such that $\varrho u/\delta \leq H$ with $1 \leq \delta \leq 1/\eta$ satisfy necessarily $u \leq H/(\varrho\eta) \leq H_{\varrho,\eta}$. Hence the maximum in (42) is attained when $u = H/(\varrho\eta)$ and $\delta = 1/\eta$, i.e. for the largest possible u , and it equals $\sigma_\mu(H/(\varrho\eta))$.

(ii) When $H > \varrho\eta H_{\varrho,\eta}$, remark that all the values $\sigma_\mu(u)$ for $\varrho u/\delta \leq H$, $1 \leq \delta \leq 1/\eta$ and $u \leq H_{\varrho,\eta}$ are less than $\sigma_\mu(H_{\varrho,\eta})$, so they will not contribute to find the maximum in (42). So, the discussion reduces to **Case 1.**, with the same conclusions.

This shows that $\tilde{D}_{\mu,\varrho,\eta}$ and $D_{\mu,\varrho,\eta}$ given by formula (7) coincide in this case.

Case 2.b. $\sigma_\mu(H_{\min}) \leq \frac{d(1-\varrho)}{1/\eta-\varrho}$ and $H_{\varrho,\eta} \geq H_\varrho$:

Recall (39). The proof is essentially the same as the previous one, except that in this case, only $H_{\varrho,\eta}$ matters:

(i) When $\varrho\eta H_{\min} \leq H \leq \varrho\eta H_{\varrho,\eta}$, by the same argument as above, all those $u \in [H_{\min}, \tau'_\mu(0)]$ such that $\varrho u/\delta \leq H$ with $1 \leq \delta \leq 1/\eta$ satisfy necessarily $u \leq H/(\varrho\eta) \leq H_{\varrho,\eta}$. Hence the maximum in (42) is $\sigma_\mu(H/(\varrho\eta))$.

(ii) When $H > \varrho\eta H_{\varrho,\eta}$, the same argument as in **Case 2.a (ii)** shows that the maximal value is $\sigma_\mu(\theta_\varrho^{-1}(H))$. \square

5.3. Estimations of the local dimensions. By analogy with Lemma 4.1 and Proposition 4.4, the following holds.

Lemma 5.3. *With probability 1:*

(1) *there exists a positive sequence $(\tilde{\varepsilon}_j)_{j \geq 1}$ such that for every $w \in \Sigma^*$,*

$$(43) \quad \mu(I_w^\varrho) 2^{-|w|\tilde{\varepsilon}_{|w|}} \leq \mathbf{M}_\varrho(I_w) \leq \mu(I_w^{\varrho\eta}).$$

(2) *for every $x \in [0, 1]$, if $\delta_x = 1$, then $\underline{\dim}(\mathbf{M}_\varrho, x) = \varrho \underline{\dim}(\mu, x)$.*

Proof. The proof is easily adapted from the ones of Lemma 4.1 and Proposition 4.4 by replacing I_w by I_w^ϱ everywhere it is necessary to (for instance, in (22) and then (23) in Proposition 4.4). \square

The elements of interest are now those points in $[0, 1]^d$ which are approximated at rate δ by dyadic numbers x_w such that $\mu(I_w^{\varrho\eta})$ (and not $\mu(I_w)$ as in Section 4.2.2) has a specific value. This explains the introduction of the property $\mathcal{P}(\varrho, H, \tilde{\xi})$ below.

Definition 5.4. *Let $H \geq 0$, $\delta \geq 1$ and $\tilde{\xi} = (\xi_j)_{j \geq 1}$ a positive sequence. The word $w \in \Sigma^*$ is said to satisfy $\mathcal{P}(\varrho, H, \tilde{\xi})$ whenever*

$$|I_w^{\varrho\eta}|^{H+\xi_{|w|}} \leq \mu(I_w^{\varrho\eta}) \leq |I_w^{\varrho\eta}|^{H-\xi_{|w|}}.$$

Then one sets

$$\mathcal{A}_{\varrho, H, \tilde{\xi}, \delta} = \bigcap_{J \geq 1} \bigcup_{j \geq J} \bigcup_{w \in \mathcal{S}_j(\eta): w \text{ satisfies } \mathcal{P}(\varrho, H, \tilde{\xi})} B(x_w, (\ell_w)^\delta).$$

The notations are consistent with those of the previous section: $\mathcal{P}(1, H, \tilde{\xi}) = \mathcal{P}(H, \tilde{\xi})$ and $\mathcal{A}_{1, H, \tilde{\xi}, \delta} = \mathcal{A}_{H, \tilde{\xi}, \delta}$. Proposition 4.6 extends to the following one.

Proposition 5.5. *With probability 1, if $\tilde{\xi}$ is a positive sequence converging to 0 and $x \in \mathcal{A}_{\varrho, H, \tilde{\xi}, \delta}$, then*

$$\varrho \eta H \leq \underline{\dim}(\mathbb{M}_\varrho, x) \leq \frac{\varrho H}{\min(\delta, 1/\eta)}.$$

Proof. Again, the proof of Proposition 4.6 can be modified by replacing I_w by I_w^η , and using $\mathcal{P}(\varrho, H, \tilde{\xi})$ instead of $\mathcal{P}(H, \tilde{\xi})$. \square

To compute the multifractal spectrum of \mathbb{M}_ϱ , the standard inequality (2) cannot be used, since the multifractal formalism fails. We rather use the following properties of the sets $\underline{E}_{\mathbb{M}_\varrho}(H)$.

Proposition 5.6. *With probability 1,*

(1) *For every $H \geq 0$, writing $\tilde{\varepsilon}$ for the constant sequence equal to ε , we have*

$$\underline{E}_{\mathbb{M}_\varrho}^{\leq}(H) \subset \bigcap_{\varepsilon > 0} \bigcup_{H' \geq 0, \delta \in [1, 1/\eta], \frac{\varrho H'}{\delta} \leq H + \varepsilon} \mathcal{A}_{\varrho, H', \tilde{\varepsilon}, \delta}.$$

(2) *For every $H \geq 0$, $\underline{E}_{\mathbb{M}_\varrho}(H) \subset \overline{E}_\mu^{\geq}(H/\varrho)$.*

The first inclusion will be used to find an upper bound for the multifractal spectrum $\sigma_{\mathbb{M}_\varrho}$ in the increasing part $H \leq \varrho \tau'_\mu(0)$, while the second one will be used in the decreasing part $H \geq \varrho \tau'_\mu(0)$.

Proof. (1) Suppose that $x \in \underline{E}_{\mathbb{M}_\varrho}(H)$. Recalling (10), there exists an increasing sequence $(J_n)_{n \geq 1}$ of positive integers such that

$$(44) \quad \lim_{n \rightarrow +\infty} \frac{\log \widetilde{\mathbb{M}}_\varrho(I_{J_n}(x))}{-J_n \log 2} = H.$$

By Lemma 3.5, there exists a sequence of positive integers $(J'_n)_{n \geq 1}$ satisfying $J_n \leq J'_n$ and $1 \leq \limsup_{n \rightarrow \infty} J'_n/J_n \leq 1/\eta$, such that for each n there exists $w_n \in \mathcal{S}_{J'_n}(\eta)$ and $\widetilde{\mathbb{M}}_\varrho(I_{J_n}(x)) = \mu(I_{w_n}^{\varrho \eta})$. Let us write $\mu(I_{w_n}^{\varrho \eta}) = |I_{w_n}^{\varrho \eta}|^{H_n}$ for some $H_n \in [\alpha_m, \alpha_M]$ (recall the uniform bound (16) for μ). One has

$$\frac{\log \widetilde{\mathbb{M}}_\varrho(I_{j_n}(x))}{-j_n \log 2} = \frac{\log \mu(I_{w_n}^{\varrho \eta})}{-J_n \log 2} = \frac{-\lfloor J'_n \varrho \eta \rfloor H_n}{-J_n} = \varrho H_n / \delta_n,$$

with $\delta_n = \frac{\varrho J_n}{\lfloor J'_n \varrho \eta \rfloor}$. By construction,

$$1 \leq \liminf_{n \rightarrow +\infty} \delta_n \leq \limsup_{n \rightarrow +\infty} \delta_n \leq 1/\eta.$$

Up to extraction of a subsequence, one may assume that the two sequences $(H_n)_{n \geq 1}$ and $(\delta_n)_{n \geq 1}$ converge respectively to $H' \in [H_{\min}, H_{\max}]$ and $\delta \in [1, 1/\eta]$. By (44), one has $\varrho H' / \delta = H$. To conclude, we remark that $|x - x_{w_n}| \leq 2 \cdot 2^{-J_n} = 2 \cdot 2^{-\delta_n \lfloor \varrho \eta J'_n \rfloor / \varrho}$. So, recalling (20), for every $\varepsilon > 0$, for every n large enough so that $|H_n - H'| \leq \varepsilon/2$, $|\delta_n - \delta| \leq \varepsilon/2$ and $|\varrho H_n / \delta - H| \leq \varepsilon/2$:

- $w_n \in \mathcal{S}_{J'_n}(\eta)$,
- $x \in B(x_{w_n}, 2 \cdot 2^{-\delta_n \lfloor \varrho \eta J'_n \rfloor / \varrho}) \subset B(x_{w_n}, \ell_{w_n}^{\delta - \varepsilon})$

- One has

$$(45) \quad \mu(I_{w_n}^{\varrho\eta}) \in [|I_{w_n}^{\varrho\eta}|^{H'-\epsilon}, |I_{w_n}^{\varrho\eta}|^{H'+\epsilon}].$$

Finally, x belongs to the set $\mathcal{A}_{\varrho, H', \tilde{\epsilon}, \delta - \epsilon}$, hence the result.

Remark 5.7. *One has $I_{w_n} \subset \mathcal{N}(I_{J_n}(x))$. Let us make denote by j_n the largest integer j such that $I_{w_n} \subset \mathcal{N}(I_j(x))$. Clearly $J_n \leq j_n \leq J'_n$ and $\widetilde{M}_\varrho(I_{j_n}(x)) = \mu(I_{w_n}^{\varrho\eta})$. It follows that $\lim_{n \rightarrow \infty} \frac{J_n}{j_n} = 1$, for otherwise $\underline{\dim}(M_\varrho, x) < H$.*

(2) With the same notations as above, for each $x \in \underline{E}_{M_\varrho}(H)$ we deduce from (45) and the doubling property of μ that $\overline{\dim}(\mu, x) \geq H' = H\delta/\varrho \geq H/\varrho$ since $\delta \geq 1$. \square

5.4. Useful bounds for estimating the Hausdorff dimensions of the sets $\mathcal{A}_{\varrho, H, \tilde{\xi}, \delta}$. To get upper and lower bounds for the multifractal spectrum σ_{M_ϱ} , we recall some results obtained in [4]. There, we studied the Hausdorff dimension of limsup sets of the form

$$\bigcap_{N \geq 1} \bigcup_{n \geq N: (r_n)^{eH + \xi_n} \leq \mu(B(t_n, r_n^\varrho)) \leq (r_n)^{eH + \xi_n}} B(t_n, r_n^\delta),$$

where $H \geq 0$, $\delta \geq 1$, $(\xi_n)_{n \geq 1}$ is a positive sequence converging to zero, μ is a Gibbs measure, and where the sequence of balls $(B(t_n, r_n))_{n \geq 1}$ satisfies some properties described in the article. This obviously looks exactly like our present situation with a proper reformulation: t_n and r_n are replaced by x_w and ℓ_w , and the scaling condition on $\mu(B(t_n, r_n^\varrho))$ is replaced by $\mathcal{P}(\varrho, H, \tilde{\xi})$ (which does not change the result, as explained in Remark 4.8).

In particular, a key property in [4] is called *weak redundancy*: the family of balls $(B(t_n, r_n))_{n \geq 1}$ is weakly redundant when calling $T_N = \{n : 2^{-N} \leq r_n \leq 2^{-N+1}\}$, there exists a sequence $(\alpha_N)_{N \geq 1}$ of positive numbers such that

- $\lim_{N \rightarrow +\infty} \frac{\alpha_N}{N} = 0$.
- each T_N can be decomposed into α_N families of pairwise disjoint families $(T_{N,i})_{i=1, \dots, \alpha_N}$ such that for every $n \neq n' \in T_{N,i}$, $B(t_n, r_n) \cap B(t_{n'}, r_{n'}) = \emptyset$.

Weak redundancy of a family of balls implies that the balls of comparable size do not overlap too much (their overlaps are controlled by the sequence $(\alpha_N)_{N \geq 1}$). This is the case for the sequence $(B(x_w, \ell_w))_{w \in \mathcal{S}_j(\eta)}$ by Lemma 3.6. Hence, we can use results of [4], whose statements are recalled now:

At first, we will need the following slight modification [4, of Theorem 2.2], which readily follows from the proof of this theorem and the doubling property of μ .

Theorem 5.8. *With probability 1, for every Gibbs capacity μ , every $H \geq 0$, $\delta \geq 1$, and $\epsilon > 0$, denoting by $\tilde{\epsilon}$ the constant sequence $(\epsilon)_{n \geq 1}$,*

$$(46) \quad \dim \mathcal{A}_{\varrho, H, \tilde{\epsilon}, \delta} \leq \min \left(\frac{d(1 - \varrho) + \varrho S_\mu(H, \epsilon)}{\delta - \epsilon}, S_\mu(H, \epsilon) \right),$$

where $S_\mu(H, \epsilon) = \sup\{\sigma_\mu(H') : H' \in [H - \epsilon, H + \epsilon]\}$.

Regarding the lower bound, since Lemma 3.5 implies that with probability 1, not only (17) holds, but for all $j \geq 1$ large enough, for all $w \in \mathcal{S}_j(\eta)$, there exists $\tilde{\mathcal{S}}_{j, \varrho}(w) \subset \mathcal{S}_j(\eta)$ of cardinality at least $5^{-1} \cdot 2^{j(1 - \varrho - \epsilon_j)}$, such that the balls $B(x_{w'}, \ell_{w'})$, $w' \in \tilde{\mathcal{S}}_{j, \varrho}(w)$, are included in $B(x_w, \ell_w)$ and are pairwise disjoint, the following result holds (recall the formula (41) for m_ϱ):

Theorem 5.9 (Corollary of Theorem 2.7 of [4]). *Let μ be a Gibbs capacity. Let $\varrho \in (0, 1)$.*

With probability 1, for all $\delta > 1$, and $H \geq 0$ such that $\sigma_\mu(H) > 0$, there exists a sequence $\tilde{\epsilon}$ decreasing to zero and a Borel probability measure $\nu_{\varrho, H, \delta}$ such that:

- $\nu_{\varrho, H, \delta}(\mathcal{A}_{\varrho, H, \tilde{\xi}, \delta}) > 0$,
- for any set E of Hausdorff dimension less than $m_{\varrho}(H, \delta)$, $\nu_{\varrho, H, \delta}(E) = 0$.

In particular, $\dim \mathcal{A}_{\varrho, H, \tilde{\xi}, \delta} \geq m_{\varrho}(H, \delta)$.

Remark 5.10. Like for the case $\varrho = 1$, Theorem 2.7 of [4] deals with $H \in (H_{\min}, H_{\max})$, but a simple adaptation of the proof makes it possible to extend the result to $H \in \{H_{\min}, H_{\max}\}$.

5.5. Upper bound for $\sigma_{M_{\varrho}}$. To check the desired upper bound $\sigma_{M_{\varrho}} \leq D_{\mu, \varrho, \eta}$, we distinguish the cases $H \leq \varrho\tau'_{\mu}(0)$ and $H \geq \varrho\tau'_{\mu}(0)$. The first case is a direct consequence of Proposition 5.2, (46), Proposition 5.6(1), and the following Lemma.

Lemma 5.11. For $\varepsilon > 0$, denote by $\tilde{\varepsilon}$ the constant sequence equal to ε . With probability 1, for all $H \geq 0$,

$$\dim \left(\bigcap_{\varepsilon > 0} \bigcup_{H' \geq 0, \delta \in [1, 1/\eta], \frac{\varrho H'}{\delta} \leq H + \varepsilon} \mathcal{A}_{\varrho, H', \tilde{\varepsilon}, \delta} \right) \leq \tilde{D}_{\mu, \varrho, \eta}(H).$$

Proof. Fix $\varepsilon \in (0, 1)$. For every integer $N \geq 1/\varepsilon$, set $P_N = \{k/N : k \in \mathbb{N}\}$. We have

$$\bigcup_{H' \geq 0, \delta \in [1, 1/\eta], \frac{\varrho H'}{\delta} \leq H + \varepsilon} \mathcal{A}_{H', \varrho, \tilde{\varepsilon}, \delta} = \bigcup_{H' \in P_N, \delta \in [1, 1/\eta] \cap P_N, \frac{\varrho H'}{\delta} \leq H + \varepsilon} \mathcal{A}_{H', \varrho, \tilde{\varepsilon}, \delta}.$$

By Lemma 3.6 the family of balls $(B(x_w, \ell_w))_{w \in \bigcup_{j \geq 1} S_j(\eta)}$ satisfies the weakly redundancy property [4, Definition 2.1]. Thus by (46),

$$\begin{aligned} \dim \bigcap_{\varepsilon > 0} \bigcup_{\substack{H' \geq 0, \delta \in [1, 1/\eta] \\ \frac{\varrho H'}{\delta} \leq H + \varepsilon}} \mathcal{A}_{H', \varrho, \tilde{\varepsilon}, \delta} &\leq \inf_{\varepsilon > 0} \liminf_{N \geq 1/\varepsilon} \sup_{\substack{H' \in P_N, \delta \in [1, 1/\eta] \cap P_N \\ \frac{\varrho H'}{\delta} \leq H + \varepsilon}} \min \left(\frac{d(1 - \varrho) + \varrho S_{\mu}(H', \varepsilon)}{\delta - \varepsilon}, S_{\mu}(H', \varepsilon) \right) \\ &= \tilde{D}_{\mu, \varrho, \eta}(H), \end{aligned}$$

the conclusion coming from (42) and the continuity of the upper-semicontinuous concave function σ_{μ} over its compact domain $[H_{\min}, H_{\max}]$. \square

For the second case $H \geq \varrho\tau'_{\mu}(0)$, combining item (2) of Proposition 5.6 and (13) applied to μ yields

$$\dim \underline{E}_{M_{\varrho}}(H) \leq \dim \overline{E}^{\geq}(H/\varrho) \leq \sigma_{\mu}(H/\varrho) = D_{\mu, \varrho, \eta}(H).$$

This completes the proof.

5.6. Lower bound for the spectrum. Here one proves that $\sigma_{M_{\varrho}} \geq D_{\mu, \varrho, \eta}$. The main idea is to find, for each H , with well-chosen parameters H' and δ , either a suitable measure of the form $\mu_{H'}$ as in Proposition 4.11, or a measure $\nu_{\varrho, H', \delta}$ from Theorem 5.9 carried by the set $\underline{E}_{M_{\varrho}}(H)$.

5.6.1. Decreasing part of the spectrum: $H \geq \varrho\tau'_{\mu}(0)$. If $H_{\min} \leq H \leq \varrho H_{\max}$, by item (2) of Lemma 5.3, one knows that $\{x \in [0, 1]^d : \dim(\mu, x) = H/\varrho\} \cap \{x \in [0, 1]^d : \delta_x = 1\} \subset \underline{E}_{M_{\varrho}}(H) \subset \underline{E}_{M_{\varrho}}(H)$. This implies that the measure $\mu_{H/\varrho}$ considered in Proposition 4.11 is supported on $\underline{E}_{M_{\varrho}}(H)$, hence $\dim \underline{E}_{M_{\varrho}}(H) \geq \sigma_{\mu}(H/\varrho)$. If $H > \varrho H_{\max}$, then $\sigma_{M_{\varrho}}(H) \geq -\infty = D_{\mu, \varrho, \eta}(H)$.

5.6.2. Increasing part of the spectrum when $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho}$.

The three phases exhibited by formula 6 correspond to various choices for H' and δ to get from Theorem 5.9 a good measure $\nu_{\varrho, H', \delta}$ supported on $\underline{E}_{M_\varrho}(H')$.

- $H \leq \varrho\eta H_\varrho$. Let $H' = H/\varrho\eta$ and $\delta = 1/\eta \geq \delta_\varrho(H')$.

Note that $\sigma_\mu(H') > 0$ or $\sigma_\mu(H') = -\infty$. If $\sigma_\mu(H') > 0$, consider the measure $\nu_{\varrho, H', \delta}$ and the set $\mathcal{A}_{\varrho, H', \delta, \tilde{\xi}}$ from Theorem 5.9 (for an appropriate sequence $\tilde{\xi}$). By Proposition 5.5,

$$(47) \quad \mathcal{A}_{\varrho, H', \delta, \tilde{\xi}} \subset \underline{E}_{M_\varrho}^\leq(H).$$

Moreover, by construction, for all $n \geq 1$, we know using the upper bound for σ_{M_ϱ} obtained in Section 5.5 that

$$\dim \underline{E}_{M_\varrho}^\leq(H - 1/n) \leq \tilde{D}_{\mu, \varrho, \eta}(H - 1/n) < D_{\mu, \varrho, \eta}(H) = \frac{d(1-\varrho) + \varrho\sigma_\mu(H/\varrho\eta)}{\delta} = m_\varrho(H, \delta),$$

the last inequality coming from the formula (6) for $D_{\mu, \varrho, \eta}$. Consequently, one has $\nu_{\varrho, H', \delta}(\mathcal{A}_{\varrho, H', \delta, \tilde{\xi}}) = 1$ and $\nu_{\varrho, H', \delta}(\underline{E}_{M_\varrho}^\leq(H - 1/n)) = 0$ for every $n \geq 1$, hence $\nu_{\varrho, H', \delta}$ is supported on the set

$$F = \mathcal{A}_{\varrho, H', \delta, \tilde{\xi}} \setminus \bigcup_{n \geq 1} \underline{E}_{M_\varrho}^\leq(H - 1/n) \subset \underline{E}_{M_\varrho}(H).$$

This shows that $\nu_{\varrho, H', \delta}(\underline{E}_{M_\varrho}(H)) = \nu_{\varrho, H', \delta}(F) = 1$, hence $\dim \underline{E}_{M_\varrho}(H) \geq D_{\mu, \varrho, \eta}(H)$.

If $\sigma_\mu(H') = -\infty$, then $\tilde{D}_{\mu, \varrho, \eta}(H) = D_{\mu, \varrho, \eta}(H) = -\infty$, so $\underline{E}_{M_\varrho}(H) = \emptyset$ by 5.6(1).

- $\varrho\eta H_\varrho \leq H \leq \theta_\varrho(H_\varrho)$. Here we set $H' = H_\varrho$ and $\delta = \delta_\varrho(H_\varrho)\theta_\varrho(H_\varrho)/H = \varrho H_\varrho/H$ and fix $\tilde{\xi}$ like in Theorem 5.9. Since $\frac{\varrho H'}{\delta} = \frac{\varrho H_\varrho}{\varrho H_\varrho/H} = H$, by Proposition 5.5, (47) still holds true, and one can imply the same arguments as when $H \leq \varrho\eta H_\varrho$. The lower bound for $\dim \underline{E}_{M_\varrho}(H)$ is then

$$m_\varrho(H', \delta) = \frac{d(1-\varrho) + \varrho\sigma_\mu(H')}{\delta} = \frac{d(1-\varrho) + \varrho\sigma_\mu(H_\varrho)}{\delta_\varrho(H_\varrho)H_\varrho/H} = \frac{\sigma_\mu(H_\varrho)}{\theta_\varrho(H_\varrho)} H$$

hence the desired value.

- $\theta_\varrho(H_\varrho) \leq H \leq \varrho\tau'_\mu(0)$. Here we set $H' = \theta_\varrho^{-1}(H)$, $\delta = \delta_\varrho(\theta_\varrho^{-1}(H))$ and fix $\tilde{\xi}$ like in Theorem 5.9. The same argument as $H \leq \varrho\eta H_\varrho$ yields the lower bound

$$m_\varrho(H', \delta) = \frac{d(1-\varrho) + \varrho\sigma_\mu(H')}{\delta} = \frac{d(1-\varrho) + \varrho\sigma_\mu(\theta_\varrho^{-1}(H))}{\delta_\varrho(\theta_\varrho^{-1}(H))} = \sigma_\mu(\theta_\varrho^{-1}(H)),$$

hence the expected value.

5.6.3. Increasing part of the spectrum when $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho}$ and $H_{\varrho, \eta} \leq H_\varrho$.

We only need to deal with the case $H \leq \varrho\eta H_{\varrho, \eta}$, since the other cases can be treated as in the previous subsection.

If $H \leq \varrho\eta H_{\varrho, \eta}$ and $\sigma_\mu(H/\varrho\eta) \geq 0$, set $H' = H/\varrho\eta$ and $\delta = 1/\eta \leq \delta_\varrho(H')$. If $\sigma_\mu(H') \geq 0$ and $\sigma_\mu(H') = 0$, then $H' = H_{\min}$. In this case, note that by item (1) of Lemma 5.3 one has $\underline{\dim}(M_\varrho, x) \geq \varrho\eta H_{\min}$ for all $x \in [0, 1]^d$. Moreover, it is not difficult to show that there exists a sequence $\tilde{\xi}$ such that $\mathcal{A}_{\varrho, H_{\min}, 1/\eta, \tilde{\xi}} \neq \emptyset$. Since by Proposition 5.5, $\mathcal{A}_{\varrho, H_{\min}, 1/\eta, \tilde{\xi}} \subset \underline{E}_{M_\varrho}^\leq(\varrho\eta H_{\min})$, this yields $\underline{E}_{M_\varrho}(\varrho\eta H_{\min}) \neq \emptyset$ and $\sigma_{M_\varrho}(\varrho\eta H_{\min}) \geq 0 = \sigma_\mu(H_{\min}) = D_{\mu, \varrho, \eta}(\varrho\eta H_{\min})$.

If $\sigma_\mu(H') > 0$, the discussion is the same as in the previous section except that here $m_\varrho(H', \delta) = \sigma_\mu(H')$ instead of $m_\varrho(H', \delta) = (1-\varrho + \varrho\sigma_\mu(H'))/\delta$.

If $\sigma_\mu(H') = -\infty$, then, as in the previous subsection, $\tilde{D}_{\mu, \varrho, \eta}(H) = D_{\mu, \varrho, \eta}(H) = -\infty$, so $\underline{E}_{M_\varrho}(H) = \emptyset$ by Proposition 5.6(1).

5.6.4. **Increasing part of the spectrum when** $\sigma_\mu(H_{\min}) > \frac{d(1-\varrho)}{1/\eta-\varrho}$ **and** $H_{\varrho,\eta} > H_\varrho$.

- $H \leq \varrho\eta H_{\varrho,\eta}$. The discussion is the same as in the previous subsection.
- $\varrho\eta H_{\varrho,\eta} \leq H \leq \varrho\tau'_\mu(0)$. The discussion is identical as in the previous subsection when $H \geq H_\varrho$, since we know from Section 5.1 that $\varrho\eta H_{\varrho,\eta} \geq H_\varrho$.

5.7. **About the multifractal formalism for M_ϱ .** In this section, we compute τ_{M_ϱ} and study the (non)-validity of the multifractal formalism for M_ϱ . First we restate item (2) of Theorem 2.4.

Proposition 5.12. *With probability 1, one has*

$$\tau_{M_\varrho}(q) = \begin{cases} d(\varrho - 1) + \varrho\tau_\mu(q) & \text{if } q < q_\varrho, \\ \eta(d(\varrho - 1) + \varrho\tau_\mu(q)) & \text{if } q \geq q_\varrho. \end{cases}$$

Proof. For $j \geq 1$, one estimates the sum $\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q$.

(i) *Lower bound for τ_{M_ϱ} :*

- **Case $q < 0$:** using the lower bound in item (1) of Lemma 5.3, one has

$$\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \leq \sum_{w \in \Sigma_j} \max(M_\varrho(I) : I \in \mathcal{N}(I_j(x)))^q \leq 3^d \sum_{w \in \Sigma_j} \mu(I_w^\varrho)^q 2^{-q|w|\tilde{\varepsilon}_{|w|}}.$$

In the last sum, each interval $I_{w'}$, for $w' \in \Sigma_{\lfloor j\varrho \rfloor}$, appears as I_w^ϱ in $2^{d(j-\lfloor j\varrho \rfloor)}$ terms. So

$$\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \leq C 2^{d(j-\lfloor j\varrho \rfloor)} \sum_{w' \in \Sigma_{\lfloor j\varrho \rfloor}} \mu(I_{w'})^q 2^{-qj\tilde{\varepsilon}_j} = C 2^{d(j-\lfloor j\varrho \rfloor)} 2^{-\lfloor j\varrho \rfloor \tau_{\mu,j}(q)}.$$

Letting j go to infinity yields $\tau_{M_\varrho}(q) \geq d(\varrho - 1) + \varrho\tau_\mu(q)$.

• **Case $q > 0$:** The same arguments as in Proposition 4.9 can be adapted. By 3.5, for every word $w \in \Sigma_j$, the value of $\widetilde{M}_\varrho(I_w)$ is reached for one word w' whose length is between j and $j/(\eta - \varepsilon_j)$, and each word $w' \in \mathcal{S}_{j'}(\eta)$ with $j \leq j' \leq j/(\eta - \varepsilon_j)$ may contribute to at most 3^d values of $\widetilde{M}_\varrho(I_w)$ for $|w| = j$. Hence the upper bound in item (1) of Lemma 5.3 gives

$$\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \mathcal{S}_{j'}(\eta)} \mu(I_w^{\varrho\eta})^q \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \Sigma_{j'}} p_w \mu(I_w^{\varrho\eta})^q$$

Taking expectation, one gets

$$\mathbb{E} \left(\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \right) \leq 3^d \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \Sigma_{j'}} 2^{-dj'(1-\eta)} \mu(I_w^{\varrho\eta})^q.$$

As before, each cube $I_{w'}$, for $w' \in \Sigma_{\lfloor \varrho\eta j' \rfloor}$ and $j \leq j' \leq j/(\eta - \varepsilon_j)$, contains at most $3^d \times 2^{d(j'-\lfloor j'\varrho\eta \rfloor)}$ dyadic cubes of generation $2^{j'}$. Thus,

$$\begin{aligned} \mathbb{E} \left(\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \right) &\leq 3^d \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} \sum_{w \in \Sigma_{\lfloor j'\varrho\eta \rfloor}} 2^{d(j'-\lfloor j'\varrho\eta \rfloor)} 2^{-dj'(1-\eta)} \mu(I_w)^q \\ (48) \qquad \qquad \qquad &\leq C \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} 2^{d\eta(1-\varrho)j'} 2^{-\lfloor j'\varrho\eta \rfloor \tau_{\mu,\lfloor j'\varrho\eta \rfloor}(q)}. \end{aligned}$$

Again, the behavior of the last sum depends on the value of $\tau_\mu(q)$.

• $0 < q < q_\rho$: since by (33) one has $d(\varrho - 1) + \varrho\tau_\mu(q_\rho) = 0$, one sees that $d(\varrho - 1) + \varrho\tau_\mu(q) < 0$. Fixing $s < d(\varrho - 1) + \varrho\tau_\mu(q)$ and $\varepsilon = (d(\varrho - 1) + \varrho\tau_\mu(q) - s)/3$. Recalling (14), for j' sufficiently large, $|\tau_{\mu, [j'\varrho\eta]}(q) - \tau_\mu(q)| \leq \varepsilon$, so that

$$\mathbb{E}\left(\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q\right) \leq C \sum_{j'=j}^{\lfloor j/(\eta-\varepsilon_j) \rfloor} 2^{-j'\eta(d(\varrho-1)+\varrho\tau_\mu(q)-\varepsilon)} \leq C 2^{-j\eta/(\eta-\varepsilon_j)((d(\varrho-1)+\varrho\tau_\mu(q)-\varepsilon))}.$$

Then, as before,

$$\mathbb{E}\left(\sum_{j \geq 1} 2^{js} \sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q\right) \leq C \sum_{j \geq 1} 2^{j(s-\eta/(\eta-\varepsilon_j)(d(\varrho-1)+\varrho\tau_\mu(q)-\varepsilon))} < \infty,$$

from which we deduce that, almost surely,

$$\sum_{j \geq 1} 2^{js} \sum_{w \in \Sigma_j} \widetilde{M}_\varrho 1(I_w)^q < \infty.$$

Consequently, $\tau_{M_\varrho}(q) \geq s$, and letting s tend to $d(\varrho - 1) + \varrho\tau_\mu(q)$ yields the result.

• $q > q_\rho$: Fix $0 < s < d(\varrho - 1) + \varrho\tau_\mu(q)$, $\varepsilon = (d(\varrho - 1) + \varrho\tau_\mu(q) - s)/3$ and j' large. Since $d(\varrho - 1) + \varrho\tau_\mu(q) > 0$, the sum in (48) is bounded above by the term $j' = j$, and

$$\mathbb{E}\left(\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q\right) \leq C 2^{-j\eta(d(\varrho-1)+\varrho\tau_\mu(q)-\varepsilon)}.$$

The same conclusion as above gives $\tau_{M_\varrho}(q) \geq \eta(d(\varrho - 1) + \varrho\tau_\mu(q))$, almost surely.

The case where $\tau_\mu(q) = 0$ is obtained by continuity.

(ii) *Upper bound for τ_{M_ϱ} :*

• Case $q \geq 0$: We show that with probability 1, for all $q \geq 0$,

$$\tau_{M_\varrho}(q) \leq \min(\eta(d(\varrho - 1) + \varrho\tau_\mu(q)), d(\varrho - 1) + \varrho\tau_\mu(q)) = \begin{cases} \eta(d(\varrho - 1) + \varrho\tau_\mu(q)) & \text{if } q \leq q_\rho \\ d(\varrho - 1) + \varrho\tau_\mu(q) & \text{if } q \geq q_\rho. \end{cases}$$

By Lemma 5.3, with probability 1, $M_\varrho(I_w) \geq \mu(I_w^\varrho) 2^{-j\tilde{\varepsilon}_j}$ for all $w \in \Sigma_j$, $j \geq 1$. Consequently, for every $q \geq 0$

$$\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \geq \sum_{w \in \Sigma_j} \mu(I_w^\varrho)^q 2^{-qj\tilde{\varepsilon}_j}.$$

By the arguments already used before, each $I_{w'}$, for $w' \in \Sigma_{\lfloor \varrho j \rfloor}$, appears $2^{d(j-\lfloor \varrho j \rfloor)}$ times in the above sum. Hence

$$\sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q \geq \sum_{w' \in \Sigma_{\lfloor \varrho j \rfloor}} \mu(I_{w'})^q 2^{d(j-\lfloor \varrho j \rfloor)} 2^{-qj\tilde{\varepsilon}_j} = 2^{-\lfloor \varrho j \rfloor \tau_{\mu, \lfloor \varrho j \rfloor}(q) + d(j-\lfloor \varrho j \rfloor) - qj\tilde{\varepsilon}_j}.$$

Letting j tend to infinity yields $\tau_{M_\varrho}(q) \leq d(\varrho - 1) + \varrho\tau_\mu(q)$ for all $q \geq 0$.

To get the other bound, let $H = \tau'_\mu(q)$. Consider the measure μ_H provided by Proposition 4.11 as in Lemma 3.5. Since μ_H is exact dimensional and supported on $E_\mu(H)$, and μ is doubling, given $\varepsilon > 0$, for J large enough, there is a subset $\Sigma_J(H, \varepsilon)$ of Σ_J of cardinality at least $2^{J(\tau_\mu^*(q) - \varepsilon)}$ such that for all $u \in \Sigma_J(H, \varepsilon)$ one has $2^{-J(H+\varepsilon)} \leq \mu(I_u) \leq 2^{-J(H-\varepsilon)}$.

We now use Lemma 3.5 and the sequence $(\varepsilon_j)_{j \geq 1}$ defined therein (which does not depend on q). With probability 1, when j is large enough, every word $v \in \Sigma_{\lfloor (\eta-\varepsilon_j)j \rfloor}$ contains at least one $w_v \in \Sigma_j$ such that $p_{w_v} = 1$. Hence, for j large enough, taking $J = \lfloor \varrho \eta j \rfloor$, fixing $u \in \Sigma_J(H, \varepsilon)$, this applies to each element $v \in \Sigma_{\lfloor (\eta-\varepsilon_j)j \rfloor}$ such that $I_v \subset I_u$: I_v contains a subcube I_{w_v} with

$w_v \in \Sigma_j$ and $p_{w_v} = 1$. In particular, $\widetilde{M}_\varrho(I_{w_v}) \geq \mu(I_{w_v}^{\varrho\eta})$. Also, $I_{w_v}^{\varrho\eta} \in \Sigma_J$, so $I_{w_v}^{\varrho\eta} = I_u$, and $\widetilde{M}_\varrho(I_v) \geq \mu(I_u) \geq 2^{-J(H+\varepsilon)} = 2^{-\varrho\eta j(H+\varepsilon)}$.

There are $2^{d(\lfloor(\eta-\varepsilon_j)j\rfloor-J)} = 2^{d(\lfloor(\eta-\varepsilon_j)j\rfloor-\lfloor\varrho\eta j\rfloor)}$ such disjoint subcubes I_v . These observations and the fact that $q \geq 0$ imply that

$$\begin{aligned} \sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q &\geq \sum_{u \in \Sigma_J(H, \varepsilon)} \sum_{v \in \Sigma_{\lfloor(\eta-\varepsilon_j)j\rfloor}: I_v \subset I_u} \sum_{w \in \Sigma_j: I_w \subset I_v \text{ and } p_{w_v}=1} \widetilde{M}_\varrho(I_w)^q \\ &\geq 2^{\lfloor\varrho\eta j\rfloor(\tau_\mu^*(q)-\varepsilon)} 2^{d(\lfloor(\eta-\varepsilon_j)j\rfloor-\lfloor\varrho\eta j\rfloor)} 2^{-\varrho\eta j(H+\varepsilon)}. \end{aligned}$$

This yields $\tau_{M_\varrho}(q) \leq \eta(d(\varrho-1) + \varrho(qH - \tau_\mu^*(q))) + O(\varepsilon)$. Since $qH - \tau_\mu^*(q) = \tau_\mu(q)$ and ε is arbitrary, we conclude that with probability 1, $\tau_{M_\varrho}(q) \leq \eta(d(\varrho-1) + \varrho\tau_\mu(q))$ for all $q \geq 0$.

• **Case $q < 0$:** Let us make two preliminary observations.

First, if $1 < \gamma < 1/\eta$, for all $j \geq 1$ and $w \in \Sigma_j$, consider the event $\mathcal{E}(I_w) = \{\exists v \in \bigcup_{k=j}^{\lfloor\gamma j\rfloor} \Sigma_k : I_v \subset 3I_w \text{ and } p_v = 1\}$. When $\mathcal{E}(I_w)$ does not hold for some $w \in \Sigma_j$, all the $\mu(I_v)$ are put to zero by the sampling process, for $I_v \subset 3I_w$ and $|v| \in \{j+1, \dots, \gamma j\}$. So intuitively, the corresponding surviving weight $\widetilde{M}_\varrho(I_w)$ will be small. By independence of the p_v 's,

$$(49) \quad \mathbb{P}(\mathcal{E}(I_w)) = 1 - \left(\prod_{k=j}^{\lfloor\gamma j\rfloor} (1 - 2^{-d(1-\eta)k})^{2^{d(k-j)}} \right)^{3^d} \leq 1 - \exp(-C2^{-d_j(1-\gamma\eta)}) \leq C2^{-d_j(1-\gamma\eta)}$$

for some $C > 0$ depending on (d, η, γ) .

Next, we are interested in counting how many times $\mathcal{E}(I_w)$ does not hold for $I_w \in \Sigma_{j'}$ inside a given cube $I \in \Sigma_J$ with $J \leq j'$. For this, fix $\varepsilon \in (0, 1)$ and $\theta > 1$, as well as a Borel probability measure ν on $[0, 1]^d$. It is easily checked that by combining (49), the Markov inequality and the Fubini-Tonelli Theorem,

$$(50) \quad \begin{aligned} &\sum_{J \geq 1} \sum_{j' \geq \theta J} \mathbb{E} \left(\nu \left(\left\{ x \in [0, 1]^d : \mathcal{L}^d(\{y \in I_J(x) : \mathcal{E}(I_{j'}(y)) \text{ holds}\}) \geq 2^{-dJ}(1 - 2^{-d(j'-J)\varepsilon}) \right\} \right) \right) \\ &\leq \sum_{J \geq 1} \sum_{j' \geq \theta J} (1 - 2^{-d(j'-J)\varepsilon})^{-1} C2^{-d_j'(1-\gamma\eta)} < \infty. \end{aligned}$$

Now, for $q < 0$, let $H = \tau'_\mu(q)$ and take for ν the measure μ_H of Proposition 4.11. By the Borel-Cantelli Lemma and (50), with probability 1, for μ_H -almost every x , for J large enough, for all $j \geq \theta J$,

$$\mathcal{L}^d(\{y \in I_J(x) : \mathcal{E}(I_{j'}(y)) \text{ does not hold}\}) \geq 2^{-dJ} 2^{-d(j'-J)\varepsilon} = 2^{d(j'-J)(1-\varepsilon)} 2^{-dj'}.$$

Hence for every $j' \geq \theta J$, there exists a subset $\Sigma_{j'}(I_J(x))$ of $\Sigma_{j'}$ of cardinality at least $2^{\lfloor d(j'-J)(1-\varepsilon) \rfloor}$ such that for all $w \in \Sigma_{j'}(I_J(x))$, $I_w \subset I_J(x)$ and $\mathcal{E}(I_w)$ does not hold.

Take $\theta = (1 + \varrho^{-1})/2$, $\varepsilon \in (0, 1 \wedge (\eta^{-1} - 1)/2)$, and $\gamma = 1/\eta - \varepsilon$. Remark that with this choice of parameter, if $J = \lfloor \varrho j \rfloor$, then $j \geq \varrho^{-1}J > \theta j$, so the previous conclusion holds for $j' = j$.

Combining what just precedes with the properties of μ_H already used in the case $q \geq 0$, one obtains that for all J large enough of the form $J = \lfloor \varrho j \rfloor$, there exists:

- a subset $\Sigma_J(H, \varepsilon)$ of Σ_J of cardinality at least $2^{J(\tau_\mu^*(q)-\varepsilon)}$ such that for all $u \in \Sigma_J(H, \varepsilon)$ one has $\mu(I_u) \leq 2^{-J(H-\varepsilon)}$,
- for all $u \in \Sigma_J(H, \varepsilon)$, there is a subset $\Sigma_j(I_u)$ of Σ_j of cardinality at least $2^{\lfloor d(j-J)(1-\varepsilon) \rfloor}$ such that for all $w \in \Sigma_j(I_u)$, $I_w \subset I_u$ and $\mathcal{E}(I_w)$ does not hold.

Hence for each $w \in \Sigma_j(I_u)$, the dyadic cubes I_v contained in $3I_w$ such that $p_v = 1$ are of generation at least $\lfloor(\eta^{-1} - \varepsilon)j\rfloor$. For such surviving vertices v , their generation $g(v)$ satisfies $\lfloor\varrho\eta g(v)\rfloor \geq \varrho\eta\lfloor(\eta^{-1} - \varepsilon)j\rfloor \geq j(\varrho - C'\varepsilon)$, for some constant C' depending on η and μ . So $I_v^{\varrho\eta} \subset$

$I_u^{(\rho-C'\varepsilon)/\varrho}$, and the doubling property (15) of μ together with (16) imply that $\mu(I_v^{\rho\eta}) \leq \mu(I_u)2^{C''j\varepsilon}$ for some constant C'' . Hence $\widetilde{M}_\varrho(I_w) \leq \mu(I_u)2^{C'j\varepsilon} \leq 2^{-J(H-\varepsilon)}2^{C''j\varepsilon}$.

Putting everything together, in the case where $q < 0$, one finally gets

$$(51) \quad \begin{aligned} \sum_{w \in \Sigma_j} \widetilde{M}_\varrho(I_w)^q &\geq \sum_{u \in \Sigma_J(H, \varepsilon)} \sum_{w \in \Sigma_j(I_u)} \widetilde{M}_\varrho(I_w)^q \\ &\geq 2^{J(\tau_\mu^*(q) - \varepsilon)} 2^{\lfloor d(j-J)(1-\varepsilon) \rfloor} (2^{-J(H-\varepsilon)} 2^{C''j\varepsilon})^q. \end{aligned}$$

Recalling that $J = \lfloor \varrho j \rfloor$, one deduces that $\tau_{M_\varrho}(q) \leq d(\varrho - 1) + \varrho(qH - \tau_\mu^*(H)) + O(\varepsilon) = d(\varrho - 1) + \varrho\tau_\mu(q) + O(\varepsilon)$. Since ε is arbitrary, this yields the desired upper bound, for all $q < 0$, almost surely.

Finally, τ_{M_ϱ} and τ_μ being continuous functions, all the previous inequalities relating these functions hold almost surely for all $q < 0$. \square

Next proposition entails the failure of the multifractal formalism over subsets of $\text{dom}(\sigma_{M_\varrho})$.

Proposition 5.13. *Recall the definition of $I_{\varrho, \eta}$ in Theorem 2.4. With probability 1, the multifractal formalism holds over $I_{\varrho, \eta}$ and it fails over $\text{dom}(\sigma_{M_\varrho}) \setminus I_{\varrho, \eta}$.*

Proof. It follows from Proposition 5.12 that, with probability 1,

$$(52) \quad \tau_{M_\varrho}^*(H) = \begin{cases} \eta d(1 - \varrho) + \eta \varrho \sigma_\mu(H/\varrho\eta) & \text{when } H \leq \varrho\eta H_\varrho, \\ \frac{\sigma_\mu(H_\varrho)}{\theta_\varrho(H_\varrho)} H & \text{when } \varrho\eta H_\varrho < H \leq \varrho H_\varrho, \\ d(1 - \varrho) + \varrho \sigma_\mu(H/\varrho) & \text{when } H > \varrho H_\varrho. \end{cases}$$

Case 1. One compares (52) with (6).

The two formulas coincide when $H_{\min} \leq H \leq \theta_\varrho(H_\varrho)$, and at $H = \varrho\tau_\mu'(0)$.

When $\theta_\varrho(H_\varrho) < H \leq \varrho H_\varrho$, $\tau_{M_\varrho}^*$ stays linear but σ_{M_ϱ} becomes strictly concave, so $\sigma_{M_\varrho} < \tau_{M_\varrho}^*$.

When $\varrho H_\varrho < H \leq \varrho\tau_\mu'(0)$, $H_\varrho \leq H/\varrho \leq \theta_\varrho^{-1}(H)$, and since the mapping (34) is decreasing on $[H_\varrho, H_{\max}]$, one has

$$\begin{aligned} \sigma_\mu(\theta_\varrho^{-1}(H)) &= \frac{d(1 - \varrho) + \varrho \sigma_\mu(\theta_\varrho^{-1}(H))}{\delta_\varrho(\theta_\varrho^{-1}(H))} = \frac{\theta_\varrho^{-1}(H)}{\delta_\varrho(\theta_\varrho^{-1}(H))} \frac{d(1 - \varrho) + \varrho \sigma_\mu(\theta_\varrho^{-1}(H))}{\theta_\varrho^{-1}(H)} \\ &< \frac{\theta_\varrho^{-1}(H)}{\delta_\varrho(\theta_\varrho^{-1}(H))} \frac{d(1 - \varrho) + \varrho \sigma_\mu(H/\varrho)}{H/\varrho}. \end{aligned}$$

But by definition of δ_ϱ , $\frac{\theta_\varrho^{-1}(H)}{\delta_\varrho(\theta_\varrho^{-1}(H))} = \frac{\theta_\varrho(\theta_\varrho^{-1}(H))}{\varrho} = H/\varrho$, so $\sigma_\mu(\theta_\varrho^{-1}(H)) < d(1 - \varrho) + \varrho \sigma_\mu(H/\varrho)$.

When $H > \varrho\tau_\mu'(0)$, by convexity $\sigma_\mu(H/\varrho) < d(1 - \varrho) + \varrho \sigma_\mu(H/\varrho)$.

Case 2.a. The formulas (52) with (7) do not coincide any more when $H_{\min} \leq H < \varrho\eta H_\varrho$, and the other regions are similar to Case 1.

Case 2.b. This case is simpler and follows from the previous ones. \square

REFERENCES

- [1] J. M. Aubry and S. Jaffard. Multifractal wavelet series. *Comm. Math. Phys.*, 227:483–514, 2002.
- [2] J. Barral and S. Seuret. Combining multifractal additive and multiplicative chaos. *Commun. Math. Phys.*, 257(2):473–497, 2005.
- [3] J. Barral and S. Seuret. From multifractal measures to multifractal wavelet series. *J. Fourier Anal. Appl.*, 11(5):589–614, 2005.
- [4] J. Barral and S. Seuret. Heterogeneous ubiquitous systems in \mathbb{R}^d and Hausdorff dimensions. *Bull. Brazilian Math. Soc.*, 38(3):467–515, 2007.
- [5] J. Barral and S. Seuret. Random sparse sampling on a Gibbs weighted tree and phase transitions. *J. Inst. Math. de Jussieu*, 19:65–116, 2020.

- [6] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, volume 470. Springer-Verlag, New York, 1975.
- [7] G. Brown, G. Michon, and J. Peyrière. On the multifractal analysis of measures. *J. Stat. Phys.*, 66:775–790, 1992.
- [8] P. Collet, J. L. Lebowitz, and A. Porzio. The dimension spectrum of some dynamical systems. *J. Stat. Phys.*, 47:609–644, 1987.
- [9] A. Dembo and O. Zeitouni. *Large deviations techniques and applications*, volume 38. Applications of Mathematics (New York). Springer-Verlag, New York, second edition, 2009.
- [10] Arnaud Durand. Random wavelet series based on a tree-indexed Markov chain. *Comm. Math. Phys.*, 283(2):451–477, 2008.
- [11] C. Esser and B. Védél. Counter-examples to the multifractal formalism. arXiv:2207.03733.
- [12] A.-H. Fan and D.-J. Feng. On the distribution of long-term time averages on symbolic space. *J. Stat. Phys.*, 99:813–856, 2000.
- [13] S. Jaffard. On lacunary wavelet series. *Ann. Appl. Probab.*, 10(1):313–329, 2000.
- [14] S. Jaffard. Wavelet techniques in multifractal analysis. In *Fractal Geometry and Applications: A Jubilee of Benoit Mandelbrot*, volume 72 of *Proc. Symposia in Pure Mathematics*, pages 91–151. AMS Providence, RI, 2004.
- [15] S. Jaffard and Y. Meyer. Wavelet methods for pointwise regularity and local oscillations of functions. *Mem. Amer. Math. Soc.*, 23:N. 587, 1996.
- [16] K.S. Lau and S.-M. Ngai. Multifractal measures and a weak separation condition. *Adv. Math.*, 141:45–96, 1999.
- [17] J. Lévy Véhel and R. Vojak. Multifractal analysis of Choquet capacities. *Adv. Appl. Math.*, 20:1–43, 1998.
- [18] Y. Meyer. *Ondelettes et opérateurs I*. Hermann, 1990.
- [19] L. Olsen. A multifractal formalism. *Adv. Math.*, 116:92–195, 1995.
- [20] Y. Pesin. *Dimension theory in dynamical systems*. Contemporary views and applications. University of Chicago Press, 1997.
- [21] D. Rand. The singularity spectrum $f(\alpha)$ for cookie-cutters. *Ergod. Th. Dynam. Sys.*, 9:527–541, 1989.
- [22] R. Riedi. An improved multifractal formalism and self-similar measures. *J. Math. Anal. Appl.*, 189:462–490, 1995.
- [23] D. Ruelle. *Thermodynamic formalism. The mathematical structures of classical equilibrium statistical mechanics*, volume 5 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley Publishing Co., Reading, Mass., 1978.

JULIEN BARRAL, LABORATOIRE D'ANALYSE, GÉOMÉTRIE ET APPLICATIONS, CNRS (UMR 7539), UNIVERSITÉ SORBONNE PARIS NORD, F-93430 VILLETANEUSE, FRANCE
Email address: barral@math.univ-paris13.fr

STÉPHANE SEURET, UNIV PARIS EST CRETEIL, UNIV GUSTAVE EIFFEL, CNRS, LAMA UMR8050, F-94010 CRETEIL, FRANCE
Email address: seuret@u-pec.fr