

THE NUMBER AND LOCATION OF EIGENVALUES FOR THE TWO-PARTICLE SCHRÖDINGER OPERATORS ON LATTICES

SAIDAKHMAT N. LAKAEV, MUKHAYYO O. AKHMADOVA

ABSTRACT. We study the Schrödinger operators $H_{\gamma\lambda\mu}(K)$, $K \in \mathbb{T}$ being a fixed (quasi)momentum of the particles pair, associated with a system of two identical bosons on the one-dimensional lattice \mathbb{Z} , where the real quantities γ , λ and μ describe the interactions between pairs of particles on one site, two nearest neighboring sites and next two neighboring sites, respectively. We found a partition of the three-dimensional space (γ, λ, μ) of interaction parameters into connected components and the exact number of eigenvalues of this operator that lie below and above the essential spectrum, in each component. Moreover, we show that for any $K \in \mathbb{T}^d$ the number of eigenvalues of $H_{\gamma\lambda\mu}(K)$ is not less than the corresponding number of eigenvalues of $H_{\gamma\lambda\mu}(0)$.

1. INTRODUCTION

Lattice models of physical systems are one of the widely used mathematical models in mathematical physics. Few-body Hamiltonians [29], among such models may be viewed as the simplest version of the corresponding Bose-Hubbard model involving a finite number of particles of a certain type. The few-body lattice Hamiltonians have been intensively studied over the past several decades [2, 3, 4, 5, 10, 11, 17, 19, 20, 21, 22, 23, 27, 25, 30].

Another reason for studying the discrete Hamiltonians is that they can provide a natural approximation for their continuous counterparts [9], which allows studying few-body systems. It is well known that the Efimov effect [8] was originally attributed to the three-body systems moving in the three-dimensional continuous space \mathbb{R}^3 . A rigorous mathematical proof of the Efimov effect has been given in [32, 35, 36, 39]. The celebrated Efimov phenomenon to take place also in the lattice three-particle systems [2, 4, 7, 19]. Discrete Schrödinger operators also represent a simple model for description of few-body systems formed by particles traveling through periodic structures, such as ultracold atoms injected into optical crystals created by the interference of counter-propagating laser beams [6, 38]. The study of ultracold few-atom systems in optical lattices have become popular in the last years due to availability of controllable parameters, such as temperature, particle masses, interaction potentials etc. (see e.g., [6, 13, 14, 28] and references therein). These possibilities give an opportunity to experimentally observe stable repulsively bound pairs of ultracold atoms ([31, 38], which is not the case for the traditional condensed matter systems, where stable composite objects are formed by means attractive forces. Lattice Hamiltonians are of particular interest in fusion physics too. For example, in [30], a one-dimensional one-particle lattice Hamiltonian has been successfully employed to show that certain class of molecules in lattice structures may enhance nuclear fusion probability.

In the continuous case, the center-of-mass motion can be separated, which is not the case for lattice few-body problems. However, for the lattice Hamiltonian H acting in the functional Hilbert

Date: April 25, 2023.

Key words and phrases. Two-particle system, discrete Schrödinger operator, essential spectrum, bound states, Fredholm determinant.

space $\mathbb{T}^{n \cdot d}$ we have a von Neumann direct integral decomposition

$$(1.1) \quad \mathbb{H} \simeq \int_{K \in \mathbb{T}^d}^{\oplus} H(K) dK,$$

where \mathbb{T}^d is the d -dimensional torus. The so called fiber Hamiltonians $H(K)$ acting on the Hilbert space $\mathbb{T}^{(n-1)d}$ nontrivially depends on the quasimomentum $K \in \mathbb{T}^d$ (see e.g., [2, 3, 33]). This decomposition allows us to reduce the problem to studying the operators $H(K)$.

In this work, we study the spectral properties of the fiber Hamiltonians $H(K)$, $K \in \mathbb{T}$ acting in the Hilbert space $L^{2,e}(\mathbb{T})$ as

$$(1.2) \quad H_{\gamma\lambda\mu}(K) := H_0(K) + V_{\gamma\lambda\mu},$$

where $H_0(K)$ is the fiber kinetic-energy operator,

$$(H_0(K)f)(p) = \mathcal{E}_K(p)f(p),$$

with

$$(1.3) \quad \mathcal{E}_K(p) := 2(1 - \cos \frac{K}{2} \cos p)$$

and $V_{\gamma\lambda\mu}$ is the combined interaction potential. The parameters γ , λ and μ , called coupling constants, describe interactions between the particles which are located on one site, on the nearest neighboring sites of the lattice and in the next nearest neighboring sites, respectively.

Within this new model, we find both the exact number of eigenvalues and their locations of the operator $H_{\gamma\lambda\mu}(0)$. We describe the mechanisms of emission and absorption of the eigenvalues of $H_{\gamma\lambda\mu}(0)$ at the thresholds of its essential (continuous) spectrum depending on the interaction parameters. Furthermore, we establish sharp lower bounds for the number of isolated eigenvalues $H_{\gamma\lambda\mu}(K)$ depending on the quasi-momentum $K \in \mathbb{T}$, which lie both below the essential spectrum and above that.

For this, we apply the results obtained for the operator $H_{\gamma\lambda\mu}(0)$ and the nontrivial dependence of the dispersion relation \mathcal{E}_K on the (quasi)momentum $K \in \mathbb{T}^d$.

We recall that the two-particle Schrödinger operator $H_\mu(K) = -\Delta + \mu V$, $\mu > 0$ on the lattice \mathbb{Z}^d associated to a system of two bosons with zero-range repulsive interactions $\mu > 0$ has been considered as a theoretical basis for explanation of the experimental results obtained in [31, 38].

Note that the continuous counterpart of the two-particle Schrödinger operators on lattices, which associated with a system of two quantum-mechanical particles on \mathbb{R}^d interacting via short-range potential $v(x)$ has isolated eigenvalues lying only below the essential spectrum fulfilling the semi-axis $[0, +\infty)$ and hence this model is well adapted to describe systems of two-particles with attractive interactions.

To study the eigenvalues of the discrete Schrödinger operator $H_{\gamma\lambda\mu}(K)$, we apply analytic function theory, namely, we investigate the corresponding Fredholm determinant $\Delta_{\gamma\lambda\mu}(K, z)$, as there is a one-to-one mapping between the sets of eigenvalues of the operator $H_{\gamma\lambda\mu}(K)$ and the zeros of $\Delta_{\gamma\lambda\mu}(K, z)$ (see [23]). Correspondingly, the change in the number of zeros of Fredholm determinant $\Delta_{\gamma\lambda\mu}(0, z)$ results in the change of the number of isolated eigenvalues of Schrödinger operator $H_{\gamma\lambda\mu}(0)$.

Our main finding is that the number of zeros of the determinant $\Delta_{\gamma\lambda\mu}(0, z)$ located below (resp. above) the essential spectrum changes if and only if the principal term $C^-(\gamma, \lambda, \mu)$ (resp. $C^+(\gamma, \lambda, \mu)$) of the asymptotics of the Fredholm determinant $\Delta_{\gamma\lambda\mu}(0, z)$ vanishes as z approaches the lower (resp. upper) threshold of the essential spectrum (see Lemma 4.3).

Using this property, we establish a partition of the three-dimensional (γ, λ, μ) -space into four disjoint connected components by means of curves $C^-(\gamma, \lambda, \mu) = 0$ or $C^+(\gamma, \lambda, \mu) = 0$. This allows us to prove that the number of zeros of the Fredholm determinant is constant in each connected component.

In [23, 25], it was studied the Schrödinger operators on the lattice \mathbb{Z}^d , $d = 1, 2$, associated to a system of two bosons with the zero-range on one site interaction ($\lambda \in \mathbb{R}$) and interactions on the nearest neighboring sites ($\mu \in \mathbb{R}$) of the d -dimensional lattice \mathbb{Z}^d . We emphasize that, our results is an extensions of [23, 25]. The authors of [23] consider the Schrödinger operators $H_{\lambda\mu}(K)$ on two-dimensional lattice \mathbb{Z}^2 . The operator $H_{\lambda\mu}(K)$ can have one or two eigenvalues, lying as below the essential spectrum, as well as above it. The connected components, which split the two-dimensional plane \mathbb{R}^2 of interaction parameters are described by means of a second order elementary curves (hyperbolas). Similar results for the number of eigenvalues of one-particle Schrödinger operators in \mathbb{Z}^d , $d \geq 1$ have been obtained, for instance in [22] with attractive interactions and $d = 3$ and also with attractive and repulsive interaction cases in [11] for all $d \geq 1$ considering only negative eigenvalues.

In the present work, the connected components are described by implicit functions (third-order polynomials) of three variables defined on \mathbb{R}^3 . This is a more challenging case as the connected components are described by the third-order surfaces.

The discrete two-particle Schrödinger operator $H_\mu(k)$ associated to a system of one and two quantum-mechanical particles on \mathbb{Z}^d interacting via short-range potentials have been studied in recent years [3, 5, 18, 20, 30].

Note that some results such as the existence of eigenvalues and their finiteness can be received for a large class of Schrödinger type operators (see e.g., [15, 16, 18, 24]). However, our results show that the study of a qualitative change in the number of eigenvalues of $H_{\gamma\mu\lambda}(K)$, even for $K = 0$, is very delicate: There is ball, with arbitrarily small radius, in the three-dimensional (γ, λ, μ) -space in which the number of eigenvalues has a jump (see Theorem 3.6).

The paper is organized as follows: The section 1 is introduction. In Section 2, we introduce the two-particle Hamiltonian $\mathbb{H}_{\gamma\lambda\mu}$ of a system of two bosons in the position and momentum space representations and also the Schrödinger operator $H_{\gamma\lambda\mu}(K)$ associated to the Hamiltonian $\mathbb{H}_{\gamma\lambda\mu}$. The main results of the paper are stated in Section 3 and their proofs are contained in Section 4.

2. THE TWO-PARTICLE HAMILTONIAN

2.1. The position-space representation. Let \mathbb{Z} be the one-dimensional lattice and $\mathbb{Z} \times \mathbb{Z}$ be cartesian square of \mathbb{Z} . Let $\ell^{2,s}(\mathbb{Z} \times \mathbb{Z})$ be the Hilbert space of square-summable symmetric functions on $\mathbb{Z} \times \mathbb{Z}$.

The free Hamiltonian $\widehat{\mathbb{H}}_0$ of a system of two identical particles (bosons), in the position space representation, is usually associated with the following self-adjoint (bounded) Toeplitz-type operator on the Hilbert space $\ell^{2,s}(\mathbb{Z} \times \mathbb{Z})$:

$$(\widehat{\mathbb{H}}_0 \widehat{f})(x_1, x_2) = \sum_{s_1 \in \mathbb{Z}} \widehat{\varepsilon}(x_1 - s_1) \widehat{f}(s_1, x_2) + \sum_{s_2 \in \mathbb{Z}} \widehat{\varepsilon}(x_2 - s_2) \widehat{f}(x_1, s_2), \quad \widehat{f} \in \ell^{2,s}(\mathbb{Z} \times \mathbb{Z})$$

where

$$(2.1) \quad \widehat{\varepsilon}(s) = \begin{cases} 2 & \text{if } |s| = 0, \\ -\frac{1}{2} & \text{if } |s| = 1, \\ 0 & \text{if } |s| > 1. \end{cases}$$

The interaction operator $\widehat{\mathbb{V}}_{\gamma\lambda\mu}$, in the position space representation, is the multiplication operator by the function $\widehat{v} \in \ell^1(\mathbb{Z})$, *i.e.*,

$$\widehat{\mathbb{V}}_{\gamma\lambda\mu}\widehat{f}(x, y) = \widehat{v}_{\gamma\lambda\mu}(x - y)\widehat{f}(x, y), \quad \widehat{f} \in \ell^{2,s}(\mathbb{Z} \times \mathbb{Z}),$$

where

$$(2.2) \quad \widehat{v}_{\gamma\lambda\mu}(s) = \begin{cases} 2\gamma & \text{if } |s| = 0, \\ \lambda & \text{if } |s| = 1, \\ \mu & \text{if } |s| = 2, \\ 0 & \text{if } |s| > 2. \end{cases}$$

The total Hamiltonian $\widehat{\mathbb{H}}_{\gamma\lambda\mu}$ of a system of two identical particles is described as a bounded self-adjoint operator on $\ell^{2,s}(\mathbb{Z} \times \mathbb{Z})$:

$$\widehat{\mathbb{H}}_{\gamma\lambda\mu} = \widehat{\mathbb{H}}_0 + \widehat{\mathbb{V}}_{\gamma\lambda\mu}, \quad \gamma, \lambda, \mu \in \mathbb{R}.$$

2.2. The two-particle Hamiltonian: the (quasi)momentum-space representation. Let $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z} \equiv (-\pi, \pi]$ be the one-dimensional torus, the Pontryagin dual group of \mathbb{Z} . Let $L^2(\mathbb{T})$ be the Hilbert space of square-integrable functions on \mathbb{T} and let

$$\mathcal{F} : \ell^2(\mathbb{Z}) \rightarrow L^2(\mathbb{T}), \quad \mathcal{F}\widehat{f}(p) = \frac{1}{2\pi} \sum_{x \in \mathbb{Z}} \widehat{f}(x)e^{ipx}.$$

is the standard Fourier transform with the inverse

$$\mathcal{F}^* : L^2(\mathbb{T}) \rightarrow \ell^2(\mathbb{Z}), \quad \mathcal{F}^*f(p) = \frac{1}{2\pi} \int_{\mathbb{T}} f(x)e^{ipx} dx.$$

The free Hamiltonian $\mathbb{H}_0 = (\mathcal{F} \otimes \mathcal{F})\widehat{\mathbb{H}}_0(\mathcal{F}^* \otimes \mathcal{F}^*)$ of a system of two identical particles, in the momentum space representation, is the multiplication operator by the function $\varepsilon(p) + \varepsilon(q)$ in the Hilbert space $L^{2,s}(\mathbb{T} \times \mathbb{T})$ of symmetric functions on the cartesian square $\mathbb{T} \times \mathbb{T}$ of the torus \mathbb{T} :

$$\mathbb{H}_0 f(p, q) = [\varepsilon(p) + \varepsilon(q)]f(p, q),$$

where the continuous function (dispersion relation) ε is given by

$$\varepsilon(p) = [\mathcal{F}\widehat{\varepsilon}](p) = 1 - \cos p, \quad p \in \mathbb{T}.$$

The interaction operator $\mathbb{V}_{\gamma\lambda\mu} = (\mathcal{F} \otimes \mathcal{F})\widehat{\mathbb{V}}_{\gamma\lambda\mu}(\mathcal{F}^* \otimes \mathcal{F}^*)$ is the integral operator of convolution type acting in $L^{2,s}(\mathbb{T} \times \mathbb{T})$ as

$$(\mathbb{V}_{\gamma\lambda\mu}f)(p, q) = \frac{1}{2\pi} \int_{\mathbb{T}} v_{\gamma\lambda\mu}(p - u)f(u, p + q - u)du,$$

where the kernel function $v_{\gamma\lambda\mu}(\cdot)$ is given by

$$v_{\gamma\lambda\mu}(p) = [\mathcal{F}\widehat{v}_{\gamma\lambda\mu}](p) = \frac{1}{2\pi} \sum_{x \in \mathbb{Z}} \widehat{v}_{\gamma\lambda\mu}(x)e^{ipx} = \frac{1}{\pi}(\gamma + \lambda \cos p + \mu \cos 2p), \quad p \in \mathbb{T}.$$

The total two-particle Hamiltonian $\mathbb{H}_{\gamma\lambda\mu}$ of a system of two identical quantum-mechanical particles interacting via a finite range attractive potentials $\widehat{v}_{\gamma\lambda\mu}$, in the momentum space representation, is the bounded self-adjoint operator acting in $L^{2,s}(\mathbb{T} \times \mathbb{T})$ as

$$\mathbb{H}_{\gamma\lambda\mu} := (\mathcal{F} \otimes \mathcal{F})\widehat{\mathbb{H}}_{\gamma\lambda\mu}(\mathcal{F} \otimes \mathcal{F})^* = \mathbb{H}_0 + \mathbb{V}_{\gamma\lambda\mu}.$$

2.3. The Floquet-Bloch decomposition of $\mathbb{H}_{\gamma\lambda\mu}$ and discrete Schrödinger operators. Since the operator $\widehat{H}_{\gamma\lambda\mu}$ commutes with the shift operators on the lattice $\mathbb{Z} \times \mathbb{Z}$, we can decompose the space $L^{2,s}(\mathbb{T} \times \mathbb{T})$ and Hamiltonian $\mathbb{H}_{\gamma\lambda\mu}$ into the von Neumann direct integrals as

$$(2.3) \quad L^{2,s}(\mathbb{T} \times \mathbb{T}) \simeq \int_{K \in \mathbb{T}}^{\oplus} L^{2,e}(\mathbb{T}) \, dK \quad \text{and} \quad \mathbb{H}_{\gamma\lambda\mu} \simeq \int_{K \in \mathbb{T}}^{\oplus} H_{\gamma\lambda\mu}(K) \, dK,$$

respectively, where $L^{2,e}(\mathbb{T})$ is the Hilbert space of square-integrable even functions on \mathbb{T} (see, e.g., [3]).

The fiber operator $H_{\gamma\lambda\mu}(K)$, $K \in \mathbb{T}$ is a self-adjoint operator defined in $L^{2,e}(\mathbb{T})$ as

$$H_{\gamma\lambda\mu}(K) := H_0(K) + V_{\gamma\lambda\mu},$$

where the unperturbed operator $H_0(K)$ is the multiplication operator by the function

$$\mathcal{E}_K(p) := 2(1 - \cos \frac{K}{2} \cos p)$$

and the perturbation $V_{\gamma\lambda\mu}$ is defined as

$$V_{\gamma\lambda\mu}f(p) = \frac{1}{\pi} \int_{\mathbb{T}} (\gamma + \lambda \cos p \cos q + \mu \cos 2p \cos 2q) f(q) \, dq.$$

In some literature, the parameter K is called *quasi-momentum* and the fiber $H_{\gamma\lambda\mu}(K)$ is called *discrete Schrödinger operator* associated to the two-particle Hamiltonian $\mathbb{H}_{\gamma\lambda\mu}$.

Using the Fourier transform, $\ell^{2,s}(\mathbb{Z} \times \mathbb{Z})$ and $\mathbb{H}_{\gamma\lambda\mu}$ can also be rewritten as decompositions

$$\ell^{2,s}(\mathbb{Z} \times \mathbb{Z}) \simeq \int_{K \in \mathbb{T}}^{\oplus} \ell^{2,e}(\mathbb{Z}) \, dK \quad \text{and} \quad \mathbb{H}_{\gamma\lambda\mu} \simeq \int_{K \in \mathbb{T}}^{\oplus} H_{\gamma\lambda\mu}(K) \, dK,$$

respectively, where $\ell^{2,e}(\mathbb{Z})$ is the Hilbert space of square-summable even functions on \mathbb{Z} and

$$(2.4) \quad \widehat{H}_{\gamma\lambda\mu}(K) := \mathcal{F}^* H_{\gamma\lambda\mu}(K) \mathcal{F} = \widehat{H}_0(K) + \widehat{V}_{\gamma\lambda\mu},$$

and

$$\widehat{H}_0(K) \widehat{f}(x) = \sum_{s \in \mathbb{Z}} \widehat{\mathcal{E}}_K(x-s) f(s), \quad \widehat{f} \in \ell^{2,e}(\mathbb{Z}),$$

with

$$\widehat{\mathcal{E}}_K(x) = 2\widehat{\mathcal{E}}(x) \cos \frac{Kx}{2}$$

and the operator $\widehat{V}_{\gamma\lambda\mu}$ acts in $\ell^{2,e}(\mathbb{Z})$ as

$$\widehat{V}_{\gamma\lambda\mu} \widehat{f}(x) = \widehat{v}_{\gamma\lambda\mu}(x) \widehat{f}(x).$$

2.4. The essential spectrum of discrete Schrödinger operators. Since $V_{\gamma\lambda\mu}$ is a finite rank operator, by well known Weyl's Theorem (see [34, Theorem XIII.14]) for any $K \in \mathbb{T}$ the essential spectrum $\sigma_{\text{ess}}(H_{\gamma\lambda\mu}(K))$ of $H_{\gamma\lambda\mu}$ coincides with the spectrum of $H_0(K)$, i.e.,

$$(2.5) \quad \sigma_{\text{ess}}(H_{\gamma\lambda\mu}(K)) = \sigma(H_0(K)) = [\mathcal{E}_{\min}(K), \mathcal{E}_{\max}(K)],$$

where

$$\begin{aligned} \mathcal{E}_{\min}(K) &:= \min_{p \in \mathbb{T}} \mathcal{E}_K(p) = 2(1 - \cos \frac{K}{2}) \geq 0 = \mathcal{E}_{\min}(0), \\ \mathcal{E}_{\max}(K) &:= \max_{p \in \mathbb{T}} \mathcal{E}_K(p) = 2(1 + \cos \frac{K}{2}) \leq 4 = \mathcal{E}_{\max}(0). \end{aligned}$$

3. MAIN RESULTS

Let $K \in \mathbb{T}$ and $n_+(H_{\gamma\lambda\mu}(K))$ resp. $n_-(H_{\gamma\lambda\mu}(K))$ be the number of eigenvalues of the operator $H_{\gamma\lambda\mu}(K)$ above resp. below its essential spectrum.

Our first main result is a generalization of Theorem 2 in [3].

Theorem 3.1. *Suppose that $H_{\gamma\lambda\mu}(0)$ has n eigenvalues below resp. above the essential spectrum for some $\gamma, \lambda, \mu \in \mathbb{R}$. Then for each $K \in \mathbb{T}$ the operator $H_{\gamma\lambda\mu}(K)$ has at least n eigenvalues below resp. above its essential spectrum. In other words,*

$$n_-(H_{\gamma\lambda\mu}(K)) \geq n_-(H_{\gamma\lambda\mu}(0))$$

and

$$n_+(H_{\gamma\lambda\mu}(K)) \geq n_+(H_{\gamma\lambda\mu}(0)).$$

Let us consider the cubic polynomial C^\pm of three-variable defined by

$$(3.1) \quad C^\pm(\gamma, \lambda, \mu) := \mp(\gamma + \lambda + \mu + \gamma\lambda\mu) + \gamma\lambda + 2\gamma\mu + \lambda\mu.$$

Lemma 3.2. *The set of points \mathbb{R}^3 satisfying the equation $C^\pm(\gamma, \lambda, \mu) = 0$ coincides with the graph of function*

$$\gamma^\pm(\lambda, \mu) = -\frac{Q_0^\pm(\lambda, \mu)}{Q_1^\pm(\lambda, \mu)},$$

where $Q_0^\pm(\lambda, \mu)$ and $Q_1^\pm(\lambda, \mu)$ are defined as

$$(3.2) \quad \begin{aligned} Q_0^\pm(\lambda, \mu) &= \mp(\lambda + \mu) + \lambda\mu, \\ Q_1^\pm(\lambda, \mu) &= \mp(\lambda + 2\mu) + 1 + \lambda\mu. \end{aligned}$$

Proof. It is sufficient to show that the following system of equations has no solutions

$$(3.3) \quad \begin{cases} Q_1^\pm(\lambda, \mu) = 0 \\ C^\pm(\gamma, \lambda, \mu) = 0. \end{cases}$$

System (3.3) obeys that

$$(3.4) \quad \begin{cases} Q_1^\pm(\lambda, \mu) = 0 \\ Q_0^\pm(\lambda, \mu) = 0 \end{cases} \Rightarrow \begin{cases} \mu = 0 \\ \mu = \pm 1. \end{cases}$$

The last system has no solutions. Therefore, (3.3) hasn't any solutions too. It leads to the proof of Lemma 3.2. \square

Lemma 3.3. *The set of points of \mathbb{R}^2 satisfying the equation $Q_1^\pm(\lambda, \mu) = 0$ coincides with the graph of the function*

$$(3.5) \quad \lambda^\pm(\mu) = -\frac{2\mu \mp 1}{1 \mp \mu}.$$

Proof. The following system of equations

$$(3.6) \quad \begin{cases} 1 \mp \mu = 0 \\ Q_1^\pm(\lambda, \mu) = 0 \end{cases}$$

is equivalent to (3.4), which has no solutions, thus, it hasn't any solutions. It yields the proof of Lemma 3.3. \square

The straight lines $\mu = -1$ and $\mu = 1$ separate the graph of the functions $\lambda^-(\cdot)$ and $\lambda^+(\cdot)$ on the (λ, μ) -plane into two (different) continuous curves $\{\tau_1^-, \tau_2^-\}$ and $\{\tau_1^+, \tau_2^+\}$:

$$\tau_1^- = \{(\lambda, \mu) \in \mathbb{R}^2 : \lambda = -\frac{1+2\mu}{1+\mu}, \mu > -1\},$$

$$\tau_2^- = \{(\lambda, \mu) \in \mathbb{R}^2 : \lambda = -\frac{1+2\mu}{1+\mu}, \mu < -1\}$$

and

$$\tau_1^+ = \{(\lambda, \mu) \in \mathbb{R}^2 : \lambda = \frac{1-2\mu}{1-\mu}, \mu < 1\},$$

$$\tau_2^+ = \{(\lambda, \mu) \in \mathbb{R}^2 : \lambda = \frac{1-2\mu}{1-\mu}, \mu > 1\}.$$

The curves $\{\tau_1^-, \tau_2^-\}$ and $\{\tau_1^+, \tau_2^+\}$ divide the (λ, μ) - plane into the domains D_1^-, D_2^-, D_3^- and D_1^+, D_2^+, D_3^+ of the functions $\lambda^-(\cdot)$ and $\lambda^+(\cdot)$, respectively:

$$(3.7) \quad D_1^- = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^-(\lambda, \mu) > 0, \mu > -1\},$$

$$D_2^- = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^-(\lambda, \mu) < 0\},$$

$$D_3^- = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^-(\lambda, \mu) > 0, \mu < -1\}$$

and

$$(3.8) \quad D_1^+ = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^+(\lambda, \mu) > 0, \mu < 1\},$$

$$D_2^+ = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^+(\lambda, \mu) < 0\},$$

$$D_3^+ = \{(\lambda, \mu) \in \mathbb{R}^2 : Q_1^+(\lambda, \mu) > 0, \mu > 1\}.$$

The following lemma summarizes the locations of the domains $D_\alpha^\pm, \alpha = 1, 2, 3$ defined in (3.7) and (3.8) and also their relations.

Lemma 3.4. *The followings are true:*

- (i) $D_3^- \subseteq D_1^+$,
- (ii) $D_3^+ \subseteq D_1^-$,
- (iii) $D_3^- \cap D_3^+ = \emptyset$,
- (iv) *For any $\alpha = 1, 2, 3$ the regions D_α^- and D_α^+ are symmetric with respect to origin.*

Proof. The definitions of $D_\alpha^\pm, \alpha = 1, 2, 3$ yield the proofs of items (i)-(iii) of Lemma 3.4 and the equality $Q_1^+(\lambda, \mu) = Q_1^-(-\lambda, -\mu)$ yields (iv) (see Figure 1). \square

Recall that, the domain of the function $\gamma^\pm(\cdot, \cdot)$ is an open set

$$\mathbb{R}^2 \setminus (\tau_1^\pm \cup \tau_2^\pm) = D_1^\pm \cup D_2^\pm \cup D_3^\pm.$$

The curves τ_1^\pm and τ_2^\pm in \mathbb{R}^2 are define the corresponding surfaces $\Upsilon_j^\pm \subset \mathbb{R}^3, j = 1, 2$:

$$\Upsilon_j^\pm := \{(\gamma, \lambda, \mu) \in \mathbb{R}^3, (\lambda, \mu) \in \tau_j^\pm\}.$$

Further, the surfaces Υ_1^\pm and Υ_2^\pm will separate the graph of the function $\gamma^\pm(\cdot, \cdot)$ into three different continuous (connected) surfaces $\Gamma_1^\pm, \Gamma_2^\pm$ and Γ_3^\pm in \mathbb{R}^3 :

$$\Gamma_j^\pm = \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : \gamma = -\frac{Q_0^\pm(\lambda, \mu)}{Q_1^\pm(\lambda, \mu)}, (\lambda, \mu) \in D_j^\pm\}, j = 1, 2, 3.$$




FIGURE 1. The domains $D_\alpha^\pm, \alpha = 1, 2, 3$ of the principle functions γ^\pm in the (λ, μ) -plane of the parameters $\lambda, \mu \in \mathbb{R}$.

The surfaces $\Gamma_1^-, \Gamma_2^-, \Gamma_3^-$ and $\Gamma_1^+, \Gamma_2^+, \Gamma_3^+$ divides the three dimensional space \mathbb{R}^3 into four separated connected components $\mathbb{G}_0^-, \mathbb{G}_1^-, \mathbb{G}_2^-, \mathbb{G}_3^-$ and $\mathbb{G}_0^+, \mathbb{G}_1^+, \mathbb{G}_2^+, \mathbb{G}_3^+$ respectively:

$$(3.9) \quad \begin{aligned} \mathbb{G}_0^- &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^-(\gamma, \lambda, \mu) > 0, (\lambda, \mu) \in D_1^-\}, \\ \mathbb{G}_1^- &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^-(\gamma, \lambda, \mu) < 0, (\lambda, \mu) \in \overline{D_1^-} \cup D_2^-\}, \\ \mathbb{G}_2^- &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^-(\gamma, \lambda, \mu) > 0, (\lambda, \mu) \in \overline{D_2^-} \cup D_3^-\}, \\ \mathbb{G}_3^- &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^-(\gamma, \lambda, \mu) < 0, (\lambda, \mu) \in D_3^-\} \end{aligned}$$

and

$$(3.10) \quad \begin{aligned} \mathbb{G}_0^+ &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^+(\gamma, \lambda, \mu) > 0, (\lambda, \mu) \in D_1^+\}, \\ \mathbb{G}_1^+ &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^+(\gamma, \lambda, \mu) < 0, (\lambda, \mu) \in \overline{D_1^+} \cup D_2^+\}, \\ \mathbb{G}_2^+ &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^+(\gamma, \lambda, \mu) > 0, (\lambda, \mu) \in \overline{D_2^+} \cup D_3^+\}, \\ \mathbb{G}_3^+ &:= \{(\gamma, \lambda, \mu) \in \mathbb{R}^3 : C^+(\gamma, \lambda, \mu) < 0, (\lambda, \mu) \in D_3^+\}. \end{aligned}$$

Theorem 3.5. *Let \mathcal{C} be one of the above connected components of the partition of the (γ, λ, μ) -space. Then for any $(\gamma, \lambda, \mu) \in \mathcal{C}$ the numbers $n_+(H_{\gamma\lambda\mu}(0))$ and $n_-(H_{\gamma\lambda\mu}(0))$ of eigenvalues of $H_{\gamma\lambda\mu}(0)$ lying, respectively, below and above the essential spectrum (2.5) remain constant.*

Now we study the number of eigenvalues of $H_{\gamma\lambda\mu}(0)$ in $(-\infty, 0)$ and $(4, +\infty)$ depending on the potential parameters γ, λ and μ .

Theorem 3.6. *For any $\alpha = 0, 1, 2, 3$ the following statements are true:*

- (i) *if $(\gamma, \lambda, \mu) \in \mathbb{G}_\alpha^-$, then $n_-(H_{\gamma\lambda\mu}(0)) = \alpha$;*
- (ii) *if $(\gamma, \lambda, \mu) \in \mathbb{G}_\alpha^+$, then $n_+(H_{\gamma\lambda\mu}(0)) = \alpha$.*

In the following lemma the locations of the connected components

$$(3.11) \quad \mathcal{C}_{\alpha\beta} = \mathbb{G}_\alpha^- \cap \mathbb{G}_\beta^+, \quad \alpha, \beta = 0, 1, 2, 3$$

are described, which divide the three dimensional space \mathbb{R}^3 into a disjoint parts.

Lemma 3.7. *The following relations are true:*

- (i) $\mathcal{C}_{30} = \mathbb{G}_3^- \cap \mathbb{G}_0^+ = \mathbb{G}_3^-$;
- (ii) $\mathcal{C}_{03} = \mathbb{G}_0^- \cap \mathbb{G}_3^+ = \mathbb{G}_3^+$;
- (iii) *If $\alpha + \beta > 3$, then $\mathcal{C}_{\alpha\beta} = \mathbb{G}_\alpha^- \cap \mathbb{G}_\beta^+ = \emptyset$.*
- (iv) *If $\alpha + \beta \leq 3$, then $\mathcal{C}_{\alpha\beta}$ is symmetric to $\mathcal{C}_{\beta\alpha}$ with respect to origin.*

Proof. The assertions (i)-(iii) follow from the definition (3.9), Lemma 3.4 and several simple inequalities. Proof of the assertion (iv) is obtained by the equality $C^+(\gamma, \lambda, \mu) = C^-(-\gamma, -\lambda, -\mu)$. \square

Theorem 3.6 and definition (3.11) of connected components $\mathcal{C}_{\alpha\beta}$ yield the following corollary.

Corollary 3.8. *Let $\gamma, \lambda, \mu \in \mathbb{R}$. If $(\gamma, \lambda, \mu) \in \mathcal{C}_{\alpha\beta}$ then*

$$n_-(H_{\gamma\lambda\mu}(0)) = \alpha \quad \text{and} \quad n_+(H_{\gamma\lambda\mu}(0)) = \beta,$$

where $\alpha, \beta = 0, 1, 2, 3$ and $\alpha + \beta \leq 3$.

Recall that by the min-max principle $H_{\gamma\lambda\mu}(K)$ can have at most three eigenvalues outside its essential spectrum.

The following theorem provides the sharp lower bound for the number of eigenvalues lying outside the essential spectrum of $H_{\gamma\lambda\mu}(K)$ depending only on γ, λ and μ .

Theorem 3.9. *Let $K \in \mathbb{T}$ and $(\gamma, \lambda, \mu) \in \mathbb{R}^3$. For all $\alpha, \beta = 0, 1, 2, 3$ satisfying the condition $\alpha + \beta \leq 3$ the following relations are true :*

- (i) *if $(\gamma, \lambda, \mu) \in \mathcal{C}_{\alpha\beta}$ and $\alpha + \beta < 3$, then $n_-(H_{\gamma\lambda\mu}(K)) \geq \alpha$, $n_+(H_{\gamma\lambda\mu}(K)) \geq \beta$;*
- (ii) *if $(\gamma, \lambda, \mu) \in \mathcal{C}_{\alpha\beta}$ and $\alpha + \beta = 3$, then $n_-(H_{\gamma\lambda\mu}(K)) = \alpha$, $n_+(H_{\gamma\lambda\mu}(K)) = \beta$.*

4. PROOF OF THE MAIN RESULTS

4.1. The discrete spectrum of $H_{\gamma\lambda\mu}(0)$. In the case of $K = 0$, the Fredholm determinant $\Delta_{\gamma\lambda\mu}(0, z)$ is easier to study. Note that the essential spectrum of Hamiltonian $H_{\gamma\lambda\mu}(0)$ coincides with the segment $[0, 4]$.

We try to find an (implicit) equation for the discrete eigenvalues of $H_{\gamma\lambda\mu}(0)$, i.e., for the non-zero solutions of equation

$$H_{\gamma\lambda\mu}(0)f = zf$$

in $z \in \mathbb{R} \setminus [0, 4]$.

We apply the Fredholm's determinants method to study the number and location of eigenvalues (see, e.g., [1, 26]). The Fredholm determinant associated to $H_{\gamma\lambda\mu}(0)$ can be written as

$$(4.1) \quad \Delta_{\gamma\lambda\mu}(0; z) = \begin{vmatrix} 1 + \gamma a(z) & \lambda b(z) & \mu c(z) \\ \gamma b(z) & 1 + \lambda d(z) & \mu e(z) \\ \gamma c(z) & \lambda e(z) & 1 + \mu f(z) \end{vmatrix},$$

where

$$(4.2) \quad \begin{aligned} a(z) &:= \int_{\mathbb{T}} \frac{2dt}{\varepsilon(t) - z}, & b(z) &:= \int_{\mathbb{T}} \frac{2 \cos t dt}{\varepsilon(t) - z}, \\ c(z) &:= \int_{\mathbb{T}} \frac{2 \cos 2t dt}{\varepsilon(t) - z}, & d(z) &:= \int_{\mathbb{T}} \frac{2 \cos^2 t dt}{\varepsilon(t) - z}, \\ e(z) &:= \int_{\mathbb{T}} \frac{2 \cos t \cos 2t dt}{\varepsilon(t) - z}, & f(z) &:= \int_{\mathbb{T}} \frac{2 \cos^2 2t dt}{\varepsilon(t) - z}. \end{aligned}$$

Functions $a(\cdot)$, $d(\cdot)$ and $f(\cdot)$ are analytic in $\mathbb{R} \setminus [0, 4]$, strictly decreasing in $\mathbb{R} \setminus [0, 4]$, positive in the interval $(-\infty, 0)$ and negative in $(4, +\infty)$. The functions $b(\cdot)$, $c(\cdot)$ and $e(\cdot)$ are analytic in $\mathbb{R} \setminus [0, 4]$ too. Their behaviour near $z = 0$ and $z = 4$ are described in the following proposition.

Proposition 4.1. *The functions defined in (4.2) have the following asymptotics*

$$\begin{aligned}
a(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}, \\
b(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} - 1 + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + 1 + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}, \\
c(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} - 2 + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + 2 + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}, \\
d(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} - 1 + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + 1 + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}, \\
e(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} - 2 + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + 2 + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}, \\
f(z) &= \begin{cases} \frac{1}{(-z)^{\frac{1}{2}}} - 2 + O(-z)^{\frac{1}{2}}, & \text{as } z \nearrow 0 \\ -\frac{1}{(z-4)^{\frac{1}{2}}} + 2 + O(z-4)^{\frac{1}{2}}, & \text{as } z \searrow 4 \end{cases}.
\end{aligned}$$

Here $(-z)^{\frac{1}{2}}$ and $(z-4)^{\frac{1}{2}}$ are denote those branches of analytic functions that are positive for $-z > 0$ and $z-4 > 0$.

Proposition 4.1 can be proved as [21, Proposition 4].

Lemma 4.2. *For all $\gamma, \lambda, \mu \in \mathbb{R}$ the determinant $\Delta_{\gamma\lambda\mu}(0; z)$ has asymptotics*

$$(4.3) \quad \Delta_{\gamma\lambda\mu}(0; z) = \begin{cases} C^-(\gamma, \lambda, \mu)(-z)^{-\frac{1}{2}} + D^-(\gamma, \lambda, \mu) + O((-z)^{\frac{1}{2}}), & \text{as } z \nearrow 0, \\ C^+(\gamma, \lambda, \mu)(z-4)^{-\frac{1}{2}} + D^+(\gamma, \lambda, \mu) + O((z-4)^{\frac{1}{2}}), & \text{as } z \searrow 4, \end{cases}$$

where C^\pm is defined in (3.1) and

$$D^\pm(\gamma, \lambda, \mu) = 1 - (\gamma\lambda + 2\lambda\mu + 4\gamma\mu) \mp (\lambda + \mu + \gamma\lambda\mu).$$

The proof of Lemma 4.2 can be derived by using the asymptotics of functions $a(\cdot), b(\cdot), c(\cdot), d(\cdot), e(\cdot)$ and $f(\cdot)$ in Proposition 4.1.

Corollary 4.3. *For all $(\gamma, \lambda, \mu) \in \mathbb{R}^3$ the relations*

- (i). $\lim_{z \rightarrow -\infty} \Delta_{\gamma\lambda\mu}(0; z) = 1,$
- (ii). $\lim_{z \nearrow 0} \sqrt{-z} \Delta_{\gamma\lambda\mu}(0; z) = C^-(\gamma, \lambda, \mu),$
- (iii). $\lim_{z \searrow 4} \sqrt{z-4} \Delta_{\gamma\lambda\mu}(0; z) = C^+(\gamma, \lambda, \mu)$

hold.

Proof of Corollary 4.3. The first item follows from the Lebesgue dominated convergence theorem. Lemma 4.2 yields the proof of other items. \square

The next lemma describes a one-to-one correspondence between the eigenvalues $H_{\gamma\lambda\mu}(0)$ and the zeros of the Fredholm determinant $\Delta_{\gamma\lambda\mu}(0; z)$.

Lemma 4.4. *The number $z \in \mathbb{R} \setminus [0, 4]$ is an eigenvalue of $H_{\gamma\lambda\mu}(0)$ if and only if it is a zero of $\Delta_{\gamma\lambda\mu}(0; \cdot)$. Moreover, in $\mathbb{R} \setminus [0, 4]$ the function $\Delta_{\gamma\lambda\mu}(0; \cdot)$ has at most three zeros.*

Proof. The first assertion follows from the theory of Fredholm determinants (see, for example, [1]). Since the operator $H_{\gamma\lambda\mu}(0)$ has rank at most three, by the minimum-max principle it has at most three eigenvalues outside the essential spectrum. So, according to the first part of the proposition, $\Delta_{\gamma\lambda\mu}(0; \cdot)$ has at most three zeros in $\mathbb{R} \setminus [0, 4]$. \square

The following lemmas determine the number and arrangement of eigenvalues of the operator $H_{\gamma\lambda 0}(0)$ that lie below the essential spectrum.

Lemma 4.5. *Let $(\gamma, \lambda) \in \mathbb{R}^2$, then the following relations hold.*

- (i) *If $\gamma + \lambda + \gamma\lambda > 0$ and $\gamma + 1 > 0$, then $H_{\gamma\lambda 0}(0)$ has no eigenvalues in $(-\infty, 0)$.*
- (ii) *If $\gamma + \lambda + \gamma\lambda < 0$ or $\gamma + \lambda + \gamma\lambda = 0$ and $\gamma + 1 > 0$, then $H_{\gamma\lambda 0}(0)$ has one eigenvalue in $(-\infty, 0)$.*
- (iii) *If $\gamma + \lambda + \gamma\lambda \geq 0$ and $\gamma + 1 \leq 0$, then $H_{\gamma\lambda 0}(0)$ has two eigenvalues in $(-\infty, 0)$.*

The following lemma provides the dependence of the number of eigenvalues of the operator $H_{\gamma\lambda 0}(0)$ in $(4, +\infty)$ on γ and λ :

Lemma 4.6. *Let $(\gamma, \lambda) \in \mathbb{R}^2$, then the following relations hold.*

- (i) *If $-\gamma - \lambda + \gamma\lambda > 0$ and $\gamma - 1 < 0$, then $H_{\gamma\lambda 0}(0)$ has no eigenvalues in $(4, +\infty)$.*
- (ii) *If $-\gamma - \lambda + \gamma\lambda < 0$ or $-\gamma - \lambda + \gamma\lambda = 0$ and $\gamma - 1 < 0$, then $H_{\gamma\lambda 0}(0)$ has one eigenvalue in $(4, +\infty)$.*
- (iii) *If $-\gamma - \lambda + \gamma\lambda \geq 0$ and $\gamma - 1 \geq 0$, then $H_{\gamma\lambda 0}(0)$ has two eigenvalues in $(4, +\infty)$.*

Lemmas 4.5 and 4.6 can be proved as in [25, Theorem 5.5].

Proof of Theorem 3.5. Let us assume, without loss of generality, that $C^-(\gamma, \lambda, \mu) < 0$ for all $(\gamma, \lambda, \mu) \in \mathcal{C}$. The definition of determinant and Lemma 4.3 yield

$$(4.4) \quad \lim_{z \rightarrow -\infty} \Delta_{\gamma\lambda\mu}(0, z) = 1, \quad \lim_{z \nearrow 0} \Delta_{\gamma\lambda\mu}(0, z) < 0.$$

Let $(\gamma_0, \lambda_0, \mu_0) \in \mathcal{C}$ be a point of \mathcal{C} and $z_0 < 0$ be a zero of the function $\Delta_{\gamma_0\lambda_0\mu_0}(z)$ of multiplicity $m \geq 1$. For each fixed $z < 0$ the determinant $\Delta_{\gamma\lambda\mu}(z)$ is a real analytic function in $(\gamma, \lambda, \mu) \in \mathcal{C}$ and for each fixed $\gamma, \lambda, \mu \in \mathbb{R}$ the function $\Delta_{\gamma\lambda\mu}(z)$ is real analytic in $z \in (-\infty, 0)$. Hence, for each $\varepsilon > 0$ there are numbers $\delta > 0$, $\eta > 0$ and an open neighborhood $W_\eta(z_0)$ of z_0 with radius η such that for all $z \in \overline{W_\eta(z_0)}$ and $(\gamma, \lambda, \mu) \in \mathcal{C}$ obeying the conditions $|z - z_0| = \eta$ and $\|(\gamma, \lambda, \mu) - (\gamma_0, \lambda_0, \mu_0)\| < \delta$ the following two inequalities $|\Delta_{\gamma_0\lambda_0\mu_0}(z)| > \eta$ and $|\Delta_{\gamma\lambda\mu}(z) - \Delta_{\gamma_0\lambda_0\mu_0}(z)| < \varepsilon$ hold. Then by Rouché's theorem the number of zeros of the function $\Delta_{\gamma\lambda\mu}(z)$ in $W_\eta(z_0)$ remains constant for all $(\gamma, \lambda, \mu) \in \mathcal{C}$ satisfying the inequality $\|(\gamma, \lambda, \mu) - (\gamma_0, \lambda_0, \mu_0)\| < \delta$. Since the root $z_0 < 0$ of the function $\Delta_{\gamma\lambda\mu}(z)$ is arbitrary in (B_1, B_2) we conclude that the number of its zeros remains constant in (B_1, B_2) for all $(\gamma, \lambda, \mu) \in \mathcal{C}$ satisfying $\|(\gamma, \lambda, \mu) - (\gamma_0, \lambda_0, \mu_0)\| < \delta$.

Further each Jordan curve $\Gamma \subset \mathcal{C}$ connecting any two points of \mathcal{C} is a *compact set*, so the number of zeros of the function $\Delta_{\gamma\lambda\mu}(z)$ lying below the bottom of the essential spectrum for any $(\gamma, \lambda, \mu) \in \Gamma$ remains constant. Therefore, Lemma 4.4 yields that the number of eigenvalues $n_-(H_{\gamma\lambda\mu}(0))$ of the operator $H_{\gamma\lambda\mu}(0)$ below the essential spectrum is constant.

The proof in the case of $n_+(H_{\gamma\lambda\mu}(0))$ is done in the same way. \square

Proof of Theorem 3.6. Let us prove Theorem 3.6 in the cases $\alpha = 0, 1, 2, 3$ successively. According to (3.9) and (3.1) we have $(0, 0, 1) \in \mathbb{G}_0^-$. The representation (4.1) of the determinant $\Delta_{\gamma\lambda\mu}(0; z)$ yields that

$$\Delta_{001}(0; z) = 1 + f(z).$$

By definition (4.2) of $f(z)$, for all $z \in (-\infty, 0)$ the inequalities $f(z) > 0$ and $\Delta_{001}(0; z) > 0$ hold, i.e., the determinant $\Delta_{001}(0; z)$ has no negative zeros in $z \in (-\infty, 0)$. Lemma 4.4 yields that the operator $H_{001}(0)$ has no eigenvalues below the essential spectrum. Theorem 3.5 gives that the operator $H_{\gamma\lambda\mu}(0)$ has no eigenvalues below the essential spectrum for all $(\gamma, \lambda, \mu) \in \mathbb{G}_0^-$.

Due to (3.9) we have $(0, -1, 0) \in \mathbb{G}_1^-$. Lemma 4.5 gives that the operator $H_{0(-1)0}(0)$ has exactly one eigenvalue. Theorem 3.5 yields that for any $(\gamma, \lambda, \mu) \in \mathbb{G}_1^-$, the operator $H_{\gamma\lambda\mu}(0)$ has exactly one eigenvalue below the essential spectrum.

We note $(-3, -3, 0) \in \mathbb{G}_2^-$. According to (iii) of Lemma 4.5 the operator $H_{(-3)(-3)0}(0)$ has two eigenvalues. Theorem 3.5 yields that for any $(\gamma, \lambda, \mu) \in \mathbb{G}_2^-$ the operator $H_{\gamma\lambda\mu}(0)$ has two eigenvalues below the essential spectrum.

Now assume that $(\gamma, \lambda, \mu) \in \mathbb{G}_3^-$.

Due to (3.9) we have

$$C^-(\gamma, \lambda, \mu) < 0, \quad (\lambda, \mu) \in D_3^-.$$

Definition 3.7 and inclusion $(\lambda, \mu) \in D_3^-$ give

$$(4.5) \quad \mu < -1, \quad Q_1^-(\lambda, \mu) > 0.$$

Inequalities (4.5) and $C^-(\gamma, \lambda, \mu) < 0$ guarantee that

$$(4.6) \quad \gamma < -\frac{Q_0^-(\lambda, \mu)}{Q_1^-(\lambda, \mu)} = \frac{\mu + 1}{Q_1^-(\lambda, \mu)} - 1 < 0.$$

According to negativity of γ the function $\Delta_{\gamma 00}(0; z) = 1 + \gamma a(z)$ is continuous and monotone decreasing in $(-\infty, 0)$.

By (3.1) and (4.6) we have that $C^-(\gamma, 0, 0) = \gamma < 0$. Corollary 4.3 yields that

$$(4.7) \quad \lim_{z \rightarrow -\infty} \Delta_{\gamma 00}(0; z) = 1 \quad \text{and} \quad \lim_{z \nearrow 0} \Delta_{\gamma 00}(0; z) = -\infty.$$

Since the function $\Delta_{\gamma 00}(0; z)$ is continuous and monotone decreasing in the interval $(-\infty, 0)$ it has exactly one zero z_{11} below the essential spectrum. Obviously

$$(4.8) \quad \begin{aligned} \Delta_{\gamma 00}(0; z) &= 1 + \gamma a(z) > 0 & \text{if } z < z_{11}, \\ \Delta_{\gamma 00}(0; z) &= 1 + \gamma a(z) < 0 & \text{if } z > z_{11}. \end{aligned}$$

Observe

$$(4.9) \quad \Delta_{\gamma\lambda 0}(0; z_{11}) = (1 + \gamma a(z_{11}))(1 + \lambda d(z_{11})) - \gamma\lambda(b(z_{11}))^2 = -\gamma\lambda(b(z_{11}))^2 < 0.$$

According to inequalities (4.5) we have

$$C^-(\gamma, \lambda, 0) = \gamma + \lambda + \gamma\lambda = Q_1(\lambda, \mu) - (\mu + 1) > 0.$$

Corollary 4.3 and inequality (4.9) give

$$\lim_{z \rightarrow -\infty} \Delta_{\gamma\lambda 0}(0; z) = 1, \quad \Delta_{\gamma\lambda 0}(0; z_{11}) < 0 \quad \text{and} \quad \lim_{z \nearrow 0} \Delta_{\gamma\lambda 0}(0; z) = +\infty.$$

Thus, the continuous function $\Delta_{\gamma\lambda 0}(0; z)$ has at least one zero in each intervals $(-\infty, z_{11})$ and $(z_{11}, 0)$. Therefore there exists real numbers satisfying the inequalities

$$(4.10) \quad z_{21} < z_{11} < z_{22} < 0$$

such that the following equalities hold:

$$\Delta_{\gamma\lambda 0}(0; z_{21}) = \Delta_{\gamma\lambda 0}(0; z_{22}) = 0.$$

Lemma 4.4 and the min-max principle yield that $H_{\gamma\lambda_0}(0)$ has at least two eigenvalues below the essential spectrum. Hence the determinant $\Delta_{\gamma\lambda_0}(0; z)$ has exactly two zeros z_{21} and z_{22} in $(-\infty, 0)$, which yields

$$(4.11) \quad \begin{aligned} (1 + \gamma a(z_{21}))(1 + \lambda d(z_{21})) &= \gamma \lambda (b(z_{21}))^2 > 0, \\ (1 + \gamma a(z_{22}))(1 + \lambda d(z_{22})) &= \gamma \lambda (b(z_{22}))^2 > 0. \end{aligned}$$

The relations (4.8), (4.10) and (4.11) yield

$$\begin{aligned} 1 + \gamma a(z_{21}) &> 0 \quad \text{and} \quad 1 + \lambda d(z_{21}) > 0, \\ 1 + \gamma a(z_{22}) &< 0 \quad \text{and} \quad 1 + \lambda d(z_{22}) < 0. \end{aligned}$$

We note that $(\gamma, \lambda, \mu) \in \mathbb{G}_3^-$ and so $C^-(\gamma, \lambda, \mu) > 0$. Applying Corollary 4.3 we have that

$$\lim_{z \rightarrow -\infty} \Delta_{\gamma\lambda\mu}(0; z) = 1 \quad \text{and} \quad \lim_{z \nearrow 0} \Delta_{\gamma\lambda\mu}(0; z) = -\infty.$$

Observe that

$$\begin{aligned} \Delta_{\gamma\lambda\mu}(0; z_{21}) &= \Delta_{\gamma\lambda_0}(0; z_{21})[1 + \mu f(z_{21})] + 2\gamma\lambda\mu b(z_{21})c(z_{21})e(z_{21}) - \\ &\quad \gamma\mu[1 + \lambda d(z_{21})]c^2(z_{21}) - \lambda\mu[1 + \gamma a(z_{21})](e(z_{21}))^2 = \\ &\quad \mu[\sqrt{-\gamma[1 + \lambda d(z_{21})]}c(z_{21}) + \sqrt{-\lambda[1 + \gamma a(z_{21})]}e(z_{21})]^2 < 0. \end{aligned}$$

Similarly can be shown

$$\Delta_{\gamma\lambda\mu}(0; z_{22}) = -\mu[\sqrt{\gamma[1 + \lambda d(z_{22})]}c(z_{22}) + \sqrt{\lambda[1 + \gamma a(z_{22})]}e(z_{22})]^2 > 0.$$

Then the relations

$$\lim_{z \rightarrow -\infty} \Delta_{\gamma\lambda\mu}(0; z) = 1, \quad \Delta_{\gamma\lambda\mu}(0; z_{21}) < 0, \quad \Delta_{\gamma\lambda\mu}(0; z_{22}) > 0, \quad \lim_{z \nearrow 0} \Delta_{\gamma\lambda\mu}(0; z) = -\infty.$$

yield the existence of zeros z_{31} , z_{32} and z_{33} of function $\Delta_{\gamma\lambda\mu}(0; z)$ satisfying the inequalities $z_{31} < z_{21} < z_{32} < z_{22} < z_{33} < 0$, i.e

$$\Delta_{\gamma\lambda\mu}(0; z_{31}) = \Delta_{\gamma\lambda\mu}(0; z_{32}) = \Delta_{\gamma\lambda\mu}(0; z_{33}) = 0.$$

Hence the function $\Delta_{\gamma\lambda\mu}(0; z)$ has three single zeros less than 0. Lemma 4.4 gives that the operator $H_{\gamma\lambda\mu}(0)$ has three eigenvalues below the essential spectrum.

The proof of item (ii) of the theorem is carried out similarly to the proof of item (i). \square

4.2. The discrete spectrum of $H_{\gamma\lambda\mu}(K)$. For every $n \geq 1$ define

$$(4.12) \quad e_n(K; \gamma, \lambda, \mu) := \sup_{\phi_1, \dots, \phi_{n-1} \in L^2(\mathbb{T})} \inf_{\psi \in [\phi_1, \dots, \phi_{n-1}]^\perp, \|\psi\|=1} (H_{\gamma\lambda\mu}(K)\psi, \psi)$$

and

$$(4.13) \quad E_n(K; \gamma, \lambda, \mu) := \inf_{\phi_1, \dots, \phi_{n-1} \in L^2(\mathbb{T})} \sup_{\psi \in [\phi_1, \dots, \phi_{n-1}]^\perp, \|\psi\|=1} (H_{\gamma\lambda\mu}(K)\psi, \psi).$$

By the minimax principle, $e_n(K; \gamma, \lambda, \mu) \leq \mathcal{E}_{\min}(K)$ and $E_n(K; \gamma, \lambda, \mu) \geq \mathcal{E}_{\max}(K)$. Since, the rank of $V_{\gamma\lambda\mu}$ does not exceed three, by choosing $\phi_1 \equiv 1$, $\phi_2(p) = \cos p$ and $\phi_3(p) = \cos 2p$ we immediately see that $e_n(K; \gamma, \lambda, \mu) = \mathcal{E}_{\min}(K)$ and $E_n(K; \gamma, \lambda, \mu) = \mathcal{E}_{\max}(K)$ for all $n \geq 4$.

Lemma 4.7. *Let $n \geq 1$. Then the maps*

$$K \in \mathbb{T} \mapsto \mathcal{E}_{\min}(K) - e_n(K; \gamma, \lambda, \mu)$$

and

$$K \in \mathbb{T} \mapsto E_n(K; \gamma, \lambda, \mu) - \mathcal{E}_{\max}(K)$$

is non-increasing in $(-\pi, 0]$ and non-decreasing in $[0, \pi]$.

Proof. For $\psi \in L^2(\mathbb{T})$, we consider

$$((H_0(K) - \mathcal{E}_{\min}(K))\psi, \psi) = \int_{\mathbb{T}} 2 \cos \frac{K}{2} (1 - \cos q) |\psi(q)|^2 dq.$$

Then, the map $K \in \mathbb{T} \mapsto ((H_0(K) - \mathcal{E}_{\min}(K))\psi, \psi)$ is non-decreasing in $(-\pi, 0]$ and is non-increasing in $[0, \pi]$. Since $V_{\gamma\lambda\mu}$ is independent of K , from the definition of $e_n(K; \gamma, \lambda, \mu)$, the map $K \in \mathbb{T} \mapsto e_n(K; \gamma, \lambda, \mu) - \mathcal{E}_{\min}(K)$ is non-decreasing in $(-\pi, 0]$ and is non-increasing in $[0, \pi]$. \square

Proof of Theorem 3.1. For any for any $K \in \mathbb{T}$ and $m \geq 1$ Lemma 4.7 gives

$$(4.14) \quad 0 \leq \mathcal{E}_{\min}(0) - e_m(0; \gamma, \lambda, \mu) \leq \mathcal{E}_{\min}(K) - e_m(K; \gamma, \lambda, \mu),$$

and

$$(4.15) \quad 0 \leq E_m(0; \gamma, \lambda, \mu) - \mathcal{E}_{\max}(0) \leq E_m(K; \gamma, \lambda, \mu) - \mathcal{E}_{\max}(K).$$

By assumption $e_n(0; \gamma, \lambda, \mu)$ is an discrete eigenvalue of $H_{\gamma\lambda\mu}(0)$ lying below the bottom $\mathcal{E}_{\min}(K)$. So $\mathcal{E}_{\min}(0) - e_n(0; \gamma, \lambda, \mu) > 0$ and hence, by (4.14) and (2.5), $e_n(K; \gamma, \lambda, \mu)$ is a discrete eigenvalue of $H_{\gamma\lambda\mu}(K)$ for any $K \in \mathbb{T}$. Since $e_1(K; \gamma, \lambda, \mu) \leq \dots \leq e_n(K; \gamma, \lambda, \mu) < \mathcal{E}_{\min}(K)$, it follows that $H_{\gamma\lambda\mu}(K)$ has at least n eigenvalue below the essential spectrum. The case of $E_n(K; \gamma, \lambda, \mu)$ is similar. \square

Proof of Theorem 3.9 is obtained by combining Theorem 3.1 and Corollary 3.8. \square

Acknowledgments. The authors acknowledge support of this research by Ministry of Innovative Development of the Republic of Uzbekistan (Grant No. FZ-20200929224).

REFERENCES

- [1] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, H. Holden: Solvable Models in Quantum Mechanics, Springer 1988.
- [2] S. Albeverio, S. N. Lakaev, Z. I. Muminov: Schrödinger operators on lattices. The Efimov effect and discrete spectrum asymptotics, Ann. Henri Poincaré. **5** (2004), 743-772.
- [3] S. Albeverio, S. N. Lakaev, K. A. Makarov, Z. I. Muminov: The Threshold Effects for the Two-particle Hamiltonians on Lattices, Comm. Math. Phys. **262** (2006), 91-115.
- [4] S. Albeverio, S. N. Lakaev, A. M. Khalkhujayev: Number of Eigenvalues of the Three-Particle Schrödinger Operators on Lattices, Markov Process. Relat. Fields. **18** (2012), 387-420.
- [5] V. Bach, W. de Siqueira Pedra, S.N. Lakaev: Bounds on the discrete spectrum of lattice Schrödinger operators. J. Math. Phys. **59:2** (2017), 022109.
- [6] I. Bloch: Ultracold quantum gases in optical lattices, Nat. Phys. **1** (2005), 23-30.
- [7] G. Dell'Antonio, Z.I. Muminov, V.M. Shermatova: On the number of eigenvalues of a model operator related to a system of three particles on lattices, Journal of Physics A: Mathematical and Theoretical, **44** (2011), 315302. DOI 10.1088/1751-8113/44/31/315302
- [8] V.N. Efimov, Yad. Fiz. **12**, 1080 (1970) [Sov. J. Nucl. Phys. **12**, 589 (1970)].
- [9] L.D. Faddeev, S.P. Merkuriev, Quantum Scattering Theory for Several Particle Systems (Dordrecht: Kluwer Academic Publishers, 1993).
- [10] P.A. Faria Da Veiga, L. Ioriatti, M. O'Carroll: Energy-momentum spectrum of some two-particle lattice Schrödinger Hamiltonians. Phys. Rev. E **66** (2002), 016130.
- [11] F. Hiroshima, Z. Muminov, U. Kuljanov: Threshold of discrete Schrödinger operators with delta-potentials on N -dimensional lattice. Linear and Multilinear Algebra (2020).
- [12] W. Hofstetter et al.: High-temperature superfluidity of fermionic atoms in optical lattices, Phys. Rev. Lett. **89** (2002), 220407.
- [13] D. Jaksch, C. Bruder, J. Cirac, C.W. Gardiner, P. Zoller: Cold bosonic atoms in optical lattices. Phys. Rev. Lett. **81** (1998), 3108-3111.
- [14] D. Jaksch, P. Zoller: The cold atom Hubbard toolbox. Ann. Phys. **315** (2005), 52-79.
- [15] M. Klaus: On the bound state of Schrödinger operators in one dimension. Ann. Phys. **108** (1977), 288-300.

- [16] M. Klaus, B. Simon: Coupling constant thresholds in non-relativistic quantum mechanics. I. Short range two body case. *Ann. Phys.* **130** (1980), 251–281.
- [17] Sh.Yu. Kholmatov, S.N. Lakaev, F.M. Almuratov: On the spectrum of Schrödinger-type operators on two dimensional lattices, *J. Math. Anal. Appl.* **504**:2 (2022), 126363
- [18] Sh.Yu. Kholmatov, S.N. Lakaev, F. Almuratov: Bound states of discrete Schrödinger operators on one and two dimensional lattices, *J. Math. Anal. Appl.* **503**:1 (2021), 125280.
- [19] S.N. Lakaev: The Efimov’s effect of the three identical quantum particle on a lattice. *Funct. Anal. Appl.* **27** (1993), 15–28
- [20] S.N. Lakaev, S.Kh. Abdulkhakimov: Threshold effects in a two-fermion system on an optical lattice. *Theoret. and Math. Phys.* **203**:2 (2020), 251–268
- [21] S.N.Lakaev, A.Boltaev,F.Almuratov: On the discrete spectra of Schrödinger-type operators on one dimensional lattices. *Lob. J. Math.* **43-3**(2022),770–783.
- [22] S.N. Lakaev, I.N. Bozorov: The number of bound states of a one-particle Hamiltonian on a three-dimensional lattice. *Theoret. and Math. Phys.* **158** (2009), 360–376.
- [23] S.N.Lakaev, Sh.Yu.Kholmatov, Sh.I.Khamidov: Bose-Hubbard model with on-site and nearest-neighbor interactions; exactly solvable case. *J. Phys. A: Math. Theor.* **54** (2021), 245201 [22 pages].
- [24] S. N. Lakaev, Sh. Kh. Kurbanov, Sh. U. Alladustov: Convergent Expansions of Eigenvalues of the Generalized Friedrichs Model with a Rank-One Perturbation, *Complex Analysis and Operator Theory* (2021) 15:121 <https://doi.org/10.1007/s11785-021-01157-9> (Scopus. IS= 1.09)
- [25] S.N. Lakaev, E. Özdemir: The existence and location of eigenvalues of the one particle Hamiltonians on lattices. *Hacetatepe J. Math. Stat.* **45** (2016), 1693–1703.
- [26] S.N.Lakaev: Some spectral properties of the generalized Friedrichs model, *Journal of Soviet Mathematics*, **45**(6) (1989), 1540–1563.
- [27] S. N. Lakaev, A. M. Khalkhuzhaev, Sh. S. Lakaev: Asymptotic behavior of an eigenvalue of the two-particle discrete Schrödinger operator. *Theoret. and Math. Phys.* **171**:3 (2012), 438–451.
- [28] M. Lewenstein, A. Sanpera, V. Ahufinger: *Ultracold Atoms in Optical Lattices: Simulating Quantum Many-body Systems*. Oxford University Press, Oxford, 2012.
- [29] D. Mattis: The few-body problem on a lattice. *Rev. Mod. Phys.* **58** (1986), 361–379.
- [30] A. K. Motovilov, W. Sandhas, and V. B. Belyaev: Perturbation of a lattice spectral band by a nearby resonance, *J. Math. Phys.* **42** (2001), 2490–2506.
- [31] C. Ospelkaus, S. Ospelkaus, L. Humbert, P. Ernst, K. Sengstock, K. Bongs: Ultracold heteronuclear molecules in a 3d optical lattice. *Phys.Rev. Lett.* **97**, 2006.
- [32] Y.N. Ovchinnikov, I.N. Sigal: Number of bound states of three-body systems and Efimov’s effect. *Ann. Phys.* **123**:2 (1979), 274–295.
- [33] M.Reed, B.Simon:Methods of modern mathematical pyhsics.III:Scattering Theory, Academic Press,N.Y.,(1978).
- [34] M.Reed, B.Simon:Methods of modern mathematical pyhsics. IV:Analyses of operators, Academic Press,N.Y.,(1978).
- [35] A.V. Sobolev: The Efimov effect. Discrete spectrum asymptotics, *Commun. Math. Phys.* **156**:1, 101–126 (1993).
- [36] H. Tamura: The Efimov effect of three-body Schrödinger operators. *J. Funct. Anal.* **95**:2 (1991), 433–459.
- [37] H. Tamura: Asymptotic distribution of negative eigenvalues for three-body systems in two dimensions: Efimov effect in the antisymmetric space, *Rev. Math. Phys.* **31**:9 (2019), 1950031.
- [38] K. Winkler, G. Thalhammer, F. Lang, R. Grimm, J. Hecker Denschlag, A.J. Daley, A. Kantian, H.P. Büchler, P. Zoller: Repulsively bound atom pairs in an optical lattice. *Nature* **441**, 853–856, 2006.
- [39] D.R. Yafaev: On the theory of the discrete spectrum of the three-particle Schrödinger operator, *Mat. Sb.* **94(136)**, 567–593, 655–656 (1974).

(Saidakhmat Lakaev) SAMARKAND STATE UNIVERSITY, 140104, SAMARKAND, UZBEKISTAN
Email address: slakaev@mail.ru

(Muhayyo Akhmadova) SAMARKAND STATE UNIVERSITY, 140104, SAMARKAND, UZBEKISTAN
Email address: mukhayyo@mail.ru