Chern-Simons Theory, Ehrhart Polynomials, and Representation Theory

Chao Ju

Berkeley Center for Theoretical Physics and Department of Physics University of California, Berkeley, CA 94720, U.S.A.

E-mail: cju19@berkeley.edu

ABSTRACT: The Hilbert space of level q Chern-Simons theory of gauge group G of the ADE type quantized on T^2 can be represented by points that lie on the weight lattice of the Lie algebra \mathfrak{g} up to some discrete identifications. Of special significance are the points that also lie on the root lattice. The generating functions that count the number of such points are quasi-periodic Ehrhart polynomials which coincide with the generating functions of SU(q)representation of the ADE subgroups of SU(2) given by the McKay correspondence. This coincidence has roots in a string/M theory construction where D3(M5)-branes are put along an ADE singularity. Finally, a new perspective on the McKay correspondence that involves the inverse of the Cartan matrices is proposed.

ARXIV EPRINT: 2304.11830

Contents

1	Introduction	1
2	String Theory and D-branes on Orbifold	3
3	Quantization of Chern-Simons Theory on T^2 and the States	4
4	A Special Class of States	6
	4.1 The Geometry and the Counting Problem	6
	4.2 Ehrhart Polynomials	8
5	Computation of $\mathbf{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$	8
	5.1 The Ω Operator	9
	5.2 Applying the Ω Operator	10
6	$\mathbf{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ From Representation Theory	12
	6.1 $SU(q)$ representations of $\Gamma(\mathfrak{su}(N)) = \mathbb{Z}_N$	14
	6.2 Computation of $\Phi_{\mathfrak{g}}(z)$	15
7	Ehrhart Polynomial for $\mathfrak{so}(2(N+2))$	16
	7.1 $SU(q)$ representation of Dic_N	16
	7.2 Equivalence to the Ehrhart polynomial	19
8	A New Perspective on the McKay Correspondence	21
9	The Physical Origin of the Duality	22
10	Conclusion	25

1 Introduction

The 3-dimensional Chern-Simons theory is a remarkably rich tool that has applications in both mathematics and physics. For example, Witten showed that Chern-Simons theory is connected to the Jones polynomial and knot invariants [1]. It is also applied in the study of S-duality [2, 3], a subject related to the geometric Langlands program [4]. In terms of the more tangible physics applications, Chern-Simons theory has been used extensively to study physics on 2-dimensional surfaces. For example, it is used to endow particles in 2+1 dimensions with fractional statistics [6, 7], so that upon exchanging two identical particles the wave function of the two particles can end up with a phase different from ± 1 . In addition, Laughlin explained fractional quantum hall effect by applying Chern-Simons theory¹ [8].

The fact that Chern-Simons theory is important for 2+1 dimensional physics is no accident. The Chern-Simons action

$$S = \frac{q}{4\pi} \int \operatorname{tr}\left(A \wedge dA + \frac{2}{3}A \wedge A \wedge A\right) \tag{1.1}$$

is the unique relativistically invariant action of the gauge field A that has only one derivative acting on A. In this paper, we take the convention that A is \mathfrak{g} valued 1-form and the trace is taken in the representation such that q is a quantized positive integer. We take \mathfrak{g} to be simply laced, namely $\mathfrak{su}(N)$, $\mathfrak{so}(2N)$, \mathfrak{e}_6 , \mathfrak{e}_7 , and \mathfrak{e}_8 . The global property of the gauge group will not concern us in this paper. The geometry we are interested in is $T^2 \times \mathbb{R}$ where \mathbb{R} is the time direction. Upon quantizing the Chern-Simons theory in this geometry, one gets a discrete set of states that can be identified with points in the set [22]

$$\frac{\Lambda_w}{W \ltimes q\Lambda_r} \tag{1.2}$$

where Λ_w , Λ_r are the weight and the root lattice of \mathfrak{g} , and W is the Weyl group.

In this paper, we study the counting of a special class of states that lie in the set

$$\frac{\Lambda_w}{W \ltimes q \Lambda_r} \cap \Lambda_r \tag{1.3}$$

where the intersection with Λ_r picks out the states in the set (1.2) that are also roots.

We will show that the counting of such states leads to a curious connection to Ehrhart polynomials, McKay correspondence, and representation theory. Ehrhart polynomials were first constructed to count lattice points in rational polytopes [18], a problem that is in general NP-hard [19] to solve by computer. Ehrhart polynomials have been found to connect different areas of mathematics such as number theory, geometry, and topology². Since closed-form Ehrhart polynomials are rare and are only for very special geometries, our computation in this paper will add more examples to the known collection of Ehrhart polynomials.

We will also show that the Ehrhart polynomials we obtain from the geometric point of view can also be obtained from a representation theory point of view. In fact, the latter comes from a dual formulation of the problem. The two approaches are connected by the McKay correspondence [23], which gives a ADE Dynkin diagram classification of discrete subgroups of SU(2) and their irreducible representations. The physics behind this dual formulation of the problem comes from a holographic system of D-branes on ADE singularity. More precisely, there is a duality between the ground state Hilbert space $SU(q) \ \mathcal{N} = 4$ Supersymmetric Yang-Mills theory on S^3/Γ and a certain subspace (1.3) of the Hilbert space of level q Chern-Simons theory on T^2 with gauge algebra $\mathfrak{g}(\Gamma)$ given by the McKay correspondence.

 $^{^{1}}$ For a review of Chern-Simons theory and fractional quantum hall effect, see [5] and the references therein.

²For an extensive introduction, see the book [20]

This paper is organized as follows. Because our construction has to do with D-branes on ADE singularity, in section 2 we will review this subject and point out the difference between our construction and the literature. In section 3, we review the quantization of Chern-Simons theory on T^2 and formulate the Hilbert space geometrically in terms of certain points on the Lie algebra lattice. To illustrate the rather abstract notation, we give an example of states of $\mathfrak{g} = \mathfrak{su}(3)$. In section 4, we pose the problem of counting the number of the special class of states in the set defined in (1.2). We shall find that the problem is the same as counting lattice points in rational polytopes, and that the generating function for counting the states is exactly the corresponding Ehrhart polynomial. In section 5, we compute the explicit form of the Ehrhart polynomial for a specific case by using the Ω operator introduced by MacMahon [24]. The general case is solved by reverse-engineering some representation theory formulas in section 6. We will see that the Ehrhart polynomial that counts the special states at level q with gauge algebra \mathfrak{g} is the same as the generating function for the SU(q) representation of the ADE subgroup given by the McKay correspondence. In section 7, we extend our result to the D-series where the gauge algebra of Chern-Simons theory is $\mathfrak{so}(2(N+2)), N \geq 1$. In section 8, we discuss some curious representation theory properties implied by the inverse of Cartan matrices modulo 1 for ADE Lie algebras by focusing on the exceptional series \mathfrak{e}_6 , \mathfrak{e}_7 , and \mathfrak{e}_8 . In section 9, we derive the string/M theory origin of the proposed duality.

The reader who is only interested in the physical aspect of this work can skip the second half of section 4 and all of section 5.

2 String Theory and D-branes on Orbifold

There is a long history of the study of string theory and D-branes on orbifold singularity. Since the particular orbifold singularity that will concern us in this paper is the ADE orbifold singularity, we will in this section do a quick review of some past research on ADE singularity and string theory. We will discuss how this work is different from past research and motivate the Chern-Simons/SYM duality mentioned in the introductory section 1.

There are roughly speaking three uses of putting string theory or D-branes on ADE singularity. The first is to obtain a more realistic compactification of string theory down to lower dimensions. The second is to use string theory to probe the topology and geometry of the ADE singularity. The third is to use string theory construction to obtain new classes of quantum field theory.

One way to obtain ADE singulairty is the "blow-down" of K3 surface [10], the only nontrivial Calabi-Yau manifold in four (real) dimensions [11]. String theory on K3 surface leads to more realistic models of string compactification because the SU(2) holonomy of K3 surface helps break half of the supersymmetry. It also leads to more examples of string dualities [11]. In fact, as we will see in section 9, the duality considered in this paper is derived using the duality between heterotic string on T^4 and IIA string on K3.

The second use partially overlaps with the first use because the low energy spectra of string theory on K3 sheds light on the topological invariants of the K3 surface [10, 11]. To probe the geometry and not just the topology of K3 surface, D-brane technologies must be

used [12]. In [12], the metric of some ALE (asymptotically locally Euclidean) space was computed by probing the space using D1 branes *transverse* to the space. The fact that the moduli space of vacua for the low energy theory coincides with the ALE space suggests that the metric of the underlying space coincides with the metric that enters the kinetic term of the low energy theory.

The third use dates back to [13], where it was mentioned that M-theory on ADE singulairty Γ leads to a seven dimensional super Yang Mills theory transverse to the 4-dimensional ADE singularity with the gauge algebra $\mathfrak{g}(\Gamma)$ McKay-dual to the ADE singularity. In [14] [15], quiver gauge theories are constructed by putting D3 branes *transverse* to the orbifold singularity. In the language of AdS-CFT correspondence, in this construction, the singularity modes out the S^5 part of the bulk geometry and keeps the AdS_5 part untouched.

The difference between this work and the previous constructions is that here, the ADE singularity is longitudinal to the D3 brane worldvolume. In other words, the bulk geometry is $AdS_5/\Gamma \times S^5$. As will be derived in section 9, the long distance limit of this holographic system will involve a Chern-Simons theory in the AdS bulk direction. Constructions similar to this work can be found in [16][17]. Differences between our construction and those in [16][17] will be explained in section 9.

3 Quantization of Chern-Simons Theory on T^2 and the States

The Chern-Simons theory with gauge algebra \mathfrak{g} on T^2 can be quantized using the standard quantization procedure by choosing a gauge $A_0 = 0$ and imposing the constraint $\delta S/\delta A_0 = 0$, where S is defined in equation (1.1). The constraint gives the condition that the connections are flat $F = dA + A \wedge A = 0$. Imposing the constraint, using the remaining freedom to gauge transform the connection A into the maximal torus of \mathfrak{g} , and imposing the canonical commutation relation, one obtains that the states are in one-to-one correspondence with points in the set [22] described in equation (1.2).

In words, states are points on the weight lattice of \mathfrak{g} with two states being identified if they differ by some combinations of the Weyl transformation and q times the root lattice translation. We shall henceforth call this set the state set. Notice that because of the identification by $q\Lambda_r$, the number of states is finite, a fact that can also be seen from the compactness of the Chern-Simons phase space on T^2 . In the large q limit, the number of states is given by

$$\frac{q^r \det C}{|W|} \tag{3.1}$$

where r is the rank of \mathfrak{g} , C is the Cartan matrix, and |W| is the size of the Weyl group. This formula can either be derived from equation (1.2) by noting that det $C = |\Lambda_w / \Lambda_r|$ or going back to canonical quantization and demanding that the phase space contains one state per 2π cell (we set $\hbar = 1$).

To give an example of the state set of $\mathfrak{su}(3)$ Chern-Simons theory at level q = 1 and q = 2, we present the figure (see Fig. 1) from $[25]^3$.

³In [25], a more pedestrian way of quantizing the Chern-Simons theory on T^2 for is used.




Figure 1. In both pictures, e_1 and e_2 are the simple roots, and d_1 and d_2 are the fundamental weights. In the top picture, three black dots represent the three unique states of the level q = 1 theory. In the bottom picture, the six black dots are the unique states of the level q = 2 theory. It is easy to convince oneself that one can reach the white dots or other weight lattice points through a combination of Weyl reflections and q times the root lattice translation on the black dots. Figure retrieved from [25].

Fig. 1 shows the unique states in the Hilbert space for the level 1 and the level 2 theory, respectively. Notice that the pattern continues to all levels: all states lie in the fundamental Weyl chamber, and that as the level increases, the number of state increases quadratically (roughly as the area of the shaded region). In fact, the number of states for the $\mathfrak{su}(3)$ theory at level q is

$$\frac{q(q+1)}{2} \tag{3.2}$$

This formula can be derived as follows. For simply laced algebra (which is our only focus in this paper), the root lattice Λ_r is the same as the coroot lattice Λ_{cr} . A theory in affine Lie algebra shows that the state set (1.2) is simply the highest weight representation of the corresponding affine Lie algebra $\tilde{\mathfrak{g}}$ at level q [26]. Let a state be labeled by the Dynkin label $(a_1, a_2, ..., a_r)$, where each a_i is some nonnegative integer. The highest weight

states satisfy the inequality

$$\sum_{i=1}^{r} c_i a_i \le q \tag{3.3}$$

where $c_i \in \mathbb{Z}_{>0}$ is the coefficient multiplying the *i*th simple root α_i in the expression for the highest root θ :

$$\theta \equiv \sum_{i=1}^{r} c_i \alpha_i \tag{3.4}$$

For the case of $\mathfrak{su}(3)$, the numbers are $c_1 = c_2 = 1$, so the number of Chern-Simons states at level q is the same as the number of solutions to the inequality

$$a_1 + a_2 \le q$$

and solving for this gives exactly the quadratic formula (3.2).

Although the example is given for $\mathfrak{su}(3)$, the reader should keep in mind the generalization of the picture to other ADE gauge algebras. In the next section, we look at a special class of states on the state lattice.

4 A Special Class of States

4.1 The Geometry and the Counting Problem

The special states we want to focus on are those that belong to the set

$$Q_q = \frac{\Lambda_w}{W \ltimes q \Lambda_r} \cap \Lambda_r \tag{4.1}$$

namely states that also roots. The physical motivation for considering such states is mentioned in [21] and is explained in the section 9. Following the $\mathfrak{su}(3)$ example in Fig. 1, at level 1 there is one state that belongs to Q_q , the state at the origin. At level 2 there are two states, the additional one being at the position $d_1 + d_2$.

The question we pose is: given \mathfrak{g} and level q, how many states are in Q_q ? There are two equivalent formulations of this counting problem. The first formulation uses the Cartan matrix and it naturally leads to the concept of Ehrhart polynomials. The second (dual) formulation uses the inverse of the Cartan matrix, and it leads to representation theory and connections to string theory. The rest of the section focuses on the first formulation of the problem.

Let the Cartan matrix of \mathfrak{g} be C. In the basis of Dynkin labels, the rows of C give the Dynkin coefficients of the simple roots. Since we are dealing with simply laced Lie algebras, $C^T = C$, so that the columns of C also give the representation of the simple roots. If $y \in Q_q$, y being the Dynkin label (as a column vector) of some state in the theory, then $y \in \Lambda_r$ by definition, so that it can be represented as a linear combination of the simple roots:

$$y = Cx$$

where $x \in \mathbb{Z}_{\geq 0}^r$ is a column of nonnegative integers. The nonnegativity of x is because states in Q_q are all dominant weights. What constraints does y have to satisfy? The first constraint on y is that the entries are nonnegative (note that $x \ge 0$ does not imply $y \ge 0$). The second constraint is the level q constraint in equation (3.3). In terms of x, the constraints read

$$\sum_{j=1}^{r} C_{ij} x_j \ge 0 \tag{4.2}$$

$$\sum_{i,j=1}^{r} c_i C_{ij} x_j \le q \tag{4.3}$$

Geometrically, the above constraints define a rational polytope, a polygon whose vertices have rational coordinates. The solutions to the above constraints are simply integer points contained inside the rational polytope. By adding slack variables, any rational polytope can be represented as a system of linear equations [20]

$$Ax' = b \tag{4.4}$$

for some matrix A, vector b, and unknowns x'.

For our problem, we need r+1 slack variables $k_1, k_2, ..., k_{r+1} \in \mathbb{Z}_{\geq 0}$, so that the system of inequalities reduces to the system of equalities:

$$\sum_{j=1}^{r} C_{ij} x_j - k_i = 0 \tag{4.5}$$

$$\sum_{i,j=1}^{r} c_i C_{ij} x_j + k_{r+1} = q \tag{4.6}$$

Stacking the vector x and k into x' = (x, k), we have the system of linear equation Ax' = b where A is r + 1 by 2r + 1 and b is r + 1 by 1:

The problem of counting states in Q_q is then transformed into counting the solutions to this system of linear equations.

4.2 Ehrhart Polynomials

What we did in the previous section is under the guise of Ehrhart polynomials [18]. They are generating functions for counting lattice points contained in polytopes. Let us formulate the theory following the notation of [20]. Let a rational polytope \mathcal{P} be specified by a system of linear equations with slack variables added:

$$\mathcal{P} = \{ x \in \mathbb{R}^d_{>0} : Ax = b \}$$

$$\tag{4.9}$$

for some integer valued matrix A and integer valued vector b. One considers the tth dilate of \mathcal{P} , defined as

$$t\mathcal{P} = \{x \in \mathbb{R}^d_{\ge 0} : Ax = tb\}$$

$$(4.10)$$

where $t \in \mathbb{Z}_{>0}$. Let $L_{\mathcal{P}}(t)$ denote the number of lattice points contained in the $t\mathcal{P}$:

$$L_{\mathcal{P}}(t) = \#\{x \in \mathbb{Z}_{\geq 0}^d : Ax = tb\}$$
(4.11)

The Ehrhart polynomial associated to the polytope ${\mathcal P}$ is defined as

$$\operatorname{Ehr}_{\mathcal{P}}(z) = 1 + \sum_{t \ge 1} L_{\mathcal{P}}(t) z^{t}$$

$$(4.12)$$

In our problem, the polytope is given as the qth dilate of

$$Ax' = \begin{pmatrix} 0\\0\\\vdots\\0\\1 \end{pmatrix}$$

with A and x' defined in the previous section (see equation (4.7) and equation (4.8)). We call this base polytope $Q_{\mathfrak{g}}$, where the dependence on the Lie algebra \mathfrak{g} is made explicit as each simply laced Lie algebra has its unique polytope. Since we are interested in computing the number of special states for each level q > 0, we want to find the number of lattice points contained in the qth dilate of $Q_{\mathfrak{g}}$ for each positive q. Therefore, the question posed in the previous section can be now phrased as finding the Ehrhart polynomial of $Q_{\mathfrak{g}}$:

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z) = ?$$

This concludes the first formulation of our problem. The dual formulation of the problem in terms of the inverse of Cartan matrices will be introduced in section 6. In the next section, we develop a formal method to compute $\operatorname{Ehr}_{\mathcal{Q}_g}(z)$ in the most general way possible.

5 Computation of $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{q}}}(z)$

In this section and the next, we compute $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ for all simply laced \mathfrak{g} . We give two approaches to this computation. The first approach uses MacMahon's Ω operator method [24]

but quickly becomes tedious when the level q becomes large. However, the merit of this approach is that it is general: it can be applied to solving for all Ehrhart polynomials given the constraints, and it will be amiss if we do not discuss the general solution. Due to the computational difficulty, we shall only use this approach to give the explicit formula for $Q_{\mathfrak{g}}$ for the case $\mathfrak{g} = \mathfrak{su}(2)$. MacMahon's method is reviewed in section 5.1 and the computation for $\mathfrak{su}(2)$ is done in section 5.2.

The second formulation of the problem uses a hint from representation theory, and leads to the expression of $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ in one full sweep. The second approach is inspired from a duality relation constructed from string theory (see section 9 and ref [16][17]), without which it is not obvious how one can make a connection of $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ to being solved by representation theory.

5.1 The Ω Operator

In computing the number of ways of partitioning some integer u into a sum of n nonnegative integers $a_1+a_2+\ldots+a_n = u$, the order of the integers in the sum does not matter. Therefore, the problem of counting n-partitions of u is quite different from the problem of counting solutions to the the equation $a_1 + a_2 + \ldots + a_n = u$, where the number of solutions is the coefficient of the x^u term in the generating function

$$\frac{1}{(1-x)^n}\tag{5.1}$$

To introduce the Ω operator, we focus on the number partition problem, and therefore we can assume an ordering of a_i to be $a_1 \ge a_2 \ge ... \ge a_n$ without loss of generality. One way to impose this ordering constraint is to consider the expression [24]

$$\Omega \frac{1}{\geq (1 - \lambda_1 x)(1 - \frac{\lambda_2}{\lambda_1} x)(1 - \frac{\lambda_3}{\lambda_2} x)...(1 - \frac{\lambda_n}{\lambda_{n-1}} x)}$$
(5.2)

where the notation Ω_{\geq} means restricting terms that have only nonnegative powers of each λ and setting each λ to be one in the end. This is easily verified by expanding each fraction in power series.

The Ω_{\geq} leads to many identities. For example, one can again expand in power series and verify that

$$\Omega \frac{1}{\geq (1 - \lambda x^{p_1})(1 - \frac{x^{p_2}}{\lambda})} = \frac{1}{(1 - x^{p_1})(1 - x^{p_1 + p_2})}$$
(5.3)

Repeated use of this identity shows that equation (5.2) is equal to

$$\frac{1}{(1-x)(1-x^2)(1-x^3)\dots(1-x^n)}$$
(5.4)

which is exactly the generating function to count the n-partition of some integer.

One can also compose the Ω_{\geq} operation. We modify the notation accordingly if there is any ambiguity with extra variables:

$$\begin{split} \Omega & \Omega \\ \lambda \ge \mu \ge \frac{1}{(1 - \lambda \mu x)(1 - \frac{y}{\lambda^2 \mu})} = \Omega \frac{1}{\lambda \ge (1 - \lambda x)(1 - \frac{xy}{\lambda})} \\ &= \frac{1}{(1 - x)(1 - x^2 y)} \end{split}$$

This expression counts the number of partition into two nonnegative integers a_1 and a_2 such that the constraint $a_1 \ge 2a_2$ is satisfied⁴.

In fact, one can also have two other operators Ω_{\leq} and $\Omega_{=}$, defined in a self-explanatory way. As we shall see, we will be interested in identities involving $\Omega_{=}$. The three operators satisfy some useful algebraic relations listed in [24], and can be used to compute $\Omega_{=}$. For example, let $F(\lambda)$ be some polynomial depending on λ as the variable used in the Ω operators, the following expression

$$\underset{=}{\Omega F(\lambda)} = \underset{\geq}{\Omega F(\lambda)} + \underset{\geq}{\Omega F(\lambda^{-1})} - F(1)$$
(5.5)

can be used to derive identities involving $\Omega_{=}$ in terms of the identities involving Ω_{-} . In the next subsection we will use an identity $\Omega_{=}$ to compute $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ for $\mathfrak{g} = \mathfrak{su}(2)$. We will also sketch the idea of computing $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ for a general simply laced \mathfrak{g} .

5.2 Applying the Ω Operator

To illustrate the use of the Ω operator, we use it to write down a formal expression for the Ehrhart polynomial $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$. Recall that the base polytope is given by

$$Ax' = \begin{pmatrix} 0\\0\\\vdots\\0\\1 \end{pmatrix}$$
(5.6)

where A (r + 1 by 2r + 1) was given in equation (4.7). Let $\mathbf{z} \equiv (z_1, z_2, ..., z_r, z)$. The Ehrhart polynomial is computed by the formal expression

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z) = \underset{z_1=z_2=}{\Omega} \dots \underset{z_r=}{\Omega} \left(\prod_{i=1}^{2r+1} \frac{1}{1 - z_1^{A_{1i}} z_2^{A_{2i}} \dots z_r^{A_{ri}} z^{A_{r+1,i}}} \right)$$
(5.7)

where a composition of $r \, \Omega$ is applied, each time restricting the polynomial to the 0th order term of some z_i . This formula looks intimidating, but can be easily derived as follows. The term involving the product is simply the generating function for counting the combinations

⁴Reason: $\Omega_{\mu\geq}$ implements the condition that ordering does not matter, and $\Omega_{\lambda\geq}$ implements the constraint $a_1 \geq 2a_2$.

of Ax'. Imposing the constraint that the right hand side of the equation has r vanishing entries means that one must restrict to the 0th order term of $z_1, ..., z_r$. Since the last entry of the column on the right hand side is 1, the number of solutions to the qth dilate of the polytope is then the the coefficient of the z^q term. This completes the argument that the formal expression in (5.7) computes the Ehrhart polynomial.

It is nice to have a formal expression like eqn. (5.7) for the Ehrhart polynomial. If one wants to compute the first few terms, a computer can easily do the job. However, we want to take a step further and obtain a closed form solution, which as we shall see exists for all simply laced \mathfrak{g} .

To illustrate how one might obtain a closed form solution using the current formalism, we first focus on the case of $\mathfrak{su}(2)$. The Cartan matrix for $\mathfrak{su}(2)$ is simply 2, the coefficient for the highest root is $c_1 = 1$. Therefore the A matrix is

$$A = \begin{pmatrix} 2 & -1 & 0 \\ 2 & 0 & 1 \end{pmatrix}$$
(5.8)

Using expression (5.7), the Ehrhart polynomial is given by

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{su}}(2)}(z) = \Omega_{z_1=} \left(\frac{1}{1 - z_1^2 z^2} \frac{1}{1 - z_1^{-1}} \frac{1}{1 - z} \right)$$
(5.9)

Using the Ω operator identity [24]

$$= \Omega_{\lambda=} \frac{1}{(1-\lambda^2 x)(1-y\lambda^{-1})} = \frac{1}{1-xy^2}$$
(5.10)

we find that the Ehrhart polynomial for $\mathfrak{g} = \mathfrak{su}(2)$ is

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{su}}(2)}(z) = \frac{1}{(1-z)(1-z^2)}$$
(5.11)

Now let us generalize to arbitrary \mathfrak{g} . The first thing to note is that one needs more general Ω identities in addition to equation (5.10), since a variable Z_i can appear more than two times in the product in equation (5.7). However, the Ω identities for a general polynomial are not documented and do not have clean solutions. This problem has been solved by [27], where the authors developed the Omega Package in Mathematica to compute Ω identities for a general polynomial

$$\frac{P(x_1, ..., x_n; \lambda_1, ...\lambda_r)}{\prod_{i=1}^n (1 - x_i \lambda_1^{v_1(i)} ... \lambda_r^{v_r(i)})}$$
(5.12)

For our problem, we can apply the program r times to eliminate $z_1, ... z_r$ to find $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ of rank r. In the next section, we will use a trick from representation theory to compute all $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ by hand, bypassing the need for the computer program computation.

6 Ehr_{Q_a}(z) From Representation Theory

To connect $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ to representation theory, we briefly summarize the McKay correspondence [23], which associates any simply laced \mathfrak{g} to some discrete subgroup $\Gamma(\mathfrak{g})$ of SU(2). More specifically,

$$\Gamma(\mathfrak{su}(N)) = \mathbb{Z}_N \tag{6.1}$$

$$\Gamma(\mathfrak{so}(2(N+2))) = \operatorname{Dic}_N \tag{6.2}$$

$$\Gamma(\mathbf{e}_i) = E_i \qquad i=6,7,8 \tag{6.3}$$

where \mathbb{Z}_N is the cyclic group of order N, Dic_N is the dicyclic (binary dihedral) group of order 4N, and E_6, E_7, E_8 are the binary tetrahedral group, binary octahedral group, and the binary icosahedral group, respectively (also called 2T, 2O, and 2I).

For some simply laced \mathfrak{g} , consider the homomorphism $\Gamma(\mathfrak{g}) \to SU(q)$, or, in other words, SU(q) representations of the group $\Gamma(\mathfrak{g})$. Let the number of Weyl-inequivalent representations⁵ $\Gamma(\mathfrak{g}) \to SU(q)$ be b_q , so that one forms the generating function to count the number of SU(q) representations

$$\Phi_{\mathfrak{g}}(z) \equiv 1 + \sum_{i=1} b_i z^i \tag{6.4}$$

We claim that this generating function is exactly the same as the Ehrhart polynomial corresponding to the same Lie algebra:

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z) = \Phi_{\mathfrak{g}}(z) \tag{6.5}$$

This beautiful equation connects geometry (left hand side) with representation theory (right hand side) via the McKay correspondence and string theory (the equality sign). The rest of this subsection is devoted to proving this equality and using this equality to compute the closed form $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ for $\mathfrak{g} = \mathfrak{su}(N)$ and $\mathfrak{g} = \mathfrak{so}(2(N+2))$. The formulae for the exceptional Lie algebras are computed in the appendix of [21]. In section 9, we will "prove" this equation using a construction in string/M theory.

The strategy we use here is to look at the dual formulation by starting with weights, expressing them in terms of the simple roots, and imposing the constraints. This reverse process is carried out using the inverse the Cartan matrix. Since the inverse of Cartan matrices in general has fractional entries, this dual formulation is less suited for geometric arguments we had in the previous sections. Instead, we will use purely algebraic arguments to prove equation (6.5).

Let $\mathfrak{g} = \mathfrak{su}(r+1)$ and the corresponding Cartan matrix be C. Start with some weight x in the state set (1.2) which we reproduce here for convenience

$$\frac{\Lambda_w}{W \ltimes q \Lambda_r} \tag{6.6}$$

⁵This means that two diagonal SU(q) matrices are identified if the diagonal elements differ by some permutation.

We can represent x as a vector in $\mathbb{Z}_{\geq 0}^r$ by using its Dynkin label $(x_1, ..., x_r)$. It satisfies the constraint

$$\sum_{i=1}^{r} x_i \le q \tag{6.7}$$

as argued in section 3.

Since the simple roots span \mathbb{R}^r , the weight x has a unique expansion in terms of the simple roots $\alpha_1, ..., \alpha_r$:

$$x = \sum_{i=1}^{r} l_i \alpha_i \tag{6.8}$$

The expansion coefficients l_j are simply given by multiplying x (as a vector) by the inverse of the Cartan matrix [26]

$$l_i = \sum_{j=1}^r C_{ij}^{-1} x_j \tag{6.9}$$

Since we want x to lie on the root lattice Λ_r as well, the only other constraint we need to impose is that each l_j is an integer

$$l_i = \sum_{j=1}^r C_{ij}^{-1} x_j \in \mathbb{Z}$$
(6.10)

The inverse of the Cartan matrix for $\mathfrak{su}(N)$ algebra has an interesting modular structure. Here, we display C^{-1} for $\mathfrak{su}(4)$ and $\mathfrak{su}(7)$:

$$C_{\mathfrak{su}(4)}^{-1} = \frac{1}{4} \begin{pmatrix} 3 & 2 & 1 \\ 2 & 4 & 2 \\ 1 & 2 & 3 \end{pmatrix}$$

$$C_{\mathfrak{su}(7)}^{-1} = \frac{1}{7} \begin{pmatrix} 6 & 5 & 4 & 3 & 2 & 1 \\ 5 & 10 & 8 & 6 & 4 & 2 \\ 4 & 8 & 12 & 9 & 6 & 3 \\ 3 & 6 & 9 & 12 & 8 & 4 \\ 2 & 4 & 6 & 8 & 10 & 5 \\ 1 & 2 & 3 & 4 & 5 & 6 \end{pmatrix}$$

$$(6.11)$$

In general, the formula for $C_{\mathfrak{su}(N)}^{-1}$ is [28]

$$C_{\mathfrak{su}(N),ij}^{-1} = \frac{1}{N} \left[\min(i,j) \times N - ij \right]$$
(6.13)

Even though we have r number of constraints from equation (6.10), we shall see that because of the peculiar property of $C_{\mathfrak{su}(N)}^{-1}$, there is effectively only one constraint, the one imposed by the last row of the matrix:

$$\frac{1}{N}\sum_{i=1}^{N-1} ix_i \in \mathbb{Z}$$
(6.14)

A quick proof that constraint (6.14) implies the rest of the constraints is as follows. The matrix elements $C_{\mathfrak{su}(N),ki}^{-1}$ of the *k*th row are

$$C_{\mathfrak{su}(N),ki}^{-1} = \begin{cases} \frac{(N-k)i}{N} & k > i\\ \frac{(N-i)k}{N} & k \le i \end{cases}$$

Using this, we obtain the constraint imposed by the kth row:

$$l_{k} = \sum_{j=1}^{N-1} C_{\mathfrak{su}(N),kj}^{-1} x_{j}$$
$$= \sum_{i=1}^{k-1} \frac{(N-k)i}{N} x_{i} + \sum_{i=k}^{N-1} \frac{(N-i)k}{N} x_{i}$$
$$= -\frac{k}{N} \sum_{i=1}^{N-1} i x_{i} \mod 1$$

Up to integers, the constraint imposed by the kth row is simply k times that of the constraint imposed by the last row. Therefore, the only unique constraint in (6.10) is the one given by the last row. In summary, there are two constraints on x:

$$\sum_{i=1}^{N-1} x_i \le q \tag{6.15}$$

$$\frac{1}{N}\sum_{i=1}^{N-1} ix_i \in \mathbb{Z}$$
(6.16)

We would now like to argue that this is exactly the same constraints satisfied by SU(q) representations of $\Gamma(\mathfrak{su}(N)) = \mathbb{Z}_N$.

6.1 SU(q) representations of $\Gamma(\mathfrak{su}(N)) = \mathbb{Z}_N$

Since finite dimensional representations are built up from irreducible representations, we first look at the irreducible representations of \mathbb{Z}_N . There are in total N 1-dimensional irreducible representations. The kth irreducible representation is given by the kth power of the Nth root of unity

$$\omega_k = \exp\left(\frac{2\pi ik}{N}\right) \tag{6.17}$$

Here, k runs from 0 to N-1, with k=0 being the trivial representation. The SU(q) representations are constructed by inserting x_k copies of the kth irreducible representation. By definition, the determinant of any SU(q) matrix must be 1, so we have

$$\prod_{k=0}^{N-1} \omega_k^{x_k} = \exp\left(\frac{2\pi i}{N} \sum_{k=0}^{N-1} k x_k\right) = 1$$
(6.18)

But this is equivalent to the constraint 6.16:

$$\frac{1}{N}\sum_{i=1}^{N-1}ix_i\in\mathbb{Z}$$

The constraint that the dimensions of the irreducible representations add up to q is

$$\sum_{i=0}^{N-1} x_i = q \tag{6.19}$$

However, treating x_0 as a slack variable, this constraint is equivalent to (6.15)

$$\sum_{i=1}^{N-1} x_i \le q$$

Therefore, we have shown that the two counting problems satisfy the same constraints and are secretly one and the same. This concludes the proof of eqn. (6.5) for the case $\mathfrak{g} = \mathfrak{su}(N)$:

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z) = \Phi_{\mathfrak{g}}(z)$$

In the next subsection, we give explicit formulae for $\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{g}}}(z)$ by computing $\Phi_{\mathfrak{g}}(z)$.

6.2 Computation of $\Phi_{\mathfrak{g}}(z)$

We want to find the generating function (6.4)

$$\Phi_{\mathfrak{g}}(z) = 1 + \sum_{i} b_i z^i$$

in which b_q counts the number of inequivalent SU(q) representations of \mathbb{Z}_N . We saw in the previous subsection that the N irreducible representations are given by the Nth roots of unity. Consider the function⁶

$$\frac{1}{(1-z)(1-wz)(1-w^2z)\dots(1-w^{N-1}z)}$$
(6.20)

The coefficient of z^q is in general a sum of n terms (n is some integer), representing n ways of building a q-dimensional representation. Each of the n terms has some coefficient w^m for some integer $m \in \mathbb{Z}$, which represents the determinant of that representation. We want to retain terms of determinant 1 only. To project out terms of non-unit determinant, we simply have to sum over w in w^m and divide by N, since we know from Fourier analysis that

$$\sum_{w} w^{m} = \begin{cases} N & m \equiv 0 \mod N \\ 0 & m \neq 0 \mod N \end{cases}$$

where the sum is taken over the Nth root of unity. Therefore, the generating function is

$$\Phi_{\mathfrak{su}(N)}(z) = 1 + \sum_{i} b_{i} z^{i} = \frac{1}{N} \sum_{i=0}^{N-1} \frac{1}{(1-z)(1-w^{i}z)(1-w^{2i}z)\dots(1-w^{(N-1)i}z)}$$
(6.21)

⁶We thank O. Ganor for pointing out this trick.

One can check that b_i is a quasi-polynomial in i, a property shared by Ehrhart polynomial. Finally, we use the theorem proved in the last subsection to give the explicit expression for the Ehrhart polynomial

$$\operatorname{Ehr}_{\mathcal{Q}_{\mathfrak{su}}(N)}(z) = \frac{1}{N} \sum_{i=0}^{N-1} \frac{1}{(1-z)(1-w^{i}z)(1-w^{2i}z)\dots(1-w^{(N-1)i}z)}$$
(6.22)

For example, let $\mathfrak{g} = \mathfrak{su}(2)$. Then

Ehr<sub>$$Q_{\mathfrak{su}(2)}(z) = \frac{1}{2} \left(\frac{1}{(1-z)(1-z)} + \frac{1}{(1-z)(1+z)} \right)$$

= $\frac{1}{(1-z)(1-z^2)}$</sub>

which agrees with what we obtained in equation (5.11) using the Ω operator calculus.

7 Ehrhart Polynomial for $\mathfrak{so}(2(N+2))$

In the previous subsections, we obtained the Ehrhart polynomial for $\mathfrak{su}(N)$ by counting the SU(q) representations of \mathbb{Z}_N . We now use the same method to compute the Ehrhart polynomial for $\mathfrak{so}(2(N+2))$, $N \geq 1$. Here, N is shifted by 2 because of convenience. The Dynkin diagram associated to $\mathfrak{so}(2(N+2))$ is D_{N+2} . According to the McKay correspondence, we should be looking for the representation of the discrete group Dic_N , the dicyclic (or binary dihedral) group of order N. Since Dic_N is less well-known than \mathbb{Z}_N , we analyze the group structure in detail and derive the irreducible representations in the next subsection. After that, we will use the irreducible representations to construct the SU(q)representation of Dic_N and obtain the generating function $\Phi(z)$. We prove that, in a spirit similar to what we did for the $\mathfrak{su}(N)$ case, the generating function $\Phi(z)$ coincides with the Ehrhart polynomial for $\mathfrak{so}(2(N+2))$.

7.1 SU(q) representation of Dic_N

The group Dic_N is defined by the following multiplication rules:

$$r^{2N} = e$$
$$s^2 = r^N$$
$$s^{-1}rs = r^{-1}$$

where e is the identity element. The reader may have noticed a similarity to the dihedral group of order 2N, identifying r with the fundamental rotation and s with the rotation. The difference here is that the reflection s does not square to the identity. Instead, it squares to a central element of the group.

To analyze the irreducible representations of this group, we need to understand the conjugacy classes. For Dic_N , there are N + 3 conjugacy classes:

$$\{e\}, \{r, r^{-1}\}, \{r^2, r^{-2}\}, \dots, \{r^{N-1}, r^{-N+1}\}, \{r^N\}, \{sr^{2k}\}, \{sr^{2k+1}\}$$
(7.1)

where for the last two conjugacy classes, k is an integer running from 0 to 2N-1. Therefore, there are N+3 irreducible representations. The affine Dynkin diagram of D_{N+2} is shown in Fig. 2 [26].

$$\begin{array}{c} & & & & \\ & & & | & (0,1) & & & | & (N+2,1) \\ & & & \circ & - & \circ & - & \circ & - & \circ & (N+1,1) \\ (1,1) & (2,2) & (3,2) & & & (N,2) \end{array}$$

Figure 2. The affine Dynkin diagram for D_{N+2} . The tuple (x, y) indicates the *x*th simple root with (co)mark *y*. The mark is the same as the comark here because the algebra is simply-laced. By the McKay correspondence, the mark also indicates the dimension of the corresponding irreducible representation. Therefore, there are four 1-dimensional irreducible representations and N-1 2-dimensional irreducible representations.

By the McKay correspondence [23] or by abelianizing the group, one sees that there are four 1-dimensional irreducible representations and N - 1 2-dimensional irreducible representations for Dic_N behave differently for N even and N odd. In the following, we restrict N to be an even integer, since the case for N odd can be treated analogously. We present the character table:

$\{e\}$	$\{r, r^{-1}\}$	$\{r^2,r^{-2}\}$	 $\{r^{N-1}, r^{-N+1}\}$	$\{r^N\}$	$\{sr^{2k}\}$	$\{sr^{2k+1}\}$
1	1	1	 1	1	1	1
1	1	1	 1	1	-1	-1
1	-1	1	 -1	1	1	-1
1	-1	1	 -1	1	-1	1
2	$w + w^{-1}$	$w^2 + w^{-2}$	 $w^{N-1} + w^{-(N-1)}$	-2	0	0
2	$w^2 + w^{-2}$	$w^4 + w^{-4}$	 $w^{2(N-1)} + w^{-2(N-1)}$	2	0	0
:		:			:	÷
2	$w^{N-1} + w^{-N+1}$	$w^{2(N-1)} + w^{-2(N-1)}$	 $w^{(N-1)^2} + w^{-(N-1)^2}$	-2	0	0

Table 1. Character table for the Dic_N group. The first four lines are the characters for the four 1-dimensional irreducible representations. Note that the first line is the trivial 1-dimensional representation. The rest of the irreducible representations are 2-dimensional. w represents the 2Nth root of unity $\exp(\pi i/N)$. One can check that the character orthogonality relation holds.

For the N-1 2-dimensional irreducible representations, we note that r and s take on the following form:

$$r = \begin{pmatrix} \exp(m\pi i/N) & 0\\ 0 & \exp(-m\pi i/N) \end{pmatrix} \quad m = 1, ..., N - 1$$
$$s = \begin{cases} \begin{pmatrix} 0 & 1\\ -1 & 0 \end{pmatrix}, m \text{ odd} \\ \begin{pmatrix} 0 & 1\\ +1 & 0 \end{pmatrix}, m \text{ even} \end{cases}$$

In particular, for the 2-dimensional irreducible representations, det r = 1, and a nonunit determinant can only come from s. The formula for s shows that the determinant of the 2-dimensional representations⁷ alternates between 1 and -1. For example, Dic₂ has one 2-dimensional irreducible representation with determinant 1, whereas Dic₄ has three 2-dimensional irreducible representations, with determinant 1, -1, 1, respectively.

We now use the above information to compute the generating function for the SU(q) representation of Dic_N . The SU(q) representation is constructed out of the N + 3 irreducible representations so that the dimensions add up to q. Let the number of the four 1-dimensional representations be $x_0, x_1, x_{N+1}, x_{N+2}$, respectively⁸. Let the number of 2-dimensional representations be labeled by $x_2, x_3, ..., x_N$. The constraint on the size of the representation is

$$x_0 + x_1 + 2x_2 + 2x_3 + \dots + 2x_N + x_{N+1} + x_{N+2} = q$$
(7.2)

In addition to the size constraint, we also have the unit-determinant constraint. From the character table, we see that a -1 determinant can only come from three columns⁹: $\{r, r^{-1}\}, \{sr^{2k}\}, \text{ and } \{sr^{2k+1}\}$. The unit-determinant constraints coming from the three columns are

$$\frac{x_{N+1} + x_{N+2}}{2} \in \mathbb{Z} \tag{7.3}$$

$$\frac{x_1 + x_3 + x_5 + \dots + x_{N-1} + x_{N+2}}{2} \in \mathbb{Z}$$
(7.4)

$$\frac{x_1 + x_3 + x_5 + \dots + x_{N-1} + x_{N+1}}{2} \in \mathbb{Z}$$
(7.5)

Note that the constraints are not independent, since (7.4) and (7.5) imply (7.3). Therefore, we effectively only have two determinant constraints.

To write down the generating function $\Phi_{\mathfrak{so}(2(N+2))}$, we can use the independent constraints (7.4) and (7.5). The idea is similar to the $\mathfrak{su}(N)$ case. Consider the function

$$\frac{1}{(1-z)(1-w_1z)(1-w_2z)(1-w_1w_2z)(1-z^2)^{N/2}(1-w_1w_2z^2)^{N/2-1}}$$
(7.6)

Here, $w_1, w_2 \in \{1, -1\}$. The first four terms represent the contributions of inserting the four 1-dimensional representations, and the last two terms represent the contributions of

⁷In our case, we define the determinant of a representation as follows. If all conjugacy classes in the representation have determinant 1, we say that the representation has determinant 1. If some conjugacy classes have non-unit determinant, we pick the determinant D that, in the polar decomposition, has the smallest angle $\theta \in [0, 2\pi)$ and call D the determinant of the representation. For example, if ω is the 3rd root of unity, $\omega = \exp(2\pi i/3)$. Suppose there are two conjugacy classes with non-unit determinant, one with determinant ω and the other ω^2 . Because ω has a smaller polar angle than ω^2 , we say that the representation has determinant ω .

⁸The numbering here is to make connection with the numbering of the nodes in the affine Dynkin diagram.

⁹Note that the entries in the character table contain the trace of the conjugacy classes. The determinant coincides with the trace for 1-dimensional representations. For 2-dimensional representations the determinant in this case was computed earlier by looking at the explicit matrix representations. There are other columns that have -1 determinant, but one can show that there are only these three independent columns to consider.

inserting the N-1 copies of the 2-dimensional representations. If we expand the fractions into a power series, a generic term would look like

$$cw_1^{k_1}w_2^{k_2}z^n$$
 (7.7)

where $c, k_1, k_2 \in \mathbb{Z}$. The $w_2^{k_2}$ term represents the determinant contribution from the two 1-dimensional representations labeled by N + 1 and N + 2. We want to retain the term that satisfies $w_2^{k_2} = 1$. This can be done by a projection similar to what we did for the $\mathfrak{su}(N)$ case, except now we need to sum over $\omega_2 \in \{1, -1\}$ and divide by 2. We also need to do a similar projection on w_1 . In fact, the two projections help us retain the terms that satisfy the constraints (7.4) and (7.5). Therefore, we need to compute

$$\Phi_{\mathfrak{so}(2(N+2))}(z) = \frac{1}{4} \sum_{w_1, w_2} \frac{1}{(1-z)(1-w_1z)(1-w_2z)(1-w_1w_2z)(1-z^2)^{N/2}(1-w_1w_2z^2)^{N/2-1}}$$

e which gives us the answer for $\Phi_{\mathfrak{so}(2(N+2))}(z)$:

$$\frac{1}{4} \left(\frac{1}{(1-z)^4 (1-z^2)^{N-1}} + \frac{2}{(1-z^2)^2 (1-z^2)^{N/2} (1+z^2)^{N/2-1}} + \frac{1}{(1-z^2)^2 (1-z^2)^{N-1}} \right)$$
(7.8)

As an example, we set N = 2, so that we are looking at the generating function for SU(q) representations of Dic₂. The first few terms of (7.8) are

$$\Phi_{\mathfrak{so}(8)}(z) = 1 + z + 5z^2 + \dots$$

which suggests that there are one SU(1) representation and five SU(2) representations. The one SU(1) representation is just the trivial representation. Even though there are four unitary 1-dimensional representations for Dic₂, only the trivial representation has unit determinant as can be seen from the character table. The reader can easily work out the five SU(2) representations, one of which comes from the unique *irreducible* 2-dimensional representation of Dic₂. The rest come from the *reducible* 2-dimensional representations by combining the 1-dimensional irreducible representations such that the determinant constraint is satisfied.

We will show in the next subsection that this is exactly the Ehrhart polynomial $\operatorname{Ehr}_{\mathcal{Q}_{so}(2(N+2))}(z)$ for the D_{N+2} polytope defined in a similar fashion as in section 4.

7.2 Equivalence to the Ehrhart polynomial

We are looking at the level q highest weight representations of $\mathfrak{so}(2(N+2))$ which also lie on the root lattice. To show that the Ehrhart polynomial coincides with the generating function obtained in the previous subsection, we simply show that the two systems have the same constraints. Let the Dynkin label of some highest weight representation be $\lambda = (x_1, x_2, ..., x_{N+2})$, each term some nonnegative integer. The level q constraint yields

$$(\lambda, \theta) = x_1 + 2x_2 + 2x_3 + \dots + 2x_N + x_{N+1} + x_{N+2} \le q$$
(7.9)

where θ is the highest root whose expansion coefficients in terms of the simple roots can be read off from the affine Dynkin diagram. By adding a slack variable to turn the inequality into an equality, we reproduce the first constraint (7.3). Next, we need to demand that the weights are expressed as integer combinations of simple roots:

$$\sum_{j=1}^{N+2} C_{ij}^{-1} x_j \in \mathbb{Z}$$
(7.10)

The inverse of the Cartan matrix for the D-series has an interesting form [29]. Since the matrix is symmetric, we only give the values for the upper half of the matrix. Let C^{-1} be the inverse of the Cartan matrix for D_{N+2} . We have

$$C_{ij}^{-1} = \begin{cases} i, & 1 \le i \le j \le N \\ i/2, & i \le N, j = N+1 \text{ or } N \\ \frac{N}{4}, & i = N+1, j = N+2 \\ \frac{N+2}{4}, & i = j = N+1 \text{ or } N+2 \end{cases}$$

We give an example of C_{ij}^{-1} for D_6 where N = 4:

$$C_{\mathfrak{so}(12)}^{-1} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1/2 & 1/2 \\ 1 & 2 & 2 & 2 & 1 & 1 \\ 1 & 2 & 3 & 3 & 3/2 & 3/2 \\ 1 & 2 & 3 & 4 & 2 & 2 \\ 1/2 & 1 & 3/2 & 2 & 3/2 & 1 \\ 1/2 & 1 & 3/2 & 2 & 1 & 3/2 \end{pmatrix}$$

Note that the N by N block is integer-valued, so it does not enter into the constraint (7.10). We only have to worry about the last two columns and the last two rows (which are the same as the last two columns by symmetry). For our case, N is an even number, so the only fraction that enter into the constraints modulo 1 is 1/2. Restricting our attention to the last two columns, we see that as the row number increases from 1 to N, the values of the last two columns alternate between being half-integer valued and integer valued. Therefore, from the first N rows of C^{-1} , we effectively get only one constraint:

$$\frac{x_{N+1} + x_{N+2}}{2} \in \mathbb{Z} \tag{7.11}$$

The constraints coming from the last two rows can be deduced similarly. Restricting ourselves to the last two rows, as the column number j increases from 1 to N, the values of $C_{N+1,j}^{-1}$ and $C_{N+2,j}^{-1}$ alternate between being half-integer valued and integer valued. Taking into account of the last two columns, the constraints are

$$\frac{x_1 + x_3 + x_5 + \dots + x_{N-1} + x_{N+2}}{2} \in \mathbb{Z}$$
(7.12)

$$\frac{x_1 + x_3 + x_5 + \dots + x_{N-1} + x_{N+1}}{2} \in \mathbb{Z}$$
(7.13)

We see that (7.9), (7.12), and (7.12) are exactly the same constraints we obtained in the last subsection. This establishes the equivalence of the generating function for SU(q)representation of Dic_N group and the Ehrhart polynomial for the $\mathfrak{so}(2(N+2))$ polytope. In fact, one can repeat the same argument to show that the equivalence holds for the exceptional Lie algebras as well. This concludes the proof of remarkable formula (6.5).

8 A New Perspective on the McKay Correspondence

Let C be some Cartan matrix of an ADE type Lie algebra of rank r. Let G_C be the discrete SU(2) subgroup corresponding to C according to the McKay correspondence. Define $[C^{-1}]$ as C^{-1} modulo 1, where modulo 1 is done element-wise. Our computation in the previous two sections shows that a great deal of information is hidden in this object. In particular, $[C^{-1}]$ can tell us about the determinant¹⁰ of the irreducible representations of the group G_C . To make it more precise, we define the following \vee operator acting on the rational numbers in the congruence class of 1 as

$$a \lor b = \begin{cases} a+b, & \text{if } a = 0 \text{ or } b = 0\\ a, & b \text{ is a nonzero integer multiple of } a\\ b, & a \text{ is a nonzero integer multiple of } b \end{cases}$$

Note that this definition comes with a priority structure: there could be cases where condition 2 and 3 are both satisfied. In that case, we stick with condition 2 and demand that $a \lor b = a$. For example, bearing in mind that we are working within the congruence class of 1, the above rules imply

$$2/3 \lor 1/3 = 2/3$$
 because $1/3$ is 2 times $2/3$
 $5/7 \lor 6/7 = 5/7$ because $6/7$ is 4 times $5/7$
 $1/4 \lor 0 = 1/4$
 $1/2 \lor 1/2 = 1/2$

As we shall see, we never have to worry about the case when a and b do not satisfy the three cases. To use the \vee operator, let us define X_i to be the *i*th row of $[C^{-1}]$. Let the operator \vee act element-wise on X_i . Define the row vector X as

$$X \equiv X_1 \lor X_2 \lor \dots \lor X_r \tag{8.1}$$

The dual group G_C has r + 1 irreducible representations, among which 1 of them is the trivial representation. We claim that the determinant of the rest of the r non-trivial representations, encoded in the row vector $D = (d_1, ..., d_r)$, can be found by

$$D = \exp 2\pi i X \tag{8.2}$$

where the exponential is taken element-wise on X, producing another row vector.

This claim can be proved by checking for all ADE Lie algebras. The information presented in the previous sections is enough for the readers to check the claim for the A and the D series. Here we will focus on the exceptional series \mathfrak{e}_6 , \mathfrak{e}_7 , and \mathfrak{e}_8 . The character tables for the dual group are worked out in [21]. We give a brief summary.

• \mathfrak{e}_6 . The dual group has six nontrivial irreducible representations, of which two have determinant $\exp(2\pi i/3)$ and two have determinant $\exp(4\pi i/3)$. The other two have determinant 1.

¹⁰The determinant of a representation is defined in footnote 7.

- \mathfrak{e}_7 . The dual group has seven nontrivial irreducible representations, of which three have determinant -1 while the rest has determinant 1.
- \mathfrak{e}_8 . All representations of the dual group have unit determinant.

Let us now compare the prediction made by equation (8.2) to the facts cited above. The most trivial case to check is \mathfrak{e}_8 , whose inverse Cartan matrix modulo 1 vanishes. In this case, $X_1 = X_2 = \ldots = X_8 = 0$, so according to the formula above, all irreducible representations of the binary icosahedral group have unit determinant.

For \mathfrak{e}_7 , $[C^{-1}]$ is

The \lor sum of all the rows gives (0, 0, 0, 1/2, 0, 1/2, 1/2). Equation (8.2) shows that the determinants are (1, 1, 1, -1, 1, -1, -1), agreeing with the facts above.

For \mathfrak{e}_6 , $[C^{-1}]$ is

$$[C_{\mathfrak{e}_6}^{-1}] = \begin{pmatrix} 1/3 \ 2/3 \ 0 \ 1/3 \ 2/3 \ 0 \\ 2/3 \ 1/3 \ 0 \ 2/3 \ 1/3 \ 0 \\ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \\ 1/3 \ 2/3 \ 0 \ 1/3 \ 2/3 \ 0 \\ 2/3 \ 1/3 \ 0 \ 2/3 \ 1/3 \ 0 \\ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \end{pmatrix}$$

The \vee sum of all the rows gives (1/3, 2/3, 0, 1/3, 2/3, 0). Equation (8.2) shows that the determinants are $(w, w^2, 1, w, w^2, 1)$, where w is the third root of unity. This also agrees with the facts cited above, and completes the proof of equation (8.2).

9 The Physical Origin of the Duality

In this section, we give an overview of the physics underlying equation (6.5). As we shall see in this section, the right hand side of equation (6.5) will be interpreted as the generating function for computing the dimension of the ground state Hilbert space of $\mathcal{N} = 4 \ SU(q)$ Yang-Mills theory on S^3/Γ . The left hand side of equation (6.5) is the generating function for computing the dimension of a certain subspace (1.3) of the Hilbert space of level q Chern-Simons theory with gauge algebra $\mathfrak{g}(\Gamma)$, where $\mathfrak{g}(\Gamma)$ is given by the McKay correspondence. The SYM theory ground states are simply the flat SU(q) Wilson lines wrapping around the Γ singularity, which in the language of representation theory translates to SU(q) representation of Γ up to conjugation and Weyl group identification. To prove (6.5), we would like to show that these two Hilbert spaces indeed have the same dimension for a given SU(q) gauge group on the SYM side and for any ADE singularity Γ . To show this, consider in IIB string theory a stack of N D3 branes that in the Euclidean signature span the 0123 directions. The ADE singularity Γ acts on the world volume¹¹ of the D3 branes \mathbb{C}^2 to make it \mathbb{C}^2/Γ . Since M-theory on T^2 is dual to IIB string theory, we lift the D3 branes to M5 branes by first taking the T-duality in the 4 direction to turn them into D4 branes and then blowing up the M-theory circle in the # direction. The situation can be summarized in Table 2.

	0	1	2	3	4	5	6	7	8	9	#
D3	Ν	Ν	Ν	Ν	D	D	D	D	D	D	х
M5	Ν	Ν	Ν	Ν	Ν	D	D	D	D	D	Ν

Table 2. The string/M theory setup. Here, D denotes "Dirichlet" and N "Neumann". To avoid confusion, we use # to deonte the 10th direction. Since the D3 brane exists only in the 9+1 dimensional universe, an "x" is put under the 10th direction to indicate that the D3 brane does not exist in that direction.

A stack of N M5 branes whose worldvolume is placed on an ADE singularity \mathbb{C}^2/Γ has a near horizon geometry $AdS_7/\Gamma \times S^4$. Let the Lie algebra corresponding to the ADE singularity be $\mathfrak{g}(\Gamma)$. The boundary theory lives on directions 012345. Let the bulk direction be 6. We claim that there exists a coupling

$$-\frac{1}{4\pi}\int \frac{C_3}{2\pi}\wedge \operatorname{tr}(F\wedge F) \tag{9.1}$$

where C_3 is the 3-form in M-theory and F is the 2-form field strength taking values in \mathfrak{g} . The integral is taken in the 456789# directions. The 5-plane transverse to the M5 branes can be decomposed into $\mathbb{R}^+ \times S^4$, where \mathbb{R}^+ is the direction into the bulk, and S^4 is the 4-sphere that surrounds the M5 branes. Integrating by parts, we obtain

$$\frac{1}{4\pi} \int_{T^2 \times \mathbb{R}^+ \times S^4} \frac{dC_3}{2\pi} \wedge \operatorname{tr}\left(A \wedge dA + \frac{2}{3}A \wedge A \wedge A\right) = \frac{N}{4\pi} \int_{T^2 \times \mathbb{R}^+} \operatorname{tr}\left(A \wedge dA + \frac{2}{3}A \wedge A \wedge A\right) \tag{9.2}$$

reproducing the Chern-Simons theory on T^2 at level N.

Before we derive this, we mention two similar constructions in the literature. Ref [16] reproduces a part of this duality by considering intersecting D4-D6 branes in type IIA string theory, so only the A-singularity is probed since D6 branes cannot create the D- or the E-singularity. Ref [17], on the other hand, considers Nakajima's instanton construction [31] on \mathbb{R}^4/Γ using a certain twisted super Yang-Mills theory without explicitly mentioning the Chern-Simons theory. In [17], it is shown that a certain combination of the middle-dimensional cohomology of U(q) n-instanton moduli space on \mathbb{R}^4/Γ has the same structure as the highest weight representation of a level q state of algebra $\mathfrak{g}(\Gamma)$ (see equation 4.42 in [17]). The link to the Chern-Simons theory is implicit in their work because of the equivalence between WZW conformal blocks and Chern-Simons theory as shown in [1]. We derive the duality using a perspective different from these two constructions. Compared

¹¹Note that this setup is different from the D-brane on orbifold setup considered in [14], where the orbifold action is along the transverse direction of the D-brane worldvolume rather than the longitudinal direction.

to [16], our construction has the virtue of incorporating the D- and the E-singularity. Compared to [17], our construction does not require twisting the super Yang-Mills theory and is easier to understand from a physicist's perspective.

The strategy we use to derive equation (9.1) is through the string-string duality. In the first step, we compactify the 4 direction and take the T-dual to end up in the Type IIA string theory. We now have N D4 branes along the 01234 directions. The ADE singularity acting on the 0123 directions can be locally thought of as a certain degeneration limit of a K3 surface. Although the latter is compact and the former is noncompact, the distinction will not be important in the following derivation. It is known that heterotic string on T^4 is dual to IIA string on K3, with the coupling constants being related by the equation

$$e^{\phi} = e^{-\phi'} \tag{9.3}$$

where unprimed quantities are for the heterotic theory and the primed quantities are for the IIA theory [30]. In addition, the NS-NS 2-forms are related in 6D as

$$dB = e^{-2\phi'} \star dB' \tag{9.4}$$

Now, both the T^4 and the K3 are along the Euclideanized 0123 directions. Recall that the NS-NS 2-form in IIA theory descends from the 3-form in M-theory as $dB' \wedge dx^{\#} = dC_3$. So if one blows up the M-theory circle and lift the above relation to 7D (i.e. adding the #direction), one gets

$$dB = e^{-2\phi'} \star dC_3 \tag{9.5}$$

We now have N M5 branes along the 01234# directions, where the 4 and the # directions are circles.

The heterotic theory has a modified action for its 3-form field strength (setting the curvature contribution to $zero^{12}$):

$$\frac{1}{2} \int_{\mathbb{R}^6} e^{-2\phi} (dB - CS, dB - CS)$$
(9.6)

where CS denotes the Chern-Simons three form and where we considered the T^4 compactified action so the integration is along the directions 456789 in the IIA theory. Notice that $e^{-2\phi} = e^{2\phi'}$ is the heterotic dilaton coupling. If we consider blowing up the M-theory circle to lift the above action to 7D, and plug in the duality relation, the above term becomes

$$\frac{1}{2} \int_{\mathbb{R}^7 = T^2 \times \mathbb{R}^5} e^{2\phi'} (e^{-2\phi'} \star dC_3 - CS, e^{-2\phi'} \star dC_3 - CS)$$
(9.7)

where the integral is taken along the directions 456789# and we split the directions along the torus (4#) and the \mathbb{R}^5 (56789) transverse to the N M5 branes (see Table 2). The \mathbb{R}^5

 $^{^{12}}$ It is amusing to note that if one were to include curvature, i.e. the gravitational Chern-Simons term, one would be able to give a string-theory derivation of the curvature counter term in the one-loop correction to the Chern-Simons action. In [1], this counter term arises out of the consideration for canceling the framing anomaly.

can be further decomposed into a radial direction times an S^4 that surrounds the N M5 branes. From this expression, we get a cross term

$$-\int (CS, \star dC_3) = \int CS \wedge \star \star dC_3 = \int CS \wedge dC_3 = N \int CS$$
(9.8)

where in the last step we integrated the 4-form fluxes along the S^4 to pick up the overall prefactor N, reproducing (9.1) and the Chern-Simons action we sought for.

In the IR limit, the Chern-Simons action is the unique action that has the lowest number of derivatives, and will characterize the ground state structure of the theory. Therefore, the ground state Hilbert space of $\mathcal{N} = 4 U(q)$ SYM on S^3/Γ is equivalent to that of the holographic dual which is the Hilbert space of level q Chern-Simons theory with gauge group $\mathfrak{g}(\Gamma)$. This Chern-Simons Hilbert space is given by the set (1.2). In this work, we considered $\mathcal{N} = 4 SU(q)$ SYM on S^3/Γ . Changing the gauge group from U(q) to SU(q)reduces the number of ground states in the SYM ground state Hilbert space. What is the corresponding reduction on the Chern-Simons side? To see this, we use the example $\Gamma = \mathbb{Z}_k$. The SU(q) SYM ground states Hilbert space admits a level-rank duality: the dimension of SU(q) ground state Hilbert on S^3/\mathbb{Z}_k is the same as that of the SU(k) ground state Hilbert space on S^3/\mathbb{Z}_q^{13} . The only nontrivial subspace of the Chern-Simons Hilbert space that is invariant under the level-rank duality is the one given by the set (1.3) [26]. This concludes the physical derivation of (6.5).

10 Conclusion

We have shown that the counting of root lattice states of level q Chern-Simons theory Hilbert space on T^2 can be solved by computing the exact generating function using either the Ω operator calculus or the duality proposed in [21] that has a similar origin to![16][17]. The reader might wonder why we framed the problem in terms of the Ehrhart polynomials. First, the number of root lattice states grows as some quasi-periodic polynomial, a property shared by Ehrhart polynomials. As we saw in section 4, the geometric formulation of the counting problem leads very naturally to the idea of Ehrhart polynomials. Second, and most importantly, recent development of mathematics and physics shows that it is always fruitful to find connections between different subfields of mathematics and physics. Since Ehrhart polynomials connect various branches of mathematics such as number theory and topology, formulating the problem using Ehrhart polynomials is an attempt at achieving more unification.

Despite being a decades-old subject, Chern-Simons theory is at the heart of the inter-connectedness explored in this paper. In fact, as shown in the last section and in [21][16][17], the Chern-Simons theory arise naturally at the long distance limit of some holographic system in string theory. Another key ingredient is the McKay correspondence [23]. In fact, we saw in section 8 that the simple Lie algebras know a lot more than the content of the McKay correspondence through the inverse of the Cartan matrices. The

¹³A quick way to show this is to use the SU(N) Ehrhart polynomials computed in (6.22) and show that the *q*th term of $\operatorname{Ehr}_{\mathfrak{su}(k)}$ is the same as the *k*th term of $\operatorname{Ehr}_{\mathfrak{su}(q)}$.

simple Lie algebras secretly know the determinant of the irreducible representations of the corresponding ADE subgroup.

Finally, we pose some open questions. In this work, we focused on a special subspace of the Chern-Simons Hilbert space on T^2 given by the set (1.3) and found that there is a description using a certain Supersymmetric Yang-Mills theory. Are there other Chern-Simons states that admit a similar dual description? Chern-Simons theory can be quantized on higher-genus Riemann surfaces. On the SYM side, compactifying the M5 branes on higher-genus Riemann surfaces will break another half of the supersymmetry. The building blocks of the Hilbert space on closed Riemann surfaces are the N_{ijk} fusion coefficients. What does the SYM dual of the fusion coefficients look like? We will tentatively propose an answer to this question in a forthcoming work. In this work, we focused on the SU(q)gauge group for the SYM theory, but one can also ponder over other compact Lie groups such as SO(q) or Sp(2q). What special states in the Chern-Simons theory would these representations correspond to? A standard way to construct SO(q) and Sp(2q) gauge groups in string theory is by using orientifolds, so we would expect that a similar brane construction using orientifolds can help shed light on this question.

Acknowledgments

We would like to thank Emil Albrychiewicz, Ori Ganor, Yasunori Nomura, and Tong Zhou for useful discussions. In particular, we thank Ori Ganor for pointing out the potential significance of the states considered in this paper.

References

- E. Witten, Quantum field theory and the Jones polynomial, Comm. Math. Phys. 121(3) (1989) 351-399.
- [2] O.J. Ganor and Y.P. Hong, Selfduality and Chern-Simons Theory, [0812.1213].
- [3] D. Gaiotto and E. Witten, Supersymmetric Boundary Conditions in N = 4 Super Yang-Mills Theory, Journal of Statistical Physics. 135 (2009) 789–855. [0804.2902].
- [4] A. Kapustin, E. Witten, Electric-Magnetic Duality And The Geometric Langlands Program, Commun. Num. Theor. Phys. 1 (2007) 1-236. [0604151].
- [5] A. Zee, Quantum Hall Fluids, Field Theory, Topology and Condensed Matter Physics 456 (2007) 99–153. [9501022].
- [6] F. Wilczek, Magnetic Flux, Angular Momentum, and Statistics, Phys. Rev. Lett. 48 (1982) 1144.
- [7] F. Wilczek and A. Zee, Linking Numbers, Spin, and Statistics of Solitons, Phys. Rev. Lett. 51 (1983) 2250.
- [8] R.B. Laughlin, Anomalous Quantum Hall Effect: an Incompressible Quantum Fluid with Fractionally Charged Excitations, Phys. Rev. Lett. 50 (1983) 1395.
- [9] E. Witten, 2+1 Dimensional Gravity as an Exactly Soluble System, Nucl. Phys. B 311 (1988) 46-78.

- [10] L. Dixon, J. Harvey, C. Vafa, E. Witten, String on Orbifolds, Nucl. Phys. B 261 (1985) 678-686.
- [11] P. Aspinwall, K3 Surfaces and String Duality, [9611137]
- [12] C. Johnson, R. Myers, Aspects of Type IIB Theory on ALE Spaces, [9610140]
- [13] E. Witten, Some Comments on String Dynamics, [9507121]
- [14] M. Douglas and G. Moore, *D-branes, Quivers, and ALE Instantons*, (1993) [9603167]
- [15] D. Diaconescu, M. Douglas, J. Gomis, Fractional Branes and Wrapped Branes, [9712230] JHEP 1998 (1998).
- [16] R. Dijkgraaf, L. Hollands, P. Sulkowski, C. Vafa, Supersymmetric Gauge Theories, Intersecting Branes and Free Fermions, JHEP 2008(2) (2008). [0709.4446]
- [17] C. Vafa, E. Witten, A Strong Coupling Test of S-Duality, Nucl. Phys. B, 431 (1994) 3-77. [9408074]
- [18] E. Ehrhart, Sur les polyèdres rationnels homothétiques à n dimensions, Comptes rendus de l'Académie des Sciences 254 (1962) 616-618.
- [19] J.A. De Loera, The Many Aspects of Counting Lattice Points in Polytopes, Mathematische Semesterberichte 52 (2005) 175–195.
- [20] M. Beck and S. Robins, Computing the Continuous Discretely, Springer (2015).
- [21] O.J. Ganor and C. Ju, work in progress.
- [22] S. Elitzur, G. Moore, A. Schwimmer, N. Seiberg, Remarks on the Canonical Quantization of the Chern-Simons-Witten Theory, Nucl. Phys. B 326 (1989) 108-134.
- [23] J. McKay, Graphs, singularities, and finite groups, Proc. Symp. Pure Math. 38 (1980) 159-162.
- [24] P.A. MacMahon, Combinatory Analysis, Volume II, Cambridge The University Press (1916).
- [25] X.G. Wen and A. Zee, Topological Degeneracy of Quantum Hall Fluids, Phys. Rev. B 58 (1998). [9711223]
- [26] V. Kac, Infinite Dimensional Lie Algebras, Cambridge University Press (1990).
- [27] G.E. Andrews, P. Paul, and A. Riese, MacMahon's Partition Analysis: The Omega Package, Europ. J. Combinatorics 22 (2001) 887-904.
- [28] E.B. Dynkin, Semisimple Subalgebras of Semisimple Lie Algebras, American Mathematical Society Translations 30 (1952) 349–462.
- [29] Y. Wei and Y.M. Zou, Inverses of Cartan Matrices of Lie Algebras and Lie Superalgebras, Linear Algebra and its Applications 521 (2017) 283-298. [1711.01294].
- [30] E. Witten, String Theory Dynamics In Various Dimensions, Nucl. Phys. B, 443 (1995) 85-126. [9503124]
- [31] H. Nakajima, Instantons on ALE Spaces, Quiver Varieties, and Kac-Moody Algebras, Duke Mathematics Journal 76(2) (1994).