Spectral analysis of N-body Schrödinger operators at two-cluster thresholds

Erik Skibsted, Matematisk Institut, Aarhus Universitet Ny Munkegade 8000 Aarhus C, Denmark

Xue Ping Wang, Laboratoire de Mathématiques Jean Leray UMR CNRS 6629, Nantes Université 44322 Nantes Cedex, France

April 25, 2023

Contents

1	Introduction, examples and notation							
	1.1	Scope and results	1					
		1.1.1 Applications to threshold scattering	5					
	1.2	Many-body Schrödinger operators	6					
		1.2.1 Principal example, dynamical nuclei	8					
	1.3	$N\text{-}\mathrm{body}$ Schrödinger operators with infinite mass nuclei 1	0					
		1.3.1 Principal example, fixed nuclei	1					
	1.4	Generalized N-body Schrödinger operators	2					
		1.4.1 Spaces and notation	4					
2	Red	uction to a one-body problem 1	7					
	2.1	An abstract reduction scheme	8					
	2.2	Non-multiple two-cluster threshold case	1					
	2.3	Multiple two-cluster threshold, $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$	3					
		2.3.1 $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$; the case $\lambda_0 = \Sigma_2$ and $\lambda_0 \notin \sigma_{pp}(H')$	7					
	2.4	The case $\lambda_0 \in \sigma_{pp}(H')$ 3	2					
		2.4.1 $\lambda_0 \in \sigma_{pp}(H')$; non-multiple case	2					
		2.4.2 $\lambda_0 \in \sigma_{\rm pp}(H')$; multiple case	3					
	2.5	Multiple two-cluster case, $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$	4					
		2.5.1 $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$; a general approach $\ldots \ldots \ldots \ldots \ldots 3$	4					
3	Spe	etral analysis of H' near λ_0 3	7					
	3.1	Mourre estimate	8					
		3.1.1 Proof of Proposition 3.3 $\ldots \ldots \ldots \ldots \ldots \ldots \ldots 4$	2					
	3.2	Multiple commutators and calculus	6					
		3.2.1 Computing a commutator $\ldots \ldots \ldots \ldots \ldots \ldots \ldots \ldots 5$	1					
		$3.2.1.1 \text{Smooth sign function} \dots \dots$	4					
	3.3	Positive commutator estimates	5					
		3.3.1 A Rellich type theorem	5					
		3.3.2 LAP bound	7					
		3.3.3 Microlocal bounds and LAP	0					
4	Rell	ich type theorems 6'	7					
	4.1	The case $\lambda_0 = \Sigma_2$ 6	7					
		4.1.1 Extended eigentransform for $\lambda_0 = \Sigma_2$	8					
		4.1.2 Negative slowly decaying effective potentials 6	8					
		4.1.3 Positive slowly decaying effective potentials 7	1					
		4.1.4 Homogeneous degree -2 effective potentials $\ldots \ldots \ldots .$	4					

	4.2	The ca 4.2.1 4.2.2 4.2.3 4.2.4 Model	ase $\lambda_0 > \Sigma_2$ Extended eigentransform for $\lambda_0 > \Sigma_2$ Negative slowly decaying effective potentials Positive slowly decaying effective potentials	81 82 87 88 89 90			
5	Res	olvent	asymptotics near a two-cluster threshold	99			
Ŭ	5.1	Rapid	ly decaying effective potentials	100			
	0.1	5.1.1	Two-cluster threshold resonances	100			
			5.1.1.1 The case $\lambda_0 = \Sigma_2$	100			
			5.1.1.2 The case $\lambda_0 > \Sigma_2$	105			
		5.1.2	Resolvent asymptotics near the lowest threshold	109			
		5.1.3	Resolvent asymptotics near higher two-cluster thresholds \ldots	124			
	5.2 Positive slowly decaying effective potentials						
	5.3	Resolv	vent asymptotics for physics models near two-cluster thresholds	145			
6	App	olicatio	ons	151			
	6.1	Negati	ive slowly decaying effective potentials $\ldots \ldots \ldots \ldots \ldots \ldots$	151			
		6.1.1	Sommerfeld's theorem	153			
		6.1.2	Elastic part of the scattering matrix at λ_0	156			
			6.1.2.1 Scattering for the one-body problem at zero energy, [DS1]	157			
			6.1.2.2 Elastic scattering for the N-body problem at λ_0	161			
			6.1.2.3 Elastic scattering at λ_0 , a 'geometric' approach	167			
			6.1.2.4 Elastic scattering at Σ_2	172			
		6.1.3	Scattering for physics models at a two-cluster threshold, case \sim				
			$\mathcal{A} = \mathcal{A}_1$	184			
	6.2	Effecti	r^{-2} potentials, atom-ion case	185			
	6.3	Non-ti	ransmission at a threshold, physics models $\dots \dots \dots \dots$	186			
		6.3.1	Proof of (6.98) , Case I)	188			
		6.3.2	Proof of (0.98) , Case II) \ldots	190			
		0.3.3	Proof of (0.98) , Case III)	195			
	64	0.3.4 Threel	All example of transmission	190			
	0.4	6 4 1	Finiteness of total cross-sections in atom-ion scattering	201			
		6.4.2	Total cross-sections at Σ_2 non-multiple two-cluster case	201			
		6.4.3	Total cross-sections at Σ_2 , non multiple two-cluster case	205			
		0.1.0	$ = \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{i=1}^{n} \sum_{i=1}^{n} \sum_{i=1}^{n} \sum_{j=1}^{n} \sum$	-00			

Bibliography

Chapter 1 Introduction, examples and notation

1.1 Scope and results

The spectral and scattering theory for the quantum mechanical one-body problem at zero energy is a well studied subject. The classical theory [JK, JN1, JN2] involves a real potential V(x) on \mathbb{R}^3 fastly decaying at least like $\mathcal{O}(|x|^{-\rho})$ for some $\rho > 2$. The slowly decaying case for which the decay rate $\rho \in (0, 2)$ requires additional conditions, roughly sign conditions [Ya1, Ya2, Na, FS, DS1, DS2]. The critical case is defined by $V(x) \approx C|x|^{-2}$, possibly with angular dependence, and the results depend on the coupling constant [Wa5, SW]. The obtained results for the above models are highly model and case sensitive and include possible existence of zero energy bound and/or resonance states as well as zero energy asymptotics of the resolvent and scattering matrix quantities (like scattering phase shifts). Thresholds of an N-body Schrödinger operator are eigenvalues of the sub-Hamiltonians. There exists much less literature on threshold spectral analysis for the N-body problem. We only mention [Wa2] on the resolvent expansion in a special case of the lowest threshold which is the bottom of the essential spectrum.

The goal of the present work is to present a systematic study of spectral and scattering theory for the quantum mechanical N-body problem at any negative twocluster threshold λ_0 , *i.e.*, λ_0 is an eigenvalue of (possibly several) sub-Hamiltonians associated with two-cluster decomposition, but not of those with three or more clusters. These restrictions on the nature of the considered threshold exclude the presence of the Efimov effect there. So for example for the (dynamical nuclei physics) 3-body problem, the threshold zero is excluded from our analysis, while all other thresholds for this model are negative of two-cluster type. (We shall later in this chapter give precise definitions.) Philosophically, the two-cluster threshold problem is amenable to simplification in terms of an effective one-body problem by the Feshbach-Grushin dimension reduction method. This is indeed realized in [Wa2] for fastly decaying pair potentials for the case of the lowest threshold $\lambda_0 = \Sigma_2$ (assumed non-multiple). However in the present work we extend the framework considerably, so that it covers the usual atom physics models (see Subsections 1.2.1 and 1.3.1) for which the slowly decaying nature of the Coulomb pair potentials requires refined analysis. Also we include the cases where the two-cluster threshold $\lambda_0 > \Sigma_2$ as well as multiple two-cluster and degenerate eigenvalue cases, which also call for refined analysis, in particular micro-local analysis.

One main ingredient which enables us to attain the goal of spectral analysis at any two-cluster threshold is the Mourre's estimate for the Hamiltonian with one threshold removed. For a given two-cluster threshold λ_0 , the restriction of the total Hamiltonian onto the orthogonal complement of the associated spectral subspace is a non-local N-body Hamiltonian for which λ_0 is no longer a threshold. We essentially prove the Mourre's estimate at λ_0 for this reduced Hamiltonian and deduce the limiting absorption principles and micro-local resolvent estimates. The limiting absorption principles are used to construct an appropriate Grushin problem such that we can reduce the two-cluster problem to an effective one-body problem near an arbitrary two-cluster threshold.

To be more concrete let us now consider the dynamical nuclei physics model from Subsection 1.2.1 (see (1.13)) and assume that the particle dimension n = 3. Let a two-cluster decomposition $a = (C_1, C_2)$ of N particles be given. We then write the full Hamiltonian as

$$H = H^1 \otimes 1 \otimes 1 + 1 \otimes H^2 \otimes 1 + 1 \otimes 1 \otimes p_a^2 + I_a$$

where H^k , k = 1, 2, are cluster-Hamiltonians (defined like for H in their center of mass frames), p_a^2 is the inter-cluster kinetic energy Hamiltonian and I_a is the *intercluster potential*. Suppose $\lambda_a = \lambda_0$ is an eigenvalue of the sub-Hamiltonian $H^a =$ $H^1 \otimes 1 + 1 \otimes H^2$ (λ_0 being of two-cluster type, see (1.27)). Picking a corresponding orthonormal basis $\varphi_1^a, \ldots \varphi_m^a \in \ker(H^a - \lambda_0) \subset L^2(\mathbf{X}^a), m = m_a$, the *effective intercluster potential* is the $m \times m$ matrix-valued function in the relative position variable of the two clusters, viz. $R = R_1 - R_2$,

$$V(R)_{kl} := \langle \varphi_k^a, I_a \varphi_l^a \rangle_{L^2(\mathbf{X}^a)} = Q_1 Q_2 \delta_{kl} |R|^{-1} + Q_{kl}(\widehat{R}) |R|^{-2} + \mathcal{O}(|R|^{-3}).$$
(1.1)

Here Q_1 and Q_2 are the total charge of the particles in the clusters C_1 and C_2 , respectively, and δ_{kl} is the Kronecker symbol. In addition we denote by Q_a the matrix-valued homogeneous potential (Q_{kl}) and $\hat{R} = R/|R|$. Let P^a denote the orthogonal (rank m) projection onto ker $(H^a - \lambda_0)$ in $L^2(\mathbf{X}^a)$. Then obviously $\Pi^a = P^a \otimes 1$ projects onto the span of functions of the form $\varphi^a \otimes f_a$, $\varphi^a \in \text{ker}(H^a - \lambda_0)$, in $L^2(\mathbf{X})$.

In terms of (1.1) a relevant classification reads:

Case 1 (slowly decaying case) $Q_1Q_2 \neq 0$.

Case 2 (critically decaying case) $Q_1Q_2 = 0$ and the function $Q_a \neq 0$.

Case 3 (fastly decaying case) $Q_1Q_2 = 0$ and the function $Q_a = 0$.

In general λ_0 might be a multiple two-cluster threshold, which might suggest that we group the set of thresholds a for which $\lambda_0 \in \sigma_{\rm pp}(H^a)$, say denoted by \widetilde{A} , into $\widetilde{A} = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3$ specified as follows.

- \mathcal{A}_1 : the effective inter-cluster interaction is to leading order attractive Coulombic, i.e. $Q_1Q_2 < 0$.
- \mathcal{A}_2 : the effective inter-cluster interaction is to leading order repulsive Coulombic, i.e. $Q_1Q_2 > 0$.

1.1. Scope and results

 \mathcal{A}_3 : the effective inter-cluster interaction is $\mathcal{O}(|x_a|^{-2})$, i.e. $Q_1Q_2 = 0$.

Clearly the elements of $\mathcal{A}_1 \cup \mathcal{A}_2$ are classified as Case 1, while the elements of \mathcal{A}_3 are classified either as Case 2 or Case 3. This motivates the splitting $\mathcal{A}_3 = \mathcal{A}_3^{cd} \cup \mathcal{A}_3^{fd}$ by specifying

$$\mathcal{A}_3^{\mathrm{cd}} = \{ a \in \mathcal{A}_3 | Q_a \neq 0 \} \text{ and } \mathcal{A}_3^{\mathrm{fd}} = \{ a \in \mathcal{A}_3 | Q_a = 0 \}$$

corresponding to Case 2 and Case 3, respectively.

For any $a \in \mathcal{A}_3$ there are computable numbers $s_a \geq 1$ and $d_a \in \mathbb{N}_0$ determined by spectral properties of the vector-valued Schrödinger operator on the unit-sphere \mathbb{S}^2 with the matrix-valued potential Q_a (see Section 4.3) such that in terms of standard weighted L^2 -space and weighted Sobolov-space notation (see Subsection 1.4.1 for definitions), referring here to our most general result (see Section 4.3):

Theorem 1.1. For any two-cluster threshold λ_0 :

1) The space of locally H^1 solutions to $(H - \lambda_0)u = 0$ in

$$\sum_{a \in \mathcal{A}_1} \Pi^a L^2_{-3/4} + \sum_{a \in \mathcal{A}_2} \Pi^a L^2_{(-3/2)^+} + \sum_{a \in \mathcal{A}_3} \Pi^a L^2_{(-\min\{3/2, s_a\})^+} + L^2_{-1/2}, \quad (1.2a)$$

say denoted by \mathcal{E} , has finite dimension.

- 2) If $\mathcal{A}_3 = \emptyset$, then $\mathcal{E} \subset H^1_{\infty}$.
- 3) The dimension of the space of resonance states

$$n_{\text{res}} = \dim \left(\mathcal{E} / \ker(H - \lambda_0)_{|H^1} \right) \le \sum_{a \in \mathcal{A}_3} d_a.$$

4) The numbers $s_a = 3/2$ and $d_a = m_a$ for any $a \in \mathcal{A}_3^{\text{fd}}$. In particular if $\mathcal{A}_3^{\text{cd}} = \emptyset$, then (1.2a) simplifies as

$$\sum_{a \in \mathcal{A}_1} \Pi^a L^2_{-3/4} + \sum_{a \in \mathcal{A}_2 \cup \mathcal{A}_3^{\text{fd}}} \Pi^a L^2_{(-3/2)^+} + L^2_{-1/2}, \qquad (1.2b)$$

and 3) reads

$$n_{\rm res} \le \sum_{a \in \mathcal{A}_3^{\rm fd}} m_a$$

One may view Theorem 1.1 as a version of the well-known Rellich theorem for non-threshold energies [AIIS, Theorem 1.4] (see also Theorem 3.12) although the above analogue at a two-cluster threshold is considerably more complex. For example the analogue of (1.2a) and (1.2b) in the continuous spectrum away from thresholds reads $L^2_{-1/2}$, and we note that for almost all (probably valid for all) such energies the space of generalized eigenfunctions in $L^2_{-1/2-\epsilon}$ (for any $\epsilon > 0$) has infinite dimension [Sk6]. In comparison in the context of Theorem 1.1 with $\tilde{\mathcal{A}} = \mathcal{A}_1 = \{a\}$ we show that the space of generalized eigenfunctions at λ_0 in $\Pi^a L^2_{-3/4-\epsilon} + L^2_{-1/2-\epsilon}$ is infinitely dimensional (see Theorem 6.13 2)). The proof of Theorem 1.1 is complicated, in fact we give a full proof only under the two technical conditions (4.68a) and (4.68b) (not to be elaborated on in this introduction), treating the general case in a somewhat sketchy fashion.

One of the threshold phenomena indicated by Theorem 1.1 is the possible existence of resonance states combined freely with the possible existence of L^2 eigenfunctions at the two-cluster threshold λ_0 . This is completely analogous to the situation for the one-body problem for fastly decaying potentials [JK] (exhibiting a somewhat similar sophisticated Rellich theorem at zero energy), giving rise to the classification into a Regular Case (where λ_0 is neither an eigenvalue nor a resonance of H) and Exceptional Cases 1,2 and 3 (see Subsection 5.1.1.1). The resolvent asymptotics at zero energy for the one-body problem is determined by this classification. It is a separate issue for us to obtain similar resolvent asymptotics at λ_0 in the present framework. However our analysis is not complete, mainly due to lack of strong decay of Coulombic potentials hampering the analysis. Of course the Regular Case is the easiest case and we shall actually treat this with $\tilde{\mathcal{A}} = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3^{\rm fd}$ (see Theorem 5.43). For the Exceptional Cases 1 and 3 (defined by the presence of a resonance) we show the following result.

Theorem 1.2 (Exceptional point of 1st or 3rd kind). Let λ_0 be any two-cluster threshold for which $n_{\text{res}} \geq 1$, i.e. λ_0 is a resonance of H. Suppose the technical conditions (4.68a) and (4.68b) (referred to above),

$$\operatorname{ran} \Pi_H \subset L_t^2 \text{ for some } t > 3/2, \tag{1.3}$$

where Π_H is the orthogonal projection onto ker $(H - \lambda_0)$ (i.e. the eigenprojection if λ_0 is an eigenvalue of H and zero otherwise), and suppose

$$\widetilde{\mathcal{A}} = \mathcal{A}_2 \cup \mathcal{A}_3^{\text{fd}}.\tag{1.4}$$

Then the following asymptotics hold for $R(\lambda_0 + z) = (H - \lambda_0 + z)^{-1}$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ in $\mathcal{Z}_{\pm} = \{\operatorname{Re} z \ge 0, \pm \operatorname{Im} z > 0\}$ and for some $\epsilon = \epsilon(s) > 0$.

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \frac{i}{\sqrt{z}} \sum_{j=1}^{n_{res}} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
(1.5)

Here $\{u_1, \ldots, u_{n_{\text{res}}}\} \subset H^1_{(-1/2)^-}$ is a basis of resonance states of H being independent of the choice of the sign of \mathcal{Z}_{\pm} .

Among the appearing conditions (1.3) is the 'unpleasant one'. It is an implicit (possibly redundant) condition appearing as an artifact of our methods. If λ_0 is exceptional point of 1st kind, (1.3) is obviously fulfilled since then $\Pi_H = 0$. Our Theorem 5.43 as well the above Theorem 1.2 require explicitly $\mathcal{A}_3^{cd} = \emptyset$.

Under a spectral condition for certain elements of $\mathcal{A}_3^{\text{cd}}$ (those for which (1.8) in Subsection 1.1.1 is violated) oscillatory behaviour of the resolvent near the twocluster threshold is expected. This is expected thanks to arguments of [SW].

1.1.1 Applications to threshold scattering

Here we briefly outline our main applications in Chapter 6. One of our results concerns the following generalization of a result from [DS1, DS2] (see also [Fr]).

Theorem 1.3. Suppose that $\lambda_0 = \Sigma_2$ is a two-cluster threshold, the technical conditions (4.68a) and (4.68b),

$$\mathcal{A} = \mathcal{A}_1,\tag{1.6}$$

and suppose the Regular Case (i.e. that λ_0 is neither an eigenvalue nor a resonance of H). Let C denote the set of scattering channels $\alpha = (a, \lambda_0, \varphi_\alpha)$ with $a \in \widetilde{\mathcal{A}}$ (note that λ_0 is a simple eigenvalue of H^a). For $\alpha, \beta \in C$ the element of the scattering matrix $S_{\beta\alpha}(\lambda)$ (modelled after [DS1, DS2]) is well-defined for λ slightly above λ_0 and possessing a strong limit as $\lambda \to (\lambda_0)_+$. Moreover the singular support of the limiting element $S_{\beta\alpha}(\lambda_0)$ fulfills

sing supp
$$S_{\beta\alpha}(\lambda_0)$$

$$\begin{cases} \subset \{(\omega, \omega') \mid \omega \cdot \omega' = -1\} & \text{for } \alpha = \beta, \\ = \emptyset & \text{for } \alpha \neq \beta. \end{cases}$$

The proof of Theorem 1.3 may be considered as an extension of the one used in [DS1, DS2] to obtain a similar 'semi-classical' result on the scattering matrix $S_{\text{cou}}(E)$ for the one-body problem with an attractive Coulomb potential. See also [Va1, Va2] for N-body scattering matrices in the short-range case.

Another result under (1.6) concerns the difference

$$S_{\alpha\alpha}(\lambda) - S_{\rm cou}(\lambda - \lambda_0); \quad \lambda \in [\lambda_0, \lambda_0 + \delta].$$

Under conditions, in particular including the non-multiple property $\# \mathcal{A} = 1$ (primarily used to simplify the presentation) however covering the case where the twocluster threshold $\lambda_0 > \Sigma_2$, we show that this difference is a 'partial smoothing operator' (see Theorem 6.10 and Remark 6.11 3)). Yet another result is a characterization of the limiting element $S_{\alpha\alpha}(\lambda_0)$ given by asymptotics in terms of appropriate 'channel quasi-modes' (see Theorem 6.13 2)).

This leads to another subject of interest, more precisely *non-transmission* at λ_0 . This is a 'geometric concept' amounting to the feature

$$\|1 - S_{\alpha\alpha}(\lambda)^* S_{\alpha\alpha}(\lambda)\| \to 0 \text{ for } \lambda \to (\lambda_0)_+.$$
(1.7)

We derive a formula under (1.6) (see Corollary 6.15) indicating that transmission does occur in this case if $\lambda_0 > \Sigma_2$ (see also Remarks 6.16 and 6.29 4)).

In contrast to the attractive slowly decaying case we do prove non-transmission in the following three cases (assuming as above in all cases the non-multiple property $\#\widetilde{\mathcal{A}} = 1$):

- I) Effective repulsive Coulombic case, i.e. $\widetilde{\mathcal{A}} = \mathcal{A}_2$.
- II) $I_a(x^a = 0) = 0$, 'above the Hardy limit' and λ_0 be regular.
- **III)** $I_a(x^a = 0) = 0$ and 'fastly decaying case', i.e. $\widetilde{\mathcal{A}} = \mathcal{A}_3^{\text{fd}}$, and λ_0 be 'maximally exceptional of 1st kind'.

A special case of II) is that $\widetilde{\mathcal{A}} = \mathcal{A}_3^{\text{fd}}$ and λ_0 neither be an eigenvalue nor a resonance. The notions in II) and III) are in general given as follows (see also Section 6.3). The phrase 'above the Hardy limit' refers to a spectral property of the vector-valued Schrödinger operator on the unit-sphere \mathbb{S}^2 with the matrix-valued potential Q_a (writing $\widetilde{\mathcal{A}} = \{a\}$), more precisely

$$\inf \sigma \left(-\Delta_{\theta} + Q_a(\theta) \right) > -1/4. \tag{1.8}$$

For III) the potential $Q_a = 0$ and 'maximally exceptional of 1st kind' refers to Exceptional Case 1 and the condition $n_{\rm res} = m_a = \dim \ker(H^a - \lambda_0)$. We note for comparison that if $m_a > 1$ and $n_{\rm res} = \{1, \ldots, m_a - 1\}$ then indeed transmission can occur for λ_0 be exceptional of 1st kind (see Subsection 6.3.4).

The last subject of interest concerns total cross-sections for atom-ion scattering. It is an observed phenomenon at the very beginning of the quantum mechanics that when there is no dipole moment for the atom, the total cross-sections are finite. A mathematical proof for this physics folklore is given in [JKW]. The operator under consideration is a special case of the dynamical nuclei physics model from Subsection 1.2.1 with the particle dimension n = 3. Assume $\lambda_0 = \Sigma_2$ is a two-cluster threshold, the technical conditions (4.68a) and (4.68b), (1.3) and $\widetilde{\mathcal{A}} = \mathcal{A}_3^{\text{fd}}$. It is known from [JKW] that for any channel $\alpha = (a, \lambda_0, \varphi_\alpha), a \in \widetilde{\mathcal{A}}$, and any incident direction $\omega \in \mathbb{S}^2$,

the total cross-section $\sigma_{\alpha}(\lambda, \omega)$

is finite for non-threshold λ 's above λ_0 . In the present work we derive bounds and asymptotics of this quantity as $\lambda \to (\lambda_0)_+$ (see Section 6.4). The result depends on whether λ_0 is regular or exceptional of 2nd kind (yielding bounded asymptotics, see Theorem 6.33) or if λ_0 is of 1st or 3rd kind (yielding $(\lambda - \lambda_0)^{-1}$ type unbounded asymptotics, see Theorem 6.34). Our proof relies on the derivation of Theorem 1.2.

1.2 Many-body Schrödinger operators

Let H denote the many-body Schrödinger operator obtained by the removal of the center of mass from the total Hamiltonian

$$\widetilde{H} = -\sum_{j=1}^{N} \frac{1}{2m_j} \Delta_{x_j} + \sum_{1 \le i < j \le N} V_{ij}(x_i - x_j), \quad x_j \in \mathbb{R}^n,$$
(1.9)

where x_j and m_j denote the position and mass of the j'th particle. The pair potentials V_{ij} are assumed to be real and relatively compact with respect to $-\Delta$ in $L^2(\mathbb{R}^n)$, and they satisfy for some $\rho > 0$ the condition

$$|V_{ij}(y)| \leq C_{ij}|y|^{-\rho}$$
 for $y \in \mathbb{R}^n$ with $|y| > R$,

for some R > 0. However we shall need some extra regularity. It is convenient to use the following condition.

Condition 1.4. There exists $\rho > 0$ such that for all pair potentials V_{ij} there is a splitting $V_{ij} = V_{ij}^{(1)} + V_{ij}^{(2)}$, where

1.2. Many-body Schrödinger operators

(1) $V_{ij}^{(1)}$ is smooth and

$$\partial_y^{\alpha} V_{ij}^{(1)}(y) = \mathcal{O}\big(|y|^{-\rho - |\alpha|}\big). \tag{1.10}$$

(2) $V_{ij}^{(2)}$ is compactly supported and

$$V_{ij}^{(2)}(-\Delta+1)^{-1}$$
 is compact on $L^2(\mathbb{R}^n_y)$. (1.11)

The Hamiltonian H is regarded as a self-adjoint operator on $L^2(\mathbf{X})$, where \mathbf{X} is the n(N-1) dimensional real vector space $\mathbf{X} := \{\sum_{j=1}^N m_j x_j = 0\}$. Let \mathcal{A} denote the set of all cluster decompositions of the N-particle system. The notation a_{\max} and a_{\min} refers to the 1-cluster and N-cluster decompositions, respectively. Let for $a \in \mathcal{A}$ the notation #a denote the number of clusters in a. For $i, j \in \{1, \ldots, N\}, i < j$, we denote by (ij) the (N-1)-cluster decomposition given by letting $C = \{i, j\}$ form a cluster and all other particles $l \notin C$ form 1-particle clusters. We write $(ij) \subset a$ if iand j belong to the same cluster in a. More general, we write $b \subset a$ if each cluster of b is a subset of a cluster of a. If a is a k-cluster decomposition, $a = (C_1, \ldots, C_k)$, we let

$$\mathbf{X}^{a} = \left\{ x \in \mathbf{X} \mid \sum_{l \in C_{j}} m_{l} x_{l} = 0, j = 1, \dots, k \right\} = \mathbf{X}^{C_{1}} \oplus \dots \oplus \mathbf{X}^{C_{k}},$$

and

$$\mathbf{X}_a = \left\{ x \in \mathbf{X} \mid x_i = x_j \text{ if } i, j \in C_m \text{ for some } m \in \{1, \dots, k\} \right\}.$$

Note that $a \subset b \Leftrightarrow \mathbf{X}^a \subset \mathbf{X}^b$. Moreover \mathbf{X}^a and \mathbf{X}_a give an orthogonal decomposition for \mathbf{X} equipped with the quadratic form

$$q(x) = \sum_{j} 2m_j |x_j|^2, \qquad x \in \mathbf{X}.$$

For $x \in \mathbf{X}$, we have the corresponding orthogonal decomposition: $x = x^a + x_a$ with $x^a = \pi^a x \in \mathbf{X}^a$ and $x_a = \pi_a x \in \mathbf{X}_a$.

With this notation, the many-body Schrödinger operator H introduced above can be written in the form

$$H = H_0 + V$$

where $H_0 = p^2$ is (minus) the Laplace-Beltrami operator on the Euclidean space (\mathbf{X}, q) and $V = V(x) = \sum_{a=(ij)\in\mathcal{A}} V_a(x^a)$ with $V_a(x^a) = V_{ij}(x_i - x_j)$ for the (N-1)-cluster decomposition a = (ij). More precisely, for example,

$$x^{(12)} = \left(\frac{m_2}{m_1 + m_2}(x_1 - x_2), -\frac{m_1}{m_1 + m_2}(x_1 - x_2), 0, \dots, 0\right).$$

We note the following geometric properties for $N \geq 3$: For all $a, b \in \mathcal{A}$ with #a = 2, #b = N - 1 and $b \not\subset a$

$$\operatorname{ran}\left(\pi^{b}\pi^{a}\right) = \operatorname{ran}\,\pi^{b},\tag{1.12a}$$

$$\pi^b : \mathbf{X}_a \to \mathbf{X}^b$$
 is bijective. (1.12b)

1.2.1 Principal example, dynamical nuclei

Consider a system of ${\cal N}$ particles interacting by Coulomb forces. The Hamiltonian then reads

$$H = -\sum_{j=1}^{N} \frac{1}{2m_j} \Delta_{x_j} + \sum_{1 \le i < j \le N} q_i q_j |x_i - x_j|^{-1}, \quad x_j \in \mathbb{R}^n, \ n \ge 3,$$
(1.13)

where x_j , m_j and q_j denote the position, mass and charge of the *j*'th particle, respectively. *H* is regarded as a self-adjoint operator in $L^2(\mathbf{X})$ (with mass center removed).

Let us consider a two-cluster decomposition $a = (C_1, C_2)$. For convenience assume $C_1 = \{1, \ldots, J\}$ and $C_2 = \{J + 1, \ldots, N\}$. We can write

$$H = H^1 \otimes 1 \otimes 1 + 1 \otimes H^2 \otimes 1 + 1 \otimes 1 \otimes p_a^2 + I_a$$

where H^k , k = 1, 2, are cluster-Hamiltonians (defined similarly in their center of mass frames) and

$$I_a = \sum_{i \in C_1, j \in C_2} q_i q_j |x_i - x_j|^{-1}.$$

To expand I_a we let for k = 1, 2

$$Q_{k} = \sum_{j \in C_{k}} q_{j}, M_{k} = \sum_{j \in C_{k}} m_{j},$$

$$R_{k} = R_{k}(x) = \sum_{j \in C_{k}} \frac{m_{j}}{M_{k}} x_{j}, \widetilde{Q}_{k} = \widetilde{Q}_{k}(x^{C_{k}}) = \sum_{j \in C_{k}} q_{j}(x_{j} - R_{k})$$

$$M = M_{1} + M_{2}, R = R_{1} - R_{2},$$

and we decompose for all $x \in \mathbf{X}$

$$x = x^{C_1} + x^{C_2} + x_a,$$

$$x^{C_1} = (x_1 - R_1, \dots, x_J - R_1, 0, \dots, 0) \in \mathbf{X}^{C_1},$$

$$x^{C_2} = (0, \dots, 0, x_{J+1} - R_2, \dots, x_N - R_2) \in \mathbf{X}^{C_2},$$

$$x_a = \left(\frac{M_2}{M}R, \dots, \frac{M_2}{M}R, -\frac{M_1}{M}R, \dots, -\frac{M_1}{M}R\right) \in \mathbf{X}_a.$$

Note that indeed the center of charge \widetilde{Q}_k is a function of x^{C_k} .

Consequently we can expand for $i \in C_1$ and $j \in C_2$

$$|x_i - x_j|^{-1} = |R|^{-1} - \frac{R}{|R|^3} \cdot \left((x_i - R_1) - (x_j - R_2) \right) + \mathcal{O}(|R|^{-3}) |x^a|^2.$$

This is in the regime $|R| \to \infty$ and $|x_i - R_1| + |x_j - R_2| \le \frac{1}{2}|R|$. Whence in turn we obtain for $|R| \to \infty$

$$I_a = Q_1 Q_2 |R|^{-1} + \frac{R}{|R|^3} \cdot \left(Q_1 \widetilde{Q}_2(x^{C_2}) - Q_2 \widetilde{Q}_1(x^{C_1}) \right) + \mathcal{O}(|R|^{-3}) |x^a|^2, \qquad (1.14)$$

which leads to various cases. We use the notation φ^k , k = 1, 2, to denote a cluster bound state (for the cluster Hamiltonian H^k) and $\langle \cdot, \cdot \rangle_k$ to denote the corresponding cluster inner product. The *effective potential*

$$V(R) := \langle \varphi^1 \otimes \varphi^2, I_a \varphi^1 \otimes \varphi^2 \rangle_{L^2(\mathbf{X}^a)}.$$

Case 1 $V \approx |R|^{-1}$: $Q_1 Q_2 \neq 0$.

Case 2 $V \approx |R|^{-2}$:

Subcase 2a $Q_1 \neq 0, Q_2 = 0 \text{ and } \langle \varphi^2, \widetilde{Q}_2 \varphi^2 \rangle_2 \neq 0.$

Subcase 2b $Q_2 \neq 0, Q_1 = 0 \text{ and } \langle \varphi^1, \widetilde{Q}_1 \varphi^1 \rangle_1 \neq 0.$

Case 3 $V = O(|R|^{-3})$:

Subcase 3a $Q_1 = Q_2 = 0.$

Subcase 3b $Q_1 \neq 0, Q_2 = 0 \text{ and } \langle \varphi^2, \widetilde{Q}_2 \varphi^2 \rangle_2 = 0.$

Subcase 3c $Q_2 \neq 0, Q_1 = 0 \text{ and } \langle \varphi^1, \widetilde{Q}_1 \varphi^1 \rangle_1 = 0.$

Note that for Subcases 2a, 3a and 3b, assuming sufficient decay of the cluster bound states, the effective potential

$$V(R) = \langle \varphi^1 \otimes \varphi^2, I_a \varphi^1 \otimes \varphi^2 \rangle_{L^2(\mathbf{X}^a)} = Q_1 \frac{R}{|R|^3} \cdot \langle \varphi^2, \widetilde{Q}_2 \varphi^2 \rangle_2 + \mathcal{O}(|R|^{-3}).$$
(1.15)

Whence indeed $V \approx |R|^{-2}$ at infinity in Subcase 2a, while indeed $V = \mathcal{O}(|R|^{-3})$ for Subcases 3a and 3b. We can argue similarly for Subcases 2b and 3c. Note also that $|R|^{-2}$ is the critical decay rate for threshold analysis, cf. [SW]. Case 1 is the slowly decaying case. In Case 1 the potential $V \approx |R|^{-1}$, and V is said to be slowly decaying. For $Q_1Q_2 < 0$ and $Q_1Q_2 > 0$ the one-body results of [FS] and [Na, Ya2] will be useful, respectively. In Case 3 the effective potential is said to be fastly decaying and other one-body results/techniques will be useful, cf. for example [JK]. Case 2 (the critical case) is different and rather 'rich'.

A detailed analysis of the structure of a class of generalized eigenfunctions at a two-cluster threshold, possibly a multiple and/or a non-simple two-cluster threshold, will be carried out for physical models in Section 4.3. (See (1.27) for the definition of a 'two-cluster threshold'.)

From the derivation it follows that it could happen that the second term $\mathcal{O}(|R|^{-3})$ of (1.15) actual has homogeneity -3 at infinity. For example this happens for Subcase 3a exactly when the moments $\widetilde{R}_1 := \langle \varphi^1, \widetilde{Q}_1 \varphi^1 \rangle_1 \neq 0$ and $\widetilde{R}_2 := \langle \varphi^2, \widetilde{Q}_2 \varphi^2 \rangle_2 \neq 0$ due to the computation for this case

$$\mathcal{O}(|R|^{-3}) = |R|^{-5} (|R|^2 \widetilde{R}_1 \cdot \widetilde{R}_2 - 3(R \cdot \widetilde{R}_1)(R \cdot \widetilde{R}_2)) + \mathcal{O}(|R|^{-4})$$

If certain 'moments' vanish for Subcases 3a and 3b the order of the second term of (1.15) is of the form $\mathcal{O}(|R|^{-4})$, cf. [JKW, Appendix A]. In Chapter 5 we shall obtain leading order resolvent expansions for Case 3 without distinguishing between whether the homogeneous -3 term vanishes or not. In Section 6.4 we shall study a case, where in fact the effective potential is (at least) of order $\mathcal{O}(|R|^{-4})$. In the same section an explicit calculation of the Hamiltonian is given in terms of so-called clustered atomic coordinates.

Strictly speaking the distinction between Cases 2 and 3 as defined above makes best sense for a simple two-cluster threshold and we will not use this classification in the non-simple case. Rather in the general possibly non-simple case one needs the following (slightly) different definition, see Section 4.3 for further details. Let λ_a be a non-threshold eigenvalue of the sub-Hamiltonian $H^a = H^1 \otimes 1 + 1 \otimes H^2$ (more precisely, we will need $\lambda_a \in \mathcal{T}_2$, see (1.27)). Picking an orthonormal basis $\varphi_1^a, \ldots \varphi_m^a \in L^2(\mathbf{X}^a), m = m_a$ (being one or possibly bigger), in the range of the corresponding eigenprojection, the *effective potential* is the $m \times m$ -matrix-valued function in the variable $R = R_1 - R_2$

$$V(R)_{kl} := \langle \varphi_k^a, I_a \varphi_l^a \rangle_{L^2(\mathbf{X}^a)} = Q_1 Q_2 \delta_{kl} |R|^{-1} + Q_{kl}(\widehat{R}) |R|^{-2} + \mathcal{O}(|R|^{-3}).$$
(1.16)

Here δ_{kl} is the Kronecker symbol and $\widehat{R} = R/|R|$. In terms of (1.16) the more general (and correct) classification reads:

Case 1 $Q_1Q_2 \neq 0.$

Case 2 $Q_1Q_2 = 0$ and the matrix-valued function $Q_a = (Q_{kl}) \neq 0$.

Case 3 $Q_1Q_2 = 0$ and the matrix-valued function $Q_a = (Q_{kl}) = 0$.

1.3 *N*-body Schrödinger operators with infinite mass nuclei

In the case of $M \geq 1$ infinite mass nuclei located at $R_m \in \mathbb{R}^n$, $m = 1, \ldots, M$, the Hamiltonian reads

$$H = -\sum_{j=1}^{N} \frac{1}{2m_j} \Delta_{x_j} + \sum_{1 \le i < j \le N} V_{ij}(x_i - x_j) + \sum_{1 \le j \le N, \ 1 \le m \le M} V_{jm}^{\text{ncl}}(x_j - R_m), \quad (1.17)$$

where we impose similar conditions on V_{jm}^{ncl} as for V_{ij} in Condition 1.4. The one-body problem N = 1 is included in (1.17) (the middle term is absent in that case). The configuration space reads $\mathbf{X} = \mathbb{R}^{nN}$, and we use the metric q as before. The 'electronelectron' interaction $V_{ii}(x_i - x_i)$ takes as before the form $V_a(x^a)$ where $x^a = \pi^a x_i$ a = (ij), is the orthogonal projection of x onto an n-dimensional subspace. Similarly the 'electron-nuclei' interaction $\sum_{1 \le m \le M} V_{jm}^{\text{ncl}}(x_j - R_m)$ takes the form $V_a(x^a)$ where again $x^a = \pi^a x$, a = a(j), is the orthogonal projection of x onto an n-dimensional subspace (let $x^a = (0, \ldots, 0, x_i, 0, \ldots, 0)$, i.e. all other coordinates than the j'th are put equal to zero). Rather than using the cluster decompositions to label a family of 'subspaces of internal motion' $\{\mathbf{X}^a\}$ similar to those considered in Section 1.2 we prefer henceforth to appeal to abstract labeling. Precisely we consider the smallest finite family $\{\mathbf{X}^a \mid a \in \mathcal{A}\}$ of subspaces of **X** which is stable under addition and which contains $\{0\}$ and the *n*-dimensional subspaces discussed above. See Section 1.4, and see [DG, Section 5.1] for a discussion of the abstract N-body problem. On the other hand there is a concrete description of the index set \mathcal{A} and this family $\{\mathbf{X}^a \mid a \in \mathcal{A}\}\$ which can be useful to have in mind: Consider $a = (C_1, \ldots, C_p)$ where the sets C_q are disjoint subsets of $\{1, \ldots, N\}$. For $p \ge 2$ and q < p we have $\#C_q \ge 2$ and we let $\mathbf{X}^{C_q} = \{x \in \mathbf{X} \mid x_j = 0 \text{ if } j \notin C_q \text{ and } \sum_{i \in C_q} m_i x_i = 0\}$. Either similarly $\mathbf{X}^{C_p} = \{x \in \mathbf{X} \mid x_j = 0 \text{ if } j \notin C_p \text{ and } \sum_{i \in C_p} m_i x_i = 0\}$ (in that case we have $\#C_p \geq 2$) or $\mathbf{X}^{C_p} = \{x \in \mathbf{X} \mid x_j = 0 \text{ if } j \notin C_p\}$. In both cases let correspondingly $\mathbf{X}^{a} = \mathbf{X}^{C_{1}} \oplus \cdots \oplus \mathbf{X}^{C_{p}}$. Moreover we supplement by writing $\mathbf{X}^{a_{\min}} = \{0\}$ where, for

example, $a_{\min} := \emptyset$. This is a concrete labeling of the family of subspaces of internal motion.

The ordering of subspaces yields an ordering of the abstract set of indices \mathcal{A} , by definition $a \subset b \Leftrightarrow \mathbf{X}^a \subset \mathbf{X}^b$. We denote $\mathbf{X} = \mathbf{X}^{a_{\max}}$ and $\mathbf{X}^a + \mathbf{X}^b = \mathbf{X}^{a \cup b}$. The orthogonal complement of \mathbf{X}^a is denoted by \mathbf{X}_a . To have a uniform language we refer to the indices $a \in \mathcal{A}$ as 'cluster decompositions'. The length of a chain of cluster decompositions $a_1 \subsetneq \cdots \subsetneq a_k$ is the number k. This chain is said to connect $a = a_1$ and $b = a_k$. The maximal length of all chains connecting a given $a \in \mathcal{A} \setminus \{a_{\max}\}$ and a_{\max} is denoted by #a. We define $\#a_{\max} = 1$ and note that $\#a_{\min} = N + 1$. We say $a \in \mathcal{A}$ is k-cluster if #a = k.

We note the following geometric properties for $N \ge 2$: For all $a, b \in \mathcal{A}$ with #a = 2, #b = N and $b \not\subset a$

$$\operatorname{ran}\left(\pi^{b}\pi^{a}\right) = \{0\} \text{ or } \operatorname{ran}\left(\pi^{b}\pi^{a}\right) = \operatorname{ran}\pi^{b}, \qquad (1.18a)$$

$$\pi^b : \mathbf{X}_a \to \mathbf{X}^b$$
 is bijective. (1.18b)

1.3.1 Principal example, fixed nuclei

Consider a system of N n-dimensional particles, $n \ge 3$, interacting by Coulomb forces. The Hamiltonian (1.17) then reads

$$H = -\sum_{j=1}^{N} \frac{1}{2m_j} \Delta_{x_j} + \sum_{1 \le i < j \le N} q_i q_j |x_i - x_j|^{-1} + \sum_{1 \le j \le N, \ 1 \le m \le M,} q_j q_m^{\text{ncl}} |x_j - R_m|^{-1}, \ (1.19)$$

where x_j , m_j and q_j denote the position, mass and charge of the *j*'th 'electron', and R_m and $q_m^{\rm ncl}$ are the position and charge of the *m*'th 'nucleus'.

Consider the two-cluster decomposition a = (C), $C = \{1, \ldots, N-1\}$, meaning $\mathbf{X}^a = \{x = (x_1, \ldots, x_N) \in \mathbf{X} = \mathbb{R}^{nN} \mid x_N = 0\}$. Letting $R = x_N$ we write $H = H^1 \otimes 1 + 1 \otimes p_R^2 + I_a$ where H^1 is the cluster-Hamiltonian (i.e. the Hamiltonian for the first N - 1 electrons) and

$$I_a = \sum_{1 \le i \le N-1} q_i q_N |x_i - R|^{-1} + \sum_{1 \le m \le M,} q_N q_m^{\text{ncl}} |R - R_m|^{-1}.$$

Introducing

$$Q = \sum_{1 \le j \le N-1} q_j + \sum_{1 \le m \le M} q_m^{\text{ncl}},$$
$$\widetilde{Q} = \widetilde{Q}(x^a) = \sum_{1 \le j \le N-1} q_j x_j,$$
$$\widetilde{Q}^{\text{ncl}} = \sum_{1 \le m \le M} q_m^{\text{ncl}} R_m,$$

the asymptotics of I_a for $|R| \to \infty$ reads

$$I_{a} = q_{N}Q|R|^{-1} + q_{N}\frac{R}{|R|^{3}} \cdot \left(\widetilde{Q}(x^{a}) + \widetilde{Q}^{\mathrm{ncl}}\right) + \mathcal{O}\left(|R|^{-3}\right)\left(1 + |x^{a}|^{2}\right).$$
(1.20)

For the expectation in a cluster bound state $\varphi = \varphi^a(x^a)$ with sufficient decay we consequently obtain the asymptotics for $|R| \to \infty$

$$\langle \varphi, I_a \varphi \rangle_{L^2(\mathbf{X}^a)} = q_N Q |R|^{-1} + q_N \frac{R}{|R|^3} \cdot \left(\langle \varphi, \widetilde{Q} \varphi \rangle_{L^2(\mathbf{X}^a)} + \widetilde{Q}^{\mathrm{ncl}} \right) + \mathcal{O}\left(|R|^{-3} \right).$$
(1.21)

This leads to various cases.

Case 1 $q_N Q \neq 0$.

Case 2
$$q_N \langle \varphi, \hat{Q} \varphi \rangle_{L^2(\mathbf{X}^a)} \neq -q_N \hat{Q}^{\text{ncl}} \text{ and } Q = 0.$$

Case 3 $q_N = 0$, or $q_N \neq 0$, Q = 0 and $\langle \varphi, \widetilde{Q} \varphi \rangle_{L^2(\mathbf{X}^a)} = -\widetilde{Q}^{\mathrm{ncl}}$.

Case 1 is the slowly decaying case, Case 2 is the critical case and Case 3 is the fastly decaying case. Strictly speaking this classification makes best sense for φ being unique, i.e. for the simple case; in the non-simple case one needs a slightly different terminology, see Subsection 1.2.1 and Section 4.3.

1.4 Generalized *N*-body Schrödinger operators

Motivated by Sections 1.2 and 1.3 we discuss the abstract N-body problem, cf. [DG,Section 5.1]. Let $\mathbf{X} \neq \{0\}$ be a real finite dimensional vector space with an inner product q. We consider a finite family $\{\mathbf{X}^a \mid a \in \mathcal{A}\}$ of subspaces of $\mathbf{X}^a \subset \mathbf{X}$ which is stable under addition and which contains $\{0\}$ and **X**. The ordering of subspaces yields an ordering of the abstract set of indices $\mathcal{A}, a \subset b \Leftrightarrow \mathbf{X}^a \subset \mathbf{X}^b$. We denote $\{0\} = \mathbf{X}^{a_{\min}}, \mathbf{X} = \mathbf{X}^{a_{\max}}$ and $\mathbf{X}^a + \mathbf{X}^b = \mathbf{X}^{a \cup b}$. The orthogonal complement of \mathbf{X}^a is denoted by \mathbf{X}_a . We refer to the indices $a \in \mathcal{A}$ as 'cluster decompositions'. The *length* of a chain of cluster decompositions $a_1 \subsetneq \cdots \subsetneq a_k$ is the number k. This chain is said to connect $a = a_1$ and $b = a_k$. The maximal length of all chains connecting a given $a \in \mathcal{A} \setminus \{a_{\max}\}$ and a_{\max} is denoted by #a. We define $\#a_{\max} = 1$ and denoting $\#a_{\min} = N + 1$ we say the family $\{\mathbf{X}^a \mid a \in \mathcal{A}\}$ is of N-body type. Note that for the setup of Sections 1.2 and 1.3 these examples are of (N-1)-body type and of N-body type, respectively. This terminology might appear slightly misleading for Section 1.2. Henceforth we shall treat the generalized N-body framework only. This would consequently apply to the many-body framework of Section 1.2 with N there replaced by N + 1. A cluster decomposition a is said to be k-cluster if #a = k.

Given the above uniform setup of structure of 'internal subspaces' we can introduce corresponding generalized Schrödinger operators. Let $-\Delta^a = (p^a)^2$ and $-\Delta_a = p_a^2$ denote (minus) the Laplacians on $L^2(\mathbf{X}^a)$ and $L^2(\mathbf{X}_a)$, respectively. Here $p^a = \pi^a p$ and $p_a = \pi_a p$ denote the internal (i.e. 'within clusters') and the inter-cluster components of the momentum operator $p = -i\nabla$, respectively. For all $a \in \mathcal{A}' := \mathcal{A} \setminus \{a_{\min}\}$, we introduce

$$H^{a} = -\Delta^{a} + V^{a}(x^{a}), V^{a}(x^{a}) = \sum_{b \subset a} V_{b}(x^{b}), H_{a} = H^{a} - \Delta_{a}, I_{a}(x) = \sum_{b \not \subset a} V_{b}(x^{b}),$$

where the potentials fulfill the condition below. We define $H^{a_{\min}} = 0$ on $L^2(\mathbf{X}^{a_{\min}}) = \mathbb{C}$ and $H = H^{a_{\max}}$ on $L^2(\mathbf{X})$.

Condition 1.5. There exists $\rho > 0$ such that for all $a \in \mathcal{A}'$ there is given a function $V_a : \mathbf{X}^a \to \mathbb{R}$ with a splitting $V_a = V_a^{(1)} + V_a^{(2)}$, where

(1) $V_a^{(1)}$ is smooth and

$$\partial_y^{\alpha} V_a^{(1)}(y) = \mathcal{O}\big(|y|^{-\rho - |\alpha|}\big). \tag{1.22}$$

(2) $V_a^{(2)}$ is compactly supported and

$$V_a^{(2)}(-\Delta_y+1)^{-1} \text{ is compact on } L^2(\mathbb{R}_y^{\dim \mathbf{X}^a}).$$
(1.23)

Condition 1.5 will be imposed throughout this work. To treat local singularities we shall impose an additional condition, depending on an $a \in \mathcal{A}$ from a given context. The condition is fulfilled for the models of Sections 1.2 and 1.3, cf. (1.12a), (1.12b), (1.18a) and (1.18b).

Consider for a given $a \in \mathcal{A}$ with #a = 2 the following properties for $b \not\subset a$:

$$\operatorname{ran}\left(\pi^{b}\pi^{a}\right) = \{0\} \text{ or } \operatorname{ran}\left(\pi^{b}\pi^{a}\right) = \operatorname{ran}\pi^{b}, \qquad (1.24a)$$

$$\pi^b : \mathbf{X}_a \to \mathbf{X}^b \text{ is onto.}$$
 (1.24b)

We note that the map in (1.24b) is necessarily injective, and hence bijective if (1.24b) is fulfilled, see (2.20).

Condition 1.6. For all $b \not\subset a$ for which the conditions (1.24a) and (1.24b) are not fulfilled, the singular part $V_b^{(2)} = 0$ and hence $V_b = V_b^{(1)} \in C^{\infty}(\mathbf{X}^b)$.

If T is a self-adjoint operator on a Hilbert space the subsets of \mathbb{C} ,

$$\sigma(T), \sigma_{\rm d}(T), \sigma_{\rm ess}(T) \text{ and } \sigma_{\rm pp}(T)$$

refer to the spectrum, the discrete spectrum, the essential spectrum and the set of eigenvalues of T, respectively.

The operator H^a is the sub-Hamiltonian associated with the cluster decomposition a and I_a is the sum of all inter-cluster interactions. The detailed expression of H^a depends on the choice of coordinates on \mathbf{X}^a . Let

$$\mathcal{T} = \mathcal{T}(H) = \bigcup_{a \in \mathcal{A}, \#a \ge 2} \sigma_{\mathrm{pp}}(H^a)$$

be the set of thresholds of H. The HVZ theorem [RS, Theorem XIII.17] gives the bottom of the essential spectrum $\Sigma_2 := \inf \sigma_{\text{ess}}(H)$ of H by the formula

$$\Sigma_2 = \min_{a \in \mathcal{A} \setminus \{a_{\max}\}} \inf \sigma(H^a) = \min_{a \in \mathcal{A}, \#a=2} \inf \sigma(H^a).$$
(1.25)

If $N \ge 2$ we also introduce

$$\Sigma_3 := \min_{a \in \mathcal{A}, \#a \ge 3} \inf \sigma(H^a).$$
(1.26)

Under Condition 1.5 it is known that non-threshold bound states decay exponentially [FH]. It is also well known [FH] and [Pe] that under rather general conditions generalized Schrödinger operators do not have positive eigenvalues and that the negative eigenvalues can at most accumulate at the thresholds from below, see also [AIIS].

The goal of the present work is to obtain spectral and scattering properties of H near a general two-cluster threshold $\lambda_0 \leq 0$, i.e.

$$\lambda_0 \in \mathcal{T}_2 := \mathcal{T} \setminus \bigcup_{a \in \mathcal{A}, \#a \ge 3} \sigma_{\mathrm{pp}}(H^a).$$
(1.27)

Note that $\lambda_0 = 0$ for N = 1, while $\lambda_0 < 0$ for $N \ge 2$. Note also that $\lambda_0 \ne \Sigma_3$ for $N \ge 2$ and that $\lambda_0 > \Sigma_3$ could occur for $N \ge 3$. Since the case N = 1 has been treated extensively in the literature we assume throughout our work that $N \ge 2$ and therefore $\lambda_0 < 0$. The special case for $N \ge 2$ when λ_0 is equal to Σ_2 and is given as the eigenvalue of a unique two-cluster sub-Hamiltonian is studied in [Wa2] for fastly decaying potentials, meaning $\rho > 2$ or bigger.

1.4.1 Spaces and notation

For given Banach (or Fréchet) spaces X and Y, the space of linear continuous operators $T: X \to Y$ is denoted by $\mathcal{L}(X, Y)$, and we abbreviate $\mathcal{L}(X) = \mathcal{L}(X, X)$.

Let H_s^k , $k, s \in \mathbb{R}$, be the weighted Sobolev space on **X** (or possibly \mathbf{X}_a for any $a \in \mathcal{A}'$) equipped with the norm

$$||u||_{k,s} := \left(\int |\langle x \rangle^s (1-\Delta)^{k/2} u|^2 \mathrm{d}x\right)^{1/2}, \quad \text{where } \langle x \rangle := (1+|x|^2)^{1/2}.$$

The space H_{-s}^{-k} can be identified as the dual space of H_s^k with the usual L^2 inner product used as 'pairing'. Let $\mathcal{L}(k,s;k',s') = \mathcal{L}(H_s^k,H_{s'}^{k'})$. It will be convenient to regard H as an operator in $\mathcal{L}(2,0;0,0)$ or as an operator in $\mathcal{L}(1,0;-1,0)$. We abbreviate $H^k = H_0^k$, $L_s^2 = H_s^0$, and $H_\infty^k = \bigcap_{s \in \mathbb{R}} H_s^k$, $H_{-\infty}^k = \bigcup_{s \in \mathbb{R}} H_s^k$, $L_\infty^2 = \bigcap_{s \in \mathbb{R}} L_s^2$ and $L_{-\infty}^2 = \bigcup_{s \in \mathbb{R}} L_s^2$. Introduce also $H_{t^+}^k = \bigcup_{s > t} H_s^k$, $H_{t^-}^k = \bigcap_{s < t} H_s^k$, $L_{t^+}^2 = \bigcup_{s > t} L_s^2$ and $L_{t^-}^2 = \bigcap_{s < t} L_s^2$ for any $k, t \in \mathbb{R}$.

We shall also use weighted Sobolev spaces of \mathbb{C}^m -valued functions/distributions on **X** indicated by similar notation, for example $H^k = H^k(\mathbf{X}; \mathbb{C}^m)$. For any complex Hilbert spaces \mathcal{H}_1 and \mathcal{H}_1 we use the notation $\mathcal{C}(\mathcal{H}_1, \mathcal{H}_2) \subset \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2)$ for the space of compact operators $T : \mathcal{H}_1 \to \mathcal{H}_2$. By standard Sobolev embedding theory, for all $k_1 > k_2$ and $s_1 > s_2$ the operator $H^{k_1}_{s_1} \ni f \to f \in H^{k_2}_{s_2}$ is compact. Let H^1_{loc} denote the set of locally H^1 functions, more precisely the set of $v \in L^2_{-\infty}$ for which $\varphi v \in H^1$ for all $\varphi \in C^{\infty}_c$.

Consider balls $B(R) = \{x \in \mathbf{X} | |x| < R\}, R \ge 1$, and the characteristic functions

$$F_0 = F(B(R_0))$$
 and $F_{\nu+1} = F(B(R_{\nu+1}) \setminus B(R_{\nu})), R_{\nu} = 2^{\nu}, \nu \in \mathbb{N}_0 := \mathbb{N} \cup \{0\},\$

where F(M) denotes the sharp characteristic function of a subset $M \subset \mathbf{X}$. We introduce the Besov spaces $\mathcal{B}_s = \mathcal{B}_s(\mathbf{X})$ and $\mathcal{B}_s^* = \mathcal{B}_s^*(\mathbf{X}) = (\mathcal{B}_s(\mathbf{X}))^*$, s > 0, as follows.

$$\mathcal{B}_{s} = \{ \psi \in L^{2}_{\text{loc}} \mid \|\psi\|_{\mathcal{B}_{s}} < \infty \}, \quad \|\psi\|_{\mathcal{B}_{s}} = \sum_{\nu=0}^{\infty} R^{s}_{\nu} \|F_{\nu}\psi\|_{L^{2}}, \\ \mathcal{B}^{*}_{s} = \{ \psi \in L^{2}_{\text{loc}} \mid \|\psi\|_{\mathcal{B}^{*}_{s}} < \infty \}, \quad \|\psi\|_{\mathcal{B}^{*}_{s}} = \sup_{\nu \ge 0} R^{-s}_{\nu} \|F_{\nu}\psi\|_{L^{2}},$$

respectively. (Note that indeed \mathcal{B}_s^* is the dual space of \mathcal{B}_s .) We define $\mathcal{B}_{s,0}^*$ to be the closure of L^2 in \mathcal{B}_s^* . Note that

$$u \in \mathcal{B}_s^* \Leftrightarrow u \in L^2_{\text{loc}} \text{ and } \sup_{R \ge 1} R^{-s} \|F(B(R))u\|_{L^2} < \infty,$$

and that

$$u \in \mathcal{B}_{s,0}^* \Leftrightarrow u \in L^2_{\text{loc}} \text{ and } \lim_{R \to \infty} R^{-s} \|F(B(R))u\|_{L^2} = 0.$$

Note the following relations between the standard weighted L^2 spaces and the Besov spaces:

$$\forall s_1 > s > 0: \quad L^2_{s_1} \subsetneq \mathcal{B}_s \subsetneq L^2_s \subsetneq L^2 \subsetneq L^2_{-s} \subsetneq \mathcal{B}^*_{s,0} \subsetneq \mathcal{B}^*_s \subsetneq L^2_{-s_1}. \tag{1.28}$$

We introduce for any $R\geq 1$

$$\chi_R(t) = \chi(t/R) \text{ for a real-valued } \chi \in C^{\infty}(\mathbb{R}) : \quad \chi(t) = \begin{cases} 1 & \text{for } t \le 4/3 \\ 0 & \text{for } t \ge 5/3 \end{cases},$$
(1.29)

and $\bar{\chi}_R = 1 - \chi_R$. We assume $-\chi' \ge 0$ and that $\sqrt{-\chi'} \in C^{\infty}$.

Chapter 2

Reduction to a one-body problem

In this chapter we show that the spectral analysis of a generalized N-body operator near a two-cluster threshold can be reduced to the analysis of a one-body operator with a non-linear spectral parameter. This is under Condition 1.5. Of course such reduction would not be needed for the one-body problem, however we recall that throughout our work we impose the condition $N \ge 2$.

The idea of the reduction goes as follows. For a given two-cluster threshold $\lambda_0 < 0$, denote by \mathcal{F} the closed subspace in $L^2(\mathbf{X})$ 'spanned by' bound states of all possible two-cluster sub-Hamiltonians with eigenvalue λ_0 . (For a precise definition/representation in a special case see (2.15) and the discussion there.) Let Π be the orthogonal projection from $L^2(\mathbf{X})$ onto \mathcal{F} , $\Pi' = 1 - \Pi$ and $H' = \Pi' H \Pi'$. We study a Grushin problem for an operator of the form

$$H(z) = \begin{pmatrix} H-z & S \\ S^* & 0 \end{pmatrix} \text{ on } L^2(\mathbf{X}) \oplus \mathcal{H}, \qquad (2.1)$$

where the space \mathcal{H} is some auxiliary one-body type space, $S : \mathcal{H} \to L^2(\mathbf{X})$ is an appropriately defined operator whose range coincides with \mathcal{F} , i.e. the range of the projection Π , and S^* is the adjoint of S. The choice of \mathcal{H} and S may vary according to various situations and is to make H(z) invertible for z near λ_0 and $\operatorname{Im} z \neq 0$. If this is realized, we set

$$H(z)^{-1} = \begin{pmatrix} E(z) & E_{+}(z) \\ E_{-}(z) & E_{\mathcal{H}}(z) \end{pmatrix}.$$
 (2.2)

Then the resolvent $R(z) = (H - z)^{-1}$ is represented as

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z).$$
(2.3)

The operator $H(z)^{-1}$ can be computed in terms of the reduced resolvent $R'(z) = (H'-z)^{-1}\Pi'$. If R'(z) has some good properties near λ_0 , the spectral analysis of H near λ_0 is then reduced to the analysis of $E_{\mathcal{H}}(z)^{-1}$ on \mathcal{H} . $E_{\mathcal{H}}(z)$ is a one-body operator with a non-linear dependence on the spectral parameter z. For example to obtain a resolvent expansion, one can then try to use known methods for one-body operators to study the asymptotics of $E_{\mathcal{H}}(z)^{-1}$ as $z \to \lambda_0$, Im z > 0.

Below we shall only give detailed analysis in some situations. In fact, the reductions given below are only useful in the case $\lambda_0 \notin \sigma_{\rm pp}(H')$. The case $\lambda_0 \in \sigma_{\rm pp}(H')$ can be treated by adding an additional finite dimensional space, as done in [Wa2, Section 4] for the lowest threshold. Before we go into details on various concrete cases we study the reduction scheme from a more general point of view. As we will see afterwards, the abstract scheme can be applied to these cases.

2.1 An abstract reduction scheme

In this section we shall describe an abstract reduction scheme that will be used many times.

Abstract framework Suppose \mathcal{H} and \mathcal{G} are given Hilbert spaces and that \mathcal{F} is a closed subspace of \mathcal{G} . Suppose $S : \mathcal{H} \to \mathcal{F} \subset \mathcal{G}$ is a bi-continuous isomorphism (i.e. S is linear, one-to-one, onto \mathcal{F} and as a map from \mathcal{H} to \mathcal{F} bi-continuous). Let Π denote the orthogonal projection in \mathcal{G} onto \mathcal{F} , and let $\Pi' = 1 - \Pi$. Suppose H is a self-adjoint operator on \mathcal{G} with $\Pi : \mathcal{D}(H) \to \mathcal{D}(H)$, and that $\Pi'H\Pi$ and $\Pi H\Pi'$ (initially defined on the domain $\mathcal{D}(H)$ of H) extend to bounded operators on \mathcal{G} . Define $H' = \Pi'H\Pi'$ with domain $\mathcal{D}(H') = \mathcal{D}(H) \cap \operatorname{ran}(\Pi') = \Pi' \mathcal{D}(H)$ and $H_{\Pi} = \Pi H\Pi$ with domain $\mathcal{D}(H_{\Pi}) = \mathcal{D}(H) \cap \operatorname{ran}(\Pi) = \Pi \mathcal{D}(H)$. Then H' and H_{Π} are self-adjoint on $\Pi'\mathcal{G}$ and \mathcal{F} , respectively (see the proof of Lemma 2.8 3)). Let $R'(z) = (H' - z)^{-1}\Pi'$ for $\operatorname{Im} z \neq 0$.

Introduce for all $z \in \mathbb{C}$ with $\operatorname{Im} z \neq 0$

$$E(z) = R'(z), \tag{2.4a}$$

$$E_{+}(z) = S - R'(z)HS,$$
 (2.4b)

$$E_{-}(z) = S^* - S^* H R'(z), \qquad (2.4c)$$

$$E_{\mathcal{H}}(z) = S^* (z - H + HR'(z)H)S.$$
(2.4d)

Obviously $E(z) \in \mathcal{L}(\mathcal{G}), E_+(z) \in \mathcal{L}(\mathcal{H}, \mathcal{G})$ and $E_-(z) \in \mathcal{L}(\mathcal{G}, \mathcal{H})$. Also note that $S^*HS = S^*H_{\Pi}S$ is self-adjoint and consequently that the operator $E_{\mathcal{H}}(z)$ of (2.4d) is a closed operator on \mathcal{H} with domain given by $S^{-1}\mathcal{D}(H_{\Pi})$.

Proposition 2.1. Under the above conditions, for $\text{Im } z \neq 0$:

- 1) $E_{\mathcal{H}}(z)^* = E_{\mathcal{H}}(\bar{z}).$
- 2) $E_{\mathcal{H}}(z)$ is an invertible operator on \mathcal{H} obeying

$$\frac{\operatorname{Im} E_{\mathcal{H}}(z)}{\operatorname{Im} z} \ge S^* S. \tag{2.5}$$

3)

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z).$$
(2.6)

Proof. The identity 1) is trivial due to the self-adjointness property stated before the proposition.

The inequality (2.5) follows from the identity

$$\operatorname{Im} E_{\mathcal{H}}(z) = \operatorname{Im} z \, S^* \big(1 + H R'(z)^* R'(z) H \big) S. \tag{2.7}$$

The invertibility property of 2) follows from a standard numerical range argument combining 1) and (2.5).

Finally the identity (2.6) follows by an elementary calculation outlined here: Apply H - z from the left on the right-hand side of the identity and write for the first term of (2.6)

$$H\Pi' = \Pi' H'\Pi' + \Pi H\Pi'. \tag{2.8}$$

The result is the expression $\Pi' + \Pi HR'(z)$. For the second term of (2.6) we substitute the expression (2.4b) and use again (2.8) (to deal with the second term of (2.4b)). We see that by applying H - z to the second term of (2.6) we obtain an operator taking values in \mathcal{F} . Consequently we may insert $\Pi = (S^*)^{-1}S^*$ to the left and then combine the factor S^* with the given calculated expression. This gives a factor $E_{\mathcal{H}}(z)E_{\mathcal{H}}(z)^{-1}$ which of course can be omitted on $\mathcal{H} = \operatorname{ran} E_{-}(z)$. Then by substituting the expression (2.4c) we conclude that the contribution from the second term of (2.6) is $\Pi - \Pi HR'(z)$. Hence the result of applying H - z from the left on the right-hand side of (2.6) is

$$\left(\Pi' + \Pi H R'(z)\right) + \left(\Pi - \Pi H R'(z)\right) = 1,$$

which coincides with the result of applying H-z from the left on the left-hand side of the identity.

Remark 2.2. There is an alternative approach for deriving the formula (2.6) based on a certain abstract Grushin problem. For a review of this method we may refer to [SZ], but for sake of completeness of presentation we outline this alternative approach here, cf. the beginning of this chapter:

We consider a Grushin problem for an operator of the form

$$H(z) = \begin{pmatrix} H-z & S \\ S^* & 0 \end{pmatrix} \text{ on } \mathcal{G} \oplus \mathcal{H}.$$
 (2.9)

To simplify the notation, we denote still by $(SS^*)^{-1} = (SS^*)^{-1}\Pi$ the extension of $(SS^*)^{-1}$ to \mathcal{G} by setting $(SS^*)^{-1} = 0$ on \mathcal{F}^{\perp} .

The invertibility of SS^* on \mathcal{F} allows us to show that H(z) is invertible on $\mathcal{G} \oplus \mathcal{H}$. In fact, one can explicitly compute its inverse. Let $T := (SS^*)^{-1}S$, and let Q(z) and B(z) be defined by

$$Q(z) = \begin{pmatrix} R'(z) & T \\ T^* & T^*(z-H)T \end{pmatrix} \text{ and } B(z) = \begin{pmatrix} \Pi H R'(z) & \Pi' H T \\ 0 & 0 \end{pmatrix}$$

Then one has, at least formally,

$$H(z)Q(z) = 1 + B(z).$$

Since $B(z)^3 = 0$ the operator 1 + B(z) is invertible and therefore H(z) has a right inverse. Similarly, one can show that H(z) has a left inverse. Consequently H(z) should be invertible with inverse

$$H(z)^{-1} = Q(z)(1 - B(z) + B(z)^2).$$
(2.10)

Write $H(z)^{-1}$ in the form

$$H(z)^{-1} = \begin{pmatrix} \widetilde{E}(z) & \widetilde{E}_{+}(z) \\ \widetilde{E}_{-}(z) & \widetilde{E}_{\mathcal{H}}(z) \end{pmatrix}.$$
 (2.11)

We have the formulas

$$\tilde{E}(z) = R'(z), \qquad (2.12a)$$

$$\widetilde{E}_{+}(z) = T - R'(z)HT, \qquad (2.12b)$$

$$\widetilde{E}_{-}(z) = T^* - T^* H R'(z),$$
(2.12c)

$$\widetilde{E}_{\mathcal{H}}(z) = T^* \big(z - H + HR'(z)H \big) T, \qquad (2.12d)$$

and

$$R(z) = \widetilde{E}(z) - \widetilde{E}_{+}(z)\widetilde{E}_{\mathcal{H}}(z)^{-1}\widetilde{E}_{-}(z).$$
(2.13)

Now (2.6) follows from (2.13) and the identity

$$T = (SS^*)^{-1}S = S(S^*S)^{-1}.$$
(2.14)

Remark 2.3. Suppose $\lambda_0 \notin \sigma(H')$ so that the operators in (2.12a)–(2.12d) have limits when taking $z \to \lambda = \lambda_0 \in \mathbb{R}$. Then we learn from the identities

$$\widetilde{E}_{-}(\lambda)(H-\lambda) \subset -\widetilde{E}_{\mathcal{H}}(\lambda)S^{*},$$
$$S\widetilde{E}_{\mathcal{H}}(\lambda) \subset -(H-\lambda)\widetilde{E}_{+}(\lambda),$$

that

$$S^* : \ker(H - \lambda) \to \ker \tilde{E}_{\mathcal{H}}(\lambda),$$
$$\tilde{E}_+(\lambda) : \ker \tilde{E}_{\mathcal{H}}(\lambda) \to \ker(H - \lambda),$$

respectively.

Combined with the identities

$$\widetilde{E}_{+}(\lambda)S^{*} \supset 1 - \widetilde{E}(\lambda)(H - \lambda),$$

$$S^{*}\widetilde{E}_{+}(\lambda) = 1,$$

we then conclude that in fact

$$\widetilde{E}_{+}(\lambda)S^{*} = 1 \text{ on } \ker(H - \lambda),$$

 $S^{*}\widetilde{E}_{+}(\lambda) = 1 \text{ on } \ker\widetilde{E}_{\mathcal{H}}(\lambda),$

i.e. $S^* : \ker(H - \lambda) \to \ker \widetilde{E}_{\mathcal{H}}(\lambda)$ is a linear isomorphism.

Whence, cf. (2.14), T^* : ker $(H - \lambda) \rightarrow$ ker $E_{\mathcal{H}}(\lambda)$ is a linear isomorphism with inverse $E_+(\lambda)$: ker $E_{\mathcal{H}}(\lambda) \rightarrow$ ker $(H - \lambda)$. We shall refer to the vector $f = T^*\phi \in$ ker $E_{\mathcal{H}}(\lambda)$ as the *eigentransform* of a given $\phi \in$ ker $(H - \lambda)$, and the equation $\phi = E_+(\lambda)f$ as the *inversion formula* for the eigentransform of ϕ . The Hilbert space setting discussed here is in our application an L^2 setting which will have extensions to Besov space settings and settings where $\lambda_0 \in \sigma(H')$, that are incompatible with the framework discussed here. These concrete extensions will be studied in Chapter 4.

2.2 Non-multiple two-cluster threshold case

Assume in this section that $\lambda_0 < 0$ is a *non-multiple two-cluster threshold* in the following sense:

Condition 2.4. There exists a unique $a_0 \in \mathcal{A} \setminus \{a_{\max}\}$ such that $\lambda_0 \in \sigma_{pp}(H^{a_0})$. This cluster decomposition is a two-cluster decomposition, i.e. $\#a_0 = 2$. Condition 1.6 is fulfilled for $a = a_0$.

Note that we do not here impose that $\lambda_0 \in \sigma_d(H^{a_0})$. Nevertheless λ_0 is not a threshold for the Hamiltonian H^{a_0} , and consequently the corresponding bound states have exponential decay, cf. [FH, AIIS]. Note that the multiplicity of λ_0 as an eigenvalue of H^{a_0} , say m, can be arbitrary. The simplest case is $\lambda_0 = \Sigma_2$ (under Condition 2.4 necessarily m = 1 in this case), and it is studied in [Wa2] for fastly decaying potentials. For $\lambda_0 > \Sigma_2$, we can use the same idea to reduce H to a one-body type operator with a non-linear spectral parameter, although additional complications arise, in particular for $\lambda_0 > \Sigma_3$.

Let $\{\varphi_1, \ldots, \varphi_m\}$ be an orthonormal basis of the eigenspace of H^{a_0} associated with λ_0 . Let Π be the projection in $\mathcal{G} := L^2(\mathbf{X})$ defined by

$$\Pi g = \sum_{j=1}^{m} \varphi_j \otimes \langle \varphi_j, g \rangle_{L^2(\mathbf{X}^{a_0})}, \ g \in \mathcal{G}.$$
(2.15)

Let $\Pi' = 1 - \Pi$ and $H' = \Pi' H \Pi'$. Note that H' is self-adjoint with domain $\mathcal{D}(H') = \mathcal{D}(H) \cap \operatorname{ran} \Pi' = \Pi' \mathcal{D}(H)$. In fact H_{a_0} is reduced by Π' ('reduced' in the sense of [Ka, Subsection V.3.9]) implying in particular that $H'_{a_0} = \Pi' H_{a_0} \Pi'$ is self-adjoint. Since $I'_{a_0} = \Pi' I_{a_0} \Pi'$ is infinitesimally small relatively to H'_{a_0} it follows from [RS, Theorem X.12] that indeed H' is self-adjoint. For Im $z \neq 0$ we set

$$R'(z) = (H' - z)^{-1} \Pi'.$$

Let $I_0 = I_{a_0}$ and abbreviate similarly $p_0 = p_{a_0}$. Since we have imposed Condition 1.6 with $a = a_0$ we can use a 'free factor' $(|p^{a_0}|^2 + 1)^{-1}$ (from Π) to conclude that

$$\Pi' I_0 \Pi \in \mathcal{L}(\mathcal{G}). \tag{2.16}$$

Let $\mathcal{H} = \mathcal{H}_{a_0} := L^2(\mathbf{X}_{a_0}; \mathbb{C}^m), \ H^2_{a_0} := H^2(\mathbf{X}_{a_0}; \mathbb{C}^m) \subset \mathcal{H}_{a_0} \text{ and the operator } S : \mathcal{H} \to \mathcal{G}$ be defined by

$$S: f = (f_1, \dots, f_m) \to Sf = \sum_{j=1}^m S_j f_j = \sum_{j=1}^m \varphi_j(x^{a_0}) f_j(x_{a_0}).$$
(2.17)

Obviously $S: H^2_{a_0} \to H^2(\mathbf{X})$ and the (L^2) adjoint $S^*: H^2 \to H^2_{a_0}$. Since S in this case is isometric the formulas (2.4a)–(2.4d) and (2.12a)–(2.12d) coincide. They read (in terms of the identity matrix $\mathbf{1}_m$ of size m)

$$E(z) = R'(z), \tag{2.18a}$$

$$E_{+}(z) = (1 - R'(z)I_0)S, \qquad (2.18b)$$

$$E_{-}(z) = S^{*}(1 - I_{0}R'(z)), \qquad (2.18c)$$

$$E_{\mathcal{H}}(z) = (z - \lambda_0) - (|p_0|^2 \mathbf{1}_m + S^* I_0 S - S^* I_0 R'(z) I_0 S), \qquad (2.18d)$$

which can be used in combination with (2.6), i.e.

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z).$$
(2.19)

We examine in the following some basic properties of $E_{\mathcal{H}}(z)$. Note that $S^*I_{a_0}S$ is a matrix-valued potential. Let us first split $I_0 = I_0^{(1)} + I_0^{(2)} = I_{a_0}^{(1)} + I_{a_0}^{(2)}$ in agreement with the splitting of Condition 1.5 and look at the contribution from $I_0^{(1)}$. Recalling the elementary geometric property

$$X_a \cap X_b = \{0\} \text{ if } \#a = 2 \text{ and } b \not\subset a,$$
 (2.20)

guaranteeing that $\pi^b : \mathbf{X}_{a_0} \to \mathbf{X}^b$ is injective for all $b \not\subset a_0$, we obtain by a Taylor expansion that the leading term has a scalar leading form. More precisely we obtain by a zero'th order Taylor expansion, by using (2.50a) (stated below) and by using the polynomial decay of the bound states φ_j that

$$S_i^* I_0^{(1)} S_j = I_0^{(1)} (0+y) \delta_{ij} + \mathcal{O}(|y|^{-\rho-1}) \text{ for } |y| \to \infty; \quad i, j \le m.$$
(2.21a)

Here the variable $y = x_{a_0}$ can be thought of as a vector in \mathbb{R}^n , abbreviating here and henceforth dim $\mathbf{X}_{a_0} = n$. We claim the following analogue of (2.21a) for $I_0^{(2)}$:

$$S_i^* I_0^{(2)} S_j - I_0^{(2)} (0+y) \delta_{ij} \in \mathcal{C} \big(H_s^2(\mathbf{X}_{a_0}), L_t^2(\mathbf{X}_{a_0}) \big) \text{ for all } s, t \in \mathbb{R}.$$
(2.21b)

To prove (2.21b) it suffices to consider the contribution from $V_b^{(2)}$ for any $b \not\subset a_0$ under the conditions (1.24a) and (1.24b), in fact with the second condition of (1.24a) fulfilled. Due to (1.24b) the restriction $V_b^{(2)}(=V_b^{(2)}(y)) \in \mathcal{C}(H_s^2(\mathbf{X}_{a_0}), L_t^2(\mathbf{X}_{a_0}))$. It is compactly supported, possibly with singularities. On the other hand $S_i^* V_b^{(2)} S_j$ is a bounded potential due to the second condition of (1.24a), and it remains to show that this potential decays faster than any negative power $\langle y \rangle^{-s}$. By the compact support property we can pick R > 1 such that

$$F(|\pi^{b} \cdot | > R)S_{i}^{*}V_{b}^{(2)}S_{j} = S_{i}^{*}F(|\pi^{b}y| > R)F(|\pi^{b}x^{a_{0}}| \ge \frac{1}{2}|\pi^{b}y|)V_{b}^{(2)}S_{j}.$$

By the polynomial decay of the cluster bound states the right-hand side has arbitrary power decay too, showing the desired decay and therefore (2.21b).

We conclude from (2.21a) and (2.21b) that

$$S_{i}^{*}I_{0}S_{j} - I_{0}^{(1)}(0+y)\delta_{ij} \in \mathcal{C}(H_{s}^{2}(\mathbf{X}_{a_{0}}), L_{t}^{2}(\mathbf{X}_{a_{0}})) \text{ for } s, t \in \mathbb{R}, t < \rho + 1 + s,$$

$$S_{i}^{*}I_{0}S_{j} - I_{0}^{(1)}(0+y)\delta_{ij} \in \mathcal{L}(H_{s}^{2}(\mathbf{X}_{a_{0}}), L_{\rho+1+s}^{2}(\mathbf{X}_{a_{0}})) \text{ for } s \in \mathbb{R}.$$
(2.22)

It remains to examine the term $S^*I_0R'(z)I_0S$ in (2.18d). Note that the two factors of I_0 freely can be changed to $\Pi I_0\Pi'$ and $\Pi'I_0\Pi$, respectively. We can then conveniently implement the following improvements of (2.16):

$$\Pi I_0 \Pi', \, \Pi' I_0 \Pi \in \mathcal{L} \left(L_s^2(\mathbf{X}), L_{\rho+1+s}^2(\mathbf{X}) \right) \text{ for all } s \in \mathbb{R}.$$

$$(2.23)$$

For these bounds it suffices to consider $\Pi' I_0 \Pi$. The contribution from $I_0^{(1)}$ is treated by a Taylor expansion as in (2.21a) (and by (2.50a)). The contribution from $I_0^{(2)}$ is treated by using a 'free factor' $(|p^{a_0}|^2 + 1)^{-1}$ (as for (2.16)) as well as using the above proof of (2.21b).

We conclude the following representation of the operator $E_{\mathcal{H}}(z)$:

$$E_{\mathcal{H}}(z) + \left(p_0^2 + I_0^{(1)}(\pi_{a_0} \cdot) + \lambda_0 - z\right) \mathbf{1}_m = \widetilde{V} - \langle y \rangle^{-\rho - 1} \widetilde{K}(z) \langle y \rangle^{-\rho - 1};$$

$$\widetilde{V} = \mathcal{O}(\langle y \rangle^{-\rho - 1}), \quad \widetilde{K}(z) = \mathcal{O}(\langle y \rangle^0);$$

here the meaning of the $\mathcal{O}(\cdot)$ notation is given more precisely by (2.22) and (2.23), respectively. Note that \widetilde{V} is multiplicative matrix-valued, while $\widetilde{K} \in \mathcal{L}(\mathcal{H})$ is nonmultiplicative with a z-dependence through the appearance of the factor R'(z). Note also that $\langle y \rangle^{\rho} I_0^{(1)}(y)$ is bounded. Although the above discussion is based on a natural operator interpretation of $E_{\mathcal{H}}(z)$ we shall prefer to use the corresponding form interpretation (see Remark 2.13 for details in a different but similar context).

The representation formula (2.19) is valid for any z with $\text{Im } z \neq 0$, but for our purposes it is only useful if one has good properties of R'(z) for z near λ_0 . When λ_0 is the lowest threshold, the essential spectrum of H' is shifted to the right and the representation (2.19) can be used in the case λ_0 is not an eigenvalue of H', cf. [Wa2, Lemma 2.1] (see Lemma 2.10 for an extension). With this assumption, the reduced resolvent R'(z) is holomorphic in a small neighbourhood of λ_0 , and so are $E_+(z), E_H(z)$ and $E_-(z)$. Moreover in this case R'(z) has uniform bounds in weighted L^2 spaces (see (2.56) for a similar assertion for the multiple case). This is helpful in the study of asymptotic expansions of $E_H(z)^{-1}$ at λ_0 and therefore in turn (by (2.19)) for expansions of R(z) near λ_0 (see Section 5.1.2 for the multiple case). In the case λ_0 happens to be an eigenvalue of H' another reduction is needed.

If $\lambda_0 > \Sigma_2$, λ_0 is in the essential spectrum of H'. Under Conditions 1.5 and 2.4 we shall show in Chapter 3 that a limiting absorption principle and microlocal resolvent estimates hold for R'(z) with z near λ_0 provided λ_0 is not an eigenvalue of H'. This leads in this case to an extension of the 'eigentransform' discussed in Remark 2.3 (see Chapter 4), and then in turn (2.19) can again be used to analyse the resolvent of H near the threshold λ_0 (see Section 5.1.3). The analysis for $\lambda_0 > \Sigma_2$ is more complicated than for $\lambda_0 = \Sigma_2$ in that, some more refined mapping properties of R'(z) near λ_0 are needed, causing in particular some 'loss of weight'.

2.3 Multiple two-cluster threshold, $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$

Consider now the case where there exist at least two two-cluster decompositions such that the two-cluster threshold $\lambda_0(< 0)$ is an eigenvalue of the corresponding sub-Hamiltonians. To simplify the presentation, let us only consider in detail the case where λ_0 is double occurring without eigenvalue multiplicity:

Condition 2.5. There exist unique $a_1, a_2 \in \mathcal{A} \setminus \{a_{\max}\}, a_1 \neq a_2$, such that $\lambda_0 \in \sigma_{pp}(H^{a_j}), j = 1, 2$. The cluster decompositions a_1 and a_2 are two-cluster decompositions, i.e. $\#a_1 = \#a_2 = 2$, and Condition 1.6 is fulfilled for both of them. The number λ_0 is a simple eigenvalue for both of the operators H^{a_1} and H^{a_2} .

Denote

$$\mathbf{X}^{j} = \mathbf{X}^{a_{j}}, \quad \mathbf{X}_{j} = \mathbf{X}_{a_{j}} \quad \text{and} \quad n_{j} = \dim \mathbf{X}_{j} \quad \text{for} \quad j = 1, 2,$$

and similarly for elements x of \mathbf{X}^{a_j} and \mathbf{X}_{a_j} , viz. $x = x^j \oplus x_j$. Let φ_j denote the corresponding (real) normalized eigenvector of H^{a_j} , j = 1, 2. Let

$$\mathcal{F}_j = \{ g = \varphi_j(x^j) f_j(x_j) \mid f_j \in L^2(\mathbb{R}^{n_j}_{x_j}) \}, \quad j = 1, 2.$$

Note that \mathcal{F}_j is a closed subspace of $\mathcal{G} := L^2(\mathbf{X})$. Let Π be the orthogonal projection in \mathcal{G} onto \mathcal{F} , which by definition is the closure of $\mathcal{F}_1 + \mathcal{F}_2$ in \mathcal{G} . Let $\mathcal{H} = L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2)$.

To construct an associated Grushin problem we need to distinguish between two cases: a) $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$ and b) $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$. Here we shall impose the condition a), i.e.

$$\mathcal{F}_1 + \mathcal{F}_2 = \mathcal{F}_1 \oplus \mathcal{F}_2. \tag{2.24}$$

We believe that this condition is always fulfilled for the finite mass many-body operators of Section 1.2, however we do not have a proof. A case where it fails for the infinite mass many-body operators of Section 1.3 will be discussed in Section 2.5. Moreover the general case of b) will be studied there.

Let $S = (S_1, S_2) : \mathcal{H} \to \mathcal{G}$ be defined by $Sf = S_1f_1 + S_2f_2$ for $f = (f_1, f_2)$ and with

$$S_j : L^2(\mathbf{X}_j) \to \mathcal{G}, \quad f_j \to S_j f_j = \varphi_j(x^j) \otimes f_j(x_j), S_j^* : \mathcal{G} \to L^2(\mathbf{X}_j), \quad f \to S_j^* f = \langle \varphi_j, f \rangle_j; \quad j = 1, 2.$$

$$(2.25)$$

Here $\langle ., . \rangle_j$ denotes the scalar product in $L^2(\mathbf{X}^j)$; the notation $\langle \cdot, \cdot \rangle$ will be used to denote the scalar product in \mathcal{G} . One has

$$S^*S = 1 + \begin{pmatrix} 0 & s_{12} \\ s_{21} & 0 \end{pmatrix} \quad \text{on } \mathcal{H} = L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2),$$

where $s_{ij} \in \mathcal{L}(L^2(\mathbf{X}_j), L^2(\mathbf{X}_i)), i \neq j$, are given by

$$s_{ij}f_j = \langle \varphi_i, \varphi_j \otimes f_j \rangle_i.$$

In the proof of Proposition 2.7 stated below we shall use that $s_{ij} \in \mathcal{C}(L^2(\mathbf{X}_j), L^2(\mathbf{X}_i))$, $i \neq j$. This property is a consequence of the following lemma.

Lemma 2.6. For all $r, t \in \mathbb{R}$

$$s_{ij} \in \mathcal{L}(L^2_r(\mathbf{X}_j), H^2_t(\mathbf{X}_i)) \cap \mathcal{L}(H^{-2}_r(\mathbf{X}_j), L^2_t(\mathbf{X}_i)).$$

Proof. Note that (2.20) implies that

$$\operatorname{ran}(\pi_b \pi^a) = \operatorname{ran}(\pi_b) \text{ if } \#b = 2 \text{ and } a \not\subset b.$$
(2.26)

We apply (2.26) to $a = a_i$ and $b = a_j$ and do integration by parts (or alternatively change variables) obtaining that

$$s_{ij} \in \mathcal{L}(L^2(\mathbf{X}_j), H^2(\mathbf{X}_i)).$$

Whence for $|\alpha| \leq 2$ we have a formula for $\partial_{x_i}^{\alpha} s_{ij}$ involving partial derivatives $\partial^{\beta} \varphi_k$, k = 1, 2 and $|\beta| \leq 2$. By using this formula, polynomial decay of $\partial^{\beta} \varphi_k$ and the bound

$$|x^i| + |x^j| \ge c|x|$$

(the latter is a consequence of (2.20)) we obtain that indeed

$$s_{ij} \in \mathcal{L}(L^2_r(\mathbf{X}_j), H^2_t(\mathbf{X}_i)).$$

Since $s_{ij} = s_{ji}^*$ also

$$s_{ij} \in \mathcal{L}(H_r^{-2}(\mathbf{X}_j), L_t^2(\mathbf{X}_i)).$$

Proposition 2.7. Suppose (2.24). Then

- 1) $\mathcal{F} = \mathcal{F}_1 \oplus \mathcal{F}_2$ (i.e. $\mathcal{F}_1 + \mathcal{F}_2$ is closed).
- 2) $S^* = 0$ on \mathcal{F}^{\perp} and $S^* : \mathcal{F} \to \mathcal{H}$ is a bi-continuous isomorphism. Similarly $S : \mathcal{H} \to \mathcal{F}$ is a bi-continuous isomorphism, and therefore $SS^* : \mathcal{F} \to \mathcal{F}$ is invertible on \mathcal{F} . One has

$$S^*(SS^*)^{-1}S = 1 \text{ on } \mathcal{H}.$$
 (2.27)

3) S^*S is invertible on \mathcal{H} and

$$S(S^*S)^{-1}S^* = \Pi \ on \ \mathcal{G}.$$
 (2.28)

Proof. 1). For $f \in \mathcal{F}_1 + \mathcal{F}_2$, one has for some $f_j \in L^2(\mathbf{X}_j)$,

$$f = \varphi_1 \otimes f_1 + \varphi_2 \otimes f_2.$$

Since $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$, this decomposition is unique. We claim that $N(f) = ||f_1|| + ||f_2||$ defines a norm on $\mathcal{F}_1 + \mathcal{F}_2$ which is equivalent with the norm of \mathcal{G} . Clearly, one has

$$\|f\| \le N(f).$$

Conversely, we want to show the existence of a constant C > 0 such that $N(f) \leq C ||f||$ on $\mathcal{F}_1 + \mathcal{F}_2$. If this is not true, there would be a sequence $g_n \in \mathcal{F}_1 + \mathcal{F}_2$ such that $||g_n|| < 1/n$ and $N(g_n) = 1$. Write g_n as

$$g_n = \varphi_1 \otimes f_{1,n} + \varphi_2 \otimes f_{2,n}$$

with $||f_{1,n}|| + ||f_{2,n}|| = 1$. Without loss of generality, we can assume that $\{f_{k,n}\}$ converges weakly to f_k in $L^2(\mathbf{X}_k)$ as $n \to \infty$. Then, one has

$$f_{1,n} = \langle \varphi_1, g_n \rangle_1 - s_{12} f_{2,n} = \mathcal{O}(n^{-1}) - s_{12} f_{2,n},$$

$$f_{2,n} = \langle \varphi_2, g_n \rangle_2 - s_{21} f_{1,n} = \mathcal{O}(n^{-1}) - s_{21} f_{1,n}.$$

Substituting the second equation into the first one, we obtain that

$$f_{1,n} = \mathcal{O}(n^{-1}) + K_1 f_{1,n}, \ K_1 = s_{12} s_{21}.$$

Since $f_{1,n}$ is weakly convergent and K_1 is compact, $K_1 f_{1,n}$ is strongly convergent. This implies that $f_{1,n}$ converges to some f_1 in $L^2(\mathbf{X}_1)$. Similarly one shows that $f_{2,n}$ converges to some f_2 in $L^2(\mathbf{X}_2)$. Since $g_n \to 0$, this implies

$$\varphi_1 \otimes f_1 + \varphi_2 \otimes f_2 = 0, \quad ||f_1|| + ||f_2|| = 1,$$

which is impossible since $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$. This proves the equivalence of the norms. It follows that $\mathcal{F}_1 + \mathcal{F}_2$ is closed.

<u>2</u>). Let $\tilde{S}: \mathcal{H} \to \mathcal{F}$ act as S, i.e. $\tilde{S}f = Sf$ for $f \in \mathcal{H}$. By the equivalence of norms shown above we see that \tilde{S} is a bi-continuous isomorphism. Clearly $S^* = 0$ on \mathcal{F}^{\perp} . On the other hand we can identify $(S^*)_{|\mathcal{F}}$ as the adjoint of the map \tilde{S} , i.e. $(\tilde{S})^*$, which is a bi-continuous isomorphism (since \tilde{S} is). In particular SS^* is invertible on \mathcal{F} . As for the last part of 2) we note that $P := S^*(SS^*)^{-1}S$ is bijective on \mathcal{H} and that it is also a projection, $P^2 = P$. Therefore P = 1, showing (2.27).

3). The invertibility of S^*S on \mathcal{H} follows from 2). Letting $Q = S(S^*S)^{-1}S^*$, we note

$$Q^2 = Q, \quad Q^* = Q, \quad \operatorname{ran}(Q) = \operatorname{ran}(S) = \mathcal{F},$$

showing $Q = \Pi$.

Put $\Pi' = 1 - \Pi$ and $H' = \Pi' H \Pi'$ with domain $\mathcal{D}(H') = \mathcal{D}(H) \cap \operatorname{ran}(\Pi') = \Pi' \mathcal{D}(H)$. We show below that indeed Π' preserves $\mathcal{D}(H)$ and that H' is self-adjoint. Let $\Pi_j = S_j S_j^*$ and $\Pi'_j = 1 - \Pi_j$; j = 1, 2. We recall that Condition 1.6 is imposed for $a = a_1$ and $a = a_2$. As in (2.16) we record that

$$\Pi'_{i}I_{j}\Pi_{j} \in \mathcal{L}(\mathcal{G}); \ j = 1, 2.$$

$$(2.29)$$

Here and henceforth we abbreviate $I_j(x) = I_{a_j}(x)$; j = 1, 2.

Lemma 2.8. Suppose (2.24). Then

1) The operators

$$\Pi - SS^* \in \mathcal{L}(\mathcal{G}, H^2(\mathbf{X})).$$
(2.30)

- 2) Π preserves $\mathcal{D}(H)$, and $\Pi'H\Pi$ and $\Pi H\Pi'$ (initially defined on $\mathcal{D}(H)$) extend to bounded operators on \mathcal{G} .
- 3) H' is self-adjoint.

Proof. 1). We shall use the identities

$$\Pi = S(S^*S)^{-1}S^* = SS^* + S\left(S^*S\right)^{-1} - 1\right)S^*,$$
(2.31a)

$$(S^*S)^{-1} - 1 = -\begin{pmatrix} 0 & s_{12} \\ s_{21} & 0 \end{pmatrix} + \begin{pmatrix} 0 & s_{12} \\ s_{21} & 0 \end{pmatrix} (S^*S)^{-1} \begin{pmatrix} 0 & s_{12} \\ s_{21} & 0 \end{pmatrix}, \quad (2.31b)$$

which follow from Proposition 2.7 and the general identity

$$(A+1)^{-1} - 1 = -A(A+1)^{-1} = -A + A(A+1)^{-1}A,$$

respectively. The statement (2.30) follows from Lemma 2.6, (2.31a) and (2.31b).

2). Writing $SS^* = \Pi_1 + \Pi_2$ it suffices (due to 1)) to show that Π_j preserves $\overline{\mathcal{D}}(H)$ and that $\Pi' H \Pi_j$ extends to a bounded operator on \mathcal{G} (note that boundedness

of $\Pi H \Pi'$ follows from boundedness of $\Pi' H \Pi$). Since Π_j reduces H_{a_j} clearly Π_j preserves $\mathcal{D}(H) = \mathcal{D}(H_{a_j})$, and since $\Pi' H \Pi_j = \Pi' \Pi'_j I_{a_j} \Pi_j \in \mathcal{L}(\mathcal{G})$, cf. (2.29), we are done.

3). Take $g \in \mathcal{D}((H')^*) \subset \Pi'\mathcal{G}$. The functional $\Pi'\mathcal{D}(H) \ni f \to \langle g, H'\Pi'f \rangle$ extends to a bounded functional, so by (2)) also $\mathcal{D}(H) \ni f \to \langle g, Hf \rangle$ extends to a bounded functional. This shows that $g \in \mathcal{D}(H^*) = \mathcal{D}(H)$. Whence $g \in \mathcal{D}(H')$.

Some parts of the previous lemma can be generalized as follows.

Lemma 2.9. Suppose (2.24). Then

1) For all $r, t \in \mathbb{R}$

 $\Pi - SS^* \in \mathcal{L}(L^2_r(\mathbf{X}), H^2_t(\mathbf{X})) \cap \mathcal{L}(H^{-2}_r(\mathbf{X}), L^2_t(\mathbf{X})).$

2) For all $t \in \mathbb{R}$ there exist extensions

$$\langle x \rangle^{\rho+1} \Pi' H \Pi, \langle x \rangle^{\rho+1} \Pi H \Pi' \in \mathcal{L}(L^2_t(\mathbf{X})).$$

Proof. For 1) we mimic the proof of Lemma 2.8. For 2) it suffices (seen by using 1)) to show that

$$\langle x \rangle^{\rho+1} \big(\Pi'_j - \Pi_i \big) I_j \Pi_j, \ \langle x \rangle^{\rho+1} \Pi_j I_j \big(\Pi'_j - \Pi_i \big) \in \mathcal{L} \big(L^2_t(\mathbf{X}) \big); \ i \neq j.$$

Next we implement (1.24a) of Condition 1.6. We obtain, cf. Lemma 2.6 and (2.23),

$$\forall r, t \in \mathbb{R} : \quad \Pi_i I_j \Pi_j, \ \Pi_j I_j \Pi_i \in \mathcal{L} \left(L_r^2(\mathbf{X}), L_t^2(\mathbf{X}) \right) \ (i \neq j), \\ \langle x \rangle^{\rho+1} \Pi'_j I_j \Pi_j, \ \langle x \rangle^{\rho+1} \Pi_j I_j \Pi'_j \in \mathcal{L} \left(L_t^2(\mathbf{X}) \right).$$

$$(2.32)$$

-			
	-		

2.3.1 $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$; the case $\lambda_0 = \Sigma_2$ and $\lambda_0 \notin \sigma_{\mathrm{pp}}(H')$

The case $\lambda_0 = \Sigma_2$ is simpler and there are better mapping properties of various operators compared to the case $\lambda_0 > \Sigma_2$. Consequently we pay special attention to the former case. We shall show the following extension of [Wa2, Lemma 2.1].

Lemma 2.10. Suppose Condition 2.5 with $\lambda_0 = \Sigma_2$ and (2.24). Then there exists $\epsilon_0 > 0$ such that the essential spectrum of $H' = \Pi' H \Pi'$ satisfies

$$\sigma_{\rm ess}(H') \subset [\lambda_0 + \epsilon_0, \infty). \tag{2.33}$$

Proof. Note

$$\lambda_0 = \Sigma_2 < \inf \sigma(H^b), \text{ for } b \notin \{a_1, a_2, a_{\max}\}.$$
(2.34)

In particular we have

$$\lambda_0 = \min \sigma_{\rm d}(H^{a_j}).$$

We introduce a auxiliary Hamiltonians (with inter-cluster momenta $p_j = p_{a_j}$)

$$\begin{split}
\breve{H} &= H' + p_1^2 \Pi_1 + p_2^2 \Pi_2, \\
\widetilde{H} &= H - \lambda_0 \big(\Pi_1 + \Pi_2 \big); \\
\mathcal{D}(\breve{H}) &= \mathcal{D}(\widetilde{H}) = \mathcal{D}(H).
\end{split}$$
(2.35a)

These operators differ by H-compact terms:

Note that the first and second terms of K_1 are *H*-compact due to Lemma 2.9. The relative compactness of the third and fourth terms of K_1 follows from rewriting $\Pi_j I_j \Pi_j = \varphi_j \otimes S_j^* I_j S_j \langle \varphi_j |$ and then invoking the complete analogue of (2.22). As for K_2 we can also use Lemma 2.9 first by replacing the factors of Π in the third term by $SS^* = \Pi_1 + \Pi_2$ and then expanding into four terms. We are left with considering the sum of cross terms

$$\Pi_1 H \Pi_2 + \Pi_2 H \Pi_1.$$

By writing

$$\Pi_{1}H\Pi_{2} = \Pi_{1}I_{1}\Pi_{2} + H_{a_{1}}\Pi_{1}\Pi_{2},
\Pi_{1}\Pi_{2} = \varphi_{1} \otimes s_{12}\langle\varphi_{2}|,$$
(2.36)

and then using (2.32) and Lemma 2.6, we conclude that the term $\Pi_1 H \Pi_2$ is *H*-compact. We can argue similarly for the term $\Pi_2 H \Pi_1$ and then conclude that also K_2 is *H*-compact.

Next, it follows from the very definition (2.35a) that \mathcal{F} reduces \tilde{H} . Whence $\sigma_{\text{ess}}(H') \subset \sigma_{\text{ess}}(\check{H})$, and consequently it suffices to show (2.33) with H' replaced by \check{H} and hence in turn with H' replaced by \tilde{H} (by the compactness shown above).

Consider a family of smooth non-negative functions $\{j_b | b \in \mathcal{A}, \#b = 2\}$ on **X** obeying that for some c > 0:

$$\sum_{\#b=2} j_b(x)^2 = 1,$$
(2.37a)

$$|x^{a}|j_{b}(x) \ge c|x|j_{b}(x) \text{ for } a \not\subset b \text{ and } |x| \ge 1,$$
(2.37b)

$$|\partial^{\alpha} j_b(x)| \le C_{\alpha} \langle x \rangle^{-|\alpha|} \text{ for all } \alpha \in \mathbb{N}_0^{\dim \mathbf{X}}.$$
(2.37c)

We have, using this family of functions,

$$H = \sum_{\#a=2} j_a H_a j_a + \sum_{\#a=2} I_a j_a^2 - \sum_{\#a=2} |\nabla j_a|^2 = \sum_{\#a=2} j_a H_a j_a + K,$$
(2.38)

where K is H-compact.

For $a \neq a_j$, j = 1, 2, it follows from (2.34) that $\inf \sigma(H_a) = \inf \sigma(H^a) \ge \lambda_0 + \epsilon'_0$ for some $\epsilon'_0 > 0$. Therefore,

$$\sum_{\#a=2, a \neq a_j, j=1,2} j_a H_a j_a \ge (\lambda_0 + \epsilon'_0) \sum_{\#a=2, a \neq a_j, j=1,2} j_a^2.$$
(2.39)

2.3.1. $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$; the case $\lambda_0 = \Sigma_2$ and $\lambda_0 \notin \sigma_{pp}(H')$

Since λ_0 is the lowest eigenvalue of H^{a_j} , j = 1, 2, there exists $\epsilon_1 > 0$ such that

$$\Pi'_j H_{a_j} \Pi'_j \ge (\lambda_0 + \epsilon_1) \Pi'_j$$

Hence for $a = a_j$, j = 1, 2, we obtain (assuming in the last step that $\lambda_0 + \epsilon_1 \leq 0$ and using that $p_j^2 \geq 0$)

$$j_{a_{j}}H_{a_{j}}j_{a_{j}} \ge (\lambda_{0} + \epsilon_{1})j_{a_{j}}\Pi'_{j}j_{a_{j}} + j_{a_{j}}H_{a_{j}}\Pi_{j}j_{a_{j}}$$

$$\ge (\lambda_{0} + \epsilon_{1})j_{a_{j}}^{2} + \lambda_{0}j_{a_{j}}\Pi_{j}j_{a_{j}}.$$
(2.40)

It is a consequence of (2.37a) and (2.37b) that the operator $\Pi_j j_{a_j}^2 - \Pi_j$ is *H*-compact. Using (2.37c) it follows by a Taylor expansion that $[j_{a_j}, \Pi_j]\langle x \rangle$ is bounded (cf. (3.42)). We conclude that also

$$K'_{j} := \lambda_0 \left(j_{a_j} \Pi_j j_{a_j} - \Pi_j \right) \text{ is } H \text{-compact.}$$

$$(2.41)$$

We write (2.40) as

$$j_{a_j}H_{a_j}j_{a_j} - \lambda_0 \Pi_j \ge (\lambda_0 + \epsilon_1)j_{a_j}^2 + K'_j, \ j = 1, 2.$$
(2.42)

Let $\epsilon_0 = \min{\{\epsilon'_0, \epsilon_1\}}$. We then deduce from (2.37a), (2.38), (2.39) and (2.42) that

$$\hat{H} \ge (\lambda_0 + \epsilon_0) + \hat{K}, \tag{2.43}$$

where $\widetilde{K} = K + K'_1 + K'_2$.

Since \tilde{K} is \tilde{H} -compact it follows from (2.43) and Weyl's theorem [RS, Theorem XIII.14] that

$$\sigma_{\rm ess}(\tilde{H}) \subset [\lambda_0 + \epsilon_0, \infty).$$

Remark 2.11. By appropriately enlarging the projection Π as to include the span of all threshold eigenstates corresponding to thresholds $\lambda^a \in [\Sigma_2, \Sigma_3 - \epsilon_0)$ for any given $\epsilon_0 > 0$, one can make sure that $\sigma_{ess}(H') \subset [\Sigma_3 - \epsilon_0, \infty)$. Here $H' = \Pi' H \Pi'$ as before, and the proof is essentially the same as the one of Lemma 2.10. This trick of 'subtracting' all low-energy 2-cluster channels is limited to energies below Σ_3 . We prefer for simplicity of presentation to treat two-cluster thresholds $\lambda_0 > \Sigma_2$ on an equal footing not distinguishing between the cases $\lambda_0 \in (\Sigma_2, \Sigma_3)$ and $\lambda_0 > \Sigma_3$. In fact our results would be the same anyway for the two cases.

Assume from this point that

$$\lambda_0$$
 is not an eigenvalue of H' . (2.44)

By Lemma 2.10 we can then chose $\delta > 0$ such that $\sigma(H') \cap \{|z - \lambda_0| \leq \delta\} = \emptyset$. Due to Proposition 2.1

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z), \qquad (2.45)$$

where the quantities (2.4a)–(2.4d) in this case are analytic in $\{|z - \lambda_0| < \delta\}$.

Let us analyse the form of $E_{\mathcal{H}}(z)$. For $f \in \mathcal{F}$, $f = \varphi_1 \otimes f_1 + \varphi_2 \otimes f_2$ and we can write

$$Hf = \lambda_0 f + (p_1^2 + I_1)(\varphi_1 \otimes f_1) + (p_2^2 + I_2)(\varphi_2 \otimes f_2),$$

recalling $p_j = p_{a_j}$ and $I_j = I_{a_j}$, j = 1, 2. Therefore, for $i, j, k \in \{1, 2\}$

$$S_k^*(H-z)S_k = p_k^2 + \lambda_0 - z + W_k(x_k), \qquad (2.46a)$$

$$S_i^*(H-z)S_j = \check{s}_{ij}(z) + W_{ij},$$
 (2.46b)

$$S_i^* H R'(z) H S_j = -K_{ij}(z),$$
 (2.46c)

where

$$\check{s}_{ij}(z) = \langle \varphi_i, \varphi_j \otimes (p_j^2 + \lambda_0 - z) \cdot \rangle_i, \qquad (2.47a)$$

$$W_{ij} = \langle \varphi_i, I_j \varphi_j \otimes \cdot \rangle_i, \qquad (2.47b)$$

$$W_k(\cdot) = W_{kk} = \langle \varphi_k, I_k \varphi_k \rangle_k(\cdot), \qquad (2.47c)$$

$$K_{ij}(z) = -\langle \varphi_i, I_i R'(z) I_j(\varphi_j \otimes \cdot) \rangle_i.$$
(2.47d)

Introducing $\mathcal{H}^2 = H^2(\mathbf{X}_1) \oplus H^2(\mathbf{X}_2)$ this yields the following expression for the operator $E_{\mathcal{H}}(z) : \mathcal{H}^2 \to \mathcal{H}$.

$$E_{\mathcal{H}}(z) = z - \lambda_0 - \begin{pmatrix} -\Delta_{x_1} + W_1(x_1) + K_{11}(z) & \check{s}_{12}(z) + W_{12} + K_{12}(z) \\ \check{s}_{21}(z) + W_{21} + K_{21}(z) & -\Delta_{x_2} + W_2(x_2) + K_{22}(z) \end{pmatrix}.$$
(2.48)

Clearly $\check{s}_{ij}(z) = s_{ij}(\lambda_0 - z + p_j^2)$ for $i \neq j$, and therefore Lemma 2.6 yields that $\check{s}_{ij}(z)$ for $i \neq j$ is bounded and in fact polynomially decreasing (uniformly in z near λ_0). Here and henceforth an operator $b_{ij} : L^2(\mathbf{X}_j) \to L^2(\mathbf{X}_i)$ is said to be polynomially decreasing if for all $r, t \in \mathbb{R}$

$$b_{ij} \in \mathcal{C}\big(H_r^2(\mathbf{X}_j), L_t^2(\mathbf{X}_i)\big) \cap \mathcal{C}\big(L_r^2(\mathbf{X}_j), H_t^{-2}(\mathbf{X}_i)\big).$$

We note that also the operators W_{12} and W_{21} are polynomially decreasing. An operator B on \mathcal{H} is said to be polynomially decreasing if its entries $b_{ij} : L^2(\mathbf{X}_j) \to L^2(\mathbf{X}_i)$ are polynomially decreasing.

We claim that also $K_{12}(z)$ and $K_{21}(z)$ are polynomially decreasing, in fact uniformly in z near λ_0 . To see this it suffices (by symmetry) to consider $K_{12}(z)$, and it suffices to show that $K_{12}(z) \in \mathcal{L}(L^2_{-s}(\mathbf{X}_2), L^2_s(\mathbf{X}_1))$ for any $s \geq 0$. So let us fix $s \geq 0$. A small consideration using the argument for (2.16) and the polynomial decay of the cluster bound states shows, that it suffices to check that

$$\langle x^1 \rangle^{-2s} \langle x \rangle^s R'(z) \langle x \rangle^s \langle x^2 \rangle^{-2s}$$
 is uniformly bounded near λ_0 . (2.49)

Recall that $\{z | |z - \lambda_0| \le \delta\}$ is included in the resolvent set of H' for a small $\delta > 0$, cf. (2.44). We also record the following elementary estimates,

$$\langle x+y\rangle^t \le 2^{|t|/2} \langle x\rangle^{|t|} \langle y\rangle^t, \tag{2.50a}$$

$$\langle x \rangle \le C(\langle x^1 \rangle + \langle x^2 \rangle) \le 2C \langle x^1 \rangle \langle x^2 \rangle.$$
 (2.50b)

In turn, by interpolation and by using (2.50b), the assertion (2.49) is a consequence of

$$\langle x^2 \rangle^t R'(z) \langle x^2 \rangle^{-t}$$
 and $\langle x^1 \rangle^{-t} R'(z) \langle x^1 \rangle^t$ are uniformly bounded; $t = 2s.$ (2.51)

We bound the first expression only. Representing, using notation of the proof of Lemma 2.10,

$$R'(z) = \breve{R}(z)\Pi', \ \breve{R}(z) = (\breve{H} - z)^{-1},$$
(2.52)

we first bound $\langle x^2 \rangle^t \breve{R}(z) \langle x^2 \rangle^{-t}$. Whence we want to bound $\langle \kappa x^2 \rangle^t \breve{R}(z) \langle \kappa x^2 \rangle^{-t}$ with $\kappa = 1$, for which it suffices to bound this quantity for any small $\kappa > 0$. We show such bound uniformly in z near λ_0 . As in the proof of Lemma 2.10

$$\check{H} = H - \lambda_0 (\Pi_1 + \Pi_2) - K_1 + K_2.$$
(2.53)

Letting $\breve{H}_{\kappa,t} = \langle \kappa x^2 \rangle^t \breve{H} \langle \kappa x^2 \rangle^{-t}$ we obtain from (2.53) that

$$\breve{H}_{\kappa,t} = \breve{H} + \mathcal{O}(\kappa)(\breve{H} - \mathrm{i}).$$

For example, seen by using Taylor expansion and (2.50a),

$$\langle \kappa x^2 \rangle^t \Pi_1 \langle \kappa x^2 \rangle^{-t} - \Pi_1 = [\langle \kappa \pi^2 x \rangle^t - \langle \kappa \pi^2 x_1 \rangle^t, \Pi_1] \langle \kappa x^2 \rangle^{-t}$$

= $\kappa \mathcal{O}(\langle \kappa \pi^2 x_1 \rangle^{t-1}) \langle \kappa x_1 \rangle^{-t}$ (2.54)
= $\mathcal{O}(\kappa).$

We can treat the terms K_1 and K_2 similarly. Therefore

$$(\breve{H}_{\kappa,t}-z)\breve{R}(z) = 1 + \mathcal{O}(\kappa)(\breve{H}-i)\breve{R}(z)$$

is invertible for $|z - \lambda_0| \leq \delta$ and for small $\kappa > 0$. This shows that $(\check{H}_{\kappa,t} - z)^{-1}$ is uniformly bounded accomplishing our first goal. The second goal, cf. (2.52), is to bound $\langle x^2 \rangle^t \Pi' \langle x^2 \rangle^{-t}$, but indeed $\langle x^2 \rangle^t \Pi \langle x^2 \rangle^{-t}$ is bounded due to Lemma 2.9 1) and (2.54). Consequently we have justified (2.51) and therefore shown that $K_{12}(z)$ is polynomially decreasing uniformly near λ_0 .

One can somewhat similarly show that

 $\langle x_j \rangle^{\rho_1} K_{jj}(z) \langle x_j \rangle^{\rho_2}$ is uniformly bounded for on $L^2(\mathbf{X}_j)$ for $\rho_1 + \rho_2 \le 2\rho + 2$. (2.55)

In fact we can use a refinement of (2.16) related to (2.23) and Lemma 2.9 2). Note here the Taylor expansion $I_j^{(1)}(x) = I_j^{(1)}(x_j) + \mathcal{O}(\langle x^j \rangle^{\rho+2}) \langle x_j \rangle^{-\rho-1}$, cf. (2.50a), which in turn leads to the following bounds for all $s \in \mathbb{R}$ and for $\rho' \in \{\rho_1, \rho_2\}$,

$$\Pi_j \langle x_j \rangle^{\rho'} I_j \Pi', \quad \Pi' I_j \langle x_j \rangle^{\rho'} \Pi_j \in \mathcal{L} \big(L^2_s(\mathbf{X}), L^2_{\rho-\rho'+1+s}(\mathbf{X}) \big).$$

We deduce (2.55) by combining these bounds with the following consequence of (2.50a) and (2.51),

$$\langle x \rangle^{s} \mathring{R}(z) \Pi' \langle x \rangle^{-s}$$
 is bounded for all $s \in \mathbb{R}$ (uniformly near λ_0). (2.56)

Due to the above discussion and (2.22) we finally obtain a simplified version of (2.48):

Proposition 2.12. Suppose Condition 2.5 with $\lambda_0 = \Sigma_2$, (2.24) and (2.44). As a bounded operator $E_{\mathcal{H}}(z) : \mathcal{H}^2 \to \mathcal{H}$ one then has

$$E_{\mathcal{H}}(z) \equiv z - \lambda_0 - \begin{pmatrix} -\Delta_{x_1} + W_1(x_1) + K_{11}(z) & 0\\ 0 & -\Delta_{x_2} + W_2(x_2) + K_{22}(z) \end{pmatrix}, \quad (2.57)$$

where " \equiv " means the equality modulo a polynomially decreasing term which depends holomorphically on z near λ_0 . We have

$$W_j(x_j) - I_j^{(1)}(x_j) \in \mathcal{L}\left(H_s^2(\mathbf{X}_j), L_{\rho+1+s}^2(\mathbf{X}_j)\right) \text{ for } s \in \mathbb{R},$$
(2.58a)

$$W_j(x_j) - I_j^{(1)}(x_j) \in \mathcal{C}\left(H_s^2(\mathbf{X}_j), L_t^2(\mathbf{X}_j)\right) \text{ for } s \in \mathbb{R}, \ t < \rho + 1 + s,$$
(2.58b)

$$K_{jj}(z) \in \mathcal{L}\left(L_s^2(\mathbf{X}_j), L_{2\rho+2+s}^2(\mathbf{X}_j)\right) \text{ for } s \in \mathbb{R}.$$
(2.58c)

Moreover the operator $K_{jj}(z)$ depends holomorphically on z, and the potential W_j is p_j^2 -compact with the singularities of located in a bounded set.

Remark 2.13. We shall prefer a version of Proposition 2.12 based on forms rather than operators, although this is not essential. Thus we consider $E_{\mathcal{H}}(z)$ as an operator $E_{\mathcal{H}}(z) : H^1(\mathbf{X}_1) \oplus H^1(\mathbf{X}_2) \to H^{-1}(\mathbf{X}_1) \oplus H^{-1}(\mathbf{X}_2)$. Now an operator $b_{ij} : L^2(\mathbf{X}_j) \to$ $L^2(\mathbf{X}_i)$ is said to be *polynomially decreasing* if for all $r, t \in \mathbb{R}$

$$b_{ij} \in \mathcal{C}(H_r^1(\mathbf{X}_j), H_t^{-1}(\mathbf{X}_i)).$$

Also we note that the expansion (2.58a) should be given the interpretation of being in the space $\mathcal{L}(H_s^1(\mathbf{X}_j), H_{\rho+1+s}^{-1}(\mathbf{X}_j))$, $s \in \mathbb{R}$, rather than in the stated space (and similarly for (2.58b)). Note for comparison that $I_j^{(1)}(x_j) \in \mathcal{L}(L_s^2(\mathbf{X}_j), L_{\rho+s}^2(\mathbf{X}_j))$, $s \in \mathbb{R}$. With these modifications (2.57) still holds. Moreover it is easy to show by a resolvent equation that the null space ker $E_{\mathcal{H}}(\lambda_0)$ is independent on whether the operator or the form interpretation of $E_{\mathcal{H}}(\lambda_0)$ is used.

2.4 The case $\lambda_0 \in \sigma_{pp}(H')$

We discuss briefly the modifications needed in the previous two sections to treat the case $\lambda_0 \in \sigma_{\rm pp}(H')$.

2.4.1 $\lambda_0 \in \sigma_{\rm pp}(H')$; non-multiple case

For simplicity we assume m = 1 in the setting of Section 2.2. The corresponding eigenfunction is denoted by φ (rather than φ_1), and we shall use the notation $\langle \varphi | \cdot \rangle_0 = \langle \varphi, \cdot \rangle_{L^2(\mathbf{X}^{a_0})}$. We follow [Wa2] assuming for simplicity that λ_0 is a simple eigenvalue of H' and introduce a corresponding normalized eigenfunction ψ , $(H' - \lambda_0)\psi = 0$. Let $I_0 = I_{a_0}$.

Now $\mathcal{H} = L^2(\mathbf{X}_{a_0}) \oplus \mathbb{C}$, and $S : \mathcal{H} \to \mathcal{G} = L^2(\mathbf{X})$ is given by

$$S(f,c) = \varphi \otimes f + c\psi$$

We let the orthogonal projection SS^* onto the range of this new S be denoted by $1-\Pi''$ (for the S in Section 2.2 the projection is denoted by $1-\Pi'$), and we introduce
$H'' = \Pi'' H \Pi''$ on the Hilbert space $\Pi'' \mathcal{G}$. By construction λ_0 is not an eigenvalue of H''. Using (2.4a)–(2.4d) as before we obtain (2.19) now with

$$-E_{\mathcal{H}}(z) = -S^* \left(z - H + HR''(z)H \right) S$$

=
$$\begin{pmatrix} \left(\lambda_0 - z + p_{a_0}^2 + \langle \varphi | I_0 | \varphi \otimes \cdot \rangle_0 - \langle \varphi | I_0 R''(z) I_0 | \varphi \otimes \cdot \rangle_0 \right) & \langle \varphi, I_0 \psi \rangle_0 \\ \langle I_0 \psi, \varphi \otimes \cdot \rangle & \lambda_0 - z \end{pmatrix}.$$

Let

$$H = H - \lambda_0 \Pi, \quad \Pi = \varphi \otimes \langle \varphi | \cdot \rangle_0,$$

$$\breve{H} = \widetilde{H} - \breve{K}; \qquad (2.59a)$$

$$\breve{K} = \Pi I_0 + I_0 \Pi - \Pi I_0 \Pi + \lambda_0 | \psi \rangle \langle \psi |.$$

Note that \check{K} is *H*-compact. This construction is motivated by the following direct sum decomposition.

$$\breve{H} = H'' + p_{a_0}^2 \Pi; \ H'' = \Pi'' H \Pi''.$$
(2.59b)

The basic structure of resolvents is given as follows.

$$R''(z) = \breve{R}(z) - (p_{a_0}^2 - z)^{-1}\Pi - z^{-1}|\psi\rangle\langle\psi|,$$

$$R''(z) = \Pi''\breve{R}(z)\Pi'' = \Pi''\breve{R}(z) = \breve{R}(z)\Pi''; \quad \breve{R}(z) = (\breve{H} - z)^{-1}.$$
(2.59c)

We shall prove that $\psi \in H^2_{\infty}(\mathbf{X})$ (see Theorem 3.12), which in turn implies 'good' properties of $\breve{R}(z)$ and Π'' near λ_0 (see Remark 3.21).

2.4.2 $\lambda_0 \in \sigma_{\rm pp}(H')$; multiple case

We adapt the setting of Section 2.3 in the case (2.44) of Subsection 2.3.1 is not fulfilled and without imposing the condition $\lambda_0 = \Sigma_2$ as in that subsection. In particular (2.24) is satisfied.

Assume for simplicity that λ_0 is a simple eigenvalue of H'. We introduce a corresponding normalized eigenfunction $(H' - \lambda_0)\psi = 0$. Now $\mathcal{H} = L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2) \oplus \mathbb{C}$, and $S : \mathcal{H} \to \mathcal{G} = L^2(\mathbf{X})$ is given by

$$S(f_1, f_2, c) = \varphi_1 \otimes f_1 + \varphi_2 \otimes f_2 + c\psi.$$

We introduce Π'' in terms of this *S* as in the previous subsection and let again $H'' = \Pi'' H \Pi''$. Note that $\Pi'' = 1 - \Pi - |\psi\rangle\langle\psi| = \Pi' - |\psi\rangle\langle\psi|$ where Π is given as in Section 2.3. We obtain (2.19) now with

$$E_{\mathcal{H}}(z) = S^* \big(z - H + HR''(z)H \big) S \big),$$

and this operator has a similar representation as in the previous subsection, now by a 3 × 3-block representation $(e_{ij})_{i,j\leq 3}$ rather than a 2 × 2-block representation as given there. Here $(e_{ij})_{i,j\leq 2}$ is given as in Proposition 2.12 (with R'(z) replaced by R''(z)), $e_{33} = z - \lambda_0$, $e_{i3} = -\langle \varphi_i, I_i \psi \rangle_i$ and $e_{3i} = e_{i3}^* = \langle e_{3i} |; i = 1, 2$. The analogue of (2.35a) reads

$$\begin{aligned}
\breve{H} &= H'' + p_1^2 \Pi_1 + p_2^2 \Pi_2, \\
\widetilde{H} &= H - \lambda_0 \big(\Pi_1 + \Pi_2 \big); \\
\mathcal{D}(\breve{H}) &= \mathcal{D}(\widetilde{H}) = \mathcal{D}(H).
\end{aligned}$$
(2.60a)

These operators differ by H-compact terms, cf. (2.35b):

$$\widetilde{H} = \widetilde{H} - K_1 + K_2;$$
(2.60b)
$$K_1 = \Pi H \Pi' + \Pi' H \Pi + \Pi_1 I_1 \Pi_1 + \Pi_2 I_2 \Pi_2 + \lambda_0 |\psi\rangle \langle\psi|,$$

$$K_2 = \Pi_1 H \Pi_1 + \Pi_2 H \Pi_2 - \Pi H \Pi.$$

The basic structure of resolvents is given as follows.

$$R''(z) = \breve{R}(z) - (p_1^2 \Pi_1 + p_2^2 \Pi_2 - z)^{-1} \Pi - z^{-1} |\psi\rangle \langle\psi|,$$

$$R''(z) = \Pi'' \breve{R}(z) \Pi'' = \Pi'' \breve{R}(z) = \breve{R}(z) \Pi''; \quad \breve{R}(z) = (\breve{H} - z)^{-1}.$$
(2.60c)

Again we have good decay properties of ψ , cf. Theorem 3.12 and Remark 3.20, which in turn yields good properties of $\breve{R}(z)$ near λ_0 , cf. Remark 3.21.

2.5 Multiple two-cluster case, $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$

It may happen that the condition (2.24) is not satisfied. We start out by presenting an example (the only one we know). It is given by an atomic type 2-body Schrödinger operator with a third particle of infinite mass fixed at the origin (fitting into the framework of N-body case of Section 1.3):

$$H = \sum_{j=1}^{2} (-\Delta_{x^j} + V_j(x^j)) + V_{12}(x^1 - x^2), \qquad (2.61)$$

where $x^j \in \mathbb{R}^n$. Here $x = (x^1, x^2)$ is used as global coordinates on \mathbb{R}^{2n} . Let λ_0 be the lowest threshold of H. Assume that this threshold is double and is attained by the lowest eigenvalue of H_j

$$H_j = -\Delta_{x^j} + V_j(x^j), \quad j = 1, 2.$$

In this case, the condition (2.24) is not satisfied. In fact, let $\varphi_j(x^j)$ be the eigenfunction of H_j associated with λ_0 . Then clearly one has

$$\mathcal{F}_1 \cap \mathcal{F}_2 = \operatorname{span}\{\varphi_1(x^1)\varphi_2(x^2)\}.$$

2.5.1 $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$; a general approach

We discuss a method which is easy to generalize to the case of an arbitrary multiplicity of the two-cluster threshold λ_0 .

We define

$$\mathcal{H} = \left\{ f \in L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2) \mid f \perp \ker S^*S \right\},\tag{2.62}$$

where the components of this $S = (S_i)$ are given by (2.25). Due to Lemma 2.6 the space ker S^*S is finite dimensional consisting of vectors with components in $H^2_{\infty}(\mathbf{X}_j)$. By assumption dim(ker S^*S) ≥ 1 . Clearly $S : \mathcal{H} \to \mathcal{F} = \mathcal{F}_1 + \mathcal{F}_2 \subset \mathcal{G} = L^2(\mathbf{X})$ is a continuous isomorphism. Arguing by contradiction as in the proof of Proposition 2.7 we deduce that in fact $S : \mathcal{H} \to \mathcal{F}$ is bi-continuous. In particular \mathcal{F} is closed in \mathcal{G} , and we let correspondingly Π , Π' , H' and R'(z) be given (as in Section 2.1). If we consider S as a map from the bigger space $L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2)$ (as in (2.62)) the notation $S^*g, g \in \mathcal{G}$, may seem ambiguous. However this is in fact not the case since then $S^*g \perp \ker S^*S$, so $S^*g \in \mathcal{H}$. For this reason the conclusion of Lemmas 2.8 and 2.9 are still valid and the formula (2.48) applies again.

As before we would like to use (2.48) in a neighbourhood of λ_0 . Let us here assume that $\lambda_0 = \Sigma_2$ and (2.44). Then of course we can let $z = \lambda_0$ in (2.48) and obtain formulas as in Proposition 2.12 and Remark 2.13. We consider correspondingly the operator $E_{\mathcal{H}}(\lambda_0)$ either as an operator mapping $(H^2(\mathbf{X}_1) \oplus H^2(\mathbf{X}_2)) \cap \mathcal{H} \to \mathcal{H}$ or as an operator mapping

$$(H^1(\mathbf{X}_1) \oplus H^1(\mathbf{X}_2)) \cap \mathcal{H} \to \{f \in H^{-1}(\mathbf{X}_1) \oplus H^{-1}(\mathbf{X}_2) \mid f \perp \ker S^*S\}.$$

If $\lambda_0 \in \sigma_{\rm pp}(H')$ we need to modify the construction of S and $E_{\mathcal{H}}(\lambda_0)$. This is doable along the lines of Subsection 2.4.2. Finally letting P_0 denote the orthogonal projection onto ker S^*S in $L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2)$ it is convenient to study $E_{\mathcal{H}}(\lambda_0) + P_0$ on $L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2)$ (rather than $E_{\mathcal{H}}(\lambda_0)$ on \mathcal{H}). This is because we have a 'good parametrix' of diag (h_1, h_2) on $L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2)$.

In Section 4.3 we shall see what the above ideas amount to in the setting of the models of physics introduced in Chapter 1. This will be a general treatment not assuming $\lambda_0 = \Sigma_2$.

Chapter 3 Spectral analysis of H' near λ_0

In the bulk of this chapter we impose Conditions 1.5 and 2.4. We shall prove various N-body resolvent estimates for the operator H' appearing in the Grushin method (or more prescisely for the operator \check{H} defined below). The analysis overlaps [AIIS], in particular it is based on an appropriate Mourre estimate. Sharing the spirit of [AIIS] our procedure avoids in any other sense the 'classical Mourre theory' [Mo, Je, JMP, Wa1]. The multiple two-cluster case can be treated in a similar way, although it is notationally more complicated, see Remarks 3.20–3.22.

Recall from Section 2.2 the notation $I_0 = I_{a_0}$ and $p_0 = p_{a_0}$. We introduce the following modifications of H (note the similarity with (2.35a) and (2.35b)),

$$\begin{split} \widetilde{H} &= H - \lambda_0 \Pi, \\ \widetilde{H} &= \widetilde{H} - \breve{K}; \\ \widetilde{K} &= \Pi I_0 + I_0 \Pi - \Pi I_0 \Pi. \end{split}$$
(3.1a)

Note that \check{K} is *H*-compact. This construction is motivated by the following direct sum decomposition.

$$\check{H} = H' + p_0^2 \Pi; \ H' = \Pi' H \Pi'.$$
 (3.1b)

The basic structure for resolvents is given as follows.

$$R'(z) = \breve{R}(z) - (p_0^2 - z)^{-1}\Pi,$$

$$R'(z) = \Pi'\breve{R}(z)\Pi' = \Pi'\breve{R}(z) = \breve{R}(z)\Pi'; \quad \breve{R}(z) = (\breve{H} - z)^{-1}.$$
(3.2a)

Note that (3.2a) yields 'good estimates' of R'(z) near λ_0 provided we can show 'good estimates' of $\check{R}(z)$ near λ_0 . The goal of this chapter is to prove the latter, which more or less correspond to (3.79) and Corollary 3.19, stated as follows: Suppose $\lambda_0 \notin \sigma_{\rm pp}(\check{H})$. Then there exists the strong weak-star limits

s-w^{*}-lim
$$\breve{R}(\lambda_0 \pm i\epsilon) = \breve{R}(\lambda_0 \pm i0) \in \mathcal{L}(\mathcal{B}_{1/2}(\mathbf{X}), \mathcal{B}_{1/2}^*(\mathbf{X})),$$
 (3.2b)

and if $\breve{R}(\lambda_0 + i0)f = \breve{R}(\lambda_0 - i0)f$ for a given $f \in L^2_s$ for some s > 1/2, then

$$\check{R}(\lambda_0 + i0)f = \check{R}(\lambda_0 - i0)f \in L^2_{s-1}.$$
(3.2c)

We shall prove a Mourre estimate for \check{H} near λ_0 and show that λ_0 cannot be an accumulating point of eigenvalues of \check{H} (which are the same as those of H'). If λ_0 is not an eigenvalue, the limiting absorption principle and some microlocal estimates hold for the resolvent of \check{H} near λ_0 . If λ_0 is an eigenvalue of \check{H} , the associated eigenfunctions are polynomially decaying. We don't assume that $\lambda_0 < \Sigma_3$.

The case of a multiple two-cluster threshold can be treated similarly and will not be discussed in detail. See Remark 3.20 for a discussion and see (4.37a)-(4.37c) for results similar to (3.2a)-(3.2c).

3.1 Mourre estimate

We shall use the vector field constructed by Graf [Gr] and the associated family of conjugate operators, cf. [Sk1]–[Sk3] and [IS1]. This vector field satisfies the following properties, cf. [Sk3, Lemma 4.3].

Lemma 3.1. There exist on **X** a smooth vector field $\tilde{\omega}$ with symmetric derivative $\tilde{\omega}_*$ and a partition of unity $\{\tilde{q}_a\}$ indexed by $a \in \mathcal{A}$ and consisting of smooth functions, $0 \leq \tilde{q}_a \leq 1$, such that for some positive constants r_1 and r_2

- (1) $\tilde{\omega}_*(x) \ge \sum_a \pi_a \tilde{q}_a$.
- (2) $\tilde{\omega}^a(x) = 0$ if $|x^a| < r_1$.
- (3) $|x^b| > r_1$ on $\operatorname{supp}(\tilde{q}_a)$ if $b \not\subset a$.
- (4) $|x^a| < r_2 \text{ on } \operatorname{supp}(\tilde{q}_a).$
- (5) For all $\alpha \in \mathbb{N}_0^{\dim \mathbf{X}}$ and $k \in \mathbb{N}_0$ there exist $C \in \mathbb{R}$:

$$\left|\partial_x^{\alpha} \tilde{q}_a\right| + \left|\partial_x^{\alpha} (x \cdot \nabla)^k \left(\tilde{\omega}(x) - x\right)\right| \le C.$$

For each $a \in \mathcal{A}$ there is a similar vector field, denoted by $\tilde{\omega}^a$, and from the construction of these vector fields there is a relationship we are going to use (see [Sk2, Appendix A] for a proof for the model of Section 1.2.1, see also [Sk6, Section 5]).

For any $\delta > 0$ there exists $\widetilde{R} = \widetilde{R}(\delta) > 1$ such that for all $a \in \mathcal{A}$

$$\widetilde{\omega}(x) = x_a + \widetilde{\omega}^a(x^a) \text{ for all } x \in \mathbf{Y},$$

$$\mathbf{Y} = \mathbf{Y}_{a,\delta,\widetilde{R}} := \{ x \in \mathbf{X} \mid |x| > \widetilde{R}, \, |x^b| > \delta |x| \text{ if } b \not\subset a \}.$$
(3.3)

Considering rescaled vector fields $\tilde{\omega}_R^a(x) := R\tilde{\omega}(x^a/R), R \ge 1$, obviously a consequence of (3.3) is the analogous result for the rescaled fields,

$$\tilde{\omega}_R(x) = x_a + \tilde{\omega}_R^a(x^a) \text{ for all } x \in \mathbf{Y}_{a,\delta,R\tilde{R}}, R \ge 1.$$
 (3.4)

Now, proceeding as in [Sk3], we introduce

$$A_R = \frac{1}{2} \left(\tilde{\omega}_R(x) \cdot p + p \cdot \tilde{\omega}_R(x) \right), \ R \ge 1,$$

and a function $d : \mathbb{R} \to \mathbb{R}$ by

$$d(\lambda) = \begin{cases} \inf_{\tau \in \mathcal{T}(\lambda)} (\lambda - \tau), \ \mathcal{T}(\lambda) \coloneqq \mathcal{T}() = \mathcal{T} \cap] - \infty, \lambda \neq \emptyset, \\ 1, \ \mathcal{T}(\lambda) = \emptyset. \end{cases}$$
(3.5)

These devices enter into the following Mourre estimate (we refer to [PSS] for another N-body Mourre estimate). We refer to [Sk3, Corollary 4.5] noting that all inputs needed for the proof are stated in Lemma 3.1.

Lemma 3.2. For all $\lambda \in \mathbb{R}$ and $\epsilon > 0$ the exists $R_0 \ge 1$ such that for all $R \ge R_0$ there is a neighbourhood \mathcal{V} of λ and a compact operator K on L^2 such that

$$f(H)i[H, A_R]f(H) \ge f(H)\{2d(\lambda) - \epsilon - K\}f(H) \text{ for all real } f \in C^{\infty}_{c}(\mathcal{V}).$$
(3.6)

We can write $\tilde{\omega} = \nabla r^2/2$ for some positive smooth function r with $r^2 - x^2$ bounded, cf. [De3].

A basic ingredient of our procedure is the operator

$$B = \frac{1}{2} \big(\omega(x) \cdot p + p \cdot \omega(x) \big) \in \mathcal{L}(H^1, L^2), \tag{3.7}$$

where $\omega = \omega_R = \tilde{\omega}_R/r_R$, $r_R(x) = Rr(\frac{x}{R})$. Note that $\tilde{\omega}_R = \nabla r_R^2/2$, and whence that $\omega_R = \nabla r_R$. We shall suppress the dependence of the parameter R (which eventually is taken as a large number, depending on λ). In particular we shall slightly abuse the notation writing for example r rather than the rescaled version r_R . Using the notation **D** for the Heisenberg derivative $i[H, \cdot]$ we note the computations $2B = \mathbf{D}r$, $A = r^{1/2}Br^{1/2}$ and (formally)

$$\mathbf{D}B = r^{-1/2} \big(\mathbf{D}A - 2B^2 \big) r^{-1/2} + \mathcal{O}(r^{-3}).$$
(3.8)

Here the function

$$\mathcal{O}(r^{-3}) = \frac{1}{2}\omega \cdot (\nabla^2 r)\omega/r^2 = r^{-3}v(x),$$

where v belongs to the algebra $\mathcal{F} = \mathcal{F}(\mathbf{X})$ of smooth functions on **X** obeying

$$\forall \alpha \in \mathbb{N}_0^{\dim \mathbf{X}} \quad \forall k \in \mathbb{N}_0 : |\partial_x^{\alpha} (x \cdot \nabla)^k v(x)| \le C_{\alpha,k}$$

Note also that the function $r^2 - x^2 \in \mathcal{F}$. Obviously $\mathcal{F}(\mathbf{X}) \supset \mathcal{F}(\mathbf{X}^a)$ for any $a \neq a_{\min}$. The exact computation of **D**A reads

$$\mathbf{D}A = 2p\tilde{\omega}_*\left(\frac{x}{R}\right)p - \left(4R^2\right)^{-1}\left(\triangle(\nabla\cdot\tilde{\omega})\right)\left(\frac{x}{R}\right) - R\tilde{\omega}\left(\frac{x}{R}\right)\cdot\nabla V.$$
(3.9)

In particular

$$\mathbf{D}A = \mathbf{i}[H, A] = \sum_{|\alpha| \le 2} v_{\alpha} p^{\alpha}; \quad v_{\alpha} \in \mathcal{F},$$

which make sense as a bounded form on $H^1 = Q(H)$. Although we define **D**A by (3.9), it can be computed as a strong limit,

$$i[H, A] = \operatorname{s-lim}_{t \to 0} t^{-1} (H e^{itA} - e^{itA} H) \in \mathcal{L}(H^1, H^{-1}).$$
(3.10)

Similarly the formal commutator in (3.8) can be computed as a strong limit in $\mathcal{L}(H^1, H^{-1})$,

$$DB = \underset{t \to 0}{\text{s-lim}} t^{-1} (H e^{itB} - e^{itB} H)$$

= $r^{-1/2} (DA - 2B^2) r^{-1/2} + r^{-3} v$
= $\sum_{|\alpha| \le 2} r^{-1} v_{\alpha} p^{\alpha}; \quad v_{\alpha} \in \mathcal{F}.$ (3.11)

The assertions (3.10) and (3.11) are standard results, which follow from mapping properties of the involved groups and the fact that the formal commutators are bounded forms on H^1 .

We will need modifications of A and B in terms of a parameter κ . This parameter is needed to control certain multiple commutators. Let

$$B_{\kappa,R} = B(\kappa^2 B^2 + 1)^{-1}, \quad A = A_{\kappa,R} = r^{1/2} B_{\kappa,R} r^{1/2}; \quad \kappa \in [0,1].$$
(3.12)

We may consider H defined in (3.1a) as a 'generalized' N-body Schrödinger operator. The set of thresholds of \tilde{H} , say denoted by $\tilde{\mathcal{T}}$, coincides with the set \mathcal{T} of thresholds of H except for having one less point. This exception follows from the identity

$$\sigma_{\rm pp}(\widetilde{H}^{a_0}) = \left(\sigma_{\rm pp}(H^{a_0}) \setminus \{\lambda_0\}\right) \cup \{0\}; \quad \widetilde{H}^{a_0} := H^{a_0} - \lambda_0 \Pi.$$

Note in particular that it follows that $\lambda_0 \notin \widetilde{\mathcal{T}}$. On the other hand there is no simple relationship between the eigenvalues of \widetilde{H} and those of H. A similar 'subtraction' of a genuine eigenprojection P of H was employed in [AHS] in which case indeed a similar relation between $\sigma_{pp}(H - \lambda_0 P)$ and $\sigma_{pp}(H)$ hold. Let in the following $\widetilde{d} : \mathbb{R} \to \mathbb{R}$ refer to the (lower) distance function defined by replacing $\mathcal{T} \to \widetilde{\mathcal{T}}$ in (3.5). Note that $\widetilde{d}(\lambda_0) > 0$.

Proposition 3.3. Suppose Conditions 1.5 and 2.4. Let $A_{\kappa,R}$ and \check{H} be given by (3.12) and (3.1a), respectively.

For all $\lambda \in \mathbb{R}$ and $\epsilon > 0$ there exist $R_0 \ge 1$ and $\kappa_0 \in (0, 1]$ such for all $R \ge R_0$, there exist a neighbourhood \mathcal{U} of λ and a compact operator K on L^2 :

$$f(\check{H})i[\check{H}, A_{\kappa,R}]f(\check{H}) \\ \geq f(\check{H})\{2\tilde{d}(\lambda) - \epsilon - K\}f(\check{H}) \text{ for all } \kappa \in [0, \kappa_0] \text{ and real } f \in C^{\infty}_{c}(\mathcal{U}).$$

$$(3.13)$$

The meaning of the appearing commutator will be explained in Subsection 3.1.1. However for $\kappa = 0$ there is an alternative interpretation to be elaborated on now. We claim that we can use (3.13) at $\lambda = \lambda_0$ and for $\kappa = 0$ to conclude, that there are at most a finite number of eigenvalues for H' in a neighbourhood of λ_0 . To see this it suffices to show that the commutator in (3.13), interpretated as the formal commutator, can be computed as a strong limit

$$\mathbf{i}[\breve{H}, A_R] = \operatorname{s-lim}_{t \to 0} t^{-1} \left(\breve{H} \mathrm{e}^{\mathrm{i}tA_R} - \mathrm{e}^{\mathrm{i}tA_R} \breve{H} \right) \in \mathcal{L}(H^2, H^{-2}).$$
(3.14)

Writing $\check{H} = H - T$, $T := \lambda_0 \Pi + \check{K}$, the part of (3.14) related to H is justified by (3.10). For the part related to the second term it suffices to show that the form, say a priori defined on $C_c^{\infty}(\mathbf{X})$, extends as follows.

Lemma 3.4. The form $\operatorname{iad}_{A_R}(T) := \operatorname{i}[T, A_R]$ extends to a bounded form on H^2 . More generally $\operatorname{iad}_{A_R}(T) \in \mathcal{L}(H^2, H^{-1}) \cap \mathcal{L}(H^1, H^{-2})$.

Proof. Write the form as

$$i[\lambda_0\Pi + \Pi I_0 + I_0\Pi - \Pi I_0\Pi, A_R] = T_1 + \dots + T_4.$$

We know that $[I_0, A_R] \in \mathcal{F} \subset \mathcal{L}(L^2)$. Let us only consider the contribution $[\Pi, A_R]I_0$ (from T_2) and only show that $[\Pi, A_R]I_0 \in \mathcal{L}(H^2, H^{-1})$. For that it suffices to show that $[\Pi, A_R] \in \mathcal{L}(L^2, H^{-1})$.

Let $1 = \chi_1 + \chi_2$ be a partition of unity on **X** given as follows. We demand that for some (small) $\sigma > 0$ and all $\check{R} \ge 1$:

(1)
$$\chi_1 \in C^{\infty}(\mathbf{X})$$
 and for all $\alpha \in \mathbb{N}_0^{\dim \mathbf{X}}$ there exists $C \in \mathbb{R}$:

$$|\langle x \rangle^{|\alpha|} \partial_x^{\alpha} \chi_1(x)| \le C.$$

(2)
$$\chi_1(x) = 1$$
 for $x \in \{ |x| \ge 2\check{R}, |x^{a_0}| \le \sigma |x| \}$ and
 $\operatorname{supp} \chi_1 \subset \{ |x| > \check{R}, |x^{a_0}| < 2\sigma |x| \} \subset \mathbf{X} \setminus \bigcup_{b \not \subset a_0} \mathbf{X}_b.$

Note that (2) implies, referring here to notation of (3.3) where $\delta = \delta(\sigma) > 0$ is taken sufficiently small (independently of \check{R}),

$$\operatorname{supp} \chi_1 \subset \mathbf{Y}_{a_0,\delta,\check{R}}.\tag{3.15}$$

We write

$$i[\Pi, A_R] = (\chi_1 + \chi_2)i[\Pi, A_R](\chi_1 + \chi_2) = \chi_1 i[\Pi, A_R]\chi_1 + S.$$
(3.16)

Now for all sufficiently big values of \check{R} (viz. $\check{R} \geq R\tilde{R} = R\tilde{R}(\delta)$) we obtain by combining (3.4) and (3.15) that

$$\tilde{\omega}_R(x) = x_{a_0} + \tilde{\omega}_R^{a_0}(x^{a_0}) \text{ for all } x \in \operatorname{supp} \chi_1.$$
(3.17)

The above construction can be done in an explicit way (including an explicit dependence of parameter R): Let us here and henceforth fix $\check{R} = R\tilde{R}(\delta)$ and choose $\chi_1(x) = \bar{\chi}_{\check{R}}(|x|)\Theta(x/|x|)$ with $\bar{\chi}_{\check{R}}$ specified in (1.29) and for a suitable real-valued smooth function Θ . We record the following improvement of (1).

(1)' For all $\alpha \in \mathbb{N}_0^{\dim \mathbf{X}}$ and $k \in \mathbb{N}$ there exists $C \in \mathbb{R}$ such that for all $R \geq 1$:

$$\left|\langle x \rangle^{|\alpha|} \partial_x^{\alpha} \chi_1(x)\right| + \left|\langle x \rangle^{|\alpha|} \partial_x^{\alpha} \left(\frac{\langle x \rangle^k}{\langle x^{a_0} \rangle^k} \bar{\chi}_{2\breve{R}}(|x|) \chi_2(x)\right)\right| \le C.$$

Since the operator A_R is a local operator the first term in (3.16) simplifies due to (3.17) as

$$\chi_1 \mathbf{i}[\Pi, A_R] \chi_1 = \chi_1 \mathbf{i}[\Pi, A_R^{a_0}] \chi_1 = \mathbf{i} \chi_1 \left(\Pi A_R^{a_0} - A_R^{a_0} \Pi \right) \chi_1 \in \mathcal{L}(L^2).$$
(3.18)

Now let us look at the term S in (3.16). It is given by

$$S = \chi_2 i[\Pi, A_R] + \chi_1 i[\Pi, A_R] \chi_2.$$
(3.19)

The two terms are treated similarly, so let us only elaborate on the first term. We write

$$A_R = r_R B_R - \frac{1}{2} |(\nabla r)(x/R)|^2,$$

and then

$$\chi_{2}[\Pi, A_{R}] = \chi_{2}\Pi A_{R} - \chi_{2}A_{R}\Pi$$

$$= K_{1}B_{R} - B_{R}K_{2} - iK_{3};$$

$$K_{1} = \chi_{2}\Pi r_{R},$$

$$K_{2} = r_{R}\chi_{2}\Pi,$$

$$K_{3} = \frac{1}{2}\chi_{2}\Pi |(\nabla r)(x/R)|^{2} + \frac{1}{2}\chi_{2}|(\nabla r)(x/R)|^{2}\Pi + (\tilde{\omega}_{R} \cdot \nabla \chi_{2})\Pi.$$
(3.20)

Noting, cf. (1)', that

$$K_1, K_2 \in \mathcal{L}(H^{-1}) \cap \mathcal{L}(L^2), K_3 \in \mathcal{L}(L^2) \text{ and } B_R \in \mathcal{L}(L^2, H^{-1}),$$
 (3.21)

we are done.

Remark 3.5. It follows from the technique of the proof that the second order commutator $\operatorname{ad}_{A_R}^2(T) \in \mathcal{L}(H^2, H^{-2})$ (and possibly no better for singular potentials). This suffices for the limiting absorption principle at λ_0 , cf. [PSS]. However higher commutators do not exist and we need refined micro-local estimates, which usually require multiple commutators. (In particular we need bounds with weights in position space.) Using the κ -distortions of A_R and B_R will allow us to treat multiple commutators. Note for example that for $\kappa > 0$ the above proof yields, at least formally, the improved result $\operatorname{ad}_{A_{\kappa,R}}(T) \in \mathcal{L}(H^1, H^{-1})$. We introduce a calculus of the κ -distortion of B which will be a major object to study. This is done in Section 3.2, and we give a number of applications of Proposition 3.3 and this calculus in Section 3.3.

3.1.1 Proof of Proposition 3.3

Introduce for $\varepsilon > 0$ the operators

$$X_{\varepsilon} = X_{\varepsilon,R} = r/(1+\varepsilon r), \quad A_{\varepsilon,\kappa,R} = X_{\varepsilon}^{1/2} B_{\kappa,R} X_{\varepsilon}^{1/2}; \quad \kappa \in [0,1].$$
(3.22)

Note that $A_{\varepsilon,\kappa,R} \in \mathcal{L}(H^2, H^1)$. Whence $i[H, A_{\varepsilon,\kappa,R}]$ is a well-defined form on H^2 . We *define* the commutators $i[H, A_{\kappa,R}]$, $i[\tilde{H}, A_{\kappa,R}]$ and $i[\check{H}, A_{\kappa,R}]$ as strong weak-star limits in $\mathcal{L}(H^2, H^{-2})$ in the following way. For technical reasons to be explained after the definitions, this works for small enough values of κ only. Let for each of the operators $H^{\#} = H$, \tilde{H} or \check{H}

$$i[H^{\#}, A_{\kappa,R}] = \operatorname{s-w}^{*}\operatorname{-lim}_{\varepsilon \to 0_{+}} i[H^{\#}, A_{\varepsilon,\kappa,R}].$$
(3.23)

It is easy to see that for $\kappa = 0$, these limiting forms exist and coincide with our previous computations, cf. (3.14) and Lemma 3.4. The case of $\kappa > 0$ requires an elaboration to be given in Lemma 3.6.

3.1.1. Proof of Proposition 3.3

An immediate virtue of (3.23) is the validity of the virial theorem. Thus for example, (3.23) combined with Proposition 3.3 yields that eigenvalues of \check{H} cannot accumulate at λ_0 . Note that this assertion does not require $\lambda_0 = \Sigma_2$ for which case in fact $\lambda_0 \notin \sigma_{\text{ess}}(\check{H})$, cf. [Wa2, Lemma 2.1] or the proof of Lemma 2.10. Proposition 3.3 will be a crucial tool for us only for $\lambda_0 > \Sigma_2$. At the level of proofs, the reader will see similarities of the proofs of Lemma 2.10 and Proposition 3.3.

We are going to use that for $\kappa > 0$

$$B_{\kappa,R} = \frac{1}{2\kappa} \left((\kappa B + i)^{-1} + (\kappa B - i)^{-1} \right).$$
(3.24)

Note that it follows from Mourre theory, [Mo], that $(\kappa B \pm i)^{-1}$ preserve H^2 provided κ is small enough. We need a uniform bound in κ and R. Thus, more precisely, we claim that there exists $\kappa'_0 \in (0, 1]$ such that for all $\kappa \in [0, \kappa'_0]$ and all $R \ge 1$ the space H^2 is preserved, and in fact

$$\sup_{R \ge 1, \kappa \le \kappa'_0} \| (\kappa B \pm i)^{-1} \|_{\mathcal{L}(H^2)} < \infty.$$
(3.25a)

We may take this constant as $\kappa'_0 = \min\{(2C)^{-1}, 1\}$, where

$$C = \sup_{R \ge 1} \| [H_0, B] (H_0 + 1)^{-1} \|_{\mathcal{L}(L^2)} < \infty.$$
(3.25b)

Note that the *R*-dependence of *B* is through $\omega_R(x) = (\nabla r)(x/R)$, which is bounded along with all derivatives. Whence the above finiteness claims hold. We also note that for any $R \ge 1$

$$\operatorname{s-lim}_{\kappa \to 0} (\kappa B \pm i)^{-1} = \mp i \text{ in } \mathcal{L}(H^2).$$
(3.25c)

Clearly there are similar properties as (3.25a) and (3.25c) with H^2 replaced by L_s^2 for any $s \ge 0$. For (3.25a) in that case, κ'_0 can be an arbitrary positive number (it does not need to be small); see the beginning of the proof of Lemma 3.9.

Lemma 3.6. The limits $i[H^{\#}, A_{\kappa,R}]$, with $H^{\#} = H$, \tilde{H} or \check{H} , are well-defined bounded form on H^2 for $\kappa \in [0, \kappa'_0]$. More generally $i[H^{\#}, A_{\kappa,R}] \in \mathcal{L}(H^2, H^{-1}) \cap \mathcal{L}(H^1, H^{-2})$.

Moreover for each of the above three forms

$$\exists C > 0 \,\forall R \ge 1 \,\forall \kappa \in [0, \kappa_0']: \quad \left\| i[H^\#, A_{\kappa, R}] - i[H^\#, A_R] \right\|_{\mathcal{L}(H^2, H^{-2})} \le C\kappa. \tag{3.26}$$

Proof. We need to examine the case $\kappa > 0$. We start by examining the quantity $i[H, A_{\kappa,R}]$.

We calculate using (3.11), (3.24) and (3.25a)

$$\begin{split} &\mathbf{i} \Big[H, A_{\varepsilon,\kappa,R} \Big] \\ &= \mathbf{i} X_{\varepsilon}^{1/2} \Big[H, B_{\kappa,R} \Big] X_{\varepsilon}^{1/2} + 2 \operatorname{Re} \left(\mathbf{i} \Big[H, X_{\varepsilon}^{1/2} \Big] B_{\kappa,R} X_{\varepsilon}^{1/2} \right) \\ &= -\frac{1}{2} \sum_{\pm} X_{\varepsilon}^{1/2} (\kappa B \pm \mathbf{i})^{-1} \mathbf{i} \Big[H, B \Big] (\kappa B \pm \mathbf{i})^{-1} X_{\varepsilon}^{1/2} + 2 \operatorname{Re} \left(\operatorname{Re} \left(X_{\varepsilon}^{-1/2} (\nabla X_{\varepsilon}) \cdot p \right) B_{\kappa,R} X_{\varepsilon}^{1/2} \right) \\ &= -\frac{1}{2} \sum_{|\alpha| \le 2, \pm} X_{\varepsilon}^{1/2} (\kappa B \pm \mathbf{i})^{-1} r^{-1} v_{\alpha} p^{\alpha} (\kappa B \pm \mathbf{i})^{-1} X_{\varepsilon}^{1/2} \\ &+ 2 \operatorname{Re} \left(r^{-1/4} (1 + \varepsilon r)^{-3/4} B_{R} (1 + \varepsilon r)^{-3/4} r^{-1/4} B_{\kappa,R} X_{\varepsilon}^{1/2} \right); \quad v_{\alpha} \in \mathcal{F}. \end{split}$$

We can let $\varepsilon \to 0$, yielding the existence of the desired limit in the H^2 -form sense. The result is

$$i \Big[H, A_{\kappa,R} \Big] = -\frac{1}{2} \sum_{|\alpha| \le 2, \pm} r^{1/2} (\kappa B \pm i)^{-1} r^{-1} v_{\alpha} p^{\alpha} (\kappa B \pm i)^{-1} r^{1/2} + 2 \operatorname{Re} \left(r^{-1/4} B_R r^{-1/4} B_{\kappa,R} r^{1/2} \right); \quad v_{\alpha} \in \mathcal{F}.$$

This expression is obviously in $\mathcal{L}(H^2, H^{-1}) \cap \mathcal{L}(H^1, H^{-2})$ (in fact in $\mathcal{L}(H^1, H^{-1})$ in this case).

Next we move multiplication operators to the middle as to become sandwiched by $(\kappa B \pm i)^{-1}$. Thereby we pick up errors of order $\mathcal{O}(\kappa)$ in $\mathcal{L}(H^2, H^{-2})$ uniformly in $R \geq 1$, cf. (3.25a) and the remark after (3.25c). This means that we only have to consider

$$-\frac{1}{2}\sum_{\pm} (\kappa B \pm i)^{-1} i [H, A] (\kappa B \pm i)^{-1}, \qquad (3.27)$$

where i[H, A] is given by (3.9). But

$$(\kappa B \pm i)^{-1} \pm i = \pm i\kappa(\kappa B \pm i)^{-1}B,$$
 (3.28)

which give an extra error $\mathcal{O}(\kappa)$ in $\mathcal{L}(H^2, H^{-2})$ when removing the factors $(\kappa B \pm i)^{-1}$ in (3.27) one by one. This proves (3.26) in the case $H^{\#} = H$.

For the commutators $i[\check{H}, A_{\kappa,R}]$ and $i[\check{H}, A_{\kappa,R}]$ we proceed similarly and by using in addition the proof of Lemma 3.4. Note that in the case of $H^{\#} = \check{H}$ the topology in (3.26) is the strongest Sobolev space topology we can use for singular potentials by this method, cf. Remark 3.5.

Lemma 3.7. For all $\lambda \in \mathbb{R}$ and $\epsilon > 0$ there exist $R_0 \ge 1$ such for all $R \ge R_0$, there exist a neighbourhood \mathcal{V} of λ and a compact operator \widetilde{K} on L^2 :

$$f(\tilde{H})i[\tilde{H}, A_R]f(\tilde{H}) \\ \ge f(\tilde{H})\{2\tilde{d}(\lambda) - \epsilon - \tilde{K}\}f(\tilde{H}) \text{ for all real } f \in C^{\infty}_{c}(\mathcal{V}).$$

$$(3.29)$$

Proof. We divide the proof into several steps.

<u>*I*</u>. We note that for any product $P_1 \cdots P_m$ of factors P_l given either as $P_l = \langle x^{a_0} \rangle$ or $P_l = A_R^{a_0}$ we have

$$P_1 \cdots P_m \Pi \in \mathcal{L} \left(L^2(\mathbf{X}^{a_0}) \right). \tag{3.30}$$

In fact since $\langle x^{a_0} \rangle^m (H^{a_0} + i)^m \Pi \in \mathcal{L}(L^2(\mathbf{X}^{a_0}))$ the result follows from the property

$$P_1 \cdots P_m (H^{a_0} + \mathbf{i})^{-m} \langle x^{a_0} \rangle^{-m} \in \mathcal{L} (L^2(\mathbf{X}^{a_0})),$$

which in turn follows from repeated commutation and the property $A_R^{a_0} \langle x^{a_0} \rangle^{-1} (H^{a_0} + i)^{-1} \in \mathcal{L}(L^2(\mathbf{X}^{a_0})).$

<u>II</u>. There is a complete analogue of Lemma 3.2 with the triple $(H, i[H, A_R], d)$ replaced by $(\tilde{H}, i[H, A_R], \tilde{d})$. It reads conveniently as follows.

For all $\lambda \in \mathbb{R}$ and $\epsilon > 0$ there exists $R_0 \ge 1$ such for all $R \ge R_0$, there exist a bounded neighbourhood \mathcal{V} of λ and a compact operator \widetilde{K}_1 on L^2 :

$$f(\widetilde{H})i[H, A_R]f(\widetilde{H}) \geq f(\widetilde{H})\{2\widetilde{d}(\lambda) - \epsilon/2 - \widetilde{K}_1\}f(\widetilde{H}) \text{ for all real } f \in C^{\infty}_{c}(\mathcal{V}).$$

$$(3.31)$$

This statement follows by mimicking the proofs of [Sk3, Corollaries 4.2 and 4.5]. One comment is due since the 'potential' Π is not local: Expanding the operator $T(\delta')$ appearing in [Sk3, (4.16)] one 'new term' is given by $T := [\Pi, j_b]$; here j_b is a partition function with similar properties as the ones in (2.37a)–(2.37c). We need to show that T is \tilde{H} -compact. If $a_0 \not\subset b$ indeed Πj_b and $j_b \Pi$, and hence also T, are H-compact, cf. Step I. If on the other hand $a_0 \subset b$ we have $a_0 = b$ and we can write $T = -\sum_{d \neq b} [\Pi, j_d]$. Since for any such term we have $a_0 \not\subset d$ the previous argument yields that $[\Pi, j_d]$ is H-compact.

<u>*III*</u>. Recalling the definition (3.1a) it remains to show the following estimate in terms of the set $\mathcal{V} = \mathcal{V}(R)$ from (3.31).

There exists a compact operator $\tilde{K}_2 = \tilde{K}_2(R)$ such that

$$1_{\mathcal{V}}(\widetilde{H})i[-\lambda_0\Pi, A_R]1_{\mathcal{V}}(\widetilde{H}) \ge -\epsilon/2 - \widetilde{K}_2.$$
(3.32)

Clearly the combination of (3.31) and (3.32) yields (3.29) with this neighbourhood \mathcal{V} and with $\tilde{K} = \tilde{K}_1 + \tilde{K}_2$. To show (3.32) we decompose as in the proof of Lemma 3.4 writing

$$i[\Pi, A] = (\chi_1 + \chi_2)i[\Pi, A](\chi_1 + \chi_2) = \chi_1 i[\Pi, A_R^{a_0}]\chi_1 + S.$$

Since

$$\|\Pi A_R^{a_0} - A_R^{a_0}\Pi\| \to 0 \text{ for } R \to \infty,$$

cf. Lemma 3.1(2) and Step I, the first term conforms with (3.32).

It remains to consider S, which is given by (3.19). We decompose as in (3.20) (treating again only the first term of S). The operators K_1 , K_2 and K_3 are not only bounded (as stated in (3.21)) but also $\langle p \rangle$ -compact. Since \mathcal{V} is bounded (3.32) follows.

Proof of Proposition 3.3. Let $\lambda \in \mathbb{R}$ and $\epsilon > 0$ be given. We need to specify $R_0 \geq 1$, $\kappa_0 \in (0, \kappa'_0]$ and for $R \geq R_0$ a neighbourhood $\mathcal{U}(R)$ of λ and a compact K(R) such that (3.13) holds.

<u>I</u>. First we use Lemma 3.7 to obtain the statement (3.29) with the factors of $f(\tilde{H})$ replaced by the factors of $f(\tilde{H})$. More precisely we use the statement with ϵ replaced by $\epsilon/2$ allowing us to pick corresponding $R_0 \geq 1$ and for $R \geq R_0$ a neighbourhood $\mathcal{V}(R)$ of λ and a compact $\tilde{K}(R)$ such that (3.29) holds. Now take for any such set $\mathcal{V}(R)$ any smaller neighbourhood of λ , denoted by $\mathcal{U}(R)$, with compact closure contained in the interior of $\mathcal{V}(R)$. We can pick a real function $\tilde{f} \in C_c^{\infty}(\mathcal{V}(R))$ such that $\tilde{f} = 1$ on $\mathcal{U}(R)$ and then estimate as follows. Note that

 $\breve{K}_0 = \breve{K}_0(R) := (H_0 + 1) (\tilde{f}(\breve{H}) - \tilde{f}(\widetilde{H}))$ is compact (cf. (3.37) given below) and that $i[\widetilde{H}, A_R] \in \mathcal{L}(H^2, H^{-2})$ (cf. Lemma 3.4). Writing then

$$\begin{aligned}
\breve{K}_1 &:= \widetilde{f}(\breve{H}) \mathrm{i}[\widetilde{H}, A_R] \widetilde{f}(\breve{H}) - \widetilde{f}(\widetilde{H}) \mathrm{i}[\widetilde{H}, A_R] \widetilde{f}(\widetilde{H}) \\
&= \breve{K}_0^* \breve{T} + \widetilde{T}^* \breve{K}_0; \\
\breve{T} &= \left(H_0 + 1\right)^{-1} \mathrm{i}[\widetilde{H}, A_R] \widetilde{f}(\breve{H}), \\
&\widetilde{T} &= \left(H_0 + 1\right)^{-1} \mathrm{i}[\widetilde{H}, A_R] \widetilde{f}(\breve{H}),
\end{aligned}$$

we conclude that \breve{K}_1 is compact. Note also that

$$\breve{K}_2(R) := \left(2\tilde{d}(\lambda) - \epsilon/2\right) \left(\tilde{f}(\breve{H})^2 - \tilde{f}(\widetilde{H})^2\right)$$

is compact.

In conclusion we take $R_0 \ge 1$, $\mathcal{U}(R)$ as described above and

$$\breve{K}_3(R) := \tilde{f}(\widetilde{H})\widetilde{K}\tilde{f}(\widetilde{H}) - \breve{K}_1 + \breve{K}_2; \quad R \ge R_0,$$

yielding

$$f(\breve{H})i[\widetilde{H}, A_R]f(\breve{H})$$

$$\geq f(\breve{H})\{2\tilde{d}(\lambda) - \epsilon/2 - \breve{K}_3\}f(\breve{H}) \text{ for all real } f \in C^{\infty}_{c}(\mathcal{U}(R)).$$

 \underline{II} . We note that

$$\check{K}_4(R) := 1_{U(R)}(\check{H})i[\check{K}, A_R]1_{U(R)}(\check{H})$$
 is compact.

We subtract these terms in the previous estimate, yielding

$$f(\breve{H})\mathbf{i}[\breve{H}, A_R]f(\breve{H})$$

$$\geq f(\breve{H})\{2\tilde{d}(\lambda) - \epsilon/2 - K\}f(\breve{H}) \text{ for all real } f \in C^{\infty}_{\mathbf{c}}(\mathcal{U}(R)),$$

where $K = K(R) = \breve{K}_3 + \breve{K}_4$.

<u>III</u>. We invoke (3.26), estimating

$$1_{U(R)}(\breve{H})(\mathrm{i}[\breve{H}, A_{\kappa, R}] - \mathrm{i}[\breve{H}, A_R])1_{U(R)}(\breve{H}) \ge -\epsilon/2,$$

for all small enough κ . In combination with the previous estimate this estimate yields (3.13).

3.2 Multiple commutators and calculus

It is convenient to introduce a concept of 'order' for certain classes of linear operators T on L^2 , cf. [GIS]. It is based on the following elementary result, cf. Lemma 3.9.

$$\forall \kappa \in (0,1], \ \forall R \ge 1: \quad B_{\kappa,R} L^2_{\infty} \subset L^2_{\infty}.$$
(3.33)

For operators S, T on L^2 we define (formally) multiple commutators by

$$\operatorname{ad}_{S}^{0}(T) = T$$
 and $\operatorname{ad}_{S}^{k}(T) = [\operatorname{ad}_{S}^{k-1}(T), S]$ for $k \in \mathbb{N}$.

Let X be multiplication by $r = r_R$ on L^2 , and recall that $\kappa'_0 \in (0, 1]$ is introduced in the discussion of (3.25a) and (3.25b). **Definition 3.8.** Let $R \ge 1$ be given. An operator T on L^2 is of κ -order $t \in \mathbb{R}$ if

- (1) $\mathcal{D}(T), \mathcal{D}(T^*) \supset L^2_{\infty},$
- (2) T and T^* leave L^2_{∞} invariant,
- (3) $\forall \kappa \in (0, \kappa'_0] \forall s \in \mathbb{R} \quad \forall k \in \mathbb{N}_0 : \quad X^{s-k-t} \mathrm{ad}^k_{B_{\kappa,R}}(T) X^{-s} \in \mathcal{L}(L^2).$

For any operator T of κ -order t the adjoint T^* also has κ -order t, and we write $T, T^* = \mathcal{O}_{\kappa}(X^t)$. The class of such operators is denoted by $\mathfrak{O}_{\kappa}(X^t)$. It is readily seen that $X^t = \mathcal{O}_{\kappa}(X^t)$, cf. (3.39) and (3.40) stated below. This is related to the fact that $\mathrm{ad}_B^k(X^k) \in \mathcal{F}$ for $k \in \mathbb{N}$, where the algebra \mathcal{F} is introduced in the discussion of (3.8). By the Leipniz rule, if S and T are of κ -order s and t, respectively, then ST is of order s + t. We do not keep track of neither the κ - nor the R-dependence, however it is important for our applications of the above concept of 'order' that $\kappa = 0$ is not included in (3). We abbreviate $\check{R}(z) = (\check{H} - z)^{-1}$ and $R_{\kappa}(z) = (B_{\kappa} - z)^{-1} = (B_{\kappa,R} - z)^{-1}$.

Lemma 3.9. Let $R \ge 1$ be given big enough. Then for any $f \in C_c^{\infty}(\mathbb{R})$, $g \in C^{\infty}(\mathbb{R})$, $v \in \mathcal{F}$, $|\alpha| \le 2$, $z \in \mathbb{C} \setminus \mathbb{R}$ and polynomials P_1, P_2 in A^{a_0} and the components of x^{a_0} and p^{a_0} , such that the total number of components of p^{a_0} is at most two for P_1 as well as for P_2 , the operators

$$g(B_{\kappa}), v, P_{1}\Pi P_{2}, I_{0}(p^{2}+1)^{-1}, p^{\alpha}\breve{R}(z), I_{0}\Pi\breve{R}(z), f(\breve{H}) \in \mathfrak{O}_{\kappa}(X^{0}).$$
 (3.34)

Proof. We prove the bounds one by one. For simplicity of presentation the above list is not stated as complete as possible. In the proof we will see that a certain algebra $\breve{\mathcal{B}} \subset \mathfrak{O}_{\kappa}(X^0)$; this algebra contains a few operators not listed above for which the assertion that they are contained in $\mathfrak{O}_{\kappa}(X^0)$ will also be useful later.

By repeated commutation we obtain for the regularization X_{ε} of (3.22) that for any $s \ge 0$, $\varepsilon > 0$ and $z \in \mathbb{C} \setminus \mathbb{R}$

r 1

$$X_{\varepsilon}^{s}(B-z)^{-1} = \sum_{k=0}^{\lfloor s \rfloor} (-1)^{k} (B-z)^{-k-1} \mathrm{ad}_{B}^{k}(X_{\varepsilon}^{s}) + (-1)^{\lfloor s \rfloor+1} (B-z)^{-\lfloor s \rfloor-1} \mathrm{ad}_{B}^{\lfloor s \rfloor+1}(X_{\varepsilon}^{s}) (B-z)^{-1}.$$
(3.35)

We multiply by X^{-s} from the right and take $\varepsilon \to 0$ observing that the resulting right-hand side is explicitly in $\mathcal{L}(L^2)$. Note that the operator $\mathrm{ad}_B^{[s]+1}(X^s) \in \mathcal{F}$, and therefore it is a bounded multiplication operator. We can argue similarly for s < 0. In particular we conclude (3.33) (using (3.24)) as well as a boundedness result remarked before Lemma 3.6. In fact it follows that

$$(\kappa B \pm \mathbf{i})^{-1} \in \mathfrak{O}_{\kappa}(X^0). \tag{3.36}$$

 $\underline{g(B_{\kappa})}$; 1st proof. In terms of an almost analytic extension \tilde{g} of a given $g \in C_{c}^{\infty}(\mathbb{R})$ ('absorbed' in a measure $\mu = \mu_{g}$ and writing z = u + iv)

$$g(B) = \frac{1}{\pi} \int_{\mathbb{C}} \left(\bar{\partial} \tilde{g} \right)(z) (B-z)^{-1} \mathrm{d}u \mathrm{d}v = \int_{\mathbb{C}} (B-z)^{-1} \mathrm{d}\mu(z).$$
(3.37)

Let for any $t \in \mathbb{R}$

$$\mathcal{G}_t = \{ g \in C^{\infty}(\mathbb{R}) | \, \forall k \in \mathbb{N}_0 : \, |g^{(k)}(x)| \le C_k \langle x \rangle^{t-k} \}.$$
(3.38)

It is well-known that (3.37) is valid in fact for $g \in \mathcal{G}_t$ with t < 0 (for any self-adjoint operator B).

Now for any given $g \in C^{\infty}(\mathbb{R})$ we write $g(B_{\kappa}) = g_{\kappa}(B) + g(0)$ with $g_{\kappa} \in \mathcal{G}_{-1}$. By using (3.37) for g_{κ} we conclude by using (3.35) with $\varepsilon = 0$ (and its adjoint version) that $g(B_{\kappa}) = g_{\kappa}(B) + g(0) \in \mathfrak{O}_{\kappa}(X^0)$.

<u> $g(B_{\kappa})$; 2nd proof</u>. Since $B_{\kappa,R}$ is bounded we can write $g(B_{\kappa,R}) = f(B_{\kappa,R})$ for some $\overline{f \in C_c^{\infty}(\mathbb{R})}$ and then apply (3.37). For $s \ge 0$ (treating only this case in (3)) we expand as in (3.35)

$$X^{s}R_{\kappa}(z) = \sum_{k=0}^{[s]} (-1)^{k}R_{\kappa}(z)^{k+1} \mathrm{ad}_{B_{\kappa}}^{k}(X^{s}) + (-1)^{[s]+1}R_{\kappa}(z)^{[s]+1} \mathrm{ad}_{B_{\kappa}}^{[s]+1}(X^{s})R_{\kappa}(z).$$

It suffices to show that

$$\operatorname{ad}_{B_{\kappa}}^{k}(X^{s})X^{k-s}, k = 0, \dots, [s], \text{ and } \operatorname{ad}_{B_{\kappa}}^{[s]+1}(X^{s}) \text{ are bounded.}$$
 (3.39)

Using (3.24) and (3.36) we compute for k = 1, ..., [s] + 1

$$\operatorname{ad}_{B_{\kappa}}^{k}(X^{s}) = \sum_{\sigma = (\sigma_{1}, \dots, \sigma_{k}) \in \{-1, 1\}^{k}} T_{\sigma} v_{\sigma} X^{s-k} T_{\sigma};$$

$$T_{\sigma} = \prod_{j=1}^{k} (\kappa B + \sigma_{j} i)^{-1}, \quad v_{\sigma} \in \mathcal{F}.$$
(3.40)

Clearly this shows that $\operatorname{ad}_{B_{\kappa}}^{[s]+1}(X^s)$ is bounded. For $k \leq [s]$ we obtain the first part of (3.39) by using that $X^{s-k}T_{\sigma}X^{k-s}$ is bounded, cf. (3.36). Consequently we are done.

 \underline{v} . We compute similarly (using the same notation)

$$\mathrm{ad}_{B_{\kappa}}^{k}(v) = \sum_{\sigma \in \{-1,1\}^{k}} T_{\sigma} v_{\sigma} X^{-k} T_{\sigma}.$$

We conclude that $v \in \mathfrak{O}_{\kappa}(X^0)$ using again (3.36).

 $P_1 \Pi P_2$. We let T be of the form $T = P_1 \Pi P_2$ and compute

$$\begin{aligned} \operatorname{ad}_{B_{\kappa}}(T) &= -\sum_{\sigma \in \{-1,1\}} (\kappa B + \sigma i)^{-1} [T, B] (\kappa B + \sigma i)^{-1}, \\ \operatorname{i}[T, B] &= r^{-1/2} \operatorname{i}[T, A] r^{-1/2} + 2 \operatorname{Re} \left(\operatorname{i} \left[T, r^{-1/2} \right] A r^{-1/2} \right) \\ &= \operatorname{i} r^{-1/2} \chi_1 \left(T A^{a_0} - A^{a_0} T \right) \chi_1 r^{-1/2} + T_1 + T_2 + T_3; \\ T_1 &= r^{-1/2} \chi_2 \operatorname{i}[T, r^{1/2} B r^{1/2}] r^{-1/2}, \\ T_2 &= r^{-1/2} \chi_1 \operatorname{i}[T, r^{1/2} B r^{1/2}] \chi_2 r^{-1/2}, \\ T_3 &= 2 \operatorname{Re} \left(\operatorname{i} \left[T, r^{-1/2} \right] r^{1/2} B \right). \end{aligned}$$

3.2. Multiple commutators and calculus

We need to compute higher order commutators as well, so we need to iterate this computation. We will demonstrate a self-similar structure which will allow us to control higher order commutators. To this end we introduce the graded algebra $\mathcal{B} = \sum_{k \in \mathbb{N}_0} \mathcal{B}_k$, where \mathcal{B}_k is given by linear combinations of any products of

$$X^{-s} \ (s \in \mathbb{R}), v \in \mathcal{F}, (\kappa B \pm i)^{-1}, T, P^{\alpha}_{\sigma}, (P^{\alpha}_{\sigma})^{*}; \quad P^{\alpha}_{\sigma} := p^{\alpha} T_{\sigma} (p^{2} + 1)^{-1} \ (|\alpha| \le 2).$$

Here T is any operator of the form $T = P_1 \Pi P_2$ and T_{σ} is given by the last equation of (3.40) for $k \in \mathbb{N}$ while $T_{\sigma} := I$ for k = 0, and for any such product the total sum of appearing exponents $s \in \mathbb{R}$ is given by k. We claim that

$$\forall S \in \mathcal{B}_k : \quad \mathrm{ad}_{B_k}(S) \in \mathcal{B}_{k+1}. \tag{3.41}$$

From this property it follows that $\operatorname{ad}_{B_{\kappa}}^{k}(S) \in \mathcal{B}_{k}$ for any $S \in \mathcal{B}_{0}$ (in particular for T), and we readily see that the powers of X can be redistributed as we want, i.e. we obtain that $\mathcal{B} \subset \mathcal{D}_{\kappa}(X^{0})$, in particular $T \in \mathcal{D}_{\kappa}(X^{0})$ follows.

To show (3.41) it suffices to consider k = 0 and then in turn only the operator $T = P_1 \Pi P_2 \in \mathcal{B}_0$. So we consider the above computation of the commutator with \mathcal{B}_{κ} . The first term is clearly in \mathcal{B}_1 since TA^{a_0} and $A^{a_0}T$ have the same form as T and the two outer factors of $r^{-1/2}$ together provides us with the extra factor X^{-1} required for the class \mathcal{B}_1 .

For the term T_1 and T_2 we also 'undo' the commutation. Whence

$$-iT_1 = r^{-1/2}\chi_2 Tr^{1/2} B - B\chi_2 r^{1/2} Tr^{-1/2} + [B,\chi_2]r^{1/2} Tr^{-1/2}$$

We use (3.28) to combine the factors of B with the factors of $(\kappa B + \sigma i)^{-1}$. The remaining terms can for any $s \in \mathbb{R}$ be written as $X^{-s}v_1 \check{T}v_2 X^{-s}$, where $v_1, v_2 \in \mathcal{F}$ and \check{T} has the same form as T (cf. (1)' in the proof of Lemma 3.4). In particular we get the extra factor X^{-1} by choosing s = 1/2. We can argue similarly for T_2 .

For the term T_3 we note the following general formula. Let $g \in \mathcal{G}_t$ for any $t \in \mathbb{R}$ and consider the commutation formula for the composition g(r),

$$i[\Pi, g(r)] = \Pi h(x) - h(x)\Pi,$$

$$h(x) = i \int_0^1 dt \, g'(r(tx^a + x_a)) \int_0^t x^a \cdot (\nabla^2 r) \big((sx^a + x_a) \big) x^a \, ds.$$
(3.42)

Here we used two zero'th order Taylor expansions, the fact that $[\Pi, g(r(x_a))] = 0$ and the property $\nabla r(x_a) \cdot x^a = 0$, cf. Lemma 3.1(2).

Applied to $g(r) = r^{-1/2}$ we conclude that

$$i[T, r^{-1/2}] = Th(x) - h(x)T, \qquad (3.43)$$

with h given by (3.42). We can write $h = \langle x^{a_0} \rangle^{7/2} r^{-3/2} v$, where $v \in \mathcal{F}$.

This representation can be refined using the formula

$$r\nabla^2 r = \nabla^2 r^2 / 2 - |\nabla r\rangle \langle \nabla r| \in \mathcal{F}.$$
(3.44)

In fact it follows from (3.42) and (3.44) that (3.43) holds with

$$h = \langle x^{a_0} \rangle^{9/2} r^{-5/2} v, \text{ where } v \in \mathcal{F}.$$
(3.45)

Thus intuitively the commutator is two inverse powers of X better. Obviously this is a general property for commutation by g(r) for $g \in \mathcal{G}_t$, and this comes in handy later, see the proof of Lemma 3.10. In any case by the appearance of at least one extra power of X^{-1} and (3.28) we see that also T_3 contributes by a term in \mathcal{B}_1 .

For an application in the next step of the proof let us note that

$$S_1 := (p^2 + 1)\Pi(p^2 + 1)^{-1} \in \mathcal{B}_0 \text{ and } S_2 := (p^2 + 1)[\Pi, B_\kappa](p^2 + 1)^{-1} \in \mathcal{B}_1.$$
(3.46)

Obviously $S_1 \in \mathcal{B}_0$, and therefore the second assertion follows from (3.41) and

$$S_2 = [S_1, B_\kappa] - [p^2, B_\kappa](p^2 + 1)^{-1}S_1 + S_1[p^2, B_\kappa](p^2 + 1)^{-1} \in \mathcal{B}_1.$$

 $\frac{I_0(p^2+1)^{-1} \text{ and } p^{\alpha}T_{\sigma_1,\ldots,\sigma_j}\breve{R}(z), \ p^{\alpha}T_{\sigma_1,\ldots,\sigma_j}\Pi\breve{R}(z); \ |\alpha| \leq 2, \ j \in \mathbb{N}_0.$ We add these operators as well as the adjoint expressions to the list of generators of algebra \mathcal{B} and denote the corresponding algebra by $\breve{\mathcal{B}} = \sum_{k \in \mathbb{N}_0} \breve{\mathcal{B}}_k$, where as before the total sum of appearing exponents $s \in \mathbb{R}$ in any term of an element of $\breve{\mathcal{B}}_k$ is given by k. We claim that

$$\forall S \in \breve{\mathcal{B}}_k : \quad \mathrm{ad}_{B_\kappa}(S) \in \breve{\mathcal{B}}_{k+1}. \tag{3.47}$$

From this property it follows that $\operatorname{ad}_{B_{\kappa}}^{k}(S) \in \check{\mathcal{B}}_{k}$ for any $S \in \check{\mathcal{B}}_{0}$ (in particular for the enlisted operators and their adjoint expressions), and again we can redistribute the powers of X as we want. Whence we obtain that $\check{\mathcal{B}} \subset \mathfrak{O}_{\kappa}(X^{0})$, in particular the fourth, fifth and sixth operators listed in (3.34) are all in $\mathfrak{O}_{\kappa}(X^{0})$ as claimed. In conclusion it suffices to show (3.47), and therefore in turn (3.47) with k = 0. As for the first term we note that $[I_{0}, B] \in X^{-1}\mathcal{F}$. Whence it suffices to consider the other terms. For the second term $S = p^{\alpha}T_{\sigma_{1},...,\sigma_{j}}\check{R}(z) = p^{\alpha}T_{\sigma}\check{R}(z)$ we compute

$$\begin{split} [S, B_{\kappa}] &= [p^{\alpha}, B_{\kappa}] T_{\sigma} \check{R}(z) - p^{\alpha} T_{\sigma} \check{R}(z) [\check{H}, B_{\kappa}] \check{R}(z) \\ &= T_1 + T_2 + T_3 + T_4; \\ T_1 &= -\sum_{\sigma' \in \{-1,1\}} (\kappa B + \sigma' i)^{-1} [p^{\alpha}, B] T_{\sigma_1, \dots, \sigma_j, \sigma'} \check{R}(z), \\ T_2 &= -p^{\alpha} T_{\sigma} \check{R}(z) [H, B_{\kappa}] \check{R}(z), \\ T_3 &= \lambda_0 p^{\alpha} T_{\sigma} \check{R}(z) [\Pi, B_{\kappa}] \check{R}(z), \\ T_4 &= p^{\alpha} T_{\sigma} \check{R}(z) [\check{K}, B_{\kappa}] \check{R}(z). \end{split}$$

Since $[p^{\alpha}, B] \in X^{-1} \sum_{|\beta| \leq 2} v_{\beta} p^{\beta}$ where $v_{\beta} \in \mathcal{F}$, the operator $T_1 \in \check{\mathcal{B}}_1$. Thanks to (3.11) we conclude that also $T_2 \in \check{\mathcal{B}}_1$. Since $\Pi \in \mathcal{B}_0$ and therefore $[\Pi, B_{\kappa}] \in \mathcal{B}_1$ (due to (3.41)) also $T_3 \in \check{\mathcal{B}}_1$. For T_4 we combine the facts that

$$I_0(p^2+1)^{-1}, (p^2+1)\breve{R}(z) \in \breve{\mathcal{B}}_0$$

and (3.46). Thus, for example (with S_2 given in (3.46))

$$[I_0\Pi, B_{\kappa}]\breve{R}(z) - [I_0, B_{\kappa}]\Pi\breve{R}(z) = (I_0(p^2 + 1)^{-1})S_2((p^2 + 1)\breve{R}(z)) \in \breve{\mathcal{B}}_1.$$

Clearly the second term to the left is in $\check{\mathcal{B}}_1$ too, completing our treatment of one of the three terms of \check{K} . This argument also works for the other terms of \check{K} .

Finally for the third term $S = p^{\alpha}T_{\sigma}\Pi \check{R}(z)$ we write

$$S = P^{\alpha}_{\sigma} S_1 \big((p^2 + 1) \breve{R}(z) \big).$$

Since $P_{\sigma}^{\alpha}, S_1 \in \mathcal{B}_0$, cf. (3.46), $[P_{\sigma}^{\alpha}S_1, B_{\kappa}] \in \mathcal{B}_1 \subset \breve{\mathcal{B}}_1$, and since $[(p^2+1)\breve{R}(z), B_{\kappa}] \in \breve{\mathcal{B}}_1$ (as proved above), it follows that indeed $[S, B_{\kappa}] \in \breve{\mathcal{B}}_1$ as we want.

 $f(\breve{H})$. We use (3.37) with *B* replaced by \breve{H} . We have proven that $R(z) = \mathcal{O}_{\kappa}(X^0)$ for any fixed non-real *z*. Whence we can compute any number of commutators with B_{κ} , redistribute powers of *X* as we want, and then estimate to see that indeed $f(\breve{H}) \in \mathcal{O}_{\kappa}(X^0)$. Note that the bounds $\|\breve{R}(z)^m\| \leq |\operatorname{Im} z|^{-m}$ come in naturally estimating integrals like (3.37) and in fact appear 'harmless' in combination with the appearing measure μ .

3.2.1 Computing a commutator

We are interested in computing commutators $i[\check{H}, P]$ where P in all relevant cases has the form

$$P = f(\check{H})h(r)g(B_{\kappa})h(r)f(\check{H}); \quad f \in C_{c}^{\infty}(\mathbb{R}), \ h, g \in C^{\infty}(\mathbb{R}), \ \text{real-valued}.$$

(We suppress the dependence of R.) Let $\tilde{f}(\lambda) = \lambda \tilde{f}(\lambda)$ for any real-valued $\tilde{f} \in C_c^{\infty}(\mathbb{R})$ chosen such that $\tilde{f} = 1$ on the support of f. Let $\check{H} = \check{f}(\check{H})$ and denote the corresponding Heisenberg derivative $i[\check{H}, \cdot]$ by $\check{\mathbf{D}}$, in particular $i[\check{H}, P] = \check{\mathbf{D}}P$. We are interested in specific choices of h and g, and for those choices P has a κ -order, say 2t. More precisely we impose the condition $h \in \mathcal{G}_t$, cf. (3.38), and conclude that the composed function h(r) has κ -order t, so that indeed P has a κ -order 2t (note that due to Lemma 3.9 the κ -order of $g(B_{\kappa})$ and \check{H} are zero). This implies that the commutator $\check{\mathbf{D}}P$ has order 2t, possibly (and indeed) smaller. We need to compute the order more exactly along with the leading order term.

Lemma 3.10. Let $R \geq 1$ be given big enough and suppose $h \in \mathcal{G}_t$. Then

$$\check{\mathbf{D}}P - L_1 - L_2 \in \mathfrak{O}_{\kappa}(X^{2t-2});$$

$$L_1 = 4f(\check{H}) \operatorname{Re}\left((\kappa^2 B^2 + 1)h'(r)B_{\kappa}g(B_{\kappa})h(r)\right)f(\check{H}), \qquad (3.48a)$$

$$L_2 = f(\check{H})h(r) \operatorname{Re}\left(g'(B_{\kappa})\check{\mathbf{D}}B_{\kappa}\right)h(r)f(\check{H}).$$

The operators $L_1, L_2 \in \mathfrak{O}_{\kappa}(X^{2t-1})$, so in particular $\check{\mathbf{D}} P \in \mathfrak{O}_{\kappa}(X^{2t-1})$.

Suppose in addition that $g' \ge 0$ and that $(g')^{1/2} \in C^{\infty}$. Then L_2 can be replaced by any of the following expressions $L_3, L_4, L_5 \in \mathfrak{O}_{\kappa}(X^{2t-1})$ for which $Com = f(\breve{H})i[\breve{H}, A_{\kappa}]f(\breve{H})$ is defined in agreement with Lemma 3.6.

$$L_{3} = h(r)\sqrt{g'(B_{\kappa})}f(\breve{H})i[\breve{H}, B_{\kappa}]f(\breve{H})\sqrt{g'(B_{\kappa})}h(r), \qquad (3.48b)$$
$$L_{4} = h(r)\sqrt{g'(B_{\kappa})}X^{-1/2}Com X^{-1/2}\sqrt{g'(B_{\kappa})}h(r)$$

$$L_{4} = h(r)\sqrt{g'(B_{\kappa})}X^{-1/2}Com X^{-1/2}\sqrt{g'(B_{\kappa})}h(r) -2h(r)\sqrt{g'(B_{\kappa})}X^{-1/2}f(\breve{H})(\kappa^{2}B^{2}+1)B_{\kappa}^{2}f(\breve{H})X^{-1/2}\sqrt{g'(B_{\kappa})}h(r),$$
(3.48c)

$$L_{5} = h(r)X^{-1/2}\sqrt{g'(B_{\kappa})}Com\sqrt{g'(B_{\kappa})}X^{-1/2}h(r) - 2h(r)X^{-1/2}\sqrt{g'(B_{\kappa})}f(\breve{H})(\kappa^{2}B^{2}+1)B_{\kappa}^{2}f(\breve{H})\sqrt{g'(B_{\kappa})}X^{-1/2}h(r).$$
(3.48d)

Proof. We write

$$\mathbf{\check{D}}P = L'_1 + L'_2;$$

$$L'_1 = 2f(\breve{H}) \operatorname{Re}\left(\left(\mathbf{\check{D}}h(r)\right)g(B_\kappa)h(r)\right)f(\breve{H}),$$

$$L'_2 = f(\breve{H})h(r)\left(\mathbf{\check{D}}g(B_\kappa)\right)h(r)f(\breve{H}).$$
(3.49)

By (3.37)

$$\widetilde{\mathbf{D}}h(r) = -\int_{\mathbb{C}} \breve{R}(z)\mathbf{i}[\breve{H}, h(r)]\breve{R}(z)\mathrm{d}\mu_{\breve{f}}(z); \qquad (3.50a)$$

$$\mathbf{i}[\breve{H}, h(r)] = 2\operatorname{Re}\left(h'(r)B\right) - \lambda_0\mathbf{i}[\Pi, h(r)] - \mathbf{i}[\breve{K}, h(r)].$$
(3.50b)

We insert (3.50b) into (3.50a) obtaining then three terms, say T_1, T_2, T_3 . We can write (seen by commutation)

$$T_1 = 2\check{f}'(\check{H})h'(r)B + O_{\kappa}(X^{t-2}) \in \mathfrak{O}_{\kappa}(X^{t-1}).$$

The terms T_2 and T_3 are treated using (3.42) and (3.44) and Lemma 3.9 (cf. the proof of discussion after (3.44)) to obtain

$$T_2, T_3 \in \mathfrak{O}_{\kappa}(X^{t-2}).$$

Next we insert the resulting formula

$$\check{\mathbf{D}}h = 2\check{f}'(\check{H})h'(r)B + \mathcal{O}_{\kappa}(X^{t-2}) \in \mathfrak{O}_{\kappa}(X^{t-1})$$
(3.51)

into the expression L'_1 , and we see that $L'_1 = L_1 + \mathcal{O}_{\kappa}(X^{2t-2})$ (note that $f\check{f}' = f$). Clearly $L_1 \in \mathfrak{O}_{\kappa}(X^{2t-1})$.

As for L'_2 we can assume that $g \in C^{\infty}_{c}(\mathbb{R})$ (since B_{κ} is bounded we can truncate g outside $\sigma(B_{\kappa})$) and use (3.37) again

$$\check{\mathbf{D}}g(B_{\kappa}) = -\int_{\mathbb{C}} R_{\kappa}(z) \bigl(\check{\mathbf{D}}B_{\kappa}\bigr) R_{\kappa}(z) \mathrm{d}\mu_{g}(z)
= g'(B_{\kappa})\check{\mathbf{D}}B_{\kappa} + \int_{\mathbb{C}} R_{\kappa}(z)^{2} [\check{\mathbf{D}}B_{\kappa}, B_{\kappa}] R_{\kappa}(z) \mathrm{d}\mu_{g}(z)
= g'(B_{\kappa})\check{\mathbf{D}}B_{\kappa} + \mathcal{O}_{\kappa}(X^{-2}) \in \mathfrak{O}_{\kappa}(X^{-1});$$
(3.52)

for the last identity we used Lemma 3.9 (and its proof). Obviously the first term to the right can be replaced by its real part. We conclude that $L'_2 = L_2 + \mathcal{O}_{\kappa}(X^{2t-2})$. Whence $L'_1 + L'_2 = L_1 + L_2 + \mathcal{O}_{\kappa}(X^{2t-2})$, showing (3.48a). Clearly $L_2 \in \mathfrak{O}_{\kappa}(X^{2t-1})$. Next to show that $L_3 - L_2 \in \mathfrak{O}_{\kappa}(X^{2t-2})$ under the extra conditions on g we

Next to show that $L_3 - L_2 \in \mathfrak{O}_{\kappa}(X^{2t-2})$ under the extra conditions on g we first note that the truncation of g outside $\sigma(B_{\kappa})$, to make it compactly supported, obviously applies to $(g')^{1/2}$ as well making again (3.37) applicable. Whence we can symmetrize L_2 , so that it suffices to show that

$$\bar{L}_{2} - \bar{L}_{3} \in \mathfrak{O}_{\kappa}(X^{2t-2});$$

$$\bar{L}_{2} = f(\breve{H})h(r)\sqrt{g'(B_{\kappa})}\bigl(\breve{\mathbf{D}}B_{\kappa}\bigr)\sqrt{g'(B_{\kappa})}h(r)f(\breve{H}),$$

$$\bar{L}_{3} = h(r)\sqrt{g'(B_{\kappa})}f(\breve{H})\bigl(\breve{\mathbf{D}}B_{\kappa}\bigr)f(\breve{H})\sqrt{g'(B_{\kappa})}h(r).$$
(3.53)

(Note that $L_3 = \overline{L}_3$.) Using (3.51) and (3.52) with \check{f} replaced by f we see that

$$[f(\breve{H}), h(r)] \in \mathfrak{O}_{\kappa}(X^{t-1}), \, [f(\breve{H}), \sqrt{g'(B_{\kappa})}] \in \mathfrak{O}_{\kappa}(X^{-1}),$$

which shows (3.53). By this proof we see that $L_3 \in \mathfrak{O}_{\kappa}(X^{2t-1})$. Alternatively we may easily check the latter property directly. (We can verify similarly that $L_4, L_5 \in \mathfrak{O}_{\kappa}(X^{2t-1})$.)

To show that $L_4 - L_3 \in \mathfrak{O}_{\kappa}(X^{2t-2})$ it suffices to show that

$$X^{1/2} f(\breve{H}) i[\breve{H}, B_{\kappa}] f(\breve{H}) X^{1/2} - f(\breve{H}) i[\breve{H}, A_{\kappa}] f(\breve{H}) + 2f(\breve{H}) (\kappa^2 B^2 + 1) B_{\kappa}^2 f(\breve{H}) \in \mathfrak{O}_{\kappa}(X^{-1}).$$
(3.54)

For the first term we substitute

$$\mathbf{i}[\breve{H}, B_{\kappa}] = \mathbf{i}[H, B_{\kappa}] - \lambda_0 \mathbf{i}[\Pi, B_{\kappa}] - \mathbf{i}[\breve{K}, B_{\kappa}].$$

As we saw in the last part of the proof of Lemma 3.9 each of these commutators contributes by an operator in $\breve{\mathcal{B}} \subset \mathfrak{O}_{\kappa}(X^0)$. This remains valid if we replace the term by

$$f(\breve{H})X^{1/2}\mathrm{i}[\breve{H}, B_{\kappa}]X^{1/2}f(\breve{H}),$$

and indeed the error (given by commuting the factors of $X^{1/2}$ and $f(\check{H})$) contributes by an operator in $\mathfrak{O}_{\kappa}(X^{-1})$. On the other hand

$$\begin{split} f(\breve{H}) \Big(X^{1/2} \mathrm{i}[\breve{H}, B_{\kappa}] X^{1/2} - \mathrm{i}[\breve{H}, A_{\kappa}] + 2(\kappa^{2}B^{2} + 1)B_{\kappa}^{2} \Big) f(\breve{H}) \\ &= f(\breve{H}) \Big(-2 \operatorname{Re} \left(\mathrm{i}[\breve{H}, X^{1/2}] B_{\kappa} X^{1/2} \right) + 2(\kappa^{2}B^{2} + 1)B_{\kappa}^{2} \Big) f(\breve{H}) \\ &= f(\breve{H}) \Big(-\operatorname{Re} \left(X^{-1/2} 2B B_{\kappa} X^{1/2} \right) + 2(\kappa^{2}B^{2} + 1)B_{\kappa}^{2} \Big) f(\breve{H}) + O_{\kappa}(X^{-1}) \\ &\in \mathfrak{O}_{\kappa}(X^{-1}). \end{split}$$

Thus (3.54) is proven.

Noting that $\operatorname{Com} \in \mathfrak{O}_{\kappa}(X^0)$ and $\left[\sqrt{g'(B_{\kappa})}, X^{-1/2}\right] \in \mathfrak{O}_{\kappa}(X^{-3/2})$ it follows that $L_5 - L_4 \in \mathfrak{O}_{\kappa}(X^{2t-2})$, which finishes the proof.

Remark 3.11. The formula (3.52) is an example of the following general commutator expansion formula for operators in $\mathfrak{O}_{\kappa}(X^t)$, cf. [GIS]. Thus for given $S \in \mathfrak{O}_{\kappa}(X^t), t \in \mathbb{R}, g \in C^{\infty}_{c}(\mathbb{R})$ and $K \geq 1$

$$[S, g(B_{\kappa})] = -\int_{\mathbb{C}} R_{\kappa}(z) \operatorname{ad}_{B_{\kappa}}(S) R_{\kappa}(z) d\mu_{g}(z) = \sum_{k=1}^{K} \frac{(-1)^{k-1}}{k!} g^{(k)}(B_{\kappa}) \operatorname{ad}_{B_{\kappa}}^{k}(S) + R(K); R(K) = (-1)^{K+1} \int_{\mathbb{C}} R_{\kappa}(z)^{K+1} \operatorname{ad}_{B_{\kappa}}^{K+1}(S) R_{\kappa}(z) d\mu_{g}(z) \in \mathfrak{O}_{\kappa}(X^{t-K-1}).$$
(3.55)

We will need better control of the error terms of Lemma 3.10 in the application in Subsection 3.3.3. Here we explain a procedure for establishing this. As noted before

the function g in the first part of the lemma may be taken compactly supported as required for (3.55). A principal goal is to refine the right-hand side of the statement

$$\check{\mathbf{D}}P - L_1 - L_5 = (L'_1 - L_1) + (L'_2 - L_5) \in \mathfrak{O}_{\kappa}(X^{2t-2}).$$

Due to (3.51) we can write

$$L'_{1} = 4f(\breve{H}) \operatorname{Re} \left(h'(r)Bg(B_{\kappa})h(r) \right) f(\breve{H}) + f(\breve{H}) \operatorname{Re} \left(\mathcal{O}_{\kappa}(X^{t-2})g(B_{\kappa})h(r) \right) f(\breve{H}) = L_{1} + f(\breve{H}) \operatorname{Re} \left(\mathcal{O}_{\kappa}(X^{t-2})g(B_{\kappa})h(r) \right) f(\breve{H}).$$

$$(3.56)$$

For L'_2 we apply (3.55) to $S = \check{f}(\check{H})$, yielding (for $K \ge 2$)

$$L'_{2} = f(\check{H})h(r)(\check{\mathbf{D}}g(B_{\kappa}))h(r)f(\check{H})$$

= $i\sum_{k=1}^{K} \frac{(-1)^{k-1}}{k!} f(\check{H})h(r)g^{(k)}(B_{\kappa})ad_{B_{\kappa}}^{k}(\check{f}(\check{H}))h(r)f(\check{H}) + O_{\kappa}(X^{2t-K-1})$
= $f(\check{H})h(r)g'(B_{\kappa})(\check{\mathbf{D}}B_{\kappa})f(\check{H}))h(r)f(\check{H}) + \sum_{k=2}^{K} S_{k} + O_{\kappa}(X^{2t-K-1}).$ (3.57)

The terms $S_k \in \mathfrak{O}_{\kappa}(X^{2t-k})$, and they have an explicit form suitable for the induction argument of Subsection 3.3.3. A consequence of (3.57) is the refined formula

$$L'_{2} = L_{2} + \sum_{k=2}^{K} \operatorname{Re}(S_{k}) + O_{\kappa}(X^{2t-K-1}).$$
(3.58)

3.2.1.1 Smooth sign function

In some of our applications of Lemma 3.10 the function g will be a 'smooth sign function' ζ_{ϵ} , cf. [AIIS]. It is constructed in terms of a cut-off function $\eta_{\epsilon} \in C^{\infty}(\mathbb{R})$ with special properties: The parameter $\epsilon > 0$ is considered small, and we define $\eta_{\epsilon}(b) = \frac{1}{\epsilon}\eta(\frac{b}{\epsilon})$, where $\eta'(b) > 0$ for |b| < 1, $\eta(b) = 0$ for $b \leq -1$ and $\eta(b) = 1$ for $b \geq 1$. We can choose η such that η' is even, $\sqrt{\eta}, \sqrt{\eta'} \in C^{\infty}(\mathbb{R})$ and for some c > 0

$$\eta'(b) \ge c \eta(b) \text{ for } b \in (-1, 1/2].$$
 (3.59)

The optimal choice of such c is not important for us since we will only need (3.59) in the following disguised form: For any $\tilde{c} > 0$ and all ϵ small enough ($\epsilon^2 \leq \frac{2}{3}c\tilde{c}$ suffices)

$$(\frac{\epsilon}{2} - b)\eta_{\epsilon}(b) \le \tilde{c}\,\eta_{\epsilon}'(b)$$
 for all $b \in \mathbb{R}$. (3.60a)

Note also that since η'_{ϵ} is even

$$1 = \epsilon \eta_{\epsilon}(b) + \epsilon \eta_{\epsilon}(-b). \tag{3.60b}$$

Let $\zeta_{\epsilon}(b) = \eta_{\epsilon}(b) - \eta_{\epsilon}(-b).$

3.3 Positive commutator estimates

We shall prove properties of possibly existing eigenfunctions of \check{H} at λ_0 . In the case where they dont exist we shall prove various resolvent estimates of \check{H} near λ_0 . Our analysis is based on Lemma 3.10 and positive commutator methods of [IS2], [AIIS] and [GIS]. For convenience we abbreviate in this section the Besov spaces with index s = 1/2 as $\mathcal{B} := \mathcal{B}_{1/2}, \ \mathcal{B}^* := \mathcal{B}^*_{1/2} \ \mathcal{B}^0_0 := \mathcal{B}^*_{1/2,0}$. (Note however that the same notation is used with a slightly different meaning in Subsection 3.3.2.)

3.3.1 A Rellich type theorem

As a first application we show the following result, largely mimicking [IS2].

Theorem 3.12. Every generalized eigenfunction in \mathcal{B}_0^* of \check{H} at λ_0 , or at any sufficiently nearby real λ'_0 , is in L^2_{∞} .

Proof. We shall use the Mourre estimate (3.13) with $\lambda = \lambda_0$, say with the positive number $\epsilon = \tilde{d}(\lambda_0)$ and $R = R_0 \geq 1$ fixed sufficiently big. In agreement with the statement $\kappa_0 > 0$ is fixed too (small), and we can freely use the estimate for $\kappa \in (0, \kappa_0]$ and for a fixed (small) neighbourhood \mathcal{U} of λ_0 and a fixed compact K. Whence we have freedom to choose $\kappa > 0$ very small whenever conveniently in the proof (note that \mathcal{U} and K do not depend on κ). Now suppose $\phi \in \mathcal{B}_0^*$ and $(\check{H} - \lambda'_0)\phi = 0$ with $|\lambda_0 - \lambda'_0|$ small, then we can write $\phi = f(H)\phi$ where $f \in C_c^{\infty}(\mathcal{U})$, $f(\lambda'_0) = 1$ and f is real.

Let $y_{\varepsilon}(r) = \frac{1}{1+\varepsilon r}$ and $x_{\varepsilon}(r) = ry_{\varepsilon}(r)$ for $\varepsilon \in [0, 1]$, and let X_{ε} and Y_{ε} be the operators of multiplication by x_{ε} and y_{ε} , respectively (this notation is consistent with (3.22)). These quantities are henceforth used with $R = R_0$ only. We note that $X := X_0$ agrees with the notation used in Section 3.2 and that $X_{\varepsilon} = XY_{\varepsilon}$. Note also that $\nabla x_{\varepsilon}(r) = y_{\varepsilon}^2 \omega$, whence for example $i[H, X_{\varepsilon}] = 2Y_{\varepsilon}BY_{\varepsilon}$.

It is convenient to introduce the following terminology for families $(T_{\varepsilon})_{0 < \varepsilon \leq 1} \subset \mathcal{D}_{\kappa}(X^t)$ of operators on L^2 , $t \in \mathbb{R}$. We say (T_{ε}) is uniformly of κ -order t if

$$\forall \kappa \in (0, \kappa'_0] \; \forall s \in \mathbb{R} \; \forall k \in \mathbb{N}_0 : \quad \sup_{\varepsilon \in (0, 1]} \|X^{s-k-t} \mathrm{ad}^k_{B_{\kappa, R_0}}(T_\varepsilon) X^{-s}\| < \infty.$$

We shall allow ourselves to write $T_{\varepsilon} = \mathcal{O}_{unf}(X^t)$ and $T_{\varepsilon} \in \mathfrak{O}_{unf}(X^t)$ to symbolize that the operator T_{ε} is member of a family of operators (T_{ε}) which is uniformly of κ -order t. If (S_{ε}) and (T_{ε}) are uniformly of κ -order s and t, respectively, then $(S_{\varepsilon}T_{\varepsilon})$ is uniformly of κ -order s + t.

<u>I</u>. We show that $\phi \in L^2_{-1/2}$. Fix any $\delta \in (0, 1/2)$. We shall consider the 'propagation observable'

$$P_{\varepsilon} = f(\breve{H}) X_{\varepsilon}^{\delta} \zeta_{\epsilon}(B_{\kappa}) X_{\varepsilon}^{\delta} f(\breve{H}), \, \varepsilon \in (0, 1].$$

Clearly $P_{\varepsilon} \in \mathfrak{O}_{\mathrm{unf}}(X^{2\delta})$. The positive parameter ϵ used here will be fixed shortly, small enough. Note that X_{ε} and P_{ε} are bounded due to the appearance of the factor y_{ε} . Eventually this factor will be removed by letting $\varepsilon \to 0$. More precisely we shall demonstrate some 'essential positivity' of $i[\check{H}, P_{\varepsilon}]$ persisting in the $\varepsilon \to 0$ limit. For any $n \in \mathbb{N}$ the function $\phi_n = \chi_n(r)\phi \in H^1$ (cf. the notation (1.29)), $(\breve{H} - \lambda'_0)\phi_n = [\breve{H}, \chi_n]\phi$ and whence the expectation (we use in general the notation $\langle T \rangle_{\phi} = \langle \phi, T \phi \rangle$)

$$\langle \mathbf{i}[\breve{H}, P_{\varepsilon}] \rangle_{\phi_n} = -2 \operatorname{Re} \langle \mathbf{i}[\breve{H}, \chi_n] P_{\varepsilon} \chi_n \rangle_{\phi}.$$
 (3.61a)

Since $\phi \in \mathcal{B}_0^*$ the term to the right vanishes as $n \to \infty$. Writing $\check{H} = H - \lambda_0 \Pi - \check{K}$ this claim is obvious for the contribution from H. For the other terms the commutator is effectively of order X^{-2} , cf. (3.42) and the subsequent discussion, whence their contributions vanish too as $n \to \infty$. It remains to study the left-hand side of (3.61a) in this limit.

Let
$$S_{\varepsilon} = Y_{\varepsilon}^{2} X_{\varepsilon}^{\delta-1} \left(\in \mathfrak{O}_{unf}(X^{\delta-1}) \right)$$
 and $\theta_{\epsilon} = \sqrt{\eta_{\epsilon}}$. We compute using Lemma 3.10
 $\langle \mathbf{i}[\breve{H}, P_{\varepsilon}] \rangle_{\phi_{n}} = \langle \breve{\mathbf{D}} P_{\varepsilon} \rangle_{\phi_{n}} = \langle L_{1} + L_{5} + \mathcal{O}_{unf}(X^{2\delta-2}) \rangle_{\phi_{n}};$
 $L_{1} = 4\delta f(\breve{H}) \operatorname{Re} \left((\kappa^{2}B^{2} + 1)S_{\varepsilon}B_{\kappa}\zeta_{\varepsilon}(B_{\kappa})X_{\varepsilon}^{\delta} \right) f(\breve{H}),$
 $X^{1/2}X_{\varepsilon}^{-\delta}L_{5}X_{\varepsilon}^{-\delta}X^{1/2} = \theta_{\epsilon}(B_{\kappa})f(\breve{H})\mathbf{i}[\breve{H}, A_{\kappa}]f(\breve{H})\theta_{\epsilon}(B_{\kappa})$
 $+ \theta_{\epsilon}(-B_{\kappa})f(\breve{H})\mathbf{i}[\breve{H}, A_{\kappa}]f(\breve{H})\theta_{\epsilon}(-B_{\kappa})$
 $- 2\theta_{\epsilon}(B_{\kappa})f(\breve{H})(\kappa^{2}B^{2} + 1)B_{\kappa}^{2}f(\breve{H})\theta_{\epsilon}(B_{\kappa})$
 $- 2\theta_{\epsilon}(-B_{\kappa})f(\breve{H})(\kappa^{2}B^{2} + 1)B_{\kappa}^{2}f(\breve{H})\theta_{\epsilon}(-B_{\kappa}).$

Noting the essential positivity of the term containing the factor $\kappa^2 B^2$ we obtain after symmetrizing

$$L_1 \ge 4\delta f(\breve{H}) Y_{\varepsilon} X_{\varepsilon}^{\delta - 1/2} B_{\kappa} \zeta_{\varepsilon}(B_{\kappa}) X_{\varepsilon}^{\delta - 1/2} Y_{\varepsilon} f(\breve{H}) + O_{\rm unf}(X^{2\delta - 2}).$$
(3.61b)

Using the Mourre estimate of the form discussed in the beginning of the proof and factors of $\tilde{f}(\check{H})$ (the latter can be inserted for free to bound $\kappa^2 B^2$), we can estimate for $\kappa > 0$ sufficiently small

$$L_{5} \geq X_{\varepsilon}^{\delta} X^{-1/2} \theta_{\epsilon}(B_{\kappa}) f(\breve{H}) \big(\tilde{d}(\lambda_{0}) - K \big) f(\breve{H}) \theta_{\epsilon}(B_{\kappa}) X^{-1/2} X_{\varepsilon}^{\delta} + X_{\varepsilon}^{\delta} X^{-1/2} \theta_{\epsilon}(-B_{\kappa}) f(\breve{H}) \big(\tilde{d}(\lambda_{0}) - K \big) f(\breve{H}) \theta_{\epsilon}(-B_{\kappa}) X^{-1/2} X_{\varepsilon}^{\delta} - 4\epsilon^{2} f(\breve{H}) X_{\varepsilon}^{\delta} X^{-1/2} \Big(\eta_{\epsilon}'(B_{\kappa}) + \eta_{\epsilon}'(-B_{\kappa}) \Big) X^{-1/2} X_{\varepsilon}^{\delta} f(\breve{H}) + \mathcal{O}_{\mathrm{unf}}(X^{2\delta-2}).$$

We shall require that $\epsilon > 0$ is so small that $8\epsilon^2 < \tilde{d}(\lambda_0)$, implying $\tilde{d}(\lambda_0) - 4\epsilon^2 > \tilde{d}(\lambda_0)/2$. We shall use (3.60a) with $\tilde{c} = \frac{\tilde{d}(\lambda_0)}{8\delta}$ and any possibly smaller ϵ , henceforth considered fixed. Now we fix a big $m \in \mathbb{N}$ such that (with this ϵ)

$$d(\lambda_0) - 4\epsilon^2 - \|K - \chi_m K \chi_m\| \ge d(\lambda_0)/2$$

and note that the contribution from the operator $\chi_m K \chi_m$ is in $\mathfrak{O}_{unf}(X^{2\delta-2})$.

Then we estimate

$$L_{5} \geq 2^{-1}\tilde{d}(\lambda_{0})f(\breve{H})X_{\varepsilon}^{\delta}X^{-1/2}\Big(\eta_{\epsilon}'(B_{\kappa}) + \eta_{\epsilon}'(-B_{\kappa})\Big)X^{-1/2}X_{\varepsilon}^{\delta}f(\breve{H}) + \mathcal{O}_{\mathrm{unf}}(X^{2\delta-2})$$
$$\geq 4\delta\tilde{c}f(\breve{H})Y_{\varepsilon}X_{\varepsilon}^{\delta-1/2}\Big(\eta_{\epsilon}'(B_{\kappa}) + \eta_{\epsilon}'(-B_{\kappa})\Big)X_{\varepsilon}^{\delta-1/2}Y_{\varepsilon}f(\breve{H}) + \mathcal{O}_{\mathrm{unf}}(X^{2\delta-2}).$$

We conclude the following lower bound by combining this bound with (3.61b) and by using (3.60a) and (3.60b),

$$L_1 + L_5 \ge 2\delta\epsilon f(\breve{H}) Y_{\varepsilon} X_{\varepsilon}^{\delta - 1/2} \big(\eta_{\epsilon}(B_{\kappa}) + \eta_{\epsilon}(-B_{\kappa}) \big) X_{\varepsilon}^{\delta - 1/2} Y_{\varepsilon} f(\breve{H}) + \mathcal{O}_{\mathrm{unf}}(X^{2\delta - 2}) = 2\delta f(\breve{H}) X_{\varepsilon}^{2\delta - 1} Y_{\varepsilon}^2 f(\breve{H}) + \mathcal{O}_{\mathrm{unf}}(X^{2\delta - 2}).$$

Whence we obtain from these arguments the uniform bound

$$\|X_{\varepsilon}^{\delta-1/2}Y_{\varepsilon}\phi\|^{2} = \lim_{n \to \infty} \|X_{\varepsilon}^{\delta-1/2}Y_{\varepsilon}f(\breve{H})\phi_{n}\|^{2} \le C\|X^{\delta-1}\phi\|^{2}.$$
 (3.62)

By letting $\varepsilon \to 0$ in (3.62) it follows that $\phi \in L^2_{\delta^{-1/2}} \subset L^2_{-1/2}$.

<u>II</u>. We show that $\phi \in L^2_{\infty}$ by a bootstrap argument. So suppose we have shown that $\phi \in L^2_{(m-1)/2-1/2}$ for an $m \in \mathbb{N}$. We did show this for m = 0 in Step I. Then we come to the conclusion that $\phi \in L^2_{m/2-1/2}$ by repeating the previous procedure using now the observable

$$P = P_{\varepsilon} = f(\breve{H}) X_{\varepsilon}^{m/2} \zeta_{\epsilon}(B_k) X_{\varepsilon}^{m/2} f(\breve{H}), \, \varepsilon > 0,$$

leading to the bound

$$\|Y_{\varepsilon}X_{\varepsilon}^{m/2-1/2}\phi\|^{2} = \lim_{n \to \infty} \|Y_{\varepsilon}X_{\varepsilon}^{m/2-1/2}f(\breve{H})\phi_{n}\|^{2} \le C\|X^{(m-1)/2-1/2}\phi\|^{2}.$$
 (3.63)

By letting $\varepsilon \to 0$ we deduce that $\phi \in L^2_{m/2-1/2}$.

3.3.2 LAP bound

We show a Besov space limiting absorption principle bound, largely mimicking [AIIS].

Theorem 3.13. Suppose λ_0 is not an eigenvalue of \hat{H} . Then there exist a neighbourhood $I \subset \mathbb{R}$ of λ_0 and C > 0 such that for all for $z \in \mathbb{C} \setminus \mathbb{R}$ with $\operatorname{Re} z \in I$ and all $\psi \in \mathcal{B}$

$$\|\check{R}(z)\psi\|_{\mathcal{B}^*} \le C \|\psi\|_{\mathcal{B}}.$$
(3.64)

To prove this bound we shall use the following weight-functions parametrized by $\nu \in \mathbb{N}_0$ and defined on \mathbb{R}_+

$$\Theta = \Theta_{\nu}(r) = 1 - \left(1 + r/2^{\nu}\right)^{-1}.$$

Noting the formula for derivatives

$$\Theta^{(k)} = (-1)^{k-1} k! 2^{-k\nu} (1 + r/2^{\nu})^{-1-k}; \quad k \ge 1,$$

we obtain the bounds

$$0 < \Theta \le \min\{1, r/2^{\nu}\}, 0 < (-1)^{k-1}\Theta^{(k)} \le k(-1)^k r^{-1}\Theta^{(k-1)} \le k! r^{-k}\Theta; \quad k \ge 2.$$
(3.65)

Below we will consider the composition $\Theta = \Theta_{\nu}(r_R)$ given in terms of the parameter R appearing in Proposition 3.3. In fact we shall apply Proposition 3.3 in essentially the same way as done in the beginning of the proof of Theorem 3.12, in particular for $R = R_0$ only. Since $\lambda_0 \notin \sigma_{\rm pp}(\check{H})$ we can now take K = 0 and therefore replace $2\tilde{d}(\lambda_0) - \epsilon - K$ by $\tilde{d}(\lambda_0)$ in (3.13). More precisely we use the Mourre estimate

(3.13) with $\epsilon = \tilde{d}(\lambda_0)/2 > 0$ and $R = R_0 \ge 1$ fixed sufficiently big. In agreement with this assertion, $\kappa_0 > 0$ is fixed too (small, in particular $\kappa_0 \le \kappa'_0$), and we can freely use the estimate for $\kappa \in (0, \kappa_0]$, for a fixed (small) open neighbourhood \mathcal{U} of λ_0 and in fact with K = 0, leaving us with the lower bound $\tilde{d}(\lambda_0)$ as claimed above. By the virial theorem $\sigma_{\rm pp}(\check{H}) \cap \mathcal{U} = \emptyset$. We fix a compact neighbourhood $I \subset \mathcal{U}$ of λ_0 such that there are no nonzero generalized eigenfunctions of \check{H} in \mathcal{B}_0^* at any point of I, which is doable thanks to Theorem 3.12. As we will see this I works in Theorem 3.13. Choose a real-valued $f \in C_c^{\infty}(\mathcal{U})$ such that f = 1 on a neighbourhood of I.

Below we use the notation \mathcal{B} and \mathcal{B}^* for the Besov spaces given in terms of r_{R_0} (rather than in terms of |x| as before), the latter written for short $r = r_{R_0}$. Of course the spaces $\mathcal{B} = \mathcal{B}(r)$ and $\mathcal{B}(|x|)$ coincide and similarly for the adjoint spaces (allowing us to change the meaning of the notation in (3.64)). We could use X for multiplication by r as in the previous subsection, however we find it more convenient to use the notation r only, even though mostly it will be the operator of multiplication by r. We will assign the notation $\mathcal{O}_{unf}(r^t)$ a different meaning (although very related) than $\mathcal{O}_{unf}(X^t)$ used in the previous subsection. More precisely we introduce the following terminology for families $(T_{\nu})_{\nu \in \mathbb{N}_0}$ of operators on L^2 .

We say $(T_{\nu}) \subset \mathfrak{O}_{\kappa}(X^t)$ is uniformly of κ -order t (for $t \in \mathbb{R}$) if

$$\forall \kappa \in (0, \kappa'_0] \; \forall s \in \mathbb{R} \; \forall k \in \mathbb{N}_0 : \\ \sup_{\nu \in \mathbb{N}_0} \| r^{s-k-t} \mathrm{ad}_{B_{\kappa,R_0}}^k(T_{\nu}) r^{-s} \| + \sup_{\nu \in \mathbb{N}_0} 2^{\nu/2} \| r^{s-k-t-1/2} \mathrm{ad}_{B_{\kappa,R_0}}^k(T_{\nu}) r^{-s} \| < \infty.$$

For the elements T_{ν} in such a family (T_{ν}) we write $T_{\nu} = \mathcal{O}_{unf}(r^t)$ and $T_{\nu} \in \mathfrak{O}_{unf}(r^t)$. Note that for example $\Theta^{1/2}$, considered as the composed function $\Theta_{\nu}^{1/2}(r(\cdot))$, is uniformly of κ -order 0, cf. (3.65). If $T_{\nu} \in \mathfrak{O}_{unf}(r^t)$ and S is any of the operators listed in Lemma 3.9 (with $R = R_0$), then (ST_{ν}) and $(T_{\nu}S)$ are also uniformly of κ -order t. More generally for any $T_{\nu} \in \mathfrak{O}_{unf}(r^t)$ and $S \in \mathfrak{O}_{\kappa}(X^s)$ the operators $ST_{\nu}, T_{\nu}S \in \mathfrak{O}_{unf}(r^{s+t})$. Similarly, if $S_{\nu} \in \mathfrak{O}_{unf}(r^s)$ and $T_{\nu} \in \mathcal{O}_{unf}(r^t)$ then $S_{\nu}T_{\nu}, T_{\nu}S_{\nu} \in \mathfrak{O}_{unf}(r^{s+t})$.

Lemma 3.14. There exists C > 0 and such that for all $z \in \mathbb{C} \setminus \mathbb{R}$ with $\operatorname{Re} z \in I$ and for all $\nu \in \mathbb{N}_0$ and $\psi \in \mathcal{B}$

$$\|\Theta'^{1/2}\phi\|^{2} \leq C(\|\phi\|_{\mathcal{B}^{*}}\|\psi\|_{\mathcal{B}} + \|\psi\|_{\mathcal{B}}^{2} + 2^{-\nu/2}\|r^{-3/4}\phi\|^{2}), \qquad (3.66a)$$

$$\|\phi\|_{\mathcal{B}^*}^2 \le C(\|\psi\|_{\mathcal{B}}^2 + \|r^{-3/4}\phi\|^2); \tag{3.66b}$$

here $\Theta = \Theta_{\nu}(r(\cdot))$ and $\phi = \check{R}(z)\psi$.

Proof. The bound (3.66b) follows from (3.66a) by taking supremum over $\nu \ge 0$. So it suffices to show (3.66a).

We consider $P_{\nu} = f(\breve{H})\Theta^{1/2}\zeta_{\epsilon}(B_{\kappa})\Theta^{1/2}f(\breve{H})$, where B_{κ} and ζ_{ϵ} are given as in the previous subsection and f is the function introduced above. Note that $P_{\nu} = \mathcal{O}_{\rm unf}(r^0)$. As before we have the freedom to choose $\kappa > 0$ sufficiently small, and again also the parameter $\epsilon > 0$ will be fixed sufficiently small.

Let $\theta_{\epsilon} = \sqrt{\eta_{\epsilon}'}$. We compute using Lemma 3.10

$$\begin{split} \langle \mathbf{i}[\breve{H}, P_{\nu}] \rangle_{\phi} &= \langle \mathbf{\check{D}} P_{\nu} \rangle_{\phi} = \langle L_{1} + L_{5} + \mathcal{O}_{\mathrm{unf}}(r^{-2}) \rangle_{\phi}; \\ L_{1} &= 2f(\breve{H}) \operatorname{Re} \left((\kappa^{2}B^{2} + 1) \frac{\Theta'}{\Theta^{1/2}} B_{\kappa} \zeta_{\epsilon}(B_{\kappa}) \Theta^{1/2} \right) f(\breve{H}), \\ r^{1/2} \Theta^{-1/2} L_{5} \Theta^{-1/2} r^{1/2} &= \theta_{\epsilon}(B_{\kappa}) f(\breve{H}) \mathbf{i}[\breve{H}, A_{\kappa}] f(\breve{H}) \theta_{\epsilon}(B_{\kappa}) \\ &+ \theta_{\epsilon}(-B_{\kappa}) f(\breve{H}) \mathbf{i}[\breve{H}, A_{\kappa}] f(\breve{H}) \theta_{\epsilon}(-B_{\kappa}) \\ &- 2\theta_{\epsilon}(B_{\kappa}) f(\breve{H}) (\kappa^{2}B^{2} + 1) B_{\kappa}^{2} f(\breve{H}) \theta_{\epsilon}(B_{\kappa}) \\ &- 2\theta_{\epsilon}(-B_{\kappa}) f(\breve{H}) (\kappa^{2}B^{2} + 1) B_{\kappa}^{2} f(\breve{H}) \theta_{\epsilon}(-B_{\kappa}). \end{split}$$

Noting the essential positivity of the term containing the factor $\kappa^2 B^2$ we obtain after symmetrizing

$$L_1 \ge 2f(\breve{H})\Theta'^{1/2}B_\kappa\zeta_\varepsilon(B_\kappa)\Theta'^{1/2}f(\breve{H}) + O_{\rm unf}(r^{-2}).$$
(3.67)

Using the Mourre estimate of the form discussed in the beginning of the proof and factors of $\tilde{f}(\check{H})$ (the latter can be inserted for free to bound $\kappa^2 B^2$), we can estimate for $\kappa > 0$ sufficiently small

$$L_5 \ge (\tilde{d}(\lambda_0) - 4\epsilon^2) f(\breve{H}) \Theta^{1/2} r^{-1/2} \Big(\eta'_{\epsilon}(B_{\kappa}) + \eta'_{\epsilon}(-B_{\kappa}) \Big) r^{-1/2} \Theta^{1/2} f(\breve{H}) + \mathcal{O}_{\rm unf}(r^{-2}).$$

We shall require that $\epsilon > 0$ is so small that $8\epsilon^2 < \tilde{d}(\lambda_0)$, implying $\tilde{d}(\lambda_0) - 4\epsilon^2 \ge \tilde{d}(\lambda_0)/2$. We shall use (3.60a) with $\tilde{c} = \tilde{d}(\lambda_0)/4$ and any possibly smaller ϵ , henceforth considered fixed. Thus by combining the above bound with (3.67) and using the bound $r\Theta' \le \Theta$ we finally obtain the following lower bound

$$L_1 + L_5 \ge \epsilon f(\breve{H}) \Theta'^{1/2} (\eta_{\epsilon}(B_{\kappa}) + \eta_{\epsilon}(-B_{\kappa})) \Theta'^{1/2} f(\breve{H}) + \mathcal{O}_{\rm unf}(r^{-2})$$

= $f(\breve{H}) \Theta' f(\breve{H}) + \mathcal{O}_{\rm unf}(r^{-2}).$

Note also the trivial bound $\langle \Theta' \rangle_{(1-f(\check{H}))\phi} \leq C \|\psi\|_{\mathcal{B}}^2$ (using that $\operatorname{Re} z \in I$) leading finally to the bound

$$\|\Theta'^{1/2}\phi\|^{2} \leq 2\langle \mathbf{i}[\breve{H}, P_{\nu}]\rangle_{\phi} + C\left(2^{-\nu/2}\|r^{-3/4}\phi\|^{2} + \|\psi\|_{\mathcal{B}}^{2}\right).$$
(3.68)

On the other hand

$$\langle \mathbf{i}[\check{H}, P_{\nu}] \rangle_{\phi} = \mathbf{i} \langle \psi, P_{\nu} \phi \rangle - \mathbf{i} \langle P_{\nu} \phi, \psi \rangle + 2(\operatorname{Im} z) \langle P_{\nu} \rangle_{\phi} \leq 2 \|P_{\nu}\|_{\mathcal{L}(\mathcal{B})} \|\psi\|_{\mathcal{B}} \|\phi\|_{\mathcal{B}^{*}} + 2 \|P_{\nu}\| |\operatorname{Im} \langle \check{H} - z \rangle_{\phi}| \leq 2 (\|P_{\nu}\|_{\mathcal{L}(\mathcal{B})} + \|P_{\nu}\|) \|\psi\|_{\mathcal{B}} \|\phi\|_{\mathcal{B}^{*}} \leq C \|\phi\|_{\mathcal{B}^{*}} \|\psi\|_{\mathcal{B}}.$$

$$(3.69)$$

The combination of (3.68) and (3.69) yields (3.66a).

Proof of Theorem 3.13. Let I be given as in Lemma 3.14. Suppose by contradiction that $z_n \to \lambda'_0 \in I$, $\|\psi_n\|_{\mathcal{B}} \to 0$ and $\|\phi_n\|_{\mathcal{B}^*} = 1$ where $\phi_n = \check{R}(z_n)\psi_n$ (here \mathcal{B} and \mathcal{B}^* are defined in terms of $r = r_{R_0}$). Fix $s \in (1/2, 3/4)$. We can assume that there exists w^{*}-lim_{$n\to\infty$} $\phi_n =: \phi \in L^2_{-s}$. By local compactness and an energy estimate we easily see (using the notation (1.29) and $\chi_m = \chi_m(r)$) that

$$\forall m \in \mathbb{N} : \quad \lim_{n \to \infty} \chi_m \phi_n = \chi_m \phi \text{ in } H^1$$

Moreover by (3.66a)

$$\exists C > 0 \,\forall m \in \mathbb{N} \,\forall \nu \in \mathbb{N}_0: \quad \|\Theta'^{1/2} \chi_m \phi\|^2 = \lim_{n \to \infty} \|\Theta'^{1/2} \chi_m \phi_n\|^2 \le C 2^{-\nu/2}.$$

From this bound we learn that $\phi \in \mathcal{B}_0^*$, and since also $(\check{H} - \lambda'_0)\phi = 0$ we obtain that $\phi = 0$ (cf. Theorem 3.12). By local compactness it then follows that $\lim_{n\to\infty} \phi_n = \phi = 0$ in $L^2_{-3/4}$, and by (3.66b) we then deduce that $\lim_{n\to\infty} \|\phi_n\|_{\mathcal{B}^*}^2 = 0$ contradicting the assumption that $\|\phi_n\|_{\mathcal{B}^*} = 1$ for all n.

3.3.3 Microlocal bounds and LAP

We prove microlocal bounds of the resolvent $\check{R}(z)$, largely mimicking [GIS] and [AIIS]. As in [GIS] we then obtain similar bounds of powers of the resolvent, and this implies LAP (the limiting absorption principle) for \check{H} near λ_0 .

We assume $\lambda_0 \notin \sigma_{\rm pp}(\check{H})$. Since we are not going to need the sharpest version of these microlocal bounds which use the optimal constant in Proposition 3.3 we will simplify the presentation and proceed exactly as in the beginning of the previous subsection, not to be repeated. (The sharper version is given by replacing the constant $\tilde{d}(\lambda_0)/2$ in Lemma 3.15 by any positive number less than $\tilde{d}(\lambda_0)$.) With the given neighbourhood $I \ni \lambda_0$ we define $I_{\pm} = \{z | \operatorname{Re} z \in I, \pm \operatorname{Im} z > 0\}$ and $I_{\mathbb{C}} = I_+ \cup I_-$.

We start by showing a version of the first step of an induction procedure of [GIS], cf. [AIIS]. Let for all $\sigma > 0$ the notation \mathcal{G}_{-}^{σ} and \mathcal{G}_{+}^{σ} signify the classes of real functions $g \in C^{\infty}(\mathbb{R})$ with support in $(-\infty, \sigma)$ and $(-\sigma, \infty)$, respectively.

Lemma 3.15. Under the above conditions let $\tilde{\sigma} > 0$ be given by $\tilde{\sigma}^2 = \tilde{d}(\lambda_0)/2$ and let $\sigma \in (0, \tilde{\sigma})$. There exists $\check{\kappa}_0 \in (0, \kappa_0]$ such that the following bounds hold for any $\chi(\pm \cdot < \sigma) \in \mathcal{G}^{\sigma}_{\pm}$ and $t \in (0, 1/2)$.

$$\forall \kappa \in (0, \check{\kappa}_0] \ \exists C > 0 \ \forall z \in I_{\pm} : \quad \|\chi(\pm B_{\kappa} < \sigma) \check{R}(z)\|_{\mathcal{L}(L^2_{1-t}, L^2_{-t})} \le C.$$
(3.70)

Proof. Let $t \in (0, 1/2)$ and $\delta = 1/2 - t$. Let $z \in I_+$, $\psi \in L^2_{1-t}$ and $\phi = \tilde{R}(z)\psi$ (we only consider I_+). We introduce for $\epsilon > 0$ (and with $R = R_0$) the following operator P of κ -order 2δ ,

$$P = f(\breve{H})X^{\delta}g_{\epsilon}(B_{\kappa})X^{\delta}f(\breve{H}); \quad g_{\epsilon}(b) = -(\sigma + 2\epsilon - b)^{2\delta}\eta_{\epsilon}^{2}(\sigma - b).$$
(3.71)

We compute

$$\frac{1}{2}(\sigma+2\epsilon-b)^{2t}g'_{\epsilon}(b) = \delta\eta^{2}_{\epsilon}(\sigma-b) + (\sigma+2\epsilon-b)(\eta_{\epsilon}\eta'_{\epsilon})(\sigma-b), \qquad (3.72)$$

and noting that $g'_{\epsilon} \ge 0$ and $(g'_{\epsilon})^{1/2} \in C^{\infty}$ we see that all of Lemma 3.10 applies. Letting $\theta_{\epsilon} = \sqrt{g'_{\epsilon}}$ we read off

$$\langle \mathbf{i}[\breve{H}, P] \rangle_{\phi} = \langle \mathbf{\check{D}}P \rangle_{\phi} = \langle L_1 + L_5 + \mathcal{O}_{\kappa}(X^{2\delta-2}) \rangle_{\phi}; L_1 = 4\delta f(\breve{H}) \operatorname{Re} \left((\kappa^2 B^2 + 1)) X^{\delta-1} B_{\kappa} g(B_{\kappa}) X^{\delta} \right) f(\breve{H}) X^t L_5 X^t = \theta_{\epsilon}(B_{\kappa}) f(\breve{H}) \mathbf{i}[\breve{H}, A_{\kappa}] f(\breve{H}) \theta_{\epsilon}(B_{\kappa}) - 2\theta_{\epsilon}(B_{\kappa}) f(\breve{H}) (\kappa^2 B^2 + 1) B_{\kappa}^2 f(\breve{H}) \theta_{\epsilon}(B_{\kappa}).$$

Letting $T = (\sigma + 2\epsilon - B_{\kappa})^{-t} \eta_{\epsilon} (\sigma - B_{\kappa}) X^{-t} f(\breve{H})$ we can estimate

$$L_1 \ge 4\delta f(\breve{H}) X^{-t} B_{\kappa} g(B_{\kappa}) X^{-t} f(\breve{H}) + \mathcal{O}_{\kappa}(X^{2\delta-2})$$
$$\ge 4\delta T^* (B_{\kappa}^2 - (\sigma + \epsilon)(\sigma + 2\epsilon)) T + \mathcal{O}_{\kappa}(X^{2\delta-2}).$$

To bound L_5 we first fix $\check{\kappa}_0 \in (0, \kappa_0]$ such with $C' = \|\tilde{f}(\check{H})B^4\tilde{f}(\check{H})\|$ and with an arbitrary sufficiently small $\epsilon' > 0$, the constant

$$c' = \tilde{d}(\lambda_0) - 2(\sigma + \epsilon')(\sigma + 2\epsilon') - 2\check{\kappa}_0^2 C' \text{ is positive.}$$
(3.73)

Then for all $\kappa \in (0, \check{\kappa}_0]$ and $\epsilon \in (0, \epsilon']$ the second term on the right-hand side of (3.72) contributes by a non-negative term and we obtain

$$L_5 \ge X^{-t} f(\breve{H}) \big(\tilde{d}(\lambda_0) - 2\kappa^2 C' - 2B_{\kappa}^2 \big) g'(B_{\kappa}) f(\breve{H}) X^{-t} + \mathcal{O}_{\kappa}(X^{2\delta-2}) \\\ge 2\delta T^* \big(\tilde{d}(\lambda_0) - 2\kappa^2 C' - 2B_{\kappa}^2 \big) T + \mathcal{O}_{\kappa}(X^{2\delta-2}).$$

Observing the cancellation of the terms containing B_{κ}^2 these bounds lead to the lower bounds

$$\check{\mathbf{D}}P \geq 2\delta T^* \big(-2(\sigma+\epsilon)(\sigma+2\epsilon) + \tilde{d}(\lambda_0) - 2\kappa^2 C' \big) T + \mathcal{O}_{\kappa}(X^{2\delta-2}) \\
\geq 2\delta c' T^* T + \mathcal{O}_{\kappa}(X^{2\delta-2}); \quad \kappa \in (0,\check{\kappa}_0], \ \epsilon \in (0,\epsilon'].$$
(3.74)

Next, introducing $S = \eta_{\epsilon}(\sigma - B_{\kappa})X^{-t}$ and using the fact that B_{κ} is bounded we obtain

$$\|Sf(\check{H})\phi\| \le C_1 \|T\phi\|.$$

Using the notation $\|\cdot\|_s = \|\cdot\|_{L^2_s}$ we also note that

$$|S(1 - f(\tilde{H})\phi|| \le C_2 ||\psi||_{\delta - 1/2}.$$

We conclude that

$$||S\phi||^2 \le C_3 (||T\phi||^2 + ||\psi||^2_{\delta - 1/2}).$$

By combining this bound with (3.74) we obtain

$$c \|S\phi\|^2 \le \langle \mathbf{\tilde{D}}P \rangle_{\phi} + C_4 (\|\phi\|^2_{\delta-1} + \|\psi\|^2_{\delta-1/2}); \quad c = 2\delta c'/C_3.$$

On the other hand for any $\varepsilon > 0$

$$\langle \mathbf{\tilde{D}} P \rangle_{\phi} = \mathbf{i} \langle \psi, P \phi \rangle - \mathbf{i} \langle P \phi, \psi \rangle + 2(\operatorname{Im} z) \langle P \rangle_{\phi} \leq C (\|S\phi\| + \|\phi\|_{\delta-3/2}) \|\psi\|_{\delta+1/2} \leq C \varepsilon (\|S\phi\|^2 + \|\phi\|_{\delta-3/2}^2) + C \varepsilon^{-1} \|\psi\|_{\delta+1/2}^2.$$

We choose $\varepsilon = c/(2C)$, yielding

$$\frac{c}{2} \|S\phi\|^2 \le C_5 \left(\|\phi\|_{\delta-1}^2 + \|\psi\|_{\delta+1/2}^2 \right).$$

Finally we invoke Theorem 3.13 and conclude (after a commutation) that for all $\kappa \in (0, \check{\kappa}_0]$ and $\epsilon \in (0, \epsilon']$

$$\|\eta_{\epsilon}(\sigma - B_{\kappa})\phi\|_{\delta^{-1/2}}^{2} \le C_{6}\|\psi\|_{\delta^{+1/2}}^{2}.$$
(3.75)

This finishes the proof since for any given function $\chi = \chi(\cdot < \sigma)$, we can write $\chi = \epsilon \chi \eta_{\epsilon}(\sigma - \cdot)$ for a small enough ϵ and then for any $\kappa \in (0, \check{\kappa}_0]$ bound

$$\|\chi(B_{\kappa} < \sigma)\phi\|_{\delta - 1/2} \le C_7 \|\eta_{\epsilon}(\sigma - B_{\kappa})\phi\|_{\delta - 1/2} \le C_8 \|\psi\|_{\delta + 1/2}.$$

Proposition 3.16. Under the conditions of Lemma 3.15 the same assertion holds for arbitrary t < 1/2 (i.e. the constraint t > 0 of the lemma is removed).

Proof. Let $J_m = \frac{1}{4}([2,3) - m)$; $m \in \mathbb{N}$. We saw in the previous proof that $\check{\kappa}_0 = \check{\kappa}_0(\sigma) \in (0, \kappa_0]$ chosen in agreement with (3.73) works for any given $\sigma \in (0, \tilde{\sigma})$, and in particular we obtained (3.70) with $t \in J_1$. We fix $\check{\kappa}_0 = \check{\kappa}_0(\sigma)$ this way along with the other positive constant $\epsilon' = \epsilon'(\sigma)$ of (3.73) and proceed to show by induction the assertion q(m):

$$\forall \sigma \in (0, \tilde{\sigma}) \, \forall t \in J_m \, \forall g \in \mathcal{G}^{\sigma}_{\pm} \, \forall \kappa \in (0, \check{\kappa}_0(\sigma)] :$$
$$\sup_{z \in I_{\pm}} \quad \|g(B_{\kappa}) \breve{R}(z)\|_{\mathcal{L}(L^2_{1-t}, L^2_{-t})} < \infty.$$

Since we have shown q(1) we can assume q(m-1) for a given $m \ge 2$ is known, and then it remains to verify q(m).

Let $t \in J_m$ and $\kappa \in (0, \check{\kappa}_0]$ be given, and introduce again $\delta = 1/2 - t$. Let $z \in I_+, \psi \in L^2_{1-t}$ and $\phi = \check{R}(z)\psi$ (we consider only I_+). We consider again (3.71) for $\epsilon \in (0, \epsilon']$. It suffices to show (3.75) for any such ϵ by the argument at the end of the proof of Lemma 3.15. For that we use the same scheme of proof as before, however since now possibly $\delta - 1 \ge -1/2$ we can not use Theorem 3.13 in the same way. Rather we need to combine the commutator expansion formula (3.55) with Lemma 3.10 which will allow us to use the induction hypothesis in combination with Theorem 3.13. This is already discussed in Remark 3.11.

First we look at the contribution to $\check{\mathbf{D}}P$ from $L'_1 - L_1 \in \mathfrak{O}_{\kappa}(X^{2\delta-2})$ of (3.56), i.e. we look at

$$Q_{2\delta-2} = f(\breve{H}) \operatorname{Re}\left(\mathcal{O}_{\kappa}(X^{\delta-2})g_{\epsilon}(B_{\kappa})X^{\delta}\right) f(\breve{H}).$$

By commutator expansion we see that with $\check{g}_{\epsilon} = \eta_{\epsilon}(\sigma + 2\epsilon - \cdot)$

$$Q_{2\delta-2} = \check{g}_{\epsilon}(B_{\kappa})Q_{2\delta-2}\check{g}_{\epsilon}(B_{\kappa}) + \mathcal{O}_{\kappa}(X^{-2})$$

= $\check{g}_{\epsilon}(B_{\kappa})X^{-t-1/2}\mathcal{O}_{\kappa}(X^{0})X^{-t-1/2}\check{g}_{\epsilon}(B_{\kappa}) + \mathcal{O}_{\kappa}(X^{-2}).$ (3.76)

Whence the induction hypothesis combined with Theorem 3.13 works upon taking $\epsilon > 0$ sufficiently small.

Next we look at the contribution to $\mathbf{D}P$ from $L'_2 - L_5 \in \mathfrak{O}_{\kappa}(X^{2\delta-2})$. As in the proof of Lemma 3.10 we split

$$L'_{2} - L_{5} = (L'_{2} - L_{2}) - (L_{3} - L_{2}) - (L_{4} - L_{3}) - (L_{5} - L_{4}) = \sum_{j=1}^{4} T_{j} \in \mathfrak{O}_{\kappa}(X^{2\delta - 2}).$$

By (3.58) the term T_1 can be represented as $Q_{2\delta-2}$ in (3.76), so we can argue in the same way for this term. By inspection of the proof of Lemma 3.10 we see that there are similar expansions for $T_j \in \mathfrak{O}_{\kappa}(X^{2\delta-2}), j = 2, 3, 4$, which allows us to treat these terms in the same way.

Finally using the auxiliary operators S and T of the proof of Lemma 3.15 we can mimic the last part of the proof using again the localization argument above to treat lower order terms, in particular various terms in $\mathfrak{O}_{\kappa}(X^{2\delta-2})$. We obtain q(m).

By the same method we can prove a two-sided estimate.

Proposition 3.17. Let $\tilde{\sigma} > 0$ be given by $\tilde{\sigma}^2 = \tilde{d}(\lambda_0)/2$ and let $\sigma \in (0, \tilde{\sigma})$. There exists $\check{\kappa}_0 > 0$ such that the following bounds hold for any s > 0 and for any pair $g_{\pm} \in \mathcal{G}_{\pm}^{\sigma}$ such that $\sup \sup g_{\pm} < \inf \sup g_{\pm}$.

$$\forall \kappa \in (0, \check{\kappa}_0] \ \exists C > 0 \ \forall z \in I_+ : \quad \|g_-(B_\kappa)\check{R}(z)g_+(B_\kappa)\|_{\mathcal{L}(L^2_{-s}, L^2_s)} \le C. \tag{3.77}$$

As in [GIS] the assertions Theorem 3.13 and Propositions 3.16 and 3.17 combine algebraically yielding bounds (including microlocal ones) of powers of the resolvent. In particular the following bounds follow, which in turn implies LAP.

Theorem 3.18. Suppose λ_0 is not an eigenvalue of H. Then

$$\forall k \in \mathbb{N} \,\forall s < 1/2 \,\exists C > 0 \,\forall z \in I_{\mathbb{C}} : \quad \|\breve{R}(z)^k\|_{\mathcal{L}(L^2_{k-s}, L^2_{s-k})} \le C. \tag{3.78}$$

In particular for any t > 1/2 the limits $\breve{R}(\lambda \pm i0) = \lim_{\epsilon \to 0_+} \breve{R}(\lambda \pm i\epsilon)$ exist in $\mathcal{L}(L^2_t, L^2_{-t})$ uniformly in $\lambda \in I$. Moreover there exists the strong weak-star limits

s-w^{*}-lim
$$\breve{R}(\lambda \pm i\epsilon) = \breve{R}(\lambda \pm i0) \in \mathcal{L}(\mathcal{B}, \mathcal{B}^*); \quad \lambda \in I.$$
 (3.79)

We will need the following application, see (4.37c).

Corollary 3.19. Suppose λ_0 is not an eigenvalue of \check{H} , s > 1/2 and that $f \in L^2_s$ is given such that $\check{R}(\lambda_0 + i0)f = \check{R}(\lambda_0 - i0)f$. Then

$$\breve{R}(\lambda_0 + i0)f = \breve{R}(\lambda_0 - i0)f \in L^2_{s-1}.$$
(3.80)

Proof. The result follows from Proposition 3.16 and Theorem 3.18 by using a suitable decomposition $1 = g_{-}(B_{\kappa}) + g_{+}(B_{\kappa}), g_{\pm} \in \mathcal{G}_{\pm}^{\sigma}$.

Remark 3.20. In the multiple case, here discussed with the assumptions of Section 2.3 only (in particular with (2.24) imposed), we need the analogues of (3.79) and Corollary 3.19, cf. (4.37a)-(4.37c). We argue by commenting on the necessary modifications that the same methods of proof work with a minimum of extra complication.

First note that the operator \breve{K} of (3.1a) needs to be replaced $\breve{K} = K_1 - K_2$ with K_1 and K_2 given by (2.35b). To simplify the form of this operator \breve{K} we introduce the notation

$$\mathcal{L}_{-\infty,\infty} = \bigcap_{r,t \in \mathbb{R}} \mathcal{L}(L_r^2(\mathbf{X}), L_t^2(\mathbf{X})).$$
(3.81)

Now using (2.32), Lemmas 2.6 and 2.9 as in the proof of Lemma 2.10 we see that

$$\begin{aligned}
\breve{K} - \breve{K}_1 - \breve{K}_2 &\in \mathcal{L}_{-\infty,\infty}; \\
\breve{K}_j &= \Pi_j I_j + I_j \Pi_j - \Pi_j I_j \Pi_j, \quad j = 1, 2.
\end{aligned}$$
(3.82)

Note for example that

$$\Pi - \Pi_1 - \Pi_2 \in (1 - \Delta)^{-1} \mathcal{L}^2_{-\infty,\infty} \subset \mathcal{L}^2_{-\infty,\infty}$$

Using (3.82) we can prove Proposition 3.3 by mimicking the proof for the nonmultiple case. In the first step of the proof of Lemma 3.7 we have the given properties for a_j and Π_j , j = 1, 2, rather than for a_0 and Π . This is all we need to repeat the proof. In Lemma 3.9 we need a similar replacement (in particular the polynomial factors in $P_1\Pi_jP_2$ are polynomials in quantities defined for a_j rather than for a_0). In the proof of Lemma 3.9 we can obviously add the class of operators in the intersection of all operators of finite κ -order to the classes \breve{B}_k . Note that $\mathcal{L}_{-\infty,\infty}$ is included in this way, and we can repeat the proof. The applications of Subsection 3.2.1 and the present section are the same as before.

- **Remarks 3.21.** 1) We discuss the extension to the case $\lambda_0 \in \sigma_{\rm pp}(H')$. We shall use Section 2.4 and Remark 3.20. Note that $\lambda_0 \notin \sigma_{\rm pp}(H'')$ and that Theorem 3.12 implies that the L^2 -eigenfunctions of H' at λ_0 are in L^2_{∞} (note also that the dimension of the corresponding eigenspace is finite, cf. Proposition 3.3). Under the assumptions of Section 2.4 the 'extra term' $\lambda_0 |\psi\rangle \langle \psi |$ of (2.59a) and (2.60b) is consequently in $\mathcal{L}_{-\infty,\infty}$. We can therefore prove Proposition 3.3 with the operator \check{H} of Section 2.4 (cf. Remark 3.20) and obtain the corresponding resolvent bounds of \check{R} from Subsections 3.3.2 and 3.3.3. In particular Corollary 3.19 also holds for the operators \check{H} and \check{R} of Section 2.4 in the case $\lambda_0 \in \sigma_{\rm pp}(H')$.
 - 2) We shall in Chapter 5 use variations of this construction. These are given by applying the theory of the present chapter to H replaced by $H_{\sigma} := H - \sigma \Pi_H$, $\sigma > 0$ small, where Π_H is the orthogonal projection onto the eigenspace of H corresponding to λ_0 (assuming that λ_0 is an eigenvalue). In the context of Chapter 5 this projection has finite rank, and if all L^2 -eigenfunctions are in L^2_{∞} the results of the present chapter generalize easily. However we do not have this good decay of the eigenfunctions for the cases considered in Chapter 5. Assuming 'sufficient decay', see for example (5.159) and (3.82), the results of the present chapter applies to some degree (sufficiently well for Chapter 5). In particular, to be concrete, there is the following version of (parts of) Theorem 3.18 and Corollary 3.19. Suppose

$$\operatorname{ran} \Pi_H \subset H_t^2 \text{ for some } t \in (1, 3/2), \tag{3.83}$$

and suppose λ_0 is not an eigenvalue of \check{H}_{σ} and that $f \in L^2_s$ for some $s \in (1/2, t - 1/2)$. Then there exist limits $\check{R}_{\sigma}(\lambda_0 + i0)f$, $\check{R}_{\sigma}(\lambda_0 - i0)f \in L^2_{(-1/2)^{-}}$. If in addition $\check{R}_{\sigma}(\lambda_0 + i0)f = \check{R}_{\sigma}(\lambda_0 - i0)f$, then

$$\ddot{R}_{\sigma}(\lambda_0 + \mathrm{i}0)f = \ddot{R}_{\sigma}(\lambda_0 - \mathrm{i}0)f \in L^2_{s-1}.$$
(3.84)

These results follow by straightforward modifications of the previous proofs. Note that all terms in the calculus involving the perturbation $-\sigma\Pi_H$ are regarded as 'errors', in fact we can use Lemma 3.10 and freely interchange $f(\check{H})$ and $\check{f}(\check{H})$ by $f(\check{H}_{\sigma})$ and $\check{f}(\check{H}_{\sigma})$, respectively, and similarly for the Mourre estimate Proposition 3.3. To obtain (3.84) under the given hypothesis and given limitations on s and t we need Lemma 3.15 for \check{R}_{σ} , and the interested reader may check that the errors from the indicated substitutions are harmless when mimicking the proof of the lemma.

In Section 5.3 a version of (3.84) is needed, but only for a s arbitrarily close 1/2.

Remark 3.22. The case when (2.24) fails was discussed in Section 2.5. One can then obtain the same results as the ones of Remarks 3.20 and 3.21 (depending on whether λ_0 is an eigenvalue of H' or not, respectively). Using Subsection 2.5.1 we can again invoke the formulas (2.35b) and (2.60b) for \check{H} (in the respectively cases) and in fact argue as before, obtaining the results.

Chapter 4

Rellich type theorems

We investigate the structure of eigenfunctions and resonances states at a two-cluster threshold under Condition 1.5 and (relevant part of) Condition 1.6. This structure depends strongly on decay properties of effective potentials. In Section 4.1 we treat the lowest threshold case which allows for somewhat stronger assertions than above this threshold. In Section 4.2 we study the latter case which demands a more careful examination of the 'eigentransform' of Remark 2.3. In Section 4.3 we study the physical models from Sections 1.2 and 1.3 by applying previously discussed methods, in particular we shall apply most of Section 4.2.

We are going to treat the multiple two-cluster case in detail using for the lowest threshold Proposition 2.12. The non-multiple two-cluster case can be treated similarly, it is notationally simpler of course, and we do not pay special attention to this case. The non-simple cases can be treated similarly. Only in Section 4.3 we do a complete study covering all cases, however for the physical models only.

Under the conditions of Proposition 2.12 (used in Section 4.2 without the lowest threshold condition) it is convenient to consider operators as quadratic forms defined on the first order Sobolev spaces, cf. Remark 2.13. For $k, s \in \mathbb{R}$, let H_s^k denote the weighted Sobolev space of order k with position-space weight $\langle x \rangle^s$ (as defined in Subsection 1.4.1) and

$$\mathcal{H}_{s,t}^{k,l} = H_s^k(\mathbf{X}_1) \oplus H_t^l(\mathbf{X}_2), \quad \mathcal{H}_s^k = \mathcal{H}_{s,s}^{k,k}, \quad \mathcal{H}^k = \mathcal{H}_0^k, \quad \mathcal{H}_s = \mathcal{H}_s^0, \quad \mathcal{H} = \mathcal{H}_0^0, \\
\mathcal{H}_{\infty}^k = \bigcap_{s \in \mathbb{R}} \mathcal{H}_s^k, \quad \mathcal{H}_{-\infty}^k = \bigcup_{s \in \mathbb{R}} \mathcal{H}_s^k, \quad \mathcal{H}_{s^+}^k = \bigcup_{t > s} \mathcal{H}_t^k, \quad \mathcal{H}_{s^-}^k = \bigcap_{t < s} \mathcal{H}_t^k.$$
(4.1)

We may write the operator $E_{\mathcal{H}}(\lambda_0)$ of Proposition 2.12 as

$$E_{\mathcal{H}}(\lambda_0) = -(p_1^2 + W_1) \oplus (p_2^2 + W_2) - V.$$
(4.2)

This is for the lowest threshold only, and the operator V is a non-local symmetric potential of order $\mathcal{O}(|x|^{-2-2\rho})$, see Proposition 2.12 and Remark 2.13. A modification of (4.2) for higher thresholds will be discussed in Section 4.2, and the 'exceptional cases' of Sections 2.4 and 2.5 will be discussed in Remarks 4.9 and 4.16. We will shortly introduce a slightly different decomposition (with corresponding different notation).

4.1 The case $\lambda_0 = \Sigma_2$

For $\rho < 2$ there are no results for the one-body problem in general. Thus for example it is not known if zero can be an eigenvalue of infinite multiplicity. However if either the potential is negative or positive at infinity like $-|x|^{-\rho}$ or $|x|^{-\rho}$, respectively, then it is known that zero is at most an eigenvalue of finite multiplicity (in fact it is not an eigenvalue in the negative case). In this section we impose the conditions of Proposition 2.12. In Subsections 4.1.2 and 4.1.3 we are going to use the proposition under additional sign conditions on the effective potentials W_1 and W_2 . In Subsection 4.1.4 the effective potentials are assumed to be homogeneous of degree -2 to leading order, which is a border line case with a rich structure. The case $\lambda_0 > \Sigma_2$ is divided into similar cases, to be treated in Section 4.1.1 and 4.2.1, respectively.

4.1.1 Extended eigentransform for $\lambda_0 = \Sigma_2$

Recall from Remark 2.3 the eigentransform and the corresponding inversion formula:

$$f = T^* u \text{ and } u = E_+(\lambda_0)f = Sf - \mathring{R}(\lambda_0)\Pi' HSf.$$

$$(4.3)$$

We can apply these relations to L^2 -eigenfunctions as well as to generalized eigenfunctions. In this extended sense the following result holds.

Lemma 4.1. For any $s \in \mathbb{R}$ the relations (4.3) obey

$$u \in H^1_s(=H^1_s(\mathbf{X})) \text{ and } (H-\lambda_0)u = 0 \Leftrightarrow f \in \mathcal{H}^1_s \text{ and } E_{\mathcal{H}}(\lambda_0)f = 0.$$
 (4.4)

Proof. Let $s \in \mathbb{R}$ be given. First we extend the boundedness result (2.56), now claiming that

$$\check{R}(z)\Pi' \in \mathcal{L}(L_s^2, H_s^2) \subset \mathcal{L}(L_s^2, H_s^1)$$
 (in particular for $z = \lambda_0$).

Noting that $\langle x \rangle^s (-\Delta - \breve{H}) \langle x \rangle^{-s}$ is ϵ -bounded relatively to $-\Delta$ the result follows by a standard computation using (2.56). By (2.31b), Lemmas 2.6, 2.9 2) and the polynomial decay of the threshold bound states it then follows that

$$S^*, T^* \in \mathcal{L}(H^1_s, \mathcal{H}^1_s) \text{ and } (1 - \check{R}(\lambda_0)\Pi'H)S \in \mathcal{L}(\mathcal{H}^1_s, H^1_s).$$
 (4.5)

In particular (4.4) holds.

The above proof is rather simple due to the fact that $\lambda_0 \notin \sigma(H')$. In Subsection 4.2.1 we derive a rather detailed version of the lemma when $\lambda_0 > \Sigma_2$. In this case the eigentransform and the corresponding inversion formula need a more subtle treatment. Moreover we remark that by ellipticity of the equation on the left-hand side of (4.4) we can replace the requirement $u \in H_s^1$ by $u \in L_s^2$ or $u \in H_s^2$ without changing the content of this side of (4.4). A similar comment is due for the right-hand side of (4.4), cf. the last part of Remark 2.13.

4.1.2 Negative slowly decaying effective potentials

Suppose $\rho < 2$ and that $I_j(x_j) = I_j(x)_{x^j=0}$, j = 1, 2, fulfill the following condition, cf. [FS]:

$$\exists R \ge 0 \quad \exists \epsilon > 0 \quad \forall y \in \mathbf{X}_j \text{ with } |y| \ge R :$$

$$I_j(y) \le -\epsilon \langle y \rangle^{-\rho} \text{ and } -2I_j(y) - y \cdot \nabla I_j(y) \ge \epsilon \langle y \rangle^{-\rho}.$$
(4.6)
We can for each j write the potential $W_j(x_j) = \widehat{W}_j(x_j) + B_j(x_j)$, where B_j is polynomially decreasing and \widehat{W}_j is smooth with $\partial_y^{\alpha} \widehat{W}_j = \mathcal{O}(\langle y \rangle^{-\rho - |\alpha|})$ and fulfills

$$\exists \epsilon > 0 \quad \forall y \in \mathbf{X}_j :$$

$$\widehat{W}_j(y) \le -\epsilon \langle y \rangle^{-\rho} \text{ and } -2\widehat{W}_j(y) - y \cdot \nabla \widehat{W}_j(y) \ge \epsilon \langle y \rangle^{-\rho}.$$
(4.7)

To see this we split each V_b in the definition of I_j as $V_b = V_b^{(1)} + V_b^{(2)}$, where the second term is compactly supported. We note that $\langle V_b^{(2)} \rangle_{\varphi_j}(x_j) = \langle \varphi_j, V_b^{(2)}(\cdot + x_j)\varphi_j \rangle_j$ is polynomially decreasing. This can be proven in the same way as we proved (2.22). It remains to consider the contribution from $V_b^{(1)}$. Now the potential $A(x_j) = \langle V_b^{(1)} \rangle_{\varphi_j}(x_j) - V_b^{(1)}(x_j)$ is C^{∞} . We note that $A(y) = \mathcal{O}(\langle y \rangle^{-\rho-1})$ follows in the same way as we proved (2.21a). In a similar fashion one checks that also $y \cdot \nabla A(y) = \mathcal{O}(\langle y \rangle^{-\rho-1})$, and due to these properties we are lead to consider $V_b^{(1)}(x_j)$, or rather $I_j^{(1)}(x_j) = \sum_b V_b^{(1)}(x_j)$. Since we know the property (4.6) for this sum we conclude a similar property for $\sum_b \langle V_b^{(1)} \rangle_{\varphi_j}$, and we can add a suitable compactly supported potential to have the bounds fulfilled for R = 0 (and some $\epsilon > 0$), cf. [FS]. This leads to a \widehat{W}_j fulfilling (4.7).

For convenience we use below the notation w_j for the potential \widehat{W}_j . Introducing the operators $h_j = p_j^2 + w_j$, j = 1, 2 we write (4.2) as

$$-E_{\mathcal{H}}(\lambda_0) = h_1 \oplus h_2 + v. \tag{4.8}$$

Note that v - V is polynomially decreasing. The operators h_1 and h_2 have a number of microlocal properties as stated in [FS, Theorems 4.1 and 4.2]. In particular there is a limiting absorption principle at zero: Let $s_0 = \frac{1}{2} + \frac{\rho}{4}$. Then for any $s > s_0$ there exist the norm-limits

$$r_j(0\pm \mathrm{i}0) = \lim_{\epsilon \to 0_+} r_j(\pm \mathrm{i}\epsilon) \in \mathcal{L}(H_s^{-1}(\mathbf{X}_j), H_{-s}^1(\mathbf{X}_j)),$$

where $r_j(z) = (h_j - z)^{-1}$. Moreover it is known from [Sk4] that

$$r_j(0\pm \mathrm{i}0) \in \mathcal{L}(\mathcal{B}_{s_0}, \mathcal{B}^*_{s_0}),\tag{4.9a}$$

stated in terms of the Besov spaces introduced in Subsection 1.4.1. We are going to use yet another property.

Suppose $r_j(0+i0)f = r_j(0-i0)f$ for a given $f \in L^2_t$ for some $t > s_0$, then

$$r_j(0 + i0)f = r_j(0 - i0)f \in L_s^2$$
 for any $s < t - 2s_0$,

cf. [FS, Theorems 4.1 (ii) and (iii)]. In fact one can strenghten the proof in [FS] and obtain the following statement:

Suppose $r_j(0+i0)f = r_j(0-i0)f$ for a given $f \in L^2_t$ for some $t > s_0$, then

$$r_j(0+i0)f = r_j(0-i0)f \in L^2_{t-2s_0}.$$
 (4.9b)

This statement is similar to Corollary 3.19, and it can be proved by improved versions of some resolvent bounds from [FS] which are one-body threshold analogues of the bounds of Proposition 3.16.

Theorem 4.2. Under the conditions of Proposition 2.12 and (4.6) for j = 1, 2,

 $\dim \ker(H - \lambda_0) < \infty.$

If $\phi \in \mathcal{B}^*_{s_0,0}(\mathbf{X})$ solves the distributional equation $(H - \lambda_0)\phi = 0$, then $\phi \in L^2_{\infty}(\mathbf{X})$.

Proof. <u>I</u>. By Remarks 2.3 and 2.13

$$\dim \ker(H - \lambda_0) = \dim \ker E_{\mathcal{H}}(\lambda_0).$$

Note that we have chosen the form interpretation of the appearing operators in agreement with Remark 2.13, but that the operator interpretation of Remark 2.3 amount to the same spaces, cf. Remark 2.13. Let $R_{\text{diag}}^{\pm} = r_1(0 \pm i0) \oplus (r_2(0 \pm i0))$ and suppose $f \in \ker E_{\mathcal{H}}(\lambda_0)$. We write the equations $R_{\text{diag}}^{\pm}E_{\mathcal{H}}(\lambda_0)f = 0$ as

$$(1+K^{\pm})f = 0, \quad K^{\pm} = R^{\pm}_{\text{diag}}v,$$
 (4.10)

where 1f arises by writing $f = (f_1, f_2) \in \mathcal{H}^1$ and using (to be shown below) that

$$r_j(0 \pm i0)h_j f_j = f_j.$$
 (4.11)

We note that

$$K^{\pm} \in \mathcal{C}(\mathcal{H}_{-s}^{1}) \text{ for } s \in (s_{0}, 2\rho + 2 - s_{0}).$$
 (4.12)

Since K^{\pm} is compact (on any such space) it follows from Fredholm theory that dim ker $E_{\mathcal{H}}(\lambda_0) < \infty$.

<u>II</u>. We prove the following more general version of (4.11): Suppose $f_j \in \mathcal{B}^*_{s_0,0}(\mathbf{X}_j)$ and that the distribution $h_j f_j \in \mathcal{B}_{s_0}(\mathbf{X}_j)$, then (4.11) holds and the function $f_j \in H^1_{-s}$ for any $s > s_0$ (for the problem above $h_j f_j \in L^2_s(\mathbf{X}_j) \subset \mathcal{B}_{s_0}(\mathbf{X}_j)$ for any s as in (4.12)). To see this we let $\chi_R(x) = \chi(|x|/R), R \ge 1$, be given in agreement with (1.29). We consider χ_R as a multiplication operator on \mathbf{X}_j . Now $r_j(0 \pm i0)h_j\chi_R f_j = \chi_R f_j$ for all R > 1. On the other hand

$$h_j \chi_R f_j = \chi_R h_j f_j - (\Delta \chi_R) f_j - 2(\nabla \chi_R) \cdot \nabla f_j.$$

For the last term

$$R^{2} \| (\nabla \chi_{R}) \cdot \nabla f_{j} \|_{L^{2}(\mathbf{X}_{j})}^{2} \\ \leq C_{1} \langle p^{2} \rangle_{\chi_{2R} \bar{\chi}_{R/2} f_{j}} \\ \leq C_{2} \left(\operatorname{Re} \langle \chi_{2R}^{2} \bar{\chi}_{R/2}^{2} f_{j}, h_{j} f_{j} \rangle + R^{-\rho} \| \chi_{4R} f_{j} \|^{2} \right)$$

$$= C_{2} \left(\mathcal{O}(R^{0}) + R^{-\rho} o(R^{2s_{0}}) \right)$$

$$= o(R^{1-\rho/2}),$$

$$(4.13)$$

yielding

$$\lim_{R \to \infty} R^{s_0} \| (\nabla \chi_R) \cdot \nabla f_j \|_{L^2(\mathbf{X}_j)} = 0$$

For the middle term

$$R^{s_0} \| (\Delta \chi_R) f_j \|_{L^2(\mathbf{X}_j)} \le C R^{2s_0 - 2} \big(R^{-s_0} \| \chi_{2R} f_j \|_{L^2(\mathbf{X}_j)} \big),$$

yielding

$$\lim_{R \to \infty} R^{s_0} \| (\Delta \chi_R) f_j \|_{L^2(\mathbf{X}_j)} = 0.$$

These computations lead to

$$f_j = \underset{R \to \infty}{\text{w}^*-\text{lim}} \ \chi_R f_j = \underset{R \to \infty}{\text{w}^*-\text{lim}} \ r_j(0 \pm i0) h_j \chi_R f_j = r_j(0 \pm i0) h_j f_j \text{ in } \mathcal{B}_{s_0}^*(\mathbf{X}_j),$$

and therefore (4.11) holds for $f_j \in \mathcal{B}^*_{s_0,0}(\mathbf{X}_j)$ obeying $h_j f_j \in \mathcal{B}_{s_0}(\mathbf{X}_j)$.

<u>III</u>. Using again the 'eigentransform' of Remark 2.3 it is readily seen that any $\phi \in \mathcal{B}^*_{s_0,0}(\mathbf{X})$ solving the distributional equation $(H - \lambda_0)\phi = 0$ corresponds to a (distributional) solution $f = (f_1, f_2), E_{\mathcal{H}}(\lambda_0)f = 0$ with $f_j \in \mathcal{B}^*_{s_0,0}(\mathbf{X}_j)$ obeying $h_j f_j \in \mathcal{B}_{s_0}(\mathbf{X}_j)$, cf. Lemma 4.1. By the previous steps we then conclude that (4.10) is fulfilled. In particular $K^+f = K^-f$. Conversely using that v is symmetric we obtain that

$$0 = \operatorname{Im}\langle f, vf \rangle = -\operatorname{Im}\langle R^+_{\operatorname{diag}}vf, vf \rangle \tag{4.14}$$

for any solution to $(1 + K^+)f = 0$, where $f = (f_1, f_2)$ obeys $f_j \in \mathcal{B}^*_{s_0}(\mathbf{X}_j)$, j = 1, 2. Whence also $(1 + K^-)f = 0$ for any such f. (Similarly a solution to $(1 + K^-)f = 0$ is also a solution to $(I + K^+)f = 0$.) Moreover $E_{\mathcal{H}}(\lambda_0)f = 0$ in the distributional sense. If $(1 + K^+)f = 0$ and $(1 + K^-)f = 0$ where $f = (f_1, f_2)$ and $f_j \in \mathcal{B}^*_{s_0}(\mathbf{X}_j)$ better decay is obtained by invoking (4.9b). Explicitly we may fix s as in (4.12) and starting with the input $f \in \mathcal{H}^0_{-s}$ conclude that $f \in \mathcal{H}^0_{-s+2\kappa}$, where $\kappa = \rho + 1 - s_0$. By iterating this argument we conclude that $f_j \in L^2_{\infty}(\mathbf{X}_j)$. Next, by the 'inversion formula' of Remark 2.3 any such f corresponds to a

$$\phi \in \ker(H - \lambda_0) \cap L^2_{\infty}(\mathbf{X}),$$

see Lemma 4.1. In particular, cf. the remark at the beginning of the paragraph, this is valid for any $\phi \in \mathcal{B}^*_{s_0,0}(\mathbf{X})$ solving $(H - \lambda_0)\phi = 0$.

4.1.3 Positive slowly decaying effective potentials

Suppose $\rho < 2$ and that $I_j(x_j) = I_j(x)|_{x^j=0}$, j = 1, 2, fulfill the following condition:

$$\exists R \ge 0 \quad \exists \epsilon > 0 \quad \exists \bar{\rho} \in [\rho, \frac{2}{3}(1+\rho)) \quad \forall y \in \mathbf{X}_j \text{ with } |y| \ge R :$$
$$I_j(y) \ge \epsilon \langle y \rangle^{-\bar{\rho}}. \tag{4.15}$$

We write for each j the potential $W_j(x_j) = \widehat{W}_j(x_j) + B_j(x_j)$, where B_j is polynomially decreasing and \widehat{W}_j is smooth with $\partial_y^{\alpha} \widehat{W}_j = \mathcal{O}(\langle y \rangle^{-\rho - |\alpha|})$ and fulfills

$$\exists \epsilon > 0 \quad \exists \bar{\rho} \in [\rho, \frac{2}{3}(1+\rho)) \quad \forall y \in \mathbf{X}_j : \quad \widehat{W}_j(y) \ge \epsilon \langle y \rangle^{-\bar{\rho}}. \tag{4.16}$$

This is seen by arguments as in Subsection 4.1.2 (using that $\bar{\rho} < 1 + \rho$). Again we shall use the notation w_j for the potential \widehat{W}_j , introduce the operators $h_j = p_j^2 + w_j$, j = 1, 2 and write (4.2) as (4.8).

We introduce the symbols $s_j = s_j(y,\xi) = (\xi_j^2 + w_j(y))^{-1}$. Note that s_j belongs to the Hörmander class $S(\langle y \rangle^{\bar{\rho}} \langle \xi \rangle^{-2}, g)$, where the metric is given by

$$g(v) = \langle y \rangle^{2(\bar{\rho} - \rho - 1)} v_y^2 + \langle y \rangle^{\bar{\rho}} v_{\xi}^2, \quad v = (v_y, v_{\xi}),$$

and note that the corresponding Weyl calculus has 'Planck constant' of size

$$\langle y \rangle^{(\bar{\rho}-\rho-1)} \langle y \rangle^{\bar{\rho}/2} = \langle y \rangle^{-t}; t = 1 + \rho - 3\bar{\rho}/2.$$

Next we may construct a parametrix to infinite order for $h_j = \operatorname{Op}^{\mathsf{w}}(\xi_j^2 + w_j(y))$ following a standard procedure. However for the theorem below we only need the first step which amounts to letting the parametrix be given by $r_j^0 = \operatorname{Op}^{\mathsf{w}}(s_j)$. We compute $r_j^0 h_j = 1 + A_j$ for some $A_j = \operatorname{Op}^{\mathsf{w}}(a_j)$ with $a_j \in S(\langle y \rangle^{-t} \langle \xi \rangle^{-1}, g)$.

Theorem 4.3. Under the conditions of Proposition 2.12 and (4.15) for j = 1, 2,

$$\dim \ker(H - \lambda_0) < \infty.$$

If $\phi \in L^2_{-\infty}(\mathbf{X})$ (or alternatively $\phi \in H^1_{-\infty}(\mathbf{X})$) solves the distributional equation $(H - \lambda_0)\phi = 0$, then $\phi \in L^2_{\infty}(\mathbf{X})$.

Proof. <u>I</u>. By Remark 2.3

$$\dim \ker(H - \lambda_0) = \dim \ker E_{\mathcal{H}}(\lambda_0).$$

Suppose $(H - \lambda_0)\phi = 0$ for $\phi \in H^1_s(\mathbf{X})$ for some $s \in \mathbb{R}$ (possibly nonzero). Due to Lemma 4.1 the eigentransform $f = T^*\phi \in \mathcal{H}^1_s$ and $E_{\mathcal{H}}(\lambda_0)f = 0$. Let $R = r_1^0 \oplus r_2^0$ and $A = A_1 \oplus A_2$. Then we write the equation $RE_{\mathcal{H}}(\lambda_0)f = 0$ for any $f \in L^2_s$ as

$$(1 + A + K)f = 0, \quad K = Rv, \tag{4.17}$$

where we use (4.8) and that for the components of $f = (f_1, f_2)$

$$r_j^0(0)h_jf_j = (1+A_j)f_j.$$

We note that $A + K \in \mathcal{C}(\mathcal{H}_s^1)$. It then follows from Fredholm theory that f belongs to a finite-dimensional subspace of \mathcal{H}_s^1 . This dimension is an upper bound of the dimension of the set of functions $\phi \in H_s^1(\mathbf{X})$ solving $(H - \lambda_0)\phi = 0$ (seen by the inversion formula). In particular the latter space is finite-dimensional. The first statement of the theorem follows by this argument for s = 0.

<u>II</u>. By an iteration procedure using (4.17), cf. the proof of Theorem 4.2, it follows that any (zero-energy) generalized eigenfunction $f \in \mathcal{H}^1_{-\infty}$ must belong to \mathcal{H}^1_{∞} . Note that we can deduce that $f \in \mathcal{H}^1_{s+t}$ given that $f \in \mathcal{H}^1_{s+(k-1)t}$, where $t = 1 + \rho - 3\bar{\rho}/2$. Since there is no limit on $k \in \mathbb{N}$) used for this argument, indeed this conclusion comes out by iteration. Consequently, thanks to Lemma 4.1, the generalized eigenfunctions of H at λ_0 in $\mathcal{H}^1_{-\infty}(\mathbf{X})$ are all in $L^2_{\infty}(\mathbf{X})$.

4.1.3. Positive slowly decaying effective potentials

The equation (4.17) might have 'spurious' solutions, i.e. solutions not corresponding to eigenfunctions of H. We can cure this 'deficiency' by showing that the exact inverse h_j^{-1} , j = 1, 2, in fact is a pseudodifferential operator. We do this below. In particular h_j^{-1} has nice commutation proporties with power-type weights in the configuration space. Such properties were proved by Yafaev [Ya2], and we are in fact going to use [Ya2]. More precisely we are going to use the assertion

$$\forall a, b \ge 0, a - b \ge \bar{\rho}: \quad B = \underset{\epsilon \to 0_+}{\text{s-lim}} \langle y \rangle^{-a} (h_j + \epsilon)^{-1} \langle y \rangle^{b} \text{ exists.}$$
(4.18)

Here we note that $B_{\epsilon} = \langle y \rangle^{-a} (h_j + \epsilon)^{-1} \langle y \rangle^b$ is bounded uniformly in small positive ϵ , which follows by keeping track of constants in the proof of [Ya2, Theorem 1]. Since $\lim_{\epsilon \to 0_+} B_{\epsilon} u$ exists if $u = \langle y \rangle^{-b} h_j v$ with $v \in \mathcal{D}(h_j)$ and the set of such u's is dense in $L^2(\mathbf{X}_j)$, indeed (4.18) follows (in fact $Bu = \langle y \rangle^{-a} h_j^{-1} \langle y \rangle^b u$ for any u of this form).

Lemma 4.4. The operators s-lim_{$\epsilon \to 0_+$} $\langle y \rangle^{-\bar{\rho}} (h_j + \epsilon)^{-1}$, j = 1, 2, (extending $\langle y \rangle^{-\bar{\rho}} h_j^{-1}$) are pseudodifferential operators with symbol in $S(\langle \xi \rangle^{-2}, g)$.

Proof. Recall $r_j^0 h_j = 1 + A_j$, where A_j has symbol $a_j \in S(\langle y \rangle^{-t} \langle \xi \rangle^{-1}, g)$. If $-1 \notin \sigma(A_j)$

$$(h_j + \epsilon)^{-1} = (1 + A_j)^{-1} r_j^0 - \epsilon (1 + A_j)^{-1} r_j^0 (h_j + \epsilon)^{-1}$$

yielding s-lim_{$\epsilon > 0$} $\langle y \rangle^{-\bar{\rho}} (h_j + \epsilon)^{-1} = \langle y \rangle^{-\bar{\rho}} (1 + A_j)^{-1} r_j^0$. (Whence formally $h_j^{-1} = (1 + A_j)^{-1} r_j^0$.) Using the Neumann series to expand $(1 + A_j)^{-1}$ and Beal's criterion, cf. [FS, Subsection 4.2], we conclude that indeed $\langle y \rangle^{-\bar{\rho}} (1 + A_j)^{-1} r_j^0$ has symbol in $S(\langle \xi \rangle^{-2}, g)$ completing the proof in this case.

To treat the general case we let $\chi_1(r) = \chi(r < 1)$ and $\chi_2(r) = \chi(r > 1)$ form a quadratic partition of unity of smooth non-negative functions on \mathbb{R} , $\chi_1(r)^2 + \chi_2(r)^2 = 1$, such that χ_1 is supported in $(-\infty, 2)$ and $\chi_1(r) = 1$ for r < 1. We introduce for l > 1 the functions $\chi_{i,l}(r) := \chi_i(r/l)$; i = 1, 2. Let

$$R_{l} = \bigoplus_{j=1}^{2} \left(\chi_{1,l}(r_{j}) h_{j}^{-1} \chi_{1,l}(r_{j}) + \chi_{2,l}(r_{j}) r_{j}^{0} \chi_{2,l}(r_{j}) \right) = \bigoplus_{j=1}^{2} r_{j,l}.$$

Here $\chi_{1,l}(r_j)h_j^{-1}\chi_{1,l}(r_j) := \text{s-lim}_{\epsilon \searrow 0} \chi_{1,l}(r_j)(h_j + \epsilon)^{-1}\chi_{1,l}(r_j)$. Using (4.18) and Beal's criterion we deduce that $\chi_{1,l}(r_j)h_j^{-1}\chi_{1,l}(r_j)$ and therefore also $r_{j,l}$ are pseudodifferential operators with symbol in $S(\langle y \rangle^{\bar{\rho}} \langle \xi \rangle^{-2}, g)$. Next we write $r_{j,l}h_j = 1 + A_{j,l}$ and note (as above) that $A_{j,l}$ has symbol $a_{j,l} \in S(\langle y \rangle^{-t} \langle \xi \rangle^{-1}, g)$. We also observe that

$$\|\chi_{2,l}(r_j)A_j\chi_{2,l}(r_j)\| \to 0 \text{ for } l \to \infty.$$

Using this and a little computation it follows that

$$||A_{j,l}|| \to 0 \text{ for } l \to \infty.$$

In particular it follows that $-1 \notin \sigma(A_{j,l})$ for l taken big enough. Consequently for any such big number $\langle y \rangle^{-\bar{\rho}} h_j^{-1} = \langle y \rangle^{-\bar{\rho}} (1 + A_{j,l})^{-1} r_{j,l}$ has symbol in $S(\langle \xi \rangle^{-2}, g)$.

We note that Lemma 4.4 can be used to replace (4.17) by the cleaner assertion

$$(1+\widetilde{K})f = 0; \quad \widetilde{K} = \widetilde{R}v, \quad \widetilde{R} = h_1^{-1} \oplus h_1^{-1},$$

$$(4.19)$$

of course given an appropriate interpretation.

4.1.4 Homogeneous degree -2 effective potentials

Suppose $\rho \ge 1/2$, the conditions of Proposition 2.12 and that $W_j = W_j(x_j)$, j = 1, 2, fulfill the following condition (recall that in general W_j is bounded outside a bounded set):

For j = 1, 2 there exists a real continuous function $q_j = q_j(y)$ on the unit sphere \mathbb{S}_j of \mathbf{X}_j and a bounded potential $B_j(x_j) = \mathcal{O}(\langle x_j \rangle^{-3})$ on \mathbf{X}_j such that

$$\exists R > 0 \quad \forall y = r\theta \in X_j \text{ with } |y| = r \ge R : W_j(y) = \frac{q_j(\theta)}{r^2} + B_j(y). \tag{4.20}$$

This condition may arise by a Taylor expansion of $I_j(x)$, j = 1, 2, as follows: Decompose

$$\sum_{b \not \subset a_j} V_b^{(1)}(x^b) = \sum_{b \not \subset a_j} V_b^{(1)}((x_j)^b) + \sum_{b \not \subset a_j} (x^j)^b \cdot \nabla V_b^{(1)}((x_j)^b) + \mathcal{O}(\langle x_j \rangle^{-\rho-2}).$$

Assume now that the first term vanishes identically and that the last term contributes to W_j by a term of order $\mathcal{O}(\langle x_j \rangle^{-3})$ (valid for $\rho \geq 1$ of course). The middle term contributes to W_j by $\sum_{b \not \in a_j} k_j^b \cdot \nabla V_b^{(1)}((x_j)^b)$, where $k_j = \langle \varphi_j, x^j \varphi_j \rangle_j$, so the content of (4.20) would in this case be the condition

$$\sum_{b \not \subset a_j} k_j^b \cdot \nabla V_b^{(1)}((x_j)^b) = \frac{q_j(\theta)}{r^2} + \mathcal{O}(r^{-3}).$$

This has relevance with nonzero q_j 's in a certain case for systems with Coulomb interactions, see Subsections 1.2.1 and 1.3.1. The case $q_j = 0$, j = 1, 2 is of course included in (4.20), and the relevance of this case for the physics models is also explained there.

For simplicity of presentation let us in the following assume R = 1. For $n_j = \dim \mathbf{X}_j \geq 2$ the spectrum of the (minus) Laplace-Beltrami operator $-\Delta_{\theta_j}$ on the unit sphere $\mathbb{S}_j \subset \mathbf{X}_j$ is known to be $\{l(l+n_j-2) \mid l \in \mathbb{N}_0\}$. For our problem it is relevant to study the spectrum of $-\Delta_{\theta_j} + q_j$. For $n_j = 1$ we define $-\Delta_{\theta_j} = 0$, so in this case the spectrum of $-\Delta_{\theta_j} + q_j$ is $\{q_j(-1), q_j(1)\}$. In any dimension it is convenient to use the following parametrization: Write each $\mu \in \sigma(-\Delta_{\theta_j} + q_j)$ as

$$\mu = \nu^2 - \frac{(n_j - 2)^2}{4},\tag{4.21}$$

where by convention $\nu \geq 0$ if $\mu \geq -\frac{(n_j-2)^2}{4}$, and $i\nu > 0$ if $\mu < -\frac{(n_j-2)^2}{4}$. Then the collection of such numbers ν is denoted by σ_j . Let $P_{j,\nu}$, $\nu \in \sigma_j$, denote the orthogonal projection onto the corresponding eigenspace. For convenience we omit in the following the subscript j. Letting ζ denote a corresponding eigenvector, the eigenvalue problem

$$\left(-\Delta + \frac{q_j(\theta)}{r^2}\right) \left(r^{\frac{1-n_j}{2}}u(r)\right) \otimes \zeta(\theta) = 0$$

reduces to the Euler equation

$$-u''(r) + \frac{\nu^2 - 1/4}{r^2} u(r) = 0.$$
(4.22)

Consider now for any $\nu \ge 0$ or $i\nu > 0$ the Dirichlet problem

$$u$$
 fulfills (4.22) for $r \ge 1$, and $u(1) = 0$. (4.23)

The regular solution is

$$\phi_{\nu}(r) = \begin{cases} \frac{r^{1/2+\nu}-r^{1/2-\nu}}{2\nu} & \text{for } \nu \neq 0, \\ r^{1/2}\ln r & \text{for } \nu = 0 \end{cases};$$

note that indeed $\phi_{\nu}(1) = 0$. The *outgoing solution* to (4.22) (also defined for $r \ge 1$) is

$$\psi_{\nu}(r) = r^{1/2-\nu};$$

note that indeed this is 'outgoing' for ν complex. These two solutions form a fundamental system, and we can define a Green's function by

$$R_{\nu}(r,r') = \phi_{\nu}(r_{<})\psi_{\nu}(r_{>}); \quad r_{<} = \min\{r,r'\}, \ r_{>} = \max\{r,r'\}.$$

Its formal adjoint $R^*_{\nu}(r, r') = \phi_{\nu}(r_{<})\overline{\psi_{\nu}(r_{>})}$ is also a Green's function. This means that for all sufficiently decaying functions v = v(r)

$$-u''(r) + \frac{\nu^2 - 1/4}{r^2} u(r) = v(r), \text{ where } u(r) = \int_1^\infty R_\nu(r, r') v(r') \, \mathrm{d}r',$$

and similarly for R_{ν}^{*} .

Let $\chi_1(r) = \chi(r < 8)$ and $\chi_2(r) = \chi(r > 8)$ form a quadratic partition of unity of smooth non-negative functions on \mathbb{R} , $\chi_1(r)^2 + \chi_2(r)^2 = 1$, such that χ_1 is supported in $(-\infty, 8)$ and $\chi_1(r) = 1$ for r < 4.

Let $w_j = W_j$, $h_j = p_j^2 + w_j$ and v = V (this notation conforms with the previous subsections). We easily check that $v \in \mathcal{L}(\mathcal{H}_s^0, \mathcal{H}_{3+s}^0)$ for $s \in \mathbb{R}$, cf. (2.58c). Let

$$G_{+} = \bigoplus_{j=1}^{2} \left(\chi_{1}(r_{j})(h_{j} - i)^{-1} \chi_{1}(r_{j}) + \sum_{\nu \in \sigma_{j}} \left(\chi_{2}(r_{j})r_{j}^{\frac{1-n_{j}}{2}} R_{j,\nu}r_{j}^{\frac{n_{j}-1}{2}} \chi_{2}(r_{j}) \right) \otimes P_{j,\nu} \right),$$

$$G_{+}^{*} = \bigoplus_{j=1}^{2} \left(\chi_{1}(r_{j})(h_{j} + i)^{-1} \chi_{1}(r_{j}) + \sum_{\nu \in \sigma_{j}} \left(\chi_{2}(r_{j})r^{\frac{1-n_{j}}{2}} R_{j,\nu}^{*}r^{\frac{n_{j}-1}{2}} \chi_{2}(r_{j}) \right) \otimes P_{j,\nu} \right),$$

$$(4.24)$$

where $r_j = |x_j|$ and $R_{j,\nu}$ has a formal interpretation as an (unbounded) operator on $L_I^2 := L^2(I; dr), I := [1, \infty)$. Phrased differently the powers of r are isometrically identifying the spaces L_I^2 and $L^2(I; r^{n_j-1}dr)$; note that the latter space appears naturally for tensor decompositions in spherical coordinates. The operators G_+ and G_+^* are parametrices for $-E_{\mathcal{H}}(\lambda_0)$ in the sense of the following lemma. Of course these operators are not the only options for a parametrix construction. Under stronger conditions, we consider in fact a simpler parametrix in Section 5.1.2.

Introduce also the numbers

$$\nu_0 = \min_{j \in \{1,2\}} \min_{\nu \in \sigma_j} \operatorname{Re} \nu, \qquad s_0 = 1 + \nu_0.$$
(4.25)

In the fastly decaying case, $q_j = 0$ for j = 1, 2, these numbers are explicit dimensional depending constants, cf. (4.67). For example $\nu_0 = \frac{1}{2}$ for $n_1 = n_2 = 3$ in this case (to be considered in Section 5.1.2).

Lemma 4.5. 1) The operators

$$G_{+}, G_{+}^{*} \in \mathcal{L}(\mathcal{H}_{s'}^{-1}, \mathcal{H}_{-s}^{1}) \text{ for } s, s' > 1 - \nu_{0} \text{ with } s + s' > 2,$$

$$(4.26)$$

$$K_{+}^{*} := -E_{\mathcal{H}}^{*}(\lambda_{0})G_{+}^{*} - 1 \in \mathcal{C}(\mathcal{H}_{s}^{-1}) \text{ for any } s \in (1 - \nu_{0}, 2 + \nu_{0}); \qquad (4.27)$$

here $E^*_{\mathcal{H}}(\lambda_0) = E_{\mathcal{H}}(\lambda_0)$ is given the distributional meaning, and we define $K^+ = (K^*_+)^* \in \mathcal{C}(\mathcal{H}^1_{-s})$. Explicitly

$$K^{+} = G_{+}v + \bigoplus_{j=1}^{2} \left(\chi_{1}(h_{j} - i)^{-1} (i\chi_{1} + [\chi_{1}, p_{j}^{2}]) + \sum_{\nu \in \sigma_{j}} \left((\chi_{2}r^{\frac{1-n_{j}}{2}}R_{j,\nu}r^{\frac{n_{j}-1}{2}}\chi_{2}) \otimes P_{j,\nu} \right) B_{j} + \sum_{\nu \in \sigma_{j}} \left(\chi_{2}r^{\frac{1-n_{j}}{2}}R_{j,\nu}r^{\frac{n_{j}-1}{2}}[\chi_{2}, p_{j}^{2}] \right) \otimes P_{j,\nu} \right).$$

$$(4.28)$$

- 2) $K^+f = -G_+E_{\mathcal{H}}(\lambda_0)f f$ for every $f \in \mathcal{H}^1_{-s_0}$ such that $E_{\mathcal{H}}(\lambda_0)f \in \mathcal{H}^{-1}_s$ for some $s > 1 \nu_0$.
- 3) For any s > 1 the operator $G_+ \in \mathcal{L}(\mathcal{H}_s^{-1}, \mathcal{H}_{-s}^1)$ is injective, and

Im
$$G_{+} = \frac{G_{+} - G_{+}^{*}}{2i} \ge \bigoplus_{j=1}^{2} \left(\chi_{1} (h_{j} + i)^{-1} (h_{j} - i)^{-1} \right) \chi_{1} \ge 0.$$
 (4.29)

Proof. <u>I</u>. Let s, s' be given as in (4.26). We show that $r^{-s}R_{j,\nu}r^{-s'}$ is bounded on L_I^2 with a bound independent of $\nu \in \sigma_j$. The Hilbert-Schmidt criterion works. We note the bounds (here for $\nu \neq 0$ only; the case $\nu = 0$ is simpler)

$$\begin{split} &\int_{1}^{\infty} \mathrm{d}r |\nu|^{-2} r^{1+2\operatorname{Re}\nu-2s} \int_{r}^{\infty} r'^{(1-2\operatorname{Re}\nu-2s')} \mathrm{d}r' \leq C |\nu|^{-2} (s' + \operatorname{Re}\nu - 1)^{-1}, \\ &\int_{1}^{\infty} \mathrm{d}r |\nu|^{-2} r^{1-2\operatorname{Re}\nu-2s} \int_{1}^{r} r'^{(1+2\operatorname{Re}\nu-2s')} \mathrm{d}r' \\ &\leq C |\nu|^{-2} \begin{cases} (s + \operatorname{Re}\nu - 1)^{-1} (s' - \operatorname{Re}\nu - 1)^{-1} & \text{for } s' > \operatorname{Re}\nu + 1, \\ (\operatorname{Re}\nu + 1 - s')^{-1} & \text{for } s' < \operatorname{Re}\nu + 1, \\ \int_{1}^{\infty} r^{1-2\operatorname{Re}\nu-2s} \ln r \, \mathrm{d}r & \text{for } s' = \operatorname{Re}\nu + 1, \end{cases}$$

This gives in particular the operator bound $\mathcal{O}(|\nu|^{-3/2})$ for $\operatorname{Re}\nu \to \infty$. Similarly $\partial_r r^{1-s} R_{j,\nu} r^{-s'}$ is bounded with the bound $\mathcal{O}(|\nu|^{-1/2})$. This leads to

$$\left(-\Delta_j + \frac{q_j(\theta)}{r^2} \right) \left(\left(\chi_2 r^{\frac{1-n_j}{2}} R_{j,\nu} r^{\frac{n_j-1}{2}} \chi_2 \right) \otimes P_{j,\nu} \right)$$

= $\left([p_j^2, \chi_2] r^{\frac{1-n_j}{2}} R_{j,\nu} r^{\frac{n_j-1}{2}} \chi_2 \right) \otimes P_{j,\nu} + \chi_2^2 \otimes P_{j,\nu} \in \mathcal{L}(L_{s'}^2, L_{-s}^2),$

where the first term is bounded by $\mathcal{O}(|\nu|^{-1/2})$. By using the ellipticity of $-\Delta_j + \frac{q_j(\theta)}{r^2}$ and interpolation we then obtain that

$$\left(\chi_2 r^{\frac{1-n_j}{2}} R_{j,\nu} r^{\frac{n_j-1}{2}} \chi_2\right) \otimes P_{j,\nu} \in \mathcal{L}(H_{s'}^{-1}, H_{-s}^1)$$

76

with the bound $\mathcal{O}(|\nu|^0)$.

Since the infinite sum defining G_+ is an orthogonal direct sum decomposition (and $\chi_1(h_j - i)^{-1}\chi_1 \in \mathcal{L}(H_{s'}^{-1}, H_{-s}^1)$, obviously) we conclude (4.26).

<u>II</u>. The compactness assertion (4.27) follows by first computing K_+^* , estimating as in Step I (for a good choice of parameters) and then invoking compactness on 'each ν -sector' (obtained by the Hilbert-Schmidt criterion) and the $\mathcal{O}(|\nu|^{-1/2})$ bound (also from I). The requirement $s \in (1 - \nu_0, 2 + \nu_0)$ is dictated by the Hilbert-Schmidt criterion. More precisely fixing any such s we need to find $s' \in \mathbb{R}$ such that, most importantly, $r^{-s'}R_{j,\nu}^*r^{-s}$ and $r^{s-3}R_{j,\nu}^*r^{-s}$ are bounded on L_I^2 . Due to (4.26) it suffices to have

$$s' > 1 - \nu_0$$
, $s + s' > 2$ and $s - 3 + s' \le 0$.

These requirements are fulfilled with s' = 3-s, and we can use that $||r^{s-3}R_{j,\nu}^*r^{-s}|| \to 0$ for $\nu \to \infty$. We have shown 1).

<u>*III*</u>. We show 2) by mimicking the proof of Theorem 4.2. For any given $f \in \mathcal{H}^1_{-s_0}$ with $E_{\mathcal{H}}(\lambda_0)f \in \mathcal{H}^{-1}_s$ for some $s > 1 - \nu_0$ we apply G_+ to $E_{\mathcal{H}}(\lambda_0)f$ and do an integration by parts. More precisely we compute as follows for any $g \in \mathcal{H}^0_{\infty}$ using the smooth cut-off argument of Step *II* of the proof of Theorem 4.2 (the integration by parts) in the third step below:

$$\langle g, G_+ E_{\mathcal{H}}(\lambda_0) f \rangle = \langle G_+^* g, E_{\mathcal{H}}(\lambda_0) f \rangle = \lim_{R \to \infty} \langle G_+^* g, \chi_R E_{\mathcal{H}}(\lambda_0) f \rangle$$

= $\langle -K_+^* g - g, f \rangle = \langle g, -K^+ f - f \rangle,$

yielding 2) by a density argument. This argument uses conveniently that $vf \in \mathcal{H}^{-1}_{(3-s_0)^-}$, which in turn follows from (2.56).

<u>*IV*</u>. For the assertion (4.29) it suffices to show that $\text{Im } R_{j,\nu} \ge 0$. For $\nu \ge 0$ the kernel of $R_{j,\nu}$ is real and symmetric, whence $\text{Im } R_{j,\nu} = 0$ in that case. If $\nu = -i\sigma$ where $\sigma > 0$ the kernel is

$$\left(\operatorname{Im} R_{j,\nu}\right)(r,r') = \sigma \phi_{\nu}(r_{<})\phi_{\nu}(r_{>}),$$

and since ϕ_{ν} is real, it then follows that $\operatorname{Im} R_{j,\nu} \geq 0$ also in that case.

If $G_+f = 0$, $f = (f_1, f_2)$, then (4.29) yields $\chi_1 f = 0$. Whence

$$\left(\chi_2 r^{\frac{1-n_j}{2}} R_{j,\nu} r^{\frac{n_j-1}{2}}\right) \otimes P_{j,\nu} \chi_2 f_j = 0 \text{ for all } \nu \in \sigma_j.$$

Since $R_{j,\nu}(r, r')$ is a Green's function it follows from these formulas that also $\chi_2 f = 0$. Therefore in turn f = 0, and 3) is shown.

Remark. The space $\mathcal{H}_{-s_0}^1$ in 2) is not optimal for the assertion. It suffices to require that $f_j \in \mathcal{B}_{s_0,0}^*(\mathbf{X}_j)$, j = 1, 2. On the other hand if one uses a slightly smaller space than below in the case $s_0 = 1$ to define the notion of a resonance, viz. $\mathcal{H}_{(-1)^+}^1 = \bigcup_{s>-1} \mathcal{H}_s^1$ rather than \mathcal{H}_{-1}^1 , one avoids the special treatment of the case $s_0 = 1$ in Theorem 4.7 3) and 4).

Definition 4.6. (1) λ_0 is called a resonance of H if the equation $(H - \lambda_0)u = 0$ admits a solution $u \in H^1_{-s_0} \setminus H^1$. Such solution is a resonance state of H. The multiplicity of the resonance λ_0 is defined as the dimension, say denoted by n_{res} , of the quotient space

$$\ker(H - \lambda_0)_{|H^1_{-s_0}} / \ker(H - \lambda_0)_{|H^1}$$

(2) 0 is called a resonance of $E_{\mathcal{H}}(\lambda_0)$ if the equation $E_{\mathcal{H}}(\lambda_0)f = 0$ has a solution $f \in \mathcal{H}^1_{-s_0} \setminus \mathcal{H}^1$. Such solution is a resonance state of $E_{\mathcal{H}}(\lambda_0)$. The multiplicity of the resonance 0 is defined as the dimension of the quotient space

$$\ker E_{\mathcal{H}}(\lambda_0)_{|\mathcal{H}^1_{-\infty}} / \ker E_{\mathcal{H}}(\lambda_0)_{|\mathcal{H}^1}.$$

It follows from Lemma 4.1 (applied with $s = s_0$ and s = 0) that λ_0 is a resonance (resp., an eigenvalue) of H if and only if 0 is a resonance (resp., an eigenvalue) of $E_{\mathcal{H}}(\lambda_0)$ and their multiplicities are the same.

We introduce for j = 1, 2,

$$\sigma_{j,k} = \sigma_j \setminus (k, \infty) \text{ and } \sigma_{j,k}^+ = \sigma_j \cap (0, k]; \quad k \in \mathbb{N}.$$

We take an orthonormal basis in ran $P_{j,\nu}$ for each $\nu \in \sigma_j$, say $\zeta_{j,\nu}^{(1)}, \ldots, \zeta_{j,\nu}^{(n_{j,\nu})}$. The set of resonances is partially determined by the set $\sigma_{j,1}^+$, as the following result shows.

Theorem 4.7. Suppose the conditions of Proposition 2.12 and (4.20) for j = 1, 2.

- 1) The dimension of the space of vectors $u \in H^1_{-s_0}$ solving $(H \lambda_0)u = 0$ is finite. If a vector $u \in H^1_{-s_0}$ obeys $(H - \lambda_0)u = 0$ then $f = T^*u \in \mathcal{H}^1_{-s_0}$ and $E_{\mathcal{H}}(\lambda_0)f = 0$, and conversely, if $f \in \mathcal{H}^1_{-s_0}$ obeys $E_{\mathcal{H}}(\lambda_0)f = 0$ then $u = E_+(\lambda_0)f \in H^1_{-s_0}$ and $(H - \lambda_0)u = 0$.
- 2) A vector $f \in \mathcal{H}^1_{-s_0}$ obeys $E_{\mathcal{H}}(\lambda_0)f = 0$ if and only if $f \in \mathcal{H}^1_{(\nu_0-1)^-}$ and $f \in \ker(1+K^+)$; here K^+ is given by (4.28) (as an operator on \mathcal{H}^1_{-s} for any $s \in (1-\nu_0, 2+\nu_0)$).
- 3) Suppose $f \in \mathcal{H}^1_{-s_0}$ and $E_{\mathcal{H}}(\lambda_0)f = 0$. In the case $s_0 = 1$ suppose in addition that $f \in \mathcal{H}^1_{-s}$ for some s < 1. Then the components of $f = (f_1, f_2)$ can be decomposed as

$$-f_{j}(r_{j}\theta_{j}) = \sum_{\nu \in \sigma_{j,1}} \sum_{k=1}^{n_{j,\nu}} l_{j,\nu,k}(f) r_{j}^{\frac{2-n_{j}}{2}-\nu} \chi_{2}(r_{j}) \zeta_{j,\nu}^{(k)}(\theta_{j}) + g_{j},$$

where $g_{j} \in L^{2}$ and
 $l_{j,\nu,k}(f) = \langle \phi_{\nu}(|y_{j}|) |y_{j}|^{-\frac{n_{j}-1}{2}} \otimes \zeta_{j,\nu}^{(k)}, \tilde{f}_{j}(y_{j}) \rangle_{L^{2}(\mathrm{d}y_{j})};$
 $\tilde{f}_{j} = \chi_{2}(vf)_{j} + \chi_{2}B_{j}f_{j} + [\chi_{2}, p_{j}^{2}]f_{j} (\in L^{2}_{3-s_{0}}).$

$$(4.30)$$

Here

$$l_{j,\nu,k}(f) = 0 \text{ for } i\nu \ge 0.$$
 (4.31)

In particular

j

$$f \in \mathcal{H} \quad \Longleftrightarrow \quad \forall j = 1, 2, \nu \in \sigma_{j,1}^+, k = 1, \dots, n_{j,\nu} :$$
$$l_{j,\nu,k}(f) = 0.$$
(4.32)

4) Suppose $u \in H^1_{-s_0}$ and $(H - \lambda_0)u = 0$. In the case $s_0 = 1$ suppose in addition that $u \in H^1_{-s}$ for some s < 1. Then

$$u \in L^{2}(\mathbf{X}) \quad \Longleftrightarrow \quad \forall j = 1, 2, \nu \in \sigma_{j,1}^{+}, k = 1, \dots, n_{j,\nu} :$$
$$l_{j,\nu,k}(T^{*}u) = 0.$$
(4.33)

In particular if $s_0 \neq 1$, then the multiplicity n_{res} of the resonance of H at λ_0 (if existing) is bounded by

$$n_{\rm res} \le \sum_{j=1}^{2} \sum_{\nu \in \sigma_{j,1}^{+}} n_{j,\nu}.$$
 (4.34)

For $s_0 = 1$ the bound (4.34) is valid provided the left-hand side is replaced by the dimension of the quotient space $\ker(H - \lambda_0)_{|H_{(-1)+}^1} / \ker(H - \lambda_0)_{|H^1}$.

Proof. <u>I</u>. For the second part of 1) we refer to Lemma 4.1. To show the first part of 1) we use this correspondence and study the equation $E_{\mathcal{H}}(\lambda_0)f = 0$ with $f \in \mathcal{H}^1_{-s_0}$. By Lemma 4.5 2) we can write $-f = G_+ E_{\mathcal{H}}(\lambda_0)f + K^+ f = K^+ f$, and we recall that $K^+ \in \mathcal{C}(\mathcal{H}^1_{-s})$ for any $s \in (1 - \nu_0, 2 + \nu_0)$. In particular $f \in \mathcal{H}^1_{-s'}, K^+ \in \mathcal{C}(\mathcal{H}^1_{-s'})$ and $f + K^+ f = 0$ for $s' = 3/2 + \nu_0$. Whence it follows from Fredholm theory that the dimension of the space of functions $f \in \mathcal{H}^1_{-s_0}$ solving $E_{\mathcal{H}}(\lambda_0)f = 0$ is finite. We have shown 1).

<u>II</u>. To show the 'only if part' of 2) we consider any $f \in \mathcal{H}^1_{-s_0} \cap \ker(1+K^+)$. Due to Step I it suffices to show that $f \in \mathcal{H}^1_{(\nu_0-1)^-}$. This property is trivially fulfilled for $s_0 = 1$. If $s_0 > 1$ we argue as follows. By using (4.28) we conclude that $f \in \mathcal{H}^1_{-s_1}$ for any $s_1 > \max\{s_0 - 1, 1 - \nu_0\}$, and we are done if $s_0 \leq 2 - \nu_0$. In general we pick the smallest $k \in \mathbb{N}$ such $s_0 - k \leq 1 - \nu_0$ and note that after k - 1 iterations of the above argument we get $f \in \mathcal{H}^{1}_{-s_k}$ for any $s_k > \max\{s_0 - k, 1 - \nu_0\} = 1 - \nu_0$.

To show the 'if part' of 2) suppose $f \in \mathcal{H}^1_{(\nu_0-1)^-} \cap \ker(1+K^+)$. We need to show that $E_{\mathcal{H}}(\lambda_0)f = 0$. By an explicit calculation using (4.28) it follows that $E_{\mathcal{H}}(\lambda_0)f = -E_{\mathcal{H}}(\lambda_0)K^+f \in \mathcal{H}^{-1}_s$ for all $s < 2 + \nu_0$, in particular for some $s > 1 - \nu_0$. Whence by Lemma 4.5 2) we have $-G_+E_{\mathcal{H}}(\lambda_0)f = f + K^+f = 0$. Invoking Lemma 4.5 3) we then conclude that $E_{\mathcal{H}}(\lambda_0)f = 0$.

<u>III</u>. We show (4.30) for any given $f \in \mathcal{H}^1_{-s_0}$ obeying $E_{\mathcal{H}}(\lambda_0)f = 0$. If $\nu_0 = 0$ by assumption $f \in \mathcal{H}^1_{-s}$ for some s < 1. If $\nu_0 > 0$ we know from Step II that $f \in \mathcal{H}^1_{-s}$ for all $s > 1 - \nu_0$. Whence

for some
$$s < 1$$
: $f \in \mathcal{H}^1_{-s}$. (4.35)

Since v and B_j appearing in the definition of $l_{j,\nu,k}(f)$ in (4.30) are of order r^{-3} , we conclude that

for some
$$s < 1$$
 and $j = 1, 2$: $\tilde{f}_j \in L^2_{3-s}$. (4.36)

On the other hand for $\nu \in \sigma_{j,1}$ the vector $\chi_2(|y_j|)\phi_{\nu}(|y_j|)|y_j|^{-\frac{n-1}{2}} \otimes \zeta_{j,\nu}^{(k)} \in H^1_{(-2)^-}$, so $l_{j,\nu,k}(f)$ is well-defined. As we saw in the proof of Lemma 4.5 the operator $r^{-s}R_{j,\nu}r^{-s'}$ is bounded on L_I^2 with a bound independent of $\nu \in \sigma_j$ provided $s, s' > 1 - \operatorname{Re}\nu$ and s + s' > 2. In particular if $\nu > 1$ we can take s = 0 and conclude boundedness of $R_{j,\nu}r^{-s'}$ for any s' > 2. In combination with (4.36) we conclude that the terms in the expansion of $-K_+f$ corresponding to $\nu > 1$ sum up to a vector in L^2 .

It remains to consider the contributions from $\nu \in \sigma_{j,1}$. We examine the term in K_+f corresponding to any fixed $j, \nu \in \sigma_{j,1}$ and $k = 1, \ldots, n_{j,\nu}$. Write it as

$$\left\langle \zeta_{j,\nu}^{(k)}, \left(\chi_2 r^{\frac{1-n_j}{2}} R_{j,\nu} r^{\frac{n_j-1}{2}} \tilde{f}_j \right) (r_j, \cdot) \right\rangle_{L^2(\mathbb{S}_j)} \otimes \zeta_{j,\nu}^{(k)}$$

= $r_j^{\frac{1-n_j}{2}} \chi_2(r_j) \left(l_{j,\nu,k}(f) \psi_{\nu}(r_j) + f_{j,\nu,k}(r_j) \right) \otimes \zeta_{j,\nu}^{(k)},$

where

$$f_{j,\nu,k}(r) = \phi_{\nu}(r) \int_{r}^{\infty} \psi_{\nu}(r') \left\langle \zeta_{j,\nu}^{(k)}, r'^{\frac{n_{j}-1}{2}} \tilde{f}_{j}(r', \cdot) \right\rangle_{L^{2}(\mathbb{S}_{j})} \mathrm{d}r' - \psi_{\nu}(r) \int_{r}^{\infty} \phi_{\nu}(r') \left\langle \zeta_{j,\nu}^{(k)}, r'^{\frac{n_{j}-1}{2}} \tilde{f}_{j}(r', \cdot) \right\rangle_{L^{2}(\mathbb{S}_{j})} \mathrm{d}r'.$$

Using the Cauchy Schwarz inequality and (4.36) we obtain that

for $r \ge 1$ and for some $s < 1 : |f_{j,\nu,k}(r)| \le Cr^{s-3/2}$.

In particular we see that the function g given by $g(r\theta) = r^{\frac{1-n_j}{2}} \chi_2(r) f_{j,\nu,k}(r) \zeta_{j,\nu}^{(k)}(\theta)$ is in L^2 .

We have shown (4.30). Obviously (4.31) and (4.32) are consequences of (4.30) and (4.35).

<u>IV</u>. As for 4) we note that (4.33) follows from 1), 3) and Lemma 4.1, and that the remaining statements of 4) follow from (4.33).

Remark 4.8. In this section we imposed the conditions of Proposition 2.12 and considered three different cases of asymptotics of the effective inter-cluster interaction W_j , j = 1, 2. For symplicity we did not consider cases where these asymptotics mix. Due to the diagonal structure exhibited in Proposition 2.12 it is easy to see that the diagonal parametrix construction works in such cases too. Thus for example it could be the case where W_1 is determined by a I_1 fulfilling (4.6) while W_2 fulfills (4.20). Then the first diagonal element of the parametrix should be the one in (4.9a) (taken with plus) and the second element should be the operator G_+ appearing in Lemma 4.5. By using an analogue of (4.14) it follows by a similar iteration scheme as the ones used before that the resonance states of $E_{\mathcal{H}}(\lambda_0)$ (defined similarly in this case) have a similar structure as in Theorem 4.7 3) and now constitute a subspace of $\mathcal{H}^{1,1}_{\infty,-s_0}$, where s_0 is defined as in (4.25) in terms of the (assumed) asymptotics of W_2 .

Remark 4.9. Since we imposed the conditions of Proposition 2.12 we did not treat the exceptional cases discussed in Sections 2.4 and 2.5. This is an additional technical issue that can be treated essentially by the same parametric construction as the one discussed above. We omit the details, referring the reader to Section 4.3 for a treatment of models of physics.

4.2 The case $\lambda_0 > \Sigma_2$

Here we investigate the case where $\lambda_0 > \Sigma_2$. We need a replacement $R'(\lambda_0) \rightarrow R'(\lambda_0 \pm i0)$ in the definition of $E_{\mathcal{H}}(\lambda_0)$ in Proposition 2.12 (or rather in Remark 2.13), giving rise to the notation $E_{\mathcal{H}}^{\pm}(\lambda_0)$. For that (and for other purposes) we need various statements from Chapter 3. The basic structure of the resolvent is

$$R'(z) = \breve{R}(z) - (p_1^2 \Pi_1 + p_2^2 \Pi_2 - z)^{-1} \Pi,$$

$$R'(z) = \Pi' \breve{R}(z) \Pi' = \Pi' \breve{R}(z) = \breve{R}(z) \Pi'; \quad \breve{R}(z) = (\breve{H} - z)^{-1}.$$
(4.37a)

We shall use the following properties that are parallel to (4.9a) and (4.9b). See Corollary 3.19 for the non-multiple two-cluster case and Remark 3.20 for the present multiple case. There exists the strong weak-star limits

s-w^{*}-lim
$$\breve{R}(\lambda_0 \pm i\epsilon) = \breve{R}(\lambda_0 \pm i0) \in \mathcal{L}(\mathcal{B}_{1/2}(\mathbf{X}), \mathcal{B}_{1/2}^*(\mathbf{X})),$$
 (4.37b)

and if $\breve{R}(\lambda_0 + i0)f = \breve{R}(\lambda_0 - i0)f$ for a given $f \in L^2_t$ for some t > 1/2, then

$$\ddot{R}(\lambda_0 + \mathrm{i}0)f = \ddot{R}(\lambda_0 - \mathrm{i}0)f \in L^2_{t-1}.$$
(4.37c)

We shall use the operators $h_j = p_j^2 + w_j$, j = 1, 2, from Subsections 4.1.2– 4.1.4. Recall that $w_j - W_j$ are real polynomially decreasing potentials with varying meaning reflecting the different hypotheses (4.6), (4.15) and (4.20), respectively. In this section we modify each of the three subsections to cover the case $\lambda_0 > \Sigma_2$. For each case we write (4.2) as

$$-E_{\mathcal{H}}^{\pm}(\lambda_0) = h_1 \oplus h_2 + v^{\pm} \in \mathcal{L}(\mathcal{H}^1, \mathcal{H}^{-1}), \qquad (4.38)$$

where the only new feature is that the previous v is replaced by operators v^{\pm} defined in terms of the two limits $R'(\lambda_0 \pm i0)$, respectively. As before they are non-local, but now no longer symmetric. Due to to (4.37b) they are roughly of order $\mathcal{O}(|x|^{-1-2\rho})$, and a priori they do not have 'good commutation properties' when commuting with multiplication operators (which is in contrast to the lowest threshold setup of Proposition 2.12). More precisely we need the following version of (2.48) and (2.57)

$$-E_{\mathcal{H}}^{\pm}(\lambda_0) \equiv h_1 \oplus h_2 + (K_{ij}^{\pm}(\lambda_0))_{i,j \le 2}, \qquad (4.39)$$

where the difference is not only polynomially decreasing, but in fact also symmetric (to be used in (4.41)). Due to (4.37b) there are bounds

$$K_{ij}^{\pm}(\lambda_0) \in \mathcal{L}\left(L_s^2(\mathbf{X}_j), L_t^2(\mathbf{X}_i)\right) \text{ for all } s > -1/2 - \rho \text{ and } t < 1/2 + \rho, \qquad (4.40)$$

and we note that

$$\mp \operatorname{Im} v^{\pm} = \mp \operatorname{Im}(K_{ij}^{\pm}(\lambda_0))_{i,j\leq 2} \ge 0,$$

$$\mp \operatorname{Im} \sum_{i,j\leq 2} \langle f_i, K_{ij}^{\pm}(\lambda_0) f_j \rangle = \pm \langle F, \operatorname{Im} R'(\lambda_0 \pm \mathrm{i}0) F \rangle;$$

$$F := \sum_{i\leq 2} F_i, \quad F_i = I_i \varphi_i \otimes f_i.$$

(4.41)

4.2.1 Extended eigentransform for $\lambda_0 > \Sigma_2$

Since the operator $\mathring{R}(\lambda_0 \pm i0)\Pi'$ cannot be expected to preserve weighted spaces for $\lambda_0 > \Sigma_2$ (as in (2.56)) we need to examine carefully the 'eigentransform' of Remark 2.3, cf. Lemma 4.1.

First we note the following modification of (4.5),

$$E_{+}^{\pm}(\lambda_{0}) = \left(1 - \ddot{R}(\lambda_{0} \pm \mathrm{i}0)\right)\Pi'(H - \lambda_{0})S \in \mathcal{L}(\mathcal{H}_{s}^{1}, H_{t}^{1})$$

provided $s > -1/2 - \rho, t < -1/2$ and $t \leq s,$ (4.42)

cf. the proof of Lemma 4.1. We shall use a Besov space version of (4.42), in fact we shall need (cf. the proof of Lemma 4.1) that

$$\langle -\Delta \rangle^{1/2} \breve{R}(\lambda_0 \pm i0) \Pi'(H - \lambda_0) S \in \mathcal{L}(\mathcal{B}, \mathcal{B}^*_{1/2}(\mathbf{X})); \mathcal{B} := \mathcal{B}_{1/2}(\mathbf{X}_1) \oplus \mathcal{B}_{1/2}(\mathbf{X}_2).$$

$$(4.43)$$

The following Lemma 4.10 holds for the cases treated in Subsections 4.1.2–4.1.4, in fact the setting is that of Proposition 2.12 (now with $\lambda_0 > \Sigma_2$). We introduce $s'_0 = 1/2 + \rho'/4$, where $0 < \rho' < 4\rho$, $\rho' \leq 2$ and

$$|W_j(y)| \le C\langle y \rangle^{-\rho'}$$
 for $|y| \ge R$; *R* large. (4.44)

If $\rho = 1$ (our main interest) the condition simplify as (4.44) for some $\rho' \leq 2$, and in that case we would choose $\rho' = 1$ and $\rho' = 2$ in Subsections 4.2.2 and 4.2.4, respectively. On the other hand Subsection 4.2.3 does not involve (4.44) and Lemma 4.10 directly, but rather a version of Lemma 4.10 stated below as Lemma 4.11. We regard Lemmas 4.10 and 4.11 as substitutes for Lemma 4.1. Recall the notation $\mathcal{H}_s = L_s^2(\mathbf{X}_1) \oplus L_s^2(\mathbf{X}_2)$.

Lemma 4.10. Suppose the conditions of Proposition 2.12 (except for assuming now $\lambda_0 > \Sigma_2$). Let $s \leq s'_0$ and $t \in (0, s'_0]$.

1) The map $T^*: \mathcal{E}_{-s}^{\mathcal{G}} \to \mathcal{E}_{-s}^{\mathcal{H}}$, where

$$\mathcal{E}_{-s}^{\mathcal{G}} := \{ u \in L_{-s}^{2}(\mathbf{X}) \mid (H - \lambda_{0})u = 0, \ \Pi' u \in \mathcal{B}_{1/2,0}^{*}(\mathbf{X}) \}, \\ \mathcal{E}_{-s}^{\mathcal{H}} := \{ f \in \mathcal{H}_{-s} \mid E_{\mathcal{H}}^{+}(\lambda_{0})f = 0 \},$$

is a well-defined linear isomorphism with inverse $E^+_+(\lambda_0): \mathcal{E}^{\mathcal{H}}_{-s} \to \mathcal{E}^{\mathcal{G}}_{-s}$.

- 2) The spaces $\mathcal{E}_{-s}^{\mathcal{H}}$ and $\{f \in \mathcal{H}_{-s} \mid E_{\mathcal{H}}^{-}(\lambda_0)f = 0\}$ coincide, and $E_{+}^{-}(\lambda_0)f = E_{+}^{+}(\lambda_0)f$ for all $f \in \mathcal{E}_{-s}^{\mathcal{H}}$.
- 3) The map T^* : ker $(H-\lambda_0) \to \ker E^+_{\mathcal{H}}(\lambda_0)|_{\mathcal{H}}$ is a well-defined linear isomorphism with inverse $E^+_+(\lambda_0)$.
- 4) The map $T^*: \mathcal{E}_{-t,0}^{\mathcal{G}} \to \mathcal{E}_{-t,0}^{\mathcal{H}}$, where

$$\mathcal{E}_{-t,0}^{\mathcal{G}} := \{ u \in \mathcal{B}_{t,0}^{*}(\mathbf{X}) \mid (H - \lambda_{0})u = 0, \ \Pi' u \in \mathcal{B}_{1/2,0}^{*}(\mathbf{X}) \}, \\ \mathcal{E}_{-t,0}^{\mathcal{H}} := \{ f \in \mathcal{B}_{t,0}^{*}(\mathbf{X}_{1}) \oplus \mathcal{B}_{t,0}^{*}(\mathbf{X}_{2}) \mid E_{\mathcal{H}}^{+}(\lambda_{0})f = 0 \},$$

is a well-defined linear isomorphism with inverse $E^+_+(\lambda_0): \mathcal{E}^{\mathcal{H}}_{-t,0} \to \mathcal{E}^{\mathcal{G}}_{-t,0}$

- 5) The spaces $\mathcal{E}_{-t,0}^{\mathcal{H}}$ and $\{f \in \mathcal{B}_{t,0}^{*}(\mathbf{X}_{1}) \oplus \mathcal{B}_{t,0}^{*}(\mathbf{X}_{2}) \mid E_{\mathcal{H}}^{-}(\lambda_{0})f = 0\}$ coincide, and $E_{+}^{-}(\lambda_{0})f = E_{+}^{+}(\lambda_{0})f$ for all $f \in \mathcal{E}_{-t,0}^{\mathcal{H}}$.
- 6) There exists $\sigma < 0$ such that for any real function $\chi_{\sigma} \in C^{\infty}(\mathbb{R})$ which is supported in $(-\infty, \sigma/2)$ and which is 1 on $(-\infty, \sigma)$, $\mathcal{E}_{-s'_0,0}^{\mathcal{G}}$ and the two spaces

$$\mathcal{E}_{-s'_{0},\pm\sigma}^{\mathcal{G},\pm} := \{ u \in \mathcal{B}_{s'_{0},0}^{*}(\mathbf{X}) \mid (H - \lambda_{0})u = 0, \ \Pi' u \in \mathcal{B}_{1/2}^{*}(\mathbf{X}), \\ \chi_{\sigma}(\pm B_{R_{0}})\Pi' u \in \mathcal{B}_{1/2,0}^{*}(\mathbf{X}) \}$$

all coincide. (Here $B = B_{R_0}$ is given by (3.7).)

Proof. <u>I</u>. Let $u \in \mathcal{E}_{-s}^{\mathcal{G}}$ be given. Then $f = T^* u \in \mathcal{H}_{-s}$ follows easily, so to show that $f \in \mathcal{E}_{-s}^{\mathcal{H}}$ it remains to show that $E_{\mathcal{H}}^{\pm}(\lambda_0)f = 0$. Note that $\Pi'(H - \lambda_0)\Pi u \in \mathcal{B}_{1/2}(\mathbf{X})$, cf. Lemma 2.9, so that $R'(\lambda_0 \pm i0)(H - \lambda_0)\Pi u$ is a well-defined element of $\mathcal{B}_{1/2}^*(\mathbf{X})$. We calculate using this fact, (2.14) and (2.31a)

$$E_{\mathcal{H}}^{\pm}(\lambda_0)f = S^*(\lambda_0 - H)ST^*u + S^*HR'(\lambda_0 \pm i0)(H - \lambda_0)ST^*u$$

$$= S^*(\lambda_0 - H)\Pi u + S^*HR'(\lambda_0 \pm i0)(H - \lambda_0)\Pi u$$

$$= S^*(\lambda_0 - H)\Pi u + S^*(H - \lambda_0)\Pi'\breve{R}(\lambda_0 \pm i0)(H - \lambda_0)\Pi u$$

$$= S^*(\lambda_0 - H)\Pi u + \lim_{R \to \infty} S^*(H - \lambda_0)\Pi'\breve{R}(\lambda_0 \pm i0)(H - \lambda_0)\Pi\chi_R u;$$

here $\chi_R(x) = \chi(|x|/R), R \ge 1$, is given in agreement with (1.29). We calculate the second term as follows using the weak-star topology on \mathcal{B}^* where $\mathcal{B} = \mathcal{B}_{1/2}(\mathbf{X}_1) \oplus \mathcal{B}_{1/2}(\mathbf{X}_2)$

$$\lim_{R \to \infty} S^* (H - \lambda_0) \Pi' \breve{R} (\lambda_0 \pm i0) (H - \lambda_0) \Pi \chi_R u$$

$$= - w_{R \to \infty}^* \lim_{R \to \infty} S^* (H - \lambda_0) \Pi' \breve{R} (\lambda_0 \pm i0) (H - \lambda_0) \Pi' \chi_R u$$

$$+ w_{R \to \infty}^* (H - \lambda_0) \Pi' \breve{R} (\lambda_0 \pm i0) (H - \lambda_0) \chi_R u$$

$$= w_{R \to \infty}^* \lim_{R \to \infty} S^* (\lambda_0 - H) \Pi' \chi_R u$$

$$+ w_{R \to \infty}^* \lim_{R \to \infty} S^* (H - \lambda_0) \Pi' \breve{R} (\lambda_0 \pm i0) [H - \lambda_0, \chi_R] u$$

$$= S^* (\lambda_0 - H) \Pi' u$$

$$+ w_{R \to \infty}^* (H - \lambda_0) \Pi' \breve{R} (\lambda_0 \pm i0) [H - \lambda_0, \chi_R] u.$$

Suppose we can show that the second term

$$\underset{R \to \infty}{\text{w}^*-\lim} S^*(H - \lambda_0) \Pi' \breve{R}(\lambda_0 \pm i0) [H - \lambda_0, \chi_R] u = 0.$$
(4.45)

Then we obtain from the above computations the desired result

$$E_{\mathcal{H}}^{\pm}(\lambda_0)f = S^*(\lambda_0 - H)\Pi u + S^*(\lambda_0 - H)\Pi' u$$
$$= S^*(\lambda_0 - H)u$$
$$= 0.$$

So it remains to show (4.45). We note that

$$-[(H - \lambda_0), \chi_R]u = 2\nabla \cdot (\nabla \chi_R) u - (\Delta \chi_R)u$$

and that for all the components ∇_k of ∇ , the operator ∇_k combines with the operator to the left in (4.45), and for this combination we may use (4.43). Keeping this 'effective boundedness' in mind we treat the first term by substituting

$$u = \Pi u + \Pi' u = \Pi_1 u + \Pi_2 u + (\Pi - SS^*)u + \Pi' u.$$

For j = 1, 2

$$\Pi' \nabla \cdot (\nabla \chi_R) \Pi_j u = \Pi' \nabla_{x_j} \cdot (\nabla \chi_R)_{|x^j=0} \Pi_j u + \mathcal{O}(R^{-2}) \Pi u = 0 + \mathcal{O}(R^{-2}) u, \quad (4.46)$$

and therefore the first and second terms do not contribute in the limit. The third term is treated by Lemma 2.9. Similarly

$$\Pi' \nabla \cdot (\nabla \chi_R) \Pi' u = \mathcal{O}(R^{-1}) \Pi' u$$

does not contribute in the limit, and obviously the term $(\Delta \chi_R)u = \mathcal{O}(R^{-2})u$ does not neither. We conclude (4.45).

<u>II</u>. First we note that $E_{\mathcal{H}}^{\pm}(\lambda_0)f$ has a well-defined distributional meaning for $f \in \mathcal{H}_{-s}$ so that $\mathcal{E}_{-s}^{\mathcal{H}}$ is well-defined. We show that $E_{\mathcal{H}}^{+}(\lambda_0)f = 0$ if and only if $E_{\mathcal{H}}^{-}(\lambda_0)f = 0$. Suppose $E_{\mathcal{H}}^{+}(\lambda_0)f = 0$. First we show that

$$\text{Im}\langle f_j, \dot{h}_j f_j \rangle = 0; \quad \dot{h}_j = -\Delta_j + W_j, \ j = 1, 2.$$
(4.47)

Note that $\check{h}_j f_j \in L^2_s$ for any $s < 1/2 + \rho$, $f_j \in L^2_{-s'_0}$ and that $s'_0 < 1/2 + \rho$, so that $\langle f_j, \check{h}_j f_j \rangle$ is well-defined. We compute

$$\chi_R \check{h}_j f_j = \check{h}_j \chi_R f_j + (\Delta \chi_R) f_j + 2(\nabla \chi_R) \cdot \nabla f_j,$$

and then

$$\operatorname{Im}\langle f_j, \check{h}_j f_j \rangle = \lim_{R \to \infty} \operatorname{Im} \langle \chi_R f_j, \chi_R \check{h}_j f_j \rangle = 2 \lim_{R \to \infty} \operatorname{Im} \langle \chi_R f_j, (\nabla \chi_R) \cdot \nabla f_j \rangle$$

Next we estimate as in (4.13)

$$\begin{aligned} R^{2} \| (\nabla \chi_{R}) \cdot \nabla f_{j} \|_{L^{2}(\mathbf{X}_{j})}^{2} \\ &\leq C_{1} \langle p^{2} \rangle_{\chi_{2R} \bar{\chi}_{R/2} f_{j}} \\ &\leq C_{2} \big(\operatorname{Re} \langle \chi_{2R}^{2} \bar{\chi}_{R/2}^{2} f_{j}, \check{h}_{j} f_{j} \rangle + R^{-\rho'} \| \chi_{4R} f_{j} \|^{2} \big) \\ &= C_{2} \big(o(R^{0}) + R^{-\rho'} o(R^{2s'_{0}}) \big) \\ &= o(R^{1-\rho'/2}), \end{aligned}$$

yielding

$$\lim_{R \to \infty} R^{s'_0} \| (\nabla \chi_R) \cdot \nabla f_j \|_{L^2(\mathbf{X}_j)} = 0$$

4.2.1. Extended eigentransform for $\lambda_0 > \Sigma_2$

Moreover

$$\lim_{R \to \infty} R^{-s'_0} \| \chi_R f_j \|_{L^2(\mathbf{X}_j)} = 0.$$

Whence we obtain (4.47) by the Cauchy-Schwarz inequality.

Next we compute by using (4.41) and (4.47) that

$$0 = \operatorname{Im}\langle f, E_{\mathcal{H}}^{+}(\lambda_{0})f \rangle = \operatorname{Im}\langle F, R'(\lambda_{0} + \mathrm{i}0)F \rangle;$$

$$F = \sum_{i \leq 2} F_{i}, \quad F_{i} = I_{i}\varphi_{i} \otimes f_{i}.$$
(4.48)

Whence also $\operatorname{Im}\langle F, R'(\lambda_0 - i0)F \rangle = 0$ and we learn that $R'(\lambda_0 + i0)F = R'(\lambda_0 - i0)F$, and therefore that also $E_{\mathcal{H}}(\lambda_0)f = 0$. We can argue similarly for the other implication.

<u>III</u>. Suppose $f \in \mathcal{E}_{-s}^{\mathcal{H}}$. We need to show that $u := E_{+}^{+}(\lambda_{0})f \in \mathcal{E}_{-s}^{\mathcal{G}}$ and that $T^{*}u = f$. Clearly $Sf \in L_{-s}^{2}(\mathbf{X})$. We noted in Step II that $R'(\lambda_{0} + i0)F = R'(\lambda_{0} - i0)F$ where F is specified in (4.48). This means that $u = E_{+}^{-}(\lambda_{0})f$, and using (4.37a)–(4.37c) we then conclude that $u \in L_{-s}^{2}(\mathbf{X})$ and $\Pi'u \in \mathcal{B}_{1/2,0}^{*}(\mathbf{X})$. To show that $u \in \mathcal{E}_{-s}^{\mathcal{G}}$ it remains to show that $(H - \lambda_{0})u = 0$. We calculate

$$(H - \lambda_0)u = (H - \lambda_0)Sf - (H - \lambda_0)R'(\lambda_0 + i0)HSf$$

= $(H - \lambda_0)Sf - \Pi'(H - \lambda_0)R'(\lambda_0 + i0)(H - \lambda_0)Sf$
 $- \Pi(H - \lambda_0)R'(\lambda_0 + i0)(H - \lambda_0)Sf$
= $(H - \lambda_0)Sf - \Pi'(H - \lambda_0)Sf - \Pi HR'(\lambda_0 + i0)HSf$
= $\Pi(H - \lambda_0)Sf - \Pi HR'(\lambda_0 + i0)HSf.$

We conclude that $(H - \lambda_0)u \in \Pi L^2_{-s}(\mathbf{X})$ (since $\Delta f \in \mathcal{H}_{-s}$) and that

$$S^*(H - \lambda_0)u = -E^+_{\mathcal{H}}(\lambda_0)f = 0,$$

and therefore (since S^* maps injectively on the space $\Pi L^2_{-s}(\mathbf{X})$) that indeed $(H - \lambda_0)u = 0$.

Next we compute

$$T^* E^+_+(\lambda_0) f = T^* S f - T^* R'(\lambda_0 + i0)(H - \lambda_0) S f$$

= f - 0 (since $T^* \Pi' = 0$)
= f.

<u>*IV*</u>. Let $u \in \mathcal{E}_{-s}^{\mathcal{G}}$ and note that we showed in Step *I* that $f := T^* u \in \mathcal{E}_{-s}^{\mathcal{H}}$. Using parts of the proof we show that $u = E_+^+(\lambda_0)f$ as follows.

$$E_{+}^{+}(\lambda_{0})T^{*}u = ST^{*}u - R'(\lambda_{0} + i0)(H - \lambda_{0})ST^{*}u$$

$$= \Pi u - R'(\lambda_{0} + i0)(H - \lambda_{0})\Pi u$$

$$= \Pi u - \lim_{R \to \infty} \Pi' \breve{R}(\lambda_{0} + i0)(H - \lambda_{0})\Pi \chi_{R}u$$

$$= \Pi u + \Pi' u$$

$$= u.$$

This finishes the proof of 1) and 2), and 3) is a special case of 1). The statements 4) and 5) are proved verbatim as 1) and 2).

<u>V</u>. Obviously $\mathcal{E}_{-s'_0,0}^{\mathcal{G}} \subset \mathcal{E}_{-s'_0,\pm\sigma}^{\mathcal{G},\pm\sigma}$. We choose $\sigma < 0$ such that $\bar{\chi}_{\sigma}(B_{R_0})\breve{R}(\lambda_0 - \mathrm{i}0) \in \mathcal{L}(\mathcal{B}_{1/2}(\mathbf{X}), \mathcal{B}^*_{1/2,0}(\mathbf{X})); \quad \bar{\chi}_{\sigma}(B_{R_0}) = (1 - \chi_{\sigma}(B_{R_0})),$

cf. Lemma 3.15. We show the opposite inclusion for $\mathcal{E}_{-s'_0,\sigma}^{\mathcal{G},+}$ only. So let $u \in \mathcal{E}_{-s'_0,\sigma}^{\mathcal{G},+}$ be given. Then we can mimic Step I and see that also in this case $f = T^* u \in \mathcal{E}_{-s'_0,0}^{\mathcal{H}}$. Note that for showing (4.45) we can use the above bound to estimate

$$\begin{split} & \underset{R \to \infty}{\text{w}^*-\lim} S^*(H - \lambda_0) \Pi' \breve{R}(\lambda_0 + \mathrm{i}0) [H - \lambda_0, \chi_R] u \\ &= -2 \underset{R \to \infty}{\text{w}^*-\lim} S^*(H - \lambda_0) \Pi' \breve{R}(\lambda_0 + \mathrm{i}0) \Pi' \nabla \cdot (\nabla \chi_R) \Big(\chi_\sigma(B_{R_0}) + \bar{\chi}_\sigma(B_{R_0}) \Big) \Pi' u \\ &= 0 + 0 \\ &= 0. \end{split}$$

We use the same argument mimicking Step IV, cf. 4). So indeed $u \in \mathcal{E}_{-s'_0,0}^{\mathcal{G}}$, and we have shown that $\mathcal{E}_{-s'_0,\sigma}^{\mathcal{G},+} = \mathcal{E}_{-s'_0,0}^{\mathcal{G}}$.

In the regime $s \in (s'_0, 1/2 + \rho)$ we can define $\mathcal{E}_{-s}^{\mathcal{H}}$ as in Lemma 4.10 1), however we are not able to conclude Lemma 4.10 2) is this case. This leads to the following definition

 $\mathcal{E}_{-s}^{\mathcal{H}} := \{ f \in \mathcal{H}_{-s} | \quad E_{\mathcal{H}}^+(\lambda_0) f = E_{\mathcal{H}}^-(\lambda_0) f = 0 \} \quad \text{for} \quad s < 1/2 + \rho,$

which is consistent with Lemma 4.10. Obviously we can also extend the definition of $\mathcal{E}_{-s}^{\mathcal{G}}$ given in Lemma 4.10 1) to any real s.

Lemma 4.11. Suppose the conditions of Proposition 2.12 (except for assuming now $\lambda_0 > \Sigma_2$). Suppose $s \leq 3/2$ and $s < 1/2 + \rho$. Then the map

 $T^*: \mathcal{E}_{-s}^{\mathcal{G}} \to \mathcal{E}_{-s}^{\mathcal{H}}$

is a well-defined linear isomorphism with inverse $E^+_+(\lambda_0) : \mathcal{E}^{\mathcal{H}}_{-s} \to \mathcal{E}^{\mathcal{G}}_{-s}$ (defined by (4.42)). Moreover $E^-_+(\lambda_0)f = E^+_+(\lambda_0)f$ for all $f \in \mathcal{E}^{\mathcal{H}}_{-s}$.

Proof. We can assume (4.44) and that $s \in (s'_0, 1/2 + \rho)$, in particular that s > 1/2. It follows from Step I of the proof of Lemma 4.10 that T^* maps into $\mathcal{E}_{-s}^{\mathcal{H}}$. Obviously $E_+^+(\lambda_0)$ maps into $L_{-s}^2(\mathbf{X})$, but to show that $E_+^+(\lambda_0)$ in fact maps into $\mathcal{E}_{-s}^{\mathcal{G}}$ we cannot use (4.48), however the following substitute works:

$$0 = \operatorname{Im}\langle f, E_{\mathcal{H}}^{+}(\lambda_{0})f - E_{\mathcal{H}}^{-}(\lambda_{0})f \rangle = 2 \operatorname{Im}\langle F, R'(\lambda_{0} + \mathrm{i}0)F \rangle;$$

$$F = \sum_{i \leq 2} F_{i}, \quad F_{i} = I_{i}\varphi_{i} \otimes f_{i}.$$
(4.49)

This shows that $R'(\lambda_0 + i0)F = R'(\lambda_0 - i0)F$ for all $f \in \mathcal{E}_{-s}^{\mathcal{H}}$, and therefore that $\Pi' E_+^+(\lambda_0) f \in \mathcal{B}_{1/2,0}^*(\mathbf{X})$ and $E_+^-(\lambda_0) f = E_+^+(\lambda_0) f$ for all $f \in \mathcal{E}_{-s}^{\mathcal{H}}$. Next we use Step III to see that also $(H - \lambda_0)E_+^+(\lambda_0)f = 0$ showing that indeed $E_+^+(\lambda_0)$ maps into $\mathcal{E}_{-s}^{\mathcal{G}}$.

Finally it follows from the last part of Step *III* and of Step *IV* that T^* and $E^+_+(\lambda_0)$ are mutually inverses, as we want.

4.2.2 Negative slowly decaying effective potentials

We aim at proving a version of Theorem 4.2 in the setting of Subsection 4.1.2 (except that now $\lambda_0 > \Sigma_2$). In particular $\rho < 2$ and we can use Lemma 4.10 with $s'_0 = s_0 = 1/2 + \rho/4$ (corresponding to taking $\rho = \rho'$).

Theorem 4.12. Under the conditions of Proposition 2.12 (except for assuming now $\lambda_0 > \Sigma_2$) and (4.6) for j = 1, 2

$$\dim \ker(H - \lambda_0) < \infty.$$

Moreover $\mathcal{E}^{\mathcal{G}}_{-s_0,0} \subset H^1_{\infty}$.

Proof. We basically mimic the proof of Theorem 4.2 using Lemma 4.10 as a substitute for Lemma 4.1.

Note that Lemma $4.10\ 3$) implies that

$$\dim \ker(H - \lambda_0) = \dim \ker E_{\mathcal{H}}^{\pm}(\lambda_0), \qquad (4.50)$$

and that the map

$$E^{\pm}_{+}(\lambda_0) : \ker E^{\pm}_{\mathcal{H}}(\lambda_0) \to \ker(H - \lambda_0)$$
 (4.51)

is an isomorphism.

We shall use the notation h_1 , h_2 and $R_{\text{diag}}^{\pm} = r_1(0 \pm i0) \oplus (r_2(0 \pm i0))$ from the proof of Theorem 4.2. Suppose $f \in \ker E_{\mathcal{H}}^{\pm}(\lambda_0)$. Then we write the equations $R_{\text{diag}}^{\pm} E_{\mathcal{H}}^{\pm}(\lambda_0) f = 0$ as

$$(1+K^{\pm})f = 0, \quad K^{\pm} = R^{\pm}_{\text{diag}}v^{\pm},$$
 (4.52)

where 1f arises by writing $f = (f_1, f_2) \in \mathcal{H}^1$ and using that

$$r_j(0\pm i0)h_jf_j = f_j.$$
 (4.53)

Note that it is shown in Step II of the proof of Theorem 4.2 that generally, if $f_j \in \mathcal{B}^*_{s_0,0}(\mathbf{X}_j)$ obeys $h_j f_j \in \mathcal{B}_{s_0}(\mathbf{X}_j)$, then (4.53) holds. In particular (4.53) holds for $f \in \ker E^{\pm}_{\mathcal{H}}(\lambda_0)$.

Next (4.40) and (4.9a) imply that

$$K^{\pm} \in \mathcal{C}(\mathcal{H}_{-s}^1) \text{ for } s \in (s_0, 1/2 + \rho).$$

$$(4.54)$$

Since K^{\pm} is compact (on any such space) it follows from Fredholm theory that dim ker $E_{\mathcal{H}}^{\pm}(\lambda_0) < \infty$, proving the first assertion of the theorem.

By Lemma 4.10 any given $\phi \in \mathcal{E}_{-s_0,0}^{\mathcal{G}}$ corresponds to the vector $f = (f_1, f_2) = T^* \phi \in \mathcal{E}_{-s_0,0}^{\mathcal{H}}$ (obeying $\phi = E_+^+(\lambda_0)f = E_+^-(\lambda_0)f$). Note also that $h_j f_j \in L_s^2(\mathbf{X}_j) \subset \mathcal{B}_{s_0}(\mathbf{X}_j)$ for any *s* given as in (4.54), so by the assertion in Step *II* of the proof of Theorem 4.2 we can conclude that (4.52) is fulfilled. In particular $K^+ f = K^- f$.

A converse assertion is true. We first observe using (4.41), that $\operatorname{Im} v^+ \leq 0$. Since $\operatorname{Im} R_{\text{diag}}^+ \geq 0$ the equation

$$0 = \operatorname{Im}\langle f, v^+ f \rangle + \operatorname{Im}\langle R^+_{\operatorname{diag}}v^+ f, v^+ f \rangle \tag{4.55}$$

for any solution to $(1 + K^+)f = 0$, where $f = (f_1, f_2) \in \mathcal{B}^*_{s_0}(\mathbf{X}_1) \oplus \mathcal{B}^*_{s_0}(\mathbf{X}_2)$, then yields that $v^-f = v^+f$ and that also $(1 + K^-)f = 0$ for any such f. (Similarly a solution to $(1 + K^-)f = 0$ is also a solution to $(I + K^+)f = 0$.) Moreover $E^{\pm}_{\mathcal{H}}(\lambda_0)f = 0$ in the distributional sense indeed yielding a converse statement.

More importantly we can improve the decay of any $f = (f_1, f_2) \in \mathcal{B}^*_{s_0}(\mathbf{X}_1) \oplus \mathcal{B}^*_{s_0}(\mathbf{X}_2)$ solving $(1 + K^+)f = 0$ (and therefore also $(1 + K^-)f = 0$) by invoking repeatedly (4.9b) and (4.37c): Since $f_j \in \mathcal{B}^*_{s_0}(\mathbf{X}_j)$, j = 1, 2, and $v^-f = v^+f$ (as noted above) we deduce from (4.37c) that $v^+f \in \mathcal{H}^{-1}_t$ for any $t < 1 + 2\rho - s_0$. Next using (4.9b) (in combination with (4.55)) we deduce that $f = -K^+f \in \mathcal{H}^1_{\epsilon-s_0}$ for any $\epsilon < \epsilon_0$, where $\epsilon_0 = 1 + 2\rho - 2s_0 = \frac{3}{2}\rho$. Repeating this argument, say k times, yields $f \in \mathcal{H}^1_{t-s_0}$ for any $t < \epsilon_0 k$, and therefore that $f \in \mathcal{H}^1_{\infty}$.

We have shown that for any given $\phi \in \mathcal{E}_{-s_0,0}^{\mathcal{G}}$ the vector $f = (f_1, f_2) = T^* \phi \in \mathcal{H}_{\infty}^1$ and therefore, due to Lemma 4.10 1), indeed $\phi = E_+^+(\lambda_0)f \in H_{\infty}^1$.

4.2.3 Positive slowly decaying effective potentials

We aim at proving a version of Theorem 4.3 in the setting of Subsection 4.1.3 (except that now $\lambda_0 > \Sigma_2$).

Theorem 4.13. Under the conditions of Proposition 2.12 and (4.15) for j = 1, 2,

$$\dim \ker(H - \lambda_0) < \infty$$

Moreover $\mathcal{E}_{-s}^{\mathcal{G}} \subset H_{\infty}^1$ for and real s such that $s \leq 3/2$ and $s < 1/2 + \rho$.

Proof. Due to Lemma 4.11 we need to study the space $\mathcal{E}_{-s}^{\mathcal{H}}$ for any such *s*. Using the notation $\widetilde{R} = h_1^{-1} \oplus h_2^{-1}$ of (4.19) as well as the notation (4.38), we write the equation $\widetilde{R}E_{\mathcal{H}}^+(\lambda_0)f = 0$ for any $f \in \mathcal{E}_{-s}^{\mathcal{H}}$ as

$$(1+\widetilde{K})f = 0; \quad \widetilde{K} = \widetilde{R}v^+. \tag{4.56}$$

Due to Lemma 4.4 we can easily check that $\widetilde{K} \in \mathcal{C}(\mathcal{H}^1_{-s})$. It then follows from Fredholm theory that f belongs to a finite-dimensional subspace of \mathcal{H}^1_{-s} , yielding in particular the first statement of the theorem by taking s = 0.

Finally we show

$$\mathcal{E}_{-s}^{\mathcal{H}} \subset \mathcal{H}_{\infty}^{1}, \tag{4.57}$$

using an iteration procedure as in the proof of Theorem 4.12. Note that due to Lemma 4.11 we can use (4.37c) and (4.56) to improve the decay. We deduce that $f \in \mathcal{H}^1_{-s+\bar{t}}$ given that $f \in \mathcal{H}^1_{-s+(k-1)\bar{t}}$ where $\bar{t} := 1 + 2\rho - \bar{\rho}$. Since there is no limit on $k \in \mathbb{N}$ used for this argument, indeed (4.57) follows by iteration. By Lemma 4.11 and (4.57) it follows that $\mathcal{E}^{\mathcal{G}}_{-s} \subset H^1_{\infty}$.

4.2.4 Homogeneous degree -2 effective potentials

We aim at proving a version of Theorem 4.7 in a setting similar to that of Subsection 4.1.4. Now of course $\lambda_0 > \Sigma_2$, and having the physics examples in mind we demand $\rho = 1$ rather than $\rho \ge 1/2$ as before. It is easy to see that our theory is void if $\rho = 1/2$ is kept. On the other hand there is something to say in the case $\rho \in (1/2, 1)$, but for simplicity of presentation we leave this case out. It turns out that the proof of Theorem 4.7 works again with only minor modifications, in particular we will need Lemma 4.10 2) with $s'_0 = 1$.

First we examine how Lemma 4.5 can be modified for $\lambda_0 > \Sigma_2$. Due to restricted mapping proporties of v^+ we can only obtain the following weakened result (by the same proof). Let $s_1 = \max\{1 - \nu_0, 1/2\}$, and recall that $s_0 = 1 + \nu_0$ in agreement with (4.25).

Lemma 4.14. 1) The operators

$$G_+, G_+^* \in \mathcal{L}(\mathcal{H}_{s'}^{-1}, \mathcal{H}_{-s}^1) \text{ for } s, s' > 1 - \nu_0 \text{ with } s + s' > 2,$$
 (4.58)

$$K_{+}^{*} := -E_{\mathcal{H}}^{+}(\lambda_{0})^{*}G_{+}^{*} - 1 \in \mathcal{C}(\mathcal{H}_{s}^{-1}) \text{ for } s \in (s_{1}, 3/2);$$

$$(4.59)$$

here $E^+_{\mathcal{H}}(\lambda_0)^*$ is given the distributional meaning, and we define $K^+ = (K^*_+)^* \in \mathcal{C}(\mathcal{H}^1_{-s})$. Explicitly

$$K^{+} = G_{+}v^{+} + \bigoplus_{j=1}^{2} \left(\chi_{1}(h_{j} - i)^{-1} (i\chi_{1} + [\chi_{1}, p_{j}^{2}]) + \sum_{\nu \in \sigma_{j}} \left((\chi_{2}r^{\frac{1-n_{j}}{2}}R_{j,\nu}r^{\frac{n_{j}-1}{2}}\chi_{2}) \otimes P_{j,\nu} \right) B_{j} + \sum_{\nu \in \sigma_{j}} \left(\chi_{2}r^{\frac{1-n_{j}}{2}}R_{j,\nu}r^{\frac{n_{j}-1}{2}}[\chi_{2}, p_{j}^{2}] \right) \otimes P_{j,\nu} \right).$$

$$(4.60)$$

- 2) $K^+f = -G_+E^+_{\mathcal{H}}(\lambda_0)f f$ for every $f \in \bigcup_{s < 3/2} \mathcal{H}^1_{-\min\{s,s_0\}}$ with $E^+_{\mathcal{H}}(\lambda_0)f \in \mathcal{H}^{-1}_{(1-\nu_0)^+}$.
- 3) For any s > 1 the operator $G_+ \in \mathcal{L}(\mathcal{H}_s^{-1}, \mathcal{H}_{-s}^1)$ is injective.

Using Lemmas 4.10 and 4.14 (as substitutes for Lemma 4.5) and Lemma 4.11 (as a substitute for Lemma 4.1) we can show the following weakened version of Theorem 4.7. We mimic the proof, yielding immediately the second assertion of Theorem 4.15 1) (stated below) from Lemma 4.11 and then in turn the first one from Lemma 4.14 2). The results 2) and 3) follow partly as before. Note however that Lemma 4.10 2) is used for the second implication of 2). Clearly 4) follows from 1), 3) and Lemma 4.11.

Theorem 4.15. Suppose the conditions of Proposition 2.12 and (4.20) for j = 1, 2. Suppose s < 3/2 and $s \le s_0$.

- 1) The dimension of the space $\mathcal{E}_{-s}^{\mathcal{G}} \subset H_{-s}^1$ is finite.
 - If a vector $u \in \mathcal{E}_{-s}^{\mathcal{G}}$, then $f = T^* u \in \mathcal{E}_{-s}^{\mathcal{H}} \subset \mathcal{H}_{-s}^1$. Conversely if $f \in \mathcal{E}_{-s}^{\mathcal{H}}$, then $u = E_+^+(\lambda_0)f \in \mathcal{E}_{-s}^{\mathcal{G}}$ and $v^+f = v^-f$.

- 2) If $f \in \mathcal{E}_{-s}^{\mathcal{H}}$, then $f \in \mathcal{H}_{(\nu_0-1)^-}^1$ and $f \in \ker(1+K^+)$; here $K^+ \in \mathcal{C}(\mathcal{H}_{-t}^1)$ is given by (4.60) and $t \in (s_1, 3/2)$ is arbitrary. Conversely if $f \in \mathcal{H}_{-1}^1$ and $f \in \ker(1+K^+)$, then $f \in \mathcal{E}_{-1}^{\mathcal{H}}$ (in particular $E_+^-(\lambda_0)f = E_+^+(\lambda_0)f$).
- 3) Suppose $f \in \mathcal{E}_{-s}^{\mathcal{H}}$. In the case $s_0 = 1$ suppose in addition that $f \in \mathcal{H}_{-t}^1$ for some t < 1. Then the components of $f = (f_1, f_2)$ can be decomposed as

$$-f_{j}(r_{j}\theta_{j}) = \sum_{\nu \in \sigma_{j,1}} \sum_{k=1}^{n_{j,\nu}} l_{j,\nu,k}(f) r_{j}^{\frac{2-n_{j}}{2}-\nu} \chi_{2}(r_{j}) \zeta_{j,\nu}^{(k)}(\theta_{j}) + g_{j},$$

where $g_{j} \in L^{2}$ and
 $l_{j,\nu,k}(f) = \langle \phi_{\nu}(|y_{j}|) |y_{j}|^{-\frac{n_{j}-1}{2}} \otimes \zeta_{j,\nu}^{(k)}, \tilde{f}_{j}(y_{j}) \rangle_{L^{2}(\mathrm{d}y_{j})};$
 $\tilde{f}_{j} = \chi_{2} (v^{+}f)_{j} + \chi_{2}B_{j}f_{j} + [\chi_{2}, p_{j}^{2}]f_{j}.$

$$(4.61)$$

Here

$$l_{j,\nu,k}(f) = 0 \text{ for } i\nu \ge 0.$$
 (4.62)

In particular

$$f \in \mathcal{H} \quad \Longleftrightarrow \quad \forall j = 1, 2, \nu \in \sigma_{j,1}^+, \, k = 1, \dots, n_{j,\nu} :$$
$$l_{j,\nu,k}(f) = 0.$$
(4.63)

4) Suppose $u \in \mathcal{E}_{-s}^{\mathcal{G}}$. In the case $s_0 = 1$ suppose in addition that $u \in H^1_{-t}$ for some t < 1. Then

$$u \in L^{2}(\mathbf{X}) \quad \Longleftrightarrow \quad \forall j = 1, 2, \nu \in \sigma_{j,1}^{+}, k = 1, \dots, n_{j,\nu} :$$
$$l_{j,\nu,k}(T^{*}u) = 0.$$
(4.64)

In particular if $s_0 \neq 1$, then

$$\dim \left(\mathcal{E}_{-1}^{\mathcal{G}} / \ker(H - \lambda_0)_{|H^1} \right) \le \sum_{j=1}^2 \sum_{\nu \in \sigma_{j,1}^+} n_{j,\nu}.$$
(4.65)

For $s_0 = 1$ the bound (4.65) is valid provided that $\mathcal{E}_{-1}^{\mathcal{G}}$ to the left is replaced by $\mathcal{E}_{-1}^{\mathcal{G}} \cap H^1_{(-1)^+}$.

Remark 4.16. We note that Remarks 4.8 and 4.9 apply equally well for the case $\lambda_0 > \Sigma_2$.

4.3 Models of physics

In this section we treat the Coulombic potential models of Subsections 1.2.1 and 1.3.1 (with $N \ge 3$ and $N \ge 2$, respectively) and demonstrate consequences of the previous sections for these models for any given two-cluster threshold $\lambda_0 < 0$. For

convenience we shall not distinguish between the cases $\lambda_0 = \Sigma_2$ and $\lambda_0 > \Sigma_2$ as done before. Whence we will be concerned with generalizing Section 4.2 only (note that the case $\lambda_0 = \Sigma_2$ can be considered as a special case of Section 4.2 although stronger results are presented in Section 4.1). As in Sections 1.2 and 1.3 we shall not consider models with spin included (see however Remark 4.19).

We consider a two-cluster threshold λ_0 for the models of Sections 1.2 and 1.3, grouping the set of thresholds *a* for which $\lambda_0 \in \sigma_{pp}(H^a)$ into \mathcal{A}_1 , \mathcal{A}_2 and \mathcal{A}_3 (all depending on λ_0) for which

 \mathcal{A}_1 : the effective inter-cluster interaction is to leading order attractive Coulombic,

 \mathcal{A}_2 : the effective inter-cluster interaction is to leading order repulsive Coulombic,

 \mathcal{A}_3 : the effective inter-cluster interaction is $\mathcal{O}(|x_a|^{-2})$.

Note that this distinction does not depend on choices of corresponding sub-Hamiltonian bound states φ^a (i.e. channels); it is determined by charges only (cf. Case 1 introduced independently in both of the above sections).

Let for $a \in \mathcal{A} := \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3$ (again we suppress the dependence of λ_0) the operator P^a be the corresponding orthogonal projection onto $\ker(H^a - \lambda_0)$ in $L^2(\mathbf{X}^a)$ and let m_a be the dimension of this space. Obviously $\Pi^a := P^a \otimes 1$ projects onto the span of functions of the form $\varphi^a \otimes f_a$, $\varphi^a \in \ker(H^a - \lambda_0)$, in $L^2(\mathbf{X})$. We identify ran P^a , say spanned by an orthonormal basis $\varphi_1^a, \ldots, \varphi_{m_a}^a$, with \mathbb{C}^{m_a} (using the basis), and similarly

$$L^{2}(\mathbf{X}_{a}, \mathbb{C}^{m_{a}}) \simeq \bigoplus_{m \leq m_{a}} L^{2}(\mathbf{X}_{a}) \ni \bigoplus_{m \leq m_{a}} f_{a,m} = f_{a}$$
$$\simeq S_{a} f_{a} := \sum_{m \leq m_{a}} \varphi_{m}^{a} \otimes f_{a,m} \in \operatorname{ran} \Pi^{a}.$$

Considering for the moment only $a \in \mathcal{A}_3$ we write $W_a := S_a^* I_a S_a = Q_a |x_a|^{-2} + B_a$, where Q_a is a $m_a \times m_a$ matrix-valued function depending only on $\theta = \hat{x}_a = |x_a|^{-1} x_a$ while $B_a = B_a(x_a) = \mathcal{O}(|x_a|^{-3})$. The 'right generalization' to the case $m_a > 1$ of the distinction between Cases 2) and 3) (discussed primarily for $m_a = 1$ in both of Sections 1.2 and 1.3) is to let Case 2) correspond to $Q_a \neq 0$ and let Case 3) correspond to $Q_a = 0$, respectively.

Let $\mathbb{S}_a = \mathbb{S}_a^{n-1}$, $n = \dim \mathbf{X}_a$, denote the unit sphere in \mathbf{X}_a . We use spherical coordinates on \mathbf{X}_a and the Laplace-Beltrami operator Δ_{θ} to express the Laplacian p_a^2 on $L^2(\mathbf{X}_a, \mathbb{C}^{m_a})$. The operator $-\Delta_{\theta} + Q_a$ on $L^2(\mathbb{S}_a, \mathbb{C}^{m_a})$ has discrete spectrum. As in (4.21) we parametrize its eigenvalues in terms of a parameter ν the collection of which is denoted be σ_a . More precisely consider for each $\mu \in \sigma(-\Delta_{\theta} + Q_a)$ the equation $\mu = \nu^2 - \frac{(n-2)^2}{4}$, where by convention $\nu \ge 0$ if $\mu \ge -\frac{(n-2)^2}{4}$, and $i\nu > 0$ if $\mu < -\frac{(n-2)^2}{4}$. Then the collection of such numbers ν is denoted by σ_a . The orthogonal projection onto the corresponding eigenspace is denoted by $P_{a,\nu}$. We take an orthonormal basis in ran $P_{a,\nu}$ for each $\nu \in \sigma_a$, say $\zeta_{a,\nu}^{(1)}, \ldots, \zeta_{a,\nu}^{(n_a,\nu)}$. Let

$$\begin{aligned} \sigma_{a,1} &= \sigma_a \setminus (1,\infty), \quad \sigma_{a,1}^+ = \sigma_a \cap (0,1], \\ \nu_a &= \min_{\nu \in \sigma_a} \operatorname{Re} \nu, \quad s_a = 1 + \nu_a, \quad d_a = \sum_{\nu \in \sigma_{a,1}^+} n_{a,\nu}. \end{aligned}$$

(

Now for the physics models we have the following result, recalling the notation $L_{t^+}^2 = L_{t^+}^2(\mathbf{X}) = \bigcup_{s>t} L_s^2(\mathbf{X})$ for any real t.

Theorem 4.17. 1) The space of locally H^1 solutions to $(H - \lambda_0)u = 0$ in

$$\sum_{a \in \mathcal{A}_1} \Pi^a L^2_{-3/4} + \sum_{a \in \mathcal{A}_2} \Pi^a L^2_{(-3/2)^+} + \sum_{a \in \mathcal{A}_3} \Pi^a L^2_{(-\min\{3/2, s_a\})^+} + L^2_{-1/2},$$

say denoted by \mathcal{E} , has finite dimension.

If $\mathcal{A}_3 = \emptyset$, then $\mathcal{E} \subset H^1_{\infty}$.

2) The number

$$\dim \left(\mathcal{E} / \ker(H - \lambda_0)_{|H^1} \right) \le \sum_{a \in \mathcal{A}_3} d_a$$

3) There exist linear functionals $\check{l}_{a,\nu,k} : \mathcal{E} \to \mathbb{C}$ defined for $a \in \mathcal{A}_3$, $\nu \in \sigma_{a,1}^+$ and $k = 1, \ldots, n_{a,\nu}$, such that for any $u \in \mathcal{E}$

$$u \in L^2 \Leftrightarrow l_{a,\nu,k}(u) = 0$$
 for all such a, ν and k .

Any $u \in \mathcal{E}$ fulfills the asymptotics

$$u - \sum_{a,\nu,k,m} \check{l}_{a,\nu,k}(u)\varphi_m^a \otimes \zeta_{a,\nu,m}^{(k)} \otimes |x_a|^{\frac{2-n}{2}-\nu} F(|x_a| > 1) \in L^2.$$
(4.66)

(Here
$$\zeta_{a,\nu}^{(k)}$$
, labels the m_a coordinates of $\zeta_{a,\nu}^{(k)}$.)

For those $a \in \mathcal{A}_3$ for which $s_a \in (1, 3/2)$ we can replace $L^2_{(-\min\{3/2, s_a\})^+}$ by $L^2_{-s_a}$ in the definition of \mathcal{E} . If $\lambda_0 = \Sigma_2$ we can to some extent use a bigger space of generalized eigenfunctions than the one considered in 1), cf. Theorems 4.3 and 4.7 (not to be elaborated on).

Before giving details of proofs we discuss the structure of (4.66) in the case the angular-depending potential Q_a vanishes (for a given $a \in \mathcal{A}_3$). Note that for $m_a = 1$ this corresponds to Case 3 of Sections 1.2 and 1.3. If $Q_a = 0$ the set $\sigma_{a,1}^+$ is given explicitly as

$$\sigma_{a,1}^{+} = \begin{cases} \left\{ \frac{1}{2} \right\}, & n = 3, \\ \left\{ 1 \right\}, & n = 4, \\ \emptyset, & n \ge 5. \end{cases}$$
(4.67)

Moreover for n = 3 and n = 4 the total multiplicity of the single point in $\sigma_{a,1}^+$ is m_a (corresponding to constant eigenfunctions $\zeta_{a,\nu}^{(k)}$, $k = 1, \ldots, m_a$). In particular we see that if $Q_a = 0$ for all $a \in \mathcal{A}_3$, then the space \mathcal{E} simplifies as the space of generalized eigenfunctions in

$$\sum_{a \in \mathcal{A}_1} \Pi^a L^2_{-3/4} + \sum_{a \in \mathcal{A}_2 \cup \mathcal{A}_3} \Pi^a L^2_{(-3/2)^+} + L^2_{-1/2},$$

and we see that λ_0 can be a 'resonance' only for n = 3 or n = 4 and then with multiplicity at most $\sum_{a \in A_3} m_a$.

4.3. Models of physics

The proof Theorem 4.17 relies strongly on Section 4.2. Clearly the assertion 2) is a consequence of 3). A complication of the proof of the assertions 1) and (4.66) is the possible existence of exceptional cases discussed previously, partly in remarks. In fact the definition of the functionals $\check{l}_{a,\nu,k}$ depends on the such cases. We shall first give the proof of Theorem 4.17 in the 'generic' case and then outline the proof for exceptional cases.

Proof of Theorem 4.17 in the 'generic' case. We consider the non-exceptional case. This amounts to proving the theorem under two separate additional conditions. We introduce $\mathcal{G} = L^2(\mathbf{X})$ and $\mathcal{F}_a = \Pi^a \mathcal{G}$, $a \in \widetilde{\mathcal{A}}$. The first assumption is

$$\mathcal{F} := \sum_{a \in \widetilde{\mathcal{A}}} \mathcal{F}_a = \sum_{a \in \widetilde{\mathcal{A}}} \oplus \mathcal{F}_a.$$
(4.68a)

Let Π be the orthogonal projection in \mathcal{G} onto \mathcal{F} (with (4.68a) \mathcal{F} is closed, cf. Proposition 2.7). The second assumption is

$$\lambda_0 \notin \sigma_{\rm pp}(H'); \quad H' = \Pi' H \Pi', \, \Pi' = 1 - \Pi.$$
 (4.68b)

Under (4.68a) and (4.68b) we let $\mathcal{H} = \sum_{a \in \widetilde{\mathcal{A}}} \oplus \mathcal{H}_a$ where $\mathcal{H}_a = L^2(\mathbf{X}_a, \mathbb{C}^{m_a}) = \bigoplus_{m \leq m_a} L^2(\mathbf{X}_a)$, and we let $S = (S_a) : \mathcal{H} \to \mathcal{F} \subset \mathcal{G}$ be given by

$$\oplus_{a\in\widetilde{\mathcal{A}}} f_a \to \sum_{a\in\widetilde{\mathcal{A}}} S_a f_a; \quad f_a = \sum_{m \le m_a} \oplus f_{a,m}, \quad S_a f_a = \sum_{m \le m_a} \varphi_m^a \otimes f_{a,m}.$$

Letting $T = (SS^*)^{-1}S$ we note that $\Pi = ST^*$.

In agreement with (2.4d) and (4.39) we consider two operators $E_{\mathcal{H}}^{\pm}(\lambda_0)$, specified by the entries

$$-E_{\mathcal{H}}^{\pm}(\lambda_0)_{ab} = S_a^*(p_b^2 + I_b)S_b + K_{ab}^{\pm}(\lambda_0) \equiv \delta_{ab}\check{h}_a + K_{ab}^{\pm}(\lambda_0); \quad a, b \in \widetilde{\mathcal{A}},$$
(4.69)

as operators from $\mathcal{H}^1 = \sum_{a \in \widetilde{\mathcal{A}}} \oplus H^1(\mathbf{X}_a, \mathbb{C}^{m_a})$ to $\mathcal{H}^{-1} = \sum_{a \in \mathcal{A}_1 \cup \mathcal{A}_2} \oplus H^{-1}(\mathbf{X}_a, \mathbb{C}^{m_a})$. Here $\check{h}_a = -\Delta_a + S_a^* I_a S_a$ and $K_{ab}^{\pm}(\lambda_0) = -S_a^* I_a R'(\lambda_0 \pm i0) I_b S_b$. As in (4.39) the difference is symmetric and polynomially decreasing. The latter notion is defined as in Remark 2.13, i.e. (b_{ab}) is polynomially decreasing if $b_{ab} \in \mathcal{C}(H_r^1(\mathbf{X}_b, \mathbb{C}^{m_b}), H_t^{-1}(\mathbf{X}_a, \mathbb{C}^{m_a}))$ for all $r, t \in \mathbb{R}$.

We note that

$$K_{ab}^{\pm}(\lambda_0) \in \mathcal{L}(L_{-s}^2(\mathbf{X}_b, \mathbb{C}^{m_b}), L_s^2(\mathbf{X}_a, \mathbb{C}^{m_a})) \text{ for all } s < 3/2.$$
(4.70a)

For $a \in \widetilde{\mathcal{A}}$ the effective inter-cluster interaction $W_a = S_a^* I_a S_a$ obeys

$$W_a = C_a |x_a|^{-1} \mathbb{1}_{\mathbb{C}^{m_a}} + \mathcal{O}(|x_a|^{-2}),$$

where $C_a \neq 0$ if and only if $a \in \mathcal{A}_1 \cup \mathcal{A}_2$. (4.70b)

Recall also

$$\forall a \in \mathcal{A}_3: \quad W_a = |x_a|^{-2}Q_a(\hat{x}_a) + B_a(x_a), \quad B_a = \mathcal{O}(|x_a|^{-3}).$$
 (4.70c)

Write $\mathcal{A}_3 = \{a_1, \ldots, a_{\bar{l}}\}$ (assuming $\mathcal{A}_3 \neq \emptyset$) and let \mathcal{T} denote the set of vectors $\bar{t} = (t_1, \ldots, t_{\bar{l}}) \in \mathbb{R}^{\bar{l}}$ such that $t_l > -\min\{3/2, s_{a_l}\}$ for all $l \leq \bar{l}$. We introduce for $r \geq -3/4, s > -3/2$ and $\bar{t} \in \mathcal{T}$ the spaces

$$\mathcal{H}_{a,s}^{k} = H_{s}^{k}(\mathbf{X}_{a}, \mathbb{C}^{m_{a}}), \quad k \in \mathbb{R}, \quad a \in \mathcal{A},$$

$$\mathcal{H}_{r,s,\bar{t}}^{k} = \bigoplus_{b \in \mathcal{A}_{1}} \mathcal{H}_{b,r}^{k} \bigoplus \bigoplus_{b \in \mathcal{A}_{2}} \mathcal{H}_{b,s}^{k} \bigoplus \bigoplus_{l \leq \bar{l}} \mathcal{H}_{a_{l},t_{l}}^{k}, \quad k \in \mathbb{R},$$

$$\mathcal{E}_{r,s,\bar{t}}^{\mathcal{H}} = \{ f \in \mathcal{H}_{r,s,\bar{t}}^{1} | \quad E_{\mathcal{H}}^{+}(\lambda_{0})f = E_{\mathcal{H}}^{-}(\lambda_{0})f = 0 \},$$

where $E_{\mathcal{H}}^{\pm}(\lambda_0)$ is given by (4.69). A similar notation may be used if $\mathcal{A}_3 = \emptyset$, for example $\mathcal{E}_{r,s}^{\mathcal{H}}$, however we prefer to keep the uniform notation $\mathcal{E}_{r,s,\bar{t}}^{\mathcal{H}}$ not distinguising between whether one (or two) of the sets \mathcal{A}_1 , \mathcal{A}_2 and \mathcal{A}_3 is (are) empty or, possibly, all three sets are non-empty. Let

$$\mathcal{E}^{\mathcal{H}} = \bigcup_{s > -3/2, \, \bar{t} \in \mathcal{T}} \quad \mathcal{E}^{\mathcal{H}}_{-3/4, s, \bar{t}}. \tag{4.71}$$

Let $E^{\pm}_{+}(\lambda_0) \in \mathcal{L}(\mathcal{E}^{\mathcal{H}}, \mathcal{E})$ be given by

$$E^{\pm}_{+}(\lambda_{0})f = Sf - \sum_{a \in \widetilde{\mathcal{A}}} R'(\lambda_{0} \pm \mathrm{i}0)I_{a}S_{a}f_{a}.$$
(4.72)

By mimicking the proof of Lemma 4.11 we can show that indeed $E^+_+(\lambda_0) : \mathcal{E}^{\mathcal{H}} \to \mathcal{E}$ and that, considered as such a map, it is a linear isomorphism with inverse $T^* : \mathcal{E} \to \mathcal{E}^{\mathcal{H}}$. Note that as before we need the property that $E^+_+(\lambda_0)f = E^-_+(\lambda_0)f$ for any $f \in \mathcal{E}^{\mathcal{H}}$.

Next we adapt the parametrix construction of Section 4.2 (combining the three different cases treated there) and convert the equation $E_{\mathcal{H}}^+(\lambda_0)f = 0, f \in \mathcal{E}^{\mathcal{H}}$, to an equation of the form $(1 + K^+)f = 0$. We can then show by iteration that f belongs to a suitable space (independent of f) on which K^+ is compact, yielding 1).

To carry this out in more details, let us define $R^+ = \bigoplus_{a \in \widetilde{\mathcal{A}}} r_a$ where r_a is the following parametrix (depending on the three cases): For $a \in \mathcal{A}_1 \cup \mathcal{A}_2$ we take r_a diagonal in \mathcal{H}_a , $r_a = \bigoplus_{m \leq m_a} r_{a,m}$. For $a \in \mathcal{A}_1$ and $m \leq m_a$ we let

$$r_{a,m} = \lim_{\epsilon \to 0_+} (h_a - i\epsilon)^{-1} \in \mathcal{L}(H_s^{-1}(\mathbf{X}_a), H_{-s}^1(\mathbf{X}_a)), \ s > 3/4,$$

where $h_a = p_a^2 + w_a$ is constructed as in Subsection 4.1.2. For $a \in \mathcal{A}_2$ we let $r_a = \bigoplus_{m \le m_a} r_{a,m}$, where $r_{a,m} = h_a^{-1}$ with $h_a = p_a^2 + w_a$ been constructed as in Subsection 4.1.3. Note that in this case $\langle \cdot \rangle^{-1} h_a^{-1}$ is a bounded pseudodifferential operator, cf. Lemma 4.4. Finally we let for $a \in \mathcal{A}_3$ the operator $r_a = G_{a+}$, where G_{a+} is constructed as in Subsection 4.1.4, i.e.

$$G_{a+} = \chi_1(\check{h}_a - \mathbf{i})^{-1}\chi_1 + \sum_{\nu \in \sigma_a} \left(\chi_2 \left|\cdot\right|^{\frac{1-n}{2}} R_{a,\nu} \left|\cdot\right|^{\frac{n-1}{2}} \chi_2\right) \otimes P_{a,\nu}.$$
 (4.73)

Here $\check{h}_a = p_a^2 + W_a$, where $W_a = S_a^* I_a S_a = |\cdot|^{-2} Q_a + \mathcal{O}(|\cdot|^{-3})$ acting as a (matrixvalued) multiplication operator on \mathcal{H}_a . The quantities Q_a , σ_a and $P_{a,\nu}$ are introduced before Theorem 4.17, and the operator $R_{a,\nu}$ is the one-dimensional Green's function studied in Subsection 4.1.4. While the parametrix r_a for $a \in \mathcal{A}_1 \cup \mathcal{A}_2$ is in some sense taken as an exact inverse, the operator $r_a = G_{a+}$ chosen for $a \in \mathcal{A}_3$ is less accurate. This is explicitly seen in (4.60) in which the second up to the fourth terms represent 'errors' not appearing for $a \in A_1 \cup A_2$.

Next we write the equation $-R^+E^+_{\mathcal{H}}(\lambda_0)f = 0$ for $f \in \mathcal{E}^{\mathcal{H}}$ as $(1+K^+)f = 0$, where integration by parts is used to produce the term 1f. This is done separately for the components of f by the arguments of Subsections 4.2.2–4.2.4. By (4.69)there are terms in K^+ involving polynomially decreasing factors and there are also terms of the form $r_a K_{ab}^{\pm}(\lambda_0)$. Effectively $K_{ab}^{\pm}(\lambda_0)$ is of order -3 when applied to an f given as above. For $a \in \mathcal{A}_3$ there are additional terms as discussed above, cf. (4.60). In any case it follows that the weighted orders of the components of $K^+ f$ tend to be at least 1/2 power better (as determined by (4.70a)-(4.70c)) than the apriory orders of the components of $f(=-K^+f)$ (at this point, note also (4.53) and its application (4.9b)). However there is an exception for this general rule due to the restriction $s > 1 - \nu_a$ imposed by mapping properties of G_{a+} , cf. (4.58). Note for example that to prove $\sum_{b} r_a K_{ab}^{\pm}(\lambda_0) f_b \in \mathcal{H}^1_{a,-s}$ for $a \in \mathcal{A}_3$ we need at least $s > 1 - \nu_a$. This is the only restriction for an iteration argument. Actually we infer from one iteration (i.e. just from the properties of K^+f) that for $a \in \mathcal{A}_1$ the component $f_a \in \mathcal{H}^1_{a,-1/4}$, that for $a \in \mathcal{A}_2$ the component $f_a \in \mathcal{H}^1_{a,-1/2}$, while $f_a \in \mathcal{H}^1_{a,\epsilon-1}$ for some positive ϵ for $a \in \mathcal{A}_3$ (noting though that the latter property is an assumption if $\nu_a = 0$). With one more iteration we then obtain that $f_a \in \mathcal{H}^1_{a,0}$ for $a \in \mathcal{A}_1 \cup \mathcal{A}_2$. If $\mathcal{A}_3 = \emptyset$ continued iteration leads to $f \in \mathcal{H}^1_{\infty}$ (with Lemma 4.11 this proves the very last part of 1)). In particular we can conclude that

$$f \in \mathcal{E}^{\mathcal{H}} \Rightarrow f \in \mathcal{H}^1_* := \mathcal{H}^1_{-1,-1,\bar{t}},$$

where all coordinates of \bar{t} are taken to be -4/3. Since $K^+ \in \mathcal{C}(\mathcal{H}^1_*)$ we have shown 1).

It remains to prove 3). We only need to construct linear functionals $l_{a,\nu,k} : \mathcal{E} \to \mathbb{C}$ such that (4.66) is fulfilled for any $u \in \mathcal{E}$. For given $u \in \mathcal{E}$ we let $f = T^*u$ and need to examine the components f_a of f that might not be in \mathcal{H}_a . As we have seen this amounts to looking at $a \in \mathcal{A}_3$ only in which case we know the analogue of (4.35) that $f_a \in \mathcal{H}^1_{a,-s}$ for some s < 1. Looking at $-f_a = (K^+f)_a$ for such a we need the first equation of (4.69) written as

$$-E_{\mathcal{H}}^{\pm}(\lambda_0)_{ab} = S_a^*(p_b^2 + I_b)S_b + K_{ab}^{\pm}(\lambda_0) = \delta_{ab}\check{h}_a + v_{ab}^+; \quad \check{h}_a = p_a^2 + W_a.$$
(4.74)

Now, by using (4.73) and the analogue of (4.60), we obtain by mimicking Step III of the proof of Theorem 4.7 that

$$-f_{a}(|x_{a}|\theta_{a}) = \sum_{\nu \in \sigma_{a,1}} \sum_{k=1}^{n_{a,\nu}} l_{a,\nu,k}(f) |x_{a}|^{\frac{2-n}{2}-\nu} \chi_{2}(|x_{a}|) \zeta_{a,\nu}^{(k)}(\theta_{a}) + g_{a},$$
where $g_{a} \in \mathcal{H}_{a}$ and
$$l_{a,\nu,k}(f) = \langle \phi_{\nu}(|y_{a}|) |y_{a}|^{-\frac{n-1}{2}} \otimes \zeta_{a,\nu}^{(k)}, \tilde{f}_{a}(y_{a}) \rangle_{\mathcal{H}_{a}};$$

$$\tilde{f}_{a} = \chi_{2} \sum_{b \in \tilde{\mathcal{A}}} v_{ab}^{+} f_{b} + \chi_{2} B_{a} f_{a} + [\chi_{2}, p_{a}^{2}] f_{a}.$$
(4.75)

Note that $\tilde{f}_a \in \mathcal{H}^0_{a,3-s}$ for some s < 1, cf. (4.36), yielding in particular that $l_{j,\nu,k}(f)$ is well-defined. Next we deduce from (4.75) that

$$l_{j,\nu,k}(f) = 0 \text{ for } i\nu \ge 0.$$
 (4.76)

Clearly (4.75) and (4.76), in combination with (4.72) and the adapted version of Lemma 4.11 (stated after (4.72)), lead to (4.66) with $\tilde{l}_{a,\nu,k}$ given by

$$\check{l}_{a,\nu,k}(u) = -l_{a,\nu,k}(T^*u); \quad u \in \mathcal{E}, \ a \in \mathcal{A}_3, \ \nu \in \sigma_{a,1}^+, \ k = 1, \dots, n_{a,\nu}.$$

Proof of Theorem 4.17 in the 'general' case, a sketch. We outline a proof of the theorem without the two additional conditions (4.68a) and (4.68b) imposed. We shall implement Remarks 4.9 and 4.16 using the previous notation as much as possible.

First, if (4.68a) is not fulfilled, we infer again that the space $\sum_{a \in \widetilde{\mathcal{A}}} \mathcal{F}_a$ is closed by considering as in Subsection 2.5 the 'restriction' of S to

$$\mathcal{H}_{0} := \left\{ f \in \sum_{a \in \widetilde{\mathcal{A}}} \oplus \mathcal{H}_{a} \mid f \perp \ker \left(S_{ab} \right)_{a,b \in \widetilde{A}} \right\}, S_{ab} = S_{a}^{*} S_{b}, \tag{4.77}$$

where as before $\mathcal{H}_a = L^2(\mathbf{X}_a, \mathbb{C}^{m_a})$. Then $S = (S_a) : \mathcal{H}_0 \to \mathcal{F}_0 := \sum_{a \in \widetilde{\mathcal{A}}} \mathcal{F}_a \subset \mathcal{G} = L^2(\mathbf{X})$ is a continuous isomorphism. In particular indeed \mathcal{F}_0 is closed, and we let Π denote the orthogonal projection onto \mathcal{F}_0 in \mathcal{G} . The quantities Π' , H' and R'(z) are introduced as in previous sections (with \mathcal{F}_0 playing the role of \mathcal{F}).

Next we would like to consider limits $R'(\lambda_0 \pm i0)$ which is doable under the condition (4.68b). However we shall allow, and here consider, the exceptional case where (4.68b) is not fulfilled, and we need a version of Section 2.4 to treat this case. By a version of Theorem 3.12, cf. Remarks 3.20, 3.21 and 3.22, λ_0 has finite multiplicity as an eigenvalue of H' with eigenfunctions in H^2_{∞} .

To simplify the presentation let us make the same assumption as in Subsection 2.4 that this eigenspace is given as $\operatorname{span}\{\psi\}$, $\|\psi\| = 1$. We introduce then $\mathcal{H} = \mathcal{H}_0 \oplus \mathbb{C}$ and $S : \mathcal{H} \to \mathcal{F} := \mathcal{F}_0 \oplus \operatorname{span}\{\psi\}$ by

$$S\tilde{f} = \sum_{a} S_{a}f_{a} + c\psi, \quad \tilde{f} = ((f_{a}), c).$$

We let Π'' denote the orthogonal projection onto the orthogonal complement of \mathcal{F} in \mathcal{G} , and let $H'' := \Pi'' H \Pi''$ and $R''(z) = (H'' - z)^{-1}$. Note that $\Pi'' = 1 - \Pi - |\psi\rangle\langle\psi| = \Pi' - |\psi\rangle\langle\psi|$. Recall then (2.4d)

$$E_{\mathcal{H}}(z) = S^* \big(z - H + H R''(z) H \big) S.$$
(4.78)

Letting $M = \#\widetilde{\mathcal{A}} + 1$ this operator has a $M \times M$ -block representation $(e_{ij})_{i,j \leq M}$. Here (listing the elements $a \in \widetilde{\mathcal{A}}$ as a_1, \ldots, a_{M-1}) $e_{MM} = z - \lambda_0$, $e_{iM} = -S_{a_i}^* H \psi = -S_{a_i}^* I_{a_i} \psi$ and $e_{Mi} = e_{iM}^* = \langle e_{Mi} |$ for $i \leq M - 1$, while for $i, j \leq M - 1$ we set $e_{ij} = S_{a_i}^* (z - H + HR''(z)H) S_{a_j}$. The analogue of (2.60a) reads

$$\begin{split} \breve{H} &= H'' + \sum_{a \in \widetilde{\mathcal{A}}} p_a^2 \Pi^a, \\ \widetilde{H} &= H - \lambda_0 \sum_{a \in \widetilde{\mathcal{A}}} \Pi^a; \\ \mathcal{D}(\breve{H}) &= \mathcal{D}(\widetilde{H}) = \mathcal{D}(H). \end{split}$$
(4.79a)

4.3. Models of physics

These operators differ by H-compact terms, cf. (2.60b),

$$\breve{H} = \widetilde{H} - K_1 + K_2;$$

$$K_1 = \Pi H \Pi' + \Pi' H \Pi + \sum_{a \in \widetilde{\mathcal{A}}} \Pi^a I_a \Pi^a + \lambda_0 |\psi\rangle \langle\psi|,$$

$$K_2 = \sum_{a \in \widetilde{\mathcal{A}}} \Pi^a H \Pi^a - \Pi H \Pi,$$
(4.79b)

and the basic structure of resolvents is given as

$$R''(z) = \breve{R}(z) - \left(\sum_{a \in \widetilde{\mathcal{A}}} p_a^2 \Pi^a - z\right)^{-1} \Pi - z^{-1} |\psi\rangle \langle\psi|,$$

$$R''(z) = \Pi'' \breve{R}(z) \Pi'' = \Pi'' \breve{R}(z) = \breve{R}(z) \Pi''; \quad \breve{R}(z) = (\breve{H} - z)^{-1}.$$
(4.79c)

Due to the good properties of ψ we can make sense to $\check{R}(\lambda_0 \pm i0)$ and then in turn to $R''(\lambda_0 \pm i0)$. In particular we can consider (4.72) with f replaced by $\tilde{f} := ((f_a), c) \in \mathcal{H}$ and $R'(\lambda_0 \pm i0)$ by $R''((\lambda_0 \pm i0))$, i.e.

$$E^{\pm}_{+}(\lambda_{0})\tilde{f} = S\tilde{f} - \sum_{a\in\tilde{\mathcal{A}}} R''(\lambda_{0}\pm \mathrm{i}0)I_{a}S_{a}f_{a}.$$
(4.80a)

Similarly, by taking limits in (4.78), we obtain

$$E_{\mathcal{H}}^{\pm}(\lambda_0) = S^* \big(\lambda_0 - H + HR''(\lambda_0 \pm i0)H\big) S = (e_{ij})_{i,j \le M, z = \lambda_0 \pm i0}.$$
 (4.80b)

The previous definition of $\mathcal{E}^{\mathcal{H}}$ given in (4.71) needs to be modified as follows. We introduce (for parameters given as before)

$$\begin{aligned} \mathcal{H}_{r,s,\bar{t}}^{k,0} &= \left\{ f \in \mathcal{H}_{r,s,\bar{t}}^k \mid f \perp \ker \left(S_{ab} \right)_{a,b \in \tilde{A}} \right\}, \\ \mathcal{E}_{r,s,\bar{t}}^{\mathcal{H}} &= \{ \tilde{f} = (f,c) \mid f \in \mathcal{H}_{r,s,\bar{t}}^{1,0}, c \in \mathbb{C}, \ E_{\mathcal{H}}^+(\lambda_0) \tilde{f} = E_{\mathcal{H}}^-(\lambda_0) \tilde{f} = 0 \}, \\ \mathcal{E}^{\mathcal{H}} &= \cup_{s > -3/2, \, \bar{t} \in \mathcal{T}} \quad \mathcal{E}_{-3/4,s,\bar{t}}^{\mathcal{H}}. \end{aligned}$$

As before we can now check that $E^+_+(\lambda_0) : \mathcal{E}^{\mathcal{H}} \to \mathcal{E}$ and that, considered as such a map, it is a linear isomorphism with inverse $T^* = S^*(SS^*)^{-1} : \mathcal{E} \to \mathcal{E}^{\mathcal{H}}$.

Next we consider the space

$$\widetilde{\mathcal{H}} = \left(\sum_{a \in \widetilde{\mathcal{A}}} \oplus \mathcal{H}_a\right) \oplus \mathbb{C}.$$

Clearly $\widetilde{\mathcal{H}} = \mathcal{H} \oplus \ker (S_{ab})_{a,b\in\widetilde{A}}$, and $E_{\mathcal{H}}^{\pm}(\lambda_0)$ acts only on the first component. Let us introduce $\widetilde{E}_{\mathcal{H}}^{\pm}(\lambda_0) = E_{\mathcal{H}}^{\pm}(\lambda_0) + P_0$, where P_0 denotes the orthogonal projection onto $\ker (S_{ab})_{a,b\in\widetilde{A}}$ in $\sum_{a\in\widetilde{A}} \oplus \mathcal{H}_a$. We can consider P_0 as acting on the second component of $\widetilde{\mathcal{H}}$ and hence write $\widetilde{E}_{\mathcal{H}}^{\pm}(\lambda_0) = E_{\mathcal{H}}^{\pm}(\lambda_0) \oplus 1$. Clearly $\ker \widetilde{E}_{\mathcal{H}}^{\pm}(\lambda_0) = \ker E_{\mathcal{H}}^{\pm}(\lambda_0)$, and similar relations hold in weighted Sobolov spaces. We are lead to consider $\widetilde{E}_{\mathcal{H}}^{\pm}(\lambda_0) = E_{\mathcal{H}}^{\pm}(\lambda_0) \oplus 1$ on spaces of the form

$$\widetilde{\mathcal{H}}_{r,s,\bar{t}}^{k} = \left(\mathcal{H}_{r,s,\bar{t}}^{k,0} \oplus \mathbb{C}\right) \oplus \ker\left(S_{ab}\right)_{a,b\in\widetilde{A}} = \mathcal{H}_{r,s,\bar{t}}^{k} \oplus \mathbb{C}.$$

With the same restrictions on the parameters we can prove a version of Lemma 4.11 (using the same proof), and by applying the parametrix construction $\bigoplus_{a \in \widetilde{\mathcal{A}}} r_a$ of the previous proof of the theorem (the regular case) extended as $R_{\text{diag}}^+ = (\bigoplus_{a \in \widetilde{\mathcal{A}}} r_a) \oplus 1_{\mathbb{C}}$ to the equation

$$\widetilde{E}_{\mathcal{H}}^{\pm}(\lambda_0)\widetilde{f} = 0, \quad \widetilde{f} := (f, c) = \left(\left(P_0'f, c \right), P_0 f \right) \in \widetilde{\mathcal{H}}_{r, s, \overline{t}}^k, \ P_0' = 1 - P_0,$$

we arrive again at Fredholm theory. More precisely, using that the operator $E_{\mathcal{H}}^{\pm}(\lambda_0) \oplus$ 0 is realized concretely by the expression on the right-hand side of (4.80b), we arrive at the equation

$$0 = R^+_{\text{diag}} \left(P_0 f + (\widetilde{E}^{\pm}_{\mathcal{H}}(\lambda_0)\widetilde{f} - P_0 f) \right) = -(1 + \widetilde{K}^+)\widetilde{f}$$

for a 'nice' operator \widetilde{K}^+ . Indeed using the specific form of this slightly modified new K^+ we can proceed as before. We omit the details.

Remark 4.18. If $\mathcal{A}_3 = \emptyset$ we can write the above operator as $\widetilde{K}^+ = R^+_{\text{diag}} \widetilde{v}^+$, where Im $\widetilde{v}^+ \leq 0$, however in general the form of \widetilde{K}^+ is more complicated. Moreover the structure of the set of solutions to the equations $(1 + K^+)f = 0$ or $(1 + \widetilde{K}^+)\widetilde{f} = 0$ studied in the above proofs appears 'cleanest' in this case in the sense that 'spurious' solutions, i.e. resonances states, do not occur. On the other hand if $\mathcal{A}_3 \neq \emptyset$ resonances states could occur, and for this reason our theory of resolvent expansion at the two-cluster threshold is more complicated for $\mathcal{A}_3 \neq \emptyset$. See Chapter 5 for a systematic study applicable to the fastly decaying case for which $\mathcal{A} = \mathcal{A}_3$ and the effective inter-cluster interaction is $\mathcal{O}(|x_a|^{-3})$ for $a \in \mathcal{A}$. Resolvent expansion at the two-cluster threshold for some non-fastly decaying cases is treated in Chapter 6, see Remark 6.1.

Remark 4.19. We did not consider particles with spin in Theorem 4.17. However the theorem applies to cases with spin because this basically amount to restricting the operator H to suitable subspaces. However this is under the assumption that λ_0 is an (un-restricted) two-cluster threshold. It could be that λ_0 is a threshold not of this type but in the restricted sense is a two-cluster threshold. To treat such situation one would need a different procedure, although it could be that a somewhat similar treatment would work. Thus for example, to indicate a possible preliminary step, an obvious procedure for modifying S for fermions would be to use an anti-symmetrization of $S_a f_a = \varphi^a \otimes f_a$ to bring this vector into the fermionic subspace of $L^2(\mathbf{X})$.

Chapter 5

Resolvent asymptotics near a two-cluster threshold

This chapter is devoted to the asymptotics of the resolvent near an arbitrary twocluster threshold λ_0 for the N-body Schrödinger operator H. As discussed in Chapter 4, for physics models, the effective potential may decay like $\mathcal{O}(|x_a|^{-1})$ (slowly decaying case), $\mathcal{O}(|x_a|^{-2})$ (critically decaying case) or $\mathcal{O}(|x_a|^{-3})$ (fastly decaying case). We only study two cases: 1) the effective potential is fastly decaying; 2) the effective potential is slowly decaying and positive outside a compact set. The main difference between these two cases is that threshold resonance may appear in the first one and it is absent in the second one. For fastly decay effective potentials, we only study in full details two situations: i) $\lambda_0 = \Sigma_2$ is a double two-cluster threshold (the case $\lambda_0 = \Sigma_2$ is a non-multiple two-cluster threshold is easier and already studied in [Wa2]); ii) $\lambda_0 > \Sigma_2$ is a non-multiple two-cluster threshold of H. We calculate the leading term of resolvent expansions according to cases the threshold is an eigenvalue or/and a resonance. For positively slowly decreasing effective potentials we only study the lowest threshold and prove Gevrey estimates in exponentially weighted spaces for the remainder in the resolvent expansion (cf. ([AW, Wa6]) for one-body operators). Our study will have applications for the physics models of Sections 1.2 and 1.3. Parts of our study easily generalize to cover interesting cases for the physics models of Sections 1.2 and 1.3; our results will be stated with sketched proofs.

This chapter is organized as follows. In Section 5.1 we study the asymptotic expansion of the resolvent $R(z) = (H - z)^{-1}$ for rapidly decreasing effective potentials. For simplicity we assume that the intercluster spaces are of dimension three and analyze the notion of two-cluster threshold resonances in Subsection 5.1.1. We give some normalization conditions for resonance states according to spectral nature of the two-cluster threshold. These conditions are useful, they allow to explicitly compute some constants in physically interesting models (cf. Chapter 6). The resolvent expansions are given in Subsection 5.1.2 for $\lambda_0 = \Sigma_2$ (when Σ_2 is a double two-cluster threshold) and in Subsection 5.1.3 for $\lambda_0 > \Sigma_2$. In Section 5.2 we study the case where the effective potential is positive outside a compact set and slowly decaying at infinity. In this case threshold resonance is absent and we prove oneterm resolvent expansions with Gevrey estimates on the remainder at the lowest threshold Σ_2 . These results may be used to show sub-exponential time-decay of local energies. In Section 5.3 we study the combination of the previous sections for the physics models of Sections 1.2 and 1.3.

5.1 Rapidly decaying effective potentials

5.1.1 Two-cluster threshold resonances

Let $\lambda_0 \in \mathcal{T}$ be a two-cluster threshold of H. We analyze in this subsection spectral properties of H at λ_0 , assuming the effective potential obtained through the Grushin method decays sufficiently fastly. In addition, we assume the intercluster configuration \mathbf{X}_a is of dimension three for two-cluster decompositions a such that λ_0 is an eigenvalue of H^a .

5.1.1.1 The case $\lambda_0 = \Sigma_2$

Consider first the case

 $\lambda_0 = \Sigma_2$ is a double two-cluster threshold in the sense of Condition 2.5. (5.1)

Whence the lowest threshold Σ_2 is a two-cluster threshold of H and an eigenvalue of exactly two sub-Hamiltonians H^{a_j} , j = 1, 2. Here of course a_1 and a_2 are two-cluster decompositions of the N-body system. The above condition implies that $\Sigma_2 < 0$, however the requirement of Condition 2.5 that $\lambda_0 (= \Sigma_2)$ be a simple eigenvalue for both H^{a_1} and H^{a_2} is automatically fulfilled. It is also a part of Condition 2.5 that that Condition 1.6 are fulfilled for $a = a_1$ and $a = a_2$. In agreement with Section 2.3, let φ_j denote a corresponding (real) normalized eigenvector of H^{a_j} , j = 1, 2. Denote

$$\mathbf{X}^{j} = \mathbf{X}^{a_{j}}, \quad \mathbf{X}_{j} = \mathbf{X}_{a_{j}}, \quad j = 1, 2,$$

and x^{j} (resp., x_{j}) variables in \mathbf{X}^{j} (resp., in \mathbf{X}_{j}). Let

$$\mathcal{F}_j = \{g = \varphi_j(x^j) f_j(x_j) \mid f_j \in L^2(\mathbf{X}_j)\}, \quad j = 1, 2.$$

 \mathcal{F}_j is a closed subspace of $L^2(\mathbf{X})$. Assume, cf. (2.24),

$$\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$$
 and $n_j := \dim \mathbf{X}_j = 3$ for $j = 1, 2$.

Recall from Proposition 2.7 that $\mathcal{F} = \mathcal{F}_1 + \mathcal{F}_2$ is a closed subspace in $L^2(\mathbf{X})$. Let Π be the orthogonal projection from $L^2(\mathbf{X})$ onto \mathcal{F} and $\Pi' = 1 - \Pi$. We borrow the notation (4.1) and denote, in particular,

$$\mathcal{H}_s^k = H_s^k(\mathbf{X}_1) \oplus H_s^k(\mathbf{X}_2), \quad \mathcal{H}^k = \mathcal{H}_0^k, \quad \mathcal{H}_s = \mathcal{H}_s^0, \quad \mathcal{H} = \mathcal{H}_0^0.$$

Let $\mathcal{L}(k, s; k', s') = \mathcal{L}(\mathcal{H}_s^k, \mathcal{H}_{s'}^{k'})$, cf. Subsection 1.4.1.

Let $S = (S_1, S_2) : \mathcal{H} \to L^2(\mathbf{X})$ be defined by $Sf = S_1f_1 + S_2f_2$ for $f = (f_1, f_2)$ and with

$$S_j: L^2(\mathbf{X}_j) \to L^2(\mathbf{X}), \quad f_j \to S_j f_j = \varphi_j(x^j) \otimes f_j(x_j).$$

Then $S_j^* : L^2(\mathbf{X}) \to L^2(\mathbf{X}_j)$ is given by $S_j^* : f \to S_j^* f = \langle \varphi_j, f \rangle_j$, for j = 1, 2. Here $\langle ., . \rangle_j$ denotes the scalar product in $L^2(\mathbf{X}^j)$; the notation $\langle \cdot, \cdot \rangle$ will be used to denote the scalar product in $L^2(\mathbf{X})$ (or in $L^2(\mathbf{X}_j)$). One has

$$S^*S = 1 + \begin{pmatrix} 0 & s_{12} \\ s_{21} & 0 \end{pmatrix} \quad \text{on } \mathcal{H} = L^2(\mathbf{X}_1) \oplus L^2(\mathbf{X}_2),$$

where $s_{ij} \in \mathcal{L}(L^2(\mathbf{X}_j), L^2(\mathbf{X}_i)), i \neq j$, are given by $s_{ij}f_j = \langle \varphi_i, \varphi_j \otimes f_j \rangle_i$. Introduce for $z \in \mathbb{C}$ with $\operatorname{Im} z \neq 0$

$$E(z) = R'(z), E_{+}(z) = S - R'(z)HS, E_{-}(z) = S^{*} - S^{*}HR'(z), E_{\mathcal{H}}(z) = S^{*}(z - H + HR'(z)H)S$$

Then (recalling from Section 2.3)

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z).$$
(5.2)

 $E_{\mathcal{H}}(z): \mathcal{H} \to \mathcal{H}$ is computed in (2.48):

$$E_{\mathcal{H}}(z) = z - \lambda_0 - \begin{pmatrix} -\Delta_{x_1} + W_1(x_1) + K_{11}(z) & \check{s}_{12}(z) + W_{12} + K_{12}(z) \\ \check{s}_{21}(z) + W_{21} + K_{21}(z) & -\Delta_{x_2} + W_2(x_2) + K_{22}(z) \end{pmatrix},$$
(5.3)

where

$$\begin{split}
\check{s}_{ij}(z) &= \langle \varphi_i, \varphi_j \otimes (\lambda_0 - z - \Delta_{x_j}) \cdot \rangle_i, \\
W_{ij} &= \langle \varphi_i, I_j \varphi_j \otimes \cdot \rangle_i, \\
W_k(\cdot) &= W_{kk} = \langle \varphi_k, I_k \varphi_k \rangle_k(\cdot), \\
K_{ij}(z) &= -\langle \varphi_i, I_i R'(z) I_j(\varphi_j \otimes \cdot) \rangle_i.
\end{split}$$

Assume now $\lambda_0 \notin \sigma_d(H')$. Then by Lemma 2.10 R'(z) is holomorphic in z for z near λ_0 and $E_{\mathcal{H}}(\lambda_0)$ is well-defined. To simplify notation, denote $P = -E_{\mathcal{H}}(\lambda_0)$ and decompose this operator as $P = P_0 + U$, where

$$P_{0} = \begin{pmatrix} P_{1,0} & 0\\ 0 & P_{2,0} \end{pmatrix} := \begin{pmatrix} -\Delta_{x_{1}} & 0\\ 0 & -\Delta_{x_{2}} \end{pmatrix},$$
(5.5a)

$$U = \begin{pmatrix} U_{11} & U_{12} \\ U_{21} & U_{22} \end{pmatrix} := \begin{pmatrix} W_1 & W_{12} + \check{s}_{12}(\lambda_0) \\ W_{21} + \check{s}_{21}(\lambda_0) & W_2 \end{pmatrix} + K(\lambda_0).$$
(5.5b)

We impose the condition $\rho > \frac{1}{2}$ (where ρ is given in (1.22)) and (4.20) with $q_j(\theta) = 0$ for j = 1, 2, and hence (cf. Proposition 2.12) that at least for $\rho_0 = 3$

$$U: \mathcal{H}_{-s}^{1} \to \mathcal{H}_{\rho_{0}-s}^{-1} \quad \text{is continuous for any } s \in [0, \rho_{0}].$$
(5.6)

This leads us finally to impose the following set of conditions (including the previous ones):

$$\dim \mathbf{X}_1 = \dim \mathbf{X}_2 = 3, \quad \mathcal{F}_1 \cap \mathcal{F}_2 = \{0\} \text{ and } \lambda_0 \notin \sigma(H'), \tag{5.7a}$$

$$\rho > \frac{1}{2}$$
 and diag $(W_1, W_2) \in \mathcal{L}(\mathcal{H}^1, \mathcal{H}_{\rho_0}^{-1})$ for some $\rho_0 \ge 3$, (5.7b)

$$2 + 2\rho \ge \rho_0. \tag{5.7c}$$

Actually (5.7c) is imposed for Section 5.1.2 only; it is a convenient assumption relevant for the case $\rho_0 > 3$. Clearly (5.6) is fulfilled (in fact the operator U(z)of Section 5.1.2 fulfills $U(z) \in \mathcal{L}(1, -s; -1, \rho_0 - s)$ for $0 \leq s \leq \rho_0$ uniformly for z near zero, generalizing (5.6)). The conditions (5.7a) and (5.7b) will be imposed throughout the section.

Recalling the notation $H_{t^-}^1 = \bigcap_{s < t} H_s^1$ and $\mathcal{H}_{t^-}^1 = \bigcap_{s < t} \mathcal{H}_s^1$ for any $t \in \mathbb{R}$, (5.7a) and (5.7b) lead to the following effective version of Definition 4.6, see Theorems 4.7 and 4.15 (and (4.67)).

- **Definition 5.1.** (1) λ_0 is a resonance of H if the equation $Hu = \lambda_0 u$ admits a solution $u \in H^1_{(-1/2)^-} \setminus H^1$. The multiplicity of the resonance λ_0 is defined as the dimension of the quotient space $\ker(H \lambda_0)_{|H^1_{(-1/2)^-}} / \ker(H \lambda_0)_{|H^1}$.
 - (2) 0 is a resonance of P if the equation Pv = 0 has a solution $v \in \mathcal{H}^{1}_{(-1/2)^{-}} \setminus \mathcal{H}^{1}$. The multiplicity of the resonance zero is defined as the dimension of the quotient space ker $P_{|\mathcal{H}^{1}_{(-1/2)^{-}}} / \ker P_{|\mathcal{H}^{1}}$.

Similarly we can define (although this will not be needed) a notion of zeroresonance for $\tilde{P} = (S^*S)^{-1}P(S^*S)^{-1}$. Since $(S^*S)^{-1}$ and S^*S are continuous on \mathcal{H}^1_s for any s, zero is a resonance of \tilde{P} if and only if it is a resonance of P and their multiplicities are the same. Following [Ne, JK] we distinguish between four cases for the threshold λ_0 according to its spectral nature as follows.

- **Case 0** A regular point: λ_0 (0, resp.) is neither an eigenvalue nor a resonance of H (P, resp.).
- **Case 1** An exceptional point of the first kind: λ_0 (0, resp.) is a resonance but not an eigenvalue of H (P, resp.).
- **Case 2** An exceptional point of the second kind: λ_0 (0, resp.) is an eigenvalue but not a resonance of H (P, resp.).
- **Case 3** An exceptional point of the third kind: λ_0 (0, resp.) is simultaneously an eigenvalue and a resonance of H (P, resp.).

From Lemma 4.1 we deduce the following result.

Lemma 5.2. λ_0 is a resonance (resp., an eigenvalue) of H if and only if zero is a resonance (resp., an eigenvalue) of P and their multiplicities are the same.

We need the following simplified version of Theorem 4.7.

Theorem 5.3. Assume the conditions (5.7a) and (5.7b).

(a) Suppose $v = (v_1, v_2) \in \mathcal{H}^1_{(-1/2)^-}$ and Pv = 0. Then one has

$$v_j(x_j) = -\frac{\langle 1, U_{j1}v_1 + U_{j2}v_2 \rangle}{4\pi |x_j|} + w_j \text{ for } |x_j| \ge 1,$$
(5.8)

where $w_j \in L^2(\mathbf{X}_j)$; j = 1, 2. Moreover

$$v \in \mathcal{H} \iff \langle 1, U_{j1}v_1 + U_{j2}v_2 \rangle = 0; \quad j = 1, 2.$$

$$(5.9)$$

5.1.1.1. The case $\lambda_0 = \Sigma_2$

(b) Suppose $u \in H^1_{(-1/2)^-}$ is a solution to the equation $(H - \lambda_0)u = 0$. Let $v = T^*u \in \mathcal{H}^1_{(-1/2)^-}$. Then u is an eigenfunction of H at λ_0 if and only if

$$\langle 1, U_{j1}v_1 + U_{j2}v_2 \rangle = 0; \quad j = 1, 2.$$
 (5.10)

Proof. Since $n_j = 3$ and we assume that $q_j(\theta) = 0$ for j = 1, 2, in Theorem 4.7 the sets $\sigma_{j,1} = \{\frac{1}{2}\}$. Whence by this theorem we need to consider the functionals $l_{j,\frac{1}{2},1}$ applied to $v \in \mathcal{H}^1_{(-1/2)^-}$ solving Pv = 0. Since $q_j = 0$,

$$\sqrt{4\pi} \, l_{j,\frac{1}{2},1}(v) = \left\langle \phi_{\frac{1}{2}}(|y_j|) | y_j |^{-1}, \Delta_{y_j}(\chi_2 v_j)(y_j) \right\rangle_{L^2(\mathrm{d}y_j)}.$$

By integrating by parts in the integral to the right of this formula we verify that

$$\sqrt{4\pi} l_{j,\frac{1}{2},1}(v) = \left\langle 1, \Delta_{y_j}(\chi_2 v_j)(y_j) \right\rangle_{L^2(\mathrm{d}y_j)} = \left\langle 1, \Delta_{y_j} v_j(y_j) \right\rangle_{L^2(\mathrm{d}y_j)} = \left\langle 1, U_{j1}v_1 + U_{j2}v_2 \right\rangle,$$

showing (5.8). Trivially (5.9) is a consequence of (5.8). The above computation and Theorem 4.7 4) yield (b). \Box

Remark 5.4. Although this was not used above let us note that the equation Pv = 0 for $v = (v_1, v_2) \in \mathcal{H}^1_{(-1/2)^-}$ (equivalently) reads

$$v = -G_0 U v, \tag{5.11}$$

where $G_0 := \text{s-lim}_{z \to 0, z \notin [0,\infty)} (P_0 - z)^{-1}$ is computed as $G_0 = G_{0,1} \oplus G_{0,2}$ with $G_{0,j}$ specified by its integral kernel

$$G_{0,j}(x_j, y_j) = \frac{1}{4\pi |x_j - y_j|}; \quad j = 1, 2.$$

This assertion follows from an integration by parts argument as in the proof of Lemma 4.5 2). We note that the operators $\langle x \rangle^{-s} G_{0,j} \langle x \rangle^{-s'}$ are Hilbert-Schmidt operators for $s, s' > \frac{1}{2}$ and s + s' > 2, cf. [JK, Lemmas 2.2]. The asymptotics (5.8) may alternatively be derived from (5.11).

The condition $\langle 1, U_{j1}v_1 + U_{j2}v_2 \rangle = 0$ for j = 1, 2 is equivalent to the condition $\langle U^*c, v \rangle = 0$ for all $c \in \mathbb{C}^2$. Since $Hu = \lambda_0 u$ if and only if Pv = 0 with $v = T^*u$, the following result is an immediate consequence of Theorem 5.3.

Corollary 5.5. Assume conditions of Theorem 5.3. Let $u \in H^1_{(-1/2)^-}$ be a solution to the equation $Hu = \lambda_0 u$. Then u is an L^2 -eigenfunction of H if and only if

$$\langle TU^*c, u \rangle = 0 \quad \text{for any } c \in \mathbb{C}^2.$$
 (5.12)

The condition (5.12) can be rewritten as the system of equations

$$\int_{\mathbf{X}} \varphi_1 I_1 u dx + \langle 1, \check{s}_{12}(T^* u)_2 \rangle = 0, \qquad (5.13a)$$

$$\int_{\mathbf{X}} \varphi_2 I_2 u dx + \langle 1, \check{s}_{21}(T^*u)_1 \rangle = 0.$$
(5.13b)

In fact, using $u = Sv - R'(\lambda_0)HSv$ with $v = T^*u$ and writing $Sv = \varphi_1v_1 + \varphi_2v_2$, one has

$$\int_{\mathbf{X}} \varphi_1 I_1 u dx = \langle I_1 \varphi_1, Sv - R'(\lambda_0) H Sv \rangle$$
$$= \langle 1, (W_1 + K_{11}(\lambda_0)) v_1 \rangle + \langle 1, (W_{12} + K_{12}(\lambda_0)) v_2 \rangle,$$

showing, by using (5.12) with c = (1, 0), that

$$\int_{\mathbf{X}} \varphi_1 I_1 u dx + \langle 1, \check{s}_{12} v_2 \rangle = \langle 1, (UT^* u)_1 \rangle = 0.$$

Similarly (5.13b) follows from the computation

$$\int_{\mathbf{X}} \varphi_2 I_2 u dx + \langle 1, \check{s}_{21} v_1 \rangle = \langle 1, (UT^* u)_2 \rangle = 0.$$

By the same computations we see that (5.13a) and (5.13b) imply (5.12).

For $s \in (\frac{1}{2}, \frac{3}{2})$ we set

$$\mathcal{K} = \ker(1 + G_0 U) \subset \mathcal{H}^1_{-s}.$$
(5.14)

This space \mathcal{K} coincides with the subspace $\{v \in \mathcal{H}^1_{(-1/2)^-} | Pv = 0\}$ and is of finite dimension. By Theorem 5.3, if zero is a resonance of P, its multiplicity is at most two. Next we introduce the constant functions

$$g_j(x_i) = \frac{1}{2\sqrt{\pi}} \text{ for } x_i \in \mathbf{X}_i; \quad i = 1, 2.$$
 (5.15a)

Set $\mathfrak{g}_1 = (g_1, 0)$ and $\mathfrak{g}_2 = (0, g_2)$. For $v \in \mathcal{K}$, define $c(v) = (c_1(v), c_2(v)) \in \mathbb{C}^2$ by

$$c_i(v) = \langle \mathfrak{g}_i, Uv \rangle = \langle g_i, U_{i1}v_1 + U_{i2}v_2 \rangle; \quad i = 1, 2.$$
(5.15b)

Theorem 5.3 implies that $v \in \mathcal{K}$ is an eigenfunction of P if and only if c(v) = 0. In case zero is a resonance of P, we use the following normalization of resonance states:

• If zero resonance is simple, we denote by $\psi_1 \in \mathcal{K}$ a resonance state such that

$$|c(\psi_1)| = 1. \tag{5.16}$$

• If zero resonance is double, we denote by ψ_1 and ψ_2 two resonance states in \mathcal{K} such that

$$c(\psi_1) = (1,0) \text{ and } c(\psi_2) = (0,1),$$
 (5.17)

respectively.

If λ_0 is a resonance of H, let κ denote its multiplicity. Then $\kappa \in \{1, 2\}$ and the corresponding normalization condition for resonance states $u_j = (1 - R'(\lambda_0)H)S\psi_j$, $1 \leq j \leq \kappa$, of H reads

$$|\langle TU^* \mathfrak{g}_1, u_1 \rangle|^2 + |\langle TU^* \mathfrak{g}_2, u_1 \rangle|^2 = 1, \quad \text{if } \kappa = 1, \quad (5.18)$$

$$\langle TU^* \mathfrak{g}_i, u_j \rangle = \delta_{ij} \quad \text{for } 1 \le i, j \le 2, \qquad \text{if } \kappa = 2.$$
 (5.19)
With this choice of resonance functions we show in Section 5.1.2, see Theorem 5.14, that if Σ_2 is a resonance of multiplicity κ but not an eigenvalue of H, then

$$R(z) = \frac{\mathrm{i}}{\sqrt{z - \lambda_0}} \left(\sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z - \lambda_0|^{\epsilon}) \right)$$
(5.20)

in $\mathcal{L}(-1, s; 1, -s)$, s > 1, for z near λ_0 and $z - \lambda_0 \notin [0, \infty)$.

The case where Σ_2 is an eigenvalue (and possibly a resonance as well) is also studied, see Theorem 5.14 when $\rho_0 > 3$. For $\rho_0 = 3$ this requires an a priori weak decay property of the corresponding eigenfunctions, see Theorem 5.16. On the other hand with this additional assumption in fact we obtain the expansion of the resolvent up to second order. Expansions to higher orders usually require strong decay of the potential, see [JK, Wa2]. In particular in our setting this would mean that ρ_0 should be sufficiently big. We shall not pursue this direction, but rather mainly restrict our study to the case $\rho_0 = 3$ which indeed has relevance for the physics models, see discussions in Sections 1.2 and 1.3.

5.1.1.2 The case $\lambda_0 > \Sigma_2$

In this Subsection we consider the case

 $\lambda_0 > \Sigma_2$ is a non-multiple two-cluster threshold in the sense of Condition 2.4.

(5.21) This means there exists a unique $a_0 \in \mathcal{A} \setminus \{a_{\max}\}$ such that $\lambda_0 \in \sigma_{pp}(H^{a_0}) \setminus \{\Sigma_2\}$, and this cluster decomposition is a two-cluster decomposition obeying Condition 1.6. As in Section 2.2 we let *m* denote the multiplicity of eigenvalue λ_0 of H^{a_0} and $\{\varphi_1, \ldots, \varphi_m\}$ be an orthonormal basis of the associated eigenspace. Let Π be the projection in $L^2(\mathbf{X})$ defined by

$$\Pi g = \sum_{j=1}^{m} \varphi_j \otimes \langle \varphi_j, g \rangle_{L^2(\mathbf{X}^{a_0})}, \ g \in L^2(\mathbf{X}).$$
(5.22)

Put $\Pi' = 1 - \Pi$, $H' = \Pi' H \Pi'$ and $R'(z) = (H' - z)^{-1} \Pi'$. Let in this Subsection

$$\mathcal{H} = L^2(\mathbf{X}_{a_0}; \mathbb{C}^m), \quad \mathcal{H}_s^k = H_s^k(\mathbf{X}_{a_0}; \mathbb{C}^m), \quad \mathcal{H}_{s^-}^k = \cap_{t < s} \mathcal{H}_t^k; \quad k, s \in \mathbb{R}.$$

The scalar product in \mathcal{H} concerns only the x_{a_0} -variable and will be denoted as $\langle \cdot, \cdot \rangle_0$, but we shall allow ourselves (slightly abusively) to use the same notation for the inner product on $L^2(\mathbf{X}_{a_0})$. Define $S : \mathcal{H} \to L^2(\mathbf{X})$ by

$$S: f = (f_1, \dots, f_m) \to Sf = \sum_{j=1}^m \varphi_j(x^{a_0}) f_j(x_{a_0}).$$
 (5.23)

From Section 2.2 we have the representation formula for the resolvent,

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z) \text{ for } \operatorname{Im} z \neq 0, \qquad (5.24)$$

where

$$E(z) = R'(z),$$

$$E_{+}(z) = (1 - R'(z)I_{0})S,$$

$$E_{-}(z) = S^{*}(1 - I_{0}R'(z)),$$

$$E_{\mathcal{H}}(z) = (z - \lambda_{0}) - (P_{0} + S^{*}I_{0}S - S^{*}I_{0}R'(z)I_{0}S),$$

where $P_0 = -\Delta_{x_{a_0}} \mathbf{1}_m$ with $\mathbf{1}_m$ being the identity matrix of size m and $I_0 =$ $\sum_{b \not\subset a} V_b(x^b)$. We assume (5.27a) stated below. Then one has the following limiting absorption principle for H',

$$\forall s > \frac{1}{2}: \quad \|\langle x \rangle^{-s} R'(\lambda \pm i0) \langle x \rangle^{-s} \| \le C_s \text{ for } \lambda \text{ near } \lambda_0.$$
 (5.25)

This follows from Theorem 3.13 for $\breve{R}(z)$, recalling $R'(z) = \breve{R}(z)\Pi'$. Decompose $P^{\pm} = -E_{\mathcal{H}}(\lambda_0 \pm \mathrm{i}0)$ as

$$P^{\pm} = P_0 + U^{\pm}, \tag{5.26a}$$

where

$$U^{\pm} = S^* I_0 S - S^* I_0 R'(\lambda_0 \pm i0) I_0 S.$$
 (5.26b)

In addition to (5.21) we assume that

$$\dim \mathbf{X}_{a_0} = 3 \text{ and } \lambda_0 \notin \sigma_{\rm pp}(H'), \tag{5.27a}$$

$$\rho \ge 1 \text{ and } S^* I_0 S \in \mathcal{L}(\mathcal{H}^1_{-\rho_1}, \mathcal{H}^{-1}_{\rho_1}) \text{ with } \rho_1 \ge \frac{3}{2},$$
(5.27b)

$$\rho \ge \rho_1 - \frac{1}{2}.\tag{5.27c}$$

Recall that $\rho > 0$ is the rate of decay of the pair potentials as specified by (1.22). The condition (5.27c) will be convenient (although not essential) in Section 5.1.3; it is not used in the present section. (Note that with (5.27c) the operator U(z) of Section 5.1.3 fulfills $U(z) \in \mathcal{L}(1, -s; -1, s)$ for $s < \rho_1$ uniformly in z near zero with Im $z \neq 0$.) Imposing only (5.27b) obviously the property (5.27c) is valid for $\rho_1 = \frac{3}{2}$. This weak version of (5.27c) will be very useful.

According to discussions in Section 2.2, (5.25) and the condition (5.27b) imply

$$U^{\pm} \in \mathcal{L}(\mathcal{H}_{-s}^1, \mathcal{H}_s^{-1}) \text{ for } s < \frac{3}{2}.$$
(5.28)

The free resolvent $r_0(z) = (P_0 - z)^{-1}$ can be expanded in appropriately weighted spaces as

$$r_0(z) = G_0 + z^{\frac{1}{2}}G_1 + zG_2 + \cdots, \qquad (5.29)$$

see (5.94), and $K^{\pm} = G_0 U^{\pm}$ are compact operators on \mathcal{H}^1_{-s} for $s \in (\frac{1}{2}, \frac{3}{2})$. Using the integral kernel of G_0 , one deduces that $\ker_{\mathcal{H}_{-s}^1}(1+K^{\pm}) \subset \mathcal{H}_{(-1/2)^{-}}^1$. Set $\mathcal{K} =$ $\ker_{\mathcal{H}_{-s}^1}(1+K^+)$ for $s \in (\frac{1}{2}, \frac{3}{2})$. Let for any $s < \frac{3}{2}$

$$\mathcal{E}_{-s}^{\mathcal{G}} = \{ u \in L^{2}_{-s}(\mathbf{X}) | (H - \lambda_{0})u = 0, \Pi' u \in \mathcal{B}^{*}_{1/2,0}(\mathbf{X}) \}.$$

Theorem 5.6. Assume conditions (5.21), (5.27a) and (5.27b), and let $s \in (\frac{1}{2}, \frac{3}{2})$.

(a) For any $v \in \mathcal{K}$

$$R'(\lambda_0 + i0)I_0Sv = R'(\lambda_0 - i0)I_0Sv \quad and \quad U^+v = U^-v \in \mathcal{H}^{-1}_{(2\rho_1 - 1/2)^-}.$$
 (5.30)
Moreover

Moreover

$$\mathcal{K} = \ker_{\mathcal{H}_{-s}^{1}}(1 + K^{-}) = \ker_{\mathcal{H}_{-s}^{1}}(P^{+}) = \ker_{\mathcal{H}_{-s}^{1}}(P^{-}).$$
(5.31)

5.1.1.2. The case $\lambda_0 > \Sigma_2$

(b) For any $v = (v_1, \cdots, v_m) \in \mathcal{K}$ and $k = 1, \dots, m$

$$v_k(x_{a_0}) = -\frac{\langle 1, U_{k_1}^{\pm} v_1 + \dots + U_{k_m}^{\pm} v_m \rangle_0}{4\pi |x_{a_0}|} + w_k(x_{a_0}) \text{ for } |x_{a_0}| \ge 1, \quad (5.32)$$

where $w = (w_1, \cdots, w_m) \in \mathcal{H}$.

(c) Let $v \in \mathcal{K}$. Then $v \in \mathcal{H}$ if and only if

$$\forall c \in \mathbb{C}^m : \quad \langle c, U^{\pm} v \rangle_0 = 0. \tag{5.33}$$

Here $c \in \mathbb{C}^m$ is considered as an element of \mathcal{H}^1_{-s} , $s > \frac{3}{2}$.

- (d) $u \in \mathcal{E}_{-s}^{\mathcal{G}}$ if and only if $u = (S R'(\lambda_0 \pm i0)I_0S)v$ for some $v \in \mathcal{K}$. In this case v is uniquely given by $v = S^*u \in \mathcal{K}$.
- (e) $u \in \mathcal{E}_{-s}^{\mathcal{G}} \cap L^2(\mathbf{X})$ if and only if $v = S^* u \in \mathcal{K} \cap \mathcal{H}$.

Proof. We may argue essentially as in Section 4.2.4, in particular using Lemmas 4.10 and 4.11, although the setup is different there (being a multiple two-cluster case). The parametrix G_0 is of course simpler to use than the more general construction used in Section 4.2.4, but the resulting null space $\ker_{\mathcal{H}_{-s}^1}(1+K^+)$ is independent of the specific choice of parametrix to define K^+ , cf. Remark 5.4. Accepting a detour to Section 4.2.4 the identification of functionals would follow like in the proof of Theorem 5.3; alternatively (b) follows from the asymptotics of G_0 , cf. Remark 5.4.

We are lead to define the following effective version of the notion of threshold resonances.

- **Definition 5.7.** (1) λ_0 is a resonance of H if the equation $Hu = \lambda_0 u$ admits a solution in $\mathcal{E}^{\mathcal{G}}_{(-1/2)^-} := \bigcap_{s>1/2} \mathcal{E}^{\mathcal{G}}_{-s}$ which is not in H^1 . The multiplicity of the resonance λ_0 is defined as the dimension of the quotient space $\mathcal{E}^{\mathcal{G}}_{(-1/2)^-} / \ker(H \lambda_0)_{|H^1}$.
 - (2) 0 is a resonance of P^{\pm} if the equation $P^+v = 0$ (and therefore $P^-v = 0$, and vice versa) has a solution $v \in \mathcal{H}^1_{(-1/2)^-} \setminus \mathcal{H}^1$. The multiplicity of the resonance zero is defined as the dimension of the quotient space ker $P^+_{|\mathcal{H}^1_{(-1/2)^-}} / \ker P^+_{|\mathcal{H}^1}$.

Note the consequence of Theorem 5.6 that if zero is a resonance of P^{\pm} then its multiplicity does not exceed m. As in Subsection 5.1.1.1 we can, based on those definitions, introduce Cases 0-3, and Theorem 5.6 then shows that the threshold spectral properties of H at λ_0 are determined by those of P^{\pm} at zero. In fact, completely parallel to the case of the lowest threshold Σ_2 , λ_0 is a regular point (resp., an exceptional point of the first kind, the second kind, the third kind) of H if and only if zero is a regular point (resp., an exceptional point of the first kind, the second kind, the third kind) of P^{\pm} . Assume zero is a resonance of P^{\pm} . Then the quotient space $\mathcal{K}/\mathcal{H} := \mathcal{K}/(\mathcal{K} \cap \mathcal{H})$ has dimension $\kappa = \dim \mathcal{K}/\mathcal{H} \ge 1$. We call κ the multiplicity of the zero resonance of P^{\pm} . Let $\mu = \dim \mathcal{K}$. For $\phi = (v_1, \cdots, v_m) \in \mathcal{K}$, define $c(\phi) \in \mathbb{C}^m$ by

$$c(\phi) = \frac{1}{2\sqrt{\pi}} \left(\langle 1, U_{11}^{\pm} v_1 + \dots + U_{1m}^{\pm} v_m \rangle_0, \dots, \langle 1, U_{m1}^{\pm} v_1 + \dots + U_{mm}^{\pm} v_m \rangle_0 \right).$$
(5.34)

Theorem 5.6 (b) shows that $\phi \in \mathcal{K}$ is a resonance state of P^{\pm} if and only if $c(\phi) \neq 0$. Clearly $c(\cdot)$ is a linear action on \mathcal{K} . It follows that a family of resonance states $\{\psi_1, \ldots, \psi_k\}$ of P^{\pm} is linearly independent in \mathcal{K}/\mathcal{H} if and only if the family of vectors $\{c(\psi_1), \ldots, c(\psi_k)\}$ is linearly independent in \mathbb{C}^m .

Proposition 5.8. (a) Assume zero is a resonance of P^{\pm} with multiplicity κ . Then there exists a basis $\{\psi_1, \dots, \psi_{\kappa}\}$ of \mathcal{K}/\mathcal{H} such that

$$(c(\psi_i), c(\psi_j)) = \delta_{ij}, \quad i, j = 1, \cdots, \kappa,$$

$$(5.35)$$

where (\cdot, \cdot) is the scalar product of \mathbb{C}^m .

(b) Assume zero is a resonance but not an eigenvalue of P^{\pm} (i.e. that $\kappa = \mu$). Then the operator Q defined by

$$Q = \sum_{j=1}^{\kappa} \langle \psi_j, \cdot \rangle \psi_j : \mathcal{H}_s^{-1} \to \mathcal{H}_{-s}^1, s > \frac{1}{2},$$
(5.36)

is independent of the choice of basis $\{\psi_1, \ldots, \psi_\kappa\}$ of \mathcal{K} verifying (5.35): If $\{\psi'_1, \ldots, \psi'_\kappa\}$ is another basis of \mathcal{K} verifying (5.35), then one has

$$\sum_{j=1}^{\kappa} \langle \psi_j, \cdot \rangle \psi_j = \sum_{j=1}^{\kappa} \langle \psi'_j, \cdot \rangle \psi'_j.$$

Proof. (a). Let $\Phi = \{\phi_1, \ldots, \phi_\kappa\}$ be a basis of \mathcal{K}/\mathcal{H} . Then the rank of the matrix $C(\Phi) = (c(\phi_1), \ldots, c(\phi_\kappa)) \in \mathcal{M}_{m \times \kappa}$ is equal to κ , where $c(\phi_j)$ is considered as the j'th column of $C(\Phi)$. Consequently $C(\Phi)^*C(\Phi)$ is positive definite. Let $M_0 \in \mathcal{M}_{\kappa \times \kappa}(\mathbb{C})$ be the positive definite Hermitian matrix obeying $M_0^2 = (C(\Phi)^*C(\Phi))^{-1}$. Set $M_0 = (m_{ij})_{1 \leq i,j \leq \kappa}$ and define

$$\psi_i = \sum_{k=1}^{\kappa} m_{ki} \phi_k, \quad i = 1, \cdots, \kappa.$$
(5.37)

Then $\{\psi_1, \ldots, \psi_\kappa\}$ is also a basis of \mathcal{K}/\mathcal{H} .

Let $C(\Psi)$ be the matrix defined in the same way as $C(\Phi)$ with $\Phi = \{\phi_1, \ldots, \phi_\kappa\}$ replaced by $\Psi = \{\psi_1, \ldots, \psi_\kappa\}$. Then

$$c(\psi_j) = \sum_{k=1}^{\kappa} m_{kj} c(\phi_k); \quad j = 1, \cdots, \kappa$$

This shows $C(\Psi) = C(\Phi)M_0$ and

$$C(\Psi)^*C(\Psi) = M_0 C(\Phi)^* C(\Phi) M_0 = 1 \text{ in } \mathcal{M}_{\kappa \times \kappa}(\mathbb{C}).$$

It follows that $\Psi = \{\psi_1, \ldots, \psi_\kappa\}$ is a basis of \mathcal{K}/\mathcal{H} verifying the normalization condition (5.35).

(b). Let $t_{ij} \in \mathbb{C}, 1 \leq i, j \leq \kappa$, be such that

$$\psi'_i = \sum_{j=1}^{\kappa} t_{ij} \psi_j, \quad i = 1, \cdots, \kappa.$$

Then $c(\psi'_i) = \sum_{j=1}^{\kappa} t_{ij} c(\psi_j)$ in \mathbb{C}^m . The condition $(c(\psi'_i), c(\psi'_j)) = \delta_{ij}$ becomes

$$(c(\psi_i'), c(\psi_j')) = \sum_{l,m=1}^{\kappa} \overline{t_{il}} t_{jm}(c(\psi_l), c(\psi_m))$$
$$= \sum_{l=1}^{\kappa} \overline{t_{il}} t_{jl} = \delta_{ij}$$

for $i, j = 1, ..., \kappa$. This means that if both $\{\psi_1, ..., \psi_\kappa\}$ and $\{\psi'_1, ..., \psi'_\kappa\}$ satisfy (5.35), the matrix $T = (t_{ij})_{1 \le i,j \le \kappa}$ is unitary. We obtain

$$\sum_{i=1}^{\kappa} \langle \psi_i', \cdot \rangle \psi_i' = \sum_{l,m=1}^{\kappa} \left(\sum_{i=1}^{\kappa} \overline{t_{il}} t_{im} \right) \langle \psi_l, \cdot \rangle \psi_m = \sum_{l,m=1}^{\kappa} \delta_{lm} \langle \psi_l, \cdot \rangle \psi_m = \sum_{l=1}^{\kappa} \langle \psi_l, \cdot \rangle \psi_l.$$

The normalization condition of the resonance states (5.35) will be used to compute the leading term of the resolvent $R(z) = (H - z)^{-1}$ for z near λ_0 and Im $z \neq 0$ in the case λ_0 is a resonance but not an eigenvalue of H. The normalization condition (5.16) can be regarded as a special case of (5.35), and Proposition 5.8 (b) is also valid for the case $\lambda_0 = \Sigma_2$ is a double two-cluster threshold.

A main result on the resolvent expansion in Subsection 5.1.3 is for the case where λ_0 is not an eigenvalue of H, see Theorem 5.24, Case 0 and Case 1. However the case where λ_0 is an eigenvalue (and possibly a resonance as well) is also studied, cf. Subsection 5.1.1.1. For the physical models we then need an additional weak decay property of the corresponding L^2 -eigenfunctions, see Theorem 5.26. On the other hand with this additional assumption we obtain the expansion of the resolvent up to second order.

5.1.2 Resolvent asymptotics near the lowest threshold

In this subsection, we keep the conditions and the notation of Subsection 5.1.1.1, in particular, $\lambda_0 = \Sigma_2$ is a double two-cluster threshold. We want to study the asymptotics of the resolvent $R(z) = (H - z)^{-1}$ near Σ_2 , using the formula (5.2) for R(z). We let $P(z) = -E_{\mathcal{H}}(\lambda_0 + z)$ and decompose P(z) as

$$P(z) = P_0 + U(z) - z$$
 and $U(z) = U + zU_1 + z^2U_2(z)$ (5.38a)

with P_0 and U defined by (5.5a) and (5.5b), respectively, and

$$U_1 = -\begin{pmatrix} 0 & \langle \varphi_1, \varphi_2 \otimes \cdot \rangle_1, \\ \langle \varphi_2, \varphi_1 \otimes \cdot \rangle_2 & 0 \end{pmatrix} - K_1(\lambda_0),$$
(5.38b)

$$U_2(z) = -\sum_{j=2}^{\infty} z^{j-2} K_j(\lambda_0),$$
(5.38c)

$$K_j(\lambda_0) = (\langle \varphi_l, I_l(R'(\lambda_0))^{j+1} I_m \varphi_m \cdot \rangle)_{1 \le l, m \le 2}; \quad j \in \mathbb{N}.$$
(5.38d)

Since $K(\lambda_0 + z)$ is holomorphic in z near zero and continuous from $\mathcal{H}^1_{-1-\rho}$ to $\mathcal{H}^{-1}_{1+\rho}$ with $\rho > 0$ given by (1.22), the above power series converges in the space $\mathcal{L}(1, -1 - \rho; -1, 1 + \rho)$ for example. Note that

$$P(z) = P - z(1 - U_1) + z^2 U_2(z); \quad P = P_0 + U.$$
(5.39)

Differently from one-body Schrödinger operators, cf. [JK], P(z) is an operator pencil depending on the spectral parameter in a non-linear way. The following result is important for the existence of an asymptotic expansion of $P(z)^{-1}$ as $z \to 0$ in the case 0 is an eigenvalue of P.

Lemma 5.9. $1 - U_1$ is positive definite on \mathcal{H} .

Proof. Note firstly that $K_1(\lambda_0) \ge 0$. In fact, for $f = (f_1, f_2) \in \mathcal{H}$, since $R'(\lambda_0)$ is a bounded self-adjoint operator,

$$\langle f, K_1(\lambda_0) f \rangle = \sum_{i,j} \langle \varphi_i \otimes f_i, I_i(R'(\lambda_0))^2 I_j \varphi_j \otimes f_j \rangle = \|R'(\lambda_0)F\|^2 \ge 0,$$

where $F = \sum_{j=1}^{2} I_j \varphi_j \otimes f_j$. One can check that

$$1 + \left(\begin{array}{cc} 0 & \langle \varphi_1, \varphi_2 \otimes \cdot \rangle_1 \\ \langle \varphi_2, \varphi_1 \otimes \cdot \rangle_2 & 0 \end{array}\right) \ge 0.$$

In fact, cf. the proof of Proposition 2.7, its expectation value on f is given by

$$||f_1||_1^2 + ||f_2||_2^2 + 2\operatorname{Re}\langle\varphi_1 \otimes f_1, \varphi_2 \otimes f_2\rangle = ||\varphi_1 \otimes f_1 + \varphi_2 \otimes f_2||^2.$$

It follows that

$$\langle (1 - U_1)f, f \rangle = \|\varphi_1 \otimes f_1 + \varphi_2 \otimes f_2\|^2 + \|R'(\lambda_0)F\|^2 \ge 0.$$
 (5.40)

Therefore, $1 - U_1$ is non-negative on \mathcal{H} and $f \in \ker(1 - U_1)$ if and only if

$$\varphi_1 \otimes f_1 + \varphi_2 \otimes f_2 = 0$$
 and $R'(\lambda_0)F = 0$

In particular, since we assume (5.7a), ker $(1 - U_1)$ is reduced to $\{0\}$ and $1 - U_1$ is positive definite on \mathcal{H} .

5.1.2. Resolvent asymptotics near the lowest threshold

Set

$$r_0(z) = (P_0 - z)^{-1}; \quad z \notin [0, \infty).$$
 (5.41)

To obtain the asymptotic expansion of $P(z)^{-1}$ for $z \notin [0, \infty)$ and z near zero, we first use the resolvent equation

$$P(z)^{-1} = W(z)^{-1} r_0(z)$$
 where $W(z) := 1 + r_0(z)U(z).$ (5.42)

Next we shall apply the Grushin method to study the factor $W(z)^{-1}$.

Let $N \in \mathbb{N}$ and $s > N + \frac{1}{2}$. Since dim $\mathbf{X}_j = 3$, the free resolvent $r_0(z)$ can be expanded in $\mathcal{L}(-1, s; 1, -s)$ as

$$r_0(z) = G_0 + \sqrt{z}G_1 + \dots + z^{\frac{N}{2}}G_N + \mathcal{O}(|z|^{\frac{N}{2}+\epsilon}),$$
(5.43)

for some $\epsilon > 0$ depending on N and s, where $G_i = \text{diag}(G_{i,1}, G_{i,2})$ for $i \leq N$ are diagonal matrices which can be calculated explicitly, cf. [JK, Lemma 2.3]. In particular the integral kernel of $G_{i,j}$ for i = 0, 1 and j = 1, 2 are given by

$$G_{0,j}(x_j, y_j) = \frac{1}{4\pi |x_j - y_j|}, \quad G_{1,j}(x_j, y_j) = \frac{i}{4\pi}; \quad j = 1, 2.$$
 (5.44)

We also recall, cf. [JK, Lemmas 2.1–2.3], that $r_0(z) \in \mathcal{L}(-1, s'; 1, -s)$ if s, s' > 1/2 and s + s' > 2 with a Hölder continuous dependence in z at z = 0. Hence

$$r_0(z) - G_0 = \mathcal{O}(|z|^{\epsilon}) \in \mathcal{L}(-1, s'; 1, -s) \text{ for } s, s' > \frac{1}{2} \text{ and } s + s' > 2.$$
 (5.45)

We consider now expansions of the operator W(z) using the expansions (5.38a), (5.43) and (5.45).

Lemma 5.10. Under the conditions (5.7a)–(5.7c) the following expansions in z (with $z \notin [0, \infty)$) hold in terms of the quantities

$$W_0 = 1 + G_0 U, \quad W_1 = G_1 U, \quad W_2 = G_2 U + G_0 U_1,$$

and for some (small) positive number ϵ (depending on given parameters s and s').

(a) For $s > \frac{1}{2}$, $s \ge s'$ and $\rho_0 - s' > \max\{\frac{1}{2}, 2 - s\}$ $W(z) = W_0 + \mathcal{O}(|z|^{\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$ (5.46a)

(b) For $s > \frac{3}{2}$, $s \ge s'$ and $\rho_0 - s' > \frac{3}{2}$

$$W(z) = W_0 + \sqrt{z}W_1 + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$$
(5.46b)

(c) For $s > \frac{5}{2}$, $s \ge s'$ and $\rho_0 - s' > \frac{5}{2}$ $W(z) = W_0 + \sqrt{z}W_1 + zW_2 + \mathcal{O}(|z|^{1+\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$ (5.46c)

The assertion (a) follows from (5.45), while (b) and (c) follow from the bounds (5.43) with N = 1 and N = 2, respectively. In all cases we use (5.38a) as well.

Higher order asymptotic expansions can also be established under stronger decay assumptions on the effective potentials.

From the identity $\langle -U \cdot, \cdot \rangle = \langle P_0 \cdot, \cdot \rangle$ on $\mathcal{K} = \ker(1 + G_0 U)$ and an integration by parts (cf. (4.13)) it follows that $\langle -U \cdot, \cdot \rangle$ is a positive quadratic form on \mathcal{K} . Let

$$\mu = \dim \mathcal{K},$$

and let $\{\phi_1, \ldots, \phi_\mu\}$ be a basis of \mathcal{K} orthonormalized with respect to $\langle -U \cdot, \cdot \rangle$:

$$\langle -U\phi_i, \phi_j \rangle = \delta_{ij}$$

This normalization is a technical tool from [Wa2] which in general does not conform with (5.16) and (5.17). We make the convention that if zero is a resonance of Pwith multiplicity κ , ϕ_j for $1 \leq j \leq \kappa$ are resonance states and ϕ_j for $\kappa < j \leq \mu$ (for $\kappa < \mu$ only of course) are eigenstates of P.

In order to obtain the expansion of $W(z)^{-1}$, consider the Grushin problem

$$\mathcal{W}(z) = \begin{pmatrix} W(z) & \mathcal{S} \\ \mathcal{S}^* & 0 \end{pmatrix} : \mathcal{H}^1_{-s} \times \mathbb{C}^\mu \to \mathcal{H}^1_{-s} \times \mathbb{C}^\mu,$$

where $s \in (\frac{1}{2}, \rho_0 - \frac{1}{2}), \mathcal{S} : \mathbb{C}^{\mu} \to \mathcal{H}^1_{-s}$ is defined by

$$\mathcal{S}c = \sum_{j=1}^{\mu} c_j \phi_j, \quad c = (c_1, \dots, c_{\mu}) \in \mathbb{C}^{\mu},$$

and

$$\mathcal{S}^* f = (\langle -U\phi_1, f \rangle \dots, \langle -U\phi_\mu, f \rangle), \quad f \in \mathcal{H}^1_{-s}.$$

Note that \mathcal{S}^* can be regarded as the formal adjoint of \mathcal{S} with respect to the form $\langle -U \cdot, \cdot \rangle$ on \mathcal{H}^1_{-s} (it is not the Hilbert space adjoint). Define for s as above the map $Q: \mathcal{H}^1_{-s} \to \mathcal{H}^1_{-s}$ by

$$Qf = \sum_{j=1}^{\mu} \langle -U\phi_j, f \rangle \phi_j.$$

Then,

$$\mathcal{SS}^* = Q \text{ on } \mathcal{H}^1_{-s} \text{ and } \mathcal{S}^*\mathcal{S} = 1 \text{ on } \mathbb{C}^{\mu},$$

in particular Q is a projection in \mathcal{H}^1_{-s} .

One can then prove (see [JK, Wa2]) that

$$\mathcal{H}_{-s}^1 = \mathcal{K} \oplus \operatorname{ran}(1 + G_0 U), \tag{5.47}$$

and that Q projects onto \mathcal{K} relatively to the direct sum decomposition (5.47), in particular ker $Q = \operatorname{ran}(1 + G_0 U)$. Then of course Q' = 1 - Q is the projection from \mathcal{H}_{-s}^1 onto $\operatorname{ran}(1 + G_0 U)$ relatively to (5.47). It follows readily that $Q'(1 + G_0 U)Q'$ is bijective on $\operatorname{ran}(1 + G_0 U)$. Since $\operatorname{ran}(1 + G_0 U)$ is closed the operator

$$D_0 := (Q'(1+G_0U)Q')^{-1}Q'$$
(5.48)

exists and is continuous on \mathcal{H}_{-s}^1 . By an argument of perturbation based on (5.46a) it follows that for |z| small enough Q'W(z)Q' is invertible on $\operatorname{ran}(1 + G_0U)$ with continuous inverse. Let

$$D(z) = (Q'W(z)Q')^{-1}Q'.$$

The following expansions hold in $\mathcal{L}(1, -s; 1, -s)$ under the specified conditions and with

$$D_0 = (Q'W_0Q')^{-1}Q', \quad D_1 = -D_0W_1D_0, \quad D_2 = -D_0W_2D_0 + D_0W_1D_0W_1D_0.$$

$$D(z) = D_0 + \mathcal{O}(|z|^{\epsilon}), \text{ if } \frac{1}{2} < s < \rho_0 - \frac{1}{2}.$$
(5.49a)

$$D(z) = D_0 + \sqrt{z}D_1 + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}), \text{ if } \frac{3}{2} < s < \rho_0 - \frac{3}{2}.$$
 (5.49b)

$$D(z) = D_0 + \sqrt{z}D_1 + zD_2 + \mathcal{O}(|z|^{1+\epsilon}), \text{ if } \frac{5}{2} < s < \rho_0 - \frac{5}{2}.$$
 (5.49c)

Note that if $\mathcal{K} = \{0\}$, we have Q' = 1 and $D(z) = W(z)^{-1}$ and the asymptotic expansion of $W(z)^{-1}$ is given by the one of D(z). In the following we treat the case $\mathcal{K} \neq \{0\}$. The assertions (5.49b) and (5.49c) are not needed for leading term expansions which is our main interest.

Using the operator D(z), we can compute the inverse of $\mathcal{W}(z)$ as

$$\mathcal{W}(z)^{-1} = \begin{pmatrix} \mathcal{E}(z) & \mathcal{E}_{+}(z) \\ \mathcal{E}_{-}(z) & \mathcal{E}_{-+}(z) \end{pmatrix},$$
(5.50)

where

$$\begin{aligned} \mathcal{E}(z) &= D(z), \quad \mathcal{E}_{+}(z) = \mathcal{S} - D(z)W(z)\mathcal{S}, \\ \mathcal{E}_{-}(z) &= \mathcal{S}^{*} - \mathcal{S}^{*}W(z)D(z), \quad \mathcal{E}_{-+}(z) = \mathcal{S}^{*}(-W(z) + W(z)D(z)W(z))\mathcal{S}. \end{aligned}$$

One obtains from (5.50) a representation of the inverse of W(z),

$$W(z)^{-1} = \mathcal{E}(z) - \mathcal{E}_{+}(z)\mathcal{E}_{-+}(z)^{-1}\mathcal{E}_{-}(z).$$
(5.51)

 $\mathcal{E}_{\pm}(z)$ and $\mathcal{E}_{-+}(z)$ can be expanded similarly as D(z), that is

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \sqrt{z}\mathcal{E}_{\pm,1} + z\mathcal{E}_{\pm,2} + \dots$$
$$\mathcal{E}_{-+}(z) = \mathcal{E}_{-+,0} + \sqrt{z}\mathcal{E}_{-+,1} + z\mathcal{E}_{-+,2} + \dots$$

More precisely one has the following result.

Lemma 5.11. Assume (5.7a)–(5.7c).

(a) One has in $\mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}^{1}_{-s})$ (for the + case) or $\mathcal{L}(\mathcal{H}^{1}_{-s}, \mathbb{C}^{\mu})$ (for the - case):

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \mathcal{O}(|z|^{\epsilon}), \text{ if } s > \frac{1}{2}.$$
 (5.52a)

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \sqrt{z}\mathcal{E}_{\pm,1} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}), \text{ if } s > \frac{3}{2}.$$
 (5.52b)

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \sqrt{z}\mathcal{E}_{\pm,1} + z\mathcal{E}_{\pm,2} + \mathcal{O}(|z|^{1+\epsilon}), \text{ if } s > \frac{5}{2} \text{ and } \rho_0 > 4.$$
 (5.52c)

Here

$$\begin{aligned} \mathcal{E}_{+,0} &= \mathcal{S}, \quad \mathcal{E}_{-,0} &= \mathcal{S}^*, \\ \mathcal{E}_{+,1} &= -D_0 W_1 \mathcal{S}, \quad \mathcal{E}_{-,1} &= -\mathcal{S}^* W_1 D_0, \\ \mathcal{E}_{+,2} &= -(D_0 W_2 + D_1 W_1) \mathcal{S}, \quad \mathcal{E}_{-,2} &= -\mathcal{S}^* (W_2 D_0 + W_1 D_1). \end{aligned}$$

(b) One has in $\mathcal{M}_{\mu \times \mu}(\mathbb{C})$:

$$\mathcal{E}_{-+}(z) = \sqrt{z}\mathcal{E}_{-+,1} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}).$$
(5.53a)

$$\mathcal{E}_{-+}(z) = \sqrt{z}\mathcal{E}_{-+,1} + z\mathcal{E}_{-+,2} + \mathcal{O}(|z|^{1+\epsilon}), \text{ if } \rho_0 > 3.$$
 (5.53b)

Here

$$\begin{aligned} \mathcal{E}_{-+,1} &= -\mathcal{S}^* W_1 \mathcal{S}, \\ \mathcal{E}_{-+,2} &= -\mathcal{S}^* (W_2 - W_1 D_0 W_1) \mathcal{S} \end{aligned}$$

Proof. (a). Recall that $\mathcal{S} \in \mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}_{-s'}^1)$ for any $s' > \frac{1}{2}$. Since $W_0 \mathcal{S} = 0$, using (5.46a)-(5.46c), the following expansions in $\mathcal{L}(\mathbb{C}^{\mu}; \mathcal{H}_{-s}^1)$ hold.

$$\forall s > \frac{1}{2}: \quad W(z)\mathcal{S} = \mathcal{O}(|z|^{\epsilon})\mathcal{S}.$$
(5.55a)

$$\forall s > \frac{3}{2}: \quad W(z)\mathcal{S} = \sqrt{z}W_1\mathcal{S} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}).$$
(5.55b)

$$\forall s > \frac{5}{2}$$
: $W(z)\mathcal{S} = (\sqrt{z}W_1 + zW_2)\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}), \quad \text{if } \rho_0 > 3.$ (5.55c)

The expansions for $\mathcal{E}_{+}(z)$ follows from (5.49a), (5.49b) and (5.55a)-(5.55c). The results for $\mathcal{E}_{-}(z)$ can be proved in a similar way.

(b). For (5.53a) we use (5.46a), (5.46b) and (5.49a). Note that indeed since $W_0 \mathcal{S} = \mathcal{S}^* W_0 = 0$ and $D(z) : \mathcal{H}_{-s}^1 \to \mathcal{H}_{-s}^1$ is uniformly bounded for any $s \in (\frac{1}{2}, \rho_0 - \frac{1}{2})$, one obtains

$$\mathcal{S}^*W(z)\mathcal{S} = \sqrt{z}\mathcal{S}^*W_1\mathcal{S} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}), \qquad (5.56a)$$

$$\mathcal{S}^*W(z)D(z)W(z)\mathcal{S} = \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}).$$
(5.56b)

Clearly (5.53a) follows from (5.56a) and (5.56b).

For (5.53b) we use (5.46b), (5.46c) and (5.49a). Note that

$$\mathcal{S}^*W(z)\mathcal{S} = \sqrt{z}\mathcal{S}^*W_1\mathcal{S} + z\mathcal{S}^*W_2\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}), \qquad (5.57a)$$

$$\mathcal{S}^*W(z)D(z)W(z)\mathcal{S} = z\mathcal{S}^*W_1D(z)W_1\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon})$$

= $z\mathcal{S}^*W_1D_0W_1\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}).$ (5.57b)

Clearly (5.53b) follows from (5.57a) and (5.57b).

If zero is an eigenvalue but not a resonance, then $\phi_j \in L^2$ for all j and S is continuous from \mathbb{C}^{μ} to \mathcal{H} , and by Theorem 5.3 the composition $W_1 \mathcal{S} = 0$.

In the case $\rho_0 > 3$ the asymptotics (5.53b) then amounts to the statement

$$\mathcal{E}_{-+}(z) = -z\mathcal{S}^* W_2 \mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}).$$
(5.58)

For $\rho_0 = 3$ the right hand sides of (5.53b) and (5.58) make sense since S maps to \mathcal{H} , however we dont know if these asymptotics still hold in that case. In fact we only know the following weaker (and too poor) assertion for $\rho_0 = 3$,

$$\forall \epsilon > 0: \quad \mathcal{E}_{-+}(z) = \mathcal{O}(|z|^{1-\epsilon}). \tag{5.59}$$

To show (5.59) we apply (5.46c) with s' = 0 and $s \in (\frac{5}{2}, 3]$ and conclude that

$$\mathcal{S}^*W(z)\mathcal{S} = z\mathcal{S}^*W_2\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}) = \mathcal{O}(|z|).$$

Next by (5.46b),

$$W(z)\mathcal{S} = \mathcal{O}(|z|^{\frac{1}{2}+\epsilon})$$

in $\mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}_{-s}^{1})$ for any $s > \frac{3}{2}$ and with $\epsilon = \epsilon(s) > 0$. We apply (5.49a) with an s taken close to $\frac{3}{2}$. Then we argue that for any small $\delta > 0$

$$\mathcal{S}^*W(z) = \mathcal{O}(|z|^{\frac{1}{2}-\kappa}) \text{ in } \mathcal{L}(\mathcal{H}^1_{-s}.\mathbb{C}^{\mu}), \ \kappa := (s - \frac{3}{2} + \delta)/2, \tag{5.60}$$

Given (5.60) it follows that

$$\mathcal{S}^*W(z)D(z)W(z)\mathcal{S} = \mathcal{O}(|z|^{1+\epsilon-\kappa})$$

showing that

$$\mathcal{E}_{-+}(z) + \mathcal{S}^* W(z) \mathcal{S} = \mathcal{O}(|z|^{1+\epsilon-\kappa}) = \mathcal{O}(|z|^{1-\kappa}).$$

In particular (5.59) holds.

The bound (5.60) follows by interpolating the bounds

$$\mathcal{S}^*W(z) = \mathcal{O}(|z|^{\epsilon'}) \text{ in } \mathcal{L}(\mathcal{H}^1_{-\frac{5}{2}+\delta}, \mathbb{C}^{\mu}),$$
$$\mathcal{S}^*W(z) = \mathcal{O}(|z|^{\frac{1}{2}+\epsilon'}) \text{ in } \mathcal{L}(\mathcal{H}^1_{-\frac{3}{2}+\delta}, \mathbb{C}^{\mu}),$$

in turn valid due to (5.46a) and (5.46b), respectively. (The interpolation requires the bounds with $\epsilon' = 0$ only.)

Calculation of $\mathcal{E}_{-+,j}$, j = 1, 2. To compute explicitly these leading terms, we distinguish between different situations according to the spectral properties of the threshold zero.

Case 1. Suppose zero is a resonance, but not an eigenvalue of P. In this case, $\mu = 1$ or 2 and

$$\mathcal{E}_{-+,1} = (\langle U\phi_i, G_1 U\phi_j \rangle)_{1 \le i,j \le \mu}.$$

If $\mu = 1$, $\mathcal{E}_{-+,1} = \langle U\phi_1, G_1U\phi_1 \rangle_0$. Note that $G_1 = \text{diag}(G_{1,1}, G_{1,2})$ with $G_{1,i}$, i = 1, 2, given by the rank-one operator

$$G_{1,i} = \mathbf{i} \langle g_i, \cdot \rangle g_i, \tag{5.61}$$

where g_i is the constant function in x_i introduced in (5.15a). Using (5.15b) we then obtain

$$G_1 U \phi_1 = \mathbf{i}(c_1(\phi_1)g_1, c_2(\phi_1)g_2)$$

and

$$\mathcal{E}_{-+,1} = \mathbf{i}(|c_1(\phi_1)|^2 + |c_2(\phi_1)|^2) = \mathbf{i}|c(\phi_1)|^2$$
(5.62)

with $c(\phi_1) = (c_1(\phi_1), c_2(\phi_1)) \in \mathbb{C}^2$. If $\mu = 2$, a similar computation gives

$$G_1 U \phi_j = i(c_1(\phi_j)g_1, c_2(\phi_j)g_2); \quad j = 1, 2.$$

Therefore

$$\langle U\phi_i, G_1U\phi_j \rangle = i(\overline{c_1(\phi_i)}c_1(\phi_j) + \overline{c_2(\phi_i)}c_2(\phi_j)).$$

It follows that

$$\mathcal{E}_{-+,1} = (\langle U\phi_i, G_1 U\phi_j \rangle)_{1 \le i,j \le 2} = \mathbf{i}B_0$$
(5.63)

where $B_0 = C_0^* C_0$ and

$$C_0 = \begin{pmatrix} c_1(\phi_1) & c_1(\phi_2) \\ c_2(\phi_1) & c_2(\phi_2) \end{pmatrix}$$
(5.64)

Summing up, one obtains in Case 1

$$\mathcal{E}_{-+,1} = \mathrm{i}B_0 \tag{5.65}$$

where B_0 is a $\mu \times \mu$ matrix given by

$$B_0 = |c(\phi_1)|^2$$
 if $\mu = 1$, and $B_0 = C_0^* C_0$ if $\mu = 2$. (5.66)

In both cases B_0 is invertible due to Theorem 5.3. The explicit formula of $\mathcal{E}_{-+,2}$ is not needed for the leading term of the resolvent expansion in Case 1.

Case 2. Suppose zero is an eigenvalue, but not a resonance of P. In this case, all ϕ_j 's are eigenfunctions and by Theorem 5.3 on the characterization of resonance states, one has $W_1 S = 0$ which implies

$$\mathcal{E}_{-+,1} = -\mathcal{S}^* W_1 \mathcal{S} = 0. \tag{5.67}$$

Assume $\rho_0 > 3$, so that (5.53b) and (5.58) apply. This means more explicitly that

$$\mathcal{E}_{-+,2} = (\langle U\phi_i, (G_2U + G_0U_1)\phi_j \rangle)_{i,j=1,\dots,\mu}$$

Moreover, since $\phi_j \in L^2$, $j = 1, \ldots, \mu$, we can check as in [JK] that

$$\langle U\phi_i, G_2 U\phi_j \rangle = \langle \phi_i, \phi_j \rangle. \tag{5.68}$$

In fact, writing

$$G_2 = z^{-1}(r_0(z) - G_0 - \sqrt{z}G_1) + \mathcal{O}(|z|^{\epsilon})$$

for Im z > 0 and z near zero, one obtains using the 1st resolvent equation that

$$G_2 U \phi_j = -r_0(z) \phi_j + \mathcal{O}(|z|^{\epsilon}) U \phi_j,$$

and

$$\langle U\phi_i, G_2 U\phi_j \rangle = -\langle r_0(\overline{z})U\phi_i, \phi_j \rangle + \mathcal{O}(|z|^{\epsilon})$$

Similarly $r_0(\overline{z})U\phi_i = -\phi_i - \overline{z}r_0(\overline{z})\phi_i$, and by taking the limit $z = i\gamma \to 0$ with $\gamma \in \mathbb{R}_+$, we indeed obtain (5.68).

It is clear that $\langle U\phi_i, G_0U_1\phi_j\rangle = -\langle \phi_i, U_1\phi_j\rangle$. Therefore in Case 2, one has

$$\mathcal{E}_{-+,1} = 0, \quad \mathcal{E}_{-+,2} = (\langle \phi_i, (1 - U_1) \phi_j \rangle)_{1 \le i,j \le \mu}.$$
 (5.69)

By Lemma 5.9, $(\langle \phi_i, (1-U_1)\phi_j \rangle)_{1 \le i,j \le \mu}$ is a positive definite matrix.

Case 3. Suppose zero is both a resonance and an eigenvalue of P. Let κ be the multiplicity of zero resonance of P. Then $\kappa = 1$ or 2 and $\kappa < \mu$. For $\kappa + 1 \le j \le \mu$, ϕ_j is an eigenfunction and therefore

$$W_1 \phi_j = G_1 U \phi_j = 0, \quad j = \kappa + 1, \cdots, \mu.$$
 (5.70)

 $\mathcal{E}_{-+,1}$ can be computed as in Case 1. One has in $\mathcal{M}_{\mu \times \mu}(\mathbb{C})$

$$\mathcal{E}_{-+,1} = \begin{pmatrix} iB_0 & 0 & \cdots & 0\\ 0 & 0 & \cdots & 0\\ \vdots & \vdots & \ddots & \vdots\\ 0 & 0 & \cdots & 0 \end{pmatrix}$$
(5.71)

where B_0 is the $\kappa \times \kappa$ matrix given by (5.66) with μ replaced by κ .

Using the method of Case 2 and taking notice of (5.70), we find that under the condition $\rho_0 > 3$

$$\mathcal{E}_{-+,2} = \begin{pmatrix} \mathcal{E}_{11}^{(2)} & \mathcal{E}_{12}^{(2)} \\ \mathcal{E}_{21}^{(2)} & \mathcal{E}_{22}^{(2)} \end{pmatrix},$$
(5.72)

where

$$\mathcal{E}_{11}^{(2)} = (\langle U\phi_i, (W_2 - W_1 D_0 W_1)\phi_j \rangle)_{1 \le i,j \le \kappa},$$

$$\mathcal{E}_{12}^{(2)} = (\langle U\phi_i, W_2\phi_j \rangle)_{1 \le i \le \kappa, \kappa+1 \le j \le \mu},$$

$$\mathcal{E}_{21}^{(2)} = (\langle U\phi_i, W_2\phi_j \rangle)_{\kappa+1 \le i \le \mu, 1 \le j \le \kappa},$$

$$\mathcal{E}_{22}^{(2)} = (\langle \phi_i, (1 - U_1)\phi_j \rangle)_{\kappa+1 \le i,j \le \mu}.$$

If 0 is an eigenvalue of P we let Π_0 denote the spectral projection in \mathcal{H} onto the zero-eigenspace of this operator. The quantity Π_0 enters for Cases 2 and 3 below.

Proposition 5.12. The following asymptotics as $z \to 0$ and $z \notin [0, \infty)$ hold in $\mathcal{L}(1, -s; 1, -s)$ for s > 1 and close to 1.

Case 1. Suppose that zero is an exceptional point of the first kind. Then

$$W(z)^{-1} = \frac{\mathrm{i}}{\sqrt{z}}Q_0 + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
 (5.73)

Here

$$Q_0 = \sum_{j=1}^{\kappa} \langle -U\psi_j, \cdot \rangle \psi_j \tag{5.74}$$

with $\psi_j \in \mathcal{K}$ such that if $\kappa = 1$, ψ_1 verifies the normalization condition (5.16) and if $\kappa = 2$, ψ_1 and ψ_2 verify the normalization condition (5.17).

Cases 2 and 3. Suppose that zero is an exceptional point of the second or the third kind, respectively, and suppose that $\rho_0 > 3$. Then

$$W(z)^{-1} = z^{-1} (\Pi_0 (1 - U_1) \Pi_0)^{-1} \Pi_0 U + \mathcal{O}(|z|^{-1+\epsilon}).$$
(5.75)

Proof. <u>Case 1</u>. For any $s \in (\frac{1}{2}, \rho_0 - \frac{1}{2})$ and for some $\epsilon > 0$

$$\begin{aligned} \mathcal{E}(z) &= D_0 + \mathcal{O}(|z|^{\epsilon}), & \text{in } \mathcal{L}(1, -s; 1, -s), \\ \mathcal{E}_{\pm}(z) &= \mathcal{E}_{\pm,0} + \mathcal{O}(|z|^{\epsilon}), & \text{in } \mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}^1_{-s}) \text{ or } \mathcal{L}(\mathcal{H}^1_{-s}, \mathbb{C}^{\mu}) \\ \mathcal{E}_{-+}(z) &= \mathrm{i}\sqrt{z}B_0 + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}) \end{aligned}$$

where B_0 is given by (5.66). Note that $\mathcal{E}_{+,0} = \mathcal{S}$, $\mathcal{E}_{-,0} = \mathcal{S}^*$ and $\mathcal{SS}^* = Q$. It follows from (5.51) that

$$W(z)^{-1} = \frac{\mathrm{i}}{\sqrt{z}} \mathcal{S}B_0^{-1} \mathcal{S}^* + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
(5.76)

If $\kappa = 1, B_0 = |c(\phi_1)|^2$. Set

$$\psi_1 = \frac{1}{|c(\phi_1)|} \phi_1. \tag{5.77}$$

Then ψ_1 verifies (5.16) and

$$\mathcal{S}B_0^{-1}\mathcal{S}^* = \frac{1}{|c(\phi_1)|^2} \langle -U\phi_1, \cdot \rangle \phi_1 = \langle -U\psi_1, \cdot \rangle \psi_1 = Q_0$$

If $\kappa = 2$, then $B_0 = C_0^* C_0$. Take $\psi_1, \psi_2 \in \mathcal{K}$ such that

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = {}^t C_0^{-1} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}.$$
 (5.78)

For $f \in \mathcal{H}_{-s}^1$, set $(v_1, v_2) = (\langle -U\phi_1, f \rangle, \langle -U\phi_2, f \rangle) \in \mathbb{C}^2$. One has

$$\mathcal{S}B_0^{-1}\mathcal{S}^*f = (\phi_1, \phi_2)C_0^{-1}C_0^{*-1} \left(\begin{array}{c} v_1\\ v_2 \end{array}\right) = u_1\psi_1 + u_2\psi_2$$

where

$$\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = C_0^{*-1} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = \begin{pmatrix} \langle -U\psi_1, f \rangle \\ \langle -U\psi_2, f \rangle \end{pmatrix}.$$

It follows that

$$\mathcal{S}B_0^{-1}\mathcal{S}^*f = \langle -U\psi_1, f \rangle \psi_1 + \langle -U\psi_2, f \rangle \psi_2 = Q_0 f$$

To show that ψ_1 and ψ_2 verify the normalization condition (5.17), set $C_0 = (c_{ij})_{1 \le i,j \le 2}$, ${}^tC_0^{-1} = (d_{ij})_{1 \le i,j \le 2}$ and $\psi_k = (\psi_{k1}, \psi_{k2})$. Then

$$c_j(\psi_k) = \langle g_j, U_{j1}\psi_{k1} + U_{j2}\psi_{k2} \rangle = \sum_{m=1}^2 d_{km}c_{jm} = \delta_{jk}; \quad j, k = 1, 2.$$

Therefore ψ_1 and ψ_2 are resonance states verifying the normalization condition (5.17), and (5.73) is proved.

<u>*Case 2*</u>. By (5.69) and Lemmas 5.9 and 5.11

$$\mathcal{E}_{-+}(z) = zM_1 + \mathcal{O}(|z|^{1+\epsilon}),$$

where the matrix $M_1 = (\langle \phi_i, (1 - U_1)\phi_j \rangle)_{1 \le i,j \le \mu}$ is positive definite. We deduce from (5.51) that

$$W(z)^{-1} = -z^{-1}\mathcal{S}M_1^{-1}\mathcal{S}^* + \mathcal{O}(|z|^{-1+\epsilon})).$$

Let $\mathcal{S}^{\#}$ be the adjoint of \mathcal{S} with respect to the scalar product $\langle \cdot, \cdot \rangle$ of \mathcal{H} . Then $\mathcal{S}^{*} = -\mathcal{S}^{\#}U$ and $M_{1} = \mathcal{S}^{\#}(1 - U_{1})\mathcal{S}$. The orthogonal projection Π_{0} of \mathcal{H} onto the zero eigenspace of P can be expressed in terms of \mathcal{S} and $\mathcal{S}^{\#}$ as

$$\Pi_0 = \mathcal{S}(\mathcal{S}\mathcal{S}^\#)^{-1}\mathcal{S}^\# \tag{5.79a}$$

and obeys

$$\Pi_0 \mathcal{S} = \mathcal{S}, \quad \mathcal{S}^{\#} \Pi_0 = \mathcal{S}^{\#} \tag{5.79b}$$

Letting $T = \mathcal{S}(\mathcal{S}^{\#}(1-U_1)\mathcal{S})^{-1}\mathcal{S}^{\#}$ we compute using (5.79a) and (5.79b)

$$-\mathcal{S}M_1^{-1}\mathcal{S}^* = TU,$$

($\Pi_0(1 - U_1)\Pi_0$) $T = \mathcal{S}(\mathcal{S}\mathcal{S}^{\#})^{-1}\mathcal{S}^{\#} = \Pi_0,$
 $T(\Pi_0(1 - U_1)\Pi_0) = \mathcal{S}(\mathcal{S}\mathcal{S}^{\#})^{-1}\mathcal{S}^{\#} = \Pi_0.$

This leads to the identity $T = (\Pi_0(1 - U_1)\Pi_0)^{-1}\Pi_0$, and consequently that

$$-\mathcal{S}M_1^{-1}\mathcal{S}^* = TU = (\Pi_0(1-U_1)\Pi_0)^{-1}\Pi_0 U,$$

which proves (5.75)

<u>*Case 3*</u>. We use again (5.53b). We want to show $\mathcal{E}_{-+}(z)^{-1}$ exists and to calculate its leading term as $z \to 0$ and $z \notin [0, \infty)$. To do this, let $M(z) = \sqrt{z} \mathcal{E}_{-+,1} + z \mathcal{E}_{-+,2} \in \mathcal{M}_{\mu \times \mu}(\mathbb{C})$. Decompose M(z) into blocks:

$$M(z) = \left(\begin{array}{cc} \sqrt{z}M_{11}(z) & zM_{12} \\ zM_{21} & zM_{22} \end{array}\right)$$

with

$$M_{11}(z) = iB_0 + \sqrt{z}\mathcal{E}_{11}^{(2)}$$

$$M_{ij} = \mathcal{E}_{ij}^{(2)} \quad \text{for } (ij) \neq (11).$$

 M_{22} is positive definite by Lemma 5.9. The diagonal part of M(z) is invertible and one has

$$M(z)$$
diag $((\sqrt{z}M_{11}(z))^{-1}, (zM_{22})^{-1}) = \begin{pmatrix} 1 & a \\ b & 1 \end{pmatrix}$

where

$$a = M_{12}M_{22}^{-1}, \quad b = b(z) = \sqrt{z}M_{21}M_{11}(z)^{-1}.$$

Since $b = \mathcal{O}(\sqrt{|z|})$, the matrix $\begin{pmatrix} 1 & a \\ b & 1 \end{pmatrix}$ is invertible for $z \notin [0, \infty)$ and |z| small, and an elementary calculation gives

$$\left(\begin{array}{cc} 1 & a \\ b & 1 \end{array}\right)^{-1} = \left(\begin{array}{cc} (1-ab)^{-1} & -a(1-ba)^{-1} \\ -b(1-ab)^{-1} & (1-ba)^{-1} \end{array}\right).$$

Consequently we obtain a formula for the inverse of M(z) for $z \notin [0, \infty)$ with |z| small, and from that we read off that

$$zM(z)^{-1} = \begin{pmatrix} \sqrt{z}M_{11}(z)^{-1} & -\sqrt{z}M_{11}(z)^{-1}M_{12}M_{22}^{-1} \\ -\sqrt{z}M_{22}^{-1}M_{21}M_{11}(z)^{-1} & M_{22}^{-1} \end{pmatrix} + \mathcal{O}(\sqrt{|z|}).$$

Recalling our choice of basis $\{\phi_1, \ldots, \phi_\mu\}$ in \mathcal{K} the map \mathcal{S} splits naturally as $\mathcal{S} = \mathcal{S}_r \oplus \mathcal{S}_e : \mathbb{C}^{\kappa} \oplus \mathbb{C}^{\mu-\kappa} \to \mathcal{H}^1_{-s}$ with the range of \mathcal{S}_r (\mathcal{S}_e , respectively) included in the zero-resonance space (the zero-eigenspace, respectively) of P. Similarly, decompose $\mathcal{S}^* = \mathcal{S}_r^* \oplus \mathcal{S}_e^*$. Then,

$$\mathcal{S}_r \mathcal{S}_r^* = Q_r, \ \mathcal{S}_r^* \mathcal{S}_r = I_\kappa, \quad \mathcal{S}_e \mathcal{S}_e^* = Q_e, \ \mathcal{S}_e^* \mathcal{S}_e = I_{\mu-\kappa},$$

where $Q_r = \sum_{j=1}^{\kappa} \langle -U\phi_j, \cdot \rangle \phi_j$ and $Q_e = \sum_{j=\kappa+1}^{\mu} \langle -U\phi_j, \cdot \rangle \phi_j$. Since $\mathcal{E}_{-+}(z) = M(z) + o(|z|)$, we obtain that

$$-\mathcal{E}_{+}(z)\mathcal{E}_{-+}(z)^{-1}\mathcal{E}_{-}(z) = \frac{1}{z} \big(-\mathcal{S}_{e}M_{22}^{-1}\mathcal{S}_{e}^{*} + \mathcal{O}(|z|^{\epsilon}) \big)$$

Since $M_{22} = (\langle (1 - U_1)\phi_i, \phi_j \rangle)_{\kappa+1 \le i, j \le \mu}$, we can verify as in Case 2 that

$$-\mathcal{S}_e M_{22}^{-1} \mathcal{S}_e^* = (\Pi_0 (1 - U_1) \Pi_0)^{-1} \Pi_0 U.$$

Whence (5.75) is proved for Case 3 also.

Recall that

$$E_{\mathcal{H}}(\lambda_0 + z)^{-1} = -W(z)^{-1}(P_0 - z)^{-1}$$
, and $G_0U = -1$ on \mathcal{K} . (5.80)

The following result follows immediately from Proposition 5.12.

Proposition 5.13. The following asymptotics hold in $\mathcal{L}(-1, s; 1, -s)$ for any s > 1 and $z \notin [0, \infty)$ with |z| small.

Case 1. Suppose zero is an exceptional point of the first kind. Let κ be the multiplicity of zero resonance. If $\kappa = 1$, let ψ_1 be a resonance state normalized according to (5.16), and if $\kappa = 2$, let ψ_1 and ψ_2 be resonance states normalized according to (5.17). Then for some $\epsilon > 0$

$$E_{\mathcal{H}}(\lambda_0 + z)^{-1} = -\frac{\mathrm{i}}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle \psi_j, \cdot \rangle \psi_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}).$$
(5.81a)

Cases 2 and 3. Suppose that zero is an exceptional point of the second or the third kind, respectively, and suppose that $\rho_0 > 3$. Then for some $\epsilon > 0$

$$E_{\mathcal{H}}(\lambda_0 + z)^{-1} = z^{-1}(\Pi_0(1 - U_1)\Pi_0)^{-1}\Pi_0 + \mathcal{O}(|z|^{-1+\epsilon}).$$
 (5.81b)

Proof. We apply Proposition 5.12, (5.80) and the leading order expansion

$$\forall s > 1: \quad (P_0 - z)^{-1} = G_0 + \mathcal{O}(|z|^{\epsilon}) \in \mathcal{L}(-1, s; 1, -s).$$
(5.82)

We are now able to give the asymptotics of the resolvent of N-body operator H near its first threshold $\lambda_0 = \Sigma_2$. Let $R(\lambda_0 + z) = (H - \lambda_0 - z)^{-1}$. By (5.2),

$$R(\lambda_0 + z) = R'(\lambda_0 + z) - (1 - R'(\lambda_0 + z)H)SE_{\mathcal{H}}(\lambda_0 + z)^{-1}S^*(1 - HR'(\lambda_0 + z)).$$
(5.83)

Here $R'(\lambda_0 + z)$ is the reduced resolvent which under the condition (5.7a) is holomorphic in z in a neighborhood of zero. The proof of Lemma 5.2 shows that ψ is a zero-resonance state of P if and only if $u = (1 - R'(\lambda_0)H)S\psi$ is an λ_0 -resonance state of H and ϕ can be recovered from u by $\phi = T^*u$, where $T^* = (S^*S)^{-1}S^*$.

If λ_0 is a resonance of H its multiplicity is denoted by κ . If λ_0 is an eigenvalue of H the spectral projection of H onto the corresponding eigenspace is denoted by Π_H . Define

$$S_I f = I_1 \varphi_1 f_1 + I_2 \varphi_2 f_2, \quad f = (f_1, f_2) \in \mathcal{H}^1_{-\infty}.$$

Theorem 5.14. Assume (5.7a) and (5.7b). The following asymptotics hold for $R(\lambda_0 + z)$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ and $z \notin [0, \infty)$, and $\epsilon = \epsilon(s) > 0$.

Case 0. Suppose λ_0 is a regular point of H. Then one has

$$R(\lambda_0 + z) = R'(\lambda_0) + (S - R'(\lambda_0)S_I)D_0G_0(S^* - S_I^*R'(\lambda_0)) + \mathcal{O}(|z|^{\epsilon}).$$
(5.84a)

Case 1. Suppose λ_0 is an exceptional point of the first kind of H. Then

$$R(\lambda_0 + z) = \frac{\mathrm{i}}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}), \qquad (5.84\mathrm{b})$$

where $u_j = S\psi_j - R'(\lambda_0)S_I\psi_j$ with ψ_j given in Proposition 5.13. The quantities u_j are resonance states of H obeying (5.18) and (5.19) for $\kappa = 1$ and $\kappa = 2$, respectively.

Cases 2 and 3. Suppose that λ_0 is an exceptional point of the second or the third kind, respectively, and suppose that $\rho_0 > 3$. Then

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \mathcal{O}(|z|^{-1+\epsilon}).$$
(5.84c)

Proof. <u>Case θ </u>. One has

$$(1 - R'(\lambda_0 + z)H)S = S - R'(\lambda_0)S_I + \mathcal{O}(|z|) = E_+(\lambda_0) + \mathcal{O}(|z|),$$

in $\mathcal{L}(\mathcal{H}_{-s}^1, \mathcal{H}_{-s}^1)$ for $0 \le s \le \frac{3}{2}$. Similarly, $S^*(1 - HR'(\lambda_0 + z)) = S^* - S_I^*R'(\lambda_0) + \mathcal{O}(z)$ in $\mathcal{L}(\mathcal{H}_s^{-1}, \mathcal{H}_s^{-1})$. One obtains then from (5.80), (5.49a), (5.82) and (5.83) that

$$R(\lambda_0 + z) = R'(\lambda_0) + (S - R'(\lambda_0)S_I)D_0G_0(S^* - S_I^*R'(\lambda_0)) + \mathcal{O}(|z|^{\epsilon})$$
(5.85)

in $\mathcal{L}(-1, s; 1, -s)$ for s > 1, proving (5.84a).

<u>*Case 1*</u>. One obtains from (5.83) and Proposition 5.13 that

$$R(\lambda_0 + z) = -E_+(\lambda_0)E_{\mathcal{H}}(\lambda_0 + z)^{-1}E_-(\lambda_0) + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon})$$

in $\mathcal{L}(-1, s; 1, -s)$ for s > 1. Let $u_j = E_+(\lambda_0)\psi_j$. Since $E_-(\lambda_0)^* = E_+(\lambda_0)$ it follows from (5.81a) that

$$E_{+}(\lambda_{0})E_{\mathcal{H}}(\lambda_{0}+z)^{-1}E_{-}(\lambda_{0}) = -\frac{\mathrm{i}}{\sqrt{z}}\sum_{j=1}^{\kappa} \langle u_{j}, \cdot \rangle u_{j} + (|z|^{-\frac{1}{2}+\epsilon}),$$

proving (5.84b).

<u>Cases 2 and 3</u>. We can apply (5.81b) and (5.83) to obtain

$$R(\lambda_0 + z) = -z^{-1}B + \mathcal{O}(|z|^{-1+\epsilon}),$$

where in terms of the spectral projection Π_0 of P (for the eigenvalue zero)

$$B = E_{+}(\lambda_{0})(\Pi_{0}(1 - U_{1})^{-1}\Pi_{0})^{-1}\Pi_{0}E_{-}(\lambda_{0})$$

It remains to check that B is equal to the spectral projection Π_H , cf. [Wa2]. Introducing $\tau = E_+(\lambda_0)\Pi_0$ it follows that $\tau^* = \Pi_0 E_-(\lambda_0)$, and by using the properties $R'(\lambda_0)S = S^*R'(\lambda_0) = 0$ and (5.40) we compute

$$\tau^* \tau = \Pi_0 (S^* (1 - HR'(\lambda_0)) (1 - R'(\lambda_0)H) S \Pi_0 = \Pi_0 (1 - U_1) \Pi_0$$

Consequently B can be written as

$$B = \tau (\tau^* \tau)^{-1} \tau^*.$$

It follows that B is an orthogonal projection with ran $B \subset \operatorname{ran} \tau$ and ker $B = \ker \tau^*$, and therefore in fact ran $B = \operatorname{ran} \tau$. Since ran τ is equal to the λ_0 -eigenspace of H, we conclude $B = \prod_H$ as wanted.

For the case where $\rho_0 = 3$ and λ_0 is an eigenvalue of H there is no information in Theorem 5.14. Furthermore we only extracted the leading term asymptotics. Assuming that all eigenfunctions of H at λ_0 have a certain (weak) power decay we can obtain a resolvent expansion for that case too, and that expansion will be to second order without further assumptions, see Theorem 5.16 stated below.

Remark 5.15. For possible higher order expansions the condition (5.7c) is needed and give limitations. For example it could be that (5.7b) is fulfilled for Coulombic systems (for which $\rho = 1$) with a big ρ_0 in some cases, but then the condition (5.7c) would read $\rho_0 \leq 4$ and we dont see how to avoid this restriction. More precisely, for example we can not improve (5.6) to be valid for any $\rho_0 > 4$ for such systems (not even with $s = \rho_0/2$ only). This is rooted in the fact that we do not know how to estimate the diagonal parts of the matrix $K(\lambda_0)$ better than $\mathcal{O}(\langle x_j \rangle^{-4})$ for $\rho = 1$. The main reason for not studying higher order resolvent expansions is that the conditions would be too strong to be interesting for Coulombic systems.

Theorem 5.16. Assume (5.7a), (5.7b) and $\rho_0 = 3$. Suppose λ_0 is an eigenvalue of H and that ran $\Pi_H \subset L_t^2$ for some t > 3/2. Then the following asymptotics hold for

 $R(\lambda_0 + z)$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ and $z \notin [0, \infty)$, and $\epsilon = \epsilon(s) > 0$.

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \frac{i}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}).$$
(5.86)

Here the second term on the right-hand side appears only if $\kappa \geq 1$, that is if λ_0 is a resonance of H, and in that case the states u_j constitute a basis of resonance states of H (recall that either $\kappa = 1$ and $\kappa = 2$). If on the other hand λ_0 is not a resonance of H, then $R(\lambda_0 + z) + z^{-1}\Pi_H$ has a limit in norm as $z \to 0$.

One can show Theorem 5.16 by mimicking the proof of the assertion on Case 1 in Theorem 5.14 with H replaced by $H_{\sigma} := H - \sigma \Pi_H$, $\sigma > 0$. Note that λ_0 is not an eigenvalue of H_{σ} . Letting $R_{\sigma}(\zeta) = (H_{\sigma} - \zeta)^{-1}$ we decompose

$$R(\lambda_0 + z) = R_{\sigma}(\lambda_0 + z) - z^{-1}\Pi_H + (z + \sigma)^{-1}\Pi_H, \qquad (5.87)$$

and conclude that it suffices to show that for a fixed small $\sigma>0$

$$R_{\sigma}(\lambda_0 + z) = \frac{\mathrm{i}}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
(5.88)

This can largely be shown as before since the last assumption in (5.7a) is valid with H' replaced by $H'_{\sigma} := \Pi' H_{\sigma} \Pi'$ for $\sigma > 0$ small (by an argument of perturbation). We can largely mimic the previous procedure, now for H_{σ} rather than for H, cf. Remark 3.21 2) which will be particularly relevant for similar constructions in the next sections. However we need to argue that the resonance states u_j in (5.88), which a priori are resonance states of H_{σ} , in fact also are resonance states of H. For that it suffices in turn to note that $\Pi_H u_j = 0$, which follows from the fact that $-\sigma \Pi_H u_j =$ $\Pi_H (H_{\sigma} - \lambda_0) u_j = 0$. If κ_{σ} denotes the dimension of the space of resonance states of H_{σ} , we see that $\kappa_{\sigma} \leq \kappa$. The converse $\kappa_{\sigma} \geq \kappa$ follows by modifying any given base of resonance states of H by projecting out the corresponding components in ran Π_H , yielding a base of resonance states of H_{σ} . Whence our use of the notation κ in (5.88), rather than the initially correct quantity κ_{σ} , is justified.

Another comment relates to (5.8). The added term $-\sigma \Pi_H$ is not local, and if we introduce the notation U_{σ} and P_{σ} for the corresponding reduced quantities, the corresponding version of (5.8) does not follow the same way unless ran $\Pi_H \subset L_t^2$ for some t > 2. (But we only assume this for some t > 3/2, so if $t \in (3/2, 2]$ a different argument is needed.) Nevertheless we have the following version of (5.9) for any $v = (v_1, v_2) \in \mathcal{H}^1_{(-1/2)^-}$ and $P_{\sigma}v = 0$:

$$v = 0 \iff \langle 1, U_{\sigma, j1} v_1 + U_{\sigma, j2} v_2 \rangle = 0; \quad j = 1, 2.$$
 (5.89)

This follows by noting that if the functionals to the right vanish on v, then the proof of (5.8) shows that $v \in \mathcal{H}^1_{(-1/2)^+}$. Then the corresponding $u \in H^1_{(-1/2)^+}$ (given by the inverse eigentransform), and this cannot hold unless $u \in L^2$ since by the above discussion $(H - \lambda_0)u = (H_{\sigma} - \lambda_0)u = 0$, and therefore we can conclude that $u \in L^2$ using Theorem 4.7. Since λ_0 is not an eigenvalue of H_{σ} it follows that u = 0 and then in turn that v = 0. With (5.89) in place we can mimic the proof of the assertion on Case 1 in Theorem 5.14 with H replaced by $H_{\sigma} = H - \sigma \Pi_H$ and conclude (5.88).

- **Remarks 5.17.** 1) We used, among others, the conditions $\lambda_0 = \Sigma_2 \notin \sigma_{\rm pp}(H')$ and $\mathcal{F}_1 \cap \mathcal{F}_2 = \{0\}$. The case $\lambda_0 \in \sigma_{\rm pp}(H')$ can be treated as in [Wa2] when Σ_2 is a non-multiple two-cluster threshold. The same statements as those of Theorem 5.14 remain true, but their proof requires a more complicated Grushin reduction, cf. Sections 2.4 and 4.3. If $\mathcal{F}_1 \cap \mathcal{F}_2 \neq \{0\}$, one can still reduce the spectral analysis of H at λ_0 to a one-body problem, cf. Sections 2.5 and 4.3. The conditions dim $\mathbf{X}_j = 3$ and $\rho_0 \geq 3$ of (5.7a) and (5.7b), as well as (5.7c), simplify the resolvent expansion at the threshold λ_0 and fit well with the physics models of Sections 1.2 and 1.3.
 - 2) The very last assertion of Theorem 5.16 only needs $\operatorname{ran} \Pi_H \subset L_t^2$ for some t > 1 (rather than this decay for some t > 3/2, as required in Theorem 5.16) due to the fact that λ_0 , under the given hypotheses, is neither an eigenvalue nor a resonance of H_{σ} .
 - 3) The cases where dim $\mathbf{X}_i \neq 3$ or the effective potential decays slowly or has a critically decaying part were studied in Chapter 4 and we obtained some threshold spectral properties of H. One may for these cases try to combine the existing results for one-body problems and the reduction made in Chapter 2 to establish the asymptotics of the resolvent of H near λ_0 . Although leaving this issue partly as an open question to the interested reader, see however Section 5.3, let us remark that the effectively slowly decaying potential cases corresponding to Subsections 4.1.2, 4.1.3, 4.2.2 and 4.2.3 are relatively easy to treat due to the fact that the threshold is not a resonance and eigenfunctions (if existing) have arbitrary polynomial decay (cf. Theorem 4.17 1) in the context of physics models). We make use of this simplicity in Chapter 6, see Remark 6.1. This is in the context of a possible higher two-cluster threshold subject to conditions of Subsections 4.1.2, 4.1.3, 4.2.2 and 4.2.3. However we do treat a special case with a critical decaying part of the effective potential, see Subsection 6.3.2. Using the theory of the present chapter we treat in Subsections 6.3.3 and 6.3.4 the fastly decaying case for the physics models with the occurrence of a resonance at a possible higher two-cluster threshold.

5.1.3 Resolvent asymptotics near higher two-cluster thresholds

Assume (5.21) and (5.27a)–(5.27c) as in Subsection 5.1.1.2. We shall tacitly use the notation from that subsection. Let $P(z) = -E_{\mathcal{H}}(\lambda_0 + z)$ for z near zero and Im $z \neq 0$. Then P(z) can be written as

$$P(z) = P_0 + U(z) - z, (5.90)$$

where $P_0 = -\Delta_{x_{a_0}} \mathbf{1}_m$ and $U(z) = S^* I_0 S - S^* I_0 R'(\lambda_0 + z) I_0 S$. Here $\mathbf{1}_m$ denotes the identity matrix in $\mathcal{M}_{m \times m}(\mathbb{C})$. Applying Theorem 3.18, one has for $\pm \operatorname{Im} z > 0$ and $j \in \mathbb{N}_0$ with $j < \rho_1$

$$U(z) = U^{\pm} + zU_1^{\pm} + \dots + z^j U_j^{\pm} + \mathcal{O}(|z|^{j+\epsilon}),$$
(5.91)

as bounded operators from \mathcal{H}_{-s}^1 to \mathcal{H}_s^{-1} for any $s < \rho_1 - j$ and for some $\epsilon > 0$ depending on s and ρ . Furthermore

$$U^{\pm} = S^* I_0 S - S^* I_0 R'(\lambda_0 \pm i0) I_0 S,$$

$$U_i^{\pm} = -S^* I_0 (R'(\lambda_0 \pm i0))^{j+1} I_0 S; \quad j \ge 1$$

See the proof of Lemma 5.19 for an elaboration for the relevant cases (we shall only need (5.91) for j = 0 and j = 1).

Recall that $\mathcal{K} = \ker_{\mathcal{H}_{-s}^1}(1 + G_0 U^{\pm})$ for $s \in (\frac{1}{2}, \frac{3}{2})$, but that this space is independent of such s as well as of the sign \pm .

Lemma 5.18. The quadratic form $q(\cdot) = \langle (1 - U_1^{\pm}) \cdot, \cdot \rangle_0$ defined on $\mathcal{K} \cap \mathcal{H}$ is positive definite and independent of sign \pm .

Proof. For $v \in \mathcal{K} \cap \mathcal{H}$ one has $\Pi' I_0 S v \in \mathcal{H}_{(3/2)^+}^{-1}$, allowing us to conclude that $R'(\lambda_0 \pm i0)^2 I_0 S v \in \mathcal{H}_{(-3/2)^-}^1$. Hence q is well-defined. Now $R'(\lambda_0 + i0) I_0 S v = R'(\lambda_0 - i0) I_0 S v$ and $R'(\lambda_0 \pm i0) I_0 S v \in L^2(\mathbf{X})$ (cf. Theorem 4.15, Theorem 5.6 (a) and the part of the proof of Theorem 5.6 based on microlocal resolvent estimates). These facts allow us to compute

$$q(v) = \langle (1 + S^* I_0 R' \lambda_0 + i0)^2 I_0 S v, v \rangle$$

= $||v||^2 + \langle R'(\lambda_0 + i0) I_0 S v, R'(\lambda_0 - i0) I_0 S v \rangle$
= $||v||^2 + ||R'(\lambda_0 + i0) I_0 S v||^2 \ge ||v||^2$,

proving that $q(\cdot)$ is positive definite on $\mathcal{K} \cap \mathcal{H}$.

We shall study the asymptotics of the resolvent $R(\lambda_0 + z) = (H - \lambda_0 - z)^{-1}$ as $z \to 0$ and $\pm \text{Im } z > 0$. Set

$$r_0(z) = (P_0 - z)^{-1}; \quad z \notin [0, \infty).$$
 (5.92)

To obtain the asymptotic expansion of $P(z)^{-1}$ for $z \notin \mathbb{R}$ and z near zero, we use as in Section 5.1.2 the resolvent equation

$$P(z)^{-1} = (1 + r_0(z)U(z))^{-1}r_0(z)$$
(5.93)

and apply the Grushin method to study $(1 + r_0(z)U(z))^{-1}$. For simplicity we shall mostly consider the case Im z > 0 only.

Let $N \ge 1$ and $s > N + \frac{1}{2}$. Similarly to the (partly scalar) case studied in Section 5.1.2, the free resolvent $r_0(z)$ can be expanded $\mathcal{L}(-1, s; 1, -s)$ as

$$r_0(z) = G_0 + \sqrt{z}G_1 + \dots + z^{\frac{N}{2}}G_N + \mathcal{O}(|z|^{\frac{N}{2}+\epsilon}),$$
(5.94)

for some $\epsilon > 0$ depending on N and s, where G_j is the $m \times m$ diagonal matrix (with operator-valued entries) whose integral kernel is given by

$$\frac{i^{j}}{4\pi}|x_{a_{0}} - y_{a_{0}}|^{j-1}\mathbf{1}_{m}; \quad j \in \mathbb{N}_{0}.$$
(5.95)

In addition to the higher order expansions (5.94) we record the zero'th order expansion (5.45) (with an obvious change of interpretation).

Since $U^+ = U^-$ on \mathcal{K} it follows that $\langle -U^+, \cdot \rangle$ is a quadratic form on \mathcal{K} . By the identity $\langle -U^+, \cdot \rangle = \langle P_0, \cdot \rangle$ and an integration by parts it follows that $\langle -U^+, \cdot \rangle$ is a positive quadratic form. Hence there exists a basis $\{\phi_1, \ldots, \phi_\mu\}$ of \mathcal{K} which is orthonormal with respect to $\langle -U^+, \cdot \rangle$. If zero is a resonance of P^+ , we denote by κ its multiplicity. We use the convention that the functions ϕ_j for $1 \leq j \leq \kappa$ are resonance states of P^+ while the other ϕ_j 's are eigenfunctions of P^+ .

With the above notation, we can apply the Grushin method to obtain asymptotic expansions of $W(z)^{-1}$ as $z \to 0$, Im z > 0, where

$$W(z) := 1 + r_0(z)U(z).$$

The leading order expansion of W(z) is given by

$$W(z) = 1 + G_0 U^+ + \mathcal{O}(|z|^{\epsilon}) \in \mathcal{L}(1, -s'; 1, -s),$$
(5.96)

for Im z > 0 and $\frac{1}{2} < s, s' < \rho_1$ and for some $\epsilon = \epsilon(s, s') > 0$. In particular this holds under the minimum conditions $\rho = 1$ and $\rho_1 = \frac{3}{2}$, see Lemma 5.19 (a) below and the outlined proof.

A higher order expansion of W(z) needs a stronger condition than $\rho_1 = \frac{3}{2}$ and therefore also $\rho > 1$. With such condition we can apply Theorem 3.18 on higher powers of the reduced resolvent and obtain the following extension of (5.96). The assertions (b) and (c) clearly exclude the interesting case $\rho = 1$. Nevertheless we will later show a higher order expansion of $W(z)\phi$ when applied to vectors $\phi \in \mathcal{K}$, even when $\rho = 1$, see the proof of Lemma 5.21. This will in fact yield a non-trivial resolvent expansion in the exceptional case of the first kind when $\rho = 1$.

Lemma 5.19. Suppose (5.21) and (5.27a)-(5.27c). In terms of the quantities

$$W_0 = 1 + G_0 U^+, \quad W_1 = G_1 U^+, \quad W_2 = G_2 U^+ + G_0 U_1^+,$$
 (5.97)

the following expansions in z (with Im z > 0) hold for some $\epsilon = \epsilon(s, s') > 0$.

(a) For $s, s' \in (\frac{1}{2}, \rho_1), s \ge s'$,

$$W(z) = W_0 + \mathcal{O}(|z|^{\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$$
(5.98a)

(b) If $\rho_1 > \frac{3}{2}$, then for $s, s' \in (\frac{3}{2}, \rho_1)$ with $s \ge s'$,

$$W(z) = W_0 + \sqrt{z}W_1 + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$$
(5.98b)

(c) If $\rho_1 > \frac{5}{2}$, then for $s \in (\frac{5}{2}, \rho_1)$, $s \ge s'$ and $s' \in (\frac{3}{2}, \rho_1 - 1)$ $W(z) = W_0 + \sqrt{z}W_1 + zW_2 + \mathcal{O}(|z|^{1+\epsilon}) \in \mathcal{L}(1, -s'; 1, -s).$ (5.98c)

Proof. By Theorem 3.18 there exists for any $j \in \mathbb{N}$ and $\epsilon > 0$ a constant C > 0 such that

$$\|\langle x \rangle^{\frac{1}{2}-j-\epsilon} R'(\lambda_0+z)^j \langle x \rangle^{\frac{1}{2}-j-\epsilon} \| \le C$$

and therefore

$$\|\langle x\rangle^{-\frac{3}{2}-\epsilon} \big(R'(\lambda_0+z) - R'(\lambda_0+\mathrm{i}0) \big) \langle x\rangle^{-\frac{3}{2}-\epsilon} \| \le C|z|$$

for z near zero with Im z > 0. Using complex interpolation we then obtain the following Hölder estimate for the reduced resolvent for $0 < r' < r \le 1$,

$$\|\langle x \rangle^{-\frac{1}{2}-r} \left(R'(\lambda_0 + z) - R'(\lambda_0 + i0) \right) \langle x \rangle^{-\frac{1}{2}-r} \| \le C_{r,r'} |z|^{r'}, \quad \text{Im} \, z > 0.$$
 (5.99a)

In particular this bound yield the expansion (5.91) with j = 0.

By a similar argument it follows that for $0 < r' < r \le 1$ and Im z > 0,

$$\|\langle x \rangle^{-\frac{3}{2}-r} \left(R'(\lambda_0 + z) - R'(\lambda_0 + i0) - zR'(\lambda_0 + i0)^2 \right) \langle x \rangle^{-\frac{3}{2}-r} \| \le C_{r,r'} |z|^{1+r'},$$
(5.99b)

implying the expansion (5.91) with j = 1.

We only prove (c) since the pattern of proof of (5.98a) and (5.98c) is the same. So suppose $\rho_1 > \frac{5}{2}$. Applying (5.94) (with N = 2) we first obtain that

$$W(z) = 1 + (G_0 + \sqrt{z}G_1 + zG_2)U(z) + \mathcal{O}(|z|^{1+\epsilon})$$

in $\mathcal{L}(1, -s'; 1, -s)$ with $s, s' \in (\frac{5}{2}, \rho_1)$. Note here that $U(z) \in \mathcal{L}(1, -s'; -1, s')$ with a uniform bound, cf. Theorems 3.13 and 3.18. Making (again) use of the boundedness of G_2 and the expansion (5.91) with j = 0 we obtain

$$G_2U(z) = G_2U^+ + \mathcal{O}(|z|^{\epsilon})$$
 in $\mathcal{L}(1, -s'; 1, -s)$.

It follows from (5.99a) that for $s \in (\frac{5}{2}, \rho_1)$ and $s' \in (2, \rho_1 - \frac{1}{2})$

$$\begin{aligned} \|\langle x_{a_0} \rangle^{-s} G_1(U(z) - U^+) \langle x_{a_0} \rangle^{s'} \| \\ &\leq C \|\langle x_{a_0} \rangle^{s-1} S^* I_0(R'(\lambda_0 + z) - R'(\lambda_0 + \mathrm{i0})) I_0 S \langle x_{a_0} \rangle^{s'} \| \\ &\leq C' \|\langle x \rangle^{-\rho_1 - \frac{1}{2} + s - 1} (R'(\lambda_0 + z) - R'(\lambda_0 + \mathrm{i0})) \langle x \rangle^{-\rho_1 - \frac{1}{2} + s'} \| \\ &\leq C'' |z|^{\frac{1}{2} + \delta}; \end{aligned}$$
(5.100)

this is for some $\delta > 0$.

It remains to show that

$$G_0U(z) = G_0U^+ + zG_0U_1^+ + \mathcal{O}(|z|^{1+\epsilon})$$
 in $\mathcal{L}(1, -s'; 1, -s)$.

This expansion for $s \in (\frac{5}{2}, \rho_1)$ and $s' \in (\frac{3}{2}, \rho_1 - 1)$ follows from (5.91) with j = 1 (justified by (5.99b)). We conclude (c).

Let $\mathcal{S}: \mathbb{C}^{\mu} \to \mathcal{H}^{1}_{-s}, s \in (\frac{1}{2}, \rho_{1})$, be defined by

$$\mathcal{S}c = \sum_{j=1}^{\mu} c_j \phi_j, \quad c = (c_1, \dots, c_{\mu}) \in \mathbb{C}^{\mu},$$

and $\mathcal{S}^* : \mathcal{H}^1_{-s} \to \mathbb{C}^\mu$ by

$$\mathcal{S}^*f = (\langle -U^+\phi_1, f \rangle \dots, \langle -U^+\phi_\mu, f \rangle), \quad f \in \mathcal{H}^1_{-s}$$

Let $Q: \mathcal{H}^1_{-s} \to \mathcal{H}^1_{-s}$ be given by

$$Qf = \sum_{j=1}^{\mu} \langle -U^+ \phi_j, f \rangle \phi_j.$$

One can then prove, cf. Section 5.1.2, that Q is a projection from \mathcal{H}_{-s}^1 onto \mathcal{K} , Q' = 1 - Q is a projection from \mathcal{H}_{-s}^1 onto $\operatorname{ran}(1 + G_0 U^+)$ and that

$$D_0^+ = (Q'(1+G_0U^+)Q')^{-1}Q'$$
(5.101)

exists and is continuous on \mathcal{H}_{-s}^1 , $s \in (\frac{1}{2}, \rho_1)$. By an argument of perturbation, it follows that for |z| small enough and $\operatorname{Im} z > 0$, Q'W(z)Q' is invertible on $\operatorname{ran}(1 + G_0U^+)$ with continuous inverse. Let

$$D(z) = (Q'W(z)Q')^{-1}Q'.$$

Then according to the varying conditions on ρ_1 in Lemma 5.19 one has the following expansions in $\mathcal{L}(1, -s; 1, -s)$:

$$D(z) = D_0 + \mathcal{O}(|z|^{\epsilon}), \text{ if } s \in (\frac{1}{2}, \rho_1).$$
 (5.102a)

$$D(z) = D_0 + \sqrt{z}D_1 + \mathcal{O}(|z|^{\frac{1}{2} + \epsilon_1}), \text{ if } \rho_1 > \frac{3}{2} \text{ and } s \in (\frac{3}{2}, \rho_1).$$
(5.102b)

Here

$$D_0 = D_0^+$$
 and $D_1 = -D_0 W_1 D_0$.

Using the operator D(z), we can establish the following representation formula for $W(z)^{-1}$ for Im z > 0:

$$W(z)^{-1} = \mathcal{E}(z) - \mathcal{E}_{+}(z)\mathcal{E}_{-+}(z)^{-1}\mathcal{E}_{-}(z), \qquad (5.103)$$

where

$$\mathcal{E}(z) = D(z), \quad \mathcal{E}_{+}(z) = \mathcal{S} - D(z)W(z)\mathcal{S},$$

$$\mathcal{E}_{-}(z) = \mathcal{S}^{*} - \mathcal{S}^{*}W(z)D(z), \quad \mathcal{E}_{-+}(z) = \mathcal{S}^{*}(-W(z) + W(z)D(z)W(z))\mathcal{S}.$$

The operators $\mathcal{E}_{\pm}(z)$ and $\mathcal{E}_{-+}(z)$ can be expanded similarly as D(z),

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \sqrt{z}\mathcal{E}_{\pm,1} + z\mathcal{E}_{\pm,2} + \dots$$
$$\mathcal{E}_{-+}(z) = \mathcal{E}_{-+,0} + \sqrt{z}\mathcal{E}_{-+,1} + z\mathcal{E}_{-+,2} + \dots$$

More precisely one has the following two lemmas on these expansions.

Lemma 5.20. Suppose (5.21) and (5.27a)–(5.27c). One has in $\mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}_{-s}^{1})$ (for the + case) or $\mathcal{L}(\mathcal{H}_{-s}^{1}, \mathbb{C}^{\mu})$ (for the - case):

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \mathcal{O}(|z|^{\epsilon}), \quad \text{if } s \in (\frac{1}{2}, \rho_1). \tag{5.104a}$$

$$\mathcal{E}_{\pm}(z) = \mathcal{E}_{\pm,0} + \sqrt{z}\mathcal{E}_{\pm,1} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}), \text{ if } \rho_1 > \frac{3}{2} \text{ and } s \in (\frac{3}{2}, \rho_1).$$
(5.104b)

Here

$$\mathcal{E}_{+,0} = \mathcal{S}, \quad \mathcal{E}_{-,0} = \mathcal{S}^*,$$

$$\mathcal{E}_{+,1} = -D_0 W_1 \mathcal{S}, \quad \mathcal{E}_{-,1} = -\mathcal{S}^* W_1 D_0.$$

The proof of Lemma 5.20 is the same as that for the corresponding assertions in Lemma 5.11 and will not be repeated here. For the leading order resolvent expansion only (5.104a) is needed.

Formally one can expand the matrix

$$\mathcal{E}_{-+}(z) = \mathcal{S}^*(-W(z) + W(z)D(z)W(z))\mathcal{S} \in \mathcal{M}_{\mu \times \mu}(\mathbb{C})$$

as

$$\mathcal{E}_{-+}(z) = \sqrt{z}\mathcal{E}_{-+,1}(z) + z\mathcal{E}_{-+,2} + \mathcal{O}(|z|^{1+\epsilon}), \qquad (5.105a)$$

where

$$\mathcal{E}_{-+,1} = -\mathcal{S}^* W_1 \mathcal{S},\tag{5.105b}$$

$$\mathcal{E}_{-+,2} = -\mathcal{S}^* (W_2 - W_1 D_0 W_1) \mathcal{S}.$$
 (5.105c)

The following lemma gives a precise meaning to this expansion. Note that (5.27c) is not imposed. Furthermore none of the assertions (5.98c) or (5.102b) is used, and (5.98b) is not used neither for treating the first case of the lemma (consequently the stated result applies for $\rho = 1$ and $\rho_1 = \frac{3}{2}$). Note that consequently the strong condition $\rho_1 > \frac{5}{2}$ of (5.98c) is not relevant for the lemma, although superficially this condition might appear necessary for justifying (5.105a)–(5.105c); we can do without (5.98c).

Lemma 5.21. Suppose (5.21), (5.27a) and (5.27b).

Case 1. If zero is an exceptional point of the first kind of P^+ , then

$$\mathcal{E}_{-+}(z) = \sqrt{z}\mathcal{E}_{-+,1} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon})$$
(5.106a)

for z near zero and Im z > 0, where the matrix $\mathcal{E}_{-+,1}$ is given by

$$\mathcal{E}_{-+,1} = iC_0^*C_0 \tag{5.106b}$$

with the j'th column of $C_0 \in \mathcal{M}_{m \times \mu}(\mathbb{C}), \ 1 \leq j \leq \mu$, specified as

$$c(\phi_j) = \frac{1}{2\sqrt{\pi}} (\langle 1, \sum_{k=1}^m U_{1k}^+ \phi_{j,k} \rangle_0, \cdots, \langle 1, \sum_{k=1}^m U_{mk}^+ \phi_{j,k} \rangle_0) \in \mathbb{C}^m.$$
(5.106c)

Here ϕ_j is denoted as $\phi_j = (\phi_{j,1}, \cdots, \phi_{j,m}) \in \mathcal{K}$.

Case 2. Assume zero is an exceptional point of the second kind of P^+ , $\rho > 1$ and $\rho_1 > \frac{3}{2}$. Then

$$\mathcal{E}_{-+}(z) = z\mathcal{E}_{-+,2} + \mathcal{O}(|z|^{1+\epsilon})$$
 (5.107a)

for z near zero and Im z > 0, where

$$\mathcal{E}_{-+,2} = (\langle (1 - U_1^+)\phi_i, \phi_j \rangle_0)_{1 \le i,j \le \mu}.$$
 (5.107b)

Case 3. Assume zero is an exceptional point of the third kind of P^+ , $\rho > 1$ and $\rho_1 > \frac{3}{2}$. Let $1 \le \kappa < \mu$ denote the multiplicity of zero resonance of P^+ . Then

$$\mathcal{E}_{-+}(z) = \sqrt{z}\mathcal{E}_{-+,1} + z\mathcal{E}_{-+,2} + \mathcal{O}(|z|^{1+\epsilon})$$
(5.108a)

for z near zero and Im z > 0, where $\mathcal{E}_{-+,1}$ and $\mathcal{E}_{-+,2}$ can be written as block matrices

$$\mathcal{E}_{-+,1} = \begin{pmatrix} iC_0^*C_0 & 0\\ 0 & 0 \end{pmatrix} \quad and \quad \mathcal{E}_{-+,2} = \begin{pmatrix} \mathcal{E}_{11}^{(2)} & \mathcal{E}_{12}^{(2)}\\ \mathcal{E}_{21}^{(2)} & \mathcal{E}_{22}^{(2)} \end{pmatrix}$$
(5.108b)

with the j'th column of $C_0 \in \mathcal{M}_{m \times \kappa}(\mathbb{C})$, $1 \leq j \leq \kappa$, equal to $c(\phi_j)$ and with $\mathcal{E}_{22}^{(2)} = (\langle (1 - U_1^+)\phi_i, \phi_j \rangle_0)_{\kappa+1 \leq i,j \leq \mu}$.

Proof. <u>Case 1</u>. We start by estimating the contribution from the second term of the representation

$$\mathcal{E}_{-+}(z) = (\langle U^+\phi_i, W(z)\phi_j - W(z)D(z)W(z)\phi_j \rangle)_{1 \le i,j \le \mu}.$$

Using the identity $\phi_j = -G_0 U^+ \phi_j$, $j \leq \mu$, one can estimate $||W(z)\phi_j||_{\mathcal{H}^1_{-s}}$ for any $s \in (1, \rho_1)$ as

$$||W(z)\phi_j||_{\mathcal{H}^1_{-s}} \le ||(r_0(z) - G_0)U(z)\phi_j||_{\mathcal{H}^1_{-s}} + ||G_0(U(z) - U^+)\phi_j||_{\mathcal{H}^1_{-s}}.$$

By the (uniform) bound $U(z)\phi_j \in \mathcal{H}^{-1}_{(3/2)^-}$ and a free resolvent bound, possibly deduced by interpolating (5.45) and (5.94) (with N = 1),

$$||(r_0(z) - G_0)U(z)\phi_j||_{\mathcal{H}^1_{-s}} \le C|z|^{\frac{1}{4}+\epsilon}.$$

From this point we fix any $s \in (1, \frac{5}{4})$. Then $1 + \rho - s > \frac{3}{4}$ and by (5.99a)

$$\|G_0(U(z) - U^+)\phi_j\|_{\mathcal{H}^1_{-s}} \le C_1 \|\langle x \rangle^{-1 - \rho + s} (R'(\lambda_0 + z) - R'(\lambda_0 + i0)) \langle x \rangle^{-1 - \rho + s} \|$$

$$\le C_2 |z|^{\frac{1}{4} + \epsilon}.$$

This proves that

$$||W(z)\phi_j||_{\mathcal{H}^1_{-s}} \le C|z|^{\frac{1}{4}+\epsilon}.$$
 (5.109a)

Similarly we can prove that

$$\|W(z)^* U^+ \phi_i\|_{\mathcal{H}_s^{-1}} \le C |z|^{\frac{1}{4} + \epsilon}.$$
(5.109b)

Due to (5.102a) the operator D(z) is uniformly bounded in $\mathcal{L}(\mathcal{H}_{-s}^1, \mathcal{H}_{-s}^1)$, and we conclude using (5.109a) and (5.109b) that

$$\mathcal{E}_{-+}(z) = (\langle U^+\phi_i, W(z)\phi_j \rangle)_{1 \le i,j \le \mu} + \mathcal{O}(|z|^{\frac{1}{2}+2\epsilon}).$$

Next we simplify the first term to the right. It follows by similar arguments that it is expanded as

$$\langle U^+ \phi_i, W(z)\phi_j \rangle_0 = (\langle U^+ \phi_i, \phi_j + r_0(z)U^+ \phi_j \rangle)_{1 \le i,j \le \mu} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}),$$

Since $U^+ = U^-$ on \mathcal{K} , it follows from Corollary 3.19 that $U^+\phi_i, U^+\phi_j \in \mathcal{H}^{-1}_{(3/2)^+}$. Then we obtain from (5.94) (with N = 1) that

$$\langle U^+\phi_i, W(z)\phi_j \rangle_0 = (\langle U^+\phi_i, \sqrt{z}G_1U^+\phi_j \rangle)_{1 \le i,j \le \mu} + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}),$$

Using the explicit formula for the integral kernel of G_1 , one obtains

$$\langle U^+\phi_i, G_1U^+\phi_j \rangle_0 = \mathbf{i}(c(\phi_i), c(\phi_j))$$

where $c(\phi_j) \in \mathbb{C}^m$ is given by (5.106c). This shows $\mathcal{E}_{-+,1} = iC_0^*C_0$, and we have proven the assertion for Case 1.

<u>Case 2</u>. If zero is an exceptional point of the second kind, then $\phi_j \in \mathcal{H}$ and Theorem 5.6 shows $W_1\phi_j = G_1U^+\phi_j = 0$ for $1 \leq j \leq \mu$. Therefore $\mathcal{E}_{-+,1} = -\mathcal{S}^*W_1\mathcal{S} = 0$.

By possibly making ρ_1 smaller we can assume that (5.27c) is fulfilled and therefore that (5.102a) is valid for some $s > \frac{3}{2}$. We invoke the implied uniform boundedness of $D(z) \in \mathcal{L}(1, -s; 1, -s)$ for such s, apply (5.98b) twice and conclude that

$$\mathcal{S}^*W(z)D(z)W(z)\mathcal{S} = \mathcal{O}(|z|^{1+\epsilon})$$

It remains to show that

$$(\langle U^+\phi_i, W(z)\phi_j \rangle)_{1 \le i,j \le \mu} = z(\langle (1-U_1^+)\phi_i, \phi_j \rangle_0)_{1 \le i,j \le \mu} + \mathcal{O}(|z|^{1+\epsilon}).$$
(5.110)

For that we first invoke (5.91) with j = 1 and write

$$(\langle U^+\phi_i, W(z)\phi_j \rangle)_{1 \le i,j \le \mu} = -\mathcal{S}^*(1+r_0(z)(U^++zU_1^+))\mathcal{S} + \mathcal{O}(|z|^{1+\epsilon}).$$

Next we note that $U^+\phi_j \in \mathcal{H}^{-1}_{(5/2)^+}$ (cf. Corollary 3.19), so that we can use the expansion (5.94) with N = 2. Finally we proceed as in the proof of (5.69) to rewrite the leading term obtained this way, concluding (5.110).

<u>Case 3</u>. If zero is an exceptional point of the third kind, we can combine methods used for Case 1 and Case 2 to show (5.108a) and (5.108b). The details are omitted. \Box

We remark that for Case 1, rank $C_0 = \kappa = \mu$ and $C_0^* C_0$ is positive definite. Moreover Lemma 5.18 ensures that the matrices $(\langle (1 - U_1^+)\phi_i, \phi_j \rangle_0)_{1 \le i,j \le \mu}$ for Case 2 and $(\langle (1 - U_1^+)\phi_i, \phi_j \rangle_0)_{\kappa+1 \le i,j \le \mu}$ for Case 3 of Lemma 5.21 are invertible. These properties are used in the following calculation of the leading term for the inverse of W(z). The proof of the proposition goes largely along the lines of that of Proposition 5.12.

If 0 is an eigenvalue of P^+ (relevant for Cases 2 and 3 below) we let Π_0 denote the spectral projection in \mathcal{H} onto the zero-eigenspace of this operator.

Proposition 5.22. Assume Conditions (5.21), (5.27a) and (5.27b). The following asymptotics hold in $\mathcal{L}(1, -s; 1, -s)$ for s > 1 sufficiently close to 1 and for |z| small with Im z > 0.

Case 1. Suppose that zero is an exceptional point of the first kind. Then

$$W(z)^{-1} = \frac{\mathrm{i}}{\sqrt{z}}Q_0 + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
 (5.111a)

Here

$$Q_0 = \sum_{j=1}^{\kappa} \langle -U^+ \psi_j, \cdot \rangle_0 \psi_j \tag{5.111b}$$

with vectors $\psi_j \in \mathcal{K}$ obeying

$$(c(\psi_j), c(\psi_k)) = \delta_{jk} \quad \text{for } j, k = 1, \cdots, \kappa.$$
(5.111c)

Here $c(\psi_j) \in \mathbb{C}^m$ is defined as $c(\phi_j)$ in Lemma 5.21 with ϕ_j replaced by ψ_j and (\cdot, \cdot) denotes the scalar product of \mathbb{C}^m .

Cases 2 and 3. Suppose that zero is an exceptional point of the second or the third kind, respectively, $\rho > 1$ and $\rho_1 > \frac{3}{2}$. Then

$$W(z)^{-1} = z^{-1} (\Pi_0 (1 - U_1^+) \Pi_0)^{-1} \Pi_0 U^+ + \mathcal{O}(|z|^{-1+\epsilon}).$$
(5.112)

Proof. The results follow from Lemmas 5.20 and 5.21 and the formula (5.103). We only study Case 1. For $s \in (1, \rho_1)$, one has for some $\epsilon > 0$

$$\begin{aligned} \mathcal{E}(z) &= D_0 + \mathcal{O}(|z|^{\epsilon}) \text{ in } \mathcal{L}(1, -s; 1, -s), \\ \mathcal{E}_{\pm}(z) &= \mathcal{E}_{\pm,0} + \mathcal{O}(|z|^{\epsilon}) \text{ in } \mathcal{L}(\mathbb{C}^{\mu}, \mathcal{H}^1_{-s}) \text{ or in } \mathcal{L}(\mathcal{H}^1_{-s}, \mathbb{C}^{\mu}) \\ \mathcal{E}_{-+}(z) &= \mathrm{i}\sqrt{z}B_0 + \mathcal{O}(|z|^{\frac{1}{2}+\epsilon}), \end{aligned}$$

where $B_0 = C_0^* C_0$ is positive definite. Therefore

$$W(z)^{-1} = \frac{\mathrm{i}}{\sqrt{z}} \mathcal{S}B_0^{-1} \mathcal{S}^* + \mathcal{O}(|z|^{-\frac{1}{2}+\epsilon}).$$
(5.113)

Decompose B_0^{-1} as $B_0^{-1} = M_0^2$ where $M_0 \in \mathcal{M}_{\kappa \times \kappa}(\mathbb{C})$ is a positive definite Hermitian matrix. Set $M_0 = (m_{ij})_{1 \le i,j \le \kappa}$ and

$$\psi_i = \sum_{k=1}^{\mu} m_{ki} \phi_k, \quad i = 1, \cdots, \kappa.$$
(5.114)

Then $\{\psi_1, \dots, \psi_\kappa\}$ is a basis of resonance functions of P^+ . Similarly to the proof of Proposition 5.8 (a) we can check that $\{\psi_1, \dots, \psi_\kappa\}$ verifies the normalization condition

$$(c(\psi_i), c(\psi_j)) = \delta_{ij}; \quad i, j \le \kappa$$

Hence the expansion (5.111a), with the specification (5.111b)-(5.111c), is proved. $\hfill \Box$

Since $E_{\mathcal{H}}(\lambda_0 + z)^{-1} = -W(z)^{-1}(P_0 - z)^{-1}$ and $G_0U = -1$ on \mathcal{K} , the following result follows immediately from Proposition 5.22 (cf. (5.45)).

Proposition 5.23. Assume (5.21), (5.27a) and (5.27b). The following asymptotics hold in $\mathcal{L}(-1, s; 1, -s)$ for s > 1 and for |z| small with Im z > 0.

Case 1. Suppose zero is an exceptional point of the first kind of P^+ . Let κ be the multiplicity of zero resonance. Then for some $\epsilon > 0$

$$E_{\mathcal{H}}(\lambda_0 + z)^{-1} = -\frac{\mathrm{i}}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle \psi_j, \cdot \rangle \psi_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}), \qquad (5.115a)$$

132

where ψ_i , $j = 1, \dots, \kappa$, are resonance states verifying (5.111c).

Cases 2 and 3. Suppose zero is an exceptional point of the second or the third kind, respectively, $\rho > 1$ and $\rho_1 > \frac{3}{2}$. Then for some $\epsilon > 0$

$$E_{\mathcal{H}}(\lambda_0 + z)^{-1} = z^{-1}(\Pi_0(1 - U_1^+)\Pi_0)^{-1}\Pi_0 + \mathcal{O}(|z|^{-1+\epsilon}).$$
(5.115b)

A main result of this section is the following. In agreement with previous usage, if λ_0 is a resonance of H its multiplicity is denoted by κ , and if λ_0 is an eigenvalue of H the spectral projection of H onto the corresponding eigenspace is denoted by Π_H .

Theorem 5.24. Assume (5.21), (5.27a) and (5.27b). The following asymptotics hold for $R(\lambda_0 + z)$ as a bounded operator from H_s^{-1} to H_{-s}^1 for any s > 1, as $z \to 0$ and $\pm \text{Im } z > 0$, and $\epsilon = \epsilon(s) > 0$.

Case 0. Suppose λ_0 is a regular point of H. Then one has

$$R(\lambda_0 + z) = R'(\lambda_0 \pm i0) + (S - R'(\lambda_0 \pm i0)I_0S)D_0G_0(S^* - S^*I_0R'(\lambda_0 \pm i0)) + \mathcal{O}(|z|^{\epsilon}).$$
(5.116)

Here the boundary value of resolvent $R'(\lambda_0 + i0)$ (resp., $R'(\lambda_0 - i0)$) is used when $\operatorname{Im} z > 0$ (resp. $-\operatorname{Im} z > 0$).

Case 1. Suppose λ_0 is an exceptional point of the first kind of H. Then

$$R(\lambda_0 + z) = \frac{\mathrm{i}}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon})$$
(5.117a)

where u_i 's are resonance states of H given by

$$u_j = (1 - R'(\lambda_0 \pm i0)I_0)S\psi_j, \quad 1 \le j \le \kappa,$$
 (5.117b)

where ψ_j 's are the resonance states of P^{\pm} given in Proposition 5.22.

Cases 2 and 3. Suppose λ_0 is an exceptional point of the second or the third kind of H, respectively, $\rho > 1$ and $\rho_1 > \frac{3}{2}$. Then

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \mathcal{O}(|z|^{-1+\epsilon}).$$
(5.118)

Mimicking the proof of Theorem 5.14 we see that Theorem 5.24 for Im z > 0 is a consequence of Proposition 5.23 and the formula (5.24). The case Im z < 0 can be proved in the same way with P^+ replaced by P^- . Note that due to Theorem 5.6 (a), the resonance states u_j given by (5.117b) are independent of the choice of the sign \pm . **Remark 5.25.** For Case 1 of Theorem 5.24 the leading term of the resolvent $R(\lambda_0 + z)$ is expressed in terms of a specific basis of resonance states $\{u_1, \ldots, u_\kappa\}$ of H such that $\psi_j = S^* u_j, j = 1, \ldots, \kappa$, are given as in Proposition 5.22. The normalization condition (5.111c) expressed in terms of the u_j 's reads

$$(c(S^*u_j), c(S^*u_k)) = \delta_{jk}, \quad j, k = 1, \dots, \kappa.$$
 (5.119)

By Proposition 5.8 (b), (5.117a) remains valid for any basis of resonance states $\{u_1, \ldots, u_{\kappa}\}$ of H verifying (5.119). Note also the formula

$$c(S^*u_j) = \frac{1}{2\sqrt{\pi}} \Big(\int_{\mathbf{X}} \varphi_1 I_0 u_j \mathrm{d}x, \dots, \int_{\mathbf{X}} \varphi_m I_0 u_j \mathrm{d}x \Big), \quad j = 1, \dots, \kappa.$$
(5.120)

For Cases 0 and 1, $\rho = 1$ is legitimate. However Cases 2 and 3 of Theorem 5.24 have the stronger condition $\rho > 1$ and therefore exclude the physics models which would imply a limitation on the application in Section 6.3, see Remark 6.29 1). However with an a priori weak decay property of the corresponding L^2 -eigenfunctions if λ_0 is an eigenvalue of H (as for Cases 2 and 3) we can almost verbatim mimic the proof of Theorem 5.16 and obtain a resolvent expansion also for $\rho = 1$. Note however that the property $\lambda_0 \notin \sigma_{\rm pp}(H'_{\sigma})$ for small σ is based on a perturbation of Proposition 3.3, cf. [AHS], and note also the relevance of Remark 3.21 2). This expansion is up to second order (at least for Case 3).

Theorem 5.26. Assume (5.21) and (5.27a), as well as (5.27b) with $\rho = 1$ and $\rho_1 = 3/2$. Suppose λ_0 is an eigenvalue of H and that $\operatorname{ran} \Pi_H \subset L_t^2$ for some t > 3/2. Then the following asymptotics hold for $R(\lambda_0 + z)$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ and $\pm \operatorname{Im} z > 0$, and $\epsilon = \epsilon(s) > 0$.

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \frac{i}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}).$$
(5.121)

Here the second term on the right-hand side appears only if $\kappa \geq 1$, that is if λ_0 is a resonance of H, and in that case $\{u_1, \ldots, u_\kappa\}$ is a basis of resonance states of H being independent of the choice of the sign \pm . If on the other hand λ_0 is not a resonance of H, then $R(\lambda_0 + z) + z^{-1}\Pi_H$ has limits in norm as $z \to 0, \pm \text{Im } z > 0$.

Remark 5.27. As in Remark 5.17 2) the condition ran $\Pi_H \subset L_t^2$ for some t > 1 suffices for the last assertion of Theorem 5.26.

5.2 Positive slowly decaying effective potentials

Low-energy scattering for one-body Schrödinger operators with positive slowly decaying potentials is studied in [Na, Ya1, Ya2]. It is shown in [Wa6] that this kind of operators satisfies Gevrey type resolvent estimates at the threshold and this can be used to establish large time asymptotics of the quantum dynamics with subexponential time decay estimates on the remainder [AW, Wa6]. In this section, we want to show similar results for the N-body Schrödinger operator at the lowest threshold $\lambda_0 = \Sigma_2$, assuming that the effective potential is positive outside a compact set and slowly decaying at infinity.

5.2. Positive slowly decaying effective potentials

Recall first from [Wa6] some results for one-body operators. The model operator H_0 in this framework is a closed second order elliptic operator of the form

$$H_0 = -\sum_{i,j=1}^n \partial_{x_i} a^{ij}(x) \partial_{x_j} + \sum_{j=1}^n b_j(x) \partial_{x_j} + v(x), \qquad (5.122)$$

where $a^{ij}(x)$, $b_j(x)$ and v(x) are complex-valued functions in \mathbb{R}^n , $n \ge 1$. We assume that a^{ij} , b_j are bounded C^1 functions with bounded derivatives and there exists c > 0 such that

$$\operatorname{Re}(a^{ij}(x)) \ge cI_n, \quad \forall x \in \mathbb{R}^n.$$
 (5.123a)

Assume also that v is relatively bounded with respect to $-\Delta$ with relative bound zero and there exist some constants $0 < \mu < 1$ and $c_0 > 0$ such that

$$|\langle H_0 u, u \rangle| \ge c_0(||\nabla u||^2 + ||\langle x \rangle^{-\mu} u||^2), \text{ for all } u \in H^2,$$
 (5.123b)

$$\sup_{x} |\langle x \rangle^{\mu} b_j(x)| < \infty, \quad j = 1, \cdots, n.$$
(5.123c)

The bound (5.123b) is called a weighted coercive condition and is essential for Gevrey estimates of the resolvent at the threshold zero.

Under the assumptions (5.123b) and (5.123c), H_0 is bijective from $D(H_0)$ to $R(H_0)$. Let $R_0(0) : R(H_0) \to D(H_0)$ be its algebraic inverse. $R_0(0)$ is a densely defined and closed operator, continuous from L_s^2 to $L_{s-2\mu}^2$ and compact from L_s^2 to $L_{s-2\mu-\epsilon}^2, \epsilon > 0$, for any $s \in \mathbb{R}$ ([Wa6, Ya2]). Thus $R_0(0)^N : L_s^2 \to L_{s-2\mu N}^2$ is bounded. If Re $H \ge 0$, one can check that the strong limit

$$\operatorname{s-lim}_{z\in\Omega(\delta),z\to0}\langle x\rangle^{-2N\mu}(R_0(z)^N-R_0(0)^N)=0,$$

where $\Omega(\delta) = \{z : |\arg z| > \frac{\pi}{2} + \delta\}$ with $\delta > 0$ small. The following Gevrey-type estimates hold for the resolvent at the threshold zero.

Theorem 5.28 ([Wa6, Theorem 2.1]). Assume conditions (5.123a), (5.123b) and (5.123c). For any a > 0 there exists $C_a > 0$ such that

$$\|\mathrm{e}^{-a\langle x\rangle^{1-\mu}}R_0(0)^N\| + \|R_0(0)^N\mathrm{e}^{-a\langle x\rangle^{1-\mu}}\| \le C_a^{N+1}N^{\gamma N},$$

uniformly in $N \ge 1$. Here $\gamma = \frac{2\mu}{1-\mu}$.

From Theorem 5.28 one deduces the following result.

Corollary 5.29 ([Wa6, Corollary 4.2]). Let $H_0 = -\Delta + v(x)$ be self-adjoint and positive and satisfy the assumptions of Theorem 5.28. Then for any a > 0 there exists $C_a > 0$ such that

$$\|e^{-a\langle x\rangle^{1-\mu}}R_0(z)^N\| + \|R_0(z)^N e^{-a\langle x\rangle^{1-\mu}}\| \le C_a^{N+1}N^{\gamma N}, \quad \forall N \ge 1,$$

uniformly in $N \ge 1$ and $z \in \Omega = \{\zeta \in \mathbb{C} : |\arg \zeta| > \delta\}$ with $\delta > 0$.

Since $R_0^{(N)}(z) = N! R_0(z)^{N+1}$, Corollary 5.29 means that $e^{-a\langle x \rangle^{1-\mu}} R_0(z)$ belongs to the Gevrey class $\mathcal{G}^{(1+\gamma)}(\Omega)$, where

$$\mathcal{G}^{(1+\gamma)}(\Omega) = \{F: \Omega \to \mathcal{L}(L^2) \mid \exists C > 0 \text{ s.t. } \|F^{(N)}(z)\| \le C^{N+1}N! N^{\gamma N}, \ \forall z \in \Omega, N \in \mathbb{N}\}$$

For n = 3, the repulsive Coulomb Hamiltonian $-\Delta + \frac{c}{|x|}$, c > 0, satisfies all conditions of Corollary 5.29 with $\mu = \frac{1}{2}$, so its resolvent belongs to $\mathcal{G}^{(3)}(\Omega)$ in exponentially weighted spaces. In [AW, Wa6], non-selfadjoint perturbations H of H_0 are studied and large time expansions are obtained for the quantum dynamics e^{-itH} and e^{-tH} as $t \to +\infty$.

To study the N-body Schrödinger operator H at its lowest threshold $\lambda_0 = \Sigma_2$, we assume

 $\lambda_0 = \Sigma_2$ is a non-multiple two-cluster threshold in the sense of Condition 2.4.

(5.124) This means there exists a unique $a_0 \in \mathcal{A} \setminus \{a_{\max}\}$ such that $\Sigma_2 \in \sigma_{pp}(H^{a_0})$, and this cluster decomposition is a two-cluster decomposition obeying Condition 1.6. We keep the notation of Section 2.2 (in the present case, m = 1, φ_1 is a normalized eigenfunction of H^{a_0} with eigenvalue Σ_2 and $\mathcal{H} = L^2(\mathbf{X}_{a_0})$). Denote $x_0 = x_{a_0} \in \mathbf{X}_{a_0}$. From Section 2.2 we have the representation formula for the resolvent,

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z)$$
 for $\text{Im } z \neq 0$,

where

$$E(z) = R'(z) = (H' - z)^{-1} \Pi',$$

$$E_{+}(z) = (1 - R'(z)I_{0})S,$$

$$E_{-}(z) = S^{*} (1 - I_{0}R'(z)),$$

$$E_{\mathcal{H}}(z) = (z - \lambda_{0}) - (-\Delta_{0} + S^{*}I_{0}S - S^{*}I_{0}R'(z)I_{0}S),$$

and $\Delta_0 = \Delta_{x_0}$, $I_0(x) = \sum_{b \not\subset a} V_b(x^b)$ and $Sf(x)(x) = \varphi_1(x^{a_0})f(x_0)$. The effective potential here is

$$S^*I_0S(x_0) = \langle I_0\varphi_1, \varphi_1 \rangle_0,$$

where $\langle \cdot, \cdot \rangle_0$ is the scalar product of $L^2(\mathbf{X}^{a_0})$.

In addition to (5.124) we assume that

$$\lambda_0 \notin \sigma_{\rm pp}(H'), \quad \rho \in (0,2), \tag{5.125a}$$

$$\exists c, R > 0 \text{ such that } S^* I_0 S \ge \frac{c}{|x_0|^{\rho}} \text{ for } |x_0| > R.$$
 (5.125b)

Conditions (5.124) and (5.125a) show that R'(z) is holomorphic for z near λ_0 . Condition (5.125b) implies that in some sense λ_0 can not be a resonance of H (cf. Lemma 5.32). Therefore we need only to distinguish the cases λ_0 be an eigenvalue of H or not. In the case λ_0 is not an eigenvalue of H we prove the following result.

Theorem 5.30. In addition to conditions (5.124), (5.125a) and (5.125b), we assume $\lambda_0 = \Sigma_2 \notin \sigma_{\rm pp}(H)$. Then for any a > 0, $e^{-a\langle x_0 \rangle^{1-\mu}} R(z)$ belongs to the Gevrey class $\mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_{\lambda_0}(\delta))$, where $\mu = \frac{\rho}{2}$ and $\Omega_{\lambda_0}(\delta) = \{z \in \mathbb{C} : |z - \lambda_0| < \delta, |\arg(z - \lambda_0)| > \delta\}$, $\delta > 0$ small.

The proof of Theorem 5.30 is divided into several steps. The main task is to prove that $E_{\mathcal{H}}(z)^{-1} \in \mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_{\lambda_0}(\delta))$ if λ_0 is not an eigenvalue of H. Set

$$E_{\mathcal{H}}(\lambda_0 + \lambda) = \lambda - (-\Delta_0 + S^* I_0 S - S^* I_0 R'(\lambda_0 + \lambda) I_0 S) := \lambda - (-\Delta_0 + U(\lambda))$$

and

$$W(\lambda) = -S^* I_0 R'(\lambda_0 + \lambda) I_0 S = -\langle \varphi_1, I_0 R'(\lambda + \lambda_0) I_0(\varphi_1 \otimes \cdot) \rangle_0.$$
 (5.126)

 $W(\lambda)$ is holomorphic for λ near 0 and satisfies

$$\|\langle x \rangle^{1+2\mu} W(\lambda) \langle x \rangle^{1+2\mu}\| \le C$$

uniformly in $|\lambda| < 2\delta$ for some $\delta > 0$ small, because $\Pi I_0 \Pi' = \mathcal{O}(\langle x_0 \rangle^{-1-2\mu})$. Under the assumption (5.125b), the results of [Wa6] can be applied to $-\Delta_0 + S^* I_0 S$. However the non-local term $W(\lambda)$ can not be treated as a perturbation in the Gevrey setting. To prove Theorem 5.30 we follow the approach of [Wa6] from the very beginning and exploit the holomorphicity of $W(\lambda)$ in λ to prove some uniform energy estimates.

For $s \in \mathbb{R}$, let φ_s be the weight function defined by

$$\varphi_s(x_0) = \left(1 + \frac{|x_0|^2}{R_s^2}\right)^{s/2}$$
 with $R_s = M \langle s \rangle^{\frac{1}{1-\mu}}$,

where M > 1 is to be chosen sufficiently large and is independent of s.

Lemma 5.31. There exist constants $M, \delta, C > 0$ such that

$$\|\langle x_0 \rangle^r \varphi_s W(\lambda) \varphi_{-s} \langle x_0 \rangle^{r'} \| \le C$$

for $r, r' \in \mathbb{R}$ with $r, r' \leq 1 + 2\mu$, for $\lambda \in \mathbb{C}$ with $|\lambda| < \delta$ and for $s \in \mathbb{R}$.

Proof. Since $[\varphi_s, \Pi] = 0$ and $\Pi I_0 \Pi' = \mathcal{O}(\langle x_0 \rangle^{-1-2\mu})$, one has

$$\|\langle x_0 \rangle^r \varphi_s W(\lambda) \varphi_{-s} \langle x_0 \rangle^{r'} \| \le C \|\varphi_s R'(\lambda_0 + \lambda) \varphi_{-s} \|.$$

Computing the commutator $[-\Delta_0, \varphi_s]$ one obtains

$$\varphi_s H' \varphi_{-s} = H' + \mathcal{O}\left(\frac{s}{M\langle s \rangle^{\frac{1}{1-\mu}}}\right) = H' + \mathcal{O}\left(\frac{1}{M}\right),$$

where the term $\mathcal{O}(\frac{1}{M})$ satisfies the bound

$$\|\mathcal{O}\left(\frac{1}{M}\right)(H'+i)^{-1}\| \le \frac{C}{M}$$

uniformly in $s \in \mathbb{R}$. Since λ_0 is in the resolvent set of H', there exists $\delta > 0$ such that $R'(\lambda_0 + \lambda)$ is a well defined holomorphic function for $|\lambda| < 2\delta$ and we can take M > 1 large so that

$$\|\mathcal{O}\left(\frac{1}{M}\right)R'(\lambda_0+\lambda)\| \le \frac{1}{2}$$

for $|\lambda| < \delta$. From the equation

$$\varphi_s R'(\lambda_0 + \lambda)\varphi_{-s} = R'(\lambda_0 + \lambda) \left(1 + \mathcal{O}\left(\frac{1}{M}\right)R'(\lambda_0 + \lambda)\right)^{-1}\Pi',$$

it follows that $\varphi_s R'(\lambda_0 + \lambda)\varphi_{-s}$ is uniformly bounded for $s \in \mathbb{R}$ and $|\lambda| < \delta$. Whence Lemma 5.31 is proven.

Let $\chi_R(x_0) = \chi_1(\frac{x_0}{R}), R \ge 1$, where $\chi_1 \in C^{\infty}$ is a cut-off such that $0 \le \chi_1 \le 1$, $\chi_1(x_0) = 0$ if $|x_0| \le 1$ and $\chi_1(x_0) = 1$ if $|x_0| \ge 2$. Set

$$F(\lambda) = -\Delta_0 + 1 - \chi_R + \chi_R U(\lambda)\chi_R$$
$$\widetilde{U}(\lambda) = U(\lambda) - (1 - \chi_R + \chi_R U(\lambda)\chi_R)$$
$$h_0 = -\Delta_0 + 1 - \chi_R + \chi_R^2 S^* I_0 S.$$

Then $E_{\mathcal{H}}(\lambda_0 + \lambda)$ can be decomposed as

$$E_{\mathcal{H}}(\lambda_0 + \lambda) = \lambda - (F(\lambda) + \tilde{U}(\lambda))$$

$$= \lambda - (h_0 + (1 - \chi_R)((1 + \chi_R)S^*I_0S - 1) + W(\lambda))$$
(5.127)

For R > 1 sufficiently large, h_0 is a one-body Schrödinger operator with globally positive and slowly decaying potential $v_0 = 1 - \chi_R + \chi_R^2 S^* I_0 S$:

$$v_0(x_0) \ge \frac{c}{\langle x_0 \rangle^{2\mu}}, \quad x_0 \in \mathbf{X}_0,$$

for some c > 0. The operator $F(0) = h_0 + \chi_R W(0) \chi_R$ is a non-local perturbation of h_0 and $F(0) \ge 0$. Note that

$$\widetilde{U}(\lambda) = (1 - \chi_R)(U(\lambda) - 1) + \chi_R U(\lambda)(1 - \chi_R).$$

Since $1 - \chi_R$ has compact support and λ_0 is in the resolvent set of H', making use of the relation

$$e^{a\langle x_0\rangle}H'e^{-a\langle x_0\rangle} = H' + \mathcal{O}(a)$$

for a > 0 small, one sees that $\widetilde{U}(\lambda)$ is exponentially decaying in the sense that

$$\|\mathrm{e}^{a\langle x_0\rangle}\widetilde{U}(\lambda)\mathrm{e}^{a\langle x_0\rangle}\| \le C \tag{5.128}$$

uniformly for λ near 0 and $|a| \leq \delta$, $\delta > 0$ small.

Lemma 5.32. Assume (5.125b) and let $\mu = \frac{\rho}{2} \in (0, 1)$. One has

$$\|\langle x_0 \rangle^{-\mu} \varphi_s u\| + \|\nabla(\varphi_s u)\| \le C \|\langle x_0 \rangle^{\mu} \varphi_s(F(\lambda) - \lambda)u\|$$
(5.129)

uniformly in $s \in \mathbb{R}$, $u \in S$ and $\lambda \in \Omega_0(\delta) = \{\zeta \in \mathbb{C} : |\zeta| < \delta, |\arg \zeta| > \delta\} \cup \{0\}$.

Proof. By construction, $h_0 = -\Delta_0 + v_0(x_0)$ satisfies the weighted coercive condition (5.123b). Moreover, for $\lambda = \tau e^{i\phi} \in \mathbb{C}$ with $\tau > 0$ and $\phi \neq 0$, $|\phi| \geq \delta > 0$, $e^{-i\phi}(h_0 - \lambda)$ also satisfied (5.123b) with a lower bounded independent of λ so long as $|\phi| \geq \delta$ and $|\tau| < \delta$ for $\delta > 0$ small. Applying [Wa6, Lemma 3.1] to h_0 and $e^{-i\phi}(h_0 - \lambda)$, we deduce

$$\|\langle x_0 \rangle^{-\mu} \varphi_s u\| + \|\nabla(\varphi_s u)\| \le C \|\langle x_0 \rangle^{\mu} \varphi_s (h_0 - \lambda) u\|$$
(5.130)

uniformly in $s \in \mathbb{R}$ and $\lambda \in \Omega_0(\delta)$.

Since $F(\lambda) = h_0 + \chi_R W(\lambda) \chi_R$, Lemma 5.31 shows that

$$\|\langle x_0 \rangle^{\mu} \varphi_s(h_0 - \lambda) u\| \le \|\langle x_0 \rangle^{\mu} \varphi_s(F(\lambda) - \lambda) u\| + CR^{-2-2\mu} \|\langle x_0 \rangle^{-\mu} \varphi_s u\|$$

uniformly in $s \in \mathbb{R}$ and $|\lambda|$ small. Now Lemma 5.32 follows if R > 1 is taken appropriately large.

5.2. Positive slowly decaying effective potentials

Lemma 5.32 shows that $G(\lambda) = (F(\lambda) - \lambda)^{-1}$ satisfies the estimate

$$\|\langle x_0 \rangle^{-\mu} \varphi_s G(\lambda) \varphi_{-s} \langle x_0 \rangle^{-\mu} \| + \| \nabla \varphi_s G(\lambda) \varphi_{-s} \langle x_0 \rangle^{-\mu} \| \le C$$

uniformly in $s \in \mathbb{R}$ and $\lambda \in \Omega_0(\delta)$. It follows that

$$\|\langle x_0 \rangle^{-2\mu} \varphi_s G(\lambda) \varphi_{-s}\| \le C' \langle s \rangle^{\gamma}, \quad \gamma = \frac{2\mu}{1-\mu}, \tag{5.131}$$

uniformly in s and λ . The following technical estimates are the main step in the proof of Gevrey estimates of $G(\lambda)$, cf. [Wa6, Theorem 3.4].

Proposition 5.33. There exist constants $C, \delta > 0$ such that for any $r \in \mathbb{R}$, $N \in \mathbb{N}$ and $\lambda \in \Omega_0(\delta)$

$$\|\langle x_0 \rangle^{-2\mu} \langle x_{N,r} \rangle^{-(2N+r)\mu} G^{(N)}(\lambda) \langle x_{N,r} \rangle^{r\mu} \| \le C^{N+1} N! \langle (2N+r)\mu \rangle^{\gamma(N+1)}.$$
 (5.132)

Here

$$x_{N,r} = \frac{x_0}{R_{N,r}} \text{ with } R_{N,r} = M \langle (2N+r)\mu \rangle^{\frac{1}{1-\mu}}$$

and $\langle x_{N,r} \rangle = (1 + |x_{N,r}|^2)^{\frac{1}{2}}.$

Proof. The case N = 0 and $r \ge 0$ follows from (5.131) with $s = r\mu$. For the general case $N \ge 1$ and $r \ge 0$, we write

$$G^{(N)}(\lambda) = (G(\lambda)^2 - G(\lambda)\chi_R W'(\lambda)\chi_R G(\lambda))^{(N-1)},$$

and prove by an induction on N that

$$\begin{aligned} \|\langle x \rangle^{-2\mu} \langle x_{N,r} \rangle^{-(2N+r)\mu} G^{(N)}(\lambda) \langle x_{N,r} \rangle^{r\mu} \| \\ &\leq C_N^{N+1} N! \langle (2N+r)\mu \rangle^{\gamma(N+1)} \end{aligned}$$

with $C_{N+1} \leq C_N(1 + \frac{c}{N^{1+\gamma}})$ for some c > 0 independent of N. The details are the same as the proof of [Wa6, Theorem 3.4] and are omitted here.

To convert polynomial weight depending on N into exponential weight independent of N, we use the following estimate.

$$\forall a > 0 \,\exists A_a > 0 : \quad \|\langle x_{N,r} \rangle^{(2N+r)\mu} \mathrm{e}^{-a\langle x_0 \rangle^{1-\mu}} \|_{L^{\infty}} \le A_a^{\max\{2N+r,1\}}, \tag{5.133}$$

uniformly in $N \in \mathbb{N}$ and $r \in \mathbb{R}$. In fact, if $2N + r \leq C$ for some constant C > 0, the left-hand side of (5.133) is uniformly bounded by some constant C_1 . For 2N + r > C with C > 1 large but fixed, consider the function

$$f(\tau) = \left\langle \frac{\tau}{R_{N,r}} \right\rangle^{(2N+r)\mu} e^{-a\tau^{1-\mu}}, \quad \tau = |x_0|.$$

One has

$$f'(\tau) = \frac{f(\tau)}{\tau^{\mu}(R_{N,r}^2 + \tau^2)} \left((2N+r)\mu\tau^{1+\mu} - (1-\mu)a(R_{N,r}^2 + \tau^2) \right).$$

Since $\mu \in]0,1[$, for each a > 0, one can find some constant A > 0 such that

$$f'(\tau) < 0,$$
 for $\tau > AR_{N,r}$.

Therefore for 2N + r > C,

$$\|\langle x_{N,r}\rangle^{(2N+r)\mu}e^{-a\langle x_0\rangle^{1-\mu}}\|_{L^{\infty}} \leq \sup_{0 \leq \tau \leq AR_{N,r}} f(\tau) \leq \langle A \rangle^{(2N+r)\mu}$$

This proves (5.133) for some appropriate constant $A_a > 0$.

Proposition 5.33 implies the following Gevrey estimates for $G(\lambda) = (F(\lambda) - \lambda)^{-1}$:

$$\|\langle x_0 \rangle^{-\tau} \mathrm{e}^{-a \langle x_0 \rangle^{1-\mu}} G^{(N)}(\lambda) \langle x_0 \rangle^{\tau} \| \le C_a^{N+1+\tau} N! \langle N+\tau \rangle^{\gamma(N+1)+\frac{\tau}{1-\mu}}$$
(5.134)

uniformly in $\tau \geq 0, N \in \mathbb{N}$ and $\lambda \in \Omega_0(\delta)$.

Lemma 5.34. Assume (5.124), (5.125a) and (5.125b). Let u be a solution to the equation $E_{\mathcal{H}}(\lambda_0)u = 0$ and $u \in \mathcal{H}_{\tau}$ for some $\tau \in \mathbb{R}$. Then there exists some positive constant b_0 such that $e^{b_0 \langle x_0 \rangle^{1-\mu}}u \in \mathcal{H}$. Here $\mathcal{H} = L^2(\mathbf{X}_0)$ and $\mathcal{H}_{\tau} = L^2(\mathbf{X}_0, \langle x_0 \rangle^{2\tau} dx_0)$.

Proof. u satisfies the equation $u = G(0)(1-\chi_R)(1-U(0))u$. Since U(0) is continuous in \mathcal{H}_s for any $s \in \mathbb{R}$, $u \in \mathcal{H}_\tau$ and $1-\chi_R$ is of compact support, $(1-\chi_R)(1-U(0))u \in \mathcal{H}_s$ for any s > 0. Proposition 5.33 with N = 0 and r < 0 shows that $u \in \mathcal{H}_\infty$. See also Subsection 4.1.3. To show the sub-exponential decay of u, we write

$$E_{\mathcal{H}}(\lambda_0) = -(h_0 + (1 - \chi_R)(S^*I_0S - 1) + W(\lambda_0)).$$

Since $h_0 \ge -\Delta_0 + \frac{c}{\langle x_0 \rangle^{2\mu}}$ for some c > 0, the following Agmon energy estimate holds true: $\exists b, C > 0$ such that

$$\|\langle x_0 \rangle^{-\mu} \mathrm{e}^{b \langle x_0 \rangle^{1-\mu}} f \|^2 + \|\nabla_{x_0} (\mathrm{e}^{b \langle x_0 \rangle^{1-\mu}} f)\|^2 \le C(|\langle \mathrm{e}^{2b \langle x_0 \rangle^{1-\mu}} h_0 f, f \rangle| + \|f\|^2) \quad (5.135)$$

for $f \in D(h_0)$ with $e^{2b\langle x_0 \rangle^{1-\mu}} h_0 f \in \mathcal{H}$ (cf. [Wa6, (5.37)]). R'(z) being holomorphic for z near λ_0 , one has for b > 0 small enough

$$\begin{aligned} |\langle e^{2b\langle x_0 \rangle^{1-\mu}} W(\lambda_0) f, f \rangle| &\leq C_1(||\langle x_0 \rangle^{-1-2\mu} e^{b\langle x_0 \rangle^{1-\mu}} f||^2 + ||f||^2) \\ &\leq \epsilon ||\langle x_0 \rangle^{-\mu} e^{b\langle x_0 \rangle^{1-\mu}} f||^2 + C_\epsilon ||f||^2 \end{aligned}$$

for any $\epsilon > 0$. We deduce from (5.135) that

$$\|\langle x_0 \rangle^{-\mu} e^{b \langle x_0 \rangle^{1-\mu}} f \|^2 + \|\nabla_{x_0} (e^{b \langle x_0 \rangle^{1-\mu}} f)\|^2 \le C(|\langle e^{2b \langle x_0 \rangle^{1-\mu}} E_{\mathcal{H}}(\lambda_0) f, f \rangle| + \|f\|^2)$$
(5.136)

with possibly another constant C. Since $u \in \mathcal{H}_{\infty}$ and $E_{\mathcal{H}}(\lambda_0)u = 0$, the above inequality applied to u shows that $e^{b_0 \langle x_0 \rangle^{1-\mu}} u \in \mathcal{H}$ for $0 < b_0 < b$.

Proof of Theorem 5.30. Since λ_0 is not an eigenvalue of H, Lemma 5.34 shows that $E_{\mathcal{H}}(\lambda_0)$ is injective in \mathcal{H}_s for any s. Writing

$$E_{\mathcal{H}}(\lambda_0 + \lambda) = -(F(\lambda) - \lambda)(1 + G(\lambda)U(\lambda))$$
(5.137)

for $\lambda \in \Omega_0(\delta)$, one sees that $1+G(0)\widetilde{U}(0)$ is injective. The mapping $\lambda \to G(\lambda)\widetilde{U}(\lambda) \in \mathcal{L}(\mathcal{H}_s)$ is a continuous and compact operator-valued. Consequently, $1+G(0)\widetilde{U}(0)$ is invertible and $(1+G(0)\widetilde{U}(0))^{-1} \in \mathcal{L}(\mathcal{H}_s)$ which implies, by continuity, $(1+G(\lambda)\widetilde{U}(\lambda))^{-1}$ exists and

$$\|(1+G(\lambda)U(\lambda))^{-1}\| \le C$$
for $\lambda \in \Omega_0(\delta)$, $\delta > 0$ small. This proves that $E_{\mathcal{H}}(\lambda_0 + \lambda)$ is invertible with the inverse given by

$$E_{\mathcal{H}}(\lambda_0 + \lambda)^{-1} = -(1 + G(\lambda)\widetilde{U}(\lambda))^{-1}G(\lambda) = -G(\lambda)(1 + \widetilde{U}(\lambda)G(\lambda)))^{-1}.$$
 (5.138)

As operator from \mathcal{H}_s to $\mathcal{H}_{s-2\mu}$, $E_{\mathcal{H}}(\lambda_0 + \lambda)^{-1}$ is uniformly bounded for $\lambda \in \Omega_0(\delta)$. Therefore the formula

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z)$$

initially valid for Im $z \neq 0$ can be extended to $z \in \Omega_{\lambda_0}(\delta)$. We conclude that H has no eigenvalue in $(\lambda_0 - \delta, \lambda_0)$, hence $\sigma_d(H)$ is finite.

E(z) and $E_{\pm}(z)$ are holomorphic for z near λ_0 . Since

$$e^{-a\langle x_0 \rangle^{1-\mu}} H' e^{a\langle x_0 \rangle^{1-\mu}} = H' + O(a)$$

 $e^{-a\langle x_0\rangle^{1-\mu}}R'(z)e^{a\langle x_0\rangle^{1-\mu}}$ is holomorphic and uniformly bounded for $|z - \lambda_0|$ small, provided that a > 0 is small. It follows that $e^{-a\langle x_0\rangle^{1-\mu}}E_+(z)e^{a\langle x_0\rangle^{1-\mu}}$ is holomorphic and uniformly bounded for $z \in \Omega_{\lambda_0}(\delta)$. Therefore to prove Theorem 5.30, it is sufficient to show that $e^{-a\langle x_0\rangle^{1-\mu}}E_{\mathcal{H}}(z)^{-1}$ belongs to $\mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_{\lambda_0}(\delta))$. For a > 0 small, $\tilde{U}(\lambda)e^{a\langle x_0\rangle^{1-\mu}}$ is holomorphic and uniformly bounded for $\lambda \in$

For a > 0 small, $U(\lambda)e^{a\langle x_0\rangle^{1-\mu}}$ is holomorphic and uniformly bounded for $\lambda \in \Omega_0(\delta)$. We conclude from (5.134) (with $\tau = 0$) that $\widetilde{U}(\lambda)G(\lambda) \in \mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_0(\delta))$ which, together with the uniform bound

$$\|(1+U(\lambda)G(\lambda))^{-1}\| \le C,$$

shows that $(1+\widetilde{U}(\lambda)G(\lambda))^{-1} \in \mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_0(\delta))$. For $z = \lambda_0 + \lambda$, it follows from (5.134) that

$$e^{-a\langle x_0\rangle^{1-\mu}} E_{\mathcal{H}}(z)^{-1} = -e^{-a\langle x_0\rangle^{1-\mu}} G(z-\lambda_0) (1+\widetilde{U}(z-\lambda_0)G(z-\lambda_0))^{-1}$$

belongs to $\mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_{\lambda_0}(\delta))$. This finishes the proof of Theorem 5.30.

Remarks 5.35. 1) Making use of (5.131) and repeating the proof of Theorem 5.30, one can prove that for any $N \in \mathbb{N}$ and $\tau \in \mathbb{R}$, one has

$$\|\langle x_0 \rangle^{-2(N+1)\mu-\tau} R^{(N)}(z) \langle x_0 \rangle^{\tau}\| \le C_{N,\tau}$$
(5.139)

uniformly for $\lambda \in \Omega_{\lambda_0}(\delta)$). This implies in particular that the limit

$$R(\lambda_0) = \lim_{\lambda \in \Omega_{\lambda_0}(\delta)), \lambda \to \lambda_0} R(\lambda)$$
(5.140)

exists in $\mathcal{L}(L^2_{\tau}, L^2_{\tau-2\mu-\epsilon})$ for any $\epsilon > 0$ and $R(\lambda_0) \in \mathcal{L}(L^2_{\tau}, L^2_{\tau-2\mu})$ for any $\tau \in \mathbb{R}$. Here and in the remaining part of this section, $L^2_r = L^2(\mathbf{X}, \langle x_0 \rangle^{2r} \mathrm{d}x)$.

2) Making use of (5.134), one can show the following improvement of Theorem 5.30: For any a > 0 and $\tau \ge 0$, there exists some constant C such that

$$\|\langle x_0 \rangle^{-\tau} \mathrm{e}^{-a \langle x \rangle^{1-\mu}} R^{(N)}(z) \langle x_0 \rangle^{\tau} \| \le C^{N+1} N! \langle N+1 \rangle^{\gamma(N+1)}$$
(5.141)

uniformly in $N \in \mathbb{N}$ and $z \in \Omega_{\lambda_0}(\delta)$.

Proposition 5.36. Assume the conditions of Theorem 5.30 and let $s > \frac{1+\mu}{2}$. Then the boundary values of the resolvent

$$R(\lambda \pm i0) = \lim_{\epsilon \to 0_+} R(\lambda \pm i\epsilon)$$

exist in $\mathcal{L}(L_s^2, L_{-s}^2)$ for $\lambda \in [\lambda_0, \lambda_0 + \delta]$, $\delta > 0$ and

$$\|\langle x_0 \rangle^{-s} R(\lambda \pm i0) \langle x_0 \rangle^{-s} \| \le C$$
(5.142)

uniformly in $\lambda \in [\lambda_0, \lambda_0 + \delta]$.

Proof. We keep the notation used before. It is known [Na] that (5.142) holds true for $r_0(z) = (h_0 - z)^{-1}$. Whence

$$\forall s > \frac{1+\mu}{2} \; \exists C > 0 : \quad \|\langle x_0 \rangle^{-s} r_0(\lambda \pm i0) \langle x_0 \rangle^{-s} \| \le C \tag{5.143}$$

uniformly in $\lambda \in [0, \delta]$. For $F(z) = h_0 + \chi_R W(z) \chi_R$, we write

$$F(z) - z = (h_0 - z)(1 + r_0(z)\chi_R W(z)\chi_R), \quad \text{Im} \, z \neq 0.$$

For $\lambda \geq 0$ small, $r_0(\lambda + i0)\chi_R W(\lambda)\chi_R$ is compact in \mathcal{H}_{-s} and continuous for $\lambda \in [0, \delta]$, $\delta > 0$. For R > 1 large, $F(0) \geq -\Delta_0 + \frac{c}{\langle x_0 \rangle^{2\mu}}$, c > 0, in sense of selfadjoint operators and Lemma 5.32 remains true if $E_{\mathcal{H}}(\lambda_0)$ is replaced by F(0). Consequently, F(0) is injective in \mathcal{H}_t for any $t \in \mathbb{R}$ because 0 is not an eigenvalue of F(0). This implies that $1 + r_0(0)\chi_R W(0)\chi_R$ is injective in \mathcal{H}_{-s} , hence $1 + r_0(0)\chi_R W(0)\chi_R$ is invertible. By the continuity in λ , we conclude that $(1 + r_0(\lambda \pm i0)\chi_R W(\lambda)\chi_R)$ is invertible in $\mathcal{L}(\mathcal{H}_{-s})$ for $\lambda > 0$ small and its inverse is continuous in $\lambda \in [0, \delta]$. Consequently the boundary values of $G(z) = (F(z) - z)^{-1}$ exist in $\mathcal{L}(\mathcal{H}_s, \mathcal{H}_{-s})$ and

$$G(\lambda \pm i0) = \left(1 + r_0(\lambda \pm i0)\chi_R W(\lambda)\chi_R\right)^{-1} r_0(\lambda \pm i0)$$

are continuous for $\lambda \in [0, \delta]$. A similar argument shows that the boundary values

$$E_{\mathcal{H}}(\lambda_0 + \lambda \pm \mathrm{i}0)^{-1} = -(1 + G(\lambda \pm \mathrm{i}0)\widetilde{U}(\lambda))^{-1}G(\lambda \pm \mathrm{i}0)$$

exist in $\mathcal{L}(\mathcal{H}_s, \mathcal{H}_{-s})$ and are continuous in $\lambda \in [0, \delta]$. Finally we obtain

$$R(\mu \pm i0) = E(\mu) - E_{+}(\mu)E_{\mathcal{H}}(\mu \pm i0)^{-1}E_{-}(\mu)$$

exist in $\mathcal{L}(L_s^2, L_{-s}^2)$ and are continuous in $\mu \in [\lambda_0, \lambda_0 + \delta]$.

Corollary 5.37. Let $e(\lambda)$ denote the spectral projector of H on $] -\infty, \lambda]$. Assume the conditions of Theorem 5.30. Then for any a > 0 and $s > \frac{1+\mu}{2}$, there exist some constants b, B > 0 such that

$$\|\mathrm{e}^{-a\langle x_0\rangle^{1-\mu}}e'(\lambda)\langle x_0\rangle^{-s}\| \le Be^{-b|\lambda-\lambda_0|^{-\frac{1}{\gamma}}}, \quad \lambda \in (\lambda_0, \lambda_0 + \delta).$$
(5.144)

Proof. Since $e'(\lambda) = \frac{1}{2\pi i} (R(\lambda + i0) - R(\lambda - i0)), ||\langle x_0 \rangle^{-s} e'(\lambda) \langle x_0 \rangle^{-s}||$ is uniformly bounded for $\lambda \in (\lambda_0, \lambda_0 + \delta)$. Iterating the first resolvent equation, one obtains for any $N \in \mathbb{N}$

$$e'(\lambda) = (\lambda - \lambda_0)^N R(\lambda_0)^N e'(\lambda).$$
(5.145)

Applying (5.141) with $\tau = s$, one deduces that for any a > 0, there exist some constants c, C > 0 such that

$$\|\mathrm{e}^{-a\langle x_0\rangle^{1-\mu}}e'(\lambda)\langle x_0\rangle^{-s}\| \le Cc^N N^{\gamma N} (\lambda - \lambda_0)^N$$
(5.146)

for all $N \in \mathbb{N}$ and $\lambda \in (\lambda_0, \lambda_0 + \delta)$. It remains to minimize the right-hand side by choosing N in terms of $\lambda - \lambda_0$ such that $N \approx A|\lambda - \lambda_0|^{-\frac{1}{\gamma}}$ as $\lambda \to \lambda_0$ for some appropriate constant A > 0. Then

$$c^{N} N^{\gamma N} |\lambda - \lambda_{0}|^{N} \approx e^{A|\lambda - \lambda_{0}|^{-\frac{1}{\gamma}} (\gamma \ln A + \ln c)}$$

$$\leq B e^{-b|\lambda - \lambda_{0}|^{-\frac{1}{\gamma}}}, \quad \lambda \in (\lambda_{0}, \lambda_{0} + \delta),$$

for some constants b, B > 0, if A > 0 is such that $\gamma \ln A + \ln c < 0$. This proves (5.144).

When λ_0 is an eigenvalue of H we prove the following analogue of Theorem 5.30.

Theorem 5.38. Assume the conditions (5.124), (5.125a) and (5.125b). Let λ_0 be an eigenvalue of H and Π_{λ_0} be the eigenprojection of H associated with λ_0 . Then one has

$$R(z) = -\frac{\Pi_{\lambda_0}}{z - \lambda_0} + R_1(z)$$
(5.147)

where, for any a > 0, $e^{-a\langle x_0 \rangle^{1-\mu}} R_1(z)$ belongs to the Gevrey class $\mathcal{G}^{(\frac{1+\mu}{1-\mu})}(\Omega_{\lambda_0}(\delta))$.

Proof. As in the proof of Theorem 5.30, we are led to study $E_{\mathcal{H}}(z)$ for $z \in \Omega_{\lambda_0}(\delta)$. We use formula (5.137) and another Grushin reduction to study $E_{\mathcal{H}}(\lambda_0 + \lambda)^{-1}$. The smoothness of $G(\lambda)$ and $\tilde{U}(\lambda)$ at $\lambda = 0$ implies that they can be expanded in appropriate spaces in powers of λ for λ near 0. Set

$$G(\lambda) = G_0 + \lambda G_1 + o(\lambda), \qquad (5.148)$$

$$\widetilde{U}(\lambda) = \widetilde{U}_0 + \lambda \widetilde{U}_1 + o(\lambda).$$
(5.149)

One has $G_j \in \mathcal{L}(\mathcal{H}_s, \mathcal{H}_{s-(j+1)\rho})$ and $\widetilde{U}_j \in \mathcal{L}(\mathcal{H}_s, \mathcal{H}_\infty)$ for any $s \in \mathbb{R}$.

Let k denote the multiplicity of eigenvalue λ_0 of H. Then 0 is an eigenvalue of $E_{\mathcal{H}}(\lambda_0)$ with multiplicity k. Let $K(z) = G(z)\widetilde{U}(z)$ and $K_0 = G_0\widetilde{U}_0$. Then K_0 is a compact operator in \mathcal{H} and ker $(1 + G_0\widetilde{U}_0) = \ker E_{\mathcal{H}}(\lambda_0)$. As in Subsection 5.1.2, we can choose a basis, $\{\phi_1, \ldots, \phi_k\}$, of ker $E_{\mathcal{H}}(\lambda_0)$ verifying

$$\langle -U_0\phi_i,\phi_j\rangle = \delta_{ij}, \quad 1 \le i,j \le k$$

and construct a Grushin problem for $1 + K(\lambda)$. Let

$$\mathcal{S}: \mathbb{C}^k \to \mathcal{H}, c = (c_1, \cdots, c_k) \to \mathcal{S}c = \sum_{j=1}^k c_j \phi.$$

Then $Q = SS^*$ is a projection onto ker $(1 + K_0)$. Let Q' = 1 - Q. One has $Q'(1 + K_0)Q'$ is invertible on ran Q' and by continuity, $Q'(1 + K(\lambda))Q'$ is invertible on ran Q' with uniformly bounded inverse for $\lambda \in \Omega_0(\delta)$, $\delta > 0$ small. Let

$$D(\lambda) = (Q'(1 + K(z)Q')^{-1}Q')^{-1}Q'$$

Since $(1 + K(\lambda)) \in \mathcal{G}^{(1+\gamma)}(\Omega_0(\delta))$ with $\gamma = \frac{2\mu}{1-\mu}$, one has $D(\lambda) \in \mathcal{G}^{(1+\gamma)}(\Omega_0(\delta))$. By studying the Grushin problem for 1 + K(z) using \mathcal{S} defined above, we obtain

$$(1 + K(\lambda))^{-1} = \mathcal{E}(\lambda) - \mathcal{E}_{+}(\lambda)\mathcal{E}_{-+}(\lambda)^{-1}\mathcal{E}_{-}(\lambda)$$
(5.150)

where

$$\begin{aligned} \mathcal{E}(\lambda) &= D(\lambda), \\ \mathcal{E}_{+}(\lambda) &= \mathcal{S} - D(\lambda)(1 + K(\lambda))\mathcal{S} \\ \mathcal{E}_{-}(\lambda) &= \mathcal{S}^{*} - \mathcal{S}^{*}(1 + K(\lambda))D(\lambda) \\ \mathcal{E}_{-+}(\lambda) &= \mathcal{S}^{*}(-(1 + K(\lambda)) + (1 + K(\lambda))D(\lambda)(1 + K(\lambda))\mathcal{S}. \end{aligned}$$

 $\mathcal{E}_{\pm}(\lambda)$ and $\mathcal{E}_{-+}(\lambda)$ all belong to Gevrey classes of order $1 + \gamma$. As in Subsection 5.2.1 Case 2, we can compute the $k \times k$ matrix $\mathcal{E}_{-+}(\lambda)$ et obtain

$$\mathcal{E}_{-+}(\lambda) = \lambda M_0 + \lambda^2 r_1(\lambda) \tag{5.151}$$

where M_0 is invertible and $r_1(\lambda) \in \mathcal{G}^{(1+\gamma)}(\Omega_0(\delta))$. This leads to

$$\mathcal{E}_{-+}(\lambda)^{-1} = \lambda^{-1}M_0^{-1} + r_2(\lambda)$$

with $r_2(\lambda) \in \mathcal{G}^{(1+\gamma)}(\Omega_0(\delta))$. By (5.138) and (5.150), we obtain

$$E_{\mathcal{H}}(\lambda_0 + \lambda)^{-1} = -G(\lambda)(1 + K(\lambda))^{-1} = \frac{C_0}{\lambda} + \widetilde{R}_1(\lambda)$$
(5.152)

in $\mathcal{L}(\mathcal{H}_s, \mathcal{H}_{s-2\mu-\epsilon})$, where $C_0 = G_0 \mathcal{E}_+(0) M_0^{-1} \mathcal{E}_-(0)$ and for a > 0, $e^{-a \langle x_0 \rangle^{1-\mu}} \widetilde{R}_1(\lambda) \in \mathcal{G}^{(1+\gamma)}(\Omega_0(\delta))$. Therefore, R(z) verifies the expansion

$$R(z) = -\frac{B_0}{z - \lambda_0} + R_1(z), \quad B_0 = E_+(\lambda_0)C_0E_-(\lambda_0)$$

One can show as in the proof of Theorem 5.30 that

$$e^{-a\langle x_0\rangle^{1-\mu}}R_1(z)\in \mathcal{G}^{(1+\gamma)}(\Omega_{\lambda_0}(\delta)).$$

One has necessarily $B_0 = \prod_{\lambda_0}$ by the spectral theorem for H.

Using Cauchy integral formula to represent e^{-tH} in terms of the resolvent, we obtain from Theorem 5.38 a large-time expansion for the heat semi-group e^{-tH} (cf. [Wa6, Theorem 2.3]).

Corollary 5.39. Assume the conditions of Theorem 5.30. For any a > 0, there exist some constants C, c > 0 such that

$$\left\| \mathrm{e}^{-a\langle x_0 \rangle^{1-\mu}} \left(\mathrm{e}^{-tH} - \sum_{\lambda \in \sigma_d(H)} \mathrm{e}^{-t\lambda} \Pi_{\lambda} - \mathrm{e}^{-t\lambda_0} \Pi_{\lambda_0} \right) \right\| \le C \mathrm{e}^{-t\lambda_0 - ct^{\frac{1-\mu}{1+\mu}}}.$$
 (5.153)

Here Π_{λ} is the eigenprojector of H associated with eigenvalue λ .

- **Remarks 5.40.** 1) For one-body operators it is shown in [AW, Wa6] that under some additional conditions the quantum dynamics e^{-itH} , when regarded as operator from L^2_{comp} to L^2_{loc} , can be expanded as $|t| \to \infty$ with the same subexponential estimates on the remainder as in Corollary 5.39. The conditions used there exclude a possible accumulation of quantum resonances towards threshold zero. It is an interesting and non-trivial open question to see if a similar result holds true for e^{-itH} in the N-body problem.
 - 2) If $\lambda_0 = \Sigma_2$ is a multiple two-cluster threshold one can apply the Grushin reduction of Section 2.3 to show that Theorems 5.30 and 5.38 still hold true. Since the proof of Theorems 5.30 and 5.38 relies heavily on the continuity of R'(z)in exponentially weighted spaces, which is deduced from the holomorphicity of R'(z) for z near λ_0 , one can not expect these results to be true for higher thresholds $\lambda_0 > \Sigma_2$.

Example 5.41. Theorems 5.30 and 5.38 can be applied to physics models with Coulomb interactions given by (1.13). Assume that the lowest threshold $\lambda_0 = \Sigma_2$ is non-multiple two-cluster and is equal to the lowest eigenvalue of a two-cluster Hamiltonian H^a with $a = (C_1, C_2)$. Let $Q_j = \sum_{k \in C_j} q_k$ be the total charge of particles in cluster C_j , j = 1, 2. Assume that

$$Q_1 Q_2 > 0.$$

Then the effective potential is positive and slowly decreasing outside a compact set and (5.125b) is satisfied with $\rho = 1$ (see (1.14)). In this case, Theorems 5.30 and 5.38 hold true in Gevrey class $\mathcal{G}^{(3)}(\Omega_{\lambda_0}(\delta))$.

5.3 Resolvent asymptotics for physics models near two-cluster thresholds

We will discuss some extensions of Section 5.1 for the physics models of Sections 1.2 and 1.3 (with $N \ge 3$ and $N \ge 2$, respectively) using the same notation as in Section 4.3. As in the previous sections of this chapter the single particle space dimension is fixed as n = 3. In agreement with the setting of Section 4.3 we shall not distinguish between the cases $\lambda_0 = \Sigma_2$ and $\lambda_0 > \Sigma_2$.

We recall that for a given two-cluster threshold λ_0 for the models of Sections 1.2 and 1.3 we group the set of two-cluster decompositions a for which $\lambda_0 \in \sigma_{pp}(H^a)$ into \mathcal{A}_1 , \mathcal{A}_2 and \mathcal{A}_3 for which

 \mathcal{A}_1 : the effective inter-cluster interaction is to leading order attractive Coulombic,

 \mathcal{A}_2 : the effective inter-cluster interaction is to leading order repulsive Coulombic,

 \mathcal{A}_3 : the effective inter-cluster interaction is $\mathcal{O}(|x_a|^{-2})$.

This distinction does not depend on choices of corresponding sub-Hamiltonian bound states φ^a (i.e. channels); it is determined by charges only (cf. Case 1 introduced independently in each of Sections 1.2 and 1.3).

Let for $a \in \mathcal{A} := \mathcal{A}_1 \cup \mathcal{A}_2 \cup \mathcal{A}_3$ the operator P^a be the corresponding orthogonal projection onto ker $(H^a - \lambda_0)$ in $L^2(\mathbf{X}^a)$ and let m_a be the dimension of this space. Obviously $\Pi^a := P^a \otimes 1$ projects onto the span of functions of the form $\varphi^a \otimes f_a$, $\varphi^a \in \text{ker}(H^a - \lambda_0)$, in $L^2(\mathbf{X})$. We identify ran P^a , say spanned by an orthonormal basis $\varphi^a_1, \ldots \varphi^a_{m_a}$, with \mathbb{C}^{m_a} (using the basis), and similarly

$$\mathcal{H}_a := L^2(\mathbf{X}_a, \mathbb{C}^{m_a}) \simeq \bigoplus_{m \le m_a} L^2(\mathbf{X}_a) \ni \bigoplus_{m \le m_a} f_{a,m} = f_a$$
$$\simeq S_a f_a := \sum_{m \le m_a} \varphi_m^a \otimes f_{a,m} \in \operatorname{ran} \Pi^a.$$

The effective potential for $a \in \mathcal{A}_3$ obeys $W_a := S_a^* I_a S_a = Q_a |x_a|^{-2} + B_a$, where Q_a is a $m_a \times m_a$ matrix-valued function depending only on $\theta = \hat{x}_a = |x_a|^{-1} x_a$ while $B_a = B_a(x_a) = \mathcal{O}(|x_a|^{-3})$. We shall here only study the case where $Q_a = 0$, meaning that the effective potential is *fastly decaying*. The 'right generalization' to the case $m_a > 1$ of the distinction between Cases 2) and 3) (discussed primarily for $m_a = 1$ in both of Sections 1.2 and 1.3) is to let Case 2) correspond to $Q_a \neq 0$ and let Case 3) correspond to $Q_a = 0$, respectively. This means that we shall not consider Case 2) defined in Section 4.3. Whence, letting $\mathcal{A}_3^{\text{fd}} = \{a \in \mathcal{A}_3 | Q_a = 0\}$, we assume that

$$\mathcal{A}_3 = \mathcal{A}_3^{\rm fd}.\tag{5.154}$$

For simplicity we shall also assume (4.68a) and (4.68b) leaving us with studying the Grushin resolvent representation (2.6) where (as in Section 4.3) $\mathcal{H} = \sum_{a \in \widetilde{\mathcal{A}}} \oplus \mathcal{H}_a$, $\mathcal{H}_a = L^2(\mathbf{X}_a, \mathbb{C}^{m_a}) = \bigoplus_{m \leq m_a} L^2(\mathbf{X}_a)$ and $S = (S_a) : \mathcal{H} \to \mathcal{F} \subset \mathcal{G} = L^2(\mathbf{X})$ is given by

$$f = \bigoplus_{a \in \widetilde{\mathcal{A}}} f_a \to Sf = \sum_{a \in \widetilde{\mathcal{A}}} S_a f_a; \quad f_a = \sum_{m \le m_a} \oplus f_{a,m}, \quad S_a f_a = \sum_{m \le m_a} \varphi_m^a \otimes f_{a,m}.$$

Let (as usual) $T = (SS^*)^{-1}S$. These operators S and T will freely be used on weighted spaces, and we also adapt the following notation of Theorem 5.14.

$$S_I f := \sum_{a \in \widetilde{\mathcal{A}}} I_a S_a f_a; \quad f = \bigoplus_{a \in \widetilde{\mathcal{A}}} f_a.$$

We write $-E_{\mathcal{H}}(\lambda_0 + z) = P(z) = P_0 + U(z) - z$, where $P_0 = \bigoplus_{a \in \widetilde{\mathcal{A}}} h_a$ with h_a specified as follows: If $a \in \mathcal{A}_1$ or $a \in \mathcal{A}_2$ we take $h_a = p_a^2 + w_a$ (acting as a diagonal operator if $m_a > 1$) where w_a is constructed as in Subsection 4.1.2 or 4.1.3, respectively. If $a \in \mathcal{A}_3^{\mathrm{fd}} = \mathcal{A}_3$ we take $h_a = p_a^2$. We define $r_0(z) = (P_0 - z)^{-1} = \bigoplus_{a \in \widetilde{\mathcal{A}}} r_a(z)$ for any $z \in \mathbb{C} \setminus \mathbb{R}$. Next we write

$$P(z)^{-1} = \left(1 + r_0(z)U(z)\right)^{-1} r_0(z) = W(z)^{-1} r_0(z), \quad z \in \mathbb{C} \setminus \mathbb{R},$$
(5.155)

as in the previous sections.

If λ_0 is regular, meaning that λ_0 is neither an eigenvalue nor a resonance of H (or, more precisely, that the set \mathcal{E} in Theorem 4.17 is the zero set), then ker $W^{\pm}(0) = 0$ where

$$W^{\pm}(0) = \lim_{\epsilon \to 0} W^{\pm}(\pm i\epsilon) = 1 + \lim_{\epsilon \to 0} r_0(\pm i\epsilon) \lim_{\epsilon \to 0} U(\pm i\epsilon) = 1 + r_0^{\pm} U_0^{\pm}$$
(5.156)

with limits taken in appropriate spaces. By Fredholm theory we would then obtain limits $\lim_{\epsilon\to 0} (P \mp i\epsilon)^{-1}$ and then in turn, due to (5.2), limits $\lim_{\epsilon\to 0} R(\lambda_0 \pm i\epsilon)$. Actually for λ_0 being regular we can take limits for $z \to 0$ in the quadrants

$$\mathcal{Z}_{\pm} = \{\operatorname{Re} z \ge 0, \pm \operatorname{Im} z > 0\}.$$

To be more precise about these assertions we first look at the three types of (diagonal) building blocks $r_a(z) = (h_a - z)^{-1}$. Recall the notions from Section 4.3,

$$\mathcal{H}_{a,s}^k = H_s^k(\mathbf{X}_a, \mathbb{C}^{m_a}) \text{ for } a \in \widetilde{\mathcal{A}}, \ k \in \mathbb{R} \text{ and } s \in \mathbb{R}.$$

If $a \in \mathcal{A}_1$ there are limits

$$r_a(0\pm \mathrm{i}0) = \lim_{z\to 0, z\in\mathcal{Z}_{\pm}} r_a(z) \text{ in } \mathcal{L}\big(\mathcal{H}_{a,s}^{-1}, \mathcal{H}_{a,-s}^1\big), \quad s > 3/4,$$

cf. Subsection 4.1.2. Similarly, if $a \in \mathcal{A}_2$ there are limits

$$r_a(0\pm \mathrm{i}0) = \lim_{z\to 0, z\in\mathcal{Z}_{\pm}} r_a(z) \text{ in } \mathcal{L}\big(\mathcal{H}_{a,s}^{-1}, \mathcal{H}_{a,-s}^1\big), \quad s > 3/4,$$

however in this case $r_a(0+i0) = r_a(0-i0)$ and the common limit coincides with the quantity h_a^{-1} of Lemma 4.4, cf. [Na] and [Ya2]. Finally, for $a \in \mathcal{A}_3^{\text{fd}}$ we know from the previous sections that

$$r_a(0\pm \mathrm{i}0) = \lim_{z\to 0, z\in\mathcal{Z}_{\pm}} r_a(z) \text{ in } \mathcal{L}\big(\mathcal{H}_{a,s}^{-1}, \mathcal{H}_{a,-s}^1\big), \quad s>1,$$

again with $r_a(0 + i0) = r_a(0 - i0)$ and in this case the common limit G_0 is given explicitly (each diagonal entry has the kernel $(4\pi)^{-1}|x_a - y_a|^{-1}$).

We are lead to consider for $k \in \mathbb{R}$, r, s < -3/4 and t < -1 the spaces

$$\mathcal{H}^k_{r,s,t} = \oplus_{b \in \mathcal{A}_1} \, \mathcal{H}^k_{b,r} \bigoplus \oplus_{b \in \mathcal{A}_2} \, \mathcal{H}^k_{b,s} \bigoplus \oplus_{b \in \mathcal{A}_3} \, \mathcal{H}^k_{b,t}.$$

We could for example fix $(r, s, t) = \overline{t} := -(1, 1, 4/3)$, and the corresponding spaces $\mathcal{H}_{\overline{t}}^k$ could then be used by considering $r_0(z) \in \mathcal{L}(\mathcal{H}_{-\overline{t}}^{-1}, \mathcal{H}_{\overline{t}}^1)$, $U(z) \in \mathcal{L}(\mathcal{H}_{\overline{t}}^1, \mathcal{H}_{-\overline{t}}^{-1})$ and therefore, in turn, $W(z) \in \mathcal{L}(\mathcal{H}_{\overline{t}}^1)$; $z \in \mathcal{Z}_{\pm}$. With these interpretations we can check that the limits in (5.156) exist. Since λ_0 is regular for H it follows that ker $W^{\pm}(0) = 0$. Since $W^{\pm}(0) \in \mathcal{C}(\mathcal{H}_{\overline{t}}^1)$ this allows us to take the $z \to 0$ limits in (5.155) as well.

We made an unnecessary simplifying assumption on the parameters, and using the spaces $\mathcal{H}_{r,s,t}^k$ we may similarly deduce the following result.

Proposition 5.42. Suppose the conditions of Theorem 4.17 and in addition (5.154), (4.68a) and (4.68b). Suppose λ_0 is regular. Then for all sufficiently big r, s < -3/4 and t < -1 there is an $\epsilon > 0$ such that the following asymptotics holds in $\mathcal{L}(\mathcal{H}^1_{r,s,t})$ for $z \to 0$ in \mathcal{Z}_{\pm} .

$$W(z)^{-1} = W^{\pm}(0)^{-1} + \mathcal{O}(|z|^{\epsilon}).$$
 (5.157a)

Similarly, in the space $\mathcal{L}(\mathcal{H}_{-r,-s,-t}^{-1},\mathcal{H}_{r,s,t}^{1})$

$$P(z)^{-1} = (P^{\pm})^{-1} + \mathcal{O}(|z|^{\epsilon}); \quad (P^{\pm})^{-1} := W^{\pm}(0)^{-1}r_0^{\pm}.$$
(5.157b)

Theorem 5.43 (Regular case). Under the conditions of Proposition 5.42 the following asymptotics hold for $R(\lambda_0 + z)$ as an operator from H_t^{-1} to H_{-t}^1 , t > 1, for $z \to 0$ in \mathcal{Z}_{\pm} and for some $\epsilon = \epsilon(t) > 0$.

$$R(\lambda_0 + z) = R'(\lambda_0 \pm i0) + (S - R'(\lambda_0 \pm i0)S_I)(P^{\pm})^{-1}(S^* - S_I^*R'(\lambda_0 \pm i0)) + \mathcal{O}(|z|^{\epsilon}).$$
(5.158)

This result is immediate from (2.6) and Proposition 5.42, however the latter results actually give more detailed information on the (anisotropic) behaviour of the resolvent. For example the 'most singular part' of $R(\lambda_0 + z)$ is given by the term $S(P - z)^{-1}S^* \in \mathcal{L}(H_t^{-1}, H_{-t}^1)$, but $S(P - z)^{-1}S^* \approx Sr_0(z)S^* \approx \Sigma_{a \in \mathcal{A}}P^a \otimes r_a(z)$ needs t > 1 only for $a \in \mathcal{A}_3^{\text{fd}}$; $r_a(z)$ is 'smaller' for $a \in \mathcal{A}_1 \cup \mathcal{A}_2$.

We can also derive a result if λ_0 is an exceptional point of the second kind, meaning that the set \mathcal{E} in Theorem 4.17 obeys $0 \neq \mathcal{E} \subset L^2$. For that we need the additional condition

$$\operatorname{ran} \Pi_H \subset L_t^2 \text{ for some } t > 1, \tag{5.159}$$

where Π_H is the eigenprojection corresponding to the eigenvalue λ_0 of H (note that $\Pi_H L^2 = \mathcal{E}$). We can argue as for the last assertion of Theorem 5.16, see also Remark 5.17 2). This means more precisely that we use the above procedure for $H_{\sigma} := H - \sigma \Pi_H, \sigma > 0$ small. Under the given hypotheses λ_0 is neither an eigenvalue nor a resonance of H_{σ} , and therefore there is an analogous version of (5.158) for H_{σ} and we can then (as in the previous sections) invoke (5.87). Hence we obtain the following result, where quantities depending on the 'potential' $-\sigma \Pi_H$ are equipped with the subscript σ .

Theorem 5.44 (Exceptional point of 2nd kind). Suppose the conditions of Theorem 4.17 and in addition (5.154), (4.68a), (4.68b) and (5.159). Suppose λ_0 is an eigenvalue but not a resonance of H. Then the following asymptotics hold for $R(\lambda_0 + z)$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ in \mathcal{Z}_{\pm} and for some $\epsilon = \epsilon(s) > 0$.

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \sigma^{-1}\Pi_H + R'_{\sigma}(\lambda_0 \pm i0) + (S - R'_{\sigma}(\lambda_0 \pm i0)S_{I_{\sigma}})(P^{\pm}_{\sigma})^{-1}(S^* - S^*_{I_{\sigma}}R'_{\sigma}(\lambda_0 \pm i0)) + \mathcal{O}(|z|^{\epsilon}).$$
(5.160)

Note that the $z^{-\frac{1}{2}}$ -term is absent in (5.160). This looks as if there is a discrepancy with the known results given in [JK, Wa2] where there are $z^{-\frac{1}{2}}$ -terms in the resolvent expansions for exceptional point of the second kind. However there is not the case, because under the decay assumption on the threshold eigenstates used in Theorem 5.44, one can check by an explicit calculation that the $z^{-\frac{1}{2}}$ -terms of [JK, Wa2] also disappear.

It remains to examine the cases where λ_0 is an exceptional point of the first or of the third kind, meaning (treating them uniformly) that the set \mathcal{E} in Theorem 4.17 is not strictly a subset of L^2 . For that we need the additional condition

$$\operatorname{ran} \Pi_H \subset L_t^2 \text{ for some } t > 3/2, \tag{5.161}$$

where Π_H is the orthogonal projection onto ker $(H - \lambda_0)$ (i.e. the eigenprojection if λ_0 is an eigenvalue of H). We know from Theorem 4.17 that the condition $\mathcal{E} \not\subset L^2$ needs $\mathcal{A}_3 \neq \emptyset$. If $\mathcal{A}_1 \neq \emptyset$ and $\mathcal{A}_3^{\rm fd} \neq \emptyset$ there is a technical problem on identifying the geometric and algebraic multiplicities for the eigenvalue -1 of certain operators

 K^{\pm} , cf. (5.47). The same method as the one used in the previous sections does not work (and in fact we dont know if the multiplicities are equal). On the other hand under the additional condition

$$\widetilde{\mathcal{A}} = \mathcal{A}_2 \cup \mathcal{A}_3^{\rm fd},\tag{5.162}$$

the analogue of (5.47) holds (for H_{σ}), by the same proof (see also Remark 3.21 2)). In this case we can mimic the proof of Theorem 5.16 and obtain the following result, which is very similar to Theorems 5.16 and 5.26.

Theorem 5.45 (Exceptional point of 1st or 3rd kind). Suppose the conditions of Theorem 4.17 and in addition (4.68a), (4.68b), (5.161) and (5.162). Suppose λ_0 is a resonance of H. Then the following asymptotics hold for $R(\lambda_0 + z)$ as an operator from H_s^{-1} to H_{-s}^1 , s > 1, for $z \to 0$ in \mathcal{Z}_{\pm} and for some $\epsilon = \epsilon(s) > 0$.

$$R(\lambda_0 + z) = -z^{-1}\Pi_H + \frac{i}{\sqrt{z}} \sum_{j=1}^{\kappa} \langle u_j, \cdot \rangle u_j + \mathcal{O}(|z|^{-\frac{1}{2} + \epsilon}).$$
(5.163)

Here $\{u_1, \ldots, u_\kappa\} \subset H^1_{(-1/2)^-}$ is a basis of resonance states of H being independent of the choice of the sign of \mathcal{Z}_{\pm} .

Remarks 5.46. The basis of resonance states can be specified in a fashion similar to normalization procedures in Theorem 5.14 and Remark 5.25, cf. (5.13a) and (5.13b) (and subsequent the computations).

We consider the imposed conditions (4.68a) and (4.68b) as technically convenient but not being crucial. The interested reader may trace the outlined proof of Theorem 4.17 for the general case where these conditions are not imposed and see how the above theory modifies.

We already discussed our need for (5.162), however we remark that (5.154) is not strictly necessary. In fact there are some results for the case where $\mathcal{A}_3 \setminus \mathcal{A}_3^{\text{fd}} \neq \emptyset$ (this is an interesting case for the physics models). Then the spherical potential Q_a should not to be 'too negative', more precisely we need that $\sigma(-\Delta_{\theta} + Q_a) \subset (-1/4, \infty)$ for all a in this set. With this extension there should be analogous results on the resolvent expansion at λ_0 , not to be elaborated on here, see [Wa5]. However we shall later do a version of Theorem 5.43 in a special case where indeed $\mathcal{A}_3 \setminus \mathcal{A}_3^{\text{fd}} \neq \emptyset$, see Subsection 6.3.2. If the above spectral condition on $a \in \mathcal{A}_3 \setminus \mathcal{A}_3^{\text{fd}}$ is not fulfilled the resolvent asymptotics would be expected to be oscillatory, see [SW] where oscillatory behaviour is detected for a one-body 'toy model'. We shall not study this case; it does not seem to be an 'easy problem'.

Chapter 6 Applications

We will give applications of the previous chapters to scattering theory. We shall primarily study the non-multiple case imposing Conditions 1.5 and 2.4. With additional efforts the multiple is treatable somewhat similarly, see Subsectons 6.1.3 and 6.4.3 for actual accounts of the multiple case. The non-multiple case appears rather complicated already, and we believe that treating only this case may be considered as 'heart of the matter'. In Sections 6.2, 6.3 and 6.4 we consider only the physics models of Subsections 1.2.1 and 1.3.1 with the particle dimension n = 3 (in Section 6.4 even more specialized).

Although most of the material presented in this chapter is new it depends on the literature, obviously most importantly for example [DS1, JKW]. Since scattering theory is an old well-studied subject the literature is large, let us here mention the related works [Bo, Do, CT, De1, De2, De3, DG, Is2, Is3, Is4, IT, Ne, Sk2, Sk4, Sk5, Sk6, Sk7, SW, Wa6, Ya1, Ya3, Ya4, Ya5]. Obviously this list is not complete.

6.1 Negative slowly decaying effective potentials

We impose the attractiveness condition (4.6) for the cluster decomposition $a = a_0$ given in Condition 2.4. More precisely suppose $\rho < 2$ and that the inter-cluster potential $I_a(x_a) = I_{a_0}(x_a) = I_0(x_a)$ fulfills the condition

$$\exists R \ge 0 \quad \exists \epsilon > 0 \quad \forall y \in \mathbf{X}_a \text{ with } |y| \ge R :$$

$$I_a(y) \le -\epsilon \langle y \rangle^{-\rho} \text{ and } -2I_a(y) - y \cdot \nabla I_a(y) \ge \epsilon \langle y \rangle^{-\rho}.$$
(6.1)

We assume

$$\lambda_0 \notin \sigma_{\rm pp}(H'),\tag{6.2}$$

or equivalently $\lambda_0 \notin \sigma_{\rm pp}(\check{H})$. If (6.2) is not fulfilled we may modify the theory to be discussed in agreement with Subsection 2.4, see Remark 4.18.

We know from Chapter 3 that the boundary values $\check{R}(\lambda \pm i0)$ are smooth (in weighted spaces) in a real neighbourhood $I \ni \lambda_0$. We take $I, R = R_0 \ge 1$ and the operator $B = B_R =: \check{B}$ exactly as done in Subsection 3.3.2 (this *B* should not to be mixed up with the operator $B_{\kappa} = B_{\kappa,R} = B(\kappa^2 B^2 + 1)^{-1}$ in the same chapter).

In addition we assume

$$\lambda_0 \notin \sigma_{\rm pp}(H). \tag{6.3}$$

If (6.3) is not fulfilled and Π_H is the corresponding eigenprojection then $\lambda_0 \notin \sigma_{\rm pp}(H-\sigma\Pi_H)$ for $\sigma > 0$, and we can consider the H' construction for $H_{\sigma} = H - \sigma\Pi_H$, say denoted by H'_{σ} , which fits well onto the framework of Chapter 3 since the eigenfunctions decay polynomially, cf. Theorems 4.2, 4.12 and 4.17 1). In fact if $\sigma > 0$ is taken small enough the condition $\lambda_0 \notin \sigma_{\rm pp}(H')$ implies the same property with H' replaced by H'_{σ} , see the discussion before Theorem 5.26. This would lead to a resolvent formula similar to (5.160). For simplicity we impose (6.2) as well as (6.3), which leads to the following analogue (6.4) of the physics models resolvent formula (5.158).

With these conditions we know from Chapters 3 and 4 (see also Remark 5.17 3)) that there exist continuous boundary values $R(\lambda \pm i0)$ (in appropriate spaces) in an interval of the form $I_{\delta}^{+} = [\lambda_0, \lambda_0 + \delta] \subset I$ with $\delta > 0$ small. In fact we have the formulas

$$R(\lambda \pm i0) = E^{\pm}(\lambda) - E^{\pm}_{+}(\lambda)E^{\pm}_{\mathcal{H}}(\lambda)^{-1}E^{\pm}_{-}(\lambda), \qquad (6.4)$$

where, abbreviating $I_0 = I_{a_0}$ and $p_0 = p_{a_0}$,

$$\begin{aligned} E^{\pm}(\lambda) &= R'(\lambda \pm i0), \\ E^{\pm}_{+}(\lambda) &= (1 - R'(\lambda \pm i0)I_0)S, \\ E^{\pm}_{-}(\lambda) &= S^*(1 - I_0R'(\lambda \pm i0)), \\ E^{\pm}_{\mathcal{H}}(\lambda) &= (\lambda - \lambda_0) - (p_0^2 + S^*I_0S - S^*I_0R'(\lambda \pm i0)I_0S). \end{aligned}$$

We also have

$$R'(\lambda \pm i0) = \breve{R}(\lambda \pm i0) - (p_0^2 - \lambda)^{-1}\Pi = \breve{R}(\lambda \pm i0)\Pi'.$$

The matrix-valued operator $W = S^*I_0S$ is well approximated by a diagonal one fulfilling a global virial condition, in fact approximated by a multiple of the identity say denoted by $w\mathbf{1}$, cf. the discussion in the beginning of Subsection 4.1.2. As in the same subsection we modify the potential V(z) correspondingly and denote the result by v(z). Let $h = (p_0^2 + w)\mathbf{1}$, $r(z) = (h - z)^{-1}$ and

$$r_{\lambda}^{\pm} = r(\lambda - \lambda_0 \pm i0) \text{ and } v_{\lambda}^{\pm} = v(\lambda \pm i0),$$

and note that

$$-E_{\mathcal{H}}^{\pm}(\lambda) = h + v_{\lambda}^{\pm} - (\lambda - \lambda_{0}),$$

$$-E_{\mathcal{H}}^{\pm}(\lambda)^{-1} = r_{\lambda}^{\pm}(\lambda) \left(\mathbf{1} + v_{\lambda}^{\pm}r_{\lambda}^{\pm}(\lambda)\right)^{-1};$$

$$v_{\lambda}^{\pm} = -S^{*}I_{0}R'(\lambda \pm i0)I_{0}S + \mathcal{O}(r^{-1-\rho}).$$

(6.5)

If $m_a := \dim \ker(H^a - \lambda_0) = 1$, $a = a_0$, we can here replace $\mathcal{O}(r^{-1-\rho})$ by $\mathcal{O}(r^{-\infty})$ which would refer to a polynomially decreasing term. In any case the term is a λ independent local potential (i.e. a function). The above expressions are substituted into (6.4) to obtain formulas for $R(\lambda \pm i0)$ to be studied.

Note that

$$(1 + v_{\lambda}^{\pm} r_{\lambda}^{\pm})^{-1} \in \mathcal{L}(L_s^2) \cap \mathcal{L}(\mathcal{B}_{s_0}); \quad s \in (s_0, 1/2 + \rho 3/4), \ s_0 = 1/2 + \rho/4.$$
 (6.6)

This is the best we can do when $m_a > 1$. If $\lambda_0 > \Sigma_2$ and $m_a = 1$ then any $s \in (s_0, 1/2 + \rho)$ works. On the other hand if $\lambda_0 = \Sigma_2$ and $m_a = 1$ then this inverse exists in $\mathcal{L}(L_s^2)$ for any $s \in (s_0, 2\rho + 2 - s_0)$, cf. (4.12). In particular we have the following formula for any $\psi \in \mathcal{B}_{s_0}$ (we abbreviate throughout this section $\mathcal{B} := \mathcal{B}_{1/2}$, $\mathcal{B}^* = \mathcal{B}_{1/2}^*$ and $\mathcal{B}_0^* := \mathcal{B}_{1/2,0}^*$),

$$R(\lambda \pm i0)\psi = S\phi_{a}^{\pm}(\lambda) + \mathring{R}(\lambda \pm i0)\psi_{a}^{\pm}(\lambda);$$

$$\phi_{a}^{\pm}(\lambda) = r_{\lambda}^{\pm}f_{a}^{\pm}(\lambda) \in \mathcal{B}_{s_{0}}^{*}(\mathbf{X}_{a}),$$

$$f_{a}^{\pm}(\lambda) = (\mathbf{1} + v_{\lambda}^{\pm}r_{\lambda}^{\pm})^{-1}S^{*}(1 - I_{0}\check{R}(\lambda \pm i0)\Pi')\psi \in \mathcal{B}_{s_{0}}(\mathbf{X}_{a}),$$

$$\psi_{a}^{\pm}(\lambda) = \Pi'(\psi - I_{0}S\phi_{a}^{\pm}(\lambda)) \in \mathcal{B}.$$
(6.7)

It follows that for any $s > s_0$ the $\mathcal{L}(L_s^2, L_{-s}^2)$ -valued functions $R(\cdot \pm i0)$ are continuous on I_{δ}^+ . For each $\lambda \in I_{\delta}^+$ the operators $R(\lambda \pm i0) \in \mathcal{L}(\mathcal{B}_{s_0}, \mathcal{B}_{s_0}^*)$. It will be convenient to isolate the 'main parts' of (6.7) writing

$$R(\lambda \pm i0) = Sr_{\lambda}^{\pm}S^* + \check{R}(\lambda \pm i0)\Pi' + \check{R}(\lambda \pm i0).$$
(6.8)

Here $\check{R}(\lambda \pm i0)$ are represented as sums of various terms. Note for example that for any $\psi \in \mathcal{B}_{s_0}$ (as above) $f_a^{\pm}(\lambda) - S^* \psi \in L_s^2$ for any $s \in (s_0, 1/2 + \rho 3/4)$.

Remark 6.1. Note that in particular (6.4) as well as (6.7) and (6.8) are valid for $\lambda = \lambda_0$. This is a consequence of the imposed regularity condition (6.3). For other models, to be treated in Section 6.3, there are similar formulas as (6.7) and (6.8). Again this requires regularity, i.e. absence of bound and resonance states at the threshold. Note also that the trick of replacing H by $H_{\sigma} = H - \sigma \Pi_{H}$ in the case (6.3) is not fulfilled is not restricted to the case of attractive slowly decaying effective potentials, i.e. the condition (6.1), but can be used as well for repulsive slowly decaying effective potentials fulfilling the following version of (4.15),

$$\exists R \ge 0 \ \exists \epsilon > 0 \ \exists \bar{\rho} \in [\rho, \frac{2}{3}(1+\rho)) \ \forall y \in \mathbf{X}_a \text{ with } |y| \ge R : \quad I_a(y) \ge \epsilon \langle y \rangle^{-\bar{\rho}}.$$

This is manifestly done already in the proof of Theorem 5.44 and as before doable thanks to the polynomial decay of the eigenfunctions. For possible threshold eigenfunctions for non-slowly decaying effective potentials (cf. the asymptotics condition (4.20)) the polynomial decay is missing making (6.3) a non-trivial assumption in such cases, see however Theorem 5.44.

6.1.1 Sommerfeld's theorem

We impose the above conditions (6.1)–(6.3) on the threshold λ_0 . First we recall the following version of the Sommerfeld's theorem above λ_0 , see [AIIS, Corollary 1.10] which extends the seminal work [Is4].

Theorem 6.2. For any $\lambda \in I_{\delta}^+ \setminus \{\lambda_0\}$ there exist $R = R_0 \ge 1$ and $\sigma > 0$ such that for any real function $\chi(\cdot < \sigma) \in C^{\infty}(\mathbb{R})$, which is supported in $(-\infty, \sigma)$ and whose derivative has compact support, and for any $\psi \in \mathcal{B}$

$$\chi(\pm \check{B} < \sigma)R(\lambda \pm i0)\psi \in \mathcal{B}_0^*; \, \check{B} = B_R.$$
(6.9)

Moreover (for each sign) $\phi = R(\lambda \pm i0)\psi (\in \mathcal{B}^*)$ solves $(H - \lambda)\phi = \psi$.

Conversely if $\phi \in L^2_{-\infty}$ solves $(H - \lambda)\phi = \psi$ for a given $\psi \in \mathcal{B}$ and (for either 'plus' or 'minus') $\chi(\pm \check{B} < \sigma)\phi \in \mathcal{B}^*_0$ for some $\sigma > 0$ and for all functions $\chi(\cdot < \sigma)$ of this type, then $\phi = R(\lambda \pm i0)\psi$ (with the same sign).

Here (most likely) $R = R(\lambda) \to \infty$ and $\sigma = \sigma(\lambda) \to 0$ for $\lambda \downarrow \lambda_0$. We state a new version of the Sommerfeld theorem, now at $\lambda = \lambda_0$, but otherwise under the same conditions. (A one-body version of the theorem at zero energy is given in [Sk4].) Recall for comparison the quantity $\mathcal{E}_{-s_0,0}^{\mathcal{G}}$ of Lemma 4.10,

$$\mathcal{E}_{-s_0,0}^{\mathcal{G}} = \{ u \in \mathcal{B}_{s_0,0}^* \mid (H - \lambda_0)u = 0, \quad \Pi' u \in \mathcal{B}_0^* \}.$$

Let

$$\check{B}_{\rho} = r^{\rho/4} \check{B} r^{\rho/4}; \quad r = r_R, \, \check{B} = B_R, \, R = R_0,$$

recalling that $R_0 \ge 1$ is chosen in agreement with our version of the Mourre estimate at λ_0 (as done in Subsection 3.3.2).

Theorem 6.3 (Sommerfeld's theorem at threshold). There exists $\sigma > 0$ such that for any real function $\chi(\cdot < \sigma) \in C^{\infty}(\mathbb{R})$, which is supported in $(-\infty, \sigma)$ and whose derivative has compact support, and for any $\psi \in \mathcal{B}_{s_0}$, the function $\phi = \phi^{\pm} = R(\lambda_0 \pm i0)\psi \in \mathcal{B}^*_{s_0}$ obeys

$$\Pi' \phi \in \mathcal{B}^*,$$

$$\chi(\pm \check{B} < \sigma) \Pi' \phi \in \mathcal{B}_0^*,$$

$$\chi(\pm \check{B}_\rho < \sigma) \Pi \phi \in \mathcal{B}_{s_0,0}^*,$$

$$(H - \lambda_0) \phi = \psi.$$
(6.10)

Conversely suppose that (for either 'plus' or 'minus') $\phi \in \mathcal{B}_{s_0}^*$ fulfills (6.10) for some $\sigma > 0$, for all functions $\chi(\cdot < \sigma)$ of this type and for a given $\psi \in \mathcal{B}_{s_0}$, then $\phi = R(\lambda_0 \pm i0)\psi$ (with the same sign).

Proof. We shall only consider the case of '+'. By Proposition 3.16, for some $\sigma > 0$

$$\chi(\check{B} < \sigma)\check{R}(\lambda_0 + \mathrm{i}0)\Pi'\psi \in \mathcal{B}_0^*.$$
(6.11)

(This is for $\lambda_0 > \Sigma_2$; if $\lambda_0 = \Sigma_2$ the statement is trivial.) Note that $\Pi' \psi \in \mathcal{B}$ and that (6.11) holds with \check{B} replaced by $B_{\kappa} = \check{B}(\kappa^2 \check{B}^2 + 1)^{-1}$ ($\kappa > 0$ small). Then (6.11) follows by using another function of the same type, say $\tilde{\chi}(\cdot < \sigma)$, such that

$$\chi(b < \sigma) = \chi(b < \sigma)\tilde{\chi}(b/(\kappa^2 b^2 + 1) < \sigma).$$

By using [FS, Theorems 4.1 and 4.2] one can show (here omitting the argument) that for all small $\sigma > 0$

$$\langle x_a \rangle^{t-s_0} \chi \big(\check{B}_{\rho} < \sigma \big) Sr^+_{\lambda_0} \langle x_a \rangle^{-t-s_0-\epsilon} f \in L^2; f \in L^2, \, t, \epsilon > 0.$$
 (6.12)

We use (6.7) writing

$$\phi^{+} = R(\lambda_{0} + i0)\psi = S\phi_{a}^{+}(\lambda_{0}) + \breve{R}(\lambda_{0} + i0)\psi_{a}^{+}(\lambda_{0}).$$
(6.13)

Now (6.10) follows by using (6.11) and (6.12) to treat the second and the first terms of (6.13), respectively.

To show the second assertion (the uniqueness part) note that $\phi^+ = R(\lambda_0 + i0)\psi$ is a particular solution of this problem. Whence we may assume that $\psi = 0$. Due to Theorems 4.2 and 4.12 it suffices to show that $\phi \in \mathcal{E}_{-s_0,0}^{\mathcal{G}}$ (here we use the same notation if $\lambda_0 = \Sigma_2$), and whence that

$$\phi \in \mathcal{B}^*_{s_0,0}, \quad \Pi' \phi \in \mathcal{B}^*_0. \tag{6.14}$$

Introduce a smooth quadratic partition of unity $1 = \chi(\cdot < \sigma)^2 + \chi(\cdot > \sigma)^2$ such that $\chi(b < \sigma) = 1$ for $b < \sigma/2$. Let for $R \ge 1$ the function $\chi_R = \chi_R(r)$ be given by (1.29). Abbreviating $\theta_R = \sqrt{-\chi'_R}$ we can then estimate

$$-\langle \chi'_R(r) \rangle_{\Pi'\phi} - \langle r^{-\rho/2} \chi'_R(r) \rangle_{\Pi\phi}$$

$$\leq \frac{2}{\sigma} (s_R + t_R) + v_R;$$

$$s_R = \langle \check{B} \rangle_{\chi(\check{B} > \sigma)\theta_R \Pi'\phi},$$

$$t_R = \langle \check{B}_\rho \rangle_{\chi(\check{B}_\rho > \sigma)r^{-\rho/4}\theta_R \Pi\phi},$$

where $v_R \to 0$ for $R \to \infty$. Next we write

$$t_R = \tilde{t}_R + \tilde{v}_R;$$

$$\tilde{t}_R = \langle \breve{B} \rangle_{\chi(\breve{B}_\rho > \sigma)\theta_R \Pi \phi};$$

and note that $\tilde{v}_R = \mathcal{O}(R^{\rho/2-1}) \to 0$ for $R \to \infty$. After further commutation we (should) obtain that

$$-\langle \chi'_{R}(r) \rangle_{\Pi'\phi} - \langle r^{-\rho/2} \chi'_{R}(r) \rangle_{\Pi\phi}$$

$$\leq \frac{2}{\sigma} (\langle \breve{B} \rangle_{\theta_{R}\Pi'\phi} + \langle \breve{B} \rangle_{\theta_{R}\Pi\phi}) + o(R^{0})$$

$$= -\sigma^{-1} \langle i[H, \chi_{R}] \rangle_{\phi} + o(R^{0}),$$

$$= 0 + o(R^{0}),$$

yielding (6.14) and therefore the uniqueness part. However we need to argue for the validity of the above estimates. In the second step we used that

$$\operatorname{Re}\Pi\theta_R\check{B}\theta_R\Pi' = \mathcal{O}(R^{-2}),$$

cf. (4.46). In the first step we used that

$$s_R - \langle \check{B} \rangle_{\theta_R \Pi' \phi} = - \langle \check{B} \rangle_{\chi(\check{B} < \sigma) \theta_R \Pi' \phi} = o(R^0),$$

$$\tilde{t}_R - \langle \check{B} \rangle_{\theta_R \Pi \phi} = o(R^0).$$

The first bound is easy since $\check{B}\Pi'\phi \in \mathcal{B}^*$ (the latter seen by an energy bound).

To get the second bound it suffices to show that also $\check{B}_{\rho}\Pi\phi \in \mathcal{B}^*_{s_0}$. Let us first note that due to the assumption $\chi(\check{B} < \sigma)\Pi'\phi \in \mathcal{B}^*_0$ we can verbatim use Step V of the proof of Lemma 4.10 to conclude that $E^+_{\mathcal{H}}(\lambda_0)f = 0$, $f = T^*\phi \in \mathcal{B}^*_{s_0}(\mathbf{X}_a)$. Next we decompose

$$\check{B}_{\rho}\Pi\phi = \check{B}_{\rho}ST^{*}\phi = S\check{B}_{a,\rho}f + \hat{\phi},
\check{B}_{a,\rho} = r_{a}^{\rho/4-1/2}\operatorname{Re}(x_{a} \cdot p_{a})r_{a}^{\rho/4-1/2}; \quad r_{a} = r(x_{a}).$$
(6.15)

By the arguments of the proof of Lemma 3.4 indeed $\hat{\phi} \in \mathcal{B}_{s_0}^*$, so it remains only to show that $\check{B}_{a,\rho}f \in \mathcal{B}_{s_0}^*(\mathbf{X}_a)$. Noting that the operator $\check{B}_{a,\rho}$ has symbol

$$\check{b}_{a,\rho} \in S(s,g); \quad s = \langle \xi/g_a \rangle^2, \quad g_a = \sqrt{-w}, \quad g = \langle x \rangle^{-2} \mathrm{d}x^2 + g_a^{-2} \mathrm{d}\xi^2, \quad (6.16a)$$

it suffices in turn to show that $g_a^{-1} \operatorname{Op}^{\mathsf{w}}(s) g_a f \in \mathcal{B}^*_{s_0}(\mathbf{X}_a)$. As in [DS1, (4.15)]

$$Op^{w}(s) - g_{a}^{-1}hg_{a}^{-1} - 2 \in S(\langle x \rangle^{\rho-2}, g) \subset S(1, g).$$
(6.16b)

By writing $h = (h + v_{\lambda_0}^+) - v_{\lambda_0}^+$ we end up with bounding $-g_a^{-2}v_{\lambda_0}^+ f$, which clearly belongs to $\mathcal{B}^*_{s_0}(\mathbf{X}_a)$. Whence $\check{B}_{a,\rho}f \in \mathcal{B}^*_{s_0}(\mathbf{X}_a)$ is proven.

Corollary 6.4. For any $\psi \in \mathcal{B}$ the function $\phi = R(\lambda_0 \pm i0)\psi$ obeys the bounds

$$\check{B}_{\rho}\Pi\phi, \ g_a^{-1}\check{B}_{\rho}g_a\Pi\phi\in\mathcal{B}_{s_0}^*.$$
(6.17)

Proof. We substitute (6.8) and use (6.15) and (6.16b). Note that $g_a^{-2}hT^*\phi \in \mathcal{B}^*_{s_0}$ since $2s_0 > \rho$.

6.1.2 Elastic part of the scattering matrix at λ_0

We will to a large degree use [DS1]. We recall from (2.17) that S is given in terms of cluster bound states $\varphi_1, \ldots, \varphi_{m_a}, m_a = \dim \ker(H^a - \lambda_0), a = a_0$. The quantity $\alpha = \alpha_j = (a, \lambda_0, \varphi_j), j \leq m_a$, is referred to as a *channel*.

Let us for convenience here assume $m_a = 1$ and denote φ_1 by φ_α (see Subsecton 6.1.3 for an example where $m_a = 2$). To make contact to [DS1] it is convenient to change notation: Recall that up to a polynomially decreasing potential $w(x_a) \approx \langle I_a^{(1)}(\cdot + x_a) \rangle_{\varphi^a}$. Let us now assume that

$$w = V_1 + V_2, (6.18)$$

where V_1 and V_2 fulfill the following conditions of [DS1]. (For Coulomb systems one can take $V_1(x) = -\gamma r^{-1}$ for $r := |x| \ge 1$ and $V_2 = \mathcal{O}(r^{-2})$.)

Let $n = \dim \mathbf{X}_a$.

Condition 6.5. The function w can be written as a sum of two real-valued measurable functions, $w = V_1 + V_2$, such that: For some $\rho \in (0, 2)$ we have

(1) V_1 is a smooth negative function that only depends on the radial variable r in the region $r \ge 1$ (that is $V_1(x) = V_1(r)$ for $r = |x_a| \ge 1$). There exists $\epsilon_1 > 0$ such that

$$V_1(r) \le -\epsilon_1 r^{-\rho}, \ r \ge 1.$$

(2) For all $\gamma \in \mathbb{N}_0^n$ there exists $C_{\gamma} > 0$ such that

$$\langle x \rangle^{\rho+|\gamma|} |\partial^{\gamma} V_1(x)| \le C_{\gamma}.$$

(3) There exists $\tilde{\epsilon}_1 > 0$ such that

$$rV_1'(r) \le -(2 - \tilde{\epsilon}_1)V_1(r), \ r \ge 1.$$
 (6.19)

(4) $V_2 = V_2(x)$ is smooth and there exists $\epsilon_2 > 0$ such that for all $\gamma \in \mathbb{N}_0^n$

$$\langle x \rangle^{\rho + \epsilon_2 + |\gamma|} |\partial^{\gamma} V_2(x)| \le C_{\gamma}.$$

The following condition will be needed (and imposed) only in the case $V_2 \neq 0$. **Condition 6.6.** Let V_1 be given as in Condition 6.5 and $\alpha := \frac{2}{2+\rho}$. There exists $\bar{\epsilon}_1 > \max(0, 1 - \alpha(\rho + 2\epsilon_2))$ such that

$$\limsup_{r \to \infty} r^{-1} V_1'(r) \left(\int_1^r (-V_1(s))^{-\frac{1}{2}} \mathrm{d}s \right)^2 < 2^{-1} (1 - \bar{\epsilon}_1^2),$$
$$\limsup_{r \to \infty} V_1''(r) \left(\int_1^r (-V_1(s))^{-\frac{1}{2}} \mathrm{d}s \right)^2 < 2^{-1} (1 - \bar{\epsilon}_1^2).$$

6.1.2.1 Scattering for the one-body problem at zero energy, [DS1]

We review a number of results from [DS1] valid under Conditions 6.5 and 6.6. (For a different approach to one-body scattering theory, see [Is1].) Recall that for any $\omega \in \mathbb{S}^{n-1}$, $\lambda \in [0, \infty)$ and x from an appropriate outgoing/incoming region there exists a solution to the system of equations

$$\ddot{y}(t) = -2\nabla w(y(t)),$$

$$\lambda = \frac{1}{4}\dot{y}(t)^{2} + w(y(t)),$$

$$y(\pm 1) = x,$$

$$\omega = \pm \lim_{t \to \pm \infty} y(t)/|y(t)|.$$
(6.20)

One obtains a family $y^{\pm}(t, x, \omega, \lambda)$ of solutions depending regularily (at least continuously) on parameters. Moreover all 'scattering orbits' are of this form. Using these solutions one can construct a solution $\phi^{\pm}(x, \omega, \lambda)$ to the eikonal equation

$$\left(\nabla_x \phi^{\pm}(x,\omega,\lambda)\right)^2 + w(x) = \lambda \tag{6.21}$$

satisfying $\nabla_x \phi^{\pm}(x, \omega, \lambda) = \frac{1}{2} \dot{y}(\pm 1, x, \omega, \lambda).$

For $R \ge 1$ and $\sigma \in (0, 2]$

$$\Gamma^{+}_{R,\sigma}(\omega) := \{ y \in \mathbb{R}^{n} \mid y \cdot \omega \ge (1-\sigma)|y|, \ |y| \ge R \}; \ \omega \in \mathbb{S}^{n-1},$$

$$\Gamma^{+}_{R,\sigma} := \{ (y,\omega) \in \mathbb{R}^{n} \times \mathbb{S}^{n-1} \mid y \in \Gamma^{+}_{R,\sigma}(\omega) \}.$$
(6.22)

Lemma 6.7. There exist $R_0 \geq 1$ and $\sigma_0 \in (0,2)$ such that for all $R \geq R_0$ and for all positive $\sigma \leq \sigma_0$ the system (6.20) is solved for all data $(x,\omega) \in \Gamma^+_{R,\sigma}$ and $\lambda \geq 0$ by a unique function $y^+(t, x, \omega, \lambda)$, $t \geq 1$, such that $y^+(t, x, \omega, \lambda) \in \Gamma^+_{R,\sigma}(\omega)$ for all $t \geq 1$. Define a vector field $F^+(x, \omega, \lambda)$ on $\Gamma^+_{R_0,\sigma_0}(\omega)$ by

$$F^{+}(x,\omega,\lambda) = \frac{1}{2}\dot{y}^{+}(t=1,x,\omega,\lambda).$$
 (6.23)

Then

$$\operatorname{rot}_x F^+(x,\omega,\lambda) = 0.$$

We define $\phi^+(x,\omega,\lambda)$ at $(x,\omega,\lambda) \in \Gamma^+_{R_0,\sigma_0} \times [0,\infty)$ by requiring $\nabla_x \phi^+ = F^+$ and $\phi^+(R_0\omega,\omega,\lambda) = \sqrt{\lambda}R_0$. We let

$$\phi^{-}(x,\omega,\lambda) := -\phi^{+}(x,-\omega,\lambda) \text{ for } x \in \Gamma^{-}_{R_{0},\sigma_{0}}(\omega) := \Gamma^{+}_{R_{0},\sigma_{0}}(-\omega).$$

For $\xi \neq 0$ we write $\xi = \sqrt{\lambda}\omega, \, \omega \in \mathbb{S}^{n-1}$, and then

$$\phi^{\pm}(x,\xi) = \phi^{\pm}(x,\omega,\lambda); \quad (x,\omega) \in \Gamma^{\pm}_{R_0,\sigma_0}.$$

We are motivated to write, slightly abusely, $(x,\xi) \in \Gamma^{\pm}_{R_0,\sigma_0}$ instead of $(x,\omega,\lambda) \in$ $\Gamma^{\pm}_{R_0,\sigma_0} \times (0,\infty)$ (in fact even for the case $\xi = 0$). Fixing $0 < \sigma < \sigma' < \sigma_0$ we introduce a smoothed out characteristic function

$$\chi_{\sigma,\sigma'}(t) = \begin{cases} 1, & \text{for } t \ge 1 - \sigma, \\ 0, & \text{for } t \le 1 - \sigma'. \end{cases}$$
(6.24a)

Next define, in terms of (6.24a) and the function $\bar{\chi}_R = 1 - \chi_R$ of (1.29),

$$a_0^{\pm}(x,\xi) := \chi_{\sigma,\sigma'}(\pm \hat{x} \cdot \hat{\xi}) \bar{\chi}_{R_0}(|x|); \quad \hat{z} = z/|z|$$

We introduce then a Fourier integral operator J_0^+ on $L^2(\mathbb{R}^n)$ by

$$(J_0^{\pm} f)(x) = (2\pi)^{-n/2} \int e^{i\phi^{\pm}(x,\xi)} a_0^{\pm}(x,\xi) \hat{f}(\xi) d\xi, \qquad (6.24b)$$

where

$$\hat{f}(\xi) := (2\pi)^{-n/2} \int e^{-ix \cdot \xi} f(x) dx$$

denotes the Fourier transform of f.

The WKB method suggests to approximate the wave operator by a Fourier integral operator J^+ on $L^2(\mathbb{R}^d)$ of the form

$$(J^{+}f)(x) = (2\pi)^{-n/2} \int e^{i\phi^{+}(x,\xi)} a^{+}(x,\xi) \hat{f}(\xi) d\xi, \qquad (6.25a)$$

where the symbol $a^+(x,\xi)$ is supported in $\Gamma^+_{R_0,\sigma_0}$ and constructed by an iterative procedure (partly recalled in Subsection 6.1.2.4) attempting to make the difference $T^+ := \mathrm{i}(hJ^+ - J^+p^2)$ small in $\Gamma^+_{R_0,\sigma}$. We have

$$(T^+f)(x) = (2\pi)^{-n/2} \int e^{i\phi^+(x,\xi)} t^+(x,\xi) \hat{f}(\xi) d\xi, \qquad (6.25b)$$

where

$$t^{+}(x,\xi) = \left((2\nabla_{x}\phi^{+}(x,\xi)) \cdot \nabla_{x} + (\triangle_{x}\phi^{+}(x,\xi)) \right) a^{+}(x,\xi) - i\triangle_{x}a^{+}(x,\xi).$$
(6.25c)

The symbols $a^+(x,\xi)$ and $a_0^+(x,\xi)$ coincide to leading order away from $\xi = 0$, more precisely

$$a^{+}(x,\xi) \approx \left(\det \nabla_{\xi} \nabla_{x} \phi^{+}(x,\xi)\right)^{1/2} a^{+}_{0}(x,\xi) = e^{\zeta^{+}(x,\xi)} a^{+}_{0}(x,\xi), \qquad (6.25d)$$

and $a^+(x,\xi)$ should be thought of as an 'improvement' of the right-hand side. For details of construction, see [DS1, Section 5]. However, since we are partly going to mimic this construction in Subsection 6.1.2.4, let us here recall that the equation

$$\left(\left(2\nabla_x \phi^+(x,\xi) \right) \cdot \nabla_x + \left(\triangle_x \phi^+(x,\xi) \right) \right) e^{\zeta^+(x,\xi)} = 0, \tag{6.25e}$$

takes (6.25c) onto the form

$$t^{+} = e^{\zeta^{+}} ((2\nabla_{x}\phi^{+}) \cdot \nabla_{x} - iA^{+})b^{+}, \quad b^{+} = e^{-\zeta^{+}}a^{+},$$

$$A^{+} = \Delta + 2(\nabla\zeta^{+}) \cdot \nabla + (\Delta\zeta^{+}) + (\nabla\zeta^{+})^{2}.$$
(6.25f)

Similar to (6.25a)–(6.25c) we introduce a Fourier integral operator J^- and T^- as

$$(J^{-}f)(x) = (2\pi)^{-n/2} \int e^{i\phi^{-}(x,\xi)} a^{-}(x,\xi) \hat{f}(\xi) d\xi,$$

$$(T^{-}f)(x) = (2\pi)^{-n/2} \int e^{i\phi^{-}(x,\xi)} t^{-}(x,\xi) \hat{f}(\xi) d\xi.$$
(6.26)

For all $\tau \in L^2(\mathbb{S}^{n-1})$ we introduce

$$(J^{\pm}(\lambda)\tau)(x) = (2\pi)^{-n/2} \int e^{i\phi^{\pm}(x,\omega,\lambda)} \tilde{a}^{\pm}(x,\omega,\lambda)\tau(\omega)d\omega,$$

$$(T^{\pm}(\lambda)\tau)(x) = (2\pi)^{-n/2} \int e^{i\phi^{\pm}(x,\omega,\lambda)} \tilde{t}^{\pm}(x,\omega,\lambda)\tau(\omega)d\omega,$$
(6.27)

where

$$\tilde{a}^{\pm}(x,\omega,\lambda) = \frac{\lambda^{(n-2)/4}}{\sqrt{2}} a^{\pm}(x,\sqrt{\lambda}\omega),$$

$$\tilde{t}^{\pm}(x,\omega,\lambda) = \frac{\lambda^{(n-2)/4}}{\sqrt{2}} t^{\pm}(x,\sqrt{\lambda}\omega).$$

The functions \tilde{a}^{\pm} and \tilde{t}^{\pm} are *continuous* in $(x, \omega, \lambda) \in \mathbb{R}^d \times \mathbb{S}^{n-1} \times [0, \infty)$, and therefore we can *define* $J^{\pm}(\lambda)$ and $T^{\pm}(\lambda)$ at $\lambda = 0$ by the expressions (6.27). These properties hinge on [DS1, Proposition 5.3] stating properties of the function

$$\tilde{\zeta}^+(x,\omega,\lambda) = \zeta^+(x,\sqrt{\lambda}\omega) - \ln \lambda^{(2-n)/4}; \quad \lambda > 0$$

In particular it follows that there exist locally uniform limits (along with derivatives)

$$\tilde{\zeta}^+(x,\omega,0) = \lim_{\lambda \to 0_+} \tilde{\zeta}^+(x,\omega,\lambda).$$
(6.28)

It will be convenient to use a splitting $T^{\pm}(\lambda) = T^{\pm}_{\rm bd}(\lambda) + T^{\pm}_{\rm pr}(\lambda)$ in agreement with a certain decomposition of $\tilde{t}^{\pm}(x, \sqrt{\lambda}\omega)$, see Lemma 6.8 and references given before the lemma. There are *wave operators*

$$W^{\pm}f = \lim_{t \to \pm \infty} \mathrm{e}^{\mathrm{i}th} J_0^{\pm} \mathrm{e}^{-\mathrm{i}tp^2} f = \lim_{t \to \pm \infty} \mathrm{e}^{\mathrm{i}th} J^{\pm} \mathrm{e}^{-\mathrm{i}tp^2} f; \ \hat{f} \in C_{\mathrm{c}}^{\infty}(\mathbb{R}^n \setminus \{0\}).$$
(6.29)

The two operators W^{\pm} extend isometrically on $L^2(\mathbb{R}^d)$ with extensions satisfying $hW^{\pm} = W^{\pm}p^2$. Moreover,

$$0 = \lim_{t \to \mp \infty} \mathrm{e}^{\mathrm{i}th} J_0^{\pm} \mathrm{e}^{-\mathrm{i}tp^2} f = \lim_{t \to \mp \infty} \mathrm{e}^{\mathrm{i}th} J^{\pm} \mathrm{e}^{-\mathrm{i}tp^2} f; \ \hat{f} \in C_{\mathrm{c}}^{\infty}(\mathbb{R}^n \setminus \{0\}).$$
(6.30)

For $\rho \in (1/2, 2)$ we may write $W^{\pm} = W^{\pm}_{dol} e^{i\psi^{\pm}_{dol}(p)}$ in terms of the familiar Dollard wave operators [Do] (cf. (6.122) in Section 6.4) and explicit real momentumdepending phase factors ψ_{dol}^{\pm} , see [DS1, Theorem 6.15].

Let Δ_{ω} denote the Laplace-Beltrami operator on the sphere \mathbb{S}^{n-1} . For $k \in \mathbb{R}$ we define the Sobolev spaces on the sphere $H^k(\mathbb{S}^{n-1}) = (1 - \Delta_{\omega})^{-k/2} L^2(\mathbb{S}^{n-1})$. Let $\mathcal{L}_{s}^{k} = \mathcal{L}_{s}^{k}(\mathbb{R}^{n}) = \mathcal{L}(H^{k}(\mathbb{S}^{n-1}), L_{s}^{2}(\mathbb{R}^{n})) \text{ and } \mathcal{L}^{k} = \mathcal{L}_{0}^{k} \text{ for any } k, s \in \mathbb{R}.$ For $\lambda > 0$ we introduce the restricted Fourier transform $\mathcal{F}_{0}(\lambda)$ as

$$\mathcal{F}_0(\lambda)f(\omega) = \frac{\lambda^{(n-2)/4}}{\sqrt{2}}\hat{f}(\sqrt{\lambda}\omega).$$
(6.31)

Let $s > \frac{1}{2}$ and $k \ge 0$. Note that $\mathcal{F}_0(\lambda) \in \mathcal{L}(L^2_{s+k}(\mathbb{R}^n), H^k(\mathbb{S}^{n-1}))$ with a continuous dependence on $\lambda > 0$. Likewise, $\mathcal{F}_0(\lambda)^* \in \mathcal{L}^{-k}_{-s-k}$ with a continuous dependence on $\lambda > 0$. Note also that the operator

$$\int_{\mathbb{R}_{+}}^{\oplus} \mathcal{F}_{0}(\lambda) \,\mathrm{d}\lambda : L^{2}(\mathbb{R}^{n}) \to \int_{\mathbb{R}_{+}}^{\oplus} L^{2}(\mathbb{S}^{n-1}) \,\mathrm{d}\lambda$$
(6.32)

is unitary, and consequently that it diagonalizes the operator p^2 . Formally, we have $J^{\pm}(\lambda) = J^{\pm} \mathcal{F}_0(\lambda)^*$ and $T^{\pm}(\lambda) = T^{\pm} \mathcal{F}_0(\lambda)^*$. The formal identity $W^{\pm}(\lambda) =$ $W^{\pm}\mathcal{F}_0(\lambda)^* = (J^{\pm} + \mathrm{i}r(\lambda \mp \mathrm{i}0)T^{\pm})\mathcal{F}_0(\lambda)^*$ leads us then to consider the wave matrices

$$W^{\pm}(\lambda) := J^{\pm}(\lambda) + \mathrm{i}r(\lambda \mp \mathrm{i}0)T^{\pm}(\lambda), \qquad (6.33)$$

which in fact belong to \mathcal{L}_{-s}^{-k} for any $k \geq 0$ for a suitable $s = s(k) > s_0$. In this space $W^{\pm}(\lambda)$ have continuous dependence of $\lambda \ge 0$ (including $\lambda = 0!$).

The scattering operator commutes with p^2 , which is diagonalized by the direct integral mapping (6.32). Because of that the general theory of decomposable operators yields a measurable family $\mathbb{R}_+ \ni \lambda \mapsto S(\lambda)$ with the scattering matrix $S(\lambda)$ being a unitary operator on $L^2(\mathbb{S}^{n-1})$ for almost all λ , and such that in terms of the mapping (6.32)

$$S \simeq \int_{\mathbb{R}_+}^{\oplus} S(\lambda) \, \mathrm{d}\lambda.$$
 (6.34)

A main result of [DS1] reads, that for $\lambda \geq 0$ the scattering matrix

$$S(\lambda) = -2\pi J^+(\lambda)^* T^-(\lambda) + 2\pi i T^+(\lambda)^* r(\lambda + i0) T^-(\lambda),$$

$$= -2\pi W^+(\lambda)^* T^-(\lambda),$$
 (6.35)

defining a unitary operator $L^2(\mathbb{S}^{n-1})$ with a strongly continuously dependence on $\lambda \geq 0$. Moreover (6.34) is true, and

$$\forall k \in \mathbb{R} \,\forall \epsilon > 0: \quad S(\lambda) \in \mathcal{L}(H^k(\mathbb{S}^{n-1}), H^{k-\epsilon}(\mathbb{S}^{n-1})), \tag{6.36}$$

depending norm-continuously on $\lambda \geq 0$. Hence in particular $S(\lambda)$ maps $C^{\infty}(\mathbb{S}^{n-1})$ into itself.

Another main result of [DS1] adopted to the setting discussed here (in particular not including a certain singular term V_3) is the following result:

Suppose in addition to Conditions 6.5 and 6.6 that $V_1(r) = -\gamma r^{-\rho}$ for $r \ge 1$. Then the kernel $S(0)(\omega, \omega')$ is smooth outside the set $\{(\omega, \omega') \mid \omega \cdot \omega' = \cos \frac{\rho}{2-\rho}\pi\}$.

6.1.2.2 Elastic scattering for the *N*-body problem at λ_0

Using the constructions J_0^{\pm} in the previous subsection for the potential $w = V_1 + V_2$ in the variable x_a and recalling $-E_{\mathcal{H}}(\lambda \pm i0) = h + v_{\lambda}^{\pm} + \lambda_0 - \lambda$, $h = p^2 + w$, we introduce wave operators

$$\lim_{t \to \pm \infty} e^{itH} (1 \otimes J_0^{\pm}) e^{-itH_a} Sf = \lim_{t \to \pm \infty} e^{itH} (1 \otimes J_0^{\pm}) S e^{-it(p_a^2 + \lambda_0)} f;$$

$$\hat{f} \in C_c^{\infty} (\mathbf{X}_a \setminus \{0\})^{m_a}.$$
(6.37)

We consider a channel $\alpha = (a, \lambda_0, \varphi_\alpha)$ (recall that this means that $H^a \varphi_\alpha = \lambda_0 \varphi_\alpha$ with $\|\varphi_\alpha\| = 1$) assuming for simplicity from this point that

$$m_a = \dim \ker(H^a - \lambda_0) = 1 \tag{6.38}$$

(making φ_{α} essentially unique). With this assumption the above limit is nothing but the *channel wave operator*

$$W_{\alpha}^{\pm} f = \lim_{t \to \pm \infty} e^{itH} (1 \otimes J_0^{\pm}) \varphi_{\alpha} \otimes e^{-it(p_a^2 + \lambda_0)} f$$

=
$$\lim_{t \to \pm \infty} e^{itH} (\varphi_{\alpha} \otimes J_0^{\pm} e^{-it(p_a^2 + \lambda_0)} f); \quad \hat{f} \in C_c^{\infty}(\mathbf{X}_a \setminus \{0\}).$$
 (6.39)

This leads us to define

$$\begin{aligned} \mathcal{F}_{\lambda_0}(\lambda)f(\omega) &= \frac{(\lambda-\lambda_0)^{(n-2)/4}}{\sqrt{2}}\hat{f}\left(\sqrt{\lambda-\lambda_0}\,\omega\right),\\ (J_N^{\pm}(\lambda)\tau)(x) &= (2\pi)^{-n/2}\int \mathrm{e}^{\mathrm{i}\phi^{\pm}(x,\omega,\lambda-\lambda_0)}\tilde{a}^{\pm}(x,\omega,\lambda-\lambda_0)\tau(\omega)\mathrm{d}\omega,\\ (T_N^{\pm}(\lambda)\tau)(x) &= (2\pi)^{-n/2}\int \mathrm{e}^{\mathrm{i}\phi^{\pm}(x,\omega,\lambda-\lambda_0)}\tilde{t}^{\pm}(x,\omega,\lambda-\lambda_0)\tau(\omega)\mathrm{d}\omega,\\ J_{\alpha}^{\pm}f &= \varphi_{\alpha}\otimes J^{\pm}f; \quad f = f(x_a),\\ T_{\alpha}^{\pm} &= \mathrm{i}\left(HJ_{\alpha}^{\pm}-J_{\alpha}^{\pm}(p_a^2+\lambda_0)\right),\\ J_{\alpha}^{\pm}(\lambda)\tau &= \varphi_{\alpha}\otimes J_N^{\pm}(\lambda)\tau,\\ T_{\alpha}^{\pm}(\lambda)\tau &= \varphi_{\alpha}\otimes T_N^{\pm}(\lambda)\tau,\\ T_{\alpha}^{\pm}(\lambda) &= \mathrm{i}I_a^{(2)}J_{\alpha}^{\pm}(\lambda) + \mathrm{i}\left(I_a^{(1)}-w\right)J_{\alpha}^{\pm}(\lambda),\\ W_{\alpha}^{\pm}(\lambda) &= J_{\alpha}^{\pm}(\lambda) + \mathrm{i}R(\lambda\mp\mathrm{i}0)T_{\alpha}^{\pm}(\lambda),\\ S_{\alpha\alpha}(\lambda) &= -2\pi J_{\alpha}^{+}(\lambda)^*T_{\alpha}^{-}(\lambda) + 2\pi\mathrm{i}T_{\alpha}^{+}(\lambda)^*R(\lambda+\mathrm{i}0)T_{\alpha}^{-}(\lambda). \end{aligned}$$

Here formally

$$J_N^{\pm}(\lambda) = J^{\pm}(\lambda - \lambda_0) = J^{\pm} \mathcal{F}_{\lambda_0}(\lambda)^* \text{ and } T_N^{\pm}(\lambda) = T^{\pm}(\lambda - \lambda_0) = T^{\pm} \mathcal{F}_{\lambda_0}(\lambda)^*,$$

where J^{\pm} is the 'improvement' of J_0^{\pm} and $T^{\pm} = i(hJ^{\pm} - J^{\pm}p_a^2)$ as defined by (6.25a)–(6.25c) and (6.26).

The first term of $\check{T}^{\pm}_{\alpha}(\lambda)$ has arbitrary polynomial decay, for example stated precisely as $\check{T}^{\pm}_{\alpha}(\lambda) \in \mathcal{L}^k_s = \mathcal{L}^k_s(\mathbf{X}) := \mathcal{L}(H^k(\mathbb{S}^{n-1}), L^2_s(\mathbf{X}))$ for any $k, s \in \mathbb{R}$, and this is also the case for $\prod (I^{(1)}_a - w) J^{\pm}_{\alpha}(\lambda)$. But $\prod' (I^{(1)}_a - w) J^{\pm}_{\alpha}(\lambda) = \prod' I^{(1)}_a \prod J^{\pm}_{\alpha}(\lambda)$ is only one power better than w, more precisely it has the form $\mathcal{O}(\langle x \rangle^{-1-\rho}) J^{\pm}_{\alpha}(\lambda)$. Thus we can record

$$\Pi \check{T}^{\pm}_{\alpha}(\lambda) = \mathcal{O}(\langle x \rangle^{-\infty}), \quad \Pi' \check{T}^{\pm}_{\alpha}(\lambda) = \mathcal{O}(\langle x \rangle^{-1-\rho}) J^{\pm}_{\alpha}(\lambda).$$
(6.40)

A similar remark is due for the 'non-restricted' quantity \check{T}^{\pm}_{α} in the formula

$$T^{\pm}_{\alpha} = \varphi_{\alpha} \otimes T^{\pm} + \check{T}^{\pm}_{\alpha}; \quad \check{T}^{\pm}_{\alpha} = I^{(2)}_{a}J^{\pm}_{\alpha} + \left(I^{(1)}_{a} - w\right)J^{\pm}_{\alpha}.$$

Moreover, since for any f with $\hat{f} \in C_c^{\infty}(\mathbf{X}_a \setminus \{0\})$ the quantity $||T_{\alpha}^{\pm} e^{-it(p_a^2 + \lambda_0)}f||$ is integrable at $\pm \infty$, the Cook argument gives the existence of the wave operator W_{α}^{\pm} . Note that this integrability may be shown by a stationary phase argument (for example by using (6.43) and a version of Lemma 6.8 3)).

Using [DS1, Appendix A] the elastic part of the scattering matrix defined by (6.39) may be shown to be given by the expression $S_{\alpha\alpha}(\lambda)$, $\lambda > \lambda_0$, introduced above. We will study some properties of this operator, which is an operator on $\mathcal{L}(\mathbb{S}^{n-1})$ with norm at most one.

We state some basic properties of $J^{\pm}(\lambda)$ and $T^{\pm}(\lambda) = T^{\pm}_{bd}(\lambda) + T^{\pm}_{pr}(\lambda)$, see [DS1, (5.8), (5.16), (5.19) and Lemmas 6.8 and 6.9] and the proof of [DS1, Theorem 6.11] (which we adapt to the present problem). First we recall some notation. The most basic one is the function

$$g(r) = g_{\lambda} = \sqrt{\lambda - \lambda_0 - V_1(r)}, \quad \lambda \ge \lambda_0,$$

which roughly controls the momentum. Next we introduce the symbols

$$a(x,\xi) = \frac{\xi^2}{g_{\lambda}(|x|)^2}, \quad b(x,\xi) = \frac{\xi}{g_{\lambda}(|x|)} \cdot F(x), \tag{6.41}$$

where F is an arbitrary (henceforth fixed) vector field on \mathbb{R}^n extending $F(x) = \hat{x} = x/r$ for $r = |x| \ge 1$. Of course the symbols a and b also have λ -dependence, but for convenience this is here and henceforth omitted in the notation. Let $\tilde{\chi}_-, \tilde{\chi}_+ \in C^{\infty}$ be non-negative functions obeying $\tilde{\chi}_- + \tilde{\chi}_+ = 1$ and

$$\operatorname{supp} \tilde{\chi}_{-} \subseteq (-\infty, 1 - \bar{\sigma}], \tag{6.42a}$$

$$\operatorname{supp} \tilde{\chi}_+ \subseteq [1 - 2\bar{\sigma}, \infty), \tag{6.42b}$$

where the number $\bar{\sigma} > 0$ needs to be taken sufficiently small, depending on the parameter σ used in the previous subsection (see (6.24a)) and properties of the phase $\phi^{\pm}(x,\xi)$. Let $\chi_{-}, \chi_{+} \in C^{\infty}$ be non-negative functions obeying $\chi_{-} + \chi_{+} = 1$ and

$$\operatorname{supp} \chi_{-} \subseteq (-\infty, 2), \tag{6.42c}$$

$$\operatorname{supp} \chi_+ \subseteq (1, \infty). \tag{6.42d}$$

Introduce then symbols

$$\chi_{1} = \chi_{+}(a),$$

$$\chi_{2}^{\pm} = \chi_{-}(a)\tilde{\chi}_{-}(\pm b),$$

$$\chi_{3}^{\pm} = \chi_{-}(a)\tilde{\chi}_{+}(\pm b).$$
(6.42e)

These symbols belong to a class of (parameter-depending) pseudodifferential operators studied in [FS, DS1]. The 'Planck constant' for this class is $\langle x \rangle^{-1} g_{\lambda}(|x|)^{-1}$, in particular at most $\langle x \rangle^{\rho/2-1}$. Note the partition of unity in terms of corresponding (right-quantized) operators

$$1 = Op^{r}(\chi_{1}) + Op^{r}(\chi_{2}^{\pm}) + Op^{r}(\chi_{3}^{\pm}).$$
(6.43)

Recall $\mathcal{L}^k := \mathcal{L}(H^k(\mathbb{S}^{n-1}), L^2(\mathbb{R}^n)), k \in \mathbb{R}.$

Lemma 6.8. Let χ_1 , χ_2^{\pm} and χ_3^{\pm} be given by (6.42e).

1) For all $k \in [0, \infty)$ and $\epsilon > 0$,

$$(\langle x \rangle g_{\lambda})^{-k} \langle x \rangle^{-1/2-\epsilon} g_{\lambda}^{1/2} J_N^{\pm}(\lambda)$$
(6.44a)

is a continuous \mathcal{L}^{-k} -valued function of $\lambda \in [\lambda_0, \infty)$. With a bounding constant independent of $\lambda \geq \lambda_0$,

$$g_{\lambda}^{1/2}J_{N}^{\pm}(\lambda) \in \mathcal{L}(L^{2}(\mathbb{S}^{n-1}),\mathcal{B}_{1/2}^{*}(\mathbb{R}^{n})).$$

2) For all $k \in \mathbb{R}$ and $\epsilon > 0$,

$$(\langle x \rangle g_{\lambda})^{-k} \langle x \rangle^{1/2 - \epsilon} g_{\lambda}^{-1/2} \operatorname{Op}^{\mathsf{r}}(\chi_{2}^{\pm}) T_{\mathrm{bd}}^{\pm}(\lambda - \lambda_{0})$$
(6.44b)

is a continuous \mathcal{L}^{-k} -valued function of $\lambda \in [\lambda_0, \infty)$.

3) For all $k, m \in \mathbb{R}$,

$$\langle x \rangle^m \operatorname{Opr}(\chi_1 + \chi_3^{\pm}) T_{\mathrm{bd}}^{\pm}(\lambda - \lambda_0), \ \langle x \rangle^m T_{\mathrm{pr}}^{\pm}(\lambda - \lambda_0) \ and \ \langle x \rangle^m \operatorname{Opr}(\chi_1) J_N^{\pm}(\lambda) \ (6.44c)$$

are continuous \mathcal{L}^{-k} -valued function of $\lambda \in [\lambda_0, \infty)$.

This lemma will be used in combination with the following excerpts of [DS1, Proposition 4.1] (adapted to the present problem). Note that $\operatorname{Op}^{l}(\chi_{2}^{\pm}) = \operatorname{Op}^{r}(\chi_{2}^{\pm})^{*}$ is given by left-quantization.

Lemma 6.9. Let Λ denote any interval of the form $\Lambda = [\lambda_0, \lambda'_0]$, and let $r_{\lambda}^{\pm} = r(\lambda - \lambda_0 \pm i0)$ for $\lambda \in \Lambda$. Then the following bounds hold uniformly in $\lambda \in \Lambda$, and the corresponding $\mathcal{L}(L^2(\mathbb{R}^n))$ -valued functions are continuous.

1) For all $\epsilon > 0$ there exists C > 0 such that

$$\|\langle x \rangle^{-\epsilon-1/2} g_{\lambda}^{1/2} r_{\lambda}^{\pm} g_{\lambda}^{1/2} \langle x \rangle^{-\epsilon-1/2} \| \le C.$$
(6.45a)

2) For all $s \ge 0$ and $0 \le \epsilon < \epsilon'$ there exists C > 0

$$\|(\langle x\rangle g_{\lambda})^{s} \langle x\rangle^{\epsilon-1/2} g_{\lambda}^{1/2} \operatorname{Op}^{l}(\chi_{2}^{\pm}) r_{\lambda}^{\pm} g_{\lambda}^{1/2} \langle x\rangle^{-\epsilon'-1/2} (\langle x\rangle g_{\lambda})^{-s} \| \leq C.$$
(6.45b)

Now taking $\tau \in H^k(\mathbb{S}^{n-1})$ with a sufficiently big k (actually any k > 0 suffices), we will show by combining (6.7) with Lemmas 6.8 and 6.9 that $S_{\alpha\alpha}(\lambda)\tau$ is a welldefined element of $L^2(\mathbb{S}^{n-1})$, in fact with a continuous dependence of $\lambda \in I_{\delta}^+$.

We insert (6.7) into the formula

$$S_{\alpha\alpha}(\lambda)\tau = -2\pi J_{\alpha}^{+}(\lambda)^{*}T_{\alpha}^{-}(\lambda)\tau + 2\pi \mathrm{i}T_{\alpha}^{+}(\lambda)^{*}R(\lambda+\mathrm{i}0)T_{\alpha}^{-}(\lambda)\tau.$$
(6.46)

Ignoring the contribution from $\check{T}_{\alpha}^{-}(\lambda)$ (its contribution is a 'partial smoothing operator' as exemplified in the beginning of the proof of Theorem 6.10) we obtain by using Lemma 6.8 that the first term $-2\pi J_{\alpha}^{+}(\lambda)^{*}T_{\alpha}^{-}(\lambda)\tau \approx -2\pi J_{N}^{+}(\lambda)^{*}T_{N}^{-}(\lambda)\tau \in L^{2}(\mathbb{S}^{n-1}).$

For the second term in (6.46) we (again) ignore terms containing $\check{T}^{\pm}_{\alpha}(\lambda)$ and consider only $\varphi_{\alpha} \otimes T^{\pm}_{N}(\lambda)$ using (6.43) to write

$$T_N^{\pm}(\lambda) = \operatorname{Op}^{\mathsf{r}}(\chi_1 + \chi_3^{\pm}) T_N^{\pm}(\lambda) + \operatorname{Op}^{\mathsf{r}}(\chi_2^{\pm}) T_N^{\pm}(\lambda) = T_{1,3}^{\pm}(\lambda) + T_2^{\pm}(\lambda).$$

By Lemma $6.8\ 3$) the first term has strong decay, so let us consider the seemingly worse term given (up to a constant) by

$$\left(\varphi_{\alpha} \otimes T_2^+(\lambda)\right)^* R(\lambda + \mathrm{i0})\varphi_{\alpha} \otimes T_2^-(\lambda)$$

= $T_2^+(\lambda)^* S^* R(\lambda + \mathrm{i0}) ST_2^-(\lambda).$

The contribution from the 'leading terms', cf. (6.8), are

$$T_2^+(\lambda)^* r_\lambda^+ T_2^-(\lambda)$$
 and $T_2^+(\lambda)^* S^* \check{R}(\lambda + \mathrm{i0}) \Pi' S T_2^-(\lambda)$,

respectively. The second term vanishes. For the first term we use Lemma 6.9 2) for the case of '+' and see that indeed $T_2^+(\lambda)^* r_{\lambda}^+ T_2^-(\lambda) \tau \approx T_N^+(\lambda)^* r_{\lambda}^+ T_N^-(\lambda) \tau$ is a well-defined element of $L^2(\mathbb{S}^{n-1})$.

Of course there are other terms to consider, and for some of those also Lemma 6.9 1) is needed. We can check all other terms (cf. the proof of Theorem 6.10) and see that they are well-defined with a continuous dependence of λ . Furthermore we can write the formula for the action by $S_{\alpha\alpha}(\lambda)$ as

$$S_{\alpha\alpha}(\lambda)\tau = -2\pi W_{\alpha}^{+}(\lambda)^{*}T_{\alpha}^{-}(\lambda)\tau, \qquad (6.47)$$

and this is a continuous $L^2(\mathbb{S}^{n-1})$ -valued function of $\lambda \in I^+_{\delta}$ for $\tau \in H^k(\mathbb{S}^{n-1})$, k > 0.

Next we will examine the degree of regularity that is needed on τ , measured by the size of k. Let

$$S_w(\lambda - \lambda_0) = -2\pi J_N^+(\lambda)^* T_N^-(\lambda) + 2\pi \mathrm{i} T_N^+(\lambda)^* r(\lambda + \mathrm{i} 0) T_N^-(\lambda); \quad \lambda \ge \lambda_0$$

Note that $S_w(\cdot)$ is the scattering matrix for the one-body problem given by (6.35). We will examine the quantity

$$\check{S}_{\alpha\alpha}(\lambda)\tau = S_{\alpha\alpha}(\lambda)\tau - S_w(\lambda - \lambda_0)\tau.$$

By the above preliminary investigation the term $S_w(\cdot)$ is the 'leading term', and we know that k = 0 works for this term although $k = \epsilon$ for any $\epsilon > 0$ is neeeded for operator-continuity, see (6.36) and the discussion there. So we expect welldefinedness of $\check{S}_{\alpha\alpha}(\lambda)\tau$ and continuous dependence of λ with less regularity imposed on τ . In fact we can show that $k = -\epsilon$ for a computable $\epsilon > 0$ works for this term, even for operator-continuity. More generally we have the following result given in terms of any $k \geq 0$ satisfying one of the options

$$\begin{cases} k+1/2+(1/2-k)\rho/2 < 1/2+\rho/2, & k \in [0,1/2], \\ k+1/2 < 1/2+\rho/2, & k > 1/2, \end{cases}$$
(6.48)

or equivalently stated, one of the options

$$\begin{cases} 0 \le k \le 1/2 \text{ and } k < \rho(4 - 2\rho)^{-1}, \\ 1/2 < k < \rho/2, \quad \rho > 1. \end{cases}$$
(6.49)

Theorem 6.10. Suppose (6.38), i.e. $m_a = 1$. For any $k \ge 0$ obeying (6.49) the operator

$$\check{S}_{\alpha\alpha}(\lambda) = S_{\alpha\alpha}(\lambda) - S_w(\lambda - \lambda_0) \in \mathcal{L}\big(H^{-k}(\mathbb{S}^{n-1}), H^k(\mathbb{S}^{n-1})\big)$$

with a continuous dependence of $\lambda \in I_{\delta}^+$.

- **Remarks 6.11.** 1) The number $s = 1/2 + \rho/2$ on the right-hand side in (6.48) is the number appearing in (6.54). The left-hand side comes from estimating the factor $\langle x \rangle^{k+1/2} g^{k-\frac{1}{2}}$; it is uniformly bounded by the power $C \langle x \rangle^c$ with c given as the expression appearing to the left in (6.48). If we replace I_{δ}^+ by $I_{\delta}^+ \setminus \{\lambda_0\}$ in Theorem 6.10 any $k < \rho/2$ suffices.
 - 2) We assumed $m_a = 1$. However the interested reader may check that if $m_a > 1$, then Theorem 6.10 remains valid. Moreover the off-diagonal elements of the scattering matrix, denoted by $S_{\beta\alpha}(\lambda)$ with $\alpha \neq \beta$, fulfill the same assertion as the one for $\check{S}_{\alpha\alpha}(\lambda)$ in Theorem 6.10.
 - 3) Under the additional condition that $\lambda_0 = \Sigma_2$ one can show that $\check{S}_{\alpha\alpha}(\lambda)$ is bounded on any of the spaces $H^l(\mathbb{S}^{n-1}), l \in \mathbb{R}$, with a continuous dependence of $\lambda \in I_{\delta}^+$, in fact it is partially smoothing. More precisely one can show in this case that for all $l \in \mathbb{R}$ the operator $\check{S}_{\alpha\alpha}(\lambda) \in \mathcal{L}(H^{l-k}, H^{l+k})$ with a continuous dependence on $\lambda \in I_{\delta}^+$. This is for any $k \geq 0$ obeying (6.49), in fact for a computable bigger k using (6.54) below for $s = 1 + \rho$ (recall from the discussion after (6.5) that the latter boundedness condition is fulfilled for $\lambda_0 = \Sigma_2$). The proof consists of combining ideas from Subsection 6.1.2.4 with [DS1, Proposition 4.1], however we shall not elaborate. In particular since $S_w(\lambda - \lambda_0)$ has a similar property up a loss of an ' ϵ -smoothness', cf. [DS1, Theorem 7.2], we conclude (more precisely stated) that for all $l \in \mathbb{R}$ and $\epsilon > 0$ the operator $S_{\alpha\alpha}(\lambda) \in \mathcal{L}(H^l, H^{l-\epsilon})$ with a continuous dependence of $\lambda \in I_{\delta}^+$. Note that this implies that $S_{\alpha\alpha}(\lambda)\tau \in C^{\infty}(\mathbb{S}^{n-1})$ for $\lambda \in I_{\delta}^+$ and $\tau \in C^{\infty}(\mathbb{S}^{n-1})$.

Proof of Theorem 6.10. We need to treat the term $-2\pi J_{\alpha}^{+}(\lambda)^{*} \check{T}_{\alpha}^{-}(\lambda)$ left out when discussing $-2\pi J_{\alpha}^{+}(\lambda)^{*} T_{\alpha}^{-}(\lambda)$ above. (Note that $-2\pi J_{\alpha}^{+}(\lambda)^{*} (\varphi_{\alpha} \otimes T_{N}^{\pm}(\lambda))$ already is subtracted in the definition of $\check{S}_{\alpha\alpha}(\lambda)$.) Writing $J_{\alpha}^{+}(\lambda)^{*} \check{T}_{\alpha}^{-}(\lambda) = J_{\alpha}^{+}(\lambda)^{*} \Pi \check{T}_{\alpha}^{-}(\lambda)$ we can invoke (6.40).

The remaining terms of $\tilde{S}_{\alpha\alpha}(\lambda)$ fall into three disjoint groups according to whether there is a dependence of:

- **a)** $\breve{R}(\lambda + i0))$, and no dependence of $r_{\lambda}^{+}(1 + v_{\lambda}^{+}r_{\lambda}^{+})^{-1}$.
- **b)** $r_{\lambda}^{+}(1+v_{\lambda}^{+}r_{\lambda}^{+})^{-1}$, and no dependence of $\breve{R}(\lambda+i0)$).
- c) $\breve{R}(\lambda + i0)$ as well as a dependence of $r_{\lambda}^{+}(1 + v_{\lambda}^{+}r_{\lambda}^{+})^{-1}$.
- a). We need to treat

$$T^+_{\alpha}(\lambda)^* \breve{R}(\lambda + \mathrm{i}0) \Pi' T^-_{\alpha}(\lambda) = \breve{T}^+_{\alpha}(\lambda)^* \Pi' \breve{R}(\lambda + \mathrm{i}0) \Pi' \breve{T}^-_{\alpha}(\lambda).$$

Due to (6.40) we need to bound

$$\langle x \rangle^{-1/2-\rho+\epsilon} J^{\pm}_{\alpha}(\lambda) \in \mathcal{L}^{-k} = \mathcal{L}^{-k}(\mathbf{X}) \quad \text{for some } \epsilon > 0.$$
 (6.50)

By Lemma $6.8\ 1$) this requires

$$\begin{cases} k + (1/2 - k)\rho/2 < \rho, & k \in (0, 1/2], \\ k < \rho, & k > 1/2, \end{cases}$$
(6.51)

which is weaker than (6.49).

b). We need to treat

$$T_{\alpha}^{+}(\lambda)^{*}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}S^{*}T_{\alpha}^{-}(\lambda)-T_{N}^{+}(\lambda)^{*}r_{\lambda}^{+}T_{N}^{-}(\lambda)$$
$$=\left(T_{\alpha}^{+}(\lambda)^{*}Sr_{\lambda}^{+}S^{*}T_{\alpha}^{-}(\lambda)-T_{N}^{+}(\lambda)^{*}r_{\lambda}^{+}T_{N}^{-}(\lambda)\right)$$
$$-T_{\alpha}^{+}(\lambda)^{*}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}v_{\lambda}^{+}r_{\lambda}^{+}S^{*}T_{\alpha}^{-}(\lambda)$$

Due to (6.40) the first term is a smoothing operator, i.e. in $\mathcal{L}(H^{-l}(\mathbb{S}^{n-1}), H^{l}(\mathbb{S}^{n-1}))$ for any $l \in \mathbb{R}$. We shall frequently in the rest of the proof use

$$1 = \langle x \rangle^{-s} \langle x \rangle^{s}; \quad s = 1/2 + \rho/2.$$
 (6.52)

For the second term it suffices, due to (6.6) and (6.52), to bound

$$\langle x \rangle^{-s} r_{\lambda}^{\pm} S^* T_{\alpha}^{\pm}(\lambda) \in \mathcal{L}^{-k}.$$
 (6.53)

Here we used that

$$v_{\lambda}^{+} \in \mathcal{L}(L_{-s}^{2}, L_{s}^{2}). \tag{6.54}$$

(We could do better using (6.38), cf. the discussion after (6.5), but we prefer to use a method that more or less obviously generalizes to the case where $m_a > 1$, c.f. Remark 6.11 2).) To show (6.53) it suffices due to (6.40) and Lemma 6.8 3) to bound

$$\langle x \rangle^{-s} r_{\lambda}^{\mp} T_2^{\pm}(\lambda) \in \mathcal{L}^{-k},$$
(6.55)

or equivalently that

$$T_N^{\pm}(\lambda)^* \operatorname{Op}^{\mathfrak{l}}(\chi_2^{\pm}) r_{\lambda}^{\pm} \in \mathcal{L}(L_s^2, H^k(\mathbb{S}^{n-1})).$$
(6.56)

Due to Lemma 6.8 it suffices in turn to show that

$$(\langle x \rangle g_{\lambda})^{k} \langle x \rangle^{\epsilon - 1/2} g_{\lambda}^{1/2} \operatorname{Op}^{l}(\chi_{2}^{\pm}) r_{\lambda}^{\pm} \in \mathcal{L}(L_{s}^{2}, L^{2}) \text{ for some } \epsilon > 0.$$

$$(6.57)$$

We need to bound

$$(\langle x \rangle g_{\lambda})^{k} \langle x \rangle^{\epsilon - 1/2} g_{\lambda}^{1/2} \operatorname{Op}^{l}(\chi_{2}^{\pm}) r_{\lambda}^{\pm} g_{\lambda}^{1/2} \langle x \rangle^{-1/2 - 2\epsilon} (\langle x \rangle g_{\lambda})^{-k} \left(g_{\lambda}^{-1/2} \langle x \rangle^{1/2 + 2\epsilon} (\langle x \rangle g_{\lambda})^{k} \langle x \rangle^{-s} \right) \in \mathcal{L}(L^{2}).$$

Using (6.45b) we need to check that the last factor is bounded. This amounts to

$$\begin{cases} k+1/2+2\epsilon+(1/2-k)\rho/2 \le s, \quad k \in (0, 1/2], \\ k+1/2+2\epsilon \le s, \quad k > 1/2, \end{cases}$$
(6.58)

which indeed is fulfilled for small $\epsilon > 0$ thanks to (6.48).

c). There are four terms to be considered given (up to the common factor $2\pi i$) by:

$$S_{1} = -T_{\alpha}^{+}(\lambda)^{*}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}S^{*}I_{a}\breve{R}(\lambda+\mathrm{i0})\Pi'T_{\alpha}^{-}(\lambda),$$

$$S_{2} = T_{\alpha}^{+}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i0})\Pi'I_{a}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}S^{*}I_{a}\breve{R}(\lambda+\mathrm{i0})\Pi'T_{\alpha}^{-}(\lambda),$$

$$S_{3} = -T_{\alpha}^{+}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i0})\Pi'I_{a}Sr_{\lambda}^{+}S^{*}T_{\alpha}^{-}(\lambda),$$

$$S_{4} = T_{\alpha}^{+}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i0})\Pi'I_{a}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}v_{\lambda}^{+}r_{\lambda}^{+}S^{*}T_{\alpha}^{-}(\lambda).$$

For S_1 we insert the decomposition (6.52) to the right of the factor r_{λ}^+ (the left one). Suppose we can show that

$$\langle x \rangle^s S^* I_a \check{R}(\lambda \pm i0) \Pi' T^{\mp}_{\alpha}(\lambda) \in \mathcal{L}^{-k},$$

then we are done using (6.53) and (6.58). But due to (6.40) this term has the form

$$\mathcal{O}(\langle x \rangle^{s-1-\rho})\breve{R}(\lambda \pm i0)\Pi'\breve{T}^{\mp}_{\alpha}(\lambda) = \mathcal{O}(\langle x \rangle^{s-1-\rho})\breve{R}(\lambda \pm i0)\mathcal{O}(\langle x \rangle^{-1-\rho})J^{\mp}_{\alpha}(\lambda), \quad (6.59)$$

for which we can apply (6.50). The argument for S_3 is the same.

We treat S_2 by inserting (6.52) to the right as well as to the left of the factor r_{λ}^+ (the left one) and then use (6.59) and Lemma 6.9 1) (to bound $\langle x \rangle^{-s} r_{\lambda}^+ \langle x \rangle^{-s}$).

We treat S_4 by inserting (6.52) to the right of the far left factor r_{λ}^+ and to the left of the far right factor r_{λ}^+ . Then we invoke (6.53) and (6.54).

6.1.2.3 Elastic scattering at λ_0 , a 'geometric' approach

In the spirit of [DS1, Section 8] we will give a 'geometric' description of the operator $S_{\alpha\alpha}(\lambda)$ studied in Subsection 6.1.2.2. To keep the discussion short we shall consider the limiting case $\lambda = \lambda_0$ only. For analogue results for (all most all) non-threshold energies in a general N-body setting we refer to [Sk6].

We recall the following construction for the one-body problem, see [DS1, Proposition 5.6], which could be a basis for discussing $\lambda > \lambda_0$ also.

Lemma 6.12. There exist $R \ge R_0$ and $\tilde{\sigma} \in (0, \sigma_0]$ such that for all $x = |x|\hat{x} \in \mathbb{R}^n$ with $|x| \ge R$ and $\lambda \ge \lambda_0$ there exists a unique $\omega \in \mathbb{S}^{n-1}$ satisfying $\omega \cdot \hat{x} \ge 1 - \tilde{\sigma}$ (equivalently, $x \in \Gamma^+_{R,\tilde{\sigma}}(\omega)$) and $\partial_{\omega}\phi^+(x, \omega, \lambda - \lambda_0) = 0$. We introduce the notation $\omega^+_{\text{crt}} = \omega^+_{\text{crt}}(x, \lambda)$ for this vector. It is smooth in x, and for some $\check{\epsilon} = \check{\epsilon}(\rho, \bar{\epsilon}_1, \epsilon_2) > 0$

$$\partial_x^{\gamma}(\omega_{\mathrm{crt}}^+ - \hat{x}) = \mathcal{O}(|x|^{-\check{\epsilon} - |\gamma|}).$$

Let

$$\phi(x,\lambda) = \phi^+(x,\omega_{\rm crt}^+(x,\lambda),\lambda-\lambda_0). \tag{6.60}$$

This function solves the eikonal equation

$$(\partial_x \phi(x,\lambda))^2 + w(x) = \lambda - \lambda_0; \quad |x| \ge R \text{ and } \lambda \ge \lambda_0.$$

In the spherically symmetric case (viz. $V_2(x) = V_2(r)$) we have $\omega_{crt}^+ = \hat{x}$ and

$$\phi(x,\lambda) = \phi_{\rm sph}(x,\lambda) := \int_{R_0}^{|x|} \sqrt{\lambda - \lambda_0 - w(r)} dr + \sqrt{\lambda - \lambda_0} R_0.$$
(6.61)

With reference to Theorem 6.3 let $\chi(\cdot < \sigma)$ be one of the functions described there (with $\sigma > 0$ sufficiently small) but taken with the additional property that $\chi(\cdot < \sigma) = 1$ on $(-\infty, \sigma/2)$. Let \check{B} and \check{B}_{ρ} be given as in Theorem 6.3, and let

$$\mathcal{V}^+_{-s_0,\sigma}(\lambda_0) = \{ u \in \mathcal{B}^*_{s_0}(\mathbf{X}) \mid \Pi' u \in \mathcal{B}^*, \quad \chi(\check{B} < \sigma) \Pi' u \in \mathcal{B}^*_0, \quad (H - \lambda_0) u = 0 \}, \\ \mathcal{R}^+_{\sigma} = \{ u \in \mathcal{B}^*_{s_0,0}(\mathbf{X}) \mid \Pi' u \in \mathcal{B}^*, \quad \chi(\check{B} < \sigma) \Pi' u \in \mathcal{B}^*_0 \}.$$

We use the notation $\hat{y} = y/|y|$ for (nonzero) vectors $y \in \mathbf{X}_a$ as well as

$$u_{a}^{+}(y) = c_{n} g_{\lambda_{0}}^{-1/2}(|y|) |y|^{-\frac{n-1}{2}} e^{i\phi(y,\lambda_{0})}, \quad u_{a}^{-}(y) = \overline{u_{a}^{+}(y)}; \quad c_{n} = e^{i\pi\frac{1-n}{4}} (4\pi)^{-1/2},$$
$$v_{a}^{\pm}(y) = \pm \langle y \rangle^{\rho/2} g_{\lambda_{0}}(|y|) u_{a}^{\pm}(y).$$

Theorem 6.13. 1) The channel wave matrix

$$W^{-}_{\alpha}(\lambda_0): L^2(\mathbb{S}^{n-1}) \to \mathcal{V}^{+}_{-s_0,\sigma}(\lambda_0) \subset \mathcal{B}^*_{s_0}$$

is a well-defined bicontinuous isomorphism. (In particular the space $\mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$ does not depend on the small $\sigma > 0$.)

2) For all $\tau \in L^2(\mathbb{S}^{n-1})$ the vectors $\tau^+ = S_{\alpha\alpha}(\lambda_0)\tau$ and $\phi_{\tau}^+ = W_{\alpha}^-(\lambda_0)\tau$ are the unique vectors in $L^2(\mathbb{S}^{n-1})$ and $\mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$, respectively, fulfilling

$$\phi_{\tau}^{+}(x) - \varphi_{\alpha}(x^{a}) \left(u_{a}^{-}(x_{a})\tau(-\widehat{x_{a}}) + u_{a}^{+}(x^{a})\tau^{+}(\widehat{x_{a}}) \right) \in \mathcal{R}_{\sigma}^{+}.$$
(6.62a)

3) For all $\tau \in L^2(\mathbb{S}^{n-1})$, and with $\tau^+ = S_{\alpha\alpha}(\lambda_0)\tau$ and $\phi_{\tau}^+ = W_{\alpha}^-(\lambda_0)\tau$ as above,

$$\left(\check{B}_{\rho}\Pi\phi_{\tau}^{+}\right)(x) - \varphi_{\alpha}(x^{a})\left(v_{a}^{-}(x_{a})\tau(-\widehat{x_{a}}) + v_{a}^{+}(x^{a})\tau^{+}(\widehat{x_{a}})\right) \in \mathcal{B}_{s_{0},0}^{*}.$$
 (6.62b)

Proof. \underline{I} . We insert (6.7) into the definition

$$W_{\alpha}^{-}(\lambda) = J_{\alpha}^{-}(\lambda) + iR(\lambda + i0)T_{\alpha}^{-}(\lambda),$$

leading to a study of

$$\widetilde{W}_{\alpha}^{-}(\lambda) := W_{\alpha}^{-}(\lambda) - \varphi_{\alpha} \otimes W_{w}^{-}(\lambda - \lambda_{0}), \qquad (6.63a)$$

where $W_w^-(\cdot)$ is the incoming wave matrix for the one-body problem as discussed in Subsection 6.1.2.1. We take $\lambda = \lambda_0$. It is checked as in the proof of Theorem 6.10 (i.e. by using Lemmas 6.8 and 6.9) and by using Proposition 3.16, that for any $k \ge 0$ fulfilling (6.49)

$$\widetilde{W}_{\alpha}^{-}(\lambda_{0})H^{-k} \subset \mathcal{B}_{s_{0}}^{*}, \quad \Pi'\widetilde{W}_{\alpha}^{-}(\lambda_{0})H^{-k} \subset \mathcal{B}^{*},
\chi(\widecheck{B} < \sigma)\Pi'\widetilde{W}_{\alpha}^{-}(\lambda_{0})H^{-k} \subset \mathcal{B}_{0}^{*}, \quad \chi(\widecheck{B}_{\rho} < \sigma)\Pi\widetilde{W}_{\alpha}^{-}(\lambda_{0})H^{-k} \subset \mathcal{B}_{s_{0},0}^{*}.$$
(6.63b)

(For the last property we can use the argument (6.64) given below.) In particular (6.63b) hold for k = 0. Combining the first three assertions of (6.63b) with [DS1, Theorem 8.2] (containing information on $W_w^-(0)$) we conclude that the range of $W_{\alpha}^-(\lambda_0)$ is a subset of $\mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$, i.e. that the map in 1) is a well-defined map. By the same argument it follows that this map is continuous.

<u>II</u>. Let any $\tau \in L^2(\mathbb{S}^{n-1})$ be given. By using the splitting (6.63a), Proposition 3.16, [DS1, Theorem 8.2] and [Sk5] we deduce that (6.62a) is fulfilled with $\phi_{\tau}^+ = W_{\alpha}^-(\lambda_0)\tau$ and for some $\tau^+ \in L^2(\mathbb{S}^{n-1})$.

Using the formula

$$S_{\alpha\alpha}(\lambda_0)^* = 2\pi W_{\alpha}^-(\lambda_0)^* T_{\alpha}^+(\lambda_0) = 2\pi \mathrm{i} W_{\alpha}^-(\lambda_0)^* (H - \lambda_0) J_{\alpha}^+(\lambda_0)$$

and [DS1, Theorem 5.7] we calculate for any $\tilde{\tau} \in C^{\infty}(\mathbb{S}^{n-1})$

$$\begin{aligned} \langle \tilde{\tau}, S_{\alpha\alpha}(\lambda_0) \tau \rangle &= -2\pi \mathrm{i} \lim_{n \to \infty} \langle J^+_{\alpha}(\lambda_0) \tilde{\tau}, [H, \chi_n(r)] W^-_{\alpha}(\lambda_0) \tau \rangle \\ &= -4\pi \lim_{n \to \infty} \langle J^+_{\alpha}(\lambda_0) \tilde{\tau}, g_{\lambda_0} \chi'_n W^-_{\alpha}(\lambda_0) \tau \rangle \\ &= \langle \tilde{\tau}, \tau^+ \rangle. \end{aligned}$$

We conclude that $\tau^+ = S_{\alpha\alpha}(\lambda_0)\tau$. (Note that with our normalization there is an extra factor $1/\sqrt{2}$ in [DS1, (3.13) and Theorem 5.7].)

<u>*III*</u>. For uniqueness, suppose that for a given $\tau \in L^2(\mathbb{S}^{n-1})$ the formula (6.62a) is fulfilled with $\tau^+ = \tau_1$ and $\phi_{\tau}^+ = \phi_1^+$ as well as for $\tau^+ = \tau_2$ and $\phi_{\tau}^+ = \phi_2^+$, then by Theorem 6.3 (applied with $\psi = 0$) $\phi_1^+ = \phi_2^+$, which in turn implies that $\tau_1 = \tau_2$. The injectivity part of the assertion 1) follows similarly.

<u>IV</u>. We show that the map in 1) maps onto (this finishes 1) by the open mapping theorem). So let $u \in \mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$ be given, then we need to find $\tau \in L^2 = L^2(\mathbb{S}^{n-1})$ such that $u = W^-_{\alpha}(\lambda_0)\tau$. Due to the assumption $\chi(\check{B} < \sigma)\Pi' u \in \mathcal{B}^*_0$ we can verbatim use Step V of the proof of Lemma 4.10 to conclude that $E^+_{\mathcal{H}}f = 0$, $f = T^*u \in \mathcal{B}^*_{s_0}(=\mathcal{B}^*_{s_0}(\mathbf{X}_a))$. The vector $\tilde{f} := (1 + r^+_{\lambda_0}v^+_{\lambda_0})f \in \mathcal{B}^*_{s_0}$ fulfills $h\tilde{f} = 0$. Now from [DS1, Theorem 8.2] we know that the map $W^-_w : L^2(\mathbb{S}^{n-1}) \to \{\tilde{f} \in \mathcal{B}^*_{s_0} \mid h\tilde{f} = 0\}$ is bijective. Whence $\tilde{f} = W^-_w(0)\tau$ for some $\tau \in L^2$. We want to show that $\check{u} := u - W^-_\alpha(\lambda_0)\tau = 0$. To do this we check the conditions of Theorem 6.3 (with $\psi = 0$). Since $u \in \mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$ also $\check{u} \in \mathcal{V}^+_{-s_0,\sigma}(\lambda_0)$, and it suffices to check that $\chi(\check{B}_{\rho} < \sigma)\Pi\check{u} \in \mathcal{B}^*_{s_0,0}$. Using (6.63a) and (6.63b) we calculate modulo $\mathcal{B}^*_{s_0,0}$

$$\begin{aligned} \chi(\check{B}_{\rho} < \sigma)\Pi\check{u} &\approx \chi(\check{B}_{\rho} < \sigma)Sf - \chi(\check{B}_{\rho} < \sigma)\Pi\varphi_{\alpha} \otimes W_{w}^{-}(0)\tau \\ &= \chi(\check{B}_{\rho} < \sigma)S\tilde{f} - \chi(\check{B}_{\rho} < \sigma)SW_{w}^{-}(0)\tau + \chi(\check{B}_{\rho} < \sigma)S(f - \tilde{f}) \\ &= \chi(\check{B}_{\rho} < \sigma)S(f - \tilde{f}). \end{aligned}$$

We let $\check{B}_{a,\rho}$ be given as in (6.15) and then write, substitute and estimate

$$\chi(\check{B}_{\rho} < \sigma)S - S\chi(\check{B}_{a,\rho} < \sigma) = -\int_{\mathbb{C}}\check{R}_{\rho}(z)[\check{B}_{\rho}S - S\check{B}_{a,\rho}]\check{R}_{a,\rho}(z)\mathrm{d}\mu_{\chi}(z)$$

by the arguments of the proof of Lemma 3.4, estimating

$$\chi(\check{B}_{\rho} < \sigma)S(f - \tilde{f}) \approx -S\chi(\check{B}_{a,\rho} < \sigma)r_{\lambda_0}^+ v_{\lambda_0}^+ f \approx 0;$$
(6.64)

in the last step we used Lemma 6.9. Whence $\check{u} = 0$, and we are done.

<u>V</u>. It remains to show 3). First we note that $\check{B}_{\rho}\Pi W_{\alpha}^{-}(\lambda_{0})\tau \in \mathcal{B}_{s_{0}}^{*}$, cf. the proofs of Theorem 6.3 and Corollary 6.4 (and (6.64)). Writing $u = W_{\alpha}^{-}(\lambda_{0})\tau$ and $f = T^{*}u \in \mathcal{B}_{s_{0}}^{*}$ we know that $E_{\mathcal{H}}^{+}f = 0$ and $\check{B}_{a,\rho}f \in \mathcal{B}_{s_{0}}^{*}$, and we need to show that

$$\check{B}_{a,\rho}f - f_2 \in \mathcal{B}^*_{s_0,0}; \quad f_2 = v_a^-(x_a)\tau(-\hat{x_a}) + v_a^+(x^a)\tau^+(\hat{x_a}). \tag{6.65}$$

From (6.62a) it follows that

$$f - f_1 \in \mathcal{B}^*_{s_0,0}; \quad f_1 = u_a^-(x_a)\tau(-\hat{x_a}) + u_a^+(x^a)\tau^+(\hat{x_a}).$$
 (6.66)

With symbols s and g_a given as in (6.16a) we can use (6.16b) to write

$$\tilde{f} := f - Af \in \mathcal{B}^*_{s_0,0}; \quad A = 2g_a^{-1} \operatorname{Op}^{\mathsf{w}}(s^{-1})g_a.$$

From the proof of Theorem 6.3 it follows that $\check{B}^2_{a,\rho}f \in \mathcal{B}^*_{s_0}$ and therefore we know that $\check{B}^2_{a,\rho}\tilde{f} \in \mathcal{B}^*_{s_0}$, and using this fact we can show that also $\check{B}_{a,\rho}\tilde{f} \in \mathcal{B}^*_{s_0,0}$. Estimating

$$R^{-2s_0} \|\chi_R \check{B}_{a,\rho} \tilde{f}\|^2 \le R^{-2s_0} \|\chi_R \tilde{f}\| \|\chi_R \check{B}_{a,\rho}^2 \tilde{f}\| + o(R^0) = o(R^0),$$

the assertion follows.

Next we substitute $f = Af + \tilde{f}$ into (6.65) and calculate modulo $\mathcal{B}^*_{s_0,0}$ using (6.66)

$$\check{B}_{a,\rho}f \approx \check{B}_{a,\rho}Af \approx \check{B}_{a,\rho}Af_1 \approx A\check{B}_{a,\rho}f_1 \approx Af_2 = f_2 - \tilde{f}_2; \quad \tilde{f}_2 = f_2 - Af_2.$$

We calculate using (6.16b)

$$\tilde{f}_2 \approx \frac{1}{2}A\left(g_a^{-1}\operatorname{Op}^{\mathsf{w}}(s)g_af_2 - 2f_2\right) \approx \frac{1}{2}Ag_a^{-2}hf_2$$

It remains to show that $Ag_a^{-2}hf_2 \in \mathcal{B}^*_{s_0,0}$. We approximate τ and τ^+ by sequences of smooth functions in $L^2(\mathbb{S}^{n-1})$ giving convergence in $\mathcal{B}^*_{s_0}$ since $Ag_a^{-2}h \in \mathcal{L}(\mathcal{B}^*_{s_0})$. Whence we can assume that $\tau, \tau^+ \in C^{\infty}$, which allow us to compute $g_a^{-2}hf_2 \in \mathcal{B}^*_{s_0,0}$ and we are done.

The following bound is a consequence of the definition of wave operators and Theorem 6.10, but for completeness of presentation we give an independent stationary proof.

Corollary 6.14. Under the same conditions as above $||S_{\alpha\alpha}(\lambda_0)|| \leq 1$.

Proof. Following the proof of Theorem 6.3 we calculate, abbreviating $\phi = W_{\alpha}^{-}(\lambda_{0})\tau$ for any given $\tau \in L^{2}(\mathbb{S}^{n-1})$,

$$0 = -\frac{1}{2} \langle \mathbf{i}[H, \chi_R] \rangle_{\phi} = \langle \check{B} \rangle_{\theta_R \Pi' \phi} + \langle \check{B} \rangle_{\theta_R \Pi \phi} + o(R^0)$$

$$\geq -\operatorname{Re} \langle \chi'_R \langle x \rangle^{-\rho/2} \Pi \phi, \check{B}_\rho \Pi \phi \rangle + o(R^0)$$

By (6.62a)

$$\Pi \phi - \varphi_{\alpha}(x^{a}) \left(u_{a}^{-}(x_{a})\tau(-\widehat{x_{a}}) + u_{a}^{+}(x^{a})\tau^{+}(\widehat{x_{a}}) \right) \in \mathcal{B}_{s_{0},0}^{*}$$

We insert this and (6.62b) and take $R \to \infty$, yielding $0 \ge -\|\tau\|^2 + \|S_{\alpha\alpha}(\lambda_0)\tau\|^2$. \Box

Let us for any $\tau \in C^{\infty}(\mathbb{S}^{n-1})$ write, using (6.7) and (6.13),

$$R(\lambda_0 + \mathrm{i0})\psi = S\phi_a^+(\lambda_0) + R'(\lambda_0 + \mathrm{i0})\psi_a^+(\lambda_0) \text{ with input } \psi \coloneqq T_\alpha^-(\lambda_0)\tau \in \mathcal{B}_{s_0},$$

$$\delta(H' - \lambda_0) = \frac{1}{2\pi\mathrm{i}} \big(R'(\lambda_0 + \mathrm{i0}) - R'(\lambda_0 - \mathrm{i0}) \big) \in \mathcal{L}(\mathcal{B}, \mathcal{B}^*).$$

Corollary 6.15. Under the above conditions and with $\psi_a^+(\lambda_0)$ defined by (6.7) with $\psi = T_{\alpha}^-(\lambda_0)\tau$ for any given $\tau \in C^{\infty}(\mathbb{S}^{n-1})$, or alternatively given by

$$\psi_a^+(\lambda_0) = \Pi'(H - \lambda_0)\Pi' R(\lambda_0 + \mathrm{i}0)T_\alpha^-(\lambda_0)\tau,$$

$$T_\alpha^-(\lambda_0)\tau = \mathrm{i}(H - \lambda_0)J_\alpha^-(\lambda_0)\tau \quad and \ \tau \in C^\infty(\mathbb{S}^{n-1}),$$

the following formula holds:

$$\|\tau\|^{2} - \|S_{\alpha\alpha}(\lambda_{0})\tau\|^{2} = \pi \langle \delta(H' - \lambda_{0}) \rangle_{\psi_{a}^{+}(\lambda_{0})}.$$
(6.67)

Proof. Recall

$$\phi := W_{\alpha}^{-}(\lambda_0)\tau = J_{\alpha}^{-}(\lambda_0)\tau + iR(\lambda_0 + i0)\psi.$$

We used in the proof of Corollary 6.14 that $\langle \check{B} \rangle_{\theta_R \Pi' \phi}$ asymptotically is non-negative. More precisely we can calculate

$$\lim_{R \to \infty} \langle \check{B} \rangle_{\theta_R \Pi' \phi} = \lim_{R \to \infty} \langle -\frac{1}{2} \mathrm{i} [H', \chi_R] \rangle_{\phi}$$
$$= \lim_{R \to \infty} \langle -\frac{1}{2} \mathrm{i} [H', \chi_R] \rangle_{R'(\lambda_0 + \mathrm{i} 0) \psi_a^+(\lambda_0)} = \pi \langle \delta(H' - \lambda_0) \rangle_{\psi_a^+(\lambda_0)}.$$

Remark 6.16. If $\lambda_0 = \Sigma_2$ the right-hand side of (6.67) vanishes for all $\tau \in C^{\infty}(\mathbb{S}^{n-1})$. However for the non-multiple case above Σ_2 we dont see any reason this be the case. Consequently the option of 'transmission' is left as a conjecture for $\lambda_0 > \Sigma_2$ (including the multiple case of Subsection 6.1.3). We shall study the problem of 'non-transmission' for the physics models at a two-cluster threshold in detail in Section 6.3.

6.1.2.4 Elastic scattering at Σ_2

We shall supplement Theorem 6.10 under the additional conditions that $\lambda_0 = \Sigma_2$ and that $V_1(r) = -\gamma r^{-\rho}$ for $r \ge 1$, by then proving that the kernel of $S_{\alpha\alpha}(\lambda_0)$ is smooth outside $\{\omega \cdot \omega' = \cos \frac{\rho}{2-\rho}\pi\}$. With the new more restrictive condition on λ_0 ,

$$\lambda_0 \in \sigma_{\rm d}(H^a). \tag{6.68}$$

With (6.68) it is possible to 'improve' on the properties of the operator $T^{\pm}_{\alpha}(\lambda_0)$ by solving the transport equations more carefully, cf. [Bo, Sk2]. Relying only on (6.68) this does not need λ_0 to be the lowest threshold. However we will need the additional property

$$\lambda_0 \notin \sigma_{\rm ess}(H'),\tag{6.69a}$$

see Lemma 6.21, which indeed is fulfilled for $\lambda_0 = \Sigma_2$. Although this condition can be weakened to $\lambda_0 < \Sigma_3$, cf. Remark 2.11, we will in this subsection assume $\lambda_0 = \Sigma_2$.

Recall that in this section we impose (6.2), so (6.69a) may be stated equivalently as

$$\lambda_0 \notin \sigma(H'). \tag{6.69b}$$

We shall show the following analogue of [DS1, Theorem 9.3].

Theorem 6.17. Suppose in addition to the conditions of Theorem 6.10 that $\lambda_0 = \Sigma_2$ and that $V_1(r) = -\gamma r^{-\rho}$ for $r \ge 1$. Then the kernel $S_{\alpha\alpha}(\lambda_0)(\omega, \omega')$ is smooth outside the set $\{(\omega, \omega') \mid \omega \cdot \omega' = \cos \frac{\rho}{2-\rho}\pi\}$.

We need various preparation partly similarly to [DS1, Section 9] to prove this result. The first result stated as Lemma 6.20 uses only the conditions of Theorem 6.10 and (6.68).

To see how (6.68) allows us to 'improve' $T^{\pm}_{\alpha}(\lambda_0) = (H - \lambda_0) J^{\pm}_{\alpha}(\lambda_0)$ we look at the term

$$(I_a^{(1)} - w)J_\alpha^{\pm}(\lambda_0) = \Pi(I_a^{(1)} - w)J_\alpha^{\pm}(\lambda_0) + \Pi'I_a^{(1)}\Pi J_\alpha^{\pm}(\lambda_0).$$

The first term has polynomial decay, viz. it belongs to $\mathcal{L}_s^k = \mathcal{L}(H^k(\mathbb{S}^{n-1}), L_s^2(\mathbf{X}))$ for any $k, s \in \mathbb{R}$, and the second term is $\mathcal{O}(\langle x \rangle^{-1-\rho}) J_{\alpha}^{\pm}(\lambda)$, cf. (6.40). We look at the leading term after a Taylor expansion

$$\Pi' I_a^{(1)} \Pi J_\alpha^{\pm}(\lambda_0) \approx \left(\Pi' x^a \varphi_\alpha \right) \otimes \left(\nabla I_a^{(1)}(x_a) J_N^{\pm}(\lambda_0) \right).$$

Let $\tilde{r}_{\alpha}(\lambda_0)$ be the reduced resolvent of H^a at λ_0 . Then we add the term

$$J_{\alpha}^{1\pm}(\lambda_0) := -\left(\tilde{r}_{\alpha}(\lambda_0)\Pi' x^a \varphi_{\alpha}\right) \otimes \left(\nabla I_a^{(1)}(x_a) J_N^{\pm}(\lambda_0)\right)$$

to $J^{\pm}_{\alpha}(\lambda_0)$ and compute for the resulting operator $\check{J}^{1\pm}_{\alpha}(\lambda_0) = J^{\pm}_{\alpha}(\lambda_0) + J^{1\pm}_{\alpha}(\lambda_0)$, observing a cancellation,

$$(H - \lambda_0) J_{\alpha}^{1\pm}(\lambda_0) = (p_a^2 + I_a) J_{\alpha}^{\pm}(\lambda_0) - (\Pi' x^a \varphi_{\alpha}) \otimes (\nabla I_a^{(1)}(x_a) J_N^{\pm}(\lambda_0)) - (p_a^2 + I_a) (\tilde{r}_{\alpha}(\lambda_0) \Pi' x^a \varphi_{\alpha}) \otimes (\nabla I_a^{(1)}(x_a) J_N^{\pm}(\lambda_0)) = \varphi_{\alpha} \otimes T_N^{\pm}(\lambda_0) \tau + \mathcal{O}(\langle x \rangle^{-2-\rho}) J_{\alpha}^{\pm}(\lambda_0) - (\tilde{r}_{\alpha}(\lambda_0) \Pi' x^a \varphi_{\alpha}) \otimes (h \nabla I_a^{(1)}(x_a) J_N^{\pm}(\lambda_0)) + \mathcal{O}(\langle x \rangle^{-2-2\rho}) J_{\alpha}^{\pm}(\lambda_0).$$
(6.70a)

The second and the fourth terms have better decay than the term $\mathcal{O}(\langle x \rangle^{-1-\rho}) J^{\pm}_{\alpha}(\lambda)$ we started out with. The third term has a different form and needs examination. We compute, using the notation (6.28) and doing for simplicity only the plus case,

$$h\nabla I_{a}^{(1)}(x_{a}) \left(J_{N}^{+}(\lambda_{0})\tau\right)(x_{a})$$

$$= \left(\mathcal{O}(\langle x_{a}\rangle^{-1-\rho})h + \mathcal{O}(\langle x_{a}\rangle^{-2-\rho})p_{a} + \mathcal{O}(\langle x_{a}\rangle^{-3-\rho})\right)J_{N}^{+}(\lambda_{0})\tau$$

$$= \int e^{i\phi^{+}(x_{a},\omega,0)}e^{\tilde{\zeta}^{+}(x_{a},\omega,0)}\mathcal{O}(\langle x_{a}\rangle^{-2-\rho^{3}/2})\tau(\omega)d\omega.$$
(6.70b)

Whence the third term is a power $\mathcal{O}(\langle x \rangle^{-1-\rho/2})$ better than what we started with.

Partly motivated by the above considerations let us introduce the spaces

$$L_{s}^{2,a} = L_{s}^{2,a}(\mathbf{X}) = \{ u \in L_{s}^{2}(\mathbf{X}) \mid \forall m \in \mathbb{N} : \langle x^{a} \rangle^{m} u \in L_{s}^{2}(\mathbf{X}) \}; \quad s \in \mathbb{R},$$

$$L_{-\infty}^{2,a} = \bigcup_{s \in \mathbb{R}} L_{s}^{2,a},$$

$$\mathcal{L}_{s}^{k,a} = \mathcal{L}(H^{k}(\mathbb{S}^{n-1}), L_{s}^{2,a}); \quad k, s \in \mathbb{R}.$$
(6.71)

We shall also need the following subclasses \mathcal{C}_s^+ , $s \in \mathbb{R}$; see (6.73) for a relationship.

Definition 6.18. Let \mathcal{F}^a be the set of functions in $L^2_{\infty}(\mathbf{X}^a)$ of the form $f^a = T\varphi_{\alpha}$, where T is any multiple product of factors of Π' , $\tilde{r}_{\alpha}(\lambda_0)$ and multiplication by components of x^a . Let $R \geq R_0$ and $\sigma' \in (0, \sigma_0)$ be given. We consider operators of the tensor product type

$$f^a \otimes J_g^+; \quad (J_g^+\tau)(x_a) = \int_{\mathbb{S}^{n-1}} e^{i\phi^+(y,\omega,0)} e^{\tilde{\zeta}^+(y,\omega,0)} g(y,\omega)\tau(\omega) d\omega, \quad y = x_a,$$

where $f^a \in \mathcal{F}^a$ and the symbol $g \in \mathcal{S}_s^+$, meaning (with reference to (6.22)) that

$$\left|\partial_{\omega}^{\delta}\partial_{y}^{\gamma}g(y,\omega)\right| \leq C_{\delta,\gamma}\langle y\rangle^{-s-|\gamma|} \text{ and } \operatorname{supp} g \subseteq \Gamma_{R,\sigma'}^{+}.$$
(6.72)

The class \mathcal{C}_s^+ , $s \in \mathbb{R}$, is the set of operators given as a finite sum of such products, where for each term the symbol $g \in \mathcal{S}_s^+$.

Due to properties of ϕ^+ and $\tilde{\zeta}^+$ it follows from the proof of [DS1, Theorem 6.5] that

$$C_s^+ \subset \mathcal{L}_{s-(1-\rho/2)k-s_0-\epsilon}^{-k,a}; \quad k \ge 0, \ \epsilon > 0.$$
 (6.73)

We also note that $J^+(0)$ is of the form J_g^+ for some symbol $g \in \mathcal{S}_0^+$, see (6.25a) and (6.27), and therefore $J_{\alpha}^+(\lambda_0) \in \mathcal{C}_0^+$.

Ideally we would like to solve $\check{T}^{\pm}_{\alpha}(\lambda_0) = (H - \lambda_0)\check{J}^{\pm}_{\alpha}(\lambda_0) \in \mathcal{L}^{k,a}_s$ for any given $k \leq 0$ and $s \geq 0$ by modifying the construction $J^{\pm}_{\alpha}(\lambda_0)$, say denoted by $\check{J}^{\pm}_{\alpha}(\lambda_0) \approx J^{\pm}_{\alpha}(\lambda_0)$. Of course like for the one-body problem we cannot do that, but we can solve transport equations in a (small) forward cone (cf. the splitting for the one-body problem, $T^{\pm}_N(\lambda) = T^{\pm}_{\mathrm{bd}}(\lambda - \lambda_0) + T^{\pm}_{\mathrm{pr}}(\lambda - \lambda_0)$, corresponding to the decomposition of $t^{\pm}(x,\omega,\lambda) = t^{\pm}(x,\sqrt{\lambda}\omega)$ in [DS1, (5.8) and (5.16)]). We shall mimic this procedure using the reduced resolvent $\tilde{r}_{\alpha}(\lambda_0)$ as an additional tool. In fact we computed above for $\check{J}^{1+}_{\alpha}(\lambda_0) := J^{+}_{\alpha}(\lambda_0) + J^{1+}_{\alpha}(\lambda_0)$

$$(H - \lambda_0) \breve{J}^{1+}_{\alpha}(\lambda_0) = \varphi_{\alpha} \otimes T^+_N(\lambda_0) + T^{1+}_{\alpha}(\lambda_0) + T^{1+}_{\alpha,\mathbf{r}}(\lambda_0),$$

where $T^{1+}_{\alpha}(\lambda_0) \in \mathcal{C}^+_{2+\rho_{3/2}}$ and (for example)

$$T^{1+}_{\alpha,\mathbf{r}}(\lambda_0) \in \mathcal{L}^{-l,a}_{2+\rho-(1-\rho/2)l-s_0-\epsilon}; \quad l \ge 0, \ \epsilon > 0.$$

The argument relied on using the first term of a Taylor expansion only, however we can do the Taylor expansion of $\Pi' I_a^{(1)} \Pi$ to any order, and each term will then contribute by a term in $\mathcal{C}_{2+\rho}^+$ except (possibly) a 'remainder term' which can be taken in the fixed space $\mathcal{L}_s^{k,a}$ of interest. We can argue the same way for the fourth term $\mathcal{O}(\langle x \rangle^{-2-2\rho}) J_{\alpha}^{\pm}(\lambda_0)$ of (6.70a). We conclude that $(H-\lambda_0) \check{J}_{\alpha}^{1+}(\lambda_0) - \varphi_{\alpha} \otimes T_N^+(\lambda_0) + T \in$ $\mathcal{L}_s^{k,a}$ for some $T \in \mathcal{C}_{2+\rho}^+$.

To improve further, for this $T \in \mathcal{C}_{2+\rho}^+$ ideally we would like to 'solve' the equation $(H - \lambda_0) J_{\alpha}^{2+}(\lambda_0) \approx T$. This would yield a better approximation by adding $J_{\alpha}^{2+}(\lambda_0)$, viz. by considering $\check{J}_{\alpha}^{2+}(\lambda_0) := J_{\alpha}^+(\lambda_0) + J_{\alpha}^{1+}(\lambda_0) + J_{\alpha}^{2+}(\lambda_0)$. We are lead to considering the following iteration scheme. Suppose that for given $m \in \mathbb{N}$ we have constructed $\check{J}_{\alpha}^{m+}(\lambda_0) := J_{\alpha}^+(\lambda_0) + J_{\alpha}^{1+}(\lambda_0) + \cdots + J_{\alpha}^{m+}(\lambda_0)$ such that

$$(H - \lambda_0) \breve{J}^{m+}_{\alpha}(\lambda_0) - \varphi_{\alpha} \otimes T^+_N(\lambda_0) + T^{m+}_{\text{pr}}(\lambda_0) - T^{m+}_{\text{bd}}(\lambda_0) \in \mathcal{L}^{k,a}_s$$
$$T^{m+}_{\text{pr}}(\lambda_0) \in \mathcal{C}^+_{s_m}; \quad s_m = (m-1)\min\{1 - \rho/2, \rho/2\} + 2 + \rho, \qquad (6.74)$$
$$T^{m+}_{\text{bd}}(\lambda_0) \in \mathcal{C}^+_{\text{bd}, s_1},$$

where $\mathcal{C}^+_{\mathrm{bd},s}$ is given as follows. The notation M° refers in general to the interior of a subset M of a topological space (below taken as $\mathbb{R}^n \times \mathbb{S}^{n-1}$).

Definition 6.19. Let $R \geq R_0$ and $\sigma' \in (0, \sigma_0)$ be given, cf. Definition 6.18, and let $\sigma \in (0, \sigma')$. Then $\mathcal{C}^+_{\mathrm{bd},s}$ is the subclass of operators $\varphi_{\alpha} \otimes J_g^+ \in \mathcal{C}_s^+$ for which the symbol $g = g_{\mathrm{bd}}^+ \in \mathcal{S}_s^+$ (as required by Definition 6.18) but in addition has the support property

$$\operatorname{supp} g_{\mathrm{bd}}^+ \subseteq \Gamma_{R,\sigma'}^+ \setminus \left(\Gamma_{2R,\sigma}^+\right)^\circ.$$
(6.75)

So far we have verified (6.74) for m = 1 only (with $T_{bd}^{1+}(\lambda_0) = 0$). Now, suppose (6.74) for a given $m \ge 1$. Then we split

$$T = T_{\mathrm{pr}}^{m+}(\lambda_0) = \Pi T_{\mathrm{pr}}^{m+}(\lambda_0) + \Pi' T_{\mathrm{pr}}^{m+}(\lambda_0),$$

and both of the terms to the right contributes to the construction of $J_{\alpha}^{(m+1)+}(\lambda_0)$ as follows: For the first term we mimic [DS1, Propositions 5.4 and 5.5] and add correspondingly a term, say $J_{\alpha,1}^{(m+1)+}(\lambda_0)$. By a Taylor expansion (as used above) we see that

$$(H - \lambda_0) J_{\alpha,1}^{(m+1)+}(\lambda_0) - \Pi T_{\rm pr}^{m+}(\lambda_0) \in \mathcal{C}_{s_{m+1}}^+ + \mathcal{C}_{\rm bd,\,s_1}^+ + \mathcal{L}_s^{k,a}; \tag{6.76}$$

we give the details below. Letting

$$J_{\alpha,2}^{(m+1)+}(\lambda_0) = \tilde{r}_{\alpha}(\lambda_0) \Pi' T_{\rm pr}^{m+}(\lambda_0)$$

we then obtain that indeed (6.74) is fulfilled with m replaced by m + 1 and with

$$\breve{J}_{\alpha}^{(m+1)+}(\lambda_0) = \breve{J}_{\alpha}^{m+}(\lambda_0) + J_{\alpha}^{(m+1)+}(\lambda_0); \quad J_{\alpha}^{(m+1)+}(\lambda_0) = J_{\alpha,1}^{(m+1)+}(\lambda_0) + J_{\alpha,2}^{(m+1)+}(\lambda_0).$$

Note that obviously $(H - \lambda_0) J_{\alpha,2}^{(m+1)+}(\lambda_0) - \Pi' T_{\text{pr}}^{m+}(\lambda_0) \in \mathcal{C}_{s_{m+1}}^+ + \mathcal{L}_s^{k,a}$, cf. (6.70a) and (6.70b).

To complete the recursive construction it remains to justify (6.76): We use the function in (6.24a), more precisely we consider 'cut-offs' $\bar{\chi}_R(r)$ and $\chi_{\sigma,\sigma'}(\hat{x} \cdot \omega)$. Defined in terms of the function $\zeta^+ = \zeta^+(x_a, \omega, 0)$ of (6.28) we let A^+ be the differential operator (in $x = x_a$)

$$A^{+} = \Delta + 2(\nabla \zeta^{+}) \cdot \nabla + (\Delta \zeta^{+}) + (\nabla \zeta^{+})^{2},$$

cf. (6.25f), and we define then for $\Pi T_{\mathrm{pr}}^{m+}(\lambda_0) = \varphi_{\alpha} \otimes J_{g_m}^+$ correspondingly $J_{\alpha,1}^{(m+1)+}(\lambda_0) = \varphi_{\alpha} \otimes J_{g_{m+1}}^+ \in \mathcal{C}_{s_m-1-\rho/2}^+$, where (with $x = x_a$)

$$g_{m+1} := -i\chi_{\sigma,\sigma'}(\hat{x}\cdot\omega)\bar{\chi}_R(r)\int_1^\infty g_m(y(t,x,\omega,0),\omega)\,\mathrm{d}t;$$

here $y(\cdot, x, \omega, 0)$ is the classical zero-energy orbit starting at x for t = 1 and with asymptotic normalized velocity $\omega = \lim_{t \to +\infty} y/|y|$, cf. (6.20). Let

$$\breve{g}_{m+1} = \mathrm{i}\chi_{\sigma,\sigma'}(\hat{x}\cdot\omega)\bar{\chi}_R(r)\int_1^\infty A^+g_m(y(t,x,\omega,0),\omega)\,\mathrm{d}t$$

We calculate

$$(H - \lambda_0)\varphi_{\alpha} \otimes J_{g_{m+1}}^+ + \Pi T_{\mathrm{pr}}^{m+}(\lambda_0) + \varphi_{\alpha} \otimes J_{\check{g}_{m+1}}^+$$

= $\varphi_{\alpha} \otimes h J_{g_{m+1}}^+ + (I_a - w)\varphi_{\alpha} \otimes J_{g_{m+1}}^+ + \Pi T_{\mathrm{pr}}^{m+}(\lambda_0) + \varphi_{\alpha} \otimes J_{\check{g}_{m+1}}^+$
 $\in \mathcal{C}_{s_{m+1}}^+ + \mathcal{C}_{\mathrm{bd},s_1}^+ + \mathcal{L}_s^{k,a};$

here we used a Taylor expansion to treat the second term (as we did in (6.70a)) and the fact that the three other terms cancel up to derivatives of the factor $\chi_1(r/R)\chi_2(\hat{x} \cdot \omega)$, cf. (6.25f). Using next that $\check{g}_{m+1} \in \mathcal{S}^+_{s_{m+1}}$ indeed (6.76) follows.

Next, rather than doing the Borel summation as for the one-body problem, for simplicity we 'terminate' the recursive construction at m = M taken so large that $\mathcal{C}_{s_{m+1}}^+ \subset \mathcal{L}_s^{k,a}$, cf. (6.73). Beforehand we treated for convenience only the plus case. Leaving it to the reader to figure out how Definitions 6.18 and 6.19 should read in the minus case, and how the above procedure correspondingly modifies, we consider henceforth the 'improved' operators $J_{\alpha,M}^{\pm}(\lambda_0) := \check{J}_{\alpha}^{M\pm}(\lambda_0)$ and

$$T^{\pm}_{\alpha,M}(\lambda_0) := (H - \lambda_0) J^{\pm}_{\alpha,M}(\lambda_0) \in \varphi_{\alpha} \otimes T^{\pm}_N(\lambda_0) + \mathcal{C}^{\pm}_{\mathrm{bd},s_1} + \mathcal{L}^{k,a}_s.$$

In turn we consider the corresponding wave matrices $W_{\alpha,M}^{\pm}(\lambda_0)$ and the scattering operator $S_{\alpha,M}(\lambda_0)$; see (6.82) below. Like for the one-body problem the latter quantites are canonical (this is stated more precisely as the last assertion in the following conclusion).

Lemma 6.20. Let $k \leq 0$, $s \geq 0$, $R \geq R_0$ and $0 < \sigma < \sigma' < \sigma_0$. Then the above recursive procedure, terminating at any sufficiently large $M \in \mathbb{N}$, yields the existence of $J^{\pm}_{\alpha,M}(\lambda_0) \in J^{\pm}_{\alpha}(\lambda_0) + \mathcal{C}^{\pm}_{2s_0}$, $g^{\pm}_{\mathrm{bd},M} \in \mathcal{S}^{\pm}_{2s_0}$ with $\operatorname{supp} g^{\pm}_{\mathrm{bd},M} \subseteq \Gamma^{\pm}_{R,\sigma'} \setminus (\Gamma^{\pm}_{2R,\sigma})^{\circ}$ and $R^{k,a\pm}_{s,M} \in \mathcal{L}^{k,a}_s$ such that

$$T^{\pm}_{\alpha,M}(\lambda_0) = (H - \lambda_0) J^{\pm}_{\alpha,M}(\lambda_0) = (2\pi)^{-n/2} \varphi_\alpha \otimes J^{\pm}_g + R^{k,a\pm}_{s,M}; \quad g = g^{\pm}_{\mathrm{bd},M}$$

(In particular $T^{\pm}_{\alpha,M}(\lambda_0) \in \mathcal{C}^{\pm}_{\mathrm{bd},2s_0} + \mathcal{L}^{k,a}_s$.)

The wave matrices $W_{\alpha,M}^{\pm}(\lambda_0)$ and the scattering operator $S_{\alpha,M}(\lambda_0)$ defined by the operators $J_{\alpha,M}^{\pm}(\lambda_0)$ and $T_{\alpha,M}^{\pm}(\lambda_0)$ coincide with $W_{\alpha}^{\pm}(\lambda_0)$ and $S_{\alpha}(\lambda_0)$, respectively.

Note that the parameter R, σ and σ' of the lemma are used to define the classes $C_{\mathrm{bd},s}^{\pm}$. The first part of the lemma is clearly a consequence of the explained construction. For the second part, note that for the cases of $W_{\alpha,M}^{-}(\lambda_0)$ and $S_{\alpha,M}$ the assertion is a consequence of the uniqueness part of Theorem 6.13 2). The identification $W_{\alpha,M}^{+}(\lambda_0) = W_{\alpha}^{+}(\lambda_0)$ follows from an analogue statement for $W_{\alpha}^{+}(\lambda_0)$ given in terms of $S_{\alpha}(\lambda_0)^*$ (not given in our presentation).

As in (6.41) the notation g means $g_{\lambda_0} = \sqrt{-V_1}$. We introduce symbols b and \bar{c} to decompose the normalized momentum $\eta := \xi/g \in \mathbf{X}'_a$ as

$$\eta = b\hat{x} + \bar{c}, \ b := \hat{x} \cdot \eta \text{ and } \bar{c} := \left(I - \left|\hat{x}\right\rangle \langle \hat{x} \right|\right) \eta.$$
 (6.77)

Of course this decomposition requires $x \neq 0$ since $\hat{x} = x/|x|$ only make sense for such x. (In the wave front set definition below this issue is handled by a cut-off $\bar{\chi}_1$, cf. (1.29). In (6.41) a slightly different b 'cured' the problem.)

On the energy shelf $\xi^2 = \gamma r^{-\rho}$ the quantity $a := b^2 + \bar{c}^2 = \xi^2/g^2 = 1$ and the Hamiltonian orbits solve the ODE on the 'reduced phase space' \mathbb{T}^* , consisting of points (\hat{x}, \bar{c}, b) ,

$$\begin{cases} \frac{\mathrm{d}}{\mathrm{d}\tau}\hat{x} = \bar{c}, \\ \frac{\mathrm{d}}{\mathrm{d}\tau}\bar{c} = -(1 - \frac{\rho}{2})b\bar{c} - \bar{c}^2\hat{x}, \\ \frac{\mathrm{d}}{\mathrm{d}\tau}b = (1 - \frac{\rho}{2})\bar{c}^2. \end{cases}$$
(6.78)

This is in the 'new time' τ given by $\frac{d\tau}{dt} = 2g/r$.

The maximal solution of (6.78) that passes $z = (\hat{x}, b, \bar{c}) \in \mathbb{T}^*$ at $\tau = 0$ is denoted by $\gamma(\tau, z)$. The quantity *a* is preserved by the flow, and the equation $\bar{c} = 0$ defines the fixed points. Away from those points

$$b(\tau) = \sqrt{a} \tanh \sqrt{a} (1 - \frac{\rho}{2})(\tau - \tau_0),$$

showing moreover that b is monotonely increasing in τ from $-\sqrt{a}$ to \sqrt{a} (away from fixed points). We introduce in terms of the variables (6.77) the 'wave front set' $WF_s^a(u)$ of a distribution $u \in L^{2,a}_{-\infty}$ as the subset of \mathbb{T}^* given by the condition

$$z_{1} = (\omega_{1}, \bar{c}_{1}, b_{1}) = (\omega_{1}, b_{1}\omega_{1} + \bar{c}_{1}) = (\omega_{1}, \eta_{1}) \notin WF_{s}^{a}(u)$$

$$\Leftrightarrow \qquad (6.79)$$

$$\exists neighbourhoods \mathcal{N}_{\omega_{1}} \ni \omega_{1}, \mathcal{N}_{\eta_{1}} \ni \eta_{1} \quad \forall \chi_{\omega_{1}} \in C_{c}^{\infty}(\mathcal{N}_{\omega_{1}}), \ \chi_{\eta_{1}} \in C_{c}^{\infty}(\mathcal{N}_{\eta_{1}}) :$$

$$Op^{w}(\chi_{z_{1}}\bar{\chi}_{1}(r))u \in L_{s}^{2,a} \text{ where } \chi_{z_{1}} = \chi_{z_{1}}(x,\xi) = \chi_{\omega_{1}}(\hat{x})\chi_{\eta_{1}}(\xi/g(r)).$$

Obviously this notion of wave front set is a (fibered) adaption of the notion of 'scattering wave front set' $WF^s_{sc}(v)$ of a distribution $v \in L^2_{-\infty}(\mathbb{R}^n)$ of [DS1, Subsection 4.2] to the present problem.

Due to (6.69b) the operators $(H' - \lambda_0)^{-1}$ and $\check{R}(\lambda_0)$ respect the above notion of 'fibered scattering wave front set'.
Lemma 6.21. For any $s \in \mathbb{R}$ and $u \in L^{2,a}_{-\infty}$ the following properties hold.

$$\Pi I_a \Pi' L_s^{2,a} \subset L_{s+1+\rho}^{2,a}, \quad \Pi' I_a \Pi L_s^{2,a} \subset L_{s+1+\rho}^{2,a}, \tag{6.80a}$$

$$WF^a_{s+1+\rho}(\Pi I_a\Pi' u) \subset WF^a_s(u), \quad WF^a_{s+1+\rho}(\Pi' I_a\Pi u) \subset WF^a_s(u), \tag{6.80b}$$

$$\breve{R}(\lambda_0)\Pi' L_s^{2,a} \subset L_s^{2,a},\tag{6.80c}$$

$$WF^a_s(\breve{R}(\lambda_0)\Pi' u) \subset WF^a_s(u).$$
 (6.80d)

Proof. For (6.80a) and (6.80b) we may use a simplified version of (2.23). The results are almost trivial since Π and Π' are operators in the x^a -coordinate while the wave front setting is defined in terms of quantization on \mathbf{X}_a .

The arguments for (2.51) works for (6.80c), so it remains to consider (6.80d), actually without the factor Π' . So for any $z_1 \notin WF_s^a(u)$ we need to estimate $\operatorname{Op}^{\mathsf{w}}(\chi_{z_1}\bar{\chi}_1(r))\check{R}(\lambda_0)u \in L_s^{2,a}$, where we can assume that the localization function χ_{z_1} is supported sufficiently close to z_1 . In particular there is a slightly bigger one, say denoted by $\tilde{\chi}_{z_1}$, i.e. $\tilde{\chi}_{z_1} = 1$ on $\operatorname{supp} \chi_{z_1}$, such that $\operatorname{Op}^{\mathsf{w}}(\tilde{\chi}_{z_1}\bar{\chi}_1(r))u \in L_s^{2,a}$. Whence it remains to show that

$$\operatorname{Op}^{\mathsf{w}}(\chi_{z_1}\bar{\chi}_1(r))\check{R}(\lambda_0)\left(1-\operatorname{Op}^{\mathsf{w}}(\tilde{\chi}_{z_1}\bar{\chi}_1(r))\right)u \in L^{2,a}_s$$

For that it suffice to show that

$$\forall s, t \in \mathbb{R} \exists r \ge 0: \quad \operatorname{Op}^{\mathsf{w}}(\chi_{z_1} \bar{\chi}_1(r)) \check{R}(\lambda_0) \left(1 - \operatorname{Op}^{\mathsf{w}}(\tilde{\chi}_{z_1} \bar{\chi}_1(r))\right) \langle x^a \rangle^{-r} \in \mathcal{L}(L^2_s, L^2_t).$$

Clearly we can assume that s < t. We recall that the symbols $\chi_{z_1}\bar{\chi}_1(r)$, $\tilde{\chi}_{z_1}\bar{\chi}_1(r)$ as well as ξ^2 belong to a class for which the corresponding 'Planck constant' is $\langle x \rangle^{\rho/2-1}$. Using this fact we can now repeatedly commute the quantization of localized symbols through factors of $\check{R}(\lambda_0)$ to extract the desired decrease. The first step consists in noting that $\operatorname{Op}^{w}(\chi_{z_1}\bar{\chi}_1(r))\left(1 - \operatorname{Op}^{w}(\tilde{\chi}_{z_1}\bar{\chi}_1(r))\right)$ has arbitrary decrease, writing

$$[\operatorname{Op}^{\mathsf{w}}(\chi_{z_1}\bar{\chi}_1(r)), \check{R}(\lambda_0)] = \check{R}(\lambda_0)[\check{H}, \operatorname{Op}^{\mathsf{w}}(\chi_{z_1}\bar{\chi}_1(r))]\check{R}(\lambda_0)$$

and computing the commutator to the right to gain at least a factor $\langle x_a \rangle^{\rho/2-1}$. In the next step we make a similar commutation with the far right factor of $R(\lambda_0)$ of a similar localized operator thereby obtaining another factor $\langle x_a \rangle^{\rho/2-1}$. Repeating the argument we produce in this fashion efficiently any power of $\langle x_a \rangle^{\rho/2-1}$, and taking r sufficient big (depending on given $s, t \in \mathbb{R}$) we can efficiently produce any power $\langle x \rangle^{-m}$, in particular an $m \geq t - s$ as wanted.

We also shall need the following lemma which is based on an extension of parts of [DS1, Propositions 4.1 and 4.8] (cf. the extension for obtaining (4.9b) and Remark 6.23 given below) and the propagation of singularities result [DS1, Proposition 9.1] (in fact (6.81c) follows by from (6.81b) and [DS1, Proposition 9.1]). The underlying commutator methods are rather robust. In particular, even though the lemma as stated has a more qualitative flavour, one can demonstrate concrete bounds for the claimed embeddings, and the stability of those bounds can be used to control a parameter dependence, cf. [DS1, Remark 9.2]. In our application below the pair of angles (ω, ω') on which the kernel of the scattering matrix depends will play the role of such a parameter.

Lemma 6.22. Suppose $u \in L_s^{2,a} = L_s^{2,a}(\mathbf{X})$ for some $s < s_0$, and suppose that for some $t > s_0$ and $\kappa \in (-1, 1)$

$$WF_t^a(u) \cap \{b < \kappa, a = 1\} = \emptyset.$$
(6.81a)

Then the following assertions hold:

1) There exists the limit

$$r_{\lambda_0}^+ u := \lim_{R \to \infty} \left(1 \otimes r_{\lambda_0}^+ \right) \left(\chi_R(r) u \right) \text{ exists in } L^{2,a}_{s-2s_0}.$$

2) The intersection

$$WF^a_{t-2s_0}(r^+_{\lambda_0}u) \cap \{b < \kappa, a = 1\} = \emptyset,$$

$$(6.81b)$$

and (more generally)

$$WF^{a}_{t-2s_{0}}(r^{+}_{\lambda_{0}}u) \cap \{a=1\}$$

$$\subset \{\gamma(\tau,z\} \mid \tau \ge 0, \ z \in WF^{a}_{t}(u)\} \cup \{b=1\}.$$
(6.81c)

Remark 6.23. We note that (6.81b) may be proven as in the proof of [DS1, Proposition 4.8 (iii)], but that [DS1, (4.48] in fact holds with $\epsilon = 0$ (this generalization of [DS1, Proposition 4.1], originating from [FS], was noted before, cf. (4.9b)). Another generalization comes about noting that the condition on χ_+ in [DS1, Proposition 4.1 (v)] is too strong. In fact the relevant property is not that $\inf \operatorname{supp}(\chi_+) > C_0$ for some 'big' $C_0 \geq 1$, but rather $\chi_+(a) = 0$ in a neighbourhood of a = 1. This generalization can be proven by using a parametrix construction of [Sk4] (cf. [Sk4, (3.2)]) replacing the positivity argument of [FS]. Note that for the same reason we dont need the constant C_0 for the other parts of [DS1, Proposition 4.1] neither.

Now, to analyse $S_{\alpha,M}(\lambda_0)(\omega, \omega')$ we write (formally)

$$S_{\alpha,M}(\lambda_0)(\omega,\omega') = -2\pi \langle j^+_{\alpha,M}(\cdot,\omega), t^-_{\alpha,M}(\cdot,\omega') \rangle + 2\pi i \langle t^+_{\alpha,M}(\cdot,\omega), R(\lambda_0 + i0) t^-_{\alpha,M}(\cdot,\omega') \rangle,$$
(6.82)

where for $x \in \mathbf{X}$

$$j_{\alpha,M}^{\pm}(x,\omega) = (2\pi)^{-n/2} \left(e^{i\phi^{\pm}} e^{\tilde{\zeta}^{\pm}} a_{\alpha,M}^{\pm} \right) (x,\omega,0),$$

$$t_{\alpha,M}^{\pm}(x,\omega) = t_{\alpha,M}^{1\pm}(x,\omega) + \left(R_{s,M}^{k,a\pm} \delta_{\omega} \right) (x),$$

$$t_{\alpha,M}^{1\pm}(x,\omega) \coloneqq (2\pi)^{-n/2} \varphi_{\alpha}(x^{a}) \left(e^{i\phi^{\pm}} e^{\tilde{\zeta}^{\pm}} g_{\mathrm{bd},M}^{\pm} \right) (x_{a},\omega,0);$$
(6.83)

here the 'symbol' $a_{\alpha,M}^{\pm}$ is taken as a vector-valued function of $y = x_a$ and ω in agreement with Lemma 6.20, while ϕ^{\pm} , $\tilde{\zeta}^{\pm}$ and the functions $g_{\mathrm{bd},M}^{\pm}$ (as introduced in Lemma 6.20) have dependence of the component x_a of x and ω only (i.e. no dependence of x^a). Thus we write $J_{\alpha,M}^{\pm}(\lambda_0) \in J_{\alpha}^{\pm}(\lambda_0) + C_{2s_0}^{\pm}$ as

$$(J_{\alpha,M}^{\pm}(\lambda_0)\tau)(x^a, x_a) = \int_{\mathbb{S}^{n-1}} j_{\alpha,M}^{\pm}(x,\omega)\tau(\omega) \,\mathrm{d}\omega$$

= $(2\pi)^{-n/2} \int_{\mathbb{S}^{n-1}} \mathrm{e}^{\mathrm{i}\phi^+(y,\omega,0)} \mathrm{e}^{\tilde{\zeta}^+(y,\omega,0)} a_{\alpha,M}^{\pm}(x^a, y,\omega)\tau(\omega) \,\mathrm{d}\omega,$

which in fact applies to the Dirac delta function $\tau = \delta_{\omega} \in \bigcap_{k < -(n-1)/2} H^k(\mathbb{S}^{n-1})$ thereby defining the 'kernel' $j_{\alpha,M}^{\pm}$. Similarly $t_{\alpha,M}^{\pm}$ is the kernel

$$t_{\alpha,M}^{\pm}(x,\omega) = \left((2\pi)^{-n/2} \varphi_{\alpha} \otimes J_{g}^{\pm} \delta_{\omega} + R_{s,M}^{k,a\pm} \delta_{\omega} \right)(x); \quad g = g_{\mathrm{bd},M}^{\pm}$$

We note the following bounds, cf. [DS1, Theorem 6.5].

$$\partial_{\omega}^{\delta} j_{\alpha,M}^{\pm}(\cdot,\omega) \in L_t^{2,a} \text{ for all } t < -(|\delta| + n/2)(1 - \rho/2) - \rho/2,$$
 (6.84a)

$$\partial_{\omega}^{\delta} t_{\alpha,M}^{1\pm}(\cdot,\omega) \in L_t^{2,a} \text{ for all } t < -(|\delta| + n/2)(1 - \rho/2) + 1 + \rho/2, \tag{6.84b}$$

$$R_{s,M}^{k,a\pm}\partial_{\omega}^{\delta}\delta_{\omega} \in L_s^{2,a} \text{ provided } k < -|\delta| - (n-1)/2.$$
(6.84c)

Let ϕ_{sph}^+ denote the solution of the eikonal equation for the potential V_1 at zero energy, cf. (6.21). It is given by

$$\phi_{\rm sph}^+(x_a,\omega) = \frac{\sqrt{\gamma}}{1-\rho/2} \left(|x_a|^{1-\rho/2} \cos(1-\rho/2)\theta - R_0^{1-\rho/2} \right), \tag{6.85}$$

where $\cos \theta = \hat{x}_a \cdot \omega$. We omit the subscript *a* writing below (including the following lemma) *x* rather than x_a . Using $x^{\perp} = \frac{\omega - \hat{x} \cos \theta}{\sin \theta}$ and $\nabla_x \theta = -\frac{x^{\perp}}{r}$, we can also compute

$$F_{\rm sph}(x,\omega) = F_{\rm sph}^+(x,\omega) := \nabla_x \phi_{\rm sph}^+(x,\omega)$$
$$= \sqrt{\gamma} r^{-\rho/2} \left(\hat{x} \cos(1-\rho/2)\theta + x^{\perp} \sin(1-\rho/2)\theta \right).$$

The lemma stated below is a straightforward generalization of [DS1, Lemma 9.4], and as in [DS1] it follows by integration by parts. Note that due to (6.84c) only the contribution from the term $t_{\alpha,M}^{1\pm}$ of $t_{\alpha,M}^{\pm}$ in (6.83) matters. Note also that (6.84a) and (6.84b) provide an a priori 'size' of the quantatives $\partial_{\omega}^{\delta} j_{\alpha,M}^{\pm}(\cdot,\omega)$ and $\partial_{\omega}^{\delta} t_{\alpha,M}^{1\pm}(\cdot,\omega)$.

Lemma 6.24. Let k < -(n-1)/2, $s \ge 0$, $R \ge R_0$ and $0 < \sigma < \sigma' < \sigma_0$ (cf. Lemma 6.20). Then for all $M \in \mathbb{N}$ taken large enough and for all $\omega \in \mathbb{S}^{n-1}$ and multiindices δ with $|\delta| < -k - (n-1)/2$ the quantities in (6.83) obey

$$WF_{s}^{a}(\partial_{\omega}^{\delta}j_{\alpha,M}^{\pm}(\cdot,\omega)) \subset \left\{ z = (\hat{x},\bar{c},b) \in \mathbb{T}^{*} \mid (6.86a) \right\}$$

$$1 - \sigma' \leq \pm \hat{x} \cdot \omega, \quad b\hat{x} + \bar{c} = \pm F_{\rm sph}(\hat{x},\pm\omega)/\sqrt{\gamma} \right\},$$

$$WF_{s}^{a}(\partial_{\omega}^{\delta}t_{\alpha,M}^{\pm}(\cdot,\omega)) \subset \left\{ z = (\hat{x},\bar{c},b) \in \mathbb{T}^{*} \mid (6.86b) \right\}$$

$$1 - \sigma' \leq \pm \hat{x} \cdot \omega \leq 1 - \sigma, \quad b\hat{x} + \bar{c} = \pm F_{\rm sph}(\hat{x},\pm\omega)/\sqrt{\gamma} \right\}.$$

Suppose that $\chi_+ \in C^{\infty}(\mathbb{R})$, $\chi'_+ \in C^{\infty}_c(\mathbb{R})$ and $\operatorname{supp} \chi_+ \subset (1, \infty)$. Under the same conditions as above the following bounds hold uniformly in $\omega \in \mathbb{S}^{n-1}$:

$$\operatorname{Op}^{\mathsf{w}}(\chi_{+}(a))\partial_{\omega}^{\delta}j_{\alpha,M}^{\pm}(\cdot,\omega), \ \operatorname{Op}^{\mathsf{w}}(\chi_{+}(a))\partial_{\omega}^{\delta}t_{\alpha,M}^{\pm}(\cdot,\omega) \in L_{s}^{2,a}.$$
(6.86c)

Now we fix (a big) $\check{k} \in \mathbb{N}$ and want to apply Lemma 6.24 under the additional conditions

$$\check{k} < -k - (n-1)/2$$
 and $s \ge s_{\check{k}} := (\check{k} + \frac{n}{2})(1 - \frac{\rho}{2}) + 1;$ (6.87)

this requires M = M(k, s) taken large. We consider with $\chi_R = \chi_R(|x_a|)$ and $|\delta|, |\delta'| \leq \check{k}$

$$\begin{aligned}
\partial_{\omega}^{\delta} \partial_{\omega'}^{\delta'} S_{\alpha,M}(\lambda_0)(\omega, \omega') \\
&= -2\pi \lim_{R \to \infty} \langle \partial_{\omega}^{\delta} j_{\alpha,M}^+(\cdot, \omega), \chi_R \partial_{\omega'}^{\delta'} t_{\alpha,M}^-(\cdot, \omega') \rangle \\
&+ 2\pi i \lim_{R \to \infty} \langle \partial_{\omega}^{\delta} t_{\alpha,M}^+(\cdot, \omega), \chi_R R(\lambda_0 + i0) \chi_R \partial_{\omega'}^{\delta'} t_{\alpha,M}^-(\cdot, \omega') \rangle
\end{aligned}$$
(6.88)

This is still formal, but suppose that we can prove that the two limits on the righthand side exist locally uniformly in the set $\{\omega \cdot \omega' \neq \cos \frac{\rho}{2-\rho}\pi\}$ for all multi-indices with $|\delta|, |\delta'| \leq \check{k}$, then Theorem 6.17 follows.

Proof of Theorem 6.17. <u>I</u>. We need for any $\check{k} \in \mathbb{N}$ (henceforth fixed) to verify the existence and continuity of $\partial_{\omega}^{\delta} \partial_{\omega'}^{\delta'} S_{\alpha,M}(\lambda_0)(\omega, \omega')$ in $\{\omega \cdot \omega' \neq \cos \frac{\rho}{2-\rho}\pi\}$ for $|\delta|, |\delta'| \leq \check{k}$. We impose the conditions of Lemma 6.24 as well as (6.87). We also assume that σ' is small (to be tacitly used for (6.89) stated below).

We look at the first term to the right in (6.88) (before taking the limit), which we claim is treatable on $\mathbb{S}^{n-1} \times \mathbb{S}^{n-1}$ (i.e. without restriction on (ω, ω')). This can be seen by a direct integration by parts, but we prefer to give a presentation that conforms with our treatment of the second term (see (6.90)). Under the above conditions it follows from Lemma 6.24 and [DS1, (3.5c)] that for $|\delta|, |\delta'| \leq \check{k}$,

$$WF^{a}_{s_{\tilde{k}}}(\partial^{\delta}_{\omega}j^{+}_{\alpha,M}(\cdot,\omega)) \cap WF^{a}_{s_{\tilde{k}}}(\partial^{\delta'}_{\omega'}t^{-}_{\alpha,M}(\cdot,\omega')) = \emptyset.$$
(6.89)

Note also that the functions $\partial_{\omega}^{\delta} j_{\alpha,M}^{+}(\cdot,\omega)$ and $\partial_{\omega'}^{\delta'} t_{\alpha,M}^{-}(\cdot,\omega')$ are in $L_{-s_{\tilde{k}}}^{2,a}$ due to (6.84a)–(6.84c). Next we insert a suitable phase space partition of unity conforming with (6.89) in the expression

$$\langle \partial_{\omega}^{\delta} j_{\alpha,M}^{+}(\cdot,\omega), \chi_R \partial_{\omega'}^{\delta'} t_{\alpha,M}^{-}(\cdot,\omega') \rangle,$$

and then we remove the factor χ_R (by letting $R \to \infty$). The 'large *a*-part' is controlled by (6.86c) (using here only that $s \ge 0$) applying a single partition function $\chi_+(a)$. The 'small *a*-part', treated as indicated above, needs many partition functions using (6.89) and a sufficient 'sharpness' of the localization of the partition to make sure that for each term the partition operator brings at least one of the above two functions to $L_{s_k}^{2,a}$. Since the other function globally is in $L_{-s_k}^{2,a}$ the Cauchy-Schwarz inequality applies. This argument is first used with the factor χ_R in place. Any commutator with this factor gives at least an extra factor $R^{\rho/2-1}$, so in the limit we get (by dominated convergence) the corresponding expression without χ_R which by the same arguments indeed is well-defined. Finally the convergence is uniform in $(\omega, \omega') \in \mathbb{S}^{n-1} \times \mathbb{S}^{n-1}$ since all involved bounds can be done uniformly. This feature relies on regularity of the classical constructions and the underlying integration by parts arguments (the feature is only partially stated in Lemma 6.24).

<u>II</u>. We look at the second term to the right in (6.88), which we claim is well-defined with the limit taken locally uniformly in $\{\omega \cdot \omega' \neq \cos \frac{\rho}{2-\rho}\pi\}$. Before taking the limit we insert the expression (6.7) (for the plus case) and get various terms (possibly

after a further expansion) to treat when applied to the function $\partial_{\omega'}^{\delta'} t_{\alpha,M}^{-}(\cdot,\omega')$ to the right.

It is convenient to introduce the following notation.

$$\begin{split} u_{\omega}^{+} &= \partial_{\omega}^{\delta} t_{\alpha,M}^{+}(\cdot,\omega), \\ u_{\omega'}^{-} &= \partial_{\omega'}^{\delta'} t_{\alpha,M}^{-}(\cdot,\omega'), \\ \mathcal{M}_{\omega}^{+} &= \Big\{ z = (\hat{x}, \bar{c}, b) \in \mathbb{T}^{*} \mid 1 - \sigma' \leq \hat{x} \cdot \omega \leq 1 - \sigma, \ b\hat{x} + \bar{c} = F_{\mathrm{sph}}(\hat{x}, \omega) / \sqrt{\gamma} \Big\}, \\ \mathcal{M}_{\omega'}^{-} &= \Big\{ z = (\hat{x}, \bar{c}, b) \in \mathbb{T}^{*} \mid 1 - \sigma' \leq -\hat{x} \cdot \omega' \leq 1 - \sigma, \ b\hat{x} + \bar{c} = -F_{\mathrm{sph}}(\hat{x}, -\omega') / \sqrt{\gamma} \Big\}, \\ \mathcal{M}_{\omega'+}^{-} &= \Big\{ \gamma(\tau, z) \in \mathbb{T}^{*} \mid \tau \geq 0, \ z \in \mathcal{M}_{\omega'}^{-} \Big\} \cup \{a, b = 1\}. \end{split}$$

We follow the indicated scheme, so suppose \hat{R} is one of the terms of an expansion of $R(\lambda_0 + i0)$. Then we need to treat

$$\lim_{R \to \infty} \langle u_{\omega}^+, \chi_R \hat{R} \chi_R \, u_{\omega'}^- \rangle.$$

We first study the expression $\langle u_{\omega}^+, \chi_R \hat{R} \chi_R u_{\omega'}^- \rangle$ without the factors χ_R . We let $t_1 = s_{\check{k}}$ and intend to find $t_2 \in \mathbb{R}$ such that

$$WF_{t_1}^{a}(u_{\omega}^{+}) \cap WF_{t_2}^{a}(\hat{R}u_{\omega'}^{-}) = \emptyset,$$

$$u_{\omega}^{+} \in L^{2,a}_{-t_2}, \quad \hat{R}u_{\omega'}^{-} \in L^{2,a}_{-t_1}.$$
(6.90)

More precisely we shall use (6.90) for all such terms \hat{R} except for a certain 'remainder term' that is treated differently (see the end of the proof).

Given (6.90) we insert a suitable phase space partition of unity. The 'large *a*-part' is treated by using (6.86c) to u_{ω}^+ (note that $s \ge t_1$ so that the Cauchy-Schwarz inequality works). For the 'small *a*-part' the partition functions there are chosen such that for each term the partition operator either brings u_{ω}^+ to $L_{t_1}^{2,a}$ or $\hat{R}u_{\omega'}^-$ to $L_{t_2}^{2,a}$. In either case the Cauchy-Schwarz inequality works to make sense to the expression. These arguments can also be done with the factors χ_R in place, and we can also control the uniformity in the angles (cf. the discussion in Step I and a remark before Lemma 6.22).

The way to prove (6.90) goes as follows. Since $t_1 = s_{\check{k}} \leq s$ it follows from (6.86b) that

$$WF^a_{t_1}(u^+_{\omega}) \subset \mathcal{M}^+_{\omega}.$$
(6.91)

Suppose

$$\hat{R}u_{\omega'}^{-} \in L^{2,a}_{-t_1},$$
 (6.92a)

and that t_2 is chosen such that

$$WF^a_{t_2}(\hat{R}u^-_{\omega'}) \cap \{a=1\} \subset \mathcal{M}^-_{\omega'+}, \tag{6.92b}$$

$$u_{\omega}^{+} \in L^{2,a}_{-t_2}.$$
 (6.92c)

Then

$$WF_{t_1}^a(u_{\omega}^+) \cap WF_{t_2}^a(\hat{R}u_{\omega'}^-) \subset \mathcal{M}_{\omega}^+ \cap \mathcal{M}_{\omega'+}^-,$$

so to finish the proof of (6.90) we just need to check that the right-hand side is empty. It follows from [DS1, (3.5d)] that $\mathcal{M}^+_{\omega} \cap \{a, b = 1\} = \emptyset$. To complete the proof we note that

$$\forall z \in \mathcal{M}_{\omega}^{+}: \quad \hat{x}(\tau, z) \to \omega \text{ for } \tau \to +\infty,$$

$$\forall z \in \mathcal{M}_{\omega'+}^{-} \setminus \{a, b = 1\}: \quad \hat{x}(\tau, z) \to -\omega' \text{ for } \tau \to -\infty.$$

But if $\omega \cdot \omega' \neq \cos \frac{\rho}{2-\rho} \pi$ we can not find $z \in \mathbb{T}^*$ obeying these asymptotics. So indeed $\mathcal{M}^+_{\omega} \cap \mathcal{M}^-_{\omega'+} = \emptyset$ and (6.90) follows.

<u>III</u>. It remains to check (6.92a)–(6.92c) for the terms \hat{R} in an expansion of $R(\lambda_0 + i0)$. Note that the parameter t_2 may depend on the particular term \hat{R} we consider.

 $\frac{\hat{R} = Sr_{\lambda_0}^+ S^*}{\text{and } 6.24. \text{ Take } t_2 = s_{\check{k}} - 2s_0 \text{ (recall } s_0 = 1/2 + \rho/4\text{)}. \text{ From (6.84b) it follows that } u_{\omega}^+ \in L_{-t}^{2,a} \text{ for } t > t_2 - 1\text{, so in particular (6.92c) is proven. To show (6.92b) we note that <math>WF_{s_{\check{k}}}^a(u_{\omega'}^-) \subset \mathcal{M}_{\omega'}^-$, cf. (6.86b), and therefore the condition (6.81a) is fulfilled for $u = \Pi u_{\omega'}^-$ with $t = s_{\check{k}}$ and $\kappa = \kappa_{\sigma}$, where

$$\kappa_{\sigma} = -\cos\left((1-\rho/2)\cos^{-1}(1-\sigma)\right).$$

Then we learn from (6.81c) that

$$WF^a_{t_2}(\hat{R}u^-_{\omega'}) \cap \{a=1\} \subset \mathcal{M}^-_{\omega'+},$$

showing (6.92b). It follows from (6.84b) and Lemma 6.22 1) that $\hat{R}u_{\omega'}^{-} \in L^{2,a}_{-t_1}$, showing (6.92a).

 $\underline{\breve{R}}(\lambda_0)\Pi'. \quad \text{Take } t_2 = t_1 = s_{\breve{k}}. \text{ By (6.80c) } u_{\omega'}^-, \breve{R}(\lambda_0)\Pi' u_{\omega'}^- \in L^{2,a}_{-t_1}. \text{ From the previous case we know that } u_{\omega}^+ \in L^{2,a}_{-t_2} \text{ and } WF^a_{t_1}(u_{\omega'}^-) \subset \mathcal{M}^-_{\omega'}. \text{ Hence by (6.80d)}$

$$WF^a_{t_2}(\check{R}(\lambda_0)\Pi' u^-_{\omega'}) \cap \{a=1\} \subset WF^a_{t_2}(u^-_{\omega'}) \subset \mathcal{M}^-_{\omega'} \subset \mathcal{M}^-_{\omega'+1}$$

We have shown (6.92a)-(6.92c) in this case.

 $\underline{\check{R}}(\lambda_0 + \mathrm{i0}), cf. (6.8).$ Take $t_2 = s_{\check{k}} - 2s_0$ as we did treating $\hat{R} = Sr_{\lambda_0}^+ S^*$ above, so (6.92c) is known. We also know that the condition (6.81a) is fulfilled for $u = u_{\omega'}^-$ with $t = s_{\check{k}}$ and $\kappa = \kappa_{\sigma}$, i.e.

$$WF^a_{s_{\mathfrak{k}}}(u_{\omega'}) \cap \{b < \kappa, a = 1\} = \emptyset.$$

$$(6.93)$$

We also note that for any $L \in \mathbb{N}$ (eventually taken large) and with

$$K = v_{\lambda_0} r_{\lambda_0}^+ = \left(-S^* I_a \breve{R}(\lambda_0) \Pi' I_a S + \mathcal{O}(r^{-1-\rho}) \right) r_{\lambda_0}^+$$

$$Sr_{\lambda_0}^+ (1 + v_\lambda r_\lambda^+)^{-1} S^* = \sum_{l=0}^{2L-1} Sr_{\lambda_0}^+ (-K)^l S^* + Sr_{\lambda_0}^+ (-K)^L (1+K)^{-1} (-K)^L S^*$$

Note that the present formula $v_{\lambda}^{+} = v_{\lambda}^{-}$ gives naturally raise to the unambiguous notation v_{λ} . Moreover the term $\mathcal{O}(r^{-1-\rho})$ is actually $\mathcal{O}(r^{-\infty})$ since $m_{a} = 1$ making $v_{\lambda_{0}} = \mathcal{O}(r^{-2-2\rho})$ (as discussed after (6.5)), however only the decay $v_{\lambda_{0}} = \mathcal{O}(r^{-1-\rho})$ is used below.

We will treat the terms in the summation essentially by the methods used above, while the last term will be treated differently. Let us first examine the contribution to $\check{R}(\lambda + i0)$ from the terms in the summation.

l=0. We consider

$$\widehat{R}_0 := -Sr^+_{\lambda_0}S^*I_a\breve{R}(\lambda_0)\Pi' - \breve{R}(\lambda_0)\Pi'I_aSr^+_{\lambda_0}S^*(1 - I_a\breve{R}(\lambda_0)\Pi').$$

By Lemma 6.22, (6.93) and (6.80a)–(6.80d) it follows that $\widehat{R}_0 u_{\omega'}^- \in L^{2,a}_{-t_1}$, i.e. (6.92a) holds for this contribution. (This is a rough bound due to the extra factor I_a .)

Again we learn from (6.81c) (cf. (6.93)) that

$$WF^a_{t_2}(R_0u^-_{\omega'}) \cap \{a=1\} \subset \mathcal{M}^-_{\omega'+},$$

showing (6.92b).

 $l \geq 1$. We consider

$$\widehat{R}_l := \left(1 - \breve{R}(\lambda_0)\Pi' I_a\right) Sr^+_{\lambda_0} (-K)^l S^* \left(1 - I_a \breve{R}(\lambda_0)\Pi'\right).$$

Recalling that $WF_{t_1}^a(u_{\omega'}^-) \subset \mathcal{M}_{\omega'}^-$ we are lead to investigate the action of $Sr_{\lambda_0}^+(-K)^l S^*$ to a vector $u \in L^{2,a}_{-\infty}$ with $WF_{t_1}^a(u) \subset \mathcal{M}_{\omega'}^-$. Each factor of K improves the weight by $\langle x \rangle^{2s_0-1-\rho} = \langle x \rangle^{-q}$, $q := \rho/2$, more precisely we obtain from Lemma 6.22 that

$$WF^{a}_{t_{1}+lq}(S(-K)^{l}S^{*}u) \cap \{a=1\} \subset \mathcal{M}^{-}_{\omega'+} \subset \{b \geq \kappa_{\sigma}, a=1\}, \\WF^{a}_{t_{2}+lq}(Sr^{+}_{\lambda_{0}}(-K)^{l}S^{*}u) \cap \{a=1\} \subset \mathcal{M}^{-}_{\omega'+}.$$

So we learn that

$$WF^a_{t_2+lq}(\widehat{R}_l u^-_{\omega'}) \cap \{a=1\} \subset \mathcal{M}^-_{\omega'+}, \tag{6.94}$$

in particular (6.92b) holds.

By Lemma 6.22 1) the global weight is improved by a factor $\langle x \rangle^{-q}$ for each action by K as long as the condition $s' < -s_0$ of the lemma is fulfilled (note that $L^{2,a}_{-s_0}$ can not be reached by the action by $r^+_{\lambda_0}$). Since $l \ge 1$ and $u^-_{\omega'} \in L^2_{-t_2}$ we then conclude that $\widehat{R}_l u^-_{\omega'} \in L^2_{-t_1}$, and hence also (6.92a) is shown.

<u>Remainder term</u>. Finally we need to examine

$$\widehat{R} := \left(1 - \breve{R}(\lambda_0)\Pi' I_a\right) Sr_{\lambda_0}^+ (-K)^L \left(1 + K\right)^{-1} (-K)^L S^* \left(1 - I_a \breve{R}(\lambda_0)\Pi'\right).$$

The operator $(1+K)^{-1} = 1 \otimes (1+K)^{-1} \in \mathcal{L}(L_s^2(\mathbf{X})), s \in (s_0, s_0+q)$, but we dont know its microlocal properties. Whence we proceed differently by introducing the vectors

$$v_{\omega'}^{-} = S(-K)^{L} S^{*} (1 - I_{a} \breve{R}(\lambda_{0}) \Pi') u_{\omega'}^{-},$$

$$v_{\omega}^{+} = S ((1 - \breve{R}(\lambda_{0}) \Pi' I_{a}) S r_{\lambda_{0}}^{+} (-K)^{L})^{*} u_{\omega}^{+}$$

and write

$$\langle u_{\omega}^{+}, \widehat{R}u_{\omega'}^{-} \rangle = \langle v_{\omega}^{+}, \left(1 + K\right)^{-1}v_{\omega'}^{-} \rangle.$$

It suffices to show that for some $s \in (s_0, s_0 + q)$ and L sufficiently large

$$v_{\omega'}^- \in L_s^{2,a}, \quad v_{\omega}^+ \in L_{-s}^{2,a}.$$
 (6.95)

To obtain the result for $v_{\omega'}^- \in L_s^{2,a}$ we use Lemma 6.22 repeatedly as explained above improving the global weight by a factor $\langle x \rangle^{-q}$ for each action by K. It follows that $v_{\omega'}^- \in L_s^{2,a}$ holds for any $s \in (s_0, s_0 + q)$ for L large enough. For the assertion $v_{\omega}^+ \in L_{-s}^{2,a}$ we invoke a parallel 'incoming' version of Lemma 6.22 for $r_{\lambda_0}^-$ (for simplicity not stated). Note that powers of $r_{\lambda_0}^-$ show up when we expand the adjoint to treat v_{ω}^+ . Since the construction of u_{ω}^+ and $u_{\omega'}^-$ appears symmetric indeed the incoming version of Lemma 6.22 applies in the same fashion to u_{ω}^+ as we have seen the lemma applies to $u_{\omega'}^-$.

6.1.3 Scattering for physics models at a two-cluster threshold, case $\widetilde{\mathcal{A}} = \mathcal{A}_1$

In the previous subsections we made several simplifying assumptions under the condition of an effective attractive slowly decaying inter-cluster potential. These were made partly for simplicity of presentation. Rather than giving a full account on how to remove these assumptions under the weakest possible conditions we shall here focus on the models of physics which we studied in Section 4.3 (and actually we shall not give details of proof below). This means that the effective inter-cluster potential here is attractive Coulombic, so in agreement with Section 4.3 we consider the case $\tilde{\mathcal{A}} = \mathcal{A}_1$. (Note that this condition includes cases of overall neutral as well as overall non-neutral systems of particles.) We shall focus on treating the multiple two-cluster case, but this will only be in the 'generic' situation studied in the first part of the proof of Theorem 4.17. Whence our main interest here is scattering for the physics models with $\tilde{\mathcal{A}} = \mathcal{A}_1$ in the simplest possible multiple case.

We consider the setting of Theorem 4.17 with $\mathcal{A} = \mathcal{A}_1$ under the two 'generic' conditions (4.68a) and (4.68b). Let us also for simplicity assume that $\#\mathcal{A}_1 = 2$, i.e. $\mathcal{A}_1 = \{a_1, a_2\}$ as in Proposition 2.12, and as for the latter result we assume for convenience that λ_0 is a simple eigenvalue for H^{a_j} ; j = 1, 2. We impose (6.3), again for simplicity of presentation only.

We are interested in the four parts of the scattering operator $S_{\beta\alpha} = (W_{\beta}^{+})^* W_{\alpha}^{-}$ defined by (6.39), where $\alpha, \beta \in \{(a_j, \lambda_0, \varphi_j) \mid j = 1, 2\}$, and particularly in the corresponding pieces of the scattering operator

$$S_{\beta\alpha}(\lambda) = -2\pi W_{\beta}^{+}(\lambda)^{*}T_{\alpha}^{-}(\lambda), \quad \lambda \in I_{\delta}^{+} = [\lambda_{0}, \lambda_{0} + \delta],$$

cf. (6.47).

We can prove a complete analogue of Theorem 6.10. Note that for each channel there is associated an effective one-body potential denoted by w_j (or w_{α}), which essentially is attractive Coulombic. We define correspondingly $\check{S}_{\alpha\alpha}(\lambda)$ as the remainder after the one-body scattering matrix is subtracted, exactly as in Theorem

184

6.10. Similarly we may define $\check{S}_{\beta\alpha}(\lambda) = S_{\beta\alpha}(\lambda)$ for $\alpha \neq \beta$. With these conventions we obtain the assertion of Theorem 6.10 (with the same value of k) for all of the four entries of $\check{S}_{\beta\alpha}(\lambda)$. The proof is essentially the same. In particular the off-diagonal parts of the scattering matrix are compact, while the same is the case for the diagonal parts only after a subtraction of a unitary operator, and this includes the threshold energy.

Corollary 6.25. Under the above conditions the diagonal parts $S_{\alpha\alpha}(\lambda_0)$ are unitary up to a compact term. In particular elastic two-cluster scattering exists in the small inter-cluster energy regime.

It is not known, even with the above simplifying assumptions, if $S_{\alpha\alpha}(\lambda_0)$ is an exact unitary operator (we can in fact not exclude that ker $S_{\alpha\alpha}(\lambda_0) \neq \{0\}$), cf. Corollary 6.15 and a brief discussion in Section 6.3. In particular this is not known for $\lambda_0 = \Sigma_2$ and this threshold being multiple (note that in this case the threshold is automatically a simple eigenvalue of the involved sub-Hamiltonians). On the other hand obviously the scattering matrix $S_{\alpha\alpha}(\lambda_0)$ in Theorem 6.10 (where λ_0 is non-multiple) is an exact unitary operator for $\lambda_0 = \Sigma_2$, cf. Remark 6.16.

As for the second main result Theorem 6.17 we need an analogue of Theorem 6.13 so that we can first 'improve' the construction of $S_{\beta\alpha}(\lambda_0)$ in the spirit of Lemma 6.20, not to be elaborated on. By mimicking Subsection 6.1.2.4 we then obtain the following result.

Theorem 6.26. Under the above conditions and with $\lambda_0 = \Sigma_2$

sing supp
$$S_{\beta\alpha}(\lambda_0)$$

$$\begin{cases} \subset \{(\omega, \omega') \mid \omega \cdot \omega' = -1\} & \text{for} \quad \alpha = \beta, \\ = \emptyset & \text{for} \quad \alpha \neq \beta. \end{cases}$$

6.2 Effective r^{-2} potentials, atom-ion case

One may consider elastic scattering in the non-slowly decaying case, as before slightly above a given two-cluster threshold λ_0 . How does $S_{\alpha\alpha}(\lambda)$ behave as $\lambda \to \lambda_0$?

Suppose n = 3 and consider the dynamical nuclei model, see Subsection 1.2.1. Consider a two-cluster decomposition $a = a_0 = (C_1, C_2)$ of N charged particles. Suppose (6.68), i.e.

$$\lambda_0 \in \sigma_{\rm d}(H^a),$$

and that the total charge of cluster C_1 vanishes:

$$Q_1 = \sum_{j \in C_1} q_j = 0$$

Let $\alpha = (a, \lambda_0, \varphi_\alpha)$ be a channel; $\varphi_\alpha = \varphi^{C_1} \otimes \varphi^{C_2} = \varphi^1 \otimes \varphi^2$. Recall from Subsection 1.2.1 the cluster charge moment

$$\langle \varphi^1, \widetilde{Q}_1 \varphi^1 \rangle_2 = \left\langle \varphi^1, \sum_{j \in C_1} q_j (x_j - R_1) \varphi^1 \right\rangle.$$

Now we can state a result from [Sk2] (see [Sk2, Theorem 2.6 (3)]):

Theorem 6.27 (Elastic two-cluster scattering away from thresholds). 1) There exist

$$W^{\pm}_{\alpha}f = \lim_{t \to \pm \infty} e^{itH} e^{-itH_a} (\varphi_{\alpha} \otimes f).$$

2) Let $S_{\alpha\alpha} = (W_{\alpha}^{+})^{*}W_{\alpha}^{-}$ and $I_{c} = (\lambda_{0}, \infty) \setminus \mathcal{T}$. Let θ denote the coordinate vector of $\omega - \omega' \in \mathbf{X}_{a}$. Then for any $\epsilon > 0$

$$S_{\alpha\alpha} \simeq \int_{I_c}^{\oplus} S_{\alpha\alpha}(\lambda) \, \mathrm{d}\lambda,$$

$$S_{\alpha\alpha}(\lambda, \cdot) \in C^{\infty}(I_c, \left(\mathbb{S}^2 \times \mathbb{S}^2\right) \setminus \{\theta = 0\}),$$

$$S_{\alpha\alpha}(\lambda, \omega, \omega') - \delta(\theta) + \pi^{-1} \frac{M_1 M_2}{M_1 + M_2} Q_2 \langle \varphi^1, \widetilde{Q}_1 \varphi^1 \rangle_2 \cdot \frac{\theta}{|\theta|^2} = \mathcal{O}_{\lambda}(|\theta|^{-\epsilon}).$$

The latter bound is λ -dependent with a locally uniform dependence. A natural question is, if (under conditions) there is a uniform bound on an interval $(\lambda_0, \lambda_0 + \delta)$, $\delta > 0$? This question depends on the resolvent behaviour at λ_0 , and Chapter 5 provides some information of relevance for this type of problem. A related problem is, if low-energy elastic scattering *exists*, cf. Corollary 6.25. In the next section we address the latter problem, not only for the above case (for which $I_a(x^a = 0) = 0$) but also for the effective repulsive Coulombic case.

6.3 Non-transmission at a threshold, physics models

We consider a non-transmission problem for the physics models of Sections 1.2 and 1.3 with n = 3. Although the statement of the problem makes sense more generally we restrict the attention to these models.

Mimicking (6.39) we introduce for a given channel $\alpha = (a, \lambda_0, \varphi_\alpha)$ (at a given two-cluster threshold λ_0) the channel wave operators

$$W_{\alpha}^{\pm}f = \lim_{t \to \pm \infty} e^{itH} \left(1 \otimes J_{0}^{\pm} \right) \varphi_{\alpha} \otimes e^{-it(p_{a}^{2} + \lambda_{0})} f$$

$$= \lim_{t \to \pm \infty} e^{itH} \left(\varphi_{\alpha} \otimes J_{0}^{\pm} e^{-it(p_{a}^{2} + \lambda_{0})} f \right); \quad \hat{f} \in C_{c}^{\infty}(\mathbf{X}_{a} \setminus \{0\}).$$
(6.96)

In the effective Coulombic cases the appearing stationary modifiers J_0^{\pm} are chosen as in (6.24b) and (6.39) (i.e. also this way for the repulsive case), and in the effective non-Coulombic case (like in Theorem 6.27) we take $J_0^{\pm} = 1$. Let $\Pi^{\alpha} = |\varphi_{\alpha}\rangle\langle\varphi_{\alpha}| \otimes 1$ and

$$\Pi^{\alpha}_{\pm} = \operatorname{s-lim}_{t \to \pm \infty} \mathrm{e}^{\mathrm{i}tH} \Pi^{\alpha} \mathrm{e}^{-\mathrm{i}tH} \mathbb{1}_{\mathbb{R} \setminus \sigma_{\mathrm{pp}}(H)}(H).$$

We are interested in 'transmission' (for example 'exchange', cf. [CT]), or rather lack of transmission between channels with the energy constraint of localization slightly above λ_0 . Such lack of transmission may be phrased mathematically as

$$\left\| \left(1 - \Pi_{+}^{\alpha} \right) \Pi_{-}^{\alpha} F_{\delta}(H - \lambda_{0}) \right\| \to 0 \text{ for } \delta \to 0_{+}.$$
(6.97)

186

6.3. Non-transmission at a threshold, physics models

Here F_{δ} denotes the characteristic function $F([0, \delta])$ on \mathbb{R} .

Note that a state $\psi = 1_{\mathbb{R}\setminus\sigma_{\mathrm{pp}}(H)}(H)\psi$ is in the range of the projection Π^{α}_{\pm} if and only if $(1 - \Pi^{\alpha})\psi(t) \to 0$ for $t \to \pm\infty$; $\psi(t) := \mathrm{e}^{-\mathrm{i}tH}\psi$. Intuitively a consequence of (6.97) is that if for a state ψ localized in energy slightly above λ_0 and effectively $\psi(t) \approx \Pi^{\alpha}\psi(t)$ for $t \to -\infty$ then also $\psi(t) \approx \Pi^{\alpha}\psi(t)$ for $t \to +\infty$. Whence, asymptotically in the low energy regime, only elastic scattering occurs and transmission to another channel than (the incoming) α does not occur.

Clearly $\Pi^{\alpha}_{\pm}W^{\pm}_{\alpha} = W^{\pm}_{\alpha}$, and by asymptotic completeness (cf. [De3]) $\Pi^{\alpha}_{\pm} \subset W^{\pm}_{\alpha}(W^{\pm}_{\alpha})^*$. Whence

$$\Pi^{\alpha}_{\pm} = W^{\pm}_{\alpha} \left(W^{\pm}_{\alpha} \right)^*.$$

Consequently we can compute

$$\left(1 - \Pi_+^{\alpha}\right)\Pi_-^{\alpha}F_{\delta}(H - \lambda_0) = \left(W_{\alpha}^- - W_{\alpha}^+ S_{\alpha\alpha}\right)F_{\delta}(p_a^2)\left(W_{\alpha}^-\right)^*.$$

Now

$$\| (W_{\alpha}^{-} - W_{\alpha}^{+} S_{\alpha\alpha}) f \|^{2} = \| f \|^{2} - \| S_{\alpha\alpha} f \|^{2},$$

showing with the above identity that a necessary and sufficient condition for (6.97) is that $S_{\alpha\alpha}(\lambda)$ is asymptotically isometric, that is

$$\|1 - S_{\alpha\alpha}(\lambda)^* S_{\alpha\alpha}(\lambda)\| \to 0 \text{ for } \lambda \to (\lambda_0)_+.$$
(6.98)

Note that in the context of Corollary 6.25 it follows from the assertion that the limiting diagonal parts $S_{\alpha\alpha}(\lambda_0) \neq 0$, however it is *not stated* that $S_{\alpha\alpha}(\lambda_0)$ are isometric, and we can not show this to be the case. See Remark 6.16 for a conjecture on transmission for the attractive Coulombic case.

We consider the physical models with n = 3, and for convenience we impose Condition 2.4 with $a_0 = a$, i.e. we consider only the non-multiple case. Then there are three cases for which we can show (6.98) (or equivalently (6.97)):

- I) Effective repulsive Coulombic case.
- **II)** $I_a(x^a = 0) = 0$, 'above the Hardy limit' and λ_0 be 'regular'.
- **III**) $I_a(x^a = 0) = 0$, 'fastly decaying case' and λ_0 be 'maximally exceptional of 1st kind'.

Note that I) corresponds to Case 1 in Sections 1.2 and 1.3 (with an additional sign condition for charges). II) corresponds to Cases 2 and 3 in the same sections, while III) corresponds to Case 3 only and includes the occurrence of resonance states (but not bound states). The terminology 'above the Hardy limit' is explained in the simple case $m_a = 1$ in Subsection 6.3.2, see (6.106), while 'regular' as in Section 5.1.1 refers to λ_0 not be neither an eigenvalue nor a resonance. In general the condition 'above the Hardy limit' is a spectral condition for the $m_a \times m_a$ matrixvalued effective potential $S_a^*I_aS_a = Q_a|x_a|^{-2} + B_a$, or rather for its leading term $Q_a(\hat{x}_a)$. By definition it means that the eigenvalues of $-\Delta_{\theta} + Q_a(\theta)$ are all strictly bigger than -1/4, cf. Section 4.3, and of course this condition is fulfilled for Case 3 (where $Q_a = 0$), but it might not be fulfilled in Case 2. The condition 'exceptional of 1st kind' refers to λ_0 be a resonance but not an eigenvalue, cf. Section 5.1.1. The added word 'maximally' refers to maximal multiplicity of the space of resonance states, i.e. $n_{\rm res} = m_a$. Note that the fastly decaying regular case is included in Case II).

We summarize our results as follows.

Theorem 6.28. For the physics models with n = 3 and with Condition 2.4 fulfilled at λ_0 for $a_0 = a$ there is no transmission assuming I), II) or III). Whence for each of these cases there is no transmission in the small inter-cluster energy regime from any given (incoming) channel $\alpha = (a, \lambda_0, \varphi_\alpha)$ to any different (outgoing) channel, *i.e.* (6.97) and (6.98) are fulfilled.

Our results are proven in subsequent subsections under the following additional (convenient but non-essential) simplifying conditions:

- a) $m_a = 1$.
- b) $\lambda_0 \notin \sigma_{\rm pp}(H) \cup \sigma_{\rm pp}(H').$
- **Remarks 6.29.** 1) We dont know if (6.98) holds in full generality for the fastly decaying case and λ_0 be an exceptional point of 2nd or 3rd kind, although we have results with a decay condition for the latter cases (for which $\lambda_0 \in \sigma_{\rm pp}(H)$). The proof for III) is based on Theorem 5.45 which applies under the decay condition ran $\Pi_H \subset L_t^2$ for some t > 3/2, where Π_H denotes the orthogonal projection onto ker $(H \lambda_0)$. Hence under this additonal condition, III) can be extended to cover the 'maximally exceptional case of 3rd kind'. Note that the (local) scattering theory of H and $H_{\sigma} := H \sigma \Pi_H$, $\sigma > 0$, are identical, and that λ_0 in this case is a 'maximally exceptional point of 1st kind' of H_{σ} .
 - 2) Similarly we dont know in full generality if the regularity condition in II) is necessary for the critical decay rate case. However for the 'exceptional case of 2nd kind' we can use Theorem 5.44 for which the decay condition ran $\Pi_H \subset L_t^2$ for some t > 1 is needed. Note that λ_0 is a regular point of H_{σ} in this case.
 - 3) If $m_a > 1$ and the word 'maximally' is omitted in Case III), i.e. $n_{\text{res}} \in \{1, \ldots, m_a 1\}$, we show in Subsection 6.3.4 that transmission does occurs.
 - 4) A key ingredient of our proof of Theorem 6.28 is various low-energy bounds on the one-body spectral density operator. These are absent for the effective attractive Coulombic case.

6.3.1 Proof of (6.98), Case I)

We consider the Case I), i.e. the repulsive Coulombic case. We use Subsections 4.1.3 and 4.2.3 and introduce a globally positive function \widehat{W} as in (4.16) with $\rho = \overline{\rho} = 1$. More precisely we here consider the non-multiple case version discussed in Remark 6.1, imposing the above conditions a) and b). As in Subsection 4.1.3 we then take $w = \widehat{W}$. We define $r_{\lambda}^{\pm} = (h_a + \lambda_0 - \lambda \mp i0)^{-1}$, $h_a = p_a^2 + w$ and $\lambda \ge \lambda_0$, and recall the bound

$$\|\langle x \rangle^{-s} r_{\lambda}^{\pm} \langle x \rangle^{-s} \| \le C, \quad s > s_0 = \frac{1}{2} + \frac{\rho}{4},$$
 (6.99)

cf. [Na] and [Ya2]. Recall also that $r_{\lambda_0}^+ = r_{\lambda_0}^-$.

Next we introduce relative wave operators

$$\Omega_{\alpha}^{\pm}f = \lim_{t \to \pm \infty} e^{itH} e^{-it(H_a + w)} (\varphi_{\alpha} \otimes f)$$

and the corresponding scattering operator $(\Omega_{\alpha}^{+})^{*}\Omega_{\alpha}^{-}$.

Using one-body Isozaki-Kitada scattering theory we can write

$$\delta(h_a - \lambda + \lambda_0) = W_1^{\pm}(\lambda - \lambda_0)W_1^{\pm}(\lambda - \lambda_0)^*, \qquad (6.100)$$

cf. [DS1]. Here $W_1^{\pm}(\cdot)$ signify wave matrices for the pair (p_a^2, h_a) corresponding to Isozaki-Kitada wave operators W_1^{\pm} , cf. [DS1]. Clearly the wave operators $W_{\alpha}^{\pm} = \Omega_{\alpha}^{\pm}W_1^{\pm}$. Let us use W_1^- to diagonalize h_a . Whence we introduce the 'identification operator' $J_{\alpha} = \varphi_{\alpha} \otimes (W_1^- \cdot)$ and the corresponding relative wave operators

$$\widetilde{W}^{\pm}_{\alpha} = \operatorname{s-lim}_{t \to \pm \infty} \, \mathrm{e}^{\mathrm{i}tH} J_{\alpha} \mathrm{e}^{-\mathrm{i}t(p_a^2 + \lambda_0)}.$$

Note that $\widetilde{W}_{\alpha}^{-} = W_{\alpha}^{-}$, while $\widetilde{W}_{\alpha}^{+} = W_{\alpha}^{+}S_{1}$ with S_{1} being the scattering operator for the pair (p_{a}^{2}, h_{a}) . Whence

$$S_{\alpha\alpha} = \left(W_{\alpha}^{+}\right)^{*} W_{\alpha}^{-} = S_{1} \left(\widetilde{W}_{\alpha}^{+}\right)^{*} \widetilde{W}_{\alpha}^{-} = S_{1} \widetilde{S}_{\alpha\alpha},$$

yielding upon diagonalizing $p_a^2 + \lambda_0$ by the Fourier transform

$$S_{\alpha\alpha}(\lambda) = S_1(\lambda - \lambda_0)\widetilde{S}_{\alpha\alpha}(\lambda).$$

(A similar factorization formula appears for a one-body problem in [Ba].) We can represent, cf. [DS1, Appendix A] or [Ya3, Section 7.3],

$$\widetilde{S}_{\alpha\alpha}(\lambda) = 1 - 2\pi J_{\alpha}(\lambda)^* T_{\alpha}(\lambda) + 2\pi \mathrm{i} T_{\alpha}(\lambda)^* R(\lambda + \mathrm{i}0) T_{\alpha}(\lambda), \qquad (6.101)$$

where $J_{\alpha}(\lambda) = \varphi_{\alpha} \otimes W_1^-(\lambda - \lambda_0)$ and $T_{\alpha}(\lambda) = i(H - \lambda_0)J_{\alpha}(\lambda) = i(I_a - w)J_{\alpha}(\lambda)$. Note that formally $J_{\alpha}(\lambda) = J_{\alpha}\mathcal{F}_{\lambda_0}(\lambda)^*$, where $\mathcal{F}_{\lambda_0}(\lambda) = \mathcal{F}_0(\lambda - \lambda_0)$ with $\mathcal{F}_0(\cdot)$ specified by (6.31). Note also that Taylor expansion yields $T_{\alpha}(\lambda) \approx \mathcal{O}(\langle x \rangle^{-2} \langle x^a \rangle^3)J_{\alpha}(\lambda) \approx \mathcal{O}(\langle x \rangle^{-2})J_{\alpha}(\lambda)$.

Now we can verify (6.98). By inserting (6.101) into (6.98) it suffices to show that

$$\|J_{\alpha}(\lambda)^* T_{\alpha}(\lambda)\| \to 0, \qquad (6.102a)$$

$$\|T_{\alpha}(\lambda)^* R(\lambda + i0) T_{\alpha}(\lambda)\| \to 0.$$
(6.102b)

To show (6.102a) we note the bound

$$\|\delta(h_a - \lambda + \lambda_0)\|_{\mathcal{L}(L^2_s, L^2_{-s})} = o((\lambda - \lambda_0)^0) \text{ for } s > s_0,$$
(6.103)

cf. [Na]. We use it with s = 1 in combination with (6.100), proving (6.102a).

To derive (6.102b) it suffices by the same argument to show the bound

$$\sup_{\lambda \in I_{\delta}^{+}} \|\langle x \rangle^{-1} R(\lambda + \mathrm{i}0) \langle x \rangle^{-1} \| < \infty,$$

which in turn follows from (6.7), (6.8) (cf. Remark 6.1) and (6.99).

6.3.2 Proof of (6.98), Case II)

For Case II) the wave operators are defined by (6.96) with $J_0^{\pm} = 1$. Note the following formula for the scattering matrix for λ slightly above λ_0 ,

$$S_{\alpha\alpha}(\lambda) = 1 - 2\pi J_{\alpha}^{+}(\lambda)^{*} T_{\alpha}^{-}(\lambda) + 2\pi \mathrm{i} T_{\alpha}^{+}(\lambda)^{*} R(\lambda + \mathrm{i} 0) T_{\alpha}^{-}(\lambda), \qquad (6.104)$$

where $J_{\alpha}^{+}(\lambda) = J_{\alpha}^{-}(\lambda) = \varphi_{\alpha} \otimes \mathcal{F}_{\lambda_{0}}(\lambda)^{*} =: J_{\alpha}(\lambda), \ \mathcal{F}_{\lambda_{0}}(\lambda) = \mathcal{F}_{0}(\lambda - \lambda_{0})$ and similarly $T_{\alpha}^{+}(\lambda) = T_{\alpha}^{-}(\lambda) = \mathrm{i}I_{a}J_{\alpha}(\lambda) =: T_{\alpha}(\lambda).$

As in Theorem 6.10 we subtract a one-body scattering matrix introducing

$$\check{S}_{\alpha\alpha}(\lambda) = S_{\alpha\alpha}(\lambda) - S_w(\lambda - \lambda_0);$$

here w is a reel effective one-body potential to be determined. Since we know that $S_w(\lambda - \lambda_0)$ is unitary we aim at showing that

$$\|\check{S}_{\alpha\alpha}(\lambda)\| \to 0 \text{ for } \lambda \to (\lambda_0)_+.$$
 (6.105)

To examine this problem we need a formula for $R(\lambda + i0)$ like the one we used in the proof of Theorem 6.10. In agreement with the condition of Case II) λ_0 is neither a resonance nor an eigenvalue of H and the 'above the Hardy limit' condition

$$\nu_0 := \min_{\nu \in \sigma} \operatorname{Re} \nu > 0 \tag{6.106}$$

is imposed. Here the set σ is given as in the analogous formula (4.25) for the multiple case, that is computed by the eigenvalues of a perturbed Laplace-Beltrami operator $-\Delta_{\theta} + q(\theta)$ as in (4.21). Alternatively stated

$$\inf \sigma \big(-\Delta_{\theta} + q(\theta) \big) > -1/4.$$

We assume that $m_a = 1$ and that $\lambda_0 \notin \sigma_{\rm pp}(H')$, cf. the conditions a) and b) stated after Theorem 6.28.

Under the condition (6.106) one could hope that the operator $(p_a^2 + W + \lambda_0 - \lambda \mp i0)^{-1}$ with $W := \langle \varphi_{\alpha}, I_a \varphi_{\alpha} \otimes \cdot \rangle$ would be a good auxillary operator for stydying asymptotics as $\lambda \to (\lambda_0)_+$, cf. Subsections 4.1.4 and 4.2.4. However zero could be an eigenvalue or a resonance for $p_a^2 + W$, spoiling this idea. To avoid this problem we are lead to modifying $W(x_a)$ by a suitable stronger decaying term (the perturbation will be of order $\mathcal{O}(|x_a|^{-3})$) to assure zero be regular for the auxillary operator. More precisely we introduce, using the same quadratic partition of unity as in (4.24),

$$h_{a} = \chi_{1}(r)p_{a}^{2}\chi_{1}(r) + \chi_{2}(r)\left(p_{a}^{2} + \frac{q(\theta)}{r^{2}}\right)\chi_{2}(r) = p_{a}^{2} + w,$$

$$w = |\nabla\chi_{1}|^{2} + |\nabla\chi_{1}|^{2} + \frac{q(\theta)}{r^{2}}\chi_{2}^{2} = W + \mathcal{O}(r^{-3}),$$

$$r_{\lambda}^{\pm} = (h_{a} - k^{2} \mp i0)^{-1}, \quad k \in (0, 1], \quad \lambda = \lambda_{0} + k^{2}.$$

Due to (6.106) $h_a \ge 0$, and in fact h_a is strictly positive (i.e. $h_a \ge 0$ and $0 \notin \sigma_{pp}(h_a)$).

We will show that h_a does not have a resonance at zero by an integration by parts argument, cf. [Ji], and we will show a number of properties of r_{λ}^{\pm} by a parametrix construction. In addition we will need properties of the restriction of the Fourier transform $\mathcal{F}_0(k^2)$. <u>1</u>. Let, in analogy with the zero-energy formulas (4.24),

$$G_k^{\pm} = \chi_1(r)(h_a \mp i)^{-1}\chi_1(r) + \sum_{\nu \in \sigma} \chi_2(r)r^{-1}R_{\nu,k}^{\pm}r\chi_2(r) \otimes P_{\nu};$$

$$R_{\nu,k}^{\pm} = (p^2 + \frac{\nu^2 - 1/4}{r^2} - k^2 \mp i0)^{-1}, \quad p = -i\partial_r, \ k > 0.$$

Here $R_{\nu,k}^{\pm} \in \mathcal{L}_{s',-s} := \mathcal{L}(L_{s'}^2(\mathbb{R}_+, \mathrm{d}r), L_{-s}^2(\mathbb{R}_+, \mathrm{d}r))$ for s, s' > 1, in fact with a bound independent of $\nu \geq \nu_0$ and $k \in (0, 1)$. For simplicity we assume here and below that $\nu_0 \in (0, 1)$. See the proof of [Ca, Proposition 4.1], and note also that this proof shows that the weighted space operator norm vanishes uniformly in k as $\nu \to \infty$. We define the adjoint $(K_k^{\pm})^* = (h_a - k^2)(G_k^{\pm})^* - 1$ and conclude that $K_0^{\pm} := \lim_{k \to 0_+} K_k^{\pm} \in \mathcal{L}(\mathcal{H}_{-1-\epsilon}^1)$ for any sufficiently small $\epsilon > 0$.

<u>2</u>. We derive bounds of the operator $R^+_{\nu,k}$ acting on functions on \mathbb{R}_+ , here assuming $\nu > 1$. Let $\epsilon \in (0, \nu_0)$. The kernel is given explicitly as

$$R_{\nu,k}^{+}(r,r') = -2^{-1}\sqrt{rr'} \int_{0}^{\infty} \exp\left(i\rho t + ik^{2}rr'/(2t) - i\pi\nu/2\right) J_{\nu}(t)t^{-1} dt; \quad \rho = \frac{r^{2} + r'^{2}}{2rr'}.$$

(See for example [Ca].) We write $1 = \chi_{2kr'}(t) + \bar{\chi}_{2kr'}(t)$, cf. the notation (1.29), and

$$\exp(\cdot) = \left(1 + \rho^2 (1 - y)^2\right)^{-1} \left(1 - i\rho(1 - y)\partial_t\right) \exp(\cdot); \quad y = \frac{k^2 (rr')^2}{(r^2 + r'^2)t^2}.$$

We split the above integral $\int_0^\infty = \int_0^\infty \cdot \chi_{2kr'}(t) dt + \int_0^\infty \cdot \overline{\chi}_{2kr'}(t) dt$ and note that for $\nu \ge \nu_0$ a Bessel function bound, cf. the proof of [Ca, Proposition 4.1], yields

$$\left| \int_{0}^{\infty} \cdot \chi_{2kr'}(t) \, \mathrm{d}t \right| \le o(\nu^{0}) \min\{(kr')^{\nu_{0}}, 1\} \le o(\nu^{0})(kr')^{\epsilon},$$

leading with the Hilbert-Schmidt criterion to the operator bound

$$\left\|\sqrt{rr'} \int_0^\infty \cdot \chi_{2kr'}(t) \,\mathrm{d}t\right\|_{\mathcal{L}_{s',-s}} \le o(\nu^0) k^{\epsilon}; \quad s > 1, \, s' > 1 + \epsilon.$$
(6.107a)

On the other hand, assuming (for (6.107b)) that $\nu > 1$,

$$\bar{\chi}_{2kr'}(t)y \le 1/4,$$

leading by integration by parts and Bessel function bounds to

$$\left| \int_{0}^{\infty} \cdot \bar{\chi}_{2kr'}(t) \,\mathrm{d}t \right| \le o(\nu^{0})\rho^{-1},\tag{6.107b}$$

$$\left\|\sqrt{rr'}\int_0^\infty \cdot \bar{\chi}_{2kr'}(t) \,\mathrm{d}t\right\|_{\mathcal{L}_{s',-s}} \le o(\nu^0); \quad s+s'>2, \, s, s'>0. \tag{6.107c}$$

Note that for (6.107c) we used (6.107b), the Hilbert-Schmidt criterion, the bound

$$r^{\kappa}\rho^{-1}r'^{-\kappa} \le 2; \quad \kappa \in [-1,1],$$

as well as the identity

$$2J'_{\nu}(t) = J_{\nu-1}(t) - J_{\nu-1}(t);$$

cf. [Ta1, (3.6.17)]. Now by combining (6.107a) and (6.107c) we can write

$$R_{\nu,k}^{+} = R_{\nu,k,1}^{+} + R_{\nu,k,2}^{+};$$

$$R_{\nu,k,1}^{+} \in \mathcal{L}_{s',-s}; \quad s > 1, \ s' > 1 + \epsilon,$$

$$\|R_{\nu,k,1}^{+}\|_{\mathcal{L}_{s',-s}} \leq o(\nu^{0})k^{\epsilon}.$$

$$R_{\nu,k,2}^{+} \in \mathcal{L}_{s',-s}; \quad s = 1 - \epsilon, \ s' > 1 + \epsilon,$$

$$\|R_{\nu,k,2}^{+}\|_{\mathcal{L}_{s',-s}} \leq o(\nu^{0}).$$
(6.108a)

<u>3</u>. We show bounds of $R_{\nu,k}^+$ for the case $\nu \leq 1$. Let again $\epsilon \in (0, \nu_0)$. Note the representation, cf. [Ta1, pp. 228–230],

$$R_{\nu,k}^+(r,r') = \frac{\pi i}{2k} (kr_{<})^{1/2} J_{\nu}(kr_{<}) (kr_{>})^{1/2} H_{\nu}^{(1)}(kr_{>}),$$

as well as the classical pointwise bounds

$$|t^{1/2} J_{\nu}(t)| \leq C_{\nu} \left(\frac{t}{\langle t \rangle}\right)^{\nu+1/2}, |t^{1/2} H_{\nu}^{(1)}(t)| \leq C_{\nu} \left(\frac{t}{\langle t \rangle}\right)^{-\nu+1/2}$$

Whence

$$\forall \nu \in (0, 1/2]: \quad |(kr_{<})^{1/2} J_{\nu}(kr_{<})(kr_{>})^{1/2} H_{\nu}^{(1)}(kr_{>})| \le C_{\nu}^{2} k \, r_{<}^{\nu+1/2} \, r_{>}^{-\nu+1/2},$$

$$\forall \nu \in (1/2, \infty): \quad |(kr_{<})^{1/2} J_{\nu}(kr_{<})(kr_{>})^{1/2} H_{\nu}^{(1)}(kr_{>})| \leq C_{\nu}^{2} k r_{<}.$$

Using the Hilbert-Schmidt criterion we obtain from these bounds:

$$\forall \nu \in (0, 1/2]: \quad R^+_{\nu,k} = R^+_{\nu,k,2} \in \mathcal{L}_{s',-s}; \quad s = 1 - \epsilon, \ s' > 1 + \epsilon, \\ \|R^+_{\nu,k,2}\|_{\mathcal{L}_{s',-s}} \le C_{\nu,s,s'}.$$
(6.108b)

$$\forall \nu \in (1/2, 1] : \quad R_{\nu,k}^+ = R_{\nu,k,2}^+ \in \mathcal{L}_{s',-s}; \quad s, s' > 1/2, \ s+s' > 2, \\ \|R_{\nu,k,2}^+\|_{\mathcal{L}_{s',-s}} \le C_{\nu,s,s'}.$$

$$(6.108c)$$

Clearly $R_{\nu,k,1}^+ = 0$ in (6.108b) and in (6.108c), and it is natural to ask if we also can take $R_{\nu,k,1}^+ = 0$ in (6.108a)? Well, we dont know. Note that it is known for the above pointwise bound of the Bessel function that the constant $C_{\nu} \to \infty$ for $\nu \to \infty$, see [La]. This is the reason we proceeded differently in Step 2 to treat the regime $\nu > 1$.

4. With reference to Step 1, we claim that

$$-1 \notin \sigma(K_0^+). \tag{6.109a}$$

This is equivalent to asserting that r_{λ}^{\pm} is regular at k = 0, so it remains to show that 0 is not a 'resonance' for h_a . We say that 0 is a resonance if there exists $0 \neq f \in \mathcal{H}_{-s_0}^1 \setminus \mathcal{H}^1$, $s_0 = 1 + \nu_0$, such that $h_a f = 0$, cf. Definition 4.6. Note that the notation \mathcal{H}_t^1 here and henceforth is used as an alternative to H_t^1 (to conform with the notation of Chapter 4). So suppose $f \in \mathcal{H}_{-s_0}^1$ obeys $h_a f = 0$. Then we learn from the proof of Theorem 4.7 using (6.108a)–(6.108c) with k = 0 that $f \in \mathcal{H}_{-s'}^1$ for any $s' > 1 - \nu_0$ (note that $K_0^+ \in \mathcal{C}(\mathcal{H}_{-s'}^1)$ for $s' \in (1 - \nu_0, 2 - \nu_0)$). In particular $f \in \mathcal{H}_{-1}^1$. Next we mimic the proof of Theorem 4.2, that is integrate by part and deduce that

$$0 = \lim_{R \to \infty} \langle \chi_R f, \chi_R h_a f \rangle = \lim_{R \to \infty} \langle \chi_R f, h_a \chi_R f \rangle \ge c ||x|^{-1} f||^2.$$

Whence f = 0.

<u>5</u>. We can obtain bounds of r_{λ}^{\pm} using (6.109a) and (6.108a)–(6.108c). Note that thanks to the time-reversal property the latter bounds also hold with $R_{\nu,k}^{+}$ replaced by $R_{\nu,k}^{-}$. Bounds of r_{λ}^{\pm} can then be derived from the formulas

$$r_{\lambda}^{+} = \left(1 + K_{k}^{+}\right)^{-1} (G_{k}^{-})^{*} = G_{k}^{+} \left(1 + (K_{k}^{-})^{*}\right)^{-1}.$$
 (6.109b)

We can here use $G_k^{\pm} = G_{k,1}^{\pm} + G_{k,2}^{\pm}$ in agreement with the splittings $R_{\nu,k}^+ = R_{\nu,k,1}^+ + R_{\nu,k,2}^+$ in (6.108a)–(6.108c). Implemented in (6.109b) leads to two formulas for $r_{\lambda}^+ = r_{\lambda,1}^+ + r_{\lambda,2}^+$ exhibiting mapping properties directly determined by (6.108a)–(6.108c).

Since we are assuming that λ_0 is not a resonance, we are lead to write

$$-E_{\mathcal{H}}^{+}(\lambda) = p_{a}^{2} + w + v_{\lambda}^{+} + \lambda_{0} - \lambda$$

and then consider

$$\left(\widetilde{K}_{k}^{+}\right)^{*} = \left(-E_{\mathcal{H}}^{+}(\lambda)^{*}\right)\left(r_{\lambda}^{+}\right)^{*} - 1.$$

By the regularity condition of Case II) the limiting operator $\widetilde{K}_0^+ := \lim_{k \to 0_+} \widetilde{K}_k^+ \in \mathcal{L}(\mathcal{H}_{-1-\epsilon}^1)$ (which exists for any sufficiently small $\epsilon > 0$) fulfills that

$$-1 \notin \sigma(\widetilde{K}_0^+). \tag{6.109c}$$

By using (6.109a)-(6.109c) in combination with (6.108a)-(6.108c) we can obtain bounds of the inverse of $E_{\mathcal{H}}^+(\lambda)$. However we also need bounds on the Fourier transform to treat (6.104).

 $\underline{6}$. Bounds on the restriction of the Fourier transform. We note the bound

$$\|\mathcal{F}_{\lambda_0}(\lambda)\langle x\rangle^{-1-\epsilon}\|_{\mathcal{L}(L^2(\mathbb{R}^3),L^2(\mathbb{S}^2))} \le C_{\epsilon,\epsilon'}(\lambda-\lambda_0)^{\epsilon'}, \quad 0<2\epsilon'<\epsilon\le 1/2,$$

which follows by using the well-known expression for the free three-dimensional resolvent kernel and the Hilbert-Schmidt criterion. In terms of the k-variable it corresponds to the first of the following bounds. For the second bound we refer to [Ag, Theorem 3.2].

$$\begin{aligned} \|\mathcal{F}_0(k^2)\langle x\rangle^{-1-\bar{\epsilon}}\|_{\mathcal{L}(L^2(\mathbf{X}),L^2(\mathbb{S}^2))} &\leq Ck^{\epsilon'}, \quad 0<\epsilon'<\bar{\epsilon}\leq 1/2, \\ \|\mathcal{F}_0(k^2)\langle x\rangle^{-s}\|_{\mathcal{L}(L^2(\mathbf{X}),L^2(\mathbb{S}^2))}^2 &\leq Ck^{-1}, \quad s>1/2, \end{aligned}$$

Interpolation of these bounds yields

$$\forall \kappa \in [0,1], \epsilon' < \bar{\epsilon} \le 1/2, \ s > 1/2: \| \mathcal{F}_0(k^2) \langle x \rangle^{-\kappa(1+\bar{\epsilon}) - (1-\kappa)s} \|_{\mathcal{L}(L^2(\mathbf{X}), L^2(\mathbb{S}^2))} \le Ck^{\kappa\epsilon' - (1-\kappa)/2}.$$

$$(6.110)$$

<u>7</u>. We can now treat (6.104). The middle term of (6.104) may be written

$$-2\pi J_{\alpha}(\lambda)^{*}T_{\alpha}(\lambda) = -2\pi \mathrm{i}\mathcal{F}_{\lambda_{0}}(\lambda) \left(w + \mathcal{O}(r^{-3})\right)\mathcal{F}_{\lambda_{0}}(\lambda)^{*}$$
$$= -2\pi \mathrm{i}\mathcal{F}_{\lambda_{0}}(\lambda)w\mathcal{F}_{\lambda_{0}}(\lambda)^{*} + o((\lambda - \lambda_{0})^{0})$$

The first term corresponds to the 'Born term' for the one-body scattering matrix with potential w, and similarly in the third term of (6.104) the contribution $2\pi i T_{\alpha}(\lambda)^* Sr_{\lambda}^+ S^* T_{\alpha}(\lambda)$, recalling $r_{\lambda}^{\pm} = (p_a^2 + w + \lambda_0 - \lambda \mp i0)^{-1}$, may be written up to a term of order $o((\lambda - \lambda_0)^0)$ as

$$2\pi \mathrm{i}\mathcal{F}_{\lambda_0}(\lambda)wr_{\lambda}^+w\mathcal{F}_{\lambda_0}(\lambda)^*.$$

The combination of this expression, the identity operator 1 and the Born term is exactly $S_w(\lambda - \lambda_0)$. Whence it suffices to show that

$$\|T_{\alpha}(\lambda)^* (R(\lambda + \mathrm{i}0) - Sr_{\lambda}^+ S^*) T_{\alpha}(\lambda)\| \to 0.$$

Using formulas as (6.7) and (6.8) (cf. Remark 6.1) it suffices to show that

$$\|T_{\alpha}(\lambda)^* \dot{R}(\lambda + i0)\Pi' T_{\alpha}(\lambda)\| \to 0, \qquad (6.111a)$$

$$||T_{\alpha}(\lambda)^* \dot{R}(\lambda + i0)T_{\alpha}(\lambda)|| \to 0.$$
(6.111b)

To prove these assertions we look at three cases a)–c) as in the proof of Theorem 6.10.

a). The asymptotics (6.111a) follows by first writing

$$T_{\alpha}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i}0)\Pi'T_{\alpha}(\lambda) = J_{\alpha}(\lambda)^{*}\mathcal{O}(\langle x \rangle^{-2})\breve{R}(\lambda+\mathrm{i}0)\Pi'\mathcal{O}(\langle x \rangle^{-2})J_{\alpha}(\lambda)$$

and then using Theorem 3.18 and (6.110).

As for (6.111b) we expand $\tilde{R}(\lambda + i0)$ as in the proof of Theorem 6.10, cf. Cases b) and c) in the proof.

b). We write

$$T_{\alpha}(\lambda)^{*}Sr_{\lambda}^{+}\left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}S^{*}T_{\alpha}(\lambda) - T_{\alpha}(\lambda)^{*}Sr_{\lambda}^{+}S^{*}T_{\alpha}(\lambda)$$
$$= -T_{\alpha}^{+}(\lambda)^{*}Sr_{\lambda}^{+}\tilde{v}_{\lambda}^{+}r_{\lambda}^{+}S^{*}T_{\alpha}^{-}(\lambda);$$
$$\tilde{v}_{\lambda}^{+} = \left(1+v_{\lambda}^{+}r_{\lambda}^{+}\right)^{-1}v_{\lambda}^{+}.$$

In this formula we insert the two representations of $r_{\lambda}^{+} = r_{\lambda,1}^{+} + r_{\lambda,2}^{+}$ from the beginning of Step 5 (replacing the factor of r_{λ}^{+} to the right with the middle expression of (6.109b) and replacing the factor of r_{λ}^{+} to the left with the third expression of (6.109b)). Then we can show the desired decay.

When we exand the two sums we obtain four terms. Let us first consider the contribution when the type $r_{\lambda,2}^+$ appear twice. We insert $1 = \langle x \rangle^s \langle x \rangle^{-s}$ with $s = 1 - \epsilon$, $\epsilon > 0$ small, next to the two factors of $r_{\lambda,2}^+$. Thus for the left factor we write $r_{\lambda,2}^+ = \langle x \rangle^s \langle x \rangle^{-s} r_{\lambda,2}^+$ and then in turn $r_{\lambda,2}^+ = r_{\lambda,2}^+ \langle x \rangle^{-1-2\epsilon} \langle x \rangle^{1+2\epsilon}$ using bounds (essentially given by (6.108a)–(6.108c)) to estimate $\langle x \rangle^{-s} r_{\lambda,2}^+ \langle x \rangle^{-1-2\epsilon}$. Similarly for

the right factor we write $r_{\lambda,2}^+ = r_{\lambda,2}^+ \langle x \rangle^{-s} \langle x \rangle^s$. In combination with (6.110) (used with a vanishing right-hand side as $k \to 0_+$) we then obtain the bound $o((\lambda - \lambda_0)^0)$.

We can obtain the same conclusion for the three other terms in a similar way. The only difference lies in the choice of parameters. Thus the worse mapping property of $r_{\lambda,1}^+$ leads to an application of (6.110) with a (small) negative power of k. However the bounds (6.108a) offer positive powers in this case, which in combination indeed leads to the bound $o((\lambda - \lambda_0)^0)$.

c). There are four terms to be considered given (up to the common factor $2\pi i$) by:

$$S_{1} = -T_{\alpha}(\lambda)^{*}Sr_{\lambda}^{+}(1+v_{\lambda}^{+}r_{\lambda}^{+})^{-1}S^{*}I_{a}\breve{R}(\lambda+\mathrm{i}0)\Pi'T_{\alpha}(\lambda),$$

$$S_{2} = T_{\alpha}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i}0)\Pi'I_{a}Sr_{\lambda}^{+}(1+v_{\lambda}^{+}r_{\lambda}^{+})^{-1}S^{*}I_{a}\breve{R}(\lambda+\mathrm{i}0)\Pi'T_{\alpha}(\lambda),$$

$$S_{3} = -T_{\alpha}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i}0)\Pi'I_{a}Sr_{\lambda}^{+}S^{*}T_{\alpha}(\lambda),$$

$$S_{4} = T_{\alpha}(\lambda)^{*}\breve{R}(\lambda+\mathrm{i}0)\Pi'I_{a}Sr_{\lambda}^{+}(1+v_{\lambda}^{+}r_{\lambda}^{+})^{-1}v_{\lambda}^{+}r_{\lambda}^{+}S^{*}T_{\alpha}(\lambda).$$

We argue similarly as for Case b) and conclude again the bound $o((\lambda - \lambda_0)^0)$, skipping the details. Whence (6.111b) is established.

6.3.3 Proof of (6.98), Case III)

We consider the assertion (6.98) for Case 3 of Sections 1.2 and 1.3 under the conditions a) and b) stated after Theorem 6.28. By definition of Case III), λ_0 is a resonance but not an eigenvalue of H. We can apply (5.117a) of Theorem 5.24 with $\rho = 1$ (or alternatively Theorem 5.45), used formally with small z > 0. More precisely we consider $R(\lambda + i0)$ with λ slightly above λ_0 , given by (5.117a). We insert this expression in (6.104) (again with the wave operators given by (6.96) with $J_0^{\pm} = 1$).

As $\lambda \to (\lambda_0)_+$ the second term on the right-hand side of (6.104) disappears in the limit, cf. (6.110). As for the third term there is a cancellation of powers of $\lambda - \lambda_0$ exactly as for the one-body resonance case of [JK]. In fact we compute as in [JK, Section 5], and by using Remark 5.25,

$$\lim_{\lambda \to (\lambda_0)_+} 2\pi \mathrm{i} T^+_{\alpha}(\lambda)^* R(\lambda + \mathrm{i} 0) T^-_{\alpha}(\lambda) = -2\langle Y_0, \cdot \rangle Y_0; \quad Y_0 = (4\pi)^{-1/2}.$$

Whence

$$\lim_{\lambda \to (\lambda_0)_+} S_{\alpha\alpha}(\lambda) = 1 - 2\langle Y_0, \cdot \rangle Y_0, \qquad (6.112)$$

and since the right-hand side is unitary, indeed (6.98) is proven.

We remark that (6.112) agrees with Levison's theorem for the one-body problem, see for example [Ne]. In the context of (5.117a) of Theorem 5.24 with multiplicity $m = m_a > 1$ and upon assuming $\kappa = n_{\rm res} = m$ (the maximality condition) we obtain the same result as (6.112), see below. We will study the case $1 \le \kappa = n_{\rm res} < m$ in the next subsection, but let us here give a formula valid in both situations.

We will use the normalization in Remark 5.25, i.e. $(c(S^*u_j), c(S^*u_k)) = \delta_{jk}$ for $j, k = 1, \ldots, \kappa$, where for any $u \in \mathcal{E}$ (with \mathcal{E} given in Theorem 4.17)

$$c_i(S^*u) = \frac{1}{2\sqrt{\pi}} \int_{\mathbf{X}} \varphi_i I_0 u \, \mathrm{d}x; \qquad i = 1, \dots, m.$$

Considering $\alpha_i = (a, \lambda_0, \varphi_i), i = 1, \dots, m$, we compute the following substitute for (6.112),

$$\lim_{\lambda \to (\lambda_0)_+} S_{\alpha_i \alpha_i}(\lambda) = 1 - 2 \sum_{j \le \kappa} |c_i(S^* u_j)|^2 \langle Y_0, \cdot \rangle Y_0.$$
(6.113)

Obviously the result (6.98) for Case III) follows from (6.113) (since $\kappa = m$ in that case), cf. (5.35).

6.3.4 An example of transmission

For Case III) treated above, but with the maximality condition replaced by $1 \leq \kappa = n_{\rm res} < m$, the limiting scattering operator might differ from (6.112). Thanks to (6.113), obviously this happens for $\alpha_i = (a, \lambda_0, \varphi_i)$ unless $\sum_{j \leq \kappa} |c_i(\psi_j)|^2 = 1$ (i.e. not smaller). The other extreme is that $\sum_{j \leq \kappa} |c_i(\psi_j)|^2 = 0$ meaning that $\lim_{\lambda \to (\lambda_0)_+} S_{\alpha_i \alpha_i}(\lambda) = 1$, which is a unitary operator. If on the other hand $0 < \sum_{j \leq \kappa} |c_i(\psi_j)|^2 < 1$, the isometry property of $\lim_{\lambda \to (\lambda_0)_+} S_{\alpha_i \alpha_i}(\lambda)$ is not fulfilled and transmission from the channel α_i will occur. We will construct such example by redefining the basis $\Phi := \{\varphi_1, \ldots, \varphi_m\}$. Whence we consider a general unitary transformation $(\varphi'_1, \ldots, \varphi'_m)^t = M(\varphi_1, \ldots, \varphi_m)^t$. We introduce the basis $\Phi' := \{\varphi'_1, \ldots, \varphi'_m\}$ and notation $S_{\Phi}, S_{\Phi'}$ and similarly for adjoints to indicate the dependence of basis.

We compute

$$S_{\Phi'}^* = \overline{M} S_{\Phi}^*. \tag{6.114}$$

Using the notation $\alpha_i = (a, \lambda_0, \varphi_i)$ and $\alpha'_i = (a, \lambda_0, \varphi'_i)$, $i = 1, \ldots, m$, we similarly compute

$$\left(S_{\alpha_i'\alpha_j'}(\lambda_0)\right)_{i,j\le m} = \left(\lim_{\lambda\to(\lambda_0)_+} S_{\alpha_i'\alpha_j'}(\lambda)\right)_{i,j\le m} = \overline{M}\left(S_{\alpha_i\alpha_j}(\lambda_0)\right)_{i,j\le m}\overline{M}^*.$$
 (6.115)

Let us on \mathcal{E} introduce the \mathbb{C}^m -valued function c_{Φ} , slightly abusing notation,

$$c_{\Phi,i}(u) = \frac{1}{2\sqrt{\pi}} \langle 1, (S_{\Phi}^* I_0 u)_i \rangle_0 = \frac{1}{2\sqrt{\pi}} \int_{\mathbf{X}} \varphi_i I_0 u \, \mathrm{d}x; \quad i = 1, \dots, m, \ u \in \mathcal{E}.$$

Using (6.114) we compute

$$c_{\Phi'}(u) = \overline{M}c_{\Phi}(u); \quad u \in \mathcal{E}.$$
(6.116)

The normalization of the resonance functions in Remark 5.25 reads with this notation

$$(c_{\Phi}(u_j), c_{\Phi}(u_k)) = \delta_{jk}; \quad j, k = 1, \cdots, \kappa$$

Let e_1, \ldots, e_m denote the canonical basis in \mathbb{C}^m , $(e_j)_i = \delta_{ij}$. We choose the unitary matrix M such that

$$\overline{M}c_{\Phi}(u_j) = e_j; \quad j \le \kappa.$$

Thanks to (6.116) this leads to

$$c_{\Phi',i}(u_j) = \delta_{ij}; \quad i \le m, j \le \kappa. \tag{6.117}$$

Using (6.117) we compute as in Subsection 6.3.3

$$\begin{split} S_{\alpha'_k \alpha'_l}(\lambda_0) &= \delta_{kl} 1 - 2 \sum_{j \le \kappa} c_{\Phi',k}(u_j) \overline{c_{\Phi',l}(u_j)} \langle Y_0, \cdot \rangle Y_0 \\ &= \delta_{kl} \begin{cases} 1 - 2 \langle Y_0, \cdot \rangle Y_0, \text{ for } k, l \le \kappa, \\ 1, \text{ otherwise.} \end{cases} \end{split}$$

In view of (6.115) this formula is an explicit diagonalization of the operator-valued matrix $(S_{\alpha_i\alpha_j}(\lambda_0))_{i,j\leq m}$. Each of the appearing diagonal elements is given by either $1 - 2 \langle Y_0, \cdot \rangle Y_0$ or 1 (depending on its location). Since both of these options are unitary we conclude that transmission from an incoming channel associated with (a, λ_0) , if occurring at all, is limited to outgoing channels in the family associated with (a, λ_0) . Moreover indeed such transmission does not occur from any of the channels $\alpha'_i = (a, \lambda_0, \varphi'_i), i = 1, \ldots, m$. However for 'mixtures' of these channels indeed transmission occurs. Thus for example considering $\alpha = (a, \lambda_0, \varphi)$ for the mixture $\varphi = \cos \theta \varphi'_1 + \sin \theta \varphi'_m$, $\cos \theta \sin \theta \neq 0$, we compute using (6.115)

$$S_{\alpha\alpha}(\lambda_0) = 1 - 2\cos^2\theta \,\langle Y_0, \cdot \rangle Y_0,$$

and this operator is not isometric.

6.4 Threshold behaviour of total cross-sections in atom-ion scattering

The scattering process for multi-particle Coulomb systems with initial two-cluster data has been studied in physics literature, both experimentally and theoretically. In particular, in the collision of a neutral cluster with a charged one (atom-ion scattering), physical pictures suggest that if the neutral sub-system has no static dipole moment, the total cross-sections would be finite. Its mathematical proof is subtle. In [ES], V. Enss and B. Simon put forward several open questions and the sixth one of them is the following:

"6. Atom-Ion Scattering. An induced polarization picture suggests that Coulomb cross-sections with one neutral and one charged cluster will be finite if the neutral system has no static dipole moment. We are unable to prove this. Can one obtain explicit bounds in such a case?"

In [JKW], the authors give an affirmative answer to this question of Enss-Simon. In fact they prove the finiteness of Coulomb total cross-sections in atom-ion scattering for non-threshold energies and study the Born-Oppenheimer approximation. Here we are interested in the threshold behaviour of total cross-sections.

Recall the well-known fact in one-body scattering theory (see for example [Ya4]) that if a bounded real potential V on \mathbb{R}^3 decays like $\mathcal{O}(|x|^{-\rho})$ for some $\rho > 2$ then

the total cross-section for the scattering process described by the couple of operators $(-\Delta, -\Delta + V(x))$ is finite, while if $V(x) \approx \frac{C}{|x|^2}$ as $|x| \to \infty$ for some $C \neq 0$ then the total cross-section is infinite. In the scattering theory for multi-particle Coulomb systems with initial two-cluster data, the inter-cluster interaction between the two clusters decays like $\mathcal{O}(|x|^{-1})$ in the general case, like $\mathcal{O}(|x|^{-2})$ if one of the clusters is neutral (atom-ion scattering) and like $\mathcal{O}(|x|^{-3})$ if both clusters are neutral (atom-atom scattering), see Subsection 1.2.1. Here $x \in \mathbb{R}^3$ denotes the relative position between mass-centers of the two clusters. For atom-ion scattering, the known results for the one-body case suggest that without an additional assumption, the total cross-section would be infinite. On the other hand with the assumption that the atom is in the fundamental state the finiteness of total cross-section follows from [JKW], since by the symmetry of Coulomb potentials there is no static dipole moment for the atom in this case, cf. Subcases 3b and 3c listed in Subsection 1.2.1. The goal of this section is to study the threshold behaviour of the total cross-section in atom-ion scattering at the lowest threshold Σ_2 .

Consider the Hamiltonian of a diatomic molecule with ${\cal N}$ electrons which can be written in the form

$$H_{\text{phys}} = \sum_{k=1}^{2} \frac{1}{2m_{k}} \left(-\Delta_{x_{k}}\right) + \sum_{j=3}^{N+2} \frac{1}{2} \left(-\Delta_{x_{j}}\right) + \frac{Z_{1}Z_{2}}{|x_{1} - x_{2}|} \qquad (6.118)$$
$$+ \sum_{k=1}^{2} \sum_{j=3}^{N+2} \frac{q_{j}Z_{k}}{|x_{j} - x_{k}|} + \sum_{3 \le l < j \le N+2} \frac{q_{l}q_{j}}{|x_{l} - x_{j}|},$$

where $x_k \in \mathbb{R}^3$, k = 1, 2, denote the position of the two nuclei with mass m_k and charge $Z_k > 0$ and $x_j \in \mathbb{R}^3$, $j = 3, \ldots, N+2$, denote the position of N electrons with mass 1 and charge $q_j \in \mathbb{R}$ (for the physical case charges are equal and negative). Planck's constant \hbar is taken to be 1 in this formula. Obviously (6.118) is a special form of (1.13).

We are interested in scattering processes where the incoming channel is a twocluster one, while the outgoing channel is arbitrary. Let $a = (C_1, C_2)$ be a two-cluster decomposition of $\{1, \ldots, N+2\}$, i.e. a partition (C_1, C_2) of the set of particle labels $\{1, \ldots, N+2\}$, where $j \in C_j$ for j = 1, 2. In order to make explicit calculations, we choose so called *clustered atomic coordinates* $(x, y) \in \mathbb{R}^3 \times \mathbb{R}^{3N}$ adapted to this cluster decomposition:

$$h = \left(\frac{1}{2M_1} + \frac{1}{2M_2}\right)^{1/2}, \quad M_k = m_k + |C'_k|, \quad C'_k = C_k \setminus \{k\}, \quad k = 1, 2,$$

$$R_{k} = \frac{1}{M_{k}} \left(m_{k} x_{k} + \sum_{j \in C'_{k}} x_{j} \right), k = 1, 2$$

$$x = R_{1} - R_{2},$$

$$y_{j} = x_{j} - x_{k}, \quad j \in C'_{k}, \quad k = 1, 2,$$

$$l(y) = \frac{1}{M_{1}} \sum_{j \in C'_{1}} y_{j} - \frac{1}{M_{2}} \sum_{j \in C'_{2}} y_{j}.$$

In the Born-Oppenheimer approximation studied in [JKW], h is regarded as a small parameter. Here h is regarded as a constant, so we set h = 1. Note that R_k is the center of mass of the cluster C_k for k = 1, 2, and that x is the relative position of these centers of mass. These coordinates are well adapted to describe two-cluster scattering of diatomic molecules. After removal of the molecular center of mass motion, the Hamiltonian H_{phys} may be written in this system of coordinates as

$$H = -\Delta_x + H_e(x), \quad H_e(x) = H^a + I_a(x),$$
 (6.119)

where the sub-Hamiltonian H^a is given by

$$H^a = H^{C_1} + H^{C_2}, (6.120)$$

with

$$H^{C_k} = \sum_{j \in C'_k} \left(-\frac{1}{2} \Delta_{y_j} + \frac{Z_k q_j}{|y_j|} \right) - \frac{1}{2m_k} \left(\sum_{j \in C'_k} \partial_{y_j} \right)^2 + \sum_{\substack{l, j \in C'_k \\ l < j'}} \frac{q_l q_j}{|y_l - y_j|},$$

and the inter-cluster interaction I_a by

$$I_a(x) = \frac{Z_1 Z_2}{|x - l(y)|} + \sum_{\substack{k \in C_1' \\ j \in C_2'}} \frac{q_k q_j}{|y_k - y_j + x - l(y)|} + \sum_{j \in C_1'} \frac{Z_2 q_j}{|y_j + x - l(y)|} + \sum_{j \in C_2'} \frac{Z_1 q_j}{|x - l(y) - y_j|}$$

 $H_e(x)$ is the electronic Hamiltonian in the Born-Oppenheimer approximation. Finally, we set

$$H_a = -\Delta_x + H^a. aga{6.121}$$

This operator as well as the full Hamiltonian H are considered as self-adjoint operators on $L^2_{x,y} = L^2(\mathbb{R}^{3(N+1)}; \mathrm{d}x\mathrm{d}y).$

For an arbitrary cluster decomposition $a = (C_1, \ldots, C_k)$ of $\{1, \ldots, N+2\}$, i.e. $C_1 \cup \cdots \cup C_k = \{1, \ldots, N+2\}$ and $C_j \cap C_k = \emptyset$, for $j \neq k$, we can, as for the case k = 2 discussed above, choose adapted coordinates (x_a, y^a) . We call H^a the sub-Hamiltonian, $x_a \in \mathbb{R}^{3(k-1)}$ the inter-cluster coordinates, y^a the intra-cluster coordinates, and $I_a(x_a, y^a)$ the inter-cluster interaction. By D_{x_a} (resp. D_{y^a}) and by $-\Delta_{x_a}$ (resp. $-\Delta_{y^a}$), we denote -i times the gradient and the Laplacian in the inter-cluster (resp. intra-cluster) coordinates. It is well known [Do, De3, Sk6] that for this Schrödinger operator H, the Dollard wave operators

$$W_{\alpha}^{\pm} = s - \lim_{t \to \pm \infty} e^{itH} e^{-it\left(-\Delta_{x_a} + \int_{\pm 1}^{t} I_a(2sD_{x_a}, 0)ds + \lambda_{\alpha}\right)} J_{\alpha}$$
(6.122)

exist for an arbitrary channel $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$, recalling that (by definition) a is an arbitrary cluster decomposition ($\neq a_{\max}$) and φ_{α} is a normalized eigenfunction of H^a with eigenvalue λ_{α} , $H^a \varphi_{\alpha} = \lambda_{\alpha} \varphi_{\alpha}$. The operator J_{α} denotes the identification operator, which is defined for any L^2 -function f of the variable x_a by

$$(J_{\alpha}f)(x_a, y_a) = f(x_a)\varphi_{\alpha}(y_a).$$
(6.123)

Furthermore, the family of wave operators $\{W_{\alpha}^{\pm}, \text{ all } \alpha\}$ is asymptotically complete [De3]. If $a = (C_1, C_2)$ is a two-cluster decomposition with one neutral cluster (an atom), say C_1 , i.e.

$$\sum_{j \in C_1'} q_j = -Z_1, \tag{6.124}$$

then for any channel $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ the wave operators simply (cf. Section 6.2) as

$$W_{\alpha}^{\pm} = s - \lim_{t \to \pm \infty} e^{itH} e^{-it\left(-\Delta_{x_{\alpha}} + \lambda_{\alpha}\right)} J_{\alpha}.$$
(6.125)

For any two channels α and β we define the associated scattering operator from channel α to channel β by

$$S_{\beta\alpha} = (W_{\beta}^{+})^{*}W_{\alpha}^{-}, \quad T_{\beta\alpha} = S_{\beta\alpha} - \delta_{\beta\alpha}, \quad (6.126)$$

where $\delta_{\beta\alpha} = 1$ if $\alpha = \beta$ and 0 otherwise.

Following [ES], we define the total scattering cross-sections as follows. For $\lambda > \lambda_{\alpha}$, we introduce the magnitude of the momentum associated with the kinetic energy of the relative motion of the two clusters in the channel α via

$$n_{\alpha}(\lambda) := (\lambda - \lambda_{\alpha})^{\frac{1}{2}}.$$
(6.127)

For $g \in C_{c}^{\infty}(I_{\alpha}; \mathbb{C})$, $I_{\alpha} = (\lambda_{\alpha}, \infty)$, and $\omega \in \mathbb{S}^{2}$, we consider the wave packet

$$\mathbb{R}^3 \ni x \mapsto g_{\omega}(x) = \tilde{g}(\omega \cdot x) \tag{6.128}$$

where

$$\tilde{g}(\nu) = \frac{1}{2\sqrt{\pi}} \int_{\mathbb{R}} e^{in_{\alpha}(\mu)\nu} \frac{g(\mu)}{n_{\alpha}(\mu)^{1/2}} d\mu.$$

The normalization is chosen such that

$$\|g\|_{L^2(\mathbb{R})} = \|\tilde{g}\|_{L^2(\mathbb{R})}.$$

Denoting by \mathcal{C} the set of all channels we want to apply the operator $T_{\beta\alpha}$, for any $\beta \in \mathcal{C}$, to the function $g_{\omega}(x)\varphi_{\alpha}(y)$. Since this function does not belong to $L^2(\mathbb{R}^{3(N+1)})$ - it decays rapidly only in the direction defined by ω - we regularize it by multiplication by a function $h_{R,\omega} \in L^{\infty}(\mathbb{R}^3)$, depending only on the variable $x - (\omega \cdot x)\omega$ transversal to the direction ω of the incident wave packet $g_{\omega}(x)$, such that pointwisely

$$\lim_{R \to \infty} h_{R,\omega} = 1. \tag{6.129}$$

For the purpose of this paper we shall specify this cut-off function to be a Gaussian, explicitly we take

$$h_{R,\omega}(x) = e^{-(x - (\omega \cdot x)\omega)^2/R}.$$
 (6.130)

Definition. ([JKW]) For $\lambda \in I_{\alpha}$ and $\omega \in \mathbb{S}^2$, we shall say that the total crosssection $\sigma_{\alpha}(\lambda, \omega)$ with the incoming channel α exists at the energy λ with the incident direction ω , if the following limit is finite:

$$\sigma_{\alpha}(\lambda,\omega) := \lim_{n \to \infty} \lim_{R \to \infty} \sum_{\beta \in \mathcal{C}, \lambda_{\beta} < \lambda} \| \mathbf{T}_{\beta\alpha} h_{R,\omega} g_{n,\omega} \varphi_{\alpha} \|^{2}, \qquad (6.131)$$

where $g_{n,\omega}$ is defined as in (6.128) with g replaced by g_n :

$$g_n(\mu) = n^{-1/2} h((\mu - \lambda)/n)$$

and h is any $C_{\rm c}^{\infty}(\mathbb{R})$ -function normalized by $\int_{\mathbb{R}} |h(\mu)|^2 d\mu = 1$.

Recall that in [ES] and [Wa3], the total cross-section is defined as a distribution in $\mu \in I_{\alpha}$ by

$$\int_{I_{\alpha}} \sigma_{\alpha}(\mu, \omega) |g(\mu)|^2 d\mu = \lim_{R \to \infty} \sum_{\beta \in \mathcal{C}} \| \mathbf{T}_{\beta \alpha} h_{R, \omega} g_{\omega} \varphi_{\alpha} \|^2, \qquad (6.132)$$

for all $g \in C_c^{\infty}(I_{\alpha}; \mathbb{C})$. Since $|g_n(\cdot)|^2$ converges to $\beta_{\lambda}(\cdot)$, the Dirac measure at λ , as $n \to \infty$, the definitions (6.131) and (6.132) coincide if the distribution defined in (6.132) can be identified with a continuous function in a neighbourhood of λ . For fastly decaying pair potentials, total cross-sections can also be defined through scattering amplitudes (see [IT]).

6.4.1 Finiteness of total cross-sections in atom-ion scattering

Hypothesis 1. Let $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ be a channel with $\lambda_{\alpha} \in \sigma_{d}(H^{a})$ and cluster decomposition $a = (C_{1}, C_{2})$ such that the cluster a_{1} is neutral (an atom), that is

$$\sum_{j \in C_1'} q_j + Z_1 = 0. (6.133)$$

Assume that $\lambda_{\alpha} = \lambda_{\alpha}^{1} + \lambda_{\alpha}^{2}$, $\varphi_{\alpha} = \varphi_{\alpha}^{1} \otimes \varphi_{\alpha}^{2}$, $H^{C_{j}}\varphi_{\alpha}^{j} = \lambda_{\alpha}^{j}\varphi_{\alpha}^{j}$ for j = 1, 2 and that λ_{α}^{1} is a simple eigenvalue of $H^{C_{1}}$ (i.e. the eigenvalue for the neutral cluster is non-degenerate).

The following result of [JKW] shows the finiteness of $\sigma_{\alpha}(\lambda, \omega)$ for λ outside the set

$$\mathcal{T}_{p} = \mathcal{T}_{p}(H) = \mathcal{T}(H) \cup \sigma_{pp}(H)$$

of thresholds and eigenvalues of H, and it provides an optical formula which is useful in many problems.

Theorem 6.30 ([JKW]). Let $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ be a scattering channel satisfying Hypothesis 1. We set

$$F(z,\omega) = \left\langle I_a e^{\omega}_{\alpha}, R(z) \left(I_a e^{\omega}_{\alpha} \right) \right\rangle, \qquad \text{Im} \, z \neq 0, \tag{6.134a}$$

where

$$e^{\omega}_{\alpha}(x,y) = \mathrm{e}^{in_{\alpha}(\lambda)\omega \cdot x}\varphi_{\alpha}(y).$$
 (6.134b)

Then, for any energy $\lambda \in I_{\alpha} \setminus \mathcal{T}_{p}$ and any incident direction $\omega \in \mathbb{S}^{2}$, the limit

$$F(\lambda + i0, \omega) = \lim_{\epsilon \to 0_+} F(\lambda + i\epsilon, \omega)$$
(6.135a)

exists and defines a continuous function of $\lambda \in I_{\alpha} \setminus \mathcal{T}_{p}$. The total scattering crosssection $\sigma_{\alpha}(\lambda, \omega)$ exists for any energy $\lambda \in I_{\alpha} \setminus \mathcal{T}_{p}$ and any incident direction $\omega \in \mathbb{S}^{2}$ and one has the optical formula

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{n_{\alpha}(\lambda)} \operatorname{Im} F(\lambda + \mathrm{i0},\omega).$$
(6.135b)

Note that under Hypothesis 1, $I_a e_{\alpha}^{\omega}$ only belongs to $L_{(1/2)^-}^2 = L_{(1/2)^-}^2(\mathbb{R}^{3+3N}; dxdy)$. The existence of the limit (6.135a) is non-trivial. Its proof given in [JKW] uses the limiting absorption principle (cf. Theorem 3.18) and phase space analysis through appropriate localizations in the relative kinetic energy of the two clusters. For the same reason, the results established in Chapter 5 on threshold resolvent asymptotics can not be applied directly, because they hold as operators from L_s^2 to L_{-s}^2 with at least s > 1. Nevertheless, as the reader will see, various ingredients from Chapter 5 can be applied to study the threshold behaviour of $\sigma_{\alpha}(\lambda, \omega)$. In this subsection we consider for convenience only the simplest case, where the channel satisfies the following condition. (In Subsection 6.4.3 we consider a more general setup.)

6.4.2 Total cross-sections at Σ_2 , non-multiple two-cluster case

Hypothesis 2. Let $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ be a scattering channel with $\lambda_{\alpha} = \lambda_0 = \Sigma_2$, the lowest threshold of H. Assume that Σ_2 is a non-multiple two-cluster threshold and $\Sigma_2 \notin \sigma(H')$, where $H' = \Pi'_{\alpha} H \Pi'_{\alpha}$ and

$$\Pi'_{\alpha}u = u - \langle \varphi_{\alpha}, u \rangle_{y} \otimes \varphi_{\alpha}, \quad u \in L^{2}(\mathbb{R}^{3+3N}_{x,y}).$$

Under Hypothesis 2, we can apply the Grushin reduction (cf. [Wa2], extended in Chapter 5) to study the behaviour of $\sigma_{\alpha}(\lambda, \omega)$ for λ near $\lambda_0 = \Sigma_2$. The more general situations where λ_{α} possibly is a multiple two-cluster threshold and/or $\lambda_{\alpha} \in \sigma(H')$ can also be studied by the methods developed in the present work (the multiple case will indeed be studied in Subsection 6.4.3). Under the Hypothesis 2, R'(z) = $(H' - z)^{-1}\Pi'_{\alpha}$ is holomorphic for z near λ_0 and one has

$$R(z) = E(z) - E_{+}(z)E_{\mathcal{H}}(z)^{-1}E_{-}(z)$$
(6.136)

where

$$E(z) = R'(z)$$

$$E_{+}(z) = S - R'(z)HS$$

$$E_{-}(z) = S^{*} - S^{*}HR'(z)$$

$$E_{\mathcal{H}}(z) = S^{*}(z - H + HR'(z)H)S$$

and $S: L^2(\mathbb{R}^3_x) \to L^2(\mathbb{R}^{3+3N}_{x,y}), (x,y) \in \mathbb{R}^3 \times \mathbb{R}^{3N}$, is defined by

$$Sf(x,y) = f(x)\varphi_{\alpha}(y), \quad f \in L^2_x.$$

Under Hypotheses 1 and 2, it follows from Theorem 6.30 that for $\lambda > \lambda_0$ close to λ_0

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{n_{\alpha}(\lambda)} \operatorname{Im} \langle I_{a} e_{\alpha}^{\omega}, R(\lambda + \mathrm{i}0)(I_{a} e_{\alpha}^{\omega}) \rangle$$

$$= -\frac{1}{n_{\alpha}(\lambda)} \operatorname{Im} \langle f(\lambda,\omega), E_{\mathcal{H}}(\lambda + \mathrm{i}0)^{-1} f(\lambda,\omega) \rangle_{x};$$

$$f(\lambda,\omega) = E_{-}(\lambda)(I_{a} e_{\alpha}^{\omega}).$$
(6.138)

Here we used the fact that $R'(\lambda + i0) = R'(\lambda)$ is self-adjoint for λ near λ_0 . One can calculate

$$f(\lambda,\omega) = \langle \varphi_{\alpha}, I_a e^{\omega}_{\alpha} \rangle_y - \langle \varphi_{\alpha}, I_a R'(\lambda) (I_a e^{\omega}_{\alpha}) \rangle_y, \qquad (6.139a)$$

$$\langle \varphi_{\alpha}, I_a e^{\omega}_{\alpha} \rangle_y = \mathcal{O}(|x|^{-4}),$$
 (6.139b)

$$\langle \varphi_{\alpha}, I_a R'(\lambda) (I_a e^{\omega}_{\alpha}) \rangle_y \in L^2_{(5/2)^-}(\mathbb{R}^3_x),$$
(6.139c)

uniformly in λ and ω . For (6.139b) we refer to [JKW, Lemma A.1]. Hence

$$f(\lambda,\omega) \in L^2_{(5/2)^-} = L^2_{(5/2)^-}(\mathbb{R}^3_x)$$
 uniformly in λ and ω .

Similarly,

$$E_{\mathcal{H}}(z) = z - \lambda_0 - (-\beta_x + U(z)),$$

where

$$U(z) = \langle \varphi_{\alpha}, I_{a}\varphi_{\alpha} \rangle_{y} - \langle \varphi_{\alpha}, I_{a}R'(z)I_{a}(\varphi_{\alpha} \otimes \cdot) \rangle_{y}$$

is an operator-valued holomorphic function for z near λ_0 . One has

$$\langle \varphi_{\alpha}, I_a \varphi_{\alpha} \rangle_y = \mathcal{O}(|x|^{-4}), \quad \langle \varphi_{\alpha}, I_a R'(z) I_a(\varphi_{\alpha} \otimes \cdot) \rangle_y \in \mathcal{L}(H_s^1, H_{s+4}^{-1}); \quad s \in \mathbb{R}.$$

Set

$$U = U(\lambda_0) = \langle \varphi_{\alpha}, I_a \varphi_{\alpha} \rangle_y - \langle \varphi_{\alpha}, I_a R'(\lambda_0) I_a(\varphi_{\alpha} \otimes \cdot) \rangle_y.$$
(6.140)

Comparing $f(\lambda_0, \omega)$ with U, one observes

$$f(\lambda_0, \omega) = E_{-}(\lambda_0)(I_a \varphi_\alpha) = U1, \qquad (6.141)$$

that is, U applied to the constant function $1 \in L^2_{(-3/2)^-}(\mathbb{R}^3_x)$.

Recall from Section 5.1.2 that the asymptotic expansion of the resolvent $(-\Delta_x - z')^{-1}$, $z' = z - \lambda_0$ with $\text{Im } z \neq 0$, in weighted L^2 -spaces is given by

$$(-\Delta_x - z')^{-1} = G_0 + \sqrt{z'}G_1 + \cdots$$

Theorem 6.31 (Regular case). Let Hypotheses 1 and 2 be satisfied. Assume that $\lambda_0 = \Sigma_2$ is a regular point of H (see Subsection 5.1.1.1). Then

$$\sigma_{\alpha}(\lambda,\omega) = 4\pi |s_{lnt}|^2 + o(1) \quad as \quad \lambda \to (\lambda_0)_+, \tag{6.142a}$$

where

$$s_{\rm lnt} = \frac{1}{4\pi} \int_{\mathbb{R}^3_x} (1 + UG_0)^{-1} E_-(\lambda_0) (I_a \varphi_\alpha) \, \mathrm{d}x.$$
 (6.142b)

Proof. Since λ_0 is a regular point of H, $(1 + G_0 U)^{-1}$ exists and is bounded from L_s^2 to L_{-s}^2 for any s > 1. Let $z' = z - \lambda_0$ and Im $z \neq 0$. One has

$$E_{\mathcal{H}}(\lambda_0 + z')^{-1} = -(1 + (-\Delta_x - z')^{-1}U(z'))^{-1}(-\Delta_x - z')^{-1} \\ = -(1 + G_0U)^{-1}G_0 + \sqrt{z'} \{(1 + G_0U)^{-1}G_1(U(1 + G_0U)^{-1}G_0 - 1)\} + o(\sqrt{z'}) \\ = -(1 + G_0U)^{-1}G_0 - \sqrt{z'}(1 + G_0U)^{-1}G_1(1 + UG_0)^{-1} + o(\sqrt{z'})$$

in $L_s^2 \to L_{-s}^2$, s > 3/2. Making use of (6.137), we obtain for $\lambda' = \lambda - \lambda_0$

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{\sqrt{\lambda'}} \operatorname{Im} \left\langle f(\lambda,\omega), \left((1+G_0U)^{-1}G_0 + \sqrt{\lambda'}(1+G_0U)^{-1}G_1(1+UG_0)^{-1} + o(\sqrt{\lambda'}) \right) f(\lambda,\omega) \right\rangle \\ = \operatorname{Im} \left\langle f(\lambda_0,\omega), (1+G_0U)^{-1}G_1(1+UG_0)^{-1}f(\lambda_0,\omega) \right\rangle_x + o(1)$$

as $\lambda' \to 0_+$. Since G_1 has the constant integral kernel $\frac{i}{4\pi}$, the leading term in the above equation is equal to

$$\frac{1}{4\pi} \Big| \int_{\mathbb{R}^3_x} (1 + UG_0)^{-1} E_-(\lambda_0) (I_a \varphi_\alpha) \, \mathrm{d}x \Big|^2,$$

proving (6.142a) and (6.142b).

Note that since

$$E_{-}(\lambda_0)(I_a\varphi_\alpha) = f(\lambda_0,\omega) \in L^2_{(5/2)^-},$$

indeed the constant s_{lnt} is well defined. In addition it can be written as

$$s_{\text{lnt}} = \frac{1}{4\pi} \int_{\mathbb{R}^3_x} (1 + UG_0)^{-1} U1 \, \mathrm{d}x.$$

Recall that in the scattering theory for the pair $(-\Delta, -\Delta + V(x))$, the scattering length is related to the low-energy limit of the total cross-section. In dimension three it is equal to

$$\frac{1}{4\pi} \int_{\mathbb{R}^3_x} (1 + VG_0)^{-1} V(x) \, \mathrm{d}x,$$

if V(x) decreases sufficiently rapidly (cf. [JK, Theorem 5.1]). Analogously to onebody scattering, s_{lnt} given in (6.142b) can be interpreted as a *scattering length* in atom-ion scattering.

Theorem 6.32 (Exceptional point of 1st kind). Let Hypotheses 1 and 2 be satisfied. Assume that $\lambda_0 = \Sigma_2$ is a resonance but not an eigenvalue of H. Then

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{\lambda - \lambda_0} (4\pi + o(1)) \quad as \quad \lambda \to (\lambda_0)_+.$$
(6.143)

Proof. In general λ_0 is a resonance (an eigenvalue, resp.) of H if and only if 0 is a resonance (an eigenvalue, resp.) of $-\Delta_x + U$, cf. Lemma 5.2. Since we assume that λ_0 is exceptional point of 1st kind the operator $E_{\mathcal{H}}(\lambda_0 + z')^{-1}$ admits the asymptotics

$$E_{\mathcal{H}}(\lambda_0 + z')^{-1} = -(1 + (-\Delta_x - z')^{-1}U(z'))^{-1}(-\Delta_x - z')^{-1} \\ = -\frac{i}{\sqrt{z'}}(\langle \psi, \cdot \rangle_x \psi + o(1)), \quad z' \to 0,$$

in $L_s^2 \to L_{-s}^2$, s > 1, where $\psi \in L_{(-1/2)^-}^2$ is a resonance state of $-\Delta + U$ satisfying

$$\frac{1}{2\sqrt{\pi}}\langle U\psi, 1\rangle = 1. \tag{6.144}$$

x

See the proof of [Wa2, Theorem 3.9] or that of Proposition 5.12. We deduce from (6.137) that for $\lambda' = \lambda - \lambda_0$,

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{\lambda'}(|\langle \psi, f(\lambda_0,\omega) \rangle|^2 + o(1)) \text{ as } \lambda' \to 0_+$$

Noting that

(6.143) follows.

It is a remarkable phenomenon that in presence of threshold resonance, total cross-sections display a universal behavior near the threshold, independent of concrete form of potentials. If $\lambda_0 = \Sigma_2$ is an eigenvalue of H, the method used here does not allow us to give the leading term of $\sigma_{\alpha}(\lambda, \omega)$, due to the lack of decay. However with some weak decay of the L^2 -eigenfunctions at λ_0 one can obtain similar results as above. We shall give elaboration in the next subsection in the more general setting of $\lambda_0 = \Sigma_2$ possibly being a multiple two-cluster threshold.

6.4.3 Total cross-sections at Σ_2 , multiple two-cluster case

In this subsection we consider the atom-ion model with $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ being a scattering channel satisfying Hypothesis 1. Rather than Hypothesis 2 we impose the following condition using notation from Sections 4.3 and 5.3. Note that the atom-ion model is a specific example covered by these sections. Although we can not use the results of Section 5.3 directly, we can use their proofs, cf. Subsection 6.4.2. Again we consider only the case $\lambda_0 = \Sigma_2$ being a two-cluster threshold. We shall impose the condition

$$\widetilde{\mathcal{A}} = \mathcal{A}_3^{\rm fd},\tag{6.145}$$

and in addition (5.161), that is

$$\operatorname{ran} \Pi_H \subset L_t^2 \text{ for some } t > 3/2, \tag{6.146}$$

where Π_H is the orthogonal projection onto ker $(H - \lambda_0)$.

Hypothesis 3. Let $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$ be a channel with $\lambda_{\alpha} = \lambda_0 = \Sigma_2$, the lowest threshold of H. Assume that Σ_2 is two-cluster, (4.68a), (4.68b), (6.145) and (6.146) (in particular $a \in \mathcal{A}_3^{fd}$).

We remark that Hypothesis 3 implies Hypothesis 1 for any channel α , and that it is more general (i.e. weaker) than the combination of Hypotheses 1 and 2 used in the previous subsection. The content of (6.145) is that for the two-cluster threshold Σ_2 it holds for any $a = (C_1, C_2) \in \widetilde{\mathcal{A}}$ (possibly not unique) that one of the clusters is neutral, say (6.124) is fulfilled for C_1 .

Theorem 6.33 (Regular or exceptional of 2nd kind case). Let Hypothesis 3 be satisfied for $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$, and assume that $\lambda_{\alpha} = \lambda_0 = \Sigma_2$ is either regular or exceptional of 2nd kind. Let $H_{\sigma} := H - \sigma \Pi_H$ with $\sigma > 0$ small. Then, using the notation (6.134b) and that of Theorem 5.44, we let

$$f_{\sigma}(\lambda,\omega) = \left(S^* - S^*_{I_{\sigma}}R'_{\sigma}(\lambda)\right)(I_{a,\sigma}e^{\omega}_{\alpha}); \quad \lambda \ge \lambda_0.$$
(6.147a)

$$S_{I_{\sigma}}^{*}(\cdot) = \sum_{\beta=(b,E_{\beta},\varphi_{\beta}): b\in\widetilde{\mathcal{A}}}^{\oplus} \langle \varphi_{\beta} | I_{b,\sigma}(\cdot) \rangle_{y_{b}}, \quad I_{b,\sigma} = I_{b} - \sigma \Pi_{H},$$

and we recall that $R'_{\sigma}(\lambda) = (\Pi' H_{\sigma} \Pi' - \lambda)^{-1} \Pi'$. Let similarly $U_{\sigma}(\lambda - \lambda_0)$ be defined by writing the effective Grushin Hamiltonian for H_{σ} as $-E_{\mathcal{H},\sigma}(\lambda_0 + z) = P_0 + U_{\sigma}(z) - z$, $z = \lambda - \lambda_0 \geq 0$, along the lines of (5.5a), (5.5b) and (5.38a). Then

$$\sigma_{\alpha}(\lambda,\omega) = 4\pi |s_{lnt}|^2 + o(1) \quad as \quad \lambda \to (\lambda_0)_+, \tag{6.147b}$$

where

$$s_{\text{lnt}} = \frac{1}{4\pi} \int_{\mathbb{R}^3_x} (1 + U_\sigma G_0)^{-1} f_\sigma(\lambda_0, \omega) \, \mathrm{d}x \in \mathbb{C}^m; \quad U_\sigma = U_\sigma(0), \, m = \# \widetilde{\mathcal{A}}. \quad (6.147\text{c})$$

Proof. We can mimic the proof of Theorem 6.31, combining Theorem 6.30 and (5.87), checking first the analogue bound

$$f_{\sigma}(\lambda,\omega) \in \mathcal{H}_{(3/2)^+} = \mathcal{H}_{(3/2)^+}(\mathbb{R}^3_x).$$

The latter assertion follows readily from (6.146) (yielding in fact that $f_{\sigma}(\lambda, \omega) \in \mathcal{H}_s$ provided $s \leq t$ and s < 5/2).

Theorem 6.34 (Exceptional point of 1st or 3rd kind). Let Hypothesis 3 be satisfied for $\alpha = (a, \lambda_{\alpha}, \varphi_{\alpha})$. Assume that $\lambda_{\alpha} = \lambda_0 = \Sigma_2$ is exceptional of 1st or 3rd kind. Let $H_{\sigma} = H - \sigma \Pi_H$ with $\sigma > 0$ small and $\{u_1, \ldots, u_{\kappa}\} \subset H^1_{(-1/2)^-}$ be a basis of resonance states of H, more precisely it is taken as a basis of resonance states of H_{σ} fulfilling the normalization

$$(c(T^*u_i), c(T^*u_j)) = \beta_{ij}, \quad i, j = 1, \cdots, \kappa,$$
 (6.148a)

cf. (6.144) and Proposition 5.8. By definition T is given by (2.14) and

$$c(v) = \frac{1}{2\sqrt{\pi}} \left(\langle 1, \Sigma_{k \le m} U_{\sigma, 1k} v_k \rangle_x, \cdots, \langle 1, \Sigma_{k \le m} U_{\sigma, mk} v_k \rangle_x \right) \in \mathbb{C}^m; \ m = \# \widetilde{\mathcal{A}}.$$
(6.148b)

Here the elements of $\widetilde{\mathcal{A}}$ are labelled by the numbers $1, \ldots, m$.

Then, with the convention that 1 labels a,

$$\sigma_{\alpha}(\lambda,\omega) = \frac{1}{\lambda-\lambda_0} \left(\sum_{j=1}^{\kappa} |\langle T^*u_j, f_{\sigma}(\lambda_0,\omega) \rangle|^2 + o(1) \right)$$

$$= \frac{4\pi}{\lambda-\lambda_0} \left(\sum_{j=1}^{\kappa} |c_1(T^*u_j)|^2 + o(1) \right) \quad as \quad \lambda \to (\lambda_0)_+.$$
 (6.149a)

In particular

$$\sigma_{\alpha}(\lambda,\omega) \leq \frac{4\pi}{\lambda - \lambda_0} (1 + o(1)) \quad as \quad \lambda \to (\lambda_0)_+, \tag{6.149b}$$

while in the maximally exceptionally case where $\kappa = m$,

$$\sigma_{\alpha}(\lambda,\omega) = \frac{4\pi}{\lambda - \lambda_0} (1 + o(1)) \quad as \quad \lambda \to (\lambda_0)_+.$$
(6.149c)

206

Proof. We obtain (6.149a) by mimicking the proof of Theorem 6.32. The analogue of (6.141) reads

$$f_{\sigma}(\lambda_0,\omega) = E_{-}(H_{\sigma} - \lambda_0)\varphi_{\alpha} = E_{-}(H_{\sigma} - \lambda_0)S_1[1]$$

= $-E_{\mathcal{H}}[\Sigma_k^{\oplus}\beta_{1k}] = (P_0 + U_{\sigma})[\Sigma_k^{\oplus}\beta_{1k}] = U_{\sigma}[\Sigma_k^{\oplus}\beta_{1k}].$

The assertions (6.149b) and (6.149c) follow from (6.149a) and the normalization condition (6.148a). $\hfill \Box$

Remark. The quantity $\sum_{j=1}^{\kappa} |c_1(T^*u_j)|^2$ of (6.149a) appears also in (6.113) and in Subsection 6.3.4 although for a different setup. This is not a coincidence, an overall conclusion may be viewed as the same: It may be interpreted as a probability for resonance induced scattering phenomena at a threshold. This impact appears with probability one in the maximally exceptionally case for both models.

Bibliography

- [AW] M. Aafarani, X.P. Wang, Gevrey estimates of the resolvent and subexponential time-decay for the heat and Schrödinger semigroups. II. J. Differential Equations 316 (2022), 387-424.
- [AIIS] T. Adachi, K. Itakura, K. Ito, E. Skibsted, New methods in spectral theory of N-body Schrödinger operators, Rev. Math. Phys. 33 (2021), 48 pp.
- [Ag] S. Agmon, A representation theorem for solutions of the Helmholtz equation and resolvent estimates for the Laplacian, Academic Press, Inc., Boston, MA, xii+694 pp., 1990, 39–76.
- [ABG] W.O. Amrein, A.M. Boutet de Monvel, V. Georgescu, Lower bounds for zero energy eigenfunctions of Schrödinger operators, Helv. Phys. Acta 57 no. 3 (1984), 301–306.
- [AHS] A. Agmon, I. Herbst and E. Skibsted, Perturbation of embedded eigenvalues in the generalized N-body problem, Comm. Math. Phys. 122 (1989), 411–438.
- [Ba] E. Balslev, Analyticity properties of eigenfunctions and scattering matrix, Comm. Math. Phys. 114 no. 4 (1988), 599–612.
- [Bo] A. Bommier, Properties of the 2 -cluster 2 -cluster scattering matrix for longrange N -body problems, Ann. Inst. H. Poincarée Phys. Théor. 59 no. 3 (1993), 237–267.
- [Ca] G. Carron, Le saut en zéro de la fonction de décalage spectrale, J. Funct. Analysis, 212(2004), 222-260.
- [CT] J.M. Combes, A. Tip, Properties of the scattering amplitude for electron-atom collisions, Ann. Inst. H. Poincarré Phys. Théor. 40 (1984), 117–139.
- [De1] J. Dereziński, Existence and analyticity of many-body scattering amplitudes at low energies, J. Math. Phys. 28 (1987), 1080–1088.
- [De2] J. Dereziński, Threshold singularities of two-cluster-two-cluster scattering amplitudes for dilation analytic potentials, J. Math. Phys. 29 (1988), 1171– 1173.
- [De3] J. Dereziński, Asymptotic completeness for N-particle long-range quantum systems, Ann. of Math. 38 (1993), 427–476.
- [DG] J. Dereziński and C. Gérard, Scattering theory of classical and quantum Nparticle systems, Texts and Monographs in Physics, Berlin, Springer 1997.

- [DS1] J. Dereziński, E. Skibsted, Quantum scattering at low energies, J. Funct. Anal., 257 (2009), 1828–1920.
- [DS2] J. Dereziński, E. Skibsted, Scattering at zero energy for attractive homogeneous potentials, Ann. H. Poincaré 10 no. 3 (2009), 549–571.
- [Do] J. Dollard, Asymptotic convergence and Coulomb interaction, J. Math. Phys. 5 (1964), 729–738.
- [ES] V.Enss, B.Simon, Finite Total Cross-Section in Nonrelativistic Quantum Mechanics. Comm. Math. Phys. 76 (1980), 177–209.
- [Fr] R. Frank A note on low energy scattering for homogeneous long-range potentials, Ann. H. Poincaré 10 no.3 (2009), 573–575.
- [FH] R. Froese and I. Herbst, Exponential bounds and absence of positive eigenvalues for N-body Schrödinger operators, Comm. Math. Phys. 87 (1982), 429–447.
- [FS] S. Fournais, E. Skibsted, Zero energy asymptotics of the resolvent for a class of slowly decaying potentials, Math. Z. 248 (2004), 593–633.
- [Gr] G.M. Graf, Asymptotic completeness for N-body short-range quantum systems: a new proof, Comm. Math. Phys. 132 (1990), 73–101.
- [GIS] C. Gérard, H. Isozaki, E. Skibsted, N-body resolvent estimates, J. Math. Soc. Japan 48 no. 1 (1996), 135–160.
- [Hö1] L. Hörmander, *The analysis of linear partial differential operators*. *I*, Berlin, Springer 1990.
- [Hö2] L. Hörmander, The analysis of linear partial differential operators. II-IV, Berlin, Springer 1983-85.
- [Is1] H. Isozaki, Differentiability of generalized Fourier transforms associated with Schrödinger operators, J. Math. Kyoto Univ. 25 no. 4 (1985), 789–806.
- [Is2] H. Isozaki, Structures of the S-matrices for three-body Schrödinger operators, Commun. Math. Phys. 146 (1992), 241–258.
- [Is3] H. Isozaki, Asymptotic properties of solutions to 3-particle Schrödinger equations, Commun. Math. Phys. 222 (2001), 371–413.
- [Is4] H. Isozaki, A generalization of the radiation condition of Sommerfeld for Nbody Schrödinger operators, Duke Math. J. 74 no. 2 (1994), 557–584.
- [IS1] K. Ito, E. Skibsted, Absence of positive eigenvalues for hard-core N-body systems, Ann. Inst. Henri Poincaré 15 (2014), 2379–2408.
- [IS2] K. Ito, E. Skibsted, Rellich's theorem and N-body Schrödinger operators, Reviews Math. Phys. 28 no. 5 (2016), 12 pp.
- [IT] H. Ito, H. Tamura, Semi-classical asymptotics for total scattering cross sections of N-body quantum systems, Osaka J. Math. 32(3) (1995), 753–781.

- [JKW] T. Jecko, M. Klein, X.P. Wang, Existence and Born-Oppenheimer asymptotics of the total scattering cross-section in ion-atom collisions, Long time behaviour of classical and quantum systems (Bologna, 1999), 220–237, Ser. Concr. Appl. Math., 1, World Sci. Publ., River Edge, NJ, 2001.
- [Je] A. Jensen, Propagation estimates for Schrödinger-type operators, Trans. Amer. Math. Soc. 291 no. 1 (1985), 129–144.
- [Ji] X. Jia, Some threshold spectral problems of Schrödinger operators, thesis, Université de Nantes, 2009.
- [JK] A. Jensen and T. Kato, Spectral properties of Schrödinger operators and timedecay of the wave functions, Duke Math. J. 46 no. 3 (1979), 583–611.
- [JMP] A. Jensen, É. Mourre, P. Perry, Multiple commutator estimates and resolvent smoothness in quantum scattering theory, Ann. Henri Poincaré 41 no. 2 (1984), 207–225.
- [JN1] A. Jensen, G. Nenciu, A unified approach to resolvent expansions at thresholds, Rev. Math. Phys., 13 (2001), 717-754.
- [JN2] A. Jensen, G. Nenciu, Erratum: "A unified approach to resolvent expansions at thresholds" [Rev. Math. Phys. 13 (2001), 717–754], Rev. Math. Phys., 2014, 675–677.
- [Ka] T. Kato, Perturbation theory for linear operators, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition.
- [La] L.J. Landau, Bessel functions: monotoicity and bounds, J. London Math. Soc.
 (2) 61 (2000), 197–215.
- [Mo] E. Mourre, Absence of singular continuous spectrum for certain selfadjoint operators, Comm. Math. Phys. 78 no. 3 (1980/81), 391–408.
- [Na] S. Nakamura, Low energy asymptotics for Schrödinger operators with slowly decreasing potentials, Comm. Math. Phys. 161 (1994), 63–76.
- [Ne] R. G. Newton, Noncentral potentials: the generalized Levinson theorem and the structure of the spectrum, J. Math. Phys., 18 (1977), 1582-1588.
- [Pe] P. Perry, Exponential bounds and semifiniteness of point spectrum for N-body Schrödinger operators, Comm. Math. Phys. 92 (1984), 481–483.
- [PSS] P. Perry, I.M. Sigal, and B. Simon, Spectral analysis of N-body Schrödinger operators, Ann. of Math. 114 no. 3 (1981), 519–567.
- [RS] M. Reed and B. Simon, Methods of modern mathematical physics I-IV, Academic Press, 1972-78.
- [Sk1] E. Skibsted, Propagation estimates for N-body Schrödinger operators, Comm. Math. Phys. 142 (1991), 67–98.
- [Sk2] E. Skibsted, Smoothness of N-body scattering amplitudes, Rev. Math. Phys. 4 no. 4 (1992), 619–658.

- [Sk3] E. Skibsted, Spectral analysis of N-body systems coupled to a bosonic field, Rev. Math. Phys. 10 (1998), 989–1026.
- [Sk4] E. Skibsted, Sommerfeld radiation condition at threshold, Comm. PDE 38 (2013), 1601–1625.
- [Sk5] E. Skibsted, Renormalized two-body low-energy scattering, Journal d'Analyse Mathématique 122 (2014), 25–68.
- [Sk6] E. Skibsted, Stationary scattering theory: the N-body long-range case, Comm. Math. Phys. (2023), https://doi.org/10.1007/s00220-023-04689-7
- [Sk7] E. Skibsted, Green functions and completeness; the 3-body problem revisited, preprint 30 May 2022, http://arxiv.org/abs/2205.15028v1.
- [SW] E. Skibsted, X. P. Wang, Two-body threshold spectral analysis, the critical case, J. Funct. Anal. 260 (2011), 1766-1794.
- [SZ] J. Sjöstrand and M. Zworski, Elementary linear algebra for advanced spectral problems, Ann. Inst. Fourier (Grenoble) 57 no. 7 (2007), 2095–2141.
- [Ta1] M. Taylor, Partial Differential Equations, Basic Theory, Springer, New York, 1996; corrected second printing 1999.
- [Ta2] M. Taylor, Partial Differential Equations II, Qualitative Studies of Linear Equations, Springer, New York, 1996; corrected second printing 1997.
- [Va1] A. Vasy, Propagation of singularities in three-body scattering. Astérisque No.
 262 (2000), vi+151 pp.
- [Va2] A. Vasy, Propagation of singularities in many-body scattering in the presence of bound states. J. Funct. Anal. 184(1) (2001), 177–272.
- [VW] A. Vasy and X.P. Wang, Smoothness and high energy asymptotics of the spectral shift function in many-body scattering, Comm. Partial Differential Equations 27 (2002), no. 11-12, 2139–2186.
- [Wa1] X.P. Wang, Microlocal resolvent estimates for N-body Schrödinger operators, J. Fac. Sci. Univ. Tokyo, Sec. IA Math. 40 (1993), 337-385.
- [Wa2] X.P. Wang, Asymptotic behavior of the resolvent of N-body Schrödinger operators near a threshold, Ann. Henri Poincaré 4 (2003), 553–600.
- [Wa3] X.P. Wang, Total Cross Sections in N-body Problems: Finiteness and High Energy Asymptotics. Comm. Math. Phys. 156 (1993), 333–354.
- [Wa4] X.P. Wang, Threshold energy resonance in geometric scattering, Proceedings of Symposium "Scattering and Spectral Theory", August 2003, Recife, Brazil. Matemática Contemporânea 26 (2004), 135–164.
- [Wa5] X.P. Wang, Asymptotic expansion in time of the Schrödinger group on conical manifolds, Ann. Inst. Fourier (Grenoble) 56 no. 6 (2006), 1903–1945.
- [Wa6] X.P. Wang, Gevrey estimates of the resolvent and sub-exponential time-decay for the heat and Schrödinger semigroups, J. Math. Pures Appl. 135 no. 9 (2020), 284-338
- [Ya1] D.R. Yafaev, The low energy scattering for slowly decreasing potentials, Comm. Math. Phys. 85 (1982), 177–196.
- [Ya2] D.R. Yafaev, Spectral properties of the Schrödinger operator with a potential having a slow falloff, Funct. Anal. and its Appl. (Steklov Math. Inst.) 16 (1982), 47–54. (English translation, 1983.)
- [Ya3] D.R. Yafaev, Scattering theory: some old and new problems, Lecture Notes in Mathematics, 1735. Springer-Verlag, Berlin, 2000. xvi+169 pp.
- [Ya4] D.R.Yafaev, The eikonal approximation and the asymptotics of the total scattering cross-section for the Schrödinger equation. Ann. Inst. Henri Poincaré, 40 (4) (1986), 397–425.
- [Ya5] D.R. Yafaev, Resolvent estimates and scattering matrix for N-body Hamiltonians, Intgr. Equat. Oper. Th. 21 (1995), 93–126.