EXTENDING CERCIGNANI'S CONJECTURE RESULTS FROM BOLTZMANN TO BOLTZMANN-FERMI-DIRAC EQUATION

THOMAS BORSONI

Sorbonne Université, CNRS, Université Paris Cité, Laboratoire Jacques-Louis Lions (LJLL), F-75005 Paris, France

ABSTRACT. We establish a connection between the relative Classical entropy and the relative Fermi-Dirac entropy, allowing to transpose, in the context of the Boltzmann or Landau equation, any entropy-entropy production inequality from one case to the other; therefore providing entropy-entropy production inequalities for the Boltzmann-Fermi-Dirac operator, similar to the ones of the Classical Boltzmann operator. We also provide a generalized version of the Csiszár-Kullback-Pinsker inequality to weighted L^p norms, $1 \le p \le 2$ and a wide class of entropies.

Contents

1. Introduction	2
1.1. The Boltzmann-Fermi-Dirac operator	2
1.2. Macroscopic quantities and equilibrium distribution	3
1.3. Fermi-Dirac entropy	4
1.4. Entropy production	6
1.5. Trend to equilibrium and Cercignani's conjecture	6
2. Main results	7
2.1. Comparison of the Fermi-Dirac and Classical relative entropies	7
2.2. Entropy-entropy production inequalities for the Boltzmann-Fermi-Dirac (BFD) equation	7
2.3. About the Csiszár-Kullback-Pinsker inequality	8
2.4. Convergence towards equilibrium	9
3. Proof of Theorem 1 and Proposition 2	10
3.1. Key element of the proof	11
3.2. Continuity at zero	13
3.3. Final proof of Theorem 1	14
3.4. Final proof of Proposition 2	15
4. Proof of Corollary 3	17
4.1. Super quadratic kernels	17
4.2. General kernels	18
4.3. Inverse result	19
5. About the Landau-Fermi-Dirac equation	20
6. L^1-L^2 weighted Csiszár-Kullback-Pinsker inequality	22
Appendix A. Similar results in the Bose-Einstein case	25
Appendix B. A general discussion about entropies and equilibria	27
Appendix C. Technical results	30
C.1. L^{∞} bound for the Fermi-Dirac statistics	31
C.2. Regularity in ε of the coefficients of the Fermi-Dirac statistics	32
Acknowledgements	34
References	34

E-mail address: thomas.borsoni@sorbonne-universite.fr. *Date*: February 9, 2024.

Key words and phrases. kinetic theory, Nordheim equation, Boltzmann equation, Boltzmann-Fermi-Dirac, Fermi-Dirac statistics, entropy, entropy methods, Cercignagni's conjecture, Csiszár-Kullback-Pinsker inequality.

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

1. INTRODUCTION

In the study of quantum dilute gases, Nordheim [25], Uehling and Uhlenbeck [28] proposed in the early 1930s a kinetic approach, with a modified Boltzmann equation - bearing the names Nordheim equation, quantum Boltzmann equation, or Boltzmann-Fermi-Dirac equation (for fermions). Quoting Villani [29, Chapter 5.3], this description "does not rest on the traditional quantum formalism, [...] it is rather a classical description of interacting particles with quantum features", which nevertheless allows to recover the expected Fermi-Dirac statistics (in the context of fermions) and Bose-Einstein statistics (in the context of bosons). In this paper, we focus on the case of fermions in which Pauli's exclusion principle is taken into account.

In the context of the study of trend to equilibrium, the use of entropy-entropy production inequalities, allowing to obtain explicit rates of convergence, has been increasing in the recent years. We mention the results of Villani [30] for the Classical Boltzmann operator, Desvillettes [13], Desvillettes and Villani [17], Carrapatoso, Desvillettes and He [10] for the Landau operator, Alonso, Bagland, Desvillettes and Lods [1, 2] and Alonso, Bagland, and Lods [3] for the Landau-Fermi-Dirac (LFD) operator. We also refer the reader to the reviews [29, 16]. Up to now, no such results were known for the Boltzmann-Fermi-Dirac (BFD) operator, and the initial goal of this work was to fill this gap. In doing so, we actually established a connection between the Classical and Fermi-Dirac cases, allowing to transpose any entropy-entropy production result from one setting to the other under suitable conditions. Entropy-entropy production estimates for the BFD operator then directly follow from the ones of Villani [30] for the Classical Boltzmann operator - the same procedure can be applied to transpose entropy-entropy production estimates for the Landau operator to ones for the LFD operator.

This paper is constructed as follows. In the rest of this section, we introduce our mathematical setting, useful definitions and further discuss our motivations. In Section 2, we state our main results, which are the connection between the Fermi-Dirac and Classical relative entropies (Theorem 1 and Proposition 2), the entropy-entropy production estimates for the BFD operator (Corollary 3), and a generalization of the Csiszár-Kullback-Pinsker inequality (Proposition 4). We provide the proof of Theorem 1 in Section 3, of Corollary 3 in Section 4, discuss the case of the Landau / LFD equation in Section 5 and provide the proof of Proposition 4 (along with a more precise result and discussions) in Section 6. We very briefly discuss the case of bosons in Appendix A. Considerations about a wider class of entropies (with a generalization of Proposition 4) are provided in Appendix B. In particular, we provide in Proposition 15 a general formula for the minimizer of a general entropy under the constraints given by general conserved quantities, under a sole condition of existence. There are links with the results of [7] where equilibria of collision kernels of type "two particles give two particles" appearing in weak turbulence theory are rigorously obtained. In fact, Proposition 15 does not replace a rigorous study of equilibria, such as in [7], but gives insights, in a general setting, on to which distributions we should be expected to be equilibria. Finally, technical results relevant to our study are proven in Appendix C.

1.1. **The Boltzmann-Fermi-Dirac operator.** The spatially homogeneous Boltzmann-Fermi-Dirac (BFD) equation writes

(1)
$$\partial_t f = Q_{\varepsilon}(f, f), \qquad f(0, \cdot) = f^{in},$$

where $f \equiv f(t, v) \geq 0$ represents a density of fermions (quantum particles of half-integer spin, e.g. electrons), depending on time $t \geq 0$ and velocity $v \in \mathbb{R}^3$, $f^{in} \equiv f^{in}(v) \geq 0$ is an initial distribution and Q_{ε} stands for the BFD operator. We refer the reader to [12, 18, 22, 24] for results on the Cauchy problem associated to (1). In this paper, we focus on the operator Q_{ε} and, if not specified otherwise, discard the time variable t, considering $f \equiv f(v) \geq 0$.

In dimension 3, the BFD operator writes, for $\varepsilon > 0$ and measurable $f : \mathbb{R}^3 \to [0, \varepsilon^{-1}]$,

(2)
$$Q_{\varepsilon}(f,f)(v) := \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \left[f' f'_* (1-\varepsilon f)(1-\varepsilon f_*) - f f_* (1-\varepsilon f')(1-\varepsilon f'_*) \right] B(v,v_*,\sigma) \,\mathrm{d}\sigma \,\mathrm{d}v_*,$$

where we used the standard shorthands $f \equiv f(v), f_* \equiv f(v_*), f' \equiv f(v'), f'_* \equiv f(v'_*)$ and

(3)
$$v' = \frac{v + v_*}{2} + \frac{|v - v_*|}{2}\sigma, \quad v'_* = \frac{v + v_*}{2} - \frac{|v - v_*|}{2}\sigma.$$

We insist that any physically meaningful solution f to (1) should be such that $1 - \varepsilon f \ge 0$. As a consequence of time micro-reversibility and Galilean invariance properties, the collision kernel $B \ge 0$ is assumed to satisfy, for $(v, v_*) \in \mathbb{R}^3 \times \mathbb{R}^3$ and $\sigma \in \mathbb{S}^2$,

(4)
$$B(v, v_*, \sigma) \equiv B(|v - v_*|, \cos \theta),$$

with $\cos \theta = \sigma \cdot \frac{v - v_*}{|v - v_*|}$. At this level of generality, no other assumption is made on *B*. Formally choosing $\varepsilon = 0$ in (2) yields the Classical Boltzmann operator

(5)
$$Q_0(f,f)(v) := \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \left[f' f'_* - f f_* \right] B(v,v_*,\sigma) \,\mathrm{d}\sigma \,\mathrm{d}v_*,$$

with the same shorthands as in (2). Now for $\varepsilon \geq 0$, define

(6)
$$\varphi_{\varepsilon}(x) := \frac{x}{1 - \varepsilon x}, \quad \text{for } x \in \mathbb{R}_+ \quad \text{s.t.} \quad 1 - \varepsilon x > 0.$$

Remark that when $\varepsilon = 0$, corresponding to the classical case, we have $\varphi_0 = \mathrm{Id}_{\mathbb{R}_+}$. Moreover, φ_{ε} is a \mathcal{C}^1 -diffeomorphism from $[0, \varepsilon^{-1})$ to \mathbb{R}_+ , with $\varphi_{\varepsilon}^{-1} = \varphi_{-\varepsilon}$.

Then, when f is such that $1 - \varepsilon f > 0$ almost everywhere, Equation (2) can be rewritten in the following form resembling the Classical Boltzmann operator (5),

(7)
$$Q_{\varepsilon}(f,f)(v) := \iint_{\mathbb{R}^3 \times \mathbb{S}^2} \left[\varphi_{\varepsilon}(f)' \varphi_{\varepsilon}(f)'_* - \varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f)_* \right] B^f_{\varepsilon}(v,v_*,\sigma) \,\mathrm{d}\sigma \,\mathrm{d}v_*$$

using the same shorthands as in (2), and

(8)
$$B_{\varepsilon}^{f}(v, v_{*}, \sigma) = (1 - \varepsilon f(v))(1 - \varepsilon f(v_{*}))(1 - \varepsilon f(v'))(1 - \varepsilon f(v'_{*})) B(v, v_{*}, \sigma)$$

The collision kernel B_{ε}^{f} straightforwardly satisfies the usual assumptions of symmetry and micro-reversibility which imply the weak form of the classical Boltzmann operator, (see for instance [12]), so that for any test function $\psi : \mathbb{R}^{3} \to \mathbb{R}$, (9)

$$\int_{\mathbb{R}^3} \psi(v) Q_{\varepsilon}(f, f)(v) \, \mathrm{d}v = -\frac{1}{4} \iiint_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} \left[\varphi_{\varepsilon}(f)' \varphi_{\varepsilon}(f)'_* - \varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f)_* \right] \left(\psi' + \psi'_* - \psi - \psi_* \right) B_{\varepsilon}^f \, \mathrm{d}\sigma \, \mathrm{d}v \, \mathrm{d}v_*,$$

still using the same shorthands as in (2) for $\varphi_{\varepsilon}(f)$ and ψ . Formally taking $\psi(v) = 1$, $\psi(v) = v_i$ for i = 1, 2, 3and $\psi(v) = \frac{1}{2}|v|^2$ in (9) yields the conservation of mass, momentum and energy

(10)
$$\int_{\mathbb{R}^3} \begin{pmatrix} 1\\v\\\frac{1}{2}|v|^2 \end{pmatrix} Q_{\varepsilon}(f,f)(v) \,\mathrm{d}v = 0.$$

In the following, we define, for $p \geq 1$ and $s \in \mathbb{R}$, the weighted L_s^p norm of some function g by

(11)
$$\|g\|_{L^p_s} := \left(\int_{\mathbb{R}^3} |g(v)|^p \left(1 + |v|^2\right)^{\frac{ps}{2}} \mathrm{d}v\right)^{\frac{1}{p}},$$

and its essential supremum by $||g||_{\infty}$.

1.2. Macroscopic quantities and equilibrium distribution. Throughout this paper, for any nonnegative $f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$, we denote $(\rho, u, T) \in \mathbb{R}^*_+ \times \mathbb{R}^3 \times \mathbb{R}^*_+$ respectively the density, mean velocity and temperature associated to f, in the sense that

(12)
$$\int_{\mathbb{R}^3} f(v) \begin{pmatrix} 1\\v\\|v|^2 \end{pmatrix} dv = \begin{pmatrix} \rho\\\rho u\\3\rho T + \rho|u|^2 \end{pmatrix}$$

We also define the minimal directional temperature of f,

(13)
$$T_*(f) := \frac{1}{\rho} \inf_{e \in \mathbb{S}^2} \int_{\mathbb{R}^3} f(v) \, (v \cdot e)^2 \, \mathrm{d}v,$$

as well as the so-called Fermi temperature associated to ρ and $\varepsilon > 0$,

(14)
$$T_F(\rho,\varepsilon) := \frac{1}{2} \left(\frac{3\varepsilon\rho}{4\pi}\right)^{\frac{4}{3}},$$

and the ratio

(15)
$$\gamma := \frac{T}{T_F(\rho,\varepsilon)},$$

with the convention $\gamma = +\infty$ when $\varepsilon = 0$. The adimensional number γ is thus the ratio between the actual temperature of the system and its Fermi temperature. In [22], Lu proved that

(16)
$$1 - \varepsilon f \ge 0 \implies \gamma \ge \frac{2}{5},$$

as well as, for any fixed $\varepsilon > 0$,

(17)
$$\left(1 - \varepsilon f \ge 0 \text{ and } \gamma = \frac{2}{5}\right) \iff f \equiv F_{\varepsilon}^{\rho} := \varepsilon^{-1} \mathbf{1}_{|\cdot -u| \le \left(\frac{3\varepsilon\rho}{4\pi}\right)^{\frac{1}{3}}},$$

(18)
$$\left(1 - \varepsilon f \ge 0 \text{ and } \gamma > \frac{2}{5}\right) \iff \text{there exists a unique } \varepsilon\text{-Fermi-Dirac distribution } \mathcal{M}_{\varepsilon}^{f} \text{ such that}$$

$$\int_{\mathbb{R}^3} \mathcal{M}^f_{\varepsilon}(v) \begin{pmatrix} 1\\v\\|v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho\\\rho u\\ 3\rho T + \rho |u|^2 \end{pmatrix},$$

where we recall that the ε -Fermi-Dirac distributions are the ones which are such that $\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})$ is a Maxwellian distribution. Equivalently, ε -Fermi-Dirac distributions are the distributions defined by

(19)
$$\mathcal{M}_{\varepsilon}^{f}(v) = \frac{e^{a_{\varepsilon} + b_{\varepsilon}|v - u_{\varepsilon}|^{2}}}{1 + \varepsilon e^{a_{\varepsilon} + b_{\varepsilon}|v - u_{\varepsilon}|^{2}}}, \qquad a_{\varepsilon} \in \mathbb{R}, \ u_{\varepsilon} \in \mathbb{R}^{3}, \ b_{\varepsilon} < 0.$$

In particular, a characterisation of the Fermi-Dirac (or of the Maxwellian, when $\varepsilon = 0$) distribution $\mathcal{M}_{\varepsilon}^{f}$ is that $\log \circ \varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})$ is a linear combination of $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^{2}$ (that is, of the conserved quantities). The interested reader shall find further discussions on this matter in Appendix B.

As proven in [22], F_{ε}^{ρ} and $\mathcal{M}_{\varepsilon}^{f}$ are the only two possible equilibria of the (spatially homogeneous) BFD dynamic (1). Contrary to $\mathcal{M}_{\varepsilon}^{f}$ (called first equilibrium in [22]), which is an attractor, the distribution F_{ε}^{ρ} (called second equilibrium in [22]) is not, and corresponds to a degenerate state, which can actually occur for very cold gases (note that (16)–(17) imply that F_{ε}^{ρ} is of "lowest" temperature).

From now on, we denote by M^g the Maxwellian distribution associated to a distribution g, in the sense that

$$M^g \equiv \mathcal{M}_0^g.$$

1.3. Fermi-Dirac entropy. We now define the ε -entropy of the distribution f.

Definition 1. We define, for any $\varepsilon \geq 0$ and $x \in \mathbb{R}_+$ such that $1 - \varepsilon x \geq 0$,

(20)
$$\Phi_{\varepsilon}(x) := \int_0^x \log \varphi_{\varepsilon}(y) \, \mathrm{d}y$$

where φ_{ε} is defined in (6). The function Φ_{ε} is well-defined as $\log \in L^1_{loc}(\mathbb{R}_+)$. For any nonnegative $f, g \in L^1_2(\mathbb{R}^3)$ such that $1 - \varepsilon f \ge 0$ and $1 - \varepsilon g \ge 0$, we define the ε -entropy of f by

(21)
$$H_{\varepsilon}(f) := \int_{\mathbb{R}^3} \Phi_{\varepsilon}(f)(v) \, \mathrm{d}v,$$

as well as the relative ε -entropy of f and g by

(22)
$$H_{\varepsilon}[f|g] := H_{\varepsilon}(f) - H_{\varepsilon}(g).$$

Remark 1. Our definition of entropy slightly differs from the usual ones, but is actually equivalent in the problems that we consider. We choose to define the entropy with (20)–(21) because, our study relying on a Taylor expansion, it is here the more natural definition. Let us detail the link between the ε -entropy and usual definitions.

• Classical entropy. If $\varepsilon = 0$, then $\Phi_0(x) = x \log x - x$ with the convention $0 \log 0 = 0$, the condition $1 - \varepsilon f \ge 0$ is always satisfied, and

$$H_0(f) = \int_{\mathbb{R}^3} (f \log f - f) \,\mathrm{d}v.$$

Note that often (consider for instance [30]), the classical entropy is defined by

$$\widetilde{H}_0(f) = \int_{\mathbb{R}^3} f \log f \, \mathrm{d}v \equiv \int_{\mathbb{R}^3} \widetilde{\Phi}_0(f) \, \mathrm{d}v,$$

with $\widetilde{\Phi}_0(x) = x \log x$.

• Fermi-Dirac entropy. If $\varepsilon > 0$, then $\Phi_{\varepsilon}(x) = x \log x + \varepsilon^{-1}(1 - \varepsilon x) \log(1 - \varepsilon x)$ with the convention $0 \log 0 = 0$, and the condition $1 - \varepsilon f \ge 0$ is satisfied if and only if $f \le \varepsilon^{-1}$. In this case, we have

$$H_{\varepsilon}(f) = \int_{\mathbb{R}^3} f \log f + \varepsilon^{-1} (1 - \varepsilon f) \log(1 - \varepsilon f) \, \mathrm{d}v.$$

Note that for instance in [1] (up to a minus sign, which is just a choice of convention), the Fermi-Dirac entropy is defined by

$$\widetilde{H}_{\varepsilon}(f) = \varepsilon^{-1} \int_{\mathbb{R}^3} (\varepsilon f) \log(\varepsilon f) + (1 - \varepsilon f) \log(1 - \varepsilon f) \, \mathrm{d}v \equiv \int_{\mathbb{R}^3} \widetilde{\Phi}_{\varepsilon}(f) \, \mathrm{d}v,$$

with
$$\Phi_{\varepsilon}(x) = \varepsilon^{-1} \left(\varepsilon x \log(\varepsilon x) + (1 - \varepsilon x) \log(1 - \varepsilon x) \right)$$

As we are only interested in the relative (Classical of Fermi-Dirac) entropy between f and its associated equilibrium distribution $\mathcal{M}_{\varepsilon}^{f}$ (defined in (18)), we see in the following Proposition 1 that only Φ_{ε}'' is actually relevant to the definition. From $\tilde{\Phi}_{\varepsilon}'' = \Phi_{\varepsilon}''$, we then obtain $\tilde{H}_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}] = H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}]$, ensuring the announced equivalence.

Proposition 1. For any $\varepsilon \ge 0$ and nonnegative $f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge 0$ and either $\gamma > \frac{2}{5}$ if $\varepsilon > 0$ or $f \in L \log L(\mathbb{R}^3)$ if $\varepsilon = 0$, denoting by $\mathcal{M}_{\varepsilon}^f$ the Fermi-Dirac (or Maxwellian if $\varepsilon = 0$) distribution associated to f, we have the following representation of the relative ε -entropy between f and $\mathcal{M}_{\varepsilon}^f$,

(23)
$$H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}] = \int_{0}^{1} (1-\tau) \left[\int_{\mathbb{R}^{3}} (f-\mathcal{M}_{\varepsilon}^{f})^{2} \Phi_{\varepsilon}^{\prime\prime} \left((1-\tau)\mathcal{M}_{\varepsilon}^{f} + \tau f \right) dv \right] d\tau$$
$$\int_{0}^{1} \int_{0}^{f(v)} f(v) - \tau$$

(24)
$$= \int_{\mathbb{R}^3} \int_{\mathcal{M}_{\varepsilon}^f(v)}^{f(v)} \frac{f(v) - x}{\varphi_{\varepsilon}(x)} \varphi_{\varepsilon}'(x) \, \mathrm{d}x \, \mathrm{d}v.$$

It holds that

(25)
$$H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}] \ge 0 \quad and \quad (H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}] = 0 \iff f = \mathcal{M}_{\varepsilon}^{f}).$$

The reader will find the proof of (23) and (25) in Appendix B - see Proposition 15. We just point out here that this comes straightforwardly from a Taylor expansion of the function Φ_{ε} , and that (25) is a direct consequence of (23) because $\Phi_{\varepsilon}'' > 0$ on $(0, \varepsilon^{-1})$. Equation (24) comes directly from $\Phi_{\varepsilon}'' = \frac{\varphi_{\varepsilon}'}{\varphi_{\varepsilon}}$ and the change of variable, for fixed $v, x = (1 - \tau) \mathcal{M}_{\varepsilon}^{f}(v) + \tau f(v)$ inside (23).

Remark 2. We point out that for any $0 < x < \varepsilon^{-1}$, we have

$$\Phi_{\varepsilon}''(x) = \frac{\varphi_{\varepsilon}'(x)}{\varphi_{\varepsilon}(x)} = \frac{1}{x(1-\varepsilon x)}$$

Hence for any $\tau \in (0,1)$, $(1-\tau)\mathcal{M}_{\varepsilon}^{f} + \tau f \in (0,\varepsilon^{-1})$, and $\Phi_{\varepsilon}''((1-\tau)\mathcal{M}_{\varepsilon}^{f} + \tau f)$ is thus well-defined. Moreover, with $\varepsilon = 0$ (Classical case), Equations (23)-(24) write

(26)
$$H_0\left[f|M^f\right] = \int_0^1 (1-\tau) \left[\int_{\mathbb{R}^3} \frac{(f-M^f)^2}{(1-\tau)M^f + \tau f} \,\mathrm{d}v\right] \mathrm{d}\tau$$

(27)
$$= \int_{\mathbb{R}^3} \int_{M^f(v)}^{J(v)} \frac{f(v) - x}{x} \, \mathrm{d}x \, \mathrm{d}v.$$

1.4. Entropy production. We finally define the entropy production $\mathcal{D}_{\varepsilon}$ of the BFD operator by

(28)
$$\mathcal{D}_{\varepsilon}(f) := -\int_{\mathbb{R}^3} \log \varphi_{\varepsilon}(f) Q_{\varepsilon}(f, f)(v) \, \mathrm{d}v$$

We get, taking $\psi = \log \varphi_{\varepsilon}(f)$ in (9),

(29)
$$\mathcal{D}_{\varepsilon}(f) = \frac{1}{4} \iiint_{\mathbb{R}^{3} \times \mathbb{R}^{3} \times \mathbb{S}^{2}} \left[\varphi_{\varepsilon}(f)' \varphi_{\varepsilon}(f)_{*}' - \varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f)_{*} \right] \log \left(\frac{\varphi_{\varepsilon}(f)' \varphi_{\varepsilon}(f)_{*}'}{\varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f)_{*}} \right) B_{\varepsilon}^{f} \, \mathrm{d}\sigma \, \mathrm{d}v \, \mathrm{d}v_{*} \ge 0.$$

Formally, a solution f to the BFD equation (1) satisfies

$$\frac{\mathrm{d}}{\mathrm{d}t}H_{\varepsilon}(f(t,\cdot)) = -\mathcal{D}_{\varepsilon}(f(t,\cdot))$$

or, using the relative entropy,

(30)
$$\frac{\mathrm{d}}{\mathrm{d}t}H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] = -\mathcal{D}_{\varepsilon}(f(t,\cdot)).$$

The nonnegativity of $\mathcal{D}_{\varepsilon}$ then ensures the second principle of thermodynamics.

1.5. Trend to equilibrium and Cercignani's conjecture. The relative entropy between a solution at a given time t and its associated equilibrium is a very useful quantity to estimate how close to equilibrium the system is, and (25) in Proposition 1 makes it a suitable tool into proving convergence towards equilibrium. Indeed, if one shows for some solution f that

$$H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] \xrightarrow[t \to \infty]{} 0,$$

then the Csiszár-Kullback-Pinsker (CKP) inequalities, discussed below in Proposition 4 and Section 6 yield

$$||f(t,\cdot) - \mathcal{M}_{\varepsilon}^{f^{in}}||_{L^{1}_{2}} \xrightarrow[t \to \infty]{} 0,$$

with a rate of convergence related to the one of the relative entropy. In the study of large-time behaviour of the solutions to the (Classical of Fermi-Dirac) Boltzmann equation, the relative entropy is thus a major tool. The idea of entropy-entropy production methods is to prove an inequality of the type

$$\mathcal{D}_{\varepsilon}(f) \ge \Theta(H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}])$$

for some suitable function Θ (typically, a linear or power law function), allowing to obtain, combined with (30) and using a Gronwall-type argument, an explicit rate of the convergence towards 0 of the relative entropy - thus of the L_2^1 distance, using the CKP inequality.

In the early 80's, Cercignani conjectured in [11], in the context of the Classical Boltzmann equation with suitable collision kernels, that for any distribution f, there exists some $\lambda > 0$ depending only on density, temperature and upper bound on the entropy of f (this is called the strong form of Cercignani's conjecture) such that

$$\mathcal{D}_0(f) \ge \lambda \,\rho \, H_0[f|M^f],$$

where we recall that ρ is the density associated to f, \mathcal{D}_0 is the entropy production in the classical case, H_0 the classical entropy and M^f the Maxwellian associated to f. It is now known that this is false in general (see the negative result of Bobylev and Cercignani [5]), however Villani proved in [30] the strong form of this conjecture when the collision kernel B is "super-quadratic", as well as a weak form $\mathcal{D}_0(f) \geq C(f) H_0[f|M^f]^{1+\delta}$, holding for all $\delta > 0$, for general collision kernels (and C(f) depending on various quantities related to f). We mention the cornerstone papers of Toscani and Villani [26, 27] on this topic, and refer the reader to [16] for a review of Cercignani's conjecture results for the Boltzmann and Landau equations.

Similar ideas hold in the study of the Landau equation (we shall talk about Cercignani's conjecture for the Landau equation), this topic is further addressed in Section 5.

The motivation of this work is to provide Cercignani's conjecture type results in the case of the Boltzmann-Fermi-Dirac equation, uniformly with respect to $\varepsilon > 0$.

The simple yet fundamental idea which is at the root of this work is the fact that, whenever $\varepsilon > 0$, assuming $1 - \varepsilon f \ge \kappa_0$ for some $\kappa_0 \in (0, 1)$ straightforwardly implies

$$B(v, v_*, \sigma) \ge B_{\varepsilon}^f(v, v_*, \sigma) \ge \kappa_0^4 B(v, v_*, \sigma),$$

so that, as log in nonincreasing,

(31)
$$\mathcal{D}_0\left(\varphi_{\varepsilon}(f)\right) \ge \mathcal{D}_{\varepsilon}(f) \ge \kappa_0^4 \mathcal{D}_0\left(\varphi_{\varepsilon}(f)\right)$$

Therefore, relatively to κ_0 , the quantity $\mathcal{D}_{\varepsilon}(f)$ (the Fermi-Dirac entropy production associated to f) behaves like $\mathcal{D}_0(\varphi_{\varepsilon}(f))$ (the Classical entropy production associated to $\varphi_{\varepsilon}(f)$). Our main results, Theorem 1 and Proposition 2, allow to state a similar property for the relative entropies. We find that, relatively to κ_0 , the quantity $H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}]$ (the Fermi-Dirac relative-entropy between f and $\mathcal{M}_{\varepsilon}^{f}$) behaves like $H_0[\varphi_{\varepsilon}(f)|\mathcal{M}^{\varphi_{\varepsilon}(f)}]$ (the Classical relative entropy between $\varphi_{\varepsilon}(f)$ and $\mathcal{M}^{\varphi_{\varepsilon}(f)}$). This entails that **any entropy-entropy production inequality in the Classical case yields the same inequality** (with an almost identical constant) **in the Fermi-Dirac case**; this corresponds to Corollary 3 (and Proposition 8 for the Landau/Landau-Fermi-Dirac adaptation). The same holds for counter-examples to entropy-entropy production inequalities.

2. Main results

In this section, we state our main results.

2.1. Comparison of the Fermi-Dirac and Classical relative entropies. Theorem 1 is the main contribution of this work, as it establishes a strong link between the different entropies, allowing to transfer "positive" Cercignani's conjecture results from the classical to the fermionic case.

Theorem 1. Lower-bound inequality. For any $\varepsilon > 0$ and nonnegative $f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ such that

$$1 - \varepsilon f > 0$$
 and $\frac{f}{1 - \varepsilon f} \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3),$

 $we\ have$

(32)
$$H_0\left[\frac{f}{1-\varepsilon f}\middle|M^{\frac{f}{1-\varepsilon f}}\right] \ge H_{\varepsilon}[f|\mathcal{M}^{f}_{\varepsilon}].$$

The hypotheses of Theorem 1 are actually the minimal ones to ensure that every term in (32) makes sense. The following Proposition 2 allows to transfer "negative" Cercignani's conjecture results from the classical to the fermionic case.

Proposition 2. Upper-bound inequality. For any $\kappa_0 \in (0, 1)$, any $\varepsilon > 0$ and nonnegative $f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that

(33)
$$1 - \varepsilon f \ge \kappa_0$$

we have

(34)
$$H_0\left[\frac{f}{1-\varepsilon f}\middle| M^{\frac{f}{1-\varepsilon f}}\right] \le C_0(\kappa_0) H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^f],$$

with $C_0(\kappa_0) = \exp\left(16\left(\kappa_0^{-1} - 1\right)\right)$.

Remark 3. Inequality (32) becomes reversed in the Bose-Einstein case. We refer the interested reader to Proposition 12 in Appendix A. In fact, we expect that both inequalities, but reversed (and probably with a different constant C_0), hold in the Bose-Einstein case, as the core of the proof would be the exact same in this case, only changing ε into $-\varepsilon$ (hence the reversal of the inequalities).

2.2. Entropy-entropy production inequalities for the Boltzmann-Fermi-Dirac (BFD) equation. As a corollary to Theorem 1 and Proposition 2, adapting the main results found in Villani [30], we obtain the following entropy-entropy production estimates.

Corollary 3. • Cercignani's conjecture result for the BFD operator with super-quadratic kernels (adaptation of [30, Theorem 2.1]). Assume that B satisfies (4) and

$$B(v, v_*, \sigma) \ge 1 + |v - v_*|^2.$$

Then, for any $\kappa_0 \in (0,1)$, $\varepsilon \geq 0$ and nonnegative $f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \geq \kappa_0$, we have

(35)
$$\mathcal{D}_{\varepsilon}(f) \ge K(f) H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}],$$

(36)
$$K(f) = \frac{2\pi}{7} \kappa_0^5 \min(1, T) \rho T_*(f),$$

where ρ , T, $T_*(f)$ are defined in (12)–(13), $\mathcal{D}_{\varepsilon}$, H_{ε} and $\mathcal{M}^f_{\varepsilon}$ are given respectively in (28)–(29), (21)–(22) and (18) (note that $T_*(f)$ can be bounded below using the entropy or an L^{∞} bound [17, Proposition 2]).

• Super-linear Cercignani's conjecture result for the BFD operator with general kernels (adaptation of [30, Theorem 4.1]). Assume the existence of $B_0 > 0$ and $\beta_+, \beta_- \ge 0$ such that

(37)
$$B(v, v_*, \sigma) \ge B_0 \min\left(|v - v_*|^{\beta_+}, |v - v_*|^{-\beta_-}\right)$$

and B satisfies (4). We consider $\kappa_0 \in (0,1), 0 \leq f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ and $\varepsilon \geq 0$ such that $1 - \varepsilon f \geq \kappa_0$ and

(38)
$$\forall v \in \mathbb{R}^3, \quad f(v) \ge K_0 e^{-A_0 |v|^{q_0}} \quad (K_0 > 0, \quad A_0 > 0, \quad q_0 \ge 2).$$

Then for any $\alpha \in (0,1)$ there exists an explicit constant $K_{\alpha}(f)$ depending on α , B_0 , an upper bound of ε and on f only through ρ , T, q_0 , and upper bounds on A_0 , $1/K_0$, $1/\kappa_0$, $||f||_{L^1_s}$ and $||f||_{H^k}$, where $s = s(\alpha, q_0, \beta_-, \beta_+)$ and $k = k(\alpha, s, \beta_-, \beta_+)$, such that

(39)
$$\mathcal{D}_{\varepsilon}(f) \ge K_{\alpha}(f) H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}]^{1+\alpha}.$$

• Counter-example to Cercignani's conjecture for the BFD operator with sub-quadratic kernels (weak adaptation of [5] and [30, Theorem 1.1]). Assume that B satisfies (4) and that there exists $B_0 > 0$ and $0 \le \beta < 2$ such that

$$\int_{\mathbb{S}^2} B(v, v_*, \sigma) \,\mathrm{d}\sigma \le B_0 \left(1 + |v - v_*|^\beta\right).$$

Then for any $\varepsilon \in (0, \frac{1}{2})$, we have

$$\inf\left\{\frac{\mathcal{D}_{\varepsilon}(f)}{H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}]}\right| f \in \mathcal{C}_{1,0,1}^{\varepsilon} \text{ and } f(v) \geq \frac{1}{2}(2\pi)^{-3/2} e^{-|v|^{2}}, \ v \in \mathbb{R}^{3}\right\} = 0,$$

with

$$f \in \mathcal{C}^{\varepsilon}_{1,0,1} \iff 0 \le f \in L^{1}_{2}(\mathbb{R}^{3}), \quad 1 - \varepsilon f \ge 0, \quad 1 \le \rho \le 1 + 2\varepsilon, \quad |u| \le 4\sqrt{3}\varepsilon, \quad \frac{1}{1 + 2\varepsilon} \le T \le 1 + 2\varepsilon.$$

We recall that ρ, u and T are defined in (12). The last part of the corollary (the counter-example) is a weak version of [5, Theorem 1], due to the presence of $C_{1,0,1}^{\varepsilon}$ instead of the more expected $C_{1,0,1}^{0}$ (the set of normalized functions) and the lack of a condition on the $\|\cdot\|_{L_s^1}$ and $\|\cdot\|_{H^k}$ norms. One could probably replace $C_{1,0,1}^{\varepsilon}$ by $C_{1,0,1}^{0}$ and add the $\|\cdot\|_{L_s^1}$ and $\|\cdot\|_{H^k}$ conditions with further technical work.

2.3. About the Csiszár-Kullback-Pinsker inequality. The Csiszár-Kullback-Pinsker (CKP) inequality is a well-known link between the total variation distance of two probability measures and their relative entropy, which is often used to transform a convergence in relative entropy into a convergence in an actual norm, such as L^1 or L_2^1 . In conducting our study, we were led to prove Proposition 4 (or the optimal Proposition 10), which are generalizations of the CKP inequality.

A generalization of the CKP inequality allowing for a weight in the total variation distance, called "weighted Csiszár-Kullback-Pinsker inequalities" was already provided by Bolley and Villani in [6, Theorem 2.1]. However our inequality does differ from theirs. On the one hand, as far as drawbacks are concerned, we need the reference measure to be an equilibrium distribution while it is not the case in [6], and our constant depends on f while it doesn't in the inequality in [6]. On the other hand, for the improvements, our constant seems more precise and a great deal smaller (possibly the constant in Proposition 10 is optimal), in the sense that choosing a weight equal to $|v|^{\alpha}$ with $\alpha > 2$ yields a finite constant in our case but an infinite one in [6, Theorem 2.1]. In particular, remark that by Jensen's inequality, $||M \, \varpi^2||_{L^1} \leq \log \int e^{\varpi^2} M \, dv$.

In another direction, in the study of the Vlasov-Poisson system, Cáceres et al. provided a generalization of the CKP inequality to L^p norms with $1 \le p \le 2$ in [8, Proposition 3.1] (specifically in the note [9]) for various different entropies, which formulation and proof is quite close to our result. However, as far as the

9

standard entropy is concerned, their result works only for p = 1, is without weights, and their constant is not optimal. I am not aware of other such generalizations of the CKP inequality in the literature.

Finally, the main idea of our proof of this inequality was already present in Jüngel's [20, proof of Theorem A.2], and he also proposed a generalization in [20, Theorem A.3], which differs from ours and holds for the L^1 norm.

In the following Proposition 4 (and in Proposition 10 in Section 6), we propose an extension the CKP inequality to weighted L^p distances for $p \in [1, 2]$, as well as to other entropies than the Classical entropy (typically, the Fermi-Dirac and the Bose-Einstein entropies), which could be seen as a extension of the previously mentionned three generalizations. We denote this extension by L^1-L^2 weighted Csiszár-Kullback-Pinsker inequality.

We moreover wish to highlight the fact that the proof of these inequalities, which is the same as in [20, Theorem A.2], but with the addition of weights and of considering L^p , $1 \le p \le 2$, relies on a Hölder inequality. In particular, our proof of the original CKP inequality relies on a Taylor expansion and a Cauchy-Schwarz

argument, instead of the tricky inequality (see for instance [19]) $x \log x - x + 1 \ge \frac{3}{2} \times \frac{(x-1)^2}{x+2}$.

For the sake of readability, we present hereafter a simplified non-optimal version of the proposition, and refer the reader to Section 6 for the refined and more general version of it, where it is also extended to the Bose-Einstein case and further discussed. We also extend it to a broader class of general entropies (Φ -entropies in the text) in Proposition 16 of Appendix B.

Proposition 4. L^1 - L^2 weighted Csiszár-Kullback-Pinsker inequality related to the Classical and Fermi-Dirac relative entropies. [simplified]

Let $\varpi : \mathbb{R}^3 \to \mathbb{R}_+$ be measurable, and $r \in [1, 2]$.

• Classical CKP inequality. For any $0 \leq f \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$, assuming that the norms below are finite, it holds

(40)
$$\left\| (f - M^{f}) \,\varpi \right\|_{L^{r}}^{2} \leq 2 \,\max \left(\left\| M^{f} \,\varpi^{2} \right\|_{L^{\frac{r}{2-r}}}, \left\| f \,\varpi^{2} \right\|_{L^{\frac{r}{2-r}}} \right) H_{0}[f|M^{f}].$$

where M^f is the Maxwellian associated to f.

• Fermi-Dirac CKP inequality. For any $\varepsilon > 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge 0$ and $\gamma > \frac{2}{5}$, assuming that the norms below are finite,

(41)
$$\left\| \left(f - \mathcal{M}_{\varepsilon}^{f} \right) \varpi \right\|_{L^{r}}^{2} \leq 2 \max \left(\left\| \mathcal{M}_{\varepsilon}^{f} \varpi^{2} \right\|_{L^{\frac{r}{2-r}}}, \left\| f \varpi^{2} \right\|_{L^{\frac{r}{2-r}}} \right) H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}],$$

where $\mathcal{M}^{f}_{\varepsilon}$ is the ε -Fermi-Dirac distribution associated to f.

In the above proposition, when r = 2, $L^{\frac{r}{2-r}}$ should be understood as L^{∞} .

2.4. **Convergence towards equilibrium.** A result like Corollary 3 constitutes the core of the proof of convergence of solutions to the Boltzmann-Fermi-Dirac equation towards equilibrium based on an entropy method. The proof is then complete provided that the solution satisfies a variety of bounds. The obtention of these bounds is an ongoing work that Bertrand Lods and I are currently conducting in the cutoff hard potentials case.

Proposition 5. Let $\varepsilon > 0$ and $f^{in} \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $0 \leq f^{in} \leq \varepsilon^{-1}$. We consider a collision kernel B satisfying the symmetry and micro-reversibility assumptions, thus having the form (4), as well as, for some $B_0 > 0$ and $\beta^+, \beta^- \geq 0$,

$$B(|v - v_*|, \cos \theta) \ge B_0 \min \left(|v - v_*|^{\beta_+}, |v - v_*|^{-\beta_-} \right).$$

We assume the existence of a solution $f \equiv f(t, v)$ to the spatially homogeneous Boltzmann-Fermi-Dirac equation (1) associated to the quantum parameter ε , the collision kernel B and the initial distribution f^{in} . Assume that there exists an explicit time $t_0 > 0$ depending only on ε , B and f^{in} such that

- $\exists \kappa_0 \in (0,1)$ depending only on ε , B and f^{in} such that $1 \varepsilon f(t,v) \ge \kappa_0$ for any $(t,v) \in [t_0, +\infty) \times \mathbb{R}^3$,
- $\exists K_0, A_0 > 0$ depending only on ε , B and f^{in} such that $f(t, v) \geq K_0 e^{-A_0|v|^2}$ for any $(t, v) \in [t_0, +\infty) \times \mathbb{R}^3$,

• $\forall k, l \in \mathbb{N}$, there exist $C_k > 0, \widetilde{C}_l > 0$ depending only on ε , B and f^{in} such that $\sup_{t \ge t_0} \|f(t, \cdot)\|_{L^1_k} \le C_k$ and $\sup_{t > t_0} \|f(t, \cdot)\|_{H^1} \le \widetilde{C}_l$.

Then for any $p \ge 1$, $l \ge 0$ and $\alpha > 0$ there exists an explicit constant $\mathbf{C}_{p,l,\alpha} > 0$ depending only on p, l, ε , B and f^{in} such that for any $t \ge t_0$,

(42)
$$\left\|f(t,\cdot) - \mathcal{M}_{\varepsilon}^{f^{in}}\right\|_{L_{l}^{p}} \leq \mathbf{C}_{p,l,\alpha} \left(1+t\right)^{-1/\alpha}.$$

Proof. The assumptions on the solution f allow to apply Corollary 3, more specifically Equation (39), yielding the fact that for any $t \ge t_0$ and $\alpha > 0$, there is an explicit constant $K_{\alpha}(f(t, \cdot))$ depending only on α , B_0 , an upper bound of ε and on $f(t, \cdot)$ only through ρ , T (its density and temperature, defined by (12)), and upper bounds on A_0 , $1/K_0$, $1/\kappa_0$, $||f(t, \cdot)||_{L^1_s}$ and $||f(t, \cdot)||_{H^k}$, where $s = s(\alpha, q_0, \beta_-, \beta_+)$ and $k = k(\alpha, s, \beta_-, \beta_+)$, such that

(43)
$$\mathcal{D}_{\varepsilon}(f(t,\cdot)) \ge K_{\alpha}(f(t,\cdot)) H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f(t,\cdot)}]^{1+\alpha}.$$

From our assumptions, we observe that $K_{\alpha}(f(t, \cdot))$ can be upper-bounded by a constant \hat{K}_{α} depending only on ε , B and f^{in} (in particular not on time t). Moreover, since f solves (1), we have for all $t \ge 0$ that $\mathcal{M}_{\varepsilon}^{f(t,\cdot)} = \mathcal{M}_{\varepsilon}^{f^{in}}$ and, at least formally, that

$$\frac{\mathrm{d}}{\mathrm{d}t}H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] = -\mathcal{D}_{\varepsilon}(f(t,\cdot)).$$

Then (43) implies

$$\frac{\mathrm{d}}{\mathrm{d}t}H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] \leq -\hat{K}_{\alpha}H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}]^{1+\alpha}.$$

Integrating this equation between t_0 and $t > t_0$ yields

$$H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] \leq \left(H_{\varepsilon}[f(t_0,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}]^{\alpha} + \alpha \hat{K}_{\alpha}(t-t_0)\right)^{-1/\alpha}$$

We then conclude that for any $t \ge t_0$, (even if $H_{\varepsilon}[f(t_0, \cdot) | \mathcal{M}_{\varepsilon}^{f^{in}}] = 0$, in which case $H_{\varepsilon}[f(t, \cdot) | \mathcal{M}_{\varepsilon}^{f^{in}}] = 0$)

(44)
$$H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] \leq \hat{C}_{\alpha} (1+t)^{-1/\alpha},$$

where $\hat{C}_{\alpha} > 0$ is explicit and depends only on α , ε , B and f^{in} . Finally, we recall the CKP inequalities provided in Proposition 4, Equation (41) in the Fermi-Dirac context, for weighted L^1 norms, implying that for any $k \ge 0$ we have for any $t \ge t_0$

$$\left\|f(t,\cdot) - \mathcal{M}_{\varepsilon}^{f^{in}}\right\|_{L_{k}^{1}}^{2} \leq 2 \max\left(\left\|\mathcal{M}_{\varepsilon}^{f^{in}}\right\|_{L_{2k}^{1}}, \left\|f(t,\cdot)\right\|_{L_{2k}^{1}}\right) H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}].$$

Since by assumption $\|f(t,\cdot)\|_{L^{1}_{2k}} \leq C_{2k}$, we conclude that there exists an explicit C'_{k} depending only on ε , *B* and f^{in} such that

$$\left\| f(t,\cdot) - \mathcal{M}_{\varepsilon}^{f^{in}} \right\|_{L^{1}_{k}}^{2} \leq C'_{k} H_{\varepsilon}[f(t,\cdot)|\mathcal{M}_{\varepsilon}^{f^{in}}] \leq C'_{k} \hat{C}_{\alpha} (1+t)^{-1/\alpha}.$$

Then (42) for the L_k^1 norm is finally obtained with $\mathbf{C}_{1,k,\alpha} = \sqrt{C'_k \hat{C}_{\alpha/2}}$ and L_l^p norms follow, as $\mathcal{M}_{\varepsilon}^{f^{in}} \in L^{\infty}(\mathbb{R}^3)$ and $f \in L^{\infty}(\mathbb{R}_+ \times \mathbb{R}^3)$.

3. Proof of Theorem 1 and Proposition 2

This section is devoted to the proof of Theorem 1 and Proposition 2. We make use of Proposition 1, which is later proven in Appendix B - see Proposition 15.

3.1. Key element of the proof. The main argument in the proofs of Theorem 1 and Proposition 2 lies in the following proposition.

Proposition 6. Let $0 \leq g \in L^1_2(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$ and define, for $\varepsilon \geq 0$,

(45)
$$R_g(\varepsilon) := H_{\varepsilon} \left[\varphi_{\varepsilon}^{-1}(g) \left| \mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)} \right] \right].$$

Then $\varepsilon \mapsto R_g(\varepsilon)$ is \mathcal{C}^1 on \mathbb{R}^*_+ , with, for any $\varepsilon > 0$,

(46)
$$R'_g(\varepsilon) = \int_{\mathbb{R}^3} \int_{M_\varepsilon(v)}^{g(v)} \frac{\partial_\varepsilon(\varphi_\varepsilon^{-1})(g(v)) - \partial_\varepsilon(\varphi_\varepsilon^{-1})(y)}{y} \, \mathrm{d}y \, \mathrm{d}v,$$

where we used the shorthand $M_{\varepsilon} = \varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)}).$

The striking point of (46) lies in the fact that it immediately proves that, for any $\varepsilon > 0$, the global monotonicity of $x \mapsto \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(x)$ translates into the same monotonicity of R_g at the point ε , as the integral in yin (46) is automatically nonnegative if $x \mapsto \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(x)$ is nondecreasing, or nonpositive if $x \mapsto \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(x)$ is nonincreasing. Moreover, this proposition generalises to any family of entropies and any conserved quantities (provided a suitable differentiability property for R_g), as all key arguments rely only the generic framework we present in Appendix B, and the specificity of the Fermi-Dirac case is used only for the study of the differentiability of R_g .

Proof. Using the relative entropy representation (24) in Proposition 1, we have, for any $\varepsilon > 0$,

$$R_g(\varepsilon) = \int_{\mathbb{R}^3} \int_{\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g(v))}(v)}^{\varphi_{\varepsilon}^{-1}(g(v))} (\varphi_{\varepsilon}^{-1}(g(v)) - x) \frac{\varphi_{\varepsilon}'(x)}{\varphi_{\varepsilon}(x)} \, \mathrm{d}x \, \mathrm{d}v$$

Changing variables $y = \varphi_{\varepsilon}(x)$, with then $dy = \varphi'_{\varepsilon}(x) dx$ and $x = \varphi_{\varepsilon}^{-1}(y)$, we obtain

$$R_g(\varepsilon) = \int_{\mathbb{R}^3} \int_{M_{\varepsilon}(v)}^{g(v)} \frac{\varphi_{\varepsilon}^{-1}(g(v)) - \varphi_{\varepsilon}^{-1}(y)}{y} \, \mathrm{d}y \, \mathrm{d}v.$$

where we denoted $M_{\varepsilon} = \varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)})$ for convenience of notations. We fix $v \in \mathbb{R}^3$ and set, for $(\varepsilon, z) \in (\mathbb{R}^*_+)^2$,

(47)
$$K(\varepsilon, z) := \int_{z}^{g(v)} \frac{\varphi_{\varepsilon}^{-1}(g(v)) - \varphi_{\varepsilon}^{-1}(y)}{y} \, \mathrm{d}y$$

Note that if g(v) = 0 then $K(\varepsilon, z) = \varepsilon^{-1} \log(1 + \varepsilon z)$. The application K is clearly \mathcal{C}^1 on $(\mathbb{R}^*_+)^2$, and straightforward computations can provide that differentiation in ε under the integral in y is valid. We find that

$$\partial_{\varepsilon} K(\varepsilon, z) = \int_{z}^{g(v)} \frac{\partial_{\varepsilon}(\varphi_{\varepsilon}^{-1}(g(v))) - \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1}(y))}{y} \, \mathrm{d}y,$$
$$\partial_{z} K(\varepsilon, z) = -\frac{\varphi_{\varepsilon}^{-1}(g(v)) - \varphi_{\varepsilon}^{-1}(z)}{z}.$$

As a result of Lemma 21 in Appendix C, the application $\varepsilon \mapsto M_{\varepsilon}(v)$ is \mathcal{C}^1 on \mathbb{R}^*_+ (as a composition of \mathcal{C}^1 applications), hence so is $\varepsilon \mapsto K(\varepsilon, M_{\varepsilon}(v))$, and we have, for any $\varepsilon > 0$ and $v \in \mathbb{R}^3$,

(48)
$$\frac{\mathrm{d}}{\mathrm{d}\varepsilon}K(\varepsilon, M_{\varepsilon}(v)) = \int_{M_{\varepsilon}(v)}^{g(v)} \frac{\partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(g(v)) - \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(y)}{y} \,\mathrm{d}y - (\partial_{\varepsilon}M_{\varepsilon}(v)) \frac{\varphi_{\varepsilon}^{-1}(g(v)) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon}(v))}{M_{\varepsilon}(v)}$$

Now note that

(49)
$$R_g(\varepsilon) = \int_{\mathbb{R}^3} K(\varepsilon, M_{\varepsilon}(v)) \,\mathrm{d}v$$

Then R_g is indeed \mathcal{C}^1 on \mathbb{R}^*_+ , and differentiation under the integral is valid, if we can prove that for any $\varepsilon > 0$, there is a $\delta \in (0, \varepsilon)$ such that

(50)
$$\int_{\mathbb{R}^3} \sup_{\varepsilon_* \in (\varepsilon - \delta, \varepsilon + \delta)} \left| \frac{\mathrm{d}}{\mathrm{d}\varepsilon_*} K(\varepsilon_*, M_{\varepsilon_*}(v)) \right| \mathrm{d}v < \infty.$$

For the sake of clarity, we continue our computations assuming (50) for now, and show (50) after we have obtained our claimed result (46).

Differentiating (49) in ε under the integral in v, using (48), yields for any $\varepsilon > 0$ (omitting to write the dependence in v for clarity)

(51)
$$R'_{g}(\varepsilon) = \int_{\mathbb{R}^{3}} \int_{M_{\varepsilon}}^{g} \frac{\partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(g) - \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(y)}{y} \, \mathrm{d}y \, \mathrm{d}v - \int_{\mathbb{R}^{3}} (\partial_{\varepsilon} M_{\varepsilon}) \times \frac{\varphi_{\varepsilon}^{-1}(g) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon})}{M_{\varepsilon}} \, \mathrm{d}v.$$

Now remark that

(52)
$$\frac{\partial_{\varepsilon} M_{\varepsilon}}{M_{\varepsilon}} = \partial_{\varepsilon} \left(\log \circ \varphi_{\varepsilon} \left(\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)} \right) \right)$$

Since $\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)}$ is a Fermi-Dirac statistics, we have, denoting its coefficients by a_{ε} , u_{ε} and b_{ε} (as in (19)), that for any $v \in \mathbb{R}^3$,

(53)
$$\log \circ \varphi_{\varepsilon} \left(\mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)} \right) (v) = \alpha_{\varepsilon} + \beta_{\varepsilon} \cdot v + b_{\varepsilon} |v|^{2},$$

where $\alpha_{\varepsilon} = a_{\varepsilon} + b_{\varepsilon}|u_{\varepsilon}|^2$ and $\beta_{\varepsilon} = -2b_{\varepsilon}u_{\varepsilon}$. Lemma 21 in Appendix C ensures that the application $\varepsilon \mapsto (a_{\varepsilon}, u_{\varepsilon}, b_{\varepsilon})$ is \mathcal{C}^1 on \mathbb{R}^*_+ , hence so is the above expression pointwise in v, and we conclude that $\frac{\partial_{\varepsilon}M_{\varepsilon}}{M_{\varepsilon}}$ is a linear combination of the functions $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$. By construction, $\varphi_{\varepsilon}^{-1}(g)$ and $\varphi_{\varepsilon}^{-1}(M_{\varepsilon}) = \mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)}$ share the same normalization in $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$, hence

$$\int_{\mathbb{R}^3} \left(\varphi_{\varepsilon}^{-1}(g) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon}) \right) \frac{\partial_{\varepsilon} M_{\varepsilon}}{M_{\varepsilon}} \, \mathrm{d}v = 0.$$

and (51) becomes the announced (46).

We now show (50), ensuring the validity of the differentiation in ε under the integral in Equation (49), by dominated convergence. The previous computations imply that for any $\varepsilon > 0$,

$$\left|\frac{\mathrm{d}}{\mathrm{d}\varepsilon} K(\varepsilon, M_{\varepsilon}(v))\right| \leq \left|\int_{M_{\varepsilon}}^{g} \frac{\partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(g) - \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(y)}{y} \,\mathrm{d}y\right| + \left|\frac{\partial_{\varepsilon} M_{\varepsilon}}{M_{\varepsilon}}(\varphi_{\varepsilon}^{-1}(g) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon}))\right|.$$

On the one hand, using (52)–(53), we get, for any $v \in \mathbb{R}^3$,

$$\left| \frac{\partial_{\varepsilon} M_{\varepsilon}}{M_{\varepsilon}} (\varphi_{\varepsilon}^{-1}(g) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon})) \right| \leq \left(|\partial_{\varepsilon} \alpha_{\varepsilon}| + |\partial_{\varepsilon} \beta_{\varepsilon}| + |\partial_{\varepsilon} b_{\varepsilon}| \right) (1 + |v|^{2}) |\varphi_{\varepsilon}^{-1}(g(v)) - \varphi_{\varepsilon}^{-1}(M_{\varepsilon}(v))|$$
$$\leq \left(|\partial_{\varepsilon} \alpha_{\varepsilon}| + |\partial_{\varepsilon} \beta_{\varepsilon}| + |\partial_{\varepsilon} b_{\varepsilon}| \right) (1 + |v|^{2}) \{g(v) + M_{\varepsilon}(v)\},$$

where for the last inequality, we used the inequalities $|x - y| \le x + y$ and $\varphi_{\varepsilon}^{-1}(x) \le x$ for $x, y \ge 0$. On the other hand, remarking that

(54)
$$\forall x \ge 0, \qquad \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(x) = \partial_{\varepsilon}\left(\frac{x}{1+\varepsilon x}\right) = -\left(\frac{x}{1+\varepsilon x}\right)^2 = -\varphi_{\varepsilon}^{-1}(x)^2,$$

we obtain, as $x \mapsto \varphi_{\varepsilon}^{-1}(x)^2$ is increasing,

$$\left|\int_{M_{\varepsilon}}^{g} \frac{\partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(g) - \partial_{\varepsilon}(\varphi_{\varepsilon}^{-1})(y)}{y} \,\mathrm{d}y\right| = \int_{M_{\varepsilon}}^{g} \frac{\varphi_{\varepsilon}^{-1}(g)^{2} - \varphi_{\varepsilon}^{-1}(y)^{2}}{y} \,\mathrm{d}y$$

In the last equation, as well as in the four following ones, we omit writing the dependence of g and M_{ε} in v for clarity. From the identity $|x^2 - y^2| = (x + y)|x - y|$, for $x, y \ge 0$, the fact that $\|\varphi_{\varepsilon}^{-1}\|_{\infty} = \varepsilon^{-1}$ and that $\varphi_{\varepsilon}^{-1}$ is increasing, we get

$$\int_{M_{\varepsilon}}^{g} \frac{\varphi_{\varepsilon}^{-1}(g)^2 - \varphi_{\varepsilon}^{-1}(y)^2}{y} \, \mathrm{d}y \le 2\varepsilon^{-1} \int_{M_{\varepsilon}}^{g} \frac{\varphi_{\varepsilon}^{-1}(g) - \varphi_{\varepsilon}^{-1}(y)}{y} \, \mathrm{d}y,$$

which, using the bound $|\varphi_{\varepsilon}^{-1}(x) - \varphi_{\varepsilon}^{-1}(y)| \le |x - y|$ for $x, y \ge 0$ and $\varepsilon > 0$, is less than

$$2\varepsilon^{-1} \int_{M_{\varepsilon}}^{g} \frac{g-y}{y} \,\mathrm{d}y$$

By direct integration, the latter equals

$$2\varepsilon^{-1} \left(g \log g - g \log M_{\varepsilon} - g + M_{\varepsilon} \right),$$

which, recalling that $M_{\varepsilon}(v) = \exp\left(a_{\varepsilon} + |v - u_{\varepsilon}|^2\right)$ and $g \ge 0$, is itself less than

$$2\varepsilon^{-1} \left(g \log g - a_{\varepsilon} g + M_{\varepsilon}\right)$$
 .

All in all, as $-a_{\varepsilon}g \leq |a_{\varepsilon}|g$, we have for any $\varepsilon > 0$ and $v \in \mathbb{R}^3$ that

(55)
$$\left| \frac{\mathrm{d}}{\mathrm{d}\varepsilon} K(\varepsilon, M_{\varepsilon}(v)) \right| \leq 2\varepsilon^{-1} \left\{ g \log g(v) + |a_{\varepsilon}|g(v) + M_{\varepsilon}(v) \right\} + \left(|\partial_{\varepsilon}\alpha_{\varepsilon}| + |\partial_{\varepsilon}\beta_{\varepsilon}| + |\partial_{\varepsilon}b_{\varepsilon}| \right) \left(1 + |v|^{2} \right) \left\{ g(v) + M_{\varepsilon}(v) \right\}.$$

Let us now fix $\varepsilon > 0$ and an arbitrary $\delta \in (0, \varepsilon)$. Since the application $\varepsilon \mapsto (\alpha_{\varepsilon}, \beta_{\varepsilon}, b_{\varepsilon})$ is \mathcal{C}^1 on \mathbb{R}^*_+ , there exists $c_0 > 0$ such that

$$\sup_{\varepsilon_* \in (\varepsilon - \delta, \varepsilon + \delta)} \left(|\partial_{\varepsilon_*} \alpha_{\varepsilon_*}| + |\partial_{\varepsilon_*} \beta_{\varepsilon_*}| + |\partial_{\varepsilon_*} b_{\varepsilon_*}| \right) \le c_0.$$

Moreover, as $\varepsilon \mapsto a_{\varepsilon}$ is continuous on \mathbb{R}_+ , there is a $c_1 > 0$ such that

$$\sup_{\varepsilon_* \in (\varepsilon - \delta, \varepsilon + \delta)} |a_{\varepsilon_*}| \le c_1$$

Finally, from Lemma 22 in Appendix C (with $\overline{\varepsilon} = \varepsilon + \delta$ in the lemma), there exist $c_2 > 0$ and $\eta > 0$ such that for any $v \in \mathbb{R}^3$,

$$\sup_{\varepsilon_* \in [0,\varepsilon+\delta]} M_{\varepsilon_*}(v) \le c_2 \, e^{-\eta |v|^2}.$$

Then (55) implies that for any $v \in \mathbb{R}^3$,

$$\sup_{\varepsilon_* \in (\varepsilon - \delta, \varepsilon + \delta)} \left| \frac{\mathrm{d}}{\mathrm{d}\varepsilon_*} K(\varepsilon_*, M_{\varepsilon_*}(v)) \right| \leq 2(\varepsilon - \delta)^{-1} \left(g \log g(v) + c_1 g(v) + c_2 e^{-\eta |v|^2} \right) + c_0 (1 + |v|^2) \left\{ g(v) + c_2 e^{-\eta |v|^2} \right\}.$$

Since $g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$, the right-hand side of the above equation is integrable, and we conclude, by dominated convergence, to the differentiability of R_g and the validity of the differentiation in ε under the integral, ending the proof.

3.2. Continuity at zero. The second and last argument required in the proofs of Theorem 1 and Proposition 2 is the continuity of $\varepsilon \mapsto R_q(\varepsilon)$ at the point $\varepsilon = 0$, which is given in the following lemma.

Lemma 7. Let $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$. Then the function $\varepsilon \in \mathbb{R}_+ \mapsto R_g(\varepsilon)$ defined in (45) is continuous at the point $\varepsilon = 0$.

Proof. We write $R_g(\varepsilon)$ as

(56)
$$R_g(\varepsilon) = H_{\varepsilon}\left(\frac{g}{1+\varepsilon g}\right) - H_{\varepsilon}\left(\mathcal{M}_{\varepsilon}^{\frac{g}{1+\varepsilon g}}\right).$$

We show that both terms, in the right-hand-side, are continuous at the point $\varepsilon = 0$. Firstly, by definition of the Fermi-Dirac entropy, it holds for any $\varepsilon > 0$ that

$$\begin{aligned} H_{\varepsilon}\left(\frac{g}{1+\varepsilon g}\right) &= \int_{\mathbb{R}^3} \left\{\frac{g}{1+\varepsilon g} \log\left(\frac{g}{1+\varepsilon g}\right) + \varepsilon^{-1} \left(1-\varepsilon \frac{g}{1+\varepsilon g}\right) \log\left(1-\varepsilon \frac{g}{1+\varepsilon g}\right)\right\} \mathrm{d}v \\ &= \int_{\mathbb{R}^3} \left\{\frac{g}{1+\varepsilon g} \log g - \varepsilon^{-1} \log(1+\varepsilon g)\right\} \mathrm{d}v. \end{aligned}$$

As $g \ge 0$, we have for any $\varepsilon > 0$ that

$$\left|\frac{g}{1+\varepsilon g}\log g\right| \le g|\log g|$$
 and $\left|-\varepsilon^{-1}\log(1+\varepsilon g)\right| \le g,$

Since by hypothesis $0 \le g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$, the function $g |\log g| + g$ is integrable, and, by dominated convergence, we then conclude to the limit

$$H_{\varepsilon}\left(\frac{g}{1+\varepsilon g}\right) \xrightarrow{\varepsilon \to 0} \int_{\mathbb{R}^3} \left\{g \log g - g\right\} \mathrm{d}v = H_0(g).$$

Secondly, going through the same computations as for the first term, we have, for any $\varepsilon > 0$,

$$H_{\varepsilon}\left(\mathcal{M}_{\varepsilon}^{\frac{g}{1+\varepsilon g}}\right) = \int_{\mathbb{R}^{3}} \left\{ \frac{M_{\varepsilon}}{1+\varepsilon M_{\varepsilon}} \log M_{\varepsilon} - \varepsilon^{-1} \log(1+\varepsilon M_{\varepsilon}) \right\} \mathrm{d}v,$$

where we denoted $M_{\varepsilon} = \varphi_{\varepsilon} \left(\mathcal{M}_{\varepsilon}^{\frac{g}{1+\varepsilon g}} \right)$. Lemma 22 in Appendix C (with $\overline{\varepsilon} = 1$ in the lemma) provides the existence of $C, \eta > 0$ such that for all $0 \le \varepsilon \le 1$,

$$|\log M_{\varepsilon}(v)| \le C(1+|v|^2)$$
 and $M_{\varepsilon}(v) \le C e^{-\eta|v|^2}$

Hence, for any $0 \leq \varepsilon \leq 1$ and $v \in \mathbb{R}^3$,

$$\left|\frac{M_{\varepsilon}(v)}{1+\varepsilon M_{\varepsilon}(v)}\log M_{\varepsilon}(v)\right| \leq C^{2} e^{-\eta|v|^{2}}(1+|v|^{2}), \text{ and } \left|\varepsilon^{-1}\log(1+\varepsilon M_{\varepsilon}(v))\right| \leq M_{\varepsilon}(v) \leq C e^{-\eta|v|^{2}}.$$

By dominated convergence, it thus comes that

$$\lim_{\varepsilon \to 0} H_{\varepsilon} \left(\mathcal{M}_{\varepsilon}^{\frac{g}{1+\varepsilon g}} \right) = \int_{\mathbb{R}^3} \lim_{\varepsilon \to 0} \left(\frac{M_{\varepsilon}}{1+\varepsilon M_{\varepsilon}} \log M_{\varepsilon} - \varepsilon^{-1} \log(1+\varepsilon M_{\varepsilon}) \right) \mathrm{d}v$$

As Lemma 20 ensures that $\lim_{\varepsilon \to 0} M_{\varepsilon}(v) = \mathcal{M}_0^g(v)$ for all $v \in \mathbb{R}^3$, the above limit becomes

$$\int_{\mathbb{R}^3} \mathcal{M}_0^g \log \mathcal{M}_0^g - \mathcal{M}_0^g \, \mathrm{d} v,$$

that is $H_0(\mathcal{M}_0^g)$, ending the proof.

We can finally conclude to the announced inequalities of Theorem 1 and Proposition 2.

3.3. Final proof of Theorem 1. Let $\varepsilon > 0$ and $0 \le f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ such that

$$g := \frac{f}{1 - \varepsilon f} \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}.$$

Proposition 6 ensures that the application R_g defined by (45) is \mathcal{C}^1 on \mathbb{R}^*_+ , and combining (46) with (54) yields, for any $\varepsilon_* > 0$,

$$R'_g(\varepsilon_*) = -\int_{\mathbb{R}^3} \int_{M_{\varepsilon_*}}^g \frac{\varphi_{\varepsilon_*}^{-1}(g)^2 - \varphi_{\varepsilon_*}^{-1}(y)^2}{y} \,\mathrm{d}y \,\mathrm{d}v.$$

As, for any $\varepsilon_* > 0$, the application $x \mapsto \varphi_{\varepsilon_*}^{-1}(x)^2$ is increasing, the integral in y is nonnegative, hence so is the integral in y and v, therefore

$$\forall \varepsilon_* > 0, \qquad \qquad R'_q(\varepsilon_*) \le 0.$$

Combining this result with Lemma 7, which states that $\varepsilon_* \mapsto R_g(\varepsilon_*)$ is continuous at the point $\varepsilon_* = 0$, R_g is then nonincreasing on \mathbb{R}_+ , which implies in particular that

$$R_g(0) \ge R_g(\varepsilon),$$

that is

$$H_0[g|M^g] \ge H_{\varepsilon}\left[\varphi_{\varepsilon}^{-1}(g) \left| \mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)} \right. \right],$$

or the announced inequality (32),

$$H_0\left[\frac{f}{1-\varepsilon f}\left|M^{\frac{f}{1-\varepsilon f}}\right|\right] \ge H_{\varepsilon}\left[f\left|\mathcal{M}_{\varepsilon}^{f}\right|\right].$$

3.4. Final proof of Proposition 2. Let us fix $\varepsilon > 0$, $\kappa_0 \in (0,1)$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$. We let

$$g := \frac{f}{1 - \varepsilon f}.$$

Then $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L^{\infty}(\mathbb{R}^3) \setminus \{0\}$, which is in particular included in $L \log L(\mathbb{R}^3)$. We apply the results of Proposition 6 and Lemma 7 with the function g, ensuring that the application R_g , defined by (45), is continuous on \mathbb{R}_+ and \mathcal{C}^1 on \mathbb{R}^*_+ with, combining (46) with (54),

$$\forall \, \varepsilon_* > 0, \qquad R'_g(\varepsilon_*) = -\int_{\mathbb{R}^3} \int_{M_{\varepsilon_*}}^g \frac{\varphi_{\varepsilon_*}^{-1}(g)^2 - \varphi_{\varepsilon_*}^{-1}(y)^2}{y} \, \mathrm{d}y \, \mathrm{d}v$$

For $\varepsilon_* > 0, v \in \mathbb{R}^3$ and any y between $M_{\varepsilon_*}(v)$ and g(v), one has (omitting the dependencies in v for clarity)

$$|\varphi_{\varepsilon_*}^{-1}(g)^2 - \varphi_{\varepsilon_*}^{-1}(y)^2| = (\varphi_{\varepsilon_*}^{-1}(g) + \varphi_{\varepsilon_*}^{-1}(y))|\varphi_{\varepsilon_*}^{-1}(g) - \varphi_{\varepsilon_*}^{-1}(y)| \le c_g(\varepsilon_*) |\varphi_{\varepsilon_*}^{-1}(g) - \varphi_{\varepsilon_*}^{-1}(y)|,$$

we denoted

$$c_g(\varepsilon_*) := \|\varphi_{\varepsilon_*}^{-1}(g)\|_{\infty} + \max\left(\|\varphi_{\varepsilon_*}^{-1}(g)\|_{\infty}, \|\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)}\|_{\infty}\right).$$

As both $z \mapsto \varphi_{\varepsilon_*}^{-1}(z)^2$ and $z \mapsto \varphi_{\varepsilon_*}^{-1}(z)$ are increasing, we thus have

$$R'_g(\varepsilon_*) \ge -c_g(\varepsilon_*) \int_{\mathbb{R}^3} \int_{M_{\varepsilon_*}}^g \frac{\varphi_{\varepsilon_*}^{-1}(g) - \varphi_{\varepsilon_*}^{-1}(y)}{y} \,\mathrm{d}y \,\mathrm{d}v = -c_g(\varepsilon_*) \,R_g(\varepsilon_*).$$

Noticing that

where w

$$c_g(\varepsilon_*) \le \|g\|_{\infty} + \max\left(\|g\|_{\infty} + \|\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)}\|_{\infty}\right)$$

we conclude to the relationship, for any $\varepsilon_* > 0$,

(57)
$$R'_{g}(\varepsilon_{*}) \geq -\left(\|g\|_{\infty} + \max\left(\|g\|_{\infty} + \|\mathcal{M}_{\varepsilon_{*}}^{\varphi_{\varepsilon_{*}}^{-1}(g)}\|_{\infty}\right)\right) R_{g}(\varepsilon_{*})$$

By hypothesis, it holds that $1 - \varepsilon f \ge \kappa_0$, hence

(58)
$$||g||_{\infty} \le \varepsilon^{-1} (\kappa_0^{-1} - 1),$$

and we have for any $\varepsilon_* > 0$ that

(59)
$$\varepsilon_* \varphi_{\varepsilon_*}^{-1}(g) = \frac{\varepsilon_* g}{1 + \varepsilon_* g} \le \left(1 + \frac{1}{\varepsilon_* \|g\|_{\infty}}\right)^{-1} \le \left(1 + \frac{\varepsilon}{\varepsilon_*} (\kappa_0^{-1} - 1)^{-1}\right)^{-1}$$

Let us use the notation

$$\gamma_{\varepsilon_*} = \frac{T_*}{T_F(\rho_*, \varepsilon_*)}$$

where ρ_* and T_* are respectively the density and temperature associated to the distribution $\varphi_{\varepsilon_*}^{-1}(g)$ defined as in (12), and $T_F(\rho_*, \varepsilon_*) = \frac{1}{2} \left(\frac{3\rho_*\varepsilon_*}{4\pi}\right)^{2/3}$ is the corresponding ε_* -Fermi temperature. Using Equation (117) in Proposition 18 in Appendix C and the fact that $\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)} \leq \varphi_{\varepsilon_*} \left(\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)}\right)$, we have, whenever $\gamma_{\varepsilon_*} \geq \gamma^{\dagger} = \left(\frac{4}{\pi}\right)^{\frac{1}{3}} \left(\frac{5}{3}\right)^{\frac{5}{3}}$,

(60)
$$\varepsilon_* \| \mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)} \|_{\infty} \leq \frac{2}{3} \left(\frac{\gamma_{\varepsilon_*}}{\gamma^{\dagger}} \right)^{-3/2}$$

We now notice that [22, Proposition 4] actually implies

$$\gamma_{\varepsilon_*} \ge \frac{2}{5} \left(\varepsilon_* \| \varphi_{\varepsilon_*}^{-1}(g) \|_{\infty} \right)^{-2/3}$$

which, combined with (59), yields

(61)
$$\gamma_{\varepsilon_*} \ge \frac{2}{5} \left(1 + \frac{\varepsilon}{\varepsilon_*} \left(\kappa_0^{-1} - 1 \right)^{-1} \right)^{2/3}.$$

We define

(62)
$$\varepsilon^{\dagger} := \left(\left(\frac{5}{2} \gamma^{\dagger} \right)^{\frac{3}{2}} - 1 \right)^{-1} (\kappa_0^{-1} - 1)^{-1} \varepsilon.$$

Then, for any $\varepsilon_* \in (0, \varepsilon^{\dagger}]$, combining (61) with (62) yields

$$\gamma_{\varepsilon_*} \ge \frac{2}{5} \left(1 + \frac{\varepsilon}{\varepsilon^{\dagger}} \left(\kappa_0^{-1} - 1 \right)^{-1} \right)^{2/3} = \gamma^{\dagger}.$$

Applying (60), along with (61) and the fact that

$$\frac{2}{3}\left(\frac{5}{2}\gamma^{\dagger}\right)^{\frac{3}{2}} = \frac{2}{3}\left(\frac{5}{2}\right)^{\frac{3}{2}}\left(\frac{4}{\pi}\right)^{\frac{1}{2}}\left(\frac{5}{3}\right)^{\frac{5}{2}} \le 11,$$

gives, for any $\varepsilon_* \in (0, \varepsilon^{\dagger}]$,

$$\|\mathcal{M}_{\varepsilon_{*}}^{\varphi_{\varepsilon_{*}}^{-1}(g)}\|_{\infty} \leq \frac{11}{\varepsilon_{*} + \varepsilon(\kappa_{0}^{-1} - 1)^{-1}} \leq 11 \,\varepsilon^{-1} \,(\kappa_{0}^{-1} - 1).$$

Therefore, recalling (58), we have for any $\varepsilon_* \in (0, \varepsilon^{\dagger}]$ that

$$\|g\|_{\infty} + \max\left(\|g\|_{\infty}, \|\mathcal{M}_{\varepsilon_{*}}^{\varphi_{\varepsilon_{*}}^{-1}(g)}\|_{\infty}\right) \leq \varepsilon^{-1} \left(\kappa_{0}^{-1} - 1\right) + 11\varepsilon^{-1} \left(\kappa_{0}^{-1} - 1\right) = 12\varepsilon^{-1} \left(\kappa_{0}^{-1} - 1\right),$$

implying, with the relationship (57), that for any $\varepsilon_* \in (0, \varepsilon^{\dagger}]$,

(63)
$$R'_g(\varepsilon_*) \ge -12\,\varepsilon^{-1}\,(\kappa_0^{-1}-1)\,R_g(\varepsilon_*)$$

Applying a Grönwall argument over (63) and using the continuity of $\varepsilon_* \mapsto R_g(\varepsilon_*)$ at the point $\varepsilon_* = 0$ provided by Lemma 7, we obtain

(64)
$$\forall \varepsilon_* \in (0, \varepsilon^{\dagger}], \qquad R_g(0) \le \exp\left(12\frac{\varepsilon_*}{\varepsilon}\left(\kappa_0^{-1}-1\right)\right) R_g(\varepsilon_*).$$

On the one hand, if $\varepsilon \leq \varepsilon^{\dagger}$, we take $\varepsilon_* = \varepsilon$ in (64) to get

(65)
$$R_g(0) \le \exp\left(12\left(\kappa_0^{-1} - 1\right)\right) R_g(\varepsilon).$$

On the other hand, if $\varepsilon \geq \varepsilon^{\dagger}$, we apply (64) with $\varepsilon_* = \varepsilon^{\dagger}$ to get

(66)
$$R_g(0) \le \exp\left(12\frac{\varepsilon^{\dagger}}{\varepsilon}(\kappa_0^{-1}-1)\right) R_g(\varepsilon^{\dagger}).$$

Since it holds for any $\varepsilon_* > 0$ that $\|\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)}\|_{\infty} \leq \varepsilon_*^{-1}$, we have in particular that, for any $\varepsilon_* \in [\varepsilon^{\dagger}, \varepsilon]$,

$$\|\mathcal{M}_{\varepsilon_*}^{\varphi_{\varepsilon_*}^{-1}(g)}\|_{\infty} \leq (\varepsilon^{\dagger})^{-1},$$

which, combined with (57) and (58), leads to

(67)
$$\forall \varepsilon_* \in [\varepsilon^{\dagger}, \varepsilon], \qquad R'_g(\varepsilon_*) \ge -\varepsilon^{-1} \left((\kappa_0^{-1} - 1) + \max\left(\kappa_0^{-1} - 1, \frac{\varepsilon}{\varepsilon^{\dagger}} \right) \right) R_g(\varepsilon_*).$$

By a Grönwall argument over (67), we obtain

(68)
$$R_g(\varepsilon^{\dagger}) \le \exp\left(\frac{\varepsilon - \varepsilon^{\dagger}}{\varepsilon} \left(\kappa_0^{-1} - 1 + \max\left(\kappa_0^{-1} - 1, \frac{\varepsilon}{\varepsilon^{\dagger}}\right)\right)\right) R_g(\varepsilon).$$

Recalling that ε^{\dagger} is defined by (62), we have

$$\frac{\varepsilon}{\varepsilon^{\dagger}} = \left(\left(\frac{5}{2} \gamma^{\dagger}\right)^{\frac{3}{2}} - 1 \right) (\kappa_0^{-1} - 1).$$

16

As $\gamma^{\dagger} = \left(\frac{4}{\pi}\right)^{\frac{1}{3}} \left(\frac{5}{3}\right)^{\frac{5}{3}}$, we have $\left(\frac{5}{2}\gamma^{\dagger}\right)^{\frac{3}{2}} - 1 \le 15$, therefore

(69)
$$\left(\kappa_0^{-1} - 1 + \max\left(\kappa_0^{-1} - 1, \frac{\varepsilon}{\varepsilon^{\dagger}}\right)\right) \le 16\left(\kappa_0^{-1} - 1\right)$$

Combining (68) with (69), we get

(70)
$$R_g(\varepsilon^{\dagger}) \le \exp\left(\frac{\varepsilon - \varepsilon^{\dagger}}{\varepsilon} \operatorname{16}\left(\kappa_0^{-1} - 1\right)\right) R_g(\varepsilon)$$

Finally, combining (66) with (70), we obtain, in this case where we assumed $\varepsilon \geq \varepsilon^{\dagger}$,

$$R_g(0) \le \exp\left(\left\{12\frac{\varepsilon^{\dagger}}{\varepsilon} + 16\frac{\varepsilon - \varepsilon^{\dagger}}{\varepsilon}\right\} (\kappa_0^{-1} - 1)\right) R_g(\varepsilon) \le \exp\left(16\left(\kappa_0^{-1} - 1\right)\right) R_g(\varepsilon).$$

We thus conclude, with (65), that in any case, we have

$$R_g(0) \le \exp\left(16\left(\kappa_0^{-1} - 1\right)\right) R_g(\varepsilon),$$

yielding the required result as $R_g(0) = H_0\left[\frac{f}{1-\varepsilon f} \left| M^{\frac{f}{1-\varepsilon f}} \right] \text{ and } R_g(\varepsilon) = H_{\varepsilon}\left[f \left| \mathcal{M}_{\varepsilon}^f \right]\right]$.

4. Proof of Corollary 3

This section is devoted to the proof of Corollary 3. Throughout this section, we use the notation

(71)
$$\begin{pmatrix} \rho_{\varepsilon} \\ \rho_{\varepsilon} u_{\varepsilon} \\ 3\rho_{\varepsilon} T_{\varepsilon} + \rho_{\varepsilon} |u_{\varepsilon}|^2 \end{pmatrix} = \int_{\mathbb{R}^3} \varphi_{\varepsilon}(f)(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv, \qquad \begin{pmatrix} \rho \\ \rho u \\ 3\rho T + \rho |u|^2 \end{pmatrix} = \int_{\mathbb{R}^3} f(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv.$$
In the following, we use the bound

In the following, we use the bound

(72)
$$1 - \varepsilon f \ge \kappa_0 \implies T_{\varepsilon} \ge \kappa_0 T.$$

This comes by first remarking that, if $1 - \varepsilon f \ge \kappa_0$,

$$3\rho_{\varepsilon}T_{\varepsilon} = \int_{\mathbb{R}^3} \varphi_{\varepsilon}(f)(v)|v - u_{\varepsilon}|^2 \,\mathrm{d}v \ge \int_{\mathbb{R}^3} f(v)|v - u_{\varepsilon}|^2 \,\mathrm{d}v = \int_{\mathbb{R}^3} f(v)|v - u - (u_{\varepsilon} - u)|^2 \,\mathrm{d}v = 3\rho T + \rho|u_{\varepsilon} - u|^2,$$
implying $\rho_{\varepsilon}T_{\varepsilon} \ge \rho T$, and the fact that

(73)
$$\rho_{\varepsilon} = \int_{\mathbb{R}^3} \varphi_{\varepsilon}(f)(v) \, \mathrm{d}v \le \kappa_0^{-1} \int_{\mathbb{R}^3} f(v) \, \mathrm{d}v = \kappa_0^{-1} \, \rho.$$

4.1. Super quadratic kernels.

Proof. Let
$$\kappa_0 \in (0,1)$$
, $\varepsilon > 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$. We have from (31) that
(74) $\mathcal{D}_{\varepsilon}(f) \ge \kappa_0^4 \mathcal{D}_0(\varphi_{\varepsilon}(f)).$

We apply [30, Theorem 2.1] to the function $\varphi_{\varepsilon}(f)$, adapted (since it is written for normalized functions) with the change of variable $w = \sqrt{T_{\varepsilon}} v + u_{\varepsilon}$, and obtain

$$\mathcal{D}_{0}(\varphi_{\varepsilon}(f)) \geq \min(1, T_{\varepsilon}) \frac{|\mathbb{S}^{2}|}{28} \left(3\rho_{\varepsilon}T_{\varepsilon} - \rho_{\varepsilon}T^{*}(\varphi_{\varepsilon}(f)) \right) H_{0} \left[\varphi_{\varepsilon}(f) \Big| M^{\varphi_{\varepsilon}(f)} \right],$$

$$\sup_{\varepsilon \in \mathcal{L}^{2}} \int_{\mathbb{S}^{2}} g(v) \left(v \cdot e \right)^{2} dv$$

where $T^*(g)$ stands for the maximal directional temperature of g, defined by $\frac{\sup_{e \in \mathbb{S}^2} \int_{\mathbb{R}^3} g(v) \, \mathrm{d} v}{\int_{\mathbb{R}^3} g(v) \, \mathrm{d} v}$. Since in particular,

$$3\rho_{\varepsilon}T_{\varepsilon} - \rho_{\varepsilon}T^*(\varphi_{\varepsilon}(f)) \ge 2\rho_{\varepsilon}T_*(\varphi_{\varepsilon}(f)),$$

where $T_*(g)$ stands for the minimal directional temperature of g, defined in (13), we obtain

(75)
$$\mathcal{D}_0(\varphi_{\varepsilon}(f)) \ge \min(1, T_{\varepsilon}) \frac{2\pi}{7} \rho_{\varepsilon} T_*(\varphi_{\varepsilon}(f)) H_0\left[\varphi_{\varepsilon}(f) \middle| M^{\varphi_{\varepsilon}(f)}\right]$$

Applying Theorem 1's lower-bound inequality (32), we obtain

(76)
$$H_0\left[\varphi_{\varepsilon}(f)\middle| M^{\varphi_{\varepsilon}(f)}\right] \ge H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^f]$$

Combining (74)–(76) yields

$$\mathcal{D}_{\varepsilon}(f) \geq \frac{2\pi}{7} \kappa_0^4 \min(1, T_{\varepsilon}) \rho_{\varepsilon} T_*(\varphi_{\varepsilon}(f)) H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^f].$$

The required result is obtained after remarking that, from (72), $T_{\varepsilon} \geq \kappa_0 T$, and

$$\rho_{\varepsilon} T_*(\varphi_{\varepsilon}(f)) = \inf_{e \in \mathbb{S}^2} \int_{\mathbb{R}^3} \varphi_{\varepsilon}(f)(v) \, (v \cdot e)^2 \, \mathrm{d}v \ge \inf_{e \in \mathbb{S}^2} \int_{\mathbb{R}^3} f(v) \, (v \cdot e)^2 \, \mathrm{d}v = \rho \, T_*(f).$$

4.2. General kernels.

Proof. Let $\kappa_0 \in (0,1)$, $\varepsilon > 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$. Again, (31) yields

$$\mathcal{D}_{\varepsilon}(f) \ge \kappa_0^4 \mathcal{D}_0(\varphi_{\varepsilon}(f)).$$

Remarking that $\varphi_{\varepsilon}(f)(v) \geq f(v) \geq K_0 e^{-A_0|v|^{q_0}}$, we apply [30, Theorem 4.1] to the function $\varphi_{\varepsilon}(f)$ and obtain for any $\alpha \in (0, 1)$ the existence of some $\overline{K_{\alpha}}(\varphi_{\varepsilon}(f))$ depending only on α , upper and lower bounds on ρ_{ε} , T_{ε} (see Remark 4 below), q_0 and upper bounds on A_0 , $1/K_0$, $\|\varphi_{\varepsilon}(f)\|_{L^1_s}$ and $\|\varphi_{\varepsilon}(f)\|_{H^k}$ where $s = s(\alpha, q_0, \beta_+, \beta_-)$ and $k = k(\alpha, s, \beta_+, \beta_-)$ such that

$$\mathcal{D}_0(\varphi_{\varepsilon}(f)) \ge \overline{K}_{\alpha}\left(\varphi_{\varepsilon}(f)\right) H_0\left[\varphi_{\varepsilon}(f) \middle| M^{\varphi_{\varepsilon}(f)} \right]^{1+\alpha}$$

Again, Theorem 1's lower-bound inequality (32) implies

$$H_0\left[\varphi_{\varepsilon}(f)\Big|M^{\varphi_{\varepsilon}(f)}\right] \ge H_{\varepsilon}[f|\mathcal{M}^f_{\varepsilon}],$$

so that

$$\mathcal{D}_{\varepsilon}(f) \ge \kappa_0^4 \overline{K}_{\alpha} \left(\varphi_{\varepsilon}(f) \right) \ H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^f]^{1+\alpha}.$$

Moreover, we can upper-bound $\|\varphi_{\varepsilon}(f)\|_{L^{1}_{s}}$ by $\kappa_{0}^{-1}\|f\|_{L^{1}_{s}}$, and $\|\varphi_{\varepsilon}(f)\|_{H^{k}}$ by a polynomial in $(\|f\|_{H^{l}})_{l\leq k}, \kappa_{0}^{-1}$ and an upper bound on ε , since for $l \geq 1$, and x such that $1 - \varepsilon x \geq \kappa_{0}$,

$$|\varphi_{\varepsilon}^{(l)}(x)| = \frac{l! \varepsilon^{l-1}}{(1-\varepsilon x)^{l+1}} \le l! \kappa_0^{-l-1} \varepsilon^{l-1}.$$

We end the proof by using

- Equations (72)–(73) stating that $\rho_{\varepsilon} \leq \kappa_0^{-1} \rho$ and $T_{\varepsilon} \geq \kappa_0 T$,
- the fact that $\rho \leq \rho_{\varepsilon}$, coming from

$$\int_{\mathbb{R}^3} f \, \mathrm{d}v \le \int_{\mathbb{R}^3} \frac{f}{1 - \varepsilon f} \, \mathrm{d}v,$$

• the fact that $T_{\varepsilon} \leq \kappa_0^{-1} T$, coming from

$$3\rho_{\varepsilon}T_{\varepsilon} + \rho_{\varepsilon}|u_{\varepsilon} - u|^2 = \int_{\mathbb{R}^3} \varphi_{\varepsilon}(f)(v) |v - u|^2 \,\mathrm{d}v \le \kappa_0^{-1} \int_{\mathbb{R}^3} f(v) |v - u|^2 \,\mathrm{d}v = \kappa_0^{-1} \,3\rho T.$$

implying $\rho_{\varepsilon}T_{\varepsilon} \le \kappa_0^{-1}\rho T$ hence $T_{\varepsilon} \le \kappa_0^{-1} \frac{\rho}{\rho_{\varepsilon}}T \le \kappa_0^{-1}T.$

Remark 4. In [30, Theorem 4.1], the constant $\overline{K}_{\alpha}(\varphi_{\varepsilon}(f))$ is said to depend on ρ_{ε} , T_{ε} . Having a look at the proof there, this dependence can be relaxed into a dependence on upper and lower bounds of those quantities.

4.3. Inverse result. Let $\varepsilon \in (0, \frac{1}{2})$, $B_0 > 0$ and $0 \le \beta < 2$. We assume that

$$\int_{\mathbb{S}^2} B(v, v_*, \sigma) \,\mathrm{d}\sigma \le B_0 \left(1 + |v - v_*|^\beta\right)$$

Relying on Bobylev and Cercignani's [5, Theorem 1], and the nice formulation of this theorem by Villani [30, Theorem 1.1], we know that there exists a family of normalized functions (g_l) , in the sense that for any l,

(77)
$$\int_{\mathbb{R}^3} g_l(v) \begin{pmatrix} 1\\v\\|v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} 1\\0\\3 \end{pmatrix},$$

such that (taking $\delta = 1 - \frac{1+2\varepsilon}{2} > 0$ in [30, Theorem 1.1])

(78)
$$\forall v \in \mathbb{R}^3, \qquad 2 \ge g_l(v) \ge \frac{1+2\varepsilon}{2} (2\pi)^{-3/2} e^{-|v|^2},$$

and

(79)
$$\frac{\mathcal{D}_0(g_l)}{H_0[g_l|M^{g_l}]} \xrightarrow[l \to \infty]{} 0.$$

We now consider

$$f_l^{\varepsilon} = \frac{g_l}{1 + \varepsilon g_l} \equiv \varphi_{\varepsilon}^{-1}(g_l).$$

As $\varphi_{\varepsilon}^{-1}$ is increasing, and from (78), we have

$$f_l^{\varepsilon}(v) \equiv \frac{g_l(v)}{1 + \varepsilon g_l(v)} \ge \frac{g_l(v)}{1 + 2\varepsilon} \ge \frac{1 + 2\varepsilon}{2(1 + 2\varepsilon)} (2\pi)^{-3/2} e^{-|v|^2} = \frac{1}{2} (2\pi)^{-3/2} e^{-|v|^2}.$$

We denote

$$\int_{\mathbb{R}^3} f_l^{\varepsilon}(v) \begin{pmatrix} 1\\v\\|v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho_{\varepsilon}^l\\ \rho_{\varepsilon}^l u_{\varepsilon}^l\\ 3\rho_{\varepsilon}^l T_{\varepsilon}^l + \rho_{\varepsilon}^l |u_{\varepsilon}^l|^2 \end{pmatrix},$$

We have, from $f_l^{\varepsilon} = \frac{g_l}{1 + \varepsilon g_l}$ and $g_l \le 2$,

$$\frac{1}{1+2\varepsilon} \int_{\mathbb{R}^3} g_l(v) \, \mathrm{d}v \le \int_{\mathbb{R}^3} f_l^{\varepsilon}(v) \, \mathrm{d}v \le \int_{\mathbb{R}^3} g_l(v) \, \mathrm{d}v$$

that is

$$\frac{1}{1+2\varepsilon} \le \rho_{\varepsilon}^{l} \le 1.$$

Secondly,

$$3\rho_{\varepsilon}^{l}T_{\varepsilon}^{l} \leq 3\rho_{\varepsilon}^{l}T_{\varepsilon}^{l} + \rho_{\varepsilon}^{l}|u_{\varepsilon}^{l}|^{2} = \int_{\mathbb{R}^{3}} f_{l}^{\varepsilon}(v) |v|^{2} \,\mathrm{d}v \leq \int_{\mathbb{R}^{3}} g_{l}(v) |v|^{2} \,\mathrm{d}v = 3,$$

that is $\rho_{\varepsilon}^{l}T_{\varepsilon}^{l} \leq 1$, hence

$$T_{\varepsilon}^{l} \leq \frac{1}{\rho_{\varepsilon}^{l}} \leq 1 + 2\varepsilon.$$

On the other hand, since g_l is normalized by (77), we have

$$3\rho_{\varepsilon}^{l}T_{\varepsilon}^{l} = \int_{\mathbb{R}^{3}} f_{l}^{\varepsilon}(v) |v - u_{\varepsilon}^{l}|^{2} dv \geq \frac{1}{1 + 2\varepsilon} \int_{\mathbb{R}^{3}} g_{l}(v) |v - u_{\varepsilon}^{l}|^{2} dv = \frac{3 + |u_{\varepsilon}^{l}|^{2}}{1 + 2\varepsilon} \geq \frac{3}{1 + 2\varepsilon},$$
 that is $\rho_{\varepsilon}^{l}T_{\varepsilon}^{l} \geq \frac{1}{1 + 2\varepsilon}$, hence

$$T^l_{\varepsilon} \geq \frac{1}{\rho^l_{\varepsilon}} \times \frac{1}{1+2\varepsilon} \geq \frac{1}{1+2\varepsilon}$$

Finally, since $f_l^{\varepsilon} = g_l - \varepsilon \frac{g_l^2}{1 + \varepsilon g_l}$, g_l is normalized by (77) and $g_l \leq 2$, we have

$$\begin{split} \rho_{\varepsilon}^{l} \left| u_{\varepsilon}^{l} \right| &= \left| \int_{\mathbb{R}^{3}} f_{l}^{\varepsilon}(v) \, v \, \mathrm{d}v \right| = \varepsilon \left| \int_{\mathbb{R}^{3}} \frac{g_{l}(v)^{2}}{1 + \varepsilon g_{l}(v)} \, v \, \mathrm{d}v \right| \leq \varepsilon \int_{\mathbb{R}^{3}} \frac{g_{l}(v)^{2}}{1 + \varepsilon g_{l}(v)} \left| v \right| \, \mathrm{d}v \leq 2\varepsilon \int_{\mathbb{R}^{3}} g_{l}(v) \left| v \right| \, \mathrm{d}v \\ &\leq 2\varepsilon \left(\int_{\mathbb{R}^{3}} g_{l}(v) \, \mathrm{d}v \right)^{1/2} \left(\int_{\mathbb{R}^{3}} g_{l}(v) \left| v \right|^{2} \, \mathrm{d}v \right)^{1/2} = 2\sqrt{3}\varepsilon, \end{split}$$

where the last inequality is a Cauchy-Schwarz argument. We then conclude, since $\varepsilon \in (0, \frac{1}{2})$, that

$$|u_{\varepsilon}^{l}| \leq \frac{1}{\rho_{\varepsilon}^{l}} 2\sqrt{3} \varepsilon \leq 2\sqrt{3} (1+2\varepsilon) \varepsilon \leq 4\sqrt{3} \varepsilon$$

All in all, we proved that for any $l, f_l^{\varepsilon} \in \mathcal{C}_{1,0,1}^{\varepsilon}$.

We now apply Equation (31) to get

$$\mathcal{D}_{\varepsilon}(f_l^{\varepsilon}) \leq \mathcal{D}_0(g_l),$$

and Proposition 2's upper-bound inequality, Equation (34), which gives, for any $\varepsilon \in (0, \frac{1}{2})$, and denoting $\kappa_0^{\varepsilon} = 1 - 2\varepsilon$ (so that $1 - \varepsilon f_l^{\varepsilon} \ge \kappa_0^{\varepsilon}$),

$$H_{\varepsilon}\left[f_{l}^{\varepsilon}\left|\mathcal{M}_{\varepsilon}^{f_{l}^{\varepsilon}}\right] \geq \frac{1}{C_{0}(\kappa_{0}^{\varepsilon})}H_{0}\left[\frac{f_{l}^{\varepsilon}}{1-\varepsilon f_{l}^{\varepsilon}}\left|M^{\frac{f_{l}^{\varepsilon}}{1-\varepsilon f_{l}^{\varepsilon}}}\right] = \frac{1}{C_{0}(\kappa_{0}^{\varepsilon})}H_{0}[g_{l}|M^{g_{l}}].$$

We finally obtain that, for any $\varepsilon \in (0, \frac{1}{2})$,

(80)
$$\frac{\mathcal{D}_{\varepsilon}\left(f_{l}^{\varepsilon}\right)}{H_{\varepsilon}\left[f_{l}^{\varepsilon}\left|\mathcal{M}_{\varepsilon}^{f_{l}^{\varepsilon}}\right]\right]} \leq \frac{1}{C_{0}(\kappa_{0}^{\varepsilon})} \times \frac{\mathcal{D}_{0}(g_{l})}{H_{0}[g_{l}|M^{g_{l}}]} \underset{l \to \infty}{\longrightarrow} 0.$$

5. About the Landau-Fermi-Dirac equation

A study of the entropy-entropy production relationship for the Landau-Fermi-Dirac equation for various cross sections was recently conducted by Alonso, Bagland, Desvillettes and Lods in [1, 2] (see also Desvillettes [15] and Alonso, Bagland, Lods [3]) and we do not (intend to) provide here any new result on this side. We refer the interested reader to these papers and the references therein for a detailed study of the Landau-Fermi-Dirac equation. Nevertheless, using the exact same strategy of transposing results from the Classical towards the Fermi-Dirac case that we used in the Boltzmann-Fermi-Dirac case (Corollary 3), we obtain an original proof of inequalities reminiscent of the ones stated in [1, 2] and [15], namely [1, Theorem 1.4], the first inequality in the proof of [2, Proposition 5.8] and [15, Proposition 2]. Let us first briefly introduce the Landau-Fermi-Dirac equation.

In dimension 3, the Landau-Fermi-Dirac operator writes, for $\varepsilon \ge 0$ and nonnegative $f \in L_2^1(\mathbb{R}^3)$ such that $1 - \varepsilon f \ge 0$,

(81)
$$Q_{\varepsilon}^{L}(f)(v) := \nabla_{v} \cdot \int_{\mathbb{R}^{3}} \Psi(|v - v_{*}|) \Pi(v - v_{*}) \Big[f_{*}(1 - \varepsilon f_{*}) \nabla f - f(1 - \varepsilon f) \nabla f_{*} \Big] dv_{*}$$

The case $\varepsilon = 0$ corresponds to the classical Landau operator. In (81), we used the common short-hands $f \equiv f(v)$ and $f_* \equiv f(v_*)$, $\Pi(z)$ denotes the orthogonal projection on $(\mathbb{R}z)^{\perp}$, whose components are

$$\Pi_{ij}(z) = \delta_{ij} - \frac{z_i z_j}{|z|^2},$$

and Ψ is sometimes called the kinetic potential. The entropy production associated to the Landau-Fermi-Dirac operator writes

(82)
$$\mathcal{D}_{\varepsilon}^{L}(f) = \frac{1}{2} \int_{\mathbb{R}^{3} \times \mathbb{R}^{3}} \Psi(|v - v_{*}|) ff_{*}(1 - \varepsilon f)(1 - \varepsilon f_{*}) \left| \Pi(v - v_{*}) \left(\frac{\nabla f}{f(1 - \varepsilon f)} - \frac{\nabla f_{*}}{f_{*}(1 - \varepsilon f_{*})} \right) \right|^{2} \mathrm{d}v \,\mathrm{d}v_{*}.$$

We notice that (82) can be rewritten (where φ_{ε} is defined in (6))

$$\mathcal{D}_{\varepsilon}^{L}(f) = \frac{1}{2} \int_{\mathbb{R}^{3} \times \mathbb{R}^{3}} \Psi(|v - v_{*}|) \varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f) \varphi_{\varepsilon}(f) (1 - \varepsilon f)^{2} (1 - \varepsilon f_{*})^{2} \left| \Pi(v - v_{*}) \left(\frac{\nabla \varphi_{\varepsilon}(f)}{\varphi_{\varepsilon}(f)} - \frac{\nabla \varphi_{\varepsilon}(f)_{*}}{\varphi_{\varepsilon}(f)_{*}} \right) \right|^{2} dv dv_{*}.$$

This directly implies that for any $\kappa_0 \in (0, 1)$, $\varepsilon > 0$ and nonnegative $f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$,

(83)
$$\mathcal{D}_0^L(\varphi_{\varepsilon}(f)) \ge \mathcal{D}_{\varepsilon}^L(f) \ge \kappa_0^4 \mathcal{D}_0^L(\varphi_{\varepsilon}(f)).$$

This inequality, similar to (31), allows to apply the same strategy as in the Boltzmann-Fermi-Dirac case. Adapting the entropy-entropy production inequalities known for the Landau equation [29, Chapter 3, Theorem 14] and [15, Remark 1], we get the following proposition. **Proposition 8.** Entropy-entropy production inequalities for the Landau-Fermi-Dirac operator (adaptation of [29, Chapter 3, Theorem 14] and [15, Remark 1]). We recall that $\mathcal{D}_{\varepsilon}^{L}$, H_{ε} and $T_{*}(f)$ are defined respectively in (82), (21)-(22) and (13).

• Over-Maxwellian case [Adaptation of [29, Chapter 3, Theorem 14, (i)]]. Assume $\Psi(|z|) \ge |z|^2$. Then, for any $\kappa_0 \in (0,1)$, $\varepsilon \ge 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$, we have

(84)
$$\mathcal{D}_{\varepsilon}^{L}(f) \ge 4 \,\kappa_{0}^{4} \,\rho \, T_{*}(f) \,H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}],$$

• Soft potentials [Adaptation of [29, Chapter 3, Theorem 14, (ii)]]. Assume $\Psi(|z|) \ge |z|^2(1+|z|)^{-\beta}$ with $\beta > 0$. Then, for any $\kappa_0 \in (0,1)$, $\varepsilon \ge 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$, for any s > 0, there exists a constant $C_s^L(f)$, explicit and depending on f only via ρ , T, κ_0 , an upper bound on ε and an upper bound on $H_{\varepsilon}(f)$ such that

(85)
$$\mathcal{D}_{\varepsilon}^{L}(f) \geq C_{s}^{L}(f) \left(\|f\|_{L_{s+2}^{1}} + \|\nabla\sqrt{f}\|_{L_{1+\frac{s}{2}}^{2}}^{2} \right)^{-\frac{\beta}{s}} H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}]^{1+\frac{\beta}{s}}$$

• Hard potentials [Adaptation of [15, Remark 1]]. Assume $\Psi(|z|) \ge |z|^{2+\beta}$ with $\beta \in (0,1]$. Then, for any $\kappa_0 \in (0,1)$, $\varepsilon \ge 0$ and $0 \le f \in L^1_2(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge \kappa_0$, there exist two explicit constants $K^L_1(f)$ and $K^L_2(f)$ which depend on β , on f only via ρ , T, κ_0 and on an upper bound on $\|f\|^2_{L^2_6}$ such that

(86)
$$\mathcal{D}^{L}_{\varepsilon}(f) \leq K_{1}^{L}(f) \implies \mathcal{D}^{L}_{\varepsilon}(f) \geq K_{2}^{L}(f) H_{\varepsilon}[f|\mathcal{M}^{f}_{\varepsilon}].$$

We briefly discuss here the qualitative difference of our results with respect to the ones in [1, 2] and [15]. The inequality we obtain in the over-Maxwellian case (84) is almost identical to the one in [1, Theorem 1.4], with however an extra quantity in the constant in their inequality. As for the soft potentials case, the inequality in the proof of [2, Proposition 5.8] involves the hard potential of order $\eta > 0$ entropy production (for some chosen $\eta > 0$), which relates in (85) with the $\|\nabla \sqrt{f}\|_{L^2_{1+\frac{8}{3}}}^2$ norm (for some chosen s > 0). We also point

out that, while [2, Proposition 5.8] holds for soft potentials of order at most $-\frac{4}{3}$, there is no such restriction in (85). As far as hard potentials are concerned, the result we obtain (which could actually be more precise, but somewhat more complicated, by using [15, Theorem 1] instead of [15, Remark 1]) is very similar to the one obtained by Desvillettes in [15, Proposition 2], with the difference that his result involves the relative Fisher information instead of the relative entropy (hence yielding a stronger result); our methodology only gives another proof to obtain it. Also, the inequality obtained in [1, Theorem 1.4] comes with a more complicated constant than in [15, Proposition 2] (similarly, than (86)), but holds without any assumption of smallness on $\mathcal{D}_{\varepsilon}^{L}(f)$. We mention that a statement of the type (86) entails an exponential decay with explicit constants (if $K_{1}^{L}(f)$ and $K_{2}^{L}(f)$ are controlled below) as shown in [15, Lemma 1]. Finally, we can also obtain inequalities for hard potentials of order $\beta > 1$ by adapting [29, Chapter 3, Theorem 14, (iii)].

Remark 5. We want to mention [13] which bounds below the entropy production with a weighted version of the Fisher information, which would lead to a weighted relative entropy with power one in the right-hand-side of (85). A similar result for hard potentials was proven in [10]. We refer the interested reader to [14] (in french) for a discussion on the entropic structure of the Landau operator. However our method of adapting results from the Classical to the Fermi-Dirac situation breaks here, as we did not prove any link between weighted entropies.

Convergence towards equilibrium. As an example of application of Proposition 8, we provide a convergence to equilibrium proposition in the hard potential case. We remind that this type of result was already obtained in [1] (and in [2] for soft potentials), although the constants that we obtain do differ, and our proof is shorter.

Proposition 9. Hard potentials case. Assume $\Psi(|z|) = |z|^{2+\beta}$ with $\beta \in (0,1]$. Consider $0 \leq f^{in} \in L^1_{s_\beta}(\mathbb{R}^3) \cap L^{\infty}(\mathbb{R}^3)$ with $s_\beta = \max\left(\frac{3\beta}{2}, 4-\beta\right)$. Then there exists $\varepsilon^{in} > 0$ depending only on f^{in} such that for any $\varepsilon \in (0, \varepsilon^{in}]$, there exists a solution f^{ε} to the homogeneous Landau-Fermi-Dirac equation (associated to the collision operator Q_{ε}^L defined in (81) and the initial distribution f^{in}) and two constants $C_1, C_2 > 0$ such that for any $t \geq 0$,

(87)
$$\|f^{\varepsilon}(t,\cdot) - \mathcal{M}^{f^{in}}_{\varepsilon}\|_{L^{1}_{2}} \leq C_{1} e^{-C_{2}t}.$$

Proof. The proof is similar to the one of Proposition 5, to which we refer the reader, and we only outline here the crucial points. First, we rely on [4, Corollary 3.7] to obtain the existence of $\varepsilon^{in} > 0$ depending only on f^{in} such that for any $\varepsilon \in (0, \varepsilon^{in}]$, the solution f^{ε} associated to LFD_{ε} satisfies

$$\sup_{t \ge 1} \left(1 - \varepsilon \| f^{\varepsilon}(t, \cdot) \|_{\infty} \right) \ge \frac{1}{2}.$$

We are now able to use (86). In order for the proof to be complete, we must ensure that the coefficient that multiplies the relative entropy can be lower bounded uniformly in time. It amounts to upper bound the L_6^2 norm of $f^{\varepsilon}(t, \cdot)$ uniformly in time. This is possible as we have a uniform-in-time L^{∞} bound and [1, Theorem 1.3] ensures the generation and uniform-in-time boundedness of all moments at some explicit time. Then the same argument as in [15, Lemma 1] provides the announced exponential decay to equilibrium.

6. L^1 - L^2 weighted Csiszár-Kullback-Pinsker inequality

In this section, we discuss an optimized version of the L^1-L^2 weighted Csiszár-Kullback-Pinsker (CKP) inequality presented in Proposition 4. For a discussion on the CKP inequality and its previous generalizations, we refer the reader to the paragraph preceding Proposition 4 in Subsection 2.3.

We highlight that the idea of the proof of (90) in the case r = 1, $\varpi = 1$, can be found in [22], and the proof of (104) (Bose-Einstein case) for r = 1, $\varpi = 1$ was also done with the same approach by Lu [23]. Notice moreover that we consider here the case when f is defined on \mathbb{R}^3 , but (89)–(90) and (104) do hold in the broader setting where f and ϖ are defined on an arbitrary measured set, as long as the equilibrium distribution is well-defined. This is further detailed in Appendix B. Finally, we inform the reader that the Bose-Einstein case is briefly discussed in Appendix A, where useful definitions may be found.

Proposition 10. L^1 - L^2 weighted Csiszár-Kullback-Pinsker inequality. [optimal]

Let $\varpi : \mathbb{R}^3 \to \mathbb{R}_+$ be measurable, and $r \in [1, 2]$. Denote

(88)
$$\Lambda(\lambda) := \begin{cases} 2 & \text{if } \lambda = 1, \\ \frac{(\lambda - 1)^2}{\lambda \log \lambda - \lambda + 1} & \text{if } \lambda \in \mathbb{R}_+ \setminus \{1\}. \end{cases}$$

• Classical CKP inequality. For any $0 \leq f \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$, assuming that the norms below are finite,

(89)
$$\|(f-M)\varpi\|_{L^{r}}^{2} \leq \|M\varpi^{2}\|_{L^{\frac{r}{2-r}}} \Lambda\left(\frac{\|f\varpi^{2}\|_{L^{\frac{r}{2-r}}}}{\|M\varpi^{2}\|_{L^{\frac{r}{2-r}}}}\right) H_{0}[f|M],$$

where we denoted $M \equiv M^f$ the Maxwellian distribution associated to f and H_0 is defined in (21)–(22) (with $\varepsilon = 0$).

• Fermi-Dirac CKP inequality. For any $\varepsilon > 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge 0$ and $\gamma > \frac{2}{5}$, assuming that the norms below are finite,

(90)
$$\|(f - \mathcal{M}) \varpi\|_{L^{r}}^{2} \leq \left\|\mathcal{M} \varpi^{2}\right\|_{L^{\frac{r}{2-r}}} \Lambda\left(\frac{\left\|f \varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}{\left\|\mathcal{M} \varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}\right) H_{\varepsilon}[f|\mathcal{M}],$$

where we denoted $\mathcal{M} \equiv \mathcal{M}_{\varepsilon}^{f}$ the ε -Fermi-Dirac distribution associated to f and H_{ε} is defined in (21)–(22).

The announced inequalities (40)–(41) of Proposition 4 are then consequence of Proposition 10 and the fact that for any $a \ge 0$ and b > 0,

(91)
$$b \Lambda\left(\frac{a}{b}\right) \le 2 \max(a, b)$$

The latter inequality can be deduced from the following ones. We claim that in fact,

(92)
$$\Lambda(\lambda) \le 1 + \lambda^{2/3}, \qquad \lambda \in [0, 1],$$

(93)
$$\Lambda(\lambda) \le \frac{2}{3}(2+\lambda), \qquad \lambda \in (1,10)$$

(94)
$$\Lambda(\lambda) \le \frac{\lambda}{\log(\lambda) - 1}, \qquad \lambda \ge 10.$$

Notice the improvement of (94) over (91) by a factor $\frac{1}{2(\log a - \log b - 1)}$ when $\lambda = \frac{a}{b}$ is large, which, even if slowly, converges towards 0 as a goes to infinity. The second inequality (93) is well-known, and is actually the one used in the standard proof of the usual CKP inequality (see for instance [19]). As for the third one (94), we have for $\lambda \geq 10$,

$$\Lambda(\lambda) = \frac{(\lambda - 1)^2}{\lambda \log \lambda - \lambda + 1} = \frac{\lambda - 1}{\frac{\lambda}{\lambda - 1} \log \lambda - 1} \le \frac{\lambda}{\log(\lambda) - 1}.$$

Finally, the proof of the first inequality (92) is trickier and was suggested to me by Matthieu Dolbeault. Changing variables $x = \lambda^{1/3}$, it is enough to show that for any $x \in [0, 1]$,

$$(1+x^2)(3x^3\log x - x^3 + 1) - (1-x^3)^2 \ge 0.$$

Developing, the above term actually equals $x^2g(x)$, where

$$g(x) = 1 + x - x^{3} - x^{4} + 3(x + x^{3})\log x.$$

We can then conclude once we prove that g is nonnegative on (0, 1). We compute its first four derivatives,

$$g'(x) = 4 - 4x^3 + (3 + 9x^2)\log x, \qquad g''(x) = \frac{3}{x} + 9x - 12x^2 + 18x\log x,$$

$$g'''(x) = 27 - \frac{3}{x^2} - 24x + 18\log x, \qquad g^{(4)}(x) = 6(-4 + x^{-3} + 3x^{-1}).$$

The function $g^{(4)}$ is clearly nonnegative on (0, 1), hence g''' is nondecreasing on (0, 1). But g'''(1) = 0, hence g''' is nonpositive on (0, 1), thus g'' is nonincreasing on (0, 1). Again, g''(1) = 0 hence g'' is nonnegative, thus g' is nondecreasing. Again, g'(1) = 0 hence g' is nonpositive, thus g is nonincreasing on (0, 1). Finally, g(1) = 0, so that g is nonnegative on (0, 1).

Let us now prove Proposition 10.

Proof. This proposition is a consequence of the general inequality (114) in Corollary 17 in Appendix B. We prove both inequalities (89)–(90) simultaneously, as the first one corresponds to the limit case $\varepsilon = 0$ of the second. Let us then consider $\varepsilon \ge 0$.

We apply Corollary 17 with $\Phi(x) \equiv \Phi_{\varepsilon}(x) = \int_0^x \log \varphi_{\varepsilon}(y) \, \mathrm{d}y, \ J = (0, \varepsilon^{-1})$ and

$$\mathcal{F} = \left\{ 0 \le g \in L_2^1(\mathbb{R}^3) \, \middle| \, 1 - \varepsilon g \ge 0, \quad \int_{\mathbb{R}^3} g(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho \\ \rho u \\ 3 \rho T + \rho |u|^2 \end{pmatrix} \right\},$$

with ρ, T and ε such that $\gamma > \frac{2}{5}$ (ensuring the existence of $\mathcal{M}_{\varepsilon}^{f}$). Then (114) writes, since $H_{\Phi_{\varepsilon}} \equiv H_{\varepsilon}$ the ε -Fermi (or Classical in the case $\varepsilon = 0$) entropy, and $\frac{1}{\Phi_{\varepsilon}''}(x) = x(1 - \varepsilon x) \leq x$,

$$\left\| (f - \mathcal{M}_{\varepsilon}^{f}) \,\varpi \right\|_{L^{r}}^{2} \leq \left(\int_{0}^{1} (1 - \tau) \,\left\| ((1 - \tau)\mathcal{M}_{\varepsilon}^{f} + \tau f) \,\varpi^{2} \right\|_{L^{\frac{r}{2-r}}}^{-1} \,\mathrm{d}\tau \right)^{-1} \,H_{\varepsilon}[f|\mathcal{M}_{\varepsilon}^{f}].$$

We focus on the integral in the variable τ . From Minkowski's inequality, we have

$$\left(\int_0^1 \frac{1-\tau}{\left\| \left((1-\tau)\mathcal{M}_{\varepsilon}^f + \tau f \right) \varpi^2 \right\|_{L^{\frac{r}{2-r}}}} \, \mathrm{d}\tau \right)^{-1} \leq \left(\int_0^1 \frac{1-\tau}{(1-\tau) \left\| \mathcal{M}_{\varepsilon}^f \varpi^2 \right\|_{L^{\frac{r}{2-r}}} + \tau \left\| f \varpi^2 \right\|_{L^{\frac{r}{2-r}}}} \, \mathrm{d}\tau \right)^{-1} \\ = \left\| \mathcal{M}_{\varepsilon}^f \varpi^2 \right\|_{L^{\frac{r}{2-r}}} \, \Lambda \left(\frac{\left\| f \varpi^2 \right\|_{L^{\frac{r}{2-r}}}}{\left\| \mathcal{M}_{\varepsilon}^f \varpi^2 \right\|_{L^{\frac{r}{2-r}}}} \right),$$

where Λ is defined in (88). Indeed, this last equality comes from the Taylor expansion of $\lambda \mapsto \lambda \log \lambda - \lambda$ around 1,

$$\lambda \log \lambda - \lambda = -1 + (\lambda - 1) \log 1 + (\lambda - 1)^2 \int_0^1 \frac{1 - \tau}{(1 - \tau) + \tau \lambda} d\tau,$$

that is

$$\int_0^1 \frac{1-\tau}{(1-\tau)+\tau\lambda} \,\mathrm{d}\tau = \begin{cases} \frac{1}{2} & \text{if } \lambda = 1, \\ \frac{\lambda \log \lambda - \lambda + 1}{(\lambda - 1)^2} & \text{if } \lambda \in \mathbb{R}_+ \setminus \{1\} \\ = \frac{1}{\Lambda(\lambda)}. \end{cases}$$

Finally, as a corollary to Proposition 10 come the standard Csiszár-Kullback-Pinsker inequalities.

Corollary 11. Standard Csiszár-Kullback-Pinsker inequalities.

For any real number $x \in \mathbb{R}$, we denote in the following $x_{+} = \max(0, x)$.

• Standard Classical CKP inequalities. For any $\alpha \ge 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$,

(95) $||f - M||_{L^1}^2 \le 2 ||M||_{L^1} H_0[f|M],$

(96)
$$\|(M-f)_+\|_{L^1_{\alpha}}^2 \le 2 \|M\|_{L^1_{2\alpha}} H_0[f|M].$$

(97)
$$\|f - M\|_{L^{1}_{2}}^{2} \leq 8 \|M\|_{L^{1}_{4}} H_{0}[f|M],$$

where we denoted $M \equiv M^f$ the Maxwellian distribution associated to f.

• Standard Fermi-Dirac CKP inequalities. For any $\varepsilon > 0$, $\alpha \ge 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \setminus \{0\}$ such that $1 - \varepsilon f \ge 0$ and $\gamma > \frac{2}{5}$,

(98) $\|f - \mathcal{M}\|_{L^1}^2 \leq 2 \|\mathcal{M}\|_{L^1} H_{\varepsilon}[f|\mathcal{M}],$

(99)
$$\|(\mathcal{M} - f)_{+}\|_{L^{1}_{\alpha}}^{2} \leq 2 \|\mathcal{M}\|_{L^{1}_{2\alpha}} H_{\varepsilon}[f|\mathcal{M}]$$

(100)
$$\|f - \mathcal{M}\|_{L_2^1}^2 \leq 8 \|\mathcal{M}\|_{L_4^1} H_{\varepsilon}[f|\mathcal{M}].$$

where we denoted $\mathcal{M} \equiv \mathcal{M}_{\varepsilon}^{f}$ the ε -Fermi-Dirac distribution associated to f.

Proof. We only prove (98)–(100) as (95)–(97) can be seen as a limit case of the previous ones when $\varepsilon \to 0$.

• Proof of (98). Applying Proposition 10, specifically Equation (90), with r = 1 and $\varpi = 1$, we obtain

$$\|f - \mathcal{M}\|_{L^1}^2 \le \|\mathcal{M}\|_{L^1} \Lambda\left(\frac{\|f\|_{L^1}}{\|\mathcal{M}\|_{L^1}}\right) H_{\varepsilon}[f|\mathcal{M}].$$

Since $||f||_{L^1} = ||M||_{L^1}$ and $\Lambda(1) = 2$, we obtain (98).

• Proof of (99). Applying Proposition 10, specifically Equation (90), with $\varpi(v) = (1+|v|^2)^{\alpha/2} \mathbf{1}_{f \leq \mathcal{M}}$ and r = 1, we obtain

$$\|(\mathcal{M}-f)_+\|_{L^1_{\alpha}}^2 \leq \|\mathcal{M}\mathbf{1}_{f\leq\mathcal{M}}\|_{L^1_{2\alpha}} \Lambda\left(\frac{\|f\mathbf{1}_{f\leq\mathcal{M}}\|_{L^1_{2\alpha}}}{\|\mathcal{M}\mathbf{1}_{f\leq\mathcal{M}}\|_{L^1_{2\alpha}}}\right) H_{\varepsilon}[f|\mathcal{M}].$$

Since $\|f \mathbf{1}_{f \leq \mathcal{M}}\|_{L^{1}_{2\alpha}} \leq \|\mathcal{M} \mathbf{1}_{f \leq \mathcal{M}}\|_{L^{1}_{2\alpha}}$ and, from (92), $\Lambda \leq 2$ on [0,1], we obtain (99) after upper bounding $\|\mathcal{M} \mathbf{1}_{f \leq \mathcal{M}}\|_{L^{1}_{2\alpha}}$ by $\|\mathcal{M}\|_{L^{1}_{2\alpha}}$.

• Proof of (100). We remark that $|f - \mathcal{M}| = f - \mathcal{M} + 2(\mathcal{M} - f)_+$, and

$$\int_{\mathbb{R}^3} (f - \mathcal{M})(v) \left(1 + |v|^2\right) \mathrm{d}v = 0,$$

so that

$$\|f - \mathcal{M}\|_{L_2^1}^2 = 4\|(\mathcal{M} - f)_+\|_{L_2^1}^2.$$

We apply (99) with $\alpha = 2$ and obtain (100).

Appendix A. Similar results in the Bose-Einstein case

An upper-bound inequality similar to (34) can also be obtained in the Bose-Einstein case. This latter case formally corresponds to taking $-\varepsilon$ instead of ε in our formulas. First define for any $x \in \mathbb{R}_+$ and $\varepsilon > 0$,

$$\varphi_{\varepsilon}^{BE}(x) := \frac{x}{1 + \varepsilon x}, \qquad \Phi_{\varepsilon}^{BE}(x) := \int_{0}^{x} \log \varphi_{\varepsilon}^{BE}(y) \, \mathrm{d}y,$$

and the Bose-Einstein entropy of $0 \leq f \in L_2^1(\mathbb{R}^3)$:

$$H_{\varepsilon}^{BE}(f) := \int_{\mathbb{R}^3} \Phi_{\varepsilon}^{BE}(f) \, \mathrm{d}v.$$

Lu proved in [21] that, under the condition

(101)
$$T \ge \frac{\zeta(\frac{5}{2})}{\zeta(\frac{3}{2})} T_c, \qquad T_c := \frac{1}{2\pi} \left(\frac{\rho \varepsilon}{\zeta(\frac{3}{2})}\right)^{2/3},$$

where ζ is the Riemann Zêta function and T_c is called the critical temperature, there exists a unique ε -Bose-Einstein statistics $\mathcal{M}_{\varepsilon}^{BE,f}$ associated to f - that is a distribution such that $\log \varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})$ is a linear combination of $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$ and sharing the same normalization in $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$ as f. We can obtain from Proposition 15 in Appendix B that, denoting $H_{\varepsilon}^{BE}[f|\mathcal{M}_{\varepsilon}^{BE,f}] :=$ $H_{\varepsilon}^{BE}(f) - H_{\varepsilon}^{BE,f}(\mathcal{M}_{\varepsilon}^{BE,f})$, we have

$$H_{\varepsilon}^{BE}[f|\mathcal{M}_{\varepsilon}^{BE,f}] = \int_{0}^{1} (1-\tau) \left(\int_{\mathbb{R}^{3}} \left(f(v) - \mathcal{M}_{\varepsilon}^{BE,f}(v) \right)^{2} \Phi_{\varepsilon}^{BE''} \left((1-\tau) \mathcal{M}_{\varepsilon}^{BE,f}(v) + \tau f(v) \right) dv \right) d\tau$$

(102)
$$= \int_{\mathbb{R}^{3}} \int_{\mathcal{M}_{\varepsilon}^{BE,f}(v)}^{f(v)} \frac{f(v) - x}{\varphi_{\varepsilon}^{BE}(x)} \varphi_{\varepsilon}^{BE'}(x) dx dv,$$

where the last equality comes from $\Phi_{\varepsilon}^{BE''} = \frac{\varphi_{\varepsilon}^{BE'}}{\varphi_{\varepsilon}^{BE}}$ and the change of variables $x = \mathcal{M}_{\varepsilon}^{BE,f}(v) + \tau(f(v) - \mathcal{M}_{\varepsilon}^{BE,f})$.

In the following Proposition, we provide a link between the relative entropies to equilibrium of the Bose-Einstein and the classical cases. Although I believe that both inequalities could be obtained, we only present here the "upper-bound" inequality as its proof is rather short. Further work may allow to obtain the lower-bound inequality, with a constant that probably depends on an L^{∞} bound on f. This constitutes another reason why we did not investigate this other inequality, as, although L^{∞} bounds are natural to use in the Fermi-Dirac context, due to Pauli's exclusion principle, they are not in the Bose-Einstein one, due to the phenomenon of condensation.

Proposition 12. Upper-bound in the Bose-Einstein case. For any $\varepsilon > 0$ and nonnegative $f \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$ which density and temperature satisfy (101), we have

(103)
$$H_0\left[\frac{f}{1+\varepsilon f}\middle|M^{\frac{f}{1+\varepsilon f}}\right] \le H_{\varepsilon}^{BE}\left[f\middle|\mathcal{M}_{\varepsilon}^{BE,f}\right].$$

Proof. The starting point of the proof are Equation (102) and the inequality $|y - z| \ge |\varphi_{\varepsilon}^{BE}(y) - \varphi_{\varepsilon}^{BE}(z)|$ for all $(y, z) \in \mathbb{R}^2_+$, yielding

$$H_{\varepsilon}^{BE}\left[f|\mathcal{M}_{\varepsilon}^{BE,f}\right] = \int_{\mathbb{R}^{3}} \int_{\mathcal{M}_{\varepsilon}^{BE,f}}^{f} \frac{f-x}{\varphi_{\varepsilon}^{BE}(x)} \varphi_{\varepsilon}^{BE'}(x) \,\mathrm{d}x \,\mathrm{d}v \ge \int_{\mathbb{R}^{3}} \int_{\mathcal{M}_{\varepsilon}^{BE,f}}^{f} \frac{\varphi_{\varepsilon}^{BE}(f) - \varphi_{\varepsilon}^{BE}(x)}{\varphi_{\varepsilon}^{BE}(x)} \,\varphi_{\varepsilon}^{BE'}(x) \,\mathrm{d}x \,\mathrm{d}v.$$

Applying the change of variables $y = \varphi_{\varepsilon}^{BE}(x)$ and using formula (27), we obtain

$$H_{\varepsilon}^{BE}\left[f|\mathcal{M}_{\varepsilon}^{BE,f}\right] \geq H_{0}\left[\varphi_{\varepsilon}^{BE}(f)|M^{\varphi_{\varepsilon}^{BE}(f)}\right] + \int_{\mathbb{R}^{3}} \int_{\varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})}^{M^{\varphi_{\varepsilon}^{BE}(f)}} \frac{\varphi_{\varepsilon}^{BE}(f) - y}{y} \, \mathrm{d}y \, \mathrm{d}v.$$

Remark that, as $\log M^{\varphi_{\varepsilon}^{BE}(f)} - \log \varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})$ is a linear combination of conserved quantities (namely, $v \mapsto 1, v \mapsto v$ and $v \mapsto |v|^2$), we have by definition of $M^{\varphi_{\varepsilon}^{BE}(f)}$ that

$$\int_{\mathbb{R}^3} \varphi_{\varepsilon}^{BE}(f) \log\left(\frac{M^{\varphi_{\varepsilon}^{BE}(f)}}{\varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})}\right) \mathrm{d}v = \int_{\mathbb{R}^3} M^{\varphi_{\varepsilon}^{BE}(f)} \log\left(\frac{M^{\varphi_{\varepsilon}^{BE}(f)}}{\varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})}\right) \mathrm{d}v,$$

so that

$$\int_{\mathbb{R}^3} \int_{\varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})}^{M^{\varphi_{\varepsilon}^{BE}(f)}} \frac{\varphi_{\varepsilon}^{BE}(f) - y}{y} \, \mathrm{d}y \, \mathrm{d}v = \int_{\mathbb{R}^3} \int_{\varphi_{\varepsilon}^{BE}(\mathcal{M}_{\varepsilon}^{BE,f})}^{M^{\varphi_{\varepsilon}^{BE}(f)}} \frac{M^{\varphi_{\varepsilon}^{BE}(f)} - y}{y} \, \mathrm{d}y \, \mathrm{d}v \ge 0,$$
option

ending the proof.

Proposition 13. Bose-Einstein CKP inequality. Let $\varpi : \mathbb{R}^3 \to \mathbb{R}_+$ be measurable, and $r \in [1,2]$. We recall the definition (88) of the function Λ ,

$$\Lambda(\lambda) := \begin{cases} 2 & \text{if } \lambda = 1, \\ \frac{(\lambda - 1)^2}{\lambda \log \lambda - \lambda + 1} & \text{if } \lambda \in \mathbb{R}_+ \setminus \{1\} \end{cases}$$

Then for any $\varepsilon > 0$ and $0 \le f \in L^1_2(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$ which density and temperature satisfy (101), assuming that the norms below are finite,

(104)
$$\|(f - \mathcal{M}) \ \varpi\|_{L^{r}}^{2} \leq \left\|\mathcal{M}(1 + \varepsilon \mathcal{M}) \ \varpi^{2}\right\|_{L^{\frac{r}{2-r}}} \Lambda\left(\frac{\left\|f(1 + \varepsilon f) \ \varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}{\left\|\mathcal{M}(1 + \varepsilon \mathcal{M}) \ \varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}\right) H_{\varepsilon}^{BE} \left[f \left|\mathcal{M}\right]$$

where we denoted $\mathcal{M} \equiv \mathcal{M}_{\varepsilon}^{BE,f}$ the ε -Bose-Einstein distribution associated to f, and $\mathcal{H}_{\varepsilon}^{BE}$ is the Bose-Einstein entropy. When r = 2, $L^{\frac{r}{2-r}}$ shall be understood as L^{∞} .

Proof. The proof is similar to the one of Proposition 10. Let $\varepsilon > 0$. We apply Corollary 17 with $\Phi(x) \equiv \Phi_{\varepsilon}^{BE}(x) = \int_{0}^{x} \log \frac{y}{1+\varepsilon y} \, \mathrm{d}y, \ J = \mathbb{R}_{+}^{*}$ and

$$\mathcal{F} = \left\{ 0 \le g \in L_2^1(\mathbb{R}^3) \left| \int_{\mathbb{R}^3} g(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho \\ \rho u \\ 3 \rho T + \rho |u|^2 \end{pmatrix} \right\},\$$

with ρ, T and ε such that (101) is satisfied, ensuring the existence of $\mathcal{M}_{\varepsilon}^{BE,f}$. In the rest of this proof, we denote for the sake of clarity $\mathcal{M} \equiv \mathcal{M}_{\varepsilon}^{BE,f}$. Then (114) writes, since $H_{\Phi_{\varepsilon}^{BE}} \equiv H_{\varepsilon}^{BE}$ the ε -Bose-Einstein entropy, and $\frac{1}{\Phi_{\varepsilon}^{BE''}}(x) = x(1 + \varepsilon x)$ is convex,

$$\|(f-\mathcal{M})\,\varpi\|_{L^r}^2 \le \left(\int_0^1 (1-\tau)\,\left\|((1-\tau)\mathcal{M}(1+\varepsilon\mathcal{M})+\tau f(1+\varepsilon f))\,\varpi^2\right\|_{L^{\frac{r}{2-r}}}^{-1}\,\mathrm{d}\tau\right)^{-1}\,H_{\varepsilon}^{BE}[f|\mathcal{M}].$$

We focus on the term with the integral in τ . From Minkowski's inequality, it is smaller than

$$\begin{split} & \left(\int_{0}^{1} \frac{1-\tau}{(1-\tau) \left\|\mathcal{M}(1+\varepsilon\mathcal{M})\,\varpi^{2}\right\|_{L^{\frac{r}{2-r}}} + \tau \left\|f(1+\varepsilon f)\,\varpi^{2}\right\|_{L^{\frac{r}{2-r}}} \,\mathrm{d}\tau\right)^{-1} \\ &= \left\|\mathcal{M}(1+\varepsilon\mathcal{M})\,\varpi^{2}\right\|_{L^{\frac{r}{2-r}}} \,\Lambda\left(\frac{\left\|f(1+\varepsilon f)\,\varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}{\left\|\mathcal{M}(1+\varepsilon\mathcal{M})\,\varpi^{2}\right\|_{L^{\frac{r}{2-r}}}}\right), \end{split}$$

where Λ is defined in (88) and appears thanks to a Taylor expansion of $\lambda \mapsto \lambda \log \lambda - \lambda$ around 1 like in the proof of Proposition 10, allowing to conclude.

From the previous proposition, we easily deduce the following standard inequalities.

Corollary 14. Standard Bose-Einstein CKP inequalities. For any $\varepsilon > 0$, $\alpha \ge 0$ and $0 \le f \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3) \setminus \{0\}$ satisfying (101),

(105)
$$\|(\mathcal{M}-f)_+\|_{L^1_{\alpha}}^2 \le 2\left(\|\mathcal{M}\|_{L^1_{2\alpha}} + \varepsilon \|\mathcal{M}\|_{L^2_{\alpha}}^2\right) H^{BE}_{\varepsilon}[f|\mathcal{M}],$$

(106)
$$\|f - \mathcal{M}\|_{L^1}^2 \leq 8 \left(\|\mathcal{M}\|_{L^1} + \varepsilon \|\mathcal{M}\|_{L^2}^2\right) \quad H_{\varepsilon}^{BE}[f|\mathcal{M}]$$

(107)
$$\|f - \mathcal{M}\|_{L^{1}_{2}}^{2} \leq 8 \left(\|\mathcal{M}\|_{L^{1}_{4}} + \varepsilon \|\mathcal{M}\|_{L^{2}_{2}}^{2}\right) H^{BE}_{\varepsilon}[f|\mathcal{M}]_{\varepsilon}^{2}$$

where we denoted for clarity $\mathcal{M} \equiv \mathcal{M}_{\varepsilon}^{BE,f}$ the ε -Bose-Einstein distribution associated to f.

We prove the above inequalities similarly as we did for Corollary 11. For (105) we apply Proposition 13 with r = 1 and $\varpi(v) = (1 + |v|^2)^{\frac{\alpha}{2}} \mathbf{1}_{f \leq \mathcal{M}}$ and notice that $\Lambda \leq 2$ on [0, 1]. We then obtain (106)–(107) by decomposing $|f - \mathcal{M}| = f - \mathcal{M} + 2(\mathcal{M} - f)_+$, using the fact that f and \mathcal{M} share the same normalization in $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$, and (105) with respectively $\alpha = 0$ and $\alpha = 2$.

APPENDIX B. A GENERAL DISCUSSION ABOUT ENTROPIES AND EQUILIBRIA

In this section we intend to provide general considerations on the entropy, which are much more general than the scope of this paper, but give a good understanding of the notions we used, and could also be helpful in the study of weak turbulence, where various kinds of unusual entropies can emerge (see [7]). Our setting is laid down quite generally. Consider a measured space $(\mathcal{E}, \mathcal{A}, \mu)$, an open interval $J \subset \mathbb{R}$ which closure we denote by \overline{J} , and a function $\Phi \in C^2(J) \cap C^0(\overline{J})$ such that $\Phi'' > 0$ on J. Remark that Φ' is then a C^1 -diffeomorphism from J onto $\Phi'(J)$.

Entropy. We define the Φ -entropy, for any $(\mathcal{A}, \operatorname{Bor}(\overline{J}))$ -measurable $f : \mathcal{E} \to \overline{J}$ such that the following integral makes sense and is finite, by

(108)
$$H_{\Phi}(f) := \int_{\mathcal{E}} \Phi(f(\zeta)) \,\mathrm{d}\mu(\zeta),$$

and we denote by E_{Φ} the set of such f. We let the relative Φ -entropy of f and g to be

(109)
$$H_{\Phi}[f|g] = H_{\Phi}(f) - H_{\Phi}(g).$$

We also define, for $(\mathcal{A}, \operatorname{Bor}(\bar{J}))$ -measurable $f, g : \mathcal{E} \to \bar{J}$, the Φ -relative-entropy (which in general differs from the relative Φ -entropy) of f and g by

(110)
$$\mathcal{H}_{\Phi}[f|g] := \int_{0}^{1} (1-\tau) \left(\int_{g \in J} (f-g)^{2} \Phi''((1-\tau)g + \tau f) \,\mathrm{d}\mu(\zeta) \right) \mathrm{d}\tau.$$

Note that $\mathcal{H}_{\Phi}[f|g]$ is possibly infinite, but always well-defined, as

$$g \in J \implies \forall \tau \in (0,1), \ (1-\tau)g + \tau f \in J,$$

and that $\mathcal{H}_{\Phi}[f|g]$ is always nonnegative. The following Proposition 15 gives a quite simple but general result linking entropies, conserved quantities and equilibrium distributions, under a sole existence assumption.

Proposition 15. Let I be a countable set, $(\phi_i)_{i \in I}$ a family of measurable real functions, $(\omega_i)_{i \in I}$ a family of real numbers, and

$$\mathcal{F} = \left\{ f \in E_{\Phi} \text{ s.t. } \forall i \in I, \int_{\mathcal{E}} |f(\zeta)| |\phi_i(\zeta)| \, \mathrm{d}\mu(\zeta) < \infty \text{ and } \int_{\mathcal{E}} f(\zeta) \, \phi_i(\zeta) \, \mathrm{d}\mu(\zeta) = \omega_i \right\}.$$

Assume that $\Phi'(J) = \mathbb{R}$, and that there exists $(\alpha_i(\omega)) \in \mathbb{R}^I$ such that $M_{\Phi}^{\mathcal{F}} \in \mathcal{F}$, where

(111)
$$M_{\Phi}^{\mathcal{F}} := (\Phi')^{-1} \left(\sum_{i \in I} \alpha_i(\omega) \phi_i \right).$$

Then the following four propositions are equivalent. Let $g \in \mathcal{F}$.

(i)
$$g \in J$$
 μ -a.e. and $\forall f \in \mathcal{F}, \ H_{\Phi}[f|g] = \mathcal{H}_{\Phi}[f|g],$
(ii) $g \in J$ μ -a.e. and $\forall f \in \mathcal{F}, \ \mathcal{H}_{\Phi}[f|g] < \infty$ and $\int_{\mathcal{E}} (f-g) \Phi'(g) d\mu(\zeta) = 0$
(iii) $H_{\Phi}(g) = \min_{h \in \mathcal{F}} H_{\Phi}(h),$
(iv) $g = M_{\Phi}^{\mathcal{F}} \quad \mu$ -a.e.

In particular, $M_{\Phi}^{\mathcal{F}}$ is the unique minimizer of H_{Φ} under the constraints of the set \mathcal{F} .

The above proposition actually proves the following (under the assumptions $\Phi'(J) = \mathbb{R}$ and of existence of $M_{\Phi}^{\mathcal{F}}$). An admissible distribution g is an equilibrium relative to the conserved quantities ϕ_i , in the sense of the minimization of the Φ -entropy [(iii)], if and only if $\Phi'(g)$ is a linear combination of the functions ϕ_i [(iv)], if and only if the relative Φ -entropy between any admissible distribution f and g is given by (110) [(i)], if and only if the quantity $\Phi'(g)$ is conserved amongst all admissible distributions [(ii)] - indeed, (ii) is equivalent, assuming the following integrals make sense, to $\forall f_1, f_2 \in \mathcal{F}, \int_{\mathcal{E}} f_1 \Phi'(g) d\mu(\zeta) = \int_{\mathcal{E}} f_2 \Phi'(g) d\mu(\zeta)$.

The reader may notice that the Classical case corresponds to the choice $\Phi(x) \equiv \Phi_0(x) = x \log x - x$, for which $(\Phi'_0)^{-1} = \exp$, hence $M^{\mathcal{F}}_{\Phi}$ is in this case a Maxwellian, since the conserved quantities (corresponding to the functions ϕ_i in the proposition) are $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^2$.

Moreover, the Fermi-Dirac case corresponds to the choice $\Phi(x) \equiv \Phi_{\varepsilon}(x) \equiv \int_{0}^{x} \log \frac{y}{1-\varepsilon y} \, \mathrm{d}y$, for which $(\Phi'_{\varepsilon})^{-1}(x) = \frac{e^{x}}{1+\varepsilon e^{x}}$, hence $M_{\Phi}^{\mathcal{F}}$ is in this case a Fermi-Dirac distribution, since again, the conserved quantities (corresponding to the functions ϕ_{i} in the proposition) are $v \mapsto 1$, $v \mapsto v$ and $v \mapsto |v|^{2}$. Proposition 1 then comes as a corollary to Proposition 15 with $\mathcal{E} = \mathbb{R}^{3}$ endowed with the Lebesgue measure, $J = (0, \varepsilon^{-1})$, $\Phi(x) \equiv \Phi_{\varepsilon}(x) \equiv \int_{0}^{x} \log \varphi_{\varepsilon}(y) \, \mathrm{d}y, M_{\Phi}^{\mathcal{F}} \equiv \mathcal{M}_{\varepsilon}^{f}$ and

$$\mathcal{F} = \left\{ 0 \le f \in L_2^1(\mathbb{R}^3) \, \middle| \, 1 - \varepsilon f \ge 0, \quad \int_{\mathbb{R}^3} f(v) \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \, \mathrm{d}v = \begin{pmatrix} \rho \\ \rho u \\ 3 \rho T + \rho |u|^2 \end{pmatrix} \right\},$$

with ρ, T and ε such that $\gamma > \frac{2}{5}$ (ensuring the existence of $\mathcal{M}_{\varepsilon}^{f}$, as proven in [22]).

Remark 6. The Bose-Einstein case is also recovered with $\Phi(x) \equiv \Phi_{\varepsilon}^{BE}(x) \equiv \int_{0}^{x} \log \frac{y}{1+\varepsilon y} \, \mathrm{d}y$, for which $(\Phi_{-\varepsilon}')^{-1}(x) = \frac{e^{x}}{1-\varepsilon e^{x}}$, hence $M_{\Phi}^{\mathcal{F}}$ is in this case a Bose-Einstein distribution (when it exists).

Proof. We start by proving (i) \iff (ii). Let $f, g \in \mathcal{F}$ such that $g \in J$ μ -almost everywhere and $\mathcal{H}_{\Phi}[f|g] < \infty$. Using a Taylor expansion followed by Fubini's Theorem, we get

$$\begin{split} H_{\Phi}[f|g] &= \int_{\mathcal{E}} (\Phi(f) - \Phi(g)) \, \mathrm{d}\mu(\zeta) = \int_{g \in J} (\Phi(f) - \Phi(g)) \, \mathrm{d}\mu(\zeta) \\ &= \int_{g \in J} \left((f - g) \Phi'(g) + (f - g)^2 \int_0^1 (1 - \tau) \Phi''((1 - \tau)g + \tau f) \, \mathrm{d}\tau \right) \, \mathrm{d}\mu(\zeta) \\ &= \int_{g \in J} (f - g) \Phi'(g) \, \mathrm{d}\mu(\zeta) + \mathcal{H}_{\Phi}[f|g], \end{split}$$

thus proving the announced equivalence. Remark that in the last equality we used the fact that $\mathcal{H}_{\Phi}[f|g] < \infty$ to ensure that $\int_{g \in J} (f - g) \Phi'(g) d\mu(\zeta)$ is well-defined. We now show $(iv) \implies (ii)$. Since $\operatorname{Im}(\Phi'^{-1}) = J$, we

do have $M_{\Phi}^{\mathcal{F}} \in J \ \mu$ -almost everywhere, and

$$\int_{M_{\Phi}^{\mathcal{F}} \in J} (f - M_{\Phi}^{\mathcal{F}}) \Phi'(M_{\Phi}^{\mathcal{F}}) d\mu(\zeta) = \int_{\mathcal{E}} (f - M_{\Phi}^{\mathcal{F}}) \sum_{i} \alpha_{i}(\omega_{i}) \phi_{i} d\mu(\zeta)$$
$$= \sum_{i} \alpha_{i}(\omega_{i}) \left(\int_{\mathcal{E}} f \phi_{i} d\mu(\zeta) - \int_{\mathcal{E}} M_{\Phi}^{\mathcal{F}} \phi_{i} d\mu(\zeta) \right) = 0,$$

where the last equality comes from the fact that both f and $M_{\Phi}^{\mathcal{F}}$ belong to \mathcal{F} . We now focus on $(ii) \implies (iv)$. Assume the existence of $g \in \mathcal{F}$ such that $g \in J$ μ -almost everywhere and

$$\forall f \in \mathcal{F}, \quad \int_{g \in J} (f - g) \, \Phi'(g) \, \mathrm{d}\mu(\zeta) = 0.$$

Since we also have $M_{\Phi}^{\mathcal{F}} \in \mathcal{F}$, then $M_{\Phi}^{\mathcal{F}} \in J$ μ -almost everywhere and we just proved that

$$\int_{M_{\Phi}^{\mathcal{F}} \in J} (g - M_{\Phi}^{\mathcal{F}}) \, \Phi'(M_{\Phi}^{\mathcal{F}}) \, \mathrm{d}\mu(\zeta) = 0,$$

allowing to deduce that

$$\int_{\mathcal{E}} (g - M_{\Phi}^{\mathcal{F}})(\Phi'(g) - \Phi'(M_{\Phi}^{\mathcal{F}})) \,\mathrm{d}\mu(\zeta) = 0$$

Since Φ' is increasing, this implies that $g = M_{\Phi}^{\mathcal{F}} \mu$ -almost everywhere. We now focus on $(iv) \implies (iii)$. Since we already proved $(iv) \implies (ii) \implies (i)$, we have for any $f \in \mathcal{F}$,

$$H_{\Phi}[f|M_{\Phi}^{\mathcal{F}}] = \mathcal{H}_{\Phi}[f|M_{\Phi}^{\mathcal{F}}] \ge 0,$$

thus

$$H_{\Phi}(f) \ge H_{\Phi}(M_{\Phi}^{\mathcal{F}}).$$

Finally, we prove $(iii) \implies (iv)$. Assume $H_{\Phi}(g) = \min_{h \in \mathcal{F}} H_{\Phi}(h)$. We just proved that $H_{\Phi}(M_{\Phi}^{\mathcal{F}}) = \min_{h \in \mathcal{F}} H_{\Phi}(h)$, hence $H_{\Phi}(g) = H_{\Phi}(M_{\Phi}^{\mathcal{F}})$ and $H_{\Phi}[g|M_{\Phi}^{\mathcal{F}}] = 0$. Since we also proved $(iv) \implies (i)$, we know that

$$\mathcal{H}_{\Phi}[g|M_{\Phi}^{\mathcal{F}}] = H_{\Phi}[g|M_{\Phi}^{\mathcal{F}}]$$

hence $\mathcal{H}_{\Phi}[g|M_{\Phi}^{\mathcal{F}}] = 0$, that is

$$\int_0^1 (1-\tau) \left(\int_{M_\Phi^{\mathcal{F}} \in J} (g - M_\Phi^{\mathcal{F}})^2 \Phi''((1-\tau)M + \tau g) \,\mathrm{d}\mu(\zeta) \right) \,\mathrm{d}\tau = 0$$

This implies, since $M_{\Phi}^{\mathcal{F}} \in J$ and $(1-\tau)M_{\Phi}^{\mathcal{F}} + \tau g \in J$ μ -almost everywhere for any $\tau \in (0, 1)$, and $\Phi'' > 0$ on J, that $g = M_{\Phi}^{\mathcal{F}} \mu$ almost everywhere.

General Csiszár-Kullback-Pinsker inequality. The famous Csiszár-Kullback-Pinsker inequality, linking the squared L^1 distance of two probabilities with their relative Classical entropy can in fact be generalized to the whole family of Φ -entropies, and for weighted L^r , $1 \le r \le 2$ distances, by the following Proposition 16, and more specifically, in the context of convergence towards equilibrium, by Corollary 17. Again, such inequalities may be useful in the study of weak turbulence [7].

Proposition 16. Let some $(\mathcal{A}, \operatorname{Bor}(\bar{J}))$ -measurable $\varpi : \mathcal{E} \to \bar{J}$ and $r \in [1, 2]$. Then for any $(\mathcal{A}, \operatorname{Bor}(\bar{J}))$ -measurable $f, g : \mathcal{E} \to \bar{J}$ such that all terms below are finite, we have

(112)
$$\|(f-g)\varpi\|_{L^{r}(g\in J)}^{2} \leq \left(\int_{0}^{1}(1-\tau)\left\|\frac{\varpi^{2}}{\Phi''((1-\tau)g+\tau f)}\right\|_{L^{\frac{r}{2-r}}(g\in J)}^{-1}\mathrm{d}\tau\right)^{-1}\mathcal{H}_{\Phi}[f|g],$$

where \mathcal{H}_{Φ} is the Φ -relative-entropy defined in (110).

Proof. Let us recall the definition of $\mathcal{H}_{\Phi}[f|g]$, that is

$$\mathcal{H}_{\Phi}[f|g] = \int_{0}^{1} (1-\tau) \left(\int_{g \in J} (f-g)^2 \Phi''((1-\tau)g + \tau f) \,\mathrm{d}\mu(\zeta) \right) \,\mathrm{d}\tau$$

We fix $\tau \in (0,1)$. Let $p = \frac{2}{r} \in [1,2]$ and $q = \frac{2}{2-r} \in [2,+\infty]$, so that $\frac{1}{p} + \frac{1}{q} = 1$. By Hölder's inequality, we have

$$\begin{split} \int_{g\in J} |f-g|^{\frac{2}{p}} \, \varpi^{\frac{2}{p}} \, \mathrm{d}\mu(\zeta) &= \int_{g\in J} |f-g|^{\frac{2}{p}} \Phi''((1-\tau)g+\tau f)^{\frac{1}{p}} \left[\frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)} \right]^{\frac{1}{p}} \, \mathrm{d}\mu(\zeta) \\ &\leq \left(\int_{g\in J} (f-g)^2 \, \Phi''((1-\tau)g+\tau f) \, \mathrm{d}\mu(\zeta) \right)^{\frac{1}{p}} \, \left\| \left[\frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)} \right]^{\frac{1}{p}} \, \right\|_{L^q(g\in J)}. \end{split}$$

Raising the above inequality to the power p, we obtain (113)

$$\left(\int_{g\in J} |f-g|^{\frac{2}{p}} \,\varpi^{\frac{2}{p}} \,\mathrm{d}\mu(\zeta)\right)^{p} \leq \left(\int_{g\in J} (f-g)^{2} \,\Phi''((1-\tau)g+\tau f) \,\mathrm{d}\mu(\zeta)\right) \,\left\|\frac{\varpi^{2}}{\Phi''((1-\tau)g+\tau f)}\right\|_{L^{\frac{q}{p}}(g\in J)},$$

where we used the fact that $\|\cdot^{\frac{1}{p}}\|_{L^q}^p = \|\cdot\|_{L^{\frac{q}{p}}}$. Since $\frac{2}{p} = r$ and $\frac{q}{p} = \frac{r}{2-r}$, (113) actually writes

$$\|(f-g)\,\varpi\|_{L^r(g\in J)}^2 \le \left(\int_{g\in J} (f-g)^2 \,\Phi''((1-\tau)g+\tau f)\,\mathrm{d}\mu(\zeta)\right) \,\left\|\frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)}\right\|_{L^{\frac{r}{2-r}}(g\in J)}.$$

If ϖ is zero μ -almost everywhere on $\{g \in J\}$, then the proposition is trivial. Else, since $\Phi'' > 0$ on $\{g \in J\}$ and $(1 - \tau)g + \tau f \in J$ on $\{g \in J\}$ for any $\tau \in (0, 1)$, we know that $\left\|\Phi''((1 - \tau)g + \tau f) \,\varpi^2\right\|_{L^{\frac{r}{2-r}}(g \in J)} > 0$. Since we assumed that the quantity

$$\left(\int_{0}^{1} (1-\tau) \left\| \frac{\varpi^{2}}{\Phi''((1-\tau)g+\tau f)} \right\|_{L^{\frac{r}{2-r}}(g\in J)}^{-1} \mathrm{d}\tau \right)^{-1}$$

is finite, we also know that for almost every $\tau \in (0,1)$ we have $\left\|\frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)}\right\|_{L^{\frac{r}{2-r}}(g\in J)} < \infty$. For these values of τ , we then have

$$\|(f-g)\,\varpi\|_{L^r(g\in J)}^2 \,\frac{1-\tau}{\left\|\frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)}\right\|_{L^{\frac{r}{2-r}}(g\in J)}} \leq (1-\tau)\left(\int_{g\in J} (f-g)^2 \,\Phi''((1-\tau)g+\tau f)\,\mathrm{d}\mu(\zeta)\right).$$

Integrating in τ yields

$$\left\| (f-g)\,\varpi \right\|_{L^r(g\in J)}^2 \left(\int_0^1 (1-\tau) \, \left\| \frac{\varpi^2}{\Phi''((1-\tau)g+\tau f)} \right\|_{L^{\frac{r}{2-r}}(g\in J)}^{-1} \, \mathrm{d}\tau \right) \le \mathcal{H}_{\Phi}[f|g].$$
 Since, again by hypothesis, the integral in τ is nonzero (its inverse is finite), we obtain (112).

Remarking that $\{M_{\Phi}^{\mathcal{F}} \in J\} = \mathcal{E}$ and that for any $f \in \mathcal{F}$, $\mathcal{H}_{\Phi}[f|M_{\Phi}^{\mathcal{F}}] = H_{\Phi}[f|M_{\Phi}^{\mathcal{F}}]$ (see Proposition 15), we straightforwardly obtain the following corollary.

Corollary 17. Let some $(\mathcal{A}, \operatorname{Bor}(\overline{J}))$ -measurable $\varpi : \mathcal{E} \to \overline{J}$ and $r \in [1, 2]$. With the same notations as in Proposition 15, assuming $M_{\Phi}^{\mathcal{F}}$ exists, we have for any $f \in \mathcal{F}$ such that the integral term below is finite,

(114)
$$\left\| (f - M_{\Phi}^{\mathcal{F}}) \varpi \right\|_{L^{r}}^{2} \leq \left(\int_{0}^{1} (1 - \tau) \left\| \frac{\varpi^{2}}{\Phi''((1 - \tau)M_{\Phi}^{\mathcal{F}} + \tau f)} \right\|_{L^{\frac{r}{2 - r}}}^{-1} \mathrm{d}\tau \right)^{-1} H_{\Phi}[f|M_{\Phi}^{\mathcal{F}}],$$

where H_{Φ} is defined in (108)–(109) and $M_{\Phi}^{\mathcal{F}}$ is the equilibrium associated to Φ and the set \mathcal{F} , defined in (111).

Appendix C. Technical results

In this section, we consider $\mathcal{M}^{f}_{\varepsilon}$, the Fermi-Dirac distribution associated to some $\varepsilon > 0$ and $0 \le f \in L^{1}_{2}(\mathbb{R}^{3})$ such that $1 - \varepsilon f \ge 0$. The existence of $\mathcal{M}^{f}_{\varepsilon}$ is provided by assuming $\gamma > \frac{2}{5}$ (see [22]), where we recall the notation

(115)
$$\gamma := \frac{T}{T_F(\rho,\varepsilon)}$$

where $T_F(\rho,\varepsilon) = \frac{1}{2} \left(\frac{3\rho\varepsilon}{4\pi}\right)^{2/3}$ is the Fermi temperature associated to ρ and ε ; and ρ, T are respectively the density and temperature associated to the distribution f, defined in (12). We also recall the notation, for $x \in [0, \varepsilon^{-1})$,

$$\varphi_{\varepsilon}(x) = \frac{x}{1 - \varepsilon x}.$$

C.1. L^{∞} bound for the Fermi-Dirac statistics. In this subsection, we provide an L^{∞} bound on the Fermi-Dirac statistics. The following result is very similar to [3, Lemma A.1].

Proposition 18. Let $\varepsilon > 0$ and $f \in L_2^1(\mathbb{R}^3)$ be a nonnegative distribution such that $1 - \varepsilon f \ge 0$ and $\gamma \ge \gamma^{\dagger}$, where γ is given by (115) and

(116)
$$\gamma^{\dagger} := \left(\frac{4}{\pi}\right)^{\frac{1}{3}} \left(\frac{5}{3}\right)^{\frac{3}{3}}.$$

Then the quantity $\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}^{f}_{\varepsilon})\|_{\infty}$ satisfies

(117)
$$\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})\|_{\infty} \leq \frac{2}{3} \left(\frac{\gamma}{\gamma^{\dagger}}\right)^{-3/2}$$

Proof. Our proof is based on [22, proof of Proposition 3]. We introduce, for $s \ge 0$,

$$I_s(t) := \int_0^\infty \frac{r^s}{1 + te^{r^2}} \,\mathrm{d}r, \qquad P(t) := I_4(t)[I_2(t)]^{-5/3}, \quad t > 0$$

It is proven in [22, proof of Proposition 3] that P is continuous and increasing on \mathbb{R}_+ , and that

$$P\left(\frac{1}{\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})\|_{\infty}}\right) = 3\rho^{-2/3} T (4\pi\varepsilon^{-1})^{2/3} \equiv \frac{3^{5/3}}{2} \gamma.$$

Let $\alpha > 0$ and

$$t = \alpha^{-1} \times \frac{3\sqrt{\pi}}{4} \gamma^{3/2}.$$

As, for any $r \ge 0$, it holds that

$$\frac{e^{-r^2}}{1+t} \le \frac{1}{1+t \, e^{r^2}} \le \frac{e^{-r^2}}{t},$$

we have

$$P(t) \le \left(\frac{1}{t} \int_0^\infty r^4 \, e^{-r^2} \, \mathrm{d}r\right) \left(\frac{1}{1+t} \int_0^\infty r^2 \, e^{-r^2} \, \mathrm{d}r\right)^{-5/3} = \frac{(1+t)^{5/3}}{t} \left(\frac{1}{2}\Gamma\left(\frac{5}{2}\right)\right) \left(\frac{1}{2}\Gamma\left(\frac{3}{2}\right)\right)^{-5/3},$$

t is

that is

$$P(t) \le \frac{(1+t)^{5/3}}{t} \times \frac{3 \times 2^{1/3}}{\pi^{1/3}}.$$

We define

$$\gamma^{\alpha} := \left(\frac{4}{3\sqrt{\pi}} \times \frac{\alpha}{\alpha^{2/5} - 1}\right)^{2/3}$$

Then, whenever $\gamma \ge \gamma^{\alpha}$, we have $t \ge \frac{1}{\alpha^{2/5} - 1}$, so that $1 + t \le \alpha^{2/5} t$, implying

$$P(t) \le t^{2/3} \times \frac{3 \times 2^{1/3} \alpha^{2/3}}{\pi^{1/3}} = \alpha^{-2/3} \times \left(\frac{3\sqrt{\pi}}{4}\right)^{2/3} \gamma \times \frac{3 \times 2^{1/3} \alpha^{2/3}}{\pi^{1/3}} = \frac{3^{5/3}}{2} \gamma = P\left(\frac{1}{\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})\|_{\infty}}\right)$$

Since P is increasing, we deduce that, whenever $\gamma \geq \gamma^{\alpha}$, we have

$$\alpha^{-1} \times \frac{3\sqrt{\pi}}{4} \gamma^{3/2} = t \le \frac{1}{\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})\|_{\infty}}$$

that is

(118)
$$\varepsilon \|\varphi_{\varepsilon}(\mathcal{M}_{\varepsilon}^{f})\|_{\infty} \leq \alpha \times \frac{4}{3\sqrt{\pi}} \gamma^{-3/2}.$$

By computing the derivative of $\alpha \mapsto \frac{\alpha}{\alpha^{2/5} - 1}$, we can minimize $\alpha \mapsto \gamma^{\alpha}$ and find that the minimum value is

$$\gamma^{\dagger} = \left(\frac{4}{\pi}\right)^{\frac{1}{3}} \left(\frac{5}{3}\right)^{\frac{5}{3}},$$

reached for $\alpha^{\dagger} = \left(\frac{5}{3}\right)^{5/2}$, and, combined with (118), this proves (117).

C.2. Regularity in ε of the coefficients of the Fermi-Dirac statistics. In this last subsection, for any $\varepsilon \geq 0$ and $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$, we denote by $\mathcal{M}_{\varepsilon} \equiv \mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g)}$ the ε -Fermi distribution associated to $\varphi_{\varepsilon}^{-1}(g)$. In particular, in the limit case $\varepsilon = 0$, \mathcal{M}_0 is the Maxwellian distribution associated to g. We also denote $a_{\varepsilon}, b_{\varepsilon}, \bar{u}_{\varepsilon}$ and $\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon}$ the quantities such that, letting

(119)
$$\mathcal{M}_{\varepsilon} \equiv \mathcal{M}_{\varepsilon}^{\frac{g}{1+\varepsilon g}}, \qquad M_{\varepsilon} = \frac{\mathcal{M}_{\varepsilon}}{1-\varepsilon \mathcal{M}_{\varepsilon}}$$

we have, for any $v \in \mathbb{R}^3$,

(120)
$$M_{\varepsilon}(v) = \exp\left(a_{\varepsilon} + b_{\varepsilon}|v - \bar{u}_{\varepsilon}|^2\right)$$

and

(121)
$$\int_{\mathbb{R}^3} \frac{g}{1+\varepsilon g} \begin{pmatrix} 1\\v\\|v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho_{\varepsilon}\\\rho_{\varepsilon} u_{\varepsilon}\\3\rho_{\varepsilon} T_{\varepsilon} + \rho_{\varepsilon} |u_{\varepsilon}|^2 \end{pmatrix}.$$

Lemma 19. Let $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$. Using the notation (121), the application $\varepsilon \mapsto (\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon})$ is continuous on \mathbb{R}_+ and \mathcal{C}^1 on \mathbb{R}^*_+ .

Proof. The continuity of $\varepsilon \mapsto (\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon})$ on \mathbb{R}_+ comes by dominated convergence, as it holds for any $\varepsilon \ge 0$ and $v \in \mathbb{R}^3$ that

$$\frac{g}{1+\varepsilon g} \, (1+|v|^2) \le g \, (1+|v|^2),$$

and by hypothesis $0 \leq g \in L_2^1(\mathbb{R}^3)$. Similarly, for any $\varepsilon > 0$ and $v \in \mathbb{R}^3$ we have

$$\left|\partial_{\varepsilon}\frac{g}{1+\varepsilon g}\right| (1+|v|^2) = \left(\frac{g}{1+\varepsilon g}\right)^2 (1+|v|^2) \le \varepsilon^{-1} g (1+|v|^2).$$

Therefore, for any $\varepsilon > 0$, we have

$$\sup_{* \in (\varepsilon/2, 3\varepsilon/2)} \left| \partial_{\varepsilon_*} \frac{g}{1 + \varepsilon_* g} \right| (1 + |v|^2) \le 2\varepsilon^{-1} g (1 + |v|^2).$$

The differentiability of $\varepsilon \mapsto (\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon})$ on \mathbb{R}^*_+ then comes by dominated convergence, as $g \in L^1_2(\mathbb{R}^3)$. \Box

The following lemmas provide the continuity of $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon}, \bar{u}_{\varepsilon})$ at the point $\varepsilon = 0$ and its differentiability on \mathbb{R}^*_+ .

Lemma 20. Let $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$. Using the notations (119)–(120), the application $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon}, \bar{u}_{\varepsilon})$ is continuous at the point $\varepsilon = 0$.

Proof. Using the notations (119) and (121), we have

(122)
$$\left| \int_{\mathbb{R}^3} M_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv - \int_{\mathbb{R}^3} \mathcal{M}_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv \right| = \left| \int_{\mathbb{R}^3} \frac{\varepsilon M_{\varepsilon}^2}{1 + \varepsilon M_{\varepsilon}} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv \right| \le \varepsilon \|M_{\varepsilon}\|_{\infty} \int_{\mathbb{R}^3} \mathcal{M}_{\varepsilon} \begin{pmatrix} 1 \\ |v| \\ |v|^2 \end{pmatrix} dv$$
$$\le \varepsilon \|M_{\varepsilon}\|_{\infty} (\rho_{\varepsilon} + 3\rho_{\varepsilon}T_{\varepsilon} + \rho_{\varepsilon}|u_{\varepsilon}|^2).$$

Recall the notation, in this case,

$$\gamma_{\varepsilon} = \frac{T_{\varepsilon}}{T_F(\rho_{\varepsilon}, \varepsilon)},$$

where $T_F(\rho_{\varepsilon},\varepsilon) = \frac{1}{2} \left(\frac{3\rho_{\varepsilon}\varepsilon}{4\pi}\right)^{2/3}$. By continuity at the point $\varepsilon = 0$ of the application $\varepsilon \mapsto (\rho_{\varepsilon}, T_{\varepsilon})$, given by Lemma 19, we have

$$\gamma_{\varepsilon} \xrightarrow{\varepsilon \to 0} +\infty.$$

Thereby, there exists $\varepsilon^* > 0$ such that $\gamma_{\varepsilon} \ge \gamma^{\dagger}$ for any $\varepsilon \in (0, \varepsilon^*)$, where γ^{\dagger} is a universal constant defined in Proposition 18. Then, from (117) in Proposition 18, for any $\varepsilon \in (0, \varepsilon^*)$,

$$\varepsilon \|M_{\varepsilon}\|_{\infty} \leq \frac{2}{3} \left(\frac{\gamma_{\varepsilon}}{\gamma_{*}}\right)^{-\frac{2}{2}}$$

which vanishes as $\varepsilon \to 0$, since γ_{ε} tends to $+\infty$ in this limit. Combining this result, the continuity of $\varepsilon \mapsto (\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon})$ at $\varepsilon = 0$, given by Lemma 19, and Equation (122), we obtain

$$\left| \int_{\mathbb{R}^3} M_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v - \int_{\mathbb{R}^3} \mathcal{M}_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v \right| \xrightarrow{\varepsilon \to 0} 0.$$

The continuity of $\varepsilon \mapsto (\rho_{\varepsilon}, u_{\varepsilon}, T_{\varepsilon})$ at $\varepsilon = 0$ being equivalent to the statement

$$\int_{\mathbb{R}^3} \mathcal{M}_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v \xrightarrow{\varepsilon \to 0} \int_{\mathbb{R}^3} \mathcal{M}_0 \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} \mathrm{d}v,$$

we finally conclude that

$$\int_{\mathbb{R}^3} M_{\varepsilon} \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv \xrightarrow{\varepsilon \to 0} \int_{\mathbb{R}^3} \mathcal{M}_0 \begin{pmatrix} 1 \\ v \\ |v|^2 \end{pmatrix} dv.$$

Both M_{ε} and \mathcal{M}_0 are gaussian distributions, which coefficients are continuously defined by the above moments, allowing to conclude to the continuity of these coefficients at the point $\varepsilon = 0$.

Lemma 21. Let $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$. Using the notations (119)–(120), the application $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon}, \bar{u}_{\varepsilon})$ is \mathcal{C}^1 on \mathbb{R}^*_+ .

Proof. Since the distribution $M_{\varepsilon}(\cdot + \bar{u}_{\varepsilon})$ is radially symmetric, so is $\mathcal{M}_{\varepsilon}(\cdot + \bar{u}_{\varepsilon})$, hence $\bar{u}_{\varepsilon} = u_{\varepsilon}$, which, by Lemma 19, is \mathcal{C}^1 on \mathbb{R}^*_+ . We then define

$$g_{\varepsilon} := g(\cdot + u_{\varepsilon}), \qquad \mathcal{N}_{\varepsilon} := \mathcal{M}_{\varepsilon}(\cdot + u_{\varepsilon}) \equiv \mathcal{M}_{\varepsilon}^{\varphi_{\varepsilon}^{-1}(g_{\varepsilon})} \text{ and } N_{\varepsilon} := M_{\varepsilon}(\cdot + u_{\varepsilon}),$$

so that, for any $\varepsilon > 0$, it holds that

$$N_{\varepsilon} = \exp\left(a_{\varepsilon} + b_{\varepsilon}|v|^2\right).$$

As in (121), we let $\rho_{\varepsilon}, T_{\varepsilon} > 0$ be such that

$$\int_{\mathbb{R}^3} \frac{g_{\varepsilon}}{1+\varepsilon g_{\varepsilon}} \begin{pmatrix} 1\\ v\\ |v|^2 \end{pmatrix} \mathrm{d}v = \begin{pmatrix} \rho_{\varepsilon}\\ 0\\ 3\rho_{\varepsilon}T_{\varepsilon} \end{pmatrix}.$$

Let us now show that $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon})$ is \mathcal{C}^1 on \mathbb{R}^*_+ . It is proven in [22, proof of Proposition 3] that, letting

$$I_s(\tau) := \int_0^\infty \frac{r^s}{1 + \tau \, e^{r^2}} \, \mathrm{d}r, \qquad P(\tau) := I_4(\tau) \, I_2(\tau)^{-5/3}, \qquad \tau \in \mathbb{R}^*_+,$$

the function P is an increasing C^1 function from \mathbb{R}^*_+ to $\left(\frac{3^{5/3}}{5}, +\infty\right)$, with P' > 0 on \mathbb{R}^*_+ . Therefore it is invertible, and P^{-1} is also C^1 . All the more, a dominated convergence argument ensures that I_2 is C^1 on \mathbb{R}^*_+ . It is moreover shown in [22, proof of Proposition 3] that

$$\left(\frac{\varepsilon}{4\pi}\right)^{2/3} P\left(\frac{1}{\varepsilon e^{a_{\varepsilon}}}\right) = \frac{3\rho_{\varepsilon}T_{\varepsilon}}{\rho_{\varepsilon}^{5/3}}, \qquad b_{\varepsilon} = -\left(\frac{4\pi}{\varepsilon\rho_{\varepsilon}}I_2\left(\frac{1}{\varepsilon e^{a_{\varepsilon}}}\right)\right)^{\frac{2}{3}}.$$

By Lemma 19, the application $\varepsilon \mapsto (\rho_{\varepsilon}, T_{\varepsilon})$ is \mathcal{C}^1 on \mathbb{R}^*_+ , hence so is the application $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon})$, as a composition of \mathcal{C}^1 applications.

Lemma 22. Let $0 \leq g \in L_2^1(\mathbb{R}^3) \cap L \log L(\mathbb{R}^3)$. Using the notation (119), for any $\overline{\varepsilon} > 0$, there exist C > 0and $\eta > 0$ such that for any $\varepsilon \in [0, \overline{\varepsilon}]$ and $v \in \mathbb{R}^3$, we have

(123)
$$M_{\varepsilon}(v) \le C e^{-\eta |v|^2}.$$

and

(124)
$$|\log M_{\varepsilon}(v)| \le C(1+|v|^2).$$

Proof. We denote

$$\underline{a} = \sup_{\varepsilon \in [0,\overline{\varepsilon}]} |a_{\varepsilon}|, \qquad \underline{b} = -\sup_{\varepsilon \in [0,\overline{\varepsilon}]} |b_{\varepsilon}|, \qquad \overline{b} = -\inf_{\varepsilon \in [0,\overline{\varepsilon}]} |b_{\varepsilon}| \quad \text{and} \quad \underline{u} = \sup_{\varepsilon \in [0,\overline{\varepsilon}]} |u_{\varepsilon}|$$

Combining the results of Lemmas 20 and 21, the application $\varepsilon \mapsto (a_{\varepsilon}, b_{\varepsilon}, u_{\varepsilon})$ is continuous on \mathbb{R}_+ , from which we deduce that $\underline{a}, \underline{b}$ and \underline{u} are finite.

Moreover, as $M_{\varepsilon} \in L^1(\mathbb{R}^3)$ for all $\varepsilon \in [0, \overline{\varepsilon}]$, the application $\varepsilon \mapsto b_{\varepsilon}$ is (strictly) negative on $[0, \overline{\varepsilon}]$, so that, as $\varepsilon \mapsto b_{\varepsilon}$ is continuous on $[0, \overline{\varepsilon}]$, we have $\overline{b} < 0$.

Therefore, for any $0 \leq \varepsilon \leq \overline{\varepsilon}$ and $v \in \mathbb{R}^3$, we have

$$|\log M_{\varepsilon}(v)| = |a_{\varepsilon} + b_{\varepsilon}|v - u_{\varepsilon}|^{2}| \le \underline{a} + 2|\underline{b}|\,\underline{u}^{2} + 2|\underline{b}|\,|v|^{2},$$

and, since $|v - u_{\varepsilon}|^2 \ge \frac{1}{2}|v|^2 - |u_{\varepsilon}|^2 \ge \frac{1}{2}|v|^2 - \underline{u}^2$ and $\overline{b} < 0$,

$$M_{\varepsilon}(v) = e^{a_{\varepsilon} + b_{\varepsilon}|v - u_{\varepsilon}|^2} \le e^{\underline{a} + \overline{b}|v - u_{\varepsilon}|^2} \le e^{\underline{a} + |\overline{b}|\underline{u}^2 + \frac{1}{2}\overline{b}|v|^2}$$

Letting $\eta = -\frac{1}{2}\overline{b}$ and $C = \max\left(\underline{a} + 2|\underline{b}|\underline{u}^2, e^{\underline{a} + |\overline{b}|\underline{u}^2}\right)$ yields the result.

Acknowledgements

I thank Bertrand Lods with whom I had fruitful discussions and Matthieu Dolbeault who suggested to me the proof of the nice inequality (92).

I have no competing interests to declare that are relevant to the content of this article.

References

- R. Alonso, V. Bagland, L. Desvillettes, and B. Lods. About the use of entropy production for the Landau-Fermi-Dirac equation. J. Stat. Phys., 183(1):Paper No. 10, 27, 2021.
- [2] R. Alonso, V. Bagland, L. Desvillettes, and B. Lods. About the Landau-Fermi-Dirac equation with moderately soft potentials. Arch. Ration. Mech. Anal., 244(3):779–875, 2022.
- [3] R. Alonso, V. Bagland, and B. Lods. Long time dynamics for the Landau-Fermi-Dirac equation with hard potentials. J. Differential Equations, 270:596–663, 2021.
- [4] R. Alonso, V. Bagland, and B. Lods. Long time dynamics for the landau-fermi-dirac equation with hard potentials. Journal of Differential Equations, 270:596–663, 2021.
- [5] A. V. Bobylev and C. Cercignani. On the rate of entropy production for the Boltzmann equation. J. Statist. Phys., 94(3-4):603-618, 1999.
- [6] F. Bolley and C. Villani. Weighted Csiszár-Kullback-Pinsker inequalities and applications to transportation inequalities. Ann. Fac. Sci. Toulouse Math. (6), 14(3):331–352, 2005.
- [7] M. Breden and L. Desvillettes. Rigorous study of the equilibria of collision kernels appearing in the theory of weak turbulence. Arch. Ration. Mech. Anal., 235(2):1151–1176, 2020.
- [8] M. J. Cáceres, J. A. Carrillo, and J. Dolbeault. Nonlinear stability in L^p for a confined system of charged particles. SIAM J. Math. Anal., 34(2):478–494, 2002.
- [9] María J. Cáceres, José A. Carrillo, and Jean Dolbeault. A generalization of the Csiszár-Kullback inequality. https://www.ceremade.dauphine.fr/~dolbeaul/Teaching/files/SNS-2012/Csiszar-Kullback.pdf.
- [10] K. Carrapatoso, L. Desvillettes, and L. He. Estimates for the large time behavior of the Landau equation in the Coulomb case. Arch. Ration. Mech. Anal., 224(2):381–420, 2017.
- [11] C. Cercignani. H-theorem and trend to equilibrium in the kinetic theory of gases. Arch. Mech. (Arch. Mech. Stos.), 34(3):231–241 (1983), 1982.
- [12] S. Chapman and T. G. Cowling. The Mathematical Theory of Non-uniform Gases. Cambridge University Press, Cambridge, 1939.
- [13] L. Desvillettes. Entropy dissipation estimates for the Landau equation in the Coulomb case and applications. J. Funct. Anal., 269(5):1359–1403, 2015.
- [14] L. Desvillettes. Structure entropique du noyau de collision de Landau. In Séminaire Laurent Schwartz-Équations aux dérivées partielles et applications. Année 2014–2015, pages Exp. No. XIV, 9. Ed. Éc. Polytech., Palaiseau, 2016.

- [15] L. Desvillettes. Some remarks about the link between the Fisher information and Landau or Landau-Fermi-Dirac entropy dissipation. preprint. ArXiv: 2401.02877, 2024.
- [16] L. Desvillettes, C. Mouhot, and C. Villani. Celebrating Cercignani's conjecture for the Boltzmann equation. *Kinet. Relat. Models*, 4(1):277–294, 2011.
- [17] L. Desvillettes and C. Villani. On the spatially homogeneous Landau equation for hard potentials. II. H-theorem and applications. Comm. Partial Differential Equations, 25(1-2):261–298, 2000.
- [18] J. Dolbeault. Kinetic models and quantum effects: a modified Boltzmann equation for Fermi-Dirac particles. Arch. Rational Mech. Anal., 127(2):101–131, 1994.
- [19] G. L. Gilardoni. On Pinsker's and Vajda's type inequalities for Csiszár's f-divergences. IEEE Trans. Inform. Theory, 56(11):5377–5386, 2010.
- [20] A. Jüngel. Entropy methods for diffusive partial differential equations, volume 804. Springer, 2016.
- [21] X. Lu. A modified Boltzmann equation for Bose-Einstein particles: isotropic solutions and long-time behavior. J. Statist. Phys., 98(5-6):1335–1394, 2000.
- [22] X. Lu. On spatially homogeneous solutions of a modified Boltzmann equation for Fermi-Dirac particles. J. Statist. Phys., 105(1-2):353–388, 2001.
- [23] X. Lu. On isotropic distributional solutions to the Boltzmann equation for Bose-Einstein particles. J. Statist. Phys., 116(5):1597–1649, 2004.
- [24] X. Lu and B. Wennberg. On stability and strong convergence for the spatially homogeneous Boltzmann equation for Fermi-Dirac particles. Arch. Ration. Mech. Anal., 168(1):1–34, 2003.
- [25] L. W. Nordhiem. On the kinetic method in the new statistics and application in the electron theory of conductivity. Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character, 119(783):689–698, 1928.
- [26] G. Toscani and C. Villani. Sharp entropy dissipation bounds and explicit rate of trend to equilibrium for the spatially homogeneous Boltzmann equation. Comm. Math. Phys., 203(3):667–706, 1999.
- [27] G. Toscani and C. Villani. On the trend to equilibrium for some dissipative systems with slowly increasing a priori bounds. J. Statist. Phys., 98(5-6):1279–1309, 2000.
- [28] E. A. Uehling and G. E. Uhlenbeck. Transport phenomena in Einstein-Bose and Fermi-Dirac gases. i. Physical Review, 43(7):552, 1933.
- [29] C. Villani. A review of mathematical topics in collisional kinetic theory. In Handbook of mathematical fluid dynamics, Vol. I, pages 71–305. North-Holland, Amsterdam, 2002.
- [30] C. Villani. Cercignani's conjecture is sometimes true and always almost true. Comm. Math. Phys., 234(3):455-490, 2003.