

# Hydrodynamics in long-range interacting systems with center-of-mass conservation

Alan Morningstar <sup>1</sup>, Nicholas O’Dea,<sup>1</sup> and Jonas Richter <sup>1,2</sup>

<sup>1</sup>*Department of Physics, Stanford University, Stanford, CA 94305, USA*

<sup>2</sup>*Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

(Dated: April 26, 2023)

In systems with a conserved density, the additional conservation of the center of mass (dipole moment) has been shown to slow down the associated hydrodynamics. At the same time, long-range interactions generally lead to faster transport and information propagation. Here, we explore the competition of these two effects and develop a hydrodynamic theory for long-range center-of-mass-conserving systems. We demonstrate that these systems can exhibit a rich dynamical phase diagram containing subdiffusive, diffusive, and superdiffusive behaviors, with continuously varying dynamical exponents. We corroborate our theory by studying quantum lattice models whose emergent hydrodynamics exhibit these phenomena.

Hydrodynamic theories are coarse-grained descriptions of the flow of conserved densities. They help us understand large-scale behavior and universality without knowing how these emerge from microscopic details [1, 2]. The price for this is the introduction of unknown macroscopic parameters such as diffusivity or viscosity, which are difficult to compute from first principles [3–8]. Understanding how hydrodynamics emerges in quantum systems is of great interest, and recent progress has been made in this direction [9–13]. It is also of interest to understand what kinds of hydrodynamic behavior can possibly arise in quantum many-body systems, including both chaotic and integrable models [14–30], as well as models with symmetries or kinetic constraints [31–37].

For highly-excited quantum chaotic lattice models, the default expectation is often that standard diffusion will emerge. However, the dynamics can be systematically slowed down by conserving higher moments of the density distribution. A common example of this is the subdiffusion that results from conserving the center of mass (COM), also known as the dipole moment [38–47]. This can occur in “tilted” systems with a strong linear potential [48–51], as realized in cold-atom setups [52–54], and is relevant also for quantum Hall systems [55–58]. The additional conservation law can not only modify the hydrodynamics but can result in a frozen phase where the state space is fragmented into dynamically disconnected sectors [59–68]. While we focus on the highly-excited dynamics of such systems, there has also been interest in their low-temperature equilibrium properties [69–81].

In contrast to constrained dynamics, faster dynamics can be achieved in systems with long-range interactions [82, 83]. Interactions that decay as a power law of the distance between particles—including Coulomb, dipolar, and van der Waals—are ubiquitous, and can nowadays be explored using, e.g., ultracold atoms [84–87], polar molecules [88–90], or trapped ions [91]. In such systems, Lieb-Robinson bounds can become superballistic, and hydrodynamics superdiffusive, depending on the power-law exponent and the dimensionality of the




FIG. 1. **(a)** We consider models involving long-range interactions and COM conservation. The dynamics are governed by pairs of equal and opposite currents (or the underlying “pair hopping” processes depicted) acting on lattice sites separated by the distances  $s \geq 0$  and  $r \geq 1$ . The strengths of the currents are suppressed by power laws  $(s + 1)^{-\alpha} r^{-\beta}$ . **[(b), (c)]** Schematic cuts through the dynamical phase diagram for the limiting cases  $r \rightarrow 1$  ( $\beta \rightarrow \infty$ ) and  $s \rightarrow 0$  ( $\alpha \rightarrow \infty$ ).

underlying lattice [92–102].

Given their competing effects, it is natural to examine the interplay between higher-order conservation laws and long-range interactions, however, this problem has only been briefly touched on in the literature [39]. Here, we bridge the gap and consider a class of models where “hops” are long-range but always occur in pairs such that the COM is conserved [Fig. 1(a)]. We develop a hydrodynamic theory of such models and map out its phase diagram [Fig. 1(b),(c)]. More specifically, our model is characterized by two power-law exponents,  $\alpha$  and  $\beta$ , that control the suppression of pair-hopping amplitudes. We calculate the dynamical exponent  $z$ , governing long-wavelength relaxation, as a function of these two exponents, as well as the scaling functions for density-density correlations. When  $\alpha$  and  $\beta$  are large enough, we recover the known  $z = 4$  subdiffusion of short-range COM-conserving systems in one dimension. When  $\alpha$  or  $\beta$  is

small enough,  $z$  continuously varies. The scaling function of the correlations also continuously varies with the exponents. We find consistent results when examining a less-tractable quantum spin-1 model (a long-range version of the systems studied in Ref. [61, 62]). We also discuss the relevance of our findings to long-range many-body quantum systems in a tilted potential.

**Setup.** As a starting point, we consider a 1D quantum spin model

$$H = \sum_x \sum_{s,r} J_{sr} (S_x^+ S_{x+r}^- S_{x+r+s}^- S_{x+s+2r}^+ + \text{h.c.}), \quad (1)$$

where  $S_x^\pm$  are raising and lowering operators at site  $x$ , and  $J_{sr}$  depend on the distances  $s \geq 0$  and  $r > 0$ . The Hamiltonian conserves the magnetization  $\sum_x S_x^z$  and its dipole moment  $\sum_x x S_x^z$ , i.e., the ‘‘center of mass’’ of the  $S^z$  density. Eq. (1) can be understood as a generalization of the short-range ‘‘pair-hopping’’ models of Refs. [61, 62, 64]. When those systems thermalize, they exhibit subdiffusion with dynamical exponent  $z = 4$  and density correlations that are nonmonotonic in space [38–40]. Here, we are interested in how the hydrodynamics is affected by long-range interactions, particularly for  $J_{sr}$  with power-law decay.

Since directly computing the emergent hydrodynamics of quantum models is generally intractable, here we also introduce a solvable hydrodynamic model that is inspired by Eq. (1). We consider a density  $n_x \in \mathbb{R}$  which is slightly perturbed from its uniform equilibrium value. The dynamics of the model are governed by pairs of equal but opposite currents that displace density from positions  $x+r$  and  $x+r+s$  to positions  $x$  and  $x+s+2r$  (and vice versa), where  $s \geq 0$  and  $r \geq 1$  [Fig. 1(a)]. Such ‘‘pair currents’’ occur in superposition for all  $s$  and  $r$ , but the magnitudes of each are driven by the density at the four positions involved. The analog to Fick’s law is that each pair current is driven with strength proportional to the curvature of  $n_x$  on those four sites [38]. The proportionality for a given  $s$  and  $r$  is denoted  $C_{sr}$  and it encodes the suppression of such pair currents as a function of  $s$  and  $r$ . The equation of motion for the density profile is

$$\begin{aligned} \dot{n}_x = \sum_{r,s} C_{sr} [ & -(n_{x-2r-s} - n_{x-r-s} - n_{x-r} + n_x) \\ & + (n_{x-r-s} - n_{x-s} - n_x + n_{x+r}) \\ & + (n_{x-r} - n_x - n_{x+s} + n_{x+s+r}) \\ & - (n_x - n_{x+r} - n_{x+s+r} + n_{x+s+2r})]. \quad (2) \end{aligned}$$

The four lines correspond to the ways in which four sites with spacings  $s$  and  $r$  can intersect site  $x$ . As a simple tractable choice, we will focus on the separable form

$$C_{sr} = (s+1)^{-\alpha} r^{-\beta}, \quad (3)$$

where  $\alpha$  and  $\beta$  are tuning parameters. Later, we will also discuss the nonseparable form of  $C_{sr}$  that arises in strongly tilted systems.




FIG. 2. The dynamical exponent  $z$ . (a) Contour plot of  $\log_2(\gamma_{2k}/\gamma_k) \cong \hat{z}$  on the  $(\alpha, \beta)$ -plane, numerically evaluated using  $k = 2\pi/3200$ . The dashed box marks  $z = 4$  subdiffusion. The dotted line along  $\beta = 2 + \alpha$  marks where the behavior of  $z$  changes from  $z = \beta - 1$  to  $z = \alpha + 1$ . (b) Demonstration of the boundary behavior  $\hat{z}(k) = 4 - \log(1/k)^{-1}$  where the horizontal axis is  $L = 2\pi/k$ , the lattice sizes used in the estimates. The vertical shift between the two curves is from different subleading (but only by a  $\log(1/k)$  factor) contributions. [(c),(d)] Cuts along the top and right edges of the contour plot. Colors (light to dark) represent estimates using  $k \geq 2\pi/3200$ , i.e., lattices of size  $L \leq 3200$ .

While this classical hydrodynamic model is our main focus, we expect that our results qualitatively carry over to the transport behavior of appropriate quantum systems. Indeed, after solving our hydrodynamic model, below we numerically study the dynamics generated by Eq. (1) for small spin-1 chains and obtain behavior consistent with our hydrodynamic theory if the pair-hopping amplitudes in the Hamiltonian are chosen as  $J_{sr} \sim \sqrt{C_{sr}}$ , in accordance with Fermi’s Golden Rule [98], cf. Fig. 4.

**Hydrodynamic Theory.** Equation (2) is a linear equation with a basis of decaying plane-wave solutions  $n_x = e^{ikx - \gamma_k t}$ , where the decay rates are  $\gamma_k = \sum_{s,r} 16C_{sr} \sin^2(\frac{kr}{2}) \sin^2(\frac{kr+ks}{2})$ . We are interested in the small- $k$  behavior of  $\gamma_k$  as a function of  $\alpha$  and  $\beta$ . In that limit, the lattice approximates a continuum such that

$$\gamma(k) = \int_1^\infty \int_0^\infty \frac{16 \sin^2(\frac{kr}{2}) \sin^2(\frac{kr+ks}{2})}{(s+1)^\alpha r^\beta} ds dr. \quad (4)$$

At small enough  $k$  we expect  $\gamma(k) \propto k^z$  where  $z$  is a dynamical exponent that could depend on  $\alpha$  and  $\beta$ . It is then useful to define an effective exponent  $\hat{z}(k) = \frac{d \log \gamma(k)}{d \log k}$ .

To get an immediate picture of the model’s behavior, in Fig. 2 we numerically evaluate  $\log_2(\gamma_{2k}/\gamma_k) \cong \hat{z}$  on

the  $(\alpha, \beta)$ -plane [Fig. 2(a)], and along two cuts through it [Fig. 2(c),(d)]. We identify three phases: **(A)**  $\alpha > 3$  and  $\beta > 5$ . The power laws are sufficiently short-ranged that  $z = 4$ , consistent with the known behavior in finite-range COM-conserving systems in one dimension [38–40, 52]. **(B)**  $\alpha > \beta - 2$  and  $\beta < 5$ . The long-range “hopping” dominates. The result is a dynamical exponent that continuously varies with  $\beta$  but not  $\alpha$ . This is similar to Refs. [97, 98] where the hydrodynamics of models with long-range hopping (but no higher-moment conservation laws) was studied. In this phase, the dynamical exponent varies within  $z \in (0, 4)$ . **(C)**  $\beta > \alpha + 2$  and  $\alpha < 3$ . The long-range “pairing” dominates, and hopping is short ranged. This again leads to a continuously varying dynamical exponent, but this time it varies with  $\alpha$  and not  $\beta$ . Notably, the fastest relaxation possible in this phase is diffusive ( $z = 2$ ). This results from the effective loss (locally) of the COM conservation that follows from allowing the paired short-ranged currents to be far away from each other.

These phases and boundaries can be studied in more detail by analyzing Eq. (4): At large  $\alpha$  and  $\beta$ , the integral is dominated by  $s, r \ll k^{-1}$  and we can expand the sin functions to leading order which yields  $\gamma(k) \propto k^4$  as expected, i.e.,  $z = 4$ . The resulting integral is only convergent when  $\alpha > 3$  and  $\beta > 5$ , consistent with Fig. 2.

For  $\alpha = \infty$  ( $s = 0$ ), the asymptotic small- $k$  behavior of Eq. (4) is

$$\gamma(k) = \begin{cases} (\beta - 5)^{-1} k^4 & \beta > 5 \\ k^4 \log(1/k) & \beta = 5 \\ (2^\beta - 8)\Gamma(1 - \beta) \sin\left(\frac{\pi\beta}{2}\right) k^{\beta-1} & \beta < 5 \end{cases}, \quad (5)$$

where  $\Gamma(\cdot)$  is the Gamma function. Thus, for  $\beta > 5$  the long-wavelength behavior is characterized by dynamical exponent  $z = 4$ , while for  $\beta < 5$  it is  $z = \beta - 1$ , in accordance with Fig. 2. At the phase boundary  $(\alpha, \beta) = (\infty, 5)$ , the dynamics incurs a logarithmic correction that corresponds to  $\hat{z}(k) = 4 - \log(1/k)^{-1}$ . This scaling of  $\hat{z}$  at the phase boundary is demonstrated in Fig. 2(b).

For  $\beta = \infty$  ( $r = 1$ ), we similarly get

$$\gamma(k) = \begin{cases} (\alpha - 3)^{-1} k^4 & \alpha > 3 \\ k^4 \log(1/k) & \alpha = 3 \\ 2[-\Gamma(1 - \alpha)] \sin\left(\frac{\pi\alpha}{2}\right) k^{\alpha+1} & \alpha < 3 \end{cases}, \quad (6)$$

that is, the behavior changes from  $z = 4$  to  $z = \alpha + 1$  at  $\alpha = 3$ , and at the boundary we obtain the same logarithmic correction.

In contrast, across  $\beta = 2 + \alpha$  [dotted line in Fig. 2(a)] there is a qualitative change of behavior, from  $z = \beta - 1$  to  $z = \alpha + 1$ , but the effective  $z$  converges to a constant without a log correction. The dashed and dotted boundaries in Fig. 2(a) are therefore distinct in that respect. It should be noted that the boundaries between




FIG. 3. Scaled density profiles obtained from time evolving  $n(x, 0) = \delta_{x,0}$  under Eq. (2) (red), and comparison with  $F_z(\xi)$ . The system size is  $L = 3200$  and the time is  $t \sim \gamma_{2\pi/L}^{-1}$  for each panel. Position  $x$  extends from  $-\frac{L}{2}$  to  $\frac{L}{2}$  but we show  $x \geq 0$ . We set  $\alpha = \infty$ , and  $\beta$  varies over panels (a)-(f). The annotated values of  $z$  are used for the scaling of the axes. Data at other late times collapses well onto the same curves, but we show curves from one time for clarity. Data is in red, and  $D^{-1/z} F_z$  is plotted in faint blue for comparison, where  $D$  is extracted from  $n(0, t) = Dt^{1/z}$ , but these are directly behind the data curves due to excellent agreement.

the “phases” are not continuous phase transitions with a diverging length scale, so there is no criticality of that type.

Distinct spatial correlations develop as a result of the various different  $z$  values. In Fig. 3 we show the late-time density profile resulting from evolving  $n(x, 0) = \delta_{x,0}$ , and compare with our theoretical understanding. At late enough times, the relaxation is governed by  $\gamma(k) \cong Dk^z$ , so  $n(x, t) \cong (Dt)^{-1/z} F_z(x/(Dt)^{1/z})$ , where  $D$  is the generalized diffusion constant (depends on  $\alpha$  and  $\beta$ ) and the only parameter controlling the scaling function  $F_z(\cdot)$  is the value of  $z$  governing the long-wavelength relaxation. More specifically, we have  $F_z(\xi) = \int_{-\infty}^{\infty} e^{i\kappa\xi - |\kappa|^z} \frac{d\kappa}{2\pi}$ . In Fig. 3 we test this scaling, with  $z$  set to its predicted asymptotic value in each panel, and find excellent agreement. When  $z = 4$  the sign of correlations is positive at smaller distances, negative at intermediate distances, and positive again at larger distances [Fig. 3(d)-(f)]. At  $z = 3$ , the correlations no longer become positive again at larger distances; they approach zero from below [Fig. 3(c)]. At  $z = 2$ , we regain standard Gaussian diffusion at long times [Fig. 3(b)], which does not have any negative lobe in the spatial profile. Finally, at  $z = 1$  we have a heavy-tailed Lorentzian profile [Fig. 3(a)].

**Dynamics of a quantum spin-1 chain.** Having established the properties of our classical hydrodynamic model, we now turn to less-tractable microscopic many-body quantum systems. Given a quantum spin Hamil-




FIG. 4. Infinite-temperature autocorrelation function for a long-range COM-conserving spin-1 chain as in Eq. (1) with  $J_{sr} \propto s^{-\alpha/2} r^{-\beta/2}$  and system size  $L = 16$ . (a) Varying  $\alpha$  for  $\beta \rightarrow \infty$ . (b) Varying  $\beta$  for  $\alpha \rightarrow \infty$ . The dashed curves indicate power laws  $\propto t^{-1/z}$  with  $z$  set according to the prediction of our hydrodynamic theory. The thin gray curves are data for smaller  $L = 14$ .

tonian  $H$  as in Eq. (1) with couplings  $J_{sr}$ , transition rates will scale as  $\propto J_{sr}^2$  due to Fermi’s Golden rule [98]. Thus, in order to compare the transport properties of  $H$  to our hydrodynamic theory, it is natural to choose  $J_{sr} \propto \sqrt{C_{sr}}$  [100, 102, 103]. More specifically, we study Eq. (1) for spin-1 and set  $J_{sr} = 1$  if  $s + 2r \leq 3$ , while  $J_{sr} \propto s^{-\alpha/2} r^{-\beta/2}$  for terms of longer range. This choice of  $J_{sr}$  avoids signatures of strong Hilbert-space fragmentation and, under  $\alpha, \beta \rightarrow \infty$ , recovers the pair-hopping model of Ref. [61].

In Fig. 4, we show the infinite-temperature autocorrelation function  $\text{tr}[S_x^z(t)S_x^z]/3^L$ , obtained using quantum typicality [104, 105], for chains with  $L = 16$ . The specific site  $x$  is irrelevant due to periodic boundaries. We consider the limiting cases of  $\beta \rightarrow \infty$ , i.e., short hopping distance  $r$  [Fig. 4 (a)], and  $\alpha \rightarrow \infty$ , i.e., short pairing distance  $s$  [Fig. 4 (b)]. In all cases, we find a hydrodynamic tail  $\propto t^{-1/z}$ , where the value of the dynamical exponent  $z$  is consistent with the prediction of our hydrodynamic theory [Fig. 1]. Given the small systems in the quantum case, a more detailed comparison is challenging and the regime of smaller  $\alpha, \beta$  is not accessible. Nevertheless, the data in Fig. 4 substantiates that our hydrodynamic model indeed correctly captures the dynamics of generic quantum systems with long-range pair-hopping processes.

**Quantum systems in a tilted lattice.** Our study of the hydrodynamics in long-range models with COM conservation is also motivated by potential experimental realizations. Consider, e.g., a long-range spin-1/2 XY model in a “tilted” potential,

$$H_{XY} = \sum_{i < j} \frac{J}{|i - j|^\nu} (\sigma_i^+ \sigma_j^- + \sigma_i^- \sigma_j^+) + F \sum_i i n_i, \quad (7)$$

where  $F$  is the tilt strength,  $\nu > 1/2$  the power-law exponent, and  $n_i = (\sigma_i^z + \mathbb{1})/2$ . Such models with power-law interactions can nowadays be achieved in experiments with Rydberg atoms, polar molecules, or trapped ions, at least for certain values of  $\nu$ , e.g., Refs. [97, 106, 107].

When  $F/J$  is large, the model has a long-lived conservation of a dressed version of  $F \sum_i i n_i$ . As we show in the SM [108] using a Schrieffer-Wolff (SW) transformation, the dynamics in the strong-tilt regime are governed by pair-hopping processes  $\sigma_i^+ \sigma_{i+r}^- \sigma_{i+r+s}^- \sigma_{i+2r+s}^+$  (and h.c.). These terms occur with amplitude

$$J_{\text{eff}} \propto \frac{J^3}{F^2} \frac{1}{r^{\nu+1} (r+s)^{\nu+1}} \left( \frac{1}{s^\nu} - \frac{1}{(2r+s)^\nu} \right), \quad (8)$$

and can be understood as third-order processes via, e.g., combining  $\sigma_{i+2r+s}^+ \sigma_{i+r+s}^-$ ,  $\sigma_{i+r+s}^+ \sigma_{i+r}^-$ , and  $\sigma_i^+ \sigma_{i+r+s}^-$ , with the intermediate states being off-shell by energies  $Fr$  and  $F(r+s)$  [108]. Therefore, strongly-tilted long-range models are indeed governed by pair-hopping analogous to our hydrodynamic model. However, we note that Eq. (8) is not separable and depends only on one tuning exponent  $\nu$ .

Similar to our results in the context of Fig. 2, we can make predictions about the hydrodynamics of the strongly-tilted XY model by studying the small- $k$  behavior of the decay rates  $\gamma(k)$ , but now for a hydrodynamic model with the more complicated pair-current strengths  $C_{sr} \propto J_{\text{eff}}^2$ . Doing so, we find that  $z = 4$  subdiffusion in fact remains stable even for originally long-range systems with small  $\nu = 1/2$ . This stability occurs because the exponents in the denominator of  $J_{\text{eff}}^2$  are larger due to the pair-hopping processes being generated at third order. Realizing the dynamical regimes with continuously-varying  $z$  found for our simplified model in Fig. 1 thus seems challenging in strongly tilted systems; see also the SM [108].

**Conclusion.** Using a hydrodynamic model inspired by pair-hopping, we have investigated the interplay of long-range interactions and COM conservation on the dissipative hydrodynamics of 1D systems. We have also corroborated the predictions of our theory using simulations of small spin-1 chains. When the interactions decay sufficiently quickly, the dynamics are subdiffusive with dynamical exponent  $z = 4$ , consistent with earlier work. As the power-law interactions are made longer-ranged, our model enters a phase where  $z$ , and the associated spatial profile of density correlations, vary continuously. One possible extreme for this is when the “pairing distance” of the pair currents is long-range but the “hopping distance” is short-range, where transport can become as fast as diffusive ( $z = 2$ ). Another extreme is where the pairing distance is short and the hopping distance is long, then  $z$  can be made arbitrarily small.

While we have discussed the connections and differences of our pair-hopping model to potential experimental realizations in strongly-tilted systems, we note that in the case of short-range systems, Ref. [52] observed and explained subdiffusive relaxation at long wavelengths even with weak lattice tilts. Indeed, as we numerically illustrate in the SM [108], taking  $F$  to be large appears not to be necessary to study the interplay of long-range

interactions and COM conservation. Nonetheless, it appears that the new regimes of  $z < 4$  found in this work will be difficult to realize using tilted systems. How to realize these regimes in experimental systems is therefore an interesting question for future research.

**Note added.** While finalizing this manuscript we became aware of related independent works by J. Gliozzi *et al.* [109] and Ogunnaike *et al.* [110] that will appear in the same arXiv posting.

**Acknowledgements.** We thank Paolo Glorioso, Vedika Khemani, Tibor Rakovszky, Pablo Sala, and Alex Schuckert for helpful discussions, and Vedika Khemani and David Huse for previous collaboration on related topics. Numerical simulations were performed on Stanford Research Computing Center’s Sherlock cluster. Jonas Richter acknowledges funding from the European Union’s Horizon Europe research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 101060162, and the Packard Foundation through a Packard Fellowship in Science and Engineering. N.O. was supported by the US Department of Energy, Office of Science, Basic Energy Sciences, under Early Career Award Nos. DE-SC0021111. A.M. was supported by the Stanford Q-FARM Bloch Postdoctoral Fellowship in Quantum Science and Engineering and the Gordon and Betty Moore Foundation’s EPIQS Initiative through Grant GBMF8686.

- 
- [1] P. M. Chaikin and T. C. Lubensky, [Principles of Condensed Matter Physics](#) (Cambridge University Press, 1995).
- [2] H. Spohn, [Large Scale Dynamics of Interacting Particles](#), Theoretical and Mathematical Physics (Springer Berlin Heidelberg, 2012).
- [3] Eyal Leviatan, Frank Pollmann, Jens H. Bardarson, David A. Huse, and Ehud Altman, “Quantum thermalization dynamics with matrix-product states,” (2017), [arXiv:1702.08894 \[cond-mat.stat-mech\]](#).
- [4] Benedikt Kloss, Yevgeny Bar Lev, and David Reichman, “Time-dependent variational principle in matrix-product state manifolds: Pitfalls and potential,” *Phys. Rev. B* **97**, 024307 (2018).
- [5] Christopher David White, Michael Zaletel, Roger S. K. Mong, and Gil Refael, “Quantum dynamics of thermalizing systems,” *Phys. Rev. B* **97**, 035127 (2018).
- [6] Tomaz Prosen and Marko Žnidarič, “Matrix product simulations of non-equilibrium steady states of quantum spin chains,” *Journal of Statistical Mechanics: Theory and Experiment* **2009**, P02035 (2009).
- [7] Tibor Rakovszky, C. W. von Keyserlingk, and Frank Pollmann, “Dissipation-assisted operator evolution method for capturing hydrodynamic transport,” *Phys. Rev. B* **105**, 075131 (2022).
- [8] Curt von Keyserlingk, Frank Pollmann, and Tibor Rakovszky, “Operator backflow and the classical simulation of quantum transport,” *Phys. Rev. B* **105**, 245101 (2022).
- [9] Subroto Mukerjee, Vadim Oganesyan, and David Huse, “Statistical theory of transport by strongly interacting lattice fermions,” *Phys. Rev. B* **73**, 035113 (2006).
- [10] Vedika Khemani, Ashvin Vishwanath, and David A. Huse, “Operator spreading and the emergence of dissipative hydrodynamics under unitary evolution with conservation laws,” *Phys. Rev. X* **8**, 031057 (2018).
- [11] Tibor Rakovszky, Frank Pollmann, and C. W. von Keyserlingk, “Diffusive hydrodynamics of out-of-time-ordered correlators with charge conservation,” *Phys. Rev. X* **8**, 031058 (2018).
- [12] Sarang Gopalakrishnan, David A. Huse, Vedika Khemani, and Romain Vasseur, “Hydrodynamics of operator spreading and quasiparticle diffusion in interacting integrable systems,” *Phys. Rev. B* **98**, 220303 (2018).
- [13] Tom Banks and Andrew Lucas, “Emergent entropy production and hydrodynamics in quantum many-body systems,” *Phys. Rev. E* **99**, 022105 (2019).
- [14] Olalla A. Castro-Alvaredo, Benjamin Doyon, and Takato Yoshimura, “Emergent hydrodynamics in integrable quantum systems out of equilibrium,” *Phys. Rev. X* **6**, 041065 (2016).
- [15] Bruno Bertini, Mario Collura, Jacopo De Nardis, and Maurizio Fagotti, “Transport in out-of-equilibrium  $xxz$  chains: Exact profiles of charges and currents,” *Phys. Rev. Lett.* **117**, 207201 (2016).
- [16] Benjamin Doyon, “Lecture notes on Generalised Hydrodynamics,” *SciPost Phys. Lect. Notes*, 18 (2020).
- [17] Marko Ljubotina, Marko Žnidarič, and Tomaž Prosen, “Spin diffusion from an inhomogeneous quench in an integrable system,” *Nature Communications* **8**, 16117 (2017).
- [18] Marko Ljubotina, Marko Žnidarič, and Tomaž Prosen, “Kardar-parisi-zhang physics in the quantum heisenberg magnet,” *Phys. Rev. Lett.* **122**, 210602 (2019).
- [19] Vir B. Bulchandani, Romain Vasseur, Christoph Karrasch, and Joel E. Moore, “Bethe-boltzmann hydrodynamics and spin transport in the  $xxz$  chain,” *Phys. Rev. B* **97**, 045407 (2018).
- [20] Vir B. Bulchandani, “Kardar-parisi-zhang universality from soft gauge modes,” *Phys. Rev. B* **101**, 041411 (2020).
- [21] Sarang Gopalakrishnan and Romain Vasseur, “Kinetic theory of spin diffusion and superdiffusion in  $xxz$  spin chains,” *Phys. Rev. Lett.* **122**, 127202 (2019).
- [22] Enej Ilievski, Jacopo De Nardis, Sarang Gopalakrishnan, Romain Vasseur, and Brayden Ware, “Superuniversality of superdiffusion,” *Phys. Rev. X* **11**, 031023 (2021).
- [23] Jacopo De Nardis, Sarang Gopalakrishnan, and Romain Vasseur, “Non-linear fluctuating hydrodynamics for  $kpz$  scaling in isotropic spin chains,” (2022), [arXiv:2212.03696 \[cond-mat.quant-gas\]](#).
- [24] Sebastian Hild, Takeshi Fukuhara, Peter Schauß, Johannes Zeiher, Michael Knap, Eugene Demler, Immanuel Bloch, and Christian Gross, “Far-from-equilibrium spin transport in heisenberg quantum magnets,” *Phys. Rev. Lett.* **113**, 147205 (2014).
- [25] Paul Niklas Jepsen, Jesse Amato-Grill, Ivana Dimitrova, Wen Wei Ho, Eugene Demler, and Wolfgang Ketterle, “Spin transport in a tunable heisenberg model realized with ultracold atoms,” *Nature* **588**, 403–407 (2020).
- [26] A. Scheie, N. E. Sherman, M. Dupont, S. E. Nagler, M. B. Stone, G. E. Granroth, J. E. Moore, and D. A.

- Tennant, “Detection of kardar–parisi–zhang hydrodynamics in a quantum heisenberg spin-1/2 chain,” *Nature Physics* **17**, 726–730 (2021).
- [27] David Wei, Antonio Rubio-Abadal, Bingtian Ye, Francisco Machado, Jack Kemp, Kritsana Srakaew, Simon Hollerith, Jun Rui, Sarang Gopalakrishnan, Norman Y. Yao, Immanuel Bloch, and Johannes Zeiher, “Quantum gas microscopy of kardar–parisi–zhang superdiffusion,” *Science* **376**, 716–720 (2022).
- [28] B. Bertini, F. Heidrich-Meisner, C. Karrasch, T. Prosen, R. Steinigeweg, and M. Žnidarič, “Finite-temperature transport in one-dimensional quantum lattice models,” *Rev. Mod. Phys.* **93**, 025003 (2021).
- [29] Vir B. Bulchandani and David A. Huse, “Hot band sound,” (2022), [arXiv:2208.13767 \[cond-mat.stat-mech\]](#).
- [30] Christopher M. Langlett and Shenglong Xu, “Noise induced universal diffusive transport in fermionic chains,” (2023), [arXiv:2304.02671 \[cond-mat.str-el\]](#).
- [31] Paolo Glorioso, Luca V. Delacrétaz, Xiao Chen, Rahul M. Nandkishore, and Andrew Lucas, “Hydrodynamics in lattice models with continuous non-Abelian symmetries,” *SciPost Phys.* **10**, 015 (2021).
- [32] Hansveer Singh, Brayden A. Ware, Romain Vasseur, and Aaron J. Friedman, “Subdiffusion and many-body quantum chaos with kinetic constraints,” *Physical Review Letters* **127**, 230602 (2021).
- [33] Maxime Dupont and Joel E. Moore, “Universal spin dynamics in infinite-temperature one-dimensional quantum magnets,” *Physical Review B* **101**, 121106 (2020).
- [34] Marko Ljubotina, Jean-Yves Desaulles, Maksym Serbyn, and Zlatko Papić, “Superdiffusive energy transport in kinetically constrained models,” *Physical Review X* **13**, 011033 (2023).
- [35] Pietro Brighi, Marko Ljubotina, and Maksym Serbyn, “Hilbert space fragmentation and slow dynamics in particle-conserving quantum east models,” (2022), [arXiv:2210.15607 \[quant-ph\]](#).
- [36] Jason Iaconis, Sagar Vijay, and Rahul Nandkishore, “Anomalous subdiffusion from subsystem symmetries,” *Physical Review B* **100**, 214301 (2019).
- [37] Jonas Richter and Arijeet Pal, “Anomalous hydrodynamics in a class of scarred frustration-free hamiltonians,” *Physical Review Research* **4**, l012003 (2022).
- [38] Alan Morningstar, Vedika Khemani, and David A. Huse, “Kinetically constrained freezing transition in a dipole-conserving system,” *Phys. Rev. B* **101**, 214205 (2020).
- [39] Andrey Gromov, Andrew Lucas, and Rahul M. Nandkishore, “Fracton hydrodynamics,” *Phys. Rev. Res.* **2**, 033124 (2020).
- [40] Johannes Feldmeier, Pablo Sala, Giuseppe De Tomasi, Frank Pollmann, and Michael Knap, “Anomalous diffusion in dipole- and higher-moment-conserving systems,” *Phys. Rev. Lett.* **125**, 245303 (2020).
- [41] Jason Iaconis, Andrew Lucas, and Rahul Nandkishore, “Multipole conservation laws and subdiffusion in any dimension,” *Phys. Rev. E* **103**, 022142 (2021).
- [42] Ansgar G. Burchards, Johannes Feldmeier, Alexander Schuckert, and Michael Knap, “Coupled hydrodynamics in dipole-conserving quantum systems,” *Phys. Rev. B* **105**, 205127 (2022).
- [43] Paolo Glorioso, Jinkang Guo, Joaquin F. Rodriguez-Nieva, and Andrew Lucas, “Breakdown of hydrodynamics below four dimensions in a fracton fluid,” *Nature Physics* **18**, 912–917 (2022).
- [44] Jinkang Guo, Paolo Glorioso, and Andrew Lucas, “Fracton hydrodynamics without time-reversal symmetry,” *Phys. Rev. Lett.* **129**, 150603 (2022).
- [45] Paolo Glorioso, Xiaoyang Huang, Jinkang Guo, Joaquin F. Rodriguez-Nieva, and Andrew Lucas, “Goldstone bosons and fluctuating hydrodynamics with dipole and momentum conservation,” (2023), [arXiv:2301.02680 \[hep-th\]](#).
- [46] Charles Stahl, Marvin Qi, Paolo Glorioso, Andrew Lucas, and Rahul Nandkishore, “Fracton superfluid hydrodynamics,” (2023), [arXiv:2303.09573 \[cond-mat.stat-mech\]](#).
- [47] Jung Hoon Han, Ethan Lake, and Sunghan Ro, “Scaling and localization in kinetically constrained diffusion,” (2023), [arXiv:2304.03276 \[cond-mat.stat-mech\]](#).
- [48] Evert van Nieuwenburg, Yuval Baum, and Gil Refael, “From bloch oscillations to many-body localization in clean interacting systems,” *Proceedings of the National Academy of Sciences* **116**, 9269–9274 (2019).
- [49] M. Schulz, C. A. Hooley, R. Moessner, and F. Pollmann, “Stark many-body localization,” *Phys. Rev. Lett.* **122**, 040606 (2019).
- [50] Stephan Mandt, Akos Rapp, and Achim Rosch, “Interacting fermionic atoms in optical lattices diffuse symmetrically upwards and downwards in a gravitational potential,” *Phys. Rev. Lett.* **106**, 250602 (2011).
- [51] Pengfei Zhang, “Subdiffusion in strongly tilted lattice systems,” *Physical Review Research* **2**, 033129 (2020).
- [52] Elmer Guardado-Sanchez, Alan Morningstar, Benjamin M. Spar, Peter T. Brown, David A. Huse, and Waseem S. Bakr, “Subdiffusion and heat transport in a tilted two-dimensional fermi-hubbard system,” *Phys. Rev. X* **10**, 011042 (2020).
- [53] Sebastian Scherg, Thomas Kohlert, Pablo Sala, Frank Pollmann, Bharath Hebbe Madhusudhana, Immanuel Bloch, and Monika Aidelsburger, “Observing non-ergodicity due to kinetic constraints in tilted fermi-hubbard chains,” *Nature Communications* **12**, 4490 (2021).
- [54] Thomas Kohlert, Sebastian Scherg, Pablo Sala, Frank Pollmann, Bharath Hebbe Madhusudhana, Immanuel Bloch, and Monika Aidelsburger, “Experimental realization of fragmented models in tilted fermi-hubbard chains,” (2021), [arXiv:2106.15586 \[cond-mat.quant-gas\]](#).
- [55] Emil J. Bergholtz and Anders Karlhede, “Half-filled lowest landau level on a thin torus,” *Phys. Rev. Lett.* **94**, 026802 (2005).
- [56] E. J. Bergholtz and A. Karlhede, “Quantum hall system in tao-thouless limit,” *Phys. Rev. B* **77**, 155308 (2008).
- [57] Emil J Bergholtz and Anders Karlhede, “‘one-dimensional’ theory of the quantum hall system,” *Journal of Statistical Mechanics: Theory and Experiment* **2006**, L04001 (2006).
- [58] Zheng-Yuan Wang, Shintaro Takayoshi, and Masaaki Nakamura, “Spin-chain description of fractional quantum hall states in the jain series,” *Phys. Rev. B* **86**, 155104 (2012).
- [59] Shriya Pai, Michael Pretko, and Rahul M. Nandkishore, “Localization in fractonic random circuits,” *Phys. Rev. X* **9**, 021003 (2019).
- [60] Shriya Pai and Michael Pretko, “Dynamical scar states

- in driven fracton systems,” *Phys. Rev. Lett.* **123**, 136401 (2019).
- [61] Pablo Sala, Tibor Rakovszky, Ruben Verresen, Michael Knap, and Frank Pollmann, “Ergodicity breaking arising from hilbert space fragmentation in dipole-conserving hamiltonians,” *Phys. Rev. X* **10**, 011047 (2020).
- [62] Vedika Khemani, Michael Hermele, and Rahul Nandkishore, “Localization from hilbert space shattering: From theory to physical realizations,” *Phys. Rev. B* **101**, 174204 (2020).
- [63] Tibor Rakovszky, Pablo Sala, Ruben Verresen, Michael Knap, and Frank Pollmann, “Statistical localization: From strong fragmentation to strong edge modes,” *Phys. Rev. B* **101**, 125126 (2020).
- [64] Sanjay Moudgalya, Abhinav Prem, Rahul Nandkishore, Nicolas Regnault, and B. Andrei Bernevig, “Thermalization and its absence within krylov subspaces of a constrained hamiltonian,” in *Memorial Volume for Shoucheng Zhang* (WORLD SCIENTIFIC, 2021) pp. 147–209.
- [65] Sanjay Moudgalya, B. Andrei Bernevig, and Nicolas Regnault, “Quantum many-body scars and hilbert space fragmentation: a review of exact results,” *Reports on Progress in Physics* **85**, 086501 (2022).
- [66] Sanjay Moudgalya and Olexei I. Motrunich, “Hilbert space fragmentation and commutant algebras,” *Phys. Rev. X* **12**, 011050 (2022).
- [67] Xiaozhou Feng and Brian Skinner, “Hilbert space fragmentation produces an effective attraction between fractons,” *Phys. Rev. Res.* **4**, 013053 (2022).
- [68] Calvin Pozderac, Steven Speck, Xiaozhou Feng, David A. Huse, and Brian Skinner, “Exact solution for the filling-induced thermalization transition in a one-dimensional fracton system,” *Phys. Rev. B* **107**, 045137 (2023).
- [69] Charles Stahl, Ethan Lake, and Rahul Nandkishore, “Spontaneous breaking of multipole symmetries,” *Phys. Rev. B* **105**, 155107 (2022).
- [70] Ethan Lake, Michael Hermele, and T. Senthil, “Dipolar bose-hubbard model,” *Phys. Rev. B* **106**, 064511 (2022).
- [71] Ethan Lake, Hyun-Yong Lee, Jung Hoon Han, and T. Senthil, “Dipole condensates in tilted bose-hubbard chains,” (2022), [arXiv:2210.02470 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2210.02470).
- [72] Ethan Lake and T. Senthil, “Non-fermi liquids from kinetic constraints in tilted optical lattices,” (2023), [arXiv:2302.08499 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2302.08499).
- [73] Philip Zechmann, Ehud Altman, Michael Knap, and Johannes Feldmeier, “Fractonic luttinger liquids and supersolids in a constrained bose-hubbard model,” (2022), [arXiv:2210.11072 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2210.11072).
- [74] Amogh Anakru and Zhen Bi, “Non-fermi liquids from dipolar symmetry breaking,” (2023), [arXiv:2304.01181 \[cond-mat.str-el\]](https://arxiv.org/abs/2304.01181).
- [75] Jian-Keng Yuan, Shuai A. Chen, and Peng Ye, “Fractonic superfluids,” *Phys. Rev. Res.* **2**, 023267 (2020).
- [76] Shuai A. Chen, Jian-Keng Yuan, and Peng Ye, “Fractonic superfluids. ii. condensing subdimensional particles,” *Phys. Rev. Res.* **3**, 013226 (2021).
- [77] Subir Sachdev, K. Sengupta, and S. M. Girvin, “Mott insulators in strong electric fields,” *Phys. Rev. B* **66**, 075128 (2002).
- [78] Susanne Pielawa, Takuya Kitagawa, Erez Berg, and Subir Sachdev, “Correlated phases of bosons in tilted frustrated lattices,” *Phys. Rev. B* **83**, 205135 (2011).
- [79] Oleg Dubinkin, Julian May-Mann, and Taylor L. Hughes, “Theory of dipole insulators,” *Phys. Rev. B* **103**, 125129 (2021).
- [80] Julian May-Mann and Taylor L. Hughes, “Topological dipole conserving insulators and multipolar responses,” *Phys. Rev. B* **104**, 085136 (2021).
- [81] Oleg Dubinkin, Julian May-Mann, and Taylor L. Hughes, “Lieb-schultz-mattis-type theorems and other nonperturbative results for strongly correlated systems with conserved dipole moments,” *Phys. Rev. B* **103**, 125133 (2021).
- [82] M. A. Baranov, M. Dalmonte, G. Pupillo, and P. Zoller, “Condensed matter theory of dipolar quantum gases,” *Chemical Reviews* **112**, 5012–5061 (2012).
- [83] Nicolo Defenu, Tobias Donner, Tommaso Macri, Guido Pagano, Stefano Ruffo, and Andrea Trombettoni, “Long-range interacting quantum systems,” (2021), [arXiv:2109.01063 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2109.01063).
- [84] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, “The physics of dipolar bosonic quantum gases,” *Reports on Progress in Physics* **72**, 126401 (2009).
- [85] Lauriane Chomaz, Igor Ferrier-Barbut, Francesca Ferlaino, Bruno Laburthe-Tolra, Benjamin L. Lev, and Tilman Pfau, “Dipolar physics: a review of experiments with magnetic quantum gases,” *Reports on Progress in Physics* **86**, 026401 (2022).
- [86] M. Saffman, T. G. Walker, and K. Mølmer, “Quantum information with rydberg atoms,” *Rev. Mod. Phys.* **82**, 2313–2363 (2010).
- [87] Antoine Browaeys and Thierry Lahaye, “Many-body physics with individually controlled rydberg atoms,” *Nature Physics* **16**, 132–142 (2020).
- [88] Lincoln D Carr, David DeMille, Roman V Krems, and Jun Ye, “Cold and ultracold molecules: science, technology and applications,” *New Journal of Physics* **11**, 055049 (2009).
- [89] Bryce Gadway and Bo Yan, “Strongly interacting ultracold polar molecules,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **49**, 152002 (2016).
- [90] Steven A. Moses, Jacob P. Covey, Matthew T. Miec-nikowski, Deborah S. Jin, and Jun Ye, “New frontiers for quantum gases of polar molecules,” *Nature Physics* **13**, 13–20 (2017).
- [91] C. Monroe, W. C. Campbell, L.-M. Duan, Z.-X. Gong, A. V. Gorshkov, P. W. Hess, R. Islam, K. Kim, N. M. Linke, G. Pagano, P. Richerme, C. Senko, and N. Y. Yao, “Programmable quantum simulations of spin systems with trapped ions,” *Rev. Mod. Phys.* **93**, 025001 (2021).
- [92] Takuro Matsuta, Tohru Koma, and Shu Nakamura, “Improving the lieb–robinson bound for long-range interactions,” *Annales Henri Poincaré* **18**, 519–528 (2017).
- [93] Luis Colmenarez and David J. Luitz, “Lieb-robinson bounds and out-of-time order correlators in a long-range spin chain,” *Phys. Rev. Res.* **2**, 043047 (2020).
- [94] Dominic V. Else, Francisco Machado, Chetan Nayak, and Norman Y. Yao, “Improved lieb-robinson bound for many-body hamiltonians with power-law interactions,” *Phys. Rev. A* **101**, 022333 (2020).
- [95] Philip Richerme, Zhe-Xuan Gong, Aaron Lee, Crystal Senko, Jacob Smith, Michael Foss-Feig, Spyridon

- Michalakis, Alexey V. Gorshkov, and Christopher Monroe, “Non-local propagation of correlations in quantum systems with long-range interactions,” *Nature* **511**, 198–201 (2014).
- [96] C. Zu, F. Machado, B. Ye, S. Choi, B. Kobrin, T. Mittiga, S. Hsieh, P. Bhattacharyya, M. Markham, D. Twitchen, A. Jarmola, D. Budker, C. R. Laumann, J. E. Moore, and N. Y. Yao, “Emergent hydrodynamics in a strongly interacting dipolar spin ensemble,” *Nature* **597**, 45–50 (2021).
- [97] M. K. Joshi, F. Kranzl, A. Schuckert, I. Lovas, C. Maier, R. Blatt, M. Knap, and C. F. Roos, “Observing emergent hydrodynamics in a long-range quantum magnet,” *Science* **376**, 720–724 (2022).
- [98] Alexander Schuckert, Izabella Lovas, and Michael Knap, “Nonlocal emergent hydrodynamics in a long-range quantum spin system,” *Phys. Rev. B* **101**, 020416 (2020).
- [99] Benedikt Kloss and Yevgeny Bar Lev, “Spin transport in a long-range-interacting spin chain,” *Physical Review A* **99**, 032114 (2019).
- [100] Tianci Zhou, Shenglong Xu, Xiao Chen, Andrew Guo, and Brian Swingle, “Operator lévy flight: Light cones in chaotic long-range interacting systems,” *Physical Review Letters* **124**, 180601 (2020).
- [101] Tomotaka Kuwahara and Keiji Saito, “Strictly linear light cones in long-range interacting systems of arbitrary dimensions,” *Physical Review X* **10**, 031010 (2020).
- [102] Jonas Richter, Oliver Lunt, and Arijeet Pal, “Transport and entanglement growth in long-range random clifford circuits,” *Physical Review Research* **5**, 1012031 (2023).
- [103] Maxwell Block, Yimu Bao, Soonwon Choi, Ehud Altman, and Norman Y. Yao, “Measurement-induced transition in long-range interacting quantum circuits,” *Physical Review Letters* **128**, 010604 (2022).
- [104] Tjark Heitmann, Jonas Richter, Dennis Schubert, and Robin Steinigeweg, “Selected applications of typicality to real-time dynamics of quantum many-body systems,” *Zeitschrift für Naturforschung A* **75**, 421–432 (2020).
- [105] Fengpin Jin, Dennis Willsch, Madita Willsch, Hannes Lagemann, Kristel Michielsen, and Hans De Raedt, “Random state technology,” *Journal of the Physical Society of Japan* **90**, 012001 (2021).
- [106] P. Scholl, H. J. Williams, G. Bornet, F. Wallner, D. Barredo, L. Henriot, A. Signoles, C. Hainaut, T. Franz, S. Geier, A. Tebben, A. Salzinger, G. Zürn, T. Lahaye, M. Weidemüller, and A. Browaeys, “Microwave engineering of programmable  $xxz$  hamiltonians in arrays of rydberg atoms,” *PRX Quantum* **3**, 020303 (2022).
- [107] Lysander Christakis, Jason S. Rosenberg, Ravin Raj, Sungjae Chi, Alan Morningstar, David A. Huse, Zoe Z. Yan, and Waseem S. Bakr, “Probing site-resolved correlations in a spin system of ultracold molecules,” *Nature* **614**, 64–69 (2023).
- [108] See Supplemental Material for details on the Schrieffer-Wolff transformation, as well as numerical results on dynamics in moderately-tilted lattice models of classical spins [111, 112].
- [109] J. Gliozzi, J. May-Mann, T. L. Hughes, and G. De Tomasi, “Hierarchical hydrodynamics in long-range multipole-conserving systems,” (2023).
- [110] Olumakinde Ogunnaike, Johannes Feldmeier, and Jong Yeon Lee, “Unifying emergent hydrodynamics and lindbladian low energy spectra across symmetries, constraints, and long-range interactions,” (2023).
- [111] Vadim Oganesyan, Arijeet Pal, and David A. Huse, “Energy transport in disordered classical spin chains,” *Physical Review B* **80**, 115104 (2009).
- [112] Dennis Schubert, Jonas Richter, Fengping Jin, Kristel Michielsen, Hans De Raedt, and Robin Steinigeweg, “Quantum versus classical dynamics in spin models: Chains, ladders, and square lattices,” *Physical Review B* **104**, 054415 (2021).



# Supplemental Material for: Hydrodynamics in long-range interacting systems with center-of-mass conservation

Alan Morningstar <sup>1</sup>, Nicholas O’Dea,<sup>1</sup> and Jonas Richter <sup>1,2</sup>

<sup>1</sup>*Department of Physics, Stanford University, Stanford, CA 94305, USA*

<sup>2</sup>*Institut für Theoretische Physik, Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

(Dated: April 26, 2023)

## I. EFFECTIVE HAMILTONIAN OF THE LONG-RANGE STRONGLY-TILTED XY MODEL

Below, we will employ a Schrieffer-Wolff transformation to derive the effective coupling strength  $J_{\text{eff}}$  for the strongly-tilted long-range XY model of the main text. Before doing so, it is however instructive to consider on a more basic level which kinds of spin-flip/hopping processes can occur in a tilted system under the constraint of energy conservation. In Fig. S1, we show examples of such energy-conserving third-order processes that effectively lead to pair-hopping processes as discussed in the main text. The effective amplitude of these processes can be obtained by multiplying the power-law interaction strength with the corresponding energy denominator for all individual virtual hops in the process, see caption of Fig. S1 for more details. As we will show below, the resulting expressions for  $J_{\text{eff}}$  agree with the effective Hamiltonian obtained using the significantly more involved Schrieffer-Wolff technique. Furthermore, we note that lower order processes, as well as higher-order processes involving more lattice sites, will not contribute as their amplitudes will cancel.

In the following, we use the Schrieffer-Wolff technique to generate an effective Hamiltonian for the tilted spin-1/2 XY chain in the limit of large  $F$ . This effective Hamiltonian describes dynamics within subspaces with the same energy under  $F \sum_i i(\sigma_i^z + \mathbb{1})/2$ . The following calculations are performed in the thermodynamic limit and were inspired by those for a tilted short-range model in [1].

It is useful in the following to rewrite the Hamiltonian as

$$\frac{H}{F} = D + \lambda T \quad (\text{S1})$$

where  $T \equiv \sum_{\delta \neq 0} C_\delta T_\delta$ ,  $C_\delta \equiv \frac{1}{|\delta|^\nu}$ ,  $T_\delta \equiv \sum_x \sigma_{x+\delta}^+ \sigma_x^-$ ,  $D \equiv F \sum_i i(\sigma_i^z + \mathbb{1})/2$ ,  $\lambda \equiv \frac{J}{F}$ .

Our aim is to construct a unitary transformation that brings this Hamiltonian to a “block diagonal” form,  $H_{\text{eff}}$ , describing dynamics in sectors labeled by eigenvalues of  $D$ .

We’ll construct this transformation and  $H_{\text{eff}}$  order-by-order in  $\lambda$ :

$$H_{\text{eff}} = e^{\lambda S} H e^{-\lambda S}, \quad (\text{S2})$$

$$H_{\text{eff}} = \sum_n \lambda^n H_{\text{eff}}^{(n)} \quad (\text{S3})$$

$$S = \sum_n \lambda^n S^{(n)} \quad (\text{S4})$$

Our aim is to find the lowest non-trivial contribution beyond  $D$ , which we’ll show to be  $\lambda^3$ . Expanding  $H_{\text{eff}} = e^{\lambda S} H e^{-\lambda S}$  to this order yields

$$\begin{aligned} \frac{H_{\text{eff}}}{F} = & D + \lambda(T + [S_0, D]) + \lambda^2 \left( \frac{1}{2} [S_0, [S_0, D]] + [S_0, T] + [S_1, D] \right) \\ & + \lambda^3 \left( \frac{1}{6} [S_0, [S_0, [S_0, D]]] + \frac{1}{2} ([S_1, [S_0, D]] + [S_0, [S_1, D]] + [S_0, [S_0, T]]) + [S_1, T] + [S_2, D] \right) + O(\lambda^4) \end{aligned} \quad (\text{S5})$$

Order  $\lambda^n$  determines the form of  $S_{n-1}$ , as the anti-hermitian  $S$  must be chosen so  $H_{\text{eff}}$  is block-diagonal order-by-order. For the sake of determining  $S_0$ , note that  $T$  is purely off-block-diagonal relative to  $D$ , since  $T_\delta$  shifts the eigenvalues of  $D$  by a factor of  $\delta$ . Thus, we must have

$$[D, S_0] = T \quad (\text{S6})$$




FIG. S1. Sketch of exemplary third-order hopping process that are central to the emergent pair-hopping in quantum lattice models with strong tilt  $F$ . **(a)** First, the right particle hops a distance  $r$ , which costs an energy  $Fr$ . Then, the left particle hops a distance  $s$  and thereafter occupies the original position of the other particle. The resulting state is off by an energy  $F(r+s)$  compared to the original configuration. Finally, the left particle hops a distance  $r+s$  to the left, resulting in the final configuration that has the same energy and the same center-of-mass as the original state. **(b)** Another process, now involving a longer hop of distance  $2r+s$ . The effective amplitude  $J_{\text{eff}}$  of these processes can be obtained by multiplying the interaction strength  $\propto |i-j|^{-\nu}$  by the energy denominator  $1/E$ : For example in the case of (a), we have  $J_{\text{eff}} = \frac{J}{r^{\nu}} \frac{1}{Fr} \frac{J}{s^{\nu}} \frac{1}{F(r+s)} \frac{J}{(r+s)^{\nu}}$ . The resulting expressions for  $J_{\text{eff}}$  agree with effective Hamiltonian obtained using the Schrieffer-Wolff transformation in Eq. (S27). Note that there exist additional third-order processes that yield a nonzero contribution, similar to the ones shown here, as well as their symmetric mirrored versions.

to ensure  $H_{\text{eff}}$  is block-diagonal at order  $\lambda$ . We note that  $[D, T_{\delta}] = \delta T_{\delta}$ , so, remembering that  $T$  is defined as  $T \equiv \sum_{\delta \neq 0} C_{\delta} T_{\delta}$ , we have that

$$S_0 = \sum_{\delta \neq 0} C_{\delta} \frac{T_{\delta}}{\delta} \quad (\text{S7})$$

is a purely off-block-diagonal solution to Eq. (S6).

From Eq. (S6), we can simplify  $H_{\text{eff}}$  to

$$\frac{H_{\text{eff}}}{F} = D + \lambda^2 \left( \frac{1}{2} [S_0, T] + [S_1, D] \right) + \quad (\text{S8})$$

$$\lambda^3 \left( \frac{1}{3} [S_0, [S_0, T]] + \frac{1}{2} ([S_1, T] + [S_0, [S_1, D]]) + [S_2, D] \right) + O(\lambda^4) \quad (\text{S9})$$

To find a solution for  $S_1$ , note first that

$$[S_0, T] = \sum_{\delta \neq 0} \sum_{\delta' \neq 0} \frac{C_{\delta}}{\delta} C_{\delta'} K_{\delta, \delta'} \quad (\text{S10})$$

where  $K$  is defined via

$$K_{\delta, \delta'} \equiv [T_{\delta}, T_{\delta'}] = \sum_i \sigma_{i+\delta+\delta'}^+ \sigma_i^- (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) \quad (\text{S11})$$

From the commutator  $[D, T_{\delta}] = \delta T_{\delta}$ , it follows that  $[D, K_{\delta, \delta'}] = (\delta + \delta') K_{\delta, \delta'}$ . That is,  $K_{\delta, \delta'}$  changes an eigenvector of  $D$  into another eigenvector with eigenvalue shifted by  $\delta + \delta'$ . This is purely off-block-diagonal except for terms with  $\delta' = -\delta$ . However, such terms with  $\delta' = -\delta$  vanish:

$$K_{\delta, -\delta} = \sum_i \sigma_i^+ \sigma_i^- (\sigma_{i+\delta}^z - \sigma_{i-\delta}^z) \quad (\text{S12})$$

$$= \sum_i \frac{1 + \sigma_i^z}{2} (\sigma_{i+\delta}^z - \sigma_{i-\delta}^z) \quad (\text{S13})$$

$$= \frac{1}{2} \sum_i (\sigma_{i+\delta}^z - \sigma_{i-\delta}^z) + \frac{1}{2} \sum_i (\sigma_i^z \sigma_{i+\delta}^z - \sigma_{i-\delta}^z \sigma_i^z) \quad (\text{S14})$$

The final expression is the sum of two telescoping series,  $K_{\delta,-\delta} = 0$ . That is,  $[S_0, T]$  is purely off-block-diagonal. As  $[S_0, T]$  is off-block-diagonal, so we must have

$$[D, S_1] = \frac{1}{2}[S_0, T] \quad (\text{S15})$$

to ensure  $H_{\text{eff}}$  is block-diagonal at order  $\lambda^2$ . Using Eq. (S10),  $[D, K_{\delta,\delta'}] = (\delta + \delta')K_{\delta,\delta'}$ , it follows that

$$S_1 = \sum_{\delta \neq 0} \sum_{\delta' \neq 0} \frac{C_\delta C_{\delta'} K_{\delta,\delta'}}{2\delta(\delta + \delta')} \quad (\text{S16})$$

is a purely off-block-diagonal solution to Eq. (S15).

From Eq. (S15), we can simplify  $H_{\text{eff}}$  to

$$\frac{H_{\text{eff}}}{F} = D + \lambda^3 \left( \frac{1}{12}[S_0, [S_0, T]] + \frac{1}{2}[S_1, T] + [S_2, D] \right) + O(\lambda^4) \quad (\text{S17})$$

$S_2$  will be chosen to cancel the off-diagonal pieces of  $\frac{1}{12}[S_0, [S_0, T]] + \frac{1}{2}[S_1, T]$ , so we have that

$$\frac{H_{\text{eff}}}{F} = D + \lambda^3 \left( \frac{1}{12}[S_0, [S_0, T]] + \frac{1}{2}[S_1, T] \right) \Big|_{BD} + O(\lambda^4) \quad (\text{S18})$$

where  $|_{BD}$  is shorthand meaning that only the block-diagonal piece is included. Using Eqs. (S7), (S10), (S16), we have

$$H_{\text{eff}}^{(3)} = \sum_{\delta \neq 0} \sum_{\delta' \neq 0} \sum_{\delta'' \neq 0} C_\delta C_{\delta'} C_{\delta''} \left( \frac{1}{4\delta(\delta + \delta')} - \frac{1}{12\delta\delta''} \right) M_{\delta,\delta',\delta''} \Big|_{BD} \quad (\text{S19})$$

where  $M_{\delta,\delta',\delta''} \equiv [K_{\delta,\delta'}, T_{\delta''}]$ . Evaluating the commutators and shifting summation indices freely yields:

$$\begin{aligned} M_{\delta,\delta',\delta''} = \sum_i & \left( \sigma_{i+\delta+\delta'+\delta''}^+ (\sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) + \sigma_{i+\delta''}^z (\sigma_{i+\delta'+\delta''}^z - \sigma_{i+\delta+\delta''}^z)) \sigma_i^- \right. \\ & \left. + 2\sigma_{i+\delta+\delta'}^+ (\sigma_{i+\delta}^+ \sigma_{i+\delta-\delta''}^- + \sigma_{i+\delta'+\delta''}^+ \sigma_{i+\delta'}^- - \sigma_{i+\delta'}^+ \sigma_{i+\delta'-\delta''}^- - \sigma_{i+\delta+\delta''}^+ \sigma_{i+\delta}^-) \sigma_i^- \right) \end{aligned} \quad (\text{S20})$$

Note that  $[D, T_\delta] = \delta T_\delta$  implies  $[D, M_{\delta,\delta',\delta''}] = (\delta + \delta' + \delta'')M_{\delta,\delta',\delta''}$ . It follows that the only block-diagonal pieces of  $M$  are those for which  $\delta'' = -(\delta + \delta')$ . That is, we have

$$H_{\text{eff}}^{(3)} = \sum_{\delta \neq 0} \sum_{\delta' \neq 0} \frac{C_\delta C_{\delta'} C_{\delta+\delta'}}{3\delta(\delta + \delta')} M_{\delta,\delta',-\delta-\delta'} \quad (\text{S21})$$

We can simplify  $M_{\delta,\delta',-\delta-\delta'}$ , starting from the expression

$$\begin{aligned} M_{\delta,\delta',-\delta-\delta'} = \sum_i & \left( \sigma_i^+ (\sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) + \sigma_{i-\delta-\delta'}^z (\sigma_{i-\delta}^z - \sigma_{i-\delta'}^z)) \sigma_i^- \right. \\ & \left. + 2\sigma_{i+\delta+\delta'}^+ (\sigma_{i+\delta}^+ \sigma_{i+2\delta+\delta'}^- + \sigma_{i-\delta}^+ \sigma_{i+\delta'}^- - \sigma_{i+\delta'}^+ \sigma_{i+\delta+2\delta'}^- - \sigma_{i-\delta'}^+ \sigma_{i+\delta}^-) \sigma_i^- \right) \end{aligned} \quad (\text{S22})$$

Note that in the first line,  $\sigma_i^-$  commutes with the terms in parentheses as each of those terms is on a different site

TABLE I. **Left:**  $r$  and  $s$  in terms of  $\delta$  and  $\delta'$  for the term  $\sum_i \sigma_{i+\delta+\delta'}^+ \sigma_{i+\delta}^+ \sigma_{i+2\delta+\delta'}^- \sigma_i^-$  and its Hermitian conjugate in  $M_{\delta,\delta',-\delta-\delta'}$  for sets partitioning the  $(\delta, \delta')$  plane. The contribution to  $D_{s,r}$  from summing over such a set is given in the rightmost column. **Right:**  $r$  and  $s$  in terms of  $\delta$  and  $\delta'$  for the term  $\sum_i \sigma_{i+\delta+\delta'}^+ \sigma_{i+\delta'}^+ \sigma_{i+2\delta'+\delta}^- \sigma_i^-$  and its Hermitian conjugate in  $M_{\delta,\delta',-\delta-\delta'}$ .

	$r$	$s$	$\frac{3}{2}D_{s,r}$		$r$	$s$	$\frac{3}{2}D_{s,r}$
$\delta > 0, \delta' > 0$	$\delta$	$\delta'$	$\frac{-1}{r^\nu s^{\nu+1} (r+s)^{\nu+1}}$	$\delta > 0, \delta' > 0$	$\delta$	$\delta'$	$\frac{-1}{r^\nu s^{\nu+1} (r+s)^{\nu+1}}$
$\delta >  \delta' , \delta' < 0$	$\delta + \delta'$	$-\delta'$	$\frac{-1}{r^{\nu+1} s^{\nu+1} (r+s)^\nu}$	$\delta >  \delta' , \delta' < 0$	$\delta + \delta'$	$-\delta'$	$\frac{-1}{r^{\nu+1} s^{\nu+1} (r+s)^\nu}$
$ \delta'  > \delta >  \delta' /2, \delta' < 0$	$-(\delta + \delta')$	$2\delta + \delta'$	$\frac{1}{r^{\nu+1} (2r+s)^{\nu+1} (r+s)^\nu}$	$ \delta'  > \delta >  \delta' /2, \delta' < 0$	$-(\delta + \delta')$	$2\delta + \delta'$	$\frac{1}{r^{\nu+1} (2r+s)^{\nu+1} (r+s)^\nu}$
$ \delta' /2 > \delta > 0, \delta' < 0$	$\delta$	$-(2\delta + \delta')$	$\frac{1}{r^\nu (2r+s)^{\nu+1} (r+s)^{\nu+1}}$	$ \delta' /2 > \delta > 0, \delta' < 0$	$\delta$	$-(2\delta + \delta')$	$\frac{1}{r^\nu (2r+s)^{\nu+1} (r+s)^{\nu+1}}$
$\delta' > 2 \delta , \delta < 0$	$-\delta$	$2\delta + \delta'$	$\frac{1}{r^\nu (2r+s)^{\nu+1} (r+s)^{\nu+1}}$	$\delta' > 2 \delta , \delta < 0$	$-\delta$	$2\delta + \delta'$	$\frac{1}{r^\nu (2r+s)^{\nu+1} (r+s)^{\nu+1}}$
$2 \delta  > \delta' >  \delta , \delta < 0$	$\delta + \delta'$	$-(2\delta + \delta')$	$\frac{1}{r^{\nu+1} (2r+s)^{\nu+1} (r+s)^\nu}$	$2 \delta  > \delta' >  \delta , \delta < 0$	$\delta + \delta'$	$-(2\delta + \delta')$	$\frac{1}{r^{\nu+1} (2r+s)^{\nu+1} (r+s)^\nu}$
$ \delta  > \delta', \delta < 0$	$-(\delta + \delta')$	$\delta'$	$\frac{-1}{r^{\nu+1} s^{\nu+1} (r+s)^\nu}$	$ \delta  > \delta', \delta < 0$	$-(\delta + \delta')$	$\delta'$	$\frac{-1}{r^{\nu+1} s^{\nu+1} (r+s)^\nu}$
$\delta < 0, \delta' < 0$	$-\delta$	$\delta'$	$\frac{-1}{r^\nu s^{\nu+1} (r+s)^{\nu+1}}$	$\delta < 0, \delta' < 0$	$-\delta$	$-\delta'$	$\frac{-1}{r^\nu s^{\nu+1} (r+s)^{\nu+1}}$

$(\delta, \delta', \delta + \delta' \neq 0)$ ; we will use such arguments freely to simplify expressions. Commuting that term over, we have

$$\begin{aligned}
& \sum_i \left( \sigma_i^+ (\sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) + \sigma_{i-\delta-\delta'}^z (\sigma_{i-\delta}^z - \sigma_{i-\delta'}^z)) \sigma_i^- \right) \\
&= \frac{1}{2} \sum_i \left( (\sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) + \sigma_{i-\delta-\delta'}^z (\sigma_{i-\delta}^z - \sigma_{i-\delta'}^z)) \right) + \frac{1}{2} \sum_i \left( \sigma_i^z (\sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) + \sigma_{i-\delta-\delta'}^z (\sigma_{i-\delta}^z - \sigma_{i-\delta'}^z)) \right) \\
&= \frac{1}{2} \sum_i \left( \sigma_{i+\delta+\delta'}^z \sigma_{i+\delta}^z + \sigma_{i-\delta-\delta'}^z \sigma_{i-\delta}^z \right) - \frac{1}{2} \sum_i \left( \sigma_{i+\delta+\delta'}^z \sigma_{i+\delta'}^z + \sigma_{i-\delta-\delta'}^z \sigma_{i-\delta'}^z \right) \\
&+ \frac{1}{2} \sum_i \left( \sigma_{i+\delta+\delta'}^z \sigma_{i+\delta}^z \sigma_i^z - \sigma_i^z \sigma_{i-\delta-\delta'}^z \sigma_{i-\delta}^z \right) - \frac{1}{2} \sum_i \left( \sigma_{i+\delta+\delta'}^z \sigma_{i+\delta'}^z \sigma_i^z - \sigma_i^z \sigma_{i-\delta-\delta'}^z \sigma_{i-\delta'}^z \right) \\
&= \sum_i \left( \sigma_{i+\delta+\delta'}^z (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) \right) \\
&= \sum_i \left( (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) \sigma_i^z \right). \tag{S23}
\end{aligned}$$

Furthermore, noticing that  $\sum_i \sigma_{i+\delta+\delta'}^+ \sigma_{i-\delta}^+ \sigma_{i+\delta'}^- \sigma_i^- = \sum_i \sigma_{i+2\delta+\delta'}^+ \sigma_i^+ \sigma_{i+\delta+\delta'}^- \sigma_{i+\delta}^- = (\sum_i \sigma_{i+\delta+\delta'}^+ \sigma_{i+\delta}^+ \sigma_{i+2\delta+\delta'}^- \sigma_i^-)^\dagger$ , and that  $M$  is antisymmetric on its first two indices, we can simplify

$$M_{\delta,\delta',-\delta-\delta'} = \sum_i (\sigma_{i+\delta}^z - \sigma_{i+\delta'}^z) \sigma_i^z + 2 \left( (\sigma_{i+\delta+\delta'}^+ \sigma_{i+\delta}^+ \sigma_{i+2\delta+\delta'}^- \sigma_i^- + h.c.) - (\delta \leftrightarrow \delta') \right) \tag{S24}$$

Our aim is to relate the hopping terms involving  $\delta$  and  $\delta'$  to  $\sum_i \sigma_i^+ \sigma_{i+r}^- \sigma_{i+r+s}^- \sigma_{i+2r+s}^+$  with  $r$  and  $s$  positive; that is, we want to find  $D_{s,r}$  in

$$\sum_{s>0} \sum_{r>0} D_{s,r} \sum_i (\sigma_i^+ \sigma_{i+r}^- \sigma_{i+r+s}^- \sigma_{i+2r+s}^+ + h.c.) \equiv \sum_{\delta \neq 0} \sum_{\delta' \neq 0} \frac{2C_\delta C_{\delta'} C_{\delta+\delta'}}{3\delta(\delta+\delta')} \sum_j \left( (\sigma_{j+\delta+\delta'}^+ \sigma_{j+\delta}^+ \sigma_{j+2\delta+\delta'}^- \sigma_j^- + h.c.) - (\delta \leftrightarrow \delta') \right) \tag{S25}$$

We do this by breaking up the sums on  $\delta$  and  $\delta'$  onto sets partitioning the  $(\delta, \delta')$  plane, each of which has a simple relationship between  $(r, s)$  and  $(\delta, \delta')$ . We write out these relationships in Table I.

Summing up all the contributions to  $D_{s,r}$  from the columns of Tables I, we have that

$$\begin{aligned}
D_{s,r} &= \frac{4}{3} \left( \frac{2}{r^{\nu+1} s^\nu (r+s)^{\nu+1}} - \frac{2}{r^{\nu+1} (2r+s)^\nu (r+s)^{\nu+1}} - \frac{1}{r^\nu s^{\nu+1} (r+s)^{\nu+1}} \right. \\
&\quad \left. + \frac{1}{r^{\nu+1} s^{\nu+1} (r+s)^\nu} - \frac{1}{r^{\nu+1} (2r+s)^{\nu+1} (r+s)^\nu} - \frac{1}{r^\nu (2r+s)^{\nu+1} (r+s)^{\nu+1}} \right) \\
&= 4 \left( \frac{1}{r^{\nu+1} s^\nu (r+s)^{\nu+1}} - \frac{1}{r^{\nu+1} (2r+s)^\nu (r+s)^{\nu+1}} \right)
\end{aligned} \tag{S26}$$

That is, we have

$$J_{\text{eff}} = \frac{4J^3}{F^2} \left( \frac{1}{r^{\nu+1} s^\nu (r+s)^{\nu+1}} - \frac{1}{r^{\nu+1} (2r+s)^\nu (r+s)^{\nu+1}} \right) \tag{S27}$$

Note that this expression can also be motivated by directly considering the kinds of virtual processes in Fig. S1.

## II. NUMERICAL RESULTS FOR SYSTEMS WITH MODERATE LATTICE TILTS

Studying the transport properties of many-body quantum systems, especially with long-range interactions, is quite challenging. Therefore, instead of a quantum system, we here consider a classical spin chain where larger system sizes are accessible. Specifically, we consider a tilted XXZ chain with open boundary conditions,

$$\begin{aligned}
H &= \sum_{\ell=1}^{L-1} \sum_{\ell'>\ell}^L \frac{J}{|\ell' - \ell|^\nu} (S_\ell^x S_{\ell'}^x + S_\ell^y S_{\ell'}^y + \Delta S_\ell^z S_{\ell'}^z) \\
&\quad + F \sum_{\ell} \ell S_\ell^z,
\end{aligned} \tag{S28}$$

where the real three-component vectors  $\mathbf{S}_\ell = (S_\ell^x, S_\ell^y, S_\ell^z)$  have unit length,  $\nu \geq 0.5$  controls the range of interactions,  $\Delta$  is an anisotropy in the  $z$ -direction, and we set  $J = 1$ .

The time evolution is governed by the classical equations of motion

$$\frac{\partial}{\partial t} \mathbf{S}_\ell = \frac{\partial H}{\partial \mathbf{S}_\ell} \times \mathbf{S}_\ell, \tag{S29}$$

and we study the infinite-temperature spin-spin correlation function

$$C(r, t) = \langle S_{\ell+r}^z(t) S_\ell^z \rangle, \tag{S30}$$

where the brackets indicate averaging over random initial spin configurations, and  $r$  is the distance between the involved spins. Numerically solving Eq. (S29) can be challenging due to the tilt  $F > 0$ , which leads to large terms that require an extremely small time step to guarantee an accurate iterative time evolution. Therefore, we here consider an ‘‘even-odd’’ time evolution inspired by Ref. [2]. In the first half time-step, the odd spins are held stationary and the even spins precess in an effective field generated by the odd spins and the local tilt, while in the second half time-step the even spins are fixed and the odd spins rotate. This approach is numerically more stable and preserves conservation laws even for larger time




FIG. S2. (a) Autocorrelation function  $C(0, t)$  in the long-range classical spin chain with  $\nu = 0.5$ , tilt  $F = 1$ , anisotropy  $\Delta = 0$ , and system size  $L = 50$ . Dashed curves indicates a power-law  $\propto t^{-1/4}$ . (b) Full spatial profile  $C(r, t)$  at different times  $t$ . Data and distance are rescaled by  $t^{1/z}$  with  $z \approx 4$ .

steps. However, the downside is that for long-range interactions as in Eq. (S28), it discards interactions between spins on sites with the same parity, i.e., we effectively consider a model where  $J = 0$  if  $\ell$  and  $\ell'$  are both even (both odd). Given the universality of hydrodynamics, we nevertheless expect that the emerging transport behavior will be the same as in the original model in Eq. (S28), at least on long time and length scales.

First, in Fig. S2, we consider the case of  $\Delta = 0$ , i.e., a classical XY chain. Choosing a moderate tilt  $F = 1$  and long-range couplings with  $\nu = 1/2$ , we find in Fig. S2 (a) that the autocorrelation function  $C(0, t)$  develops a hydrodynamic tail that is consistent with a dynamical exponent  $z = 4$ ,  $C(0, t) \propto t^{-1/4}$ . Correspondingly, the full spatial profiles  $C(r, t)$  for different times  $t$  collapse onto each other when data and distances are appropriately rescaled by  $t^{1/z}$  [Fig. S2 (b)]. Moreover,  $C(r, t)$  exhibits anticorrelated regions where  $C(r, t) < 0$ , in good agreement with  $z = 4$  subdiffusion in short-range models. These numerical results for the classical spin chain are consistent with our findings for the strongly-tilted XY quantum spin chain discussed in the main text, where the Schrieffer-Wolff transformation was used to predict the stability of  $z = 4$  down to small  $\nu$ . In particular, the data in Fig. S2 suggests that the same holds true even for weak lattice tilts.

Finally, let us consider the case  $\Delta \neq 0$ . Specifically, we here choose  $\Delta = 1.5$  for which it is well established that




FIG. S3. **(a)** Autocorrelation function  $C(0, t)$  in the long-range classical spin chain with  $\nu = 2, 1, 0.5$ , tilt  $F = 2$ , anisotropy  $\Delta = 1.5$ , and system size  $L = 100$ . Dashed curves are power-law fits  $\propto t^{-1/z}$ . **[(b), (c), (d)]** Full spatial profile  $C(r, t)$  at different times  $t$  for the three values of  $\nu$ . Data and distance are rescaled by  $t^{1/z}$  leading to convincing data collapses. The values of  $z$  used for the data collapse are extracted from the fits in (a) and are indicated in the panels.

the short-range model ( $\nu \rightarrow \infty$ ) exhibits diffusive spin transport at zero tilt  $F = 0$  [3]. In Fig. S3 (a), we show  $C(0, t)$  for  $\nu = 2, 1, 0.5$  and fixed tilt  $F = 2$ . We observe that  $C(0, t)$  exhibits a power-law tail  $\propto t^{-1/z}$ , where  $z$  now appears to vary with  $\nu$  (in contrast to the case of  $\Delta = 0$  in Fig. S2). Specifically, we find subdiffusion with  $z \approx 4$  at  $\nu = 2$ , analogous to tilted short-range models, and  $z \approx 3$  at  $\nu = 1$ . At  $\nu = 0.5$ ,  $z \approx 2.3$  appears to approach the diffusive value, although our numerically extracted value of  $z$  is still slightly higher.

The full spatial profiles  $C(r, t)$  for the three values of  $\nu$  are found to collapse onto single curves when rescaled by  $t^{1/z}$  using the corresponding values of  $z$  [Fig. S3 (b)-(d)]. Moreover, the shapes of  $C(r, t)$  in Fig. S3 are qualitatively similar to the behavior of  $n(x, t)$  in the classical hydrodynamic model discussed in the main text. Thus, in contrast to the tilted XY model, the numerical results for the classical XXZ chain in Fig. S3 suggest that it might indeed be possible to realize a regime with continuously varying  $z$  in tilted systems if one additionally includes  $zz$  interactions. We leave further exploration of this for future work.

- [1] Sanjay Moudgalya, Abhinav Prem, Rahul Nandkishore, Nicolas Regnault, and B. Andrei Bernevig, “Thermalization and its absence within krylov subspaces of a constrained hamiltonian,” in [Memorial Volume for Shoucheng Zhang](#) (WORLD SCIENTIFIC, 2021) pp. 147–209.
- [2] Vadim Oganesyan, Arijeet Pal, and David A. Huse, “En-

- ergy transport in disordered classical spin chains,” [Physical Review B](#) **80**, 115104 (2009).
- [3] Dennis Schubert, Jonas Richter, Fengping Jin, Kristel Michielsen, Hans De Raedt, and Robin Steinigeweg, “Quantum versus classical dynamics in spin models: Chains, ladders, and square lattices,” [Physical Review B](#) **104**, 054415 (2021).