A generalized fractional Laplacian Xiangcheng Zheng^{*}

V.J. Ervin[†]

Hong Wang[‡]

May 3, 2024

Abstract

In this article we show that the fractional Laplacian in \mathbb{R}^2 can be factored into a product of the divergence operator, a Riesz potential operator, and the gradient operator. Using this factored form we introduce a generalization of the fractional Laplacian, involving a matrix K(x), suitable when the fractional Laplacian is applied in a non homogeneous medium. For the case of K(x) a constant, symmetric positive definite matrix we show that the fractional Poisson equation is well posed, and determine the regularity of the solution in terms of the regularity of the right hand side function.

Key words. Fractional Laplacian, Riesz potential operator, Jacobi polynomials, spherical harmonics

AMS Mathematics subject classifications. 35S15, 42C10, 35B65, 33C55

1 Introduction

In recent years there has been considerable interest in studying nonlocal differential operators. The motivation for this work has been to develop tools that can be used to more accurately model phenomena such as anomalous diffusion, which cannot be suitable modeled using local operators. One such nonlocal operator which has received much attention is the Fractional Laplace operator (a.k.a., the Fractional Laplacian). To infer the Fractional Laplacian refers to a single operator is somewhat misleading as there are several operators that come under this general heading. For example, for $0 < \alpha < 2$, on a bounded domain three such operators are [2]:

(i) The Spectral Fractional Laplacian

The Spectral Fractional Laplacian, $(-\Delta)_S^{\frac{\alpha}{2}}$, is defined in terms of the eigenvalues and eigenfunctions of the Laplace operator with homogeneous Dirichlet boundary conditions on Ω . Let $\{\psi_k\}_{k\in\mathbb{N}}$ denote

^{*}School of Mathematics, Shandong University, Jinan 250100, China. email: xzheng@sdu.edu.cn.

[†]School of Mathematical and Statistical Sciences, Clemson University, Clemson, South Carolina 29634-0975, USA. email: vjervin@clemson.edu.

[‡]Department of Mathematics, University of South Carolina, Columbia, South Carolina 29208, USA. email: hwang@math.sc.edu.

an orthonormal basis for $L^2(\Omega)$ satisfying, for $\lambda_k \in \mathbb{R}^+$,

$$-\Delta \psi_k = \lambda_k \psi_k \text{ in } \Omega,$$

$$\psi_k = 0 \text{ on } \partial \Omega.$$

Then $(-\Delta)_S^{\frac{\alpha}{2}}$ is defined for $u \in C_0^{\infty}(\Omega)$ as

$$(-\Delta)_{S}^{\frac{\alpha}{2}} u = \sum_{k=1}^{\infty} \langle u, \psi_{k} \rangle \lambda_{k}^{\frac{\alpha}{2}} \psi_{k},$$

and by extension for $u \in H_0^{\frac{\alpha}{2}}(\Omega)$.

(ii) The Integral Fractional Laplacian

The Integral Fractional Laplacian, $(-\Delta)^{\frac{\alpha}{2}}$, is defined for $support(u) \subset \Omega \subset \mathbb{R}^d$ as

$$(-\Delta)^{\frac{\alpha}{2}} u(x) = \frac{1}{|\gamma_d(-\alpha)|} \lim_{\epsilon \to 0} \int_{\mathbb{R}^d \setminus B(x,\epsilon)} \frac{u(x) - u(y)}{|x - y|^{d + \alpha}} dy, \quad x \in \Omega,$$
(1.1)

where $\gamma_d(\alpha) := 2^{\alpha} \pi^{d/2} \Gamma(\frac{\alpha}{2}) / \Gamma((d-\alpha)/2)$, and $B(x,\epsilon)$ denotes the ball centered at x with radius ϵ .

(iii) The Regional Fractional Laplacian

The Regional Fractional Laplacian is defined in a similar manner to (1.1) but where the integration is restricted to Ω (i.e., $\mathbb{R}^d \setminus B(x, \epsilon)$ replaced by $\Omega \setminus B(x, \epsilon)$).

Another approach to defining the Fractional Laplacian, introduced by Caffarelli and Silvestre [4], is to consider the Fractional Laplacian as the limit of a local operator problem defined on a semi-infinite cylinder $\Omega \times (0, \infty)$.

Two recent papers which give insight into the relationship between various Fractional Laplace operators and their properties are [10, 13].

Related to the Integral Fractional Laplacian is the Riesz Potential operator, which is defined for $\alpha \in (0, d)$ by [6]

$$(-\Delta)^{-\frac{\alpha}{2}} u(x) = \frac{1}{\gamma_d(\alpha)} \int_{\mathbb{R}^d} \frac{u(x-y)}{|y|^{d-\alpha}} dy.$$

For suitably nice functions (see [6, Proposition 1]) the Integral Fractional Laplacian is the inverse of the Riesz Potential operator.

In many physical applications "motion" occurs as a result of a local imbalance. For example, Fourier's Law and Fick's Law of Diffusion state that the local flux is proportional to the gradient of the underlying quantity. For usual diffusion the impact of the local imbalance is local. In the case of anomalous diffusion the impact of the local imbalance is nonlocal. For the usual diffusion of the quantity w, the local flux in \mathbb{R}^d is modeled as $K(x)\nabla w(x)$, and the spatial diffusion operator as $\nabla \cdot K(x)\nabla w(x)$, where $K(x) \in \mathbb{R}^{d \times d}$ is a material parameter. (In the heat equation K(x) represents the thermal conductivity of the medium.)

In order to include a material parameter into an anomalous diffusion model, consider (for $\mathbf{q}(x)$ denoting flux)

usual diffusion

$$\nabla \cdot \mathbf{q}(x) = -\nabla \cdot K(x) \nabla w(x) \qquad \qquad \nabla \cdot \mathbf{q}(x) = -\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} K(x) \nabla w(x). \qquad (1.2)$$

For I the identity matrix in $\mathbb{R}^{d \times d}$ and $k \in \mathbb{R}$, for f a sufficiently nice function, a Fourier transform argument shows that

$$k \left(-\Delta\right)^{\frac{\alpha}{2}} f = -\nabla \cdot \left(-\Delta\right)^{\frac{\alpha-2}{2}} k \mathbf{I} \nabla f.$$
(1.3)

The motivation for the work presented herein was to investigate a nonlocal fractional Laplace operator involving a material parameter K(x). As a first step in this direction, we consider K(x) to be a constant coefficient, symmetric, positive definite matrix, and the domain Ω to be the unit disk in \mathbb{R}^2 . More specifically, by the symmetry of the domain Ω , without loss of generality, we consider K(x) to be a diagonal matrix with positive entries. A corollary of our investigations (see Corollary 4.1) shows that (1.3) holds for a family of functions having an edge singular behavior consistent with that expected of the solution of the Integral Fractional Laplacian.

The problem investigated in this paper is: Given f(x), $K(x) = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix}$, $k_1, k_2 \in \mathbb{R}^+$, determine $\widetilde{u}(x) = (1 - r^2)^{\frac{\alpha}{2}}_+ u(x)$ satisfying

$$\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} K(x) \nabla \widetilde{u}(x) = f(x), \quad x \in \Omega,$$
(1.4)

$$\widetilde{u}(x) = 0, \quad x \in \mathbb{R}^2 \setminus \Omega.$$
 (1.5)

Note that (1.4) is different from the Riesz space fractional operator defined by (see [17])

$$-k_1 \frac{\partial^{\alpha} u}{\partial |x_1|^{\alpha}} - k_2 \frac{\partial^{\alpha} u}{\partial |x_2|^{\alpha}} = f(x_1, x_2), \quad (x_1, x_2) \in \Omega.$$

This difference is apparent when noting that $(-\Delta)^{\frac{\alpha-2}{2}}$ · is not a one dimensional fractional integral operator. For $\Omega \subset \mathbb{R}^1$ a detailed analysis of (1.4)–(1.5) is given in [18].

Our analysis of (1.4)–(1.5) builds upon a result proven by Dyda et al. in [6]. For $L^2_{\beta}(\Omega)$ denoting the weighted L^2 space with domain Ω , an orthogonal basis for $L^2_{\frac{\alpha}{2}}(\Omega)$ is $\left\{V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2-1)\right\}$, for $l, n = 0, 1, 2, \ldots, \infty, \mu \in \{-1, 1\}$, where $V_{l,\mu}(x)$ denotes the solid harmonic polynomials in \mathbb{R}^2 , and $P_n^{(a,b)}(t)$ the Jacobi polynomials (see Section 2). From [6, Theorem 3],

$$(-\Delta)^{\frac{\alpha}{2}} (1-r^2)^{\frac{\alpha}{2}}_+ V_{l,\mu}(x) P_n^{\frac{\alpha}{2},l}(2r^2-1) = \frac{2^{\alpha} \Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_n^{\frac{\alpha}{2},l}(2r^2-1).$$
(1.6)

A similar result to (1.6) in the \mathbb{R}^1 setting is now well known and has been used to investigate the existence, uniqueness and regularity of fractional order differential equations, as well as to develop spectral approximation methods for such problems [7, 9, 14, 15, 18].

In our analysis herein we show:

- 1. A similar result to (1.6) holds for the Riesz Potential operator, $(-\Delta)^{\frac{\alpha-2}{2}}$ (see Theorem 2.1).
- 2. For $u \in L^2_{\frac{\alpha}{2}}(\Omega)$, then $\nabla(1-r^2)^{\frac{\alpha}{2}}u \in (1-r^2)^{\frac{\alpha}{2}-1} \begin{bmatrix} \operatorname{span}\left\{V_{l,\mu}(x)P_n^{\frac{\alpha}{2}-1,l}(2r^2-1)\right\} \\ \operatorname{span}\left\{V_{l,\mu}(x)P_n^{\frac{\alpha}{2}-1,l}(2r^2-1)\right\} \end{bmatrix}$ (see

Theorem 3.1). Notable is that this mapping holds when then the gradient is expressed in

terms of the Cartesian coordinate system (i.e., in terms of \mathbf{i} and \mathbf{j}), and not when written in the polar system (i.e., in terms of \mathbf{r} and φ).

3. (i) For $\tilde{u}(x) = (1 - r^2)_+^{\frac{\alpha}{2}} u(x)$, and u(x) expressed as a linear combination of the basis functions $\left\{ V_{l,\mu}(x) P_n^{\frac{\alpha}{2},l}(2r^2 - 1) \right\}$, substituting into (1.4) and equating coefficients with the f(x) (similarly expressed in terms of the basis), results in an infinite system of coupled equations for the unknown coefficients (see (4.8)-(4.19)).

(ii) When the equations are appropriately ordered and rescaled, and the unknown coefficients renumbered and rescaled the infinite system of equations decouple into tridiagonal block matrices of sizes $1, 1, 2, 2, \ldots, n, n, \ldots$ where the block matrices are uniformly invertible (see (4.31)-(4.33)).

- 4. The Integral Fractional Laplacian can be factored into a product of the divergence operator, a Riesz Potential operator and the gradient operator (see Corollary 4.1). (Note that this result has been proven for the one-dimensional case (see, for example [3, 12]), while, to the best of our knowledge, is not available in the literature for the two-dimensional operator.)
- 5. The dependence of the regularity of the solution of (1.4),(1.5) on the regularity of the RHS, f(x).

This paper is organized as follows. In the next section we introduce the Jacobi polynomials and the solid harmonic polynomials. The combination of these polynomials form a basis for the weighted $L^2_{\beta}(\Omega)$ space, which plays a key role in our analysis. The needed properties of $\nabla \cdot$ and $(-\Delta)^{\frac{\alpha-2}{2}}$. applied to functions $(1-r^2)^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2-1)$ are derived in Section 3. Using these results, in Section 4 we establish the existence and uniqueness of solution, \tilde{u} , to (1.4), (1.5). The dependence of the regularity of \tilde{u} on the regularity of the RHS, f(x), is given in Section 5. Contained in the Appendix are some ancillary results used in the analysis.

2 Preliminaries

In this section we present definitions and notation used in this article. We follow the definitions and notation for the hypergeometric function used by Dyda, Kuznetsov and Kwaśnicki in [6].

2.1 Hypergeometric Function

For p, q nonnegative integers with $p \leq q + 1$, and $\mathbf{a} = (a_1, \ldots, a_p) \in \mathbb{C}^p$, $\mathbf{b} = (b_1, \ldots, b_q) \in \mathbb{C}^q$, with none of the b_j nonpositive integers, the generalized hypergeometric function is defined by

$${}_{p}F_{q}\left(\begin{array}{c} \mathbf{a} \\ \mathbf{b} \end{array} \middle| r\right) := \sum_{n=0}^{\infty} \frac{\prod\limits_{j=1}^{p} a_{j}^{(n)}}{\prod\limits_{j=1}^{q} b_{j}^{(n)}} \frac{r^{n}}{n!}, \qquad (2.1)$$

where $c^{(n)} = c(c+1)\cdots(c+n-1)$ denoting the rising factorial, with $c^{(0)} = 1$.

Note that ${}_{p}F_{q}\begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix}$ is invariant under the permutation of the components of \mathbf{a} or the components of \mathbf{b} .

The regularized hypergeometric function, ${}_{p}\mathbf{F}_{q}(\cdot)$ is defined as

$${}_{p}\mathbf{F}_{q}\left(\begin{array}{c}\mathbf{a}\\\mathbf{b}\end{array}\middle|r\right) := \frac{1}{\prod\limits_{j=1}^{q}\Gamma(b_{j})}{}_{p}F_{q}\left(\begin{array}{c}\mathbf{a}\\\mathbf{b}\end{array}\middle|r\right)$$
(2.2)

2.2 Jacobi Polynomials

The Jacobi polynomials are defined as

$$P_n^{(a,b)}(t) := \frac{\Gamma(a+1+n)}{n!} \, {}_2\mathbf{F}_1\left(\begin{array}{c} -n, \ 1+a+b+n \\ a+1 \end{array} \middle| \frac{1-t}{2} \right) \tag{2.3}$$

$$= \frac{(-1)^{n} \Gamma(b+1+n)}{n!} {}_{2}\mathbf{F}_{1} \left(\begin{array}{c} -n, \ 1+a+b+n \\ b+1 \end{array} \middle| \frac{1+t}{2} \right)$$
(2.4)

$$= (-1)^{n} \frac{\Gamma(n+1+b)}{n! \,\Gamma(n+1+a+b)} \sum_{j=0}^{n} (-1)^{j} \, 2^{-j} \, \binom{n}{j} \frac{\Gamma(n+j+1+a+b)}{\Gamma(j+1+b)} \, (1+t)^{j} \,.$$
(2.5)

Note that $P_0^{(a,b)}(t) = 1$.

In case a, b > -1 the Jacobi polynomials satisfy the following orthogonality property.

$$\int_{-1}^{1} (1-t)^{a} (1+t)^{b} P_{j}^{(a,b)}(t) P_{k}^{(a,b)}(t) dt = \begin{cases} 0, & k \neq j \\ ||P_{j}^{(a,b)}||^{2}, & k = j \end{cases},$$

where $||P_{j}^{(a,b)}||| = \left(\frac{2^{(a+b+1)}}{(2j+a+b+1)} \frac{\Gamma(j+a+1)\Gamma(j+b+1)}{\Gamma(j+1)\Gamma(j+a+b+1)}\right)^{1/2}.$ (2.6)

We have the following differentiation properties

$$\frac{d^{j}}{dt^{j}}P_{k}^{(a,b)}(t) = \frac{\Gamma(a+b+k+1+j)}{2^{j}\Gamma(a+b+k+1)}P_{k-j}^{(a+j,b+j)}(t), \qquad (2.7)$$

$$\frac{d}{dt}\left((1-t)^{\alpha/2}P_k^{(\frac{\alpha}{2},m)}(t)\right) = -(k+\frac{\alpha}{2})(1-t)^{\frac{\alpha}{2}-1}P_k^{(\frac{\alpha}{2}-1,m+1)}(t) \quad (\text{see } [8, (E.1)]), \qquad (2.8)$$

$$\frac{d}{dt}\left((1+t)^m P_k^{(\frac{\alpha}{2},m)}(t)\right) = (k+m)(1+t)^{m-1} P_k^{(\frac{\alpha}{2}+1,m-1)}(t).$$
(2.9)

2.3 Solid Harmonic Polynomials

The solid harmonic polynomials in \mathbb{R}^d are the polynomials in d variables which satisfy Laplace's equation.

In \mathbb{R}^2 the first few (linearly independent) solid harmonic polynomials are:

1,
$$x_1$$
, x_2 , x_1x_2 , $x_1^2 - x_2^2$,...

In \mathbb{R}^3 the first few (linearly independent) solid harmonic polynomials are:

1,
$$x_1$$
, x_2 , x_3 , x_1x_2 , x_1x_3 , x_2x_3 , $x_1^2 - x_2^2$, $x_1^2 - x_3^2$, ...

The solid harmonic polynomials of degree $l \ge 0$ form a finite dimensional vector space, having dimension

$$M_{d,l} := \frac{d+2l-2}{d+l-2} \begin{pmatrix} d+l-2\\ l \end{pmatrix}$$

In \mathbb{R}^2 the solid harmonic polynomials of degree l can be conveniently written in polar coordinates, $((r, \varphi) : 0 \le r < \infty, 0 \le \varphi < 2\pi)$, as $\{r^l \cos(l\varphi), r^l \sin(l\varphi)\}$. Note that $M_{2,l} = 2$.

2.4 Function Spaces

The (weighted) $L^2(\Omega)$ space plays a central role in the analysis. For weight function $\beta(\cdot)$, $\beta(x) > 0$, $x \in \Omega$, associated with $L^2_{\beta}(\Omega)$ we have the inner product and norm

$$(f,g)_{\beta} := \int_{\Omega} \beta(x) f(x) g(x) d\Omega, \quad \|f\|_{L^{2}_{\beta}(\Omega)} := (f,f)_{\beta}^{1/2}.$$

In \mathbb{R}^d , d = 2, 3, $\Omega = \{x : ||x|| < 1\}$, i.e., the unit ball, a basis for $L^2_{\beta}(\Omega)$ is given as a product of the solid harmonic polynomials and Jacobi polynomials.

For $x = (r, \varphi) \in \Omega \subset \mathbb{R}^2$, let $\omega^{\gamma} := (1 - r^2)^{\gamma}$.

2.4.1 In \mathbb{R}^2

Let
$$V_{l,1}(x) := r^l \cos(l\varphi), \quad l = 0, 1, 2, \dots$$
 and $V_{l,-1}(x) := r^l \sin(l\varphi), \quad l = 1, 2, \dots$

We also use the following notation

$$V_{l,\mu^*}(x) = \begin{cases} V_{l,-1}(x) & \text{if } \mu = 1, \\ V_{l,1}(x) & \text{if } \mu = -1. \end{cases}$$

Additionally, for a linear operator \mathcal{F}_{\cdot} , we use $\mathcal{F}(V_{l,\mu}(x)) = (\pm)V_{l,\sigma}(x)$ to denote

$$\mathcal{F}(V_{l,\mu}(x)) = (\pm)V_{l,\sigma}(x) = \begin{cases} +V_{l,\sigma}(x) & \text{if } \mu = 1, \\ -V_{l,\sigma}(x) & \text{if } \mu = -1. \end{cases}$$

For example,

$$\frac{\partial}{\partial \varphi} V_{l,\mu}(x) = (\mp) V_{l,\mu^*}(x)$$

In \mathbb{R}^2 an orthogonal basis for $L^2_{\beta}(\Omega)$ is [5, 11]

$$\left\{ \bigcup_{l=0}^{\infty} \bigcup_{n=0}^{\infty} \left\{ V_{l,1}(x) P_n^{(\beta,l)}(2r^2 - 1) \right\} \right\} \cup \left\{ \bigcup_{l=1}^{\infty} \bigcup_{n=0}^{\infty} \left\{ V_{l,-1}(x) P_n^{(\beta,l)}(2r^2 - 1) \right\} \right\}.$$
 (2.10)

For notation brevity we denote the basis in (2.10) as

$$\bigcup_{l=0}^{\infty} \bigcup_{n=0}^{\infty} \bigcup_{\mu=1,-1} \left\{ V_{l,\mu}(x) P_n^{(\beta,l)}(2r^2 - 1) \right\}$$

where we implicit assume that the terms $V_{0,-1}(x) P_n^{(\beta,l)}(2r^2 - 1)$, n = 0, 1, ... are omitted from the set.

2.4.2 Action of the Riesz potential operator

The following theorem, used in our analysis below, extends a result in [6] from the fractional Laplace operator to the Riesz potential operator.

Theorem 2.1 For $\delta = d + 2l$, s an integer, $\frac{\alpha}{2} - s > -1$, $f(x) = (1 - |x|^2)_+^{\frac{\alpha}{2} - s} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2} - s, \frac{\delta}{2} - 1)}(2|x|^2 - 1)$,

$$(-\Delta)^{\frac{\alpha-2}{2}} f(x) = (-1)^{1-s} 2^{\alpha-2} \frac{\Gamma(n+1-s+\frac{\alpha}{2})\Gamma(n-1+\frac{\delta+\alpha}{2})}{\Gamma(n+1)\Gamma(n+1-s+\frac{\delta}{2})} V_{l,\mu}(x) P_{n+1-s}^{(\frac{\alpha}{2}-2+s,\frac{\delta}{2}-1)}(2|x|^2-1).$$
(2.11)

The proof of Theorem 2.1 begins by transforming f(x), via a hypergeometric function, to a product of $V_{l,\mu}(x)$ and a Meijer G-function. The operator $(-\Delta)^{\frac{\alpha-2}{2}}$ is then applied and the subsequent results transformed using properties of the Meijer G-function and the hypergeometric function to (2.11). As this is the only place in the paper where hypergeometric functions and Meijer G-functions are used we present the proof of Theorem 2.1 in the Appendix.

3 Mapping properties of $\nabla \cdot$ and $(-\Delta)^{\frac{\alpha-2}{2}}$

In this section we investigate the mapping properties of $\nabla \cdot$ and $(-\Delta)^{\frac{\alpha-2}{2}}$ with the domain $\omega^{\frac{\alpha}{2}} \otimes L^2_{\frac{\alpha}{2}}(\Omega)$. Functions $g \in L^2_{\frac{\alpha}{2}}(\Omega)$ are conveniently represented using polar coordinates but, surprisingly, the mapping properties of $\nabla \cdot$ and $(-\Delta)^{\frac{\alpha-2}{2}}$ are far more revealing when written in a Cartesian framework (i.e., in terms of $\frac{\partial g}{\partial x}$ and $\frac{\partial g}{\partial y}$. To obtain this representation we firstly compute $\frac{\partial g}{\partial r}$ and $\frac{\partial g}{\partial \varphi}$, and then take an appropriate combination of these terms to obtain $\frac{\partial g}{\partial x}$ and $\frac{\partial g}{\partial y}$.

Theorem 3.1 Let $f(x) = (1 - r^2)^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1)$. Then for $n \ge 0$ and $l \ge 1$,

$$\frac{\partial f}{\partial x} = -(n + \frac{\alpha}{2})(1 - r^2)^{\frac{\alpha}{2} - 1} V_{l+1,\mu}(x) P_n^{(\frac{\alpha}{2} - 1, l+1)}(2r^2 - 1)
- (n+1)(1 - r^2)^{\frac{\alpha}{2} - 1} V_{l-1,\mu}(x) P_{n+1}^{(\frac{\alpha}{2} - 1, l-1)}(2r^2 - 1),$$

$$\frac{\partial f}{\partial y} = -(\pm)(n + \frac{\alpha}{2})(1 - r^2)^{\frac{\alpha}{2} - 1} V_{l+1,\mu^*}(x) P_n^{(\frac{\alpha}{2} - 1, l+1)}(2r^2 - 1)$$
(3.1)

$$- (\mp)(n+1)(1-r^2)^{\frac{\alpha}{2}-1}V_{l-1,\mu^*}(x)P_{n+1}^{(\frac{\alpha}{2}-1,l-1)}(2r^2-1).$$
(3.2)

For the special case l = 0 and $\mu = 1$, we have

$$\frac{\partial f}{\partial x} = -2(n+\frac{\alpha}{2})(1-r^2)^{\frac{\alpha}{2}-1}V_{1,1}(x)P_n^{(\frac{\alpha}{2}-1,1)}(2r^2-1), \qquad (3.3)$$

$$\frac{\partial f}{\partial y} = -2(n+\frac{\alpha}{2})(1-r^2)^{\frac{\alpha}{2}-1}V_{1,-1}(x)P_n^{(\frac{\alpha}{2}-1,1)}(2r^2-1).$$
(3.4)

The proof of Theorem 3.1 is relatively long, using several of the recurrence formulas for Jacobi polynomials. So as not to unnecessarily distract from the existence and uniqueness of solution result for (1.4), (1.5), we give the proof of Theorem 3.1 in the Appendix.

Theorem 3.2 Let
$$f(x) = (1 - r^2)^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1)$$
. Then for $n \ge 0$ and $l \ge 1$
 $(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial f}{\partial x} = C_2 \left(n + \frac{\alpha}{2} + l\right) V_{l+1,\mu}(x) P_n^{(\frac{\alpha}{2} - 1, l+1)}(2r^2 - 1)$
 $+ C_2 \left(n + l + 1\right) V_{l-1,\mu}(x) P_{n+1}^{(\frac{\alpha}{2} - 1, l-1)}(2r^2 - 1),$ (3.5)
 $(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial f}{\partial y} = C_2 (\pm) \left(n + \frac{\alpha}{2} + l\right) V_{l+1,\mu^*}(x) P_n^{(\frac{\alpha}{2} - 1, l+1)}(2r^2 - 1)$

$$= C_2(\pm)(n + \frac{1}{2} + l) V_{l+1,\mu^*}(x) I_n^{(\alpha)} (2l - 1) + C_2(\mp)(n + l + 1) V_{l-1,\mu^*}(x) P_{n+1}^{(\frac{\alpha}{2} - 1, l - 1)}(2r^2 - 1),$$
(3.6)

where
$$C_2 = -2^{\alpha - 2} \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + \frac{\alpha}{2} + l)}{\Gamma(n + 1) \Gamma(n + 2 + l)}$$
. (3.7)

For the special case l = 0 and $\mu = 1$, we have

$$(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial f}{\partial x} = 2C_2 \left(n + \frac{\alpha}{2}\right) V_{1,1}(x) P_n^{(\frac{\alpha}{2} - 1, 1)} (2r^2 - 1), \tag{3.8}$$

$$(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial f}{\partial y} = 2C_2 \left(n + \frac{\alpha}{2}\right) V_{1,-1}(x) P_n^{\left(\frac{\alpha}{2} - 1, 1\right)} (2r^2 - 1).$$
(3.9)

Proof: The stated results follow from applying Theorem 2.1 to the four relations in Theorem 3.1.

Corollary 3.1 Let $f(x) = V_{l,\mu}(x)P_n^{(\gamma,l)}(2r^2-1)$. Then for $n \ge 1$ and $l \ge 1$, $\frac{\partial f}{\partial x} = (n+l)V_{l-1,\mu}(x)P_n^{(\gamma+1,l-1)}(2r^2-1) + (n+\gamma+l+1)V_{l+1,\mu}(x)P_{n-1}^{(\gamma+1,l+1)}(2r^2-1)$, (3.10)

and

$$\frac{\partial f}{\partial y} = (\mp)(n+l) V_{l-1,\mu^*}(x) P_n^{(\gamma+1,l-1)}(2r^2-1) + (\pm)(n+\gamma+l+1) V_{l+1,\mu^*}(x) P_{n-1}^{(\gamma+1,l+1)}(2r^2-1).$$
(3.11)

For n = 0 and $l \ge 1$,

$$\frac{\partial f}{\partial x} = l V_{l-1,\mu}(x) P_0^{(\gamma,l)}(2r^2 - 1) \quad and \quad \frac{\partial f}{\partial y} = (\mp) l V_{l-1,\mu^*}(x) P_0^{(\gamma,l)}(2r^2 - 1).$$

For $n \geq 1$ and l = 0, $\mu = 1$

$$\frac{\partial f}{\partial x} = 2(n+\gamma+1) V_{1,1}(x) P_{n-1}^{(\gamma+1,1)}(2r^2-1) \quad and \quad \frac{\partial f}{\partial y} = 2(n+\gamma+1) V_{1,-1}(x) P_{n-1}^{(\gamma+1,1)}(2r^2-1) .$$

For $n = 0$ and $l = 0, \ \mu = 1$

$$\frac{\partial f}{\partial x} = 0 \quad and \quad \frac{\partial f}{\partial y} = 0.$$

Proof: Using (2.7) and $\frac{\partial f}{\partial x} = \cos(\varphi) \frac{\partial f}{\partial r} - \sin(\varphi) \frac{1}{r} \frac{\partial f}{\partial \varphi}$,

$$\frac{\partial f}{\partial x} = \cos(\varphi) \left(l V_{l,\mu}(x) r^{-1} P_n^{(\gamma,l)}(2r^2 - 1) + 2(n + \gamma + l + 1) V_{l,\mu}(x) r P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) \right) - \sin(\varphi) \left(\left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x) \right) r^{-1} P_n^{(\gamma,l)}(2r^2 - 1) \right).$$

Then, using (C.5), (C.6) and (C.7), (C.8),

$$\begin{aligned} \frac{\partial f}{\partial x} &= l V_{l-1,\mu}(x) P_n^{(\gamma,l)}(2r^2 - 1) + (n + \gamma + l + 1) V_{l-1,\mu}(x) r^2 P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) \\ &+ (n + \gamma + l + 1) V_{l+1,\mu}(x) P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) \\ &= (n+l) V_{l-1,\mu}(x) P_n^{(\gamma+1,l-1)}(2r^2 - 1) + (n + \gamma + l + 1) V_{l+1,\mu}(x) P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) \end{aligned}$$

where, in the last step, we have used (A.5).

Next, using
$$\frac{\partial f}{\partial y} = \sin(\varphi) \frac{\partial f}{\partial r} + \cos(\varphi) \frac{1}{r} \frac{\partial f}{\partial \varphi},$$

$$\frac{\partial f}{\partial y} = \sin(\varphi) \left(l V_{l,\mu}(x) r^{-1} P_n^{(\gamma,l)}(2r^2 - 1) + 2(n + \gamma + l + 1) V_{l,\mu}(x) r P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) \right)$$

$$+ \cos(\varphi) \left(\left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x) \right) r^{-1} P_n^{(\gamma,l)}(2r^2 - 1) \right).$$

Using (C.12), (C.13) and (C.14), (C.15),

$$\frac{\partial f}{\partial y} = l(\mp)V_{l-1,\mu^*}(x) P_n^{(\gamma,l)}(2r^2 - 1) + (n + \gamma + l + 1)(\mp)V_{l-1,\mu^*}(x) r^2 P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1) + (n + \gamma + l + 1)(\pm)V_{l+1,\mu^*}(x) P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1)$$

$$= (\mp)(n+l) V_{l-1,\mu^*}(x) P_n^{(\gamma+1,l-1)}(2r^2 - 1) + (\pm)(n + \gamma + l + 1) V_{l+1,\mu^*}(x) P_{n-1}^{(\gamma+1,l+1)}(2r^2 - 1)$$

where, again in the last step we have used (A.5).

The proofs of the other statements follow in a similar manner and are therefore omitted for brevity.

4 Existence and Uniqueness of Solution

In this section we investigate the existence and uniqueness of solution to the following problem. **Problem**:

Given f and $k_1, k_2 \in \mathbb{R}^+$, with $K(x) = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix}$, find $\widetilde{u} = \omega^{\frac{\alpha}{2}} u$ such that

$$\mathcal{L}(\widetilde{u})(x) := -\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} K(x) \nabla \widetilde{u}(x) = f(x), \quad x \in \Omega,$$
(4.1)

subject to
$$\widetilde{u} = 0$$
 on $\mathbb{R}^2 \setminus \Omega$. (4.2)

We assume u(x) can be expressed as

$$u(x) = \sum_{l \ge 1, n \ge 0, \mu \in \{1, -1\}} u_{l,n,\mu} V_{l,\mu}(x) P_n^{\left(\frac{\alpha}{2}, l\right)}(2r^2 - 1) + \sum_{n \ge 0} \frac{u_{0,n,1}}{2} V_{0,1}(x) P_n^{\left(\frac{\alpha}{2}, 0\right)}(2r^2 - 1)$$
(4.3)

and

$$f(x) = \sum_{l \ge 1, n \ge 0, \mu \in \{1, -1\}} f_{l,n,\mu} V_{l,\mu}(x) P_n^{\left(\frac{\alpha}{2}, l\right)}(2r^2 - 1) + \sum_{n \ge 0} f_{0,n,1} V_{0,1}(x) P_n^{\left(\frac{\alpha}{2}, 0\right)}(2r^2 - 1)$$

for coefficients $u_{l,n,\mu}$, $f_{l,n,\mu} \in \mathbb{R}$. (Here we divide the coefficients $\{u_{0,n,1}\}_{n\geq 0}$ by 2 in order that the matrices $A_{3,4,m}$ and $A_{3,5,m}$ in (4.32) and (4.33) are symmetric. Note that this factor of 2 results from the case of l = 0, $\mu = 1$ in Theorems 3.1 and 3.2.)

Using Theorem 3.2 and Corollary 3.1, $\mathcal{L}(\tilde{u})$ can be expressed in terms of the basis $\left\{V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2-1)\right\}_{n,l,\mu}$. Equating the coefficients of $\mathcal{L}(\tilde{u})$ with those of f we obtain a linear system of equations for the unknown coefficients $u_{l,n,\mu}$.

Using Theorem 3.2, $k_1(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial \tilde{u}}{\partial x} = T_1 + T_2$, where $T_1 = -2^{\alpha-2} k_1 \sum_{\substack{n \ge 0, \ k = 1 \\ l \ge 1, \mu = -1}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+2+l)} V_{l+1,\mu}(x) P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1)$ $= -2^{\alpha-2} k_1 \sum_{\substack{n \ge 0, \ k = 1, \mu = 1 \\ l \ge 2, \mu = -1}} u_{l-1,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2}-1,l)}(2r^2-1),$ and

$$T_{2} = -2^{\alpha-2} k_{1} \sum_{\substack{n \ge 0, \left\{ \begin{array}{l} l \ge 1, \mu = 1\\ l \ge 2, \mu = -1 \end{array} \\ n \ge 1, \left\{ \begin{array}{l} l \ge 0, \mu = 1\\ l \ge 1, \mu = -1 \end{array} \right.} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} V_{l-1,\mu}(x) P_{n+1}^{\left(\frac{\alpha}{2}-1, l-1\right)}(2r^{2}-1) \\ u_{l+1,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n)\Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2}-1, l\right)}(2r^{2}-1) \end{array}$$

Using Corollary 3.1,

$$\begin{aligned} \frac{\partial}{\partial x}T_{1} &= -2^{\alpha-2}k_{1}\sum_{\substack{n=0, \left\{\begin{array}{l}l\geq 1, \mu=1\\l\geq 2, \mu=-1\end{array}}} u_{l-1,0,\mu}\frac{\Gamma(0+1+\frac{\alpha}{2})\Gamma(0+\frac{\alpha}{2}+l)}{\Gamma(0+1)\Gamma(0+1+l)} l V_{l-1,\mu}(x) P_{0}^{(\frac{\alpha}{2}-1,l)}(2r^{2}-1) \right.} \\ &- 2^{\alpha-2}k_{1}\sum_{\substack{n\geq 1, \left\{\begin{array}{l}l\geq 1, \mu=1\\l\geq 2, \mu=-1\end{aligned}}} u_{l-1,n,\mu}\frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} \left((n+l)V_{l-1,\mu}(x) P_{n}^{(\frac{\alpha}{2},l-1)}(2r^{2}-1)\right) \\ &+ (n+\frac{\alpha}{2}+l)V_{l+1,\mu}(x) P_{n-1}^{(\frac{\alpha}{2},l+1)}(2r^{2}-1)\right). \end{aligned}$$

Recalling that $P_0^{(a,b)}(t) = 1$, we obtain

$$\frac{\partial}{\partial x}T_{1} = -2^{\alpha-2}k_{1}\sum_{\substack{n=0, \left\{\begin{array}{l}l\geq 0, \mu=1\\l\geq 1, \mu=-1\end{array}}}u_{l,0,\mu}\frac{\Gamma(1+\frac{\alpha}{2})\Gamma(1+\frac{\alpha}{2}+l)}{\Gamma(1)\Gamma(1+l)}V_{l,\mu}(x)P_{0}^{(\frac{\alpha}{2},l)}(2r^{2}-1)\right.}\\ - 2^{\alpha-2}k_{1}\sum_{\substack{n\geq 1, \left\{\begin{array}{l}l\geq 0, \mu=1\\l\geq 1, \mu=-1\end{array}}}u_{l,n,\mu}\frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)}V_{l,\mu}(x)P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1)\\ - 2^{\alpha-2}k_{1}\sum_{\substack{n\geq 0, \left\{\begin{array}{l}l\geq 2, \mu=1\\l\geq 3, \mu=-1\end{array}}}u_{l-2,n+1,\mu}\frac{\Gamma(n+2+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2)\Gamma(n+1+l)}V_{l,\mu}(x)P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1).\end{array}\right.}\right.}$$

$$(4.4)$$

Next,

$$\begin{split} \frac{\partial}{\partial x} T_{2} &= -2^{\alpha-2} k_{1} \sum_{n \geq 1, l=0, \mu=1} u_{1,n-1,1} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+\frac{\alpha}{2})}{\Gamma(n) \Gamma(n+1)} 2(n+\frac{\alpha}{2}) V_{1,1}(x) P_{n-1}^{(\frac{\alpha}{2},1)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 1, \left\{ \begin{array}{c} l \geq 1, \mu=1\\ l \geq 1, \mu=-1 \end{array}} u_{l+1,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n) \Gamma(n+1+l)} \left((n+l) V_{l-1,\mu}(x) P_{n}^{(\frac{\alpha}{2},l-1)}(2r^{2}-1)\right) \\ &+ (n+\frac{\alpha}{2}+l) V_{l+1,\mu}(x) P_{n-1}^{(\frac{\alpha}{2},l+1)}(2r^{2}-1) \right) \\ &= -2^{\alpha-2} k_{1} \sum_{n \geq 0, l=1, \mu=1} u_{1,n,1} \frac{2\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+2+\frac{\alpha}{2})}{\Gamma(n+1) \Gamma(n+2)} V_{1,1}(x) P_{n}^{(\frac{\alpha}{2},1)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 1, \left\{ \begin{array}{c} l \geq 0, \mu=1\\ l \geq 1, \mu=-1 \end{array}} u_{l+2,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 1, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{(\frac{\alpha}{2},l)}(2r^{2}-1) \\ &- 2^{\alpha-2} k_{1} \sum_{n \geq 0, \left\{ \begin{array}{c} l \geq 2, \mu=-1\\ l \geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1+\frac{\alpha}{2}+l)} U_{l,\mu}(2r^{2}-1) \\$$

Again, using Theorem 3.2, $k_2(-\Delta)^{\frac{\alpha-2}{2}} \frac{\partial \tilde{u}}{\partial y} = T_3 + T_4$, where

$$T_{3} = -2^{\alpha-2} k_{2} \sum_{\substack{n \ge 0, \ k = 1 \\ l \ge 1, \ \mu = -1}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+2+l)} (\pm) V_{l+1,\mu^{*}}(x) P_{n}^{(\frac{\alpha}{2}-1,l+1)}(2r^{2}-1)$$

$$= -2^{\alpha-2} k_{2} \sum_{\substack{n \ge 0, \ k = 1 \\ l \ge 2, \ \mu = -1}} u_{l-1,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} (\pm) V_{l,\mu^{*}}(x) P_{n}^{(\frac{\alpha}{2}-1,l)}(2r^{2}-1),$$

and

$$T_{4} = -2^{\alpha-2} k_{2} \sum_{\substack{n \ge 0, \left\{ \begin{array}{l} l \ge 2, \mu = 1\\ l \ge 1, \mu = -1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} (\mp)V_{l-1,\mu^{*}}(x) P_{n+1}^{(\frac{\alpha}{2}-1,l-1)}(2r^{2}-1) \right.} \\ = -2^{\alpha-2} k_{2} \sum_{\substack{n \ge 1, \left\{ \begin{array}{l} l \ge 1, \mu = 1\\ l \ge 0, \mu = -1 \end{array}} u_{l+1,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2})\Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n)\Gamma(n+1+l)} (\mp)V_{l,\mu^{*}}(x) P_{n}^{(\frac{\alpha}{2}-1,l)}(2r^{2}-1) \right.} \end{array}$$

Using Corollary 3.1,

$$\begin{split} \frac{\partial}{\partial y} T_{3} &= -2^{\alpha-2} k_{2} \sum_{\substack{n=0, \left\{ \begin{array}{l} l \geq 1, \mu = 1\\ l \geq 2, \mu = -1 \end{array}}} u_{l-1,0,\mu} \frac{\Gamma(0+1+\frac{\alpha}{2}) \Gamma(0+\frac{\alpha}{2}+l)}{\Gamma(0+1) \Gamma(0+1+l)} l V_{l-1,\mu}(x) P_{0}^{\left(\frac{\alpha}{2}-1,l\right)}(2r^{2}-1) \right.} \\ &- 2^{\alpha-2} k_{2} \sum_{\substack{n \geq 1, \left\{ \begin{array}{l} l \geq 1, \mu = 1\\ l \geq 2, \mu = -1 \end{array}}} u_{l-1,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} \left((n+l) V_{l-1,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l-1\right)}(2r^{2}-1) \right.} \\ &- \left(n+\frac{\alpha}{2}+l\right) V_{l+1,\mu}(x) P_{n-1}^{\left(\frac{\alpha}{2},l-1\right)}(2r^{2}-1) \right) \\ &= -2^{\alpha-2} k_{2} \sum_{\substack{n=0, \left\{ \begin{array}{l} l \geq 0, \mu = 1\\ l \geq 1, \mu = -1 \end{array}}} u_{l,0,\mu} \frac{\Gamma(1+\frac{\alpha}{2}) \Gamma(1+\frac{\alpha}{2}+l)}{\Gamma(1) \Gamma(1+l)} V_{l,\mu}(x) P_{0}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \right.} \\ &- 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 1, \left\{ \begin{array}{l} l \geq 0, \mu = 1\\ l \geq 1, \mu = -1 \end{array}}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 1, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 3, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 3, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 3, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 3, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 3, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ \end{array} \right)} \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 2, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} V_{l,\mu}(x) P_{n}^{\left(\frac{\alpha}{2},l\right)}(2r^{2}-1) \\ \end{array} \right)} \\ &+ 2^{\alpha-2} k_{2} \sum_{\substack{n\geq 0, \left\{ \begin{array}{l} l \geq 2, \mu = 1\\ l \geq 2, \mu = -1 \end{array}}} u_{l,2,n+1,\mu} \frac{\Gamma(n+2+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+2) \Gamma(n+1+l)} U_{l,\mu$$

Next,

$$\begin{split} \frac{\partial}{\partial y} T_4 &= -2^{\alpha-2} k_2 \sum_{n\geq 1, l=0, \mu=-1} u_{1,n-1,-1} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+\frac{\alpha}{2})}{\Gamma(n) \Gamma(n+1)} (+) 2(n+\frac{\alpha}{2}) V_{1,-1}(x) P_{n-1}^{\left(\frac{\alpha}{2},1\right)}(2r^2-1) \\ &- 2^{\alpha-2} k_2 \sum_{n\geq 1, \left\{ \begin{array}{c} l\geq 1, \mu=1\\ l\geq 1, \mu=-1 \end{array}} u_{l+1,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+\frac{\alpha}{2}+l)}{\Gamma(n) \Gamma(n+1+l)} \left((-)(n+l) V_{l-1,\mu}(x) P_n^{\left(\frac{\alpha}{2},l-1\right)}(2r^2-1) \right) \\ &+ \left(n+\frac{\alpha}{2}+l\right) V_{l+1,\mu}(x) P_{n-1}^{\left(\frac{\alpha}{2},l+1\right)}(2r^2-1) \right) \\ &= -2^{\alpha-2} k_2 \sum_{n\geq 0, l=1, \mu=-1} u_{1,n,-1} \frac{2\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+2+\frac{\alpha}{2})}{\Gamma(n+1) \Gamma(n+2)} V_{1,-1}(x) P_n^{\left(\frac{\alpha}{2},l\right)}(2r^2-1) \\ &+ 2^{\alpha-2} k_2 \sum_{n\geq 1, \left\{ \begin{array}{c} l\geq 0, \mu=1\\ l\geq 0, \mu=-1 \end{array}} u_{l+2,n-1,\mu} \frac{\Gamma(n+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n) \Gamma(n+1+l)} V_{l,\mu}(x) P_n^{\left(\frac{\alpha}{2},l\right)}(2r^2-1) \\ &- 2^{\alpha-2} k_2 \sum_{n\geq 0, \left\{ \begin{array}{c} l\geq 2, \mu=-1\\ l\geq 2, \mu=-1 \end{array}} u_{l,n,\mu} \frac{\Gamma(n+1+\frac{\alpha}{2}) \Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1) \Gamma(n+1+l)} V_{l,\mu}(x) P_n^{\left(\frac{\alpha}{2},l\right)}(2r^2-1) . \end{array} \right) \\ &= (4.7) \end{split}$$

As mentioned at the beginning of this section, we obtain the determining equations for the unknown coefficients $u_{l,n,\mu}$ by equating the coefficients of the basis functions $\left\{V_{l,\mu}(x) P_n^{\left(\frac{\alpha}{2},l\right)}(2r^2-1)\right\}_{n,l,\mu}$ for $\mathcal{L}(\widetilde{u}) = -\nabla \cdot K(x)(-\Delta)^{\frac{\alpha-2}{2}})\nabla \widetilde{u}$ with those of f.

To obtain the determining equations it is convenient to group the associated basis functions with the index sets $\{(n,l)\}_{n\geq 0, l\geq 0}$ for $\mu = 1$, and $\{(n,l)\}_{n\geq 0, l\geq 1}$ for $\mu = -1$, separately. We then partition the index sets into 6 pieces as follows (see Figures 4.1 and 4.2):

For
$$\mu = 1$$
: $R1 = \{(0,0)\};$
 $R2 = \{(0,1)\};$
 $R3 = \{(0,l)\}_{l \ge 2};$
 $R4 = \{(n,0)\}_{n \ge 1};$
 $R5 = \{(n,1)\}_{n \ge 1};$
 $R6 = \{(n,l)\}_{n \ge 1, l \ge 2};$

For
$$\mu = -1$$
: $R1 = \{(0,1)\};$
 $R2 = \{(0,2)\};$
 $R3 = \{(0,l)\}_{l \ge 3};$
 $R4 = \{(n,1)\}_{n \ge 1};$
 $R5 = \{(n,2)\}_{n \ge 1};$
 $R6 = \{(n,l)\}_{n \ge 1, l \ge 3}.$





Figure 4.1: Partition of the index set $\{(n,l)\}_{n>0,l>0}$ for $\mu = 1$ into $R1, R2, \ldots, R6$.



Combining (4.4)-(4.7) and equating with the coefficients of $f(x) = \sum_{n \, l \, \mu} f_{l,n,\mu} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2-1)$ we obtain the following determining system of equations for the unknowns $u_{l,n,\mu}$.

For $\mu = 1$:

$$R1 (n = 0, l = 0): \quad 2^{\alpha - 2} (k_1 + k_2) \frac{\Gamma(1 + \frac{\alpha}{2}) \Gamma(1 + \frac{\alpha}{2})}{\Gamma(1) \Gamma(1)} u_{0,0,1} = f_{0,0,1}$$

$$(4.8)$$

$$R2\left(n=0,l=1\right): \quad 2^{\alpha-2}\left(2k_{1}+\left(k_{1}+k_{2}\right)\right) \frac{\Gamma\left(1+\frac{\alpha}{2}\right)\Gamma\left(2+\frac{\alpha}{2}\right)}{\Gamma(1)\Gamma(2)}u_{1,0,1} = f_{1,0,1} \tag{4.9}$$

$$R3\left\{\left(n=0,l\right)\right\}_{l>2}:$$

$$AS\{(n = 0, l)\}_{l \ge 2}:$$

$$2^{\alpha - 2} \left(2(k_1 + k_2) \frac{\Gamma(1 + \frac{\alpha}{2}) \Gamma(1 + \frac{\alpha}{2} + l)}{\Gamma(1) \Gamma(1 + l)} u_{l,0,1} + (k_1 - k_2) \frac{\Gamma(2 + \frac{\alpha}{2}) \Gamma(1 + \frac{\alpha}{2} + l)}{\Gamma(2) \Gamma(1 + l)} u_{l-2,1,1} \right)$$

$$= f_{l,0,1}$$
(4.10)

$$R4 \{(n, l = 0)\}_{n \ge 1} :$$

$$2^{\alpha - 2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2})}{\Gamma(n) \Gamma(n + 1)} u_{2,n-1,1} + (k_1 + k_2) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2})}{\Gamma(n + 1) \Gamma(n + 1)} u_{0,n,1} \right)$$

$$= f_{0,n,1} \quad (4.11)$$

$$R5 \{(n, l = 1)\}_{n \ge 1}:$$

$$2^{\alpha - 2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 2 + \frac{\alpha}{2})}{\Gamma(n) \Gamma(n + 2)} u_{3,n-1,1} + (2k_1 + (k_1 + k_2)) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 2 + \frac{\alpha}{2})}{\Gamma(n + 1) \Gamma(n + 2)} u_{1,n,1} \right)$$

$$= f_{1,n,1} \qquad (4.12)$$

$$R6 \{(n,l)\}_{n \ge 1, l \ge 2} :$$

$$2^{\alpha-2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n) \Gamma(n + 1 + l)} u_{l+2,n-1,1} + 2(k_1 + k_2) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n + 1) \Gamma(n + 1 + l)} u_{l,n,1} + (k_1 - k_2) \frac{\Gamma(n + 2 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n + 2) \Gamma(n + 1 + l)} u_{l-2,n+1,1} \right) = f_{l,n,1}. \quad (4.13)$$

For $\mu = -1$:

$$R1 (n = 0, l = 1): \quad 2^{\alpha - 2} (2k_2 + (k_1 + k_2)) \frac{\Gamma(1 + \frac{\alpha}{2})\Gamma(2 + \frac{\alpha}{2})}{\Gamma(1)\Gamma(2)} u_{1,0,-1} = f_{1,0,-1}$$
(4.14)

$$R2(n = 0, l = 2): \quad 2^{\alpha - 2} \ 2(k_1 + k_2) \frac{\Gamma(1 + \frac{\alpha}{2})\Gamma(3 + \frac{\alpha}{2})}{\Gamma(1)\Gamma(3)} u_{2,0,-1} = f_{2,0,-1}$$

$$R3\{(n = 0, l)\}_{l > 2}:$$

$$(4.15)$$

$$2^{\alpha-2} \left(2(k_1+k_2) \frac{\Gamma(1+\frac{\alpha}{2})\Gamma(1+\frac{\alpha}{2}+l)}{\Gamma(1)\Gamma(1+l)} u_{l,0,-1} + (k_1-k_2) \frac{\Gamma(2+\frac{\alpha}{2})\Gamma(1+\frac{\alpha}{2}+l)}{\Gamma(2)\Gamma(1+l)} u_{l-2,1,-1} \right) = f_{l,0,-1}$$
(4.16)

$$R4 \{(n, l = 1)\}_{n \ge 1} :$$

$$2^{\alpha - 2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 2 + \frac{\alpha}{2})}{\Gamma(n) \Gamma(n + 2)} u_{3,n-1,-1} + (2k_2 + (k_1 + k_2)) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 2 + \frac{\alpha}{2})}{\Gamma(n + 1) \Gamma(n + 2)} u_{1,n,1} \right)$$

$$= f_{1,n,-1} \quad (4.17)$$

$$R5 \{(n, l = 2)\}_{n \ge 1} :$$

$$2^{\alpha - 2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 3 + \frac{\alpha}{2})}{\Gamma(n) \Gamma(n + 3)} u_{4, n - 1, -1} + 2(k_1 + k_2) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 3 + \frac{\alpha}{2})}{\Gamma(n + 1) \Gamma(n + 3)} u_{2, n, -1} \right)$$

$$= f_{2, n, -1} \qquad (4.18)$$

$$R6 \{(n,l)\}_{n \ge 1,l \ge 3} :$$

$$2^{\alpha-2} \left((k_1 - k_2) \frac{\Gamma(n + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n) \Gamma(n + 1 + l)} u_{l+2,n-1,-1} + 2(k_1 + k_2) \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n+1) \Gamma(n+1+l)} u_{l,n,-1} + (k_1 - k_2) \frac{\Gamma(n + 2 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2} + l)}{\Gamma(n+2) \Gamma(n+1+l)} u_{l-2,n+1,-1} \right) = f_{l,n,-1} \quad (4.19)$$

4.1 Determining the coefficients $u_{l,n,\mu}$

In this section we discuss the solution of $\{u_{l,n,\mu}\}$ satisfying (4.8)-(4.19). We focus our attention on the coefficients $\{u_{l,n,1}\}$, i.e. equations (4.8)-(4.13). The coefficients $\{u_{l,n,-1}\}$ are determined in an analogous manner.

Firstly we introduce a rescaling of the equations. Let,

$$d_{l,n} = \frac{\Gamma(n+1+\frac{\alpha}{2})}{\Gamma(n+1)} u_{l,n,1}, \quad \text{and} \quad \tilde{f}_{l,n} = 2^{-(\alpha-2)} \frac{\Gamma(n+1+l)}{\Gamma(n+1+\frac{\alpha}{2}+l)} f_{l,n,1}.$$
(4.20)

Then, after simplifying, equations (4.8)-(4.13) become:

$$R1 (n = 0, l = 0): (k_1 + k_2) d_{0,0} = f_{0,0}$$
(4.21)

$$R2(n = 0, l = 1): \quad (2k_1 + (k_1 + k_2)) d_{1,0} = \tilde{f}_{1,0}$$

$$\sim$$
(4.22)

$$R3\{(n=0,l)\}_{l\geq 2}: \quad 2(k_1+k_2)d_{l,0} + (k_1-k_2)d_{l-2,1} = f_{l,0}$$

$$(4.23)$$

$$R4\{(n, l=0)\}_{n\geq 1}: \quad (k_1 - k_2) d_{2,n-1} + (k_1 + k_2) d_{0,n} = f_{0,n}$$

$$(4.24)$$

$$R5\{(n,l=1)\}_{n\geq 1}: (k_1-k_2)d_{3,n-1} + (2k_1+(k_1+k_2))d_{1,n} = f_{1,n}$$
(4.25)

$$R6\{(n,l)\}_{n\geq 1,l\geq 2}: (k_1-k_2) d_{l+2,n-1} + 2(k_1+k_2) d_{l,n} + (k_1-k_2) d_{l-2,n+1} = f_{l,n}. \quad (4.26)$$

The coupling stencil.

Consider the "stencil" given in Figure 4.3. This stencil represents the coupling of the unknowns, $d_{l,n}$, in equations (4.21)-(4.26). (For the coupling stencil, when l, n < 0 the $d_{l,n}$ coefficient is ignored.)



Figure 4.3: Stencil illustrating the coupling of the unknowns $d_{l,n}$.

Equations (4.21) and (4.22) represent two 1×1 linear system of equations for $d_{0,0}$ and $d_{0,1}$. From R3, l = 2 and n = 0 gives the equation

$$2(k_1 + k_2) d_{2,0} + (k_1 - k_2) d_{0,1} = \tilde{f}_{2,0}.$$
(4.27)

Following the coupling of the unknowns from (4.27) (cf. coefficient stencil), the equation corresponding to l = 0, n = 1 (in R4) is

$$(k_1 - k_2) d_{2,0} + (k_1 + k_2) d_{0,1} = f_{0,1}.$$
(4.28)

Returning to R3. For l = 3 and n = 0 we have

$$2(k_1 + k_2) d_{3,0} + (k_1 - k_2) d_{1,1} = \tilde{f}_{3,0}.$$
(4.29)

Again, following the coupling of the unknowns from (4.29) (cf. coefficient stencil), the equation corresponding to l = 1, n = 1 (in R5) is

$$(k_1 - k_2) d_{3,0} + (2k_1 + (k_1 + k_2) d_{1,1} = \tilde{f}_{1,1}.$$
(4.30)

Equations (4.27)-(4.30) are two 2×2 linear systems of equations for $d_{2,0}$, $d_{0,1}$, and $d_{3,0}$, $d_{1,1}$, respectively. Continuing in this manner, i.e., in R3 corresponding to (l, n) = (2m, 0) (and (l, n) = (2m+1, 0)), $m \ge 0$, and following the coupling of the equations we terminate in R4 corresponding to (l, n) = (0, m) (in R5 with (l, n) = (1, m)) yielding an $(m+1) \times (m+1)$ decoupled linear system of equations. Hence, when the equations are appropriately assembled, we obtain decoupled linear system of equations of sizes: 1, 1, 2, 2, 3, 3, ..., m, m,

To further illustrate this point, we rename/renumber the unknowns $d_{l,n} \to e_j$ (and rhs: $\tilde{f}_{l,n} \to b_j$) in the order described above, see Figure 4.4.

nl	0	1	2	3	4	5	6	•••	n	$\mathbf{h}^{\mathbf{l}}$	0	1	2	3	4	5	6	•••
0	d _{0,0}	d _{0,1}	d _{0,2}	d _{0,3}	d _{0,4}	d _{0,5}	d _{0,6}	•••	_	0	e ₁	e ₂	e ₃	e ₅	e ₇	e ₁₀	e ₁₃	
1	d _{1,0}	d _{1,1}	d _{1,2}	d _{1,3}	d _{1,4}	d _{1,5}	d _{1,6}	•••		1	e ₄	e ₆	e ₈	e ₁₁	e ₁₄			
2	d _{2,0}	d _{2,1}	d _{2,2}	d _{2,3}	d _{2,4}	d _{2,5}	d _{2,6}	•••	_	2	e ₉	e ₁₂	e ₁₅					
3	d _{3,0}	d _{3,1}	d _{3,2}	d _{3,3}	d _{3,4}	d _{3,5}	d _{3,6}	•••	_	3	e ₁₆							
4	d _{4,0}	d _{4,1}	d _{4,2}	d _{4,3}	d _{4,4}	d _{4,5}	d _{4,6}	•••		4								
÷	:	:	:	•	•	:	:			:								

Figure 4.4: Renumbering of the unknowns $d_{l,n} \rightarrow e_j$.

Algebraically, the rename/renumbering $d_{l,n} \to e_j$ is given by

if *l* is even:
$$(l,n) \longrightarrow j = \frac{(l+2n)^2 + 2(l+4n+2)}{4}$$
,
if *l* is odd: $(l,n) \longrightarrow j = \frac{(l+2n)^2 + 2(l+4n+2) + 1}{4}$.

As discussed above, the linear system of equations for $\mathbf{e} = \{e_j\}_{j \ge 1}$, $A\mathbf{e} = \mathbf{b}$, has the following block diagonal structure.

$$A = \begin{bmatrix} k_1 + k_2 \\ 2k_1 + (k_1 + k_2) \\ A_{3,4,2} \\ A_{3,5,2} \\ & \ddots \\ & A_{3,5,m} \\ & & A_{3,5,m} \\ & & \ddots \\ & & & \ddots \end{bmatrix}$$
(4.31)

The matrices $A_{3,4,m}$ and $A_{3,5,m}$ are $m \times m$, symmetric, tridiagonal matrices given by

$$A_{3,4,m} = \begin{bmatrix} 2(k_1 + k_2) & (k_1 - k_2) \\ (k_1 - k_2) & 2(k_1 + k_2) & (k_1 - k_2) \\ & \ddots & & \\ & & \ddots & \\ & & & (k_1 - k_2) & 2(k_1 + k_2) & (k_1 - k_2) \\ & & & & (k_1 - k_2) & (k_1 + k_2) \end{bmatrix}$$
(4.32)

and

$$A_{3,5,m} = \begin{bmatrix} 2(k_1 + k_2) & (k_1 - k_2) \\ (k_1 - k_2) & 2(k_1 + k_2) & (k_1 - k_2) \\ & \ddots & & \\ & & \ddots & \\ & & & (k_1 - k_2) & 2(k_1 + k_2) & (k_1 - k_2) \\ & & & & (k_1 - k_2) & 2k_1 + (k_1 + k_2) \end{bmatrix}$$
(4.33)

The matrix $A_{3,4,m}$ represents the system of equations corresponding to unknowns $\mathbf{e}_{3,4,m} = \{e_j\}, j = m(m-1)+1, \ldots, m^2$, and $A_{3,5,m}$ represents the system of equations corresponding to unknowns $\mathbf{e}_{3,5,m} = \{e_j\}, j = m^2 + 1, \ldots, m(m+1).$

As $A_{3,4,m}$ and $A_{3,5,m}$ are real, symmetric matrices, and hence have real eigenvalues (singular values), a simple application of Gerschgorin's theorem establishes that there exists constants c_{min} and c_{max} such that the minimum and maximum eigenvalues of the matrices satisfy $0 < c_{min} \leq \lambda_{min}, \lambda_{max} \leq c_{max} < \infty$. Hence $A_{3,4,m}$ and $A_{3,5,m}$ are uniformly invertible with

$$\|\mathbf{e}_{3,4,m}\|_2 \le \|A_{3,4,m}^{-1}\|_2 \|\mathbf{b}_{3,4,m}\|_2 \le c_{min}^{-1} \|\mathbf{b}_{3,4,m}\|_2$$

and, likewise, $\|\mathbf{e}_{3,5,m}\|_2 \leq c_{min}^{-1} \|\mathbf{b}_{3,5,m}\|_2$.

We summarize the above in the following theorem.

Theorem 4.1 For $k_1, k_2 \in \mathbb{R}^+$, with $K(x) = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix}$, and $f(x) \in L^2_{\frac{\alpha}{2}}(\Omega)$ there exists a unique solution $\widetilde{u}(x) = \omega^{\frac{\alpha}{2}} u(x)$, with u(x) given by (4.3), satisfying

$$\mathcal{L}(\widetilde{u})(x) := -\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} K(x) \nabla \widetilde{u}(x) = f(x), \quad x \in \Omega,$$
(4.34)

subject to
$$\tilde{u} = 0$$
 on $\partial \Omega$, (4.35)

where the equal sign in (4.34) refers to the equality of the coefficients of the basis functions $\{V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2-1)\}_{l,n,\mu}$.

In the next section we further investigate the relationship between the coefficients $u_{l,n,\mu}$ and $f_{l,n,\mu}$, i.e., how the regularity of the solution \tilde{u} depends on the regularity of the rhs f.

We conclude this section with the following corollary that verifies equation (1.3) for $\tilde{g}(x) = \omega^{\frac{\alpha}{2}} g(x)$, with $g(x) \in L^{2}_{\frac{\alpha}{2}}(\Omega)$.

Corollary 4.1 For $\tilde{g}(x) = \omega^{\frac{\alpha}{2}} g(x)$, with $g(x) \in L^2_{\frac{\alpha}{2}}(\Omega)$, we have for $k \in \mathbb{R}$

$$k(-\Delta)^{\frac{\alpha}{2}} \widetilde{g}(x) = -\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} k \mathbf{I} \nabla \widetilde{g}(x).$$
(4.36)

Proof: Without loss of generality, we assume that k = 1. We verify (4.36) by showing that it holds on a basis for $L^2_{\frac{\alpha}{2}}(\Omega)$. Specifically, by showing that for $l, n \ge 0, \mu \in \{1, -1\}$

$$(-\Delta)^{\frac{\alpha}{2}} \omega^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) = -\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} k \mathbf{I} \nabla \omega^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1).$$

From [6, Theorem 3],

$$(-\Delta)^{\frac{\alpha}{2}} \omega^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) = \lambda_{l,n} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1),$$

where $\lambda_{l,n} = 2^{\alpha} \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)}.$

For $l \ge 1, n \ge 0, \mu \in \{1, -1\}$

$$-\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} \mathbf{I} \nabla \omega^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) = c_{l,n,\mu} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1),$$

where $c_{l,n,\mu}$ is the coefficient of $u_{l,n,\mu}$ (for $k_1 = k_2 = 1$) in equations (4.9), (4.10), (4.12) - (4.19), i.e.,

$$c_{l,n,\mu} = 2^{\alpha-2} 2^2 \frac{\Gamma(n+1+\frac{\alpha}{2})\Gamma(n+1+\frac{\alpha}{2}+l)}{\Gamma(n+1)\Gamma(n+1+l)} = \lambda_{l,n}$$

For $l = 0, n \ge 0, \mu = 1$

$$-\nabla \cdot (-\Delta)^{\frac{\alpha-2}{2}} \mathbf{I} \nabla \omega^{\frac{\alpha}{2}} V_{l,\mu}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) = 2 c_{0,n,1} V_{0,1}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1)$$

where $c_{0,n,1}$ is the coefficient of $u_{0,n,1}$ (for $k_1 = k_2 = 1$) in equations (4.8) and (4.11), i.e.,

$$2 c_{0,n,1} = 2 2^{\alpha - 2} 2 \frac{\Gamma(n + 1 + \frac{\alpha}{2}) \Gamma(n + 1 + \frac{\alpha}{2})}{\Gamma(n + 1) \Gamma(n + 1)} = \lambda_{0,n}$$

5 Regularity analysis of the solution

In Section 4 we established that given $f(x) \in L^2_{\frac{\alpha}{2}}(\Omega)$ there exists a unique $\tilde{u}(x) = \omega^{\frac{\alpha}{2}} u(x)$, with u(x) given by (4.3), satisfying (4.34), (4.35). In this section we analyze the influence of the regularity of f(x) on the regularity of u(x).

Following [8] we introduce the weighted function space $\mathbf{B}_{\beta}^{s_1,s_2}(\Omega)$ for any $s_1, s_2 > 0$ as

$$\mathbf{B}_{\beta}^{s_{1},s_{2}}(\Omega) := \left\{ v \, | \, v \in L_{\beta}^{2}(\Omega) \text{ and } |v|_{\mathbf{B}_{\beta}^{s_{1},s_{2}}(\Omega)} < \infty \right\} \,,$$

where the semi-norm $|\cdot|_{\mathbf{B}^{s_1,s_2}_{\beta}(\Omega)}$ is defined by

$$|v|^2_{\mathbf{B}^{s_1,s_2}_{\beta}(\Omega)} = \sum_{l,n,\mu} \left(l^{2s_1} + n^{2s_2} \right) \, v^2_{l,n,\mu} \, h^2_{l,n} \, .$$

The norm in this space is defined as $||v||_{\mathbf{B}^{s_1,s_2}_{\beta}(\Omega)}$, where

$$\|v\|_{\mathbf{B}^{s_1,s_2}_{\beta}(\Omega)}^2 = \sum_{l,n,\mu} \left(1 + l^{2s_1} + n^{2s_2}\right) v_{l,n,\mu}^2 h_{l,n}^2.$$

In determining the coefficients of u(x), $u_{l,n,\mu}$, in Section 4 the following (invertible) transformations were made:

$$e_{j} \leftarrow d_{l,n} \leftarrow \frac{\Gamma(n+1+\frac{\alpha}{2})}{\Gamma(n+1)} u_{l,n,\mu} \\ b_{j} \leftarrow \widetilde{f}_{l,n} \leftarrow 2^{-(\alpha-2)} \frac{\Gamma(n+1+l)}{\Gamma(n+1+\frac{\alpha}{2}+l)} f_{l,n,\mu}$$
 for $\mu = 1$ and $\mu = 2$. (5.1)

Similar to the discussion in Section 4.1, we focus our attention on $\{u_{l,n,1}\}$ and $\{f_{l,n,1}\}$, with the reliance of $\{u_{l,n,-1}\}$ on $\{f_{l,n,-1}\}$ following in an analogous manner. To this end, let

$$u_1(x) = \sum_{l \ge 1, n \ge 0} u_{l,n,1} V_{l,1}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) + \sum_{n \ge 0} \frac{u_{0,n,1}}{2} V_{0,1}(x) P_n^{(\frac{\alpha}{2},0)}(2r^2 - 1), \qquad (5.2)$$

$$f_1(x) = \sum_{l \ge 0, n \ge 0} f_{l,n,1} V_{l,1}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1) \text{ and } d(x) = \sum_{l \ge 0, n \ge 0} d_{l,n} V_{l,1}(x) P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1), \quad (5.3)$$

where $\{d_{l,n}\}$ is determined by (4.20)-(4.26).

We first relate the regularity of d(x) to that of $f_1(x)$ and then relate the regularity of $u_1(x)$ to d(x)(and subsequently to $f_1(x)$).

From Section 4.1 we have that the unknowns $\{e_j\}_{j\geq 1}$ are determined by solving the sequence of linear systems of equations

$$A_{3,4,m}\mathbf{e}_{3,4,m} = \mathbf{b}_{3,4,m}, \quad \text{and} \quad A_{3,5,m}\mathbf{e}_{3,5,m} = \mathbf{b}_{3,5,m}, \quad \text{for } m = 1, 2, \dots,$$
 (5.4)

where the mxm matrices $A_{3,4,m}$ and $A_{3,5,m}$ are symmetric, positive definite with $c_{max}^{-1} \leq \|A_{3,4,m}^{-1}\|_2$, $\|A_{3,5,m}^{-1}\|_2 \leq c_{min}^{-1}$. Hence it immediately follows that

$$\sum_{j=1}^{\infty} e_j^2 \lesssim \sum_{j=1}^{\infty} b_j^2 \iff \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} d_{l,n}^2 \lesssim \sum_{l=1}^{\infty} \sum_{n=1}^{\infty} \widetilde{f}_{l,n}^2 = \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} \left(2^{-(\alpha-2)} \frac{\Gamma(n+1+l)}{\Gamma(n+1+\frac{\alpha}{2}+l)} f_{l,n,1} \right)^2.$$

However, to determine which $\mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2}(\Omega)$ space d(x) lies in we need to be able to bound

$$\sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1 + l^{2s_1} + n^{2s_2}) d_{l,n}^2 h_{l,n}^2.$$
(5.5)

Returning to (5.4), for W a diagonal matrix, with $w_{ii} > 0$, note that

$$W A_{3,k,m} W^{-1} W \mathbf{e}_{3,k,m} = W \mathbf{b}_{3,k,m}, \ k \in \{4,5\}.$$

With $\mathcal{A} = W A_{3,k,m} W^{-1}$, note that \mathcal{A} is symmetric, positive definite, and has the same eigenvalues as $A_{3,k,m}$. Thus, $c_{max}^{-1} \leq ||\mathcal{A}^{-1}||_2 \leq c_{min}^{-1}$, and $||W\mathbf{e}_{3,k,m}||_2^2 \leq ||W\mathbf{b}_{3,k,m}||_2^2$. This enables us to introduce the weights needed in (5.5).

Lemma 5.1 For $f_1(x)$ and d(x) defined in (5.2) and (5.3), if $f_1(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2}(\Omega)$ then $d(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1+\frac{\alpha}{2},s_2+\frac{\alpha}{2}}(\Omega)$.

Proof: From above, we have that

$$\begin{split} \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) d_{l,n}^2 h_{l,n}^2 &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \tilde{f}_{l,n}^2 h_{l,n}^2 \\ &= \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \left(2^{-(\alpha-2)} \frac{\Gamma(n+1+l)}{\Gamma(n+1+\frac{\alpha}{2}+l)} f_{l,n,1}\right)^2 h_{l,n}^2 \\ \text{(using Stirling's formula)} &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \left((n+1+l)^{-\frac{\alpha}{2}} f_{l,n,1}\right)^2 h_{l,n}^2 \\ &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) (1+l+n)^{-\alpha} f_{l,n,1}^2 h_{l,n}^2 \\ &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2(t_1-\frac{\alpha}{2})}+n^{2(t_2-\frac{\alpha}{2})}) f_{l,n,1}^2 h_{l,n}^2 . \end{split}$$
Thus, if $f_1(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2}(\Omega)$ then $d(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1+\frac{\alpha}{2},s_2+\frac{\alpha}{2}}(\Omega).$

The connection between the regularity of $u_1(x)$ and d(x) is given in the next lemma.

Lemma 5.2 For $u_1(x)$ and d(x) defined in (5.2) and (5.3), if $d(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2}(\Omega)$ then $u_1(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2+\frac{\alpha}{2}}(\Omega)$.

Proof: Using (5.1)

$$\begin{split} \|u_1\|_{\mathbf{B}_{\frac{\alpha}{2}}^{t_1,t_2}(\Omega)}^2 &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \, u_{l,n,1}^2 \, h_{l,n}^2 \\ &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \, \left(\frac{\Gamma(n+1)}{\Gamma(n+1+\frac{\alpha}{2})} \, d_{l,n}\right)^2 \, h_{l,n}^2 \\ (\text{using Stirling's formula}) &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \, \left((n+1)^{-\frac{\alpha}{2}} d_{l,n}\right)^2 \, h_{l,n}^2 \\ &= \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2t_2}) \, (1+n)^{-\alpha} \, d_{l,n}^2 \, h_{l,n}^2 \\ &\lesssim \sum_{l=1}^{\infty} \sum_{n=0}^{\infty} (1+l^{2t_1}+n^{2(t_2-\frac{\alpha}{2})}) \, d_{l,n}^2 \, h_{l,n}^2 \, . \end{split}$$

Thus, if $d(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2}(\Omega)$ then $u_1(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1,s_2+\frac{\alpha}{2}}(\Omega)$.

We are now in the position to state the regularity result for the solution $\tilde{u}(x)$ satisfying (4.34), (4.35).

Theorem 5.1 For $k_1, k_2 \in \mathbb{R}^+$, with $K(x) = \begin{bmatrix} k_1 & 0 \\ 0 & k_2 \end{bmatrix}$, and $f(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1, s_2}(\Omega)$ there exists a unique solution $\widetilde{u}(x) = \omega^{\frac{\alpha}{2}} u(x)$ satisfying (4.34), (4.35), with $u(x) \in \mathbf{B}_{\frac{\alpha}{2}}^{s_1 + \frac{\alpha}{2}, s_2 + \alpha}(\Omega)$.

Proof: The result follows from Lemmas 5.1 and 5.2 applied to $\{f_{l,n,1}\}, \{u_{l,n,1}\}, \{u_{l,n,2}\}, \{u_{l,n,2}\}, \{u_{l,n,2}\}$.

Remark: For the special case of $k_1 = k_2$ Theorem 5.1 is consistent with Theorem 3.2 in [8].

6 Concluding remarks

In this article we have introduced a generalization of the fractional Laplacian, (1.2), and showed that for K(x) a constant, symmetric positive definite matrix the associated fractional Poisson equation (in \mathbb{R}^2) is well posed. Additionally we have related the regularity of the solution u(x) to the regularity of the RHS function f(x). We are currently working to extend these results to a general positive definite matrix K(x), and to extend the results to $\Omega \subset \mathbb{R}^3$.

Acknowledgements

(

The third author was partially supported by the National Science Foundation under Grant No. DMS-2012291.

A Recurrence formulas for Jacobi polynomials

In this section we give a number of recurrence formulas for Jacobi polynomials used in the analysis (see [1, pg. 782]).

$$\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + 1\right)\left(1 - x\right)P_{n}^{(\alpha+1,\beta)}(x) = \left(n + \alpha + 1\right)P_{n}^{(\alpha,\beta)}(x) - \left(n + 1\right)P_{n+1}^{(\alpha,\beta)}(x)$$
(A.1)

$$\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + 1\right)\left(1 + x\right)P_n^{(\alpha,\beta+1)}(x) = \left(n + \beta + 1\right)P_n^{(\alpha,\beta)}(x) + \left(n + 1\right)P_{n+1}^{(\alpha,\beta)}(x)$$
(A.2)

$$(2n + \alpha + \beta) P_n^{(\alpha - 1, \beta)}(x) = (n + \alpha + \beta) P_n^{(\alpha, \beta)}(x) - (n + \beta) P_{n-1}^{(\alpha, \beta)}(x)$$
(A.3)

$$(2n + \alpha + \beta) P_n^{(\alpha,\beta-1)}(x) = (n + \alpha + \beta) P_n^{(\alpha,\beta)}(x) + (n + \alpha) P_{n-1}^{(\alpha,\beta)}(x)$$
(A.4)

Lemma A.1 For $l \ge 1$ and $n \ge 1$,

$$l P_n^{(\gamma,l)}(t) + \left(\frac{1+t}{2}\right) (n+\gamma+l+1) P_{n-1}^{(\gamma+1,l+1)}(t) = (n+l) P_n^{(\gamma+1,l-1)}(t).$$
(A.5)

Proof: Using (A.2)

$$\begin{split} l \, P_n^{(\gamma,l)}(t) \; &+\; \frac{(n+\gamma+l+1)}{2} \, (1+t) \, P_{n-1}^{(\gamma+1,l+1)}(t) \\ &=\; l \, P_n^{(\gamma,l)}(t) \; +\; \frac{(n+\gamma+l+1)}{2 \, (n+\gamma+l+1)} \left[(n+l) \, P_{n-1}^{(\gamma+1,l)}(t) \; +\; n \, P_n^{(\gamma+1,l)}(t) \right] \\ &=\; \frac{l}{(2n+\gamma+l+1)} \left[(n+\gamma+l+1) \, P_n^{(\gamma+1,l)}(t) \; -\; (n+l) \, P_{n-1}^{(\gamma+1,l)}(t) \right] \; (\text{using (A.3)}) \\ &+\; \frac{(n+\gamma+l+1)}{(2n+\gamma+l+1)} \left[n \, P_n^{(\gamma+1,l)}(t) \; +\; (n+l) \, P_{n-1}^{(\gamma+1,l)}(t) \right] \\ &=\; \frac{(n+l)}{(2n+\gamma+l+1)} \left[(n+\gamma+l+1) \, P_n^{(\gamma+1,l)}(t) \; +\; (n+\gamma+1) \, P_{n-1}^{(\gamma+1,l)}(t) \right] \\ &=\; \frac{(n+l)}{(2n+\gamma+l+1)} \left[(2n+\gamma+l+1) \, P_n^{(\gamma+1,l-1)}(t) \; (\text{using (A.4)}) \right] \\ &=\; (n+l) \, P_n^{(\gamma+1,l-1)}(t) \, . \end{split}$$

B Proof of Theorem 2.1

In this section we give a proof of Theorem 2.1. For brevity, we use the notation used in [6].

Below the function $G_{pq}^{mn}\begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} denotes the Meijer$ *G* $-function, and <math>{}_{p}\mathbf{F}_{q}(\cdot)$ the regularized hypergeometric function.

The following five conditions are used in some of the statements involving the Meijer G-function.

The following result is used in the analysis. It summarizes Theorems 1 and 2 in [6].

Theorem B.1 [6, Theorem 1 and 2] Let V(x) be a solid harmonic polynomial of degree $l \ge 0$ and $\delta = d + 2l$. Assume that $-d < \alpha < 0$ or $\alpha > 0$ and parameters m, n, p, q, \mathbf{a} , and \mathbf{b} satisfy Condition A, as well as $2(1 - \bar{a}) > -\alpha + l, \quad -2b < d + l.$

$$Define \ f(x) := V(x) \ G_{pq}^{mn} \left(\begin{array}{c} \mathbf{a} \\ \mathbf{b} \end{array} \middle| |x|^2 \right). \ Then$$

$$(-\Delta)^{\alpha/2} \ f(x) = 2^{\alpha} V(x) \ G_{p+2q+2}^{m+1n+1} \left(\begin{array}{ccc} 1 - \frac{\delta + \alpha}{2} \,, & \mathbf{a} - \frac{\alpha}{2} \,, & -\frac{\alpha}{2} \\ 0 \,, & \mathbf{b} - \frac{\alpha}{2} \,, & 1 - \frac{\delta}{2} \end{array} \middle| |x|^2 \right)$$
(B.1)

for all $x \neq 0$. The same statement holds under Conditions B, C, and D provided

 $2\overline{\lambda} > -\alpha + l, \quad 2\underline{\lambda} < d + l;$

if Condition B is satisfied, we additionally require that $\nu > 0$ and either $|x| \neq 1$ or $\nu > 1 + \alpha$.

For the case $\alpha > 0$, the results extends to x = 0 whenever both f and $(-\Delta)^{\alpha/2} f$ are continuous at 0.

Proof of Theorem 2.1: Proceeding in a similar manner to the proof of Theorem 3 in [6], let $V(x) = V_{l,\mu}(x)$, and using (40a) in [6] and equation 8.4.49.22 in [16]

$$\begin{split} f(x) &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) \left(1-|x|^2\right)_{+}^{\frac{\alpha}{2}-s} {}_2\mathbf{F}_1 \left(\begin{array}{c} -n \ , \ \frac{\delta+\alpha}{2}+n-s \ |1-|x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) G_{22}^{20} \left(\begin{array}{c} 1+\frac{\alpha}{2}-s+n \ , \ 1-\frac{\delta}{2}-n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) G_{22}^{20} \left(\begin{array}{c} 1+\frac{\alpha}{2}-s+n \ , \ 1-\frac{\delta}{2}-n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) G_{22}^{20} \left(\begin{array}{c} 1+\frac{\alpha}{2}-s+n \ , \ 1-\frac{\delta}{2}-n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s)}{\Gamma(n+1)} V(x) (-1)^n G_{22}^{11} \left(\begin{array}{c} 1-\frac{\delta}{2}-n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s+n \ , \ 1+\frac{\alpha}{2}-s+n \ ||x|^2 \right) \\ &= \frac{\Gamma(n+1+\frac{\alpha}{2}-s+n \ , \ 1+\frac{\alpha}{2}-s+n \ , \$$

Using Theorem B.1,

$$\begin{aligned} (-\Delta)^{\frac{\alpha-2}{2}} V(x) G_{22}^{11} \left(\begin{array}{ccc} 1 - \frac{\delta}{2} - n, & 1 + \frac{\alpha}{2} - s + n \\ 0, & 1 - \frac{\delta}{2} \end{array} \middle| |x|^2 \right) \\ &= 2^{\alpha-2} V(x) G_{44}^{22} \left(\begin{array}{ccc} 2 - \frac{\delta+\alpha}{2}, & 2 - \frac{\delta+\alpha}{2} - n, & 2 - s + n, & 1 - \frac{\alpha}{2} \\ 0, & 1 - \frac{\alpha}{2}, & 2 - \frac{\delta+\alpha}{2}, & 1 - \frac{\delta}{2} \end{array} \middle| |x|^2 \right) \\ &= 2^{\alpha-2} V(x) G_{44}^{22} \left(\begin{array}{ccc} 2 - \frac{\delta+\alpha}{2}, & 2 - \frac{\delta+\alpha}{2} - n, & 2 - s + n, & 1 - \frac{\alpha}{2} \\ 1 - \frac{\alpha}{2}, & 0, & 2 - \frac{\delta+\alpha}{2}, & 1 - \frac{\delta}{2} \end{array} \middle| |x|^2 \right) \quad (\text{ using the defn. of } G_{44}^{22}(\cdot)) \\ &= 2^{\alpha-2} V(x) G_{33}^{12} \left(\begin{array}{ccc} 2 - \frac{\delta+\alpha}{2}, & 2 - \frac{\delta+\alpha}{2} - n, & 2 - s + n \\ 0, & 2 - \frac{\delta+\alpha}{2}, & 1 - \frac{\delta}{2} \end{array} \middle| |x|^2 \right) \quad \text{using [6, (22)]} \\ &= 2^{\alpha-2} V(x) G_{33}^{12} \left(\begin{array}{ccc} 2 - \frac{\delta+\alpha}{2}, & 2 - \frac{\delta+\alpha}{2} - n, & 2 - s + n \\ 0, & 1 - \frac{\delta}{2}, & 2 - \frac{\delta+\alpha}{2} \end{array} \middle| |x|^2 \right) \quad (\text{ using the defn. of } G_{33}^{12}(\cdot)) \\ &= 2^{\alpha-2} V(x) G_{32}^{12} \left(\begin{array}{ccc} 2 - \frac{\delta+\alpha}{2} - n, & 2 - s + n \\ 0, & 1 - \frac{\delta}{2}, & 2 - \frac{\delta+\alpha}{2} \end{array} \middle| |x|^2 \right) \quad \text{using [6, (21)].} \end{aligned}$$

$$(B.3)$$

Next, using [6, (23)], and [6, (40b)]

$$G_{22}^{11} \begin{pmatrix} 2 - \frac{\delta+\alpha}{2} - n, & 2-s+n \\ 0, & 1-\frac{\delta}{2} \end{pmatrix} |x|^2 = \frac{\Gamma(n-1+\frac{\delta+\alpha}{2})}{\Gamma(n+2-s)} {}_2\mathbf{F}_1 \begin{pmatrix} n-1+\frac{\delta+\alpha}{2}, & -1+s-n \\ \frac{\delta}{2} \end{pmatrix} = \frac{\Gamma(n-1+\frac{\delta+\alpha}{2})}{\Gamma(n+2-s)} (-1)^{n+1-s} \frac{\Gamma(n+2-s)}{\Gamma(n+1-s+\frac{\delta}{2})} P_{n+1-s}^{(\frac{\alpha}{2}-2+s,\frac{\delta}{2}-1)} (2|x|^2 - 1).$$
(B.4)

Combining (B.2)-(B.4) we obtain (2.11).

C Proof of Theorem 3.1

Proof of Theorem 3.1: Recall that in \mathbb{R}^2 , $\nabla \cdot = \frac{\partial}{\partial r} \cdot \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial}{\partial \varphi} \cdot \hat{\boldsymbol{\varphi}}$.

Using (2.8),

$$\frac{d}{dr}\left((1-r^2)^{\alpha/2}P_n^{(\frac{\alpha}{2},l)}(2r^2-1)\right) = -2(n+\frac{\alpha}{2})r\left(1-r^2\right)^{\alpha/2-1}P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1).$$
(C.1)

Also,

$$\frac{\partial}{\partial r} V_{l,\mu}(x) = l r^{-1} V_{l,\mu}(x) . \qquad (C.2)$$

Combining (C.1) and (C.2) with the definition of ∇ we obtain

$$\nabla f(x) = \left(l V_{l,\mu}(x) r^{-1} (1 - r^2)^{\alpha/2} P_n^{\left(\frac{\alpha}{2},l\right)} (2r^2 - 1) - 2(n + \frac{\alpha}{2}) V_{l,\mu}(x) r (1 - r^2)^{\alpha/2 - 1} P_n^{\left(\frac{\alpha}{2} - 1, l + 1\right)} (2r^2 - 1) \right) \hat{\mathbf{r}} + \left(\left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x) \right) r^{-1} (1 - r^2)^{\alpha/2} P_n^{\left(\frac{\alpha}{2},l\right)} (2r^2 - 1) \right) \hat{\boldsymbol{\varphi}}.$$
(C.3)

Using $\mathbf{i} = \cos(\varphi) \hat{\mathbf{r}} - \sin(\varphi) \hat{\boldsymbol{\varphi}}$, from (C.3) we obtain

$$\frac{\partial f}{\partial x} = \cos(\varphi) \left(l V_{l,\mu}(x) r^{-1} (1-r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)} (2r^2-1) - 2(n+\frac{\alpha}{2}) V_{l,\mu}(x) r (1-r^2)^{\alpha/2-1} P_n^{(\frac{\alpha}{2}-1,l+1)} (2r^2-1) \right) - \sin(\varphi) \left(\left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x) \right) r^{-1} (1-r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)} (2r^2-1) \right).$$
(C.4)

With a view to combining the $(1 - r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1)$ terms in (C.4), let $H_1 = \cos(\varphi) l V_{l,\mu}(x) r^{-1} - \sin(\varphi) \left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x)\right) r^{-1}$. Suppose $V_{l,\mu}(x) = r^l \cos(l\varphi)$. Then $\frac{\partial}{\partial \varphi} V_{l,\mu}(x) = -r^l l \sin(l\varphi)$. In this case, $H_1 = l r^{l-1} (\cos(l\varphi) \cos(\varphi) + \sin(l\varphi) \sin(\varphi)) = l r^{l-1} \cos((l-1)\varphi) = l V_{l-1,\mu}(x)$. (C.5)

Alternatively, suppose $V_{l,\mu}(x) = r^l \sin(l\varphi)$. Then $\frac{\partial}{\partial \varphi} V_{l,\mu}(x) = r^l l \cos(l\varphi)$, and $H_1 = l r^{l-1} (\sin(l\varphi) \cos(\varphi) - \cos(l\varphi) \sin(\varphi)) = l r^{l-1} \sin((l-1)\varphi) = l V_{l-1,\mu}(x)$. (C.6)

The coefficient of $-2(n+\frac{\alpha}{2})(1-r^2)^{\alpha/2-1}P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1)$ in (C.4) is $H_2 := \cos(\varphi) V_{l,\mu}(x) r$. Suppose $V_{l,\mu}(x) = r^l \cos(l\varphi)$. Then, $H_2 = r^{l+1} \cos(l\varphi) \cos(\varphi) = \frac{1}{2}r^{l+1} (\cos((l-1)\varphi) + \cos((l+1)\varphi))$ $= \frac{1}{2}r^2r^{l-1}\cos((l-1)\varphi) + \frac{1}{2}r^{l+1}\cos((l+1)\varphi)$ $= \frac{1}{2}r^2V_{l-1,\mu}(x) + \frac{1}{2}V_{l+1,\mu}(x)$. (C.7) Alternatively, suppose $V_{l,\mu}(x) = r^l \sin(l\varphi)$. Then,

$$H_{2} = r^{l+1} \sin(l\varphi) \cos(\varphi) = \frac{1}{2} r^{l+1} (\sin((l-1)\varphi) + \sin((l+1)\varphi))$$

= $\frac{1}{2} r^{2} r^{l-1} \sin((l-1)\varphi) + \frac{1}{2} r^{l+1} \sin((l+1)\varphi)$
= $\frac{1}{2} r^{2} V_{l-1,\mu}(x) + \frac{1}{2} V_{l+1,\mu}(x).$ (C.8)

Rewriting (C.4) using (C.5), (C.6) and (C.7), (C.8) we have

$$\frac{\partial f}{\partial x} = l V_{l-1,\mu}(x) (1-r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)}(2r^2-1) - (n+\frac{\alpha}{2}) V_{l-1,\mu}(x) r^2 (1-r^2)^{\alpha/2-1} P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1) - (n+\frac{\alpha}{2}) V_{l+1,\mu}(x) (1-r^2)^{\alpha/2-1} P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1).$$
(C.9)

Let $H_3 = l(1-r^2) P_n^{(\frac{\alpha}{2},l)}(2r^2-1) - (n+\frac{\alpha}{2}) r^2 P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1)$. With the substitution $t = 2r^2 - 1$,

$$H_{3} = \frac{l}{2} (1-t) P_{n}^{(\frac{\alpha}{2},l)}(t) - \frac{(n+\frac{\alpha}{2})}{2} (1+t) P_{n}^{(\frac{\alpha}{2}-1,l+1)}(t)$$

$$= \frac{l}{(2n+\frac{\alpha}{2}+l+1)} \left((n+\frac{\alpha}{2}) P_{n}^{(\frac{\alpha}{2}-1,l)}(t) - (n+1) P_{n+1}^{(\frac{\alpha}{2}-1,l)}(t) \right) \quad (\text{using (A.1)})$$

$$- \frac{(n+\frac{\alpha}{2})}{(2n+\frac{\alpha}{2}+l+1)} \left((n+l+1) P_{n}^{(\frac{\alpha}{2}-1,l)}(t) + (n+1) P_{n+1}^{(\frac{\alpha}{2}-1,l)}(t) \right) \quad (\text{using (A.2)})$$

$$= - \frac{(n+\frac{\alpha}{2})}{(2n+\frac{\alpha}{2}+l+1)} (n+1) P_{n}^{(\frac{\alpha}{2}-1,l)}(t) - \frac{(n+1)}{(2n+\frac{\alpha}{2}+l+1)} (n+\frac{\alpha}{2}+l) P_{n+1}^{(\frac{\alpha}{2}-1,l)}(t)$$

$$= - \frac{(n+1)}{(2n+\frac{\alpha}{2}+l+1)} \left((n+\frac{\alpha}{2}+l) P_{n+1}^{(\frac{\alpha}{2}-1,l)}(t) + (n+\frac{\alpha}{2}) P_{n}^{(\frac{\alpha}{2}-1,l)}(t) \right)$$

$$= - (n+1) P_{n+1}^{(\frac{\alpha}{2}-1,l-1)}(t) \quad (\text{using (A.4)}).$$
(C.10)

Substituting (C.10) into (C.9) and rearranging we obtain (3.1).

The expression for $\frac{\partial f}{\partial y}$ is obtained in a similar manner. Using $\mathbf{j} = \sin(\varphi)\hat{\mathbf{r}} + \cos(\varphi)\hat{\varphi}$, from (C.3) we obtain

$$\frac{\partial f}{\partial y} = \sin(\varphi) \left(l \, V_{l,\mu}(x) \, r^{-1} (1-r^2)^{\alpha/2} \, P_n^{(\frac{\alpha}{2},l)}(2r^2-1) - 2(n+\frac{\alpha}{2}) \, V_{l,\mu}(x) \, r \, (1-r^2)^{\alpha/2-1} \, P_n^{(\frac{\alpha}{2}-1,\,l+1)}(2r^2-1) \right) \\
+ \, \cos(\varphi) \left(\left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x) \right) \, r^{-1} (1-r^2)^{\alpha/2} \, P_n^{(\frac{\alpha}{2},l)}(2r^2-1) \right).$$
(C.11)

With a view to combining the $(1 - r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)}(2r^2 - 1)$ terms in (C.11), let $H_1 = \sin(\varphi) l V_{l,\mu}(x) r^{-1} + \cos(\varphi) \left(\frac{\partial}{\partial \varphi} V_{l,\mu}(x)\right) r^{-1}$. Suppose $V_{l,\mu}(x) = r^l \cos(l\varphi)$. In this case,

$$H_1 = l r^{l-1} \left(\cos(l\varphi) \sin(\varphi) - \sin(l\varphi) \cos(\varphi) \right) = -l r^{l-1} \sin((l-1)\varphi) = -l V_{l-1,\mu^*}(x) \cdot (C.12)$$

Alternatively, suppose $V_{l,\mu}(x) = r^l \sin(l\varphi)$. Then

$$H_1 = lr^{l-1}(\sin(l\varphi)\sin(\varphi) + \cos(l\varphi)\cos(\varphi)) = lr^{l-1}\cos((l-1)\varphi) = lV_{l-1,\mu^*}(x).$$
(C.13)

The coefficient of $-2(n+\frac{\alpha}{2})(1-r^2)^{\alpha/2-1}P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1)$ in (C.4) is $H_2 := \sin(\varphi)V_{l,\mu}(x)r$. Suppose $V_{l,\mu}(x) = r^l \cos(l\varphi)$. Then,

$$H_{2} = r^{l+1} \cos(l\varphi) \sin(\varphi) = -\frac{1}{2} r^{l+1} (\sin((l-1)\varphi) - \sin((l+1)\varphi))$$

$$= -\frac{1}{2} r^{2} r^{l-1} \sin((l-1)\varphi) + \frac{1}{2} r^{l+1} \sin((l+1)\varphi)$$

$$= -\frac{1}{2} r^{2} V_{l-1,\mu^{*}}(x) + \frac{1}{2} V_{l+1,\mu^{*}}(x).$$
(C.14)

Alternatively, suppose $V_{l,\mu}(x) = r^l \sin(l\varphi)$. Then,

$$H_{2} = r^{l+1} \sin(l\varphi) \sin(\varphi) = \frac{1}{2} r^{l+1} \left(\cos((l-1)\varphi) - \cos((l+1)\varphi) \right)$$

$$= \frac{1}{2} r^{2} r^{l-1} \cos((l-1)\varphi) - \frac{1}{2} r^{l+1} \cos((l+1)\varphi)$$

$$= \frac{1}{2} r^{2} V_{l-1,\mu^{*}}(x) - \frac{1}{2} V_{l+1,\mu^{*}}(x).$$
(C.15)

Rewriting (C.11) using (C.12), (C.13) and (C.14), (C.15) we have

$$\frac{\partial f}{\partial y} = l(\mp) V_{l-1,\mu^*}(x) (1-r^2)^{\alpha/2} P_n^{(\frac{\alpha}{2},l)}(2r^2-1) - (n+\frac{\alpha}{2}) (\mp) V_{l-1,\mu^*}(x) r^2 (1-r^2)^{\alpha/2-1} P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1) - (n+\frac{\alpha}{2}) (\pm) V_{l+1,\mu^*}(x) (1-r^2)^{\alpha/2-1} P_n^{(\frac{\alpha}{2}-1,l+1)}(2r^2-1).$$
(C.16)

Using (C.10) and rearranging we obtain (3.2). The proofs of (3.3) and (3.4) follow in a similar and much simpler manner (as $V_{0,1} = 1$), and are thus omitted.

References

- M. Abramowitz and I.A. Stegun. Handbook of mathematical functions with formulas, graphs, and mathematical tables, volume 55 of National Bureau of Standards Applied Mathematics Series. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] G. Acosta and J.P. Borthagaray. A fractional Laplace equation: Regularity of solutions and finite element approximations. SIAM J. Numer. Anal., 55(2):472–495, 2017.
- [3] G. Acosta, J.P. Borthagaray, O. Bruno, and M. Maas. Regularity theory and high order numerical methods for the (1-d)-fractional Laplacian. *Math. Comp.*, 87:1821–1857, 2018.

- [4] L. Caffarelli and L. Silvestre. An extension problem related to the fractional Laplacian. Comm. Partial Differential Equations, 32(7-9):1245–1260, 2007.
- [5] C.F. Dunkl and Y. Xu. Orthogonal polynomials of several variables, volume 155 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, second edition, 2014.
- B. Dyda, A. Kuznetsov, and M. Kwaśnicki. Fractional Laplace operator and Meijer G-function. Constr. Approx., 45(3):427–448, 2017.
- [7] V.J. Ervin. Regularity of the solution to fractional diffusion, advection, reaction equations in weighted Sobolev spaces. J. Differential Equations, 278:294–325, 2021.
- [8] Z. Hao, H. Li, Z. Zhang, and Z. Zhang. Sharp error estimates of a spectral Galerkin method for a diffusion-reaction equation with integral fractional Laplacian on a disk. *Math. Comp.*, 90(331):2107–2135, 2021.
- [9] Z. Hao and Z. Zhang. Optimal Regularity and Error Estimates of a Spectral Galerkin Method for Fractional Advection-Diffusion-Reaction Equations. SIAM J. Numer. Anal., 58(1):211–233, 2020.
- [10] M. Kwaśnicki. Ten equivalent definitions of the fractional Laplace operator. Fract. Calc. Appl. Anal., 20(1):7–51, 2017.
- [11] H. Li and Y. Xu. Spectral approximation on the unit ball. SIAM J. Numer. Anal., 52(6):2647– 2675, 2014.
- [12] Y. Li. On the decomposition of solutions: from fractional diffusion to fractional Laplacian. Fract. Calc. Appl. Anal., 24(5):1571–1600, 2021.
- [13] A Lischke, G. Pang, M. Gulian, F. Song, C. Glusa, X. Zheng, Z. Mao, W. Cai, M.M. Meerschaert, M. Ainsworth, and G.Em Karniadakis. What is the fractional Laplacian? A comparative review with new results. J. Comput. Phys., 404:109009, 62, 2020.
- [14] Z. Mao, S. Chen, and J. Shen. Efficient and accurate spectral method using generalized Jacobi functions for solving Riesz fractional differential equations. *Appl. Numer. Math.*, 106:165–181, 2016.
- [15] Z. Mao and G.Em Karniadakis. A spectral method (of exponential convergence) for singular solutions of the diffusion equation with general two-sided fractional derivative. *SIAM J. Numer. Anal.*, 56(1):24–49, 2018.
- [16] A.P. Prudnikov, Yu.A. Brychkov, and O.I. Marichev. Integrals and series. Vol. 3. Gordon and Breach Science Publishers, New York, 1990. More special functions, Translated from the Russian by G. G. Gould.
- [17] F. Zeng, F. Liu, C. Li, K. Burrage, I. Turner, and V. Anh. A Crank-Nicolson ADI spectral method for a two-dimensional Riesz space fractional nonlinear reaction-diffusion equation. *SIAM J. Numer. Anal.*, 52(6):2599–2622, 2014.
- [18] X. Zheng, V.J. Ervin, and H. Wang. Optimal Petrov-Galerkin spectral approximation method for the fractional diffusion, advection, reaction equation on a bounded interval. J. Sci. Comput., 86(3):Paper No. 29, 22, 2021.