

A note on local search for hitting sets

Zdeněk Dvořák*

Abstract

Let π be a property of pairs (G, Z) , where G is a graph and $Z \subseteq V(G)$. In the *minimum π -hitting set problem*, given an input graph G , we want to find a smallest set $X \subseteq V(G)$ such that X intersects every set $Z \subseteq V(G)$ such that (G, Z) has the property π . An important special case is that π is satisfied by (G, Z) exactly if $G[Z]$ is isomorphic to one of graphs in a finite set \mathcal{F} ; in this *minimum \mathcal{F} -hitting set problem*, X needs to hit all appearances of the graphs from \mathcal{F} as induced subgraphs of G . In this note, we show that the local search argument of Har-Peled and Quanrud gives a PTAS for the minimum \mathcal{F} -hitting set problem for graphs from any class with polynomial expansion. Moreover, we argue that the local search argument applies more generally to all properties π such that one can test whether X is a π -hitting set in polynomial time and $G[Z]$ has bounded diameter whenever (G, Z) satisfies π ; this is a common generalization of the minimum \mathcal{F} -hitting set problem and minimum r -dominating set problem. Finally, we note that the analogous claim also holds for the dual problem of finding the maximum number of disjoint sets Z such that (G, Z) has the property π ; this generalizes maximum F -matching, maximum induced F -matching, and maximum r -independent set problems.

Let H and G be graphs. If a graph isomorphic to H is obtained from a subgraph of G by contracting pairwise vertex-disjoint subgraphs, each of radius at most r , we say that H is an r -shallow minor of G . The *density* of a graph H is $|E(H)|/|V(H)|$. For a function $f : \mathbb{Z}_0^+ \rightarrow \mathbb{R}_0^+$, a graph G has

*Charles University, Prague, Czech Republic. E-mail: rakdver@iuuk.mff.cuni.cz. Supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 810115).

expansion bounded by f if for every non-negative integer r , every r -shallow minor of G has density at most $f(r)$. A class of graphs \mathcal{G} has expansion bounded by f if all graphs from \mathcal{G} do. We say \mathcal{G} has *bounded expansion* if \mathcal{G} has expansion bounded by some function f , and *polynomial expansion* if \mathcal{G} has expansion bounded by a polynomial.

Let π be a property of pairs (G, Z) , where G is a graph and $Z \subseteq V(G)$ is non-empty. The *diameter* of π is the maximum diameter of $G[Z]$ over all pairs (G, Z) satisfying the property π , or ∞ if there exist such pairs with the diameter of $G[Z]$ arbitrarily large (this includes the case that $G[Z]$ is disconnected for some (G, Z) satisfying π). For a graph G , a set $X \subseteq V(G)$ is a π -*hitting set* if X intersects every $Z \subseteq V(G)$ such that (G, Z) satisfies π . Let $\gamma_\pi(G)$ denote the minimum size of a π -hitting set in G . For a graph class \mathcal{G} , we say that π -*hitting set is efficiently testable in \mathcal{G}* if there exists a polynomial-time algorithm that for a graph $G \in \mathcal{G}$ and a set $X \subseteq V(G)$ decides whether X is a π -hitting set. Let us mention two important examples:

- For a finite set \mathcal{F} of graphs, if π is the property that $G[Z]$ is isomorphic to a graph in \mathcal{F} , we obtain the \mathcal{F} -*hitting set* problem of finding a smallest set $X \subseteq V(G)$ such that $G - X$ is \mathcal{F} -*free*, i.e., does not contain any induced subgraph isomorphic to an element of \mathcal{F} . The diameter of π is equal to the maximum of diameters of the graphs in \mathcal{F} .
- Let r be a positive integer, and let π be satisfied by (G, Z) iff there exists $z \in V(G)$ such that Z consists exactly of vertices at distance at most r from z in G . Then X is a π -hitting set iff it is an r -*dominating set*, i.e., every vertex of G is at distance at most r from X . The diameter of π is at most $2r$.

In both cases, π -hitting set is efficiently testable in all graphs.

For any π such that π -hitting set is efficiently testable and a positive integer c , one can use the following c -*local search* heuristic to find a (hopefully small) π -hitting set:

1. Let $X = V(G)$.
2. Iterate over all sets $Y \subseteq V(G)$ of size at most c such that $|X \Delta Y| < |X|$:
 - If $X \Delta Y$ is a π -hitting set, then let $X := X \Delta Y$ and repeat the step 2.

Of course, there are in general no guarantees that the result of local search is close to optimal. However, Har-Peled and Quanrud [4] proved the following claim for minimum vertex cover ($\{K_2\}$ -hitting set) and for minimum r -dominating set: For every class of graphs \mathcal{G} with polynomial expansion and every $\varepsilon > 0$, there exists c_ε such that c_ε -local search results in a π -hitting set of size at most $(1 + \varepsilon)\gamma_\pi(G)$ for every graph $G \in \mathcal{G}$. That is, local search gives a PTAS when restricted to any graph class with polynomial expansion. The main result of this note is the following common generalization of these results.

Theorem 1. *Let \mathcal{G} be a class of graphs with polynomial expansion. For every property π of finite diameter and every $\varepsilon > 0$, there exists c such that c -local search results in a π -hitting set of size at most $(1 + \varepsilon)\gamma_\pi(G)$.*

It is also natural to consider the dual notion. A set \mathcal{P} of pairwise disjoint subsets of vertices of G is a π -packing if (G, P) has the property π for every $P \in \mathcal{P}$; it is an *induced π -packing* if additionally no edge of G has ends in distinct elements of \mathcal{P} . For example:

- For a graph F , if π is the property that $G[P]$ is isomorphic to F , we obtain the F -matching (and *induced F -matching*) problem of finding the maximum number of disjoint (and non-adjacent) copies of F in the input graph.
- Let r be a positive integer, and let π be satisfied by (G, Z) iff there exists $z \in V(G)$ such that Z consists exactly of vertices at distance at most r from z in G . Then a π -packing corresponds to a $2r$ -independent set (a set of vertices is t -independent if the distance between any distinct elements is greater than t), and an induced π -packing corresponds to a $(2r + 1)$ -independent set.

Let $\alpha_\pi(G)$ denote the maximum size of a π -packing and $\alpha'_\pi(G)$ the maximum size of an induced π -packing in G . For a graph class \mathcal{G} , we say that *(induced) π -packing extension is tractable in \mathcal{G}* if for every positive integer c , there exists a polynomial-time algorithm that for a graph $G \in \mathcal{G}$ and a set $T \subseteq V(G)$ finds an (induced) π -packing \mathcal{P}_0 of size at least c and such $T \cap \bigcup \mathcal{P}_0 = \emptyset$, or decides no such \mathcal{P}_0 exists. If π has this property, we can run the following maximization variant of the c -local search algorithm:

1. Let $\mathcal{P} = \emptyset$.

2. Iterate over all $Y \subseteq \mathcal{P}$ of size less than c :

- If $\mathcal{P} \setminus Y$ can be extended to an (induced) π -packing \mathcal{P}' of size greater than $|\mathcal{P}|$, then let $\mathcal{P} := \mathcal{P}'$ and repeat the step 2.

Step 2 is implemented by testing whether there exists an (induced) π -packing \mathcal{P}_0 of size $|Y| + 1$ disjoint from $\bigcup_{P \in \mathcal{P} \setminus Y} P$ (or this set and its neighbors in the induced case). As our second result, we note that this algorithm again gives a PTAS in any class of graphs with polynomial expansion; the argument is less involved and essentially implicitly appears in [4].

Theorem 2. *Let \mathcal{G} be a class of graphs with polynomial expansion. For every property π of finite diameter and every $\varepsilon > 0$, there exists c such that the (induced) maximization c -local search results in an (induced) π -packing of size at least $(1 - \varepsilon)\alpha_\pi(G)$ in the non-induced case and at least $(1 - \varepsilon)\alpha'_\pi(G)$ in the induced case.*

The time complexity of the algorithms from Theorems 1 and 2 in a straightforward implementation is $O(n^{c+1})$ times the complexity of testing or extension, and thus the exponent of the time complexity depends on the desired precision; i.e., we obtain a PTAS, not an FPTAS. However, in all the examples that we considered, the property π is *FO-definable* in the following sense: There exists a formula φ in the first-order logic (allowing quantification over vertices, but not over sets of vertices or over edges) using the adjacency predicate, with free variables x_1, \dots, x_m and z such that (G, Z) with Z non-empty has the property π if and only if for some $v_1, \dots, v_m \in G$,

$$Z = \{v \in V(G) : G \models \varphi(v_1, \dots, v_m, v)\}.$$

E.g., for the property π used to define r -dominating and $2r$ -independent set problems, we can set $\varphi(x_1, z) \equiv \text{dist}_r(x_1, z)$, where

$$\text{dist}_r(x, y) \equiv (\exists p_0, \dots, p_r) x = p_0 \wedge y = p_r \wedge \bigwedge_{i=1}^r (p_{i-1} = p_i \vee p_{i-1}p_i \in E).$$

In this case, we can express the testing and extension in first-order logic. Let us demonstrate this on the most difficult example, the step 2 of the c -local search for maximum π -packing. Let \mathcal{P} be the current π -packing. For each element P of \mathcal{P} , color the edges of a BFS spanning tree of $G[P]$ red; we will access this information about edge colors using a new binary predicate R .

Let $\text{dist}_r^R(x, y)$ be the predicate defined analogously to $\text{dist}_r(x, y)$, but with E replaced by R . Let T be the unary predicate interpreted as $\bigcup \mathcal{P}$. We can delete at most $c - 1$ elements from \mathcal{P} and add c other elements to obtain a π -packing exactly if the following formula is satisfied:

$$\begin{aligned}
& (\exists r_1, \dots, r_{c-1}, x_1^1, \dots, x_m^1, \dots, x_1^c, \dots, x_m^c) \\
& \bigwedge_{j=1}^c (\exists z) \varphi(x_1^j, \dots, x_m^j, z) \\
& \wedge \bigwedge_{j=1}^c (\forall z) \left(\varphi(x_1^j, \dots, x_m^j, z) \wedge z \in T \Rightarrow \bigvee_{i=1}^{c-1} r_i \in T \wedge \text{dist}_{2r}^R(z, r_i) \right) \\
& \wedge \bigwedge_{1 \leq j_1 < j_2 \leq c} (\forall z_1, z_2) \left(\varphi(x_1^{j_1}, \dots, x_m^{j_1}, z_1) \wedge \varphi(x_1^{j_2}, \dots, x_m^{j_2}, z_2) \Rightarrow z_1 \neq z_2 \right).
\end{aligned}$$

The elements to be removed from \mathcal{P} are those intersected by r_1, \dots, r_{c-1} , while those added are given by the formula φ with parameters x_1^j, \dots, x_m^j for $j \in \{1, \dots, c\}$.

- The second line expresses that for each j , the formula φ parameterized by the m -tuple x_1^j, \dots, x_m^j defines a non-empty set Z_j .
- The third line states that Z_j intersects T only in the elements containing r_1, \dots, r_{c-1} that are to be removed from \mathcal{P} .
- The final line describes that the sets Z_1, \dots, Z_m are disjoint.

Dvořák, Král' and Thomas [1] described a data structure that for each graph from a fixed class with bounded expansion after a linear-time preprocessing allows one in constant time perform the following operations:

- Recolor an edge.
- Change the membership of a vertex in a set interpreting an unary predicate.
- Test whether a fixed first-order sentence ψ is satisfied, and
- if so, find a satisfying assignment for the initial segment of existential quantifiers of ψ .

With this data structure, we conclude that for a FO-definable property of bounded diameter, n -vertex graph G from any fixed class with bounded expansion, and any fixed c , we can run

- c -local search for minimum π -hitting in time $O(n)$, and
- (induced) c -local search for maximum π -packing in time $O(n^2)$.

The reason for the quadratic time in the second case is due to the need to recolor the red edges after each iteration, which may result in $O(n)$ time in case the sets of the (induced) π -packing have unbounded size. In case they have bounded size, e.g., for the maximum (induced) F -matching problem, the time complexity becomes $O(n)$. Moreover, even if the elements of the packing have unbounded size, it may be possible to use a different way how to describe the packing in the first-order logic (e.g., in the case of maximum t -independent set, we can just mark the vertices forming the set), again resulting in total time $O(n)$.

1 Tools

To prove Theorem 1, we use several standard tools from the theory of classes with bounded expansion.

Generalized coloring numbers

Let \prec be a linear ordering of vertices of a graph G . For a non-negative integer r , a vertex u is *weakly* (r, \prec) -*reachable* from a vertex $v \succeq u$ if there exists a path P from v to u in G of length at most r such that all vertices in $V(P) \setminus \{u\}$ are greater than u in the ordering \prec . Let $L_{r, \prec}(v)$ denote the set of all vertices that are weakly (r, \prec) -reachable from v , and let $R_{r, \prec}(u)$ denote set of all vertices v from which u is weakly (r, \prec) -reachable. The *weak r -coloring number* of (G, \prec) is the maximum of $|L_{r, \prec}(v)|$ over all vertices $v \in V(G)$. The *weak r -coloring number* $wcol_r(G)$ of G is the minimum weak r -coloring number of (G, \prec) over all linear orderings \prec of $V(G)$.

Theorem 3 (Zhu [5]). *For every class \mathcal{G} with bounded expansion and every $r \geq 0$, there exists an integer b such that $wcol_r(G) \leq b$ for every $G \in \mathcal{G}$.*

Expansion of shallow packings

A collection \mathcal{C} of subsets of vertices of a graph G is (ω, t) -*shallow* if every vertex of G appears in at most ω elements of \mathcal{C} and if $G[\mathcal{C}]$ has radius at most t for every $C \in \mathcal{C}$. The *packing graph* $G[\mathcal{C}]$ is defined as the graph with vertex set \mathcal{C} where C_1 and C_2 are adjacent if there exist $v_1 \in C_1$ and $v_2 \in C_2$ such that $v_1 = v_2$ or $v_1v_2 \in E(G)$. In particular, an r -shallow minor is the packing graph of a $(1, r)$ -shallow collection.

Theorem 4 (Har-Peled and Quanrud [4]). *Let G be a graph with expansion bounded by a function f . For every (ω, t) -shallow collection \mathcal{C} of subsets of vertices of G , the graph $G[\mathcal{C}]$ has expansion bounded by the function*

$$f'(r) = 5\omega^2(2t + 1)^2(2r + 1)^2 f((2t + 1)r + t).$$

Corollary 5. *For every class \mathcal{G} with polynomial expansion and all non-negative integers ω and t , there exists a class \mathcal{G}' with polynomial expansion such that the packing graph of any (ω, t) -shallow collection of subsets of vertices of a graph $G \in \mathcal{G}$ belongs to \mathcal{G}' .*

Polynomial expansion and strongly sublinear separators

A *balanced separator* in a graph G is a set $S \subseteq V(G)$ such that each component of $G - S$ has at most $\frac{2}{3}|V(G)|$ vertices. For a function $s : \mathbb{Z}^+ \rightarrow \mathbb{R}_0^+$, a class of graphs \mathcal{G} has s -*separators* if for every $G \in \mathcal{G}$, every subgraph H of G has a balanced separator of size at most $s(|V(H)|)$. A class \mathcal{G} has *strongly sublinear separators* if there exists $\varepsilon > 0$ and a function $s(n) = O(n^{1-\varepsilon})$ such that \mathcal{G} has s -separators.

Theorem 6 (Dvořák and Norin [2]). *A class of graphs has strongly sublinear separators if and only if it has polynomial expansion.*

Let \mathcal{K} be a system of subsets of vertices of a graph G . We say that \mathcal{K} is a *cover* of G if $\bigcup_{K \in \mathcal{K}} G[K] = G$. For $K \in \mathcal{K}$, let ∂K be the set of vertices of K that belong to more than one element of \mathcal{K} . The *excess* of a cover \mathcal{K} is $\sum_{K \in \mathcal{K}} |\partial K|$. The following lemma is folklore (and the proof can be found e.g. in [3]).

Lemma 7. *Let \mathcal{G} be a class of graphs with strongly sublinear separators. There exists a polynomial p such that for every $\varepsilon > 0$, every graph $G \in \mathcal{G}$ has a cover with excess at most $\varepsilon|V(G)|$ consisting of sets of size at most $p(1/\varepsilon)$.*

2 The proof

The proof of Theorem 1 follows the idea of [4], with the following crucial difference: Instead of comparing the actual solution with the optimal one, we compare it with a suitably chosen enlargement of the optimal solution. For a set O of vertices of a graph G , a linear ordering \prec of $V(G)$, and a non-negative integers r and m , we say a vertex v is (r, \prec, m, O) -rich if $|R_{r, \prec}(v) \cap O| \geq m$.

Observation 8. *Let G be a graph, O a set of vertices of G , r and m non-negative integers, and \prec a linear ordering of vertices of G . Let O' be the set of all (r, \prec, m, O) -rich vertices. If the weak r -coloring number of (G, \prec) is at most b , then*

$$|O'| \leq \frac{b}{m}|O|.$$

Proof. By the definition of richness, there are at least $m|O'|$ pairs of vertices (v, u) such that $v \in O'$ and $u \in R_{r, \prec}(v) \cap O$. However, if $u \in R_{r, \prec}(v)$, then $v \in L_{r, \prec}(u)$, and $|L_{r, \prec}(u)| \leq b$ by the assumptions. Hence, the number of such pairs is also at most $b|O|$. \square

Let us consider the setting of Observation 8, and let A be another set of vertices of G . Let $\mathcal{C}_{r, \prec, m, O, A} = \{C_v : v \in A \cup O \cup O'\}$ be a system of subsets of vertices of G defined as follows:

- For $v \in O'$, let $C_v = R_{r, \prec}(v)$.
- For $v \in O \setminus O'$, let

$$C_v = \bigcup_{x \in L_{r, \prec}(v) \setminus O'} R_{r, \prec}(x).$$

- For $v \in A \setminus (O \cup O')$, let $C_v = \{v\}$.

Lemma 9. *Let G be a graph, A and O sets of vertices of G , r and m non-negative integers, and \prec a linear ordering of vertices of G . Let O' be the set of all (r, \prec, m, O) -rich vertices. Let H be the packing graph of $\mathcal{C} = \mathcal{C}_{r, \prec, m, O, A}$. If F is a subgraph of G of diameter at most r and $V(F) \cap O \neq \emptyset$, then there exists $u \in V(F) \cap (O \cup O')$ such that $C_u C_v \in E(H)$ for every $v \in V(F) \cap A \setminus (O \cup O')$. Moreover, if the weak r -coloring number of (G, \prec) is at most b , then \mathcal{C} is $(bm + 1, 2r)$ -shallow.*

Proof. Let x be the smallest vertex of $V(F)$ in the ordering \prec . Since F has diameter at most r , for every $y \in V(F)$, there exists a path from y to x of length at most r in F , and by the choice of x , the vertices of this path appear after x in the ordering \prec . Consequently $y \in R_{r,\prec}(x)$ and $x \in L_{r,\prec}(y)$.

If $x \in O'$, let $u = x$, otherwise choose u as an arbitrary vertex of $V(F) \cap O$. In the former case, we have $V(F) \subseteq R_{r,\prec}(u) = C_u$. In the latter case, we have $x \in L_{r,\prec}(u) \setminus O'$, and thus $V(F) \subseteq R_{r,\prec}(x) \subseteq C_u$. Hence, $C_u C_v \in E(H)$ for every $v \in V(F) \cap A \setminus (O \cup O')$.

Next, let us bound the diameter of the sets in \mathcal{C} . For any $z \in V(G)$ and for every vertex w of $R_{r,\prec}(z)$, there is a path of length at most r from w to z with all vertices appearing after z in \prec ; observe that the vertices of this path also belong to $R_{r,\prec}(z)$. Therefore, every vertex of $R_{r,\prec}(z)$ is at distance at most r from z in $G[R_{r,\prec}(z)]$, and thus $G[R_{r,\prec}(z)]$ has diameter at most $2r$. For $v \in O \cup O'$, we conclude that C_v is the union of vertex sets of graphs of diameter at most $2r$, each of them containing v , and thus $G[C_v]$ has radius at most $2r$. For $v \in A \setminus (O \cup O')$, $G[C_v]$ has radius 0.

Finally, let us argue that no vertex belongs to too many of the sets in \mathcal{C} . If a vertex z belongs to C_v for $v \in O'$, then $z \in R_{r,\prec}(v)$, and thus v one of at most b vertices in $L_{r,\prec}(z)$. If z belongs to C_v for $v \in O \setminus O'$, then $z \in R_{r,\prec}(x)$ for some $x \in L_{r,\prec}(v) \setminus O'$; there are at most b choices for x in $L_{r,\prec}(z)$, and for fixed $x \notin O'$, there are less than m vertices $v \in O$ such that $x \in L_{r,\prec}(v)$ (or equivalently, $v \in R_{r,\prec}(x) \cap O$). Finally, vertex z belongs to C_v for at most one $v \in A \setminus (O \cup O')$. Therefore, z belongs to C_v for at most $b + b(m - 1) + 1$ vertices $v \in A \cup O \cup O'$. Therefore, \mathcal{C} is $(bm + 1, 2r)$ -shallow. \square

We are now ready to prove the main result.

Proof of Theorem 1. Without loss of generality, assume that $\varepsilon \leq 1$. Let r be the diameter of π . By Theorem 3, there exists b such that $\text{wcol}_r(G) \leq b$ for every $G \in \mathcal{G}$. Let $m = \lceil 3b/\varepsilon \rceil$. Let \mathcal{G}' be the class of graphs with polynomial expansion given by Corollary 5 with $\omega = bm + 1$ and $t = 2r$. Let p be the polynomial from Lemma 7 for this class \mathcal{G}' . Let us define $c = \lceil p(18/\varepsilon) \rceil$.

Consider any graph $G \in \mathcal{G}$. Let O be a π -hitting set in G of size $\gamma_\pi(G)$, and let A be the π -hitting set returned by the c -local search algorithm. Let O' be the set of all (r, \prec, m, O) -rich vertices in G ; by Observation 8, we have $|O'| \leq \frac{b}{m}|O| \leq \frac{\varepsilon}{3}|O|$. Let $\mathcal{C} = \mathcal{C}_{r,\prec,m,O,A}$; by Lemma 9, \mathcal{C} is $(bm + 1, 2r)$ -shallow, and by Corollary 5, the packing graph $H = G[\mathcal{C}]$ belongs to \mathcal{G}' . For any $Y \subseteq V(H)$, let $\overline{Y} = \{v \in V(G) : C_v \in Y\}$. By Lemma 7, there exists

a cover \mathcal{K} of H with excess at most $\frac{\varepsilon}{18}|V(H)| \leq \frac{\varepsilon}{18}(|A| + |O| + |O'|) \leq \frac{\varepsilon}{6}|A|$ such that each element of \mathcal{K} has size at most $p(18/\varepsilon) \leq c$.

Consider any $K \in \mathcal{K}$, and let $A' = (A \setminus (\overline{K \setminus \partial K})) \cup ((O \cup O') \cap \overline{K})$. We claim that A' is a π -hitting set. Indeed, suppose for a contradiction that there exists $Z \subseteq V(G)$ such that (G, Z) satisfies the property π , but $A' \cap Z = \emptyset$. Since A is a π -hitting set, there exists $v \in A \cap Z$, and since $v \notin A'$, we have $v \in (\overline{K \setminus \partial K}) \setminus (O \cup O')$. Since O is a π -hitting set, we have $Z \cap O \neq \emptyset$, and by Lemma 9 applied to $F = G[Z]$, there exists $u \in Z \cap (O \cup O')$ such that $C_u C_v \in E(H)$. Since \mathcal{K} is a cover of H , there exists $K' \in \mathcal{K}$ such that $C_u C_v \in E(H[K'])$. Since $C_v \in K \setminus \partial K$, it follows that $K' = K$, and thus this implies $C_u \in K$. Therefore, we have $u \in (O \cup O') \cap \overline{K}$, and $u \in A' \cap Z$, which is a contradiction.

Note that $A \Delta A' \subseteq \overline{K}$, and thus $|A \Delta A'| \leq c$. Hence, the c -local search algorithm considered the solution A' and did not improve A to A' , which implies that $|A'| \geq |A|$. We conclude that

$$|(O \cup O') \cap \overline{K}| \geq |A \cap (\overline{K \setminus \partial K})|$$

for every $K \in \mathcal{K}$. Therefore, denoting by s the excess of \mathcal{K} , we have

$$\begin{aligned} |A| &\leq \sum_{K \in \mathcal{K}} |A \cap \overline{K}| \leq s + \sum_{K \in \mathcal{K}} |A \cap (\overline{K \setminus \partial K})| \\ &\leq s + \sum_{K \in \mathcal{K}} |(O \cup O') \cap \overline{K}| \leq 2s + \sum_{K \in \mathcal{K}} |(O \cup O') \cap (\overline{K \setminus \partial K})| \\ &\leq 2s + |O| + |O'| \leq \frac{\varepsilon}{3}|A| + (1 + \varepsilon/3)|O|. \end{aligned}$$

Therefore,

$$|A| \leq \frac{1 + \varepsilon/3}{1 - \varepsilon/3} |O| \leq (1 + \varepsilon)|O|,$$

as required. \square

The proof for the packing version is much simpler.

Proof of Theorem 2. Without loss of generality, assume that $\varepsilon \leq 1$. Let r be the diameter of π . Let \mathcal{G}' be the class of graphs with polynomial expansion given by Corollary 5 with $\omega = 2$ and $t = r$. Let p be the polynomial from Lemma 7 for this class \mathcal{G}' . Let us define $c = \lceil p(4/\varepsilon) \rceil$.

Consider any graph $G \in \mathcal{G}$. Let O be an (induced) π -packing in G of size $\alpha_\pi(G)$ (or $\alpha'_\pi(G)$), and let A be the (induced) π -packing returned by the

c -local search algorithm. Let $\mathcal{C} = A \cup O$, and note that \mathcal{C} is $(2, r)$ -shallow. By Corollary 5, the packing graph $H = G[\mathcal{C}]$ belongs to \mathcal{G}' . By Lemma 7, there exists a cover \mathcal{K} of H with excess at most $\frac{\varepsilon}{4}|V(H)| \leq \frac{\varepsilon}{4}(|A| + |O|) \leq \frac{\varepsilon}{2}|O|$ such that each element of \mathcal{K} has size at most $p(4/\varepsilon) \leq c$.

Consider any $K \in \mathcal{K}$, and let $A' = (A \setminus K) \cup (O \cap (K \setminus \partial K))$. We claim that A' is an (induced) π -packing. Indeed, since both A and O have this property, the only way how this could be false is if there existed $P \in A \setminus K$ and $P' \in O \cap (K \setminus \partial K)$ that intersect (or contain adjacent vertices in the induced case). However, then $PP' \in E(H)$, which is a contradiction since $P' \in K \setminus \partial K$ and $P \notin K$. Observe also that $|A'| \leq |A|$, as otherwise the c -local search algorithm would extend $A \setminus K$ to a solution larger than A (if $|A'| > |A|$, then $K \not\subseteq A$, and thus $A \setminus K$ is obtained from A by deleting less than c elements).

We conclude that

$$|A \cap K| \geq |O \cap (K \setminus \partial K)|$$

for every $K \in \mathcal{K}$. Therefore, denoting by s the excess of \mathcal{K} , we have

$$\begin{aligned} |A| &\geq \sum_{K \in \mathcal{K}} |A \cap (K \setminus \partial K)| \geq -s + \sum_{K \in \mathcal{K}} |A \cap K| \\ &\geq -s + \sum_{K \in \mathcal{K}} |O \cap (K \setminus \partial K)| \geq -2s + \sum_{K \in \mathcal{K}} |O \cap K| \\ &\geq -2s + |O| \geq (1 - \varepsilon)|O|, \end{aligned}$$

as required. □

Acknowledgements

I would like to thank Abhiruk Lahiri for discussions about these problems.

References

- [1] DVOŘÁK, Z., KRÁL', D., AND THOMAS, R. Testing first-order properties for subclasses of sparse graphs. *Journal of the ACM (JACM)* 60, 5 (2013), 36.

- [2] DVOŘÁK, Z., AND NORIN, S. Strongly sublinear separators and polynomial expansion. *SIAM Journal on Discrete Mathematics* 30 (2016), 1095–1101.
- [3] HAR-PELED, S., AND QUANRUD, K. Notes on approximation algorithms for polynomial-expansion and low-density graphs. *arXiv 1603.03098* (2016).
- [4] HAR-PELED, S., AND QUANRUD, K. Approximation algorithms for polynomial-expansion and low-density graphs. *SIAM Journal on Computing* 46, 6 (2017), 1712–1744.
- [5] ZHU, X. Colouring graphs with bounded generalized colouring number. *Discrete Math.* 309, 18 (2009), 5562–5568.