

DEFORMATION AND K -THEORETIC INDEX FORMULAE ON BOUNDARY GROUPOIDS

YU QIAO AND BING KWAN SO

ABSTRACT. Boundary groupoids were introduced by the second author, which can be used to model many analysis problems on singular spaces. In order to investigate index theory on boundary groupoids, we introduce the notion of a *deformation from the pair groupoid*. Under the assumption that a deformation from the pair groupoid $M \times M$ exists for Lie groupoid $\mathcal{G} \rightrightarrows M$, we construct explicitly a deformation index map relating the analytic index on \mathcal{G} and the index on the pair groupoid. We apply this map to boundary groupoids of the form $\mathcal{G} = M_0 \times M_0 \sqcup G \times M_1 \times M_1 \rightrightarrows M = M_0 \sqcup M_1$, where G is an exponential Lie group, to obtain index formulae for (fully) elliptic (pseudo)-differential operators on \mathcal{G} , with the aid of the index formula by M. J. Pflaum, H. Posthuma, and X. Tang. These results recover and generalize our previous results for renormalizable boundary groupoids via the method of renormalized trace.

CONTENTS

1. Introduction	2
1.1. Structure of the paper	4
2. Preliminaries	5
2.1. Lie groupoids, Lie algebroids, Pseudodifferential operators on Lie groupoids, and groupoid C^* -algebras	5
2.2. Invariant submanifolds and composition series	6
2.3. Boundary groupoids and submanifold groupoids	7
2.4. The tangent groupoid and the adiabatic groupoid	9
3. The deformation groupoid and the deformation index map	10
3.1. Construction of deformations from the pair groupoid	10
3.2. Obstruction to the existence of deformations from the pair groupoid	10
3.3. The deformation index map	11
4. Fredholm and K -theoretic index of (fully) elliptic operators on boundary groupoids	13
4.1. The odd co-dimension case	14
4.2. The even co-dimension case	15
4.3. The Fredholm index for fully elliptic operators	16
References	16

Date: April 27, 2023.

Mathematics Subject Classification (2020): 58J20, 46L80, 19K56, 58H15.

Keywords: K -theoretic index, index map, deformation from the pair groupoid, Lie algebroid, boundary groupoid, submanifold groupoid.

1. INTRODUCTION

The Atiyah-Singer index theorem is one of the greatest mathematics achievements in the twentieth century, which states that the analytic index of an elliptic differential operator is equal to its topological counterpart.

There have been a plenty of results generalizing the Atiyah-Singer index theorem to other pseudo-differential calculi, constructed for different purposes [1, 3, 8, 21, 24, 31, 37]. These pseudo-differential calculi can be realized as groupoid pseudo-differential calculi on certain groupoid $\mathcal{G} \rightrightarrows M$ with M compact. Then the sub-algebras of operators of order zero and $-\infty$ can be completed to C^* -algebras $\mathfrak{U}(\mathcal{G})$ and $C^*(\mathcal{G})$, respectively, which yields the short exact sequence

$$(1) \quad 0 \rightarrow C^*(\mathcal{G}) \rightarrow \mathfrak{U}(\mathcal{G}) \rightarrow \mathfrak{U}/C^*(\mathcal{G}) \rightarrow 0.$$

For any elliptic operator of order zero, applying K -theory functor, we can define its analytic index as the image in $K_0(C^*(\mathcal{G}))$ under the boundary map. For continuous family groupoids, the analytic index is similarly defined by Lauter, Monthubert, and Nistor in [26].

In the 1990s, Connes gives a ‘‘Lie groupoid proof’’ of the Atiyah-Singer index theorem in his famous book [18, Section 2]. As the starting point, given a closed manifold M , Connes constructed the so-called *tangent groupoid*, which is a kind of *deformation* of the pair groupoid $M \times M$. This construction induces a map (which we shall denote by $\text{ind}_{\mathcal{T}(M \times M)}$ below) from the K_0 -group of the tangent bundle to

$$K_0(C^*(M \times M)) \cong K_0(\mathcal{K}).$$

Applying a mapping cone arguments as described in, for example, [1, Section 3] and [13, Section 4.1], one proves that the index map corresponding to (1) is equal to $\text{ind}_{\mathcal{T}(M \times M)}$ composed with the principal symbol. Next, Connes constructed a topological index map, by embedding M into \mathbb{R}^N for some large enough N , and considering Thom isomorphism and Morita equivalence (between groupoids). Then he showed that these two index maps coincide.

Connes’ proof of the Atiyah-Singer index theorem triggers a large number of subsequent works in index theory through Lie groupoids. For instance, based on the analysis and index problems on manifolds with corners, many groupoids were constructed [1, 10, 12, 21, 28, 32, 36]. These constructions and index theorems depend heavily on the existence of boundary defining functions and embedding the manifold under question to a cube instead of \mathbb{R}^N . Meanwhile, Androulidakis and Skandalis associate the holonomy groupoid to a singular foliation and investigated corresponding properties [4], which opens the door to explore singular foliations via Lie groupoids. Alternatively, one can study the simplified index problem on Lie groupoids by pairing with cohomology [11, 17, 38, 39, 40].

To study the Fredholm index of fully elliptic operators on manifolds with boundary, Carrillo-Rouse, Lescure, and Monthubert replace the tangent bundle in the adiabatic groupoid construction by some ‘‘non-commutative tangent bundle’’ [11]. In that case, it is (the C^* -algebra of) the continuous family groupoid

$$(M_1 \times M_1 \times \mathbb{R}) \times (0, 1) \sqcup TM \times \{0\} \rightrightarrows M \times [0, 1] \subset \mathcal{T}(\mathcal{G}).$$

Similar arguments are utilized by Debord, Lescure, and Nistor for the case of conical pseudo-manifolds [20].

In this paper, we consider index formulae for pseudo-differential on boundary groupoids of the form (with two orbits)

$$(2) \quad \mathcal{G} := M_0 \times M_0 \sqcup G \times M_1 \times M_1 \rightrightarrows M,$$

where $M_0 = M \setminus M_1$ and the isotropy subgroup G is an exponential Lie group of dimension equal to the codimension of the manifold M_1 in M . These Lie groupoids

are often holonomy groupoids integrating some singular foliations [2, 3, 5, 6, 19]. In this case, it is not clear whether an embedding analogous to the manifold with corners exists. Fortunately, the K -theory of these Lie groupoid C^* -algebras is computed in [13], namely

$$\begin{aligned} K_0(C^*(\mathcal{G})) &\cong \mathbb{Z}, & K_1(C^*(\mathcal{G})) &\cong \mathbb{Z} & \text{if } M_1 \text{ of odd codimension } \geq 3, \\ K_0(C^*(\mathcal{G})) &\cong \mathbb{Z} \oplus \mathbb{Z}, & K_1(C^*(\mathcal{G})) &\cong \{0\} & \text{if } M_1 \text{ of even codimension.} \end{aligned}$$

Hence in order to derive an index formula for elliptic operators (or just their principal symbols), it suffices to produce one integer in the odd codimension case and two integers in the even codimension case, which would completely describe its index in $K_0(C^*(\mathcal{G}))$. Moreover, the pushforward induced by *extension map*

$$K_0(C^*(M_0 \times M_0)) \xrightarrow{\varepsilon_{\mathcal{G}, M_0}} K_0(C^*(\mathcal{G}))$$

is an isomorphism in the odd case and is injective in the even case. This implies the Fredholm index of a fully elliptic operator on \mathcal{G} is determined by its $K_0(C^*(\mathcal{G}))$ index.

In the special case when a renormalized trace can be defined, we computed the index [41] using arguments similar to that of [34]. We found that in the odd case with codimension ≥ 3 , the η -term of the renormalized index formula vanishes, hence both the Fredholm and $K_0(C^*(\mathcal{G}))$ index is just given by the Atiyah-Singer integral. Moreover, one could expect a deeper description of the relationship between the isomorphism $K(C^*(M \times M)) \cong K(C^*(\mathcal{G}))$ and the vanishing of eta term.

In this paper, we take a different approach. We shall introduce the notion of a *deformation from the pair groupoid* (See Definition 3.1), where we deform the pair groupoid $M \times M$ to our desired groupoid $\mathcal{G} \rightrightarrows M$. Here, we would like to point out two major differences between our definition and that for the tangent groupoid and the adiabatic groupoid, even for the construction of the deformation to the normal cone [22, 23, 24]:

- (1) For the tangent groupoid, the fiber at $t = 0$ is the tangent bundle of M ; whereas, for a deformation from the pair groupoid, the fiber at $t = 0$ is the groupoid \mathcal{G} , usually *not* a vector bundle.
- (2) Given a closed manifold M , there always is the associated tangent groupoid. But a deformation from the pair groupoid for $\mathcal{G} \rightrightarrows M$ may not exist. (Thus, we exploit a little obstruction theory for the existence of a deformation from the pair groupoid.)

Hence, unlike the tangent groupoid and adiabatic groupoid, our idea seems to go backwards. That is, given a Lie groupoid $\mathcal{G} \rightrightarrows M$, we ask if $\mathcal{G} \rightrightarrows M$ can be obtained by deforming the pair groupoid $M \times M$, in other words, we are looking for a bigger Lie groupoid \mathcal{D} which realizes such deformation.

For a Lie groupoid $\mathcal{G} \rightrightarrows M$, once a deformation \mathcal{D} from the pair groupoid $M \times M$ exists, we are able to construct an index map

$$\text{ind}_{\mathcal{D}} : K_0(C^*(\mathcal{G})) \rightarrow K_0(C^*(M \times M \rightrightarrows M)) \cong \mathbb{Z},$$

which satisfies the following property.

Theorem 1.1. *With the notion as above, the composition*

$$K_0(C^*(M_0 \times M_0)) \cong \mathbb{Z} \xrightarrow{\varepsilon_{\mathcal{G}, M_0}} K_0(C^*(\mathcal{G})) \xrightarrow{\text{ind}_{\mathcal{D}}} K_0(C^*(M \times M)) \cong \mathbb{Z}$$

is an isomorphism.

With this theorem at hand, we establish the following main theorem.

Theorem 1.2 (Theorem 3.9). *Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid and A the associated Lie algebroid of \mathcal{G} . Suppose that a deformation \mathcal{D} from the pair $M \times M$ exists. Then one has the commutative diagram*

$$(3) \quad \begin{array}{ccc} K_0(C^*(A)) & \xrightarrow{\text{ind}_{\mathcal{T}(\mathcal{G})}} & K_0(C^*(\mathcal{G})) \\ \cong \downarrow \text{Y} & & \downarrow \text{Y} \\ K_0(C^*(TM)) & \xrightarrow{\text{ind}_{\mathcal{T}(M \times M)}} & K_0(C^*(M \times M)), \end{array} \quad \begin{array}{c} \text{ind}_{\mathcal{D}} \\ \text{Y} \end{array}$$

where the top map is just the analytic index map constructed via the adiabatic groupoid and the bottom map is the Atiyah-Singer index.

The above theorems can be used to simplify index problems on boundary groupoids to those on the pair groupoid, which enables us to apply the pairing considered in [40] to obtain our index formulae.

Theorem 1.3 (Theorems 4.3 and 4.6). *Suppose that $\mathcal{G} \rightrightarrows M$ is a boundary groupoid of the form (2), and a deformation from the pair groupoid exists for \mathcal{G} . Let Ψ be an elliptic pseudo-differential operator on \mathcal{G} . One has the index formulae:*

(1) *if $\dim G \geq 3$ is odd, then*

$$\text{ind}(\Psi) = \text{ind}_{\mathcal{T}(M \times M)}(\partial[\sigma(\Psi)]) = \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^!TM} \rangle;$$

(2) *if $\dim G$ is even, then*

$$\begin{aligned} \text{ind}(\Psi) &= \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^!TM} \rangle \\ &\quad \oplus \int_{T^*M_1 \times \mathfrak{g}} \langle \hat{A}(T^*M_1 \oplus \mathfrak{g}) \wedge \text{ch}(\sigma[\Psi]|_{M_1}), \Omega'_{\pi^!TM_1} \rangle. \end{aligned}$$

1.1. Structure of the paper. Section 2 is devoted to reviewing basic definitions and facts related to Lie groupoids, such as Lie algebroids, boundary groupoids, submanifold groupoids, the tangent groupoid. In Section 3, given a Lie groupoid $\mathcal{G} \rightrightarrows M$, we define the notion of a *deformation from the pair groupoid* to \mathcal{G} , show that such deformation exists for submanifold groupoids, and briefly discuss obstruction to existence of such deformation groupoids. Then, under the assumption that a deformation \mathcal{D} from the pair groupoid to \mathcal{G} exists, we construct explicitly an index map

$$\text{ind}_{\mathcal{D}} : K_0(C^*(\mathcal{G})) \rightarrow K_0(C^*(M \times M)) \cong \mathbb{Z},$$

which is similar to that of the tangent (or adiabatic) groupoid, and show that it is compatible with the analytic index map, hence implies Theorem 1.2. Finally in Section 4, after reviewing the Atiyah-Singer formula appearing in [40], we combine these results with Theorem 1.2 to establish index formulae for (fully) elliptic (pseudo)-differential operators on boundary groupoids of the form

$$\mathcal{G} := M_0 \times M_0 \sqcup G \times M_1 \times M_1 \rightrightarrows M,$$

where G is an exponential Lie group. These index formulae are essentially the Atiyah-Singer formula, hence we give a K -theoretic proof of the results in [41] without renormalized trace.

Acknowledgment. We would like to thank Prof. Xiang Tang for explaining his paper [40] to us. Qiao was partially supported by the NSFC (grant nos. 11971282, 12271323).

2. PRELIMINARIES

2.1. Lie groupoids, Lie algebroids, Pseudodifferential operators on Lie groupoids, and groupoid C^* -algebras. We review first basic knowledge of Lie groupoids, Lie algebroids, pseudodifferential calculus on Lie groupoids, and groupoid C^* -algebras [3, 26, 27, 29, 30, 33, 37, 42, 44].

Definition 2.1. A Lie groupoid $\mathcal{G} \rightrightarrows M$ consists of the following data:

- (i) manifolds M , called the space of units, and \mathcal{G} ;
- (ii) a unit map (inclusion) $\mathbf{u} : M \rightarrow \mathcal{G}$;
- (iii) submersions $\mathbf{s}, \mathbf{t} : \mathcal{G} \rightarrow M$, called the source and target maps respectively, satisfying

$$\mathbf{s} \circ \mathbf{u} = \text{id}_M = \mathbf{t} \circ \mathbf{u};$$

- (iv) a multiplication map $\mathbf{m} : \mathcal{G}^{(2)} := \{(g, h) \in \mathcal{G} \times \mathcal{G} : \mathbf{s}(g) = \mathbf{t}(h)\} \rightarrow \mathcal{G}$, $(g, h) \mapsto gh$, which is associative and satisfies

$$\mathbf{s}(gh) = \mathbf{s}(h), \quad \mathbf{t}(gh) = \mathbf{t}(g), \quad g(\mathbf{u} \circ \mathbf{s}(g)) = g = (\mathbf{u} \circ \mathbf{t}(g))g;$$

- (v) an inverse diffeomorphism $\mathbf{i} : \mathcal{G} \rightarrow \mathcal{G}$, $g \mapsto g = g^{-1}$, such that $\mathbf{s}(g^{-1}) = \mathbf{t}(g)$, $\mathbf{t}(g^{-1}) = \mathbf{s}(g)$, and

$$gg^{-1} = \mathbf{u}(\mathbf{t}(g)), \quad g^{-1}g = \mathbf{u}(\mathbf{s}(g)).$$

Remark 2.2. In this paper, all groupoids are assumed to be Hausdorff.

Definition 2.3. A *homomorphism* between Lie groupoids $\mathcal{G} \rightrightarrows M_1$ and $\mathcal{H} \rightrightarrows M_2$ is by definition a functor $\phi : \mathcal{G} \rightarrow \mathcal{H}$ which is smooth both on the unit space M_1 and on \mathcal{G} . Two Lie groupoids \mathcal{G} and \mathcal{H} are said to be *isomorphic* if there are homomorphisms $\phi : \mathcal{G} \rightarrow \mathcal{H}$ and $\psi : \mathcal{H} \rightarrow \mathcal{G}$ such that $\psi \circ \phi$ and $\phi \circ \psi$ are identity homomorphisms on \mathcal{G} and \mathcal{H} respectively.

Lie groupoids are closely related with Lie algebroids. Here we recall the definition.

Definition 2.4. A *Lie algebroid* A over a manifold M is a vector bundle A over M , together with a Lie algebra structure on the space $\Gamma(A)$ of the smooth sections of A and a bundle map $\nu : A \rightarrow TM$, extended to a map between sections of these bundles, such that

- (i) $\nu([X, Y]) = [\nu(X), \nu(Y)]$;
- (ii) $[X, fY] = f[X, Y] + (\nu(X)f)Y$,

for all smooth sections X and Y of A and any smooth function f on M . The map ν is called the *anchor*.

Give a Lie groupoid \mathcal{G} with units M , we can associate a Lie algebroid $A(\mathcal{G})$ to \mathcal{G} as follows. (For more details, see [29, 30].) The \mathbf{s} -vertical subbundle of $T\mathcal{G}$ for $\mathbf{s} : \mathcal{G} \rightarrow M$ is denoted by $T^{\mathbf{s}}(\mathcal{G})$ and called simply the \mathbf{s} -vertical bundle for \mathcal{G} . It is an involutive distribution on \mathcal{G} whose leaves are the components of the \mathbf{s} -fibers of \mathcal{G} . (Here involutive distribution means that $T^{\mathbf{s}}(\mathcal{G})$ is closed under the Lie bracket, i.e. if $X, Y \in \mathfrak{X}(\mathcal{G})$ are sections of $T^{\mathbf{s}}(\mathcal{G})$, then the vector field $[X, Y]$ is also a section of $T^{\mathbf{s}}(\mathcal{G})$.) Hence we obtain

$$T^{\mathbf{s}}\mathcal{G} = \text{Ker } \mathbf{s}_* = \bigcup_{x \in M} T\mathcal{G}_x \subset T\mathcal{G}.$$

The *Lie algebroid* of \mathcal{G} , denoted by $A(\mathcal{G})$ (or simply A sometimes), is defined to be $T^{\mathbf{s}}(\mathcal{G})|_M$, the restriction of the \mathbf{s} -vertical tangent bundle to the set of units M . In this case, we say that \mathcal{G} *integrates* $A(\mathcal{G})$.

Example 2.5. The Lie algebroid of the pair groupoid $M \times M$ is the tangent bundle TM with the usual Lie bracket on vector fields and the anchor map is the identity.

Two Lie groupoids may share the same Lie algebroid. For instance, both the pair groupoid $M \times M$ and the fundamental groupoid associated to M integrate to the tangent Lie algebroid TM .

Let $E \rightarrow M$ be a vector bundle. Recall [37] that an m -th order pseudo-differential operator on \mathcal{G} is a right invariant, smooth family $P = \{P_x\}_{x \in M}$, where each P_x is an m -th order classical pseudo-differential operator on sections of $\mathfrak{t}^*E \rightarrow \mathfrak{s}^{-1}(x)$. We denote by $\Psi^m(\mathcal{G}, E)$ (resp. $D^m(\mathcal{G}, E)$) the algebra of uniformly supported, order m classical pseudo-differential operators (resp. differential operators).

Recall [26] that one defines the strong norm for $P \in \Psi^0(\mathcal{G}, E)$

$$\|P\| := \sup_{\rho} \|\rho(P)\|,$$

where ρ ranges over all bounded $*$ -representations of $\Psi^0(\mathcal{G}, E)$ satisfying

$$\|\rho(P)\| \leq \sup_{x \in M} \left\{ \int_{\mathfrak{s}^{-1}(x)} |\kappa_P(g)| \mu_x, \int_{\mathfrak{s}^{-1}(x)} |\kappa_P(g^{-1})| \mu_x \right\},$$

whenever $P \in \Psi^{-\dim M-1}(\mathcal{G}, E)$ with (continuous) kernel κ_P .

Definition 2.6. The C^* -algebras $\mathfrak{U}(\mathcal{G})$ and $C^*(\mathcal{G})$ are defined to be the completion of $\Psi^0(\mathcal{G}, E)$ and $\Psi^{-\infty}(\mathcal{G}, E)$ respectively with respect to the strong norm $\|\cdot\|$.

One also defines the reduced C^* -algebras $\mathfrak{U}_r(\mathcal{G})$ and $C_r^*(\mathcal{G})$ by completing $\Psi^0(\mathcal{G}, E)$ and $\Psi^{-\infty}(\mathcal{G}, E)$ respectively with respect to the reduced norm

$$\|P\|_r := \sup_{x \in M} \{\|P_x\|_{L^2(\mathfrak{s}^{-1}(x))}\}.$$

Recall that if the strong and reduced norm coincide, then \mathcal{G} is called (*metrically*) *amenable*, which is the case for the groupoids we shall consider.

Example 2.7. Let M be a smooth manifold. Then the tangent bundle TM has a Lie groupoid structure, by regarding its fibers as a bundle of Lie groups. Its reduced groupoid C^* -algebra $C_r^*(TM)$ can be canonically identified with $C_0(T^*M)$ via Fourier transform, where T^*M is the cotangent bundle of M . This example can be generalized to vector bundles.

2.2. Invariant submanifolds and composition series. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid.

Definition 2.8. Let S be any subset of M . We denote by

$$\mathcal{G}_S := \mathfrak{s}^{-1}(S) \cap \mathfrak{t}^{-1}(S)$$

the *reduction* of \mathcal{G} to S . The reduction \mathcal{G}_S is a sub-groupoid of \mathcal{G} . In particular, if $S = \{x\}$, then $\mathcal{G}_x := \mathcal{G}_S$ is called the *isotropy group* at x .

If $S \subseteq M$ is an embedded submanifold such that $\mathfrak{s}^{-1}(S) = \mathfrak{t}^{-1}(S)$, we say that $S \subset M$ is an *invariant submanifold*.

Definition 2.9. Given a closed invariant submanifold S of \mathcal{G} . For any groupoid pseudo-differential operator $P = \{P_x\}_{x \in M} \in \Psi^m(\mathcal{G}, E)$, we define the restriction

$$P|_{\mathcal{G}_S}(P) := \{P_x\}_{x \in \mathcal{G}_S} \in \Psi^m(\mathcal{G}_S, E).$$

Restriction extends to a map from $\mathfrak{U}(\mathcal{G})$ to $\mathfrak{U}(\mathcal{G}_{M'})$ and also from $C^*(\mathcal{G})$ to $C^*(\mathcal{G}_{M'})$. We denote both such restriction maps, and also the induced K -group homomorphisms, by $r_{\mathcal{G}, S}$.

Notations 2.10. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid, U be an open subset of M . Then $\mathcal{G}_U := \mathfrak{s}^{-1}(U) \cap \mathfrak{t}^{-1}(U) \rightrightarrows U$ is an open sub-groupoid of \mathcal{G} . Any elements in $C^*(\mathcal{G}_U)$ extends to $C^*(\mathcal{G})$ by 0. We denote such extension map by $\varepsilon_{\mathcal{G},U}$. It is a homomorphism of C^* algebras and hence induces a map from $K^\bullet(C^*(\mathcal{G}_U))$ to $K^\bullet(C^*(\mathcal{G}))$, which we shall still denote by $\varepsilon_{\mathcal{G},U}$.

Now suppose we are given a groupoid $\mathcal{G} \rightrightarrows M$ (M not necessarily compact), and closed invariant submanifolds

$$M = \bar{M}_0 \supset \bar{M}_1 \supset \cdots \supset \bar{M}_r.$$

For simplicity we shall denote $\bar{\mathcal{G}}_i := \mathcal{G}_{\bar{M}_i}$. Recall that SA' denotes the sphere sub-bundle of the dual of the Lie algebroid $A(\mathcal{G})$ of \mathcal{G} .

Definition 2.11. Let $\sigma : \Psi^m(\mathcal{G}) \rightarrow C^\infty(SA')$ denotes the principal symbol map. For each $i = 1, \dots, r$, define the *joint symbol maps*

$$(4) \quad \mathbf{j}_i : \Psi^m(\mathcal{G}) \rightarrow C^\infty(SA') \oplus \Psi^m(\bar{\mathcal{G}}_i), \quad \mathbf{j}_i(P) := (\sigma(P), P|_{\bar{\mathcal{G}}_i}).$$

The map \mathbf{j}_i extends to a homomorphism from $\mathfrak{U}(\mathcal{G})$ to $C_0(SA^*) \oplus \mathfrak{U}(\bar{\mathcal{G}}_i)$.

We say that $P \in \Psi^m(\mathcal{G})$ is *elliptic* if $\sigma(P)$ is invertible, and it is called *fully elliptic* if $\mathbf{j}_1(P)$ is invertible (which implies P is elliptic).

Definition 2.12. Denote by $\mathcal{J}_0 := \overline{\Psi^{-1}(\mathcal{G})} \subset \mathfrak{U}(\mathcal{G})$, and $\mathcal{J}_i \subset \mathfrak{U}(\mathcal{G}), i = 1, \dots, r$ the null space of \mathbf{j}_{r-i+1} .

By construction, it is clear that

$$\mathcal{J}_0 \supset \mathcal{J}_1 \supset \cdots \supset \mathcal{J}_r.$$

Also, any uniformly supported kernels in $\Psi^{-\infty}(\mathcal{G}_{\bar{M}_i \setminus \bar{M}_j}, E|_{\bar{M}_i \setminus \bar{M}_j})$, can be extended to a kernel in $\Psi^{-\infty}(\mathcal{G}_{\bar{M}_i}, E|_{\bar{M}_i})$ by zero. This induces a $*$ -algebra homomorphism from $C^*(\mathcal{G}_{\bar{M}_i \setminus \bar{M}_j})$ to $C^*(\mathcal{G}_{\bar{M}_i})$. We shall use the following key fact.

Lemma 2.13. [26, Lemma 2 and Theorem 3] *One has short exact sequences*

$$(5) \quad 0 \rightarrow \mathcal{J}_{i+1} \rightarrow \mathcal{J}_i \rightarrow C^*(\mathcal{G}_{\bar{M}_i \setminus \bar{M}_{i+1}}) \rightarrow 0,$$

$$(6) \quad 0 \rightarrow C^*(\mathcal{G}_{\bar{M}_i \setminus \bar{M}_j}) \rightarrow C^*(\bar{\mathcal{G}}_i) \rightarrow C^*(\bar{\mathcal{G}}_j) \rightarrow 0, \quad \forall j > i.$$

2.3. Boundary groupoids and submanifold groupoids. In this paper we are interested in some more specific classes of groupoids. To begin with, let us recall the definition of boundary groupoids in [44, 43].

Definition 2.14. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid with M compact. We say that \mathcal{G} is a boundary groupoid if:

- (i) the singular foliation defined by the anchor map $\nu : A \rightarrow TM$ has finite number of leaves $M_0, M_1, \dots, M_r \subset M$ (which are invariant submanifolds), such that $\dim M = \dim M_0 > \dim M_1 > \cdots > \dim M_r$;
- (ii) For all $k = 0, 1, \dots, r$, $\bar{M}_k := M_k \cup \cdots \cup M_r$ are closed submanifolds of M ;
- (iii) For $k = 0$, $\mathcal{G}_0 := \mathcal{G}_{M_0}$ is the pair groupoid, and for $k = 1, 2, \dots, r$, we have $\mathcal{G}_k := \mathcal{G}_{M_k} \cong G_k \times M_k \times M_k$ for some Lie group G_k ;
- (iv) For each $k = 0, 1, \dots, r$, there exists an unique sub-bundle $\bar{A}_k \subset A|_{\bar{M}_k}$ such that $\bar{A}_k|_{M_k} = \ker(\nu|_{M_k}) (= \mathfrak{g}_k \times M_k)$.

Boundary groupoids are closely related to Fredholm groupoids and blowup groupoids. Roughly speaking, Fredholm groupoids are those on which Fredholmness of a pseudodifferential operator is completely characterized by its ellipticity and invertibility at the boundary. For the definition and basic properties of Fredholm groupoids, one may consult [14, 15, 16]. The basic result relevant to our discussion is that boundary groupoids are often amenable and Fredholm groupoids.

Lemma 2.15. (See [27, Lemma 7]) For any boundary groupoid of the form $\mathcal{G} = (M_0 \times M_0) \cup (\mathbb{R}^q \times M_1 \times M_1)$,

$$C^*(\mathcal{G}) \cong C_r^*(\mathcal{G}).$$

In other words, \mathcal{G} is (metrically) amenable.

Moreover, since the additive group \mathbb{R}^q is amenable, the groupoid of the form $\mathbb{R}^q \times M_1 \times M_1$ is (topologically) amenable. By [16, Theorem 4.3], we have the following proposition.

Proposition 2.16. The groupoid $\mathcal{G} = (M_0 \times M_0) \cup (\mathbb{R}^q \times M_1 \times M_1)$ is a Fredholm groupoid.

Given a boundary groupoid, one naturally considers the sequence of invariant submanifolds

$$M \supset \bar{M}_1 \supset \cdots \supset \bar{M}_r,$$

where \bar{M}_i is given to be (2) of Definition 2.14. We have the short exact sequence

$$(7) \quad 0 \rightarrow C^*(\mathcal{G}_{\bar{M}_i \setminus \bar{M}_j}) \rightarrow C^*(\bar{\mathcal{G}}_i) \rightarrow C^*(\bar{\mathcal{G}}_j) \rightarrow 0, \quad \forall j > i,$$

which induces the K -theory six-terms exact sequences:

$$(8) \quad \begin{array}{ccccc} K_1(C^*(\mathcal{G}_{M_i})) & \longrightarrow & K_1(C^*(\bar{\mathcal{G}}_i)) & \longrightarrow & K_1(C^*(\bar{\mathcal{G}}_{i+1})) \\ \uparrow & & & & \downarrow \\ K_0(C^*(\bar{\mathcal{G}}_{i+1})) & \longleftarrow & K_0(C^*(\bar{\mathcal{G}}_i)) & \longleftarrow & K_0(C^*(\mathcal{G}_{M_i})) \end{array}$$

If $r = 1$, and G_1 is solvable, connected and simply connected (i.e. exponential), then the system (8) is greatly simplified – to begin with, there is only one exact sequence. Indeed, Carrillo-Rouse and the second author computed the K -theory of boundary groupoids with $r = 1$ in [13], namely:

$$\begin{array}{ll} K_0(C^*(\mathcal{G})) \cong \mathbb{Z}, & K_1(C^*(\mathcal{G})) \cong \mathbb{Z}, & \text{if } \dim G \geq 3 \text{ odd;} \\ K_0(C^*(\mathcal{G})) \cong \{0\}, & K_1(C^*(\mathcal{G})) \cong \mathbb{Z} \oplus \mathbb{Z}, & \text{if } \dim G \text{ even.} \end{array}$$

Next, we recall *submanifold groupoids*, which form a sub-class of boundary groupoids in [13].

Example 2.17 (Submanifold groupoids). Suppose that M_1 is a closed embedded sub-manifold of M of codimension $q \geq 2$. Let $f \in C^\infty(M)$ be any smooth function which vanishes on M_1 and strictly positive over $M_0 = M \setminus M_1$ (if M is connected then $M_0 = M \setminus M_1$ is still connected and f must either be strictly positive or strictly negative on M_0). Then there exists a Lie algebroid structure over

$$A_f := TM \rightarrow M$$

with anchor map

$$(z, w) \mapsto (z, f(z) \cdot w), \quad \forall (z, w) \in TM,$$

and Lie bracket on sections

$$[X, Y]_{A_f} := f[X, Y] + (X \cdot f)Y - (Y \cdot f)X.$$

This Lie algebroid is almost injective (since the anchor is injective, in fact an isomorphism, over the open dense subset $M_0 = M \setminus M_1$), and hence it integrates to a Lie groupoid that is a quasi-graphoid (see [19, Theorems 2 and 3]).

By [35], A_f integrates to a Lie groupoid of the form

$$\mathcal{G} = M_0 \times M_0 \sqcup \mathbb{R}^q \times M_1 \times M_1 \rightrightarrows M = M_0 \sqcup M_1.$$

This groupoid is called a submanifold groupoid.

Example 2.18 (Renormalizable boundary groupoids). For even more specific examples of submanifold groupoids, in [41], we considered

$$f = r^N$$

for some fixed even integers N , where $r := d(M_1, \cdot)$ is the Riemannian distance function from M_1 with respect to some metric. The resulting groupoid is called a *renormalizable boundary groupoid*. In [41], we showed that the η -term for renormalizable groupoids vanishes in the index formula via the method of renormalized trace.

2.4. The tangent groupoid and the adiabatic groupoid. Let us recall that for a closed manifold M , Connes [18] constructed the tangent groupoid, which is a Lie groupoid of the form

$$\mathcal{T}(M \times M) := M \times M \times (0, 1] \sqcup TM \times \{0\} \rightrightarrows M \times [0, 1],$$

where the differentiable structure of $\mathcal{T}(M \times M)$ is defined by fixing some Riemannian metric on M and then using the following maps from some open subsets of $TM \times [0, 1]$ to $\mathcal{T}(M \times M)$

$$\begin{aligned} \mathbf{x}(p, X, \epsilon) &:= (p, \exp_p(-\epsilon X), \epsilon), \text{ for } \epsilon > 0, \\ \mathbf{x}(p, X, 0) &:= (p, X, 0), \end{aligned}$$

as charts.

The tangent groupoid construction generalizes to any Lie groupoid \mathcal{G} . One considers the adiabatic groupoid

$$\mathcal{T}(\mathcal{G}) := \mathcal{G} \times (0, 1] \sqcup A \times \{0\} \rightrightarrows M \times [0, 1],$$

where A is the Lie algebroid of \mathcal{G} . Debord and Skandalis [24] further generalized the construction by replacing the set of units $M \subset \mathcal{G}$ with arbitrary sub-groupoid $\mathcal{H} \subset \mathcal{G}$, and considered gluing $\mathcal{N}_{\mathcal{H}}^{\mathcal{G}}$, the normal bundle of \mathcal{H} in \mathcal{G} , to $\mathcal{G} \times \mathbb{R} \setminus \{0\}$. The resulting object is naturally a Lie groupoid, which they call the deformation to the normal cone.

For simplicity we return to the case of the adiabatic groupoid $\mathcal{T}(\mathcal{G})$. One naturally constructs the index map

$$(9) \quad \text{ind}_{\mathcal{T}(\mathcal{G})} := r_{\mathcal{T}(\mathcal{G}), M \times \{1\}} \circ r_{\mathcal{T}(\mathcal{G}), M \times \{0\}}^{-1},$$

where $r_{\mathcal{T}(\mathcal{G}), M \times \{1\}} : C^*(\mathcal{T}(\mathcal{G})) \rightarrow C^*(\mathcal{G})$ and $r_{\mathcal{T}(\mathcal{G}), M \times \{0\}} : C^*(\mathcal{T}(\mathcal{G})) \rightarrow C^*(A)$ are respectively the restriction maps to the sub-groupoids of $\mathcal{T}(\mathcal{G})$ over $M \times \{1\}$ and $M \times \{0\}$.

Let Ψ be any classical elliptic pseudo-differential operator on $\mathcal{G} \rightrightarrows M$. Its principal symbol $\sigma(\Psi)$ is an invertible element in $C(SA')$, where A' is the dual bundle of A and SA' denotes the sphere bundle of A' . We identify the algebra of Schwartz functions on A' with the (fiber-wise) convolution algebra $C^*(A)$ through Fourier transform to obtain

$$K_0(C(A')) \cong K_0(C^*(A)),$$

and let

$$\partial : K_1(SA') \rightarrow K_0(C(A')) \cong K_0(C^*(A))$$

be the connecting map induced by the short exact sequence

$$0 \rightarrow C(A') \rightarrow C(\bar{A}') \rightarrow (SA') \rightarrow 0.$$

Then by [13], the analytic index of Ψ is given by

$$\text{ind}_{\mathcal{T}(\mathcal{G})}(\partial[\sigma(\Psi)]) \in C^*(\mathcal{G}),$$

which, if Ψ is of order 0, coincides with the image of $[\Psi] \in K_1(\mathfrak{U}(\mathcal{G}))$ under the connecting map induced by the short exact sequence

$$0 \rightarrow C^*(\mathcal{G}) \rightarrow \mathfrak{U}(\mathcal{G}) \rightarrow \mathfrak{U}/C^*(\mathcal{G}) \rightarrow 0.$$

3. THE DEFORMATION GROUPOID AND THE DEFORMATION INDEX MAP

Motivated by the tangent groupoid construction in the previous section, we introduce the following definition, which plays a central role in the paper.

Definition 3.1. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. A deformation from the pair groupoid $M \times M$ to \mathcal{G} is a Lie groupoid $\mathcal{D} \rightrightarrows M \times [0, 1]$ with (groupoid) isomorphisms

$$\begin{aligned} \mathcal{D}_{M \times (0,1]} &\cong M \times M \times (0, 1], \\ \mathcal{D}_{M \times \{0\}} &\cong \mathcal{G}. \end{aligned}$$

Example 3.2. Let M be a closed manifold. Regard $TM \rightarrow M$ as a Lie groupoid, then Connes' tangent groupoid, $\mathcal{T}(M \times M)$ gives a deformation from the pair groupoid to TM .

3.1. Construction of deformations from the pair groupoid. The following fact is another motivation to introduce the above definition.

Proposition 3.3. *There exists a deformation from the pair groupoid for submanifold groupoids defined in Example 2.17.*

Proof. We use the same notation as in Example 2.17. To construct the groupoid $\mathcal{D} \rightrightarrows M \times [0, 1]$, we begin with defining the Lie algebroid of \mathcal{D} . We set $\tilde{A}_f \rightarrow M \times [0, 1]$ to be the vector bundle pullback of A_f . Sections of \tilde{A}_f are just vector fields on $M \times [0, 1]$ which are tangential to the M direction. Hence one can define the anchor map

$$\nu((z, s), w) = ((z, s), (f(z) + s) \cdot w), \quad \forall (z, w) \in TM, s \in [0, 1],$$

and Lie bracket on sections by

$$[X, Y]_{\tilde{A}_f} := (f + s)[X, Y] + (X \cdot f)Y - (Y \cdot f)X.$$

The Lie algebroid \tilde{A}_f is isomorphic to A_f on $M \times \{0\}$ and TM on $M \times \{s\}$ for each $s > 0$, respectively. Using [19] and [35, Theorem 1], one observes that \tilde{A}_f integrates to a Lie groupoid of the form

$$\mathcal{G} \sqcup (M \times M \times (0, 1]) \rightrightarrows M \times [0, 1],$$

which we denote by \mathcal{D} . □

3.2. Obstruction to the existence of deformations from the pair groupoid.

In this subsection we point out some notable necessary condition for \mathcal{G} in order for a deformation from the pair groupoid to exist.

Proposition 3.4. *Suppose that $\mathcal{G} \rightrightarrows M$ is a Lie groupoid such that a deformation from the pair groupoid exists. Denote by A the Lie algebroid of \mathcal{G} . Then we have*

$$A \cong TM,$$

as vector bundles.

Proof. Let $\mathcal{D} \rightrightarrows M \times [0, 1]$ be a deformation from the pair groupoid. Denote its Lie algebroid by $\tilde{A} \rightarrow M \times [0, 1]$. Then one has

$$\tilde{A}|_{M \times \{0\}} \cong A, \quad \tilde{A}|_{M \times \{1\}} \cong TM.$$

Hence an isomorphism can be constructed by, say, fixing a connection $\tilde{\nabla}$ on \tilde{A} and then parallel transport along the family of curves $(p, t), t \in [0, 1]$ for each $p \in M$. □

The existence of a deformation from the pair groupoid also imposes necessary conditions on the structural vector fields of \mathcal{G} (i.e. the image of the anchor map). For simplicity, we consider the case when M_1 is a single point $\{p\}$. Suppose a deformation \mathcal{D} from the pair groupoid for \mathcal{G} exists. Denote the Lie algebroids of \mathcal{G} and \mathcal{D} by A and \tilde{A} respectively, and their anchor maps by ν and respectively $\tilde{\nu}$. Then it is obvious that

$$\begin{aligned}\tilde{A}|_{M \times (0,1]} &\cong TM \times (0,1], \\ \tilde{A}|_{M \times \{0\}} &\cong A.\end{aligned}$$

Suppose that s is a nowhere vanishing local section of A around p , such that $\nu(s)(p) = 0$. Then the section s extends to some nowhere vanishing local section \tilde{s} of \tilde{A} over an open neighborhood of $\{p\} \times [0,1]$. Because $\tilde{\nu}$ is injective except at $(p,0)$, it follows that $\tilde{\nu}(\tilde{s})$ is a family of local vector fields on M parameterized by $[0,1]$, which is non-vanishing on $M \times \{t\}$, $1 \geq t > 0$ and has an isolated zero at p on $M \times \{0\}$. This implies that the Hopf-Poincaré index of $\nu(s) = \tilde{\nu}(\tilde{s})|_{M \times \{0\}}$ equals zero. Hence, we are able to make the following counterexample which shows that a boundary groupoid may not possess a deformation from the pair groupoid.

Example 3.5. There exist Lie algebroids spanned by structural vector fields of non-zero degrees. For example, on \mathbb{R}^2 , the vector fields

$$x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} \quad \text{and} \quad y \frac{\partial}{\partial x} - x \frac{\partial}{\partial y}$$

have isolated zero of degree 1, and commute each other. Moreover, their flows give an action of the abelian Lie group \mathbb{R}^2 on itself. Thus the tangent Lie algebroid of the action groupoid $\mathbb{R}^2 \rtimes \mathbb{R}^2$ is spanned by these vector fields, which contradicts to the above necessary condition that the Hopf-Poincaré index is equal to zero.

Remark 3.6. The above example exhibits that there are certain obstructions to the existence of such deformation for boundary groupoids. At present, besides Proposition 3.4, we do not know of any other necessary conditions or sufficient conditions to guarantee the existence of such deformations in general, which will be left as future work.

3.3. The deformation index map. In this subsection, we always assume that a deformation from the pair groupoid

$$\mathcal{D} := \mathcal{G} \sqcup (M \times M \times (0,1]) \rightrightarrows M \times [0,1]$$

exists for the given Lie groupoid $\mathcal{G} \rightrightarrows M$.

We shall use \mathcal{D} to construct an index, similar to [18, 45]. Observe that \mathcal{D} has closed saturated subgroupoids $\mathcal{D}|_{M \times \{0\}} \cong \mathcal{G}, \mathcal{D}|_{M \times \{1\}} \cong M \times M$. Because the sequence

$$\begin{aligned}K^\bullet(C^*(\mathcal{D}|_{M \times (0,1]})) &\cong K^\bullet(C^*(M \times M \times (0,1])) \cong 0 \xrightarrow{\varepsilon_{\mathcal{D}, M \times (0,1]}} K^\bullet(C^*(\mathcal{D})) \\ &\xrightarrow{r_{\mathcal{D}, M \times \{0\}}} K^\bullet(C^*(\mathcal{G})) \xrightarrow{\partial} K^{\bullet+1}(C^*(\mathcal{D}|_{M \times (0,1]}))\end{aligned}$$

is exact, therefore the map $r_{\mathcal{D}, M \times \{0\}}$ is invertible. Hence it is natural to introduce the following index map.

Definition 3.7. The deformation index map is defined to be

$$\text{ind}_{\mathcal{D}} := r_{\mathcal{D}, M \times \{1\}} \circ r_{\mathcal{D}, M \times \{0\}}^{-1} : K^\bullet(C^*(\mathcal{G})) \rightarrow K^\bullet(C^*(M \times M)).$$

From the definition, the following theorem is immediate, and implies Theorem 1.1.

Theorem 3.8. *One has*

$$\text{ind}_{\mathcal{D}} \circ \varepsilon_{\mathcal{G}, M_0} = \varepsilon_{M \times M, M_0}.$$

Proof. Observe that

$$M_0 \times M_0 \times [0, 1] = \mathcal{D}_{M_0 \times [0, 1]}$$

is an open subgroupoid. For any $u \in C^*(M_0 \times M_0)$, let $\tilde{u} \in M_0 \times M_0 \times [0, 1]$ be the pullback of s by the projection to the $M_0 \times M_0$ factor (i.e., \tilde{u} is just u on each $M_0 \times M_0 \times \{s\}$). Extend \tilde{u} to \mathcal{D} and get $\varepsilon_{\mathcal{D}, M_0 \times [0, 1]}(\tilde{u})$. When restricted to $s = 0$ and $s = 1$, it is clear that $\varepsilon_{\mathcal{D}, M_0 \times [0, 1]}(\tilde{u})$ equals

$$\varepsilon_{\mathcal{G}, M_0}(u) \quad \text{and} \quad \varepsilon_{M \times M, M_0}(u),$$

respectively. Passing to K -theory, one obtains

$$r_{\mathcal{D}, M \times \{0\}} \circ \varepsilon_{\mathcal{D}, M_0 \times [0, 1]}([\tilde{u}]) = \varepsilon_{\mathcal{G}, M_0}([u]),$$

and

$$r_{\mathcal{D}, M \times \{1\}} \circ \varepsilon_{\mathcal{D}, M_0 \times [0, 1]}([\tilde{u}]) = \varepsilon_{M \times M, M_0}([u]),$$

for any class $[u] \in K^\bullet(M_0 \times M_0)$. Hence, we have

$$\begin{aligned} \text{ind}_{\mathcal{D}} \circ \varepsilon_{\mathcal{G}, M_0}([u]) &= r_{\mathcal{D}, M \times \{1\}} \circ r_{\mathcal{D}, M \times \{0\}}^{-1} \circ r_{\mathcal{D}, M \times \{0\}} \circ \varepsilon_{\mathcal{D}, M_0 \times [0, 1]}([\tilde{u}]) \\ &= \varepsilon_{M \times M, M_0}([u]), \end{aligned} \quad \square$$

which completes the proof.

We turn to the proof of Theorem 1.2, which is reformulated as follows.

Theorem 3.9. *Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. Suppose there exists a deformation groupoid \mathcal{D} from the pair groupoid $M \times M$. Then one has the commutative diagram*

$$(10) \quad \begin{array}{ccc} K_0(C^*(A)) & \xrightarrow{\text{ind}_{\mathcal{T}(\mathcal{G})}} & K_0(C^*(\mathcal{G})) \\ \cong \downarrow & & \downarrow \text{ind}_{\mathcal{D}} \\ K_0(C^*(TM)) & \xrightarrow{\text{ind}_{\mathcal{T}(M \times M)}} & K_0(C^*(M \times M)), \end{array}$$

where the top map is just the analytic index map constructed via the adiabatic groupoid and the bottom map is the Atiyah-Singer index.

Proof. Let $\mathcal{T}(\mathcal{D}) \rightrightarrows (M \times [0, 1]) \times [0, 1]$ be the adiabatic groupoid of \mathcal{D} . Recall that as a set $\mathcal{T}(\mathcal{D}) = \mathcal{D} \times (0, 1] \sqcup \tilde{A}$. Hence $\mathcal{T}(\mathcal{D})$ restricted to $(M \times \{0\}) \times [0, 1]$ and $(M \times \{1\}) \times [0, 1]$ are respectively

$$\mathcal{T}(\mathcal{G}) \rightrightarrows M \times [0, 1] = \mathcal{G} \times (0, 1] \sqcup A,$$

$$\text{and } \mathcal{T}(M \times M) \rightrightarrows M \times [0, 1] = M \times M \times (0, 1] \sqcup TM,$$

the adiabatic groupoid of \mathcal{G} (respectively the adiabatic groupoid of $M \times M$). One can further restrict $\mathcal{T}(\mathcal{G})$ and $\mathcal{T}(M \times M)$ to $M \times \{0\}$ or $M \times \{1\}$, resulting in the commutative diagram

$$(11) \quad \begin{array}{ccccc} C^*(A) < & \xrightarrow{r_{\mathcal{T}(\mathcal{G}), M \times \{0\}}} & C^*(\mathcal{T}(\mathcal{G})) & \xrightarrow{r_{\mathcal{T}(\mathcal{G}), M \times \{1\}}} & > C^*(\mathcal{G}) \\ \wedge & & \wedge & & \wedge \\ r_{\tilde{A}, M \times \{0\}} & & r_{\mathcal{T}(\mathcal{D}), (M \times \{0\}) \times [0, 1]} & & r_{\mathcal{D}, M \times \{0\}} \\ C^*(\tilde{A}) < & \xrightarrow{r_{\mathcal{T}(\mathcal{D}), (M \times [0, 1]) \times \{0\}}} & C^*(\mathcal{T}(\mathcal{D})) & \xrightarrow{r_{\mathcal{T}(\mathcal{D}), (M \times [0, 1]) \times \{1\}}} & > C^*(\mathcal{D}) \\ \wedge & & \wedge & & \wedge \\ r_{\tilde{A}, M \times \{1\}} & & r_{\mathcal{T}(\mathcal{D}), (M \times \{1\}) \times [0, 1]} & & r_{\mathcal{D}, M \times \{1\}} \\ C^*(TM) < & \xrightarrow{r_{\mathcal{T}(M \times M), M \times \{0\}}} & C^*(\mathcal{T}(M \times M)) & \xrightarrow{r_{\mathcal{T}(M \times M), M \times \{1\}}} & > C^*(M \times M). \end{array}$$

Then we pass to the corresponding K -group maps. Observe that $r_{\mathcal{T}(\mathcal{D}), (M \times \{0\}) \times [0,1]}$ fits in the six-term exact sequence

$$\begin{aligned} K^\bullet(C^*(\mathcal{T}(\mathcal{D})_{(M \times (0,1]) \times [0,1]})) &\xrightarrow{\varepsilon_{\mathcal{T}(\mathcal{D}), (M \times (0,1]) \times [0,1]}} K^\bullet(C^*(\mathcal{T}(\mathcal{D}))) \\ &\xrightarrow{r_{\mathcal{T}(\mathcal{D}), (M \times \{0\}) \times [0,1]}} K^\bullet(C^*(\mathcal{T}(\mathcal{G}))) \rightarrow \cdots, \end{aligned}$$

and moreover we have

$$\mathcal{T}(\mathcal{D})_{(M \times (0,1]) \times [0,1]} \cong T(\mathcal{D}_{M \times (0,1]}) \cong \mathcal{T}(M \times M) \times (0, 1],$$

whose convolution C^* -algebra is contractible. Therefore the map $r_{\mathcal{T}(\mathcal{D}), (M \times \{0\}) \times [0,1]}$ is invertible. Similarly, the maps $r_{\mathcal{T}(\mathcal{G}), M \times \{0\}}$, $r_{\tilde{A}, M \times \{0\}}$, $r_{\mathcal{T}(\mathcal{D}), (M \times [0,1]) \times \{0\}}$, $r_{\mathcal{T}(M \times M), M \times \{0\}}$ and $r_{\mathcal{D}, M \times \{0\}}$ are all isomorphisms between corresponding K -groups.

Recall (9) that

$$\begin{aligned} \text{ind}_{\mathcal{T}(\mathcal{G})} &:= r_{\mathcal{T}(\mathcal{G}), M \times \{1\}} \circ r_{\mathcal{T}(\mathcal{G}), M \times \{0\}}^{-1}, \\ \text{ind}_{\mathcal{T}(M \times M)} &:= r_{\mathcal{T}(M \times M), M \times \{1\}} \circ r_{\mathcal{T}(M \times M), M \times \{0\}}^{-1} \end{aligned}$$

are just Connes' analytic index maps. Lastly, using the same pull-back arguments as the proof of Theorem 3.8, one sees that the K -theory map

$$r_{\tilde{A}, M \times \{1\}} \circ r_{\tilde{A}, M \times \{0\}}^{-1}$$

is just equivalent to the isomorphism in Proposition 3.4. Hence one ends up with the commutative diagram

$$(12) \quad \begin{array}{ccc} K_0(C^*(A)) & \xrightarrow{\text{ind}_{\mathcal{T}(\mathcal{G})}} & K_0(C^*(\mathcal{G})) \\ \cong \downarrow & & \downarrow \text{ind}_{\mathcal{D}} \\ K_0(C^*(TM)) & \xrightarrow{\text{ind}_{\mathcal{T}(M \times M)}} & K_0(C^*(M \times M)), \end{array}$$

which completes the proof. \square

4. FREDHOLM AND K -THEORETIC INDEX OF (FULLY) ELLIPTIC OPERATORS ON BOUNDARY GROUPOIDS

In this section, we combine the map in Theorem 1.2 and the Atiyah-Singer index formula to compute the index. Before doing that, we briefly recall the Atiyah-Singer index formula, in particular, we shall use the version appearing in [40, Theorem 5.1].

Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid and A the associated Lie algebroid of \mathcal{G} . Define the line bundle

$$L := \wedge^{\text{top}} T^*M \otimes \wedge^{\text{top}} A.$$

We suppose further that \mathcal{G} is unimodular, i.e., there exists an invariant nowhere vanishing section Ω of L . Then the characteristic map χ_Ω defines a map from the groupoid cohomology to cyclic cohomology by [40, Equation (5)]

$$\begin{aligned} &\chi_\Omega(\varphi_1 \otimes \cdots \otimes \varphi_k)(a_0 \otimes \cdots \otimes a_k) \\ &:= \int_M \left(\int_{g_0 \cdots g_k = 1_x} a_0(g_0) \varphi_1(g_1) a_1(g_1) \cdots \varphi_k(g_k) a_k(g_k) \right) \Omega(x) \end{aligned}$$

(on the level of cochains). Then one combines χ_Ω with the canonical pairing between cyclic homology and cohomology, the Connes-Chern character map, which maps from $K_0(\Psi_c^{-\infty}(\mathcal{G}))$ to the cyclic homology of $\Psi_c^{-\infty}(\mathcal{G})$, and the van Est map, to obtain a pairing

$$\langle [\varphi], \alpha \rangle_\Omega \in \mathbb{C}$$

for any $[\varphi] \in K_0(\Psi_c^{-\infty}(\mathcal{G}))$, $\alpha \in H^\bullet(A, \mathbb{C})$.

Then the main result of [40, Theorem 5.1] states that such pairing can be computed by the following formula.

Lemma 4.1. *For any elliptic pseudo-differential operator Ψ and $\alpha \in H^{2k}(A, \mathbb{C})$, we have*

$$\langle \text{ind}(\Psi), \alpha \rangle_{\Omega} = (2\pi\sqrt{-1})^{-k} \int_{A'} \langle \pi^* \alpha \wedge \hat{A}(A') \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^! A} \rangle,$$

where $\hat{A}(A') \in H^{\bullet}(TM)$ is the A -hat genus, $\text{ch} : K_0(C(A')) \rightarrow H^{\text{even}}(TM)$ is the Chern Character (which can be defined by, say, the Chern-Weil construction), and $\pi^! A$ is the pull-back Lie algebroid of A along the projection $\pi : A' \rightarrow M$.

Let us consider the particular case when $\mathcal{G} = M \times M$ is the pair groupoid, hence $\mathcal{A} = TM$, and L is trivial. One obtains an invariant nowhere vanishing section Ω of L by considering,

$$(13) \quad \Omega := C dx^1 \wedge \cdots \wedge dx^{\dim M} \otimes \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^{\dim M}}$$

locally, for any constant $C \neq 0$. Also, the constant function 1 obviously defines a class $[1] \in H^0(A, \mathbb{C})$. Applying Lemma 4.1, one gets

$$(14) \quad \langle \text{ind}(\Psi), 1 \rangle_{\Omega} = \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^! TM} \rangle.$$

Since $\text{ind}(\Psi)$ is an integer, by choosing an appropriate normalization C for Ω , we have

$$\text{ind}(\Psi) = \langle \text{ind}(\Psi), 1 \rangle_{\Omega},$$

and the identity (14) simplifies to

$$(15) \quad \text{ind}(\Psi) = \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^! TM} \rangle.$$

Remark 4.2. In order to apply Lemma 4.1, one needs a non-trivial \mathcal{G} -invariant Lie algebroid class. By [25] the construction of such class is non-trivial, because the Lie algebroid associated to a boundary groupoid is in general not unimodular. However, in the following subsections we shall use a deformation from the pair groupoid to simplify the problem to that of a regular groupoid, which is clearly unimodular, so that the Lemma 4.1 can be applied.

In the sequel, we focus on boundary groupoids of the form

$$(16) \quad \mathcal{G} := M_0 \times M_0 \sqcup G \times M_1 \times M_1 \rightrightarrows M = M_0 \sqcup M_1,$$

where G is an exponential Lie group, and assume that a deformation from the pair groupoid exists for \mathcal{G} .

4.1. The odd co-dimension case. Recall that if the isotropy subgroup G is exponential and of odd dimension ≥ 3 , then

$$(17) \quad K_0(M_0 \times M_0) \cong \mathbb{Z} \xrightarrow{\varepsilon_{\mathcal{G}, M_0}} C^*(\mathcal{G}) \cong \mathbb{Z}$$

is an isomorphism. Identifying $K_0(M_0 \times M_0) \cong K_0(M \times M)$, one sees from Theorem 1.1 that $\text{ind}_{\mathcal{D}}$ is the inverse of $\varepsilon_{\mathcal{G}, M_0}$.

Given any elliptic pseudodifferential operator Ψ , in order to compute the integer

$$\text{ind}(\Psi) = \text{ind}_{\mathcal{T}(\mathcal{G})}(\partial[\sigma(\Psi)]),$$

we apply Theorem 1.2 to obtain

$$(18) \quad \text{ind}(\Psi) = \text{ind}_{\mathcal{T}(\mathcal{G})}(\partial[\sigma(\Psi)]) = \text{ind}_{\mathcal{T}(M \times M)}(\partial[\sigma(\Psi)]),$$

where we regard $\partial[\sigma] \in K_0(C^*(TM))$ on the rightmost expression. Observe that $\text{ind}_{\mathcal{T}(M \times M)}(\partial[\sigma])$ is just the analytic index of the pair groupoid, Hence we apply Lemma 4.1 (with the normalization of (15)) to conclude the following theorem.

Theorem 4.3. *Suppose that $\mathcal{G} \rightrightarrows M$ is a boundary groupoid of the form (16) with $\dim G \geq 3$ odd, and a deformation from the pair groupoid exists for \mathcal{G} . Let Ψ be an elliptic pseudo-differential operator on \mathcal{G} . One has the index formula*

$$(19) \quad \text{ind}(\Psi) = \text{ind}_{\mathcal{T}(M \times M)}(\partial[\sigma(\Psi)]) = \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^*TM} \rangle.$$

Remark 4.4. By Proposition 3.3, we see that a deformation from the pair groupoid always exists for submanifold groupoids. Because renormalizable boundary groupoids form a special class of submanifold groupoids, Theorem 4.3 applies and implies that the index is given only by the Atiyah-Singer term. Comparing with the results of [8, 9] (for higher co-dimension) and also [41], one sees that the η -term vanishes for elliptic pseudo-differential operators on \mathcal{G} . Therefore we give a completely new proof that generalizes these previous results.

Example 4.5. Recall [27, Section 6] that for any Lie groupoid the vertical de Rham operator d is an (invariant) groupoid differential operator. Its vector representation computes its Lie algebroid cohomology. Fixing a Riemannian metric on A , one can then define the formal adjoint d^* of d , which leads one to construct the Euler operator $d + d^*$, and also the signature operator. More generally, one can construct generalized Dirac operators.

For illustration here, we only consider the case when the Lie algebroid A is spin, and \mathcal{G} is of the form (16) with M_1 of odd codimension, where a deformation from the pair groupoid exists. Let S_A be the spinor bundle of A . Following [27, Section 6], one constructs the (groupoid) Dirac operator D_A associated with the Levi-Civita connection. On the other hand, Lemma 3.4 implies that M is also a spin manifold, and the spinor bundle S_{TM} is isomorphic to S_A . Hence it is standard to construct the (groupoid) Dirac operator D associated with the Levi-Civita connection. One sees from its explicit construction that the principal symbol of D_A is equal to that of D (however D_A and D are very different as groupoid differential operators). Hence by Theorem 3.9, the index of D_A is the same as the index of D . Theorem 4.3 then gives an explicit formula for $\text{ind}(D_A)$. Lastly recall [7, Section 3] that one can generalize the construction of the Dirac operator by tensoring S_A with a vector bundle W , and in this case the Chern form $\text{ch}(\sigma[D_A])$ can be written explicitly using the twisting curvature.

4.2. The even co-dimension case. On the other hand, if G is of even dimension, one has short exact sequence

$$(20) \quad 0 \rightarrow K_0(M_0 \times M_0) \cong \mathbb{Z} \xrightarrow{\varepsilon_{\mathcal{G}, M_0}} K_0(C^*(\mathcal{G})) \xrightarrow{r_{\mathcal{G}, M_1}} K_0(C^*(\mathcal{G}_{M_1})) \cong \mathbb{Z} \rightarrow 0$$

and Theorem 1.1 canonically identifies

$$K_0(C^*(\mathcal{G})) \cong \mathbb{Z} \oplus \mathbb{Z}$$

via $\text{ind}_{\mathcal{D}} \oplus r_{\mathcal{G}, M_1}$. For the $\text{ind}_{\mathcal{D}}$ component, the arguments for the odd case apply without any change, and the result is the same Atiyah-Singer integral (19). To compute

$$r_{\mathcal{G}, M_1} \circ \text{ind}_{\mathcal{T}(\mathcal{G})}(\partial[\sigma]),$$

we observe that restriction of $\mathcal{T}(\mathcal{G})$ to $M_1 \times [0, 1]$ is just $\mathcal{T}(\mathcal{G}_1)$ which is the adiabatic groupoid of $\mathcal{G}_1 = G \times M_1 \times M_1$, hence we obtain the commutative diagram

$$(21) \quad \begin{array}{ccccc} C^*(A) < & \xrightarrow{r_{\mathcal{T}(\mathcal{G}), M \times \{0\}}} & C^*(\mathcal{T}(\mathcal{G})) & \xrightarrow{r_{\mathcal{T}(\mathcal{G}), M \times \{1\}}} & > C^*(\mathcal{G}) \\ & & & & \\ & \xrightarrow{r_{A, M_1}} & & \xrightarrow{r_{\mathcal{T}(\mathcal{G}), M_1 \times [0, 1]}} & \\ C^*(A|_{M_1}) < & \xrightarrow{r_{\mathcal{T}(\mathcal{G}_1), M_1 \times \{0\}}} & C^*(\mathcal{T}(\mathcal{G}_1)) & \xrightarrow{r_{\mathcal{T}(\mathcal{G}_1), M \times \{1\}}} & > C^*(\mathcal{G}_1), \end{array}$$

which implies

$$r_{\mathcal{G}, M_1} \circ \text{ind}_{\mathcal{T}(\mathcal{G})}(\partial[\sigma]) = \text{ind}_{\mathcal{T}(\mathcal{G}_1)}(\partial[\sigma]|_{M_1}).$$

The right hand side of the above can, in turn be computed by Lemma 4.1 with $\mathcal{G}_1 = M_1 \times M_1 \times G$ as the groupoid:

$$(22) \quad \text{ind}_{\mathcal{T}(\mathcal{G}_1)}(\partial[\sigma]|_{M_1}) = \int_{T^*M_1 \times \mathfrak{g}} \langle \hat{A}(T^*M_1 \oplus \mathfrak{g}) \wedge \text{ch}(\sigma[\Psi]|_{M_1}), \Omega'_{\pi^!TM_1} \rangle,$$

where $\Omega' \in \Gamma^\infty(\wedge^{\text{top}}\mathfrak{g} \otimes \wedge^{\text{top}}T^*M_1 \otimes \wedge^{\text{top}}TM_1)$ is a suitably normalized, invariant nowhere vanishing section defined in the same manner as Equation (13), and \mathfrak{g} is the Lie algebra of the isotropy group G .

To conclude, we have arrived at an index formula for elliptic pseudodifferential operators on \mathcal{G} .

Theorem 4.6. *Suppose that $\mathcal{G} \rightrightarrows M$ is a boundary groupoid of the form (16) with $\dim G$ even, and a deformation from the pair groupoid exists for \mathcal{G} . Let Ψ be an elliptic pseudo-differential operator on \mathcal{G} . One has the index formula*

$$\begin{aligned} \text{ind}(\Psi) &= \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^!TM} \rangle \\ &\quad \oplus \int_{T^*M_1 \times \mathfrak{g}} \langle \hat{A}(T^*M_1 \oplus \mathfrak{g}) \wedge \text{ch}(\sigma[\Psi]|_{M_1}), \Omega'_{\pi^!TM_1} \rangle \in \mathbb{Z} \oplus \mathbb{Z} \cong K_0(C^*(\mathcal{G})). \end{aligned}$$

4.3. The Fredholm index for fully elliptic operators. Lastly, recall the definition of fully elliptic operators in Definition 2.11. We suppose in this subsection that Ψ is a fully elliptic operator on $\mathcal{G} \rightrightarrows M$, where \mathcal{G} is of the form (16) .

Corollary 4.7. *The Fredholm index of Ψ is*

$$\text{ind}_F(\Psi) = \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^!TM} \rangle.$$

Proof. Recall that Ψ is invertible modulo $C^*(M_0 \times M_0) \cong \mathcal{K}$ and its Fredholm index lies in $K_0(C^*(M_0 \times M_0))$. By Theorems 1.1, 1.2 and [13] we have

$$\begin{aligned} \text{ind}_F(\Psi) &= \text{ind}_{\mathcal{D} \circ \varepsilon_{\mathcal{G}, M_0}}(\text{ind}_F(\Psi)) \\ &= \text{ind}_{\mathcal{D}}(\text{ind}_{\mathcal{T}(\mathcal{G})}(\sigma(\Psi))) \\ &= \int_{T^*M} \langle \hat{A}(T^*M) \wedge \text{ch}(\sigma[\Psi]), \Omega_{\pi^!TM} \rangle. \quad \square \end{aligned}$$

Remark 4.8. One can replace the pair groupoid in Definition 3.1 by other groupoids. Most of the arguments in Section 4 still works in a more general setting. It would be interesting to see which class of groupoids can arise from this kind of (non-trivial) deformations, and what index formulae one can obtain.

REFERENCES

- [1] I. Akrouf and P. Carrillo Rouse. Longitudinal b-operators, blups and index theorems. arXiv:1711.11197v3, 2018.
- [2] B. Ammann, R. Lauter, and V. Nistor. On the geometry of Riemannian manifolds with a Lie structure at infinity. *Int. J. Math. Math. Sci.*, (1-4):161–193, 2004.
- [3] B. Ammann, R. Lauter, and V. Nistor. Pseudodifferential operators on manifolds with a Lie structure at infinity. *Ann. of Math. (2)*, 165(3):717–747, 2007.
- [4] I. Androulidakis and G. Skandalis. The holonomy groupoid of a singular foliation. *J. Reine Angew. Math.*, 626:1–37, 2009.
- [5] I. Androulidakis and G. Skandalis. The analytic index of elliptic pseudodifferential operators on a singular foliation. *J. K-Theory*, 8(3):363–385, 2011.
- [6] I. Androulidakis and G. Skandalis. Pseudodifferential calculus on a singular foliation. *J. Non-commut. Geom.*, 5(1):125–152, 2011.

- [7] N. Berline, E. Getzler, and M. Vergne. *Heat kernels and Dirac operators*. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004. Corrected reprint of the 1992 original.
- [8] K. Bohlen. Groupoids and singular manifolds. arXiv:1601.04166, 2016.
- [9] K. Bohlen and E. Schrohe. Getzler rescaling via adiabatic deformation and a renormalized index formula. *J. Math. Pures Appl. (9)*, 120:220–252, 2018.
- [10] P. Carrillo Rouse and J.-M. Lescure. Geometric obstructions for Fredholm boundary conditions for manifolds with corners. *Ann. K-Theory*, 3(3):523–563, 2018.
- [11] P. Carrillo Rouse, J.-M. Lescure, and B. Monthubert. A cohomological formula for the Atiyah-Patodi-Singer index on manifolds with boundary. *J. Topol. Anal.*, 6(1):27–74, 2014.
- [12] P. Carrillo Rouse and B. Monthubert. An index theorem for manifolds with boundary. *C. R. Math. Acad. Sci. Paris*, 347(23-24):1393–1398, 2009.
- [13] P. Carrillo Rouse and B. K. So. K -theory and index theory for some boundary groupoids. *Results Math.*, 75(4):Paper No. 172, 20, 2020.
- [14] C. Carvalho, V. Nistor, and Y. Qiao. Fredholm criteria for pseudodifferential operators and induced representations of groupoid algebras. *Electron. Res. Announc. Math. Sci.*, 24:68–77, 2017.
- [15] C. Carvalho, V. Nistor, and Y. Qiao. Fredholm conditions on non-compact manifolds: theory and examples. In *Operator theory, operator algebras, and matrix theory*, volume 267 of *Oper. Theory Adv. Appl.*, pages 79–122. Birkhäuser/Springer, Cham, 2018.
- [16] R. Côme. The Fredholm property for groupoids is a local property. *Results Math.*, 74(4):Paper No. 160, 33, 2019.
- [17] A. Connes. Sur la théorie non commutative de l'intégration. In *Algèbres d'opérateurs (Sém., Les Plans-sur-Bex, 1978)*, volume 725 of *Lecture Notes in Math.*, pages 19–143. Springer, Berlin, 1979.
- [18] A. Connes. *Noncommutative geometry*. Academic Press, Inc., San Diego, CA, 1994.
- [19] C. Debord. Holonomy groupoids of singular foliations. *J. Differential Geom.*, 58(3):467–500, 2001.
- [20] C. Debord, J.-M. Lescure, and V. Nistor. Groupoids and an index theorem for conical pseudo-manifolds. *J. Reine Angew. Math.*, 628:1–35, 2009.
- [21] C. Debord, J.-M. Lescure, and F. Rochon. Pseudodifferential operators on manifolds with fibred corners. *Ann. Inst. Fourier (Grenoble)*, 65(4):1799–1880, 2015.
- [22] C. Debord and G. Skandalis. Adiabatic groupoid, crossed product by \mathbb{R}_+^* and pseudodifferential calculus. *Adv. Math.*, 257:66–91, 2014.
- [23] C. Debord and G. Skandalis. Pseudodifferential extensions and adiabatic deformation of smooth groupoid actions. *Bull. Sci. Math.*, 139(7):750–776, 2015.
- [24] C. Debord and G. Skandalis. Blow-up constructions for Lie groupoids and a Boutet de Monvel type calculus. *Münster J. Math.*, 14(1):1–40, 2021.
- [25] S. Evens, J.-H. Lu, and A. Weinstein. Transverse measures, the modular class and a cohomology pairing for Lie algebroids. *Quart. J. Math. Oxford Ser. (2)*, 50(200):417–436, 1999.
- [26] R. Lauter, B. Monthubert, and V. Nistor. Pseudodifferential analysis on continuous family groupoids. *Doc. Math.*, 5:625–655, 2000.
- [27] R. Lauter and V. Nistor. Analysis of geometric operators on open manifolds: a groupoid approach. In *Quantization of singular symplectic quotients*, volume 198 of *Progr. Math.*, pages 181–229. Birkhäuser, Basel, 2001.
- [28] P.-Y. Le Gall and B. Monthubert. K -theory of the indicial algebra of a manifold with corners. *K-Theory*, 23(2):105–113, 2001.
- [29] K. Mackenzie. *Lie groupoids and Lie algebroids in differential geometry*, volume 124 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1987.
- [30] K. Mackenzie. *General theory of Lie groupoids and Lie algebroids*, volume 213 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2005.
- [31] B. Monthubert. Pseudodifferential calculus on manifolds with corners and groupoids. *Proc. Amer. Math. Soc.*, 127(10):2871–2881, 1999.
- [32] B. Monthubert and V. Nistor. A topological index theorem for manifolds with corners. *Compos. Math.*, 148(2):640–668, 2012.
- [33] B. Monthubert and F. Pierrot. Indice analytique et groupoïdes de Lie. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(2):193–198, 1997.
- [34] S. Moroianu and V. Nistor. Index and homology of pseudodifferential operators on manifolds with boundary. In *Perspectives in operator algebras and mathematical physics*, volume 8 of *Theta Ser. Adv. Math.*, pages 123–148. Theta, Bucharest, 2008.
- [35] V. Nistor. Groupoids and the integration of Lie algebroids. *J. Math. Soc. Japan*, 52(4):847–868, 2000.
- [36] V. Nistor. Analysis on singular spaces: Lie manifolds and operator algebras. *J. Geom. Phys.*, 105:75–101, 2016.

- [37] V. Nistor, A. Weinstein, and P. Xu. Pseudodifferential operators on differential groupoids. *Pacific J. Math.*, 189(1):117–152, 1999.
- [38] M. Pflaum, H. Posthuma, and X. Tang. The index of geometric operators on Lie groupoids. *Indag. Math. (N.S.)*, 25(5):1135–1153, 2014.
- [39] M. Pflaum, H. Posthuma, and X. Tang. The transverse index theorem for proper cocompact actions of Lie groupoids. *J. Differential Geom.*, 99(3):443–472, 2015.
- [40] M. J. Pflaum, H. Posthuma, and X. Tang. The localized longitudinal index theorem for Lie groupoids and the van Est map. *Adv. Math.*, 270:223–262, 2015.
- [41] Y. Qiao and B. K. So. Renormalized index formulas for elliptic differential operators on boundary groupoids. To appear in the J. of Operator Theory. arXiv:2108.04592.
- [42] J. Renault. *A groupoid approach to C^* -algebras*, volume 793 of *Lecture Notes in Mathematics*. Springer, Berlin, 1980.
- [43] B. K. So. Pseudo-differential operators, heat calculus and index theory of groupoids satisfying the Lauter-Nistor condition. PhD thesis, The University of Warwick, 2010.
- [44] B. K. So. On the full calculus of pseudo-differential operators on boundary groupoids with polynomial growth. *Adv. Math.*, 237:1–32, 2013.
- [45] V. F. Zenobi. Adiabatic groupoid and secondary invariants in K-theory. *Adv. Math.*, 347:940–1001, 2019.

SCHOOL OF MATHEMATICS AND STATISTICS, SHAANXI NORMAL UNIVERSITY, XI'AN, 710119, SHAANXI, CHINA

Email address: yqiao@snnu.edu.cn

SCHOOL OF MATHEMATICS, JILIN UNIVERSITY, CHANGCHUN, 130023, JILIN, CHINA

Email address: bkso@graduate.hku.hk