KHOVANOV HOMOLOGY OF POSITIVE LINKS AND OF L-SPACE KNOTS

MARC KEGEL, NAAGESWARAN MANIKANDAN, LEO MOUSSEAU, AND MARITHANIA SILVERO

Abstract. We determine the structure of the Khovanov homology groups in homological grading 1 of positive links. More concretely, we show that the first Khovanov homology is supported in a single quantum grading determined by the Seifert genus of the link, where the group is free abelian and of rank determined by the Seifert graph of any of its positive link diagrams. In particular, for a positive link, the first Khovanov homology is vanishing if and only if the link is fibered. Moreover, we extend these results to (p, q) -cables of positive knots whenever $q \geq p$. We also show that several infinite families of Heegaard Floer L-space knots have vanishing first Khovanov homology. This suggests a possible extension of our results to L-space knots.

1. INTRODUCTION

In this article, we study the Khovanov homology of positive links and their cables. More precisely, we focus on those Khovanov homology groups in homological grading 1.

Theorem 1.1 If L is a positive link, then its Khovanov homology groups in homological grading 1 fulfill the following

$$
Kh^{1,j}(L) = \begin{cases} \mathbb{Z}^{p_1(L)} & \text{if } j = 2 - \chi(L), \\ 0 & \text{otherwise,} \end{cases}
$$

where $\chi(L)$ denotes the Euler characteristic^{[1](#page-0-0)} of a genus minimizing Seifert surface of L and $p_1(L)$ denotes the first cyclomatic number of the reduced Seifert graph of any positive diagram of L.

For the precise definition of $p_1(L)$ we refer to Section [2.](#page-4-0) In particular, it follows that $p_1(L)$ is an invariant of positive links. We deduce from Theorem [1.1](#page-0-1) that the first Khovanov homology detects fiberedness among positive links.

Theorem 1.2 A positive link L is fibered if and only if its first Khovanov homology vanishes, i.e. $Kh^{1,*}(L)=0$.

In general, Khovanov homology does not detect fiberedness. For example, K9a12 is fibered and $K11n83$ is not fibered, but both knots share the same Khovanov homology (after mirroring one of them) [\[LM23\]](#page-23-0). Conceptual similar results to Theorem [1.2](#page-0-2) were also obtained in [\[Sto05,](#page-23-1) [Buc22,](#page-22-0) [Buc23\]](#page-22-1) for the Jones polynomial.

Date: April 27, 2023.

²⁰²⁰ Mathematics Subject Classification. 57K10; 57K14, 57K16, 57K18, 57K32.

Key words and phrases. Khovanov homology, positive knots, fibered knots, L-space knots. ¹If L is a knot then $\chi(L) = 1 - 2g(L)$, where $g(L)$ denotes the Seifert genus of L.

However, our results are about the homological gradings in Khovanov homology while their results about the Jones polynomials can be interpreted as results on the quantum grading in Khovanov homology and thus are in some sense orthogonal to the results presented here. On the other hand, it was shown by Stošić [\[Sto10\]](#page-23-2) that braid positive links have trivial first Khovanov homology. Theorem [1.2](#page-0-2) generalizes Stošić's result since braid positive links are necessarily fibered and positive. Nevertheless, there exist fibered and positive links that are not braid positive. Among the prime knots with at most 12 crossings, there are exactly 16 such knots, the simplest one being the knot $K10n7$ [\[LM23\]](#page-23-0). We present in Proposition [4.4](#page-11-0) an infinite family of fibered, positive knots that are not braid positive.

Corollary 1.3 If K_0 is ribbon concordant to a positive knot K_1 , then $Kh^{1,j}(K_0)$ is trivial if $j \neq 2g(K_1) + 1$ and free abelian of rank at most $p_1(K_1)$ if $j = 2g(K_1) + 1$. If K_1 is, in addition, fibered then the first Khovanov homology of K_0 is vanishing.

Proof. This follows directly from a result of Levine–Zemke [\[LZ19\]](#page-23-3) saying that a ribbon concordance induces a grading preserving injective map on Khovanov ho- \Box mology.

1.1. Previously known positivity obstructions from Khovanov homology. Other obstructions of the Khovanov homology groups of positive links were known before. In [\[Kho03,](#page-23-4) [PS14\]](#page-23-5) it was shown that for a positive link L it holds

$$
Kh^{i,j}(L) = 0 \text{ if } i < 0 \text{ or } j < -\chi(L),
$$

\n
$$
Kh^{0,j}(L) = \begin{cases} \mathbb{Z} & \text{if } j = -\chi(L) \text{ and } 2 - \chi(L), \\ 0 & \text{otherwise}, \end{cases}
$$

\n
$$
Kh^{i,-\chi(L)}(L) = 0 \text{ if } i \neq 0,
$$

\n
$$
Kh^{i,2-\chi(L)}(L) = 0 \text{ if } i \neq 0 \text{ or } 1.
$$

We refer to Figure [1](#page-1-0) for a schematic picture of the Khovanov homology of a positive link. Some of these results can also be recovered from the main results of [\[PS20,](#page-23-6) [SS22\]](#page-23-7). In particular, we observe that Khovanov homology detects the genus among positive links. On the other hand, for every $k > 0$ there exists a fibered positive knot K such that $Kh^{i}(K)$ is non-trivial for every $i = 0, 2, 3, 4, ..., k$.

i \dot{i}	$i<0$			i>1
	O	$\overline{0}$	0	?
		$\mathbb Z$	$\mathbb{Z}^{p_1(L)}$	
$\frac{2-\chi(L)}{-\chi(L)}$		$\mathbb Z$		
		0		$\left($

FIGURE 1. The Khovanov homology of a positive link L.

1.2. Khovanov homology of cables. The skein long exact sequence for Khovanov homology together with Theorems [1.1](#page-0-1) and [1.2](#page-0-2) allow us to prove that the Khovanov homology of certain positive cables of positive knots looks like the Khovanov homology of a positive knot. Here we emphasize that such a cable (even if its cabling slopes are positive) is in general not a positive link. We refer to Remark [5.2](#page-13-0) for examples of this phenomenon.

Theorem 1.4 If K is a positive knot, then for every $q \ge p \ge 2$ the Khovanov homology of its (p, q) -cable $K_{p,q}$ fulfills

$$
Kh^{i,j}(K_{p,q}) = 0 \tif \t i < 0,
$$

\n
$$
Kh^{0,j}(K_{p,q}) = \begin{cases} \mathbb{Z} & \text{if } j = 1 - \chi(K_{p,q}) \pm 1, \\ 0 & \text{otherwise}, \end{cases}
$$

\n
$$
Kh^{1,j}(K_{p,q}) = \begin{cases} \mathbb{Z}^{p_1(K)} & \text{if } j = 2 - \chi(K_{p,q}), \\ 0 & \text{otherwise}. \end{cases}
$$

In the special case when K is fibered, the above theorem implies the following.

Corollary 1.5 If K is positive and $q \ge p \ge 2$, then $Kh^{1,*}(K_{p,q})$ is trivial if and only if K is fibered.

1.3. Khovanov homology of L-space knots. Another interesting class of knots is given by L-space knots. Here an L-space knot is a knot that admits a positive surgery to a Heegaard Floer L-space [\[OS05\]](#page-23-8). L-space knots are necessarily fibered [\[Ni07\]](#page-23-9) and strongly quasipositive [\[Hed10\]](#page-22-2). Moreover, many L-space knots are braid positive [\[ABG](#page-22-3)⁺21, [BK23\]](#page-22-4) and thus are fibered and positive. By Theorem [1.2](#page-0-2) these fibered positive L-space knots have Khovanov homology $Kh^{0,2g\pm 1} = \mathbb{Z}$ and all other Khovanov homology groups of homological degree at most 1 are trivial.

However, there exist L-space knots that are not braid positive [\[BK23\]](#page-22-4) and not positive (see Remark [5.2\)](#page-13-0). Currently, the only L-space knots that are not known to be braid positive are certain cables of L-space knots [\[ABG](#page-22-3)⁺21] and an infinite family of hyperbolic L-space knots from [\[BK23\]](#page-22-4). In Section [5](#page-12-0) we prove that large infinite subclasses of these L-space knots also have Khovanov homology $Kh^{0,2g\pm 1} = \mathbb{Z}$ and all other Khovanov homology groups of homological degree at most 1 are trivial. In particular, we show that all currently known hyperbolic L-space knots have such Khovanov homology groups. As a consequence of Theorem [1.4](#page-2-0) we get the following result.

Corollary 1.6 If K is an L-space knot which is a cable of a positive knot, then

 $Kh^{0,2g(K)\pm 1}(K)=\mathbb{Z}$ and $Kh^{i,j}(K)=0$ for all other groups with $i\leq 1$.

Proof. By [\[Hom11\]](#page-22-5) it is known that if K is a cable L-space knot then its cable parameters fulfill $q \ge p \ge 2$. Since L-space knots are fibered [\[Ni07\]](#page-23-9), Theorem [1.4](#page-2-0) implies the result implies the result.

In light of these results, we conjecture that L-space knots have Khovanov homology that looks like the Khovanov homology of a positive knot.

Conjecture 1.7 If K is an L-space knot, then

 $Kh^{0,2g(K)\pm 1}(K)=\mathbb{Z}$ and $Kh^{i,j}(K)=0$ for all other groups with $i\leq 1$.

We remark that the result of Levine–Zemke [\[LZ19\]](#page-23-3) gives a potential way to study this conjecture. If we can show that any L -space knot K is ribbon concordant to a positive, fibered (or a braid positive) knot, then $Kh^{0,2g(K)\pm 1}(K)$ is 0 or Z and all other Khovanov homology groups in homological grading at most 1 vanish. On the other hand, we can search for a counterexample by finding a ribbon concordance from a knot K_0 with Khovanov homology that is not of the form of a positive knot to an L-space knot K_1 .

When not explicitly stated we always use Khovanov homology over the integers. However, all our results for Khovanov homology hold also true (by exactly the same proofs) with all possible coefficient groups.

Remark 1.8 Our main results extend to other homology theories.

- (1) Since odd Khovanov homology agrees with the usual Khovanov homology over \mathbb{Z}_2 -coefficients [\[ORS13\]](#page-23-10), we can use our main results and the universal coefficient theorem to deduce that Theorems [1.1](#page-0-1) and [1.2](#page-0-2) hold also true for odd Khovanov homology groups with rational coefficients. In fact, experimental data suggest that they hold true for any coefficient group. The odd Khovanov homology groups fit into the same long exact sequence as the usual Khovanov homology groups [\[ORS13\]](#page-23-10) and thus Theorem [1.4](#page-2-0) and Corollary [1.6](#page-2-1) hold true (by the same proofs) for odd Khovanov homology with rational coefficients as well.
- (2) If L is a fibered, positive link, then all the Khovanov–Rozansky $\mathfrak{sl}(n)$ ho-mology groups [\[KR08\]](#page-23-11) in homological grading 1 vanish, i.e. $KhR_n^{1,j}(L) = 0$ for all $j \in \mathbb{Z}$ and all $n \geq 2$. For that, we observe that the proof of Theorem [1.2](#page-0-2) (and possibly also that of Theorem [1.1\)](#page-0-1) can be adapted to work for Khovanov–Rozansky homologies as outlined in [\[Sto10\]](#page-23-2). By using [\[Kan22\]](#page-23-12) the statement from Corollary [1.3](#page-1-1) also generalizes to Khovanov–Rozansky homology.

1.4. Experimental data. Theorems [1.1](#page-0-1) and [1.2](#page-0-2) can be seen as obstructions for a knot to be positive. From the behavior of Khovanov homology under mirroring a knot, we see that this also gives an obstruction for a knot being negative. This obstruction seems to be strong, since for low-crossing knots, Khovanov homology detects positivity: Among the 2977 prime knots with at most 12 crossings, exactly 246 are positive or negative [\[LM23\]](#page-23-0). Theorems [1.1](#page-0-1) and [1.2](#page-0-2) together with the obstructions from Section [1.1](#page-1-2) obstruct all the other 2731 knots from being positive or negative [\[LM23\]](#page-23-0).

From this perspective, the more interesting class of knots is given by the census knots [\[Dun20b\]](#page-22-6), i.e. the hyperbolic knots whose complements can be triangulated by at most 9 ideal tetrahedra. Thus these knots have simple complements. However, the simplest known diagrams of some census knots have almost 300 crossings. In $[BBD^+]$ $[BBD^+]$ the Khovanov homology (with rational coefficients) for exactly 672 of the 1267 census knots was computed. Among those 672 knots, we found positive or negative diagrams for 334 [\[BBD](#page-22-7)⁺]. To obstruct positivity (or negativity) for the remaining 338 knots we applied the obstruction from Khovanov homology. Again this obstruction turns out to be strong, although not perfect. We can obstruct all but 9 of the remaining knots from being positive or negative. We refer to Figure [2](#page-4-1) for an example. The data mentioned above can be accessed at [\[KMMS\]](#page-23-13).

FIGURE 2. A diagram of the mirror of the census knot $o9_34097$ and its Khovanov homology table. Since it is not fibered, Theorem [1.2](#page-0-2) implies that it is not positive.

Acknowledgment. This project was initiated during a discussion between M.K. and M.S. at the semester program "Braids" (Feb 1- May 6, 2022) at the Institute for Computational and Experimental Research in Mathematics (ICERM). We thank ICERM for the invitation and financial support. We also thank Lukas Lewark for explaining to us why the $(2, 1)$ -cable of the trefoil is a strongly quasipositive, fibered knot with non-vanishing first Khovanov homology, which was the initiation for writing Section [5.](#page-12-0) The authors wish to thank Chris Wendl, for his hospitality when M.S. visited the HU Berlin and to the group P20-01109 at the University of Seville, where part of this research was conducted.

N.M. is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project ID: 390685689).

M.S. is partially supported by Spanish Research Project PID2020-117971GB-C21, by IJC2019-040519-I, funded by MCIN/AEI/10.13039/501100011033 and by P20-01109 (JUNTA/FEDER).

2. Preliminaries

Let D be a link diagram and let $cr(D) = \{c_1, \ldots, c_n\}$ be the set of its crossings. A state s assigns a marker 0 or 1 to each crossing of D, that is $s: cr(D) \rightarrow \{0,1\}.$ Let $S(D)$ be the collection of 2^n possible states of D. We write s_0 for the state assigning a 0-marker to every crossing.

Given $s \in \mathcal{S}(D)$, the resolution sD corresponds to the diagram obtained after smoothing every crossing $c_i \in cr(D)$ according to its marker $s(c_i)$ following Figure [3\(](#page-5-0)a).

The resolution sD consists of a collection of $|sD|$ circles embedded in the plane together with some 0- and 1-chords (segments connecting two circles in the place where there was a crossing). See Figure [4\(](#page-5-1)b).

Definition 2.1 Given a state $s \in S(D)$, the *state graph* $G_s(D)$ is a planar graph obtained by collapsing each circle of sD to a vertex so that each chord in sD becomes an edge in $G_s(D)$. The circles and chords of sD are in bijection with the vertices and edges of $G_s(D)$. See Figure [4\(](#page-5-1)c).

FIGURE 3. The smoothing of a crossing according to its 0 or 1 marker and the sign (positive or negative) of a crossing are shown in (a) and (b), respectively.

FIGURE 4. A positive diagram D, its resolution s_0D , and the associated state graph $G_0(D)$ are shown in (a), (b), and (c), respectively.

The graph $G_s(D)$ may contain loops and multi-edges (i.e. edges connecting the same pair of vertices, also called parallel edges). Given a graph G, its associated *reduced graph* G^{red} is the graph obtained from G by identifying parallel edges. We write $G_0(D)$ to denote the state graph associated with the state s_0 .

Recall that every crossing in an oriented diagram can be classified into positive or negative, according to the convention shown in Figure $3(b)$. An oriented link is positive if it admits a positive diagram, i.e. a diagram with no negative crossings. Observe that the 0-smoothing of a positive crossing agrees with its Seifert smoothing (that is, the only smoothing preserving orientation), and therefore the Seifert graph of a positive diagram D coincides with $G_0(D)$.

Definition 2.2 The cyclomatic number of a graph G is the minimal number of edges that one must delete to transform G into a forest. It can be computed as $p(G) = e - v + k$, where v, e and k denote the number of vertices, edges and connected components of G , respectively. Given a positive link L , we define its cyclomatic number $p_1(L)$ as the cyclomatic number of the reduced Seifert graph associated with any positive diagram representing L. This number is independent of the positive diagram (compare to Theorem [1.1](#page-0-1) and [\[PS20\]](#page-23-6)).

Theorem 2.3 (Stoimenow [\[Sto05\]](#page-23-1), cf. [\[Fut13,](#page-22-8) [Buc22\]](#page-22-0)) A positive diagram D represents a fibered link if and only if its reduced Seifert graph $G_0(D)^{red}$ is a tree.

As an example, we refer to Figure [4](#page-5-1) where the reduced Seifert graph $G_0(D)^{red}$ is a tree with 3 leaves and thus D represents a fibered link.

3. Khovanov homology

In this section, we briefly review the definition of Khovanov homology, following [\[BN02\]](#page-22-9). Given a diagram D, we first define a complex $\llbracket D \rrbracket$ which, after some grading shiftings, becomes the Khovanov complex \mathcal{C}_D .

Let D be an oriented link diagram with n ordered crossings c_1, \ldots, c_n . This induces an identification of the set $\mathcal{S}(D)$ of states with the elements in $\{0,1\}^n$ defined by $s = (s_1, ..., s_n) \in \{0, 1\}^n$, with $s(c_j) = s_j$. Write $|s| = \sum_{j=1}^n s_j$.

Consider the unit cube of dimension n , where each vertex consists of a state of D and there is an edge from the state s to the state s' iff they are identical except in their k^{th} coordinate, where $s_k = 0$ and $s'_k = 1$. Let V be the graded vector space with two basis elements, v_+ and v_- of degrees 1 and -1 , respectively. We associate to each vertex of the cube $s \in 2^n$ $s \in 2^n$ $s \in 2^n$ the graded vector space² $V_s(D) = V^{\otimes |sD|} \{|s|\}.$ Define a complex $[[D]]$ whose r^{th} chain group is defined as $[[D]]^r = \bigoplus_{|s|=r} V_s(D)$.

We define the differential $d^r: [D] \to [D]^{r+1}$. Let ξ be an edge of the cube oriented from a state s to a state s' satisfying $s(j) = s'(j)$ for all $j \neq k$, $s(k) = 0$ and $s'(k) = 1$, for $1 \leq j, k \leq n$. Define the height $|\xi|$ of the edge ξ as $|\xi| = |s|$ and its sign as $sgn(\xi) = \sum_{j \leq k} s(j)$, with k as before.

Resolutions sD and $s'D$ are identical except in a neighborhood of the k^{th} crossing, where either one circle in sD splits into two circles in $s'D$ or two circles in sD merge into one in $s'D$; we call the associated transformations splitting and merging, respectively. The map $d_{\xi}: V_s(D) \longrightarrow V_{s'}(D)$ is defined as the identity on the tensor factors corresponding to the circles that are preserved and as either a multiplication $m: V \otimes V \to V$ or a comultiplication $\Delta: V \to V \otimes V$ on the factors corresponding to the circles involved in a merging or a splitting, respectively. These linear maps are defined by:

> $m(v_+ \otimes v_+) = v_+$ $\Delta(v_+) = v_+ \otimes v_- + v_- \otimes v_+$ $m(v_+ \otimes v_-) = v_ \Delta(v_-) = v_- \otimes v_$ $m(v_-\otimes v_+) = v_$ $m(v_-\otimes v_-)=0$

The differential of the complex is defined as $d^r = \sum_{|\xi|=r} (-1)^{\text{sgn}(\xi)} d_{\xi}$. With the chain groups and differential defined above, $\llbracket D \rrbracket$ is indeed a chain complex. We write $H^r(D)$ for its r^{th} homology group.

The *height shift* operation [p] on a chain complex $A: \ldots \to A^r \stackrel{d^r}{\to} A^{r+1} \to \infty$ defined as $\mathcal{A}[p] = \mathcal{B}$, where $B^r = A^{r-p}$, with all differentials shifted accordingly.

Definition 3.1 Given a link diagram D with n_+ positive and n_- negative crossings, the Khovanov complex \mathcal{C}_D is defined as $[[D][-n_-]\{n_+ - 2n_-\}$. We denote its i^{th} homology group by $Kh^{i}(D)$.

Notice that each element in $[[D]]$ has two gradings: r coming from the chain group and a second grading q coming from the internal degree as an element in V. If we denote $\llbracket D \rrbracket^{r,q}$ the subset of elements of $\llbracket D \rrbracket^{r}$ with second grading q, then $\llbracket D \rrbracket^r = \bigoplus_{q \in \mathbb{Z}} \llbracket D \rrbracket^{r,q}$ and $Kh^{i,j}(D) = H^{r,q}(D)$, with $r = i + n_-$ and $q =$ $j - n_+ + 2n_$. We call such i and j homological and quantum gradings, respectively.

²Given a graded vector space $W = \bigoplus_m W_m$, the graded dimension of W is defined as the power series qdim $W = \sum_m q^m \dim W_m$. We denote by $\{l\}$ the *degree shift* operation on W by setting $W\{l\}_m = W_{m-l}$, so qdim $W\{l\} = q^l$ qdim W.

Theorem 3.2 [\[Kho00,](#page-23-14) [BN02\]](#page-22-9) For every $i, j \in \mathbb{Z}$, the isomorphism types of the groups $Kh^{i,j}(D)$ are link invariants. These groups are called Khovanov homology groups of the link.

Later, we will need the behavior of Khovanov homology under the addition of a split unknot component, which we record here.

Lemma 3.3 Let $L \sqcup \bigcirc$ denote the split union of a link L with an unknot. Then for every $i \in \mathbb{Z}$

 $Kh^{i}(L \sqcup \bigcirc) = Kh^{i}(L) \oplus Kh^{i}(L).$

In particular, $Kh^{i}(L) = 0$ implies that $Kh^{i}(L \sqcup \bigcirc) = 0$.

Proof. The result follows from the Künneth formula in Khovanov homology [\[Kho00\]](#page-23-14) together with the fact that $Kh^{i}(\bigcirc)$ equals \mathbb{Z}^{2} if $i = 0$, and is vanishing in all other gradings. \square

4. Khovanov homology of positive links

This section is devoted to proving Theorems [1.1](#page-0-1) and [1.2.](#page-0-2) We also show that the latter is a proper generalization of the result in [\[Sto10,](#page-23-2) Theorem 3.1] by presenting an infinite family of fibered, positive knots which are not closures of positive braids. In this section, we assume all positive links to be non-split without loss of generality (otherwise, just consider each split component independently and apply [\[Kho00,](#page-23-14) Proposition 33]). We will deduce Theorems [1.1](#page-0-1) and [1.2](#page-0-2) from the following more technical statement.

Theorem 4.1 Let D be a reduced positive diagram of a link L and let D' be another positive diagram of a link L' such that $G_0(D') = G_0(D)^{red}$. Then $Kh^1(L) =$ $Kh^1(L').$

We show in Lemma [4.2](#page-9-0) below that such a diagram D' always exists. However, neither D' nor L' are unique, since the Seifert circles and the chords between them might be nested and linked. But Theorem [4.1](#page-7-0) tells us that the first Khovanov homology does not depend on that extra information; however the exact quantum grading does, cf. Theorem [1.1.](#page-0-1) For a concrete example we refer to Figure [5.](#page-9-1)

Proof of Theorem [4.1.](#page-7-0) We will analyze the first homology group $H^1(D)$ of the complex $[\![D]\!]$. For that, it will be enough to consider d^1 : $[\![D]\!]^1 \to [\![D]\!]^2$. First, we describe these chain groups and d^1 in more detail. Since our statement is independent of the quantum grading we can, for ease of notation, omit the degree shift operation on the chain groups.

Consider $G = G_0(D)$ and write $E = \{e_1, \ldots, e_n\}$ for its edges, where e_j is associated with the crossing c_j , and $l = |s_0D|$ for its number of vertices. The states contributing to $llbracket D \rrbracket^1$ are those assigning a 1-marker to a single crossing, and therefore

$$
\llbracket D \rrbracket^1 = \bigoplus_{i=1}^n A_i
$$

where the summand A_i is associated to the state s^i assigning a 0-marker to every crossing but c_i . Since D is positive and therefore 0-adequate, we get that A_i is isomorphic to $V^{\otimes (l-1)}$, for every $1 \leq i \leq n$.

Generators in $\llbracket D \rrbracket^2$ are those states assigning exactly two 1-markers to the crossings of D , and therefore

$$
\llbracket D \rrbracket^2 = \bigoplus_{\substack{i,j=1 \\ i
$$

where the summand A_{ij} corresponds to the state s^{ij} assigning a 1-marker exactly to the crossings c_i and c_j . If e_i and e_j are parallel (i.e., they connect the same pair of vertices) in G then A_{ij} is isomorphic to $V^{\otimes l}$, for every $1 \leq i < j \leq n$.

The differential d^1 : $\llbracket D \rrbracket^1 \to \llbracket D \rrbracket^2$ is given as direct sum of maps δ_{kij} : $A_k \to A_{ij}$ corresponding to the edges of the cube of resolutions. Whenever $k \neq i, j$, it turns out that δ_{kij} is the zero map (equivalently, there is no edge connecting the states s^k and s^{ij} in the cube). Given two parallel edges e_i and e_j in G, with $i < j$, consider the following diagram.

Here π_i , π_j , and π_{ij} denote the projections to the summands A_i , A_j , and A_{ij} . Observe that the smoothed diagrams s^iD , s^jD and $s^{ij}D$ are identical in $(l-2)$ Seifert circles. Order the circles in these smoothed diagrams such that the p^{th} circle coincides in the three of them, for $1 \le p \le l-2$. These orderings induce preferred isomorphisms denoted by ϕ_i , ϕ_j , and ϕ_{ij} . For $k = i, j$ we define

$$
\overline{\delta}_{kij} := \phi_{ij} \circ \delta_{kij} \circ \phi_k^{-1}.
$$

By construction, the diagram above commutes. The map δ_{iij} (resp. δ_{jij}) is given by the identity map on all factors except the $(l-1)^{th}$ one, corresponding to the last circle in the chosen order, where it acts as the comultiplication map $-\Delta$ (resp. Δ). Recall that the sign is determined by the condition $i < j$. Thus we conclude

$$
\delta_{iij}: v_1 \otimes \cdots \otimes v_{l-1} \longmapsto -v_1 \otimes \cdots \otimes v_{l-2} \otimes \Delta(v_{l-1}),
$$

$$
\overline{\delta}_{jij}: v_1 \otimes \cdots \otimes v_{l-1} \longmapsto v_1 \otimes \cdots \otimes v_{l-2} \otimes \Delta(v_{l-1}).
$$

Now let x be an element in the kernel of d^1 . Since for $k \neq i, j$ the maps δ_{kij} are zero and Δ is injective it follows from the commutativity of the diagram that

$$
\phi_i \circ \pi_i(x) = \phi_j \circ \pi_j(x).
$$

Thus we have shown that the kernel of d^1 is contained in the subgroup given by elements $(x_1, \ldots, x_n) \in [D]$ ¹ such that $\phi_r(x_r) = \phi_s(x_s)$ if e_r is parallel to e_s . Since the ϕ_* are isomorphisms, identifying coordinates associated with parallel edges preserves the first homology groups. Thus we get that $H^1(D) = H^1(D')$, where D' is a diagram satisfying $G_0(D') = G^{red}$.

In the following two lemmata, we show that D' from Theorem [4.1](#page-7-0) can be chosen so that it satisfies some extra conditions.

Lemma 4.2 The positive diagram D' of L' in Theorem [4.1](#page-7-0) can be chosen to be alternating and such that its signature^{[3](#page-9-2)} is $\sigma(L') = p_1(L)$.

Proof. Since $G_0(D)$ is the Seifert graph of D, it is planar and bipartite (and so is $G_0(D)^{red}$). We first construct a positive diagram D' such that $G_0(D') = G_0(D)^{red}$. To do so, just choose a planar embedding of $G_0(D)^{red}$ and replace every vertex with a circle and every edge connecting two vertices by a positive crossing connecting the two corresponding circles. By construction, the obtained diagram D' is positive and its Seifert graph equals $G_0(D)^{red}$. Since $G(D')$ is bipartite a simple combinatorial argument shows that D' is alternating. We refer to Figure [5](#page-9-1) for an example.

In order to compute the signature of L' we assume without loss of generality that D' is reduced. We use the result in [\[Tra04\]](#page-23-15) stating that any link L_0 obtained from a link L by performing a Seifert smoothing of a positive crossing in a reduced alternating diagram of L satisfies $\sigma(L) = \sigma(L_0) + 1$. Since D' is alternating, performing a Seifert smoothing in any subset of its crossings produces an alternating diagram. Moreover, the effect of such a smoothing of a crossing corresponds to deleting the associated edge in the Seifert graph. Hence, there exists a sequence of $p_1(L)$ Seifert smoothings transforming D' into a diagram whose Seifert graph is a tree and thus represents the unknot. Since the signature of the unknot is trivial, we conclude that $\sigma(L)$ $) = p_1(L).$

FIGURE 5. A positive diagram of knot $K12n110$, its Seifert smoothing, and its reduced Seifert graph with $p_1 = 1$ are shown in (a) , (b) , and (c) , respectively. The diagram D' from Lemma [4.2](#page-9-0) has the same reduced Seifert graph. In (d) we show D' after removing nugatory crossings, from which we see that it represents the mirror of $L4a1\{0\}$. We compute that $Kh^1(K12n110) = \mathbb{Z}$ is supported in quantum grading 7 and $Kh^{1}(m(L4a1\{0\})) = \mathbb{Z}$ is supported in quantum grading 2.

³Here we use the convention that the signature of a positive link is positive.

Lemma 4.3 If D' is a diagram constructed as in the proof of Lemma [4.2](#page-9-0) representing a link L' , then

$$
Kh^{1,p_1(L)+3}(L') = 0.
$$

Proof. Without loss of generality, we can assume D' to be reduced. We will prove the result by induction on $p_1(L) = p_1(D')$. If $p_1(D') = 0$, the diagram D' represents the unknot with vanishing first Khovanov homology. If $p_1(D') > 0$, we consider one of its crossings v and consider its 0-smoothing D_0 and its 1-smoothing D_1 . The long exact sequence in Khovanov homology [\[Vir04\]](#page-23-16), cf. Lemma 2.2 in [\[MPS](#page-23-17)+18], yields an exact sequence of the form

$$
Kh^{\frac{w_1-w-1}{2}+1,\frac{3(w_1-w)-1}{2}+p_1(D')+3}(D_1)\to Kh^{1,p_1(D')+3}(D')\to Kh^{1,p_1(D_0)+3}(D_0),
$$

where w and w_1 denote the writhes of D' and D_1 respectively.

To show the vanishing of the middle group it suffices to demonstrate that the groups on the left and right vanish. For that, we observe that D_0 is again a positive and alternating diagram with $p_1(D_0) = p_1(D') - 1$ and such that its reduced Seifert graph agrees with its Seifert graph. Thus by induction hypothesis $Kh^{1,p_1(D_0)+3}(D_0)=0.$

 D_1 is in general not positive but it is always alternating and in particular it is adequate. The graph $G_0(D_1)$ is obtained from the Seifert graph of D' by deleting one edge and identifying the corresponding vertices. Thus $G_0(D_1)$ is not bipartite. Then we can apply Remark 6.4 from [\[PS20\]](#page-23-6) to deduce that

$$
Kh^{\frac{w_1-w-1}{2}+1,\frac{3(w_1-w)-1}{2}+p_1(D')+3}(D_1)=0.
$$

We are now ready to prove Theorems [1.1](#page-0-1) and [1.2.](#page-0-2)

Proof of Theorem [1.1.](#page-0-1) From Lemma [4.2](#page-9-0) and Theorem [4.1](#page-7-0) we conclude that there exists a reduced alternating and positive diagram D' of a link L' with signature $\sigma(L') = p_1(L)$ such that $Kh^1(L) = Kh^1(L')$. Since L' is alternating, its Khovanov homology is thin and based around the signature [\[Lee05\]](#page-23-18), i.e.

$$
Kh^{1,j}(L') = 0 \text{ if } j \neq p_1(L) + 1, p_1(L) + 3.
$$

From Lemma [4.3](#page-10-0) we also know that $Kh^{1,p_1(L)+3}(L')$ is trivial and thus the only potential non-trivial Khovanov homology group of L' in homological degree 1 has quantum grading $p_1(L) + 1$. To compute this group, we observe that the minimal quantum grading from [\[PS20\]](#page-23-6) is $e' - v' = p_1(L) - 1$, where e' and v' denote the number of edges and vertices in the Seifert graph $G_0(D')$. Since any Seifert graph is bipartite, it follows from [\[PS20,](#page-23-6) Remark 6.4] that $Kh^{1,p_1(L)+1}(L')$ is isomorphic to $\mathbb{Z}^{p_1(L)}$. In conclusion, we have computed that

$$
Kh^{1,j}(L') = \begin{cases} \mathbb{Z}^{p_1(L)} & \text{if } j = p_1(L) + 1, \\ 0 & \text{otherwise.} \end{cases}
$$

From Theorem [4.1](#page-7-0) we conclude directly that $Kh^1(L) = \mathbb{Z}^{p_1(L)}$. To get the statement in Theorem [1.1](#page-0-1) about the quantum grading we apply again [\[PS20,](#page-23-6) Remark 6.4] to the positive diagram D to deduce that $Kh^{1,2-\chi(L)}(L) = \mathbb{Z}^{p_1(L)}$ and thus all other Khovanov homology groups in homological degree 1 are trivial. \Box

Proof of Theorem [1.2.](#page-0-2) If L is fibered, then Theorem [2.3](#page-5-2) implies that diagram D' from Theorem [4.1](#page-7-0) can be chosen to be a diagram of the unknot, and therefore $Kh^{1}(L) = Kh^{1}(D')$ is trivial.

Conversely, if $Kh^1(L)$ is non-trivial, then Theorem [1.1](#page-0-1) implies that $p_1(L)$ is not vanishing and therefore from Theorem [2.3](#page-5-2) we get that L is not fibered. \square

We conclude this section by presenting an infinite family of fibered, positive knots that cannot be written as closures of positive braids.

Proposition 4.4 Let L_n be the oriented link represented by the diagram D_n shown in Figure [6,](#page-11-1) for $n \in \mathbb{N}$. Then L_n is fibered and positive, but cannot be realized as the closure of a positive braid, for $n \geq 1$.

FIGURE 6. The diagram D_n , where the box labeled n denotes n positive half-twists, and its Seifert graph $G_n = G_0(D_n)$. L_2 and L_4 correspond to the knots $K10n7$ and $K12n105$, respectively. L_0 is the torus knot $T(2, 5)$, which is braid positive.

Proof. The diagram D_n has reduced Seifert graph a tree and therefore L_n is fibered and positive (Theorem [2.3\)](#page-5-2). In order to show that L_n is not braid positive, we show that Ito's normalized HOMFLYPT polynomial H_{L_n} of L_n contains negative coefficients. The statement follows from the result that H_L is non-negative if L is the closure of a positive braid [\[Ito22,](#page-22-10) Theorem 1.1].

First, observe that L_n is a knot if n is even; otherwise, it is a non-split link of two components (its linking number equals $(n+3)/2$). Applying Ito's skein relation [\[Ito22,](#page-22-10) Equation (1)] to one of the *n* crossings in the box leads to the following:

$$
\widehat{H}_{L_n}(\alpha, z) = \begin{cases} \widehat{H}_{L_{n-2}}(\alpha, z) + \widehat{H}_{L_{n-1}}(\alpha, z), & \text{if } n \text{ is even,} \\ \widehat{H}_{L_{n-2}}(\alpha, z) + z^2 \widehat{H}_{L_{n-1}}(\alpha, z), & \text{if } n \text{ is odd.} \end{cases}
$$

Set $\widehat{H}_{L_n}(\alpha, z) = \sum_{i \in \mathbb{Z}} P_{L_n}^{(i)}$ $L_n^{(i)}(\alpha) \cdot z^i$. We will prove that $P_{L_n}^{(0)}$ $L_n^{(0)}$, which we denote by P_n , is not positive. The surface S_n obtained by applying the Seifert algorithm to D_n realizes the 3-genus of L_n and $\chi(S_n) = \chi(G_n) = -n-3$, and therefore we get (via the HOMFLYPT function at SageMath and Ito's normalization) for the first links in the family:

$$
P_1 = 1 - 2\alpha^2 - \alpha^3,
$$

\n
$$
P_2 = 4 + 2\alpha - 2\alpha^2 - \alpha^3.
$$

These computations can be accessed at [\[KMMS\]](#page-23-13). By induction, the above skein relation leads to

$$
P_n = \begin{cases} (3+k) + 2\alpha - 2k\alpha^2 - k\alpha^3, & \text{if } n = 2k, \\ 1 - 2\alpha^2 - \alpha^3, & \text{if } n \text{ is odd,} \end{cases}
$$

which are non-positive polynomials for any $n > 0$.

5. Khovanov homology of L-space knots and cables

In this section, we study the Khovanov homology of fibered strongly quasipositive knots, of L-space knots, and of cable links. In particular, we show that Theorems [1.1](#page-0-1) and [1.2](#page-0-2) do not extend to fibered strongly quasipositive knots. On the other hand, we prove that Theorem [1.2](#page-0-2) extends to many infinite families of L-space knots, giving support for an affirmative answer to Conjecture [1.7.](#page-2-2)

Thurston classified knots into three disjoint classes: torus knots, satellite knots, and hyperbolic knots [\[Thu86\]](#page-23-19). Torus knots are fibered, and the (strongly) quasipositive torus knots are exactly the positive torus knots $T(p,q)$, $p/q > 0$ [\[Hed10\]](#page-22-2), which are also known to be braid positive L-space knots [\[Mos71\]](#page-23-20). So for torus knots, these four positivity notions agree.

Satellite knots are those whose complement contains an incompressible, nonboundary parallel torus. Among them, a natural class is given by cable knots; we shift our attention to them in Section [5.1.](#page-12-1) In Section [5.2](#page-20-0) we focus on hyperbolic knots.

5.1. Cable knots. Given a knot K, its (p, q) -cable $K_{p,q}$ is the satellite knot with pattern the torus knot $T(p,q)$ and companion K. Given a diagram D of K, the standard diagram $D_{p,q}$ of $K_{p,q}$ is obtained by taking p parallel blackboard copies of D and adding $n = pw - q$ negative $(1/p)$ -twists to the p-parallel strands, where w is the writhe of D (see Figure [7](#page-12-2) for an example). We emphasize that the standard diagram $D_{p,q}$ depends on the diagram D of K we start with, while the isotopy class of $K_{p,q}$ depends only on the isotopy class of K. Since $K_{p,q} = K_{-p,-q}$ we can assume in the following that $p \geq 2$.

FIGURE 7. Diagram D of the positive trefoil knot is shown in (a). Standard diagrams of its cables $D_{3,9}$ and $D_{3,5}$ are shown in (b) and (c), respectively.

It is well understood which cable knots are fibered, braid positive, positive, strongly quasipositive, or L-space knots, provided one knows if K has these properties.

Theorem 5.1 Let K be a knot.

- (1) $K_{p,q}$ is fibered if K is fibered.
- (2) Let K be fibered and strongly quasipositive. Then $K_{p,q}$ is strongly quasipositive if and only if $q > 0$.
- (3) $K_{p,q}$ is an L-space knot if and only if K is an L-space knot and $q \geq$ $p(2g(K) - 1)$, where $g(K)$ denotes the 3-genus of K.
- (4) Let K be a positive (braid positive) knot. Then $K_{p,q}$ is positive (braid positive) if $q \geq p w(K)$, where $w(K)$ denotes the minimal writhe of a positive (braid positive) diagram of K.

Proof. (1) is a well-known statement. An explanation together with beautiful visualizations can be found at [\[Bak\]](#page-22-11). (2) follows from Proposition 2.4 and Theorem 2.11 at [\[Hed08\]](#page-22-12). (3) is proved by Hedden [\[Hed09\]](#page-22-13) and Hom [\[Hom11\]](#page-22-5). (4) is again a well-known statement. Let D be a positive (braid positive) diagram of K with writhe $w(D)$. If $q \geq p w(D)$, then any crossing in $D_{p,q}$ is positive.

Remark 5.2 Part (4) of Theorem [5.1](#page-13-1) does not give any statements about the positivity (braid positivity) of the cables of K for $q < p w(K)$. In fact, it turns out that for a given positive (braid positive) knot K, the cables of K with $0 \leq q <$ $p w(K)$ are often not positive (braid positive). This can be checked by computing standard obstructions for positivity and braid positivity.

In [\[KMMS\]](#page-23-13) we show that all $(2, q)$ -cables of $K = T(2, 3), T(2, 5), T(3, 4), T(3, 5),$ $T(4, 3)$, and $T(4, 5)$ with $0 \le q < 2w(K)$ have non-positive normalized HOM-FLYPT polynomial and thus are not braid positive by [\[Ito22\]](#page-22-10). In general, it is shown in [\[HK\]](#page-22-14) that $K_{p,q}$ is not a braid positive knot if $q < p(2g(K) - 1)$.

Sometimes it is also possible to obstruct positivity. If D is a reduced positive diagram with c crossings of a fibered knot K then it has minimal Seifert genus and $2q(K) - 1 = c - |s_0D|$. Since D is and its reduced Seifert graph is a tree $c \geq 2(|s_0D|-1)$, and thus $4g(K) \geq c$. Using this obstruction we see for example that the $(2, 3)$ -cable of $T(2, 3)$ is an L-space knot that is not positive (since it has Seifert genus 3 and crossing number 15 [\[Dun20a,](#page-22-15) Table 8]).

Theorem [5.1](#page-13-1) allows us to present examples of cable knots that are fibered and strongly quasipositive whose first Khovanov homology is not vanishing. The simplest such example we know is the $(2, 1)$ -cable of the $(2, 3)$ -torus knot which was communicated to us by Lukas Lewark.

Theorem 5.3 For any $1 \le g \le 200$

$$
Kh^{1,j}(T(2,2g+1)_{2,1}) = \begin{cases} \mathbb{Z} & \text{if } j = 3; \\ 0 & \text{otherwise.} \end{cases}
$$

In particular, for $1 \leq g \leq 200$ the knot $T(2, 2g + 1)_{2,1}$ is strongly quasipositive, fibered and its Khovanov homology in homological degree 1 does not vanish.

Proof. It follows from Theorem [5.1](#page-13-1) (1) and (2) that $T(2, 2g + 1)_{2,q}$ is fibered and strongly quasipositive for any $g, q > 0$. We use KnotJob [\[Sch\]](#page-23-21) and the KnotTheory package [\[BNMea\]](#page-22-16) for the computations of the Khovanov homologies. The computations can be accessed here [\[KMMS\]](#page-23-13).

The above computational results strongly suggest that Theorem [5.3](#page-13-2) holds actually true for all $g \in \mathbb{N}$. However, we are not aware of general computations of the Khovanov homology of cables. Nevertheless, we now proceed with the proof

of Theorem [1.4](#page-2-0) showing that the Khovanov homology of certain cables of positive knots behaves like the Khovanov homology of a positive knot.

In the proof of Theorem [1.4](#page-2-0) we use the skein long exact sequence in Khovanov homology [\[Vir04\]](#page-23-16). We adapt Lemma 2.2 and Theorem 3.1 from $[MPS^+18]$ $[MPS^+18]$ to the case when a 0-smoothing on a negative crossing transforms a diagram into a diagram of the unknot.

Lemma 5.4 Let v be a negative crossing in an oriented link diagram D , and consider the diagram D_0 (resp. D_1) obtained from D by smoothing the crossing v following a 0-marker (resp. 1-marker). Assume D_1 is oriented in the only way preserving the orientation of D , and orient D_0 in such a way that the orientation of the components not involved in v is preserved. Write w and w_0 for the writhes of D and D_0 , respectively. Then there is a long exact sequence in Khovanov homology of the form

$$
\cdots \to Kh^{i,j+1}(D_1) \to Kh^{i,j}(D) \to Kh^{\frac{w_0-w+1}{2}+i,\frac{3(w_0-w)+1}{2}+j}(D_0)
$$

$$
\to Kh^{i+1,j+1}(D_1) \to \cdots
$$

Proof. The proof is similar to that of $[MPS^+18$ $[MPS^+18$, Lemma 2.2.

Corollary 5.5 In the setting of Lemma [5.4,](#page-14-0) assume that D_0 is a diagram of the unknot, and set $u := \frac{w - w_0 - 1}{2}$. Then the Khovanov homologies of the links presented by D and D_1 are related as follows.

• If $(i, j) \notin \{(u, 3u + 2 \pm 1), (u + 1, 3u + 2 \pm 1)\}\$ then

$$
Kh^{i,j}(D) = Kh^{i,j+1}(D_1).
$$

• The other Khovanov homology groups fit into the following two exact sequences

$$
0 \to Kh^{u,3u+3\pm 1}(D_1) \to Kh^{u,3u+2\pm 1}(D) \to \mathbb{Z}
$$

$$
\to Kh^{u+1,3u+3\pm 1}(D_1) \to Kh^{u+1,3u+2\pm 1}(D) \to 0.
$$

Proof. The statement follows from Lemma [5.4](#page-14-0) and the fact that the Khovanov homology of the unknot is $\mathbb Z$ in gradings $(i, j) = (0, \pm 1)$ and zero for all other gradings. gradings. \square

We will deduce Theorem [1.4](#page-2-0) from a more general result about the family of twisted cables link, that we introduce now: Given $0 \le m \le p-1$, the *twisted cable* $K_{p,q;m}$ is the satellite link of K with pattern given by the braid word^{[4](#page-14-1)}

$$
w_{p,q} * [-1,-2,\ldots,-m],
$$

where $w_{p,q}$ the standard p-stranded braid word of $T(p,q)$. As for genuine cables, we get standard diagrams $D_{p,q;m}$ of twisted cables, as shown in Figure [8.](#page-15-0) It is clear that $K_{p,q;0} = K_{p,q} = K_{p,q+1;p-1}$. We write $K_{p,q;m}^{\circ}$ for any link obtained as the disjoint union of $K_{p,q;m}$ and a finite number (possibly zero) of unknots.

⁴Here we represent a standard Artin generator $\sigma_i^{\pm 1}$ by $\pm i$.

FIGURE 8. Part of a diagram $D_{p,q;m}$ representing a twisted cable link. The diagram also contains p-blackboard copies of D that are not depicted. In this particular example, $p = 7$, $n = 3$, $m = 4$, with $n = pw(D) - q.$

Theorem 5.6 Let K be a positive knot with writhe w. Then, for $q = pw-n \ge p \ge 2$, $n > 0$, and $1 \le m \le p-1$ we have:

$$
Kh^{i,*}(K_{p,q;m}) = 0 \text{ for every } i < 0, \text{ and}
$$

\n
$$
Kh^{0,j}(K_{p,q;m}) = \begin{cases} \mathbb{Z} & \text{if } j = 1 - p\chi(K) + q(p-1) - m \pm 1, \\ 0 & \text{else,} \end{cases}
$$

\n
$$
Kh^{1,j}(K_{p,q;m}) = \begin{cases} \mathbb{Z}^{p_1(K)} & \text{if } j = 2 - p\chi(K) + q(p-1) - m, \\ 0 & \text{else.} \end{cases}
$$

Proof. Let D be a positive diagram of K with writhe w. First, we recall that $D_{p,q}$ for $q \geq pw$ is a positive diagram. By analyzing its Seifert smoothing we see that every circle in s_0D yields p circles in $s_0D_{p,q}$. A simple analysis on the 0-chords shows that $G_0(D_{p,q})^{red}$ has the same cyclomatic number as $G_0(D)^{red}$ and thus $p_1(K_{p,q}) = p_1(K)$ for all $q \geq p$.

Now, we consider lexicographic order on the triples (p, n, m) and proceed by induction. We start with the base case $p = 2$. Given $n > 0$, consider the knot $K_{2,2w-n}$. The writhe of its standard diagram $D_{2,2w-n}$ is $4w-n$, and performing a 0-smoothing at any of its negative crossings leads to a diagram of the unknot with writhe $w_0 = n - 1$, while performing a 1-smoothing at the same crossing leads to the standard diagram of $K_{2,2w-n+1}$. Thus we can apply Corollary [5.5](#page-14-2) to deduce that

$$
Kh^{i,j}(K_{2,2w-n}) = Kh^{i,j+1}(K_{2,2w-n+1})
$$

whenever i is not equal to $2w - n = q$ or $2w - n + 1 = q + 1$. Inductively this yields the theorem for $p = 2$, since $K_{2,2w}$ is positive with $p_1(K_{2,2w}) = p_1(K)$ and thus the statement follows from Theorem [1.1](#page-0-1) and [\[Kho03,](#page-23-4) Proposition 8].

For the induction step, we assume that the statement is true for all (p', n', m') (p, n, m) with $p \geq 3$. Consider the diagram $D_{p, pw-n,m}$ with writhe $\overline{w} = p^2w - n(p - \overline{w})$ 1)−m. Let v be the highest negative crossing in the last partial twist (see Figure [8\)](#page-15-0). Then the 1-smoothing D_1 at v is the standard diagram of $K_{p,pw-n;m-1}$.

Claim: The 0-smoothing of $D_{p,pw-n;m}$ at v leads to a diagram D_0 with writhe w₀ satisfying the following conditions (that we will prove in Lemmas [5.7](#page-16-0) and [5.8](#page-18-0) below):

- (1) $w_0 \overline{w} + 3 < 0$.
- (2) D_0 is either a (non-positive) diagram of a positive link or a (non-standard) diagram of $K^{\circ}_{p',p'w-n',m'}$ with $(p',n',m') < (p,n,m)$.

We can then apply Lemma [5.4](#page-14-0) to see that the Khovanov homology groups of these knots fit into the following exact sequence

 $Kh^{i'-1,j'-1}(D_0) \to Kh^{i,j+1}(K_{p,pw-n;m-1}) \to Kh^{i,j}(K_{p,pw-n;m}) \to Kh^{i',j'}(D_0)$ with $i' = (w_0 - \overline{w} + 1)/2 + i$. The first condition in the claim implies that $i' < 0$ if $i \leq 1$. If D_0 represents a positive link, then we get from [\[Kho03,](#page-23-4) Proposition 8] that $Kh^{i',j'}(D_0)$ is vanishing. Otherwise, the second condition in the claim together with Lemma [3.3](#page-7-1) allows us to apply the inductive hypothesis to deduce that $Kh^{i',j'}(D_0)$ is vanishing. Analogously, we conclude that $Kh^{i'-1,j'-1}(D_0)$ is trivial. Thus we conclude that $Kh^{i,j}(K_{p,pw-n;m}) = Kh^{i,j+1}(K_{p,pw-n;m-1})$ and the theorem follows by induction since $K_{p,pw}$ is positive.

Lemma 5.7 If we orient the diagram D_0 in the proof of Theorem [5.6](#page-15-1) in such a way that the orientations of the components not involved in v are preserved, then $w(D_0) - w(D_{p, pw-n; m}) + 3 < 0.$

Proof. Figure [9](#page-16-1) shows the 0-smoothing D_0 of a standard diagram $D_{p,pw-n;m}$ of a twisted cable. We add the dotted line $\mathcal L$ to simplify diagrammatic descriptions.

FIGURE 9. Part of diagram D_0 together with the dotted grey line $\mathcal L$ illustrating the proofs of Lemmas [5.7](#page-16-0) and [5.8.](#page-18-0)

Write $n = pk + t$, with $0 \le t < p$. We set:

- x: number of strands crossing $\mathcal L$ oriented from right to left.
- x_m : number of strands crossing $\mathcal L$ oriented from right to left in position i, with $i \in \{2, \ldots m\}$. The labeling of position starts from below, as in Figure [9.](#page-16-1)
- t_d : number of strands crossing $\mathcal L$ at position i oriented from right to left, with $p - t + 1 \leq i \leq p$.
- u: it takes the value 1 (resp. -1) if the situation is as in Figure [10\(](#page-17-0)a) $(resp. Figure 10(b)).$ $(resp. Figure 10(b)).$ $(resp. Figure 10(b)).$

We compute the writhe w_0 of D_0 in terms of the above parameters. To do so, we split its crossings into four classes $(I - IV)$ and compute their writhes $(w_I - w_{IV})$ separately, so that $w_0 = w_1 + w_{II} + w_{III} + w_{IV}$.

- I: contains those crossings coming from the positive diagram D (i.e. not created by any of the twists). Each crossing in D contributes to the writhe of w_I with $((p-x) - x)^2$. Hence, the total contribution is

$$
w_{\rm I} = w(p-2x)^2.
$$

The next two classes contain those crossings created by the $n = pk + t$ twists.

FIGURE 10. The two possible situations determining the value of the paramenter u: if D_0 is oriented as in (a) then $u = 1$; in case (b) then $u = -1$.

- II: contains those crossings involved in the pk twists which are farther from \mathcal{L} . These twists can be subdivided into k blocks, each of them containing p consecutive twists (i.e. each block contains exactly one full twist). Fix a strand and compute the sign of the crossings created when it wraps around the other strands: if it crosses \mathcal{L} from right to left (resp. from left to right), then it creates $(p-x)$ positive and $(x-1)$ negative crossings (resp. x positive and $(p - x - 1)$ negative crossings). Hence, we get

$$
w_{\text{II}} = k \Big(x \big((p-x) - (x-1) \big) + (p-x) \big(x - (p-x-1) \big) \Big) = (n-t) \big(1 - p + 4x - 4x^2/p \big).
$$

- III: contains those crossings involved in the t twists which are closest to the dotted line. A similar reasoning as before leads to

$$
w_{\text{III}} = (t_d((p-x)-(x-1)) + (t-t_d)(x-(p-x-1))) = 2t_d(p-2x) + t(2x-p+1).
$$

- IV: contains those crossings created by the partial twist related to m. There are two possible situations depicted in Figure [10,](#page-17-0) from which we read-off that

$$
w_{\text{IV}} = u(2x_m - m + 1).
$$

Therefore, we get:

 $\beta := w_0 - \overline{w} + 3$

 $= 4x(k - w)(p - x) + 2xt + 2t_d(p - 2x) + u(2x_m - m + 1) + m + 3.$

Next, we bound some of the terms in β . Since $0 \le t \le p-1$ and $q \ge p$, we get:

(1) $pk + t = n = pw - q \leq p(w - 1)$, which implies $k \leq w - 1$,

with strict inequality if $t \neq 0$.

From $0 \leq x_m \leq m-1$ we deduce that

$$
u(2x_m - m + 1) \le m - 1.
$$

We consider first the case when $(p - 2x) > 0$ (we defer the other case to the end of the proof). A combinatorial argument shows that $t_d \leq x - 1$ and therefore

$$
2t_d(p - 2x) \le 2(x - 1)(p - 2x).
$$

Substitution of the previous bounds and $m \leq p-1$ we bound β as:

$$
\beta \le 2(xt - xp + 2x + m - p + 1) \le 2x(t - p + 2).
$$

Now, we consider different situations depending on the value of t:

• If $0 < t < p-1$, then [\(1\)](#page-17-1) is a strict inequality, and therefore

$$
\beta < 2x(t - p + 2) < 0.
$$

- If $t = 0$, then $\beta \leq 2x(2 p)$, which is strictly negative, since $p \geq 3$.
- If $t = p 1$, then [\(1\)](#page-17-1) becomes $k \leq w 2$ and analogous reasoning leads to

$$
\beta \le 2x(2(x - p) + 1) < 0,
$$

where the last inequality comes from $p > x > 0$.

The case when $(p - 2x) \le 0$ leads to $2t_d(p - 2x) \le 0$, and the proof holds in a nilar way. similar way.

Recall, that the diagram D_0 in the proof of Theorem [5.6](#page-15-1) is obtained from a standard diagram $D_{p,pw-n;m}$ of a twisted cable by smoothing the negative crossing v.

Lemma 5.8 There exists exactly one component C in D_0 that is involved in the smoothing at v. If we preserve the orientation of all other components then there exists a choice of orientation on C such that D_0 is either a (non-positive) diagram of a positive link or a (non-standard) diagram of $K^{\circ}_{p',p'w-n';m'}$ with (p',n',m') (p, n, m) .

Proof. There are only two arcs with vertical tangencies in Figure [9,](#page-16-1) and they are the ones obtained when smoothing v b so they belong to the same component C . We will show that we can orient that component such that D_0 satisfies the statement of the theorem.

First, we study the case when $m \neq 1, p - 1$. Starting from the diagram D_0 depicted in Figure [9,](#page-16-1) push the marked arc to the right through the whole diagram till the white and black dots intersects $\mathcal L$ at positions a and b. It is clear that $b = a + 1$. We consider positions always modulo p. Depending on the value of a there are 6 possible situations depicted in Figure [11.](#page-18-1)

FIGURE 11. The six possible cases illustrating proof of Lemma [5.8](#page-18-0) depending on the values of a (assuming original orientation of $D_{p,q;m}$ is from left to right).

In the cases β and ϵ we continue pushing the arc to the right through the diagram until we reach one of the other cases. We readily see that (after choosing the orientation on C shown in Figure [11\)](#page-18-1) the diagrams in cases C and D represent standard diagrams of twisted cables of K with $p' < p$.

Next, we consider case A . Let r denote the number of negative twists in the associated diagram D_A . If $r = 0$, then D_A oriented as in Figure [11](#page-18-1) is positive. Otherwise, we perform the isotopy shown in Figure $12(a)$ –(f) to see that it represents a twisted cable. Observe that it may be needed to push the red box several times through the whole diagram before reaching Figure [12\(](#page-19-0)d).

FIGURE 12. Isotopies illustrating proof of Lemma [5.8](#page-18-0) in the case A . From (b) to (d) the red box is pushed to the left. From (e) the blue box is pushed to the right through the diagram to obtain (f).

In case F, if $r = 0$ then we rotate the diagram by π (after choosing the orientation on C shown in Figure [11\)](#page-18-1) to get the standard diagram of a twisted cable of K , with $p' < p$. The case $r > 0$ works similarly as case A by pushing the red box to the right.

The cases when $m = 1$ or $m = p - 1$ can be handled with the same method.
Serve that in these cases an additional unknotted split component may arise. Observe that in these cases an additional unknotted split component may arise.

Proof of Theorem [1.4.](#page-2-0) Let D be a positive diagram of K with writhe $w > 0$. The-orem [5.1](#page-13-1) implies that $K_{p,q}$ is fibered and if $q \geq p w$ it is also positive, and thus the statement follows from Theorem [1.1](#page-0-1) by observing that $p_1(K_{p,q}) = p_1(K)$.

For $q < pw$ the statement follows from Theorem [5.6](#page-15-1) by using Schubert's cabling formula for the genus [\[Sch53\]](#page-23-22) to compute $-\chi(K_{p,q}) = -p\chi(K) + q(p-1)$. \Box

Remark 5.9 In the proof of Theorem [1.4](#page-2-0) we require the (p, q) -cables to satisfy $q \ge$ $p \geq 2$ so we can achieve the inequality for β in Lemma [5.7.](#page-16-0) However, computational experiments show that for $p, w \leq 30$ and $q \geq 3$ it holds that $\beta < 0$ and therefore Theorem [1.4](#page-2-0) holds true in that cases as well (see [\[KMMS\]](#page-23-13) for the computations). This suggests that Theorem [1.4](#page-2-0) is true for all $q \geq 3$.

Proof of Corollary [1.5.](#page-2-3) If K is fibered then $K_{p,q}$ is also fibered by Theorem [5.1\(](#page-13-1)1) and thus Theorems [1.4](#page-2-0) and [2.3](#page-5-2) imply the result. Conversely, if $Kh^1(K_{p,q}) = 0$ then Theorem [1.4](#page-2-0) implies that $p_1(K) = 0$ and thus Theorem [2.3](#page-5-2) implies that K is fibered. \square

In further search for a counterexample to Conjecture [1.7,](#page-2-2) it is natural to consider cables of more general L-space knots. Possible candidates to be counterexamples are iterated cables of L-space knots.

Theorem 5.10 Let K be the $(2, 3)$ -cable of $T(2, 3)$. Then, for every $q \geq 2$, the Khovanov homology groups (with \mathbb{Z}_2 -coefficients) of $K_{2,q}$ fulfill

$$
Kh^{i,j}(K_{2,q}) = \begin{cases} \mathbb{Z}_2 & \text{if } (i,j) = (0, 1 - \chi(K_{2,q}) \pm 1), \\ 0 & \text{for all other groups with } i \leq 1. \end{cases}
$$

In particular, any 2-stranded cable of $T(2,3)_{2,3}$ that is an L-space knot has Khovanov homology with \mathbb{Z}_2 -coefficients of the above form.

To prove Theorem [5.10](#page-20-1) we need the following result for 2-cables.

Theorem 5.11 Let K be a link. Then for every $q \geq 2$ the first Khovanov homologies (with arbitrary coefficients) of their 2-cables are related by

$$
Kh^{i,j}(K_{2,q}) = Kh^{i,j+2-q}(K_{2,2}) \text{ for all } i \le 1.
$$

Proof. Let D be a knot diagram of K with writhe w. The diagram $D_{2,q}$ has writhe $4w(D) + n$, with $n = 2w - q$ its number of positive half-twists. If $n < 0$, let v be a crossing involved in a negative half-twist (if $n \geq 0$, just introduce a positive and a negative crossing close to the last positive twist). Applying a 0-smoothing to v produces a diagram of the unknot with writhe $w_0 = -n - 1$, while a 1-smoothing of v gives rise to $D_{2,q+1}$. Thus Corollary [5.5](#page-14-2) (which holds for arbitrary coefficient groups) applies and we get

$$
Kh^{i,j}(K_{2,q}) = Kh^{i,j+1}(K_{2,q+1})
$$

for all $i \neq q, q + 1$.

Proof of Theorem [5.10.](#page-20-1) Since K is not positive (see Remark [5.2\)](#page-13-0) Theorem [1.4](#page-2-0) does not apply here. In [\[KMMS\]](#page-23-13) we also show that $K_{2,q}$ is not braid positive for small values of q (and conjecturally it is not positive for all $q > 0$), so we cannot apply Theorem [1.1](#page-0-1) here.

Nevertheless, by Theorem [5.11](#page-20-2) it is enough to compute the first Khovanov homology of one 2-stranded cable of K . However, the computations over the integers or over Q were not terminating on our machines. But we could compute the Khovanov homology with \mathbb{Z}_2 -coefficients using [\[BNMea\]](#page-22-16) from which we read off that its Khovanov homology has the claimed form (the data can be accessed at [\[KMMS\]](#page-23-13)).

Since $T(2,3)_{2,3}$ has Seifert genus 3, Theorem [5.1](#page-13-1) (3) implies that if $K_{2,q}$ is an L-space knot then $q > 10$ and thus the second claim holds.

5.2. Hyperbolic knots. We found hyperbolic knots that are fibered and strongly quasipositive whose first Khovanov homology is non-vanishing. The idea to construct these examples was to start with a quasipositive Seifert surface of a fibered strongly quasipositive knot and plumb a Hopf band to it (recall that Hopf plumbing preserves fiberedness and strongly quasipositivity), so that the new resulting knot is hyperbolic.

Example 5.12 Let K be the hyperbolic knot given by the closure of the braid shown in Figure [13.](#page-21-0) From the knot Floer chain complex of K (computed via Szabo's program $[Sz]$) we deduce that K is fibered and its 3-genus agrees with $\tau(K)$. Therefore K is strongly quasipositive by [\[Hed10\]](#page-22-2). However, $Kh^{1,*}(K) \neq 0$, as shown in Table [14.](#page-21-1)

FIGURE 13. The braid from Example [5.12,](#page-20-3) picture created via KLO [\[Swe\]](#page-23-24).

ι \jmath	θ	$\mathbf{1}$	$\overline{2}$	3	$\overline{4}$	5	6	$\overline{7}$	8	9	10	11	12	13	14
31															$\mathbb Z$
29														$\mathbb Z$	\mathbb{Z}_2
27													\mathbb{Z}^2	$\mathbb{Z} \oplus \mathbb{Z}_2$	
$25\,$												\mathbb{Z}^2	$\mathbb{Z}\oplus(\mathbb{Z}_2)^2$		
23											$\mathbb{Z} \oplus \mathbb{Z}_2$	$\mathbb{Z}^2 \oplus (\mathbb{Z}_2)^2$			
21										$\mathbb{Z}^2\oplus\mathbb{Z}_2$	$\mathbb{Z}^2\oplus(\mathbb{Z}_2)^2$				
19								\mathbb{Z}^2	\mathbb{Z}	$\mathbb{Z}\oplus(\mathbb{Z}_2)^3$					
17						$\mathbb Z$	z	$(\mathbb{Z}_2)^3$	$\mathbb{Z}^2\oplus\mathbb{Z}_2$						
$15\,$						\mathbb{Z}^2	$\overline{\mathbb{Z}^2\oplus(\mathbb{Z}_2)^2}$	$\mathbb{Z} \oplus \mathbb{Z}_2$							
$13\,$				\mathbb{Z}	\mathbb{Z}^3	$\mathbb{Z} \oplus \mathbb{Z}_2$	\mathbb{Z}_2								
11				$\mathbb{Z} \oplus \mathbb{Z}_2$	$\mathbb{Z}^2\oplus(\mathbb{Z}_2)^2$										
9			\mathbb{Z}^2	$\overline{\mathbb{Z}^2\oplus\mathbb{Z}_2}$											
$\overline{7}$	$\mathbb Z$		$\mathbb{Z} \oplus \mathbb{Z}_2$												
$5\,$	$\mathbb Z$	$\mathbb Z$													

Figure 14. Khovanov homology of the hyperbolic knot given as the closure of the braid in Figure [13.](#page-21-0)

For L-space knots, the situation is more complicated. Currently, all hyperbolic L-space knots that are not known to be braid positive are contained in an infinite family $\{K_n\}_{n>0}$ from [\[BK23\]](#page-22-4), where the knot K_n is given as the closure of the 4-braid

$$
\beta_n = [(2, 1, 3, 2)^{2n+1}, -1, 2, 1, 1, 2].
$$

In $[BK23]$ it is shown that all K_n are hyperbolic L-space knots. Since the normalized HOMFLYPT polynomial of K_1 is not positive, it follows that K_1 is not braid positive. When $n > 1$ the braid-positivity status of K_n is currently unclear.

Theorem 5.13 For every $n \geq 1$, the Khovanov homology of K_n satisfies

$$
Kh^{i,j}(K_n) = \begin{cases} \mathbb{Z} & \text{if } (i,j) = (0, 4+8n \pm 1), \\ 0 & \text{for all other groups with } i \le 1. \end{cases}
$$

The Khovanov homology table of K_1 is shown in Figure [15.](#page-22-17)

Proof. Let D_n be the diagram obtained as the closure of β_n , whose writhe is $w_n =$ $8n+7$. The 0-smoothing of the unique negative crossing v in D_n produces a diagram of the unknot with writhe $w_0 = 0$, while the 1-smoothing of v produces a braid positive diagram $(D_n)_1$, and thus its first Khovanov homology vanishes. Applying Corollary [5.5,](#page-14-2) where $u = \frac{w_n - w_0 - 1}{2} = 4n + 3 \ge 7$, we get

$$
Kh^{i,j}(K_n) = Kh^{i,j+1}((D_n)_1) \text{ for all } i \leq 6.
$$

The fact that $\chi((D_n)_1) = -4 - 8n$ and the Khovanov homology obstructions for braid positive links imply the result. braid positive links imply the result.

REFERENCES

- [Kan22] S. Kang, Link homology theories and ribbon concordances, Quantum Topol. 13 (2022), 183–205. MR [4404800](http://www.ams.org/mathscinet-getitem?mr=4404800)
- [Kho00] M. Khovanov, A categorification of the Jones polynomial, Duke Math. J. 101 (2000), 359–426. MR [1740682](http://www.ams.org/mathscinet-getitem?mr=1740682)
- [Kho03] , Patterns in knot cohomology I, Experiment. Math. 12 (2003), 365-374. MR [2034399](http://www.ams.org/mathscinet-getitem?mr=2034399)
- [KMMS] M. Kegel, N. Manikandan, L. Mousseau, and M. Silvero, Code and data to accompany this paper, Available at, [https://www.mathematik.hu-berlin.de/~kegemarc/](https://www.mathematik.hu-berlin.de/~kegemarc/KHposfib/KHposfib.html) [KHposfib/KHposfib.html](https://www.mathematik.hu-berlin.de/~kegemarc/KHposfib/KHposfib.html).
- [KR08] M. Khovanov and L. Rozansky, Matrix factorizations and link homology, Fund. Math. 199 (2008), 1–91. MR [2391017](http://www.ams.org/mathscinet-getitem?mr=2391017)
- [Lee05] E. S. Lee, An endomorphism of the Khovanov invariant, Adv. Math. 197 (2005), 554–586. MR [2173845](http://www.ams.org/mathscinet-getitem?mr=2173845)
- [LM23] C. Livingston and A. Moore, KnotInfo: Table of knot invariants, 2023, [http://www.](http://www.indiana.edu/~knotinfo) [indiana.edu/~knotinfo](http://www.indiana.edu/~knotinfo).
- [LZ19] A. S. Levine and I. Zemke, Khovanov homology and ribbon concordances, Bull. Lond. Math. Soc. 51 (2019), 1099–1103. MR [4041014](http://www.ams.org/mathscinet-getitem?mr=4041014)
- [Mos71] L. Moser, Elementary surgery along a torus knot, Pacific J. Math. 38 (1971), 737–745. MR [383406](http://www.ams.org/mathscinet-getitem?mr=383406)
- [MPS+18] S. Mukherjee, J. H. Przytycki, M. Silvero, X. Wang, and S. Y. Yang, Search for torsion in Khovanov homology, Exp. Math. 27 (2018), 488–497. MR [3894728](http://www.ams.org/mathscinet-getitem?mr=3894728)
- [Ni07] Y. Ni, Knot Floer homology detects fibred knots, Invent. Math. 170 (2007), 577–608. MR [2357503](http://www.ams.org/mathscinet-getitem?mr=2357503)
- [ORS13] P. S. Ozsváth, J. Rasmussen, and Z. Szabó, Odd Khovanov homology, Algebr. Geom. Topol. 13 (2013), 1465–1488. MR [3071132](http://www.ams.org/mathscinet-getitem?mr=3071132)
- [OS05] P. Ozsváth and Z. Szabó, On knot Floer homology and lens space surgeries, Topology 44 (2005), 1281–1300. MR [2168576](http://www.ams.org/mathscinet-getitem?mr=2168576)
- [PS14] J. H. Przytycki and R. Sazdanović, Torsion in Khovanov homology of semi-adequate links, Fund. Math. 225 (2014), 277–304. MR [3205574](http://www.ams.org/mathscinet-getitem?mr=3205574)
- [PS20] J. H. Przytycki and M. Silvero, Geometric realization of the almost-extreme Khovanov homology of semiadequate links, Geom. Dedicata 204 (2020), 387-401. MR [4056710](http://www.ams.org/mathscinet-getitem?mr=4056710)
- [Sch] D. Schütz, Knotjob, Available at [https://www.maths.dur.ac.uk/users/dirk.](https://www.maths.dur.ac.uk/users/dirk.schuetz/knotjob.html) [schuetz/knotjob.html](https://www.maths.dur.ac.uk/users/dirk.schuetz/knotjob.html).
- [Sch53] H. Schubert, Knoten und Vollringe, Acta Math. 90 (1953), 131–286. MR [72482](http://www.ams.org/mathscinet-getitem?mr=72482)
- [SS22] R. Sazdanović and D. Scofield, *Extremal Khovanov homology and the girth of a knot*, J. Knot Theory Ramifications 31 (2022), Paper No. 2250083, 26. MR [4520782](http://www.ams.org/mathscinet-getitem?mr=4520782)
- [Sto05] A. Stoimenow, On polynomials and surfaces of variously positive links, J. Eur. Math. Soc. (JEMS) 7 (2005), 477–509. MR [2159224](http://www.ams.org/mathscinet-getitem?mr=2159224)
- [Sto10] M. Stošić, On conjectures about positive braid knots and almost alternating torus knots, J. Knot Theory Ramifications 19 (2010), 1471–1486. MR [2746199](http://www.ams.org/mathscinet-getitem?mr=2746199)
- [Swe] F. Swenton, KLO (Knot-Like Objects), <http://KLO-Software.net>.
- [Sza] Z. Szabó, Knot floer homology calculator, Available at [https://web.math.princeton.](https://web.math.princeton.edu/~szabo/HFKcalc.html) [edu/~szabo/HFKcalc.html](https://web.math.princeton.edu/~szabo/HFKcalc.html).
- [Thu86] W. P. Thurston, Hyperbolic structures on 3-manifolds. I. Deformation of acylindrical manifolds, Ann. of Math. (2) 124 (1986), 203–246. MR [855294](http://www.ams.org/mathscinet-getitem?mr=855294)
- [Tra04] P. Traczyk, A combinatorial formula for the signature of alternating diagrams, Fund. Math. 184 (2004), 311-316. MR [2128055](http://www.ams.org/mathscinet-getitem?mr=2128055)
- [Vir04] O. Viro, Khovanov homology, its definitions and ramifications, Fund. Math. 184 (2004), 317–342. MR [2128056](http://www.ams.org/mathscinet-getitem?mr=2128056)

HUMBOLDT UNIVERSITÄT ZU BERLIN, GERMANY Email address: kegemarc@hu-berlin.de, kegelmarc87@gmail.com

HUMBOLDT UNIVERSITÄT ZU BERLIN, GERMANY $\boldsymbol{Email}_address:$ naageswaran.manikandan@hu-berlin.de

HUMBOLDT UNIVERSITÄT ZU BERLIN, GERMANY Email address: leo.mousseau@t-online.de

Universidad de Sevilla, Spain Email address: marithania@us.es